

Global variability of high nutrient low chlorophyll regions using neural networks and wavelet coherence analysis.

Gotzon Basterretxea¹, Joan S. Font-Muñoz¹, Ismael Hernández-Carrasco², Sergio A. Sañudo-Wilhelmy³

¹ Department of Marine Ecology, Instituto Mediterráneo de Estudios Avanzados, IMEDEA (UIB-CSIC), Miquel Marqués 21, 07190 Esporles, Illes Balears, Spain.

² Department of Oceanography and Global Change, Instituto Mediterráneo de Estudios Avanzados, IMEDEA (UIB-CSIC), Miquel Marqués 21, 07190 Esporles, Illes Balears, Spain.

³ Department of Biological Sciences and Department of Earth Sciences, University of Southern California, Marine Biology and Biological Oceanography, Los Angeles, California 90089-0371, United States.

Correspondence to: Gotzon Basterretxea (gotzon@imedea.uib-csic.es)

Abstract. We examine 20 years of monthly global ocean color data and modeling outputs of nutrients using self-organizing map (SOM) analysis to identify characteristic spatial and temporal patterns of High Nutrient Low Chlorophyll (HNLC) regions and their association with different climate modes. The global nitrate to chlorophyll ratio threshold of $\text{NO}_3:\text{Chl} > 17$ ($\text{mmol NO}_3/\text{mg Chl}$) is estimated to be a good indicator of the distribution limit of this unproductive biome that, on average, covers $92 \times 10^6 \text{ km}^2$ (~25% of the ocean). The trends in satellite-derived surface chlorophyll (0.6 ± 0.4 to $2 \pm 0.4\% \text{ yr}^{-1}$) suggest that HNLC regions in polar and subpolar areas have experienced an increase in phytoplankton biomass over the last decades but much of this variation, particularly in the Southern Ocean, is produced by a climate-driven transition in 2009-2010. Indeed, since 2010, the extent of the HNLC zones has decreased at the poles (up to 8%) and slightly increased at the equator ($< 0.5\%$). Our study finds that chlorophyll variations in HNLC regions ~~respond to major climate variability signals~~ Chlorophyll variations in HNLC regions respond to major climate variability signals such as El Niño Southern Oscillation (ENSO) and Meridional Overturning Circulation (MOC) at both short (2-4 years) and long (decadal) timescales. These results suggest global coupling in the functioning of distant biogeochemical regions.

1 Introduction

High nutrient low chlorophyll (HNLC) areas are ocean regions where primary production should be potentially high but phytoplankton biomass remains relatively low and constant despite the perennial nutrient availability for growth (Martin and Fitzwater, 1988; Chisholm and Morel, 1991). They are interesting regions because they challenge the accepted paradigm of a positive relation between macronutrient concentrations and phytoplankton biomass in open waters but, most importantly, because they represent an important fraction of the global ocean carbon budgets/inventories and, therefore, their extent influences the potential withdrawal of atmospheric CO_2 to the deep ocean (Martin et al., 1990;

~~de~~ Baar et al., 1995; Boyd et al., 2005). It is estimated that HNLC biomes roughly cover between 20 and 30% of the world's oceans (~~Pitchford and Brintley, 1999; Tyrrel et al., 2005~~) comprising three major ocean areas; the Subarctic North Pacific (SNP), the Eastern Equatorial Pacific (EEP), and most of the Southern Ocean (SO) (Martin, 1990; Coale et al., 1996; Parekh et al., 2005).

Because nitrogen is the mineral nutrient needed in greatest abundance by phytoplankton and owing to its generalized depletion in surface waters over much of the oceans, it is considered a key limiting nutrient for ocean production. In HNLC regions, where nitrogen is in excess, other non-exclusive factors such as rapid top-down control by zooplankton grazing, low irradiance, limitations by silicic acid availability, and/or iron (Fe) limitation, have been hypothesized to explain the persistently low chlorophyll (Chl). While these factors may contribute in different degrees to the observed low Chl and determine the phytoplankton dynamics in HNLC regions (see Chavez et al., 1991; Cullen, 1995; Coale et al., 1996; Dugdale and Wilkerson, 1998; Landry et al., 2011), it is generally acknowledged that Fe availability is central to the productivity of HNLC regions (Boyd et al., 2007). All HNLC regions share a chronic Fe-depletion in surface waters and experimental results show highly positive productivity responses to Fe addition (Martin et al., 1994; Boyd et al., 2000, 2004; Tsuda et al., 2003; ~~Coale et al., 2004~~). Indeed, ~~iron is required in largest amounts than any of~~ Fe requirements are the largest among the trace metals for several metabolic processes, and not surprisingly, it has been considered the ultimate limiting nutrient (Moore and Doney, 2007). This has led to propose a conceptual model of phytoplankton nutrient limitation in the modern ocean based on two functioning regimes: one in which the supply of nutrients is relatively slow and nitrogen availability limits productivity, and a complementary regime, with enhanced nutrient macronutrient supply, where Fe ~~often~~ limits productivity (Moore et al., 2013).

Iron limitation influences the uptake of nitrogen thereby explaining the unused nitrate concentrations in HNLC regions. Indeed, it has been proposed that a delicate balance between nitrogen and Fe availability modulates phytoplankton growth and that co-limitation is rather ubiquitous in the sea (Bryant, 2003; Browning et al., 2017). Other elements and compounds such as B-vitamins, which are also scarce in Fe-limited areas, can also be co-limiting factors for phytoplankton growth in these regions (e.g. Koch et al., 2011; Bertrand et al., 2012). For example, it has been experimentally shown that the addition of Fe and B12 to Antarctic phytoplankton assemblages can synergistically increase phytoplankton growth (Bertrand et al., 2011; Cohen et al., 2017).

Despite their relevance for global ocean productivity and carbon fluxes, HNLC regions remain loosely defined and knowledge of their temporal and spatial variability and trends is limited. Moreover, their response in a global warming scenario is uncertain. Only general aspects such as expected shifts in phytoplankton community composition or changes in Fe-cycling rates have been addressed to date (Fu et al., 2016; Lauderdale et al., 2020). The original description of HNLC systems by Minas et al. (1986)

referred to a slowly growing phytoplankton standing stock despite the presence of high nutrient concentrations. However, there are no rigid criteria accurately defining the functioning of these ecosystems. Several ecosystem characteristics such as species composition, biological ecosystem structure, carbon utilization pathways, and response to climate change also differ between the HNLC and other ecosystems, reflecting differences in the limiting factor (~~e.g.~~ Falkowski et al., 1998; Ono et al., 2008).

Of particular interest are the aspects related to the reduced variability and high permanence (i.e. temporal persistence) typically characterizing large HNLC regions. These features are distinctive from those of highly variable systems, which may temporarily present HNLC conditions. For example, some light-limited regions in high latitudes may present low productivity and enhanced nutrients during winter but it responds to a transient situation that does not correspond to the generally accepted HNLC paradigm. Similarly, high nutrients and low Chl have been observed at the end of the spring bloom in some productive systems (Nielsdóttir et al., 2009; Birchill et al., 2019) and some areas located in coastal upwelling regions (Hutchins et al., 1998, 2002; Firme et al., 2003; Eldridge et al., 2004). While complying with the necessary conditions of high nutrient and low Chl, it is uncertain whether these ephemeral systems share structural and functioning similitudes functional similarities with the large HNLC regions.

At a time when understanding biogeochemical responses to large-scale forcings, including climate change, has become a scientific priority, it seems appropriate to revisit some concepts of the functioning of HNLC regions. Their extent and variability are indicative of the dynamic changes in the bidirectional interrelationships of phytoplankton with the environment and with other organisms at large scales. Most of the information on the long-term variations of HNLC regions is depicted from global studies suggesting that their productivity is declining varying as a consequence of global warming and that they experience prominent interannual to decadal fluctuations superimposed on these long-term trends (i.e. Boyce et al., ~~2010~~2014; Martinez et al., 2020). Available evidence suggests that some HNLC regions may be decreasing in size as a result of increased ocean stratification (Ono et al., 2008). More recently, Yasunaka et al., (2016), determined that surface trends of phosphate and silicate in the North Pacific are associated with the shoaling of the mixed layer, reporting that surface nutrient concentration was correlated with the North Pacific Gyre Oscillation (NPGO). ~~Some studies have shown that oligotrophic areas in the northern hemisphere are expanding~~ between 0.8 and 4% per year, with a faster increase in winter months ((e.g. Polovina et al., 2008). ~~However; however,~~ with some exceptions (e.g.i.e. Radenac et al., 2012; Yasunaka et al., 2014), specific long-term studies on HNLC regions are scarce and knowledge ~~on~~of their variability ~~is on~~ the global ocean scale and their responseresponses to climate change remain uncertain.

The objective of the present study is to provide a quantitative assessment of the large-scale patterns of variability of the three major HNLC regions (SNP, EEP, and SO) and their relationship with the main

modes of climate variability. Systematically determining the boundaries of these HNLC regions has remained elusive since it requires coherent information on nutrients and Chl fields. The present study is based on the analysis of 20-year time series of monthly global ocean color data and nutrient concentrations from a biogeochemical model using machine learning techniques and wavelets analysis. First, based on the statistical analysis of global NO₃:Chl ratios, we determine a robust quantitative criterion to objectively define HNLC regions. Then we characterize the temporal variability patterns of HNLC regions based on their NO₃ and Chl concentrations by using the Self-Organizing Map (SOM) technique. We use the herein-established statistical criterion to assess the spatial variations of HNLC regions over the study period unveiled from the SOM analysis in the spatial domain of NO₃:Chl ratios. Finally, through a combined SOM-wavelet coherence analysis (WCA), we quantify the spectral power and the dynamic relationship between the observed Chl variability and two main global-scale forcings; El Niño Southern Oscillation (ENSO); and Meridional Overturning Circulation (MOC). We show that the combination of WCA with SOM-derived characteristic time-series is an especially suitable tool for the analysis of driver-response relationships in the ocean.

2 Materials and Methods

2.1 Ocean color data

We employ 20 years of monthly global composites of satellite Chl Level-3 products, derived from merging SeaWiFS, MERIS, MODIS AQUA, and VIIRS sensors using a GSM algorithm (Maritorena and Siegel, 2005), obtained from GlobColour data set (www.globcolour.info). The chlorophyll product is spatially gridded, and the weighted average of the different merged Level-2 products is then calculated. The composite consists of a rectangular regular map product in degrees with a spatial resolution of 0.25° (i.e. around 28 km at the equator ~~that varies with the latitude~~) and covers the period from January 1998 to December 2017. We excluded results in the Arctic Ocean and the coastal Southern Ocean due to the interference of ice cover and prolonged gaps in the data. A total of 654395 pixels were considered in the analysis. We are aware that the consistency of merged multi-mission ocean color satellite series may suffer from some limitations influencing long trend analysis (MélinaMélina et al., 2017). However, no significant increase or decrease is observed in the first-order trends of GlobColour data in more recent studies (e.g. Moradi, 2021). Therefore, while recognizing that some differences in regional and seasonal biases may occur in unified data products and, acknowledging that discontinuities and trends of the median with time should be interpreted carefully according to the sensors used (Garnesson et al., 2019), merged Chl can be generally considered a good indicator of the magnitude of the overall phytoplankton trends.

2.2 Nitrate data

Since nutrient observations are still too scarce to allow obtaining time-resolved global-scale fields, we used global NO₃ obtained from the biogeochemical hindcast model provided by Mercator-ocean (<http://marine.copernicus.eu>, see Fig. S1). ~~Data on climate indices were obtained from available databases. Bi-monthly Multivariate El Niño Southern Oscillation Index (MEI.v2), hereafter ENSO index, was obtained from the National Oceanic and Atmospheric Administration National Center for Environmental Prediction website (<https://www.esrl.noaa.gov/psd/enso/mei/>). MOC data (Smeed et al., 2019; Moat et al., 2022) for the period (2004-2018) was obtained from the RAPID-WATCH MOC monitoring project (www.rapid.ac.uk/rapidmoc). It consists of monthly mean fields of several biogeochemical variables at 0.25° horizontal resolution over the global ocean~~ obtained using the PISCES model (Aumont et al., 2015). The model is forced by daily mean fields of ocean, sea ice, and atmospheric conditions. Ocean and sea ice forcings are obtained from the numerical simulation FREEGLORYS2V4 produced at Mercator-Ocean and the source of atmospheric forcings is the ERA-Interim reanalysis produced at ECMWF. Initial conditions are set from the World Ocean Atlas 2013 climatology. A complete model description can be found at (<http://cmems-resources.cls.fr/documents/>).

2.3 Climatological data

We compared available observational nutrient data (NO₃) from the ~~upper 20 m of the water column sea surface (mean 0-20 m)~~, obtained by merging bottle cast data from the World Ocean Database (WOD18, Boyer et al., 2018; <https://www.nodc.noaa.gov/>), with model results. Generally, we found good agreement between nitrate in situ data and model results ($r=0.98$). Main deviations occur in the Southern Ocean where NO₃ concentrations are ~~slightly overestimated (up to 7.2 mmol m⁻³) and in some coastal areas affected by river runoff.~~—.

2.3 Climatological data

~~Data on climate indices were obtained from available databases. Bi-monthly Multivariate El Niño Southern Oscillation Index (MEI.v2), hereafter ENSO index, was obtained from the National Oceanic and Atmospheric Administration National Center for Environmental Prediction website (<https://www.esrl.noaa.gov/psd/enso/mei/>). MOC data (Moat et al., 2022; Smeed et al., 2019) for the period (2004-2018) was obtained from the RAPID-WATCH MOC monitoring project (www.rapid.ac.uk/rapidmoc).~~

2.4 Identification of HNLC regions

Presently, the best approximation to define the global distribution of HNLC regions in the world ocean is the use of NODC maps of surface nutrients (<https://www.nodc.noaa.gov/>). However, excess nutrient availability by itself does not necessarily reflect HNLC conditions. In situ experiments are capable to discern Fe limitation conditions but a more manageable metric to assess the limits on the spatial extent of HNLC regions is required, in particular for remote sensing applications, as well as for allowing objective comparison between different environmental scenarios and studies.

To obtain a quantitative criterion for the definition of HNLC regions, we analyze the values of NO₃:Chl ratios (mmol/mg) obtained from the SOM analysis on the time domain over the global ocean throughout the 20 years of data to identify a common statistical behavior representing—_HNLC conditions. In particular, we ~~analyze~~use the probability density function (*pdf*) of the extracted SOM NO₃:Chl temporal patterns to identify a threshold for defining HNLC conditions (P_{HNLC}). ~~We use the changes~~Changes in the trend of the standard deviation calculated for each bin of the *pdf* function are employed to set the~~a~~ threshold ratio. To calculate the total extent of each region (km²) the spatial area of each pixel was calculated, by considering its latitude.

2.5 Time and space domain SOM analyses

We use SOM (Kohonen, 1982) to elucidate spatial and temporal patterns in the complex relationship between nutrients and phytoplankton. SOM is a subtype of artificial neural network that uses an unsupervised machine learning algorithm to process and extract hidden structures in large datasets. The SOM algorithm is mainly based on a training process through which an initial neural network is transformed by iteratively presenting the input data. In this study, the architecture of the neural network is set in a sheet hexagonal map lattice of neurons, or units, to have equidistant neurons, and to avoid anisotropy artifacts. Each neuron is represented by a weight vector with a number of components that is equal to the dimension of the input data vector, i.e. number of rows or columns in the Chl and NO₃ matrices, depending on whether the analysis is performed either in the temporal or in the spatial domain. We use an initial network composed of units of random values (random initialization). In each successive iteration during the training process, the neuron with the greatest similarity (excited neuron), called Best Matching Unit (BMU), is updated by replacing their values with the Chl and NO₃ values of the input sample data. The similarity is estimated by computing the Euclidean distance between the components of the input sample and the components of the weight vector of the unit. The unit most similar to the input sample is the one with the minimum distance. In the learning process, Chl and NO₃ values of the topological neighboring neurons of the excited neuron (BMU) are also updated replacing their values with values determined by a Gaussian neighborhood function. In these computations, we use the imputation batch training algorithm (Vatanen et al., 2015) where the SOM assumes that a single sample of data (input vector) contributes to the creation of more than one pattern, as the whole neighborhood

around the best-matching pattern is also updated in each step of training. This yields a more detailed assimilation of particular features appearing on neighboring patterns. A final neural network with the NO₃:Chl patterns is obtained after repeating the training process until a stable convergence of the map is obtained.

For typical satellite datasets, the SOM can be applied to both space and time domains. By applying the SOM in the spatial domain, one can extract characteristic spatial patterns of the input data. If transposing the input data matrix and applying the SOM in the time domain, one can extract characteristic temporal patterns, i.e., the characteristic time series. Since each of these time series represents the temporal variability of a particular region, this method can be used to identify regions of differentiated variability on a map. The SOM, when applied to both space and time domains of the same data (called "dual SOM" analysis by Liu et al., 2016), provides a powerful tool for diagnosing ocean processes from ~~such~~ these different perspectives. In this study, we focus on the second type. We have addressed the analysis separately in the time and space domains of the log-transformed NO₃ and Chl datasets. In the time domain, we implement a [4x3] joint-SOM analysis of NO₃ and Chl using as input weight vectors concatenating the time-series of NO₃ and Chl at each pixel, so each neuron corresponds to a characteristic joint NO₃ and Chl temporal pattern over the total period of data. Since each pixel has an associated characteristic time series, we can obtain the location of a particular temporal pattern by computing the BMU for each pixel, providing a map of regions of differentiated NO₃:Chl temporal variability. For the analysis herein presented only the regions with NO₃:Chl > P_{HNL}C are considered (regions R1 to R5).

An obstacle to the temporal domain analysis on a global scale is the opposed seasonality in both earth's hemispheres. The algorithm classifies the time series at each grid point attending to the period of the signal but does not consider time lags between the time series. Hence, pixels located either in the northern or in the southern hemisphere displaying a similar significant period in the NO₃ and Chl temporal variability are classified in the same regional pattern even if they are in antiphase when the signals are seasonally lagged (6 months delayed). Regionalization is spatially coherent but the seasonal variation in the characteristic pattern that represents the neuron mixes the phenological patterns of both hemispheres. Therefore, to properly analyze the properties and trends of each of the classified regions, we have calculated the mean features of the regions by segregating the grid points corresponding to each pattern obtained from the SOM analysis into the northern, equator, and southern hemispheres (see scheme in Fig.1). Linear trends of NO₃ and Chl concentrations in each region are estimated by decomposing the NO₃:Chl time series in a seasonal signal plus a residual component and applying Theil-Sen slope adjustment (Sen, 1968) of the residuals of the deseasonalized series. Correlation analyses were performed using the Pearson Product Moment correlation computing best-fit linear trends using regression analysis.

The SOM analysis in the spatial domain [3x3] array, is addressed by using as input data weighted vectors consisting of spatial distributions over the global ocean of NO₃:Chl ratios at a particular time. The selection of the number of neurons depends on the complexity of the data, on the features to be examined in the dataset, and on the minimization of the errors. In this case, the resulting neurons after the training loop unveil the characteristic patterns describing the spatial variability of the HNLC regions on a global scale. Then, when computing the BMU for each time we designate the extracted characteristic spatial pattern that better describes the spatial distribution of NO₃:Chl ratios (P1 to P9) at each time, obtaining the time evolution of the characteristic spatial patterns throughout the considered period.

Because the SOM is based on the similarity computed from the Euclidean distance between samples, the input vectors of the different variables are normalized to the same range, before initializing the SOM computations. This guarantees a consistent comparison of the weights of the components when computing the distance of two vectors.

The size of the neural network (number of neurons) depends on the number of samples and on the complexity of the patterns. An optimal choice is important to maximize the quality of the SOM. In the present study, the map size is set to be [4 x 3] with 12 neurons for the time domain analysis, and a [3 x 3] neural network is used in the spatial domain. Using larger map sizes, the patterns are slightly more detailed, and more regions of a particular variability emerge, but the occurrence of the probability of the patterns decreases, without affecting the results noticeably (Basterretxea et al., 2018; [Hernandez-Hernández-Carrasco and Orfila, 2018](#)). If a reduced neural map, such as [2 x 2] is used, patterns are concentrated together with the occurrence probability in a few rough patterns but increasing, in this case, the topological error.

SOM computations have been performed using the MATLAB© toolbox of SOM v.2.0 ([Vesanto Vesanto et al., 1999](#)) provided by the Helsinki University of Technology (<http://www.cis.hut.fi/somtoolbox/>). Further information on SOM analysis is provided in the supplementary materials.

2.6 Combined SOM - wavelet coherence analysis

Joint SOM-wavelet power spectral analysis was demonstrated by Liu et al. (2016) in the study of characteristic time series of sea level variations in different regions of [the Gulf of Mexico](#). Here [in this study](#), we expand it further to combined SOM-wavelet coherence analysis to assess the response of HNLC regions to global forcings we use an approach based on the wavelet coherence analysis (WCA) between two time-series (Grinsted et al., 2004; see Supplementary Material for further details). WCA characterizes cross-correlations by identifying the main frequencies, phase differences, and time intervals over which the relationship between the variability of HNLC regions and the main global forcings considered in this

study, ENSO and MOC indexes, is strong. To do so, we first analyze the variability in both frequency and time of the characteristic time series of NO₃:Chl in the different HNLC regions extracted by the time domain SOM computations and the time series of the global forcings using the continuous wavelet transform (CWT).

Cross-wavelet transform (XWT) characterizes the association between the CWT of two signals, providing information on the common power and relative phase in the frequency-time domain of two time-series. By applying the XWT to the NO₃:Chl ratios and climate forcings, we determine the cyclic changes in each of the HNLC regions and their relationship with the global forcings mentioned above. Finally, we quantify the correlation between the continuous wavelet transform of two signals using the wavelet coherence analysis (WCA). In the time-frequency space the wavelet coherence coefficient R^2 is calculated as the squared absolute value of the smoothed cross-wavelet spectrum normalized by the product of the smoothed wavelet individual spectra for each scale (Torrence and Compo, 1998; Torrence and Webster, 1999; Grinsted et al., 2004). R^2 is interpreted as a localized correlation coefficient in the frequency-time domain and it takes values between 0 (no correlation) and 1 (perfect correlation). The statistical significance level of the wavelet coherence is estimated using Monte Carlo methods as described in Grinsted et al. (2004).— We use the MATLAB software package (Grinsted et al., 2004) for wavelet coherence analysis. It should be noted that cross-wavelet analysis does not establish causative relationships but only allows identifying possible linkages between variables through the synchrony of their time series.

3 Results

3.1 Global characterization of HNLC regions

The mean pattern of global ocean color data for the 20 years analyzed reveals the well-known contrast in phytoplankton biomass between the highly productive areas located in high latitudes and coastal upwelling regions, and the low-latitude oceanic waters, where mean values are $<0.1 \text{ mg m}^{-3}$ (Fig. 2). Low Chl regions generally correspond with low surface NO₃ concentrations whereas the opposite relationship (high nitrate and high chlorophyll) is more exceptional. Indeed, nutrient-rich productive waters are mainly restricted to shelf regions (coastal upwelling regions and shelf seas), or to the vicinity of islands (i.e. Falkland Islands) and other topographical features where multiple and overlapping sources of other elements, such as trace metals, are abundant (e.g. Pollard et al., 2009; Boyd and Ellwood, 2010). As shown in figure 2, only in the North Atlantic, the Bering Sea, and the eastern region of the Antarctic Peninsula, Chl is enhanced. Conversely, a large part of surface ocean waters, particularly in the Southern

Ocean and in the Equatorial Pacific, correspond to regions of relatively low Chl concentrations but with excess nitrate (i.e. $>4 \text{ mmol m}^{-3}$).

The analysis of the normalized *pdf* of the $\text{NO}_3:\text{Chl}$ extracted from the temporal SOM analysis (shown in Fig. S2) provides a good discrimination criterion to define HNLC regions (shown in Fig. S2). As shown in figure 2b, the normalized *pdf* of the $\text{NO}_3:\text{Chl}$ ratio displays a marked bimodal distribution with the main mode centered at low $\text{NO}_3:\text{Chl}$ ($\sim 5 \text{ mmol mg}^{-1}$). The second mode, which corresponds to high nutrient-low chlorophyll regions, is characterized by mean and standard deviation values of $\mu=24.1$ and $\sigma=6.7 \text{ mmol mg}^{-1}$, respectively. A critical $\text{NO}_3:\text{Chl}$ ratio bounds the lower limit of this distribution and can be estimated as $\mu-\sigma=17.4 \text{ mmol mg}^{-1}$. Consistently, the *pdf* bulk analysis of its associated standard deviation (*std*) function also reveals a clear critical value located where the value of the slope varies (Fig. 2c). Both analyses allow establishing a solid statistical criterion to infer a minimum value of $\text{NO}_3:\text{Chl}_{\text{HNLC}}=17 \text{ mmol mg}^{-1}$ for delimiting HNLC regions from other ocean regions. It is worth mentioning that while the *pdf* of the $\text{NO}_3:\text{Chl}$ values obtained from the SOM analysis shows a bimodal mode, the bulk *pdf* of the raw $\text{NO}_3:\text{Chl}$ values (i.e. without performing a SOM analysis) is unimodal. This suggests that the SOM technique is able to unravel relevant structures in the data that cannot be identified using classical approaches.

From the 12 characteristic time patterns of $\text{NO}_3:\text{Chl}$ variability obtained in the [4 x 3] SOM analysis, five display $\text{NO}_3:\text{Chl}$ exceeding P_{HNLC} all the times (not partially) throughout the entire study period (Fig. 3a). These associated subregions (R1 to R5) match with the three traditionally reported HNLC regions (Fig. 3b). In these regions surface chlorophyll rarely exceeds 0.8 mg m^{-3} and the mean values range between 0.21 mg m^{-3} and 0.5 mg m^{-3} (Table 1). The global extent of these 5 SOM-identified HNLC subregions encompasses 25% of the ocean, being the SO by far the broadest region (18% of the ocean), whereas SNP and EEP respectively occupy some 4% and 3% of the ocean. Besides the obvious absence of HNLC regions in the northern and central Atlantic, some latitudinal asymmetries are observed in the distribution of these regions. For example, the SO region extends to lower latitudes than the SNP (i.e. $\sim 40^\circ \text{ S}$), loosely coinciding with the South Antarctic Zone limit (SAZ; Orsi et al., 1995). Likewise, consistent with previous studies of this region (Radenac et al., 2012), the EEP displays a larger extent in the Southern Hemisphere (Fig. 3b).

The global pattern obtained from the SOM analysis reflects a clear latitudinal zonation which is mainly due to latitudinal variations in nutrient availability; while chlorophyll concentration doubles along the latitudinal gradient (R1 to R5), NO_3 increases up to 7-fold (see Table 1). It is noteworthy that nutrient concentrations are generally lower in the SNP (i.e. $<17 \text{ mmol m}^{-3}$) than in the SO_2 , while Chl is comparatively higher (see Table 1). Indeed, R1 in SNP only achieves the $\text{NO}_3:\text{Chl}$ criterion for HNLC regions during some periods. This region exhibits distinctive eastern and western provinces, which are

consistent with previous studies describing the western region as more productive and variable (Imai et al., 2002; Harrison et al., 2004).

Major differences among the characteristic NO₃:Chl patterns in the defined subregions are not only indebted to variations in mean values but, also, to the intensified seasonal variability in higher latitudes. For example, seasonality in Chl is particularly evident in R5 and, less so, in other polar subregions (Fig. 3b). Conversely, the seasonal component of variability in the EEP is masked by the intense short-term variability.

An interesting feature depicted from the temporal SOM analysis is the positive trend in Chl experienced in the HNLC regions located in polar areas, suggesting an increase in their productivity. Decadal tendencies are in the range of 0.04 to 0.06 mg m⁻³ decade⁻¹ in the most productive subregions (R2 to R5 in SO and R3 in SNP) but become negligible at the equator (Table 1). A regional average indicates a Chl increase of 0.6% yr⁻¹ in the SNP and a 1.9% yr⁻¹ in the SO. Nevertheless, in the case of the SO, positive trends are highly influenced by a positive Chl shift occurring at the end of 2010 (see Fig. 3 where mean Chl increased by 24% (see next section)).

3.2 Spatial variability of HNLC regions

The set of 9 coherent spatial patterns resulting from the SOM analysis in the space domain and their respective probabilities of occurrence are shown in figure 4. The organization of the maps in the figure reveals a hierarchical classification of the maps or spatial patterns. The most differentiated patterns, also displaying the highest probability of occurrence (the probability of finding a pattern similar to the input data), are located in the corners of the neural network and transitional stages connecting these patterns fill the center. For example, along the top and left side patterns (P1, P2, P3, and P4), generally occurring during winter (see Fig. 4 and 5b), the SNP extends over a larger region compared to P7, P8, and P9, which display a 3% decrease from the mean extent. Conversely, Fe limitation in the SO, as inferred from high NO₃:Chl ratios, is markedly enhanced towards the top and right side of the figure (P4, P5, P7, P8, and P9). The extent of the EEP region displays little variation. It should be noted that HNLC spatial extent and NO₃ concentrations are not necessarily coupled since the boundaries also depend on Chl concentration values. In addition, patterns in the proximity of the Antarctic continent are, in some cases, not well-defined during winter due to ice cover in this region.

Figure 5 displays the time-series of the BMUs and the monthly frequency of occurrence for each pattern. The main feature observed is the marked seasonality in the patterns shown in figure 4. The patterns with the highest probability of occurrence, P3 and P9 (100% in April and 70% in July respectively), represent spring and summer situations in the northern hemisphere. P4 and P8 characterize transitions toward these

patterns. Other patterns such as P6 and P2 (mostly occurring in winter and summer) are rarer but become more frequent after 2010 (Fig. 5a). As discussed below, this variation in HNLC regional patterns (i.e. ~~P3 substitutes P1P1 is not longer observed~~) suggests an abrupt and major transition towards more productive HNLC regions (higher Chl is observed).

From the nine spatial patterns shown in figure 4, we estimated the seasonal and interannual variation in the extent of the HNLC regions (Fig. 6). Note that this regional partitioning is made on a global scale with global criteria and therefore leads to a large-scale smoothing, which could impact the values of the variation of the areas. However, as this signal smoothing is common to all the areas, this should not have any effect on the regional comparison of the area variation. The magnitude of these variations remarkably contrasts between the equatorial and polar regions. While the extent of the EEP only varies by 8.9% seasonally, changes in SNP extent can exceed 100% (Figure 6). The peak in extent for the SNP corresponds to the boreal spring (63% of the mean value in April). In the case of the SO, seasonality is mainly driven by changes related to the ice limit in high latitudes. Indeed, the extent of the HNLC region in the austral winter is <20% of the mean annual extent.

A remarkably good inverse correlation ($r=-0.97$, $n=20$) is observed between the interannual variations in the extent of EEP and the SO, and a weaker though significant relationship exists between the SNP and the EEP ($r=-0.50$, $n=20$). Therefore, as the extent of HNLC in polar regions contracts (~~biomass increases~~), the equatorial region expands and vice versa. All three regions exhibit a shift in their annual extent after 2010 (Fig. 6). Both the SNP and the SO decrease after this year (5% and 2.6%) whereas the extent of the EEP slightly increases (0.4%).

—3.3 Climate drivers of HNLC region temporal variations

During the period considered here, 1997/98 2009/10, and 2015/16 were particularly strong El Niño events whereas weaker episodes occurred in 2002/03, 2004/05, and 2006/07 (E1, E2, and E3 in Fig. 7a). The 1997/98 and 2009/10 El Niño events were followed by fast transitions to La Niña phase (L1, L2 in Fig. 7a).

The WCA between NO₃:Chl ratio in each HNLC region and ENSO are shown in figures 7b to ~~47d~~. Generally, small coherence structures are observed at semiannual periods; ~~however, in particular, in the SO. However,~~ the main coherence pattern corresponds to a band extending in the 2 to 4 years in the SNP and > 2yr in the EEP. This coherence between NO₃:Chl and ENSO in the 2-4 year period is particularly clear after 2005 (Fig. 7b). In the EEP, the coherence between both series expands to periods >4 years but, unlike in the SNP region where the NO₃:Chl ratio is in-phase with ENSO signal, the signals are strongly anticorrelated in this case (anti-phase: relative phase of 180° between both signals).

~~As in the case of ENSO, the MOC presents strong seasonal and interannual variations but it is also expected to play a more active role at longer timescales (i.e. decadal and multidecadal; Buckley and Marshall, 2016).~~

Figure 7e shows the MOC transport index (hereafter MOI) measured at 26.5°N (Smeed et al., 2019). MOI displays intense interannual variability however, a clear long-term variation is observed. Transport exceeds ~~17 Sv~~ until 2009, but it weakens during 2010, stabilizing thereafter. ~~Generally, the MOI displays intense interannual variability, and~~ As shown in figures 7f to 7h coherence with NO₃:Chl ratios is strongest at interannual time scales (1-1.5 yr; Fig. 7f to h). At this timescale, it influences NO₃:Chl ratios in the three HNLC regions yet it is more intense in the SO. —) when MOC is debilitated.

4 Discussion

4.1 Global characterization of HNLC regions

In the present study, we have addressed the extent of the HNLC regions, their long-term variability, and the potential drivers of these variations. Despite the relevance of precise characterization of the extent of this biome for the understanding of physical ocean processes (i.e. upwelling), and the estimation of the amount of carbon drawn into the ocean by phytoplankton, objectively determining the boundaries of HNLC regions has remained elusive as it requires coherent information from both nutrient and Chl. We demonstrate that a statistical approach, based on a threshold in the distribution of the global NO₃:Chl ratios (P_{HNLC}), can robustly delineate these regions: (P_{HNLC}). This is also evidenced in Table 1 where the characteristics of each cluster obtained with SOM analysis are remarkably constant (i.e. low *std*).

As in precedent studies (e.g. Moore et al., 2013), we assume that excess NO₃ in surface oceanic areas is indicative of Fe limitation. This avoids relying on the scarce information available on Fe-stress or in more complex ecosystem modeling approaches. Inference of phytoplankton Fe-stress from satellite ocean color data has been attempted but it is a methodology that still presents large uncertainties (Browning et al., 2014). Furthermore, while bioavailable Fe is known to be the primary limiting factor in this biome, the establishment of HNLC conditions is influenced by various other factors such as light availability, grazing pressure, rate of Fe-remineralization, and community structure, highlighting the diverse interrelations among these factors. Despite these drawbacks, the herein-developed method provides results consistent with previous descriptions of the large-scale spatial patterns of HNLC regions, mostly based on NO₃ fields (i.e. Archer and Johnson, 2000; Ono et al., 2008; Fu and Wang, 2022). Also, the proposed method for biome definition may introduce a bias in that the resulting spatial fields are smoother compared to those based on Fe-limitation, which is due to the greater variability of Fe concentrations compared to NO₃ fields.

The P_{HNLC} obtained from the *pdf* distribution of the $\text{NO}_3:\text{Chl}$ ratios represents a statistical threshold that integrates complex biological processes, including competition for resources, grazing, changes in species composition, nutrient uptake rates, Fe-regulated algal photochemistry, etc. Unlike Redfield or C:Chl ratios which respond to physiological factors within phytoplankton cells, P_{HNLC} can be considered an environmental indicator of changes in the structure and functioning of marine phytoplankton.

According to our analysis, some 25% of the ocean (18% of the Earth's surface) corresponds to unproductive HNLC waters. With 83% of the global HNLC biome extent (Table 2), the SO is the largest region and it is the only one presenting clear latitudinal variation in the characteristic Chl patterns (Table 2 and Fig. 3). This is consistent with available descriptions of the physical and chemical properties of the SO which tend to be across latitude due to the meridional structure of the MOC and because of the rapid zonal redistribution imposed by circumpolar currents (Orsi et al., 1995). The SNP and the EEP each constitute ~8% of the total HNLC. However, while the EEP remains relatively stable ($cv=5$; Table 2) the SNP can change up to 2-fold (Fig. 4 and Table 2).

Our analysis reveals marked decadal tendencies in Chl in the most productive subregions, ranging between 0.04 to 0.06 $\text{mgChl m}^{-3} \text{ decade}^{-1}$, and negligible trends in the EEP (Table 1). Positive phytoplankton Chl trends in high latitudes and no significant tendencies in the equator at the 95% interval have been previously reported by Hammond et al. (2017). Indeed, climate change projections for the 21st century predict declines in global marine net primary production but increasing Southern Ocean productivity (Hauck et al., 2015; Moore et al., 2018). Nevertheless, it is noteworthy that our analysis shows that trends in some regions of the southern hemisphere are influenced by a Chl shift occurring in 2009-2010 (particularly R4 and R5). ~~It is unlikely that this change~~ The possible source of this shift in overall phytoplankton biomass is more extensively discussed below. However, it is unlikely to be related to satellite data merging. SeaWiFS ended operations in 2010 and MERIS sensor ceased in April 2012; however, a decrease in biomass would be expected from these changes (Garnesson et al., 2019). ~~The possible source of this shift in phytoplankton biomass is more extensively discussed below.~~

4.2 Spatial variability of HNLC regions

SOM analysis allowed the characterization of the seasonal and interannual variability of HNLC regions (Fig 5 and 6). The SO and the SNP are regions of seasonal extremes in productivity as a consequence of the large fluctuations in the environment that they experience. Important seasonal variability is observed in the SO, which is attributed to light limitation during winter and to variations in iron stocks occurring in surface waters (Tagliabue et al., 2014). Therefore, while presenting HNLC characteristics for most of the year, the SO exhibits distinct Chl variability patterns that are well captured in the SOM regionalization

and the characteristic patterns of each subregion. In the case of the SO, the changes in the extent are reduced (20% from mean values) which ~~is attributed~~we attribute to the large-scale nature of the physical processes regulating the productivity of this region ~~(Cullen, 1991)~~. Mid-depth and deep ocean waters communicate with the ocean surface after following long a circuit route driven by ocean overturning circulation (Lumpkin and Speer, 2007). As reported by Smeed et al. (2018), the variability of the MOC flow system has an important decadal component associated with thermohaline forcing. This long-term component of variability could dominate over higher frequency variability but the length of the observational record of the AMOC is still insufficient to resolve variations at this scale.

Variability patterns in the SNP, are attributed both to marked seasonality in the local forcings and ~~to~~ fluctuations in the regional circulation patterns ~~and the consequent advective Fe enrichment processes from river discharges of the surrounding marginal regions in the North Pacific (Cummins and Freeland 2007; Takeda, 2011; Lauderdale et al., 2020; Nishioka et al., 2020). In the Northern Pacific, surface Fe availability is driven by vertical mixing processes which closely relate to seasonal variations in weather conditions. Nishioka et al. (2021) discuss in detail the sources of phytoplankton variability in this region, distinguishing three cases of biological response to Fe in this region. The annual bloom in open waters would be controlled by the sedimentary Fe supplied from the continental marginal regions and circulated laterally through the intermediate layer (e.g. Lam et al., 2006; Cummins and Freeland, 2007; Takeda, 2011). These nutrients are upwelled to the surface by several mixing processes (winter mixing and eddy diffusive mixing), including interactions of tidal currents with the rough topography (Nishioka et al., 2007; 2020). Upwelling is triggered by severe winter cyclones that generate enough Ekman pumping to maintain high nutrient concentrations in the near-surface waters (Gargett, 1991; Harrison). Massive spring phytoplankton blooms that occur around the coastal boundary current would be the result of the large amounts of Fe originating from coastal shelf areas (Cullen et al., 2004, 2009). Finally, intense sporadic and patchy phytoplankton production is occasionally observed in both spring and summer in response to the atmospheric dust Fe supply.~~

Marked differences in variability are observed between the subpolar regions and the EEP, where seasonality is weak. Nevertheless, it should be noted that the EEP is a peculiar region that integrates subregions with 6-month out-phased seasonal variations (north and south of the equator). On average, the south-equatorial region contributes more (62% of the total extent) to the mean extent of the EEP whereas the Northern subregion determines most of the observed variability (Fig. S3).

At interannual scales, variations in the extent of HNLC regions ~~is~~are more modest (up to 5%; Figure 6) but all three regions are notably correlated, suggesting that their interannual variations are driven by global-scale processes. In particular, the SO and the EEP are highly correlated ($r=0.97$). ~~This~~A coupling between Southern Ocean productivity, and equatorial productivity, was suggested by Dugdale et al.

(2002) based on the observed nutrient ratios. They proposed that both regions were connected, out of phase, by the meridian SubArctic Mode Water. Nevertheless, the seaways in the Pacific circulation pathways are complex, and determining the overturning signature is challenging. Some studies have suggested that temperature anomalies subducted into the pycnocline at subtropical latitudes may not reach the Equator with any appreciable amplitude (Schneider et al., 1999a, 1999). However, mass water balances in the equatorial Pacific reveal that the strength of the equatorial upwelling is related to variations in the Pacific overturning (PMOC) (McPhaden and Zhang, 2002), and therefore, an influence on EEP extent is expected. Likewise, PMOC extends to the North Pacific; however, it could be presumed. Nevertheless, it should be reminded that the extent of the HNLC regions is not necessarily solely dictated by upwelling intensity but it is also defined by the interaction with adjacent circulation patterns (i.e. the subtropical gyres in this case).

The pathways of PMOC in the North Pacific are structured in a multicell configuration. The SNP does not ventilate the deep ocean at significant rates, and the PMOC cell at this latitude rather corresponds to an independently functioning intermediate water cell (Warren et al., 1983). PMOC in this region is reportedly weak (1-4 Sv) and extends no further than 50°N (Ishizaki, 1994; Yaremchuk et al., 2001). However, there is evidence showing the response of the SNP to changes in PMOC (Burls et al., 2017; Holzer et al., 2021). For example, it has been observed in TOPEX altimeter data that meridional overturn transport influences the basin-scale baroclinic circulation in the SNP (Kuragano and Kamauchi, 2004). This would explain the lower, yet significant, correlation with the variations at the exhibited by SO and EEP at interannual scales (Fig. 6). Indeed, the shift observed in 2009-2010 is common to all three regions (albeit with a different sign).

The causes of the drastic 2009-2010 variation in the extent of HNLC regions are uncertain but they can plausibly be related to changes in the strength of meridional circulation and the consequent atmospheric and oceanic variations. For example, Moat et al. (2020) suggest a previously unsuspected role for the AMOC in climate variability during the 2010 event which coincided with a cold winter in Europe. Several ocean scale changes that may be related to the observed changes in the HNLC region extent. Several ocean scale changes have been reported in the period 2009-2011. For example, rapid warming, salinification, and a concurrent dissolved oxygen decline have been observed at BATS during the 2010s (Bates and Johnson, 2020). There is also evidence indicating that a decadal intensification of Pacific trade winds weakened in 2011 (Bordbar et al., 2019). Trade wind intensity variability in the equatorial Pacific region is associated with SST anomalies, weakening of the equatorial divergence, and changes in upper-ocean thermal structure (England et al., 2014; Bordbar et al., 2017). The relationship between the equatorial wind intensity and the equatorial undercurrent strength is also well established (McPhaden, 1993). These atmospheric changes, affecting the upwelling of Fe in the EEP (and indirectly

to other oceanic regions), agree with the hypothesis of Winckler et al. (2016) suggesting that ocean dynamics, not dust deposition, control the equatorial Pacific productivity. ~~In our case, we observe a reduction of the extent of the HNLC region in the EEP during an enhanced wind intensity period (before 2011) and an expansion thereafter (Fig. 6). Connections of this mechanism to high latitude HNLC regions reveal large scale adjustments with consequences in global ocean productivity.~~

4.3 Drivers of HNLC region variability

~~ENSO is the primary source of the interannual variability in the EEP and its occurrence is related to the decline in NO₃ supply. The EEP the period of strongest seasonal upwelling is in the boreal summer (e.g. Philander and Chao, 1991), which is associated with an expansion of the HNLC in Figure 6a. We also observe an expansion of the extent of the HNLC region in the EEP after 2021 (Fig. 6). Changes in these forcings could explain variations in the seasonal variability but the long-term changes in HNLC extent would require more a permanent variation in the stoichiometric equilibria or in the functioning of the marine woodwebs.~~

4.3 Drivers of HNLC region variability

ENSO is the primary source of the interannual variability in the EEP and its occurrence is related to the decline in nutrient supply. For example, the deepening of the thermocline during ENSO is associated with the depression of the EUC that transports Fe across the basin from the western Pacific (Gordon et al., 1997) also reducing nutrient supply and phytoplankton productivity (Chavez et al., 1999; Strutton and Chavez, 2000). Sub-decadal fluctuations in Chl in the EEP region displaying a good correlation with the ENSO index have been reported before (Oliver and Irwin, 2008; Boyce et al., 2010). By contrast, the SO only shows weak evidence of this relationship which suggests that ENSO is not a major forcing driving the variability of NO₃:Chl in this region. This is consistent with reports from Ayers and Strutton (2013) who did not find a significant relationship between nutrients in this region and ENSO.

It can be argued that differences in the response to ENSO are due to the different nature of the forcings driving nutrient supply in each region. While the EEP and the SNP seem to respond to ENSO-related changes in wind forcing, NO₃:Chl ratios in the SO are more stable and respond to annual and semi-annual variations. The coupling between the EEP and the SNP dynamics has been reported before. Qiu (2002) observed progressive shoaling of the Alaska gyre caused by a strengthening of the cyclonic circulation. The interannual variability of this gyre was connected to ENSO-related sea surface height anomalies originating in the tropics. Several large-scale climate pattern indexes are invoked to explain physical and

biological fluctuations in the SNP. For example, Di Lorenzo et al., (2008) defined the North Pacific Gyre Oscillation (NPGO), which explains strongly correlated fluctuations of salinity, nutrients, and chlorophyll related to the circulation in the North Pacific gyre. It is beyond the scope of the present work to assess the relationships of all these indexes with the variability in the extent of HNLC regions. Nevertheless, the proposed climate indexes for the Pacific present a high relationship among them, which highlights the strong dynamical linkages between tropical and extratropical modes of climate variability in the Pacific basin, and the important role played by ENSO (Di Lorenzo et al., 2013).

~~The interpretation of the influence of MOC on global ocean productivity is more challenging. A stronger MOC should result in the faster upwelling of macronutrients and Fe at faster rates high latitudes as well as in increased Ekman transport of nutrients equatorward and subsequent subduction (Ayers and Strutton, 2013). This is observed in the high positive cross-wavelet correlation at 1–1.5 yr in the EEP region. In addition, a clear variation in the coherence phase is observed, being 90° (3–4 months) in the SNP, in phase for the EEP, and 270° (9–11 months) out of phase in the SO, suggesting a meridional propagation of the MOC effect. Figure 7e also reveals a decline of the MOC around 2010 that has remained ~15% below its pre-2010 level since then (~17 Sv; Ayala-Solares et al., 2018; Caesar et al., 2018). This trend has been attributed to climate warming and the consequent changes in the hydrological cycle, including sea-ice loss and accelerated melting of the Greenland Ice Sheet, causing further freshening of the northern Atlantic (Bakker et al., 2016; Böning et al., 2016). It has been proposed that AMOC weakening will affect large regions of the world's upper oceans that are currently supplied with nutrients by the South Antarctic Mode Water (Schmittner, 2005).~~

~~Weakened MOC after 2010 and, the particularly low value during that year (Fig. 7e), is coincident with the shift in the extent of the HNLC regions shown in Figure 5, suggesting that weaker MOC is related to increases in the extent of the EEP and contraction of the SNP and the SO HNLC regions. In the case of the EEP, it could be proposed that a larger equatorial area with Fe deficiency would be associated with a decline in upwelling intensity. It is estimated that the slowing down of the overturning circulation in the Pacific Ocean from the 1970s to the 1990s generated a decrease in upwelling of about 25% in an equatorial strip between 9° N and 9° S (McPhaden and Zhang, 2002). However, detailed information on long-term PMOC variability and its influence on EEP biogeochemistry is scarce. Alternatively, the larger HNLC region in the EEP could also be explained by the depletion of Fe in the source waters feeding the EUC. Consequently, a weakened MOC should plausibly favor Fe recycling rates, as reported by KauppRafter et al., (2014, 2017).~~

At high latitudes, the weakening of the AMOC, is coherent with a decrease in the extent of the SNP and the SO (Fig. 5). This anomaly, starting in 2009–2010, is a global feature also reflected in the intertropical convergence zone (ITCZ) time series (see, Green et al., 2017; IbanezIbanez et al., 2017), suggesting a

strong atmosphere-ocean coupling with impact on ocean productivity. It is not clear how a reduced flow would favor the increase in biomass in high latitudes. ~~While it is plausible that in the case of abrupt and/or permanent variations of AMOC primary production, and hence phytoplankton biomass, will be reduced, it is less clear how present-day variations influence phytoplankton biomass.~~ It has been proposed that a reduced AMOC from increased precipitation and melting sea ice, could contribute to reduced vertical mixing which may increase productivity in polar regions (Riebesell et al., 2009). Other studies (e.g., ~~Martinez-Garcia~~ Martínez-García et al., 2009), showed a relationship between AMOC and Chl variations, mainly due to the interaction of the main pycnocline and the upper ocean seasonal mixed layer. In addition, some paleoclimatic studies have demonstrated that AMOC weakening can increase the productivity north from the Polar Front, but only if an increase in the atmospheric soluble Fe flux occurs (Muglia et al., 2018). Paleooceanographical records showing a strong correlation between proxies of aeolian Fe flux and productivity have been reported in this region (Kumar et al., 1995; ~~Martínez-García~~ García et al., 2009) but, in present times, dust deposition in this area is—_notably different (i.e. ~~McConnell~~ McConnell et al., 2007) and this effect is unlikely to be important at the time scales considered here. Complex ecosystem processes including competition for Fe with bacteria, Fe remineralization rates, and organic complexation processes could determine the phytoplankton response under future scenarios. Further, biomass building up is not only driven by nutrient availability. Changes in biomass can be produced by variations in the thermocline depth affecting the vertical distribution of phytoplankton. Nevertheless, variations in phytoplankton composition, physiological adjustments in cellular pigmentation, and grazing could also modulate Chl variability. Indeed, the prevailing foodweb structure may play an important role in Fe fertilization (Schmidt et al., 2016). At larger scales, there are still unresolved questions about the couplings occurring at different temporal scales. For example, MOC variations are known to interact with ENSO ~~variability influencing its amplitude and variance~~ (Dong et al., 2006; Dong and Sutton, 2007; Timmermann et al., 2007).

5. Conclusions

Variations in the boundaries of the HNLC regions can provide an integrative view of how climate scale ocean variations influence ocean productivity. We established a statistical criterion to identify HNLC regions from global Chl and NO₃ data that sets the basis for systematic analyses of HNLC regions and their response to climate variations.

Our results suggest that while local-scale processes can determine certain aspects of the productivity of HNLC regions, their long-term patterns are strongly influenced by variations in global atmospheric and oceanic circulation. We observed significant interannual variations in the extent of HNLC (up to 5% in

Fig. 6), which are associated with anomalies in global ocean circulation patterns (i.e Fig. 7e). Accordingly, our findings suggest a scenario in which HNLC regions are vulnerable to interbasin teleconnections rather than local forcings. These general patterns may be modulated by feedback between different forcing mechanisms. ~~For instance, there is a positive correlation between PMOC variability and El Niño-Southern Oscillation (Tandon et al. 2020).~~ Furthermore, our analysis reveals a shift in phytoplankton biomass and HNLC variation patterns occurring at the end of 2010, which evidences the occurrence of fast transitions in ocean biogeochemistry. The underlying drivers of these regime shifts and the resulting biological responses to these ocean-scale changes require further investigation since they are a fundamental aspect of long-term variations in marine ecosystem functioning.

Finally, the present study highlights the importance of maintaining long and coherent datasets beyond satellite-borne information to be able to disentangle the different components of variability, particularly at long timescales, and to evaluate the impact of climate change on marine ecosystems. Most of the geochemical information at this scale (i.e. nutrient and Fe fields) will probably require further global sampling programs and refined modeling.

Author contributions

Data were processed and analyzed mainly by G.B., J.S.F, and I.H-C. Writing by G. B., J.S.F, and I. H-C. and S.A.S. The authors declare no competing financial interests.

Funding sources and data references

This work was partially supported by SIFOMED grant (CTM2017-83774-P) from Ministerio de Ciencia, Innovación y Universidades, the Agencia Estatal de Investigación (AEI), and the Fondo Europeo de Desarrollo Regional (FEDER, UE). G. Basterretxea was supported by Salvador de Madariaga PRX18/00056 scholarship. J.S Font-Muñoz acknowledges funding by an individual postdoctoral fellowship “Margalida Comas” (PD/018/2020) from Govern de les Illes Balears and Fondo Social Europeo. I. Hernandez-Carrasco acknowledges financial support from the project TRITOP (grant UIB2021-PD06) funded by Universitat de les Illes Balears and FEDER (EU).

All data included in the present study is accessible from the following publicly available repositories: CARINA (https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/CARINA/about_carina.html), COPERNICUS (<https://marine.copernicus.eu/>), ECMWF (<https://www.ecmwf.int/>), GEOTRACES (<https://www.geotraces.org/>), GlobColour (www.globcolour.info), MERCATOR (<https://cmems-resources.cls.fr/>), NERC (<https://nerc.ukri.org>) and NOAA (<https://www.node.noaa.gov/><https://www.nodc.noaa.gov/>).

References

- Archer, D., ~~E.~~ and Johnson, ~~K.S.K.~~: A model of the iron cycle in the ocean. ~~Global Biogeochemical, Global Biogeochem.~~ Cycles, 14, 269–279. doi: 10.1029/1999GB900053–279, 2000.
- Aumont, O., Éthé, C., Tagliabue, A., Bopp, L., & Gehlen, M.: PISCES-v2: an ocean biogeochemical model for carbon and ecosystem studies. ~~Geoscientific Model Development, 8, 2465–2513, doi.org: 10.5194/gmd-8-2465-2015, Geosci. Model Dev. Discuss., 8, 1375–1509, 2015.~~
- Ayers, ~~J.M., Strutton, P.G.J. M. and Strutton, P. G.:~~ Nutrient variability in ~~Subantarctic Mode Waters~~subantarctic mode waters forced by the ~~Southern Annular Mode and ENSO.~~ ~~Geophysical Research Letters, 40, 3419–3423. doi: 10.1002/gfl.50638, 2013~~
- Ayala Solares, J.R., Wei, H.L., Digg, G.R.: The variability of the Atlantic meridional circulation since 1980, as hindcast by a data-driven nonlinear systems model. *Acta Geophysica*, 66, 1–13. doi: 10.1007/s11600-018-0165-7, 2018.
- Bakker, P., Schmittner, A., Lenaerts, J.T.M., Abe-Ouchi, A.: Fate of the Atlantic Meridional Overturning Circulation: strong decline under continued warming and Greenland melting. *Geophysical Research Letters*, 43, 12252–12260. doi: 10.1002/2016GL070457, 2016
- ~~southern annular mode and ENSO, Geophys. Res. Lett., 40, 3419–3423, 2013.~~
- De Baar, H. J. W., de Jong, J. T. M., Bakker, D. C. E., Löscher, B. M., Veth, C., Bathmann, U., and Smetacek, V.: Importance of iron for plankton blooms and carbon dioxide drawdown in the Southern Ocean, *Nature*, 373, 412–415, 1995.
- Basterretxea, G., Font-Muñoz, J. S., Salgado-Hernanz, P. M., Arrieta, J., & Hernández-Carrasco, I.: Patterns of chlorophyll interannual variability in Mediterranean biogeographical regions. ~~Remote sensing of environment, Remote Sens. Environ., 215, 7–17, 2018~~
- ~~Bates N.R., Johnson, R. https://doi.org/10.1016/j.rse.2018.05.027, 2018.~~
- Bates, N. R. and Johnson, R. J.: Acceleration of ocean warming, salinification, deoxygenation and acidification in the surface subtropical North Atlantic Ocean. ~~Communications Earth & Environment, 1, 33, doi:10.1038/s43247-020-00030-5, Commun. Earth Environ., 1, 33, 2020.~~
- ~~Bertrand E.M, E., Saito M.A, M., Lee P.A, P., Dunbar R.B., et al, R., Sedwick, P., and DiTullio, G.: Iron limitation Limitation of a springtime bacterial and phytoplankton community Springtime Bacterial and Phytoplankton Community in the Ross Sea: implications for vitamin b12 nutrition. *Frontiers in Microbiology*, 2, 160. doi: Implications for Vitamin B12 Nutrition, *Front. Microbiol.*, 2, 160, https://doi.org/10.3389/fmicb.2011.00160,–2011.~~
- ~~Bertrand, E.M.E. M., Allen, A.E.A. E., Dupont, C.L.C. L., Norden-Krichmar, T.M., et al.: Influence of cobalamin scarcity on diatom molecular physiology and identification of a cobalamin-acquisition protein. *Proceedings of the National Academy of Sciences*, 109 (26), E1762–71. doi: 10.1073/pnas.1201731109, 2012.~~

- Böning, C.W., Behrens, E., Biastoch, A., Getzlaff, K., Bamber, J. L.: Emerging impact of Greenland meltwater on deepwater formation in the North Atlantic Ocean. *Nature Geoscience*, 9, 523–527. doi: 10.1038/ngeo2740, 2016
- T. M., Bai, J., Valas, R. E., and Saito, M. A.: Influence of cobalamin scarcity on diatom molecular physiology and identification of a cobalamin acquisition protein, *Proc. Natl. Acad. Sci.*, 109, E1762–E1771, 2012.
- Birchill, A. J., Hartner, N. T., Kunde, K., Siemering, B., Daniels, C., Gonzalez-Santana, D., Milne, A., Ussher, S. J., Worsfold, P. J., and Leopold, K.: The eastern extent of seasonal iron limitation in the high latitude North Atlantic Ocean, *Sci. Rep.*, 9, 1–12, 2019.
- Bordbar, M.H.M. H., Martin, T., Latif, M., Park and Park, W.: Role of internal variability in recent decadal to multidecadal tropical Pacific climate changes. *Geophysical Research Letters*, 44, 4246–4255. doi: 10.1002/2016GL072355, *Geophys. Res. Lett.*, 44, 4246–4255, 2017.
- Bordbar, M.H.M. H., England, M.H., M. H., Sen Gupta, A., Santoso, A., et al Taschetto, A. S., Martin, T., Park, W., and Latif, M.: Uncertainty in near-term global surface warming linked to tropical Pacific climate variability. *Nature communications*, 10, 1990 doi: 10.1038/s41467-019-09761-2, *Nat. Commun.*, 10, 1990, 2019.
- Boyce, D.G.D. G., Lewis, M.R., M. R., and Worm, B.: Global phytoplankton decline over the past century. *Nature*. 466(7306), 591–596. doi: 10.1038/nature09268, *Nature*, 466, 591–596, 2010.
- Boyce, D. G., Dowd, M., Lewis, M. R., and Worm, B.: Estimating global chlorophyll changes over the past century, *Prog. Oceanogr.*, 122, 163–173, 2014.
- Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean, *Nat. Geosci.*, 3, 675–682, 2010.
- Boyd, P. W., Watson, A. J., Law, C. S., Abraham, E.R., et al E. R., Trull, T., Murdoch, R., Bakker, D. C. E., Bowie, A. R., Buesseler, K. O., and Chang, H.: A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization. *Nature*, 407, 695–702. doi:10.1038/35037500, 2000.
- Boyd, P. W., Law, C. S., Wong, C. S., Nojiri, Y. et al, Tsuda, A., Levasseur, M., Takeda, S., Rivkin, R., Harrison, P. J., Strzepek, R., Gower, J., McKay, M., Abraham, E., Arychuk, M., Barwell-Clarke, J., Crawford, W., Crawford, D., Hale, M., Harada, K., Johnson, K., Kiyosawa, H., Kudo, I., Marchetti, A., Miller, W., Needoba, J., Nishioka, J., Ogawa, H., Page, J., Robert, M., Saito, H., Sastri, A., Sherry, N., Soutar, T., Sutherland, N., Taira, Y., Whitney, F., Wong, S.-K. E., and Yoshimura, T.: The decline and fate of an iron-induced subarctic phytoplankton bloom. *Nature*, *Nature*, 428, 549–553. doi: 553, <https://doi.org/10.1038/nature02437>, 2004.
- Boyd, P.W., Strzepek P. W., Strzepek, R., Takeda, S., Jackson, G. et al.: Mesoscale iron enrichment experiments 1993–2005: Synthesis and future directions. *Science*. 315, 612–617. doi: 10.4319/lo.2005.50.6.1872, Wong, C. S., McKay, R. M., Law, C., Kiyosawa, H., Saito, H., and Sherry, N.: The evolution and termination of an iron-induced mesoscale bloom in the northeast subarctic Pacific, *Limnol. Oceanogr.*, 50, 1872–1886, 2005.
- Boyd, P.W. P. W., Jickells, T., Law C.S., C. S., Blain, S. et al.: The evolution and termination of an iron induced mesoscale bloom in the northeast subarctic Pacific. *Limnology and Oceanography*, 50, 1872–1886, 2007.

Boyd, P.W., Ellwood, M.J.: The biogeochemical cycle of iron in the ocean. *Nature Geoscience*, 3, 675–682. doi: 10.1038/ngeo964, 2010.

Boyer, Tim P.; García, Hernán E.; Locarnini, Ricardo A.; Zweng, Melissa M.; Mishonov, Alexey V.; Reagan, James R.; Weathers, Katharine A.; Baranova, Olga K.; Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J., de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard, R., Rivkin, R. B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A. J.: Mesoscale Iron Enrichment Experiments 1993-2005: Synthesis and Future Directions, *Science* (80-.), 315, 612 LP – 617, <https://doi.org/10.1126/science.1131669>, 2007.

Boyer, T. P., García, H. E., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Reagan, J. R., Weathers, K. A., Baranova, O. K., Paver, Christopher C. R.; Seidov, Dan; Smolyar, Igor V. (2018), and Seidov, D.: World Ocean Atlas, 2018. [Nitrate,WOD]. NOAA National Centers for Environmental Information. Dataset. https://www.ncei.noaa.gov/archive/accession/NCEI_WOA18. Accessed [15/02/2022].

Browning, T.J., Bouman, H.A., Moore C.M

Browning, T. J., Bouman, H. A., and Moore, C. M.: Satellite-detected fluorescence: Decoupling nonphotochemical quenching from iron stress signals in the South Atlantic and Southern Ocean. *Global biogeochemical cycles*, 28, 510–524. doi:10.1002/2013GB00477, *Global Biogeochem. Cycles*, 28, 510–524, 2014.

Browning, T. J., Achterberg, E.P.E. P., Rapp, I., Engel, A., et al Bertrand, E. M., Tagliabue, A., and Moore, C. M.: Nutrient co-limitation at the boundary of an oceanic gyre. *Nature*, *Nature*, 551, 242–246. doi:10.1038/nature24063–246, 2017.

Bryant, D. A.: The beauty in small things revealed. *Proceedings of the National Academy of Sciences*, 100, 9647–9649. doi: /10.1073/pnas.1834558100, 2003.

Buckley, M.W., Marshall, J.: Observations, inferences, and mechanisms of Atlantic Meridional Overturning Circulation variability: A review. *Reviews of Geophysics*, 54, 54–63. doi: 10.1002/2015RG000493, 2016, *Proc. Natl. Acad. Sci.*, 100, 9647–9649, 2003.

Burls, N. J., Fedorov, A. V., Sigman, D. M., Jaccard, S. L., Tiedemann, R., & Haug, G. H.: Active Pacific meridional overturning circulation (PMOC) during the warm Pliocene. *Science Advances*, 3(9), e1700156, 2017

Caesar, L., Rahmstorf, S., Robinson, A., Feulner G., Saba, V.: Observed fingerprint of a weakening Atlantic Ocean overturning circulation. *Nature* 556, 191–196. doi: 10.1038/s41586-018-0006-5, 2018.

Chavez, F.P., Buck, K.R., Coale, K.H., Martin, J.H., et al.: Growth rate, grazing, sinking, and iron limitation of equatorial Pacific phytoplankton. *Limnology and Oceanography*, 36, 1816–1833. doi: 10.4319/lo.1991.36.8.1816, 1991.

Chisholm S.W., Morel F.M.M.: What controls phytoplankton production in nutrient rich areas of the open sea? American Society of Limnology and Oceanography Symposium – 22–24 February 1991 San Marcos, California – Preface. *Limnology and Oceanography*, 36, 1507–1965, 1991.

~~Coale, K.H., Johnson K.S., Fitzwater S.E., Gordon R.M., et al. [Sci. Adv.](#), 3, e1700156, 2017.~~

~~Chavez, F. P., Buck, K. R., Coale, K. H., Martin, J. H., DiTullio, G. R., Welschmeyer, N. A., Jacobson, A. C., and Barber, R. T.: Growth rates, grazing, sinking, and iron limitation of equatorial Pacific phytoplankton, [Limnol. Oceanogr.](#), 36, 1816–1833, 1991.~~

~~Chavez, F. P., Strutton, P. G., Friederich, G. E., Feely, R. A., Feldman, G. C., Foley, D. G., and McPhaden, M. J.: Biological and chemical response of the equatorial Pacific Ocean to the 1997-98 El Niño, [Science \(80-. \)](#), 286, 2126–2131, 1999.~~

~~Chisholm, S. W. and Morel, F. M. M.: What controls phytoplankton production in nutrient-rich areas of the open sea?, 1991.~~

~~Coale, K. H., Johnson, K. S., Fitzwater, S. E., Gordon, R. M., Tanner, S., Chavez, F. P., Ferioli, L., Sakamoto, C., Rogers, P., and Millero, F.: A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean. ~~[Nature.](#)~~ [Nature](#), 383, 495–501. doi: 10.1038/383495a0, 1996.~~

~~Coale, K.H., Johnson K.S., Chavez F.P. 501, 1996.~~

~~Coale, K. H., Johnson, K. S., Chavez, F. P., Buesseler, K., et al O., Barber, R. T., Brzezinski, M. A., Cochlan, W. P., Millero, F. J., Falkowski, P. G., and Bauer, J. E.: Southern Ocean iron enrichment experiment: ~~Carbon~~ carbon cycling in high and low Si waters. [Science](#). 304, 408–414. doi: 10.1126/science.1089778, 2004.~~

~~Cohen, N.R., Ellis, K.A. Burns, W.G., Lampe, R.H., et al and low-Si waters, [Science \(80-. \)](#), 304, 408–414, 2004.~~

~~Cohen, N. R., A. Ellis, K., Burns, W. G., Lampe, R. H., Schuback, N., Johnson, Z., Sañudo-Willhelmy, S., and Marchetti, A.: Iron and vitamin interactions in marine diatom isolates and natural assemblages of the Northeast Pacific Ocean. ~~[Limnology and Oceanography](#)~~, 62, 2076–2096. doi: 10.1002/lno.10552, [Limnol. Oceanogr.](#), 62, 2076–2096, 2017.~~

~~Cullen, J.J. J.: Status of the iron hypothesis after the open-ocean enrichment experiment. ~~[Limnology and Oceanography](#)~~, 40, 1336–1343. doi: 10.4319/lc.1995.40.7.1336, 1995.~~

~~Cullen, J. J.: Hypotheses to explain high nutrient conditions in the open sea. [Limnology and Oceanography](#), 36(8), 1578–1599, 1991~~

~~Cummins P.F, Freeland H.J.: Variability of the North Pacific Current and its bifurcation, [Progress in Oceanography](#), 75, 253–265, doi: 10.1016/j.pocean.2007.08.006, 2007~~

~~de Baar, H.J.W., de Jong, J.T.M., Bakker, D.C.E., Löseher, B.M., et al.: Importance of iron for plankton blooms and carbon dioxide drawdown in the Southern Ocean. [Nature](#). 373, 412–415. doi: 10.1038/373412a0, 1995.~~

~~Di Lorenzo E., Schneider N., Cobb K.M., Chhak K, et al.: North Pacific Gyre Oscillation links ocean climate and ecosystem change. [Geophysical Research Letters](#), 35, L08607. doi:10.1029/2007GL032838, 2008.~~

- Di Lorenzo, E., Combes, V., Keister, J.E., Strub, P.T., et al.: Synthesis of Pacific Ocean climate and ecosystem dynamics. *Oceanography*, 26, 68–81, doi: 10.5670/oceanog.2013.76, 2013, *Limnol. Oceanogr.*, 1995.
- Cullen, J. T., Chong, M., and Ianson, D.: British Columbian continental shelf as a source of dissolved iron to the subarctic northeast Pacific Ocean, *Global Biogeochem. Cycles*, 23, 2009.
- Cummins, P. F. and Freeland, H. J.: Variability of the North Pacific Current and its bifurcation, *Prog. Oceanogr.*, 75, 253–265, 2007.
- Dong, B. and Sutton, R. T.: Enhancement of ENSO variability by a weakened Atlantic thermohaline circulation in a coupled GCM, *J. Clim.*, 20, 4920–4939, 2007.
- Dong, B., Sutton, R. T., & Scaife, A. A.: Multidecadal modulation of El Niño–Southern Oscillation (ENSO) variance by Atlantic Ocean sea surface temperatures. *Geophysical Research Letters*, 33(8), 2006
- Dong, B., & Sutton, R. T.: Enhancement of ENSO variability by a weakened Atlantic thermohaline circulation in a coupled GCM. *Journal of Climate*, 20(19), 4920–4939, 2007
- Dugdale R.C., Wischmeyer A.G., Wilkerson F.P., Barber R.T., et al, *Geophys. Res. Lett.*, 33, 2006.
- Dugdale, R. C. and Wilkerson, F. P.: Silicate regulation of new production in the equatorial Pacific upwelling, *Nature*, 391, 270–273, 1998.
- Dugdale, R. C., Wischmeyer, A. G., Wilkerson, F. P., Barber, R. T., Chai, F., Jiang, M.-S., and Peng, T.-H.: Meridional asymmetry of source nutrients to the equatorial Pacific upwelling ecosystem and its potential impact on ocean–atmosphere CO₂ flux; a data and modeling approach, *Deep Sea Research Res. Part II: Topical Studies in Oceanography*, 49, 2513–2531, 2002
- Dugdale, R.C., Wilkerson F.P.: Silicate regulation of new production in the equatorial Pacific upwelling. *Nature*. 391, 270–273. doi: 10.1038/34630, 1998.
- Eldridge, M.L., Trick, C., Alm, MB, DiTullio, et al *Top. Stud. Oceanogr.*, 49, 2513–2531, 2002.
- Eldridge, M. L., Trick, C. G., Alm, M. B., DiTullio, G. R., Rue, E. L., Bruland, K. W., Hutchins, D. A., and Wilhelm, S. W.: Phytoplankton community response to a manipulation of bioavailable iron in HNLC waters of the subtropical Pacific Ocean. *Aquatic Microbial Ecology*, 35, 79–91. doi: 10.3354/ame035079, *Aquat. Microb. Ecol.*, 35, 79–91, 2004.
- England, M.H.M. H., McGregor, S., Spence, P., Meehl, G. A., et al Timmermann, A., Cai, W., Gupta, A. Sen, McPhaden, M. J., Purich, A., and Santoso, A.: Recent intensification of wind-driven circulation in the Pacific and the ongoing warming hiatus. *Nature Climate Change*, 4, 222–227. doi: 10.1038/nclimate2106, *Nat. Clim. Chang.*, 4, 222–227, 2014.
- Falkowski, P.G.P. G., Barber, R.T., R. T., and Smetacek, V.: Biogeochemical controls and feedbacks on ocean primary production. *Science*. 281, 200–206. doi: Controls and Feedbacks on Ocean Primary Production, Science (80-), 281, 200–206, <https://doi.org/10.1126/science.281.5374.200>, 1998.

Firme, ~~G.F.G. F.~~, Rue, ~~E.L.E. L.~~, Weeks, ~~D.A.D. A.~~, Bruland, ~~K.W.K. W.~~, and Hutchins, D. A.: Spatial and temporal variability in phytoplankton iron limitation along the California coast and consequences for Si, N, and C biogeochemistry. ~~Global Biogeochemical~~, Global Biogeochem. Cycles, 17, 1016. doi: ~~10.1029/2001GB001824~~, 2003.

~~Fu, W., Randerson, J.T., Moore, J.K~~2003.

Fu, W. and Wang, W.: Biogeochemical Equilibrium Responses to Maximal Productivity in High Nutrient Low Chlorophyll Regions, J. Geophys. Res. Biogeosciences, 127, e2021JG006636, 2022.

Fu, W., Randerson, J. T., and Moore, J. K.: Climate change impacts on net primary production (NPP) and export production (EP) regulated by increasing stratification and phytoplankton community structure in the CMIP5 models. Biogeosciences, 13, 5151–5170. doi: ~~10.5194/bg-13-5151-2016~~, 2016, 2016.

~~Fu, W., Wang, W. L.:~~ Biogeochemical equilibrium responses to maximal productivity in high nutrient low chlorophyll regions. Journal of Geophysical Research: Biogeosciences, 127, e2021JG006636. <https://doi.org/10.1029/2021JG006636>, 2022

~~Gargett, A. E.:~~ Physical processes and the maintenance of nutrient-rich euphotic zones. Limnology and Oceanography, 36(8), 1527–1545, 1991

~~Garnesson P., Mangin A., Fanton d'Andon O., Demaria J., Bretagnon M.:~~ The CMEMS GlobColour chlorophyll a product based on satellite observation: multi sensor merging and flagging strategies. Ocean Sci., 15 (3), pp. 819–830, 2019–5170, 2016.

Garnesson, P., Mangin, A., Fanton d'Andon, O., Demaria, J., and Bretagnon, M.: The CMEMS GlobColour chlorophyll a product based on satellite observation: Multi-sensor merging and flagging strategies, Ocean Sci., 15, 819–830, 2019.

Gordon, R. M., Coale, K. H., and Johnson, K. S.: Iron distributions in the equatorial Pacific: Implications for new production, Limnol. Oceanogr., 42, 419–431, 1997.

~~Green, B., Marshall, J., &and Donohoe, A.:~~ Twentieth century correlations between extratropical SST variability and ITCZ shifts. ~~Geophysical Research Letters~~, Geophys. Res. Lett., 44, 9039–9047, doi: ~~10.1002/2017GL075044~~, 2017.

~~Grinsted, A., Moore, J.C.,J. C., and Jevrejeva, S.:~~ Application of the cross wavelet transform and wavelet coherence to geophysical time series. Nonlinear Processes in Geophysics, 11, 561–566. doi: Nonlinear Process. Geophys., 11, 561–566, <https://doi.org/10.5194/npg-11-561-2004>, 2004.

~~Hammond, M.L.M. L., Beaulieu, C., Sahu, S.K.S. K., and Henson, S.A.S. A.:~~ Assessing trends and uncertainties in satellite-era ocean chlorophyll using space-time modeling. ~~Global Biogeochemical-time modeling~~, Global Biogeochem. Cycles, 31, 1103–1117. doi: ~~10.1002/2016GB005600~~, 2017.

~~Harrison, P. J., Whitney, F. A., Tsuda, A., Saito, H., &and Tadokoro, K.:~~ Nutrient and plankton dynamics in the NE and NW gyres of the subarctic Pacific Ocean. ~~Journal of Oceanography~~, J. Oceanogr., 60, 93–117, 2004.

Hauck, J., Völker, C., Wolf-Gladrow, D. A., Laufkötter, C., Vogt, M., Aumont, O., [Bopp, L., Buitenhuis, E. T., Doney, S. C., and Dunne, J.](#): On the Southern Ocean CO₂ uptake and the role of the biological carbon pump in the 21st century. ~~*Global Biogeochemical Cycles*, 29(9), 1451–1470, 2015.~~ [Global Biogeochem. Cycles](#), 29(9), 1451–1470, 2015.

Hernández-Carrasco, I., [and](#) Orfila, A.: The Role of an Intense Front on the Connectivity of the Western Mediterranean Sea: The Cartagena-Tenes Front. ~~*Journal of Geophysical Research: Oceans*, 123(6), J. Geophys. Res. Ocean.~~, 123, 4398–4422, 2018.

Holzer, M., DeVries, T., [and](#) de Lavergne, C.: Diffusion controls the ventilation of a Pacific Shadow Zone above abyssal overturning. *Nat. Commun.*, 12, 4348, 2021

~~Howarth D.A.: An analysis of the variability of cyclones around Antarctica and their relationship to sea-ice extent. *Annals of the Association of American Geographers*, 73, 519–537. doi:10.1111/j.1467-8306.1983.tb01856.x, 1983~~

~~Hutchins, D.A., DiTullio, G.R., Zhang, Y., Bruland, K.W.: An iron limitation mosaic in the California upwelling regime. *Limnology and Oceanography*, 43, 1037–1054. doi:10.4319/lo.1998.43.6.1037, 1998.~~

~~Hutchins D.A., Hare C.E., Weaver R.S., Zhang Y., et al.: Phytoplankton iron limitation in the Humboldt Current and Peru Upwelling. *Limnology and Oceanography*, 47, 997–1011. doi: 10.4319/lo.2002.47.4.0997, 2002.~~

~~Ibanhez, J.S.P.~~

[Hutchins, D. A., DiTullio, G. R., Zhang, Y., and Bruland, K. W.](#): An iron limitation mosaic in the California upwelling regime. *Limnol. Oceanogr.*, 43, 1037–1054, 1998.

[Hutchins, D. A., Hare, C. E., Weaver, R. S., Zhang, Y., Firme, G. F., DiTullio, G. R., Alm, M. B., Riseman, S. F., Maucher, J. M., and Geesey, M. E.](#): Phytoplankton iron limitation in the Humboldt Current and Peru Upwelling. *Limnol. Oceanogr.*, 47, 997–1011, 2002.

[Ibánhez, J. S. P.](#), Flores, M., [and](#) Lefèvre, N.: Collapse of the tropical and subtropical North Atlantic CO₂ sink in boreal spring of 2010. *Sci. Rep.*, 7, 41694. doi: 10.1038/srep41694-9, 2017.

Imai, K., Nojiri, Y., Tsurushima, N., [and](#) Saino, T.: Time series of seasonal variation of primary productivity at station KNOT (44N, 155E) in the sub-arctic western North Pacific. *Deep Sea Research Part II: Topical Studies in Oceanography*, 49, 5395–5408. doi: 10.1016/S0967-0645(02)00198-4, 2002.

~~Kaupp, L. J., Measures, C. I., Selph, K. E., & Mackenzie, F. T.: The distribution of dissolved Fe and Al in the upper waters of the Eastern Equatorial Pacific. *Deep Sea Research Part II: Topical Studies in Oceanography*, 58(3–4), 296–310, 2011. *Top. Stud. Oceanogr.*, 49, 5395–5408, 2002.~~

Ishizaki, H.: A simulation of the abyssal circulation in the North Pacific Ocean. Part II: Theoretical rationale. ~~*Journal of physical oceanography*, 24(9), 1941–~~ *J. Phys. Oceanogr.*, 24, 1941–1954, 1994.

Koch, F., Marcoval, M. A., Panzeca, C., Bruland, K. W., ~~et al~~ Sañudo-Wilhelmy, S. A., and Gobler, C. J.: The ~~effectseffect~~ of vitamin B12 on phytoplankton growth and community structure in the Gulf of Alaska. Limnology and Oceanography, 3, 1023–1034. doi: 10.4319/lo.2011.56.3.1023, 2011.

~~Kohonen, T.: Self-organized formation of topologically correct feature maps. Biol. Cybern. 43, 59–69. doi:10.1007/BF00337288, Limnol. Oceanogr., 56, 1023–1034, 2011.~~

Kohonen, T.: Analysis of a simple self-organizing process, Biol. Cybern., 44, 135–140, 1982.

~~Kumar, N., Anderson R.F., Mortlock R.A., Froelich P.N., R. F., Mortlock, R. A., Froelich, P. N., Kubik, P.,~~ Dittrich-Hannen, B., and Suter, M.: Increased biological productivity and export production in the glacial Southern Ocean. Nature. Ocean, Nature, 378, 675–680. doi: 10.1038/378675a0, 1995

~~Kuragano, T., & Kamachi, M.: Balance of volume transports between horizontal circulation and meridional overturn in the North Pacific subarctic region. Journal of oceanography, 60, 439–451, 2004~~

Landry M.R., Selph K.E., Taylor A.G., 1995.

Lam, P. J., Bishop, J. K. B., Henning, C. C., Marcus, M. A., Waychunas, G. A., and Fung, I. Y.: Wintertime phytoplankton bloom in the subarctic Pacific supported by continental margin iron, Global Biogeochem. Cycles, 20, 2006.

Landry, M. R., Selph, K. E., Taylor, A. G., Décima, M., ~~et al~~ Balch, W. M., and Bidigare, R. R.: Phytoplankton growth, grazing and production balances in the HNLC equatorial Pacific. Deep-sea Res. II, 58, 524–535. doi: 10.1016/j.dsr2.2010.08.011, Deep Sea Res. Part II Top. Stud. Oceanogr., 58, 524–535, 2011.

Lauderdale, ~~J.M.J. M.~~, Braakman, R., Forget, G., Dutkiewicz, S., and Follows, ~~M.J. M. J.~~: Microbial feedbacks optimize ocean iron availability. Proceedings of the National Academy of Sciences, 117(9), 4842–4849. doi: 10.1073/pnas.1917277117, Proc. Natl. Acad. Sci., 117, 4842–4849, 2020.

Liu, Y., ~~R.H. Weisberg~~ Weisberg, R. H., Vignudelli, S. Vignudelli, and ~~G.T. Mitchum~~ Mitchum, G. T.: Patterns of the Loop Current loop current system and regions of sea surface height variability in the eastern Gulf of Mexico revealed by the self-organizing maps, Journal of Geophysical Research Oceans, 121, 2347–2366, <http://dx.doi.org/10.1002/2015JC011493>, 2016. J. Geophys. Res. Ocean., 121, 2347–2366, 2016.

Di Lorenzo, E., Schneider, N., Cobb, K. M., Franks, P. J. S., Chhak, K., Miller, A. J., McWilliams, J. C., Bograd, S. J., Arango, H., and Curchitser, E.: North Pacific Gyre Oscillation links ocean climate and ecosystem change, Geophys. Res. Lett., 35, 2008.

Di Lorenzo, E., Combes, V., Keister, J. E., Strub, P. T., Thomas, A. C., Franks, P. J. S., Ohman, M. D., Furtado, J. C., Bracco, A., and Bograd, S. J.: Synthesis of Pacific Ocean climate and ecosystem dynamics, 26, 68–81, 2013.

Lumpkin, R., ~~& and~~ Speer, K.: Global ocean meridional overturning. Journal of Physical Oceanography, 37(10), J. Phys. Oceanogr., 37, 2550–2562, 2007.

Maritorena, S., ~~& and~~ Siegel, D. A.: Consistent merging of satellite ocean color data sets using a bio-

optical model. [Remote Sensing of Environment, 94\(4\), Remote Sens. Environ., 94, 429–440, 2005.](#)

[Martin, J.H., Fitzwater, S.E. H.: Glacial-interglacial CO2 change: The iron hypothesis, Paleoclimatology, 5, 1–13, 1990.](#)

[Martin, J. H. and Fitzwater, S. E.: Iron deficiency limits phytoplankton growth in the northeast north-east Pacific subarctic. Nature, Nature, 331, 341–343. doi: https://doi.org/10.1038/331341a0, 1988.](#)

[Martin, J.H.: Glacial-interglacial CO2 change: the iron hypothesis. Paleoclimatology, 5, 1–13. doi: 10.1029/PA005i001p00001, 1990.](#)

[Martin, J.H., Fitzwater, S.E., Gordon, R.M.: Iron deficiency limits phytoplankton growth in Antarctic waters. Global Biogeochemical Cycles, 4, 5–12. doi:10.1029/GB004i001p00005. J. H., Fitzwater, S. E., and Gordon, R. M.: Iron deficiency limits phytoplankton growth in Antarctic waters, Global Biogeochem. Cycles, 4, 5–12, 1990.](#)

[Martin, J.H. H., Coale, K.S. H., Johnson, K.S. S., Fitzwater, S.E. et al.: S. E., Gordon, R. M., Tanner, S. J., Hunter, C. N., Elrod, V. A., Nowicki, J. L., and Coley, T. L.: Testing the iron hypothesis in ecosystems of the equatorial Pacific Ocean. Nature, Nature, 371, 123–129. doi:10.1038/371123a0, 1994.](#)

[Martínez-García-García, A., Rosell-Melé-Melé, A., Geibert, W., Gersonde, R., et al. Masqué, P., Gaspari, V., and Barbante, C.: Links between iron supply, marine R-productivity, sea surface temperature, and CO2 over the last 1.1 Ma. Paleoclimatology, 24, PA1207. doi: 10.1029/2008PA001657, 2009, Paleoclimatology, 24, 2009.](#)

[Martinez, E., Gorgues, T., Lengaigne, M., Fontana, C., Sauzède, R., Menkes, C., Uitz, J., Di Lorenzo, E., and Fablet, R.: Reconstructing global chlorophyll-a variations using a non-linear statistical approach, Front. Mar. Sci., 7, 464, 2020.](#)

[McConnell, J. R., Arístarain, A. J., Banta, J. R., Edwards, P. R., et al. and Simões, J. C.: 20th-Century doubling in dust archived in an Antarctic Peninsula ice core parallels climate change and desertification in South America. Proceedings of the National Academy of Sciences, 104, 5743–5748. doi: 10.1073/pnas.0607657104. McPhaden, M.J., Proc. Natl. Acad. Sci., 104, 5743–5748, 2007.](#)

[McPhaden, M. J.: Trade wind fetch-related variations in Equatorial Undercurrent equatorial undercurrent depth, speed, and transport. Journal of Geophysical Research: Oceans, 98\(C2\), 2555–2559. doi: 10.1029/92JC02683, J. Geophys. Res. Ocean., 98, 2555–2559, 1993.](#)

[McPhaden, M. J., & Zhang, D.: Slowdown of the meridional overturning circulation in the upper Pacific Ocean. Nature, 415\(6872\), 603–608, 2002.](#)

[Mélin, F., Vantrepotte, V., Chuprin, A., Grant, M., Jackson, T., & Sathyendranath, S.: Assessing the fitness-for-purpose of satellite multi-mission ocean color climate data records: A protocol applied to OC-CCI chlorophyll-a data. Remote Sensing of Environment, Remote Sens. Environ., 203, 139–151, 2017.](#)

[Minas, H. J., Minas, M., and Packard, T.T. T.: Productivity in upwelling areas deduced from hydrographic and chemical fields. Limnology and Oceanography, 31, 1182–1206. doi: 10.4319/lo.1986.31.6.1182](#)

1, *Limnol. Oceanogr.*, 31, 1182–1206, 1986.

Moat B.I., B., Frajka-Williams, E., Smeed D.A., D., Rayner, D., Johns W.E., Baringer M.O., Johns, W. E., Baringer, M. O., Volkov, D., and Collins, J.: Atlantic meridional overturning circulation observed by the RAPID-MOCHA WBTS (RAPID Meridional Overturning Circulation and Heatflux Array Western Boundary Time Series) array at 26N from 2004 to 2020. British Oceanographic Data Centre—Natural Environment Research Council, UK. doi:10.5285/e91b10af-6f0a-7fa7-e053-6e86abe05a09, 2022

Moat, B. I., Smeed, D. A., Frajka-Williams, E., Desbruyères, D. G., Beaulieu, C., Johns, W. E., ... & Bryden, H. L.: Pending recovery in the strength of the meridional overturning circulation at 26 N. *Ocean Science*, 16(4), 863–874, 2020

Moore, J.K., Doney, S.C., 2007. RAPID-MOC time series April 2004 to December 2020 (v2020. 2), 2022.

Moore, C. M., Mills, M. M., Arrigo, K. R., Berman-Frank, I., Bopp, L., Boyd, P. W., Galbraith, E. D., Geider, R. J., Guieu, C., and Jaccard, S. L.: Processes and patterns of oceanic nutrient limitation, *Nat. Geosci.*, 6, 701–710, 2013.

Moore, J. K. and Doney, S. C.: Iron availability limits the ocean nitrogen inventory stabilizing feedbacks between marine denitrification and nitrogen fixation. *Global Biogeochemical Cycles*, 21, GB2001. doi:10.1029/2006GB002762, 2013

Moore, C.M., Mills, M.M., Arrigo, K.R., Berman Frank, I., et al.: Processes and patterns of oceanic nutrient limitation. *Nature Geoscience*, 6, 701–710. doi: 10.1038/ngeo1765, 2013

Moore, J.K., Fu, W., Primeau, F., Britten, G.L. et al.: Sustained climate warming drives declining marine biological productivity. *Science*, 359 (6380), 1139–1143. doi: 10.1126/science.aao6379, 2017.

Moore, J. K., Fu, W., Primeau, F., Britten, G. L., Lindsay, K., Long, M., Doney, S. C., Mahowald, N., Hoffman, F., and Randerson, J. T.: Sustained climate warming drives declining marine biological productivity. *Science* (80-.), 359, 1139–1143, 2018.

Moradi, M.: Evaluation of merged multi-sensor ocean-color chlorophyll products in the Northern Persian Gulf. *Continental Shelf Research*, 221, 104415, 2021.

Muglia, J., Skinner, L. C., and Schmittner, A.: Weak overturning circulation and high Southern Ocean nutrient utilization maximized glacial ocean carbon. *Earth and Planetary Science Letters*, 496, 47–56, 2018.

Nielsdóttir, M.C.M. C., Moore, C.M.C. M., Sanders, R., Hinz D.J. et al, D. J., and Achterberg, E. P.: Iron limitation of the postbloom phytoplankton communities in the Iceland Basin, *Global Biogeochemical Cycles*, 23, GB3001, doi:10.1029/2008GB003410, 2009.

Nishioka, J., Ono, T., Saito, H., Nakatsuka, T., Takeda, S., Yoshimura, T., Suzuki, K., Kuma, K., Nakabayashi, S., and Tsumune, D.: Iron supply to the western subarctic Pacific: Importance of iron export from the Sea of Okhotsk, *J. Geophys. Res. Ocean.*, 112, 2007.

Nishioka, J., ~~Obata, H., Ogawa, H., Ono, K., et al.~~ Yamashita, Y., Lee, K., Takeda, S., and Yasuda, I.: Subpolar marginal seas fuel the North Pacific through the intermediate water at the termination of the global ocean circulation. Proc. Natl. Acad. Sci., 117(23), pp.12665–12673, 10.1073/pnas.2000658117, 117, 12665–12673, 2020.

Nishioka, J., et al.: ~~Iron supply to the western subarctic Pacific: Importance of iron export from the Sea of Okhotsk~~, J. Geophys. Res., 112, C10012, doi:10.1029/2006JC004055, 2007 Obata, H., Hirawake, T., Kondo, Y., Yamashita, Y., Misumi, K., and Yasuda, I.: A review: iron and nutrient supply in the subarctic Pacific and its impact on phytoplankton production, J. Oceanogr., 77, 561–587, 2021.

Oliver, M. J. and Irwin, A. J.: Objective global ocean biogeographic provinces, Geophys. Res. Lett., 35, 2008.

Ono, T., Shiomoto, A., ~~Saino~~ and Saino, T.: Recent decrease of summer nutrients concentrations and future possible shrinkage of the subarctic North Pacific high ~~nutrient low chlorophyll region~~. Global Biogeochemical Cycles, 22, GB3027. doi:10.1029/2007GB003092, 2008.

~~Oliver, M. J., & Irwin, A. J.~~: Objective global ocean biogeographic provinces. Geophysical research letters, 35(15), 2018 nutrient low-chlorophyll region, Global Biogeochem. Cycles, 22, 2008.

Orsi, A. H., Whitworth III, T., and Nowlin Jr, W. D.: On the meridional extent and fronts of the Antarctic Circumpolar Current. Part I Oceanogr. Res. Papers, 42, 641–673. doi: 10.1016/0967-0637(95)00021-W, 1995.

Parekh, P., Follows, ~~M.J.~~ M. J., and Boyle, ~~E.A.E.~~ E. A.: Decoupling of iron and phosphate in the global ocean. Global Biogeochemical, Global Biogeochem. Cycles, 19, GB2020, doi:10.1029/2004GB002280, 2005.

~~Pitchford, J.W., Brintley, J.~~: Iron limitation, grazing pressure and oceanic high nutrient low chlorophyll (HNLC) regions. Journal of Plankton Research, 21, 525–547. doi: 10.1093/plankt/21.3.525, 1999 2005.

Philander, S. G. H. and Chao, Y.: On the contrast between the seasonal cycles of the equatorial Atlantic and Pacific Oceans, J. Phys. Oceanogr., 21, 1399–1406, 1991.

Pollard, R. T., Salter, I., Sanders, R. ~~et al.~~ J., Lucas, M. I., Moore, C. M., Mills, R. A., Statham, P. J., Allen, J. T., Baker, A. R., and Bakker, D. C. E.: Southern Ocean deep-water carbon export enhanced by natural iron fertilization. Nature, 457, 577–580, doi: 10.1038/nature07716, 2009.

Polovina, ~~J.J.~~ J. J., Howell, ~~E.A.E.~~ E. A., and Abecassis, M.: ~~Ocean's~~ Ocean's least productive waters are expanding. Geophysical Research Letters, 35. doi: 10.1029/2007GL031745, Geophys. Res. Lett., 35, 2008.

Qiu, B.: Large-scale variability in the midlatitude subtropical and subpolar North Pacific Ocean: Observations and causes. Journal of Physical Oceanography, 32, 353–375. doi: 10.1175/1520-0485(2002)032<0353:LSVITM>2.0.CO;2, J. Phys. Oceanogr., 32, 353–375, 2002.

Radenac, M.-H., Léger, F., Singh, A., and Delcroix, T.: Sea surface chlorophyll signature in the tropical Pacific during eastern and central Pacific ENSO events. Journal of Geophysical Research: Oceans, 117, C04007. doi: 10.1029/2011JC007841, 2012.

Riebesell, U., Körtzinger, A., Oeschlies, A.: Sensitivities of marine carbon fluxes to ocean change. Proc. Natl. Acad. Sci. USA. doi: 10.1073/pnas.0813291106, 2009

Serno, S., Winckler G., Anderson, R.F., Hayes, C.T. et al.: Eolian dust input to the Subarctic North Pacific. Earth and Planetary Science Letters. 387, 252–263. doi:10.1016/j.epsl.2013.11.008, 2014

Sen, P.K.: Estimates of the regression coefficient based on Kendall's Tau Pranab Kumar Sen. Journal of the American Statistical Association. 63, 1379–1389, 1968, J. Geophys. Res. Ocean., 117, 2012.

Rafter, P. A., Sigman, D. M., and Mackey, K. R. M.: Recycled iron fuels new production in the eastern equatorial Pacific Ocean, Nat. Commun., 8, 1100, 2017.

Riebesell, U., Körtzinger, A., and Oeschlies, A.: Sensitivities of marine carbon fluxes to ocean change, Proc. Natl. Acad. Sci., 106, 20602–20609, 2009.

Schmidt, K., Schlosser, C., Atkinson, A., Fielding, S., Venables, H. J., Waluda, C. M., &and Achterberg, E. P.: Zooplankton gut passage mobilizes lithogenic iron for ocean productivity. Current Biology, 26(19), 2667–2673, 2016

Schmittner, A.: Decline of the marine ecosystem caused by a reduction in the Atlantic overturning circulation. Nature. 434(7033), 628–33. doi: 10.1038/nature03476, 2005, Curr. Biol., 26, 2667–2673, 2016.

Schneider, N., Miller, A. J., Alexander, M. A., &and Deser, C.: Subduction of decadal North Pacific temperature anomalies: Observations and dynamics. Journal of Physical Oceanography, 29(5), J. Phys. Oceanogr., 29, 1056–1070, 1999.

Sen, P. K.: Estimates of the regression coefficient based on Kendall's tau, J. Am. Stat. Assoc., 63, 1379–1389, 1968.

Smed, D. A., Josey, S. A., Beaulieu, C., Johns, W. E., Moat, B. I., Frajka-Williams, E., et al Rayner, D., Meinen, C. S., Baringer, M. O., and Bryden, H. L.: The North Atlantic Oceanic Ocean is in a state of reduced overturning. Geophysical Research Letters, 45, 1527–1533. https://doi.org/10.1002/2017GL076350 overturning, Geophys. Res. Lett., 45, 1527–1533, 2018.

Smed, D. A., Moat, B. I., Rayner, D., Johns, W. E. et al, Baringer M. O., Volkov D. L., Frajka-Williams E.: Atlantic meridional overturning circulation observed by the RAPID-MOCHA-WBTS (RAPID-Meridional Overturning Circulation and Heatflux Array-Western Boundary Time Series) array at 26N from 2004 to 2018. British Oceanographic Data Centre—Natural Environment Research Council, National Oceanography Centre, NERC, UK. doi:10.5285/8cd7e7bb-9a20-05d8-e053-6c86abc012c2, 2019.

Strutton, P. G. and Chavez, F. P.: Primary productivity in the equatorial Pacific during the 1997–1998 El Niño, J. Geophys. Res. Ocean., 105, 26089–26101, 2000.

- Tagliabue, A., Aumont, O., and Bopp, L.: The impact of different external sources of iron on the global carbon cycle. Geophysical Research Letters, 41, 920–926. doi: 10.1002/2013GL059059, Geophys. Res. Lett., 41, 920–926, 2014.
- Takeda, S.: Iron and phytoplankton growth in the Subarctic North Pacific. Aqua-BioScience Monographs, 4, 41–93, 2011.
- Tandon, N. F., Saenko, O. A., Cane, M. A., & Kushner, P. J.: Interannual variability of the global meridional overturning circulation dominated by Pacific variability. Journal of Physical Oceanography, 50(3), 559–574, 2020
- Timmermann, A., Okumura, Y., An, S.-I., Clement, A., Dong, B., Guilyardi, E., et al, Terrapub, 2011.
- Timmermann, A., Okumura, Y., An, S.-I., Clement, A., Dong, B., Guilyardi, E., Hu, A., Jungclaus, J. H., Renold, M., and Stocker, T. F.: The influence of a weakening of the Atlantic meridional overturning circulation on ENSO. Journal of climate, 20(19), J. Clim., 20, 4899–4919, 2007.
- Torrence, C., and Compo, G. P.: A practical guide to wavelet analysis. Bulletin of the American Meteorological society, 79, 61–78. doi: 10.1175/1520-0477, Bull. Am. Meteorol. Soc., 79, 61–78, 1998.
- Torrence, C., and Webster, P. J.: Interdecadal changes in the ENSO–monsoon system. Journal of Climate, 12, 2679–2710. doi: 10.1175/1520-0442, J. Clim., 12, 2679–2690, 1999.
- Tsuda, A., Takeda, S., Saito, H., Nishioka, J., et al Nojiri, Y., Kudo, I., Kiyosawa, H., Shiimoto, A., Imai, K., and Ono, T.: A mesoscale iron enrichment in the western subarctic Pacific induces a large centric diatom bloom. Science, 300, 958–961. doi:10.1126/science.1082000 (80-), 300, 958–961, 2003.
- Tyrrell Vatanen, T., Merico, A., Waniek, J.J., Wong, et al.: Effect of seafloor depth on phytoplankton blooms in high nitrate, low chlorophyll (HNLC) regions. Journal of Geophysical Research: Biogeosciences, 110, G02007, doi: 10.1029/2005JG000041, 2005.
- Vatanen, T., Osmala Osmala, M., Raiko, T., Lagus, K., Sysi-Aho, M., Orešič, M., Raiko Honkela, T., Lagus, K. et al and Lähdesmäki, H.: Self-organization and missing values in SOM and GTM. Neurocomputing, 147, 60–70. doi: 10.1016/j.neucom.2014.02.061, 2015.
- Vesanto, J., Himberg, J., Alhoniemi, E., and Parhankangas, J.: Self-organizing map in Matlab: the SOM Toolbox. In, in: Proceedings of the Matlab DSP conference, Espoo, Finland, Comsol Oy 16–17, 1999.
- Warren, B. A.: Why is no deep water formed in the North Pacific?. Journal of Marine Research, 41(2), J. Mar. Res., 41, 327–347, 1983.
- Winckler, G., Anderson, R. F., Jaccard, S. L., and Marcantonio, F.: Ocean dynamics, not dust, have controlled equatorial Pacific productivity over the past 500,000 years. Proceedings of the National Academy of Sciences, 113, 6119–6124. doi:10.1073/pnas.1600616113, Proc. Natl. Acad. Sci., 113, 6119–6124, 2016.
- Yaremchuk, M., Bindoff, N. L., Schröter, J., Nechaev, D., &and Rintoul, S. R.: On the zonal and meridional circulation and ocean transports between Tasmania and Antarctica. Journal of Geophysical Research: Oceans, 106(C2), J. Geophys. Res. Ocean., 106, 2795–2814, 2001.

Yasunaka, S., Nojiri, Y., Nakaoka, S., Ono, T., ~~et al.~~, [Whitney, F. A., and Telszewski, M.](#): Mapping of sea surface nutrients in the ~~North Pacific Basin~~ [North Pacific Basin-wide](#) distribution and seasonal to interannual variability. ~~journal of geophysical research: oceans, 119, 7756–7771. doi: 10.1002/2014JC010318, J. Geophys. Res. Ocean., 119, 7756–7771,~~ 2014.

Yasunaka, S., Ono, T., Nojiri, Y., Whitney, ~~F.A., et al.~~ [F. A., Wada, C., Murata, A., Nakaoka, S., and Hosoda, S.](#): Long-term variability of surface nutrient concentrations in the North Pacific, ~~Geophysical Research Letters, 43, 3389–3397. doi:10.1002/2016GL068097~~ [Geophys. Res. Lett., 43, 3389–3397,](#) 2016.

Tables

Table 1. Mean \pm std characteristics of each of the SOM-defined HNLC subregions (R1 to R5). NO₃ (mmol m⁻³) and Chl (mg m⁻³) values are respectively from model and satellite data. Decadal chlorophyll trends (Δ Chl, mgChl m⁻³ decade⁻¹) are calculated from the mean time-series of monthly deseasonalized chlorophyll.

Region	Subregion	NO ₃ (μ M)	Chl (mg m ⁻³)	NO ₃ :Chl (mmol NO ₃ mg Chl ⁻¹)	Δ Chl (mg m ³ decade ⁻¹)
SNP	R1	4.51 \pm 1.02	0.31 \pm 0.07	15 \pm 3	+0.05
	R2	8.05 \pm 0.88	0.36 \pm 0.07	23 \pm 6	+0.26
	R3	15.52 \pm 2.27	0.49 \pm 0.16	35 \pm 15	+0.43
EEP	R1	4.04 \pm 0.77	0.22 \pm 0.02	18 \pm 3	+0.01
	R2	6.63 \pm 1.42	0.39 \pm 0.05	20 \pm 4	+0.08
SO	R1	4.13 \pm 1.05	0.22 \pm 0.06	20 \pm 4	+0.24
	R2	9.11 \pm 1.23	0.31 \pm 0.06	31 \pm 9	+0.42
	R3	15.73 \pm 1.07	0.32 \pm 0.10	55 \pm 17	+0.47
	R4	23.26 \pm 1.06	0.26 \pm 0.16	104 \pm 32	+0.62
	R5	29.18 \pm 1.57	0.43 \pm 0.92	103 \pm 54	+0.46

Table 2. Basic statistics of the annual extent of each of the SOM-defined HNLC subregions during the analyzed period (1998-2017). Maximum variation (Max. Variation) is calculated as the difference between the maximum and the minimum extent.

Regio	Mean extent±S.D.	% of total	Min-Max	Max. Variation	C.V.
n	(x10 ⁶ km ²)	extent	(x10 ⁶ km ²)	(x10 ⁶ km ²)	%
SNP	7.7±3.6	8.4	3.8-15.9	12.1	47
EEP	7.8±0.4	8.4	7.3-8.4	1.1	5
SO	76.5±0.9	83.2	61.3-86.8	25.5	12

Figures




Figure 1. Scheme of a characteristic pattern and its distribution map obtained from SOM time-domain analysis, and global and regional mean series calculated for each of the HNLC regions.









Figure 2. a) Global mean satellite Chl concentrations (mg m^{-3}) for the period 1998-2018 and superimposed mean surface NO_3 contour lines from modeling data. b) Normalized probability density function of the values of the $\text{NO}_3:\text{Chl}$ ratio obtained from the SOM temporal patterns. Green and blue lines show the fit

to a normal distribution for the first and second pdf modes, respectively. c) Standard deviation of the probability density function (*pdf*) of the NO₃:Chl (mmol/mg) monthly ratios obtained for the 20 years analyzed. Note that the y-axis scale is logarithmic. The critical point ratio $P_{\text{HNLC}} = 17 \text{ mmolNO}_3 \text{ mgChl}^{-1}$ delimits HNLC regions from macronutrient limited regions.







Figure 3. a) Characteristic temporal variability patterns of NO₃:Chl ratios (N1-N12) as unveiled by the 4x3 SOM analysis in the time domain. Red dashed lines indicate the P_{HNLC} value b) Coherent regions of HNLC variability (SNP, EEP, and SO) and corresponding subregions (R1 to R5) associated with the SOM temporal patterns exhibiting only NO₃:Chl values larger than P_{HNLC} all the times throughout the entire analyzed period (i.e. identified with N9, N6, N2, N5 and N1). Patterns corresponding to a subregion in the northern and southern hemispheres present a similar pattern although seasonally lagged (6-month delay). Insets show the time series of the averaged Chl over the corresponding subregion (complete map of regions of NO₃:Chl variability and corresponding temporal variability patterns are shown in Fig. S1).

The red line represents the 24-month filtered series and the blue line indicates the trend (values shown in Table 1).

Figure 4. Characteristic spatial patterns (P1 to P9) of HNLC regions as defined by $\text{NO}_3:\text{Chl}$ ratios $> P_{\text{HNLC}}$ obtained from monthly data. The value on top of each pattern indicates its probability of occurrence over the 20-year period analyzed.



To preserve the topology, the SOM algorithm introduces some patterns with zero probability of occurrence, such as P5. The colorbar indicates the different $\text{NO}_3:\text{Chl}$ ranges represented.












Figure 5. a) Time evolution of the spatial patterns as defined by the Best Matching Units (BMUs) for the period of 1998-2018. The blue dashed line indicates the regime shift occurring after 2010. b) Monthly frequency of occurrence of the spatial patterns identified in Figure 4.



Figure 6. Seasonal (left) and interannual variations (right) in the spatial extent of the three HNLC regions, represented as a percentage of variation from the mean extent of each region. Variations are referred to the mean extent of each region. Dark and light-colored bars indicated positive and negative values, respectively. The blue dashed lines indicate the regime shift occurring after 2010.



Year




Figure 7. a) ENSO (MEI v2) index. E1-E3 indicate intense El Niño episodes and L1 and L2 mark strong La Niña periods. b, c, d display the cross-wavelet coherence between NO₃:Chl ratio and ENSO for each of the HNLC regions. e) Meridional overturning (AMOC) volume transport for the period (2004-2017) measured at 26.5°N (Smeed et al., 2016). The red line shows interannual component is obtained by filtering the data with a 540-day low-pass filter after the removal of the mean seasonal cycle. f, g, and h display the cross-wavelet coherence between NO₃:Chl ratio and AMOC for each of the HNLC regions. The thick black contours in the cross-wavelet coherence figures designate the 95% confidence levels and

the cone of influence where edge effects are not negligible is shown as a lighter shade. The arrows indicate the phase relationship between the signals with the horizontal component indicating in-phase (rightward) or out-of-phase (leftward) and the vertical component indicating a 90° phase difference lagging (upward) or leading (downward). Period units are months.