

An analogues based forecasting system for Mediterranean marine litter concentration

Gabriel Jordà^{1,2,*} and Javier Soto-Navarro^{3,4,*}

¹Centre Oceanogràfic de les Balears, Spanish Institute of Oceanography (CN-IEO/CSIC).
5 Mallorca, 07015, Spain.

²University of the Balearic Islands (UIB). Mallorca, 07122, Spain.

³Physical Oceanography Group of the University of Málaga (GOFIMA). Málaga, 29071, Spain.

⁴Institute of Oceanic Engineering of the University of Málaga (IIO). Málaga, 29071, Spain.

*These authors contributed equally to this work

10 *Correspondence to:* Javier Soto-Navarro (javersoto@uma.es)

Abstract

In this work we explore the performance of a statistical forecasting system for marine litter (ML) concentration in the Mediterranean Sea. In particular, we assess the potential skills of a system based on the analogues method. The system uses a historical database of ML concentration
15 simulated by a high resolution realistic model and is trained to identify meteorological situations in the past that are similar to the forecasted ones. Then, the corresponding ML concentrations of the past analogue days are used to construct the ML concentration forecast. Due to the scarcity of observations, the forecasting system has been validated against a synthetic reality (i.e. the outputs from a ML modelling system). Different approaches have been tested to refine the system and the
20 results show that using integral definitions for the similarity function, based on the history of the meteorological situation, improves the system performance. We also find that the system accuracy depends on the domain of application being better for larger regions. Also, the method performs well to capture the spatial patterns but performs worse to capture the temporal variability, specially the extreme values. Despite the inherent limitations of using a synthetic reality to
25 validate the system, the results are promising and the approach has potential to become a suitable cost effective forecasting method for ML concentration.

1. Introduction

The ubiquity of the plastic waste pollution in seas and oceans worldwide raises great concern in the society and the scientific community, as it poses a significant environmental and
30 socioeconomic threat (UNEP, 2009). In consequence, the analysis of the impacts of marine litter (ML) pollution on the marine life and ecosystems has become a hot topic on marine research in recent years (Maximenko et al., 2019; Van Sebille et al., 2020; Lebreton et al., 2019; Lebreton

and Andrady, 2019; Soto-Navarro et al., 2021). ML particles accumulate both in shallow and deep waters, and particularly in enclosed basins such as the Mediterranean Sea (Soto-Navarro et al., 35 2020; Cózar et al., 2015), where the observed concentrations are in the same range of those measured in the great plastic patches formed in the subtropical gyres of the open oceans (Cózar et al., 2015; Law et al., 2014; Van Sebille et al., 2015). Moreover, risk analyses have shown that marine organisms in the Mediterranean basin can be highly impacted by ML pollution (Compa et al., 2019; Soto-Navarro et al., 2021). The starting point to analyze those impacts and to establish 40 suitable mitigation strategies is to understand the spatial distribution and temporal evolution of the ML particles. Unfortunately, to carry on that analysis solely based on observations is not feasible. The large spatial and temporal heterogeneities of the field campaigns, along with the lack of standardized observational protocols, do not allow a synoptic representation of the ML distribution (see Maximenko et al. (2019) for a thorough analysis of the ML observations 45 problems and proposed improvements). For these reasons, numerical modeling emerges as a fundamental tool to achieve a synoptic description of ML dispersion patterns and as the base for the forecasting systems that would reproduce its spatial variability and time evolution.

ML forecasting systems are usually based on the combination of two different numerical models (Lebreton et al., 2012; Van Sebille et al., 2015; Maximenko et al., 2012). On the one hand, an 50 ocean circulation forecasting system is implemented to provide ocean currents. On the other hand, a lagrangian model uses those currents to simulate the advection and diffusion of passive particles in the ocean that mimic the evolution of ML. In the Mediterranean, several studies using this methodology have been carried out using current fields from high resolution regional models covering the whole basin (Liubartseva et al., 2018; Macias et al., 2019; Mansui et al., 2015; Soto- 55 Navarro et al., 2020) or specific regions such as the Adriatic, the Tyrrhenian or the Aegean (Politikos et al., 2017; Fossi et al., 2017; Liubartseva et al., 2016; Palatinus et al., 2019). This modelling approach is considered to be the most accurate choice for ML forecasting (Van Sebille et al., 2020) provided the ML inputs are correctly prescribed (Liubartseva et al. (2018) , Soto-Navarro et al. (2020)).

60 The downside of developing a forecasting system based on the direct modelling approach is that it involves a high technical complexity and computational cost. In order to overcome this limitations, it might be possible to develop a fast and light forecasting system based on statistical methods. One choice would be the so called Statistical Downscaling Methods (SDMs) which relies on determining statistical relationships between large scale variables (usually atmospheric 65 patterns) and local variables. They are broadly used in atmospheric modelling to forecast the evolution of local variables from large scale atmospheric models. The advantage of the SDMs is that the mathematical relationship derived by the model between the local and the large scale variables is valid not only for the present climate, but can also be used to estimate the future

evolution of the local variables. In summary, the SDMs provide a simplified ‘static’ methodology
70 to forecast the evolution of local variables without the need of running a complex dynamical
models. There are numerous downscaling methodologies based on different statistical properties.
Among them, the analogues method (Lorenz, 1969) is the most broadly used due to its simplicity
and accuracy (Grouillet et al., 2016). This technique assumes that similar (or analogues)
atmospheric patterns over a given region, represented by large scale atmospheric variables or
75 predictors, lead to similar local meteorological outcomes (or predictands) in a particular location.
This assumption provides a simple algorithm to downscale the local occurrence of the variable of
interest from a given large scale atmospheric pattern (see section 2.1 for a detailed description).
In general, it has been shown that the analogues method performs as well as other more
sophisticated downscaling techniques (Zorita and von Storch, 1999), indicating that it is an
80 efficient alternative for many downscaling problems. Its main advantages are that is non-
parametric (i.e. no assumptions are made about the distribution of the variables used as
predictors), non-linear (i.e. it can take into account the non-linearity of the relationships between
predictors and predictands), and it is spatially coherent (i.e., preserves the spatial covariance
structure of the local variables). The analogues method has been satisfactorily applied in the
85 Mediterranean region not only for the downscaling of meteorological or hydrological variables
such as precipitation or river runoff (Grouillet et al., 2016; Wu et al., 2012; Caillouet et al., 2016),
but also for the reconstruction of sea surface temperature in the glacial period (Hayes et al., 2005),
the assimilation of satellite derived sea surface height (Lopez-Radcenco et al., 2019) and the
projection of complex climatic impact indices such as the fire weather index or the physiological
90 equivalent temperature (Casanueva et al., 2014).

In this study, we explore the feasibility of a ML concentration forecasting system based on the
analogues method. In particular, the surface ML concentration is linked to the atmospheric
patterns during a reference period. Then, during the forecasting phase, the forecasted atmospheric
situation is compared to those realized during the reference period to identify analogue situations.
95 The ML concentration during those analogues situations is considered to be a good approximation
of the ML concentration that will occur during the forecasted date. As this is a new approach
never tested before for ML dispersion, the first step has been to run several tests to fine-tune the
methodology and to characterize its limits of validity. Ideally, the tuning and validation of the
method should had been done using in-situ observations but, unfortunately, the available ML
100 concentration datasets are too scarce and this was not possible. Therefore, in this exploratory
study, we have used numerically simulated ML concentration fields for the development and
validation of the system.

The rest of the paper is organized as follows. In section 2, the statistical method, the datasets used
and the different choices tested are introduced. In section 3, the model results are presented and


105 discussed, and, finally, some conclusions about the capabilities of this new approach are outlined
in section 4.

2. Data and methods

2.1. The analogues method

The implementation of the analogues method requires two sets of data. First, we need a reference
110 dataset of the variables that describe the atmospheric patterns over the region of study, the so
called predictors (\mathbf{X}). The second reference dataset consists on spatial patterns of the variable of
interest for the same period for which the predictors are available. In our case, those predictands
(\mathbf{Y}) would be the ML concentration fields. Once they are defined, the methodology is based on
the assumption that if two predictors are similar ($\mathbf{X}_1 \sim \mathbf{X}_2$) the corresponding predictands would
115 also be similar ($\mathbf{Y}_1 \sim \mathbf{Y}_2$). Then, in order to obtain a forecast of the ML concentration for a given
date (\mathbf{Y}_{fct}), what we can do is to use instead a forecast of the predictor for the same date (\mathbf{X}_{fct}). In
particular, we look for K analogue dates within the reference period ($t_{an,k}$) in which the predictor
patterns are similar to the forecasted one ($\mathbf{X}(t_{an,k}) \approx \mathbf{X}_{fct}$). Then the value of the variable of interest
is estimated as a function of the predictands corresponding to the selected analogue dates $\mathbf{Y}_{fct} = \mathbf{f}$
120 ($\mathbf{Y}(t_{an,k})$). A scheme of the model algorithm is shown in figure 1a.

In our case, the predictors used to characterize the atmospheric conditions will be the Sea Level
Pressure (SLP) and the wind speed (U_{10}, V_{10}). These two variables have been successfully used to
forecast ocean surface dynamics (Wang et al., 2010; Martínez-Asensio et al., 2016), so it is
reasonable to think that they may be also good to forecast ML concentration as it is mainly driven
125 by ocean currents. The reference dataset for the atmospheric situation is obtained from an
atmospheric reanalysis (see section 2.3). Regarding the reference dataset for the predictand, we
use the ML concentration outputs from the modeling system developed by Soto-Navarro et al.
(2020) and described in section 2.4.



130 **Figure 1. a) Scheme of the functioning of the analogues method. b) example of the JM_1 cost function. The vertical red line marks the date forecasted (t_{fcst} , in the example 26/04/2006). The thin black line is the JM_1 cost function for the whole period, in the Mediterranean Sea region. Blue dots and vertical dashed lines indicate the analogue dates selected ($t_{an,k}$, see text for details).**

2.2. Algorithm implementation

135 The first step to implement the analogues method is to define a cost function, JM , that measures the similarity between different meteorological situations. Then, for the forecast day (t_{fcst}) we estimate how close is the meteorological situation of that day to the rest of the days in the reference dataset by computing JM for the whole reference period. Those days with the lowest JM values are selected as the analogue days ($\{t_{an,k}\}$, see figure 1b for an example). For the definition of JM ,
 140 the most popular choice is to use the Euclidean distance or root mean square error difference (RMSED) (Zorita et al., 1995; Cubasch et al., 1996; Gutiérrez et al., 2013), although other metrics based on different statistics can also be used. Here we have tested 4 different definitions for the cost function JM :

$$JM_1 = \sqrt{\left((SLP(t) - SLP(t_{fcst}))^2 \right)} \quad (1)$$

$$145 \quad JM_2 = \sqrt{\left((u_{10}(t) - u_{10}(t_{fcst}))^2 + (v_{10}(t) - v_{10}(t_{fcst}))^2 \right)} \quad (2)$$

$$JM_3 = JM_1(t) / \langle JM_1(t) \rangle + JM_2(t) / \langle JM_2(t) \rangle \quad (3)$$

$$JM_4 = \sum_{t_{fcst} - \Delta t}^{t_{fcst}} JM_3(t) \quad (4)$$

So, the similarity between meteorological situations is assessed either in terms of the sea level pressure (*SLP*, JM_1), the 10-m winds (U_{10} , V_{10} ; JM_2), a normalized combination of both (JM_3) or the cumulated values of JM_3 during a period (Δt) before the reference day (JM_4). In our case, Δt has been set to 7 days. Note that the horizontal bars indicate spatial averages for (JM_1) and (JM_2), while $\langle \rangle$ in (JM_3) denotes temporal mean.

In a second step, we identify the analogue dates as those with the lowest values of JM . We keep those dates in which JM is lower than the 1% percentile of all JM . Then, the ML concentration maps (C) obtained in the reference dataset for those days are combined to produce the forecast concentration map (C_{fcst}). In our case we use the median to reduce the influence of extreme concentration values close to ML sources:

$$C_{fcst} = median(C\{t_{an}\}) \quad (5)$$

2.3 Reanalysis data for the atmospheric fields

The period considered for the implementation of the analogues method is 2003 – 2013, which coincides with the period simulated by the ML dispersion model (as described in the following section). The climatic dataset necessary for the model reference period is based on the ERA5 reanalysis dataset, available at the Copernicus Climate Change Service (C3S) web platform (<https://climate.copernicus.eu/climate-reanalysis>). All the information regarding the ERA5 characteristics can be found on the C3S website.

Two variables have been considered for the characterization of atmospheric patterns forcing the ML dispersion: the wind speed at 10 meters height (U_{10} , V_{10}) and the sea level pressure (SLP). Daily mean values of these variables over the Mediterranean Sea were downloaded and processed for the whole period. The spatial resolution of the atmospheric data is 0.25° (~ 25 km) and cover the whole Mediterranean basin and the region of the North Atlantic adjacent to the Iberian Peninsula. Figure 2 shows as an example the average SLP for year 2013 in the selected domain.




Figure 2. Average SLP (in Pa) for the year 2013 in the region, computed from the ERA5 dataset. The red line at the Strait of Sicily marks the boundary between the Western and Eastern basins. The red rectangles limit the sub-basins of the Baleric Islands, The Gulf of Lions and the Aegean Sea, where specific analyses were carried out.

2.4 ML concentration data

The ML concentration data is obtained from the simulations performed by Soto-Navarro et al., (2020), as they are considered to be among the most realistic for the Mediterranean Sea. Due to the relevance of the quality of the ML concentration data, some details on the modelling system are presented below and more information can be found in Soto-Navarro et al., (2020).

The system is based on two components, a regional high resolution circulation model (RCM) reproducing the 3D current velocity field in the Mediterranean (NEMOMED36), and a lagrangian model that simulates the evolution of floating particles (Ichthyop 3.3).

The hydrodynamical model used to simulate the Mediterranean current field is an implementation on the NEMO model, with a spatial resolution of 1/36 degrees (~ 3 km) in a domain that covers the whole Mediterranean. The atmospheric forcing is a dynamical downscaling performed by the APEGE-Climate model using spectral nudging, namely ARPERA (Herrmann and Somot, 2008). Note that the forcing of NEMOMED36 (ARPERA) is not the same that the one used to characterize the meteorological situations (ERA5). Although both datasets are very similar, they are not exactly the same, thus mimicking the inaccuracies that atmospheric forecasts will inherently have.

The Individual Based Model (IBM) Ichthyop 3.3 (<http://www.ichthyop.org/>) is used to determine the 3D trajectories of the virtual ML particles from the NEMOMED36 current field. In the coastlines and the domain's boundaries the configuration of the model is set as "bouncing", meaning that the particles rebound back to the sea when reaching coastal pixels or the boundary of the domain. Therefore, no beaching scheme is implemented. Following the estimations of

Jambeck et al., (2015), a total input of 100k tons of plastic per year into the whole Mediterranean Sea is set in the model. This total amount is distributed in three different types of sources: cities, rivers and maritime traffic or ships - lanes, according to the ratio 50:30:20% respectively. The modelling period covers ten years, between 2003 and 2013. Due to computational limitations, it has been divided in 120 simulations, each one running one year and starting the first day of each month. A total of 41872 particles are released every month, which for the complete experiment makes a total of more than 5 million particles. The initial concentrations at the different source location are represented in figure 3a. The experiments were carried out using particles with positive (floating), neutral, and negative (sinking) buoyancy. In this study, only the results for floating ML particles have been used. Soto-Navarro et al. (2020) showed that the dispersion patterns for floating and neutral particles are very similar, hence the results described below can be considered valid also for neutral particles.

The results of the numerical experiments are processed to produce average ML concentration maps over the Mediterranean basin. These maps are computed by dividing the Mediterranean basin in a regular grid of 0.25° x 0.25° cells. The average concentration is estimated as the number of particles in each cell, divided by the cell surface, at each time step. Figure 3b shows the average ML concentration in the Mediterranean for the whole simulated period.

215




Figure 3. a) Spatial distribution of initial marine litter concentrations (in kg/km²) for the three simulations. Circle filled points indicate cities, diamonds indicate rivers and points over the sea indicate the ship lines. b) Average ML concentration of neutral particles (kg/km²) for the period 2003 – 2013.

220 2.5 Experiments

As mentioned before, there are no suitable observational datasets to validate the forecasting system. Homogenized datasets covering a long period of time would be required for this task. Although there are some efforts to develop new databases (Maximenko et al., 2019), up to our knowledge, there are no such datasets in the Mediterranean yet. Thus, in order to have a first
225 assessment of the quality of this methodology we have to use the concentration ML maps from the database as a “virtual reality” and compare the forecast (C_{fct}) with the C in the database for the forecast date ($C(t_{fct})$). We are aware that this may produce overoptimistic results and this issue will be discussed below.

To define the forecast day, we pick any date from the reference period and forecast the ML for
230 that day using all the data available except for a week before and after of the forecast day to avoid spurious good results due to autocorrelation. This has been repeated for all the dates in the reference period (3650) and several statistical metrics have been computed to assess the skills of the method.

To test if the model shows different skills depending on the domain of application, we have
235 applied the method to seven different regions: the whole Mediterranean, the eastern and western basins, and in the Gulf of Lions, the region around the Balearic Islands, the Adriatic Sea and the Aegean Sea (see figure 2). In each case, the analogue days have been defined using only data on the selected region.

Additionally, we have tested if the skill of the method depends on the time scales of the ML
240 concentration variability. So, in addition to use the ML concentration dataset, we have used two filtered versions of it, separating those processes above and below 15 days ($C_{hi-freq}$ and $C_{lo-freq}$).

Finally, for completeness, we propose three additional models for the forecasting. First, we
forecast the concentration change in 7 days ($\Delta_{7d}C$). The underlying idea is that the meteorological situation could be a better predictor of the rate of change than of the absolute value (e.g. winds
245 may determine the changes in the concentration rather than the absolute value). The second one is to simply assume 7-days persistence as the forecasting model (I.e. we assume $C(t_{ref}) = C(t_{ref-7\ days})$). This model will tell us if having a good observational characterization of the ML concentration would be a good predictor of what will be the situation one week later. The last one is a combination of the previous two: we add the forecast of the concentration change to the 7
250 days persistence ($C(t_{ref}) = C(t_{ref-7\ days}) + \Delta_{7d}C$). In other words, we test if combining a good observational characterization of the ML concentration with an analogues-based forecast of the concentration change can improve the results.

In summary, we have tested 4 configurations of the model over 7 different regions to forecast C , $C_{hi-freq}$ and $C_{lo-freq}$

255 2.6 Quality assessment

Several diagnostics are used to characterize the quality of the forecasts in the different experiments. The first one is the root median square error (RMEDSE):

$$RMEDSE = \sqrt{\text{median}((C_{an} - C_{ref})^2)} \quad (6)$$

We have chosen this parameter instead of the root mean square error to reduce the overall impact
260 of outliers linked to very high concentration values close to ML sources. Complementary we also compute the temporal correlation ρ :

$$\rho = \frac{\text{Cov}(C_{an}, C_{ref})}{\sigma_{C_{an}} \sigma_{C_{ref}}} \quad (7)$$

where Cov represents the covariance and σ the standard deviation. Additionally, we compute the
RMEDSE ratio (RR) which is defined as the ratio between the RMEDSE of the forecast (eq. 6)
265 and the RMEDSE computed using all the days in the database, RMEDALL:

$$RR = RMEDSE / RMEDALL \quad (8)$$

The lower the value of RR is, the better the forecast is. Values of RR close to 1 means that the
quality would be the same than using any random day, so the forecast is not providing any new
information. RMEDSE, ρ and RR are computed spatially and/or temporally.

270 3. Results

3.1 Time variability

The temporal correlation and the RR of the ML concentration reconstruction using different cost
functions and forecasting models are presented in Figures 4 and 5. The spatial patterns of the
correlation are very consistent among the different combinations. The fields are relatively patchy
275 with the highest values in the eastern basin, close to the Turkey coasts, in the Gulf of Gabes, in
the west of Sardinia and towards the north of the Balearic Islands. Conversely, the minimum
correlation values are found in the Alboran Sea, the Algerian basin and the Gulf of Lions. The
RR maps are very consistent showing lower values where/when the correlation is higher and
values closer to 1 where/when the correlation is lower.

280 Concerning the different cost functions used to identify the analogue situations, the performances
using only SLP (JM_1) or only wind (JM_2) are very similar. Using both variables the quality
slightly increases (JM_3) and becomes significantly better when using the 7-days average (JM_4).

For model 1 (forecasting the concentration), the averaged correlation using each cost function is 0.24, 0.25, 0.28 and 0.35 while the averaged RR is 0.93, 0.93, 0.90 and 0.86, respectively. The forecasting of the concentration change is worse for all cost functions, with averaged correlation values ranging from 0.08 to 0.19 and RR ranging from 1.00 to 0.98. In the light of these results, from now on, we will only consider the results of the analogues-based forecast models that use the cost function JM_4 (i.e. the one considering the 7-day averaged differences). Using it for forecasting the ML concentration we obtain correlation values ranging from 0.20 to up to 0.60 depending on the region. When forecasting the ML concentration change the values range from non-significant to 0.40 (see Figure 4).

Using 7-days persistence to forecast the ML concentration (Model 3, see Figure 4) the results largely improve. They show correlations that range from 0.20 in the Alboran Sea and the Gulf of Lions to 0.82 around Cyprus, with an average value of 0.60. The RR reaches values as low as 0.4 with an average value of 0.79. Finally, combining both methodologies in Model 4 provides the best results. Combining the 7-days persistence with the analogues-based forecast of the concentration change increases the forecasting skills. In this case the averaged correlation is 0.62 and the averaged RR 0.79.




Figure 4. Temporal correlation of the forecasts using different models and cost functions with the reference dataset. Each column corresponds to a different forecasting model: the analogues-based forecast of the concentration (model 1), the analogues-based forecast of the concentration changes in 7 days (model 2), the persistence (model 3), and the persistence in combination with the forecast of the concentration change in 7 days (model 4). Each row corresponds to the different cost functions used to identify the analogues (see text for details). Note that all panels in the third column are the same, as in Model 3 no cost function is used.




Figure 5. Same than Figure 5 but for the RMEDSE ratio. Values close to 1 (white) indicate the forecast brings little improvement with respect to use a random day.

For completeness, we also include an example of the concentration time series for the reference and models 1, 3 and 4 for a point where the forecasts perform well (Figure 6a). It can be seen that Model 1 is well correlated with the reference, showing a good chronology of events although being unable of capturing the concentration peaks. During those periods, the analogues-based forecast largely underestimates the reference values. Models 3 and 4 show almost identical good results, as far as persistence is enough to capture most of the variability. The underlying reason for this success is that, at this location, the changes of ML concentration are relatively slower, so assuming persistence can be a good predictor. For comparison, the time series for a point where the models perform poorly are shown in Figure 6b. In this case, the analogues-based forecast is unable to capture any variability and it basically produces the mean value. The other two models are able to follow the variability, although in this case the skills are lower than in the previous case. The reason is that in this point the ML concentration varies more rapidly, so assuming the persistence is not as good predictor as it was in the previous location.





Figure 6. Time series of ML concentration (in kg/km²) for (top) a location where the analogues-based forecast works well and (bottom) a location where it performs worse. The plots show the reference values, the analogues-based forecast, the persistence and the persistence in combination with the forecast of the concentration change.

3.2 Spatial Variability

A complementary view of the performance of the different forecasting models can be obtained looking at the ML concentration anomalies (i.e. with respect to the temporal mean) in given dates. In Figure 7, we show the results for a date when models show good agreement with the reference (spatial correlation values are 0.70, 0.76 and 0.78 for models 1, 3 and 4, respectively). All three models are able to identify the areas of high and low concentration. Maximum values in the north of the Balearic Islands, the Gulf of Gabes and south of Italy and minimum values in the Adriatic Sea, the Algerian basin and the easternmost part of the Med are well captured. The analogues-based forecast (Model 1) shows smoother patterns with less low extremes. This is in good agreement with what has been seen in the time series in Figure 6, suggesting that this model has difficulties to capture very high concentration values. Regarding the persistence-based models, for this particular date, they perform very well capturing not only the large scale patterns but also the local features. Looking at a date when the performance is lower something interesting appears. Although the spatial correlation of Model 1 is not significant (Figure 8b), the large scale features seem to be well captured. However, the small scale features are clearly not captured which degrades the spatial correlation. This would also support the previous finding reinforcing the idea that the analogues-based forecast performs better for the large scale features. In places or dates where/when the small scale features become dominant, the performance of the model drops.



345 **Figure 7. Maps of ML concentration anomaly for a date where the analogues-based forecast performs well (a) Reference (b) Analogues-based forecast (c) Persistence (d) Persistence in combination with the forecast of the concentration change in 7 days.**





Figure 8. Like Figure 7 but for a situation where the forecasts perform worse.

350 The time series of the spatial correlation and spatial RR at each time step are presented in Figure 9. The results are similar for the three models forecasting the ML concentration (Models 1, 3 and 4). The skills of the forecasts show a high temporal variability with correlation values ranging from 0.5 to almost 1 and an averaged value of 0.78, 0.81 and 0.84, respectively. For RR the values range from 0.3 to more than 1 with an average value of 0.76, 0.79 and 0.71, respectively. This

355 diagnostic also confirms that the best model is the one combining the persistence with the forecast of the concentration change.



360 **Figure 9. Time series of (Top) Spatial Correlation and (Bottom) Spatial RR. for the analogues-based forecast (Model 1), the persistence (Model 3), and the persistence in combination with the forecast of the concentration change (Model 3).**

3.3 Regional dependence of the forecasting skills

The methodology has also been applied to different domains. That is, the cost function, JM , has been computed in the regions defined in figure 2 and the validation has been performed looking only at the ML concentration in those regions. In general, better results are obtained when the analogues-based forecasts are applied to a larger region (see Table 1 and Table 2). For instance, the analogues-based forecast (Model 1) provides modest results, with correlations of 0.31 and 0.35 and RR of 0.92 and 0.86 for the eastern and western Mediterranean, respectively. At local scale the correlation ranges between 0.29 and 0.34 and the RR ranges between 0.80 and 0.94. The analogues-based forecast for the concentration change (Model 2) shows lower skills with correlation below 0.23 and RR above 0.98 in all regions. Both models show better performance forecasting the low-frequency component than the high-frequency one. The correlation of Model 1 forecasts in the different region ranges between 0.31 and 0.40 for the low frequency while it ranges between 0.15 and 0.22 for the high frequency. Consistent results are found when looking at the RR and Model 2 forecasts.

375 The 7-days persistence (Model 3) shows to be a good predictor for the full signal and the low-frequency component while it struggles to capture the high-frequency variability, as expected. Provided that the low-frequency part of the signal is what dominates the ML concentration variability, this model shows good skills for the full signal with correlations in all regions ranging from 0.55 to 0.64 and RR ranging from 0.75 to 0.82.

380 The best results for the forecast of the ML concentration are obtained combining the persistence with the analogues-based forecast of the 7-days concentration change (Model 4). The averaged temporal correlation is over 0.54 in all regions reaching a value of 0.65 when applied to the Western Mediterranean, while RR is below 0.80 and reaches 0.76 for the whole Mediterranean.

Correlation	Full				High-frequency				Low-frequency			
	M1	M2	M3	M4	M1	M2	M3	M4	M1	M2	M3	M4
Mediterranean	0,35	0,19	0,60	0,62	0,19	0,19	-0,13	-0,08	0,40	0,25	0,95	0,96
East Med	0,31	0,19	0,55	0,57	0,19	0,18	-0,13	-0,08	0,34	0,22	0,95	0,95
West Med	0,35	0,16	0,64	0,65	0,15	0,15	-0,12	-0,08	0,41	0,24	0,96	0,96
Gulf of Lions	0,29	0,20	0,51	0,54	0,20	0,19	-0,16	-0,10	0,31	0,19	0,94	0,94
Balearic Islands	0,36	0,18	0,60	0,62	0,17	0,17	-0,14	-0,09	0,40	0,20	0,95	0,96
Adriatic Sea	0,31	0,23	0,53	0,56	0,22	0,21	-0,13	-0,07	0,28	0,17	0,94	0,94
Aegean Sea	0,34	0,21	0,55	0,59	0,22	0,20	-0,12	-0,06	0,39	0,19	0,94	0,94

385 **Table 1. Regionally averaged temporal correlation of the different forecasting models (M1-M4) applied in different regions (see Figure 2). The models have been applied to forecast the full signal of ML concentration, the high frequency component (period < 15 days) and the low frequency component (period > 15 days).**

RR	Full				High-frequency				Low-frequency			
	M1	M2	M3	M4	M1	M2	M3	M4	M1	M2	M3	M4
Mediterranean	0,86	0,98	0,79	0,76	0,97	0,98	1,47	1,44	0,82	0,92	0,26	0,24
East Med	0,92	0,98	0,82	0,80	0,98	0,98	1,48	1,45	0,89	0,94	0,28	0,27
West Med	0,86	1,00	0,75	0,74	0,98	0,99	1,47	1,45	0,79	0,93	0,25	0,23
Gulf of Lions	0,80	0,98	0,80	0,79	0,97	0,98	1,49	1,46	0,79	0,96	0,26	0,25
Balearic Islands	0,94	0,98	0,76	0,75	0,98	0,99	1,48	1,46	0,91	0,96	0,27	0,25
Adriatic Sea	0,84	0,98	0,79	0,77	0,98	0,99	1,46	1,44	0,88	0,98	0,29	0,28
Aegean Sea	0,85	1,00	0,80	0,75	0,97	0,99	1,46	1,45	0,82	0,97	0,27	0,26

Table 2. Same as Table 1 but for the RMEDSE ratio.

390 The spatial diagnostics have also been computed applying the models to different domains (Table 3). In this case, the analogues-based forecast of concentration (M1) show average spatial correlations higher than 0.62 when applied to any region reaching up to 0.94 in the Aegean Sea. Also, the analogues-based forecast of concentration change (M2) shows significant average correlations ranging between 0.19 and 0.30. 7-days persistence (M3) is again improving the results although the combination of persistence and the analogues-based forecast of concentration change (M4) is the best model when applied in any region. Average correlation ranges between 395 0.67 and 0.96 and RR is lower than 0.83 everywhere.

	Correlation				RR			
	M1	M2	M3	M4	M1	M2	M3	M4
Mediterranean	0,75	0,23	0,83	0,84	0,86	1,00	0,80	0,75
East Med	0,70	0,23	0,75	0,76	0,89	0,97	0,85	0,82
West Med	0,76	0,22	0,84	0,85	0,86	1,00	0,78	0,72
Gulf of Lions	0,65	0,19	0,66	0,67	0,83	0,98	0,85	0,83
Balearic Islands	0,62	0,20	0,72	0,73	0,91	0,98	0,76	0,74
Adriatic Sea	0,64	0,21	0,67	0,69	0,85	1,00	0,74	0,73
Aegean Sea	0,94	0,30	0,96	0,96	0,86	1,00	0,78	0,78

Table 3. Temporally averaged regional correlation and RR of the different forecasting models (M1-M4) applied in different regions (see Figure 2).

400

It is worth mentioning that we have also tested other options for the cost function like using different temporal averages or using correlation as similarity metrics but no significant differences have been found. Also, we have tried to change the criterion to define the analogue days. Instead of identifying as analogues those days with JM lower than the 1% percentile of the whole JM time series, we have used less restrictive criteria (5% or 10%). In both cases the results worsened.

405

4. Discussion and Conclusions

The analogues-based forecasting technique has been applied to ML concentration for the first time, up to our knowledge. It has proven to be very inexpensive and relatively easy to set-up, so it is an alternative to direct modelling worth to be considered. A key step in the set-up is to select a suitable cost function and the best threshold to identify the analogue meteorological situations. In our case, it seems that using integral definitions for the cost function improve the results. In other words, it is better to identify the analogue days based on the history of the meteorological situation. Probably, using a different averaging time for each domain would allow increasing the skills of the analogues-based model. However, this fine tuning is out of the scope of this paper, as far as there are no suitable observations to validate it, as it will be discussed later.

410

415

The quality of the analogues-based forecasts depends on the region of application. Our results suggest that the larger the region of application the better, as we get better results for the whole Mediterranean or for the East/West basins than in smaller local areas. A hypothesis for explaining this result is that using the atmospheric situation as a predictor may not be suitable to capture small scale features (e.g. those related to ocean currents or the interaction with coastlines). Further tests including other predictors could be done to refine the method, including ocean currents, for instance.

420

Another important point is that the method struggles to capture the extreme values as it produces smooth spatio-temporal patterns of ML concentration. Therefore, in locations or regions where

425 short intense events or small scale features dominate the variability, the method performs worse. This is also one of the reasons why the temporal skills are relatively low (i.e. temporal correlation and RR, see section 3.1). Conversely, if instead of the time variability, what are aimed at are the spatial structures, the method shows high skills being able to locate relative maximum and minimum (see section 3.2).

430 We have also shown that persistence is a very good predictor almost everywhere. This is because the ML concentration changes relatively slowly (i.e. the system has a several days memory), at least at the spatial scales solved by the reference dataset. This means that if reliable information was available (e.g. from a monitoring program), this could be used as a first guess of the ML concentration several days later. Complementary, the analogues-based method has also been
435 applied to forecast concentration change. In this case the results were significantly poorer both to capture the time and spatial variability. However, it can be useful to improve the persistence-based forecasts.

Regarding the reliability of the analogues based forecasts that could be generated from this reference dataset, its quality would directly depend on the accuracy of the reference dataset. In
440 our case this dataset comes from the outputs of a realistic modelling (Soto-Navarro et al., 2020). However, the model may have some shortcomings as its spatial resolution, beaching parameterization or realism of ML sources. Consequently, the forecasts would be, in the best case, as good as the model outputs are. Therefore, it would have been better to validate the different forecasting models against actual observations. Unfortunately, the lack of observations with a
445 suitable spatial and temporal coverage prevents from doing it. In the future, it would be worth setting up a monitoring program with enough spatial and temporal resolution that would allow generating a comprehensive enough reference dataset. This dataset could be used to train the analogues-based forecasting system and to validate other existing systems.

In any case, it is worth noting that the validation of the methodology can be considered as robust.
450 For that purpose, it is not required that the reference dataset is an accurate representation of the actual ML concentration. Only the statistics of the ML concentration spatiotemporal evolution has to be reproduced. And in that sense, the model integrates the effects of a realistic atmospheric forcing and a realistic ocean current field. So, it is expected that the statistics of the ML concentration field is realistic enough. This extent should also be confirmed by a comprehensive
455 observational dataset, at least in certain regions.

In conclusion, the analogues-based model presented here has potential to become a suitable cost effective forecasting method for ML concentration. It could be easily implemented in any region of the world where a realistic reference dataset is available. In those regions where the large scale

ML concentration patterns dominate the variability the method will probably work better than in
460 regions where the variability is dominated by small scale structures.

5. Code and data availability

The code and data required to implement the model described in the paper and to reproduce the
results can be publicly accessed at Jordà and Soto-Navarro, (2022). Additionally, the atmospheric
fields can be downloaded from the Copernicus portal (<https://climate.copernicus.eu/climate-reanalysis>).
465 [reanalysis](https://climate.copernicus.eu/climate-reanalysis)).

6. Author contribution

Both authors (GJ and JSN) have contributed equally to the design of the study, the coding of the
modelling system, the performance of the simulations, the analysis of the results and the
preparation and revision of the manuscript.

7. Competing interest

The authors declare that they have no conflict of interest.

8. Acknowledgments

Acknowledgements to the EU-Interreg MPAs Plastic Busters Project: preserving serving
biodiversity from plastics in Mediterranean Marine Protected Areas, co-financed by the European
475 Regional Development Fund (grant agreement No 4MED17_3.2_M123_027). Also to the Junta
de Andalucía funded “Origen y evolución de la basura marina en la costa andaluza
(OBAMARAN)” (ref. Proy_Exel_00344). The authors also thank Mr. Paul Dupin for his help in
the first steps of the model development.

9. References

480 Caillouet, L., Vidal, J. P., Sauquet, E., and Graff, B.: Probabilistic precipitation and temperature
downscaling of the Twentieth Century Reanalysis over France, *Clim. Past*, 12, 635–662,
<https://doi.org/10.5194/cp-12-635-2016>, 2016.

Casanueva, A., Frías, M. D., Herrera, S., San-Martín, D., Zaninovic, K., and Gutiérrez, J. M.:
Statistical downscaling of climate impact indices: testing the direct approach, *Clim. Change*, 127,
485 547–560, <https://doi.org/10.1007/s10584-014-1270-5>, 2014.

Compa, M., Alomar, C., Wilcox, C., van Sebille, E., Lebreton, L., Hardesty, B. D., and Deudero,
S.: Risk assessment of plastic pollution on marine diversity in the Mediterranean Sea, *Sci. Total
Environ.*, 678, 188–196, <https://doi.org/10.1016/j.scitotenv.2019.04.355>, 2019.

- Cózar, A., Sanz-Martín, M., Martí, E., González-Gordillo, J. I., Ubeda, B., Á.gálvez, J., Irigoien, X., and Duarte, C. M.: Plastic accumulation in the mediterranean sea, *PLoS One*, 10, 1–12, <https://doi.org/10.1371/journal.pone.0121762>, 2015.
- Cubasch, U., Von Storch, H., Waszkewitz, J., and Zorita, E.: Estimates of climate change in Southern Europe derived from dynamical climate model output, *Clim. Res.*, 7, 129–149, <https://doi.org/10.3354/cr007129>, 1996.
- 495 Fossi, M. C., Romeo, T., Bainsi, M., Panti, C., Marsili, L., Campani, T., Canese, S., Galgani, F., Druon, J.-N., Airoidi, S., Taddei, S., Fattorini, M., Brandini, C., and Lapucci, C.: Plastic Debris Occurrence, Convergence Areas and Fin Whales Feeding Ground in the Mediterranean Marine Protected Area Pelagos Sanctuary: A Modeling Approach, *Front. Mar. Sci.*, 4, 1–15, <https://doi.org/10.3389/fmars.2017.00167>, 2017.
- 500 Grouillet, B., Ruelland, D., Ayar, P. V., and Vrac, M.: Sensitivity analysis of runoff modeling to statistical downscaling models in the western Mediterranean, *Hydrol. Earth Syst. Sci.*, 20, 1031–1047, <https://doi.org/10.5194/hess-20-1031-2016>, 2016.
- Gutiérrez, J. M., San-Martín, D., Brands, S., Manzanos, R., and Herrera, S.: Reassessing statistical downscaling techniques for their robust application under climate change conditions, *J. Clim.*, 26, 505 171–188, <https://doi.org/10.1175/JCLI-D-11-00687.1>, 2013.
- Hayes, A., Kucera, M., Kallel, N., Sbaifi, L., and Rohling, E. J.: Glacial Mediterranean sea surface temperatures based on planktonic foraminiferal assemblages, *Quat. Sci. Rev.*, 24, 999–1016, <https://doi.org/10.1016/j.quascirev.2004.02.018>, 2005.
- Herrmann, M. J. and Somot, S.: Relevance of ERA40 dynamical downscaling for modeling deep convection in the Mediterranean Sea, *Geophys. Res. Lett.*, 35, 1–5, 510 <https://doi.org/10.1029/2007GL032442>, 2008.
- Jambeck, J., Geyer, R., Wilcox, C., Siegler, T. R., Perryman, M., Andrady, A., Narayan, R., and Law, K. L.: Plastic waste inputs from land into the ocean, *Science* (80-), 347, <https://doi.org/10.1126/science.1260352>, 2015.
- 515 Jordà, G. and Soto-Navarro, J.: An analogue based forecasting system for Mediterranean marine litter concentration - Code and simulations dataset, <https://doi.org/10.5281/zenodo.7104402>, 2022.
- Law, K. L., Morét-Ferguson, S. E., Goodwin, D. S., Zettler, E. R., Deforce, E., Kukulka, T., and Proskurowski, G.: Distribution of surface plastic debris in the eastern pacific ocean from an 11-

- 520 year data set, *Environ. Sci. Technol.*, 48, 4732–4738, <https://doi.org/10.1021/es4053076>, 2014.
- Lebreton, L. and Andrady, A.: Future scenarios of global plastic waste generation and disposal, *Palgrave Commun.*, 5, 1–11, <https://doi.org/10.1057/s41599-018-0212-7>, 2019.
- Lebreton, L., Egger, M., and Slat, B.: A global mass budget for positively buoyant macroplastic debris in the ocean, *Sci. Rep.*, 9, 1–10, <https://doi.org/10.1038/s41598-019-49413-5>, 2019.
- 525 Lebreton, L. C. M., Greer, S. D., and Borrero, J. C.: Numerical modelling of floating debris in the world's oceans, *Mar. Pollut. Bull.*, 64, 653–661, <https://doi.org/10.1016/j.marpolbul.2011.10.027>, 2012.
- Liubartseva, S., Coppini, G., Lecci, R., and Creti, S.: Regional approach to modeling the transport of floating plastic debris in the Adriatic Sea, *Mar. Pollut. Bull.*, 103, 115–127, <https://doi.org/10.1016/j.marpolbul.2015.12.031>, 2016.
- 530 Liubartseva, S., Coppini, G., Lecci, R., and Clementi, E.: Tracking plastics in the Mediterranean: 2D Lagrangian model, *Mar. Pollut. Bull.*, 129, 151–162, <https://doi.org/10.1016/j.marpolbul.2018.02.019>, 2018.
- Lopez-Radcenco, M., Pascual, A., Gomez-Navarro, L., Aissa-El-Bey, A., Chapron, B., and Fablet, R.: Analog Data Assimilation of Along-Track Nadir and Wide-Swath SWOT Altimetry Observations in the Western Mediterranean Sea, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 12, 2530–2540, <https://doi.org/10.1109/JSTARS.2019.2903941>, 2019.
- Lorenz, E. N.: Atmospheric Predictability as Revealed by Naturally Occurring Analogues, *J. Atmos. Sci.*, 26, 636–646, [https://doi.org/10.1175/1520-0469\(1969\)26<636:APARBN>2.0.CO;2](https://doi.org/10.1175/1520-0469(1969)26<636:APARBN>2.0.CO;2), 1969.
- 540 0469(1969)26<636:APARBN>2.0.CO;2, 1969.
- Macias, D., Cózar, A., Garcia-Gorriz, E., González-Fernández, D., and Stips, A.: Surface water circulation develops seasonally changing patterns of floating litter accumulation in the Mediterranean Sea. A modelling approach, *Mar. Pollut. Bull.*, 149, 110619, <https://doi.org/10.1016/j.marpolbul.2019.110619>, 2019.
- 545 Mansui, J., Molcard, A., and Ourmières, Y.: Modelling the transport and accumulation of floating marine debris in the Mediterranean basin, *Mar. Pollut. Bull.*, 91, 249–257, <https://doi.org/10.1016/j.marpolbul.2014.11.037>, 2015.
- Martínez-Asensio, A., Marcos, M., Tsimplis, M. N., Jordà, G., Feng, X., and Gomis, D.: On the ability of statistical wind-wave models to capture the variability and long-term trends of the North

550 Atlantic winter wave climate, *Ocean Model.*, 103, 177–189,
<https://doi.org/10.1016/j.ocemod.2016.02.006>, 2016.

Maximenko, N., Hafner, J., and Niiler, P.: Pathways of marine debris derived from trajectories of Lagrangian drifters, *Mar. Pollut. Bull.*, 65, 51–62,
<https://doi.org/10.1016/j.marpolbul.2011.04.016>, 2012.

555 Maximenko, N., Corradi, P., Law, K. L., Van Sebille, E., Garaba, S. P., Lampitt, R. S., Galgani, F., Martinez-Vicente, V., Goddijn-Murphy, L., Veiga, J. M., Thompson, R. C., Maes, C., Moller, D., Löscher, C. R., Addamo, A. M., Lamson, M. R., Centurioni, L. R., Posth, N. R., Lumpkin, R., Vinci, M., Martins, A. M., Pieper, C. D., Isobe, A., Hanke, G., Edwards, M., Chubarenko, I. P., Rodriguez, E., Aliani, S., Arias, M., Asner, G. P., Brosich, A., Carlton, J. T., Chao, Y., Cook, A.-
560 M., Cundy, A. B., Galloway, T. S., Giorgetti, A., Goni, G. J., Guichoux, Y., Haram, L. E., Hardesty, B. D., Holdsworth, N., Lebreton, L., Leslie, H. A., Macadam-Somer, I., Mace, T., Manuel, M., Marsh, R., Martinez, E., Mayor, D. J., Le Moigne, M., Molina Jack, M. E., Mowlem, M. C., Obbard, R. W., Pabortsava, K., Robberson, B., Rotaru, A.-E., Ruiz, G. M., Spedicato, M. T., Thiel, M., Turra, A., and Wilcox, C.: Toward the Integrated Marine Debris Observing System,
565 *Front. Mar. Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00447>, 2019.

Palatinus, A., Kovač Viršek, M., Robič, U., Grego, M., Bajt, O., Šiljić, J., Suaria, G., Liubartseva, S., Coppini, G., and Peterlin, M.: Marine litter in the Croatian part of the middle Adriatic Sea: Simultaneous assessment of floating and seabed macro and micro litter abundance and composition, *Mar. Pollut. Bull.*, 139, 427–439, <https://doi.org/10.1016/j.marpolbul.2018.12.038>,
570 2019.

Politikos, D. V., Ioakeimidis, C., Papatheodorou, G., and Tsiaras, K.: Modeling the Fate and Distribution of Floating Litter Particles in the Aegean Sea (E. Mediterranean), *Front. Mar. Sci.*, 4, 1–18, <https://doi.org/10.3389/fmars.2017.00191>, 2017.

Van Sebille, E., Chris, W., Laurent, L., Nikolai, M., Britta Denise, H., Jan, A. van F., Marcus, E.,
575 David, S., Francois, G., and Kara Lavender, L.: A global inventory of small floating plastic debris, *Environ. Res. Lett.*, 10, 124006, 2015.

Van Sebille, E., Aliani, S., Law, K. L., Maximenko, N., Alsina, J. M., Bagaev, A., Bergmann, M., Chapron, B., Chubarenko, I., Cózar, A., Delandmeter, P., Egger, M., Fox-Kemper, B., Garaba, S. P., Goddijn-Murphy, L., Hardesty, B. D., Hoffman, M. J., Isobe, A., Jongedijk, C. E., Kaandorp,
580 M. L. A., Khatmullina, L., Koelmans, A. A., Kukulka, T., Laufkötter, C., Lebreton, L., Lobelle, D., Maes, C., Martinez-Vicente, V., Morales Maqueda, M. A., Poulain-Zarcos, M., Rodríguez, E., Ryan, P. G., Shanks, A. L., Shim, W. J., Suaria, G., Thiel, M., Van Den Bremer, T. S., and

Wichmann, D.: The physical oceanography of the transport of floating marine debris, *Environ. Res. Lett.*, 15, <https://doi.org/10.1088/1748-9326/ab6d7d>, 2020.

585 Soto-Navarro, J., Jordà, G., Deudero, S., Alomar, C., Amores, Á., and Compa, M.: 3D hotspots of marine litter in the Mediterranean: A modeling study, *Mar. Pollut. Bull.*, 155, 111159, <https://doi.org/10.1016/j.marpolbul.2020.111159>, 2020.

Soto-Navarro, J., Jordà, G., Compa, M., Alomar, C., Fossi, M. C., and Deudero, S.: Impact of the marine litter pollution on the Mediterranean biodiversity: A risk assessment study with focus on
590 the marine protected areas, *Mar. Pollut. Bull.*, 165, 112169, <https://doi.org/https://doi.org/10.1016/j.marpolbul.2021.112169>, 2021.

UNEP: Marine Litter, A Global Challenge, UNEP, Nairobi, 232 pp., 2009.

Wang, X. L., Swail, V. R., and Cox, A.: Dynamical versus statistical downscaling methods for ocean wave heights, *Int. J. Climatol.*, 30, 317–332, <https://doi.org/10.1002/joc.1899>, 2010.

595 Wu, W., Liu, Y., Ge, M., Rostkier-Edelstein, D., Descombes, G., Kunin, P., Warner, T., Swerdlin, S., Givati, A., Hopson, T., and Yates, D.: Statistical downscaling of climate forecast system seasonal predictions for the Southeastern Mediterranean, *Atmos. Res.*, 118, 346–356, <https://doi.org/10.1016/j.atmosres.2012.07.019>, 2012.

Zambianchi, E., Trani, M., and Falco, P.: Lagrangian Transport of Marine Litter in the
600 Mediterranean Sea, *Front. Environ. Sci.*, 5, 1–15, <https://doi.org/10.3389/fenvs.2017.00005>, 2017.

Zorita, E. and von Storch, H.: The Analog Method as a Simple Statistical Downscaling Technique: Comparison with More Complicated Methods, *J. Clim.*, 12, 2474–2489, [https://doi.org/10.1175/1520-0442\(1999\)012<2474:TAMAAS>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<2474:TAMAAS>2.0.CO;2), 1999.

605 Zorita, E., Hughes, J. P., Lettermaier, D. P., and von Storch, H.: Stochastic Characterization of Regional Circulation Patterns for Climate Model Diagnosis and Estimation of Local Precipitation, *Journal Clim.*, 8, 1023–1042, 1995.