

Snow sensitivity to temperature and precipitation change during compound cold-hot and wet-dry seasons in the Pyrenees.

Josep Bonsoms¹, Juan I. López-Moreno², Esteban Alonso-González³

¹ Department of Geography, Universitat de Barcelona, Barcelona, Spain.

² Instituto Pirenaico de Ecología (IPE-CSIC), Campus de Aula Dei, Zaragoza, Spain.

³ Centre d'Etudes Spatiales de la Biosphère (CESBIO), Université de Toulouse, CNES/CNRS/IRD/UPS, Toulouse, France.

Corresponding author: Juan I. López-Moreno (nlopez@ipe.csic.es)

1 **Abstract.** The Mediterranean basin has experienced one of the highest warming rates on earth
2 during the last few decades, and climate projections predict water-scarcity in the future. Mid-
3 latitude Mediterranean mountain areas, such as the Pyrenees, play a key role in the hydrological
4 resources for the highly populated lowland areas. However, there are still large uncertainties about
5 the impact of climate change on snowpack in the high mountain ranges of this region. Here, we
6 perform a snow sensitivity to temperature and precipitation change analysis of the Pyrenean
7 snowpack (1980 – 2019 period) using five key snow-climatological indicators. We analyzed snow
8 sensitivity to temperature and precipitation during four different compounds weather conditions
9 (cold-dry [CD], cold-wet [CW], warm-dry [WD], and warm-wet [WW]) at low elevations (1500
10 m), mid-elevations (1800 m), and high elevations (2400 m) in the Pyrenees. In particular, we
11 forced a physically based energy and mass balance snow model (FSM2), with validation by
12 ground-truth data, and applied this model to the entire range, with forcing of perturbed reanalysis
13 climate data for the period 1980 to 2019 as the baseline. The FSM2 model results successfully
14 reproduced the observed snow depth (HS) values ($R^2 > 0.8$), with relative root-mean square error
15 and mean absolute error values less than 10% of the observed HS values. Overall, the snow
16 sensitivity to temperature and precipitation change decreased with elevation and increased
17 towards the eastern Pyrenees. When the temperature increased progressively at 1°C intervals, the
18 largest seasonal HS decreases from the baseline were at +1°C ~~(47% at low elevation, 48% at~~
19 ~~mid elevation, and 25% at high elevation)~~. A 10% increase of precipitation counterbalanced the
20 temperature increases ($\leq 1^\circ\text{C}$) at high elevations during the coldest months, because temperature
21 was far from the isothermal 0°C conditions. The maximal seasonal HS and peak HS max
22 reductions were during WW seasons, and the minimal reductions were during CD seasons. During
23 WW (CD) seasons, the seasonal HS decline per °C was 37% (28 %) at low elevations, 34% (30%)
24 at mid elevations, and 27% (22 %) at high elevations. Further, the peak HS date was on average
25 anticipated 2, 3 and 8 days at low, mid and high elevation, respectively. Results suggest snow

26 sensitivity to temperature and precipitation change will be similar at other mid-latitude mountain
27 areas, where snowpack reductions will have major consequences on the nearby ecological and
28 socioeconomic systems.

29

30 **Keywords:** Snow, Climate change, Sensitivity, Alpine, Mediterranean Mountains, Mid-latitude,
31 Pyrenees.

32

33 **1 Introduction**

34

35 Snow is a key element of the Earth's climate system (Armstrong and Brun, 1998) because it cools
36 the planet (Serreze and Barry, 2011) by altering the Surface Energy Balance (SEB), increasing
37 the albedo, and modulating surface and air temperatures (Hall, 2004). Northern-Hemispheric
38 snowpack patterns have changed rapidly during recent decades (Hammond et al., 2018; Hock et
39 al., 2019; Notarnicola et al., 2020). It is crucial to improve our understanding of the timing of
40 snow ablation and snow accumulation due to changing climate conditions because snowpack
41 affects many nearby social and environmental systems. From the hydrological point of view, snow
42 melt controls mountain runoff rate during the spring (Barnett et al., 2005; Adams et al., 2009;
43 Stahl et al., 2010), river flow magnitude and timing (Morán-Tejeda et al., 2014; Sanmiguel-
44 Vallelado et al., 2017), water infiltration and groundwater storage (Gribovszki et al., 2010; Evans
45 et al., 2018), and transpiration rate (Cooper et al., 2020). The presence and duration of snowpack
46 affects terrestrial ecosystem dynamics because snow ablation date affects photosynthesis
47 (Woelber et al., 2018), forest productivity (Barnard et al., 2018), freezing and thawing of the soil
48 (Luetschg et al., 2008; Oliva et al., 2014), and thickness of the active layer in permafrost
49 environments (Hrbáček et al., 2016; Magnin et al., 2017). Snowpack also has remarkable
50 economic impacts. For example, the snowpack at high elevations and surrounding areas
51 determines the economic success of many mountain ski-resorts (Scott et al., 2003; Pons et al.,
52 2015; Gilaberte-Búrdalo et al., 2017). Changes in the snowpack of mountainous regions also
53 influence associated lowland areas because it affects the availability of snow meltwater that is
54 used for water reservoirs, hydropower generation, agriculture, industries, and other applications
55 (e.g., Sturm et al., 2017; Beniston et al., 2018).

56

57 Mid-latitude snowpacks have among the highest snow sensitivities worldwide (Brown and Mote,
58 2009; López-Moreno et al., 2017; 2020b). In regions at high latitudes or high elevations,
59 increasing precipitation can partly counterbalance the effect of increases of temperature on snow

60 cover duration (Brown and Mote, 2009). Climate warming decreases the maximum and seasonal
61 snow depth (HS), the snow water equivalent (SWE) (Trujillo and Molotch, 2014; Alonso-
62 González et al., 2020a; López-Moreno et al., 2013; 2017), and the fraction total precipitation as
63 snowfall (snowfall ratio; e.g., Mote et al., 2005; Lynn et al., 2020; Jeenings and Molotoch, 2020;
64 Marshall et al., 2019), and also delays the snow onset date (Beniston, 2009; Klein et al., 2016).
65 However, warming can slow the early snow ablation rate on the season (Pomeroy et al., 2015;
66 Rasouli et al., 2015; Jennings and Molotch, 2020; Bonsoms et al., 2022; Sanmiguel-Vallelado et
67 al., 2022) because of the earlier HS and SWE peak dates (Alonso-González et al., 2022), which
68 coincide with periods of low solar radiation (Pomeroy et al., 2015; Musselman et al., 2017a).

69

70 The Mediterranean basin is a region that is critically affected by climate change (Giorgi, 2006)
71 being densely populated (>500 million inhabitants) and affected by an intense anthropogenic
72 activity. Warming of the Mediterranean basin will accelerate for the next decades, and
73 temperatures will continue to increase in this region during the warm months (Knutti and
74 Sedlacek, 2013; Lionello and Scarascia 2018; Cramer et al., 2018; Evin et al., 2021; Cos et al.,
75 2022), increasing atmospheric evaporative demands (Vicente-Serrano et al., 2020), drought
76 severity (Tramblay et al., 2020), leading to water-scarcity over most of this region (García-Ruiz
77 et al., 2011). Mediterranean mid-latitude mountains, such as the Pyrenees, where this research
78 focuses, are the main runoff generation zones of the downstream areas (Viviroli and Weingartner,
79 2004) and provide most of the water used by major cities in the lowlands (Morán-Tejeda et al.,
80 2014).

81

82 Snow patterns in the Pyrenees have high spatial diversity (Alonso-González et al., 2019), due to
83 internal climate variability of mid-latitude precipitation (Hawkins and Sutton 2010; Deser et al.,
84 2012), high interannual and decadal variability of precipitation in the Iberian Peninsula (Esteban-
85 Parra et al., 1998; Peña-Angulo et al., 2020) as well as the abrupt topography and the different
86 mountain exposure to the Atlantic air masses (Bonsoms et al., 2021a). Thus, snow accumulation
87 per season is almost twice as much in the northern slopes as in the southern slopes (Navarro-
88 Serrano and López-Moreno, 2017), and there is a high interannual variability of snow in regions
89 at lower elevations (Alonso-González et al., 2020a) and in the southern and eastern regions of the
90 Pyrenees (Salvador-Franch et al., 2014; Salvador-Franch et al., 2016; Bonsoms et al., 2021b).
91 Since the 1980s, the energy available for snow ablation has significantly increased in the Pyrenees
92 (Bonsoms et al., 2022), and winter snow days and snow accumulation have non-statically
93 significantly increased (Buisan et al., 2016; Serrano-Notivoli et al., 2018; López-Moreno et al.,
94 2020a; Bonsoms et al., 2021a) due to the increasing frequency of positive west and south-west

95 advections (Buisan et al., 2016). 21st century climate projections for Pyrenees anticipate a
96 temperature increase of more than 1°C to 4°C (relative to 1986–2005), and an increase (decrease)
97 of precipitation by about 10% for the eastern (western) regions during winter and spring (Amblar-
98 Frances et al., 2020). Therefore, changes in snow patterns in regions with high elevations are
99 uncertain because winter snow accumulation is affected by precipitation (López-Moreno et al.,
100 2008) and Mediterranean basin winter precipitation projections have uncertainties up to 80% of
101 the total variance (Evin et al., 2021).

102

103 Previous studies in the central Pyrenees (López-Moreno et al., 2013), Iberian Peninsula Mountain
104 ranges (Alonso-González et al., 2020a), and mountain areas that have Mediterranean climates
105 (López-Moreno et al., 2017) demonstrated that snowpack sensitivity to changes in climate ~~are~~
106 ~~mostly~~are mostly controlled by elevation. Despite the impact of climate warming in mountain
107 hydrological processes, there is limited understanding of the snow sensitivity to temperature and
108 precipitation changes and seasonality of mid-latitude Mediterranean mountain snowpacks. Some
109 studies reported different snowpack sensitivities during wet and dry years (López-Moreno et al.,
110 2017; Musselman et al., 2017b; Rasouli et al., 2022; Roche et al., 2018). However, the sensitivity
111 of snow during periods when there are seasonal compound weather (temperature and
112 precipitation) conditions has not yet been analyzed. The high interannual variability of the
113 Pyrenean snowpack, which is expected to increase according to climate projections (López-
114 Moreno et al., 2008), indicates a need to examine snowpack sensitivity to temperature and
115 precipitation change focusing on the year-to-year variability. Warm seasons in the Mediterranean
116 basin require special attention because these are likely to increase in the future (e.g., Vogel et al.,
117 2019; De Luca et al., 2020; Meng et al., 2022). Further, the occurrence of different HS trends at
118 mid- and high-elevation areas of this range (López-Moreno et al., 2020a) suggest that elevation
119 and spatial factors contribute to the wide variations of the sensitivity of snow to the climate.

120

121 Therefore, the main objective of this research is to quantify snow (accumulation, ablation, and
122 timing) sensitivity to temperature and precipitation change during compound temperature and
123 precipitation seasons in the Pyrenees.


124

125 **2 Geographical area and climate setting**

126

127 The Pyrenees is a mountain range located in the north of the Iberian Peninsula (south Europe;
128 42°N-43°N to 2°W-3°E) that is aligned east-to-west between the Atlantic Ocean and the

129 Mediterranean Sea. The highest elevations are in the central region (Aneto, 3404 m) and
130 elevations decrease towards the west and east (Figure 1). The Mediterranean basin, including the
131 Pyrenees, is in a transition area, and is influenced by the continental climate and the subtropical
132 temperate climate. Precipitation is mostly driven by large-scale circulation patterns (Zappa et al.,
133 2015; Borgli et al., 2019), the jet-stream oscillation during winter (Hurrell, 1995), and land-sea
134 temperature differences (Tuel and Eltahir, 2020). During the summer, the northward movement
135 of the Azores high pressure region brings stable weather, and precipitation is mainly convective
136 at that time (Xercavins, 1985). Precipitation is highly variable depending on mountain exposure
137 to the main circulation weather types; it ranges from about 1000 mm/year to about 2000 mm/year
138 (in the mountain summits), with lower levels in the east and south (Cuadrat et al., 2007). There is
139 a slight disconnection of the general climate circulation towards the eastern Pyrenees, where the
140 Mediterranean climate and East Atlantic/West Russia (EA-WR) oscillations have greater effects
141 on snow accumulation (Bonsoms et al., 2021a). In the southern, western, and central massifs of
142 the range, the Atlantic climate and the negative North Atlantic Oscillation (NAO) phases regulate
143 snow accumulation (W and SW wet air flows; López-Moreno, 2005; López-Moreno and Vicente-
144 Serrano, 2007; Buisan et al., 2016; Alonso-González et al., 2020b). In the northern slopes, the
145 positive phases of the Western Mediterranean Oscillation (WeMO) linked with NW and N
146 advectations trigger the most episodes of snow accumulation (Bonsoms et al., 2021a). The seasonal
147 snow accumulation in the northern slopes is almost double the amount (about 500 cm more) as in
148 the southern slopes at an elevation of about 2000 m (Bonsoms et al., 2021a). The
149 temperature/elevation gradient is about 0.55°C/100 m (Navarro-Serrano and López-Moreno,
150 2018) and the annual 0°C isotherm is at about 2750 to 2950 m (López-Moreno and García-Ruiz,
151 2004). Net radiation and latent heat flux governs the energy available for snow ablation; the
152 former heat flux increases at high elevations and the latter towards the east (Bonsoms et al., 2022).
153
154



155

156 **Figure 1** (a) Study area. Pyrenean massifs grouped by sectors for (b) low, (c) mid and (d) high
 157 elevation. The white dots indicate the locations of the automatic weather stations (AWS) shown
 158 at Table 1. Massifs delimitation is based on the spatial regionalization of the SAFRAN system,
 159 which groups massifs according to topographical and meteorological characteristics (modified
 160 from Durand et al., 1999).

161

162 3 Data and methods

163

164 3.1 Snow model

165

166 Snowpack was modelled using a physical-based snow model, the Flexible Snow Model (FSM2;
 167 Essery, 2015). This model resolves the SEB and mass balance to simulate the state of the
 168 snowpack. FSM2 is open access and available at <https://github.com/RichardEssery/FSM2> (last
 169 access 16 December 2022). Previous studies tested the FSM2 (Krinner et al., 2018), and its
 170 application in different forest environments (Mazzoti et al., 2021), and hydro-climatological
 171 mountain zones such the Andes (Urrutia et al., 2019), Alps (Mazzoti et al., 2020), Colorado
 172 (Smyth et al., 2022), Himalayas (Pritchard et al., 2020), Iberian Peninsula Mountains (Alonso-
 173 González et al., 2020a; Alonso-González et al., 2022), Lebanese mountains (Alonso-González et
 174 al., 2021), providing confidential results. The FSM2 requires forcing data of precipitation, air
 175 temperature, relative humidity, surface atmospheric pressure, wind speed, incoming shortwave
 176 radiation (SW_{inc}), and incoming long wave radiation (LW_{inc}). We have evaluated different FSM2
 177 model configurations (not shown) without remarkable differences in the accuracy and

178 performance metrics. Thus, the FSM2 configuration included in this work estimates snow cover
 179 fraction based on a linear function of HS and albedo based on a prognostic function, with increases
 180 due to snowfall and decreases due to snow age. Atmospheric stability is calculated as function of
 181 the Richardson number. Snow density is calculated as a function of viscous compaction by
 182 overburden and thermal metamorphism. Snow hydrology is estimated by gravitational drainage,
 183 including internal snowpack processes, runoff, refreeze rates, and thermal conductivity. Table S1
 184 summarizes the FSM2 configuration and the FSM2 compile numbers.

185

186 3.2 Snow model validation

187 FSM2 configuration was validated by in situ snow records of four automatic weather stations
 188 (AWSs) that were at high elevations in the Pyrenees. Precipitation in mountainous and windy
 189 regions is usually affected by undercatch (Kochendorfer et al., 2020). Thus, the instrumental
 190 records of precipitation were corrected for undercatch by applying an empirical equation validated
 191 for the Pyrenees (Buisan et al., 2019). Precipitation type was classified by a threshold method
 192 (Musselman et al., 2017b; Corripio et al., 2017): snow when the air temperature was below 1°C
 193 and rain when the air temperature was above 1°C, according to previous research in the study area
 194 (Corripio et al., 2017). The LW_{inc} heat flux of the AWSs (Table 1) were estimated as previously
 195 described (Corripio et al., 2017). Due to the wide instrumental data coverage (99.3% of the total
 196 dataset), gap-filling was not performed. The HS records were measured each 30 min using an
 197 ultrasonic snow depth sensor. The meteorological data used in the validation process were
 198 provided and managed by the local meteorological service of Catalonia
 199 ([https://www.meteo.cat/wpweb/serveis/formularis/peticio-dinformes-i-dades-
 200 meteorologiques/peticio-de-dades-meteorologiques/](https://www.meteo.cat/wpweb/serveis/formularis/peticio-dinformes-i-dades-meteorologiques/peticio-de-dades-meteorologiques/); data requested: 14/01/2021). Quality-
 201 checking of the data was performed using an automatic error filtering process in combination with
 202 a climatological, spatial, and internal coherency control defined by the SMC (2011).

203

204

Table 1. Characteristics of the four AWSs.

Area	Code	Lat/Lon°	Elevation (m)	Atlantic Ocean, Distance (km)	Mediterranean Sea, Distance (km)	Validation period (years)	Years
Central-Pyrenees, Northern slopes	A1	42.77/0.73	2228	200	190	2004–2020	16
	A2	42.61/0.98	2266	225	170	2001–2020	19
Eastern Pyrenees, Southern slopes	A3	42.46/1.78	2230	295	115	2005–2020	15

Eastern Pre-Pyrenees, Northern slopes	A4	42.29/1.71	2143	300	110	2009–2020	11
--	----	------------	------	-----	-----	-----------	----

205

206 Model accuracy was estimated based on the mean absolute error (MAE) and the root mean square

207 error (RMSE), and model performance was estimated by the coefficient of determination (R^2).

208 The MAE and the RMSE indicate the mean differences of the modelled and observed values.

209

210 3.3 Atmospheric forcing data

211

212 We forced the FSM2 with the open access climate reanalysis dataset provided by Vernay et al.

213 (2021), which consists of the modelled values from the SAFRAN meteorological analysis. The

214 FSM2 was run at an hourly resolution for each massif, each elevation range, and each climate

215 baseline and perturbation scenario from 1980 to 2019. The SAFRAN system provides data for

216 homogeneous meteorological and topographical mountain massifs every 300 m, from 0 to 3600

217 m (Durand et al., 1999; Vernay et al., 2021). We analyzed three elevation bands: low (1500 m),

218 middle (1800 m), and high (2400 m). Precipitation type was classified using the same threshold

219 approach used for model validation. Atmospheric emissivity was derived from the SAFRAN

220 LW_{inc} and air temperature. SAFRAN was forced using numerical weather prediction models

221 (ERA-40 reanalysis data from 1958 to 2002 and ARPEGE from 2002 to 2020). Meteorological

222 data were calibrated, homogenized, and improved by in situ meteorological observations data

223 assimilation (Vernay et al., 2021). Durand et al. (1999; 2009a; 2009b) provided further technical

224 details of the SAFRAN system. Previous studies used the SAFRAN system for the long-term HS

225 trends (López-Moreno et al., 2020), extreme snowfall (Roux et al., 2021), and snow ablation

226 analysis (Bonsoms et al., 2022). SAFRAN system has been extensively validated for the

227 meteorological modelling of continental Spain (Quintana-Seguí et al., 2017), France (Vidal et al.,

228 2010) or alpine snowpack climate projections (Verfaillie et al., 2018), among other works.

229

230 3.4 Snow sensitivity to temperature and precipitation change analysis

231

232 Snow sensitivity to temperature and precipitation change was analyzed using a delta-change

233 methodology (López-Moreno et al., 2008; Beniston et al., 2016; Musselman et al., 2017b; Marty

234 et al., 2017; Alonso-González et al., 2020a; Sanmiguel-Valladolid et al., 2022). In this method, air

235 temperature and precipitation were perturbed for each massif and elevation range based the

236 historical period (1980–2019). Air temperature was increased from 1 to 4°C at 1°C intervals,

237 assuming an increase of LW_{inc} accordingly (Jennings and Molotch, 2020). Precipitation was

238 changed from -10% to $+10\%$ at 10% intervals, in accordance with climate model uncertainties
239 and the maximum and minimum precipitation projections for the Pyrenees (Amblar-Frances et
240 al., 2020).

241

242 **3.5 Snow climate indicators**

243

244 Snow sensitivity to temperature and precipitation change was analyzed using five key indicators:
245 (i) seasonal average HS, (ii) seasonal maximum absolute HS peak (peak HS max), (iii) date of the
246 maximum HS (peak HS date), (iv) number of days with HS > 1 cm on the ground (snow duration),
247 and (v) daily average snow ablation per season (snow ablation, hereafter). Snow ablation was
248 calculated as the difference between the maximum daily HS recorded on two consecutive days
249 (Musselman et al., 2017a), and only days with decreases of 1 cm or more were recorded. Some
250 seasons had more than one peak HS; for this reason, peak HS date was determined after applying
251 a moving average of 5-days. All indicators were computed according to massif and elevation
252 range.

253

254 **3.6 Definitions of compound temperature and precipitation seasons**

255

256 The snow season was from October 1 to June 30 (inclusive). Snow duration was defined by snow
257 onset and snow ablation dates in situ observations (Bonsoms, 2021a), and results from the
258 baseline scenario snow duration presented in this work. A “compound temperature and
259 precipitation season” (season type) was assessed based on each massif and the elevation historical
260 climate record (1980–2019) using a joint quantile approach (Beniston and Goyette, 2007;
261 Beniston, 2009; López-Moreno et al., 2011a). Compound season types were defined according to
262 López-Moreno et al. (2011a), based on the seasonal 40th percentiles (T40 for temperature and P40
263 for precipitation) and the seasonal 60th percentiles (T60 and P60). There were four types of
264 seasons based on seasonal temperature (T_{season}) and seasonal precipitation (P_{season}) data:

265 Cold and Dry (CD): $T_{\text{season}} \leq T40$ and $P_{\text{season}} \leq P40$;

266 Cold and Wet (CW): $T_{\text{season}} \leq T40$ and $P_{\text{season}} \geq P60$;

267 Warm and Dry (WD): $T_{\text{season}} > T40$ and $P_{\text{season}} \leq P40$;

268 Warm and Wet (WW): $T_{\text{season}} > T60$ and $P_{\text{season}} > P60$.

269 All remaining seasons were classified as having average (Avg) temperature and precipitation.

270 Note that the number of compound season type is different depending on the Pyrenees massif
271 (Figure S1). However, by applying the joint-quantile approach described, we are comparing the
272 snow sensitivity to temperature and precipitation change between similar climate conditions,
273 independently where each compound season type was recorded.

274

275 3.7 Spatial regionalization

276

277 We have examined spatial differences in the snow sensitivity to temperature and precipitation
278 change by compound season types. ~~Following previous studies, massifs were grouped into four~~
279 ~~sectors by applying a Principal Component Analysis (PCA) (i.e., López-Moreno et al., 2020b;~~
280 ~~Matiu et al., 2020, among others). We applied a PCA over HS data for each month, year, massif,~~
281 ~~and elevation. Massifs were grouped into four sectors depending on the maximum correlation to~~
282 ~~PC1 and PC2 scores (see Figures S2). Massifs were grouped into four sectors by applying a~~
283 ~~Principal Component Analysis (PCA) of HS data (i.e., López-Moreno et al., 2020b; Matiu et al.,~~
284 ~~2020) and for each elevation depending on PC1 and PC2 scores. PCA scores are shown at Figure~~
285 ~~S2, whereas~~ the number of season types per sector are shown at Figure S3 and the spatial
286 regionalization is presented at Figure 1.

287

288

289 4. Results

290


291 We validated the FSM2 at Section 4.1. Subsequently, we analyzed the snow sensitivity to
292 temperature and precipitation change based on five snow climate indicators, namely the seasonal
293 HS, peak HS max, peak HS date, snow duration and snow ablation. Compound season types show
294 similar relative importance on the snow sensitivity to temperature and precipitation change
295 regardless of the Pyrenean sector. For this reason, our results have been focused on seasonal snow
296 changes due to increments of temperature, elevation, and compound season type. These are the
297 key factors that ruled the snow sensitivity to temperature and precipitation change, and an accurate
298 analysis is provided at Section 4.2. Spatial differences on the snow sensitivity to temperature and
299 precipitation change during compound season types are examined at Section 4.3.

300


301 4.1 Snow model validation

302

303 Our snow model validation analysis (Figures 2 and 3) confirmed that FSM2 accurately reproduces
304 the observed HS values. On average, the FSM2 had a R^2 greater than 0.83 for all stations. In
305 general, the snow model slightly overestimated the maximum HS values. The highest R^2 values
306 were at A4 and A2 ($R^2 = 0.85$ in both stations), and the lowest were at A3 and A1 ($R^2 = 0.79$ and
307 $R^2 = 0.82$, respectively). The highest accuracy was at A4 (RMSE = 18.5 cm, MAE = 8.9 cm), and
308 the largest errors were at A2 (RMSE = 45.8 cm, MAE = 29.0 cm).




309



310

311 **Figure 2.** Time series of the observed (gray) and modelled (black) HS values at the four AWSs.

312



313

314

315 **Figure 3.** Regression analysis of observed (x-axis) and simulated (y-axis) HS values.

316

317 **4.2 Snow sensitivity to temperature and precipitation change**


318

319 We then determined seasonal HS profiles for each perturbed climate scenario and compound
 320 season type (Figure 4). The results show a non-linear response between seasonal HS loss and

321 temperature increase. When the temperature increased at 1°C intervals, the largest relative
 322 seasonal HS decrease from the baseline was at + 1°C for all elevations and all compound season
 323 type. High elevation areas had lower seasonal HS variability between compound season types
 324 than low elevations (Figure S4). At low elevations, snow was- greater during CW seasons than
 325 other seasons. All the snowpack-perturbed scenarios indicated that snowpack decreased for all
 326 elevations under warming climate scenarios. Snowpack sensitivity to temperature and
 327 precipitation change depended on the compound season type (Figures 5 and 6). At low elevations,
 328 the seasonal changes in HS ranged from -37% (WW) to -28% (CD) per °C increase. For mid-
 329 elevation ranges, there were no remarkable differences among compound season types (Table 2),
 330 and the seasonal HS changes ranged from -34% (WW) to -30% (CW) per °C increase. Low and
 331 mid-elevations had greater snowpack reductions than high elevations. In the latter, a 10% increase
 332 of precipitation counterbalanced a temperature increase of about 1°C, and there were no
 333 remarkable differences in the seasonal HS from the baseline scenario especially in the coldest
 334 months of the season (Figure S5 and Figure S6). The maximum seasonal HS sensitivity to
 335 temperature and precipitation was during WD seasons (27%/°C), and the minimum was during
 336 CW seasons (-22%/°C).

337

338



339

340 **Figure 4.** Anomalies of seasonal HS for low, mid and high elevation (rows), compound season type
 341 (columns), and different temperature increases (colors).


342 **Table 2.** Average and seasonal HS and peak HS sensitivity to temperature and precipitation
 343 change during the four different compound temperature and precipitation seasons at three
 344 different elevations.

Season type	%HS/ °C			%peak HS max/°C		
	Low	Mid	High	Low	Mid	High
Avg.	-33	-33	-25	-20	-20	-16
CD	-28	-30	-22	-17	-17	-14
CW	-33	-32	-22	-22	-20	-15
WD	-32	-30	-27	-19	-16	-16
WW	-37	-34	-26	-24	-24	-16

345
 346 At low and mid elevations, the peak HS max was greatest during WW seasons (-24%/°C) and
 347 lowest during the CD and WD seasons (-17%/°C for both). At high elevations, there were no
 348 clear differences in the peak HS max for the different seasons. The maximum peak HS max was
 349 during WD seasons (-16%/°C) and the minimum was during CD seasons (-14%/°C).

350
 351 We also determined average seasonal snow duration for each elevation range and compound
 352 season type for different temperature increases (Table 3 and Figure 5c). The minimum snow
 353 duration was during CW seasons (-13%/°C at low elevations, -10%/°C at mid-elevations, -5%/°C
 354 at high elevations). At low elevations, the snow duration was most sensitive during WW seasons
 355 (-17%/°C). On the contrary, at mid-elevations and high elevations, the snow duration was most
 356 sensitive during WD seasons (-13%/°C at mid-elevations and -8%/°C at high elevations).

357



358

359 **Figure 5.** Anomalies of seasonal HS (a), peak HS max (b) and snow duration (c) for different
 360 temperature increases relative to baseline at three different elevations during the four different
 361 compound season types. The solid black lines within each boxplot are the average. Lower and
 362 upper hinges correspond to the 25th and 75th percentiles, respectively. The whisker is a horizontal
 363 line at 1.5 interquartile range of the upper quartile and lower quartile, respectively. Dots represent
 364 the outliers. Data is grouped by season, compound season type, increment of temperature,
 365 precipitation variation, elevation, and massif.
 366

367 The peak HS date occurred earlier due to warming, independently of precipitation changes.
 368 During WD seasons, the peak HS date per °C was anticipated by 3 days at low elevations, 3 days
 369 at mid-elevations, and 6 days at high elevations; during CD seasons, the peak HS date per °C was
 370 anticipated by 4 days at low elevations, 5 days at mid-elevations, and 9 days at high elevations.
 371 In low and mid elevation areas, if the temperature increase was no more than about 1°C above

372 baseline, there was little change in the peak HS date (Figure 6). In addition, the minimum peak
373 HS date change is found during WW seasons (Table 3), because the snowpack would be scarce
374 at those times, and there were no defined peaks (Figure S4).

375

376 We determined the snow ablation sensitivity to temperature and precipitation change in response
377 to different temperature increases at different elevations and during different compound season
378 types. The results show there were low differences in absolute snow ablation values in a warmer
379 climate (Figure 7). At low elevations, the average snow ablation sensitivity to temperature and
380 precipitation change in all four compound seasons was 12%/°C (Table 3). At mid-elevations and
381 high elevations, the maximum snow ablation sensitivity to temperature and precipitation change
382 was during dry seasons; WD seasons had a snow ablation sensitivity to temperature and
383 precipitation change of 13%/°C at mid-elevations and 10%/°C at high elevations. On the other
384 hand, the minimum values for mid-elevations were during WW seasons, when the snow ablation
385 sensitivity to temperature and precipitation change was 8%/°C; the minimum values at high
386 elevations were during CW seasons, when was 5%/°C.

387

388



389

390 **Figure 6.** Difference (days) from baseline Peak HS date at three different elevations and during
 391 the four different temperature (colors) and precipitation shifts (columns) for each season (boxes).


392

393 **Table 3.** Snow duration, snow ablation, and peak HS date sensitivity to temperature and
 394 precipitation change during the four different compound season types.

Season Type	Snow duration (%/°C)			Snow ablation (%/°C)			Peak HS date (days/°C)		
	Low	Mid	High	Low	Mid	High	Low	Mid	High
Avg.	-15	-12	-6	12	11	7	-1	-3	-7
CD	-13	-11	-5	12	13	8	-4	-5	-9
CW	-13	-10	-5	12	10	5	-2	-3	-13
WD	-16	-13	-8	12	13	10	-3	-3	-6
WW	-17	-13	-7	12	8	7	-1	0	-3

395

396




397
 398 **Figure 7.** Absolute snow ablation values (cm/day) (y-axis) at three different elevations during
 399 four different compound temperature and precipitation for baseline and different increments of
 400 temperature (x-axis). seasons.

401
 402 **4.3 Spatial patterns**

403
 404 PCA analysis reveals four Pyrenean sectors, namely northern-western (NW), northern-eastern
 405 (NE), southern-western (SW), and southern-eastern (SE). No remarkable differences between
 406 sectors are found in the relative importance of each compound season type in the snow sensitivity
 407 to temperature and precipitation change (Figure 8). Snow sensitivity to temperature and
 408 precipitation change absolute values are generally lower at northern slopes (NW and NE) than at
 409 the southern slopes (SW and SE) (Figure S7 and Figure S8). In detail, seasonal HS ranged from
 410 $-26\%/^{\circ}\text{C}$ during CD (NW) to $-36\%/^{\circ}\text{C}$ during WW (SE). Similarly, the maximum peak HS max
 411 sensitivity to temperature and precipitation change was at SE during WW seasons ($25\%/^{\circ}\text{C}$) and
 412 the minimum was during CD seasons at NW ($15\%/^{\circ}\text{C}$). The snow duration sensitivity to
 413 temperature and precipitation change increased during WW seasons, and the maximum changes
 414 were at SE ($-16\%/^{\circ}\text{C}$); in contraposition, the lowest sensitivity to temperature and precipitation
 415 change are found at NW, during CD and CW seasons ($-8\%/^{\circ}\text{C}$, in both seasons). Snow ablation
 416 sensitivity to temperature and precipitation change increases towards the eastern Pyrenees,
 417 particularly during WD seasons ($14\%/^{\circ}\text{C}$ and $13\%/^{\circ}\text{C}$ for NE and SE, respectively). Finally, no
 418 remarkable peak HS date differences are observed between sectors and maximum values are

419 found during CD and CW seasons, when the peak HS date is anticipated ≥ 5 per $^{\circ}\text{C}$ for all
 420 sectors.
 421



422
 423 **Figure 8.** Average snow sensitivity to temperature and precipitation change (y-axis) grouped by
 424 sector (x-axis), season type (color bars) and snow climate indicator (boxes).

425
 426 **5. Discussion**

427
 428 The spatial and temporal patterns of snow in the Pyrenees are highly variable, and climate
 429 projections indicate that extreme events will likely increase during future decades (Meng et al.,
 430 2022). Therefore, we analyzed factors that affect the snowpack sensitivity to temperature and
 431 precipitation change gain insight into how future climate changes may affect the snow regime.
 432

433 **5.1 Snow sensitivity to temperature and precipitation change and relationship with**
 434 **historical and future snow trends**

435
 436 **5.1.1 Snow accumulation phase**

437

438 The snow losses due warming that we described here are mainly associated with increases in the
439 rain/snowfall ratio (Figure S9), changes in the snow onset and offset dates (Figure S4), and
440 increases in the energy available for snow ablation during the later months of the snow season, as
441 it was previously reported by literature (e.g., Pomeroy et al., 2015; Lynn et al., 2020; Jennings
442 and Molotch, 2020). At high elevations, a trend of increasing precipitation (+10%) could
443 counterbalance temperature increases ($<1^{\circ}\text{C}$; Figure S5), consistent with the results previously
444 reported for specific sites of the central Pyrenees (Izas, 2000m; López-Moreno et al., 2008).
445 Rasouli et al. (2014) also found that a 20% increase of precipitation could compensate for 2°C
446 increase of temperature in subarctic Canada. A climate sensitivity analysis in the western
447 Cascades (western USA) found that increases of precipitation due to warming modulated the
448 snowpack accumulation losses by about $5\%/1^{\circ}\text{C}$ (Minder, 2010). These results are consistent with
449 recent data that examined snow above 1000 m in the Pyrenees, which found that an increase in
450 the frequency of west circulation weather types since the 1980s increased the HS (Serrano-
451 Notivoli et al., 2018; López-Moreno et al., 2020), snow accumulation (Bonsoms et al., 2021a),
452 and changes in winter snow days (Buisan et al., 2016). There are similar trends in the Alps, with
453 an increase of extreme (exceeding the 100-year return level) snowfall above 3000 m during recent
454 decades (Roux et al., 2021) and increases in extreme winter precipitation (Rajczak and Schär,
455 2017).

456

457 **5.1.2 Snow ablation phase**

458

459 Climate warming leads to a cascade of physical changes in the SEB that increase snow ablation
460 near the 0°C isotherm. On overall, the snow ablation showed low to inexistent changes due to
461 warming. Comparison between low and high elevations indicated slightly faster snow ablation at
462 high elevations (Figure 7). This higher rate of snow ablation per season at high elevations (which
463 have deeper snowpacks) are probably because the snow there lasts until late spring, when more
464 energy is available for snow ablation (Bonsoms et al., 2022). Temperature increase does not imply
465 remarkable changes in snow ablation per season because warming decreases the magnitude of the
466 snowpack (seasonal HS and peak HS max) and triggers an earlier onset of snowmelt (Wu et al.,
467 2018). The earlier peak HS date (Table 3 and Figure 6) implies lower rates of net shortwave
468 radiation, because snow melting starts earlier in warmer climates (Pomeroy et al., 2015),
469 coinciding with the shorter days and lower solar zenith angle (Lundquist et al., 2013; Sanmiguel-
470 Vallelado et al., 2022). Our results agree with the slow snow melt rates reported in the Northern
471 Hemisphere from 1980 to 2017 (Wu et al., 2018). The results of previous studies were similar for
472 subarctic Canada (Rasouli et al., 2014) and western USA snowpacks (Musselman et al., 2017b),
473 but Arctic sites had faster melt rates (Krogh and Pomeroy, 2019).

474

475 **5.1.3 Snow sensitivity to temperature and precipitation change and snowpack projections**

476

477 Our results suggest that warming had a non-linear effect on snowpack reduction. Our largest snow
478 losses were for seasonal HS when the temperature increased by 1°C above baseline. At low and
479 mid elevations, the average seasonal HS decrease was more than 40% for all compound season
480 types, and the maximum sensitivity was during WW seasons. Previous research in the Pyrenees
481 and other mid-latitude mountain ranges reported similar results. A study in the central Pyrenees
482 reported the peak SWE was 29%/°C, whereas snow season duration decreased by about 20 to 30
483 days at about 2000 m (López-Moreno et al., 2013). The average peak HS max at high elevations
484 in the Pyrenees (-16 %/°C; Figure 6 and Table 2), was similar to the average peak SWE sensitivity
485 (-15%/°C) reported in the Iberian Peninsula mountains at 2500 m (Alonso-González et al.,
486 2020a). These results are also consistent with climate projections for this mountain range. In
487 particular, for a 2°C or more increase of temperature, the snow season declined by 38% at the
488 lowest ski resorts (~1500 m) in the SE Pyrenees (Pons et al., 2015). However, high emission
489 climate scenarios projected an increase in the frequency and intensity of high snowfall at high
490 elevations (López-Moreno et al., 2011b). Snow sensitivity in the easternmost areas could decline
491 during the winter because of a trend for an increase of about 10% in precipitation in this area
492 (Amblar-Francés et al., 2020). Our projected changes in the Pyrenean snowpack dynamics are
493 similar to the expected snow losses in other mountain ranges. For example, a study of the Atlas
494 Mountains of northern Africa concluded that snowpack decreases were greater in the lowlands
495 and projected seasonal SWE declines of 60% under the RCP4.5 scenario and 80% under the
496 RCP8.5 scenario for the entire range (Tuel et al., 2022). A study in the Washington Cascades
497 (western USA) found that snowpack decline was 19 to 23% per 1°C (Minder, 2010), similar to
498 the values in the present study at high elevations. A study of the French Alps (Chartreuse, 1500
499 m) found that seasonal HS decreases on the order of 25% for a 1.5°C increase and 32% for a 2°C
500 increase of global temperature above the pre-industrial years (Verfaillie et al., 2018). A study of
501 the Swiss Alps reported a snowpack decrease of about 15%/°C (Beniston, 2003); in the same
502 alpine country, another study predicted the seasonal HS will decrease by more than 70% in
503 massifs below 1000 m in all future climate projections (Marty et al., 2017). The largest snow
504 reductions will likely occur during the periods between seasons (Steger et al., 2013; Marty et al.,
505 2017). Nevertheless, at high elevations, snow climate projections found no significant trend for
506 maximum HS until the end of the 21st century above 2500 m in the eastern Alps (Willibald et al.,
507 2021), suggesting that internal climate variability is a major source of uncertainty of SWE
508 projections at high elevations (Schirmer et al., 2021).

509

510 **5.2 Influence of compound temperature and precipitation seasons**

511

512 We found that the maximum sensitivities of seasonal HS and peak HS max to temperature and
513 precipitation change were during WW seasons at low and mid-elevations and during WD seasons
514 at high elevations. Brown and Mote (2008) analyzed the sensitivity of snow to climate changes
515 in the Northern Hemisphere and found maximal SWE sensitivities in mid-latitude maritime
516 winter climate areas, and minimal SWE sensitivities to temperature and precipitation change in
517 dry and continental zones, consistent with our results. López-Moreno et al. (2017) also found
518 greater decreases of SWE in wet and temperate Mediterranean ranges than in drier regions.
519 Furthermore, Rasouli et al. (2022) studied the northern North American Cordillera and found
520 higher snowpack sensitivities to temperature and precipitation change in wet basins than dry
521 basins. Our maximum snow ablation relative change over the baseline scenario occurred during
522 WD seasons, in accordance with Musselman et al. (2017b), who found a higher snowmelt rate
523 during dry years in the western USA. Low and mid-elevations are highly sensitive to WW seasons
524 because wet conditions favor decreases in the seasonal HS due to advection from sensible heat
525 fluxes. The temperature in the Pyrenees is still cold enough to allow snowfall at high elevations
526 during WW seasons, and for this reason we found maximal sensitivities to temperature and
527 precipitation change during WD seasons. Reductions of snowfall in alpine regions can be
528 compensated in a warmer scenario, because warm and wet snow is less susceptible to blowing
529 wind transport and losses from sublimation (Pomeroy and Li, 2000; Pomeroy et al., 2015). During
530 spring, snow runoff could be also greater in wet climates due to rain-on-snow events (Corripio et
531 al., 2017), coinciding with the availability of more energy for snow ablation.

532

533 **5.3 Spatial and elevation factors controlling snow sensitivity to temperature and** 534 **precipitation change**

535

536 Comparison between Pyrenean sectors (Figure 8) reveals no remarkable differences in the relative
537 importance of each compound season type in the snow sensitivity to temperature and precipitation
538 change. This is because by applying a joint-quantile approach for each massif and elevation, we
539 are comparing similar climate seasons between sectors, regardless of the number of compound
540 season types recorded in each massif during the baseline period (Figure S1 and S3). The highest
541 absolute snow sensitivity to temperature and precipitation change values is found in the SE
542 Pyrenees. This is consistent with the snow accumulation and ablation patterns previously reported
543 in this region (Lopez-Moreno, 2005; Navarro-Serrano et al., 2018; Alonso-González et al., 2020a;
544 Bonsoms et al., 2021a; Bonsoms et al., 2021b; Bonsoms et al., 2022). The Atlantic climate has
545 less of an influence in the SE sector, and in situ observations indicated there was about half of the

546 seasonal snow accumulation amounts as in northern and western areas at the same elevation
547 (>2000 m; Bonsoms et al., 2021a). The snow in the SE Pyrenees is more sensitive to temperature
548 and precipitation change because these massifs are exposed to higher turbulence and radiative
549 heat fluxes (Bonsoms et al., 2022), and 0°C isotherm is closer. Similar conclusions are found for
550 low elevations, where the results show an upward displacement of the snow line due to warming.
551 Previous studies described the sensitivity of the snow pattern to elevation at specific stations of
552 the central Pyrenees (López-Moreno et al., 2013; 2017), Iberian Peninsula mountains (Alonso-
553 González et al., 2020a), and other ranges such as the Cascades (Jefferson, 2011; Sproles et al.,
554 2013), the Alps (Marty et al., 2017), and western USA (Pierce et al., 2013; Musselman et al.,
555 2017b). In these regions, the models suggest larger snowpack reductions due to warming at
556 subalpine sites than at alpine sites (Jennings and Molotch, 2020) due to closer isothermal
557 conditions (Brown and Mote, 2009; Lopez-Moreno et al., 2017; Mote et al., 2018).

558

559 **5.4 Environmental and socioeconomic implications**

560

561 Our results indicated there will be an increase of snow ablation days and imply a disappearance
562 of the typical sequence of snow accumulation and snow ablation seasons. Climate warming
563 triggers the simultaneous occurrence of several periods of snowfall and melting, snow droughts
564 during the winter, and ephemeral snowpacks between seasons. These expected decreases in snow
565 will likely have important impacts on the ecosystem. During spring, a snow cover cools the soil
566 (Luetschg et al., 2008), delays the initiation of freezing (Oliva et al., 2014), functions as a thick
567 active layer (Hrbáček et al., 2016), and protects alpine rocks from exposure to solar radiation and
568 high air temperatures (Magnin et al., 2017). Due to warming temperatures, the remaining glaciers
569 in the Pyrenees are shrinking and are expected to disappear before the 2050s (Vidaller et al.,
570 2021). The shallower snowpack that we identified in this work will increase the vulnerability of
571 glaciers, because snow has a higher albedo than dark ice and debris-covered glaciers and functions
572 as a protective layer for glaciers (Fujita and Sakai, 2014).

573

574 The earlier onset of snowmelt suggested by our results, which is greater at low and mid-elevations
575 during WD seasons, is in line with previous global studies that reported earlier streamflow due to
576 earlier runoff dates (Adam et al., 2009; Stewart, 2009), and with a study of changes in the Iberian
577 Peninsula River flows (Morán-Tejeda et al., 2014). Overall, our results are consistent with the
578 slight decrease of the river peak flows that have occurred in the southern slopes of the Pyrenees
579 since the 1980s (Sanmiguel-Vallelado et al., 2017). The reductions of seasonal HS that we
580 identified, suggest that snowmelt-dominated stream flows will likely shift to rainfall dominated
581 regimes. Although high elevation meltwater might increase and contribute to earlier groundwater

582 recharging (Evans et al., 2018), the increased evapotranspiration in the lowlands (Bonsoms et al.,
583 2022) could counter this effect, so there is no net change in downstream areas (Stahl et al., 2010).
584 Snow ephemerality triggers lower spring and summer flows (Barnett et al., 2005; Adam et al.,
585 2009; Stahl et al., 2010) and has an impact on the hydrological management strategies. Winter
586 snow accumulation affects hydrological availability during the months when water and
587 hydroelectric demands are higher. This is because reservoirs store water during periods of peak
588 flows (winter and spring), and release water during the driest season in the lowlands (summer)
589 (Morán-Tejeda et al., 2014). Recurrent snow-scarce seasons may intensify these hydrological
590 impacts and lead to competition for water resources among different ecological and
591 socioeconomic systems. The economic viability of mountain ski-resorts in the Pyrenees depends
592 on a regular deep snow cover (Gilaberte-Burdalo et al., 2014; Pons et al., 2015), but this is highly
593 variable, especially at low and mid-elevations. The expected increase in snow-scarce seasons that
594 we identified here is consistent with climate projections for this region, which suggest that no
595 Pyrenean ski resorts will be viable under RCP 8.5 scenario by the end of the 21st century (Spandre
596 et al., 2019).

597

598 **5.5 Limitations and uncertainties**

599 The meteorological input data that we used to model snow were estimated for flat slopes and the
600 regionalization system we used was based on the SAFRAN system. According to this system, a
601 mountain range is divided into massifs with homogeneous topography. The SAFRAN system has
602 negative biases in shortwave radiation, a temperature precision of about 1 K, and biases in the
603 accumulated monthly precipitation of about 20 kg/m² (Vernay et al., 2021). Our estimates of snow
604 sensitivity to temperature and precipitation change were based on the delta-approach, which
605 considers changes in temperature and precipitation based on climate projections for the Pyrenees
606 (Amblar-Francés et al., 2020), but assumes that the meteorological patterns of the reference period
607 will be constant over time. In this work we used a physical-based snow model since it provides
608 better results for future snow climate change estimations than degree-day models (Carletti et al.,
609 2022). The FSM2 is a physics-based model of intermediate complexity, and the estimates of snow
610 densification are simpler than those from more complex models of snowpack. However, a more
611 complex model does not necessarily provide better performance in terms of snowpack and runoff
612 estimation (Magnusson et al., 2015). The FSM2 configuration implemented in this work includes
613 snow meltwater retention, snowpack refreezing and snow albedo based on snow age, which are
614 the physical parameters included in the best-performing snow models according to Essery et al.
615 (2013). Snow model sensitivity studies reveal that intermediate complexity models exhibit similar

616 snow depth accuracies than most complex multi-layer snow models, as well as robust
617 performances across seasons (Terzago et al., 2020).

618

619 **6 Conclusions**

620

621 Our study assessed the impact of temperature and precipitation change on the Pyrenean snowpack
622 during compound cold-hot and wet-dry seasons, using a physical-based snow model that was
623 forced by reanalysis data. We determined the snow sensitivity to temperature and precipitation
624 change using five key indicators of snow accumulation and snow ablation. The lowest snow
625 sensitivity to temperature and precipitation change was at high elevations of the NW Pyrenees
626 and increased at lower elevations and in the SE slopes. An increase of 1°C at low and mid
627 elevation regions led to remarkable decreases in the seasonal HS and snow duration. However, at
628 high elevations, precipitation plays a key role, and temperature is far from the isothermal 0°C
629 during the middle of winter. In this region, a 10% increase of precipitation, as suggested by ~~many~~
630 ~~climate projections~~ [the Spanish Meteorological Agency \(AEMET\)](#) over the eastern regions of this
631 range, could compensate for temperature increases on the order of about < 1°C. The impact of
632 climate warming depends on the combination of temperature and precipitation during compound
633 seasons. Our analysis of seasonal HS and peak HS max indicated the greatest declines were during
634 WW seasons and the smallest declines were during CD seasons, independently of the Pyrenean
635 sector. For snow duration, however, the highest (lowest) sensitivity to temperature and
636 precipitation change is found during WD (CW) seasons. Similarly, snow ablation had slightly
637 greater sensitivities to temperature and precipitation change during WD seasons, in that snow
638 ablation variation is less than 10% and the peak HS date occurred about 5 days earlier per °C. Our
639 findings thus provide evidence that the Pyrenean snowpack is highly sensitive to climate warming
640 and suggest that the snowpacks of other mid-latitude mountain ranges may also show similar
641 response to warming.

642

643 **Data availability**

644

645 Snow model (FSM2) is open access and provided by Essery (2015) and available at
646 <https://github.com/RichardEssery/FSM2> (last access 16 December 2022). Climate forcing data is
647 provided by Vernay et al. (2021), through AERIS ([https://www.aeris-data.fr/landing-](https://www.aeris-data.fr/landing-page/?uuid=865730e8-edeb-4c6b-ae58-80f95166509b#v2020.2)
648 [page/?uuid=865730e8-edeb-4c6b-ae58-80f95166509b#v2020.2](https://www.aeris-data.fr/landing-page/?uuid=865730e8-edeb-4c6b-ae58-80f95166509b#v2020.2); last access 16 December 2022).
649 Data of this work is available upon request (contact: josepbonsoms5@ub.edu).

650

651 **Acknowledgements**

652

653 This work falls within the research topics examined by the research group “Antarctic, Artic,
654 Alpine Environments-ANTALP” (2017-SGR-1102) funded by the Government of Catalonia,
655 HIDROIBERNIEVE (CGL2017-82216-R) and MARGISNOW (PID2021-124220OB-100), from
656 the Spanish Ministry of Science, Innovation and Universities. JB is supported by a pre-doctoral
657 University Professor FPI grant (PRE2021-097046) funded by the Spanish Ministry of Science,
658 Innovation and Universities. The authors are grateful to Marc Oliva, who reviewed an early
659 version of this manuscript. We acknowledge the SAFRAN data provided by Météo-France –
660 CNRS and the CNRM Centre d’Etudes de la Neige, through AERIS.

661

662 **Author contributions**

663

664 JB analyzed the data and wrote the original draft. JB, JILM and EAG contributed to the
665 manuscript design and draft editing. JB, JILM and EAG read and approved the final manuscript.

666

667 **References**

668

669 Adam, J. C., and Hamlet, A. F.: Implications of Global Climate Change for Snowmelt Hydrology
670 in the Twenty First Century, *Hydrological Processes*, 23(7), 962-972,
671 <https://doi.org/10.1002/hyp.7201>, 2009.

672 Alonso-González, E., Gutmann, E., Aalstad, K., Fayad, A., Bouchet, M., and Gascoin, S.:
673 Snowpack dynamics in the Lebanese mountains from quasi-dynamically downscaled ERA5
674 reanalysis updated by assimilating remotely sensed fractional snow-covered area, *Hydrol. Earth
675 Syst. Sci.*, 25, 4455–4471, <https://doi.org/10.5194/hess-25-4455-2021>, 2021.

676 Alonso-González, E., López-Moreno, J.I., Navarro-Serrano, F., Sanmiguel-Valladolid, A.,
677 Aznárez-Balta, M., Revuelto, J., and Ceballos, A.: Snowpack Sensitivity to Temperature,
678 Precipitation, and Solar Radiation Variability over an Elevational Gradient in the Iberian
679 Mountains, *Atmos. Res.*, 243, 104973 <https://doi.org/10.1016/j.atmosres.2020.104973>, 2020a.

680 Alonso-González, E., López-Moreno, J.I., Navarro-Serrano, F., Sanmiguel-Valladolid, A.,
681 Revuelto, J., Domínguez-Castro, F., and Ceballos, A.: Snow climatology for the mountains in the
682 Iberian Peninsula using satellite imagery and simulations with dynamically downscaled reanalysis
683 data, *International Journal of Climatology*, 40(1), 477–491, <https://doi.org/10.1002/joc.6223>,
684 2019.

685 Alonso-González, E., López-Moreno, J.I., Navarro-Serrano, F.M., and Revuelto, J.: Impact of
686 North Atlantic oscillation on the snowpack in Iberian Peninsula mountains, *Water*, 12, 105–276,
687 <https://doi.org/10.3390/w12010105>, 2020b.

688 Alonso-González, E., Revuelto, J., Fassnacht, S.R., and López-Moreno, J.I.: Combined influence
689 of maximum accumulation and melt rates on the duration of the seasonal snowpack over
690 temperate mountains, *Journal of Hydrology*, 608, 127574,
691 <https://doi.org/10.1016/j.jhydrol.2022.127574>, 2022.

692 Amblar-Francés, M.P., Ramos-Calzado, P., Sanchis-Lladó, J., Hernanz-Lázaro, A., Peral- García,
693 M.C., Navascués, B., Dominguez-Alonso, M., and Rodríguez-Camino, E.: High resolution
694 climate change projections for the Pyrenees region, *Adv. Sci. Res.*, 17, 191–208,
695 <https://doi.org/10.5194/asr-17-191-2020>, 2020.

696 Armstrong, A. and Brun, E.: *Snow and Climate, Physical Processes, Surface Energy Exchange*
697 *and Modeling*, Cambridge University press, 222 pp., 1998.

698 Barnard, D. M., Knowles, J. F., Barnard, H. R., Goulden, M. L., Hu, J., Litvak, M. E., and
699 Molotch, N. P.: Reevaluating growing season length controls on net ecosystem production in
700 evergreen conifer forests, *Scientific Reports*, 8(1), 17973, [https://doi.org/10.1038/s41598-018-](https://doi.org/10.1038/s41598-018-36065-0)
701 [36065-0](https://doi.org/10.1038/s41598-018-36065-0), 2018.

702 Barnett, T. P., Adam, J. C., and Lettenmaier, D. P.: Potential impacts of a warming climate on
703 water availability in snow-dominated regions, *Nature*, 438(7066), 303–309,
704 <https://doi.org/10.1038/nature04141>, 2005.

705

706 Beniston, M., and Stoffel, M.: Rain-on-snow events, floods and climate change in the Alps: events
707 may increase with warming up to 4°C and decrease thereafter, *Sci. Total Environ.*, 571, 228–36,
708 <https://doi.org/10.1016/j.scitotenv.2016.07.146>, 2016.

709 Beniston, M.: Trends in joint quantiles of temperature and precipitation in Europe since 1901 and
710 projected for 2100, *Geophysical Research Letters*, 36, L07707,
711 <https://doi.org/10.1029/2008GL037119>, 2009.

712 Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A.,
713 Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J.I., Magnusson, J.,
714 Marty, C., Morán-Tejeda, E., Morin, S., Naaim, M., Provenzale, A., Rabatel, A., Six, D., Stötter,
715 J., Strasser, U., Terzago, S., and Vincent, C.: The European mountain cryosphere: a review of its
716 current state, trends, and future challenges, *The Cryosphere*, 12, 759–794,
717 <https://doi.org/10.5194/tc-12-759-2018>, 2018.

718 Beniston, M., and Goyette, S.: Changes in variability and persistence of climate in Switzerland:
719 exploring 20th century observations and 21st century simulations, *Global and Planetary Change*,
720 57, 1–20, <https://doi.org/10.1016/j.gloplacha.2006.11.004>, 2007.

721 Beniston, M.; Keller, F.; Ko, B., and Goyette, S.: Estimates of snow accumulation and volume in
722 the Swiss Alps under changing climatic conditions, *Theor. Appl. Climatol.*, 76, 125–140.
723 <https://doi.org/10.1007/S00704-003-0016-5>, 2003.

724 Bonsoms, J., González, S., Prohom, M., Esteban, P., Salvador-Franch, F., López- Moreno, J.I.,
725 and Oliva, M.: Spatio-temporal patterns of snow in the Catalan Pyrenees (SE Pyrenees, NE
726 Iberia), *Int. J. Climatol.*, 41 (12), 5676–5697, <https://doi.org/10.1002/joc.7147>, 2021a.

727 Bonsoms, J., López-Moreno, J.I., González, S., and Oliva, M.: Increase of the energy available for
728 snow ablation and its relation with atmospheric circulation, *Atmospheric Research*, 275, 106228,
729 <https://doi.org/10.1016/j.atmosres.2022.106228>, 2022.

730 Bonsoms, J., Salvador-Franch, F., and Oliva, M.: Snowfall and snow cover evolution in the
731 Eastern Pre-Pyrenees (NE Iberian Peninsula), *Cuad. Investig. Geogr.*, 47 (2), 291–307,
732 <https://doi.org/10.18172/cig.4879>, 2021b.

733 Brown, R.D. and Mote, P.W.: The response of Northern Hemisphere snow cover to a changing
734 climate, *Journal of Climate*, 22(8), 2124–2145, <https://doi.org/10.1175/2008JCLI2665.1>, 2009.

735 Buisan, S., Collado Aceituno, J. L. and Tierra, J.: ¿Se mide bien la precipitación en forma de
736 nieve?, <https://doi.org/10.31978/639-19-010-0.095>, 2019.

737 Buisan, S.T., López-Moreno, J.I., Saz, M.A. and Kochendorfer, J.: Impact of weather type
738 variability on winter precipitation, temperature and annual snowpack in the Spanish Pyrenees,
739 *Climate Research*, 69(1), 79–92. <https://doi.org/10.3354/cr01391>, 2016.

740 Carletti, F., Michel, A., Casale, F., Bocchiola, D., Lehning, M., and Bavay, M.: A comparison of
741 hydrological models with different level of complexity in Alpine regions in the context of climate
742 change, *Hydrol. Earth Syst. Sci. Discuss.* <https://doi.org/10.5194/hess-26-3447-2022>, 2022.

743 Cooper, A. E., Kirchner, J. W., Wolf, S., Lombardozi, D. L., Sullivan, B. W., Tyler, S. W., &
744 Harpold, A. A.: Snowmelt causes different limitations on transpiration in a Sierra Nevada conifer
745 forest, *Agricultural and Forest Meteorology*, 291, 108089.
746 <https://doi.org/10.1016/j.agrformet.2020.108089>, 2020.

747 Corripio, J., and López-Moreno, J.I.: Analysis and predictability of the hydrological response of
748 mountain catchments to heavy rain on snow events: a case study in the Spanish Pyrenees,
749 *Hydrology*, 4(2), 20, <https://doi.org/10.3390/hydrology4020020>, 2017.

750 Cos, J., Doblas-Reyes, F., Jury, M., Marcos, R., Bretonnière, P.-A., and Samsó, M.: The
751 Mediterranean climate change hotspot in the CMIP5 and CMIP6 projections, *Earth Syst. Dynam.*,
752 13, 321–340, <https://doi.org/10.5194/esd-13-321-2022>, 2022.

753 Cramer W, Guiot J, Fader M, Garrabou J, Gattuso J-P, Iglesias A, Lange MA, Lionello P, Llasat
754 MC, Paz S, Peñuelas J, Snoussi M, Toreti A, Tsimplis MN, and Xoplaki E.: Climate change and
755 interconnected risks to sustainable development in the Mediterranean, *Nat. Clim. Chang.* 8(11),
756 972–980, <https://doi.org/10.1038/s41558-018-0299-2>, 2018.

757 Cuadrat, J., Saz, M.A., Vicente-Serrano, S.: *Atlas climático de Aragón*. Gobierno de Aragón,
758 Zaragoza, 222 pp. 2007.

759 De Luca, P., Messori, G., Faranda, D., Ward, P. J., and Coumou, D.: Compound warm-dry and
760 cold-wet events over the Mediterranean, *Earth System Dynamics*, 11, 793–805,
761 <https://doi.org/10.5194/esd-11-793-2020>, 2020.

762 Deser, C., Phillips, A., Bourdette, V., and Teng, H.: Uncertainty in climate change projections:
763 the role of internal variability, *Climate dynamics*, 38(3), 527–546, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-010-0977-x)
764 010-0977-x, 2012.

765 Durand, Y., Giraud, G., Brun, E., Méridol, L., and Martin, E.: A computer-based system
766 simulating snowpack structures as a tool for regional avalanche forecasting, *J. Glaciol.*, 45, 469–
767 484, <https://doi.org/10.1017/S0022143000001337>, 1999.

768 Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Méridol, L., and Lesaffre, B.: Reanalysis
769 of 47 Years of Climate in the French Alps (1958–2005): Climatology and Trends for Snow Cover,
770 *J. Appl. Meteorol. Clim.*, 48, 2487–2512, <https://doi.org/10.1175/2009JAMC1810.1>, 2009a.

771 Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Méridol, L., and Lesaffre, B.: Reanalysis
772 of 44 Yr of Climate in the French Alps (1958–2002): Methodology, Model Validation,
773 Climatology, and Trends for Air Temperature and Precipitation., *J. Appl. Meteorol. Clim.*, 48,
774 429–449, <https://doi.org/10.1175/2008JAMC1808.1>, 2009b.

775 Essery, R.: A factorial snowpack model (FSM 1.0), *Geoscientific Model Development*, 8(12),
776 3867–3876, <https://doi.org/10.5194/gmd-8-3867-2015>, 2015.

777 Essery, R., Morin, S., Lejeune, Y., and Ménard, C.: A comparison of 1701 snow models using
778 observations from an alpine site, *Adv. Water Res.*, 55, 131–
779 148, <https://doi.org/10.1016/j.advwatres.2012.07.013>, 2013.

780 Esteban-Parra, M.J, Rodrigo, F.S. and Castro-Diez, Y.: Spatial and temporal patterns of
781 precipitation in Spain for the period 1880-1992, *Int. J. Climatol.*, 18, 1557–74, 1998.

782 Evans, S.G., Ge, S., Voss, C.I. and Molotch, N.P. The role of frozen soil in groundwater discharge
783 predictions for warming alpine watersheds, *Water Resour. Res.*, 54, 1599–1615.
784 <https://doi.org/10.1002/2017WR022098>, 2018.

785 Evin, G.; Somot, S.; Hingray, B. Balanced estimate and uncertainty assessment of European
786 climate change using the large EURO-CORDEX regional climate model ensemble, *Earth Syst.*
787 *Dyn. Discuss.*, 12(4), 1543–1569, <https://doi.org/10.5194/esd-12-1543-2021>, 2021.

788 Fujita, K. and Sakai, A.: Modelling runoff from a Himalayan debris-covered glacier, *Hydrol.*
789 *Earth Syst. Sci.*, 18, 2679–2694, <https://doi.org/10.5194/hess-18-2679-2014>, 2014.

790 García-Ruiz, J. M., López-Moreno, J. I., Vicente-Serrano, S. M., Lasanta-Martínez, T. and
791 Beguería, S. Mediterranean water resources in a global change scenario, *Earth Sci. Rev.*, 105(3–
792 4), 121–139, <https://doi.org/10.1016/j.earscirev.2011.01.006>, 2011.

793 Gilaberte-Burdalo, M., Lopez-Martin, F., M. R. Pino-Otin, M., and Lopez-Moreno, J.: Impacts of
794 climate change on ski industry, *Environ. Sci. Pol.*, 44, 51–
795 61, <https://doi.org/10.1016/j.envsci.2014.07.003>, 2014.

796 Giorgi, F.: Climate change hot-spots, *Geophysical Research Letters*, 33: L08707,
797 <https://doi.org/10.1029/2006GL025734>, 2006.

798 Gribovski, Z., Szilágyi, J., and Kalicz, P.: Diurnal fluctuations in shallow groundwater levels
799 and streamflow rates and their interpretation – A review, *J. Hydrol.*, 385, 371–
800 383, <https://doi.org/10.1016/j.jhydrol.2010.02.001>, 2010.

801 Hall, A.: Role of surface albedo feedback in climate. *J. Clim.*, 17, 1550-1568, 2004.

802 Hammond, J. C., Saavedra, F. A. and Kampf, S. K.: Global snow zone maps and trends in snow
803 persistence 2001–2016, *Int. J. Climatol.*, 38, 4369–4383, <https://doi.org/10.1002/joc.5674>, 2018.

804 Hawkins, E., and Sutton, R.: The potential to narrow uncertainty in projections of regional
805 precipitation change, *Clim Dyn.*, <https://doi.org/10.1007/s00382-010-0810-6>, 2010.

806 Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., Jachson, M., K`a`ab, A.,
807 Kang, S., Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., Steltzer, H., High mountain
808 areas. In: Portner, H.-O., Roberts, D.C., Masson- Delmotte, V., et al. (Eds.), *IPCC Special Report*
809 *on the Ocean and Cryosphere in a Changing Climate*. [https://www.ipcc.ch/srocc/chapter/chapter-](https://www.ipcc.ch/srocc/chapter/chapter-2/)
810 *2/*, 2019.

811 Hrbáček, F., Láska, K., and Engel, Z.: Effect of snow cover on the active-layer thermal regime
812 — a case study from James Ross Island, Antarctic Peninsula, *Permafrost and Periglac. Process.*,
813 27, 307– 315, <https://doi.org/10.1002/ppp.1871>, 2016.

814 Hurrell, J. W.: Decadal trends in the North Atlantic oscillation: Regional temperatures and
815 precipitation, *Science*, 269, 676–679, <https://doi.org/10.1126/science.269.5224.676>, 1995.

816 Jefferson, A. J.: Seasonal versus transient snow and the elevation dependence of climate
817 sensitivity in maritime mountainous regions, *Geophys. Res. Lett.*, 38, L16402,
818 <https://doi.org/10.1029/2011GL048346>, 2011.

819 Jennings, K.S., and Molotch, N.P.: Snowfall fraction, cold content, and energy balance changes
820 drive differential response to simulated warming in an alpine and subalpine snowpack. *Front.*
821 *Earth Sci*, 8, 2296-6463, <https://doi.org/10.3389/feart.2020.00186>, 2020.

822 Klein, G., Vitasse, Y., Rixen, C., Marty, C., and Rebetez, M.: Shorter snow cover duration since
823 1970 in the Swiss Alps due to earlier snowmelt more than to later snow onset, *Clim. Chang.* 139,
824 637–649. <https://doi.org/10.1007/s10584-016-1806-y>, 2016.

825 Knutti, R. and Sedlacek, J.: Robustness and uncertainties in the new CMIP5 climate model
826 projections, *Nature Climate Change*, 3, 369–373, <https://doi.org/10.1038/nclimate1716>, 2013.

827 Kochendorfer, J., M.E. Earle, D. Hodyss, A. Reverdin, Y. Roulet, R. Nitu, R. Rasmussen, S.
828 Landolt, S. Buisan, and Laine, T.: Undercatch adjustments for tipping bucket gauge
829 measurements of solid precipitation, *J. Hydrometeor.*, 21, 1193–1205,
830 <https://doi.org/10.1175/JHM-D-19-0256.1>, 2020.

831 Krinner, G., Derksen, C., Essery, R., Flanner, M., Hagemann, S., Clark, M., Hall, A., Rott, H.,
832 Brutel-Vuilmet, C., Kim, H., Ménard, C. B., Mudryk, L., Thackeray, C., Wang, L., Arduini, G.,
833 Balsamo, G., Bartlett, P., Boike, J., Boone, A., Chéruy, F., Colin, J., Cuntz, M., Dai, Y.,
834 Decharme, B., Derry, J., Ducharme, A., Dutra, E., Fang, X., Fierz, C., Ghattas, J., Gusev, Y.,
835 Haverd, V., Kontu, A., Lafaysse, M., Law, R., Lawrence, D., Li, W., Marke, T., Marks, D.,
836 Ménégoz, M., Nasonova, O., Nitta, T., Niwano, M., Pomeroy, J., Raleigh, M. S., Schaedler, G.,
837 Semenov, V., Smirnova, T. G., Stacke, T., Strasser, U., Svenson, S., Turkov, D., Wang, T.,
838 Wever, N., Yuan, H., Zhou, W., and Zhu, D.: ESM-SnowMIP: assessing snow models and
839 quantifying snow-related climate feedbacks, *Geosci. Model Dev.*, 11, 5027–
840 5049, <https://doi.org/10.5194/gmd-11-5027-2018>, 2018.

841 Krogh, S.A., and Pomeroy, J.W.: Impact of Future Climate and Vegetation on the Hydrology of
842 an Arctic Headwater Basin at the Tundra–Taiga Transition, *J. Hydrometeorol.*, 20, 197–215.
843 <https://doi.org/10.1175/JHM-D-18-0187.1>, 2019.

844 Lionello, P. and Scarascia, L.: The relation between climate change in the Mediterranean region
845 and global warming, *Reg. Environ. Change*, 18, 1481–1493, [https://doi.org/10.1007/s10113-018-](https://doi.org/10.1007/s10113-018-1290-1)
846 1290-1, 2018.

847 López Moreno, J.I., and Garcia Ruiz, J.M.: Influence of snow accumulation and snowmelt on
848 streamflow in the Central Spanish Pyrenees, *International. J. Hydrol. Sci.*, 49, 787–802,
849 <https://doi.org/10.1623/hysj.49.5.787.55135>, 2004.

850 López-Moreno, J.I.: Recent variations of snowpack depth in the central Spanish Pyrenees, *Arct.*
851 *Antarct. Alp. Res.*, 37, 253–260, [https://doi.org/10.1657/1523-0430\(2005\)037, 2005](https://doi.org/10.1657/1523-0430(2005)037, 2005).

852 López-Moreno, J.I., Gascoin, S., Herrero, J., Sproles, E.A., Pons, M., Alonso-González, E.,
853 Hanich, L., Boudhar, A., Musselman, K.N., Molotch, N.P., Sickman, J., and Pomeroy, J.:
854 Different sensitivities of snowpacks to warming in Mediterranean climate mountain areas,
855 *Environ. Res. Lett.*, 12 (7), 074006, <https://doi.org/10.1088/1748-9326/aa70cb>, 2017.

856 Lopez-Moreno, J.I., Goyette, S., Beniston, M., and Alvera, B.: Sensitivity of the snow energy
857 balance to climate change: Implications for the evolution of snowpack in Pyrenees in the 21st
858 century, *Climate Research* 36(3), 203–217, <https://doi.org/10.3354/cr00747>, 2008.

859 López-Moreno, J.I., Goyette, S., Vicente-Serrano, S.M., and Beniston, M.: Effects of climate
860 change on the intensity and frequency of heavy snowfall events in the Pyrenees, *Clim. Chang.*,
861 105, 489–508. <https://doi.org/10.1007/s10584-010-9889-3>, 2011b.

862 López-Moreno, J.I., Pomeroy, J.W., Alonso-González, E., Morán-Tejeda, E., and Revuelto, J.:
863 Decoupling of warming mountain snowpacks from hydrological regimes, *Environ. Res. Lett.*, 15,
864 11–15, <https://doi.org/10.1088/1748-9326/abb55f>, 2020a.

865 López-Moreno, J.I., Pomeroy, J.W., Revuelto, J., and Vicente-Serrano, S.M.: Response of snow
866 processes to climate change: spatial variability in a small basin in the Spanish Pyrenees, *Hydrol.*
867 *Process.*, 27, 2637–2650. <https://doi.org/10.1002/hyp.9408>, 2013.

868 López-Moreno, J.I., and Vicente-Serrano, S.M.: Atmospheric circulation influence on the
869 interannual variability of snowpack in the Spanish Pyrenees during the second half of the
870 twentieth century, *Nord. Hydrol.*, 38 (1), 38–44, <https://doi.org/10.2166/nh.2007.030>, 2007.

871 López-Moreno, J.I., Vicente-Serrano S.M., Morán-Tejeda E., Lorenzo J., Kenawy, A. and
872 Beniston, M.: NAO effects on combined temperature and precipitation winter modes in the
873 Mediterranean mountains: Observed relationships and projections for the 21st century, *Global*
874 *and Planetary Change*, 77, 72–66, <https://doi.org/10.1016/j.gloplacha.2011.03.003>, 2011a.

875 López-Moreno, J.I., Soubeyroux, J.M., Gascoin, S., Alonso-González, E., Durán- Gómez, N.,
876 Lafaysse, M., Vernay, M., Carmagnola, C., and Morin, S.: Long-term trends (1958–2017) in snow
877 cover duration and depth in the Pyrenees, *Int. J. Climatol.*, 40, 6122–6136,
878 <https://doi.org/10.1002/joc.6571>, 2020b.

879 López-Moreno J.I, Revuelto, J, Gilaberte, M., Morán-Tejeda, E., Pons, M., Jover, E., Esteban, P.,
880 García, C., and Pomeroy, J.W.: The effect of slope aspect on the response of snowpack to climate

881 warming in the Pyrenees, *Theoretical and Applied Climatology*, 117, 1–13,
882 <https://doi.org/10.1007/s00704-013-0991-0>, 2013.

883 López-Moreno, J, Pomeroy, J.W, Revuelto, J., Vicente-Serrano, S.M. Response of snow
884 processes to climate change: spatial variability in a small basin in the Spanish Pyrenees,
885 *Hydrological Processes*, 27(18), 2637–2650, <https://doi.org/10.1002/hyp.9408>, 2013

886 López-Moreno, J.; Boike, J.; Sanchez-Lorenzo, A. and Pomeroy, J.: Impact of climate warming
887 on snow processes in Ny-Ålesund, a polar maritime site at Svalbard, *Glob. Planet. Chang.*, 146,
888 10–21, <https://doi.org/10.1016/j.gloplacha.2016.09.006>, 2016.

889 Luetschg, M., Lehning, M., and Haeberli, W.: A sensitivity study of factors influencing warm/thin
890 permafrost in the Swiss Alps, *J. Glaciol.*, 54, 696–704.
891 <https://doi.org/10.3189/002214308786570881>, 2008

892 Lundquist, J.D., Dickerson-Lange, S.E., Lutz, J.A., and Cristea, N.C.: Lower forest density
893 enhances snow retention in regions with warmer winters: a global framework developed from
894 plot-scale observations and modeling, *Water Resour. Res.*, 49, 6356–6370.
895 <https://doi.org/10.1002/wrcr.20504>, 2013.

896 Lynn, E., Cuthbertson, A., He, M., Vasquez, J. P., Anderson, M. L., Coombe, P., et al. Technical
897 note: Precipitation-phase partitioning at landscape scales to regional scales, *Hydrology and Earth
898 System Sciences*, 24(11), 5317–5328, <https://doi.org/10.5194/hess-24-5317-2020>, 2020.

899 Magnin, F., Westermann, S., Pogliotti, P., et al.: Snow control on active layer thickness in steep
900 alpine rock walls (Aiguille du Midi, 3842 ma.s.l., Mont Blanc massif), *Catena*, 149, 648–662,
901 <https://doi.org/10.1016/j.catena.2016.06.006>, 2017.

902 Marshall, A. M., Link, T. E., Abatzoglou, J. T., Flerchinger, G. N., Marks, D. G., and Tedrow,
903 L.: Warming alters hydrologic heterogeneity: Simulated climate sensitivity of hydrology-based
904 microrefugia in the snow-to-rain transition zone, *Water Resources Research*, 55, 2122–2141,
905 <https://doi.org/10.1029/2018WR023063>, 2019.

906 Marty, C., Schlögl, S., Bavay, M., and Lehning, M.: How much can we save? Impact of different
907 emission scenarios on future snow cover in the Alps, *The Cryosphere*, 11, 517–529,
908 <https://doi.org/10.5194/tc-11-517-2017>, 2017.

909 Mazzotti, G., Essery, R., Moeser, D., and Jonas, T.: Resolving small-scale forest snow patterns
910 with an energy balance snow model and a 1-layer canopy, *Water Resources Research*, 56,
911 e2019WR026129, <https://doi.org/10.1029/2019WR026129>, 2020.

912 Mazzotti, G., Webster, C., Essery, R., and Jonas, T.: Increasing the physical representation of
913 forest-snow processes in coarse-resolution models: Lessons learned from upscaling hyper-

914 resolution simulations, *Water Resources Research*, 57(5), e2020WR029064, <https://doi.org/10.1029/2020WR029064>, 2021.

915

916 Meng, Y., Hao, Z., Feng, S., Zhang, X., Hao, F.: Increase in compound dry-warm and wet-warm
917 events under global warming in CMIP6 models, *Global and Planetary Change*, 210, 103773,
918 <https://doi.org/10.1016/j.gloplacha.2022.103773>, 2022.

919 Minder, J. R.: The Sensitivity of Mountain Snowpack Accumulation to Climate Warming, *Journal*
920 *of Climate*, 23(10), 2634–2650, <https://doi.org/10.1175/2009JCLI3263.1>, 2010.

921 Morán-Tejeda, E., Lorenzo-Lacruz, J., López-Moreno, J.I., Rahman, K. and Beniston, M.:
922 Streamflow timing of mountain rivers in Spain: Recent changes and future projections, *J. Hydrol.*
923 517, 1114–1127, <https://doi.org/10.1016/j.jhydrol.2014.06.053>, 2014.

924 Mote, P. W., Li, S., Lettenmaier, D. P., Xiao, M., and Engel, R.: Dramatic declines in snowpack
925 in the western US, *npj Clim. Atmos. Sci.*, 1, 2, <https://doi.org/10.1038/s41612-018-0012-1>, 2018.

926 Mote, P.W., A.F. Hamlet, M.P. Clark, and D.P. Lettenmaier.: Declining mountain snowpack in
927 western North America, *Bull. Am. Meteorol. Soc.*, 86, 39–49, [https://doi.org/10.1175/BAMS-86-](https://doi.org/10.1175/BAMS-86-1-39)
928 1-39, 2005.

929 Musselman, K., Clark, M., Liu, C., Ikeda, K., and Rasmussen, R.: Slower snowmelt in a warmer
930 world, *Nat. Clim. Change*, 7, 214–219, <https://doi.org/10.1038/NCLIMATE3225>, 2017a.

931 Musselman, K. N., Molotch, N. P., and Margulis, S. A.: Snowmelt response to simulated warming
932 across a large elevation gradient, southern Sierra Nevada, California, *Cryosphere*, 11, 2847–2866,
933 <https://doi.org/10.5194/tc-11-2847-2017>, 2017b.

934 Navarro-Serrano, F. and López-Moreno, J.I.: Spatio-temporal analysis of snowfall events in the
935 Spanish Pyrenees and their relationship to atmospheric circulation, *Cuad. Invest. Geogr.*, 43 (1),
936 233–254, <https://doi.org/10.18172/cig.3042>, 2017.

937 Notarnicola, C.: Hotspots of snow cover changes in global mountain regions over 2000–2018,
938 *Remote Sensing of Environment*, 243, 111781, <https://doi.org/10.1016/j.rse.2020.111781>, 2020.

939 Magnusson, J., Wever, N., Essery, R., Helbig, N., Winstral, A., and Jonas, T.: Evaluating snow
940 models with varying process representations for hydrological applications, *Water Resour. Res.*,
941 51, 2707–2723, <https://doi.org/10.1002/2014WR016498>, 2015.

942 Matiu, M., Crespi, A., Bertoldi, G., Carmagnola, C.M., Marty, C., Morin, S., Schöner, W., Cat
943 Berro, D., Chiogna, G., De Gregorio, L., Kotlarski, S., Majone, B., Resch, G., Terzago, S., Valt,
944 M., Beozzo, W., Cianfarra, P., Gouttevin, I., Marcolini, G., Notarnicola, C., Petitta, M., Scherrer,
945 S.C., Strasser, U., Winkler, M., Zebisch, M., Cicogna, A., Cremonini, R., Debernardi, A., Faletto,
946 M., Gaddo, M., Giovannini, L., Mercalli, L., Soubeyroux, J.-M., Susnik, A., Trenti, A., Urbani,
947 S., Weilguni, V., 2020. Observed snow depth trends in the European Alps 1971 to 2019. *Cryoph*
948 1-50. <https://doi.org/10.5194/tc-2020>, 2020.

949 Oliva, M., Gómez Ortiz, A., Salvador, F., Salvà, M. Pereira, P., and Geraldès, M.: Long-term soil
950 temperature dynamics in the Sierra Nevada, Spain. *Geoderma* 235-236, 170-181,
951 <https://doi.org/10.1016/j.geoderma.2014.07.012>, 2014.

952 Peña-Angulo, D., Vicente-Serrano, S., Domínguez-Castro, F., Murphy, C., Reig, F., Trambly,
953 Y., Trigo, R., Luna, M.Y., Turco, M., Noguera, I., Aznárez-Balta, M., Garcia-Herrera, R., Tomas-
954 Burguera, M. and Kenawy, A.: Long-term precipitation in Southwestern Europe reveals no clear
955 trend attributable to anthropogenic forcing, *Environmental Research Letters*, 15 (9), 094070,
956 <https://doi.org/10.1088/1748-9326/ab9c4f>, 2020.

957 Pierce, D. and Cayan, D.: The uneven response of different snow measures to human-induced
958 climate warming, *Journal of Climate*, 26, 4148–4167, [https://doi.org/10.1175/JCLI-D-12-](https://doi.org/10.1175/JCLI-D-12-00534.1)
959 [00534.1](https://doi.org/10.1175/JCLI-D-12-00534.1), 2013.

960 Pomeroy, J. W., and L. Li.: Prairie and arctic areal snow cover mass balance using a blowing
961 snowmodel, *J. Geophys. Res.*, 105(D21), 26619– 26634, <https://doi.org/10.1029/2000JD900149>,
962 2000.

963 Pomeroy, J. W., Fang, X and Rasouli, K.: Sensitivity of snow processes to warming in the
964 Canadian Rockies. 72nd Eastern Snow Conf., Sherbrooke, QC, Canada, Eastern Snow
965 Conference, 22–33, 2015.

966 Pons, M., López-Moreno, J., Rosas-Casals, M., and Jover, E.: The vulnerability of Pyrenean ski
967 resorts to climate-induced changes in the snowpack, *Climatic Change*, 131, 591–
968 605, <https://doi.org/10.1007/s10584-015-1400-8>, 2015.

969 Pritchard, D. M. W., Forsythe, N., O'Donnell, G., Fowler, H. J., and Rutter, N.: Multi-physics
970 ensemble snow modelling in the western Himalaya, *The Cryosphere*, 14(4), 1225–1244,
971 <https://doi.org/10.5194/tc-14-1225-2020>, 2020.

972 Quintana-Seguí, P., Turco, M., Herrera, S., and Miguez-Macho, G.: Validation of a new
973 SAFRAN-based gridded precipitation product for Spain and comparisons to Spain02 and ERA-
974 Interim, *Hydrol. Earth Sys. Sci.*, 21, 2187–2201, <https://doi.org/10.5194/hess-21-2187-2017>,
975 2017.

976 Rajczak, J. and Schär, C.: Projections of Future Precipitation Extremes Over Europe: A
977 Multimodel Assessment of Climate Simulations, *J. Geophys. Res.-Atmos.*, 122, 773–
978 710, <https://doi.org/10.1002/2017JD027176>, 2017.

979 Rasouli, K., J. W. Pomeroy, J. R. Janowicz, S. K. Carey, and T. J. Williams.: Hydrological
980 sensitivity of a northern mountain basin to climate change, *Hydrol. Processes*, 28, 4191–5208,
981 <https://doi.org/10.1002/hyp.10244>, 2014.

982 Rasouli, K.R., Pomeroy, J.W., and Marks, D.G.: Snowpack sensitivity to perturbed climate in a
983 cool mid-latitude mountain catchment, *Hydrol. Process.*, 29, 3925–3940.
984 <https://doi.org/10.1002/hyp.10587>, 2015.

985 Rasouli, K.R., Pomeroy, J.W., and Whietfiled, P.H.: The sensitivity of snow hydrology to changes
986 in air temperature and precipitation in three North American headwater basins, *J. Hydrol.*, 606,
987 127460, <https://doi.org/10.1016/j.jhydrol.2022.127460>, 2022

988 Roche, J.W., Bales, R.C., Rice, R., Marks, D.G.: Management Implications of Snowpack
989 Sensitivity to Temperature and Atmospheric Moisture Changes in Yosemite National Park. *J. Am.*
990 *Water Resour. Assoc.*, 54 (3), 724–741, <https://doi.org/10.1111/1752-1688.12647>, 2018.

991 Roux, E., Evin, G., Eckert, N., Blanchet, J., and Morin, S.: Elevation-dependent trends in extreme
992 snowfall in the French Alps from 1959 to 2019, *The Cryosphere*, 15, 4335–4356,
993 <https://doi.org/10.5194/tc-15-4335-2021>, 2021.

994 Salvador - Franch, F., Salvà, G., Vilar, F., and García, C.: Contribución al análisis nivométrico
995 del Pirineo Oriental: La Molina, período 1956 - 1996. En: X Congreso Internacional AEC: Clima,
996 sociedad, riesgos y ordenación del territorio, pp. 365-375, Alicante.
997 <http://hdl.handle.net/10045/58002>, 2016.

998 Salvador Franch, F., Salvà, G., Vilar, F., and García, C.: Nivometría y perfiles de innivación en
999 Núria (1970 m, Pirineo Oriental): 1985 - 2013. En: IX Congreso de la AEC, pp. 729 -738,
1000 Almería, <http://hdl.handle.net/20.500.11765/8229>, 2014.

1001 Sanmiguel-Vallelado, A., Morán-Tejeda, E., Alonso-González, E., and López-Moreno, J. I.:
1002 Effect of snow on mountain river regimes: An example from the Pyrenees, *Frontiers of Earth*
1003 *Science*, 11(3), 515–530. <https://doi.org/10.1007/s11707-016-0630-z>, 2017.

1004 Sanmiguel-Vallelado, A., McPhee, J., Esmeralda, P., Morán-Tejeda, E., Camarero, J., López-
1005 Moreno, J.I. Sensitivity of forest-snow interactions to climate forcing: Local variability in a
1006 Pyrenean valley, *Journal of Hydrol.*, <https://doi.org/10.1016/j.jhydrol.2021.127311>, 2022.

1007 Schirmer, M., Winstral, A., Jonas, T., Burlando, P., and Peleg, N.: Natural climate variability is
1008 an important aspect of future projections of snow water resources and rain-on-snow events, *The*
1009 *Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2021-276>, 2021.

1010 Scott, D, McBoyle G, and Mills B.: Climate change and the skiing industry in southern Ontario
1011 (Canada): exploring the importance of snowmaking as a technical adaptation, *Clim Res.*, 23(2),
1012 171–181, <https://doi.org/10.3354/CR023171>, 2003.

1013 Serrano-Notivoli, R., Buisan, S.T., Abad-Pérez, L.M., Sierra-Alvarez, E., Rodríguez-Ballesteros,
1014 C., López-Moreno, J.I. and Cuadrat, J.M.: Tendencias recientes en precipitación, temperatura y
1015 nieve de alta montaña en los Pirineos (Refugio de Góriz, Huesca). In: *El clima: aire, agua, tierra*
1016 *y fuego*. Madrid, Spain: Asociación Española de Climatología y Ministerio para la Transición

1017 Ecológica – Agencia Estatal de Climatología y Ministerio para la Transición Ecológica – Agencia
1018 Estatal de Meteorología, pp. 267, 1060–280, 2018.

1019 Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research
1020 synthesis, *Glob. Planet. Change*, 77, 85–96, <https://doi.org/10.1016/j.gloplacha.2011.03.004>,
1021 2011.

1022 Servei Meteorològic de Catalunya (SMC). Les estacions meteorològiques automàtiques (EMA).
1023 [https://static-m.meteo.cat/wordpressweb/wp-](https://static-m.meteo.cat/wordpressweb/wp-content/uploads/2014/11/18120559/Les_Estacions_XEMA.pdf)
1024 [content/uploads/2014/11/18120559/Les_Estacions_XEMA.pdf](https://static-m.meteo.cat/wordpressweb/wp-content/uploads/2014/11/18120559/Les_Estacions_XEMA.pdf)(accessed March 1, 2022). 2011.

1025 Smyth, E. J., Raleigh, M. S., and Small, E. E.: The challenges of simulating SWE beneath forest
1026 canopies are reduced by data assimilation of snow depth, *Water Resources Research*, 58,
1027 e2021WR030563, <https://doi.org/10.1029/2021WR030563>, 2022.

1028 Spandre, P., François, H., Verfaillie, D., Pons, M., Vernay, M., Lafaysse, M., George, E., and
1029 Morin, S. Winter tourism under climate change in the Pyrenees and the French Alps: relevance
1030 of snowmaking as a technical adaptation, *The Cryosphere*, 13, 1325–1347,
1031 <https://doi.org/10.5194/tc-13-1325-2019>, 2019.

1032 Sproles, E.A, Nolin, A.W, Rittger, K, and Painter, T. H.: Climate change impacts on maritime
1033 mountain snowpack in the Oregon Cascades, *Hydrology and Earth System Sciences*, 17(7), 2581–
1034 2597, <https://doi.org/10.5194/hess-17-2581-2013>, 2013.

1035 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L.M., van Lanen, H.A.J., Sauquet, E., Demuth, S.,
1036 Fendekova, M., and Jódar, J.: Streamflow trends in Europe: evidence from a dataset of near-
1037 natural catchments, *Hydrol. Earth. Syst. Sci.*, 14, 2367-2382, [https://doi.org/10.5194/hess-14-](https://doi.org/10.5194/hess-14-2367-2010)
1038 [2367-2010](https://doi.org/10.5194/hess-14-2367-2010), 2010.

1039 Steger, C., Kotlarski, S., Jonas, T., and Schär, C.: Alpine snow cover in a changing climate: A
1040 regional climate model perspective, *Clim. Dynam.*, 41, 735–754, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-012-1545-3)
1041 [012-1545-3](https://doi.org/10.1007/s00382-012-1545-3), 2013.

1042 Stewart I.T.: Changes in snowpack and snowmelt runoff for key mountain regions. *Hydrological*
1043 *Processes*, 23, 78–94, <https://doi.org/10.1002/hyp.7128>, 2009.

1044 Sturm, M., M. A. Goldstein, and C. Parr. Water and life from snow: A trillion dollar science
1045 question, *Water Resour. Res.*, 53, 3534– 3544, <https://doi.org/10.1002/2017WR020840>, 2017.

1046 Terzago, S., Andreoli, V., Arduini, G., Balsamo, G., Campo, L., Cassardo, C., Cremonese, E.,
1047 Dolia, D., Gabellani, S., Hardenberg, J., Morra di Cella, U., Palazzi, E., Piazza, G., Pogliotti, P.,
1048 and Provenzale, A.: Sensitivity of snow models to the accuracy of meteorological forcings in
1049 mountain environments, *Hydrol. Earth Syst. Sci.*, 24, 4061–4090, [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-24-4061-2020)
1050 [24-4061-2020](https://doi.org/10.5194/hess-24-4061-2020), 2020.

1051 Tramblay, Y.; Koutroulis, A.; Samaniego, L.; Vicente-Serrano, S.M.; Volaire, F.; Boone, A.; Le
1052 Page, M.; Llasat, M.C.; Albergel, C.; Burak, S.; et al.: Challenges for drought assessment in the
1053 Mediterranean region under future climate scenarios, *Earth Sci. Rev.*, 210, 103348,
1054 <https://doi.org/10.1016/j.earscirev.2020.103348>, 2020.

1055 Trujillo, E., and N. P. Molotch.: Snowpack regimes of the Western United States, *Water Resour.*
1056 *Res.*, 50(7), 5611–5623, <https://doi.org/10.1002/2013WR014753>, 2014.

1057 Tuel, A. and Eltahir, E. A. B.: Why Is the Mediterranean a Climate Change Hot Spot?, *J. Climate*,
1058 33, 5829–5843. <https://doi.org/10.1175/jcli-d-19-0910.1>, 2020.

1059 Tuel, A., El Moçayd, N., Hasnaoui, M. D., and Eltahir, E. A. B.: Future projections of High Atlas
1060 snowpack and runoff under climate change, *Hydrol. Earth Syst. Sci.*, 26, 571–588,
1061 <https://doi.org/10.5194/hess-26-571-2022>, 2022.

1062 Urrutia, J., Herrera, C., Custodio, E., Jódar, J., and Medina, A.: Groundwater recharge and
1063 hydrodynamics of complex volcanic aquifers with a shallow saline lake: Laguna Tuyajto, Andean
1064 Cordillera of northern Chile, *Sci. Total Environ.*, 697, 134116,
1065 <https://doi.org/10.1016/j.scitotenv.2019.134116>, 2019.

1066 Verfaillie, D., Lafaysse, M., Déqué, M., Eckert, N., Lejeune, Y., and Morin, S.: Multi-component
1067 ensembles of future meteorological and natural snow conditions for 1500 m altitude in the
1068 Chartreuse mountain range, Northern French Alps, *The Cryosphere*, 12, 1249–
1069 1271, <https://doi.org/10.5194/tc-12-1249-2018>, 2018.

1070 Vernay, M., Lafaysse, M., Monteiro, D., Hagenmuller, P., Nheili, R., Samacoits, R., Verfaillie,
1071 D., and Morin, S.: The S2M meteorological and snow cover reanalysis over the French
1072 mountainous areas, description and evaluation (1958–2020), *Earth Syst. Sci. Data Discuss*,
1073 <https://doi.org/10.5194/essd-2021-249>, 2021.

1074 Vicente-Serrano, S.M., McVicar, T., Miralles, D., Yang, Y. and Tomas-Burguera, M.:
1075 Unravelling the Influence of Atmospheric Evaporative Demand on Drought under Climate
1076 Dynamics, *Climate Change in press*, 11(2), 1757-7780, <https://doi.org/10.1002/wcc.632>, 2020.

1077 Vidal, J.-P., Martin, E., Franchistéguy, L., Baillon, M., and Soubeyroux, J.-M.: A 50-year high-
1078 resolution atmospheric reanalysis over France with the Safran system. *International Journal of*
1079 *Climatology*, 30(11), 1627–1644, <https://doi.org/10.1002/joc.2003>, 2010.

1080 Vidaller, I., Revuelto, J., Izagirre, E., Rojas-Heredia, F., Alonso-González, E., Gascoin, S., René
1081 P., Berthier, E., Rico, I., Moreno, A., Serrano, E., Serreta, A., and López-Moreno, J. I.: Toward
1082 an ice-free mountain range: Demise of Pyrenean glaciers during 2011–2020, *J. Geophys. Res.*
1083 *Lett.*, 48, e2021GL094339, <https://doi.org/10.1029/2021GL094339>, 2021.

1084 Viviroli, D., and Weingarter, R.: The hydrological significance of mountains: from regional to
1085 global scale, *Hydrology and Earth System Sciences*, 8, 1016–1029, [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-8-1017-2004)
1086 8-1017-2004, 2004.

1087 Vogel, J., Paton, E., Aich, V., and Bronstert, A.: Increasing Compound Warm Spells and Droughts
1088 in the Mediterranean Basin, *Weather Clim. Extrem.*, 32, 100312, [https://doi.](https://doi.org/10.1016/j.wace.2021.100312)
1089 [org/10.1016/j.wace.2021.100312](https://doi.org/10.1016/j.wace.2021.100312), 2021.

1090 Willibald, F., Kotlarski, S., Grêt-Regamey, A., and Ludwig, R.: Anthropogenic climate change
1091 versus internal climate variability: impacts on snow cover in the Swiss Alps, *The Cryosphere*, 14,
1092 2909–2924, [https://doi.org/10.5194/tc-14-](https://doi.org/10.5194/tc-14-2909-2020) 2909-2020, 2020.

1093 Woelber, B., Maneta, M. P., Harper, J., Jencso, K. G., Gardner, W. P., Wilcox, A. C., and López-
1094 Moreno, J.I.: The influence of diurnal snowmelt and transpiration on hillslope throughflow and
1095 stream response, *Hydrol. Earth Syst. Sci.*, 22, 4295–4310, [https://doi.org/10.5194/hess-22-4295-](https://doi.org/10.5194/hess-22-4295-2018)
1096 2018, 2018.

1097 Xercavins - Comas, A. Els climes del Pirineu Oriental: des de les terres gironines fins a la
1098 Catalunya Nord. Andorra, *Documents d'Anàlisi Geogràfica*, 7, 81-102, 1985.

1099 Zappa, G., Hoskins, B. J., and Shepherd, T. G.: The dependence of wintertime Mediterranean
1100 precipitation on the atmospheric circulation response to climate change, *Environ. Res. Lett.*, 10
1101 (10), 104012, <https://doi.org/10.1088/1748-9326/10/10/104012>, 2015.

1102

1103