

# **Snow sensitivity to temperature and precipitation change during compound cold-hot and wet-dry seasons in the Pyrenees.**

Josep Bonsoms<sup>1</sup>, Juan I. López-Moreno<sup>2</sup>, Esteban Alonso-González<sup>3</sup>

<sup>1</sup> Department of Geography, Universitat de Barcelona, Barcelona, Spain.

<sup>2</sup> Instituto Pirenaico de Ecología (IPE-CSIC), Campus de Aula Dei, Zaragoza, Spain.

<sup>3</sup> Centre d'Etudes Spatiales de la Biosphère (CESBIO), Université de Toulouse, CNES/CNRS/IRD/UPS, Toulouse, France.

Corresponding author: Juan I. López-Moreno (nlopez@ipe.csic.es)

1 **Abstract.** The Mediterranean basin has experienced one of the highest warming rates on earth  
2 during the last few decades, and climate projections predict water-scarcity in the future. Mid-  
3 latitude Mediterranean mountain areas, such as the Pyrenees, play a key role in the hydrological  
4 resources for the highly populated lowland areas. However, there are still large uncertainties about  
5 the impact of climate change on snowpack in the high mountain ranges of this region. Here, we  
6 perform a snow sensitivity to temperature and precipitation change analysis of the Pyrenean  
7 snowpack (1980 – 2019 period) using five key snow-climatological indicators. We analyzed snow  
8 sensitivity to temperature and precipitation during four different compounds weather conditions  
9 (cold-dry [CD], cold-wet [CW], warm-dry [WD], and warm-wet [WW]) at low elevations (1500  
10 m), mid-elevations (1800 m), and high elevations (2400 m) in the Pyrenees. In particular, we  
11 forced a physically based energy and mass balance snow model (FSM2), with validation by  
12 ground-truth data, and applied this model to the entire range, with forcing of perturbed reanalysis  
13 climate data for the period 1980 to 2019 as the baseline. The FSM2 model results successfully  
14 reproduced the observed snow depth (HS) values ( $R^2 > 0.8$ ), with relative root-mean square error  
15 and mean absolute error values less than 10% of the observed HS values. Overall, the snow  
16 sensitivity to temperature and precipitation change decreased with elevation and increased  
17 towards the eastern Pyrenees. When the temperature increased progressively at 1°C intervals, the  
18 largest seasonal HS decreases from the baseline were at +1°C. A 10% increase of precipitation  
19 counterbalanced the temperature increases ( $\leq 1^\circ\text{C}$ ) at high elevations during the coldest months,  
20 because temperature was far from the isothermal 0°C conditions. The maximal seasonal HS and  
21 peak HS max reductions were during WW seasons, and the minimal reductions were during CD  
22 seasons. During WW (CD) seasons, the seasonal HS decline per °C was 37% (28 %) at low  
23 elevations, 34% (30%) at mid elevations, and 27% (22 %) at high elevations. Further, the peak  
24 HS date was on average anticipated 2, 3 and 8 days at low, mid and high elevation, respectively.  
25 Results suggest snow sensitivity to temperature and precipitation change will be similar at other

26 mid-latitude mountain areas, where snowpack reductions will have major consequences on the  
27 nearby ecological and socioeconomic systems.

28

29 **Keywords:** Snow, Climate change, Sensitivity, Alpine, Mediterranean Mountains, Mid-latitude,  
30 Pyrenees.

31

## 32 **1 Introduction**

33

34 Snow is a key element of the Earth's climate system (Armstrong and Brun, 1998) because it cools  
35 the planet (Serreze and Barry, 2011) by altering the Surface Energy Balance (SEB), increasing  
36 the albedo, and modulating surface and air temperatures (Hall, 2004). Northern-Hemispheric  
37 snowpack patterns have changed rapidly during recent decades (Hammond et al., 2018; Hock et  
38 al., 2019; Notarnicola et al., 2020). It is crucial to improve our understanding of the timing of  
39 snow ablation and snow accumulation due to changing climate conditions because snowpack  
40 affects many nearby social and environmental systems. From the hydrological point of view, snow  
41 melt controls mountain runoff rate during the spring (Barnett et al., 2005; Adams et al., 2009;  
42 Stahl et al., 2010), river flow magnitude and timing (Morán-Tejeda et al., 2014; Sanmiguel-  
43 Vallelado et al., 2017), water infiltration and groundwater storage (Gribovszki et al., 2010; Evans  
44 et al., 2018), and transpiration rate (Cooper et al., 2020). The presence and duration of snowpack  
45 affects terrestrial ecosystem dynamics because snow ablation date affects photosynthesis  
46 (Woelber et al., 2018), forest productivity (Barnard et al., 2018), freezing and thawing of the soil  
47 (Luetschg et al., 2008; Oliva et al., 2014), and thickness of the active layer in permafrost  
48 environments (Hrbáček et al., 2016; Magnin et al., 2017). Snowpack also has remarkable  
49 economic impacts. For example, the snowpack at high elevations and surrounding areas  
50 determines the economic success of many mountain ski-resorts (Scott et al., 2003; Pons et al.,  
51 2015; Gilaberte-Búrdalo et al., 2017). Changes in the snowpack of mountainous regions also  
52 influence associated lowland areas because it affects the availability of snow meltwater that is  
53 used for water reservoirs, hydropower generation, agriculture, industries, and other applications  
54 (e.g., Sturm et al., 2017; Beniston et al., 2018).

55

56 Mid-latitude snowpacks have among the highest snow sensitivities worldwide (Brown and Mote,  
57 2009; López-Moreno et al., 2017; 2020b). In regions at high latitudes or high elevations,  
58 increasing precipitation can partly counterbalance the effect of increases of temperature on snow  
59 cover duration (Brown and Mote, 2009). Climate warming decreases the maximum and seasonal

60 snow depth (HS), the snow water equivalent (SWE) (Trujillo and Molotch, 2014; Alonso-  
61 González et al., 2020a; López-Moreno et al., 2013; 2017), and the fraction total precipitation as  
62 snowfall (snowfall ratio; e.g., Mote et al., 2005; Lynn et al., 2020; Jeenings and Molotoch, 2020;  
63 Marshall et al., 2019), and also delays the snow onset date (Beniston, 2009; Klein et al., 2016).  
64 However, warming can slow the early snow ablation rate on the season (Pomeroy et al., 2015;  
65 Rasouli et al., 2015; Jennings and Molotch, 2020; Bonsoms et al., 2022; Sanmiguel-Vallelado et  
66 al., 2022) because of the earlier HS and SWE peak dates (Alonso-González et al., 2022), which  
67 coincide with periods of low solar radiation (Pomeroy et al., 2015; Musselman et al., 2017a).

68

69 The Mediterranean basin is a region that is critically affected by climate change (Giorgi, 2006)  
70 being densely populated (>500 million inhabitants) and affected by an intense anthropogenic  
71 activity. Warming of the Mediterranean basin will accelerate for the next decades, and  
72 temperatures will continue to increase in this region during the warm months (Knutti and  
73 Sedlacek, 2013; Lionello and Scarascia 2018; Cramer et al., 2018; Evin et al., 2021; Cos et al.,  
74 2022), increasing atmospheric evaporative demands (Vicente-Serrano et al., 2020), drought  
75 severity (Tramblay et al., 2020), leading to water-scarcity over most of this region (García-Ruiz  
76 et al., 2011). Mediterranean mid-latitude mountains, such as the Pyrenees, where this research  
77 focuses, are the main runoff generation zones of the downstream areas (Viviroli and Weingartner,  
78 2004) and provide most of the water used by major cities in the lowlands (Morán-Tejeda et al.,  
79 2014).

80

81 Snow patterns in the Pyrenees have high spatial diversity (Alonso-González et al., 2019), due to  
82 internal climate variability of mid-latitude precipitation (Hawkins and Sutton 2010; Deser et al.,  
83 2012), high interannual and decadal variability of precipitation in the Iberian Peninsula (Esteban-  
84 Parra et al., 1998; Peña-Angulo et al., 2020) as well as the abrupt topography and the different  
85 mountain exposure to the Atlantic air masses (Bonsoms et al., 2021a). Thus, snow accumulation  
86 per season is almost twice as much in the northern slopes as in the southern slopes (Navarro-  
87 Serrano and López-Moreno, 2017), and there is a high interannual variability of snow in regions  
88 at lower elevations (Alonso-González et al., 2020a) and in the southern and eastern regions of the  
89 Pyrenees (Salvador-Franch et al., 2014; Salvador-Franch et al., 2016; Bonsoms et al., 2021b).  
90 Since the 1980s, the energy available for snow ablation has significantly increased in the Pyrenees  
91 (Bonsoms et al., 2022), and winter snow days and snow accumulation have non-statically  
92 significantly increased (Buisan et al., 2016; Serrano-Notivoli et al., 2018; López-Moreno et al.,  
93 2020a; Bonsoms et al., 2021a) due to the increasing frequency of positive west and south-west  
94 advections (Buisan et al., 2016). 21<sup>st</sup> century climate projections for Pyrenees anticipate a

95 temperature increase of more than 1°C to 4°C (relative to 1986–2005), and an increase (decrease)  
96 of precipitation by about 10% for the eastern (western) regions during winter and spring (Amblar-  
97 Frances et al., 2020). Therefore, changes in snow patterns in regions with high elevations are  
98 uncertain because winter snow accumulation is affected by precipitation (López-Moreno et al.,  
99 2008) and Mediterranean basin winter precipitation projections have uncertainties up to 80% of  
100 the total variance (Evin et al., 2021).

101

102 Previous studies in the central Pyrenees (López-Moreno et al., 2013), Iberian Peninsula Mountain  
103 ranges (Alonso-González et al., 2020a), and mountain areas that have Mediterranean climates  
104 (López-Moreno et al., 2017) demonstrated that snowpack sensitivity to changes in climate are  
105 mostly controlled by elevation. Despite the impact of climate warming in mountain hydrological  
106 processes, there is limited understanding of the snow sensitivity to temperature and precipitation  
107 changes and seasonality of mid-latitude Mediterranean mountain snowpacks. Some studies  
108 reported different snowpack sensitivities during wet and dry years (López-Moreno et al., 2017;  
109 Musselman et al., 2017b; Rasouli et al., 2022; Roche et al., 2018). However, the sensitivity of  
110 snow during periods when there are seasonal compound weather (temperature and precipitation)  
111 conditions has not yet been analyzed. The high interannual variability of the Pyrenean snowpack,  
112 which is expected to increase according to climate projections (López-Moreno et al., 2008),  
113 indicates a need to examine snowpack sensitivity to temperature and precipitation change  
114 focusing on the year-to-year variability. Warm seasons in the Mediterranean basin require special  
115 attention because these are likely to increase in the future (e.g., Vogel et al., 2019; De Luca et al.,  
116 2020; Meng et al., 2022). Further, the occurrence of different HS trends at mid- and high-elevation  
117 areas of this range (López-Moreno et al., 2020a) suggest that elevation and spatial factors  
118 contribute to the wide variations of the sensitivity of snow to the climate.

119

120 Therefore, the main objective of this research is to quantify snow (accumulation, ablation, and  
121 timing) sensitivity to temperature and precipitation change during compound temperature and  
122 precipitation seasons in the Pyrenees.

123


## 124 **2 Geographical area and climate setting**

125

126 The Pyrenees is a mountain range located in the north of the Iberian Peninsula (south Europe;  
127 42°N-43°N to 2°W-3°E) that is aligned east-to-west between the Atlantic Ocean and the  
128 Mediterranean Sea. The highest elevations are in the central region (Aneto, 3404 m) and

129 elevations decrease towards the west and east (Figure 1). The Mediterranean basin, including the  
130 Pyrenees, is in a transition area, and is influenced by the continental climate and the subtropical  
131 temperate climate. Precipitation is mostly driven by large-scale circulation patterns (Zappa et al.,  
132 2015; Borgli et al., 2019), the jet-stream oscillation during winter (Hurrell, 1995), and land-sea  
133 temperature differences (Tuel and Eltahir, 2020). During the summer, the northward movement  
134 of the Azores high pressure region brings stable weather, and precipitation is mainly convective  
135 at that time (Xercavins, 1985). Precipitation is highly variable depending on mountain exposure  
136 to the main circulation weather types; it ranges from about 1000 mm/year to about 2000 mm/year  
137 (in the mountain summits), with lower levels in the east and south (Cuadrat et al., 2007). There is  
138 a slight disconnection of the general climate circulation towards the eastern Pyrenees, where the  
139 Mediterranean climate and East Atlantic/West Russia (EA-WR) oscillations have greater effects  
140 on snow accumulation (Bonsoms et al., 2021a). In the southern, western, and central massifs of  
141 the range, the Atlantic climate and the negative North Atlantic Oscillation (NAO) phases regulate  
142 snow accumulation (W and SW wet air flows; López-Moreno, 2005; López-Moreno and Vicente-  
143 Serrano, 2007; Buisan et al., 2016; Alonso-González et al., 2020b). In the northern slopes, the  
144 positive phases of the Western Mediterranean Oscillation (WeMO) linked with NW and N  
145 advections trigger the most episodes of snow accumulation (Bonsoms et al., 2021a). The seasonal  
146 snow accumulation in the northern slopes is almost double the amount (about 500 cm more) as in  
147 the southern slopes at an elevation of about 2000 m (Bonsoms et al., 2021a). The  
148 temperature/elevation gradient is about 0.55°C/100 m (Navarro-Serrano and López-Moreno,  
149 2018) and the annual 0°C isotherm is at about 2750 to 2950 m (López-Moreno and García-Ruiz,  
150 2004). Net radiation and latent heat flux governs the energy available for snow ablation; the  
151 former heat flux increases at high elevations and the latter towards the east (Bonsoms et al., 2022).

152  
153



154

155 **Figure 1** (a) Study area. Pyrenean massifs grouped by sectors for (b) low, (c) mid and (d) high  
 156 elevation. The white dots indicate the locations of the automatic weather stations (AWS) shown  
 157 at Table 1. Massifs delimitation is based on the spatial regionalization of the SAFRAN system,  
 158 which groups massifs according to topographical and meteorological characteristics (modified  
 159 from Durand et al., 1999).

160

### 161 3 Data and methods

162

#### 163 3.1 Snow model

164

165 Snowpack was modelled using a physical-based snow model, the Flexible Snow Model (FSM2;  
 166 Essery, 2015). This model resolves the SEB and mass balance to simulate the state of the  
 167 snowpack. FSM2 is open access and available at <https://github.com/RichardEssery/FSM2> (last  
 168 access 16 December 2022). Previous studies tested the FSM2 (Krinner et al., 2018), and its  
 169 application in different forest environments (Mazzoti et al., 2021), and hydro-climatological  
 170 mountain zones such the Andes (Urrutia et al., 2019), Alps (Mazzoti et al., 2020), Colorado  
 171 (Smyth et al., 2022), Himalayas (Pritchard et al., 2020), Iberian Peninsula Mountains (Alonso-  
 172 González et al., 2020a; Alonso-González et al., 2022), Lebanese mountains (Alonso-González et  
 173 al., 2021), providing confidential results. The FSM2 requires forcing data of precipitation, air  
 174 temperature, relative humidity, surface atmospheric pressure, wind speed, incoming shortwave  
 175 radiation ( $SW_{inc}$ ), and incoming long wave radiation ( $LW_{inc}$ ). We have evaluated different FSM2  
 176 model configurations (not shown) without remarkable differences in the accuracy and

177 performance metrics. Thus, the FSM2 configuration included in this work estimates snow cover  
 178 fraction based on a linear function of HS and albedo based on a prognostic function, with increases  
 179 due to snowfall and decreases due to snow age. Atmospheric stability is calculated as function of  
 180 the Richardson number. Snow density is calculated as a function of viscous compaction by  
 181 overburden and thermal metamorphism. Snow hydrology is estimated by gravitational drainage,  
 182 including internal snowpack processes, runoff, refreeze rates, and thermal conductivity. Table S1  
 183 summarizes the FSM2 configuration and the FSM2 compile numbers.

184

### 185 **3.2 Snow model validation**

186 FSM2 configuration was validated by in situ snow records of four automatic weather stations  
 187 (AWSs) that were at high elevations in the Pyrenees. Precipitation in mountainous and windy  
 188 regions is usually affected by undercatch (Kochendorfer et al., 2020). Thus, the instrumental  
 189 records of precipitation were corrected for undercatch by applying an empirical equation validated  
 190 for the Pyrenees (Buisan et al., 2019). Precipitation type was classified by a threshold method  
 191 (Musselman et al., 2017b; Corripio et al., 2017): snow when the air temperature was below 1°C  
 192 and rain when the air temperature was above 1°C, according to previous research in the study area  
 193 (Corripio et al., 2017). The  $LW_{inc}$  heat flux of the AWSs (Table 1) were estimated as previously  
 194 described (Corripio et al., 2017). Due to the wide instrumental data coverage (99.3% of the total  
 195 dataset), gap-filling was not performed. The HS records were measured each 30 min using an  
 196 ultrasonic snow depth sensor. The meteorological data used in the validation process were  
 197 provided and managed by the local meteorological service of Catalonia  
 198 ([https://www.meteo.cat/wpweb/serveis/formularis/peticio-dinformes-i-dades-](https://www.meteo.cat/wpweb/serveis/formularis/peticio-dinformes-i-dades-meteorologiques/peticio-de-dades-meteorologiques/)  
 199 [meteorologiques/peticio-de-dades-meteorologiques/](https://www.meteo.cat/wpweb/serveis/formularis/peticio-dinformes-i-dades-meteorologiques/peticio-de-dades-meteorologiques/); data requested: 14/01/2021). Quality-  
 200 checking of the data was performed using an automatic error filtering process in combination with  
 201 a climatological, spatial, and internal coherency control defined by the SMC (2011).

202

203

**Table 1.** Characteristics of the four AWSs.

| Area                              | Code | Lat/Lon°   | Elevation (m) | Atlantic Ocean, Distance (km) | Mediterranean Sea, Distance (km) | Validation period (years) | Years |
|-----------------------------------|------|------------|---------------|-------------------------------|----------------------------------|---------------------------|-------|
| Central-Pyrenees, Northern slopes | A1   | 42.77/0.73 | 2228          | 200                           | 190                              | 2004–2020                 | 16    |
|                                   | A2   | 42.61/0.98 | 2266          | 225                           | 170                              | 2001–2020                 | 19    |
| Eastern Pyrenees, Southern slopes | A3   | 42.46/1.78 | 2230          | 295                           | 115                              | 2005–2020                 | 15    |

|  |    |            |      |     |     |           |    |
|--|----|------------|------|-----|-----|-----------|----|
| Eastern Pre-Pyrenees,<br>Northern slopes | A4 | 42.29/1.71 | 2143 | 300 | 110 | 2009–2020 | 11 |
|--|----|------------|------|-----|-----|-----------|----|

204

205 Model accuracy was estimated based on the mean absolute error (MAE) and the root mean square  
 206 error (RMSE), and model performance was estimated by the coefficient of determination ( $R^2$ ).  
 207 The MAE and the RMSE indicate the mean differences of the modelled and observed values.

208

### 209 **3.3 Atmospheric forcing data**

210

211 We forced the FSM2 with the open access climate reanalysis dataset provided by Vernay et al.  
 212 (2021), which consists of the modelled values from the SAFRAN meteorological analysis. The  
 213 FSM2 was run at an hourly resolution for each massif, each elevation range, and each climate  
 214 baseline and perturbation scenario from 1980 to 2019. The SAFRAN system provides data for  
 215 homogeneous meteorological and topographical mountain massifs every 300 m, from 0 to 3600  
 216 m (Durand et al., 1999; Vernay et al., 2021). We analyzed three elevation bands: low (1500 m),  
 217 middle (1800 m), and high (2400 m). Precipitation type was classified using the same threshold  
 218 approach used for model validation. Atmospheric emissivity was derived from the SAFRAN  
 219  $LW_{inc}$  and air temperature. SAFRAN was forced using numerical weather prediction models  
 220 (ERA-40 reanalysis data from 1958 to 2002 and ARPEGE from 2002 to 2020). Meteorological  
 221 data were calibrated, homogenized, and improved by in situ meteorological observations data  
 222 assimilation (Vernay et al., 2021). Durand et al. (1999; 2009a; 2009b) provided further technical  
 223 details of the SAFRAN system. Previous studies used the SAFRAN system for the long-term HS  
 224 trends (López-Moreno et al., 2020), extreme snowfall (Roux et al., 2021), and snow ablation  
 225 analysis (Bonsoms et al., 2022). SAFRAN system has been extensively validated for the  
 226 meteorological modelling of continental Spain (Quintana-Seguí et al., 2017), France (Vidal et al.,  
 227 2010) or alpine snowpack climate projections (Verfaille et al., 2018), among other works.

228

### 229 **3.4 Snow sensitivity to temperature and precipitation change analysis**

230

231 Snow sensitivity to temperature and precipitation change was analyzed using a delta-change  
 232 methodology (López-Moreno et al., 2008; Beniston et al., 2016; Musselman et al., 2017b; Marty  
 233 et al., 2017; Alonso-González et al., 2020a; Sanmiguel-Vallelado et al., 2022). In this method, air  
 234 temperature and precipitation were perturbed for each massif and elevation range based the  
 235 historical period (1980–2019). Air temperature was increased from 1 to 4°C at 1°C intervals,  
 236 assuming an increase of  $LW_{inc}$  accordingly (Jennings and Molotch, 2020). Precipitation was



237 changed from  $-10\%$  to  $+10\%$  at  $10\%$  intervals, in accordance with climate model uncertainties  
238 and the maximum and minimum precipitation projections for the Pyrenees (Amblar-Frances et  
239 al., 2020).

240

### 241 **3.5 Snow climate indicators**

242

243 Snow sensitivity to temperature and precipitation change was analyzed using five key indicators:  
244 (i) seasonal average HS, (ii) seasonal maximum absolute HS peak (peak HS max), (iii) date of the  
245 maximum HS (peak HS date), (iv) number of days with HS  $> 1$  cm on the ground (snow duration),  
246 and (v) daily average snow ablation per season (snow ablation, hereafter). Snow ablation was  
247 calculated as the difference between the maximum daily HS recorded on two consecutive days  
248 (Musselman et al., 2017a), and only days with decreases of 1 cm or more were recorded. Some  
249 seasons had more than one peak HS; for this reason, peak HS date was determined after applying  
250 a moving average of 5-days. All indicators were computed according to massif and elevation  
251 range.

252

### 253 **3.6 Definitions of compound temperature and precipitation seasons**

254

255 The snow season was from October 1 to June 30 (inclusive). Snow duration was defined by snow  
256 onset and snow ablation dates in situ observations (Bonsoms, 2021a), and results from the  
257 baseline scenario snow duration presented in this work. A “compound temperature and  
258 precipitation season” (season type) was assessed based on each massif and the elevation historical  
259 climate record (1980–2019) using a joint quantile approach (Beniston and Goyette, 2007;  
260 Beniston, 2009; López-Moreno et al., 2011a). Compound season types were defined according to  
261 López-Moreno et al. (2011a), based on the seasonal 40<sup>th</sup> percentiles (T40 for temperature and P40  
262 for precipitation) and the seasonal 60<sup>th</sup> percentiles (T60 and P60). There were four types of  
263 seasons based on seasonal temperature (T<sub>season</sub>) and seasonal precipitation (P<sub>season</sub>) data:

264 Cold and Dry (CD):  $T_{\text{season}} \leq T40$  and  $P_{\text{season}} \leq P40$ ;

265 Cold and Wet (CW):  $T_{\text{season}} \leq T40$  and  $P_{\text{season}} \geq P60$ ;

266 Warm and Dry (WD):  $T_{\text{season}} > T40$  and  $P_{\text{season}} \leq P40$ ;

267 Warm and Wet (WW):  $T_{\text{season}} > T60$  and  $P_{\text{season}} > P60$ .

268 All remaining seasons were classified as having average (Avg) temperature and precipitation.  
269 Note that the number of compound season type is different depending on the Pyrenees massif  
270 (Figure S1). However, by applying the joint-quantile approach described, we are comparing the  
271 snow sensitivity to temperature and precipitation change between similar climate conditions,  
272 independently where each compound season type was recorded.

273

### 274 **3.7 Spatial regionalization**

275

276 We have examined spatial differences in the snow sensitivity to temperature and precipitation  
277 change by compound season types. Following previous studies, massifs were grouped into four  
278 sectors by applying a Principal Component Analysis (PCA) (i.e., López-Moreno et al., 2020b;  
279 Matiu et al., 2020, among others). We applied a PCA over HS data for each month, year, massif,  
280 and elevation. Massifs were grouped into four sectors depending on the maximum correlation to  
281 PC1 and PC2 scores (see Figures S2). The number of season types per sector are shown at Figure  
282 S3 and the spatial regionalization is presented at Figure 1.

283

## 284 **4. Results**

285

286 We validated the FSM2 at Section 4.1. Subsequently, we analyzed the snow sensitivity to  
287 temperature and precipitation change based on five snow climate indicators, namely the seasonal  
288 HS, peak HS max, peak HS date, snow duration and snow ablation. Compound season types show  
289 similar relative importance on the snow sensitivity to temperature and precipitation change  
290 regardless of the Pyrenean sector. For this reason, our results have been focused on seasonal snow  
291 changes due to increments of temperature, elevation, and compound season type. These are the  
292 key factors that ruled the snow sensitivity to temperature and precipitation change, and an accurate  
293 analysis is provided at Section 4.2. Spatial differences on the snow sensitivity to temperature and  
294 precipitation change during compound season types are examined at Section 4.3.


295

### 296 **4.1 Snow model validation**

297

298 Our snow model validation analysis (Figures 2 and 3) confirmed that FSM2 accurately reproduces  
299 the observed HS values. On average, the FSM2 had a  $R^2$  greater than 0.83 for all stations. In  
300 general, the snow model slightly overestimated the maximum HS values. The highest  $R^2$  values  
301 were at A4 and A2 ( $R^2 = 0.85$  in both stations), and the lowest were at A3 and A1 ( $R^2 = 0.79$  and  
302  $R^2 = 0.82$ , respectively). The highest accuracy was at A4 (RMSE = 18.5 cm, MAE = 8.9 cm), and  
303 the largest errors were at A2 (RMSE = 45.8 cm, MAE = 29.0 cm).

304




305

306 **Figure 2.** Time series of the observed (gray) and modelled (black) HS values at the four AWSs.

307

308



309

310 **Figure 3.** Regression analysis of observed (x-axis) and simulated (y-axis) HS values.

311


312 **4.2 Snow sensitivity to temperature and precipitation change**

313

314 We then determined seasonal HS profiles for each perturbed climate scenario and compound  
315 season type (Figure 4). The results show a non-linear response between seasonal HS loss and  
316 temperature increase. When the temperature increased at 1°C intervals, the largest relative  
317 seasonal HS decrease from the baseline was at + 1°C for all elevations and all compound season  
318 type. High elevation areas had lower seasonal HS variability between compound season types  
319 than low elevations (Figure S4). At low elevations, snow was greater during CW seasons than  
320 other seasons. All the snowpack-perturbed scenarios indicated that snowpack decreased for all  
321 elevations under warming climate scenarios. Snowpack sensitivity to temperature and  
322 precipitation change depended on the compound season type (Figures 5 and 6). At low elevations,  
323 the seasonal changes in HS ranged from -37% (WW) to -28% (CD) per °C increase. For mid-  
324 elevation ranges, there were no remarkable differences among compound season types (Table 2),  
325 and the seasonal HS changes ranged from -34% (WW) to -30% (CW) per °C increase. Low and  
326 mid-elevations had greater snowpack reductions than high elevations. In the latter, a 10% increase  
327 of precipitation counterbalanced a temperature increase of about 1°C, and there were no  
328 remarkable differences in the seasonal HS from the baseline scenario especially in the coldest  
329 months of the season (Figure S5 and Figure S6). The maximum seasonal HS sensitivity to  
330 temperature and precipitation was during WD seasons (27%/°C), and the minimum was during  
331 CW seasons (-22%/°C).

332

333



334

335 **Figure 4.** Anomalies of seasonal HS for low, mid and high elevation (rows), compound season type  
 336 (columns), and different temperature increases (colors).

337

338 **Table 2.** Average and seasonal HS and peak HS sensitivity to temperature and precipitation  
 339 change during the four different compound temperature and precipitation seasons at three  
 different elevations.

| Season type | %HS/ °C |     |      | %peak HS max/°C |     |      |
|-------------|---------|-----|------|-----------------|-----|------|
|             | Low     | Mid | High | Low             | Mid | High |
| Avg.        | -33     | -33 | -25  | -20             | -20 | -16  |
| CD          | -28     | -30 | -22  | -17             | -17 | -14  |
| CW          | -33     | -32 | -22  | -22             | -20 | -15  |
| WD          | -32     | -30 | -27  | -19             | -16 | -16  |
| WW          | -37     | -34 | -26  | -24             | -24 | -16  |


340

341 At low and mid elevations, the peak HS max was greatest during WW seasons (-24%/°C) and  
 342 lowest during the CD and WD seasons (-17%/°C for both). At high elevations, there were no  
 343 clear differences in the peak HS max for the different seasons. The maximum peak HS max was  
 344 during WD seasons (-16%/°C) and the minimum was during CD seasons (-14%/°C).

345

346 We also determined average seasonal snow duration for each elevation range and compound  
 347 season type for different temperature increases (Table 3 and Figure 5c). The minimum snow  
 348 duration was during CW seasons (-13%/°C at low elevations, -10%/°C at mid-elevations, -5%/°C

349 at high elevations). At low elevations, the snow duration was most sensitive during WW seasons  
 350 ( $-17\%/^{\circ}\text{C}$ ). On the contrary, at mid-elevations and high elevations, the snow duration was most  
 351 sensitive during WD seasons ( $-13\%/^{\circ}\text{C}$  at mid-elevations and  $-8\%/^{\circ}\text{C}$  at high elevations).  
 352



353  
 354 **Figure 5.** Anomalies of seasonal HS (a), peak HS max (b) and snow duration (c) for different  
 355 temperature increases relative to baseline at three different elevations during the four different  
 356 compound season types. The solid black lines within each boxplot are the average. Lower and  
 357 upper hinges correspond to the 25th and 75th percentiles, respectively. The whisker is a horizontal  
 358 line at 1.5 interquartile range of the upper quartile and lower quartile, respectively. Dots represent  
 359 the outliers. Data is grouped by season, compound season type, increment of temperature,  
 360 precipitation variation, elevation, and massif.  
 361

362 The peak HS date occurred earlier due to warming, independently of precipitation changes.  
363 During WD seasons, the peak HS date per °C was anticipated by 3 days at low elevations, 3 days  
364 at mid-elevations, and 6 days at high elevations; during CD seasons, the peak HS date per °C was  
365 anticipated by 4 days at low elevations, 5 days at mid-elevations, and 9 days at high elevations.  
366 In low and mid elevation areas, if the temperature increase was no more than about 1°C above  
367 baseline, there was little change in the peak HS date (Figure 6). In addition, the minimum peak  
368 HS date change is found during WW seasons (Table 3), because the snowpack would be scarce  
369 at those times, and there were no defined peaks (Figure S4).

370

371 We determined the snow ablation sensitivity to temperature and precipitation change in response  
372 to different temperature increases at different elevations and during different compound season  
373 types. The results show there were low differences in absolute snow ablation values in a warmer  
374 climate (Figure 7). At low elevations, the average snow ablation sensitivity to temperature and  
375 precipitation change in all four compound seasons was 12%/°C (Table 3). At mid-elevations and  
376 high elevations, the maximum snow ablation sensitivity to temperature and precipitation change  
377 was during dry seasons; WD seasons had a snow ablation sensitivity to temperature and  
378 precipitation change of 13%/°C at mid-elevations and 10%/°C at high elevations. On the other  
379 hand, the minimum values for mid-elevations were during WW seasons, when the snow ablation  
380 sensitivity to temperature and precipitation change was 8%/°C; the minimum values at high  
381 elevations were during CW seasons, when was 5%/°C.

382

383



384

385 **Figure 6.** Difference (days) from baseline Peak HS date at three different elevations and during  
 386 the four different temperature (colors) and precipitation shifts (columns) for each season (boxes).

387


388 **Table 3.** Snow duration, snow ablation, and peak HS date sensitivity to temperature and  
 389 precipitation change during the four different compound season types.

| Season Type | Snow duration (%/°C) |     |      | Snow ablation (%/°C) |     |      | Peak HS date (days/°C) |     |      |
|-------------|----------------------|-----|------|----------------------|-----|------|------------------------|-----|------|
|             | Low                  | Mid | High | Low                  | Mid | High | Low                    | Mid | High |
| Avg.        | -15                  | -12 | -6   | 12                   | 11  | 7    | -1                     | -3  | -7   |
| CD          | -13                  | -11 | -5   | 12                   | 13  | 8    | -4                     | -5  | -9   |
| CW          | -13                  | -10 | -5   | 12                   | 10  | 5    | -2                     | -3  | -13  |
| WD          | -16                  | -13 | -8   | 12                   | 13  | 10   | -3                     | -3  | -6   |
| WW          | -17                  | -13 | -7   | 12                   | 8   | 7    | -1                     | 0   | -3   |

390

391





392

393 **Figure 7.** Absolute snow ablation values (cm/day) (y-axis) at three different elevations during  
 394 four different compound temperature and precipitation for baseline and different increments of  
 395 temperature (x-axis). seasons.


396

### 397 4.3 Spatial patterns

398

399 PCA analysis reveals four Pyrenean sectors, namely northern-western (NW), northern-eastern  
 400 (NE), southern-western (SW), and southern-eastern (SE). No remarkable differences between  
 401 sectors are found in the relative importance of each compound season type in the snow sensitivity  
 402 to temperature and precipitation change (Figure 8). Snow sensitivity to temperature and  
 403 precipitation change absolute values are generally lower at northern slopes (NW and NE) than at  
 404 the southern slopes (SW and SE) (Figure S7 and Figure S8). In detail, seasonal HS ranged from  
 405  $-26\%/^{\circ}\text{C}$  during CD (NW) to  $-36\%/^{\circ}\text{C}$  during WW (SE). Similarly, the maximum peak HS max  
 406 sensitivity to temperature and precipitation change was at SE during WW seasons ( $25\%/^{\circ}\text{C}$ ) and  
 407 the minimum was during CD seasons at NW ( $15\%/^{\circ}\text{C}$ ). The snow duration sensitivity to  
 408 temperature and precipitation change increased during WW seasons, and the maximum changes  
 409 were at SE ( $-16\%/^{\circ}\text{C}$ ); in contraposition, the lowest sensitivity to temperature and precipitation  
 410 change are found at NW, during CD and CW seasons ( $-8\%/^{\circ}\text{C}$ , in both seasons). Snow ablation  
 411 sensitivity to temperature and precipitation change increases towards the eastern Pyrenees,  
 412 particularly during WD seasons ( $14\%/^{\circ}\text{C}$  and  $13\%/^{\circ}\text{C}$  for NE and SE, respectively). Finally, no  
 413 remarkable peak HS date differences are observed between sectors and maximum values are

414 found during CD and CW seasons, when the peak HS date is anticipated  $\geq 5$  per  $^{\circ}\text{C}$  for all  
 415 sectors.  
 416



417  
 418 **Figure 8.** Average snow sensitivity to temperature and precipitation change (y-axis) grouped by  
 419 sector (x-axis), season type (color bars) and snow climate indicator (boxes).

420

## 421 5. Discussion

422

423 The spatial and temporal patterns of snow in the Pyrenees are highly variable, and climate  
 424 projections indicate that extreme events will likely increase during future decades (Meng et al.,  
 425 2022). Therefore, we analyzed factors that affect the snowpack sensitivity to temperature and  
 426 precipitation change gain insight into how future climate changes may affect the snow regime.

427

### 428 5.1 Snow sensitivity to temperature and precipitation change and relationship with 429 historical and future snow trends

430

#### 431 5.1.1 Snow accumulation phase

432

433 The snow losses due warming that we described here are mainly associated with increases in the  
434 rain/snowfall ratio (Figure S9), changes in the snow onset and offset dates (Figure S4), and  
435 increases in the energy available for snow ablation during the later months of the snow season, as  
436 it was previously reported by literature (e.g., Pomeroy et al., 2015; Lynn et al., 2020; Jennings  
437 and Molotch, 2020). At high elevations, a trend of increasing precipitation (+10%) could  
438 counterbalance temperature increases (<1°C; Figure S5), consistent with the results previously  
439 reported for specific sites of the central Pyrenees (Izas, 2000m; López-Moreno et al., 2008).  
440 Rasouli et al. (2014) also found that a 20% increase of precipitation could compensate for 2°C  
441 increase of temperature in subarctic Canada. A climate sensitivity analysis in the western  
442 Cascades (western USA) found that increases of precipitation due to warming modulated the  
443 snowpack accumulation losses by about 5%/1°C (Minder, 2010). These results are consistent with  
444 recent data that examined snow above 1000 m in the Pyrenees, which found that an increase in  
445 the frequency of west circulation weather types since the 1980s increased the HS (Serrano-  
446 Notivoli et al., 2018; López-Moreno et al., 2020), snow accumulation (Bonsoms et al., 2021a),  
447 and changes in winter snow days (Buisan et al., 2016). There are similar trends in the Alps, with  
448 an increase of extreme (exceeding the 100-year return level) snowfall above 3000 m during recent  
449 decades (Roux et al., 2021) and increases in extreme winter precipitation (Rajczak and Schär,  
450 2017).

451

### 452 **5.1.2 Snow ablation phase**

453

454 Climate warming leads to a cascade of physical changes in the SEB that increase snow ablation  
455 near the 0°C isotherm. On overall, the snow ablation showed low to inexistent changes due to  
456 warming. Comparison between low and high elevations indicated slightly faster snow ablation at  
457 high elevations (Figure 7). This higher rate of snow ablation per season at high elevations (which  
458 have deeper snowpacks) are probably because the snow there lasts until late spring, when more  
459 energy is available for snow ablation (Bonsoms et al., 2022). Temperature increase does not imply  
460 remarkable changes in snow ablation per season because warming decreases the magnitude of the  
461 snowpack (seasonal HS and peak HS max) and triggers an earlier onset of snowmelt (Wu et al.,  
462 2018). The earlier peak HS date (Table 3 and Figure 6) implies lower rates of net shortwave  
463 radiation, because snow melting starts earlier in warmer climates (Pomeroy et al., 2015),  
464 coinciding with the shorter days and lower solar zenith angle (Lundquist et al., 2013; Sanmiguel-  
465 Vallelado et al., 2022). Our results agree with the slow snow melt rates reported in the Northern  
466 Hemisphere from 1980 to 2017 (Wu et al., 2018). The results of previous studies were similar for  
467 subarctic Canada (Rasouli et al., 2014) and western USA snowpacks (Musselman et al., 2017b),  
468 but Arctic sites had faster melt rates (Krogh and Pomeroy, 2019).

469

### 470 **5.1.3 Snow sensitivity to temperature and precipitation change and snowpack projections**

471

472 Our results suggest that warming had a non-linear effect on snowpack reduction. Our largest snow  
473 losses were for seasonal HS when the temperature increased by 1°C above baseline. At low and  
474 mid elevations, the average seasonal HS decrease was more than 40% for all compound season  
475 types, and the maximum sensitivity was during WW seasons. Previous research in the Pyrenees  
476 and other mid-latitude mountain ranges reported similar results. A study in the central Pyrenees  
477 reported the peak SWE was 29%/°C, whereas snow season duration decreased by about 20 to 30  
478 days at about 2000 m (López-Moreno et al., 2013). The average peak HS max at high elevations  
479 in the Pyrenees (−16 %/°C; Figure 6 and Table 2), was similar to the average peak SWE sensitivity  
480 (−15%/°C) reported in the Iberian Peninsula mountains at 2500 m (Alonso-González et al.,  
481 2020a). These results are also consistent with climate projections for this mountain range. In  
482 particular, for a 2°C or more increase of temperature, the snow season declined by 38% at the  
483 lowest ski resorts (~1500 m) in the SE Pyrenees (Pons et al., 2015). However, high emission  
484 climate scenarios projected an increase in the frequency and intensity of high snowfall at high  
485 elevations (López-Moreno et al., 2011b). Snow sensitivity in the easternmost areas could decline  
486 during the winter because of a trend for an increase of about 10% in precipitation in this area  
487 (Amblar-Francés et al., 2020). Our projected changes in the Pyrenean snowpack dynamics are  
488 similar to the expected snow losses in other mountain ranges. For example, a study of the Atlas  
489 Mountains of northern Africa concluded that snowpack decreases were greater in the lowlands  
490 and projected seasonal SWE declines of 60% under the RCP4.5 scenario and 80% under the  
491 RCP8.5 scenario for the entire range (Tuel et al., 2022). A study in the Washington Cascades  
492 (western USA) found that snowpack decline was 19 to 23% per 1°C (Minder, 2010), similar to  
493 the values in the present study at high elevations. A study of the French Alps (Chartreuse, 1500  
494 m) found that seasonal HS decreases on the order of 25% for a 1.5°C increase and 32% for a 2°C  
495 increase of global temperature above the pre-industrial years (Verfaille et al., 2018). A study of  
496 the Swiss Alps reported a snowpack decrease of about 15%/°C (Beniston, 2003); in the same  
497 alpine country, another study predicted the seasonal HS will decrease by more than 70% in  
498 massifs below 1000 m in all future climate projections (Marty et al., 2017). The largest snow  
499 reductions will likely occur during the periods between seasons (Steger et al., 2013; Marty et al.,  
500 2017). Nevertheless, at high elevations, snow climate projections found no significant trend for  
501 maximum HS until the end of the 21st century above 2500 m in the eastern Alps (Willibald et al.,  
502 2021), suggesting that internal climate variability is a major source of uncertainty of SWE  
503 projections at high elevations (Schirmer et al., 2021).

504

### 505 **5.2 Influence of compound temperature and precipitation seasons**

506

507 We found that the maximum sensitivities of seasonal HS and peak HS max to temperature and  
508 precipitation change were during WW seasons at low and mid-elevations and during WD seasons  
509 at high elevations. Brown and Mote (2008) analyzed the sensitivity of snow to climate changes  
510 in the Northern Hemisphere and found maximal SWE sensitivities in mid-latitude maritime  
511 winter climate areas, and minimal SWE sensitivities to temperature and precipitation change in  
512 dry and continental zones, consistent with our results. López-Moreno et al. (2017) also found  
513 greater decreases of SWE in wet and temperate Mediterranean ranges than in drier regions.  
514 Furthermore, Rasouli et al. (2022) studied the northern North American Cordillera and found  
515 higher snowpack sensitivities to temperature and precipitation change in wet basins than dry  
516 basins. Our maximum snow ablation relative change over the baseline scenario occurred during  
517 WD seasons, in accordance with Musselman et al. (2017b), who found a higher snowmelt rate  
518 during dry years in the western USA. Low and mid-elevations are highly sensitive to WW seasons  
519 because wet conditions favor decreases in the seasonal HS due to advection from sensible heat  
520 fluxes. The temperature in the Pyrenees is still cold enough to allow snowfall at high elevations  
521 during WW seasons, and for this reason we found maximal sensitivities to temperature and  
522 precipitation change during WD seasons. Reductions of snowfall in alpine regions can be  
523 compensated in a warmer scenario, because warm and wet snow is less susceptible to blowing  
524 wind transport and losses from sublimation (Pomeroy and Li, 2000; Pomeroy et al., 2015). During  
525 spring, snow runoff could be also greater in wet climates due to rain-on-snow events (Corripio et  
526 al., 2017), coinciding with the availability of more energy for snow ablation.

527

### 528 **5.3 Spatial and elevation factors controlling snow sensitivity to temperature and** 529 **precipitation change**

530

531 Comparison between Pyrenean sectors (Figure 8) reveals no remarkable differences in the relative  
532 importance of each compound season type in the snow sensitivity to temperature and precipitation  
533 change. This is because by applying a joint-quantile approach for each massif and elevation, we  
534 are comparing similar climate seasons between sectors, regardless of the number of compound  
535 season types recorded in each massif during the baseline period (Figure S1 and S3). The highest  
536 absolute snow sensitivity to temperature and precipitation change values is found in the SE  
537 Pyrenees. This is consistent with the snow accumulation and ablation patterns previously reported  
538 in this region (Lopez-Moreno, 2005; Navarro-Serrano et al., 2018; Alonso-González et al., 2020a;  
539 Bonsoms et al., 2021a; Bonsoms et al., 2021b; Bonsoms et al., 2022). The Atlantic climate has  
540 less of an influence in the SE sector, and in situ observations indicated there was about half of the

541 seasonal snow accumulation amounts as in northern and western areas at the same elevation  
542 (>2000 m; Bonsoms et al., 2021a). The snow in the SE Pyrenees is more sensitive to temperature  
543 and precipitation change because these massifs are exposed to higher turbulence and radiative  
544 heat fluxes (Bonsoms et al., 2022), and 0°C isotherm is closer. Similar conclusions are found for  
545 low elevations, where the results show an upward displacement of the snow line due to warming.  
546 Previous studies described the sensitivity of the snow pattern to elevation at specific stations of  
547 the central Pyrenees (López-Moreno et al., 2013; 2017), Iberian Peninsula mountains (Alonso-  
548 González et al., 2020a), and other ranges such as the Cascades (Jefferson, 2011; Sproles et al.,  
549 2013), the Alps (Marty et al., 2017), and western USA (Pierce et al., 2013; Musselman et al.,  
550 2017b). In these regions, the models suggest larger snowpack reductions due to warming at  
551 subalpine sites than at alpine sites (Jennings and Molotch, 2020) due to closer isothermal  
552 conditions (Brown and Mote, 2009; Lopez-Moreno et al., 2017; Mote et al., 2018).

553

#### 554 **5.4 Environmental and socioeconomic implications**

555

556 Our results indicated there will be an increase of snow ablation days and imply a disappearance  
557 of the typical sequence of snow accumulation and snow ablation seasons. Climate warming  
558 triggers the simultaneous occurrence of several periods of snowfall and melting, snow droughts  
559 during the winter, and ephemeral snowpacks between seasons. These expected decreases in snow  
560 will likely have important impacts on the ecosystem. During spring, a snow cover cools the soil  
561 (Luetschg et al., 2008), delays the initiation of freezing (Oliva et al., 2014), functions as a thick  
562 active layer (Hrbáček et al., 2016), and protects alpine rocks from exposure to solar radiation and  
563 high air temperatures (Magnin et al., 2017). Due to warming temperatures, the remaining glaciers  
564 in the Pyrenees are shrinking and are expected to disappear before the 2050s (Vidaller et al.,  
565 2021). The shallower snowpack that we identified in this work will increase the vulnerability of  
566 glaciers, because snow has a higher albedo than dark ice and debris-covered glaciers and functions  
567 as a protective layer for glaciers (Fujita and Sakai, 2014).

568

569 The earlier onset of snowmelt suggested by our results, which is greater at low and mid-elevations  
570 during WD seasons, is in line with previous global studies that reported earlier streamflow due to  
571 earlier runoff dates (Adam et al., 2009; Stewart, 2009), and with a study of changes in the Iberian  
572 Peninsula River flows (Morán-Tejeda et al., 2014). Overall, our results are consistent with the  
573 slight decrease of the river peak flows that have occurred in the southern slopes of the Pyrenees  
574 since the 1980s (Sanmiguel-Vallelado et al., 2017). The reductions of seasonal HS that we  
575 identified, suggest that snowmelt-dominated stream flows will likely shift to rainfall dominated  
576 regimes. Although high elevation meltwater might increase and contribute to earlier groundwater

577 recharging (Evans et al., 2018), the increased evapotranspiration in the lowlands (Bonsoms et al.,  
578 2022) could counter this effect, so there is no net change in downstream areas (Stahl et al., 2010).  
579 Snow ephemerality triggers lower spring and summer flows (Barnett et al., 2005; Adam et al.,  
580 2009; Stahl et al., 2010) and has an impact on the hydrological management strategies. Winter  
581 snow accumulation affects hydrological availability during the months when water and  
582 hydroelectric demands are higher. This is because reservoirs store water during periods of peak  
583 flows (winter and spring), and release water during the driest season in the lowlands (summer)  
584 (Morán-Tejeda et al., 2014). Recurrent snow-scarce seasons may intensify these hydrological  
585 impacts and lead to competition for water resources among different ecological and  
586 socioeconomic systems. The economic viability of mountain ski-resorts in the Pyrenees depends  
587 on a regular deep snow cover (Gilaberte-Burdalo et al., 2014; Pons et al., 2015), but this is highly  
588 variable, especially at low and mid-elevations. The expected increase in snow-scarce seasons that  
589 we identified here is consistent with climate projections for this region, which suggest that no  
590 Pyrenean ski resorts will be viable under RCP 8.5 scenario by the end of the 21<sup>st</sup> century (Spandre  
591 et al., 2019).

592

### 593 **5.5 Limitations and uncertainties**

594 The meteorological input data that we used to model snow were estimated for flat slopes and the  
595 regionalization system we used was based on the SAFRAN system. According to this system, a  
596 mountain range is divided into massifs with homogeneous topography. The SAFRAN system has  
597 negative biases in shortwave radiation, a temperature precision of about 1 K, and biases in the  
598 accumulated monthly precipitation of about 20 kg/m<sup>2</sup> (Vernay et al., 2021). Our estimates of snow  
599 sensitivity to temperature and precipitation change were based on the delta-approach, which  
600 considers changes in temperature and precipitation based on climate projections for the Pyrenees  
601 (Amblar-Francés et al., 2020), but assumes that the meteorological patterns of the reference period  
602 will be constant over time. In this work we used a physical-based snow model since it provides  
603 better results for future snow climate change estimations than degree-day models (Carletti et al.,  
604 2022). The FSM2 is a physics-based model of intermediate complexity, and the estimates of snow  
605 densification are simpler than those from more complex models of snowpack. However, a more  
606 complex model does not necessarily provide better performance in terms of snowpack and runoff  
607 estimation (Magnusson et al., 2015). The FSM2 configuration implemented in this work includes  
608 snow meltwater retention, snowpack refreezing and snow albedo based on snow age, which are  
609 the physical parameters included in the best-performing snow models according to Essery et al.  
610 (2013). Snow model sensitivity studies reveal that intermediate complexity models exhibit similar

611 snow depth accuracies than most complex multi-layer snow models, as well as robust  
612 performances across seasons (Terzago et al., 2020).

613

## 614 **6 Conclusions**

615

616 Our study assessed the impact of temperature and precipitation change on the Pyrenean snowpack  
617 during compound cold-hot and wet-dry seasons, using a physical-based snow model that was  
618 forced by reanalysis data. We determined the snow sensitivity to temperature and precipitation  
619 change using five key indicators of snow accumulation and snow ablation. The lowest snow  
620 sensitivity to temperature and precipitation change was at high elevations of the NW Pyrenees  
621 and increased at lower elevations and in the SE slopes. An increase of 1°C at low and mid  
622 elevation regions led to remarkable decreases in the seasonal HS and snow duration. However, at  
623 high elevations, precipitation plays a key role, and temperature is far from the isothermal 0°C  
624 during the middle of winter. In this region, a 10% increase of precipitation, as suggested by the  
625 Spanish Meteorological Agency (AEMET) over the eastern regions of this range, could  
626 compensate for temperature increases on the order of about < 1°C. The impact of climate warming  
627 depends on the combination of temperature and precipitation during compound seasons. Our  
628 analysis of seasonal HS and peak HS max indicated the greatest declines were during WW seasons  
629 and the smallest declines were during CD seasons, independently of the Pyrenean sector. For  
630 snow duration, however, the highest (lowest) sensitivity to temperature and precipitation change  
631 is found during WD (CW) seasons. Similarly, snow ablation had slightly greater sensitivities to  
632 temperature and precipitation change during WD seasons, in that snow ablation variation is less  
633 than 10% and the peak HS date occurred about 5 days earlier per °C. Our findings thus provide  
634 evidence that the Pyrenean snowpack is highly sensitive to climate warming and suggest that the  
635 snowpacks of other mid-latitude mountain ranges may also show similar response to warming.

636

## 637 **Data availability**

638

639 Snow model (FSM2) is open access and provided by Essery (2015) and available at  
640 <https://github.com/RichardEssery/FSM2> (last access 16 December 2022). Climate forcing data is  
641 provided by Vernay et al. (2021), through AERIS ([https://www.aeris-data.fr/landing-  
642 page/?uuid=865730e8-edeb-4c6b-ae58-80f95166509b#v2020.2](https://www.aeris-data.fr/landing-page/?uuid=865730e8-edeb-4c6b-ae58-80f95166509b#v2020.2); last access 16 December 2022).  
643 Data of this work is available upon request (contact: josepbonsoms5@ub.edu).

644

## 645 **Acknowledgements**



646

647 This work falls within the research topics examined by the research group “Antarctic, Artic,  
648 Alpine Environments-ANTALP” (2017-SGR-1102) funded by the Government of Catalonia,  
649 HIDROIBERNIEVE (CGL2017-82216-R) and MARGISNOW (PID2021-124220OB-100), from  
650 the Spanish Ministry of Science, Innovation and Universities. JB is supported by a pre-doctoral  
651 University Professor FPI grant (PRE2021-097046) funded by the Spanish Ministry of Science,  
652 Innovation and Universities. The authors are grateful to Marc Oliva, who reviewed an early  
653 version of this manuscript. We acknowledge the SAFRAN data provided by Météo-France –  
654 CNRS and the CNRM Centre d'Etudes de la Neige, through AERIS.

655

#### 656 **Author contributions**

657

658 JB analyzed the data and wrote the original draft. JB, JILM and EAG contributed to the  
659 manuscript design and draft editing. JB, JILM and EAG read and approved the final manuscript.

660

#### 661 **References**

662

663 Adam, J. C., and Hamlet, A. F.: Implications of Global Climate Change for Snowmelt Hydrology  
664 in the Twenty First Century, *Hydrological Processes*, 23(7), 962-972,  
665 <https://doi.org/10.1002/hyp.7201>, 2009.

666 Alonso-González, E., Gutmann, E., Aalstad, K., Fayad, A., Bouchet, M., and Gascoin, S.:  
667 Snowpack dynamics in the Lebanese mountains from quasi-dynamically downscaled ERA5  
668 reanalysis updated by assimilating remotely sensed fractional snow-covered area, *Hydrol. Earth  
669 Syst. Sci.*, 25, 4455–4471, <https://doi.org/10.5194/hess-25-4455-2021>, 2021.

670 Alonso-González, E., López-Moreno, J.I., Navarro-Serrano, F., Sanmiguel-Valladolid, A.,  
671 Aznárez-Balta, M., Revuelto, J., and Ceballos, A.: Snowpack Sensitivity to Temperature,  
672 Precipitation, and Solar Radiation Variability over an Elevational Gradient in the Iberian  
673 Mountains, *Atmos. Res.*, 243, 104973 <https://doi.org/10.1016/j.atmosres.2020.104973>, 2020a.

674 Alonso-González, E., López-Moreno, J.I., Navarro-Serrano, F., Sanmiguel-Valladolid, A.,  
675 Revuelto, J., Domínguez-Castro, F., and Ceballos, A.: Snow climatology for the mountains in the  
676 Iberian Peninsula using satellite imagery and simulations with dynamically downscaled reanalysis  
677 data, *International Journal of Climatology*, 40(1), 477–491, <https://doi.org/10.1002/joc.6223>,  
678 2019.

679 Alonso-González, E., López-Moreno, J.I., Navarro-Serrano, F.M., and Revuelto, J.: Impact of  
680 North Atlantic oscillation on the snowpack in Iberian Peninsula mountains, *Water*, 12, 105–276,  
681 <https://doi.org/10.3390/w12010105>, 2020b.

682 Alonso-González, E., Revuelto, J., Fassnacht, S.R., and López-Moreno, J.I.: Combined influence  
683 of maximum accumulation and melt rates on the duration of the seasonal snowpack over  
684 temperate mountains, *Journal of Hydrology*, 608, 127574,  
685 <https://doi.org/10.1016/j.jhydrol.2022.127574>, 2022.

686 Amblar-Francés, M.P., Ramos-Calzado, P., Sanchis-Lladó, J., Hernanz-Lázaro, A., Peral- García,  
687 M.C., Navascués, B., Dominguez-Alonso, M., and Rodríguez-Camino, E.: High resolution  
688 climate change projections for the Pyrenees region, *Adv. Sci. Res.*, 17, 191–208,  
689 <https://doi.org/10.5194/asr-17-191-2020>, 2020.

690 Armstrong, A. and Brun, E.: *Snow and Climate, Physical Processes, Surface Energy Exchange*  
691 *and Modeling*, Cambridge University press, 222 pp., 1998.

692 Barnard, D. M., Knowles, J. F., Barnard, H. R., Goulden, M. L., Hu, J., Litvak, M. E., and  
693 Molotch, N. P.: Reevaluating growing season length controls on net ecosystem production in  
694 evergreen conifer forests, *Scientific Reports*, 8(1), 17973, [https://doi.org/10.1038/s41598-018-](https://doi.org/10.1038/s41598-018-36065-0)  
695 [36065-0](https://doi.org/10.1038/s41598-018-36065-0), 2018.

696 Barnett, T. P., Adam, J. C., and Lettenmaier, D. P.: Potential impacts of a warming climate on  
697 water availability in snow-dominated regions, *Nature*, 438(7066), 303–309,  
698 <https://doi.org/10.1038/nature04141>, 2005.

699

700 Beniston, M., and Stoffel, M.: Rain-on-snow events, floods and climate change in the Alps: events  
701 may increase with warming up to 4°C and decrease thereafter, *Sci. Total Environ.*, 571, 228–36,  
702 <https://doi.org/10.1016/j.scitotenv.2016.07.146>, 2016.

703 Beniston, M.: Trends in joint quantiles of temperature and precipitation in Europe since 1901 and  
704 projected for 2100, *Geophysical Research Letters*, 36, L07707,  
705 <https://doi.org/10.1029/2008GL037119>, 2009.

706 Beniston, M., Farinotti, D., Stoffel, M., Andreassen, L. M., Coppola, E., Eckert, N., Fantini, A.,  
707 Giacona, F., Hauck, C., Huss, M., Huwald, H., Lehning, M., López-Moreno, J.I., Magnusson, J.,  
708 Marty, C., Morán-Tejeda, E., Morin, S., Naaim, M., Provenzale, A., Rabatel, A., Six, D., Stötter,  
709 J., Strasser, U., Terzago, S., and Vincent, C.: The European mountain cryosphere: a review of its  
710 current state, trends, and future challenges, *The Cryosphere*, 12, 759–794,  
711 <https://doi.org/10.5194/tc-12-759-2018>, 2018.

712 Beniston, M., and Goyette, S.: Changes in variability and persistence of climate in Switzerland:  
713 exploring 20th century observations and 21st century simulations, *Global and Planetary Change*,  
714 57, 1–20, <https://doi.org/10.1016/j.gloplacha.2006.11.004>, 2007.

715 Beniston, M.; Keller, F.; Ko, B., and Goyette, S.: Estimates of snow accumulation and volume in  
716 the Swiss Alps under changing climatic conditions, *Theor. Appl. Climatol.*, 76, 125–140.  
717 <https://doi.org/10.1007/S00704-003-0016-5>, 2003.

718 Bonsoms, J., González, S., Prohom, M., Esteban, P., Salvador-Franch, F., López- Moreno, J.I.,  
719 and Oliva, M.: Spatio-temporal patterns of snow in the Catalan Pyrenees (SE Pyrenees, NE  
720 Iberia), *Int. J. Climatol.*, 41 (12), 5676–5697, <https://doi.org/10.1002/joc.7147>, 2021a.

721 Bonsoms, J., López-Moreno, J.I., González, S, and Oliva, M.: Increase of the energy available for  
722 snow ablation and its relation with atmospheric circulation, *Atmospheric Research*, 275, 106228,  
723 <https://doi.org/10.1016/j.atmosres.2022.106228>, 2022.

724 Bonsoms, J., Salvador-Franch, F., and Oliva, M.: Snowfall and snow cover evolution in the  
725 Eastern Pre-Pyrenees (NE Iberian Peninsula), *Cuad. Investig. Geogr.*, 47 (2), 291–307,  
726 <https://doi.org/10.18172/cig.4879>, 2021b.

727 Brown, R.D. and Mote, P.W.: The response of Northern Hemisphere snow cover to a changing  
728 climate, *Journal of Climate*, 22(8), 2124–2145, <https://doi.org/10.1175/2008JCLI2665.1>, 2009.

729 Buisan, S., Collado Aceituno, J. L. and Tierra, J.; ¿Se mide bien la precipitación en forma de  
730 nieve?, <https://doi.org/10.31978/639-19-010-0.095>, 2019.

731 Buisan, S.T., López-Moreno, J.I., Saz, M.A. and Kochendorfer, J.: Impact of weather type  
732 variability on winter precipitation, temperature and annual snowpack in the Spanish Pyrenees,  
733 *Climate Research*, 69(1), 79–92. <https://doi.org/10.3354/cr01391>, 2016.

734 Carletti, F., Michel, A., Casale, F., Bocchiola, D., Lehning, M., and Bavay, M.: A comparison of  
735 hydrological models with different level of complexity in Alpine regions in the context of climate  
736 change, *Hydrol. Earth Syst. Sci. Discuss.* <https://doi.org/10.5194/hess-26-3447-2022>, 2022.

737 Cooper, A. E., Kirchner, J. W., Wolf, S., Lombardozzi, D. L., Sullivan, B. W., Tyler, S. W., &  
738 Harpold, A. A.: Snowmelt causes different limitations on transpiration in a Sierra Nevada conifer  
739 forest, *Agricultural and Forest Meteorology*, 291, 108089.  
740 <https://doi.org/10.1016/j.agrformet.2020.108089>, 2020.

741 Corripio, J., and López-Moreno, J.I.: Analysis and predictability of the hydrological response of  
742 mountain catchments to heavy rain on snow events: a case study in the Spanish Pyrenees,  
743 *Hydrology*, 4(2), 20, <https://doi.org/10.3390/hydrology4020020>, 2017.

744 Cos, J., Doblas-Reyes, F., Jury, M., Marcos, R., Bretonnière, P.-A., and Samsó, M.: The  
745 Mediterranean climate change hotspot in the CMIP5 and CMIP6 projections, *Earth Syst. Dynam.*,  
746 13, 321–340, <https://doi.org/10.5194/esd-13-321-2022>, 2022.

747 Cramer W, Guiot J, Fader M, Garrabou J, Gattuso J-P, Iglesias A, Lange MA, Lionello P, Llasat  
748 MC, Paz S, Peñuelas J, Snoussi M, Toreti A, Tsimplis MN, and Xoplaki E.: Climate change and  
749 interconnected risks to sustainable development in the Mediterranean, *Nat. Clim. Chang.* 8(11),  
750 972–980, <https://doi.org/10.1038/s41558-018-0299-2>, 2018.

751 Cuadrat, J., Saz, M.A., Vicente-Serrano, S.: *Atlas climático de Aragón*. Gobierno de Aragón,  
752 Zaragoza, 222 pp. 2007.

753 De Luca, P., Messori, G., Faranda, D., Ward, P. J., and Coumou, D.: Compound warm-dry and  
754 cold-wet events over the Mediterranean, *Earth System Dynamics*, 11, 793–805,  
755 <https://doi.org/10.5194/esd-11-793-2020>, 2020.

756 Deser, C., Phillips, A., Bourdette, V., and Teng, H.: Uncertainty in climate change projections:  
757 the role of internal variability, *Climate dynamics*, 38(3), 527-546, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-010-0977-x)  
758 010-0977-x, 2012.

759 Durand, Y., Giraud, G., Brun, E., Mérindol, L., and Martin, E.: A computer-based system  
760 simulating snowpack structures as a tool for regional avalanche forecasting, *J. Glaciol.*, 45, 469–  
761 484, <https://doi.org/10.1017/S0022143000001337>, 1999.

762 Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Mérindol, L., and Lesaffre, B.: Reanalysis  
763 of 47 Years of Climate in the French Alps (1958–2005): Climatology and Trends for Snow Cover,  
764 *J. Appl. Meteorol. Clim.*, 48, 2487–2512, <https://doi.org/10.1175/2009JAMC1810.1>, 2009a.

765 Durand, Y., Giraud, G., Laternser, M., Etchevers, P., Mérindol, L., and Lesaffre, B.: Reanalysis  
766 of 44 Yr of Climate in the French Alps (1958–2002): Methodology, Model Validation,  
767 Climatology, and Trends for Air Temperature and Precipitation., *J. Appl. Meteorol. Clim.*, 48,  
768 429–449, <https://doi.org/10.1175/2008JAMC1808.1>, 2009b.

769 Essery, R.: A factorial snowpack model (FSM 1.0), *Geoscientific Model Development*, 8(12),  
770 3867–3876, <https://doi.org/10.5194/gmd-8-3867-2015>, 2015.

771 Essery, R., Morin, S., Lejeune, Y., and Ménard, C.: A comparison of 1701 snow models using  
772 observations from an alpine site, *Adv. Water Res.*, 55, 131–  
773 148, <https://doi.org/10.1016/j.advwatres.2012.07.013>, 2013.

774 Esteban-Parra, M.J, Rodrigo, F.S. and Castro-Diez, Y.: Spatial and temporal patterns of  
775 precipitation in Spain for the period 1880-1992, *Int. J. Climatol.*, 18, 1557–74, 1998.

776 Evans, S.G., Ge, S., Voss, C.I. and Molotch, N.P. The role of frozen soil in groundwater discharge  
777 predictions for warming alpine watersheds, *Water Resour. Res.*, 54, 1599–1615.  
778 <https://doi.org/10.1002/2017WR022098>, 2018.

779 Evin, G.; Somot, S.; Hingray, B. Balanced estimate and uncertainty assessment of European  
780 climate change using the large EURO-CORDEX regional climate model ensemble, *Earth Syst.*  
781 *Dyn. Discuss.*, 12(4), 1543–1569, <https://doi.org/10.5194/esd-12-1543-2021>, 2021.

782 Fujita, K. and Sakai, A.: Modelling runoff from a Himalayan debris-covered glacier, *Hydrol.*  
783 *Earth Syst. Sci.*, 18, 2679–2694, <https://doi.org/10.5194/hess-18-2679-2014>, 2014.

784 García-Ruiz, J. M., López-Moreno, J. I., Vicente-Serrano, S. M., Lasanta-Martínez, T. and  
785 Beguería, S. Mediterranean water resources in a global change scenario, *Earth Sci. Rev.*, 105(3–  
786 4), 121–139, <https://doi.org/10.1016/j.earscirev.2011.01.006>, 2011.

787 Gilaberte-Burdalo, M., Lopez-Martin, F., M. R. Pino-Otin, M., and Lopez-Moreno, J.: Impacts of  
788 climate change on ski industry, *Environ. Sci. Pol.*, 44, 51–  
789 61, <https://doi.org/10.1016/j.envsci.2014.07.003>, 2014.

790 Giorgi, F.: Climate change hot-spots, *Geophysical Research Letters*, 33: L08707,  
791 <https://doi.org/10.1029/2006GL025734>, 2006.

792 Gribovszki, Z., Szilágyi, J., and Kalicz, P.: Diurnal fluctuations in shallow groundwater levels  
793 and streamflow rates and their interpretation – A review, *J. Hydrol.*, 385, 371–  
794 383, <https://doi.org/10.1016/j.jhydrol.2010.02.001>, 2010.

795 Hall, A.: Role of surface albedo feedback in climate. *J. Clim.*, 17, 1550-1568, 2004.

796 Hammond, J. C., Saavedra, F. A. and Kampf, S. K.: Global snow zone maps and trends in snow  
797 persistence 2001–2016, *Int. J. Climatol.*, 38, 4369–4383, <https://doi.org/10.1002/joc.5674>, 2018.

798 Hawkins, E., and Sutton, R.: The potential to narrow uncertainty in projections of regional  
799 precipitation change, *Clim Dyn.*, <https://doi.org/10.1007/s00382-010-0810-6>, 2010.

800 Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., Jachson, M., K'a'ab, A.,  
801 Kang, S., Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., Steltzer, H., High mountain  
802 areas. In: Portner, H.-O., Roberts, D.C., Masson- Delmotte, V., et al. (Eds.), *IPCC Special Report*  
803 *on the Ocean and Cryosphere in a Changing Climate*. [https://www.ipcc.ch/srocc/chapter/chapter-](https://www.ipcc.ch/srocc/chapter/chapter-2/)  
804 [2/](https://www.ipcc.ch/srocc/chapter/chapter-2/), 2019.

805 Hrbáček, F., Láska, K., and Engel, Z.: Effect of snow cover on the active-layer thermal regime  
806 — a case study from James Ross Island, Antarctic Peninsula, *Permafrost and Periglac. Process.*,  
807 27, 307– 315, <https://doi.org/10.1002/ppp.1871>, 2016.

808 Hurrell, J. W.: Decadal trends in the North Atlantic oscillation: Regional temperatures and  
809 precipitation, *Science*, 269, 676–679, <https://doi.org/10.1126/science.269.5224.676>, 1995.

810 Jefferson, A. J.: Seasonal versus transient snow and the elevation dependence of climate  
811 sensitivity in maritime mountainous regions, *Geophys. Res. Lett.*, 38, L16402,  
812 <https://doi.org/10.1029/2011GL048346>, 2011.

813 Jennings, K.S., and Molotch, N.P.: Snowfall fraction, cold content, and energy balance changes  
814 drive differential response to simulated warming in an alpine and subalpine snowpack. *Front.*  
815 *Earth Sci*, 8, 2296-6463, <https://doi.org/10.3389/feart.2020.00186>, 2020.

816 Klein, G., Vitasse, Y., Rixen, C., Marty, C., and Rebetez, M.: Shorter snow cover duration since  
817 1970 in the Swiss Alps due to earlier snowmelt more than to later snow onset, *Clim. Chang.* 139,  
818 637–649. <https://doi.org/10.1007/s10584-016-1806-y>, 2016.

819 Knutti, R. and Sedlacek, J.: Robustness and uncertainties in the new CMIP5 climate model  
820 projections, *Nature Climate Change*, 3, 369–373, <https://doi.org/10.1038/nclimate1716>, 2013.

821 Kochendorfer, J., M.E. Earle, D. Hodyss, A. Reverdin, Y. Roulet, R. Nitu, R. Rasmussen, S.  
822 Landolt, S. Buisan, and Laine, T.: Undercatch adjustments for tipping bucket gauge  
823 measurements of solid precipitation, *J. Hydrometeorol.*, 21, 1193–1205,  
824 <https://doi.org/10.1175/JHM-D-19-0256.1>, 2020.

825 Krinner, G., Derksen, C., Essery, R., Flanner, M., Hagemann, S., Clark, M., Hall, A., Rott, H.,  
826 Brutel-Vuilmet, C., Kim, H., Ménard, C. B., Mudryk, L., Thackeray, C., Wang, L., Arduini, G.,  
827 Balsamo, G., Bartlett, P., Boike, J., Boone, A., Chéruy, F., Colin, J., Cuntz, M., Dai, Y.,  
828 Decharme, B., Derry, J., Ducharne, A., Dutra, E., Fang, X., Fierz, C., Ghattas, J., Gusev, Y.,  
829 Haverd, V., Kontu, A., Lafaysse, M., Law, R., Lawrence, D., Li, W., Marke, T., Marks, D.,  
830 Ménégoz, M., Nasonova, O., Nitta, T., Niwano, M., Pomeroy, J., Raleigh, M. S., Schaedler, G.,  
831 Semenov, V., Smirnova, T. G., Stacke, T., Strasser, U., Svenson, S., Turkov, D., Wang, T.,  
832 Wever, N., Yuan, H., Zhou, W., and Zhu, D.: ESM-SnowMIP: assessing snow models and  
833 quantifying snow-related climate feedbacks, *Geosci. Model Dev.*, 11, 5027–  
834 5049, <https://doi.org/10.5194/gmd-11-5027-2018>, 2018.

835 Krogh, S.A., and Pomeroy, J.W.: Impact of Future Climate and Vegetation on the Hydrology of  
836 an Arctic Headwater Basin at the Tundra–Taiga Transition, *J. Hydrometeorol.*, 20, 197–215.  
837 <https://doi.org/10.1175/JHM-D-18-0187.1>, 2019.

838 Lionello, P. and Scarascia, L.: The relation between climate change in the Mediterranean region  
839 and global warming, *Reg. Environ. Change*, 18, 1481–1493, <https://doi.org/10.1007/s10113-018->  
840 1290-1, 2018.

841 López Moreno, J.I., and Garcia Ruiz, J.M.: Influence of snow accumulation and snowmelt on  
842 streamflow in the Central Spanish Pyrenees, *International. J. Hydrol. Sci.*, 49, 787–802,  
843 <https://doi.org/10.1623/hysj.49.5.787.55135>, 2004.

844 López-Moreno, J.I.: Recent variations of snowpack depth in the central Spanish Pyrenees, *Arct.*  
845 *Antarct. Alp. Res.*, 37, 253–260, <https://doi.org/10.1657/1523-0430> (2005)037, 2005.

846 López-Moreno, J.I., Gascoin, S., Herrero, J., Sproles, E.A., Pons, M., Alonso-González, E.,  
847 Hanich, L., Boudhar, A., Musselman, K.N., Molotch, N.P., Sickman, J., and Pomeroy, J.:  
848 Different sensitivities of snowpacks to warming in Mediterranean climate mountain areas,  
849 *Environ. Res. Lett.*, 12 (7), 074006, <https://doi.org/10.1088/1748-9326/aa70cb>, 2017.

850 Lopez-Moreno, J.I., Goyette, S., Beniston, M., and Alvera, B.: Sensitivity of the snow energy  
851 balance to climate change: Implications for the evolution of snowpack in Pyrenees in the 21st  
852 century, *Climate Research* 36(3), 203–217, <https://doi.org/10.3354/cr00747>, 2008.

853 López-Moreno, J.I., Goyette, S., Vicente-Serrano, S.M., and Beniston, M.: Effects of climate  
854 change on the intensity and frequency of heavy snowfall events in the Pyrenees, *Clim. Chang.*,  
855 105, 489–508. <https://doi.org/10.1007/s10584-010-9889-3>, 2011b.

856 López-Moreno, J.I., Pomeroy, J.W., Alonso-González, E., Morán-Tejeda, E., and Revuelto, J.:  
857 Decoupling of warming mountain snowpacks from hydrological regimes, *Environ. Res. Lett.*, 15,  
858 11–15, <https://doi.org/10.1088/1748-9326/abb55f>, 2020a.

859 López-Moreno, J.I., Pomeroy, J.W., Revuelto, J., and Vicente-Serrano, S.M.: Response of snow  
860 processes to climate change: spatial variability in a small basin in the Spanish Pyrenees, *Hydrol.*  
861 *Process.*, 27, 2637–2650. <https://doi.org/10.1002/hyp.9408>, 2013.

862 López-Moreno, J.I., and Vicente-Serrano, S.M.: Atmospheric circulation influence on the  
863 interannual variability of snowpack in the Spanish Pyrenees during the second half of the  
864 twentieth century, *Nord. Hydrol.*, 38 (1), 38–44, <https://doi.org/10.2166/nh.2007.030>, 2007.

865 López-Moreno, J.I., Vicente-Serrano S.M., Morán-Tejeda E., Lorenzo J., Kenawy, A. and  
866 Beniston, M.: NAO effects on combined temperature and precipitation winter modes in the  
867 Mediterranean mountains: Observed relationships and projections for the 21st century, *Global*  
868 *and Planetary Change*, 77, 72-66, <https://doi.org/10.1016/j.gloplacha.2011.03.003>, 2011a.

869 López-Moreno, J.I., Soubeyroux, J.M., Gascoin, S., Alonso-González, E., Durán- Gómez, N.,  
870 Lafaysse, M., Vernay, M., Carmagnola, C., and Morin, S.: Long-term trends (1958–2017) in snow  
871 cover duration and depth in the Pyrenees, *Int. J. Climatol.*, 40, 6122–6136,  
872 <https://doi.org/10.1002/joc.6571>, 2020b.

873 López-Moreno J.I, Revuelto, J, Gilaberte, M., Morán-Tejeda, E., Pons, M., Jover, E., Esteban, P.,  
874 García, C., and Pomeroy, J.W.: The effect of slope aspect on the response of snowpack to climate

875 warming in the Pyrenees, *Theoretical and Applied Climatology*, 117, 1–13,  
876 <https://doi.org/10.1007/s00704-013-0991-0>, 2013.

877 López-Moreno, J, Pomeroy, J.W, Revuelto, J., Vicente-Serrano, S.M. Response of snow  
878 processes to climate change: spatial variability in a small basin in the Spanish Pyrenees,  
879 *Hydrological Processes*, 27(18), 2637–2650, <https://doi.org/10.1002/hyp.9408>, 2013

880 López-Moreno, J.; Boike, J.; Sanchez-Lorenzo, A. and Pomeroy, J.: Impact of climate warming  
881 on snow processes in Ny-Ålesund, a polar maritime site at Svalbard, *Glob. Planet. Chang.*, 146,  
882 10–21, <https://doi.org/10.1016/j.gloplacha.2016.09.006>, 2016.

883 Luetschg, M., Lehning, M., and Haeberli, W.: A sensitivity study of factors influencing warm/thin  
884 permafrost in the Swiss Alps, *J. Glaciol.*, 54, 696–704.  
885 <https://doi.org/10.3189/002214308786570881>, 2008

886 Lundquist, J.D., Dickerson-Lange, S.E., Lutz, J.A., and Cristea, N.C.: Lower forest density  
887 enhances snow retention in regions with warmer winters: a global framework developed from  
888 plot-scale observations and modeling, *Water Resour. Res.*, 49, 6356–6370.  
889 <https://doi.org/10.1002/wrcr.20504>, 2013.

890 Lynn, E., Cuthbertson, A., He, M., Vasquez, J. P., Anderson, M. L., Coombe, P., et al. Technical  
891 note: Precipitation-phase partitioning at landscape scales to regional scales, *Hydrology and Earth  
892 System Sciences*, 24(11), 5317–5328, <https://doi.org/10.5194/hess-24-5317-2020>, 2020.

893 Magnin, F., Westermann, S., Pogliotti, P., et al.: Snow control on active layer thickness in steep  
894 alpine rock walls (Aiguille du Midi, 3842 ma.s.l., Mont Blanc massif), *Catena*, 149, 648–662,  
895 <https://doi.org/10.1016/j.catena.2016.06.006>, 2017.

896 Marshall, A. M., Link, T. E., Abatzoglou, J. T., Flerchinger, G. N., Marks, D. G., and Tedrow,  
897 L.: Warming alters hydrologic heterogeneity: Simulated climate sensitivity of hydrology-based  
898 microrefugia in the snow-to-rain transition zone, *Water Resources Research*, 55, 2122–2141,  
899 <https://doi.org/10.1029/2018WR023063>, 2019.

900 Marty, C., Schlögl, S., Bavay, M., and Lehning, M.: How much can we save? Impact of different  
901 emission scenarios on future snow cover in the Alps, *The Cryosphere*, 11, 517–529,  
902 <https://doi.org/10.5194/tc-11-517-2017>, 2017.

903 Mazzotti, G., Essery, R., Moeser, D., and Jonas, T.: Resolving small-scale forest snow patterns  
904 with an energy balance snow model and a 1-layer canopy, *Water Resources Research*, 56,  
905 e2019WR026129, <https://doi.org/10.1029/2019WR026129>, 2020.

906 Mazzotti, G., Webster, C., Essery, R., and Jonas, T.: Increasing the physical representation of  
907 forest-snow processes in coarse-resolution models: Lessons learned from upscaling hyper-



908 resolution simulations, *Water Resources Research*, 57(5), e2020WR029064, <https://doi.org/10.1029/2020WR029064>, 2021.

909

910 Meng, Y., Hao, Z., Feng, S., Zhang, X., Hao, F.: Increase in compound dry-warm and wet-warm  
911 events under global warming in CMIP6 models, *Global and Planetary Change*, 210, 103773,  
912 <https://doi.org/10.1016/j.gloplacha.2022.103773>, 2022.

913 Minder, J. R.: The Sensitivity of Mountain Snowpack Accumulation to Climate Warming, *Journal*  
914 *of Climate*, 23(10), 2634-2650, <https://doi.org/10.1175/2009JCLI3263.1>, 2010.

915 Morán-Tejeda, E., Lorenzo-Lacruz, J., López-Moreno, J.I., Rahman, K. and Beniston, M.:  
916 Streamflow timing of mountain rivers in Spain: Recent changes and future projections, *J. Hydrol.*  
917 517, 1114–1127, <https://doi.org/10.1016/j.jhydrol.2014.06.053>, 2014.

918 Mote, P. W., Li, S., Lettenmaier, D. P., Xiao, M., and Engel, R.: Dramatic declines in snowpack  
919 in the western US, *npj Clim. Atmos. Sci.*, 1, 2, <https://doi.org/10.1038/s41612-018-0012-1>, 2018.

920 Mote, P.W., A.F. Hamlet, M.P. Clark, and D.P. Lettenmaier.: Declining mountain snowpack in  
921 western North America, *Bull. Am. Meteorol. Soc.*, 86, 39–49, [https://doi.org/10.1175/BAMS-86-](https://doi.org/10.1175/BAMS-86-1-39)  
922 1-39, 2005.

923 Musselman, K., Clark, M., Liu, C., Ikeda, K., and Rasmussen, R.: Slower snowmelt in a warmer  
924 world, *Nat. Clim. Change*, 7, 214–219, <https://doi.org/10.1038/NCLIMATE3225>, 2017a.

925 Musselman, K. N., Molotch, N. P., and Margulis, S. A.: Snowmelt response to simulated warming  
926 across a large elevation gradient, southern Sierra Nevada, California, *Cryosphere*, 11, 2847–2866,  
927 <https://doi.org/10.5194/tc-11-2847-2017>, 2017b.

928 Navarro-Serrano, F. and López-Moreno, J.I.: Spatio-temporal analysis of snowfall events in the  
929 Spanish Pyrenees and their relationship to atmospheric circulation, *Cuad. Invest. Geogr.*, 43 (1),  
930 233–254, <https://doi.org/10.18172/cig.3042>, 2017.

931 Notarnicola, C.: Hotspots of snow cover changes in global mountain regions over 2000–2018,  
932 *Remote Sensing of Environment*, 243, 111781, <https://doi.org/10.1016/j.rse.2020.111781>, 2020.

933 Magnusson, J., Wever, N., Essery, R., Helbig, N., Winstral, A., and Jonas, T.: Evaluating snow  
934 models with varying process representations for hydrological applications, *Water Resour. Res.*,  
935 51, 2707–2723, <https://doi.org/10.1002/2014WR016498>, 2015.

936 Matiu, M., Crespi, A., Bertoldi, G., Carmagnola, C.M., Marty, C., Morin, S., Schöner, W., Cat  
937 Berro, D., Chiogna, G., De Gregorio, L., Kotlarski, S., Majone, B., Resch, G., Terzago, S., Valt,  
938 M., Beozzo, W., Cianfarra, P., Gouttevin, I., Marcolini, G., Notarnicola, C., Petitta, M., Scherrer,  
939 S.C., Strasser, U., Winkler, M., Zebisch, M., Cicogna, A., Cremonini, R., Debernardi, A., Faletto,  
940 M., Gaddo, M., Giovannini, L., Mercalli, L., Soubeyroux, J.-M., Susnik, A., Trenti, A., Urbani,  
941 S., Weilguni, V., 2020. Observed snow depth trends in the European Alps 1971 to 2019. *Cryoph*  
942 1-50. <https://doi.org/10.5194/tc-2020>, 2020.

943 Oliva, M., Gómez Ortiz, A., Salvador, F., Salvà, M. Pereira, P., and Gerales, M.: Long-term soil  
944 temperature dynamics in the Sierra Nevada, Spain. *Geoderma* 235-236, 170-181,  
945 <https://doi.org/10.1016/j.geoderma.2014.07.012>, 2014.

946 Peña-Angulo, D., Vicente-Serrano, S., Domínguez-Castro, F., Murphy, C., Reig, F., Trambly,  
947 Y., Trigo, R., Luna, M.Y., Turco, M., Noguera, I., Aznárez-Balta, M., Garcia-Herrera, R., Tomas-  
948 Burguera, M. and Kenawy, A.: Long-term precipitation in Southwestern Europe reveals no clear  
949 trend attributable to anthropogenic forcing, *Environmental Research Letters*, 15 (9), 094070,  
950 <https://doi.org/10.1088/1748-9326/ab9c4f>, 2020.

951 Pierce, D. and Cayan, D.: The uneven response of different snow measures to human-induced  
952 climate warming, *Journal of Climate*, 26, 4148–4167, [https://doi.org/10.1175/JCLI-D-12-](https://doi.org/10.1175/JCLI-D-12-00534.1)  
953 00534.1, 2013.

954 Pomeroy, J. W., and L. Li.: Prairie and arctic areal snow cover mass balance using a blowing  
955 snowmodel, *J.Geophys.Res.*, 105( D21), 26619– 26634, <https://doi.org/10.1029/2000JD900149>,  
956 2000.

957 Pomeroy, J. W., Fang, X and Rasouli, K.: Sensitivity of snow processes to warming in the  
958 Canadian Rockies. 72nd Eastern Snow Conf., Sherbrooke, QC, Canada, Eastern Snow  
959 Conference, 22–33, 2015.

960 Pons, M., López-Moreno, J., Rosas-Casals, M., and Jover, E.: The vulnerability of Pyrenean ski  
961 resorts to climate-induced changes in the snowpack, *Climatic Change*, 131, 591–  
962 605, <https://doi.org/10.1007/s10584-015-1400-8>, 2015.

963 Pritchard, D. M. W., Forsythe, N., O'Donnell, G., Fowler, H. J., and Rutter, N.: Multi-physics  
964 ensemble snow modelling in the western Himalaya, *The Cryosphere*, 14(4), 1225–1244,  
965 <https://doi.org/10.5194/tc-14-1225-2020>, 2020.

966 Quintana-Seguí, P., Turco, M., Herrera, S., and Miguez-Macho, G.: Validation of a new  
967 SAFRAN-based gridded precipitation product for Spain and comparisons to Spain02 and ERA-  
968 Interim, *Hydrol. Earth Sys. Sci.*, 21, 2187–2201, <https://doi.org/10.5194/hess-21-2187-2017>,  
969 2017.

970 Rajczak, J. and Schär, C.: Projections of Future Precipitation Extremes Over Europe: A  
971 Multimodel Assessment of Climate Simulations, *J. Geophys. Res.-Atmos.*, 122, 773–  
972 710, <https://doi.org/10.1002/2017JD027176>, 2017.

973 Rasouli, K., J. W. Pomeroy, J. R. Janowicz, S. K. Carey, and T. J. Williams.: Hydrological  
974 sensitivity of a northern mountain basin to climate change, *Hydrol. Processes*, 28, 4191–5208,  
975 <https://doi.org/10.1002/hyp.10244>, 2014.

976 Rasouli, K.R., Pomeroy, J.W., and Marks, D.G.: Snowpack sensitivity to perturbed climate in a  
977 cool mid-latitude mountain catchment, *Hydrol. Process.*, 29, 3925–3940.  
978 <https://doi.org/10.1002/hyp.10587>, 2015.

979 Rasouli, K.R., Pomeroy, J.W., and Whietfiled, P.H.: The sensitivity of snow hydrology to changes  
980 in air temperature and precipitation in three North American headwater basins, *J. Hydrol.*, 606,  
981 127460, <https://doi.org/10.1016/j.jhydrol.2022.127460>, 2022

982 Roche, J.W., Bales, R.C., Rice, R., Marks, D.G.: Management Implications of Snowpack  
983 Sensitivity to Temperature and Atmospheric Moisture Changes in Yosemite National Park. *J. Am.*  
984 *Water Resour. Assoc.*, 54 (3), 724–741, <https://doi.org/10.1111/1752-1688.12647>, 2018.

985 Roux, E., Evin, G., Eckert, N., Blanchet, J., and Morin, S.: Elevation-dependent trends in extreme  
986 snowfall in the French Alps from 1959 to 2019, *The Cryosphere*, 15, 4335–4356,  
987 <https://doi.org/10.5194/tc-15-4335-2021>, 2021.

988 Salvador - Franch, F., Salvà, G., Vilar, F., and García, C.: Contribución al análisis nivométrico  
989 del Pirineo Oriental: La Molina, período 1956 - 1996. En: X Congreso Internacional AEC: Clima,  
990 sociedad, riesgos y ordenación del territorio, pp. 365-375, Alicante.  
991 <http://hdl.handle.net/10045/58002>, 2016.

992 Salvador Franch, F., Salvà, G., Vilar, F., and García, C.: Nivometría y perfiles de innivación en  
993 Núria (1970 m, Pirineo Oriental): 1985 - 2013. En: IX Congreso de la AEC, pp. 729 -738,  
994 Almería, <http://hdl.handle.net/20.500.11765/8229>, 2014.

995 Sanmiguel-Vallelado, A., Morán-Tejeda, E., Alonso-González, E., and López-Moreno, J. I.:  
996 Effect of snow on mountain river regimes: An example from the Pyrenees, *Frontiers of Earth*  
997 *Science*, 11(3), 515–530. <https://doi.org/10.1007/s11707-016-0630-z>, 2017.

998 Sanmiguel-Vallelado, A., McPhee, J., Esmeralda, P., Morán-Tejeda, E., Camarero, J., López-  
999 Moreno, J.I. Sensitivity of forest-snow interactions to climate forcing: Local variability in a  
1000 Pyrenean valley, *Journal of Hydrol.*, <https://doi.org/10.1016/j.jhydrol.2021.127311>, 2022.

1001 Schirmer, M., Winstral, A., Jonas, T., Burlando, P., and Peleg, N.: Natural climate variability is  
1002 an important aspect of future projections of snow water resources and rain-on-snow events, *The*  
1003 *Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2021-276>, 2021.

1004 Scott, D, McBoyle G, and Mills B.: Climate change and the skiing industry in southern Ontario  
1005 (Canada): exploring the importance of snowmaking as a technical adaptation, *Clim Res.*, 23(2),  
1006 171–181, <https://doi.org/10.3354/CR023171>, 2003.

1007 Serrano-Notivoli, R., Buisan, S.T., Abad-Pérez, L.M., Sierra-Alvarez, E., Rodríguez-Ballesteros,  
1008 C., López-Moreno, J.I. and Cuadrat, J.M.: Tendencias recientes en precipitación, temperatura y  
1009 nieve de alta montaña en los Pirineos (Refugio de Góriz, Huesca). In: *El clima: aire, agua, tierra*  
1010 *y fuego*. Madrid, Spain: Asociación Española de Climatología y Ministerio para la Transición

1011 Ecológica – Agencia Estatal de Climatología y Ministerio para la Transición Ecológica – Agencia  
1012 Estatal de Meteorología, pp. 267, 1060–280, 2018.

1013 Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research  
1014 synthesis, *Glob. Planet. Change*, 77, 85–96, <https://doi.org/10.1016/j.gloplacha.2011.03.004>,  
1015 2011.

1016 Servei Meteorològic de Catalunya (SMC). Les estacions meteorològiques automàtiques (EMA).  
1017 [https://static-m.meteo.cat/wordpressweb/wp-](https://static-m.meteo.cat/wordpressweb/wp-content/uploads/2014/11/18120559/Les_Estacions_XEMA.pdf)  
1018 [content/uploads/2014/11/18120559/Les\\_Estacions\\_XEMA.pdf](https://static-m.meteo.cat/wordpressweb/wp-content/uploads/2014/11/18120559/Les_Estacions_XEMA.pdf)(accessed March 1, 2022). 2011.

1019 Smyth, E. J., Raleigh, M. S., and Small, E. E.: The challenges of simulating SWE beneath forest  
1020 canopies are reduced by data assimilation of snow depth, *Water Resources Research*, 58,  
1021 e2021WR030563, <https://doi.org/10.1029/2021WR030563>, 2022.

1022 Spandre, P., François, H., Verfaillie, D., Pons, M., Vernay, M., Lafaysse, M., George, E., and  
1023 Morin, S. Winter tourism under climate change in the Pyrenees and the French Alps: relevance  
1024 of snowmaking as a technical adaptation, *The Cryosphere*, 13, 1325–1347,  
1025 <https://doi.org/10.5194/tc-13-1325-2019>, 2019.

1026 Sproles, E.A, Nolin, A.W, Rittger, K, and Painter, T. H.: Climate change impacts on maritime  
1027 mountain snowpack in the Oregon Cascades, *Hydrology and Earth System Sciences*, 17(7), 2581–  
1028 2597, <https://doi.org/10.5194/hess-17-2581-2013>, 2013.

1029 Stahl, K., Hisdal, H., Hannaford, J., Tallaksen, L.M., van Lanen, H.A.J., Sauquet, E., Demuth, S.,  
1030 Fendekova, M., and Jódar, J.: Streamflow trends in Europe: evidence from a dataset of near-  
1031 natural catchments, *Hydrol. Earth. Syst. Sci.*, 14, 2367-2382, [https://doi.org/10.5194/hess-14-](https://doi.org/10.5194/hess-14-2367-2010)  
1032 [2367-2010](https://doi.org/10.5194/hess-14-2367-2010), 2010.

1033 Steger, C., Kotlarski, S., Jonas, T., and Schär, C.: Alpine snow cover in a changing climate: A  
1034 regional climate model perspective, *Clim. Dynam.*, 41, 735–754, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-012-1545-3)  
1035 [012-1545-3](https://doi.org/10.1007/s00382-012-1545-3), 2013.

1036 Stewart I.T.: Changes in snowpack and snowmelt runoff for key mountain regions. *Hydrological*  
1037 *Processes*, 23, 78–94, <https://doi.org/10.1002/hyp.7128>, 2009.

1038 Sturm, M., M. A. Goldstein, and C. Parr. Water and life from snow: A trillion dollar science  
1039 question, *Water Resour. Res.*, 53, 3534– 3544, <https://doi.org/10.1002/2017WR020840>, 2017.

1040 Terzago, S., Andreoli, V., Arduini, G., Balsamo, G., Campo, L., Cassardo, C., Cremonese, E.,  
1041 Dolia, D., Gabellani, S., Hardenberg, J., Morra di Cella, U., Palazzi, E., Piazzzi, G., Pogliotti, P.,  
1042 and Provenzale, A.: Sensitivity of snow models to the accuracy of meteorological forcings in  
1043 mountain environments, *Hydrol. Earth Syst. Sci.*, 24, 4061–4090, [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-24-4061-2020)  
1044 [24-4061-2020](https://doi.org/10.5194/hess-24-4061-2020), 2020.

1045 Tramblay, Y.; Koutroulis, A.; Samaniego, L.; Vicente-Serrano, S.M.; Volaire, F.; Boone, A.; Le  
1046 Page, M.; Llasat, M.C.; Albergel, C.; Burak, S.; et al.: Challenges for drought assessment in the  
1047 Mediterranean region under future climate scenarios, *Earth Sci. Rev.*, 210, 103348,  
1048 <https://doi.org/10.1016/j.earscirev.2020.103348>, 2020.

1049 Trujillo, E., and N. P. Molotch.: Snowpack regimes of the Western United States, *Water Resour.*  
1050 *Res.*, 50(7), 5611–5623, <https://doi.org/10.1002/2013WR014753>, 2014.

1051 Tuel, A. and Eltahir, E. A. B.: Why Is the Mediterranean a Climate Change Hot Spot?, *J. Climate*,  
1052 33, 5829–5843. <https://doi.org/10.1175/jcli-d-19-0910.1>, 2020.

1053 Tuel, A., El Moçayd, N., Hasnaoui, M. D., and Eltahir, E. A. B.: Future projections of High Atlas  
1054 snowpack and runoff under climate change, *Hydrol. Earth Syst. Sci.*, 26, 571–588,  
1055 <https://doi.org/10.5194/hess-26-571-2022>, 2022.

1056 Urrutia, J., Herrera, C., Custodio, E., Jódar, J., and Medina, A.: Groundwater recharge and  
1057 hydrodynamics of complex volcanic aquifers with a shallow saline lake: Laguna Tuyajto, Andean  
1058 Cordillera of northern Chile, *Sci. Total Environ.*, 697, 134116,  
1059 <https://doi.org/10.1016/j.scitotenv.2019.134116>, 2019.

1060 Verfaillie, D., Lafaysse, M., Déqué, M., Eckert, N., Lejeune, Y., and Morin, S.: Multi-component  
1061 ensembles of future meteorological and natural snow conditions for 1500 m altitude in the  
1062 Chartreuse mountain range, Northern French Alps, *The Cryosphere*, 12, 1249–  
1063 1271, <https://doi.org/10.5194/tc-12-1249-2018>, 2018.

1064 Vernay, M., Lafaysse, M., Monteiro, D., Hagenmuller, P., Nheili, R., Samacoïts, R., Verfaillie,  
1065 D., and Morin, S.: The S2M meteorological and snow cover reanalysis over the French  
1066 mountainous areas, description and evaluation (1958–2020), *Earth Syst. Sci. Data Discuss*,  
1067 <https://doi.org/10.5194/essd-2021-249>, 2021.

1068 Vicente-Serrano, S.M., McVicar, T., Miralles, D., Yang, Y. and Tomas-Burguera, M.:  
1069 Unravelling the Influence of Atmospheric Evaporative Demand on Drought under Climate  
1070 Dynamics, *Climate Change in press*, 11(2), 1757-7780, <https://doi.org/10.1002/wcc.632>, 2020.

1071 Vidal, J.-P., Martin, E., Franchistéguy, L., Baillon, M., and Soubeyroux, J.-M.: A 50-year high-  
1072 resolution atmospheric reanalysis over France with the Safran system. *International Journal of*  
1073 *Climatology*, 30(11), 1627–1644, <https://doi.org/10.1002/joc.2003>, 2010.

1074 Vidaller, I., Revuelto, J., Izagirre, E., Rojas-Heredia, F., Alonso-González, E., Gascoin, S., René  
1075 P., Berthier, E., Rico, I., Moreno, A., Serrano, E., Serreta, A., and López-Moreno, J. I.: Toward  
1076 an ice-free mountain range: Demise of Pyrenean glaciers during 2011–2020, *J. Geophys. Res.*  
1077 *Letts.*, 48, e2021GL094339, <https://doi.org/10.1029/2021GL094339>, 2021.

1078 Viviroli, D., and Weingarter, R.: The hydrological significance of mountains: from regional to  
1079 global scale, *Hydrology and Earth System Sciences*, 8, 1016–1029, [https://doi.org/10.5194/hess-](https://doi.org/10.5194/hess-8-1017-2004)  
1080 8-1017-2004, 2004.

1081 Vogel, J., Paton, E., Aich, V., and Bronstert, A.: Increasing Compound Warm Spells and Droughts  
1082 in the Mediterranean Basin, *Weather Clim. Extrem.*, 32, 100312, [https://doi.](https://doi.org/10.1016/j.wace.2021.100312)  
1083 [org/10.1016/j.wace.2021.100312](https://doi.org/10.1016/j.wace.2021.100312), 2021.

1084 Willibald, F., Kotlarski, S., Grêt-Regamey, A., and Ludwig, R.: Anthropogenic climate change  
1085 versus internal climate variability: impacts on snow cover in the Swiss Alps, *The Cryosphere*, 14,  
1086 2909–2924, [https://doi.org/10.5194/tc-14-](https://doi.org/10.5194/tc-14-2909-2020) 2909-2020, 2020.

1087 Woelber, B., Maneta, M. P., Harper, J., Jencso, K. G., Gardner, W. P., Wilcox, A. C., and López-  
1088 Moreno, J.I.: The influence of diurnal snowmelt and transpiration on hillslope throughflow and  
1089 stream response, *Hydrol. Earth Syst. Sci.*, 22, 4295–4310, [https://doi.org/10.5194/hess-22-4295-](https://doi.org/10.5194/hess-22-4295-2018)  
1090 2018, 2018.

1091 Xercavins - Comas, A. Els climes del Pirineu Oriental: des de les terres gironines fins a la  
1092 Catalunya Nord. Andorra, Documents d'Anàlisi Geogràfica, 7, 81-102, 1985.

1093 Zappa, G., Hoskins, B. J., and Shepherd, T. G.: The dependence of wintertime Mediterranean  
1094 precipitation on the atmospheric circulation response to climate change, *Environ. Res. Lett.*, 10  
1095 (10), 104012, <https://doi.org/10.1088/1748-9326/10/10/104012>, 2015.

1096

1097