

# Rain process models and convergence to point processes

Scott Hottovy<sup>1</sup> and Samuel N. Stechmann<sup>2</sup>

<sup>1</sup>Department of Mathematics, United States Naval Academy, Annapolis, Maryland, USA

<sup>2</sup>Department of Mathematics & Department of Atmospheric and Oceanic Sciences, University of Wisconsin-Madison, Madison, Wisconsin, USA

**Correspondence:** Scott Hottovy (hottovy@usna.edu)


**Abstract.** A variety of stochastic models have been used to describe time series of precipitation or rainfall. Since many of these stochastic models are simplistic, it is desirable to develop connections between the stochastic models and the underlying physics of rain. Here, convergence results are presented for such a connection between two stochastic models: (i) a stochastic moisture process as a physics-based description of atmospheric moisture evolution, and (ii) a point process for rainfall time series as spike trains. The moisture process has dynamics that switch after the moisture hits a threshold, which represents the onset of rainfall and thereby gives rise to an associated rainfall process. This rainfall process is characterized by its random holding times for dry and wet periods. On average, the holding times for the wet periods are much shorter than the dry, and, in the limit of short wet periods, the rainfall process converges to a point process that is a spike train. Also, in the limit, the underlying moisture process becomes a threshold model with a teleporting boundary condition. To establish these limits and connections, formal asymptotic convergence is shown using the Fokker-Planck equation, which provides some intuitive understanding. Also, rigorous convergence is proved in mean-square with respect to continuous functions, of the moisture process, and convergence in mean-square with respect to generalized functions, of the rain process.

## 1 Introduction

Time series of precipitation or rainfall display highly irregular behavior, as illustrated in Fig. 1, and many valuable models have been based on stochastic processes. A variety of different stochastic models have been used, including renewal processes, Markov chains, Poisson processes, and point processes (Green, 1964; Katz, 1977; Richardson, 1981; Smith and Karr, 1983; Foufoula-Georgiou and Lettenmaier, 1987; Rodriguez-Iturbe et al., 1988; Cowpertwait et al., 1996; Wilks and Wilby, 1999). The many applications of these models include weather forecasting, stochastic weather generation, climate impact assessment, climate model downscaling, hydrological modeling, ecological modeling, and agricultural modeling.

Commonly, stochastic models for rainfall are empirical—i.e., based mainly on fitting the model behavior to match observational rainfall data—rather than based mainly on the underlying physical laws. Nevertheless, it is desirable to relate the stochastic models to physical principles, to the extent possible. Here, we investigate such a relation.

In particular, the goal of the present paper is to prove a connection between (i) a point-process description of rainfall time series and (ii) a physics-based model for the stochastic evolution of moisture. At first glance, the point-process model appears to be somewhat disconnected from basic physical laws based on mass, momentum, and energy. However, the point-process



**Figure 1.** Sample precipitation time series from observations at (a) Manus Island and (b) Nauru Island reproduced from Fig 3. of Abbott et al. (2016) with permission from the authors. The latter two panels are stochastic model simulations of (c) the rain rate process  $\sigma^\epsilon(t)$  with finite rain rate  $r$  and (d)  $\sigma(t)$  as the point process.

model can be seen to arise from the underlying evolution of moisture (which is the mixing ratio of water vapor in the air) (Abbott et al., 2016). Here, this connection is demonstrated via formal asymptotics on the Fokker–Planck equation, and proved rigorously in the mean-square sense.

To be more specific, a point process model of rainfall can be viewed as a spike train, as in Fig. 1d, where a rainfall event is an instantaneous spike. The point process could be defined and characterized by the random waiting time,  $\tau^d$ , of the duration of the “dry spell” in between rain events. As an empirical model of rainfall, one could estimate the probability density function (pdf) of  $\tau^d$  based on observational data (Peters et al., 2010; Deluca and Corral, 2014). For such an empirical approach, one could use data of rainfall time series alone, without appealing to any physical laws or any other type of observational data (humidity, wind speed, etc.). Similarly, beyond point processes, one could use a renewal process as a model of rainfall time

35 series, as in Fig. 1c, by introducing a finite (and possibly random) time  $\tau^r$  for the duration of the rain event. Again, as in the case of a point process, one could use a renewal process as an empirical model, based on data of rainfall time series alone, without appealing to any physical laws or any other type of observational data. However, it would be desirable to show that the point process and renewal process models can also arise from more physically based underpinnings.

Here, as mentioned above, a point process model of precipitation will be linked to the evolution of moisture, to provide a  
 40 more physically based foundation of the point process model. The moisture model used here is a continuous-time stochastic process for  $q(t)$ , which represents the amount of water vapor in a column of the atmosphere, as an anomaly from a baseline level, at time  $t$  (Stechmann and Neelin, 2011, 2014; Hottovy and Stechmann, 2015b; Abbott et al., 2016; Neelin et al., 2017). For example, the anomaly  $q(t) = 0$  corresponds to 62 mm of moisture in the column and  $q(t) = b = 3$  mm will be an upper threshold of 65 mm of moisture in the column. The  $q(t)$  process is governed by the stochastic differential equations (SDEs)

$$45 \quad dq(t) = \begin{cases} m dt + D_0 dW_t & \text{for } \sigma(t) = 0 \\ -r dt + D_1 dW_t & \text{for } \sigma(t) = r \end{cases}, \quad q(0) = 0, \quad \sigma(0) = 0, \quad (1)$$

where  $m$  and  $r$  are the moistening and rain rates respectively, and  $D_0$  and  $D_1$  are the constant diffusion coefficients which capture the fluctuations of moisture during the respective states. The quantity  $\sigma(t)$  is an indicator function for rain, and the dynamics of  $\sigma(t)$  switch from 0 to  $r$  when  $q(t)$  reaches a fixed threshold  $b > 0$ . For instance, supposing that  $(q(0), \sigma(0)) = (0, 0)$ , then  $\sigma(t) = 0$  until the time  $t_1 = \inf\{t \geq 0 : q(t) = b\}$ , at which time the value of  $\sigma$  switches to  $\sigma(t) = r$ . Then  $\sigma(t)$   
 50 switches back to zero at a later time when the moisture has been depleted to the lower threshold ( $q(t) = 0$ ). Figure 2a,b shows a realization of the processes  $q(t)$  and  $\sigma(t)$ . The process  $\sigma(t)$  can be viewed as a renewal process, with random durations  $\tau^d$  and  $\tau^r$  of dry spells and rain events, respectively, although  $\sigma(t)$  is not just a stand-alone renewal process, since it arises from the underlying dynamics of moisture  $q$ . The moisture in the column can potentially reach negative values, i.e.  $q(t) < -62$ , on very rare occasions. This effect is due to the idealized nature of the model.

55 The threshold behavior of (1) is a fundamental feature of the moisture–rainfall relationship that is seen in nature (Peters and Neelin, 2006; Deluca et al., 2015), and it is a basic aspect of many more complex moisture models and convective parameterizations as well (Lin and Neelin, 2000; Frierson et al., 2004; Khouider and Majda, 2005; Khouider et al., 2010; Hottovy and Stechmann, 2015a; Stechmann and Hottovy, 2016; Ahmed and Neelin, 2019; Mueller and Stechmann, 2020; Huang et al., 2022). Sometimes the threshold is also called a trigger (Hernandez-Duenas et al., 2019). The threshold can be viewed as the  
 60 threshold for the release of moist convective instability, and the moisture  $q$  is used as the physical quantity that governs the onset of moist convection. In this way, (1) is a physically based model of atmospheric moisture, and, from it, one can obtain a rainfall time series as a secondary or auxiliary quantity.

The main purpose of the paper is to define and show convergence of the threshold model in (1) as  $r \rightarrow \infty$  to a point process model of rainfall. For example, on the level of renewal processes,  $\tau^r \rightarrow 0$  and thus  $\sigma(t)$  converges to a process that is zero  
 65 everywhere and has spikes at infinity after random durations of length  $\tau^d$ . That is,  $\sigma(t)$  converges to a Dirac delta process. However,  $\sigma(t)$  is right continuous and has left hand limits, whereas the spike train is not. Thus the mode of convergence is not clear. For  $q(t)$  the limit is also unclear, but will be redefined in a way to show convergence with respect to the topology on

continuous functions with the uniform metric. In this study, the limiting processes are defined (in Section 2) and convergence is shown both heuristically (for the Fokker-Planck equation) and rigorously.

70 Some of the novel aspects of this work are as follows. The limit jump process  $q(t)$  has an associated Fokker-Planck equation that is derived using a matched asymptotic method. The resulting Fokker-Planck equation has a peculiar boundary flux condition which defines a “teleporting” boundary condition of  $q(t)$ . The processes are decoupled into evaporating and precipitating processes. Only after this decoupling can convergence of the evaporation processes be shown rigorously with respect to the uniform metric on the space of continuous functions. Also, the rain process  $\sigma(t)$  is shown to converge rigorously with respect  
75 to the generalized function space. This proof shows convergence of a renewal process to a delta process. Furthermore, the proof shows what kinds of bounds are needed for the rain event times  $\tau^r$  in order for integrated convergence to hold.

The convergence results shown here have the potential to impact various other fields. Many fields of study use similar renewal processes to model different types of phenomena (Cox, 1962). The connections to rain models were made above. In addition, there has been work in queuing theory to approximate point processes with renewal processes (e.g. Whitt (1982);  
80 Bhat (1994)), and using threshold triggers in financial models (Lejay and Pigato, 2019). Thresholds arise in many applications of piecewise dynamical systems where the threshold marks a change in the dynamics, as in Fillipov dynamics and hybrid switching diffusions (Filippov, 2013; Simpson and Kuske, 2014). The limiting process is similar to a stochastic resetting process studied in Evans and Majumdar (2011); Evans et al. (2020). Here the process stochastically resets to  $q = 0$  after a random hitting time  $\tau^d$  which depends on the process. Another interesting connection is with neuron stochastic integrate and  
85 fire models (see Sacerdote and Giraud (2013) for a review). The moisture process with a finite rain rate is similar to a Wiener Process model of a single Neuron with refractoriness. A similar model was studied in Albano et al. (2008) where the refractory time was constant. Here, the refractory time is random and coincides with the rain duration time  $\tau^r$ . Thus the work here is applicable to understanding the differences in using a model without refractoriness versus a model with a short, possible random, refractory time.

90 The structure of the paper is as follows. The processes for moisture and rain are defined in Section 2. The modes of convergence are discussed in Section 3. The heuristic convergence with the Fokker-Planck equation is shown in Section 3.1. Rigorous convergence of the moistening process  $E^\epsilon$  to  $E$  is shown with respect to  $L^2$  in Section 3.2 and the rain process  $\sigma^\epsilon$  is shown to converge to the sum of delta distributions  $\sigma$  with respect to generalized functions in Section 3.3. Some important statistics as well as an analysis of the differences when using the two processes are shown in Section 4. The results are summarized in  
95 Section 5. Technical details of the proofs and derivations are given in the Appendix.

## 2 Model Description

In this section the moisture and precipitation processes are defined. First the underlying moisture process of the renewal rain process is defined. The processes are defined with a small parameter  $\epsilon$  with the limit as  $\epsilon \rightarrow 0$  in mind.

The moisture process  $q^\epsilon(t) \in \mathbb{R}$  is defined as the solution to the stochastic differential equation (SDE),

$$100 \quad dq^\epsilon(t) = \begin{cases} m dt + D_0 dW_t & \text{for } \sigma^\epsilon(t) = 0 \\ -\frac{r}{\epsilon} dt + D_1 dW_t & \text{for } \sigma^\epsilon(t) = \frac{r}{\epsilon} \end{cases}, \quad q^\epsilon(0) = 0, \quad \sigma^\epsilon(0) = 0, \quad (2)$$

where  $m$  and  $r/\epsilon$  are the moistening and rain rates, and  $0 < D_0 \leq D_1$  are the fluctuations of moisture during the respective states. Here the small parameter  $\epsilon$  is essentially the ratio of moistening and rain rates (times a constant  $O(1)$  factor). In other words,  $\epsilon$  is the value which makes  $m/r$  order 1. In the tropics the rain rate is often seen to have values in excess of 15 mm h<sup>-1</sup> (e.g. see Figure 1 panels a) and b)). See Figure 7a) in Holloway and Neelin (2010) for an estimate of a typical  
105 moistening rate in the tropics. There the moistening rate is roughly 0.4 mm h<sup>-1</sup>. The rain process,  $\sigma^\epsilon(t) \in \{0, r/\epsilon\}$  is defined as follows: since  $\sigma^\epsilon(0) = 0$ , let  $\mathcal{T}_1^\epsilon \equiv \inf\{t > 0 | q^\epsilon(t) = b\}$ . Then  $\sigma^\epsilon(t) = 0$  for  $t \in [0, \mathcal{T}_1^\epsilon)$ . Next let  $\mathcal{T}_2^\epsilon \equiv \inf\{t > \mathcal{T}_1^\epsilon | q^\epsilon(t) = 0\}$ , and  $\sigma^\epsilon(t) = r/\epsilon$  for  $t \in [\mathcal{T}_1^\epsilon, \mathcal{T}_2^\epsilon)$ . This process repeats up to an arbitrary final time  $T$ . Define the time intervals  $\tau_i^{d,\epsilon}$  and  $\tau_i^{r,\epsilon}$  as

$$\tau_1^{d,\epsilon} = \mathcal{T}_1^\epsilon - 0, \quad (3a)$$

$$\tau_1^{r,\epsilon} = \mathcal{T}_2^\epsilon - \mathcal{T}_1^\epsilon, \quad (3b)$$

$$110 \quad \tau_2^{d,\epsilon} = \mathcal{T}_3^\epsilon - \mathcal{T}_2^\epsilon, \quad (3c)$$

$$\tau_2^{r,\epsilon} = \mathcal{T}_4^\epsilon - \mathcal{T}_3^\epsilon, \quad (3d)$$

and so on. These are the duration times for dry and rain events respectively.

The associated processes, as  $\epsilon \rightarrow 0$ , are defined as  $q(t)$  and  $\sigma(t)$  for the moisture and rain processes. (It would perhaps be appropriate to denote the limiting processes as  $q^0(t)$  and  $\sigma^0(t)$ , to indicate that they arise from  $q^\epsilon(t)$  and  $\sigma^\epsilon(t)$  in the limit  
115  $\epsilon \rightarrow 0$ . However, we will drop the superscript 0 from  $q^0(t)$  and  $\sigma^0(t)$  to ease notation.) The moisture process is the solution to the SDE,

$$dq(t) = m dt + D_0 dW_t, \quad q < b, \quad q(0) = 0, \quad (4)$$

with the unusual boundary condition as follows: Let the usual stopping time be  $\mathcal{T}_1 = \inf\{t > 0 | q(t) = b\}$ . Then at time  $t = \mathcal{T}_1$  the process  $q(t)$  jumps or “teleports” to  $q = 0$ . However, the function is defined as both 0 and  $b$  at  $\mathcal{T}_1$ . For convention, the  
120 process is defined as cadlag (continuous from the right with left hand limits), i.e.

$$\lim_{t \rightarrow (\mathcal{T}_1)^-} q(t) = b, \quad \lim_{t \rightarrow (\mathcal{T}_1)^+} q(t) = 0, \quad q(\mathcal{T}_1) = 0. \quad (5)$$

Then the process starts over using the dynamics of (4) until  $\mathcal{T}_2 = \inf\{t > \mathcal{T}_1 | q(t) = b\}$ , and the process repeats. The time intervals

$$\tau_i^d = \mathcal{T}_{i+1} - \mathcal{T}_i, \quad (6)$$

125 are the dry event durations. The rain event duration, on the other hand, is not defined for this limiting process, since rain events are instantaneous in the intense-rain-rate limit of  $\epsilon \rightarrow 0$ .

Example time series of the processes are shown in Figure 2. The processes with finite rain rate  $r/\epsilon$  for  $\epsilon > 0$  are shown in panels (a) and (b). Panel (a) is the moisture process  $q^\epsilon(t)$  defined in equation (2). The rain rate process is shown in panel (b) and takes the value  $r/\epsilon$  when  $q^\epsilon(t)$  reaches level  $b$  for the first time (panel (a) in black) and resets to zero when  $q^\epsilon(t)$  reaches zero (panel (a) in gray). This process repeats. The limiting processes are shown in panels (c) and (d). Panel (c) shows the limiting moisture process  $q(t)$  defined in equation (4) and panel (d) shows the rain process defined in equation (7). The moisture process is a Brownian motion with positive drift until reaching level  $b$ . When  $q(t) = b$ , the process  $\sigma(t)$  takes an infinite value and the moisture process is reset to zero.

From the definition of  $\tau_i^d$  above, the rain point process  $\sigma(t)$  is defined as

$$135 \quad \sigma(t) = b \sum_{i=1}^{\mathcal{N}(T)} \delta(t - \mathcal{T}_i), \quad (7)$$

where  $\mathcal{N}(T)$  is the random variable of the number of times the process  $q(t)$  reaches  $b$  in time  $T$ . The quantity  $b$  arises because the moisture process  $q^\epsilon$  loses moisture at a rate of  $r/\epsilon$  per time, on average. The moisture process  $q(t)$  loses all the moisture built up (which is an amount  $b$ ) instantaneously.

Note that  $q^\epsilon(t)$  has continuous paths while  $q(t)$  has jump discontinuities. Thus any mode of convergence between  $q^\epsilon$  and  $q$  with an associated metric (e.g. uniform or Skorohod) will fail (Kelley, 2017). Nevertheless, there is another way to define both  $q^\epsilon$  and  $q$  in which convergence with respect to  $L^2$  with the uniform metric on the space of continuous functions ( $\mathcal{C}[0, T]$ ) can be shown. To do so,  $q^\epsilon(t)$  is decomposed into an evaporating process,  $E^\epsilon(t)$ , and precipitating process  $P^\epsilon(t)$ . These processes are defined as

$$dE_t^\epsilon = \begin{cases} m dt + D_0 dW_t & \text{for } \sigma_t^\epsilon = 0 \\ 0 & \text{for } \sigma_t^\epsilon = r/\epsilon \end{cases}, \quad \text{and} \quad dP_t^\epsilon = \begin{cases} 0 & \text{for } \sigma_t^\epsilon = 0 \\ -\frac{r}{\epsilon} dt + D_1 dW_t & \text{for } \sigma_t^\epsilon = r/\epsilon \end{cases}. \quad (8)$$

Thus the moisture process  $q^\epsilon(t)$  is written as

$$q^\epsilon(t) = E^\epsilon(t) + P^\epsilon(t).$$

145 In the limit, the jumps will be captured in the  $P^\epsilon$  process. In the following section it will be shown (see Section 3.2) that  $E^\epsilon \rightarrow E$ , where  $E(t)$  is defined as the solution to the SDE


$$dE(t) = m dt + D_0 dW_t, \quad E(0) = 0. \quad (9)$$

Furthermore, the spike times of the  $\sigma$  process, which was defined above in (7), could now also be defined in terms of the  $E(t)$  process as  $\mathcal{T}_i = \inf\{t > 0 | E(t) = ib, i \in \mathbb{N}\}$ , i.e. the first passage time of Brownian motion with drift to  $ib$ .

### 150 3 Convergence to a Point Process

In this section convergence is shown both heuristically (e.g. Section 3.1) and rigorously (e.g. Sections 3.2 and 3.3).

Note that the simplest ideas of convergence break down when considering path-wise convergence of  $q^\epsilon$  to  $q$  and  $\sigma^\epsilon$  to  $\sigma$ . This is because  $q^\epsilon$  is a continuous process for all  $\epsilon > 0$ , whereas  $q$  is a process with jumps; and  $\sigma^\epsilon$  is left continuous with right-hand limits, whereas  $\sigma$  no longer is left continuous. Thus, there is no topology with associated metric  $d$  such that  $q^\epsilon \rightarrow q$  with



**Figure 2.** Realizations are plotted of the processes (a)  $q^\epsilon(t)$ , (b)  $\sigma^\epsilon(t)$  for rain rate  $r/\epsilon$  defined in equation (2) with  $\epsilon > 0$  and, on the other hand, the limiting ( $\epsilon \rightarrow 0$ ) processes (c)  $q(t)$  and (d)  $\sigma(t)$  defined in equation (4) and (7) respectively. Here  $m = r = 0.4$  mm/h with  $\epsilon = 0.01$ . Thus the effective rain rate is 40 mm/hr

155 respect to  $d$  (Kelley, 2017). However, one could try to show that  $q^\epsilon$  converges in a notion weaker than the Skorohod topology; see Kurtz (1991) for these conditions. Such convergence would happen in a topology which does not have an associated metric (see Jakubowski et al. (1997)). This approach is not pursued here as it is technical and does not give any insight to the model or approximation.

160 Instead, we pursue convergence in the following senses. The next three subsections prove convergence of the various processes introduced in Section 2. In Section 3.1 the Fokker-Planck equation for  $q^\epsilon$  is shown to converge (formally) to a Fokker-

Planck equation for  $q$ . This derivation gives rise to an interesting partial differential equation (PDE) with unusual “teleporting” boundary conditions. In Section 3.2 convergence in paths is shown for  $E^\epsilon$  to  $E$  with respect to the uniform metric for continuous functions on  $[0, T]$ . In Section 3.3 convergence is shown for  $\sigma^\epsilon$  to  $\sigma$  with respect to generalized functions. This norm is necessary because  $\sigma$  is a sum of Dirac delta functions. In addition, this convergence is natural to consider for applications where the errors are analyzed between using  $\sigma^\epsilon$  and a point process ( $\sigma$ ) in, for example, a climate model or as a model for observational time series.

### 3.1 Fokker-Planck Equation

In this section, we derive the Fokker-Planck equation of (4) by taking the formal asymptotic limit, as  $\epsilon \rightarrow 0$ , of the Fokker-Planck equation of (2). This mode of convergence provides some intuition for the behavior in the  $\epsilon \rightarrow 0$  limit.

The Fokker-Planck equation for (2) (see Hottovy and Stechmann (2015b)) is composed of two densities. These densities are denoted  $\rho_0$  and  $\rho_1$  for the dry state ( $\sigma^\epsilon = 0$ ) and the rain state ( $\sigma^\epsilon = 1$ ), respectively. These densities evolve according to the following Fokker-Planck equations:

$$\partial_t \rho_0 = -m \partial_q \rho_0 + \frac{D_0^2}{2} \partial_q^2 \rho_0 - \delta(q) f_1, \quad -\infty < q < b, t \geq 0, \quad (10)$$

$$\partial_t \rho_1 = \frac{r}{\epsilon} \partial_q \rho_1 + \frac{D_1^2}{2} \partial_q^2 \rho_1 + \delta(q-b) f_0, \quad 0 < q < \infty, t \geq 0, \quad (11)$$

where the fluxes  $f_i$  are defined as

$$f_0(q, t) = m \rho_0(q, t) - \frac{D_0^2}{2} \partial_q \rho_0(q, t) \quad (12a)$$

$$f_1(q, t) = -\frac{r}{\epsilon} \rho_1(q, t) - \frac{D_1^2}{2} \partial_q \rho_1(q, t), \quad (12b)$$

and with the following conditions,

$$\rho_0(b, t) = \rho_1(0, t) = 0, \quad (13)$$

$$\int_{-\infty}^{\infty} \rho_0(q, t) + \rho_1(q, t) dq = 1, \quad t \geq 0, \quad (14)$$

which are absorbing boundary conditions and the normalization condition, respectively. This implies that once the particle reaches  $q = b$  (or  $q = 0$ ) the particle is removed and added to the state  $\sigma = 1$  (or  $\sigma = 0$ ) (Gardiner, 2004). Thus the particle reaches  $q = b$  but can not be found there  $\rho_0(b, t) = 0$ . Note that the flux terms in (12a) and (12b) contain  $\rho_0, \rho_1$  terms and its derivative. If the derivatives are zero then the absorbing boundary conditions would imply the Dirac delta coupling terms in (10) and (11) are zero. However, this is not the case. The Fokker-Planck equation must be solved on three separate intervals of  $(-\infty, 0]$ ,  $[0, b]$  and  $[b, \infty)$ . This leads to points of non-differentiability for  $\rho_0$  and  $\rho_1$  at  $q = 0$  and  $q = b$ . For an example, see the stationary solutions in Section 4 and Figure 3.

To obtain the equations (10) and (11) a further approximation is used. In Hottovy and Stechmann (2015b),  $q(t)$  is modeled similar to equation (2) except the  $\sigma$  process switches states at a random time with rate  $\lambda$  after  $q(t)$  reaches the threshold.



190 Call this process  $q^\lambda(t)$ . The Fokker-Planck type equation for  $q^\lambda(t)$  uses terms from the Master Equation (see Gardiner (2004) Section 3.5). The Fokker-Planck in equation (10) and (11) is derived by taking an asymptotic of the equations for  $q^\lambda$  and taking a limit as  $\lambda \rightarrow \infty$ . That is, as the random switching time becomes small.

One interesting property of these Fokker-Planck equations is the appearance of Dirac-delta source terms, which represent transitions between the dry state and rain state. For instance, in (10), a Dirac delta source term arises at  $q = 0$ , and it represents  
 195 the transition from the rain state ( $\sigma^\epsilon = 1$ ) to the dry state ( $\sigma^\epsilon = 0$ ) when the (raining) moisture process reaches the lower threshold at  $q = 0$ . The magnitude of this Dirac delta source term is  $-f_1|_{q=0}$ , which is the outward flux of  $\rho_1$  at the lower threshold,  $q = 0$ , as defined from (11) and (12b).

The proposed limit as  $\epsilon \rightarrow 0$  for the Fokker-Planck equation is

$$\partial_t \rho_0 = -m \partial_q \rho_0 + \frac{D_0^2}{2} \partial_q^2 \rho_0 + f_0|_{q=b} \delta(q), \quad -\infty < q < b, t \geq 0, \quad (15)$$

$$200 \quad \rho_1 = 0. \quad (16)$$

with the following conditions,

$$\rho_0(b, t) = 0 \quad (17)$$

$$\int_{-\infty}^b \rho_0(q, t) dq = 1. \quad (18)$$

This Fokker-Planck equation is much different than the coupled system (10)-(11). For example, the coupled system is defined  
 205 for all  $-\infty < q < \infty$  whereas (15) is only defined for  $-\infty < q \leq b$ . This is because the boundary  $q = b$  become impassable due to the teleporting boundary; or, in physical terms, the rain rate is so strong that the moisture  $q$  moves above the threshold  $b$  by only a small  $O(\epsilon)$  amount that vanishes as  $\epsilon \rightarrow 0$ . The restriction of  $q$  can be seen in the stationary densities in Section 4. There, the stationary density for state  $\sigma^\epsilon(t) = 1$  decays to zero quickly. Another an interesting property of this Fokker-Planck equation: the absorbing boundary condition at  $q = b$  in (17) is actually coupled to a Dirac-delta source at  $q = 0$  in the Fokker-  
 210 Planck equation in (15). In this coupling, the flux  $f_0|_{q=b}$  of absorption at the boundary is also equal to the magnitude of the source  $f_0|_{q=0} \delta(q)$  which inserts mass at  $q = 0$ . Therefore, when the process is absorbed at  $q = b$ , it is re-inserted at  $q = 0$ , and in this way it represents a teleporting boundary condition.

The convergence of the time-dependent Fokker-Planck equation (10)-(11) is shown through an asymptotic expansion in the Appendix. The analysis is shown for the full time-dependent Fokker-Planck equation to show that the transient solutions to  
 215 these equations also converge as  $\epsilon \rightarrow 0$ . The method used is a matched asymptotic expansion argument (Bender and Orszag, 2013). This argument shows how the interesting delta function condition of the probability flux arises in (15).

### 3.2 Pathwise Convergence

Rigorous mathematical convergence is now considered. For this section and next, a useful lemma is first stated and proved. In essence, the lemma states that, for a finite time interval  $[0, T]$ , it is (exponentially) unlikely that a large number of rain events  
 220 will occur.

**Lemma 1.** Let  $\mathcal{N}^\epsilon(T)$  be the number of rain events for the  $q^\epsilon$  process defined in (4). Then for  $0 < s < \min\{rb/\epsilon D_2^2, mb/D_1^2\}$

$$P(\mathcal{N}^\epsilon(T) = N) \leq \exp \left\{ sT - \frac{Nmb}{D_1^2} \left( \sqrt{1 + \frac{2D_1^2 s}{m^2}} - 1 \right) \right\}.$$

The proof of the lemma is contained in Appendix. This lemma shows that the probability of  $N$  events decays exponentially in  $N$ . With this lemma, pathwise convergence is now considered. Recall from the discussion at the beginning of the section that we consider convergence not for  $q^\epsilon$  but for the evaporating process  $E^\epsilon$ . Convergence from  $E^\epsilon$  to  $E$  is shown in  $L^2(\Omega)$  with respect to the uniform metric on the space of continuous functions  $C[0, T]$ .

**Theorem 1.** Let  $q_t^\epsilon$  be defined as

$$q_t^\epsilon = E_t^\epsilon + P_t^\epsilon$$

225 where  $E_t^\epsilon, P_t^\epsilon$  are solutions to the SDEs in (8). Furthermore let  $E_t$  be defined as the solution to (9). Then

$$\lim_{\epsilon \rightarrow 0} E \left[ \left( \sup_{0 \leq t \leq T} |E_t^\epsilon - E_t| \right)^2 \right] = 0. \quad (19)$$

This theorem shows that the moistening process  $E^\epsilon$  converges to the process  $E$  as  $\epsilon \rightarrow 0$ . Furthermore the moistening process  $E$  contains all the dynamics of the joint  $(E, \sigma)$  process. The rigorous proof is shown in Appendix. The proof relies on the processes  $E^\epsilon$  and  $E$  being driven by the same white noise process. Thus they converge to each other by showing that the  
230 first two moments of  $\tau^{r, \epsilon}$  converge to zero (see equations (A27)).

### 3.3 Distributional Convergence

In this subsection  $L^2(\Omega)$  convergence of  $\sigma^\epsilon$  to  $\sigma$  is shown with respect to a generalized function norm. This norm is considered here due to the nature of the delta function. It is also a natural norm to consider as it is an integrated error. That is, this norm considers the accumulation of errors after running the model for time  $T > 0$ .

235 **Theorem 2.** Let  $\phi : [0, T] \rightarrow \mathbb{R}$  be a test function in  $C_c^\infty(0, T)$ . Let  $\sigma^\epsilon(t)$  and  $\sigma(t)$  be defined as in (2) and (7), respectively. Then

$$\lim_{\epsilon \rightarrow 0} E[|\langle \sigma^\epsilon(t), \phi(t) \rangle - \langle \sigma(t), \phi(t) \rangle|^2] = 0, \quad (20)$$

where

$$\langle f(t), g(t) \rangle = \int_0^T f(t)g(t) dt. \quad (21)$$

240 The technical details of the proof are given in Appendix. The procedure is similar to the proof of Theorem 1. However, here the case for different numbers of rain events in the time interval  $0 \leq t \leq T$  for the processes  $\sigma$  and  $\sigma^\epsilon$  must be considered in a different way (see equation (A45)). This leads to needing estimates on the first four moments of the event duration  $\tau^{r, \epsilon}$ .

## 4 Statistics and Applications

In this section, important statistics and applications of the processes  $(q^\epsilon, \sigma^\epsilon)$  and  $(q, \sigma)$  are presented. These statistics show the differences between the processes as well as give motivation for the approximations. These include the stationary Fokker-Planck solution, the rain and dry event distributions, rain fraction, and an application of Theorem 2.

### 4.1 Stationary Fokker-Planck Equation

Here the analytical solutions to the stationary Fokker-Planck equation are given. The stationary Fokker-Planck equation for the process  $q^\epsilon$  is

$$0 = -m\partial_q \rho_0^\infty + \frac{D_0^2}{2} \partial_q^2 \rho_0^\infty - \delta(q) f_1, \quad -\infty < q < b, \quad (22)$$

$$0 = \frac{r}{\epsilon} \partial_q \rho_1^\infty + \frac{D_1^2}{2} \partial_q^2 \rho_1^\infty + \delta(q-b) f_0, \quad 0 < q < \infty, \quad (23)$$

with the conditions

$$\rho_0(b, t) = \rho_1(0, t) = 0, \quad (24)$$

$$\int_{-\infty}^{\infty} \rho_0(q, t) + \rho_1(q, t) dq = 1, \quad t \geq 0, \quad (25)$$

The analytical solutions are found in Hottovy and Stechmann (2015a) and are reproduced here. They are


$$\rho_1^\infty(q) = \frac{1}{b} \frac{m}{r/\epsilon + m} \left\{ 1 - \exp\left(-\frac{2r}{\epsilon D_1^2} q\right) \right\}, \quad \text{for } 0 \leq q \leq b \quad (26)$$

$$\rho_1^\infty(q) = \frac{1}{b} \frac{m}{r/\epsilon + m} \left\{ 1 - \exp\left(-\frac{2r}{\epsilon D_1^2} b\right) \right\} \exp\left(-\frac{2r}{\epsilon D_1^2} (q-b)\right), \quad \text{for } 0 \leq q \leq b, \quad (27)$$

and similarly for  $\rho_0^\infty$ . The densities are plotted in Figure 3. The black curve is the  $\rho_0^\infty$  density for one value of  $\epsilon = 1$ . For the other masses  $\rho_0^\infty$  changes very little (order  $\epsilon$ ) and are not shown. The density  $\rho_1^\infty$  is plotted in grey for various values of  $\epsilon$ . The density is scaled by  $\epsilon^{-1}$ . The dashed gray line is the limiting shape. Thus as  $\epsilon \rightarrow 0$  the density is a uniform distribution on the interval  $[0, b]$  that tends to zero. Note that the absorbing boundary conditions from (24) are satisfied ( $\rho_0^\infty(b) = 0$  and  $\rho_1^\infty(0) = 0$ ). However their derivatives (the fluxes) are non-zero. Thus the Dirac delta terms in the Fokker-Planck equation are non-zero.

### 4.2 Event Duration

Another statistic studied here is the event duration probability density. This density gives information on the probability of a dry/rain event lasting time  $t$  minutes. For the process  $q^\epsilon$  both the dry and rain states are Brownian motions with drift ( $m$  for dry and  $r/\epsilon$  for rain). Thus the event duration densities are the first passage to  $q = b$  densities for Brownian motion with drift (Gardiner, 2004, Section 5.5.1). These densities were found in Stechmann and Neelin (2014) and are reproduced here. The



**Figure 3.** Stationary probability density functions (pdf) for the  $\rho_0^\infty$  density (black line) and  $\epsilon\rho_1^\infty$  density for values of  $\epsilon = 1, 0.5, 0.25,$  and  $0.1$  (gray lines). The dashed line is the limit shape of the  $\rho_1^\infty$  density scaled by  $\epsilon^{-1}$ .

event duration density for a rain event is

$$270 \quad \rho_{1t} = \frac{b}{\sqrt{2\pi D_1^2}} \exp\left\{\frac{rb}{\epsilon D_1^2}\right\} \exp\left\{\frac{-b^2}{2D_1^2 t}\right\} \exp\left\{\frac{-r^2 t}{2\epsilon^2 D_1^2}\right\} t^{-3/2}, \quad (28)$$

and similarly for the dry event duration  $\rho_{0t}$ . The density for rain events changes with  $\epsilon$  while the dry event density does not. The rain event duration has cutoffs for short and long times. They are

$$\text{short time cutoff: } \frac{b^2}{2D_1^2} \quad \text{long time cutoff: } \frac{2\epsilon^2 D_1^2}{r^2}.$$


The short time cutoff is independent of  $\epsilon$  while the long time cutoff tends to zero in  $O(\epsilon^2)$ . This implies that for smaller  $\epsilon$ , the extreme events are being cutoff quickly. Hence, in the  $\epsilon \rightarrow 0$  limit, extreme rainfall events do not occur. In order to preserve extreme rainfall events in a point-process model of rainfall, the rain rate amplitude would need to be modeled as a stochastic process, since rain event durations are assumed to be short (see Section 4.5).

275 These densities are plotted in Figure 4 on a log-log scale. The dry density  $\rho_{0t}$  is the black curve. The rain event duration density  $\rho_{1t}$  is plotted in gray for various values of  $\epsilon$ . The dashed gray line is a density for small  $\epsilon = 0.01$ .

### 4.3 Average Cloudiness

The average cloudiness is the fraction of time that the stationary process is in the rain state  $\sigma > 0$ . It is defined as,

$$E[\sigma/(r/\epsilon)] = \int_0^\infty \rho_1^\infty(q) dq$$



**Figure 4.** Event duration probability density functions (pdf) for dry events ( $\rho_{0t}$  black line) and rain events ( $\rho_{1t}$  gray lines) for  $\epsilon = 1, 0.5, 0.25,$  and  $0.1$ . The dashed line is for  $\epsilon = 0.01$ .

where  $\rho_1^\infty$  is the solution to the steady state Fokker-Planck equation (23). For the process  $\sigma^\epsilon$  it is

$$E[\sigma^\epsilon / (r/\epsilon)] = \frac{1}{m + r/\epsilon}.$$

Futhermore, the variance of the average cloudiness is

$$\text{var}(\sigma^\epsilon / (r/\epsilon)) = \frac{1}{m + r/\epsilon} \left( 1 - \frac{1}{m + r/\epsilon} \right) = \frac{m + r/\epsilon - 1}{(m + r/\epsilon)^2}.$$

The average cloudiness is zero when using the point process model  $\sigma$ . However, for  $\sigma^\epsilon$  as  $\epsilon$  tends to zero, the average cloudiness and its variance are  $O(\epsilon)$  and nonzero.

#### 4.4 Total Rainfall

For an example of using the results of Theorem 2, consider calculating the total rainfall at a specific grid point of a GCM. Define the rainfall time series at a point as

$$A(t) = \sigma(t)\phi(t)$$

where  $\sigma$  encodes the rainfall rate, and  $\phi(t) = 1$  when there is a cloud (or rainfall) and  $\phi(t) = 0$  otherwise. One question is: what is the impact on total rainfall in a simulation of time  $T$  between the rain processes  $\sigma^\epsilon$  and  $\sigma$ ?

Theorem 2 states that the integrated difference of the total rainfall tends to zero. That is,

$$\lim_{\epsilon \rightarrow 0} \int_0^T |A^\epsilon(t) - A(t)| dt = \int_0^T |\sigma^\epsilon(t)\phi(t) - \sigma(t)\phi(t)| dt.$$

The proof of the theorem in the Appendix yields more information than that. For example, if it is known that there are  $N$  rain events after time  $T$ , then estimate (A45), yields

$$\int_0^T |A^\epsilon(t) - A(t)| dt \leq \sum_{i=1}^N \left( \left( \frac{r}{\epsilon} \right)^2 K^2 E[|\tau_i^{r,\epsilon}|^4] + E \left[ \left| \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} - b \right) \phi(\mathcal{T}_{2i-1}^\epsilon) \right|^2 \right] + E [|\phi(\mathcal{T}_{2i-1}^\epsilon) - \phi(\mathcal{T}_i)|^2] \right), \quad (29)$$

where the probability on  $N$  events are one and where the final remainder term has been dropped (for clarity of this calculation). Here  $K$  is a global bound on the function  $\phi$ . For this example  $K = 1$ . From the estimates in the proof, the right hand side is bounded by terms which have the first four moments of the event duration time  $\tau^{r,\epsilon}$ , the global bound  $K$  of  $\phi$ , and the number of rain events. Thus, Theorem 2 and its proof give details on the differences for key atmospheric elements when using the rain processes  $\sigma^\epsilon$  and  $\sigma$ .

#### 4.5 Modifications for the point process model

The issue of how to use finite-event-duration model to inform parameter selection is discussed in this subsection. There are two potential points of concerns with the point process model. One is that the point process model  $\sigma(t)$  has rain events of duration zero, whereas the rain process model  $\sigma^\epsilon(t)$  has rain events of duration  $\tau^{\epsilon,r}$ . Thus for time  $T$ ,

$$T = \sum_{i=1}^{\mathcal{N}(T)} \tau_i^d + \tau_i^{\epsilon,r},$$

where  $\mathcal{N}(T)$  is the random number of rain events in  $T$  time. For finite  $\epsilon > 0$  the value of  $\mathcal{N}(T)$  is larger than for the point process model, on average.

There are many potential solutions for the issue of zero event times for the point process model. One example is to modify dry duration pdf to account for small but finite size of rain events. That is, let  $\tau^r$  be a random variable which models a finite size rain event. For example see equation (28). Define

$$\tau^{d'} = \tau^d + \tau^r,$$

to be a new dry event random variable. This new event distribution would account for the rain event within the dry event. As for the moisture process  $q$ , once the threshold of  $q = b$  is met, the process then holds at  $b$  for a random time of  $\tau^r$ . Then the process would jump to  $q = 0$ .

Another potential modification is the definition of rain amount. For the model with finite  $\epsilon > 0$ , for each event the model rains  $b$  amount over a random time  $\tau^{r,\epsilon}$ . For the point process model, the model rains  $b$  amount instantaneously. Here the discrepancy between the two models is captured in Theorem 2 and the example of total rainfall in the example in subsection 3.3. A possible modification to the model would be to use a finite-event-duration model to help assign a random magnitude to each point-process model event. For example, let  $b_i$  be random variables with some distribution from a finite-event-duration model which accounts for random rain amounts. Then the point process model can be modified from Eq. (7) as

$$\sigma(t) = \sum_{i=1}^{\mathcal{N}(T)} b_i \delta(t - \mathcal{T}_i). \quad (30)$$

## 5 Conclusions

In this paper, a threshold model for moisture and rain was shown to converge to a point process and related processes, and to converge for various modes of convergence. By demonstrating this type of convergence, the simple ideas of a point-process model of rainfall, which at first may appear to be only an empirical model, can be linked with underlying physical processes and evolution of moisture.

Here convergence for the moisture processes was defined and shown for the Fokker-Planck equation as well as the paths of the processes. Furthermore, the convergence of the rain process were shown in mean square difference with respect to the space of generalized functions.

Using a point process to approximate rainfall allows simplification for computation and exact formulas. For example, the autocorrelation function is known in the case of point processes as shown in Abbott et al. (2016). Furthermore, point processes have been studied extensively in the neural science literature (Sacerdote and Giraud, 2013) and many statistics have been derived. Some examples of exact statistics are shown. These examples have interesting characteristics as the small parameter  $\epsilon$  tends to zero.

The proofs shown here are revealing on their own, and they demonstrate further details of the convergence. The Fokker-Planck derivation in Section 3.1 shows that the density for the moisture in the rain state tends to zero while the flux term remains nonzero, allowing for the “teleporting” boundary condition that arises for the limiting moisture process. For the convergence of paths of moisture shown in Theorem 1, the moisture process must first be decoupled into a moistening and precipitating process. Then the moistening process is shown to converge (Theorem 1) while the precipitating process contains all of the discontinuities. Finally, the proof of convergence of the rain processes in Theorem 2 gives estimates that would be useful for determining the error rates for using the point process approximation. This is done in an example of total rainfall.

### Appendix A: Proofs from Section 3

The rigorous mathematical proofs and formal asymptotic analysis for the results presented in Section 3 are given in this appendix.

#### A1 Derivation of the Fokker-Planck Equation

The Fokker-Planck equation for the process  $q^\epsilon$  is

$$\partial_t \rho_0 = -m \partial_q \rho_0 + \frac{D_0^2}{2} \partial_q^2 \rho_0 - \delta(q) f_1, \quad -\infty < q < b, t \geq 0, \quad (\text{A1})$$

$$\partial_t \rho_1 = \frac{r}{\epsilon} \partial_q \rho_1 + \frac{D_1^2}{2} \partial_q^2 \rho_1 + \delta(q-b) f_0, \quad 0 < q < \infty, t \geq 0, \quad (\text{A2})$$

To derive the limiting ( $\epsilon \rightarrow 0$ ) Fokker–Planck equation, the analysis follows the procedure of matched asymptotic expansions (see, e.g., Bender and Orszag (2013)). Consider two regions  $[0, \epsilon]$  and  $[\epsilon, \infty)$ . Let  $\rho_{1,B}$  be the density in the first region, which is a boundary layer region and  $\rho_{1,A}$  be the density away from this region. Since the Fokker-Planck equations have parameter  $\epsilon$ , let  $\rho_{1,B}$  have the asymptotic expansion of the form,

$$\rho_{1,B} = \rho_{1,B}^0 + \epsilon \rho_{1,B}^1 + O(\epsilon^2),$$

and let  $\rho_{1,A}$  have the asymptotic expansion

$$\rho_{1,A} = \rho_{1,A}^0 + \epsilon \rho_{1,A}^1 + O(\epsilon^2).$$

. These expansions are stopped at the order  $\epsilon$  level. This is due to the higher order terms not having an impact on the limiting equation. In the end, it will be shown that  $\rho_{1,A}^0 = \rho_{1,B}^0$ . For the region away from the boundary the density is  $\rho_{1,A}^1 = \frac{1}{\tau} f_0(b, t)$ .

335 This allows for the teleporting boundary condition of  $f_0(b, t)\delta(q)$ .

First to show  $\rho_{1,B} = 0$ , consider equation (A2) with the rescaled variable  $\tilde{q} = \frac{1}{\epsilon}q$ . This yields the equation

$$\partial_t \rho_{1,B} = \frac{r}{\epsilon^2} \partial_{\tilde{q}} \rho_{1,B} + \frac{D_1^2}{2\epsilon^2} \partial_{\tilde{q}}^2 \rho_{1,B} \quad (\text{A3})$$

Substituting this expansion into equation (A3) yields, at order  $\epsilon^{-2}$  and order  $\epsilon^{-1}$ , respectively,

$$O(\epsilon^{-2}): 0 = r \partial_{\tilde{q}} \rho_{1,B}^0 + \frac{D_1^2}{2} \partial_{\tilde{q}}^2 \rho_{1,B}^0, \quad (\text{A4a})$$

$$340 \quad O(\epsilon^{-1}): 0 = r \partial_{\tilde{q}} \rho_{1,B}^1 + \frac{D_1^2}{2} \partial_{\tilde{q}}^2 \rho_{1,B}^1. \quad (\text{A4b})$$

Note that the  $O(1)$  equation is not written. At this order and higher, there are iterative PDEs written for  $\rho_{1,B}^i$  for  $i \geq 2$ . These terms will converge to zero at a rate  $O(\epsilon^i)$  and thus are not considered here. By solving the order  $\epsilon^{-2}$  equation in (A4a) and applying the absorbing boundary condition at  $\tilde{q} = 0$ , one arrives at

$$\rho_{1,B}^0 = C_1(t) \left( 1 - \exp \left[ -\frac{2r}{D_1^2} \tilde{q} \right] \right). \quad (\text{A5})$$

345 The order  $\epsilon^{-1}$  equation in (A4b) has essentially the same solution as above, and, after applying the absorbing boundary condition, one finds

$$\rho_{1,B}^1 = C_2(t) \left( 1 - \exp \left[ -\frac{2r}{D_1^2} \tilde{q} \right] \right). \quad (\text{A6})$$

Now consider the interval away from the boundary  $[\epsilon, \infty)$ . Let  $\rho_{1,A}$  be the density in this region. The equation in this region  
350 is

$$\partial_t \rho_{1,A} = \frac{r}{\epsilon} \partial_q \rho_{1,A} + \frac{D_1^2}{2} \partial_q^2 \rho_{1,A} + \delta(q-b) f_0(b, t). \quad (\text{A7})$$



Note that the  $\delta$  term acts on  $f_0$  which is a function of  $\rho_0$ . The asymptotic expansion is for  $\rho_1$  only in the  $[\epsilon, \infty)$  region, and thus the density  $\rho_0$  is an order one term. Substituting the expansion into equation (A7) gives the following equations, separated into their orders of  $\epsilon$ ,

$$355 \quad O(\epsilon^{-1}) : 0 = r \partial_q \rho_{1,A}^0 \quad (\text{A8a})$$

$$O(1) : \partial_t \rho_{1,A}^0 = r \partial_q \rho_{1,A}^1 + \frac{D_1^2}{2} \partial_q^2 \rho_{1,A}^0 + \delta(q-b) f_0(b,t). \quad (\text{A8b})$$

The order  $\epsilon^{-1}$  equation in (A8a) has the solution

$$\rho_{1,A}^0 = C_3(t). \quad (\text{A9})$$

Note that  $\rho_{1,A}$  is a density and thus  $\rho_{1,A}^0$  must be integrable on  $[O(\epsilon), \infty)$ . Thus  $C_3(t) = 0$  and

$$360 \quad \rho_{1,A}^0 = 0. \quad (\text{A10})$$

From the order one equation in (A8b), by substituting in  $\rho_{1,A}^0 = 0$ , we arrive at

$$\rho_{1,A}^1 = \begin{cases} C_4(t) & \text{for } \epsilon \leq q < b \\ C_4(t) - \frac{1}{r} f_0(b,t), & \text{for } q \geq b \end{cases} \quad (\text{A11})$$

Note that the constant of integration in each interval of  $b$  must be the same. Otherwise, the magnitude of the  $\delta$  function in (A8b) would not be correct. The density  $\rho_{1,A}^1$  must be integrable, which implies that

$$365 \quad C_4(t) = \frac{1}{r} f_0(b,t). \quad (\text{A12})$$

It is assumed that the matching between the  $A$  and  $B$  solutions must occur at an intermediate location or overlapping region.

That is, for values of  $q = \epsilon^{1/2}$ ,

$$\rho_{1,B}^0(\epsilon^{1/2}, t) = \rho_{1,A}^0(\epsilon^{1/2}, t)$$

and

$$\rho_{1,B}^1(\epsilon^{1/2}, t) = \rho_{1,A}^1(\epsilon^{1/2}, t).$$

The first equation implies that  $C_1(t) = 0$  and  $\rho_{1,B}^0 = \rho_{1,A}^0 = 0$ . In the limit as  $\epsilon \rightarrow 0$  the second equation yields

$$C_2(t) = \frac{1}{r} f_0(b,t). \quad (\text{A13})$$

Thus the densities are

$$\rho_1^0 = 0 \quad (\text{A14})$$

370 and

$$\rho_1^1 = \begin{cases} \frac{1}{r} f_0(b,t) \left(1 - \exp\left[-\frac{2r}{D_1^2} q\right]\right) & 0 \leq q \leq O(\epsilon) \\ \frac{1}{r} f_0(b,t) & O(\epsilon) \leq q \leq b \\ 0 & b < q \end{cases} \quad (\text{A15})$$

Note the flux of  $\rho_1$  at  $q = 0$  is, to leading order, in terms of  $\rho_1^1$ ,

$$f_1(0, t) = r\rho_1^1(0, t) + \epsilon \frac{D_1^2}{2} \partial_q \rho_1^1(0, t). \quad (\text{A16})$$

Using the asymptotic formula for  $\rho_1^1$  yields

$$375 \quad f_1(0, t) = \frac{D_1^2}{2} \epsilon \left\{ \frac{1}{r} f_0(b, t) \left( \frac{2r}{D_1^2 \epsilon} \right) \right\} = f_0(b, t). \quad (\text{A17})$$

Consequently, while the rain-state density itself is small (i.e.,  $\rho_1^0 = 0$ ), the flux  $f_1$  of the rain state is  $O(1)$ , and its value  $f_1(0, t)$  at the threshold  $q = 0$  represents an  $O(1)$  flux from the rain state to the dry state.

Thus the Fokker-Planck type equation for  $q(t)$  is

$$\partial_t \rho_0 = -m \partial_q \rho_0 + \frac{D_0^2}{2} \partial_q^2 \rho_0 + f_0|_{q=b} \delta(q), \quad -\infty < q < b, t \geq 0, \quad (\text{A18})$$

$$380 \quad \rho_1 = 0. \quad (\text{A19})$$

with the following conditions,

$$\rho_0(b) = 0, \quad (\text{A20})$$

$$\int_{-\infty}^b \rho_0(q, t) dq = 1. \quad (\text{A21})$$

## 385 **A2 Proof of Lemma 1**

*Proof of Lemma 1.* Note that the process  $\mathcal{N}^\epsilon(T)$  is a renewal process. It is defined by the interarrival times,

$$S_n = \tau_n^{d, \epsilon} + \tau_n^{r, \epsilon}, \quad n \geq 1, \quad (\text{A22})$$

where  $\tau_i^{d, \epsilon}$  ( $\tau_i^{r, \epsilon}$ ) is the duration for the  $i$ th dry (rain) event of the  $\sigma^\epsilon$  process. Note that  $S_n$  is used instead of  $\mathcal{T}_n - \mathcal{T}_{n-2}$  to align with common notation of renewal processes. The distributions of  $\tau_i^d$  and  $\tau_i^{d, \epsilon}$  are the same and are independent of  $\epsilon$ , while  $\tau_i^{r, \epsilon}$  depends on  $\epsilon$ . For the lemma, the quantity of interest is the probability of having  $N$  rain events in time  $T$ , which is defined as

$$P(\mathcal{N}^\epsilon(T) = N) = P(S_1 + S_2 + \cdots + S_N \leq T, \quad S_1 + S_2 + \cdots + S_N + S_{N+1} > T). \quad (\text{A23})$$

The probability on the right hand side is estimated crudely by only considering one of the two events. Note that  $S_1, S_2, S_3, \dots, S_n$  are IID random variables with  $E[S_1] = E[\tau^{d, \epsilon} + \tau^{r, \epsilon}]$ , and  $\sigma^2 = \text{Var}(S_1) < \infty$ , so that

$$P(\mathcal{N}^\epsilon(T) = N) \leq P(S_1 + S_2 + \cdots + S_N \leq T). \quad (\text{A24})$$

395 The above probability is estimated by using a variant of the Chernoff bound (Hoeffding, 1994). That is,

$$P(S_1 + S_2 + \cdots + S_N \leq T) \leq \exp(sT) \prod_{i=0}^n E[e^{-sS_i}], \quad (\text{A25})$$

for any  $s > 0$ , where  $E[e^{-sS_i}] = M_{S_i}(s)$  is the moment generating function for the random variable  $S_i$ . The moment generating function can be factored due to independence of  $\tau_i^{r,\epsilon}$  and  $\tau_i^{d,\epsilon}$ :

$$M_{S_i}(s) = M_{\tau_i^{r,\epsilon}}(s)M_{\tau_i^{d,\epsilon}}(s). \quad (\text{A26})$$

400 These moment generating functions are computed explicitly from the distributions found in Hottovy and Stechmann (2015b).

They are,

$$M_{\tau_i^{r,\epsilon}} = \int_0^\infty e^{-st} \rho_r(t) dt = \exp \left\{ \frac{-rb}{\epsilon D_2^2} \left( \sqrt{1 + \frac{2D_2^2 s \epsilon^2}{r^2}} - 1 \right) \right\}, \quad (\text{A27})$$

$$M_{\tau_i^{d,\epsilon}} = \int_0^\infty e^{-st} \rho_d(t) dt = \exp \left\{ \frac{-mb}{D_1^2} \left( \sqrt{1 + \frac{2D_1^2 s}{m^2}} - 1 \right) \right\}, \quad (\text{A28})$$

which are defined for  $s < \min\{rb/\epsilon D_2^2, mb/D_1^2\}$ . Chernoff's bound then yields

$$405 P(\mathcal{N}^\epsilon(T) = N) \leq P(S_1 + S_2 + \dots + S_n \leq T) \quad (\text{A29})$$

$$\leq \exp(sT) \prod_{i=0}^N E[e^{-sS_i}] \quad (\text{A30})$$

$$= \exp \left\{ sT - \frac{Nrb}{\epsilon D_2^2} \left( \sqrt{1 + \frac{2D_2^2 \epsilon^2 s}{r^2}} - 1 \right) - \frac{Nmb}{D_1^2} \left( \sqrt{1 + \frac{2D_1^2 s}{m^2}} - 1 \right) \right\} \quad (\text{A31})$$

$$\leq \exp \left\{ sT - \frac{Nmb}{D_1^2} \left( \sqrt{1 + \frac{2D_1^2 s}{m^2}} - 1 \right) \right\}. \quad (\text{A32})$$

□

### 410 A3 Proof of Theorem 1

*Proof of Theorem 1.* To begin, note that the SDEs for  $E^\epsilon$  and  $E$  (see (8)) only differ when  $\sigma^\epsilon(t) = r/\epsilon$ . Thus, the solutions to the SDEs give the formula

$$|E^\epsilon(t) - E(t)| = \left| \sum_{i=1}^{\mathcal{N}^\epsilon(T)} \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} m dt + \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} D_0 dW_t \right|, \quad (\text{A33})$$

where  $\mathcal{N}^\epsilon(T)$  is the number of rain events for  $T < \infty$  and  $\epsilon > 0$  fixed. Note that interval  $[\mathcal{T}_{2i-1}^\epsilon, \mathcal{T}_{2i}^\epsilon]$  has been written as  
 415  $[\mathcal{T}_{2i-1}^\epsilon, \mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}]$  to emphasize the rain event duration  $\tau_i^{r,\epsilon}$ . To proceed, the number of rain events is conditioned to be  $N$ . Note that  $m > 0$  and the stochastic integral is a martingale and Doob's maximal inequality yields,

$$E \left[ \left( \sup_{0 \leq t \leq T} |E^\epsilon(t) - E(t)| \right)^2 \right] \leq \sum_{N=1}^{\infty} 4E \left[ \left| \sum_{i=1}^N \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} m dt + \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} D_0 dW_t \right|^2 \middle| \mathcal{N}^\epsilon(T) = N \right] P(\mathcal{N}^\epsilon(T) = N). \quad (\text{A34})$$

By Lemma 1, the sum above converges due to the fast decay of  $P(\mathcal{N}^\epsilon(T) = N)$  as  $N \rightarrow \infty$ . Applying the Cauchy-Schwarz inequality to the sum and the Itô isometry to the stochastic integral yields

$$420 \quad E \left[ \left( \sup_{0 \leq t \leq T} |E^\epsilon(t) - E(t)| \right)^2 \right] \leq \sum_{N=1}^{\infty} C_N E [m^2 |\tau^{r,\epsilon}|^2 + D_0^2 |\tau^{r,\epsilon}| \mid \mathcal{N}^\epsilon(T) = N] P(\mathcal{N}^\epsilon(T) = N), \quad (\text{A35})$$

where  $\tau^{r,\epsilon}$  denotes the general event duration, which has the same distribution as all of the IID  $\tau_i^{r,\epsilon}$ . This sum converges due to the fast decay of  $P(\mathcal{N}^\epsilon(T) = N)$  as shown in Eq.(A32).

To finish the proof the following moments of  $\tau_i^{r,\epsilon}$  are used. The integrals can be computed exactly using the densities for  $\tau_i^{r,\epsilon}$  found in Hottovy and Stechmann (2015b). They are

$$425 \quad E[\tau_i^{r,\epsilon}] = \frac{b\epsilon}{r}, \quad E[|\tau_i^{r,\epsilon}|^2] = \frac{bD^2\epsilon^3}{r^3} + \frac{b^2\epsilon^2}{r^2}. \quad (\text{A36})$$

Thus the limit is

$$\lim_{\epsilon \rightarrow 0} E \left[ \left( \sup_{0 \leq t \leq T} |E^\epsilon(t) - E(t)| \right)^2 \right] \leq \lim_{\epsilon \rightarrow 0} \sum_{N=1}^{\infty} C_N E [m^2 |\tau^{r,\epsilon}|^2 + D_0^2 |\tau^{r,\epsilon}| \mid \mathcal{N}^\epsilon(T) = N] P(\mathcal{N}^\epsilon(T) = N), \quad (\text{A37})$$

$$\leq \sum_{N=1}^{\infty} \lim_{\epsilon \rightarrow 0} C_N \left[ m^2 \left( \frac{bD^2\epsilon^3}{r^3} + \frac{b^2\epsilon^2}{r^2} \right) + D_0^2 \left( \frac{b\epsilon}{r} \right) \right] P(\mathcal{N}^\epsilon(T) = N) \quad (\text{A38})$$

$$= 0, \quad (\text{A39})$$

430 where Tonelli's theorem allows the limit as  $\epsilon \rightarrow 0$  to exchange with the infinite sum. This completes the proof.  $\square$

#### A4 Proof of Theorem 2

*Proof of Theorem 2.* To prove the theorem, the expectation is conditioned on the number of events  $\mathcal{N}^\epsilon(T)$ , as was done in the previous section. Thus the expectation is

$$435 \quad E[\langle \sigma^\epsilon(t) - \sigma(t), \phi(t) \rangle^2] \quad (\text{A40})$$

$$= \sum_{N=1}^{\infty} E \left[ \left( \sum_{i=1}^N \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} \sigma^\epsilon(t) \phi(t) dt - \int_0^T b\delta(t - \mathcal{T}_i) \phi(t) dt \right. \right. \\ \left. \left. - \sum_{i=N+1}^{\mathcal{N}(T)} \int_0^T b\delta(t - \mathcal{T}_i) \phi(t) dt \right)^2 \middle| \mathcal{N}^\epsilon(T) = N \right] P(\mathcal{N}^\epsilon(T) = N)$$

where  $\mathcal{N}(T)$  is the number of dry events for the  $\sigma(t)$  process up to time  $T$ . Again, because of the decay of  $P(\mathcal{N}^\epsilon(T) = N)$  as  $N \rightarrow \infty$  given in Lemma 1, the infinite sum converges.

To estimate the quantity in (A40), one rain event is considered and the Cauchy-Schwarz bound will be used. Consider the

440  $i$ th rain event,

$$\begin{aligned}
& \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} \sigma^\epsilon(t) \phi(t) dt - b\phi(\mathcal{T}_i) \\
&= \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} \frac{r}{\epsilon} \phi(t) - \frac{r}{\epsilon} \phi(\mathcal{T}_{2i-1}^\epsilon) + \frac{r}{\epsilon} \phi(\mathcal{T}_{2i-1}^\epsilon) dt + b\phi(\mathcal{T}_{2i-1}^\epsilon) - b\phi(\mathcal{T}_{2i-1}^\epsilon) - b\phi(\mathcal{T}_i) \\
&= \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} \frac{r}{\epsilon} (\phi(t) - \phi(\mathcal{T}_{2i-1}^\epsilon)) dt + \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} - b \right) \phi(\mathcal{T}_{2i-1}^\epsilon) + b(\phi(\mathcal{T}_{2i-1}^\epsilon) - \phi(\mathcal{T}_i)).
\end{aligned} \tag{A41}$$

The function  $\phi(t)$  is smooth on  $[0, T]$  and thus is locally Lipschitz. Let the Lipschitz constant be  $K > 0$ . Then, along with the  
445 triangle inequality,

$$\left| \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} \sigma^\epsilon(t) \phi(t) dt - b\phi(\mathcal{T}_i) \right| \tag{A42}$$

$$\leq \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} \frac{r}{\epsilon} K |t - \mathcal{T}_{2i-1}^\epsilon| dt + \left| \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} - b \right) \phi(\mathcal{T}_{2i-1}^\epsilon) \right| + |\phi(\mathcal{T}_{2i-1}^\epsilon) - \phi(\mathcal{T}_i)| \tag{A43}$$

$$\leq \frac{r}{\epsilon} K |\tau_i^{r,\epsilon}|^2 + \left| \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} - b \right) \phi(\mathcal{T}_{2i-1}^\epsilon) \right| + |\phi(\mathcal{T}_{2i-1}^\epsilon) - \phi(\mathcal{T}_i)|, \tag{A44}$$

where the last inequality results from  $t - \mathcal{T}_{2i-1}^\epsilon$  being an increasing function on  $[\mathcal{T}_{2i-1}^\epsilon, \mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}]$ . Using the inequality  
450 above, along with the Cauchy-Schwarz inequality, the quantity in (A40) is bounded by

$$\begin{aligned}
& \sum_{N=1}^{\infty} E \left[ \left( \sum_{i=1}^N \int_{\mathcal{T}_{2i-1}^\epsilon}^{\mathcal{T}_{2i-1}^\epsilon + \tau_i^{r,\epsilon}} \sigma^\epsilon(t) \phi(t) dt - \int_0^T b\delta(t - \mathcal{T}_i) \phi(t) dt \right. \right. \\
& \quad \left. \left. - \sum_{i=N+1}^{\mathcal{N}(T)} \int_0^T b\delta(t - \mathcal{T}_i) \phi(t) dt \right)^2 \middle| \mathcal{N}^\epsilon(T) = N \right] P(\mathcal{N}^\epsilon(T) = N) \\
& \leq \sum_{N=1}^{\infty} \sum_{i=1}^N \left( \left( \frac{r}{\epsilon} \right)^2 K^2 E[|\tau_i^{r,\epsilon}|^4] + E \left[ \left| \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} - b \right) \phi(\mathcal{T}_{2i-1}^\epsilon) \right|^2 \right] + E[|\phi(\mathcal{T}_{2i-1}^\epsilon) - \phi(\mathcal{T}_i)|^2] \right) P(\mathcal{N}^\epsilon(T) = N) \\
& \quad + \sum_{N=1}^{\infty} E \left[ \left( \sum_{i=N+1}^{\mathcal{N}(T)} \int_0^T b\delta(t - \mathcal{T}_i) \phi(t) dt \right)^2 \middle| \mathcal{N}^\epsilon(T) = N \right],
\end{aligned} \tag{A45}$$

455 where all expectations are conditional on  $\mathcal{N}^\epsilon(T) = N$ .

To finish the theorem the following moments of  $\tau_i^{r,\epsilon}$  are used

$$E[\tau_i^{r,\epsilon}] = \frac{b\epsilon}{r}, \quad E[|\tau_i^{r,\epsilon}|^2] = \frac{bD^2\epsilon^3}{r^3} + \frac{b^2\epsilon^2}{r^2}, \quad E[|\tau_i^{r,\epsilon}|^4] = \frac{b^4\epsilon^4}{r^4} + 6\frac{b^3D_1^2\epsilon^5}{r^5} + 15\frac{b^2D_1^4\epsilon^6}{r^6} + 15\frac{bD_1^6\epsilon^7}{r^7}. \tag{A46}$$

Thus the first term in (A45) is

$$\left(\frac{r}{\epsilon}\right)^2 E[|\tau_i^{r,\epsilon}|^4] = O(\epsilon^2). \quad (\text{A47})$$

460 The second term in (A45) is

$$E \left[ \left| \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} - b \right) \phi(\mathcal{T}_{2i-1}^\epsilon) \right|^2 \right] = E \left[ \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} - b \right)^2 \right] E [\phi(\mathcal{T}_{2i-1}^\epsilon)^2] \quad (\text{A48})$$

$$= E \left[ \left( \left( \frac{r}{\epsilon} \tau_i^{r,\epsilon} \right)^2 - 2b \frac{r}{\epsilon} \tau_i^{r,\epsilon} + b^2 \right) \right] E [\phi(\mathcal{T}_{2i-1}^\epsilon)^2] \quad (\text{A49})$$

$$= O(\epsilon) \quad (\text{A50})$$

where the expectation turns into a product because  $\tau_i^{r,\epsilon}$  and  $\mathcal{T}_{2i-1}^\epsilon$  are independent. For the third term of (A45), the Lipschitz  
465 condition is used to write

$$E [|\phi(\mathcal{T}_{2i-1}^\epsilon) - \phi(\mathcal{T}_i)|^2] \leq K^2 E [|\mathcal{T}_{2i-1}^\epsilon - \mathcal{T}_i|^2]. \quad (\text{A51})$$

Note that the stopping times can be written in terms of the moistening processes in the following way:

$$\mathcal{T}_{2i-1}^\epsilon = \frac{1}{m} \int_0^{\mathcal{T}_{2i-1}^\epsilon} m \, dt, \quad (\text{A52})$$

$$= \frac{1}{m} \sum_{j=1}^{2i-1} \int_{\mathcal{T}_{j-1}^\epsilon}^{\mathcal{T}_j^\epsilon} m \, dt \quad (\text{A53})$$

$$470 \quad = \frac{1}{m} \underbrace{\sum_{j=1}^i \int_{\mathcal{T}_{2j-2}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon} m \, dt}_{\sigma^\epsilon=0} + \sum_{j=1}^i \underbrace{\int_{\mathcal{T}_{2j-1}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon + \tau_i^{r,\epsilon}} 1 \, dt}_{\sigma^\epsilon=1} \quad (\text{A54})$$

$$= \frac{1}{m} \left( \sum_{j=1}^i \int_{\mathcal{T}_{2j-2}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon} m \, dt + \int_{\mathcal{T}_{2j-2}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon} D_0 \, dW_t \right) - \int_0^{\mathcal{T}_{2i-1}^\epsilon} \frac{D_0}{m} \, dW_t$$

$$+ \sum_{j=1}^i \left( \int_{\mathcal{T}_{2j-1}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon + \tau_i^{r,\epsilon}} 1 \, dt + \int_{\mathcal{T}_{2j-1}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon + \tau_i^{r,\epsilon}} \frac{D_0}{m} \, dW_t \right) \quad (\text{A55})$$

$$= \frac{1}{m} E_{\mathcal{T}_{2i-1}^\epsilon}^\epsilon + \sum_{j=1}^i \tau_j^{r,\epsilon} + \sum_{j=1}^i \int_{\mathcal{T}_{2j-1}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon + \tau_i^{r,\epsilon}} \frac{D_0}{m} \, dW_t - \int_0^{\mathcal{T}_{2i-1}^\epsilon} \frac{D_0}{m} \, dW_t, \quad (\text{A56})$$

where  $\mathcal{T}_0^\epsilon \equiv 0$ . Similarly

$$475 \quad \mathcal{T}_i = \frac{1}{m} \int_0^{\mathcal{T}_i} m dt \tag{A57}$$

$$= \frac{1}{m} \left( \int_0^{\mathcal{T}_i} m dt + \int_0^{\mathcal{T}_i} D_0 dW_t \right) - \int_0^{\mathcal{T}_i} \frac{D_0}{m} dW_t \tag{A58}$$

$$= \frac{1}{m} E_{\mathcal{T}_i} - \int_0^{\mathcal{T}_i} \frac{D_0}{m} dW_t, \tag{A59}$$

where the Wiener process is the same realization as in (A56). The definition of the stopping times  $\mathcal{T}_{2i-1}$  and  $\mathcal{T}_i$  imply

$$E_{\mathcal{T}_{2i-1}^\epsilon} = E_{\mathcal{T}_i} = bi.$$

Thus the difference in stopping times is

$$|\mathcal{T}_{2i-1}^\epsilon - \mathcal{T}_i|^2 = \left| \frac{1}{m} (E_{\mathcal{T}_{2i-1}^\epsilon} - E_{\mathcal{T}_i}) - \int_{\mathcal{T}_i}^{\mathcal{T}_{2i-1}^\epsilon} \frac{D_0}{m} dW_t + \sum_{j=1}^i \tau_j^{r,\epsilon} + \sum_{j=1}^i \int_{\mathcal{T}_{2j-1}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon + \tau_j^{r,\epsilon}} \frac{D_0}{m} dW_t \right|^2 \tag{A60}$$

$$480 \quad \leq \left| \int_{\mathcal{T}_i}^{\mathcal{T}_{2i-1}^\epsilon} \frac{D_0}{m} dW_t \right|^2 + \sum_{j=1}^i |\tau_j^{r,\epsilon}|^2 + \sum_{j=1}^i \left| \int_{\mathcal{T}_{2j-1}^\epsilon}^{\mathcal{T}_{2j-1}^\epsilon + \tau_j^{r,\epsilon}} \frac{D_0}{m} dW_t \right|^2. \tag{A61}$$

where the triangle inequality has been used. Taking the expected value and using the Itô isometry yields

$$E[|\mathcal{T}_{2i-1}^\epsilon - \mathcal{T}_i|^2] \leq \frac{D_0^2}{m^2} E[\mathcal{T}_{2i-1}^\epsilon - \mathcal{T}_i] + \sum_{j=1}^i E[|\tau_j^{r,\epsilon}|^2] + \frac{D_0^2}{m^2} \sum_{j=1}^i E[\tau_j^{r,\epsilon}] \tag{A62}$$

$$= \frac{D_0^2}{m^2} E \left[ \sum_{j=1}^i \tau_j^{d,\epsilon} + \tau_j^{r,\epsilon} - \tau_j^d \right] + \sum_{j=1}^i E[|\tau_j^{r,\epsilon}|^2] + \frac{D_0^2}{m^2} \sum_{j=1}^i E[\tau_j^{r,\epsilon}] \tag{A63}$$

Note that  $\tau_i^{d,\epsilon}$  and  $\tau_i^d$  are IID random variables with the same distribution and thus the expectations cancel. For the remaining  
485 terms, the moments of  $\tau_j^{r,\epsilon}$  in equation (A46) are used to give

$$K^2 E[|\mathcal{T}_{2i-1}^\epsilon - \mathcal{T}_i|^2] \leq K^2 \left( i \frac{D_0^2}{m^2} E[\tau^{r,\epsilon}] + i \frac{D_0^2}{m^2} E[\tau^{r,\epsilon}] + i E[|\tau^{r,\epsilon}|^2] \right) = O(\epsilon), \tag{A64}$$

which completes the consideration of the third term of (A45).

For the last “remainder” term in (A45), the expectation is conditioned on both  $\mathcal{N}(T)$  and  $\mathcal{N}^\epsilon(T)$ . That is,

$$\sum_{N=1}^{\infty} CE \left[ \left( \sum_{i=N+1}^{\mathcal{N}(T)} \int_0^T b\delta(t - \mathcal{T}_i)\phi(t) dt \right)^2 \middle| \mathcal{N}^\epsilon(T) = N \right] P(\mathcal{N}^\epsilon(T) = N) \quad (\text{A65})$$

$$490 \quad = CE \left[ \left( \sum_{i=\mathcal{N}^\epsilon(T)+1}^{\mathcal{N}(T)} \int_0^T b\delta(t - \mathcal{T}_i)\phi(t) dt \right)^2 \right] \quad (\text{A66})$$

$$= \sum_{N=1}^{\infty} \sum_{M=1}^{\infty} CE \left[ \left( \sum_{i=N+1}^{N+M} \int_0^T b\delta(t - \mathcal{T}_i)\phi(t) dt \right)^2 \middle| \mathcal{N}^\epsilon(T) = N, \mathcal{N}(T) = N + M \right] \cdot P(\mathcal{N}^\epsilon(T) = N, \mathcal{N}(T) = N + M) \quad (\text{A67})$$

If  $\mathcal{N}^\epsilon(T) \geq \mathcal{N}(T)$ , then there is no sum and the term is zero. If  $\mathcal{N}^\epsilon(T) \neq \mathcal{N}(T)$ , then the processes  $E_t^\epsilon$  and  $E_t$  from Section 3.3 must be at least  $b$  units apart. Thus by theorem 1,

$$495 \quad P(\mathcal{N}^\epsilon(T) \neq \mathcal{N}(T)) = P(|E^\epsilon(t) - E(t)| > b). \quad (\text{A68})$$

Furthermore, convergence in expectation ( $L^2$ ) implies convergence in probability. Therefore,

$$\lim_{\epsilon \rightarrow 0} P(|E^\epsilon(t) - E(t)| > b) = 0. \quad (\text{A69})$$

Putting this together with the above estimate yields

$$\lim_{\epsilon \rightarrow 0} \sum_{N=1}^{\infty} CE \left[ \left( \sum_{i=N+1}^{\mathcal{N}(T)} \int_0^T b\delta(t - \mathcal{T}_i)\phi(t) dt \right)^2 \middle| \mathcal{N}^\epsilon(T) = N \right] P(\mathcal{N}^\epsilon(T) = N) \quad (\text{A70})$$

$$500 \quad \leq \lim_{\epsilon \rightarrow 0} \sum_{N=1}^{\infty} \sum_{M=1}^{\infty} CE \left[ \left( \sum_{i=N+1}^{N+M} \int_0^T b\delta(t - \mathcal{T}_i)\phi(t) dt \right)^2 \middle| \mathcal{N}^\epsilon(T) = N, \mathcal{N}(T) = N + M \right] P(|E^\epsilon(t) - E(t)| > b) \quad (\text{A71})$$

$$= 0, \quad (\text{A72})$$

by using Tonelli’s theorem to exchange the sums and limit. Thus all of the terms in (A45) have been shown to converge to 0 as  $\epsilon \rightarrow 0$ , so that, returning to (A40) and taking the limit, we have

$$\lim_{\epsilon \rightarrow 0} E[\langle \sigma^\epsilon(t) - \sigma(t), \phi(t) \rangle^2] = 0 \quad (\text{A73})$$

505 and the proof is completed.  $\square$

*Author contributions.* Both the authors contributed to the final draft of the work. Additionally, SH contributed to the formal analysis and writing the original draft preparation, SNS contributed to conceptualization and writing – review & editing.



*Competing interests.* There are no competing interests present for either author.

*Acknowledgements.* The research of author SH is partially supported by the National Science Foundation under Grant DMS-1815061.

## 510 References

- Abbott, T. H., Stechmann, S. N., and Neelin, J. D.: Long temporal autocorrelations in tropical precipitation data and spike train prototypes, *Geophysical Research Letters*, 43, 11–472, 2016.
- Ahmed, F. and Neelin, J. D.: Explaining scales and statistics of tropical precipitation clusters with a stochastic model, *Journal of the Atmospheric Sciences*, 76, 3063–3087, 2019.
- 515 Albano, G., Giorno, V., Nobile, A. G., and Ricciardi, L. M.: Modeling refractoriness for stochastically driven single neurons, *Scientiae Mathematicae Japonicae*, 67, 173–190, 2008.
- Bender, C. M. and Orszag, S. A.: *Advanced mathematical methods for scientists and engineers I: Asymptotic methods and perturbation theory*, Springer Science & Business Media, 2013.
- Bhat, V. N.: Renewal approximations of the switched Poisson processes and their applications to queueing systems, *Journal of the Operational*  
520 *Research Society*, 45, 345–353, 1994.
- Cowpertwait, P., O’Connell, P., Metcalfe, A., and Mawdsley, J.: Stochastic point process modelling of rainfall. I. Single-site fitting and validation, *Journal of Hydrology*, 175, 17–46, 1996.
- Cox, D. R.: *Renewal theory*, Methuen, London, 1962.
- Deluca, A. and Corral, Á.: Scale invariant events and dry spells for medium-resolution local rain data, *Nonlin. Processes Geophys.*, 21,  
525 555–567, 2014.
- Deluca, A., Moloney, N. R., and Corral, Á.: Data-driven prediction of thresholded time series of rainfall and self-organized criticality models, *Physical review E*, 91, 052 808, 2015.
- Evans, M. R. and Majumdar, S. N.: Diffusion with stochastic resetting, *Physical review letters*, 106, 160 601, 2011.
- Evans, M. R., Majumdar, S. N., and Schehr, G.: Stochastic resetting and applications, *Journal of Physics A: Mathematical and Theoretical*,  
530 53, 193 001, 2020.
- Filippov, A. F.: *Differential equations with discontinuous righthand sides: control systems*, vol. 18, Springer Science & Business Media, 2013.
- Foufoula-Georgiou, E. and Lettenmaier, D. P.: A Markov renewal model for rainfall occurrences, *Water Resour. Res.*, 23, 875–884, 1987.
- Frierson, D. M. W., Majda, A. J., and Pauluis, O. M.: Large scale dynamics of precipitation fronts in the tropical atmosphere: a novel  
535 relaxation limit, *Commun. Math. Sci.*, 2, 591–626, 2004.
- Gardiner, C. W.: *Handbook of stochastic methods: for physics, chemistry & the natural sciences*, vol. 13 of *Springer Series in Synergetics*, Springer–Verlag, Berlin, 2004.
- Green, J. R.: A model for rainfall occurrence, *J. Roy. Statist. Soc. Ser. B*, 26, 345–353, 1964.
- Hernandez-Duenas, G., Smith, L. M., and Stechmann, S. N.: Weak-and strong-friction limits of parcel models: Comparisons and stochastic  
540 convective initiation time, *Q. J. Roy. Met. Soc.*, 145, 2272–2291, <https://doi.org/10.1002/qj.3557>, 2019.
- Hoeffding, W.: Probability inequalities for sums of bounded random variables, in: *The Collected Works of Wassily Hoeffding*, pp. 409–426, Springer, 1994.
- Holloway, C. E. and Neelin, J. D.: Temporal relations of column water vapor and tropical precipitation, *J. Atmos. Sci.*, 67, 1091–1105, 2010.
- Hottovy, S. A. and Stechmann, S. N.: A spatiotemporal stochastic model for tropical precipitation and water vapor dynamics, *J. Atmos. Sci.*,  
545 72, 4721–4738, <https://doi.org/10.1175/JAS-D-15-0119.1>, 2015a.

- Hottovy, S. A. and Stechmann, S. N.: Threshold models for rainfall and convection: Deterministic versus stochastic triggers, *SIAM J. Appl. Math.*, 75, 861–884, <https://doi.org/10.1137/140980788>, 2015b.
- Huang, T., Stechmann, S. N., and Torchinsky, J. L.: Framework for idealized climate simulations with spatiotemporal stochastic clouds and planetary-scale circulations, *Physical Review Fluids*, 7, 010 502, 2022.
- 550 Jakubowski, A. et al.: A non-Skorohod topology on the Skorohod space, *Electronic journal of probability*, 2, 1997.
- Katz, R. W.: Precipitation as a chain-dependent process, *Journal of Applied Meteorology*, 16, 671–676, 1977.
- Kelley, J. L.: *General topology*, Courier Dover Publications, 2017.
- Khouider, B. and Majda, A. J.: A non-oscillatory balanced scheme for an idealized tropical climate model: Part I: Algorithm and validation, *Theor. Comp. Fluid Dyn.*, 19, 331–354, 2005.
- 555 Khouider, B., Biello, J. A., and Majda, A. J.: A stochastic multicloud model for tropical convection, *Comm. Math. Sci.*, 8, 187–216, 2010.
- Kurtz, T. G.: Random time changes and convergence in distribution under the Meyer-Zheng conditions, *The Annals of probability*, pp. 1010–1034, 1991.
- Lejay, A. and Pigato, P.: A threshold model for local volatility: evidence of leverage and mean reversion effects on historical data, *International Journal of Theoretical and Applied Finance*, 22, 1950 017, 2019.
- 560 Lin, J. and Neelin, J.: Influence of a stochastic moist convective parameterization on tropical climate variability, *Geophys. Res. Lett.*, 27, 3691–3694, <https://doi.org/10.1029/2000GL011964>, 2000.
- Mueller, E. A. and Stechmann, S. N.: Shallow-cloud impact on climate and uncertainty: A simple stochastic model, *Mathematics of Climate and Weather Forecasting*, 6, 16–37, 2020.
- Neelin, J. D., Sahany, S., Stechmann, S. N., and Bernstein, D. N.: Global warming precipitation accumulation increases above the current-c  
565 limate cutoff scale, *Proc. Natl. Acad. Sci. USA*, 114, 1258–1263, <https://doi.org/10.1073/pnas.1615333114>, 2017.
- Peters, O. and Neelin, J. D.: Critical phenomena in atmospheric precipitation, *Nature Physics*, 2, 393–396, 2006.
- Peters, O., Deluca, A., Corral, A., Neelin, J. D., and Holloway, C. E.: Universality of rain event size distributions, *J. Stat. Mech.*, 2010, P11 030, 2010.
- Richardson, C. W.: Stochastic simulation of daily precipitation, temperature, and solar radiation, *Water resources research*, 17, 182–190,  
570 1981.
- Rodriguez-Iturbe, I., Cox, D. R., and Isham, V.: A point process model for rainfall: further developments, *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 417, 283–298, 1988.
- Sacerdote, L. and Giraud, M. T.: Stochastic integrate and fire models: a review on mathematical methods and their applications, in: *Stochastic biomathematical models*, pp. 99–148, Springer, 2013.
- 575 Simpson, D. and Kuske, R.: Stochastically perturbed sliding motion in piecewise-smooth systems, *Discrete & Continuous Dynamical Systems-B*, 19, 2889, 2014.
- Smith, J. A. and Karr, A. F.: A point process model of summer season rainfall occurrences, *Water Resources Research*, 19, 95–103, 1983.
- Stechmann, S. N. and Hottovy, S.: Cloud regimes as phase transitions, *Geophysical Research Letters*, 43, 6579–6587, 2016.
- Stechmann, S. N. and Neelin, J. D.: A stochastic model for the transition to strong convection, *J. Atmos. Sci.*, 68, 2955–2970, 2011.
- 580 Stechmann, S. N. and Neelin, J. D.: First-passage-time prototypes for precipitation statistics, *J. Atmos. Sci.*, 71, 3269–3291, 2014.
- Whitt, W.: Approximating a point process by a renewal process, I: Two basic methods, *Operations Research*, 30, 125–147, 1982.
- Wilks, D. S. and Wilby, R. L.: The weather generation game: a review of stochastic weather models, *Progress in physical geography*, 23, 329–357, 1999.