



# Seasonal Controls on Isolated Convective Storm Drafts, Precipitation Intensity, and Life Cycle As Observed During GoAmazon2014/5

Scott E. Giangrande<sup>1</sup>, Thiago Biscaro<sup>2</sup>, John M. Peters<sup>3</sup>

<sup>1</sup>Environmental and Climate Sciences Department, Brookhaven National Laboratory, Upton, NY, US

5 <sup>2</sup>Meteorological Satellites and Sensors Division, National Institute for Space Research, Cachoeira Paulista, São Paulo, 12630000, Brazil

<sup>3</sup>Department of Meteorology and Atmospheric Science, The Pennsylvania State University, University Park, PA, US

*Correspondence to:* Scott E. Giangrande (sgrande@bnl.gov)

10 **Abstract.** Isolated deep convective cloud life cycle and seasonal changes in storm properties are observed for daytime events during the DOE-ARM GoAmazon2014/5 campaign to understand controls on storm behavior. Storm life cycles are documented using surveillance radar from initiation through maturity and dissipation. Vertical air velocity estimates are obtained from radar wind profiler overpasses, with the storm environment informed by radiosondes.

15 Dry season storm conditions favored reduced morning shallow cloud coverage and larger low level convective available potential energy (CAPE) than wet season counterparts. The typical dry season storm reached its peak intensity and size earlier in its life cycle compared to wet season cells. These cells exhibited updrafts in core precipitation regions ( $Z > 35$  dBZ) to above the melting level, and persistent downdrafts aloft within precipitation adjacent to their cores. Moreover, dry season cells recorded more intense updrafts to earlier life cycle stages, and a higher incidence of strong updrafts (i.e.,  $> 5 \text{ m s}^{-1}$ ) at low levels. In contrast, wet season storms were longer-lived and featured a higher incidence of moderate (i.e.,  $2\text{-}5 \text{ m s}^{-1}$ ) updrafts aloft. These storms also favored a shift in their most intense properties to later life cycle stages. Strong downdrafts were far less frequent within wet season cells aloft, indicating a potential systematic difference in downdraft behaviors between the seasons. Results from a stochastic parcel model suggest that dry season cells may expect stronger updrafts at low levels because of larger low level CAPE in the dry season. Wet season cells anticipate strong updrafts aloft because of larger free-tropospheric relative humidity and reduced entrainment-driven dilution. The enhanced dry season downdrafts are attributed to 25 increased evaporation, dry air entrainment-mixing, and negative buoyancy in regions adjacent to sampled dry season cores.

## 1 Introduction

Deep Convective Clouds (DCC) play a critical role in regulating the global energy cycle through their extensive cloud coverage and the exchange of latent heat. DCCs are a primary focus of weather and climate model improvement because of their critical role in the global climate system. It is crucial to understand how these storms will respond to potential climate change, due to the high socioeconomic impacts associated with severe weather, heavy rainfall, and lightning (e.g., Trapp et 30



al., 2009; Diffenbaugh et al., 2013; Sillman et al., 2013; Seeley and Romps 2015; Feng et al., 2016; Prein et al., 2017). Nevertheless, gaps remain in our understanding of the factors that regulate DCC size, frequency, and updraft or precipitation intensity. These gaps are partially attributed to a lack of DCC dynamical and microphysical observations, a problem tied to the difficulty in sampling storms that have intense vertical velocities, long and complex life cycles, and are organized on scales  
35 larger than individual updrafts.

As home to frequent convective clouds, the Amazon basin has been at the forefront of impactful DCC studies (e.g., Williams et al., 2002; Andreae et al., 2004; Koren et al., 2008; Rosenfeld et al., 2008; Wang et al., 2016; Fan et al., 2018). The Amazon forest is the largest tropical rain forest on the planet, and this setting promotes diverse clouds influenced by a range of land surface and aerosol conditions that vary according to seasonal regimes, and having behaviors that span tropical, oceanic  
40 and continental characteristics. Amazon cloud conditions are interconnected to shifts in the synoptic-scale thermodynamic conditions and coupled local-scale feedbacks (e.g., Fu et al., 1999; Machado et al., 2004; Li and Fu, 2004; Misra, 2008), which is a significant challenge to climate modeling (e.g., Richter and Xie, 2008; Nobre et al., 2009; Yin et al., 2013). Given this important global setting, multi-agency campaigns have long-targeted this region for DCC studies (e.g., Williams et al., 2002; Petersen et al., 2002; Machado et al., 2014; 2018; Adams et al., 2013; 2017; Martin et al., 2017).

Our focus is on isolated diurnal DCCs that are ubiquitous to the humid Amazon basin, where low wind shear promotes short-lived and slow-moving storms. In these settings, DCCs often span their entire life cycle under the umbrella of a single surveillance radar O[300km]. Cloud regimes in the Amazon are commonly divided into two seasons, the “wet season,” and “dry season.” There are distinct meteorological differences between these environments, including shifts in the convective available potential energy (CAPE) calculated over different depths, and changes in free tropospheric relative humidity (e.g.,  
50 Giangrande et al., 2020). Hence, these conditions may provide a natural laboratory for assessing the impact of bulk environmental shifts on convective cloud characteristics. Identifying and explaining these differences is a primary objective of the present article.

To accomplish this objective, we employ radar cell tracking concepts as have been well-established with recent emphasis on larger, longer-lived cells and mesoscale convective system (MCS) studies (e.g., Maddox 1980; Williams and  
55 Houze 1987; Rosenfeld 1987; Dixon and Wiener 1993; Machado et al., 1998; Göke et al., 2007; Feng et al., 2012; Borque et al., 2014; Limpert et al., 2015; Fridlind et al., 2019; Feng et al., 2019; Hu et al., 2019; Tian et al., 2022). This study draws from a unique Amazon dataset collected during the 2-year US Department of Energy (DOE) ARM Observations and Modeling of the Green Ocean Amazon campaign (GoAmazon2014/5; Martin et al., 2017; Giangrande et al., 2017) that featured surveillance radar coupled with atmospheric profiling capabilities. Although previous Amazon studies have documented  
60 seasonal-composite cloud properties (e.g., Machado et al., 2004; Ghate and Kollias 2016; Giangrande et al., 2016; Biscaro et al., 2021; Tian et al., 2021), few adopt a cell life cycle viewpoint as enabled by radar cell tracking. A unique aspect to this study is its emphasis on a set of radar-tracked cells that overpass the ARM profiling equipment, yielding direct observations of vertical hydrometeor, and by proxy, air motions. This coupled use of profiling-based vertical air velocity information builds on recent Amazon studies that have been integral to the understanding of DCC dynamics (e.g., Cifelli et al., 2002; Anderson



65 et al., 2005; Giangrande et al., 2016; Wang et al., 2019; 2020). Our approach advances on previous observational works (e.g., Byers and Braham, 1948; LeMone and Zipser, 1980; May and Rajopadhyaya, 1999; Giangrande et al., 2013; Kumar et al., 2015; Schiro et al., 2018; Wang et al., 2020) by analyzing the evolution of draft properties throughout the DCC life cycle.

## 2. Dataset and Methods

70 The data for this study were collected during the GoAmazon2014/5 field campaign that deployed from January 2014 to December 2015. The main site for the deployment was in the city of Manacapuru (Herein “T3”; 3.213S, 60.598W), at a distance 70 km west of Manaus, Brazil. The datasets were collected by the U.S. Department of Energy Atmospheric Radiation Measurement (ARM; Ackerman and Stokes, 2003; Mather and Voyles, 2013) Mobile Facility 1 (AMF1; Miller et al., 2016). The GoAmazon2014/5 AMF1 details, including cumulative campaign instrument summaries and other larger-scale regime  
75 breakdowns, are found in Giangrande et al. (2017; 2020). In addition to in situ datasets obtained by the AMF1 at T3, this study uses data collected by the nearby Manaus CENSIPAM (Amazonian Protection System) weather radar (Herein “SIPAM”; Saraiva et al., 2016). These radar data serve as the input for a cell tracking algorithm that documents storms that overpassed the site. All events require daytime convective initiation that follows an available morning radiosonde (Sect. 2.1.2). We adopt a definition of isolated cells that requires the SIPAM was able to track a longer-lived (> 50 minute) DCC in its coverage  
80 domain without an obvious split/merger (tracking criteria, Sect. 2.2.2). A final requirement was that cells overpass profiling instrumentation at the ARM T3 location (e.g., Sect. 2.1.1, 2.1.2). A listing of the events is located in Tables 1 and 2.

### 2.1 ARM AMF1 Datasets

#### 2.1.1 Radar Wind Profiler and Vertical Air Motion Retrievals

Vertical air velocity profiles were retrieved from a 1290 MHz ARM Radar Wind Profiler (RWP) located at the T3  
85 site. During GoAmazon2014/5, the RWP operated in a precipitation mode (200m gate spacing, 10 deg beamwidth) wherein time-height (at approx. 6s update) collections were interwoven with boundary layer wind modes (e.g., Tridon et al., 2013). These precipitation modes collect radar moments for the signal-to-noise ratio SNR and mean Doppler velocity (O[1 km] horizontal resolution at 6 km altitude). Reflectivity factor was estimated from the SNR and calibrated (within 1-2 dBZ) using a collocated disdrometer (e.g., Wang et al., 2018).

90 The vertical air velocity is retrieved following Giangrande et al. (2013, 2016) and recent Amazon studies (e.g., Wang et al., 2019; 2020). The techniques assume the vertical air motion is the difference between the mean Doppler velocity and a hydrometeor fall speed (estimated, as a function of  $Z$ ). For retrievals at the native RWP resolutions, the approach is typically accurate within  $O[1-2 \text{ ms}^{-1}]$  in convective cores (e.g., Heymsfield et al., 2010). Fall speed assumptions use a power-law relationship of the form  $V_f = aZ^b$ , where  $Z$  is the reflectivity factor in linear  $[\text{mm}^6/\text{m}^3]$  units. Fall speeds are subsequently  
95 corrected for changes in air density aloft (e.g., Foote and Du Toit, 1969).



For this study, we adopt a fall speed correction that follows results found in Giangrande et al. (2016). Specifically, Amazon convection was suggested as favoring higher-density graupel or frozen drops above the melting level, associated with faster fall speeds closer to that of rain than lower density ice hydrometeors (i.e.,  $V_f = aZ^b$ , where  $a = 2.7$ , and  $b = 0.1$ ). Our approach was to extend rain coefficients to all hydrometeors above the melting level (approx. 5 km AGL) in DCC contexts.

100 This approach is consistent with previous RWP studies that routinely apply rain relationships in DCC cores having  $Z > 35$  dBZ where higher density hydrometeors are expected. Our simplification is in applying this fall speed assumption for retrievals to a wider range of adjacent isolated convective (reflectivity,  $25 < Z < 35$  dBZ) conditions, including in the vicinity of the melting level, where slower-falling lower-density graupel, ice, or aggregates are also not expected. One caveat is that this choice may overestimate fall speed corrections to regions aloft (i.e.,  $> 7-8$  km AGL) if the convective ice hydrometeors density

105 decreases in these contexts with altitude (e.g., Protat and Williams, 2011), and this may bias RWP retrievals at higher altitudes towards relative updrafts. Our results consider draft properties contingent on different  $Z$  thresholds (i.e.,  $Z > 25$  dBZ,  $Z > 35$  dBZ) to differentiate behaviors that may shift when using this simplified approach.

Velocity profiles are summarized using normalized velocity cumulative frequency with altitude displays (CFADs, Yuter and Houze 1995). CFADs are drawn from the nearest 5-minutes to the associated RWP storm overpass as viewed by

110 SIPAM radar and centered on the time of the highest RWP echo top height (ETH) for that overpass. This choice also minimizes individual events disproportionately contributing to our summary plots (i.e., cells may remain over the RWP for extended periods that include multiple radar volumes). We include only those retrievals associated with  $Z$  values exceeding the matching SIPAM cell tracking threshold (e.g.,  $> 25$  dBZ). These choices limit our analysis to precipitation regions for these events (e.g.,  $R > 0.5$  mm  $\text{hr}^{-1}$ ). CFAD velocity properties above 10 km are not included owing to RWP sampling limitations aloft (sampling

115 quantity, fall speed and resolution). Similarly, we require greater than 250 retrievals at a given altitude to include that altitude on summary CFADs. Finally, RWP retrieval interpretation is tied to the representativeness of narrow field of view / vertically-pointing observations (i.e., “chording”, Jorgensen et al., 1985; Borque et al., 2014). It is known that even fortuitous DCC samples may underestimate extremes owing to randomness and/or natural variability. Previous studies indicate that similar retrievals may exhibit expected low-biased updraft magnitudes exceeding 30% (e.g., Jorgensen et al., 1985; Wang et al., 2020).

120

### 2.1.2 AMF1 Radiosonde, Surface and Profiling Instruments

Events were associated with a clear 12 GMT (08 LT) radiosonde that preceded convective initiation time for a tracked cell that overpassed the T3 site. A clear radiosonde was defined as one without precipitation at the T3 location within 30 minutes of the launch. The was confirmed by checking the SIPAM radar for a lack of echoes in the vicinity of the T3 site. We

125 computed mean-layer convective available potential energy (MLCAPE) and convective inhibition (MLCIN) using radiosonde profiles by lifting an air parcel with the average properties of the lowest 1 km of the atmosphere adiabatically (with mixed phase between  $T = 273.15\text{K}$  and  $T = 263.15\text{K}$ ). This follows methods described in Peters et al. (2022).

Additional instruments were available to investigate the pre-convective storm environments to possibly identify discrepancies in the boundary layer and its evolution. This study draws from the ARM surface meteorology station at T3 for



130 temperature measurements, and daytime planetary boundary layer (PBL) height as estimated by a collocated ceilometer. Diurnal composites for the cloud cover at T3 (cloud frequency of occurrence) are estimated by the multi-sensor ARM W-band Cloud Radar (WACR) Active Remote Sensing of Clouds (ARSCL) value-added product (e.g., Clothiaux et al., 2000).

## 2.2 Surveillance Radar and Radar Cell Tracking

### 135 2.2.1 SIPAM Radar

The SIPAM S-Band (2.2 GHz) radar is a single polarization Doppler weather radar performing a volume scan with 17 elevations (lowest: 0.9 degree, highest: 19 degrees) every 12 minutes, with a 1.98° beamwidth and radial (gate) resolution of 500 m. The SIPAM is located in the city of Manaus and has a 240 km radius coverage area. Clutter corrected reflectivity factor data were gridded into a 2 x 2 km horizontal, 3 km level constant altitude plan position indicator (CAPPI). These CAPPIs  
140 serve as input for our tracking algorithm (Sect. 2.2.2), as well as associated life cycle characterization of the convective cells.

### 2.2.2 Tracking Method and Definitions

The tracking algorithm is based on an area overlap approach, following the ForTraCC methods described by Vila et al. (2008). The main improvement is that the timestep between two adjacent radar reflectivity factor CAPPIs is automatically  
145 detected in our current implementation, which allows for non-uniform radar timesteps. The algorithm works by comparing two successive radar CAPPI fields. A first step is to identify areas with contiguous reflectivity values above a certain threshold. We consider two thresholds, a 25 dBZ and a 35 dBZ threshold; these values are consistent with a light rain lower bound and one typical of a tropical “convective” radar threshold (Anagnostou et al., 2004; Steiner et al., 1995; Wang et al., 2018).

Cells are defined using gridded CAPPI pixel clusters, and clusters smaller than 10 pixels (40 km<sup>2</sup>) were excluded to  
150 avoid noise contamination. A subsequent step verifies which cluster has an area that overlaps with the previous radar field. If a cluster at a given time matches a cluster at the previous time (defined by a minimum 20% overlap area), the cluster is said to be the continuation of that cell, and repeating this process generates the trackable cell records. Once done, we sub-select all storms that overpassed the RWP T3 site. Events were sorted for overpasses associated with storms having life cycles > 50 minutes (i.e., a minimum of 5 SIPAM scans). The resulting set was sorted by season, with cells exhibiting split/merge  
155 characteristics in their tracked evolution removed. This process led to 24 event-cells identified during the Amazon wet season, and 19 for the dry season (Tables 1 and 2, respectively).

Overall, our typical wet season storm was longer-lived than its dry season counterpart, however there was modest overlap for most tracked cell behaviors. The mean lifetime for these cells was 131 minutes (standard deviation = 61, median = 120), with the mean wet season cell lasting 141 minutes (55, 132), and a mean dry season cell lifetime of 117 minutes (66,  
160 96). As these times are based on the  $Z > 25$  dBZ threshold, total cloud lifetimes will exceed those of radar precipitation echoes. Separately, the life cycle timings for  $Z > 35$  dBZ echoes were similar across seasons, with an approximate mean of 90 minutes and a standard deviation of 30 minutes. The average cell in our composites using the  $Z > 25$  dBZ threshold initiates at a time



of 13:42 LT (standard deviation approx. 2 hours), with the mean dry season storm initiating by 13:38 LT, and a mean wet season cell onset by 13:46 LT.

165

### 3. Composite Seasonal Thermodynamic and Diurnal Conditions

Amazon regimes are defined using calendar definitions (DJFMA for “wet”; JJAS for “dry”). Events radiosonde properties are not consistent with those of transitional environments that may promote more intense convective updrafts or storm electrification (e.g., Williams et al., 2002; Giangrande et al., 2016; 2020). Single column model forcing larger-scale tendencies (e.g., Tang et al., 2016) and/or reanalysis fields are not shown, but are consistent with seasonal environments reported in previous studies (e.g., Giangrande et al., 2020).

170

#### 3.1 Composite Thermodynamic Profiles and Event Convective Parameter Summaries

In Figure 1, we plot composite radiosondes for our events. Overall, the behaviors are similar to previous studies drawn from 12 UTC GoAmazon2014/5 radiosondes (e.g., Giangrande et al., 2020). The main shift between seasonal profiles is associated with the drier mid-to-upper levels observed for the dry season. Each composite indicates a low-level capping or remnant nocturnal temperature inversion that may act to inhibit daytime shallow cumulus and/or promote deeper cloud modes when convection initiates.

175

A breakdown of event convective parameters (see also, Tables 1 and 2) is as follows: Dry season low-level (0-6 km) MLCAPE values are larger than in wet season events, with a mean MLCAPE value =  $239 \text{ J kg}^{-1}$ , as compared to a mean wet season MLCAPE =  $152 \text{ J kg}^{-1}$ . This difference is statistically significant at the 95th confidence level based on a student’s T-test (herein, “significant”). The dry season profile MLCAPE values are also larger, MLCAPE =  $1506 \text{ J kg}^{-1}$  (dry) versus MLCAPE =  $1337 \text{ J kg}^{-1}$  (wet), however this difference is not statistically significant. Insignificant seasonal differences are found in low level wind shear (not shown) and MLCIN. As expected, mean relative humidity RH values in the lower free troposphere are significantly larger in the wet season (78%) than the dry season (56%).

180

#### 185 3.2 Composite Boundary Layer and Diurnal Cloud Development

In Figure 2, we plot composite diurnal cloud and boundary layer properties to inform on pre-DCC onset differences between wet and dry events. In the upper panels (Figure 2ab), we plot the ARSCL cloud frequency of occurrence for the event-hours around radiosonde launch through convective initiation (typically, prior to 14 LT). In the lower panels, we plot T3 surface temperature and the ceilometer-estimated PBL heights. Solid lines in Figure 2cd are the composite mean behaviors, while shaded regions are the event standard deviation.

190

Morning shallow to mid-level clouds are more common for our typical wet season event when compared with composite dry season cases. This observation is not surprising, and consistent with previous studies that infer higher humidity as a control for increased cloudiness. The reduction of dry season cloud cover is also consistent with a more rapid PBL height



increase that follows sunrise than in the wet season (Figure 2d), which (presumably) results from greater insolation in the dry  
195 season. The largest PBL height discrepancies build prior to 12 LT, preceding the transition to congestus or deeper cloud modes.  
This transition is also suggested by cloud radar profiling in having more frequent cloud radar echoes to higher altitudes (by 12  
LT for wet, slightly later for dry). In short, both seasonal composites indicate similar tendencies for the daytime shallower  
cloud mode (echo top heights < 2 km) shortly after 10 LT, with a transition towards deeper clouds (echo top height  $\geq$  4 km)  
after 12 LT. However, we observe an earlier presence of upper-level (anvil) cloud signatures coupled with an absence of mid-  
200 level clouds (Figure 2b) in the dry season, which suggests that a more rapid transition to deeper convection occurs in the dry  
season.

The PBL evolution in the dry season also suggests a more rapid onset to deeper convection. The physical arguments  
that support this include the higher morning MLCAPE (at similar or reduced MLCIN) coupled with building PBL instability  
during the pre-convective hours from an increase in incoming solar radiation (reduced cloud frequency, slightly higher surface  
205 temperature). While complete surface flux measurements were unavailable, the authors speculate dry season conditions may  
favor a higher Bowen ratio (i.e., reduced soil moisture, humidity) and stronger generation of turbulent boundary layer growth  
(leading to the observed higher PBL height). Note, while our subset of radar-tracked cells exhibited similar onset timing, wet  
season cells were longer-lived (using the  $Z > 25$  dBZ threshold). As we plot in Figure 3, echo statistics drawn from the larger  
SIPAM domain cell tracking populations from these events suggests our wet (solid lines) and dry (dashed lines) events  
210 typically favored a similar frequency of occurrence and diurnal timing for  $Z > 35$  dBZ convective “core” echoes. However,  
dry events exhibited an earlier and more frequent occurrence of intense convective echoes  $Z > 45$  dBZ, consistent with  
arguments for stronger dry season cells overall, and a more rapid deep transition and/or increased anvil cloud presence. In  
contrast, wet season events suggest earlier  $Z > 25$  dBZ populations associated with precipitating congestus, or extended  
congestus-to-deep cloud transitioning.

215

#### 4. Regime-based Amazon Storm Life Cycle, Precipitation and Draft Comparisons

This section presents composite radar-tracked storm properties, and discusses the potential connections between those  
characteristics and seasonal environmental controls. As DCC intensity and life cycle may be defined in several ways (i.e.,  
rainfall, updraft maximum), we compare storm life cycle properties as viewed by surveillance radar (precipitation quantities)  
220 to fortuitous profiler overpass observations (dynamics). To conclude the section, the results of a simple Amazon updraft model  
sensitivity test (Sect. 4.3, 4.4) are provided to lend possible physical explanation for observed draft differences.

##### 4.1 Time-varying Surveillance Radar Behaviors

In Figure 4, we plot seasonal life cycle composites for precipitation properties of our tracked storms. These depictions  
apply a normalized cell lifetime for compositing purposes, where 0 represents the echo onset time, and 1 represents the final  
225 time a qualifying echo was observed. Expectedly, cell areas associated with the lower  $Z > 25$  dBZ threshold (Figure 4a) convey



larger and longer-lived precipitation tracks that encompass those associated with the more stringent  $Z > 35$  dBZ threshold (Figure 4b). Most tracked cells (19 for wet, 12 for dry) within the lower threshold set are contained within the higher threshold tracking set (i.e., exceeding 10 pixels for multiple scans exceeding 50 minutes). Nevertheless, each event recorded a maximum  $Z > 35$  dBZ at multiple points during its evolution. In Figure 4cd, we plot the mean  $Z$  associated with the tracked cells, and in  
230 Figure 4e we plot the corresponding maximum  $Z$  composite properties for those cells. The distribution of the cell overpass times relative to the normalized life cycle stage is found in Figure 4f.

The plot of seasonal cell properties for precipitation area coverage is found in Figure 4ab. Initially, composite cell properties with  $Z > 25$  dBZ (Figure 4a) display similar increases in coverage throughout earlier stages (normalized time  $< 0.4$ ). However, dry season cells typically remain similar-sized in light rain / periphery area coverage for times  $> 0.4$ , whereas wet  
235 season cells continue to increase in such coverage until a peak at normalized time  $\cong 0.7$ . Adopting a more stringent  $Z > 35$  dBZ convective echo threshold (Figure 4b), composite area properties are more consistent across the seasons, albeit representing a shorter-lived subset of the previous tracking. For the wet season, there are hints that  $Z > 35$  dBZ echo regions eventually outgrow those of the dry season, yet most cells quickly dissipate at later-relative stages (times  $> 0.7$ ). Overall, composites suggest that dry season cells are relatively compact and intense, occupied by higher  $Z > 35$  dBZ echoes and  
240 retaining modest precipitation intensity for much of their lifetimes. These findings are consistent with previous GoAmazon2014/5 studies by Giangrande et al. (2020) that proposed that drier mid-levels in the drier seasons may limit periphery precipitation (i.e., enhanced evaporation), whereas wet season cells may exhibit more resilient periphery precipitation.

Seasonal differences are also revealed when considering radar properties that are more directly associated with  $Z$   
245 magnitude (Figure 4cde). Dry season composites skew their strongest  $Z$  signatures to earlier life cycle stages, often with maximum behaviors found prior to normalized life cycle time  $< 0.2$  (i.e., within the first few qualifying radar volumes). An early storm intensification is consistent with arguments from the previous section indicating increased PBL instability during the dry season, reduced MLCIN, and higher low level MLCAPE. Nevertheless, composite dry season cell areas remain relatively unvarying after an initial intensification period throughout a lengthy portion of their normalized lifetime. In contrast,  
250 wet season composites indicate a gradual increase in  $Z$  and areal precipitation coverage, with peak  $Z$  (normalized time  $\cong 0.4$ ) preceding an associated peak in precipitation coverage (i.e., normalized time  $\cong 0.7$ ). Composite wet season storms appear to achieve similarly intense  $Z$  cores to our sampled dry counterparts at later moments in cell life cycle. This result may not be surprising since our events share statistically similar CAPE values and these comparisons target longer-lived cells that conditionally may favor the more intense behaviors from the wet season.

## 255 4.2 Composite Overpass Profiler Behaviors

In Figure 5, we plot cumulative overpass vertical air velocity retrievals contingent on season (left panels “wet”, middle panels “dry”, right panel “dry-wet difference”) and according to multiple RWP retrieval thresholds ( $> 25$  dBZ top,  $> 35$  dBZ bottom). Median vertical air velocity (solid lines) and 5th/95th percentiles (dashed lines) are overlaid on the CFADs as





reference for the extreme instantaneous observations from these events. Overall, composite velocity CFADs indicate  
260 downdrafts are common at low levels, but CFADs transition towards more prominent updraft observations aloft (peak updrafts  
 $\cong 10$  m/s, Figure 5abc).

With respect to updraft observations, the  $Z > 25$  dBZ CFADs suggest dry season maximal updrafts are more intense,  
although the relative enhancement is modest  $O[2-3 \text{ ms}^{-1}]$  and primarily observed at low levels below the melting level (to  
approx. 6 km). The dry season samples in these  $Z > 25$  dBZ CFADs displays are favoring more frequent and modest downdrafts  
265 aloft (to be discussed). However, updraft extremes aloft indicate dry season observations are recording updrafts of comparable  
intensity to our wet season samples; the most intense (95th percentile) updraft retrievals for both seasons are  $O[10 \text{ ms}^{-1}]$ .  
Potential physical reasons for observed seasonal updraft profile characteristics, comparable magnitudes aloft, and shifts therein  
will be discussed in Sect. 4.3.

For downdraft observations, the most consistent downdrafts we observe were associated with regions below the  
270 melting level (precipitation-driven). Interestingly, downdrafts are observed to higher altitudes, but the most frequent and  
vigorous ( $> 5 \text{ ms}^{-1}$ ) are found within the dry season events. As will be confirmed with our subsequent  $Z > 35$  dBZ CFAD  
discussions, the majority of these dry season downdraft retrievals aloft are associated with weaker  $Z < 35$  dBZ echoes found  
towards the peripheries of the more intense cores. Previously, Giangrande et al. (2016) investigated the GoAmazon2014/5  
RWP dataset and suggested that strong downdrafts aloft may provide indirect evidence for stronger updrafts (under higher  
275 CAPE/CIN and/or dry season conditions). Their interpretation was that compensating toroidal-like circulations associated with  
stronger updrafts that were not directly observed may promote those stronger downdrafts aloft. While not stated by those  
authors, greater precipitation/condensate loading associated with stronger updrafts may also contribute to stronger downdrafts  
using those arguments. As with their efforts, we did not observe significantly stronger updrafts aloft, but found that the  
observed shift in downdraft likelihood and intensity was a dry season phenomenon. A discussion on possible physical reasons  
280 for these downdrafts is found in Sect. 4.4 to complement updraft discussions in Sect. 4.3.

In Figure 5def, cumulative CFAD plots shift towards prominent updraft signatures when we emphasize only those  
observations from the more intense  $Z > 35$  dBZ “core” precipitation instances from these same overpasses. Moreover, higher  
reflectivity regions aloft are consistently, and increasingly to higher altitude, associated with updrafts. Physically, one  
interpretation is that larger or more intense precipitation signatures (lofted, larger or more dense hydrometeors) aloft are also  
285 those conditionally associated with stronger updrafts overall. For the wet season in particular, maximum updraft signatures  
consistently peak above the melting level, with extreme values  $O[10 \text{ ms}^{-1}]$  similar to those retrieved during the dry season  
events (above 6-7 km). Interestingly, both seasons record few observations of intense downdrafts within these higher or core  
 $Z > 35$  dBZ regions aloft. Stronger dry season downdrafts are observed below the melting level, similar to Amazon MCS  
studies by Wang et al. (2019).

290 When interpreting cumulative CFAD results above, cell maturity of the corresponding overpasses is not explicitly  
revealed by these displays. In Figures 6 and 7, we plot CFADs contingent on cell overpass ETH, where retrievals before/after  
an ETH of 10 km is used as a proxy for relative storm maturity. Overall, low level precipitation-driven downdraft signatures



for both seasons are more prominent and extend further above the melting level for our higher ETH observations. As in cumulative CFADs, dry season overpasses indicate stronger updrafts, but these stronger updrafts are primarily found below 6  
295 km, and for the  $ETH < 10$  km retrieval instances we associate with developing cloud life cycle stages. To later  $ETH > 10$  km stages, dry season retrievals are dominated by downdrafts (frequency), with strong downdraft motions observed  $O[5 \text{ ms}^{-1}]$  aloft. In contrast, wet season CFADs suggest a strengthening and more prevalent updrafts aloft to the later  $ETH > 10$  km stage observations.

The ETH displays filtered by  $Z > 35$  dBZ regions display a much clearer association between the presence of larger  
300  $Z$  values reaching higher altitudes and stronger updraft observations (both seasons). However, our CFADs suggest that similarly-intense reflectivity factors aloft (i.e., sampling 35 dBZ to 6 km) may be associated with a fairly wide range of updraft intensity contingent on the season and/or where those observations fall within the storm life cycle. Physically, these complications follow from storms having updrafts that are at times less impeded by precipitation, but any transition to stronger updrafts may also be convolved within increasing precipitation (or  $Z$ ). Nevertheless, the  $Z > 35$  dBZ CFADs reinforce that  
305 strong  $Z$  signatures to higher altitudes may be attributed to strong updrafts, and increasingly exclusive to updrafts above the melting level.

Quartile breakdowns of storm life cycle for RWP retrievals are plotted in Figure 8. Dry season vertical air velocity retrievals suggest that the strongest upwards air motions are confined to the earliest life cycle stages, consistent with surveillance radar signatures for dry season storm intensification. Wet season quartile breakdowns reveal a gradual shift in  
310 draft characteristics towards more intense air motions by the middle quartiles (or associated peaks in  $Z$  and  $ETH$ ), also in alignment with previous surveillance properties. The evidence for more intense dry season upwards air motions at the low levels does not appear confined to any particular life cycle stage. However, stronger updrafts aloft are found with increasing time for the wet season, and more prominent downdrafts aloft are found with increasing time for the dry season. Late cell phase samples (time  $> 0.75$ ) are unavailable for the dry season, but late stage wet season samples imply a higher frequency of  
315 observations associated with downdrafts below the melting level, and a shifting for the observations of updrafts further aloft (i.e., possible signatures for remnant anvil / dissipating cloud air motions).

#### 4.3 Physical Reasons for Seasonal Differences in Updraft Behavior

We explore the physical reasons for the differences in draft behaviors evident in Figs. 4-8 by applying a simplified updraft model to the 12 UTC sounding from each case. Because the entrainment rates in the observed updrafts are not known,  
320 we aim to simulate ascending parcels with a range of entrainment rates that encompasses what we might reasonably expect in weakly sheared tropical convection. To generate this range of entrainment rates, we use a stochastic parcel model (SPM) that is formulated in a similar manner to the eddy diffusivity/mass flux shallow convective scheme described in Suselj et al. (2013, 2019). A detailed technical description of the model is available in Appendix A. In short, we simulate 100 different parcels in each sounding. We assume entrainment in each parcel occurs in a series of discrete stochastic mixing events that follow a  
325 Poisson distribution, with the peak in the distribution corresponding to a typical fractional entrainment rate in tropical deep



convection of  $2 \times 10^{-4} \text{ m}^{-1}$  (e.g., Romps and Kuang 2010). Finally, we produce histograms at each height for the vertical air velocities among those parcels to compare this SPM output to observed CFAD characteristics from the previous sections. These histograms were generated by dividing the vertical velocity versus height parameter space into  $1 \text{ m s}^{-1}$  and 250 m bins respectively, and summing over a given season (i.e., wet or dry) all the points along SPM parcel paths that fell into each bin. We divided by the number of events in that season, and applied a Gaussian filter with a radius of influence of  $5 \text{ m s}^{-1}$  and 1250 m with respect to vertical velocity and height, respectively. For the model outputs we plot in Figure 9, those parcels that did not reach 5 km were excluded to maintain our focus on DCCs.

In Figures 9ab, we plot summary velocity profile behaviors from the multiple realizations that start from the morning radiosondes for each wet (Figure 9a) and dry (Figure 9b) event. Maximum parcel heights for SPM parcels were in the 10-13 km range (Figures 9ab), which is generally consistent with observed echo top heights (e.g., Wang et al., 2018). This suggests that the entrainment rates of SPM parcels were reasonably consistent with those in the observed storms, since entrainment strongly regulates cloud depth. Peak vertical velocities are 50-100% larger than what was observed by the RWPs. This discrepancy between peak SPM vertical velocities and observations is at least partially explained by an expected underestimation of the extreme/peak updraft velocities by RWP sampling (as was previously noted).

We subtracted the wet season histogram from the dry season histogram in Figure 9c to plot seasonal differences in parcel behavior, where blue (red) values indicate the SPM outputs a higher incidence for more intense wet (dry) season updrafts. More intense dry season updrafts are prevalent in model realizations at the lower levels, attributed to the higher low level MLCAPE. This result is consistent with RWP observations that also suggest more intense low level updrafts for dry season samples. Strong (i.e.,  $5\text{-}10 \text{ m s}^{-1}$ ) updrafts become more prominent aloft ( $> 6\text{-}7 \text{ km}$ ) in the wet season and more comparable to those in the dry season. These comparatively stronger updrafts aloft between the wet and dry season model realizations is also consistent with the shift in our RWP difference fields (i.e., Figs. 7f and 9c).

In Figures 9d, we plot SPM results in the form of dry-wet differences as from Figure 9c, but after re-running the realizations for each case and replacing all of the RH profiles (above 2 km) with an average RH over all dry season cases. The motivation for these tests was an attempt to remove RH considerations, thus possibly highlighting residual differences resulting from the different CAPE profiles. Given the more prominent dry season updraft realizations to higher levels, one implication from this test is that if seasonal RH considerations are removed, stronger updrafts dominate the dry season realizations. This may suggest that the lower RH mitigates the intensity of dry season updrafts, or equivalently, that the larger RH of the wet season is essential to its larger incidence of deeper updrafts.

#### 4.4 Physical Reasons for Seasonal Differences in Downdraft Behavior

To conclude our analysis, we provide a plausible explanation for why downdrafts were more intense aloft in the dry season than observed for the wet season. Recent Amazon MCS observations from Wang et al. (2019; 2020) indicate drier dry season low to mid-level conditions favor stronger downdrafts and/or higher downdraft origin heights. One hypothesis for our isolated cell events is that mixtures between drafts and environmental parcels are more negatively buoyant in the dry season.



Consequently, these parcels will experience more intense downward accelerations. To evaluate this, we leveraged the parcel  
360 properties simulated by the SPM in the previous subsection.

For each case, we selected the SPM parcel at each height with the median moist static energy (MSE). This parcel was  
defined as the “updraft parcel”, for which we recorded the MSE, water vapor  $q_v$ , and condensate  $q_c$  mixing ratios of this  
parcel. For these tests, we assumed that the MSE and  $q_c$  and of the updraft parcel mix linearly with the environment ( $q_c$  is  
0 in the environment), and consider mixtures with fractions of updraft air ranging from 0.1 to 0.9 at intervals of 0.1 (i.e., the  
365 environment composes the other fraction of this mixture). Using this range of mixtures and assuming saturation, we solved for  
the parcel temperature and buoyancy for each updraft air fraction. Then, we recorded the average buoyancy of all negatively  
buoyant mixtures at each height, which gives a vertical profile of negative buoyancy for each event. We expect that mixtures  
of the updraft and the environment are more negatively buoyant in dry season than in wet season events, because of the smaller  
free-tropospheric relative humidity in the former. We further assume that these mixtures between an updraft and its  
370 environment are responsible for initiating downdrafts, and that strongly negatively buoyant mixtures will initiate stronger  
downdrafts than their less negatively buoyant counterparts.

As we plot in Figure 10, the resulting dry season buoyancy profiles are more negative than wet season buoyancy  
profiles between 2 km and 8 km. The difference is statistically significant between 4 and 6 km, with the dry season buoyancy  
being a factor of 1.5 to 2 more negative than the wet season negative buoyancy. Note, the calculations in Figure 10 apply to  
375 updraft mixtures. However, we speculate that different mixtures of the cloud’s surrounding environmental air mixed with  
detained updraft air or downdrafts would behave similarly, where mean dry-season drafts would exhibit greater negative  
buoyancy compared to wet-season drafts.

An alternative interpretation for the observational differences is rooted in RWP sampling as related to the seasonal  
differences in cell areal precipitation characteristics. Recall, dry season downdrafts aloft were most frequently observed in 25  
380  $< Z < 35$  dBZ samples and at later life cycle stages; these observations may be those preferentially collected near the edges of  
dry season cells that we found to be more compact than their wet season counterparts. This compact nature was partially  
attributed to evaporation and/or mixing with the drier RH environment limiting cell growth, potentially prioritizing these  
observations to locations where these processes are occurring. Intrinsically, observations from compact cells are also collected  
closer in proximity to cell cores and stronger air motions. Equivalently, wet season observations having similar 25  $< Z < 35$   
385 dBZ may include retrievals embedded within resilient and/or wider-spread precipitation areas and regions that are more  
insulated from the cloud edge, and that are consequently less prone to being associated with downdrafts. A comprehensive  
exploration of all downdraft possibilities is beyond the scope of our study, but will be examined in future research.

## 5. Summary of Key Findings

390 This study investigates daytime DCC observations to document changes in storm characteristics contingent on larger-  
scale shifts between the Amazon wet and dry seasons. Our focus is on the use of surveillance weather radar cell tracking and



coupled profiler-based vertical air velocity observations. Overall, the Amazon offers a unique natural laboratory for these studies, providing the frequent DCCs necessary for documenting storm life cycle in the manner presented. Observations of this kind are rare, but critical for high-resolution cloud model development that have added new capabilities for forward radar operators, yet lack coupled microphysical/dynamical observations (e.g., Stein et al., 2015).

The key findings of this study are as follows:

- Dry season cells show more intense drafts and precipitation properties compared to wet season storms, but reduced convective area coverage.
- These dry storms rapidly developed and achieved peak intensity at early life cycle stages, potentially due to higher low level MLCAPE and/or reduced morning cloud cover in the dry season.
- Wet season storms were longer-lived, achieving modest precipitation intensity and attaining larger convective area coverage  $Z > 35$  dBZ to dry season counterparts, and achieving their most intense precipitation and updrafts later in their life cycle.
- Dry season updraft profiles exhibited stronger updrafts at lower altitudes, and stronger downdrafts aloft than wet season storms. However, wet season storms exhibited a higher incidence of moderate-to-strong updrafts aloft than in the dry season, and less intense and/or frequent downdrafts overall for our sampling conditions.
- The stronger updrafts at low levels in the dry season are attributed to the larger low-level CAPE in the storm environment. Whereas, a higher prevalence of updrafts aloft in the wet season resulted from larger environmental RH and less entrainment-driven dilution of updraft buoyancy.
- Stronger downdrafts aloft in the dry season were attributed to the smaller environmental RH and the associated increased likelihood of evaporation and negative buoyancy within the mixtures of updraft and environmental air that initiate downdrafts.

Finally, our results put forward practical connections between quantities such as radar reflectivity and updraft intensity. These ideas are of interest for proxy retrievals of storm dynamics (updraft intensity, mass flux) from spaceborne platforms that can fill gaps in oceanic, remote, or similarly-challenged regions (e.g., Jeyaratnam et al., 2021). For example, we observe a strong association between the earlier occurrence and deeper  $Z > 35$  dBZ regions aloft with the presence of stronger updrafts. These connections are not perfect, but may be physically intuitive; intense updrafts are those that likely generate more intense precipitation, while necessary to loft larger hydrometeors associated with larger reflectivity. While column reflectivity echo heights or integrated reflectivity measures (e.g., Kumar et al., 2016) are informative, our studies suggest adding life cycle guidance for proxy velocity or mass flux retrievals should help improve those methods.



## Appendix A: Stochastic Parcel Model Formulation

The SPM uses dry static energy (DSE) and moist static energy (MSE) as prognostic thermodynamic variables, which we define  
425 as:

$$(1) \quad DSE = c_p T + gz,$$

$$(2) \quad MSE = c_p T + L_v q_v - L_i \omega q_c + gz,$$

where  $c_p = 1005 \text{ J kg}^{-1} \text{ K}^{-1}$  is the heat capacity of dry air,  $T$  is updraft temperature,  $L_v = 2,501,000 \text{ J kg}^{-1}$  and  $L_i =$   
330,000  $\text{ J kg}^{-1}$  are the latent heats of vaporization and freezing respectively (approximated with their empirical values at  
430 273.15 K),  $q_v$  and  $q_c$  are the updraft's water vapor and condensate mass fractions respectively, and  $g$  is gravity. The  
dimensionless parameter  $\omega$  discriminates liquid from ice. It is set to 0 when  $T > 273.15 \text{ K}$ , 1 when  $T < 253.15 \text{ K}$ , and  
linearly transitions from 0 to 1 over the temperature range between 273.15 K and 253.15 K. Next, we define the updraft  
kinetic energy  $K$  as:

$$(3) \quad K = \frac{w^2}{2},$$

435 where  $w$  is vertical velocity. Finally, we define the saturation water vapor mass fraction  $q^*$  as:

$$(4) \quad q^* = \frac{R_d}{R_v} \frac{611.0}{p} e^{\frac{L_v}{R_v} \left( \frac{1}{T} - \frac{1}{273.15} \right)},$$

where  $R_d = 287 \text{ J kg}^{-1} \text{ K}^{-1}$  and  $R_v = 461 \text{ J kg}^{-1} \text{ K}^{-1}$  are the dry and moist specific gas constants respectively.

During the sub-saturated part of ascent (i.e,  $q_v < q^*$ ), the prognostic thermodynamic equations are:

$$(5) \quad \frac{dDSE}{dz} = -\varepsilon(DSE - DSE_0),$$

$$440 \quad (6) \quad \frac{dq_v}{dz} = -\varepsilon(q_v - q_0).$$

Once a parcel achieves saturation, they become:

$$(7) \quad \frac{dMSE}{dz} = -\varepsilon(MSE - MSE_0),$$

$$(8) \quad \frac{dq_c}{dz} = -\frac{dq^*}{dz} - \varepsilon q_c,$$

$$(9) \quad q_v = q^*.$$

445 The prognostic equation for  $k$  at all levels is:

$$(10) \quad \frac{dk}{dz} = g \frac{T - T_0}{T_0} + g \left( \frac{R_v}{R_d} - 1 \right) (q_v - q_0) - g q_c - \left( \varepsilon + \frac{3c_d}{8L} \right) k.$$

Variables with a subscript "0" represent the updraft background environment (in this case, the radiosonde profile),  $\varepsilon$  is a  
fractional entrainment inverse length scale,  $c_d$  is a drag coefficient that is set to 0.5 based on Morrison and Peters (2018), and  
 $L$  is a length scale that represents the updraft radius (given a value below). The last term in eq. (10) represents the effects of  
450 momentum entrainment (via  $\varepsilon$ ) and form drag on ascending cloud elements (via the  $\frac{3c_d}{8L}$  term).

We simulate 100 updrafts per sounding, wherein parcels within updrafts are subject to discrete Poisson-process entrainment  
events as they ascend. Hence,  $\varepsilon$  is defined as:



$$(11) \quad \varepsilon = \frac{0.2}{\Delta z} \varphi \left( \frac{\Delta z}{L} \right),$$

where  $\varphi$  is the Poisson,  $\Delta z$  is the vertical grid spacing of the discretized model, and  $L$  is once again the length scale that  
455 represents the updraft radius. As stated in the main text, we set  $L$  to 1000 m. Our conclusions were relatively unchanged by  
variations in  $L$  from 500 m to 1500 m.

Our model is vertically integrated with a simple first-order up-wind Euler scheme, with a initial  $w$  ranging from 0.5-  
1.5  $\text{m s}^{-1}$ ,  $T'$  ranging from 0.5-1.5 K,  $q'_v$  ranging from 0.5 to 1.5  $\text{g kg}^{-1}$  (where a ' denotes a departure from the value at the  
lowest level of the sounding), and a vertical grid spacing of 100 m. Vertical integration was stopped in each updraft at the first  
460 instance of  $k < 0$ , and the vertical grid point below this level was defined as the updraft top. Using all updrafts simulated  
among all dry and wet season events, histograms were created at each grid height of  $w$ , binned at 1  $\text{m s}^{-1}$  intervals.

#### Code and Data Availability.

465 All ARM data, including RWP (<https://doi.org/10.5439/1025128>, Coulter et al., 2009), ARSCL  
(<https://doi.org/10.5439/1027282>, Giangrande and Johnson, 2003), SONDE (<https://doi.org/10.5439/1021460>, Holdridge et  
al., 1994), and other datasets used in this study, can be downloaded at <http://www.arm.gov> (last access: 10 Aug 2022). These  
are associated with several standard ARM raw streams, value-added products (VAP), and GoAmazon2014/5 PI datasets.

#### 470 Author Contributions.

SEG, TB, and JMP designed the research, performed the research, and wrote the paper.

#### Competing interests.

The authors declare that they have no conflict of interest.

475

#### Acknowledgements.

This study was supported by the U.S. Department of Energy (DOE) Atmospheric System Research (ASR) Program. This paper  
has been authored by an employee of Brookhaven Science Associates, LLC, under contract DE-SC0012704 with the U.S.  
DOE. The publisher by accepting the paper for publication acknowledges that the United States Government retains a  
480 nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this paper, or allow others  
to do so, for United States Government purposes. This work was also supported by the DOE ARM Program and its AMF3  
Site Science project, which is supported by the Office of Biological and Environmental Research in the DOE, Office of  
Science, and through the U.S. DOE contract no. DE-SC0012704 to Brookhaven National Laboratory. We also acknowledge  
FAPESP (São Paulo Research Foundation) projects 2009/15235-8 and 2015/14497-0. We would like to thank CENSIPAM



485 (Centro Gestor e Operacional do Sistema de Proteção da Amazônia) for providing the Manaus SIPAM radar data. The authors  
also wish to thank David Mechem (KU) for his thoughtful comments on this work.

## References

- Ackerman, T. P. and Stokes, G. M.: The Atmospheric Radiation Measurement Program, *Physics Today*, 56(1), 38–44,  
490 doi:10.1063/1.1554135, 2003.
- Adams, D. K., Gutman, S., Holub, K., and Pereira, D.: GNSS Observations of Deep Convective timescales in the Amazon,  
*Geophys. Res. Lett.*, 40, 2818–2823, <https://doi.org/10.1002/grl.50573>, 2013.
- Adams, D. K., Barbosa, H. M. J., and Gaitán De Los Ríos, K. P.: A Spatiotemporal Water Vapor–Deep Convection Correlation  
Metric Derived from the Amazon Dense GNSS Meteorological Network, *Mon. Weather Rev.* 145, 279–288,  
495 <https://doi.org/10.1175/MWR-D-16-0140.1>, 2017.
- Anagnostou, E. N.: A convective/stratiform precipitation classification algorithm for volume scanning weather radar  
observations, *Meteorological Applications*, 11, 4, 291–300, doi:10.1017/S1350482704001409, 2004.
- Anderson, N. F., Grainger, C. A., & Stith, J. L. (2005). Characteristics of Strong Updrafts in Precipitation Systems over the  
Central Tropical Pacific Ocean and in the Amazon, *Journal of Applied Meteorology*, 44(5), 731–738.
- 500 Atmospheric Radiation Measurement (ARM): Climate Research Facility: Balloon-Borne Sounding System (SONDE), 3.21297  
S 60.5981 W: ARM Mobile Facility (MAO) Manacapuru, Amazonas, Brazil; AMF1 (M1), compiled by: Holdridge, D.,  
Kyrouac, J., and Coulter, R., Atmospheric Radiation Measurement (ARM) Climate Research Facility Data Archive, Oak  
Ridge, Tennessee, USA, Data set accessed at 2018-07-01 <https://doi.org/10.5439/1025284>, 1993.
- Biscaro, T. S., Machado, L. A. T., Giangrande, S. E., and Jensen, M. P.: What drives daily precipitation over the central  
505 Amazon? Differences observed between wet and dry seasons, *Atmos. Chem. Phys.*, 21, 6735–6754,  
<https://doi.org/10.5194/acp-21-6735-2021>, 2021.
- Bryan, G. H., and Fritsch, J. M.: A Benchmark Simulation for Moist Nonhydrostatic Numerical Models. *Monthly Weather  
Review* 130, 12, 2917–2928, [https://doi.org/10.1175/1520-0493\(2002\)130<2917:ABSFMN>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<2917:ABSFMN>2.0.CO;2), 2002.
- Borque, P., Kollias, P., & Giangrande, S. (2014). First Observations of Tracking Clouds Using Scanning ARM Cloud Radars,  
510 *Journal of Applied Meteorology and Climatology*, 53(12), 2732–2746.
- Cifelli, R., W. A. Petersen, L. D. Carey, S. A. Rutledge, and M. A. F. da Silva Dias, 2002: Radar observations of the kinematic,  
microphysical, and precipitation characteristics of two MCSs in TRMM LBA. *J. Geophys. Res.*, 107, 8077,  
doi:10.1029/2000JD000264.
- Dixon, M., and G. Wiener, 1993: TITAN: Thunderstorm Identification, Tracking, Analysis, and Nowcasting—A radar-based  
515 methodology. *J. Atmos. Oceanic Technol.*, 10, 785–797, doi:10.1175/1520-0426(1993)010<0785:TTITAA>2.0.CO;2.





- Feng, Z. and coauthors. More frequent intense and long-lived storms dominate the springtime trend in central US rainfall. *Nat. Commun.* 7, 13429 (2016).
- Feng, Z., Houze, R. A., Jr., Leung, L. R., Song, F., Hardin, J. C., Wang, J., Gustafson, W. I., Jr., & Homeyer, C. R. (2019). Spatiotemporal Characteristics and Large-Scale Environments of Mesoscale Convective Systems East of the Rocky Mountains, *Journal of Climate*, 32(21), 7303-7328.
- 520 Foote, G. B., & Du Toit, P. S. (1969). Terminal Velocity of Raindrops Aloft, *Journal of Applied Meteorology and Climatology*, 8(2), 249-253.
- Fridlind, A. M., van Lier-Walqui, M., Collis, S., Giangrande, S. E., Jackson, R. C., Li, X., Matsui, T., Orville, R., Picel, M. H., Rosenfeld, D., Ryzhkov, A., Weitz, R., and Zhang, P.: Use of polarimetric radar measurements to constrain simulated convective cell evolution: a pilot study with Lagrangian tracking, *Atmos. Meas. Tech.*, 12, 2979–3000, <https://doi.org/10.5194/amt-12-2979-2019>, 2019.
- 525 Gbate, V. P., & Kollias, P. (2016). On the Controls of Daytime Precipitation in the Amazonian Dry Season, *Journal of Hydrometeorology*, 17(12), 3079-3097.
- Giangrande, S. E., Collis, S., Straka, J., Protat, A., Williams, C., and Krueger, S.: A Summary of Convective-Core Vertical Velocity Properties Using ARM UHF Wind Profilers in Oklahoma. *Journal of Applied Meteorology and Climatology*, 52, 10, 2278-2295, <https://doi.org/10.1175/JAMC-D-12-0185.1>, 2013.
- 530 Giangrande, S. E., Feng, Z., Jensen, M. P., Comstock, J. M., Johnson, K. L., Toto, T., Wang, M., Burleyson, C., Bharadwaj, N., Mei, F., Machado, L. A. T., Manzi, A. O., Xie, S., Tang, S., Silva Dias, M. A. F., de Souza, R. A. F., Schumacher, C. and Martin, S. T.: Cloud characteristics, thermodynamic controls and radiative impacts during the Observations and Modeling of the Green Ocean Amazon (GoAmazon2014/5) experiment, *Atmospheric Chemistry and Physics*, 17(23), 14519–14541, doi:10.5194/acp-17-14519-2017, 2017.
- Giangrande, S. E., Wang, D., and Mechem, D. B.: Cloud regimes over the Amazon Basin: perspectives from the GoAmazon2014/5 campaign, *Atmospheric Chemistry and Physics*, 20, 7489–7507, <https://doi.org/10.5194/acp-20-7489-2020>, 2020.
- 540 Göke, S., H. T. Ochs, and R. M. Rauber, 2007: Radar analysis of precipitation initiation in maritime versus continental clouds near the Florida coast: Inferences concerning the role of CCN and giant nuclei. *J. Atmos. Sci.*, 64, 3695–3707, doi:10.1175/JAS3961.1.
- Hu, J., Rosenfeld, D., Ryzhkov, A., Zrnic, D., Williams, E., Zhang, P., et al. (2019). Polarimetric radar convective cell tracking reveals large sensitivity of cloud precipitation and electrification properties to CCN. *Journal of Geophysical Research: Atmospheres*, 124, 12, 194–12, 205. <https://doi.org/10.1029/2019JD030857>
- 545 Jeyaratnam, J., Luo, Z. J., Giangrande, S. E., Wang, D., & Masunaga, H. (2021). A satellite-based estimate of convective vertical velocity and convective mass flux: Global survey and comparison with radar wind profiler observations. *Geophysical Research Letters*, 48, e2020GL090675. <https://doi.org/10.1029/2020GL090675>



- Jorgensen, D. P., Zipser, E. J., & LeMone, M. A. (1985). Vertical Motions in Intense Hurricanes, *Journal of Atmospheric Sciences*, 42(8), 839-856.
- 550 Kumar, V. V., Jakob, C., Protat, A., Williams, C. R., & May, P. T. (2015). Mass-Flux Characteristics of Tropical Cumulus Clouds from Wind Profiler Observations at Darwin, Australia, *Journal of the Atmospheric Sciences*, 72(5), 1837-1855.
- Limpert, G., Houston, A., and Lock, N.: The advanced algorithm for tracking objects (AALTO), *Meteor. Apps.*, 22, 694–704, <https://doi.org/10.1002/met.1501>, 2015.
- 555 Machado, L. A., W. B. Rossow, R. L. Guedes, and A. W. Walker, 1998: Life cycle variations of mesoscale convective systems over the Americas. *Mon. Wea. Rev.*, 126, 1630–1654, doi:10.1175/1520-0493(1998)126<1630:LCVOMC>2.0.CO;2.
- Machado, L. A. T., H. Laurent, N. Dessay, and I. Miranda (2004), Seasonal and diurnal variability of convection over the Amazonia: A comparison of different vegetation types and large scale forcing, *Theor. Appl. Climatol.*, 78, 61–77, doi:10.1007/s00704-004-0044-9.
- 560 Machado, L. A. T., Silva Dias, M. A. F., Morales, C., Fisch, G., Vila, D., Albrecht, R., Goodman, S. J., Calheiros, A. J. P., Biscaro, T., Kummerow, C., Cohen, J., Fitzjarrald, D., Nascimento, E. L., Sakamoto, M. S., Cunningham, C., Chaboureau, J.-P., Petersen, W. A., Adams, D. K., Baldini, L., Angelis, C. F., Sapucci, L. F., Salio, P., Barbosa, H. M. J., Landulfo, E., Souza, R. A. F., Blakeslee, R. J., Bailey, J., Freitas, S., Lima, W. F. A., and Tokay, A.: The CHUVA Project: How Does Convection Vary across Brazil?, *Bull. Am. Meteorol. Soc.*, 95, 1365–1380, <https://doi.org/10.1175/BAMS-D-13-00084.1>, 2014.
- 565 Machado, L. A. T., Calheiros, A. J. P., Biscaro, T., Giangrande, S., Silva Dias, M. A. F., Cecchini, M. A., Albrecht, R., Andreae, M. O., Araujo, W. F., Artaxo, P., Borrmann, S., Braga, R., Burleyson, C., Eichholz, C. W., Fan, J., Feng, Z., Fisch, G. F., Jensen, M. P., Martin, S. T., Pöschl, U., Pöhlker, C., Pöhlker, M. L., Ribaud, J.-F., Rosenfeld, D., Saraiva, J. M. B., Schumacher, C., Thalman, R., Walter, D., and Wendisch, M.: Overview: Precipitation characteristics and sensitivities to environmental conditions during GoAmazon 2014/2015 and ACRIDICON-CHUVA, *Atmos. Chem. Phys.*, 18, 6461–6482, <https://doi.org/10.5194/acp-18-6461-2018>, 2018.
- 570 Maddox, R. A., 1980: Mesoscale convective complexes. *Bull. Amer. Meteor. Soc.*, 61, 1374–1387, doi:10.1175/1520-0477(1980)061<1374:MCC>2.0.CO;2.
- Martin, S. T., Artaxo, P., Machado, L., Manzi, A. O., Souza, R. A., Schumacher, C., Wang, J., Biscaro, T., Brito, J., Calheiros, A., Jardine, K., Medeiros, A., Portela, B., de Sá, S. S., Adachi, K., Aiken, A. C., Albrecht, R., Alexander, L., Andreae, M. O.,
- 575 Barbosa, H. M., Buseck, P., Chand, D., Comstock, J. M., Day, D. A., Dubey, M., Fan, J., Fast, J., Fisch, G., Fortner, E., Giangrande, S., Gilles, M., Goldstein, A. H., Guenther, A., Hubbe, J., Jensen, M., Jimenez, J. L., Keutsch, F. N., Kim, S., Kuang, C., Laskin, A., McKinney, K., Mei, F., Miller, M., Nascimento, R., Pauliquevis, T., Pekour, M., Peres, J., Petäjä, T., Pöhlker, C., Pöschl, U., Rizzo, L., Schmid, B., Shilling, J. E., Dias, M. A., Smith, J. N., Tomlinson, J. M., Tóta, J., and Wendisch, M.: The Green Ocean Amazon Experiment (GoAmazon2014/5) Observes Pollution Affecting Gases, Aerosols,
- 580 Clouds, and Rainfall over the Rain Forest, *Bull. Am. Meteorol. Soc.*, 98, 981–997, <https://doi.org/10.1175/BAMS-D-15-00221.1>, 2017.



- Mather, J. H. and Voyles, J. W.: The ARM Climate Research Facility: A Review of Structure and Capabilities. *Bulletin of the American Meteorological Society*, 94(3), 377-392, doi:10.1175/BAMS-D-11-00218.1, 2013.
- Morrison, H., & Peters, J. M. (2018). Theoretical Expressions for the Ascent Rate of Moist Deep Convective Thermals, *Journal of the Atmospheric Sciences*, 75(5), 1699-1719.
- 585 Nobre, P., Malagutti, M., Urbano, D. F., De Almeida, R. A. F., and Giarolla, E.: Amazon deforestation and climate change in a coupled model simulation, *J. Climate*, 22, 5686–5697, 2009.
- Peters, J. M., Mulholland, J. P., & Chavas, D. R. (2022). Generalized Lapse Rate Formulas for Use in Entraining CAPE Calculations, *Journal of the Atmospheric Sciences*, 79(3), 815-836.
- 590 Petersen, W. A., Nesbitt, S. W., Blakeslee, R. J., Cifelli, R., Hein, P., & Rutledge, S. A. (2002). TRMM Observations of Intraseasonal Variability in Convective Regimes over the Amazon, *Journal of Climate*, 15(11), 1278-1294.
- Prein, A. F. et al. Increased rainfall volume from future convective storms in the US. *Nat. Clim. Change* 7, 880–884 (2017).
- Protat, A., & Williams, C. R. (2011). The Accuracy of Radar Estimates of Ice Terminal Fall Speed from Vertically Pointing Doppler Radar Measurements, *Journal of Applied Meteorology and Climatology*, 50(10), 2120-2138.
- 595 Rosenfeld, D., 1987: Objective method for analysis and tracking of convective cells as seen by radar. *J. Atmos. Oceanic Technol.*, 4, 422–434, doi:10.1175/1520-0426(1987)004<0422:OMFAAT>2.0.CO;2.
- Saraiva, I., Silva Dias, M. A. F., Morales, C. A. R., and Saraiva, J. M. B.: Regional Variability of Rain Clouds in the Amazon Basin as Seen by a Network of Weather Radars, *Journal of Applied Meteorology and Climatology*, 55(12), <https://doi.org/10.1175/JAMC-D-15-0183.1>, 2657-2675, 2016.
- 600 Schiro, K.A., F. Ahmed, S. E. Giangrande, J. D. Neelin, GoAmazon2014/5 campaign points to deep-inflow approach to deep convection across scales, *Proceedings of the National Academy of Sciences* May 2018, 115 (18) 4577-4582; DOI: 10.1073/pnas.1719842115
- Schumacher, R.S., Rasmussen, K.L. The formation, character and changing nature of mesoscale convective systems. *Nat Rev Earth Environ* 1, 300–314 (2020). <https://doi.org/10.1038/s43017-020-0057-7>
- 605 Silva Dias, M. A. F., and Coauthors, 2002a: Cloud and rain processes in a biosphere-atmosphere interaction context in the Amazon Region. *J. Geophys. Res.*, 107 .8072, doi:10.1029/2001JD000335.
- Silva Dias, M. A. F., and Coauthors, 2002b: A case study of convective organization into precipitating lines in the Southwest Amazon during the WETAMC and TRMM-LBA. *J. Geophys. Res.*, 107 .8078, doi:10.1029/2001JD000375.
- Stein, T. H. M., Hogan, R. J., Clark, P. A., Halliwell, C. E., Hanley, K. E., Lean, H. W., Nicol, J. C., and Plant, R. S.: The DYMEDS project: A statistical approach for the evaluation of convective storms in high-resolution NWP models, *B. Amer. Meteorol. Soc.*, 96, 939–951, <https://doi.org/10.1175/BAMS-D-13-00279.1>, 2015.
- 610 Steiner, M., Houze, R. A., Jr., and Yuter, S. E.: Climatological Characterization of Three-Dimensional Storm Structure from Operational Radar and Rain Gauge Data. *Journal of Applied Meteorology and Climatology* 34, 9, 1978-2007, [https://doi.org/10.1175/1520-0450\(1995\)034<1978:CCOTDS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1995)034<1978:CCOTDS>2.0.CO;2), 1995.



- 615 Sušelj, K., Teixeira, J., & Chung, D. (2013). A Unified Model for Moist Convective Boundary Layers Based on a Stochastic Eddy-Diffusivity/Mass-Flux Parameterization, *Journal of the Atmospheric Sciences*, 70(7), 1929-1953.
- Sušelj, K., Kurowski, M. J., & Teixeira, J. (2019). On the Factors Controlling the Development of Shallow Convection in Eddy-Diffusivity/Mass-Flux Models, *Journal of the Atmospheric Sciences*, 76(2), 433-456.
- Tang, S., Xie, S., Zhang, Y., Zhang, M., Schumacher, C., Upton, H., Jensen, M. P., Johnson, K. L., Wang, M., Ahlgrimm, M.,  
620 Feng, Z., Minnis, P., and Thieman, M.: Large-scale vertical velocity, diabatic heating and drying profiles associated with seasonal and diurnal variations of convective systems observed in the GoAmazon2014/5 experiment, *Atmos. Chem. Phys.*, 16, 14249–14264, <https://doi.org/10.5194/acp-16-14249-2016>, 2016.
- Tian, Y., Zhang, Y., Klein, S. A., & Schumacher, C. (2021). Interpreting the diurnal cycle of clouds and precipitation in the ARM GoAmazon observations: Shallow to deep convection transition. *Journal of Geophysical Research: Atmospheres*, 126,  
625 e2020JD033766. <https://doi.org/10.1029/2020JD033766>.
- Tian, Y., Zhang, Y., & Klein, S. A. (2022). What determines the number and the timing of pulses in afternoon precipitation in the Green Ocean Amazon (GoAmazon) observations? *Geophysical Research Letters*, 49, e2021GL096075. <https://doi.org/10.1029/2021GL096075>.
- Vila, D. A., Machado, L. A. T., Laurent, H., and Velasco, I.: Forecast and Tracking the Evolution of Cloud Clusters (ForTraCC)  
630 Using Satellite Infrared Imagery: Methodology and Validation. *Weather and Forecasting* 23, 2, 233-245, <https://doi.org/10.1175/2007WAF2006121.1>, 2008.
- Wang, D., Giangrande, S. E., Bartholomew, M. J., Hardin, J., Feng, Z., Thalman, R., and Machado, L. A. T.: The Green Ocean: precipitation insights from the GoAmazon2014/5 experiment, *Atmos. Chem. Phys.*, 18, 9121–9145, <https://doi.org/10.5194/acp-18-9121-2018>, 2018.
- 635 Wang, D., Giangrande, S. E., Schiro, K., Jensen, M. P., and Houze, R. A.: The characteristics of tropical and midlatitude mesoscale convective systems as revealed by radar wind profilers, *J. Geophys. Res.-Atmos.*, 124, 4601–4619, <https://doi.org/10.1029/2018JD030087>, 2019.
- Wang, D., Giangrande, S. E., Feng, Z., Hardin, J. C., and Prein, A. F.: Updraft and Downdraft Core Size and Intensity as Revealed by Radar Wind Profilers: MCS Observations and Idealized Model Comparisons, *J. Geophys. Res.-Atmos.*, 125,  
640 e2019JD031774, <https://doi.org/10.1029/2019JD031774>, 2020.
- Williams, E., and Coauthors, 2002: Contrasting convective regimes over the Amazon: Implications for cloud electrification. *J. Geophys. Res.*, 107 .8082, doi:10.1029/2001JD000380.
- Williams, M., and R. A. Houze, 1987: Satellite-observed characteristics of winter monsoon cloud clusters. *Mon. Wea. Rev.*, 115, 505–519, doi:10.1175/1520-0493(1987)115<0505:SOCOWM>2.0.CO;2.
- 645 Yin, L., Fu, R., Shevliakova, E., and Dickinson, R.: How well can CMIP5 simulate precipitation and its controlling processes over tropical South America?, *Clim. Dynam.*, 41, 3127–3143, <https://doi.org/10.1007/s00382-012-1582-y>, 2013.

<https://doi.org/10.5194/egusphere-2022-877>

Preprint. Discussion started: 4 October 2022

© Author(s) 2022. CC BY 4.0 License.



Yuter, S. E., and R. A. Houze Jr. (1995), Three-dimensional kinematic and microphysical evolution of Florida cumulonimbus. Part II: Frequency distribution of vertical velocity, reflectivity, and differential reflectivity, *Mon. Weather Rev.*, 123, 1941–1963.

650



**Table 1. Wet Season Events and Event Details.**

| <u>Date</u> | <u>Initial Radar Echo [LT]</u> | <u>Overpass Time [LT]</u> | <u>Event Duration [min]</u> | <u>MLCAPE [J kg<sup>-1</sup>]</u> | <u>0-6 km MLCAPE [J kg<sup>-1</sup>]</u> | <u>MLCIN [J kg<sup>-1</sup>]</u> | <u>2-6 km mean RH [%]</u> |
|-------------|--------------------------------|---------------------------|-----------------------------|-----------------------------------|--|----------------------------------|---------------------------|
| 20140310    | 1900                           | 2036                      | 156                         | 1800                              | 174                                      | -65                              | 78                        |
| 20140326    | 1524                           | 1536                      | 120                         | 1068                              | 110                                      | -43                              | 69                        |
| 20140331    | 1336                           | 1512                      | 132                         | 1273                              | 112                                      | -84                              | 76                        |
| 20140420    | 1424                           | 1500                      | 108                         | 2333                              | 330                                      | -4                               | 78                        |
| 20140426    | 1312                           | 1312                      | 192                         | 1079                              | 62                                       | -86                              | 87                        |
| 20141202    | 1324                           | 1400                      | 84                          | 1980                              | 261                                      | -28                              | 61                        |
| 20141217    | 1324                           | 1424                      | 132                         | 961                               | 58                                       | -73                              | 73                        |
| 20141219    | 1348                           | 1412                      | 204                         | 1739                              | 210                                      | -13                              | 71                        |
| 20141221    | 1500                           | 1536                      | 240                         | 1887                              | 173                                      | -23                              | 59                        |
| 20141223    | 1048                           | 1148                      | 156                         | 2086                              | 267                                      | -16                              | 77                        |
| 20141227    | 1200                           | 1312                      | 288                         | 1149                              | 210                                      | -36                              | 76                        |
| 20141228    | 1612                           | 1748                      | 132                         | 1435                              | 241                                      | -27                              | 69                        |
| 20141231    | 1136                           | 1212                      | 60                          | 1157                              | 161                                      | -53                              | 70                        |
| 20150106    | 1100                           | 1124                      | 132                         | 696                               | 134                                      | -59                              | 89                        |
| 20150118    | 1224                           | 1224                      | 84                          | 621                               | 37                                       | -117                             | 85                        |
| 20150224    | 1424                           | 1536                      | 156                         | 1751                              | 260                                      | -14                              | 92                        |
| 20150302    | 1500                           | 1612                      | 168                         | 652                               | 15                                       | -182                             | 83                        |
| 20150303    | 1548                           | 1612                      | 72                          | 1292                              | 132                                      | -39                              | 86                        |
| 20150314    | 1548                           | 1700                      | 84                          | 1094                              | 93                                       | -41                              | 84                        |
| 20150322    | 1048                           | 1112                      | 132                         | 1293                              | 142                                      | -44                              | 87                        |
| 20150323    | 1212                           | 1224                      | 96                          | 725                               | 16                                       | -173                             | 88                        |
| 20150401    | 1336                           | 1336                      | 216                         | 815                               | 86                                       | -83                              | 79                        |
| 20150412    | 1124                           | 1224                      | 108                         | 2183                              | 312                                      | -9                               | 88                        |



|             |             |             |            |             |            |            |           |
|-------------|-------------|-------------|------------|-------------|------------|------------|-----------|
| 20150415    | 1624        | 1624        | 132        | 1006        | 60         | -70        | 85        |
| <b>Mean</b> | <b>1346</b> | <b>1429</b> | <i>141</i> | <b>1337</b> | <b>152</b> | <b>-58</b> | <b>78</b> |



655 **Table 2. Dry Season Events and Event Details.**

| <u>Date</u> | <u>Initial Radar Echo [LT]</u> | <u>Overpass Time [LT]</u> | <u>Event Duration [min]</u> | <u>MLCAPE [J kg<sup>-1</sup>]</u> | <u>0-6 km MLCAPE [J kg<sup>-1</sup>]</u> | <u>MLCIN [J kg<sup>-1</sup>]</u> | <u>2-6 km mean RH [%]</u> |
|-------------|--------------------------------|---------------------------|-----------------------------|-----------------------------------|--|----------------------------------|---------------------------|
| 20140612    | 1900                           | 1924                      | 48                          | 713                               | 91                                       | -103                             | 60                        |
| 20140625    | 1348                           | 1400                      | 36                          | 1200                              | 196                                      | -21                              | 51                        |
| 20140712    | 1348                           | 1348                      | 96                          | 492                               | 100                                      | -81                              | 76                        |
| 20140717    | 1648                           | 1712                      | 72                          | 1351                              | 196                                      | -33                              | 49                        |
| 20140718    | 1100                           | 1112                      | 120                         | 1715                              | 261                                      | -15                              | 55                        |
| 20140809    | 1324                           | 1348                      | 60                          | 1377                              | 226                                      | -36                              | 56                        |
| 20140811    | 1112                           | 1112                      | 84                          | 1262                              | 166                                      | -84                              | 59                        |
| 20140815    | 1412                           | 1448                      | 96                          | 2101                              | 360                                      | -23                              | 37                        |
| 20140907    | 1348                           | 1348                      | 96                          | 1759                              | 224                                      | -32                              | 59                        |
| 20140909    | 1436                           | 1448                      | 72                          | 1380                              | 199                                      | -37                              | 76                        |
| 20140913    | 1412                           | 1436                      | 156                         | 1545                              | 226                                      | -41                              | 57                        |
| 20140916    | 1612                           | 1636                      | 96                          | 1939                              | 390                                      | -44                              | 33                        |
| 20140922    | 1024                           | 1100                      | 120                         | 2411                              | 520                                      | -19                              | 35                        |
| 20150607    | 1100                           | 1248                      | 300                         | 2029                              | 298                                      | -23                              | 88                        |
| 20150607    | 1112                           | 1124                      | 108                         | 2029                              | 298                                      | -23                              | 88                        |
| 20150610    | 1148                           | 1224                      | 264                         | 1174                              | 252                                      | -52                              | 78                        |
| 20150614    | 1148                           | 1200                      | 168                         | 1314                              | 206                                      | -23                              | 68                        |
| 20150806    | 1436                           | 1436                      | 84                          | 1896                              | 264                                      | -33                              | 60                        |
| 20150904    | 1624                           | 1700                      | 144                         | 2270                              | 361                                      | -23                              | 60                        |






|             |             |             |            |             |            |            |           |
|-------------|-------------|-------------|------------|-------------|------------|------------|-----------|
| <b>Mean</b> | <b>1338</b> | <b>1401</b> | <i>117</i> | <i>1506</i> | <i>239</i> | <i>-38</i> | <i>56</i> |
|-------------|-------------|-------------|------------|-------------|------------|------------|-----------|




660




**Figure 1: Composite radiosonde skew- $T$  Log- $P$  diagrams for the Amazon (a) wet season and (b) dry season launches (launched at 12 UTC, prior to convective cells). Shading represents the standard deviation of events. Temperature values are displayed in red and dewpoint temperature in green.**

665




**Figure 2: Diurnal cycle for cloud frequency as a function of height at the T3 site during our (a) wet and (b) dry season isolated cell events, respectively. (c) Wet (blue) and (d) dry (red) season diurnal cycle plots for the surface temperature and PBL height for the same convective events. Lines are event-mean values, while shading represents the standard deviation.**



675


**Figure 3: Diurnal cycle of the frequency of occurrence for select SIPAM radar reflectivity factor levels for the selected wet and dry season events.**



680

**Figure 4: Composite Amazon wet (blue) and dry (red) season cell tracking properties for dataset events. Time is normalized according to the difference between the first and last radar cell echoes exceeding the specified Z threshold. Lines represent the event-mean values, while shaded regions are the standard deviation. (a,b) Cell area according to a specified Z threshold of 25 dBZ, 35 dBZ.**


685



690 **Figure 5: Cell overpass cumulative vertical air velocity retrievals (CFADs) for wet season (left panels), dry season (middle panels), and wet minus dry difference fields (right panels). Upper CFAD panels include overpass retrievals having  $Z > 25$  dBZ, whereas lower panels are retrievals drawn from more intense  $Z > 35$  dBZ regions.**




695



**Figure 6: Cell overpass vertical air velocity retrievals (CFADs,  $Z > 25$  dBZ) for wet season (left panels), dry season (middle panels), and wet minus dry difference fields (right panels). Upper panels are cumulative CFADs for  $ETH < 10$  km, whereas lower panels are for the  $ETH > 10$  km retrievals.**

700

705



710

Figure 7: As in Figure 6, but for RWP retrievals having  $Z > 35$  dBZ.





715






Figure 8: Quartile life cycle breakdowns for overpass vertical air velocity retrievals (CFAD,  $Z > 25$  dBZ). Upper CFAD

720 panels are for the wet season events, whereas lower panels are for the dry season events.



725

**Figure 9: Histograms of vertical velocity from all SPM runs using (a) wet season radiosonde profiles and (b) dry season profiles. (c) A plot of wet season minus dry season histograms, where positive values are blue and negative values are red. (d) Same as (c), but with all skew-Ts given the average RH profile from all dry season cases above 2 km.**



730

**Figure 10: Profiles of negative buoyancy (x axis, m s<sup>-2</sup>) resulting from mixtures of updraft and environmental air, computed using the procedure described in Sect. 4.4. Blue profiles show the average over all wet season events, and red profiles show the average over all dry season events. Circles correspond to heights where the difference between the two curves was statistically significant.**

735