

# Moist available potential energy of the mean state of the atmosphere and the thermodynamic potential for warm conveyor belts and convection

Charles G. Gertler<sup>1</sup>, Paul A. O’Gorman<sup>1</sup>, and Stephan Pfahl<sup>2</sup>

<sup>1</sup>Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

<sup>2</sup>Institute of Meteorology, Freie Universität Berlin, 12165 Berlin, Germany

**Correspondence:** Charles G. Gertler (charles.gertler@gmail.com)

**Abstract.** Much of our understanding of the atmospheric circulation comes from relationships between aspects of the circulation and the mean state of the atmosphere. In particular, the concept of mean available potential energy (MAPE) has been used previously to relate the strength of the extratropical storm tracks to the zonal-mean temperature and humidity distributions. Here, we calculate for the first time the MAPE of the zonally varying (i.e. three dimensional) time-mean state of the atmosphere including the effects of latent heating. We further calculate a local MAPE by restricting the domain to an assumed eddy size, and we partition this local MAPE into convective and nonconvective components. Local convective MAPE maximizes in the subtropics and midlatitudes, in many cases in regions of the world that are known to have intense convection. Local nonconvective MAPE has a similar spatial pattern to the Eady growth rate, although local nonconvective MAPE has the advantage that it takes into account latent heating. Furthermore, the maximum potential ascent associated with local nonconvective MAPE is related to the frequency of warm conveyor belts (WCBs) which are ascending air streams in extratropical cyclones with large impacts on weather. This maximum potential ascent can be calculated based only on mean temperature and humidity, and WCBs tend to start in regions of high maximum potential ascent on a given day. These advances in the use of MAPE are expected to be helpful to connect changes in the mean state of the atmosphere, such as under global warming, to changes in important aspects of the extratropical circulation.

## 1 Introduction

The extratropical atmospheric circulation is dominated by baroclinic eddies that draw their energy from the mean available potential energy (MAPE) of the atmosphere. MAPE is the maximum amount of kinetic energy that could be released by reversible moist adiabatic motion starting from the mean state of the atmosphere, and it is calculated as the difference between the integrated enthalpy of the mean state and a hypothetical reference state (Lorenz, 1955, 1978). The reference state is the minimum-enthalpy state that can be attained through reversible moist adiabatic rearrangements. MAPE accounts for the competing effects of the static stability, meridional temperature gradient, and moisture content on the energy available to circulations including extratropical cyclones.

Previous work has shown that the overall strength of the extratropical storm tracks as measured by eddy kinetic energy (EKE) is related to MAPE calculated from the zonal-mean state of the atmosphere. EKE scales linearly with MAPE in simulations with idealized climate models over a wide range of climates (Schneider and Walker, 2008; O’Gorman and Schneider, 2008; O’Gorman, 2011), in simulations with comprehensive climate models of global warming (O’Gorman, 2010) and solar geoengineering (Gertler et al., 2020), over the observed seasonal cycle (O’Gorman, 2010), and in the observed weakening trend of the Northern Hemisphere summertime storm track over recent decades (Gertler and O’Gorman, 2019). In targeted idealized simulations with isolated thermal forcings, the linear relation between EKE and MAPE can break down, but it still performs better in predicting changes in EKE than changes in meridional temperature gradient or static stability alone (Yuval and Kaspi, 2017). Thus MAPE calculated from the zonal-mean state has proven useful to connect aspects of storm-track behavior to the mean temperature and humidity.

MAPE is most often calculated over a hemisphere and allowing for any type of circulation, but restrictions have also been imposed on the parcel rearrangements in order to isolate particular contributions to MAPE in the atmosphere and ocean. For the atmosphere, a nonconvective MAPE has been proposed which does not permit vertical reordering of air originating at a given latitude, thereby limiting the release of convective instability (O’Gorman, 2010; Gertler and O’Gorman, 2019; Gertler et al., 2020). Nonconvective MAPE is thought to be more strongly related to large-scale EKE because kinetic energy generated by convective instability may be rapidly dissipated locally before any upscale energy transfer occurs (O’Gorman, 2010). A corollary to nonconvective MAPE, convective MAPE is defined as the difference between MAPE and nonconvective MAPE and has been interpreted as the energy available for moist convection driven by larger-scale circulations such as extratropical cyclones or other baroclinic storms (Gertler and O’Gorman, 2019). For the ocean, an eddy-size constrained MAPE has been proposed in which no parcel may be displaced by a horizontal distance larger than the local eddy size (Su and Ingersoll, 2016). This eddy-size constraint takes into account that baroclinic eddies are generated by local parcel movement at the eddy length scale, and the spatial pattern of available potential energy density with the eddy-size constraint matches the spatial pattern of EKE in the Southern Ocean (Su and Ingersoll, 2016).

In this paper, we calculate for the first time MAPE based on the three-dimensional state of the atmosphere including moisture, and we calculate the nonconvective and convective contributions to this three-dimensional MAPE. We also want to consider a local version of MAPE, but when we applied the eddy-size constraint approach of Su and Ingersoll (2016) to the atmosphere, the global optimization was overwhelmed by ascent in the deep tropics, where enthalpy is generally largest. As an alternative approach, we introduce a local MAPE that is calculated on a subdomain chosen to have a horizontal extent similar to that of a typical extratropical cyclone. By restricting parcel motions to a subdomain, local MAPE imposes a horizontal constraint on parcel motions that is similar to the eddy-size constraint of Su and Ingersoll (2016) but doesn’t allow the ascent to be dominated by the tropics, and we investigate the properties of local MAPE including both nonconvective and convective local MAPE.

We find that local convective MAPE maximizes in subtropical and midlatitude regions, in many cases in the regions of the world with intense convection (Brooks et al., 2003; Zipser et al., 2006). This spatial distribution stands in contrast to the climatology of convective available potential energy (CAPE) which maximizes in the deep tropics (Riemann-Campe et al., 2009). Convective MAPE is calculated from the three-dimensional state of the atmosphere including horizontal gradients

of temperature and moisture. Thus it is physically plausible that local convective MAPE could have relevance to organized convective storms which are known to depend on vertical wind shear (which is proportional to horizontal temperature gradients through thermal wind balance) and moisture availability in addition to vertical measures of conditional instability like CAPE (Liu et al., 2020). To investigate this connection further, we make a comparison of the spatial distribution of convective MAPE to the occurrence of intense convection events as measured by the Global Precipitation Mission (GPM) core satellite (Hou et al., 2014) using a database described in Liu and Zipser (2015).

We furthermore explore how the parcel rearrangements involved in calculating local MAPE can provide insight into the dynamics of the extratropical storm tracks. We show here that these parcel rearrangements can be used to identify a thermodynamic potential for warm conveyor belts (WCBs). WCBs are strongly ascending air streams that originate in the atmospheric boundary layer in the warm sector of an extratropical cyclone and typically move polewards while ascending to levels near the tropopause. WCBs play a large role in cloud and precipitation formation and in transport of energy and air pollution (Browning, 1990; Stohl et al., 2002), and they are responsible for many extreme precipitation events in the extratropics (Pfahl et al., 2014). Most extratropical cyclones are associated with a WCB in Northern Hemisphere winter, while the fraction of cyclones associated with a WCB is much lower in Northern Hemisphere summer when the atmosphere is moist but EKE is weak, and cyclones near the coast of Antarctica are rarely associated with a WCB presumably because of low humidity in that region (Eckhardt et al., 2004). Here, we make use of the WCB climatology of Madonna et al. (2014) which identifies WCBs as Lagrangian trajectories of air parcels that occur in the vicinity of an extratropical cyclone and experience ascent of more than 600 hPa within 2 days. This is a relatively strict criterion that identifies strong ascent involving moist processes. We demonstrate that regions of frequent WCBs correspond to regions of large ascent between the mean state of the atmosphere and the reference state in the calculation of nonconvective local MAPE, thus making a link between WCB formation and the mean temperature and moisture of the atmosphere.

The paper proceeds as follows. In Section 2, we describe the reanalysis data, GPM database, and WCB climatology that are used in the paper. In Section 3, we describe the methods used to calculate the three-dimensional MAPE of the atmosphere, including the imposition of nonconvective and local restrictions. In Section 4, we present results for three-dimensional atmospheric MAPE with and without the nonconvective and local restrictions, including a comparison to the Eady growth rate and the occurrence of intense convection. In Section 5, we relate the parcel ascent in the calculation of nonconvective local MAPE to a WCB climatology and to daily WCB activity. In Section 6, we discuss the results, present brief conclusions, and propose avenues for future work.

## 2 Reanalysis data, GPM database, and WCB climatology

All MAPE calculations and the comparison to the Eady growth rate are based on the ERA-Interim reanalysis dataset, a global atmospheric reanalysis produced by the European Center for Medium-Range Weather Forecasts (Dee et al., 2011). We use temperature, relative humidity, wind, and surface pressure data. In most cases, we use monthly data at a horizontal grid spacing of  $0.75^\circ$  by  $0.75^\circ$  and at 37 pressure levels. However, for daily MAPE, we use six-hourly data at a horizontal grid spacing of

2.5° by 2.5° and at 37 pressure levels. Unless otherwise noted, all time means are over 1979-2018. Data at levels below the surface are removed.

To compare local convective MAPE to the regions of frequent intense convection, we use the GPM precipitation feature database introduced by Liu and Zipser (2015). Specifically we use precipitation features based on Ku radar retrievals over the  
95 years 2015-2021, and we plot features with high 40dBz echo top heights to identify the occurrence of intense convection.

In addition, we compare to the WCB climatology developed by Madonna et al. (2014) which has been extended in time since publication to cover the time period 1979-2018. WCB trajectories are identified in ERA-interim as strongly ascending air parcels (600 hPa in 2 days) that occur in the vicinity of an extratropical cyclone. A trajectory is deemed to be within the vicinity of an extratropical cyclone if it is within the horizontal area bounded by the outermost closed contour of sea level  
100 pressure for at least one 6-hourly time step during the 2-day period of strong ascent.

### 3 Calculation of MAPE

This section describes the methods we use to calculate MAPE of the three-dimensional time-mean state of the atmosphere. The main challenge is to calculate the parcel rearrangement leading to the reference state (i.e., the minimum enthalpy state), and the MAPE is then calculated for a given domain as the integrated enthalpy of the mean state minus that of the reference  
105 state. We also consider two constraints on the parcel rearrangement. The first constraint, used in calculating nonconvective MAPE, restricts the release of convective instability. The second constraint, used in calculating a local MAPE, restricts parcel movements to a local subdomain centered on a given surface location with a nominal radius in the horizontal chosen to represent a characteristic eddy length scale.

One challenge in using MAPE is that the reference state can be difficult to calculate accurately when atmospheric moisture is  
110 taken into account. For the case of a dry atmosphere, Lorenz (1955) derived an approximate formulation of the available potential energy using an assumption of small isentropic slope that is reasonably accurate in the free troposphere but problematic in the boundary layer. In this approximate dry formulation, the dominant contribution to MAPE is from the zonal and time mean rather than stationary eddies (Oort et al., 1989; Peixoto and Oort, 1992), and this has been used to justify the two-dimensional calculation of MAPE based on the zonal mean state. For the case of a moist atmosphere, latent heating needs to be accounted  
115 for in the calculation of MAPE (Lorenz, 1978), and the determination of the reference state using a parcel swapping algorithm is more difficult than in the case of a dry atmosphere (Lorenz, 1979; Randall and Wang, 1992). Important advances were made recently when the first exact calculations of MAPE were made using the Munkres algorithm for the ocean (Hieronymus and Nycander, 2015) and moist atmosphere (Stansifer et al., 2017). A MAPE calculation is said to be exact in this context when the exact MAPE is found for a given mean state and set of thermodynamic assumptions. Stansifer et al. (2017) also introduced  
120 an inexact divide-and-conquer algorithm which is fast and gives an accurate answer in most situations (Stansifer et al., 2017; Harris and Tailleux, 2018). In this paper, we frame the MAPE calculation following the formulation used for the ocean by Su and Ingersoll (2016) to take advantage of having many parcels at each pressure level. We solve for the reference state using a standard integer linear programming algorithm which is faster than Munkres for this problem and also exact. However, we



also make use of the inexact divide-and-conquer algorithm for nonconvective MAPE because the integer linear programming  
 125 algorithm does not always converge when the nonconvective restriction is included. What follows in this section is a discussion  
 of the exact approach to calculating MAPE used here. Readers interested only in the results should proceed to Section 4.

### 3.1 Dividing the atmosphere into equal-mass parcels


The global atmosphere is divided into parcels of equal mass to simplify the parcel rearrangement. In previous studies of MAPE  
 based on the zonal mean (which we will refer to as zonal-mean MAPE), the domain in latitude and pressure was divided  
 130 into equal mass parcels by using uniform spacing in pressure and a uniform spacing in a cosine-weighted latitude coordinate  
 which gives equal area divisions in the horizontal (Lorenz, 1979; O’Gorman, 2010; Gertler and O’Gorman, 2019; Gertler  
 et al., 2020). This approach has the disadvantage that the spacing in latitude increases towards the poles, and for the three-  
 dimensional calculation of MAPE it leads to strongly varying aspect ratios of grid cells that are particularly problematic for  
 the calculation of local MAPE at high latitudes. Instead, we generate a uniform grid on the sphere (i.e., with equal-area grid  
 135 cells in the horizontal) following the approach of Rosca (2010) as described in detail in Appendix A. An example of such a  
 horizontal grid is shown in Figure 1.

Some previous studies of zonal-mean MAPE have used a staggered grid which reduces the problem to one dimension in  
 pressure (Lorenz, 1979; O’Gorman, 2010; Gertler and O’Gorman, 2019), but it is equally simple and more efficient to employ  
 the same pressure coordinates in all columns (Su and Ingersoll, 2016), and that is the approach used here, with uniform spacing  
 140 in pressure to give equal mass parcels.

All results (except for sensitivity tests) are based on 15 evenly-spaced pressure levels in the vertical, and any parcels below  
 the surface are excluded. We generally use a global grid with 3200 surface grid cells (a  $40 \times 40$  grid in each hemisphere), except  
 in the case of non-convective and convective MAPE of extratropical domains ( $20^\circ$ - $90^\circ$ ) in Table 2 which are calculated using  
 a global grid with 800 surface grid cells (a  $20 \times 20$  grid in each hemisphere) due to computational expense.

### 145 3.2 Determining the reference (minimum-enthalpy) state

We consider  $n$  parcels of equal mass  $M$ , each with an initial temperature ( $T_i$ ), pressure ( $p_i$ ), and relative humidity ( $r_i$ ) where  
 $i = 1, 2, \dots, n$ . The parcels can occur at  $s$  distinct pressure levels,  $p_k^L$ , where  $k = 1, 2, \dots, s$  and the superscript  $L$  refers to level.  
 The number of parcels at each level is denoted by  $m_k$ , and  $m_k$  varies by level due to surface topography. Note that, by  
 construction,  $\sum_{k=1}^s m_k = n$ . A matrix is defined which contains the enthalpy values of all parcels adiabatically and reversibly  
 150 displaced to every pressure level,  $\mathbf{h} = [h_{i,k}] (i = 1, 2, \dots, n; k = 1, 2, \dots, s)$ , where  $h_{i,k}(T_i, p_i, r_i, p_k^L)$  is the specific enthalpy of  
 parcel  $i$  when it has been displaced to pressure  $p_k^L$ . In calculating the enthalpy of parcels at different pressures, we use the  
 saturation vapor pressure formulae over ice and liquid described in Simmons et al. (1999), but follow Wang and Randall  
 (1994) to merge ice and liquid phases. We then define the binary matrix  $\mathbf{x} = [x_{i,k}] (i = 1, 2, \dots, n; k = 1, 2, \dots, s)$  that maps the  
 mean state onto the reference state, where  $x_{i,k} = 0$  or 1, and  $x_{i,k} = 1$  indicates that parcel  $i$  is located at pressure  $p_k^L$  in the  
 155 reference state. A given rearrangement has a unique  $\mathbf{x}$  matrix, and the total enthalpy of the reference state is  $M \sum_{i=1}^n \sum_{k=1}^s h_{i,k} x_{i,k}$ .



**Figure 1.** Example of a uniform global grid created using the methodology introduced in Rosca (2010). There are 3200 grid points based on a  $40 \times 40$  grid in each hemisphere. The red circle is an example subdomain for the local MAPE calculation with a geodesic radius of 15 degrees (1670 kilometers) centered on the location marked with a blue asterisk and including all grid points colored in red. Except where noted, MAPE results in figures and tables are based on this horizontal grid at 15 vertical pressure levels.

The sum over each row of  $\mathbf{x}$  must equal one to ensure that each parcel is only assigned one location in the rearrangement state, and the sum over each column of  $\mathbf{x}$  must be the same as the number of parcels at the corresponding pressure level ( $m_k$ ). Therefore, to solve for the reference state, one finds  $\mathbf{x}$  which minimizes the total enthalpy given these conditions. Following Su and Ingersoll (2016), the problem can be stated as follows:

Given an  $n \times s$  matrix  $\mathbf{h}$ , find an  $n \times s$  matrix  $\mathbf{x}$  to minimize 
$$\sum_{i=1}^n \sum_{k=1}^s h_{i,k} x_{i,k}, \quad (1)$$
 where  $x_{i,k} = 0$  or 1, subject to  $\sum_{k=1}^s x_{i,k} = 1$  for any  $i$  and  $\sum_{i=1}^n x_{i,k} = m_k$  for any  $k$ .

Su and Ingersoll (2016) recognized that this problem is an example of a minimum-cost flow problem. It is also an integer linear programming problem, and we solve it here using a standard implementation of the dual-simplex algorithm (Koberstein, 2008). To frame the problem for integer linear programming, the  $\mathbf{h}$  and  $\mathbf{x}$  matrices are vectorized to  $\mathbf{h}_{vec}$  and  $\mathbf{x}_{vec}$ , and we minimize  $\mathbf{h}_{vec} \cdot \mathbf{x}_{vec}$  over  $\mathbf{x}_{vec}$ , subject to the constraint that  $\mathbf{A} \cdot \mathbf{x}_{vec} = \mathbf{b}$ ,  $\mathbf{A}$  and  $\mathbf{b}$  being a matrix and vector, respectively, that express the constraints in (1).

We also solve for the reference state using the divide-and-conquer algorithm described in Stansifer et al. (2017). While the divide-and-conquer algorithm is slower than the integer linear programming approach, the algorithm used to solve the integer linear programming approach occasionally does not converge under the nonconvective condition imposed on the parcel rearrangement described below. The divide-and-conquer algorithm is a recursive algorithm that builds a low-enthalpy reference

170 state by dividing the atmospheric domain into smaller subdomains. At each division, the pressure-derivative of enthalpy at the middle pressure of the subdomain is used to order parcels within that subdomain from top to bottom, and then the top and bottom halves are assigned to new subdomains.

Previous work has shown that the reference pressure for the minimum enthalpy state of the zonal-mean atmosphere for the moist MAPE calculation has a convective “bubble” at low latitudes that ascends discontinuously to the upper troposphere, 175 while for nonconvective MAPE the reference pressure is always continuous as a function of latitude and pressure, by definition (see, e.g., Figure 2 of Gertler and O’Gorman (2019)). A similar area of low-latitude ascent in the three-dimensional moist MAPE calculation can be seen in Figure 2, but now resolved in latitude and longitude.

### 3.3 Calculation of nonconvective and convective MAPE


For nonconvective MAPE, the reference state is calculated under the condition that parcels originating in a given column may 180 not change their vertical ordering, thereby restricting the release of convective instability. We calculate nonconvective MAPE using the divide-and-conquer algorithm, and this is done by modifying the algorithm such that at each step of the recursion, parcels from a given initial column may not change their vertical ordering in the parcel rearrangement of each subdomain. This ensures that parcels do not “leapfrog” in pressure over parcels from the same initial column, and the reference pressure is a monotonic function of pressure in a given column. The nonconvective MAPE is then defined as the integrated enthalpy of the 185 mean state minus that of the reference state restricted in this way.

The convective MAPE is defined as the difference between the MAPE and the nonconvective MAPE. When calculating convective MAPE, both the MAPE and the nonconvective MAPE are calculated using divide and conquer in the case of the extratropical domains ( $20^{\circ}$ - $90^{\circ}$  in latitude), but for local MAPE we use divide and conquer for nonconvective MAPE and integer linear programming for MAPE because divide and conquer sometimes gave negative values in the tropics (see section 190 4.3.2 for further discussion).

### 3.4 Calculation of local MAPE

We perform a series of MAPE calculations within local subdomains whose horizontal extent is set to a nominal length scale of an extratropical cyclone. This approach captures the energy available locally to be converted into kinetic energy in baroclinic storms such as extratropical cyclones, but note that it does not account for horizontal energy fluxes into and out of these systems 195 such as the fluxes that occur in downstream development (Chang, 1992; Chang et al., 2002), and thus we expect regions of high local MAPE to correspond to regions of large generation of EKE, but we do not expect local MAPE to capture the downstream structure of the storm track.

As before, the atmosphere is divided into parcels of equal mass using a uniform grid on the sphere and equal divisions in pressure. At each surface location, a subdomain is defined as including only parcels that fall within a horizontal distance less 200 than a specified eddy length scale of that location. The MAPE at a given surface location is defined as the MAPE over the subdomain centered in the horizontal on that location. An example of such a subdomain is shown in Figure 1. We consider the



**Figure 2.** Visualization of vertical parcel motions in global three-dimensional MAPE calculations for different seasons. Shown is the original pressure minus reference pressure for parcels originating at (a,b) 864 hPa and (c, d) 321 hPa for (a, c) DJF and (b, d) JJA. Positive values imply ascent from the mean state to the reference state, and negative values imply descent from the mean state to the reference state.

MAPE and its nonconvective and convective components in each subdomain, and record the reference pressures of parcels in each MAPE calculation.

Using local MAPE as an example, we illustrate the sensitivity of the MAPE calculation to key parameters and the choice of  
 205 algorithm in Appendix B.

## 4 Three-dimensional MAPE of the atmosphere

### 4.1 Extratropical and global MAPE

The seasonal MAPE over 1979-2018 is shown in Table 1 for northern and southern extratropical domains (20° -90° latitude) and for the global domain. We focus on the extratropical domains but give results for the global domain for reference. The results  
 210 in Table 1 are based on a global uniform grid with 3200 equal-area surface grid cells (a 40 × 40 grid in each hemisphere), and 15 evenly spaced pressure levels from 1000 hPa to 50 hPa, with parcels below the surface removed. The results are generally similar beyond a resolution of about 800 surface grid cells and 10 pressure levels. The grid cells in this global grid that fall

**Table 1.** Extratropical and global MAPE values in  $\text{J kg}^{-1}$  based on the time mean over 1979-2018 for different seasons. Results are given for three-dimensional MAPE, zonal-mean MAPE, and their difference. Zonal-mean MAPE is calculated by replacing all values for temperature and humidity at a given latitude and pressure with the zonal-mean values at that latitude and pressure. The difference is calculated as the three-dimensional MAPE minus zonal-mean MAPE as a percentage of the three-dimensional MAPE. All results in this table are calculated with integer linear programming.

Season	Domain	MAPE ( $\text{J kg}^{-1}$ )	Zonal-Mean MAPE ( $\text{J kg}^{-1}$ )	Difference (%)
DJF	20°-90° N	373.6	335.3	10.2
	20°-90° S	301.2	284.1	5.6
	90° S -90° N	480.4	452.8	5.7
MAM	20°-90° N	303.0	285.5	5.8
	20°-90° S	333.0	324.7	2.5
	90° S -90° N	450.6	435.9	3.3
JJA	20°-90° N	215.2	166.8	22.5
	20°-90° S	361.0	351.1	2.8
	90° S -90° N	433.8	407.6	6.1
SON	20°-90° N	324.8	305.7	5.9
	20°-90° S	341.8	330.1	3.4
	90° S -90° N	443.2	423.3	4.5

within the extratropical domains are used to calculate the extratropical MAPE values. A zonal-mean MAPE on the same domain is also calculated by replacing all the temperature and humidity values with their zonal mean values at that latitude and pressure. We use this approach to calculating zonal-mean MAPE in which the only inputs are zonal-mean temperature and humidity but the calculation remains three-dimensional because it allows for a fair comparison with fully three-dimensional MAPE in the presence of topography that varies in longitude.

MAPE is larger in winter than summer (Table 1), implying that the effect of the stronger meridional temperature gradient in winter outweighs the lower specific humidity. The three-dimensional MAPE is always larger than the zonal-mean MAPE, and this is as expected because three-dimensional MAPE includes both zonal and meridional gradients of temperature and moisture. The difference between the three-dimensional MAPE and the zonal-mean MAPE, which can be interpreted as the contribution from zonal asymmetries, is always largest in the Northern Hemisphere extratropics, with particularly large contributions in DJF and JJA. The contribution from zonal asymmetries in the Northern Hemisphere extratropics in JJA of 22.5% is substantially larger than what was found previously for dry MAPE for the full NH (7.5% based on Table 14.1 of Peixoto and Oort (1992)).

Figure 2 illustrates the vertical motion of parcels in the global three-dimensional MAPE calculations for DJF and JJA, showing tropical and subtropical ascent and extratropical descent. Tropical and subtropical ascent is strongest at low levels in the summer hemisphere and over land, and extratropical descent is strongest at the upper levels in the winter hemisphere.

**Table 2.** Extratropical three-dimensional MAPE, nonconvective MAPE, and convective MAPE in  $\text{J kg}^{-1}$  over 1979-2018. The convective fraction is shown in parentheses as a percentage of full MAPE. Results in this table are based on a lower-resolution horizontal grid with 1800 surface grid cells (a  $30 \times 30$  grid in each hemisphere), and the (full) MAPE values differ slightly from those in Table 1 because of the lower resolution and because divide and conquer is used here rather than linear integer programming in Table 1.

Season	Domain	MAPE ( $\text{J kg}^{-1}$ )	Nonconvective MAPE ( $\text{J kg}^{-1}$ )	Convective MAPE ( $\text{J kg}^{-1}$ )
DJF	20°-90° N	372.5	366.9	5.7 (1.5%)
	20°-90° S	300.7	285.2	15.4 (5.1%)
MAM	20°-90° N	303.8	296.1	7.7 (2.5%)
	20°-90° S	333.2	318.8	14.4 (4.3%)
JJA	20°-90° N	216.7	197.3	19.4 (9.0%)
	20°-90° S	359.1	353.4	5.7 (1.6%)
SON	20°-90° N	326.1	309.8	16.4 (5.0%)
	20°-90° S	341.4	335.8	5.6 (1.7%)


Nonconvective MAPE and convective MAPE, calculated using divide and conquer, are presented in Table 2, and these are calculated on a lower-resolution grid ( $30 \times 30$  grid in the horizontal in each hemisphere) due to the greater computational requirement when the non-convective constraint is imposed and when using the divide-and-conquer algorithm. Using this grid and the divide-and-conquer algorithm for full three-dimensional MAPE gives very similar values to those from the higher-resolution exact calculation using integer linear programming (compare the extratropical MAPE values in Tables 1 and 2).

Nonconvective MAPE restricts vertical reordering of parcels starting in a column, and thus it is always equal or smaller in magnitude to the MAPE. However, MAPE and nonconvective MAPE are similar in magnitude with the largest values in Northern Hemisphere DJF and smallest values in Northern Hemisphere JJA. Convective MAPE is largest in Northern Hemisphere JJA, when it is also the largest percentage of overall MAPE, and smallest in Northern Hemisphere DJF and Southern Hemisphere JJA and SON. The relative breakdown between convective and nonconvective MAPE across seasons and hemispheres are similar for zonal-mean MAPE as for three-dimensional MAPE (not shown).

## 4.2 Local MAPE

Next the local restriction is imposed on the three-dimensional MAPE calculation. We use the same global grid as for the full three-dimensional MAPE calculation with a  $40 \times 40$  grid in the horizontal in each hemisphere. Each subdomain for the local MAPE is centered in the horizontal at one surface grid cell and has a geodesic radius of 15 degrees (1670 kilometers). This radius was chosen as representative of the length scale of midlatitude eddies.


The local MAPE based on the climatological mean over 1979-2018 is shown in Figure 3. Local MAPE represents the energy available for local parcel rearrangement at the eddy length scale, and large values correspond to extratropical regions with strong baroclinic instability. In the Northern Hemisphere, the eastern sides of continents and the western side of ocean basins



**Figure 3.** Local MAPE based on the climatological mean over 1979-2018 for (a) DJF and (b) JJA. MAPE is calculated on subdomains of geodesic radius 15 degrees (1670 kilometers) centered in the horizontal on 3200 surface grid cells.

are particular hotspots in winter due to strong horizontal gradients of temperature in these regions, while local MAPE is higher in the interior of continents in summer possibly due to a lower static stability over land. The Northern Hemisphere has the highest values in the Western Pacific in winter, while the Southern Hemisphere shows more consistently high values throughout the midlatitudes in winter. By restricting to subdomains that are roughly of the size of a typical extratropical cyclone, we have effectively introduced more dynamics into the calculation of MAPE and reduced its value to be more realizable and similar in magnitude to kinetic energy. For example, in DJF, the local MAPE averaged over the northern extratropics is  $96.8 \text{ J kg}^{-1}$  as compared to  $373.6 \text{ J kg}^{-1}$  for MAPE calculated for the northern extratropics without the local constraint.

The reference pressure distribution for each subdomain may be examined for insight into the characteristics of the local parcel rearrangements. In particular, the difference between the original pressure and reference pressure of a parcel may be




**Figure 4.** Maximum potential ascent in the local MAPE calculation based on the climatological mean over 1979-2018 for (a) DJF and (b) JJA. MAPE is calculated as in Figure 3, and the maximum potential ascent is calculated at each surface location as the maximum difference between pressure and reference pressure for any parcel in the column at that location over all the MAPE calculations that include that location.

interpreted as the ascent or descent that gives rise to the potentially largest energy release on the eddy length scale. Parcels in the column above a given surface location are included in multiple MAPE calculations for subdomains that overlap with that surface location. Therefore, we calculate the maximum ascent of any parcel over all subdomains that include that surface location, and this can be thought of as the maximum potential ascent due to baroclinic instability and convection at that location.

260 Figure 4 shows the maximum potential ascent in each season, with maximum values near the equator and high values over midlatitude land in summer. Much of this is convective ascent, and the nonconvective and convective components of local MAPE are next examined separately.





**Figure 5.** Nonconvective local MAPE (color shading) and Eady growth rate averaged in the vertical over 500-750 hPa (black contours of 0.5 and 0.7 day<sup>-1</sup>) based on the climatological mean over 1979-2018 in (a) DJF and (b) JJA.

### 4.3 Nonconvective and convective local MAPE

#### 4.3.1 Nonconvective local MAPE

265 The nonconvective local MAPE is also calculated using the same grid and subdomains as for the full local MAPE, but using the divide-and-conquer algorithm rather than integer linear programming. Nonconvective local MAPE must be smaller or equal to local MAPE, but Figure 5 shows that the overall pattern and magnitude of nonconvective local MAPE are similar to those of local MAPE in Figure 3.

Nonconvective local MAPE has high values in the entrance regions of the storm tracks, and we interpret it as a measure of  
 270 the available energy for the generation of large-scale eddies through moist baroclinic instability. Dry MAPE scales roughly

as the vertical integral of the square of the Eady growth rate (O’Gorman and Schneider, 2008), and Figure 5 shows that the nonconvective local MAPE has a similar spatial pattern to the Eady growth rate averaged in the vertical over 500-750hPa. The (maximum) Eady growth rate,  $\sigma$ , at a given vertical level is calculated as

$$\sigma = 0.31 \frac{f}{N} \left\| \frac{\partial \mathbf{u}_h}{\partial z} \right\|, \quad (2)$$


275 where  $f$  is the Coriolis parameter,  $N$  is the buoyancy frequency,  $\mathbf{u}_h$  is the horizontal wind, and  $z$  is height. Local MAPE is complementary to the Eady growth rate as a measure of local baroclinic instability in that (1) it provides a measure of energy that can be compared to EKE in terms of fractional changes over the seasonal cycle or for climate change, and (2) it includes the effects of moisture which can be important in terms of the response to climate change (although we note the Eady growth rate could also be modified to account for moisture through an effective static stability (O’Gorman, 2011)).

### 280 4.3.2 Convective local MAPE

Figure 6 shows convective local MAPE which is defined as the difference between full local MAPE and nonconvective local MAPE. The full local MAPE is calculated using the integer linear programming approach because the divide-and-conquer algorithm gives some negative values near the equator. Negative values of MAPE are physically impossible, but they can sometimes occur with the divide-and-conquer algorithm for conditionally unstable atmospheres because the divide-and-conquer algorithm is not exact (Harris and Tailleux, 2018). Thus, integer linear programming is used to calculate the full local MAPE while divide and conquer is used to calculate the nonconvective local MAPE, and we caution that this mixing of algorithms could affect the results for convective local MAPE in localized regions near the equator (see Appendix B, section B3).

Figure 6 shows that convective MAPE has large values in certain regions in the subtropics and middle latitudes, including over Argentina, Southern Africa and Australia in DJF, and the central and southeastern United states in JJA. There are also high values over some subtropical ocean regions. This spatial distribution is in contrast to the climatology of convective available potential energy (CAPE) which maximizes in the deep tropics (Riemann-Campe et al., 2009). Convective MAPE behaves differently from CAPE because convective MAPE takes account of horizontal temperature and moisture gradients in addition to vertical instability, and these horizontal gradients are small near the equator. Note also that convective MAPE values are smaller than typical CAPE values because MAPE is normalized by the full mass of the atmosphere whereas CAPE is normalized by only the mass of the air that is rising. Unlike the convective MAPE, the maximum potential ascent in the calculation of local MAPE does have large values near the equator (Figure 4). However, large displacements in the vertical do not imply large generation of kinetic energy if the buoyancy is low, explaining how convective MAPE values can remain small in that region.

Interestingly, there is some correspondence of the regions of large convective MAPE with regions of intense organized convection such as Argentina in DJF and parts of the United States in JJA (Brooks et al., 2003; Zipser et al., 2006). Gertler and O’Gorman (2019) previously interpreted convective MAPE as the energy available for convection associated with midlatitude cyclones, but the spatial distribution in Figure 6 suggests it may also be relevant for organized convective storms in baroclinic environments. This is physically plausible because organized convective systems are influenced by horizontal gradients of temperature and moisture as well as their vertical gradients (noting that vertical shear is related to horizontal temperature



**Figure 6.** Convective local MAPE based on the climatological mean (color shading) over 1979-2018 and occurrence of intense convection events (black dots) over 2015-2021 for (a) DJF and (b) JJA. Convective MAPE is calculated as the difference in MAPE and nonconvective MAPE in the local subdomains. Intense convection events are identified using GPM as precipitation features with a maximum 40dBZ echo top height that exceeds 14km over land and 9km over ocean (see text for further details).

gradients by thermal wind balance) and these factors are taken into account in convective MAPE which depends on the three-dimensional state of the atmosphere.

To explore this correspondence further, we make a comparison of the spatial distribution of convective MAPE to the occurrence of intense convection events from GPM over the years 2015-2021 in a database described in Liu and Zipser (2015). Figure 6 shows all precipitation features with a maximum 40dBZ echo top height from the Ku band radar that exceeds 14km over land and 9km over ocean. The reason to use different thresholds over land and ocean is that the convective MAPE does not capture the difference in strength of convection between land and ocean which is possibly due to aerosols and the diurnal

cycle (Pan et al., 2022), and we want to be able to show events over both land and ocean to compare to the distribution of convective MAPE. Lower thresholds such as 12km over land and 8km over ocean give a higher number of events but a similar spatial distribution. Note that the sampling by the GPM core satellite increases towards the poleward limits of the orbit at 65S and 65N, but plotting the events should be sufficient as a simple indicator of where intense convection occurs, and the seasonal and spatial distribution of intense convection is similar to what has been found in previous studies (Zipser et al., 2006; Liu and Zipser, 2015). The comparison with convective MAPE shows some similarities including regions of intense convection and high convective MAPE over Argentina and South Africa in DJF, North America and the Bay of Bengal in JJA, and over the subtropical oceans in both seasons. One notable exception is intense convection over central Africa in a region with little convective MAPE. In addition, there are some features that are displaced poleward in convective MAPE compared to the intense convection (e.g., over West Africa in DJF) which may be because local convective MAPE is plotted at the center of the local domain whereas the convective rising air in the MAPE calculation tends to originate more equatorward of this location.

Clearly, intense convection is affected by other variables in addition to the thermodynamic state of the atmosphere (e.g., aerosols, lifting by topography, surface heat and moisture fluxes), but convective MAPE is interesting as a candidate measure for the energy available to organized convective systems, and this could be investigated in future work using, for example, idealized simulations.

## 5 Application to Warm Conveyor Belts (WCBs)

### 5.1 Climatological maximum potential ascent and WCBs

Maximum potential ascent determined by the local MAPE framework can provide insight into WCB activity given that WCBs are instances of strong atmospheric ascent associated with extratropical cyclones. Ascent in WCBs is strongly diabatic and thus it is important that we are using a moist MAPE that accounts for latent heating. High-resolution simulations and satellite observations suggest that WCBs trajectories are mostly slantwise but do include instances of more vertical convective ascent embedded in the slower slantwise ascent (Rasp et al., 2016; Oertel et al., 2019, 2020). However, we focus here on ascent in calculations of nonconvective MAPE (excluding convective ascent) rather than full MAPE because we find it gives a stronger connection with WCBs, and this is presumably because the full MAPE calculation also includes convective ascent not associated with extratropical cyclones (e.g., at low latitudes).

Figure 7 shows the maximum potential ascent for the local nonconvective MAPE described in Section 4.3.1. The strong ascent in equatorial regions for the full local MAPE in Figure 4 is absent for the nonconvective local MAPE in Figure 7, suggesting that the equatorial ascent is indeed dominated by convection. By contrast, maximum potential ascent for nonconvective MAPE reaches local maxima in the extratropics.


The maximum potential ascent for nonconvective local MAPE is also compared to the climatological frequency of WCB starting points in Figure 7. WCB starting points are identified as the first point of the 48-hour ascent in the climatology of Madonna et al. (2014), and their frequency is plotted as the percentage of 6-hourly timesteps in which a WCB originates at a given location. The maximum potential ascent based on the climatological mean temperature and humidity is comparable

in magnitude to, but somewhat smaller than, the WCB ascent threshold of 600 hPa, but larger values of maximum potential ascent can occur on shorter timescales as shown below. The regions of higher maximum potential ascent in the nonconvective MAPE of the climatological atmosphere are very similar to the regions of heightened WCB activity, albeit with some regions of mismatch in the Western Pacific. Thus, the moist MAPE framework of Lorenz (1978, 1979) applied locally and excluding convection can identify regions of WCB activity based only on climatological mean temperature and humidity. Furthermore, the maximum potential ascent is providing distinct information from the Eady growth rate or non-convective MAPE, since the spatial distributions and seasonal cycles of non-convective MAPE and Eady growth rate (Figure 5) have differences from those of maximum potential ascent and WCB starting points (Figure 7). For example these distributions are quite different in the Southern Hemisphere in DJF.

## 5.2 Daily maximum potential ascent and WCBs


The ability of the local nonconvective MAPE of the climatological atmosphere to identify regions of climatological WCB activity raises questions about the ability of this approach to predict WCB activity from temperature and humidity on shorter timescales. To investigate this, the local nonconvective MAPE and reference pressures are calculated using mean temperature and humidity values on individual days (the average of four six-hourly fields), and compared to the WCB activity over the same days. We refer to this as MAPE since it is based on the mean of temperature and humidity fields over a day, but it could also be thought of as simply the daily available potential energy. Only one randomly-chosen year, 1985, is considered due to computational expense. Figure 8 shows results for example days near the beginning of the four seasons. Grid cells with one or more WCB starting points (it is not uncommon for multiple WCBs to start at the same location on a given day) tend to fall in regions of large maximum potential ascent. However, large maximum potential ascent does not always lead to WCB starting points on a given day because WCB activity also requires the presence of an extratropical cyclone on that day. In this way, this concept is similar to the maximum potential intensity for tropical cyclones introduced by Emanuel (1988), as it demonstrates the thermodynamic potential for an event which may not be realized at all times. There is considerable spatial structure to the maximum potential ascent on a given day suggesting that extratropical cyclones themselves may be contributing to the spatial structure at this timescale.

To illustrate the statistical relationship between maximum potential ascent and WCB formation, the maximum potential ascent in daily local nonconvective MAPE calculations is calculated for each grid cell and each day in 1985, and these values are then linearly interpolated to the same grid as the WCB climatology. To properly compare to the climatology, which only considers cyclones poleward of 20 degrees, we also only consider grid cells poleward of 20 degrees. The median of the maximum potential ascent over grid cells and days where at least one WCB starting point occurs is 570 hPa, which is substantially higher than the overall median of 330hPa and very close to the ascent threshold of 600hPa used in the WCB climatology. Figure 9a shows that the probability density functions of maximum potential ascent are quite distinct for cases with and without WCB starting points.. Figure 9b shows the probability of WCB formation for grid cells and days that exceed a given level of maximum potential ascent. 10% of grid cells with maximum potential ascent above 600 hPa have at least one WCB starting point, and 25% of grid cells with maximum potential ascent above 720 hPa have at least one WCB starting point. These percentages



**Figure 7.** Maximum potential ascent in local nonconvective MAPE based on the climatological mean over 1979-2018 (color shading) and WCB starting point frequency over 1979-2018 (black contours of 1% and 2%) for (a) DJF and (b) JJA. Nonconvective MAPE is calculated as in Figure 5. Maximum potential ascent is calculated at each surface location as the maximum of pressure minus reference pressure for any parcel in the column at that surface location over all the nonconvective MAPE calculations that include that location. The WCB starting point frequency is shown as the percentage of six-hourly time steps in which a WCB begins at a given location in the climatology of Madonna et al. (2014)

may be compared with the baseline of 2% of grid cells over the extratropics ( $20^{\circ}$  to  $90^{\circ}$ ) that have at least one WCB starting point. Furthermore, the probability of WCB formation increases monotonically and very nonlinearly with maximum potential ascent. The probability of WCB formation based on high maximum potential ascent is even higher in storm track regions. For example, in a region of the Pacific from  $160^{\circ}$  to  $190^{\circ}$  in longitude and  $40^{\circ}$  to  $60^{\circ}$  N in latitude over which extratropical cyclones occur with roughly 30% frequency in the climatology of Wernli and Schwierz (2006), roughly 50% of days and grid cells in




**Figure 8.** Maximum potential ascent in the local nonconvective MAPE (color shading) and WCB starting points (black contours) for example days in the year 1985. The black contours surround areas of at least one WCB starting point on the given day in the dataset of Madonna et al. (2014).

1985 with maximum potential ascent exceeding 700 hPa have at least one WCB starting point, compared to 8% probability of WCB starting points for all days and grid cells in that region in 1985.

385 The results above indicate that maximum potential ascent from daily local nonconvective MAPE calculations can predict with some skill the formation of WCBs using only the regional temperature and humidity fields. In conjunction with information on cyclone occurrence, this maximum potential ascent may have value in understanding and forecasting of WCB formation, as well as understanding of the potential effects of climate change on WCBs.

## 6 Discussion and Conclusions

390 We have calculated the three-dimensional moist MAPE of the atmosphere for the first time, and we have also introduced a new approach to calculating a local MAPE on the characteristic length scale of extratropical cyclones. Combined with a decomposition of MAPE into convective and nonconvective components, we show that MAPE provides a number of useful connections between the mean state of the atmosphere and the dynamics of extratropical cyclones, warm conveyor belts, and possibly also convective storms in baroclinic environments.



**Figure 9.** Statistical relation of WCB starting points to maximum potential ascent from daily local nonconvective MAPE calculations in the year 1985. (a) Probability density functions of the maximum potential ascent for all grid cells and days (blue) and for grid cells and days restricted to incidences of at least one WCB starting point (red). (b) Probability of at least one WCB starting point for grid cells and days above a given maximum potential ascent. For comparison, the probability of at least one WCB starting point for all days and grid cells in the extratropics (20° to 90°) is 2% (dotted line).

395 By including gradients of temperature and moisture in the zonal direction and by better accounting for topography which is difficult to deal with in a zonal-mean calculation, three-dimensional moist MAPE provides a more accurate accounting of the energy available to circulations as compared to previous calculations based on the zonal mean. Local three-dimensional MAPE identifies regions where baroclinic eddies grow, broadly consistent with the Eady growth rate, but including the effects of latent heating and giving an energetic measure which could be useful for connecting to eddy kinetic energy. Differences between pressure and reference pressure in the local MAPE calculation characterize the thermodynamic potential for ascent. The maximum potential ascent is largest in the tropics for the full MAPE but in midlatitudes for the non-convective MAPE. The convective component of the local MAPE (i.e. the difference in the full and nonconvective MAPE) maximizes in the subtropics and midlatitudes (unlike CAPE which maximizes near the equator) and has some similarities in spatial and seasonal pattern to the occurrence of intense convection. Previous work has empirically connected organized convection to combinations of vertical shear and CAPE (Brooks et al., 2003) and it is possible that convective MAPE could provide a unified energetic measure relevant for organized convection in a baroclinic environment and this could be investigated in future work. Future work could also investigate the extent to which there are connections between changes in EKE and local MAPE and changes in organized convective storms and convective MAPE under climate-change scenarios.

405

We have also demonstrated that the maximum potential ascent in the calculation of nonconvective local MAPE is related to WCB formation. This is a new connection between the mean state of the atmosphere and important high-frequency weather events. In particular, WCBs are crucial components of extratropical cyclones and every-day weather in the midlatitudes, and

410



they play an important role in cloud formation, precipitation, and transport of air pollution (Browning, 1990; Stohl et al., 2002; Pfahl et al., 2014). The maximum potential ascent based on the climatological mean temperature and humidity identifies regions of heightened WCB formation, and this link could be used to better understand the geographical and seasonal distribution of WCBs. Furthermore, regions of high maximum potential ascent on individual days are more likely to form WCBs than other regions. It is possible that the link between maximum potential ascent and WCB formation on the daily timescale could be helpful for forecasting of WCBs, but establishing this would require a detailed study of statistics such as the conditional probability of the occurrence of a WCB given the presence of both a cyclone and strong potential ascent versus only the presence of a cyclone. In addition there may be a useful link between maximum potential ascent and WCB formation on longer subseasonal timescales. Two other areas of future work related to maximum potential ascent seem promising. First, it could be investigated whether there is a simple analytical approximation for the maximum potential ascent as a function of mean temperature and humidity fields. Ideally this would be comparably simple to the expression for the Eady growth rate but capable of giving the spatial and seasonal distribution of the maximum potential ascent. Second, it would be interesting to investigate changes in maximum potential ascent in simulations of climate change to provide a link between mean warming and moistening of the atmosphere and changes in WCB formation.

*Code and data availability.* A directory including all analysis code used in this study is published online at <https://doi.org/10.5281/zenodo.5826260>. ERA interim data can be access from ECMWF at <https://apps.ecmwf.int/datasets/data/interim-full-daily/levtype=sfc/>. GPM precipitation feature database can be accessed from [atmos.tamucc.edu/trmm/data/gpm/level\\_2/kurpf/](https://atmos.tamucc.edu/trmm/data/gpm/level_2/kurpf/).

## Appendix A: Uniform grid generation

We create a uniform grid on the sphere following the methodology of Rosca (2010). The strategy used is to create an equally spaced grid on a square, and then to project it onto each hemisphere by using area-preserving bijections that map from a square to a disk and then from the disk to a hemisphere. The square is of edge  $e$ , and the sphere of radius  $r$ , where  $e = r\sqrt{2\pi}$ . Further details and proof of this methodology can be found in Rosca (2010).

The grid on the square is created using an identical set of Cartesian coordinates in  $x$  and  $y$ . For convenience,  $L$  is defined such that  $e = 2L$ . The number of grid cells in each direction is denoted  $N$ . The spacing,  $d$ , is then  $e/N$ . The grid cells are centered at grid points starting at  $-L + d/2$  and extending with a spacing of  $d$  to  $L - d/2$ . To project this grid onto the Southern Hemisphere and preserve area, we then apply the following formulas to project  $(a, b)$ , a set of  $x$  and  $y$  coordinates, into  $(A, B, C)$ , a set of  $x$ ,  $y$ , and  $z$  coordinates.

1. For  $0 \leq |b| \leq |a| \leq L$ ,

$$(A, B, C) = \left( \frac{2a}{\pi} \sqrt{\pi - \frac{a^2}{r^2}} \cos \frac{b\pi}{4a}, \frac{2a}{\pi} \sqrt{\pi - \frac{a^2}{r^2}} \sin \frac{b\pi}{4a}, \frac{2a^2}{\pi r} - r \right); \quad (\text{A1})$$

2. For  $0 \leq |a| \leq |b| \leq L$ ,

$$(A, B, C) = \left( \frac{2b}{\pi} \sqrt{\pi - \frac{b^2}{r^2}} \sin \frac{a\pi}{4b}, \frac{2b}{\pi} \sqrt{\pi - \frac{b^2}{r^2}} \cos \frac{a\pi}{4b}, \frac{2b^2}{\pi r} - r \right). \quad (\text{A2})$$

For the Northern Hemisphere, the formulas are the same as in Equations A1 and A2, except that  $C$  is of opposite sign. These Cartesian coordinates can be easily transformed to latitude and longitude on the sphere of radius  $r$ .

445 Figure 1 shows an example of such a grid which we employ in the analysis and which has  $40 \times 40$  grid cells in each hemisphere.

## Appendix B: Sensitivities of calculation of local MAPE

We focus on the calculation of local MAPE as an example, and we describe its sensitivity to resolution, assumed eddy size and calculation method.

### 450 B1 Sensitivity to resolution

Low horizontal resolutions are not sufficient to capture the characteristic patterns of local MAPE (Figure B1a) or maximum potential ascent (Figure B2a). However, at resolutions at and beyond roughly 1800 surface grid cells (a  $30 \times 30$  grid in each hemisphere) and at and beyond ten vertical levels, the results converge (Figures B1b,c and B2b,c).

### B2 Sensitivity to the eddy size


455 Unlike for horizontal resolution, as the radius of the subdomains used to calculate local MAPE increases, the results do not converge. Indeed, one would expect the magnitude of MAPE to increase as the subdomain increases in size, consistent with the scaling of dry zonal-mean MAPE per unit mass with the square of the length of the domain considered (O’Gorman and Schneider, 2008). However, the spatial pattern of local MAPE does appear to converge for geodesic radii greater than roughly 10 degrees even though the MAPE value continues to increase as the radius increase (Figure B3). Interestingly, the spatial  
460 pattern of maximum potential ascent seems to converge at a larger radius, with noticeable differences between the 10 degree and 15 degree calculations, but similar spatial patterns of maximum potential ascent for the 15 degree and 20 degree calculations (Figure B4).

### B3 Sensitivity to the method of calculation


In most of the atmosphere, and in particular in nearly all of the extratropics, the divide-and-conquer algorithm and integer linear  
465 programming give nearly identical values of local MAPE and maximum potential ascent (Figure B5 and Figure B6). However, in some regions in the tropics (where the MAPE is small), the divide-and-conquer algorithm can be very inaccurate as a fraction of the exact MAPE given by integer linear programming. Indeed, the divide-and-conquer algorithm sometimes even returns a negative value for MAPE (black contours in Figure B5), which is physically impossible but has been previously shown to

sometimes occur for certain conditionally unstable columns with the divide-and-conquer algorithm (Harris and Tailleux, 2018).


470 Figure B5 demonstrates this issue in the tropics for DJF and differences are similar across all seasons. Similarly, in some areas in the tropics, the exact calculation predicts large maximum potential ascent that is not captured by the divide-and-conquer algorithm (Figure B6). These inaccuracies of the divide-and-conquer algorithm do not affect the main conclusions of this paper.




**Figure B1.** Local MAPE based on the DJF climatological mean over 1979-2018 for grids of varying horizontal resolution: (a) 200 surface grid cells (a  $10 \times 10$  grid in each hemisphere), (b) 1800 surface grid cells (a  $30 \times 30$  grid in each hemisphere), and (c) 5000 surface grid cells (a  $50 \times 50$  grid in each hemisphere). All results shown in this figure use 15 pressure levels and subdomains of geodesic radius 15 degrees (1670 kilometers).




**Figure B2.** Maximum potential ascent in local MAPE based on the DJF climatological mean over 1979-2018 for grids of varying horizontal resolution: (a) 200 surface grid cells (a  $10 \times 10$  grid in each hemisphere), (b) 1800 surface grid cells (a  $30 \times 30$  grid in each hemisphere), and (c) 5000 surface grid cells (a  $50 \times 50$  grid in each hemisphere). All results shown in this figure use 15 pressure levels and subdomains of geodesic radius 15 degrees (1670 kilometers).



**Figure B3.** Local MAPE based on the DJF climatological mean over 1979-2018 using subdomains of varying geodesic radii: (a) 5 degrees (560 km), (b) 10 degrees (1110 km), (c) 15 degrees (1670 km), and (d) 20 degrees (2230 km). All results in this figure use 1800 surface grid cells (a  $30 \times 30$  grid in each hemisphere) and 15 pressure levels.




**Figure B4.** Maximum potential ascent in local MAPE based on the DJF climatological mean over 1979-2018 using subdomains of varying geodesic radii: (a) 5 degrees (560 km), (b) 10 degrees (1110 km), (c) 15 degrees (1670 km), and (d) 20 degrees (2230 km). All results in this figure use 1800 surface grid cells (a  $30 \times 30$  grid in each hemisphere) and 15 pressure levels.



**Figure B5.** Sensitivity to calculation method for the local MAPE based on the DJF climatological mean over 1979-2018. Shown is MAPE calculated using (a) the exact integer linear programming approach, (b) the approximate divide-and-conquer algorithm, and (c) the difference reported as integer linear programming minus divide and conquer. Black contours in (c) indicate regions where the divide-and-conquer algorithm gives negative values.





**Figure B6.** Sensitivity to calculation method for the maximum potential ascent in local MAPE based on the DJF climatological mean over 1979-2018. Shown is maximum potential ascent calculated using (a) the exact integer linear programming approach, (b) the approximate divide-and-conquer algorithm, and (c) the difference reported as integer linear programming minus divide and conquer.

*Author contributions.* C.G.G., P.A.O’G., and S.P. conceived of and designed the research and contributed to the editing of the text. C.G.G.  
475 performed the research, produced all figures, and drafted the paper.

*Competing interests.* S.P. serves as executive editor of Weather and Climate Dynamics.

*Acknowledgements.* We are grateful to H. Wernli, for helpful discussions and for first pointing out the possibility of a connection between WCBs and parcel rearrangements in MAPE. We are also grateful to R.G. Prinn for helpful insight. We thank L. Back, R. van der Drift, M. Velez Pardo, and two anonymous reviewers for helpful comments. C.G.G. was supported by the Industry and Foundation sponsors of the  
480 MIT Joint Program on the Science and Policy of Global Change, NASA grant NNX16AC98G to MIT, and the National Science Foundation Graduate Research Fellowship Program under NSF Grant 1122374. P.A.O’G. acknowledges support from NSF Grant AGS 1749986. S.P. acknowledges support from Deutsche Forschungsgemeinschaft (DFG) through grant CRC 1114 “Scaling Cascades in Complex Systems”, Project Number 235221301, Project (C06).

## References

- 485 Brooks, H. E., Lee, J. W., and Craven, J. P.: The spatial distribution of severe thunderstorm and tornado environments from global reanalysis data, *Atmospheric Research*, 67, 73–94, 2003.
- Browning, K. A.: Organization of clouds and precipitation in extratropical cyclones, *Extratropical Cyclones: the Erik Palmen Memorial Volume*, pp. 129–153, 1990.
- Chang, E. K. M.: Downstream Development of Baroclinic Waves as Inferred from Regression Analysis, *Journal of the Atmospheric Sciences*, 50, 2038–2053, [https://doi.org/10.1175/1520-0469\(1993\)050<2038:DDOBWA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<2038:DDOBWA>2.0.CO;2), 1992.
- 490 Chang, E. K. M., Lee, S. Y., and Swanson, K. L.: Storm track dynamics, *Journal of Climate*, 15, 2163–2183, [https://doi.org/10.1175/1520-0442\(2002\)015<02163:std>2.0.co;2](https://doi.org/10.1175/1520-0442(2002)015<02163:std>2.0.co;2), 2002.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haim-berger, L., Healy, S. B., Hersbach, H., Holm, E. V., Isaksen, I., Kallberg, P., Koehler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thepaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 495 Eckhardt, S., Stohl, A., Wernli, H., James, P., Forster, C., and Spichtinger, N.: A 15-year climatology of warm conveyor belts, *Journal of climate*, 17, 218–237, 2004.
- 500 Emanuel, K. A.: The maximum intensity of hurricanes, *J. Atmos. Sci.*, 45, 1143–1155, 1988.
- Gertler, C. G. and O’Gorman, P. A.: Changing available energy for extratropical cyclones and associated convection in Northern Hemisphere summer, *Proceedings of the National Academy of Sciences*, 116, 4105–4110, <https://doi.org/10.1073/pnas.1812312116>, 2019.
- Gertler, C. G., O’Gorman, P. A., Kravitz, B., Moore, J., Phipps, S. J., and Watanabe, S.: Weakening of the extratropical storm tracks in idealized solar geoengineering scenarios, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2020GL087348>, 2020.
- 505 Harris, B. L. and Tailleux, R.: Assessment of algorithms for computing moist available potential energy, *Quarterly Journal of the Royal Meteorological Society*, 144, 1501–1510, <https://doi.org/10.1002/qj.3297>, 2018.
- Hieronymus, M. and Nycander, J.: Finding the Minimum Potential Energy State by Adiabatic Parcel Rearrangements with a Nonlinear Equation of State: An Exact Solution in Polynomial Time, *Journal of Physical Oceanography*, 45, 1843–1857, <https://doi.org/10.1175/jpo-d-14-0174.1>, 2015.
- 510 Hou, A. Y., Kakar, R. K., Neeck, S., Azarbarzin, A. A., Kummerow, C. D., Kojima, M., Oki, R., Nakamura, K., and Iguchi, T.: The global precipitation measurement mission, *Bulletin of the American meteorological Society*, 95, 701–722, 2014.
- Koberstein, A.: Progress in the dual simplex algorithm for solving large scale LP problems: techniques for a fast and stable implementation, *Computational Optimization and Applications*, 41, 185–204, 2008.
- 515 Liu, C. and Zipser, E. J.: The global distribution of largest, deepest, and most intense precipitation systems, *Geophysical Research Letters*, 42, 3591–3595, 2015.
- Liu, N., Liu, C., Chen, B., and Zipser, E.: What are the favorable large-scale environments for the highest-flash-rate thunderstorms on Earth?, *Journal of the Atmospheric Sciences*, 77, 1583–1612, 2020.
- Lorenz, E. N.: Available potential energy and the maintenance of the general circulation, *Tellus*, 7, 157–167, 1955.
- 520 Lorenz, E. N.: Available energy and maintenance of a moist circulation, *Tellus*, 30, 15–31, 1978.

- Lorenz, E. N.: Numerical evaluation of moist available energy, *Tellus*, 31, 230–235, 1979.
- Madonna, E., Wernli, H., Joos, H., and Martius, O.: Warm conveyor belts in the ERA-Interim dataset (1979-2010). Part I: climatology and potential vorticity evolution, *J Clim*, 27, 3–26, <https://doi.org/10.1175/jcli-d-12-00720.1>, 2014.
- Oertel, A., Boettcher, M., Joos, H., Sprenger, M., Konow, H., Hagen, M., and Wernli, H.: Convective activity in an extratropical cyclone and its warm conveyor belt - a case-study combining observations and a convection-permitting model simulation, *Quarterly Journal of the Royal Meteorological Society*, 145, 1406–1426, <https://doi.org/10.1002/qj.3500>, 2019.
- Oertel, A., Boettcher, M., Joos, H., Sprenger, M., and Wernli, H.: Potential vorticity structure of embedded convection in a warm conveyor belt and its relevance for large-scale dynamics, *Weather and Climate Dynamics*, 1, 127–153, <https://doi.org/10.5194/wcd-1-127-2020>, 2020.
- O’Gorman, P. A.: Understanding the varied response of the extratropical storm tracks to climate change, *Proceedings of the National Academy of Sciences*, 107, 19 176–19 180, <https://doi.org/10.1073/pnas.1011547107>, 2010.
- O’Gorman, P. A.: The effective static stability experienced by eddies in a moist atmosphere, *Journal of the Atmospheric Sciences*, 68, 75–90, <https://doi.org/10.1175/2010jas3537.1>, 2011.
- O’Gorman, P. A. and Schneider, T.: Energy of midlatitude transient eddies in idealized simulations of changed climates, *Journal of Climate*, 21, 5797–5806, <https://doi.org/10.1175/2008jcli2099.1>, 2008.
- Oort, A. H., Ascher, S. C., Levitus, S., and Peixoto, J. P.: New estimates of the available potential energy in the world ocean, *Journal of Geophysical Research-Oceans*, 94, 3187–3200, <https://doi.org/10.1029/JC094iC03p03187>, 1989.
- Pan, Z., Mao, F., Rosenfeld, D., Zhu, Y., Zang, L., Lu, X., Thornton, J. A., Holzworth, R. H., Yin, J., Efraim, A., et al.: Coarse sea spray inhibits lightning, *Nature Communications*, 13, 4289, 2022.
- Peixoto, J. P. and Oort, A. H.: *Physics of climate*, American Institute of Physics, New York, 1992.
- Pfahl, S., Madonna, E., Boettcher, M., Joos, H., and Wernli, H.: Warm Conveyor Belts in the ERA-Interim Dataset (1979-2010). Part II: Moisture Origin and Relevance for Precipitation, *Journal of Climate*, 27, 27–40, <https://doi.org/10.1175/jcli-d-13-00223.1>, 2014.
- Randall, D. A. and Wang, J.: The moist available energy of a conditionally unstable atmosphere†, *J Atmos Sci*, 49, 240–255, [https://doi.org/10.1175/1520-0469\(1992\)049<0240:tmaeo>2.0.co;2](https://doi.org/10.1175/1520-0469(1992)049<0240:tmaeo>2.0.co;2), 1992.
- Rasp, S., Selz, T., and Craig, G. C.: Convective and slantwise trajectory ascent in convection-permitting simulations of midlatitude cyclones, *Mon Weather Rev*, 144, 3961–3976, <https://doi.org/10.1175/mwr-d-16-0112.1>, 2016.
- Riemann-Campe, K., Fraedrich, K., and Lunkeit, F.: Global climatology of Convective Available Potential Energy (CAPE) and Convective Inhibition (CIN) in ERA-40 reanalysis, *Atmospheric Research*, 93, 534–545, <https://doi.org/10.1016/j.atmosres.2008.09.037>, 2009.
- Rosca, D.: New uniform grids on the sphere, *Astronomy & Astrophysics*, 520, 4, <https://doi.org/10.1051/0004-6361/201015278>, 2010.
- Schneider, T. and Walker, C. C.: Scaling laws and regime transitions of macroturbulence in dry atmospheres, *Journal of the Atmospheric Sciences*, 65, 2153–2173, <https://doi.org/10.1175/2007jas2616.1>, 2008.
- Simmons, A. J., Untch, A., Jakob, C., Kållberg, P., and Undèn, P.: Stratospheric water vapour and tropical tropopause temperatures in ECMWF analyses and multi-year simulations, *Q J R Meteorol Soc*, 125, 353–386, <https://doi.org/10.1002/qj.49712555318>, 1999.
- Stansifer, E. M., O’Gorman, P. A., and Holt, J. I.: Accurate computation of moist available potential energy with the Munkres algorithm, *Quarterly Journal of the Royal Meteorological Society*, 143, 288–292, <https://doi.org/10.1002/qj.2921>, 2017.
- Stohl, A., Eckhardt, C., Forster, C., James, P., and Spichtinger, N.: On the pathways and timescales of intercontinental air pollution transport, *Journal of Geophysical Research*, 107, 4684, <https://doi.org/10.1029/2001JD001396>, 2002.

- Su, Z. and Ingersoll, A. P.: On the minimum potential energy state and the eddy size-constrained APE density, *Journal of Physical Oceanography*, 46, 2663–2674, <https://doi.org/10.1175/jpo-d-16-0074.1>, 2016.
- 560 Wang, J. and Randall, D. A.: The moist available energy of a conditionally unstable atmosphere. Part II. Further analysis of GATE data, *J Atmos Sci*, 51, 703–710, [https://doi.org/10.1175/1520-0469\(1994\)051<0703:tmaeo>2.0.co;2](https://doi.org/10.1175/1520-0469(1994)051<0703:tmaeo>2.0.co;2), 1994.
- Wernli, H. and Schwierz, C.: Surface cyclones in the ERA-40 dataset (1958-2001). Part I: Novel identification method and global climatology, *Journal of the Atmospheric Sciences*, 63, 2486–2507, <https://doi.org/10.1175/jas3766.1>, 2006.
- Yuval, J. and Kaspi, Y.: The effect of vertical baroclinicity concentration on atmospheric macroturbulence scaling relations, *J Atmos Sci*, 74, 1651–1667, <https://doi.org/10.1175/jas-d-16-0277.1>, 2017.
- 565 Zipser, E. J., Cecil, D. J., Liu, C., Nesbitt, S. W., and Yorty, D. P.: Where are the most intense thunderstorms on Earth?, *Bulletin of the American Meteorological Society*, 87, 1057–1072, 2006.