

Forcing and impact of the Northern Hemisphere continental snow cover in 1979-2014

Guillaume Gastineau¹, Claude Frankignoul^{1,2}, Yongqi Gao^{3,†}, Yu-Chiao Liang⁴, Young-Oh Kwon², Annalisa Cherchi^{5,6}, Rohit Ghosh^{7,8}, Elisa Manzini⁷, Daniela Matei⁷, Jennifer Mecking⁹, Lingling Suo³,
5 Tian Tian¹⁰, Shuting Yang¹⁰ and Ying Zhang¹¹

¹UMR LOCEAN, Sorbonne Université/IRD/MNHM/CNRS, IPSL, Paris, 75005, France

²Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, U.S.A.

³Nansen Environmental and Remote Sensing Center and Bjerknes Center for Climate Research, Bergen 5006, Norway

10 ⁴Department of Atmospheric Sciences, National Taiwan University.

⁵National Research Council of Italy, Institute of the Atmospheric Science and Climate (CNR-ISAC), Bologna, Italy.

⁶Istituto Nazionale di Geofisica e Vulcanologia, Bologna, Italy.

⁷Max Planck Institute for Meteorology, Hamburg, Germany.

15 ⁸Department of Meteorology, University of Reading, United Kingdom.

⁹National Oceanography Centre, Southampton, United Kingdom.

¹⁰Danish Meteorological Institute, Copenhagen, Denmark.

¹¹Nansen-Zhu International Research Center, Institute of Atmospheric Physics, Chinese Academy of Sciences, 100029 Beijing, People's Republic of China.

20 [†]Deceased, July 23rd 2021.

Correspondence to: Guillaume Gastineau (guillaume.gastineau@sorbonne-universite.fr)

Abstract.

The ~~main drivers of role of surface ocean anomalies for~~ the continental Northern Hemisphere snow cover ~~are~~ investigated ~~in , together with~~ the ~~1979-2014 period.interactions between snow cover and~~
25 ~~atmosphere~~. Four observational datasets ~~are used~~ and two large multi-model ensembles of atmosphere-only simulations ~~are used~~, with prescribed sea surface temperature (SST) and sea ice concentration (SIC). A first ensemble uses observed interannually varying SST and SIC conditions for 1979-2014, while a second ensemble is identical except for SIC where a repeated climatological cycle is used. SST and external forcing typically explain 10 to 25% of the snow cover variance in model simulations,
30 with a dominant forcing from the tropical and North Pacific SST ~~during this period. In terms of the climate influence, while no robust influence of the SIC is found. In observations, the Ural blocking is the main~~

~~driver of the snow cover anomalies, both observations and models show no robust links between the November and April snow cover variability and the atmospheric circulation one month later. Onover Eastern Eurasia, while the other hand, North Atlantic Oscillation (NAO) dominates the first mode of Eurasian snow cover variabilityforeing in January. In November and more robustly in January, with more extended snow dipolar anomalies of snow cover over Eurasia, with positive anomalies over Europe and negative anomalies over Southern Siberia, also precede the Arctic Oscillation (AO) by one month. In models, snow cover over western Eurasia, is found to precede by one month an atmospheric circulation pattern similar to a negative Arctic Oscillation (AO). A decomposition of the variability in the model simulations shows that this relationship is mainly due to internal climate variability. DetailedJanuary also precedes by one or two months a negative AO phase. The detailed outputs from one of the models indicatesuggest that both the Westernwestern Eurasia snow cover anomalies are preceded by a negative AO phase accompanied by a Ural blocking pattern and a stratospheric polar vortex weakening. The link between the AO and the snow cover variability is strongly related to the concomitant role of the stratosphericare generated by Ural blocking, and that both snow cover and polar vortex, with the Eurasian snow cover acting as a positive feedback for the AO variability in winter. No robust influence of the SIC variability is found, as the sea ice loss in these simulations only drives an insignificant fraction of the snow cover anomalies, with few agreements among models. anomalies act to generate the AO one or two months later.~~

50 **1 Introduction**

Understanding the origin and impact of snow variability is important for many activities such as agriculture, tourism, management of freshwater resources, or road maintenance. It is also essential for the evolution, and understanding of midlatitude and subarcticmany ecosystems. Snow is an important element for the climate as the high albedo of snow leads to increased reflected shortwave radiation at the surface with a direct influence on the earth's radiative budget. The small thermal conductivity of the snow pack also insulates the soil from the cold winter atmosphere and plays an important role in the stability of the permafrost (Pulliainen et al., 2017).

Snow over land accumulates from snowfall events and is melted by surface air temperatures above the freezing point. The variability of snow cover and snow depth is therefore modulated by the midlatitude and polar atmospheric variability. Winter atmospheric variability is large and is mostly unpredictable beyond a week or two as it owes its existence to internally-driven atmospheric processes (Feldstein, 2000; Deser et al., 2012). However, other processes influence the atmospheric variability at low frequency, which leads to the potential predictability of winter climate at the seasonal time scale (Scaife et al., 2014).

Tropical surface anomalies can strongly alter the large-scale atmospheric circulation and influence the extra-tropical regions through atmospheric teleconnections. In particular, the El Niño-Southern Oscillation (ENSO) has a large influence over North America through the Pacific-North American (PNA) pattern (Wallace and Gutzel, 1981; Lau 1997), and also over Europe (Mathieu et al. 2004; Lopez-Parages et al. 2016). The PNA can in turn modify the snow depth, as found in observations (Ge and Gong, 2009).

Extra-tropical surface anomalies may also drive the winter atmosphere, through the influence of extra-tropical sea surface temperature (SST; see the review of Kushnir et al. 2002; Gastineau and Frankignoul, 2015), sea-ice (Deser et al., 2007; Honda et al., 2009; Garcia-Serrano et al., 2015; King et al., 2016), and snow cover (Cohen and Entekhabi, 1997; Gastineau et al., 2017; see the review of Henderson et al., 2018). Lastly, troposphere-stratosphere coupling in winter can also enhance and increase the persistence of lead ~~to more persistent~~ atmospheric modes (Perlwitz and Graf; 1995; Baldwin and Dunkerton, 1999; Scaife et al., 2014).

The land snow cover is also ~~largely~~ affected by ~~climate change caused by~~ external forcings such as the increasing concentration of greenhouse gases, the evolution of aerosol or ozone concentration ~~or ozone~~, and land use change. The snow cover extent was found to decrease over the last decades (~~Gulev et al., 2021~~; Déry and Brown, 2007; Gulev et al., 2021), although observational data shows a large spread in fall and early winter (Brown and Derksen, 2013; Mudryck et al., 2017). Mudryk et al. (2020) reported negative trends below $-50 \times 10^3 \text{ km}^2$ per year over 1981-2018 in November, December, March and May.) Recent observational estimates also found a decreasing trend of the snow mass over North America but an insignificant decrease over Eurasia (Pulliainen et al., 2020). Detection-attribution studies have

attributed the decrease of ~~the~~ snow cover to human activities (Paik and Min, 2020; Guo et al., 2021), but the specific role of the different drivers is unknown. Furthermore, the atmosphere-ocean general circulation models (AOGCM) from CMIP5 (Coupled Model Intercomparison Project phase 5) underestimate the land snow extent, while they overestimate the snow mass (Derksen and Brown, 2012; 90 Mudryk et al., 2020). Even if the snow cover extent is better simulated in CMIP6 (Coupled model Intercomparison Project phase 6) models (Mudryk et al., 2020), global climate models mostly use highly simplified snow physics (Krinner et al. 2018). The simulation of snow cover anomalies over land, therefore, remains a challenge as it involves the large-scale circulation together with the parametrized atmospheric precipitation and land surface processes. In the present study, we will further assess the 95 influence of external forcing, SST and sea ice concentration (SIC) anomalies on the snow cover.

Land snow variability also influences the climate. Cohen and Entekhabi, (1997) found that when the snow cover over eastern Siberia is anomalously large in October, negative phases of the Arctic Oscillation (AO) are more frequent during the following months. This was confirmed by Saito and Cohen (2003) and Cohen 100 et al. (2014). Using an extended observational record, Gastineau et al. (2017) found a similar relationship, albeit between November snow cover and the subsequent December and January AO. They also found that concomitant sea ice anomalies reinforced the atmospheric response to snow cover anomalies. These relationships suggest that snow cover anomalies can influence the mid-latitude atmospheric circulation in the same way as SST or SIC anomalies. The pathway of the snow influence involves an amplification of 105 the climatological tropospheric stationary wave associated to a lower-troposphere cooling as the snow cover increases (Cohen et al., 2014), as found for the 2017/2018 winter (Lü et al., 2020). Such amplification was suggested to lead to stratospheric warming which can result in more frequent negative AO events through downward propagation (Baldwin and Dukerton, 1999). Sensitivity simulations using models with prescribed snow cover also revealed a consistent AO-like atmospheric response to more 110 extensive Eurasian snow cover (Gong et al., 2003; Fletcher et al., 2009). Such influence is consistent with changes in subseasonal~~seasonal~~ forecast skill when modifying the initialization of the snow cover in 2004-2009 (Orsolini et al., 2013) or in 2009-2010 (Orsolini et al. 2016), even if this influence is not found systematically from other periods and different models (Garfinkel et al., 2020). The statistical

relationships found in observations are stronger than, but consistent with ~~those the ones produced~~ in some
115 of the AOGCM simulations from CMIP5 climate models (Gastineau et al., 2017). Liang et al. (2021)
proposed that the apparent underestimation of the atmospheric response to sea-ice anomalies in the
Barents-Kara Seas in CMIP6 atmosphere-only simulations was in part due to the lack of consistency
between sea ice and snow cover anomalies when the former was prescribed. ~~Indeed~~~~For instance~~, Ural
blocking increases the eastern Siberian snow cover while it decreases the Barents-Kara SIC (Gastineau et
120 al., 2017; Peings 2019). Both the increased Siberian snow cover and Barents-Kara sea ice loss are found
to lead to negative AO-like anomalies in the following months (Gastineau et al., 2017; Simon et al. 2020).
This may result in a larger AO response than ~~the one~~ expected from the sea ice alone, as proposed by
Cohen et al. (2014). Hence, atmosphere-only simulations using prescribed sea-ice anomalies but with
prognostic snow cover cannot simulate the synchronization of sea ice and snow, and the atmospheric
125 response to SIC anomalies could not be reinforced by the snow cover anomalies, unlike in observations.
~~Lastly, a heavy spring snow cover was found to increase the soil moisture over Siberia, and the
Mongolian/Tibetan plateau, which can lead to an abnormal cooling of the land surface. This can alter the
atmospheric circulation related to monsoon through the modification of the land/sea contrasts (Barnett et
al., 1989).~~
130 In the present study, we will ~~also~~ further assess the drivers and large-scale impacts of snow cover
anomalies, focusing on early winter, winter and early spring. We use a large ensemble of atmosphere-
only simulations to characterize the ~~main drivers and impacts of~~ snow cover variability in the Northern
Hemisphere. To sample the uncertainties of the observations, we analyze four observational products.
Section 2 presents the data and methods. Section 3 discusses the influence of the observed SST and SIC
135 anomalies on continental snow. In Section 4, we investigate the internal variability of the snow cover and
its influence on the atmosphere. Discussion and conclusions are given in the last section.

2. Data and methods

2.1 Observations

140 Several snow datasets ~~are have been~~ used to sample ~~some of~~ the observational uncertainty. ~~We (Brown et al., 2010). For this study we~~ use the monthly snow water equivalent (SWE) depth and snow cover of ERA5-land in 1981-2014 (Muñoz-Sabater (C3S, 2019; Muñoz-Sabater et al., 2021), resulting from the ECMWF land-surface H-TESEL model forced with ERA5 atmospheric reanalysis (Hersbach et al., 2020). ~~for 1981-2014.~~ We also use the monthly snow diagnostics from MERRA2 (GMAO, 2015; Gelaro et al., 2017) for the same period. The NOAA climate data record (CDR) of Northern Hemisphere weekly snow cover extent dataset (Robinson et al., 2021) is retrieved from the National Center for Environmental Information and is aggregated into monthly time series for the in 1981-2014 period. The . ~~We also use the~~ monthly SWE from GlobSnow v3 (Pulliainen et al., 2020) is used snow depth in 1980-2014, where the missing data in December 1981 was interpolated linearly ~~in~~ between November 1981 and January 1982.

150 Lastly, we use the ~~daily monthly~~ CanSISE SWE in 1981-2010 observation-based ensemble of the Northern Hemisphere (Mudryk et al., 2015; Mudryk and Derksen, 2017). ~~.) snow depth and snow cover in 1981-2010;~~ which is based on five products: GlobSnow v2, ERA-Interim/Land reanalysis, MERRA reanalysis, Crocus (Brun et al., 2011) and GLDAS version 2 (Rodell et al., 2004). The CanSISE product also provides a spread based on the range (maximum minus minimum) of these five products. A snow cover from

155 CanSICE is then estimated from the SWE using a threshold of 7 mm. If the daily SWE depth is lower (larger) than 7 mm, then it is assumed that the snow cover is zero (1). A minimum and maximum snow cover is also estimated with the same procedure using the SWE and its spreads, assuming the spread is centered on the mean SWE. The SWE and snow cover from CanSISE are then aggregated into monthly means.

160

The atmospheric 2m geopotential, air temperature, ~~zonal wind~~, and sea level pressure (SLP) fields are retrieved from ERA5 reanalysis (Hersbach et al., 2019 C3S, 2017; Hersbach et al., 2020).

All data is regridded with bilinear interpolation into a 1.26°x2.5° regular grid before analysis. Coastal regions are masked if the fraction of land is below 50%. ~~Some~~~~In some~~ products, such as GlobSnow, have missing data over mountain regions. Therefore, mountain and ice cap regions are masked in all data ~~where the CanSISE data is missing.~~

2.2 Model simulations

170 We use the outputs of the two multi-model land-atmosphere simulation ensembles discussed in Liang et al. (2020, 2021). These simulations used as boundary conditions the SST and SIC provided by the HighResMIP panel of CMIP6 (Haarsma et al., 2016) and atmospheric concentration of aerosol, greenhouse gases, and ozone from CMIP6 (Eyring et al., 2016) in the 1979-2014 period. We use the outputs of eight models where the snow depth was saved and distributed (Table 1). The ensemble ALL uses interannually-varying daily SST and SIC. The other ensemble, called ~~NoSICvar~~~~NoSIC~~, is identical but uses a repeated 1979-2014 climatological SIC in the Arctic, with adjustment of the associated local SST (Hurrell et al., 2008). The climate sensitivity to SIC anomalies is provided by the difference ALL minus ~~NoSICvar~~~~NoSIC~~. As noted in Liang et al. (2020) and ~~in~~ Table 1, the experimental protocol has some small differences for each model, but these deviations are unlikely to affect the results substantially.

180 The number of members varies among models from 10 to 30, while the horizontal resolution varies from about 60 km to 150 km. The large diversity of models allows us to study the model dependence. However, for comparison with observations, these ensembles of atmosphere-only models have limitations associated with the lack of active two-way coupling with sea ice and SST, uncertainties in the SST and SIC forcing, and simplified sea ice physics, ~~(for instance the sea ice thickness is constant)~~, as discussed

185 in Liang et al. (2021).

We use the monthly ~~500-hPa geopotential~~, SLP, 2m air temperature, and snow depth in all models. For ~~LMDZOR6~~~~LMDZ6~~ and CMCC, the snow depth was converted into snow water equivalent (SWE) depth, assuming a constant snow density of ~~240330~~ kg m⁻³, as found in observations (Sturm et al. 2010). The snow cover is a diagnostic variable in many models and was not available for four models (see Table 1):

190

EC-Earth3, ECHAM6, HadGEM3 and IAP4. Lacking a better formulation, we calculate the snow cover from the SWE using a threshold of 7.5 mm. If the monthly SWE depth is lower (larger) than 7.5 mm, then it is assumed that the snow cover is zero (1). This estimation is based on LMDZOR6~~LMDZ6~~, where we found that a reasonable snow cover extent is obtained with the 7.5 mm threshold when using monthly outputs. This procedure is similar to that of Krinner et al (2018), except they used a threshold of 5 mm.

The detailed outputs from LMDZOR6 are also used with the monthly geopotential height at 500-hPa and 50-hPa, and the daily air temperature. For this model, the wave activity flux from Plumb (1985) is also calculated from daily geopotential height, zonal wind and meridional wind at 500-hPa and 250-hPa.

200

All data sets were regridded with bilinear interpolation into the regular grid $1.26^\circ \times 2.5^\circ$ (~150 km) before analysis. Coastal regions and grid points with complex orography were masked consistently in all models using the observational mask. Multi-model ensemble means (MMM) are constructed by giving the same weight to each ensemble member, which largely removes the influence of internal atmospheric variability.

205 **2.3 Methods**

We study the effects of SST, SIC and external forcing in driving snow cover anomalies with an analysis of variance (ANOVA) with two factors, also known as two-way ANOVA. The ANOVA is a statistical analysis method for comparing the means of various samples and investigating the influence of one or several categorical independent variables, called factors, on one continuous variable (Von Storch and Zwier, 1999). Here the ANOVA is applied in a balanced design to the land snow from the ALL and NoSICvar~~NoSIC~~ ensembles, separately for each individual model and for each calendar month. The first factor is the simulated year, called t , which varies from 1979 to 2014. The second factor is the ensemble, called e , and represents the ALL versus NoSICvar~~NoSIC~~ ensembles. The interaction between the year and the ensemble is called $t:e$. In the analysis, the sum of squares quantifies the variance associated with each factor. The ANOVA then compares such variance to the residual variance to test the effect of the factors. The corresponding p-value indicates if the effect of the factors (t , e and $t:e$) isare statistically

215

significant. Hereafter, we show such p-values, together with the ratio of the sum of squares over the total variance to quantify the variance explained by each factor.

220 The statistical model of the ANOVA decomposes the snow cover anomalies of a calendar month in each year and ensemble, called X , by:

$$X(t, e) = \mu + \beta_t(t) + \beta_e(e) + \beta_{t:e}(t, e) + \varepsilon, \quad (2)$$

where μ , the theoretical mean of X , corresponds to the seasonal mean of the calendar month. β_t is a different constant for each year, β_e is a constant for each ensemble, $\beta_{t:e}$ is an interaction term different for each year and ensemble, and ε is a gaussian noise. If the ANOVA is significant for the factor t , then at least one of the β_t is significantly different from zero. It implies that the time-varying prescribed boundary conditions have an influence on the snow cover in both ALL and ~~NoSICvarNoSIC~~, which should result from time-varying SST or external forcings, as they both can influence the atmosphere and land. Similarly, the effect of time-varying SIC is accounted for by the second factor e . If the second factor is significant, ~~meaning~~ with at least one of the β_e different from zero, it demonstrates an influence of varying sea-ice concentrations on the mean land snow. Lastly, if at least one of the interaction terms, $\beta_{t:e}$, is significant, it suggests that the influence of SIC is time-dependent. The ANOVA is repeated for each calendar month.

235 The main drivers of snow cover ~~and snow depth~~ are characterized using Empirical Orthogonal Functions (EOFs). The EOFs of the Northern Hemisphere snow cover are calculated north of 30°N, while the domain for Eurasian snow cover EOFs is (0°E-180°E, 30°N-90°N). Three EOF analyses are performed using the year-to-year time series corresponding to each calendar month separately. ~~The~~A first EOF analysis ~~performed~~ is based on the MMM calculated from the ALL experiments. ~~The~~~~Sueh~~ EOF ~~pattern~~ is ~~denoted~~ ~~asealled~~ EOF_{BC}, where BC stands for boundary conditions and indicates the driving effect of the prescribed SST, SIC and external forcings (concentration of greenhouse gases, aerosol and ozone). As the forcing from sea ice concentration is weak (Liang et al., 2021), the EOFs are almost identical when using ~~NoSICvarNoSIC~~ instead of ALL. For instance, the pattern correlation between the first EOF_{BC} (EOF_{1BC}) of ALL and that of ~~NoSICvarNoSIC~~ is 0.95, 0.93 and 0.98 for November, January and April,

245 respectively. EOF_{1BC} therefore mainly quantifies the main pattern of variability induced by the SST and external forcing. The corresponding principal components (PCs) are denoted PC_{BC}. A second EOF analysis, called EOF_{SIC}, is identical but performed on the difference between the MMM of ALL and ~~NoSICvarNoSIC~~, to highlight the effect of the SIC variability. The corresponding principal components (PCs) are denoted PC_{SIC}. ~~Hereafter, all principal components are normalized, and the EOFs are illustrated using their regression onto the standardized PC.~~

250

Hereafter, all principal components are standardized, and the EOFs are illustrated using their regression onto the standardized PC. The sign convention is that a positive PC corresponds to an EOF with positive loading over eastern Europe (20°E-70°E 55°N-70°N).

255

Lastly, ~~we investigate~~ the internal land-atmosphere variability is investigated in the model simulations with a third EOF analysis. The internal-~~snow~~ variability is investigated after removing the ensemble mean of the snow evolution that mostly reflects the effect of SST, SIC, and external forcing. ~~We Therefore, we~~ conduct an EOF analysis separately for each model using all the members of ALL and ~~NoSICvarNoSIC~~ concatenated after the removal of their respective ensemble means. This third analysis provides EOF_{Int}, and PC_{Int} as spatial patterns and time series, respectively. The relevance of this analysis might be limited when the ensemble size is small (only 10 members for some models), as the ensemble means are more affected by internal variability.

260

265 In addition, various fields, such as the surface air temperature, SLP, geopotential height, and zonal wind, are regressed onto PC_{BC}, PC_{SIC} and PC_{Int}. The p-values of the univariate regression slopes are given by a Student's *t*-test. The year-to-year autocorrelations for separate calendar months are typically insignificant between 0 and 0.05 (not shown). The only exception is for April, where such autocorrelation is significant over Scandinavia and the East European Plain, but it remains modest with maximum values at 0.08.

270 Hence, we did not account for a reduction in the degree of freedom due to year-to-year correlation.

The ANOVA, the retrieval of EOF_{Int} and the regression analyses using PC_{Int} are performed separately for each model, but the figures provide the mean for the eight models, ~~denoted by multi-model mean (MMM),~~ using a weight proportional to the ensemble size of each model ensemble, referred to as the multi-model mean (MMM). This avoids giving too much weight to models with only 10 ensemble members. We indicate grid points where the sign of anomalies is the same in seven models out of eight. This corresponds to a probability of 6.2% when considering that the sign of the anomaly has a probability of 50% in the models, as deduced for the Binomial probability distribution. ~~We also~~Additionally, we indicate the grid points where the p-value is below 5% in at least five models out of eight.

280

3. Simulated Northern Hemisphere snow cover and depth

3.1 Climatology

First, we briefly assess the Northern Hemisphere land snow simulated in the eight models. The mean seasonal cycle of land snow extent and snow mass is first calculated over North America and Eurasia in 1979-2014. The snow extent over North America (0°N-90°N 180°W 0°E) and Eurasia (0°N-90°N 0°E 180°E) has a maximum in January-February (Fig. 1a-b, black lines). November and April are associated with the start and the end of the season with extensive land snow coverage, respectively. ~~The mean~~ seasonal cycle of the Eurasian snow area is well represented by all the models (Fig. 1b, color lines). The differences between the models are within the range of uncertainty between the observational data sets, except for ~~ECHAM6, which underestimates the snow cover throughout the cold season. We note that~~ EC-Earth3 that simulates a slower snow cover decrease in spring. The snow cover area over North America (Fig. 1a) is also well captured by models, but it is overestimated in EC-Earth3 and ~~again~~ underestimated by ECHAM6. We also calculate the standard deviation obtained from year-to-year time series for each month. The interannual variability in models also agrees with that found in observations (Fig. S1).

290 There is less agreement on the snow mass (Fig. 1c-d). First, the snow mass estimations from observations show a large spread that is maximum from February to May. Then, ~~LMDZOR6~~LMDZ6 and CMCC both ~~largely~~ overestimate the snow mass in Eurasia and North America from December to March. NorESM

and CESM2 only overestimate the snow mass over North America, especially from February to March. Other models simulate snow masses within the spread of observational products. In conclusion, the models reproduce the observed snow cover seasonality, but tend to overestimate the snow mass, except ECHAM6, which simulates a realistic snow mass, but underestimates the snow cover. However, the formula used to convert snow water equivalent into snow cover might not apply to ECHAM6. These conclusions are in agreement with the similar analysis of CMIP6 AOGCMs from Mudryk et al. (2020) or Zhong et al. (2022). Therefore, the use of atmosphere-only simulations does not significantly reduce the land snow biases compared to AOGCM simulations.

The location of the snow cover biases of each model compared to CanSISE is illustrated for January in Fig. 2. We chose here CanSISE as a reference as it is based on an ensemble of observations. Most models simulate more snow cover than observational products from the Tibetan plateau to Eastern Siberia and too little snow cover over southwestern Eurasia. No apparent snow biases are found over the fully snow-covered domain between Eastern Europe and Central Asia. Over North America, there is generally more snow in models than in observations over the Rocky Mountains, and a few models also underestimate the snow cover over Northeastern Canada. Given the large uncertainty of the observational products over Mountain regions, more observations would be needed to fully confirm the biases over these regions. The snow water equivalent in models (Fig. 3) shows a generally positive bias over land with no consistent large-scale pattern in LMDZOR6LMDZ6, CMCC, NorESM, and CESM2. TheSuch positive bias is negativereduced in EC-Earth3 and HadGEM3. In ECHAM6 and IAP4.1, with an underestimation, there is a weak overestimation of the SWEsnow water equivalent over the East European Plain and. An overestimation is also simulated in IAP4.1, but over North-Eastern Siberia, respectively. EC-Earth3 and HadGEM3 agree with the observations.

Hereafter, we focus on the land snow variability in November, January, and April, which represent the start, the maximum, and the end of the period with large snow coverage in the Northern Hemisphere (Fig. 1).

3.2 Assessment of snow cover anomalies

Figure 4a-b shows the time series of the anomalous snow cover area in winter, defined by the average from November to March, in the MMM of ALL. The anomalies are defined with respect to the 1979-2014 climatology. The observational values are also shown but using a different scale to facilitate the comparison. Indeed, the decreasing trend of the MMM snow cover area is roughly half of that observed over Eurasia and North America (see also red and grey symbols in Fig. 5b). In addition, the year-to-year variability of ALL is much smaller than that observed. This presumably reflects that the observations correspond to one realization while the model ensemble means are averaged over 10 to 30 members, depending on the model, which strongly reduces the impact of internal variability. This suggests either that half of the observed trends is due to internal variability, or that the influence of SST or external forcing is underestimated by half in the models. Similar results are found for the snow mass, but the decreasing trend over Eurasia is much weaker in models than in observations (Fig. 4c-d and Fig. 5a). The timing of some of the minima and maxima are consistent, as in 2007/2008 for the snow extent over Eurasia (Fig. 4a), in 1993 and 1999-2000 for the snow extent over North America (Fig. 4b), or in 1987, 1998 and 2010 for the North American snow mass (Fig. 4d). This is consistent with the strong relationship between El Niño events and positive phases Pacific North American (PNA) pattern, which are associated with warm anomalies and decreasing snow depth over western North America (Ge and Gong, 2009).

The correlation between the ALL MMM and ERA5-Land snow cover area is 0.66 (0.59) over Eurasia (North America), while it is 0.69 (0.60) over Eurasia (North America) for snow mass, which demonstrates a dominant influence of the boundary conditions. After removing the linear trend from every time series, the correlations are smaller, but they remain significant, except for snow mass over Eurasia (compare red and yellow symbols in Fig. 5d).

345

The overall impact of the sea-ice variations on the snow cover area and snow mass is limited, as shown by the differences between the MMM of ALL and ~~NoSICvarNoSIC~~ over North America (Fig. 4f-h). ~~The time series in Fig. 4f-h show), which have~~ no clear trend and are not significantly related to observations at the 5% level (Fig. ~~5d, except5d~~). ~~However, a significant correlation of 0.43 is obtained~~ for the Eurasian

350 snow ~~extent~~area over Eurasia (Fig. 4e), which is significantly correlated with observations (R=0.43)
even remains significant for detrended time series (R=0.38).

The correlations of land snow area and mass between observation and models remain significant over
Eurasia and North America separately for November, January and April in Fig. 5c-d for ALL (red and
355 orange symbols), which confirms the robust influence of boundary conditions. The 1980-2014 linear
trends of snow cover and snow depth are also assessed ~~for the 3 months~~ in Fig. 5a-b, using the MMMs,
and calculating their statistical significance with a Student's *t*-test, as detailed in Liang et al. (2021).

Over Eurasia and North America, the trends in ALL models for November and, January, and April (Figs.
5a-b, red circle and plus) are consistent with that of the whole winter (NDJFM). However, in April, the
360 snow depth underestimated trend over Eurasia in ALL is more negative, while the snow extent trend over
North America is smaller and insignificant. The comparison between the observed CanSISE (Figs. 5a-b,
grey circle and plus) and simulated trend reveals important differences, as a dominant negative trend is
observed in November and January for snow extent and mass, respectively. A comparison of the trends
obtained in other observational datasets (Figs. S2 and S3) shows a large spread in trend estimates, with
365 increasing snow extent in fall and early winter in the NOAA-CDR compared to observations, while ERA5-
Land, MERRA2 and CanSISE show a decreasing snow extent, as found previously (Brown and Derksen,
2013; Mudryck et al., 2017). The reason for this difference between the NOAA-CDR and multi-
observation products is unknown (Mudryck et al., 2017). This suggests that observational uncertainties
are important, especially in fall, as reported by Fox-Kemper et al. (2021). ~~is mainly due to November.~~

370 The trend maps obtained from ALL MMM in November, January, or April (Figs. 5e-g) reveal an
important large-scale decrease in snow cover, maximum at the edges of the snow-covered domain. ~~(Figs.~~
~~5e-g).~~

The influence of sea-ice is investigated with the trends of the MMM of ALL minus ~~NoSICvar~~NoSIC, and
375 the correlation between ~~observations~~observation and the MMM of ALL minus ~~NoSICvar~~NoSIC in Fig.
5a-d (blue and sky-blue symbols). The trends of ALL minus ~~NoSICvar~~NoSIC are negative and significant
for snow mass only, which may reflect an influence of the sea-ice loss reducing the snow mass. For the

snow cover, the correlation and trend are mainly small and insignificant. The trend maps (Fig. 5h-j) show a weak but significant decreasing trend for ~~November~~, January, or April in Southern Scandinavia extending eastward into Eurasia. In November, a ~~with an additional~~ decreasing trend is located east of Scandinavia, downstream of the Barents Sea. Such a location is in agreement with the large oceanic heat release expected from the observed Barents-Kara sea ice loss (Deser et al., 2015). Another decreasing snow cover trend is also simulated over north-eastern Canada in November. ~~(Fig. 5h-j).~~

385 3.3 Role of the boundary conditions in driving snow cover

The influence of boundary conditions is quantified using an ANOVA (see section 2.3). We first applied the ANOVA separately at each grid point. The effect of SST and external forcing represented by the factor t is found to be dominant. The snow cover variance fraction associated with this factor is significant in all models in November, January, or April (Figs. 6a-c). It is largest over the mid-latitude edges of the snow-
390 cover in Eurasia or North America, and may reach 15% over the Tibetan plateau, although models and observations have large uncertainties there (Mudryk et al., 2020). A large variance fraction ($>5\%$) is also simulated over the Rockies and Scandinavia. In November, the variance fraction is ~ 4 to 5% over Northern Canada and a latitudinal band from Scandinavia to Eastern Siberia. By January, the ~~mean~~ variance fraction at the edge of the snow cover domain increases on average to $\sim 6\%$ and shifts toward
395 eastern Europe and from the Caspian Sea to eastern China. In April, the large variance fraction is more important (~ 6 -10%) but more localized from the East European Plain to southern Siberia and over Northern Canada. The results are summarized by an ANOVA using the Eurasia or North America snow area instead of grid point values in Fig. 6d-f. Models show a significant influence of SST and external forcing (indicated by SST/ext in Fig. 6d-f) with 10% to 25% of the variance explained over both domains,
400 despite important differences between models.

The snow cover variance associated with the varying sea ice concentration is given by the factors e and $t:e$ (see section 2.3 for details) representing the influence of the SIC on the time-mean and time-varying snow, respectively. The variance fractions show no clear agreement between models and are largely

405 insignificant at most grid points (not shown). The results are summarized in Fig. 6d-f for the snow cover area over Eurasia and Northern America. The sea ice only explains 1% to 5% of the variance, as found in the interaction $t:e$ and the impact on the time mean snow cover is below 0.3% in most models. In most models, the ANOVA test is not significant for these two factors. We conclude that sea ice does not have a robust influence on the snow cover in our simulations. Using ~~SWEsnow-water-equivalent~~ instead of
410 snow cover yields similar results (not shown).

3.4 Assessment of the role of the boundary conditions for snow cover

To assess the main patterns of simulated year-to-year snow cover variability, we first investigate ~~the main pattern of variability in~~ observations, ~~with~~using separate EOFs (north of 30°N) for each calendar month. Figure 7 shows the first two snow cover EOFs in ERA5-Land in, from top to bottom, November, January, and April, as well as their pattern correlation with the corresponding EOFs obtained from the three other
415 observational data sets (right). In November, the first EOF (EOF1) shows anomalies of the same sign with large loading over the East European Plain, eastern Eurasia, and central North America, near the edges of the mean snow-covered area. The second EOF (EOF2) is a dipolar pattern with large loading over eastern Europe and small loading with the opposite sign between the Aral Sea and the Baikal Lake and in western North America. In January, EOF1 is dominated by a large loading over Europe, while
420 EOF2 is dominated by strong anomalies over North America. In addition, EOF1 and EOF2 display small anomalies near the edge of the Eurasian snow-covered domain. In April, EOF1 is a dipole with large loading over North America, and anomalies with an opposite sign between the Caspian Sea and the Baikal Lake. EOF2 only shows large loading over the East European Plain. CanSISE provides very similar snow cover EOFs to ERA5-Land in all months. The EOF1 patterns in MERRA-2 and NOAA-CDR are also
425 similar in January and April, but they are different in November. The EOF2 patterns in MERRA-2 and NOAA-CDR mainly disagree with ERA5-Land. This suggests that the observational uncertainty is large in November but that the EOF1 pattern is otherwise rather robust.

430 To emphasize the role of external forcing, ~~SSTSIC~~ and ~~SICSSST~~ for the simulated snow cover changes, the first EOF of the Northern Hemisphere snow cover in the MMM of ALL, called EOF1_{BC} is shown in

Figs. 8a, 8d and 8g. The corresponding principal components, hereafter $PC1_{BC}$, all show a positive trend (not shown), so that the $EOF1_{BC}$ resembles the maps of the land snow trend from ALL in 1981-2014 (compare Fig. 5e-g and Fig. 8a,d,g). November $EOF1_{BC}$ (Fig. 8a) also shows anomalies of the same sign
435 over North America and from Scandinavia to eastern Siberia, at the edges of the mean snow-covered region, and near the Tibetan plateau. In January (Fig. 8d), the pattern is again a monopole, but it is centered over a band from Europe to East Asia as the snow-covered domain is broader than in November. In April (Fig. 8g), the loading over Eurasia is similar to that found in November, but the anomalies over North America are opposite. $EOF1_{BC}$ explains ~~from~~ 36% to 48% of the variance in November, January, and
440 April.

The EOF associated with the boundary conditions and external forcings can be compared with the observed snow cover EOF. The first two EOFs in ERA5-Land (Fig. 7, left) have some similarities with the $EOF1_{BC}$ patterns in Figs. 8a,d,g. However, the loading is more localized in ERA5-Land, without a
445 clear location at the edge of the snow-covered domain. Moreover, in ERA5-Land EOF1 (EOF2) only explains ~~from~~ 15% to 21% (12% to 13%) of the variance. The somewhat different patterns and the smaller explained variance may reflect in part the large influence of internal atmospheric variability in observations. To reveal the links between the observed snow cover variability and the influence of SST and external forcing, $PC1_{BC}$ is compared to the observed snow cover PC1 and PC2. The correlation
450 between $PC1_{BC}$ and the observed PC1 (Fig. 8c,f,i, left, blue bars) is only significant when using ERA5-Land and CanSISE in November and when using ERA-Land and NOAA-CDR in April, and it is not significant at all in January. Moreover, the correlation between $PC1_{BC}$ and the observed PC2 (orange bars) is only 5% significant in November for NOAA-CDR. After removing a linear trend from all the time series (not shown), the correlations with $PC1_{BC}$ that largely stem from SST forcing are smaller, and they
455 only remain significant with ERA5-Land PC1 in November ($R=0.36$), and with ERA5-Land ($R=0.47$) and NOAA-CDR ($R=0.48$) PC1 in April. This confirms the larger effect of changing SST and external forcing in November and April, although the EOF analysis ~~isn~~ not robust among the observational dataset. There is no significant correlation in January, presumably because the internal variability is larger.

460 The snow cover EOFs solely related to the sea ice are calculated from the MMM difference of ALL minus
| ~~NoSICvar~~~~NoSIC~~ (Figs. 8b,e,h). The variance fraction explained by the first EOF_{SIC} is between 10% and
13%. The absolute variance explained by EOF_{SIC} (not shown) is 6 to 13 times smaller than the one
explained by EOF_{1BC}, which confirms that sea ice has a much smaller influence than SST or external
forcing. In November, EOF_{1SIC} is a dipole with the same sign between the East European Plain and North
465 America, and an opposite sign over eastern Siberia. In January, EOF_{1SIC} is a dipole with the opposite
signs between North America and Eurasia. The pattern in April is reminiscent of EOF_{1BC}, but with smaller
anomalies. EOF_{1SIC} can hardly be related to the observed snow cover variability, as the correlations
between PC_{1SIC} and the observed PCs are weak. This confirms that the sea ice cover has little or no
impact.

470

The SSTs related to the snow cover changes are given by the regression of the SST anomalies from ALL
| onto PC_{1BC}, ~~using SST anomalies with respect to the 1979–2014 climatology~~. The regressions can be
interpreted as the SST patterns contributing to the snow cover anomalies of EOF_{1BC}, or the SST pattern
responding to the external forcing that affected the snow cover changes. ~~The SIC anomalies might also~~
| ~~play a secondary role~~. We note warm SST anomalies in the western equatorial Pacific and the central
475 midlatitude North Pacific in all months (Fig. 9a-c), consistent with the extended negative snow cover
anomalies of EOF_{1BC} (Fig. 8a,d,g). This might reflect the SST trends observed during that period, which
were found to result from the combination of external forcing superimposed on the changes of the Pacific
decadal variability (Dai and Bloecker, 2018; Gastineau et al. 2019). In April, there is a cold anomaly in
480 the Central and Eastern equatorial Pacific, with an extension toward the eastern North Pacific. This
horseshoe pattern resembles a La Niña pattern and its extension toward mid-latitudes, called the
Interdecadal Pacific Oscillation (Newman et al., 2016). We also note warm SST anomalies over the North
Atlantic in November, which remain similar, but with smaller amplitude, when removing the linear trends
(not shown), suggesting a ~~dominant~~ role of interannual or decadal North Atlantic variability.

485

To understand these links, the SLP anomalies in ALL MMM associated to PC_{1BC} are shown in Fig. 9d-f.
In April, the SST is associated with a weakened Aleutian Low in the North Pacific and a PNA pattern,

which are consistent with the expected pattern associated with cold equatorial Pacific SST anomalies. The weakened Aleutian Low leads to cold air advection over western North America, which explains the positive snow cover anomalies in that region (Fig. 8g). Therefore, the SST over the Pacific Ocean plays a significant role in the April snow cover over North America. The SLP anomalies are otherwise not significant for November and January. We also note positive SST anomalies ~~in~~ into the Atlantic and Pacific oceans, which might reflect the positive SST trend observed during that period at those locations.

To investigate the role of sea-ice driven variability, we calculated the regression onto the $PC1_{SIC}$ index, using the prescribed SIC in ALL, and the MMM difference of SLP from ALL minus ~~NoSICvar~~ $NoSIC$. $PC1_{SIC}$ has a decreasing trend in November (not shown) and it reflects small but significant negative SIC anomalies in the Barents, Labrador, and Bering Seas (Fig. 9g). In November, the ~~associated~~ SLP anomaly is ~~negative~~ low over the negative sea ice anomalies, as expected from the warming in the atmospheric planetary boundary layer, as shown in previous studies (Peings and Magnusdottir, 2014; Liang et al. 2021). The November SLP pattern (Fig. 9g) is different from a negative North Atlantic Oscillation (NAO) or a Ural blocking pattern as previously found as a response to sea ice loss (Mori et al., 2014; Kug et al., 2015; Nakamura et al., 2015; King et al., 2016; Smith et al., 2022), but it is consistent with previous studies using atmosphere-only experiments (Ogawa et al., 2018; Liang et al. 2021). In January and April, the sea ice anomalies are very small. In January, the SLP pattern is insignificant, but in April, there is a cyclonic anomaly around 50°N over the Atlantic Ocean, a small cyclonic anomaly over the Mongolian Plateau, and a weakening of the Aleutian Low. ~~This might be related to the impact of sea ice on the snow cover and soil moisture in spring (Barnett et al. 1989).~~ Positive SLP anomalies are also located over the Kara Sea, but ~~these anomalies~~ cannot be linked to the snow cover anomaly in a simple way. In all cases, removing the linear trends lead to different results (not shown), so the relationships shown in Fig. 9g-i mostly reflect the impact of the sea ice declining trend.

4. Internal variability and climate impacts of snow cover anomalies

4.1 Internal variability of snow cover

515 The impact of internal atmospheric variability on snow cover anomalies is investigated over Eurasia. We
focus on Eurasia as it is the largest continent, and snow cover was previously suggested to influence the
Northern Hemisphere atmospheric circulation (Cohen and Entekhabi, 1999; Cohen et al. 2014; Gastineau
et al. 2017). In the observational datasets, we remove a quadratic trend from all variables, which should
remove a large part of the changes linked to the long-term evolution of external forcing, and then
520 quantifies the internal variability, which is largely due to atmosphere-land processes, in addition to a
residual influence of SST and SIC. The first EOF of the detrended snow cover in ERA5-Land is shown
in Figs. 10a, 10e and 10i. The patterns are somewhat similar to those obtained over the Northern
Hemisphere before detrending (Fig. 7), but the EOF1 over Eurasia in November and April corresponds
to the EOF2 of the Northern Hemisphere before detrending. The similarity in the EOF1 pattern obtained
525 in January in models and observations (Figs. 7d and 10e) suggests that internal variability is dominant
during that month, as opposed to November and April where forcings are important. The regressions of
the SLP and surface air temperature (Figs. 10b,f,j) onto the PC1 at no lag illustrate in January the influence
of the NAO on the snow cover with cold air advection increasing the European snow cover. In November
and April, the snow cover is linked to a trough over the Ural region, with cold (warm) air advection over
530 its western (eastern) flank. The ~~trough~~~~troughs~~ over Central Eurasia ~~is~~~~are~~ part of a wave-like perturbation
between the Atlantic Ocean and Eurasia. In November, the trough extends to eastern Siberia and is
associated with a warming and a snow cover decrease in Central Siberia. These patterns are similar in
January when using the other three snow cover datasets (see Figs. 10d and 10h). In November, the patterns
remain similar except for MERRA-2. In April, the patterns are different (Fig. 10i) ~~and when~~ the SLP
535 anomalies are smaller (Fig. 10j+0j).

To investigate the internal atmospheric-land variability in the model simulations, we repeat the same
analysis except that we remove from snow cover, SLP, and surface air temperature the effect of boundary
conditions by removing for each model the model ensemble mean of the ALL and NoSICvar~~NoSIC~~

540 simulations from the individual members. This removes ~~most a large part~~ of the variability driven by external forcing, SST and SIC. ~~The main-EOF_{Int} is patterns are~~ calculated for each model separately and then, as ~~it is they are~~ found to be similar among models, averaged among the eight models using the ensemble size as a weight (Fig. 11a-f). This robustness is shown by the large pattern correlation of EOF1_{Int} (blue bars) and EOF2_{Int} (orange bars) of each model and those of the multi-model average, with a pattern correlation largest when the model has a large ensemble of 30 members (indicated by black stars in the x-axis numbers of Fig. 11g-i). In November, January, and April, a large monopole with positive snow cover anomalies over the edge of the snow-covered domain appears as the first mode (Fig. 11a-c), with between 9% and 30% of the variance explained. The maximum loading of the monopole is located in Westernwestern Eurasia in November. It shifts to Eastern Europe in January, and moves eastward toward,
550 ~~except in April where it is over~~ Central Siberia in April. The. A dipole is found as a second mode is a dipole, with ~~positive anomalies positioned on from Scandinavia to the~~ northwestern and southeastern endsBlack Sea, and negative anomalies from the Caspian Sea to the eastern coast of the EOF1 patterns; ~~its Siberia. The variance explained~~ variance by the second mode ranges between 6% and 18%.

555 The SLP and surface air temperature associated with the first mode are calculated by a regression onto the PC1_{Int} of the snow cover in each model, which is then averaged similarly among all models. The snow EOF1_{Int} in November, January, and April is always associated at no lag with an anticyclone located at the north or the north-west of the positive snow cover anomalies (Fig. 12a-c), as expected from cold air advection which increases snow cover. However, the location and amplitude of the cold surface air
560 temperature anomalies are not always matching the snow cover anomalies. For instance, the cold anomalies are strongest in January with a widespread cooling extending from western Europe to far eastern Siberia, but the associated snow cover anomalies have the same amplitude in all three months, and are only located over eastern Europe in January. In November, the anticyclone is centered over the Nordic seas (Fig. 12a), while it extends from the Nordic Seas to Northern Siberia in January (Fig. 12b).
565 In April, the snow anomalies cover the southern edge of the snow cover domain, and the anticyclone is centered over the eastern Arctic (Fig. 12c). EOF2_{Int} is associated with a trough over the Ural Mountains

in November and April, while the NAO and the associated temperature anomalies over southern Siberia dominate in January.

570 The comparison between Fig. 10 for observations and Figs. 11-12 for models suggests that the models reproduce fairly well the main mode of the observed variability. In both cases, the NAO is the dominant mode of variability during January. In January, the analysis of the second EOF from detrended ~~found in~~ observations (not shown) also shows that it is, ~~although the~~ associated with patterns similar with that found in models (Figs. 11e and 12e). During November and April, the observed dominant mode of 575 variability is a blocking pattern with a trough over the Ural region. This pattern also occurs in the models, but with less variance, as it is reproduced by EOF2 rather than EOF1. However, there is an important spread among observations in November and April, and is smaller in models. However, the analysis of observation is based on detrended time series that still include the contribution from interannual SST variability and short-term fluctuations in the external forcing.

580 **4.2 Climate influence of snow cover anomalies**

Because of the limited intrinsic atmospheric persistence, the SLP and temperature lagging the snow cover by one month should reflect the snow cover influence in the absence of other concomitant forcings. It is illustrated for observations in Fig. 10c,g,k and for models in Figs. 13-14. In observations, the November EOF1 is followed by negative SLP anomalies over the polar cap and some weak positive anomalies over 585 Western~~western~~ Europe and the Bering Sea. This SLP~~The pattern~~ shares some similarities with that found at no lag, but with a smaller amplitude.~~broadly resembles the positive Arctic Oscillation, which could be linked to the November snow cover influence investigated in Gastineau et al. (2017).~~ In January, the negative NAO anomalies associated in phase with EOF1 are reduced by half but they persist~~persist~~ the following month. In April, no clear~~significant~~ SLP pattern emerges at a lag of one month. In all cases, the 590 surface temperature is cold (warm) over positive (negative) snow cover anomalies, as expected, although their statistical significance is limited.

In models, no robust SLP or temperature anomalies follow the November snow cover EOF1_{Int} with a lag of one or two months (Fig. 13a,d). On the other hand, the January EOF1_{Int} (Fig. 13b) is followed by strong temperature and SLP anomalies at lag 1 (in February) with a strong anticyclonic anomaly over the polar cap extending toward central Eurasia, which is associated with cold continental air advection over northern Eurasia. The patterns are somewhat similar to those shown at lag 0 for January (Fig. 12b). Since they remain largely similar at lag 2 (Fig. 13e), albeit with smaller amplitude, this suggests that snow cover anomalies act as a positive feedback and amplify the AO one (in February) or two (in March) months later. However, it might also reflect an unusually large atmospheric persistence or the presence of a concomitant forcing. The May SLP and temperature anomalies lagging the EOF1_{Int} in April by one month (Fig. 13c) are also similar to the unlagged patterns, albeit smaller, and the air temperature remains cold over the land surfaces covered with positive snow anomalies in April. However, the anomalies are negligible at lag 2 (Fig. 13f). The pattern lagging the snow cover by one month is robust among models in January (Fig. 13h), but the agreement decreases in April (Fig. 13i) and more so in November (Fig. 13g). For the pattern following the January EOF1_{Int} by two months, the agreement among models is smaller than at lag 1, but it remains high in models with 30 members.

The temperature and SLP anomalies lagging the snow cover EOF2_{Int} by one month (Fig. 14a-c) are smaller than for EOF1_{Int} and they vanish at lag 2 (not shown). One month after January EOF2_{Int} there are substantial temperature and SLP perturbations (Fig. 14b), resembling a negative AO pattern as was the case of the forcing pattern (Fig. 12e), but weaker. Its polar center is rather located over Svalbard, and it is less associated with an intensification of the Siberian anticyclone. Negative SLP anomalies are also found over East Asia. On the other hand, only small SLP and surface air temperature anomalies follow the November and April EOF2_{Int}.

The comparison of Fig. 10 with [Figs. 12 and Fig. 13](#) shows a different relationship in models and observations for snow cover in November and April. However, in January, a more extended snow cover over Europe is followed in both models and observation by anticyclonic anomalies over Iceland and negative pressure anomalies over the mid-latitude Atlantic Ocean, as well as cold air temperature

advection toward Europe. As the same relationship is found when the January land snow leads the atmosphere by two months in models, this might indicate a large-scale atmospheric response to the snow cover anomalies. However, an anomalous persistence of ~~the~~ atmospheric anomalies can also be caused by troposphere-stratosphere interactions, increasing the memory of the atmosphere. This hypothesis is investigated in the next section.

4.3 Role of the stratosphere for the January snow cover variability

To understand the mechanism behind the statistical relationship between the January snow cover and the atmosphere, we focus on LMDZOR6 (the ~~#5~~-model ~~#5~~ in Figs. 11g-i, 12g-i, and 13g-i.), as daily outputs and three-dimensional atmospheric fields are available. LMDZOR6 reproduces the links between the January snow cover EOF1_{Int} and the atmosphere one or two months later ~~as~~ shown in Fig. 13, with nearly identical regression maps onto PC1_{Int} ~~when using only LMDZOR6 data~~ (not shown). Figure 15 shows the lag regression of the daily polar cap temperature (north of 60°N) onto the PC1_{Int} of the January snow cover. A significant lower-stratosphere warming is simulated from November to March. At 50-hPa, the temperature anomaly increases from 0.15°C in November to 0.3°C in December and 0.65°C in January. The stratospheric warming and ~~the~~ polar vortex weakening thus precede the January snow cover anomalies. However, the stratospheric temperature shows another maximum anomaly of 1.0.9°C in February, suggesting that the snow cover ~~has~~ might have intensified the polar vortex weakening. The ~~50-hPa geopotential height anomalies associated to the influence of Eurasian~~ snow cover ~~show that~~ the polar vortex weakening is widespread and affects the whole polar cap (Fig. S4).

~~The wave activity flux was calculated to investigate the propagation of the atmosphere could be induced by an amplification of the climatological stationary waves (see Section 2.2). The regression of the wave activity flux on PC1_{Int} shows an amplified upward component planetary wave pattern (Garfinkel et al., 2010), through an intensification of the wave activity flux ridge over eastern Eurasia and the western North Pacific (Fig. S5). This Eastern Europe. Such a planetary wave propagates upward into the stratosphere and may indeed~~ weaken the polar vortex within 10 to 20 days. It is well established that such polar vortex

weakening might then lead to a negative AO in the troposphere, with a lag of a few weeks (Baldwin and Dunkerton, 1999), and episodic downward propagations are ~~indeed~~ visible in Fig. 15 from ~~mid-the-end~~ of January to ~~March~~early April. However, the role of the snow cover anomalies in this mechanism remains to be established.

To do so, we consider in the same model (e.g. LMDZOR6) an index of the stratospheric polar vortex, ~~defined. We define this index~~ as the standardized January polar cap (north of 60°N) temperature anomalies at 50-hPa, ~~hereinafter~~ called PCT₅₀. ~~We also define an February AO index with~~We then remove the ~~first PC of the SLP north of 20°N. The~~ ensemble mean of that model ~~is then removed~~ from all time series. ~~The correlation between~~ PCT₅₀ and ~~PC1_{Int} is weak (0.09) but significant. Both the January PCT₅₀ and PC1_{Int} are significantly correlated to the February AO, but the variance of the AO that is explained by PCT₅₀ is larger than the one associated with the snow cover index, PC1_{Int} (Fig. 15b). Both polar vortex and snow cover remain significantly related to the AO when using a multi-variate regression (not shown). A similar analysis using composites confirms this result and shows that the relationships between both indices and the AO are mostly linear (Fig. S6). Therefore, the snow cover has a significant influence on the AO, but this influence is smaller than the one associated with the polar vortex.~~

~~atmospheric fields.~~ Using ~~then~~ regression onto PCT₅₀, we found that the snow cover anomalies precede the stratospheric warming by 1 or 2 months (~~Fig. 16a,d~~), but not by more. (~~Fig. 16a~~). The same regression using SLP (Fig. ~~16b,e~~16d) shows a dominant Ural blocking pattern in December increasing the snow cover anomalies by easterly cold air advection, and preceding by one month the January polar vortex anomaly. The Ural blocking is associated with negative SLP anomalies over ~~Western~~western Europe and the Aleutians, ~~and~~ thus projects onto the negative AO phase. ~~Ural blocking can also amplify the stationary wave pattern and warm the stratosphere through intensified heat flux due to planetary waves (Martius et al., 2009; Peings, 2019).~~ The regressions are weaker for lag -2, with only a small significant anticyclone east of Scandinavia. The Ural blocking pattern over northern Siberia is consistent with positive snow anomalies south of it. ~~The (Fig. 16b,e). These~~ regressions ~~obtained using PCT₅₀ (Fig. 16b,e)~~ resemble the one obtained with PC1_{Int} (Fig. 16c,f), even if the later anomalies are shifted toward the North Atlantic.

680 ~~In summary, the~~~~This suggests that both~~ snow cover and ~~the~~ polar vortex have a common driver, namely Ural tropospheric blocking. ~~The snow cover and the polar vortex also have a similar influence on the AO one month later. Therefore, the~~, ~~so that the lead-lag~~ relationship between January snow cover and the troposphere in February or March must be interpreted with caution, ~~as~~~~and~~ causality cannot be firmly established. However, the polar vortex anomalies in Fig. 15 show a clear amplification in February, following the January snow cover anomalies. ~~Although the snow cover influence is smaller than the one of the polar vortex, it remains significant when removing the concomitant effect of the polar vortex with a multivariate regression using snow cover and polar vortex indices.~~ This suggests that snow cover anomalies ~~might~~ act as a positive feedback ~~for the AO variability, as they~~, ~~and~~ amplify the combined negative AO and Ural blocking pattern.

5. Discussion and conclusion

690 The land snow over the Northern Hemisphere is investigated in four observational datasets and ~~in~~ two large multi-model ensembles of atmosphere-only experiments, one with prescribed SST, sea ice and external forcing during the 1979-2014 period, and the other in which sea ice variations are replaced by their climatology. Although models ~~simulate different mean states for~~~~have biases in representing the~~ snow cover, the ~~observations show an important spread as found by Mudryck et al. (2017), and both simulated trend and mean state are within the observational spread.~~~~observed snow cover trend is overall well reproduced.~~ In the multi-model ensemble mean, the trend is mainly driven by the external forcings and the associated SST warming. ~~The~~, ~~while~~ sea ice loss ~~only~~ drives a small and insignificant fraction of snow cover ~~trends. The sea ice loss only produces a decreasing snow cover trend located south and downstream of Scandinavia in January and April, and east of Scandinavia downstream of the Barents Sea in November.~~~~anomalies.~~ Cohen et al. (2014), among others, proposed that snow anomalies might amplify or damp the mid-latitude atmospheric circulation response to sea ice loss, but in our experiments, the SIC has little influence in driving the snow cover. However, the lack of two-way coupling between atmosphere, sea-ice and the ocean in our experiments does not allow reproducing realistic links between

snow and sea ice anomalies. Analogous analyses need to be conducted in AOGCMs to further investigate the links between SIC and snow cover.

705

The 1979-2014 investigated period shows a transition from a warm to a cold IPO phase and from a cold to a warm Atlantic multidecadal variability (AMV) phase (Luo et al. 2022). Our analysis suggests that the IPO has a consistent influence among models on the North American snow cover, through the PNA teleconnection pattern. The results for the Eurasian snow cover are ambiguous and cannot be specifically attributed to the SST anomalies or the external forcings. The investigation of simulations focusing on the role of external forcing, such as those realized in the RFMIP (Pincus et al., 2016) panel of CMIP6 would be necessary to distinguish between them. Sensitivity simulations using the observed changes in the AMV and IPO would also be helpful. In addition, including the period after 2014 would be important to understand the climate impacts of the observed reduction in spring snow cover (Mudryk et al. 2020). The simulations investigated here end in 2014 and cannot be used to fully understand this change. A similar multi-model investigation with simulation extended to present-day could be ~~pursued~~pursue in future works.

715

The snow cover variability due to internal atmosphere-land variability is analyzed in the large ensembles of simulation after removing the ensemble mean separately in each model. Lacking a better method, the internal variability is quantified in observations by removing the quadratic trend. We performed an EOF analysis of the Eurasian snow cover and used regression of atmospheric fields on the associated PCs to investigate the atmosphere-snow coupling. We found that the models reproduce well the main atmospheric modes responsible for the forcing of the snow cover in observations, with Ural blocking anomalies embedded with wave-like anomalies leading to dipolar snow cover anomalies in early spring and early winter. In mid-winter, the NAO has the dominant influence, increasing the snow cover over ~~eastern~~-Europe and inducing a widespread Eurasian cooling for negative NAO phases. In both observations and models~~In observation~~, we found no robust circulation pattern following that the November snow anomalies. This seems to contradict~~leads to AO-like anomalies by one month. Such a relationship is not reproduced in models, seemingly contradicting~~ the results of Gastineau et al. (2017),

725

730

~~where~~when CMIP5 preindustrial control simulations were found to partly reproduce the observed influence of November Eurasian snow. This emphasizes the need to better explore the observational uncertainties, using various data sets. ~~influence. However, as mentioned above,~~ sea ice concentration and SST anomalies are prescribed in our atmosphere-only simulations and cannot respond to the atmospheric forcing ~~as. This is different~~ in observations or coupled simulations, where snow cover, sea ice and SST are driven by the atmosphere, and provide concomitant forcings. Both models and observations show that January eastern European snow cover anomalies are linked to AO-like anomalies one month later. The relationship remains significant at a lag of two months in models. The outputs from one of the models reveal that the Ural blocking pattern acts as a common driver for the Eastern Europe snow and the polar vortex anomalies. A stratospheric warming (cooling) event and a polar vortex weakening (strengthening) areis, therefore, found to precede by two months the snow cover and negative (positive) AO changes in January. The stratospheric warming is found to explain a larger AO variance fraction than the snow cover. The lead-lag relationship between snow cover and atmosphere~~This lead-lag relationship~~ might only stem from the internal atmospheric variability, as proposed by Blackport and Screen (2021) for the case of the sea-ice atmosphere interaction that shares a large similarity with the interaction discussed here. However, we note that polar vortex anomalies (Fig. 15) are reinforced in February, one month after snow cover anomalies, and are much weaker before January. Such asymmetry suggests that the snow cover ~~intensifies acts as a positive feedback onto the combined AO-Ural blocking pattern, intensifying~~ the stratospheric warming and the resulting AO-like anomalies produced by downward propagation one month later. This mechanism is supported by the intensification of the upward propagating planetary waves following a larger snow cover extent in January, as found by Lü et al. (2020). This suggests a two-way coupling between the snow cover and the internal atmospheric variability~~atmosphere~~, where the snow cover anomalies amplify the AO-Ural blocking anomalies that generated them, ~~acting. This coupling with snow cover might act~~ as a positive feedback ~~in~~for the internal variability of the land-troposphere-stratosphere system. This conclusion agrees with the results from subseasonal forecasts (Orsolini et al., 2015; Garfinkel et al., 2020), where the snow cover through troposphere-stratosphere interaction reinforces and prolongs the NAO.

760 Further investigating causality and the specific role of the snow cover~~Investigating these hypotheses~~ would require sensitivity experiments controlling the land snow cover, or the use of specific statistics to investigate the causation (San Liang, 2014; Runge et al., 2015).

Code and data availability

765 This study is based on publicly available data for observations. The climate model simulations are available upon request from the authors of this study. Matlab code for data analysis and scripts used to generate the map (ferret and python) can be obtained upon request from the corresponding author. The Plumb wave activity flux is calculated using the code from Kazuaki Nishii (<http://www.atmos.rcast.u-tokyo.ac.jp/nishii/index.html>).

770 **Supplement**

The supplement related to this article is available online at:

Author contributions

775 All authors contributed to the realization of the model simulations. GG conceived this study, ~~performed~~performed the analysis, and wrote the first text version. All authors provided edits, comments, and inputs for writing the final text.

Competing interests

The authors declare that they have no conflict of interest.

780

Acknowledgments

We acknowledge support by the Blue-Action Project (European Union's Horizon 2020 research and innovation programme, #727852, <http://www.blue-action.eu/index.php?id=3498>). The LOCEAN-IPSL group was granted access to the HPC resources of TGCC under the allocation A5-017403 and A7-017403
785 made by GENCI. GG and CF were funded by the JPI climate/JPI Ocean ROADMAP project (ANR-19-

JPOC-003). E.M. and D.M. acknowledge the support of the German Federal Ministry of Education and Research through the JPI Climate/JPI Oceans NextG-Climate Science-ROADMAP (FKZ: 01LP2002A) Project. The NorESM2-CAM6 simulations were performed on resources provided by UNINETT Sigma2 - the National Infrastructure for High Performance Computing and Data Storage in Norway (nn2343k, 790 NS9015K). Y-OK, CF and Y-CL are supported by the US National Science Foundation Office of Polar Program (OPP-1736738, OPP-2106190).

References

- Baldwin, M.P. and Dunkerton, T.J.: Propagation of the Arctic Oscillation from the stratosphere to the 795 troposphere. *Journal of Geophysical Research: Atmospheres*, 104(D24), 30937–30946. <https://doi.org/10.1029/1999JD900445>, 1999.
- ~~Barnett, T. P., Dümenil, L., Schlese, U., Roeckner, E., & Latif, M.: The effect of Eurasian snow cover on regional and global climate variations. *Journal of the Atmospheric Sciences*, 46(5), 661–686, 1989~~
- Blackport, R., & Screen, J. A.: Observed statistical connections overestimate the causal effects of arctic 800 sea ice changes on midlatitude winter climate. *Journal of Climate*, 34(8), 3021–3038, 2021.
- Brown, R., Derksen, C., & Wang, L.: A multi-data set analysis of variability and change in Arctic spring snow cover extent, 1967–2008. *Journal of Geophysical Research: Atmospheres*, 115(D16). <https://doi.org/10.1029/2010JD013975>, 2010.
- Brun, E., Six, D., Picard, G., Vionnet, V., Arnaud, L., Bazile, E., Boone, A., Bouchard, A., Genthon, C., 805 Guidard, V., Moigne, P. L., Rabier, F., & Seity, Y.: Snow/atmosphere coupled simulation at Dome C, Antarctica. *Journal of Glaciology*, 57(204), 721–736. Cambridge Core. <https://doi.org/10.3189/002214311797409794>, 2011.
- ~~Brown, R. D., & Derksen, C. (2013). Is Eurasian October snow cover extent increasing?. *Environmental Research Letters*, 8(2), 024006. <https://doi.org/10.1088/1748-9326/8/2/024006>~~
- 810 Cohen, J., & Entekhabi, D.: Eurasian snow cover variability and northern hemisphere climate predictability. *Geophysical Research Letters*, 26(3), 345–348. <https://doi.org/10.1029/1998GL900321>, 1999.

Cohen, J., Furtado, J. C., Jones, J., Barlow, M., Whittleston, D., & Entekhabi, D.: Linking Siberian Snow Cover to Precursors of Stratospheric Variability. *Journal of Climate*, 27(14), 5422–5432.

815 <https://doi.org/10.1175/JCLI-D-13-00779.1>, 2014.

[Copernicus Climate Change Service \(C3S\): ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service Climate Data Store \(CDS\), May 8th 2020, https://cds.climate.copernicus.eu/cdsapp#!/home, 2017.](https://cds.climate.copernicus.eu/cdsapp#!/home)

820 [Copernicus Climate Change Service \(C3S\): C3S ERA5-Land reanalysis. Copernicus Climate Change Service, April 9th 2020. https://cds.climate.copernicus.eu/cdsapp#!/home, 2019.](https://cds.climate.copernicus.eu/cdsapp#!/home)

Dai, A., Bloecker, C.E.: Impacts of internal variability on temperature and precipitation trends in large ensemble simulations by two climate models. *Clim Dyn* 52, 289–306. <https://doi.org/10.1007/s00382-018-4132-4>, 2019.

825 Derksen, C., & Brown, R.: Spring snow cover extent reductions in the 2008–2012 period exceeding climate model projections. *Geophysical Research Letters*, 39(19) <https://doi.org/10.1029/2012GL053387>, 2012.

Déry, S. J., & Brown, R. D.: Recent Northern Hemisphere snow cover extent trends and implications for the snow-albedo feedback. *Geophysical Research Letters*, 34(22), 2007.

830 Deser, C., Phillips, A., Bourdette, V., & Teng, H.: Uncertainty in climate change projections: The role of internal variability. *Climate Dynamics*, 38(3), 527–546, 2012.

Deser, C., Tomas, R. A., & Peng, S.: The Transient Atmospheric Circulation Response to North Atlantic SST and Sea Ice Anomalies. *Journal of Climate*, 20(18), 4751–4767. <https://doi.org/10.1175/JCLI4278.1>, 2007.

835 [Deser, C., R. A. Tomas, and L. Sun, 2015: The Role of Ocean–Atmosphere Coupling in the Zonal-Mean Atmospheric Response to Arctic Sea Ice Loss. *J. Climate*, 28, 2168–2186, https://doi.org/10.1175/JCLI-D-14-00325.1.](https://doi.org/10.1175/JCLI-D-14-00325.1)

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.*, 9(5), 1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.

- 840 Feldstein, S. B.: The Timescale, Power Spectra, and Climate Noise Properties of Teleconnection
Patterns. *Journal of Climate*, 13(24), 4430–4440. [https://doi.org/10.1175/1520-0442\(2000\)013<4430:TTPSAC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<4430:TTPSAC>2.0.CO;2), 2000.
- Fletcher, C. G., Hardiman, S. C., Kushner, P. J., & Cohen, J.: The Dynamical Response to Snow Cover
Perturbations in a Large Ensemble of Atmospheric GCM Integrations. *Journal of Climate*, 22(5), 1208–
845 1222. <https://doi.org/10.1175/2008JCLI2505.1>, 2009.
- [Fox-Kemper, B., H.T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S.S. Drijfhout, T.L. Edwards, N.R. Golledge, M. Hemer, R.E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I.S. Nurhati, L. Ruiz, J.-B. Sallée, A.B.A. Slangen, and Y. Yu, 2021: Ocean, Cryosphere and Sea Level Change. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1211–1362, \[https://doi.org/10.1017/9781009157896.011_2021\]\(https://doi.org/10.1017/9781009157896.011_2021\)](#)
- 850 [García-Serrano, J., Frankignoul, C., Gastineau, G., & de la Cámara, A.: On the Predictability of the Winter Euro-Atlantic Climate: Lagged Influence of Autumn Arctic Sea Ice. *Journal of Climate*, 28\(13\), 5195–5216. <https://doi.org/10.1175/JCLI-D-14-00472.1>, 2015.](#)
- 855 [Garfinkel, C. I., Hartmann, D. L., & Sassi, F.: Tropospheric Precursors of Anomalous Northern Hemisphere Stratospheric Polar Vortices, *Journal of Climate*, 23\(12\), 3282-3299. <https://doi.org/10.1175/2010JCLI3010.1>, 2010.](#)
- [Garfinkel, C. I., Schwartz, C., White, I. P., & Rao, J. \(2020\). Predictability of the early winter Arctic oscillation from autumn Eurasian snowcover in subseasonal forecast models. *Climate Dynamics*, 55, 961-974. <https://doi.org/10.1007/s00382-020-05305-3>](#)
- 860 [Gastineau, G., & Frankignoul, C.: Influence of the North Atlantic SST Variability on the Atmospheric Circulation during the Twentieth Century. *Journal of Climate*, 28\(4\), 1396–1416. <https://doi.org/10.1175/JCLI-D-14-00424.1>, 2015.](#)
- 865 [Gastineau, G., García-Serrano, J., & Frankignoul, C.: The Influence of Autumnal Eurasian Snow Cover on Climate and Its Link with Arctic Sea Ice Cover. *Journal of Climate*, 30\(19\), 7599–7619. <https://doi.org/10.1175/JCLI-D-16-0623.1>, 2017.](#)

- Gastineau, G., Friedman, A.R., Khodri, M. *et al.*: Global ocean heat content redistribution during the 1998–2012 Interdecadal Pacific Oscillation negative phase. *Clim Dyn* **53**, 1187–1208.
- 870 <https://doi.org/10.1007/s00382-018-4387-9>, 2019.
- Ge, Y., & Gong, G.: North American snow depth and climate teleconnection patterns. *Journal of Climate*, *22*(2), 217–233. <https://doi.org/10.1175/2008JCLI2124.1>, 2009.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, 875 V., Conaty, A., da Silva, A. M., Gu, W., ... Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2). *Journal of Climate*, *30*(14), 5419–5454. <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Global Modeling and Assimilation Office (GMAO): MERRA-2 tavgM_2d_Ind_Nx: 2d, Monthly mean, Time-Averaged, Single-Level, Assimilation, Land Surface Diagnostics V5.12.4, Greenbelt, MD, USA, 880 Goddard Earth Sciences Data and Information Services Center (GES DISC), Accessed: [Oct. 28th 2022], <https://doi.org/10.5067/8S35XF81C28F>, 2015.
- Gong, G., Entekhabi, D., & Cohen, J.: Modeled Northern Hemisphere Winter Climate Response to Realistic Siberian Snow Anomalies. *Journal of Climate*, *16*(23), 3917–3931. [https://doi.org/10.1175/1520-0442\(2003\)016<3917:MNHWCR>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<3917:MNHWCR>2.0.CO;2), 2003.
- 885 Gulev, S.K., P.W. Thorne, J. Ahn, F.J. Dentener, C.M. Domingues, S. Gerland, D. Gong, D.S. Kaufman, H.C. Nnamchi, J. Quaas, J.A. Rivera, S. Sathyendranath, S.L. Smith, B. Trewin, K. von Schuckmann, and R.S. Vose: Changing State of the Climate System. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United 890 Kingdom and New York, NY, USA, pp. 287–422, doi:10.1017/9781009157896.004, 2021.
- Guo, H., Yang, Y., Zhang, W., Zhang, C., & Sun, H.: Attributing snow cover extent changes over the Northern Hemisphere for the past 65 years. *Environmental Research Communications*, *3*(6), 061001, 2021.
- Haarsma, R. J., Roberts, M. J., Vidale, P. L., Senior, C. A., Bellucci, A., Bao, Q., Chang, P., Corti, S., 895 Fučkar, N. S., Guemas, V., von Hardenberg, J., Hazeleger, W., Kodama, C., Koenigk, T., Leung, L. R.,

- Lu, J., Luo, J.-J., Mao, J., Mizielinski, M. S., ... von Storch, J.-S.: High Resolution Model Intercomparison Project (HighResMIP v1.0) for CMIP6. *Geosci. Model Dev.*, 9(11), 4185–4208. <https://doi.org/10.5194/gmd-9-4185-2016>, 2016.
- Henderson, G. R., Peings, Y., Furtado, J. C., & Kushner, P. J.: Snow–atmosphere coupling in the Northern Hemisphere. *Nature Climate Change*, 8(11), 954–963, 2018.
- 900 [Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J.-N. \(2019\): ERA5 monthly averaged data on single levels from 1959 to present. Copernicus Climate Change Service \(C3S\) Climate Data Store \(CDS\). \(Accessed on 08-May-2020\), https://doi.org/10.24381/cds.fl7050d7](https://doi.org/10.24381/cds.fl7050d7)
- 905 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., ... Thépaut, J.-N.: The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>, 2020.
- Honda, M., Inoue, J., & Yamane, S. (2009). Influence of low Arctic sea-ice minima on anomalously cold Eurasian winters. *Geophysical Research Letters*, 36(8). <https://doi.org/10.1029/2008GL037079>
- 910 Hurrell, J. W., Hack, J. J., Shea, D., Caron, J. M., & Rosinski, J.: A New Sea Surface Temperature and Sea Ice Boundary Dataset for the Community Atmosphere Model. *Journal of Climate*, 21(19), 5145–5153. <https://doi.org/10.1175/2008JCLI2292.1>, 2008.
- King, M. P., Hell, M., & Keenlyside, N.: Investigation of the atmospheric mechanisms related to the autumn sea ice and winter circulation link in the Northern Hemisphere. *Climate Dynamics*, 46(3), 1185–1195. <https://doi.org/10.1007/s00382-015-2639-5>, 2016.
- Krinner, G., Derksen, C., Essery, R., Flanner, M., Hagemann, S., Clark, M., ... & Zhu, D.: ESM-SnowMIP: assessing snow models and quantifying snow-related climate feedbacks. *Geoscientific Model Development*, 11(12), 5027-5049. <https://doi.org/10.5194/gmd-11-5027-2018>, 2018.
- 920 Kushnir, Y., Robinson, W. A., Bladé, I., Hall, N. M. J., Peng, S., & Sutton, R.: Atmospheric GCM Response to Extratropical SST Anomalies: Synthesis and Evaluation. *Journal of Climate*, 15(16), 2233–2256. [https://doi.org/10.1175/1520-0442\(2002\)015<2233:AGRTES>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<2233:AGRTES>2.0.CO;2), 2002.

- Kug, JS., Jeong, JH., Jang, YS. *et al.*: Two distinct influences of Arctic warming on cold winters over North America and East Asia. *Nature Geosci* **8**, 759–762 <https://doi.org/10.1038/ngeo2517>, 2015.
- 925 Lau, N.-C.: Interactions between Global SST Anomalies and the Midlatitude Atmospheric Circulation. *Bulletin of the American Meteorological Society*, 78(1), 21–34. [https://doi.org/10.1175/1520-0477\(1997\)078<0021:IBGSAA>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<0021:IBGSAA>2.0.CO;2), 1997.
- Liang, Y.-C., Frankignoul, C., Kwon, Y.-O., Gastineau, G., Manzini, E., Danabasoglu, G., Suo, L., Yeager, S., Gao, Y., Attema, J. J., Cherchi, A., Ghosh, R., Matei, D., Mecking, J. V., Tian, T., & Zhang, Y.: Impacts of Arctic Sea Ice on Cold Season Atmospheric Variability and Trends Estimated from Observations and a Multimodel Large Ensemble. *Journal of Climate*, 34(20), 8419–8443. <https://doi.org/10.1175/JCLI-D-20-0578.1>, 2021.
- 930 Liang, Y.-C., Kwon, Y.-O., Frankignoul, C., Danabasoglu, G., Yeager, S., Cherchi, A., Gao, Y., Gastineau, G., Ghosh, R., Matei, D., Mecking, J. V., Peano, D., Suo, L., & Tian, T.: Quantification of the Arctic Sea Ice-Driven Atmospheric Circulation Variability in Coordinated Large Ensemble Simulations. *Geophysical Research Letters*, 47(1), e2019GL085397. <https://doi.org/10.1029/2019GL085397>, 2020.
- López-Parages, J., Rodríguez-Fonseca, B., Dommenges, D., & Frauen, C.: ENSO influence on the North Atlantic European climate: A non-linear and non-stationary approach. *Climate Dynamics*, 47(7), 2071–2084. <https://doi.org/10.1007/s00382-015-2951-0>, 2016.
- 940 [Lü, Z., Li, F., Orsolini, Y. J., Gao, Y., & He, S. \(2020\). Understanding of European Cold Extremes, Sudden Stratospheric Warming, and Siberian Snow Accumulation in the Winter of 2017/18, *Journal of Climate*, 33\(2\), 527-545. https://doi.org/10.1175/JCLI-D-18-0861.1](https://doi.org/10.1175/JCLI-D-18-0861.1)
- Luo, B., Luo, D., Dai, A., Simmonds, I., & Wu, L.: The modulation of Interdecadal Pacific Oscillation and Atlantic Multidecadal Oscillation on winter Eurasian cold anomaly via the Ural blocking change. *Climate Dynamics*, 1-24. <https://doi.org/10.1007/s00382-021-06119-7>, 2022.
- [Martius, O., Polvani, L. M., and Davies, H. C. \(2009\), Blocking precursors to stratospheric sudden warming events, *Geophys. Res. Lett.*, 36, L14806, https://doi.org/10.1029/2009GL038776.](https://doi.org/10.1029/2009GL038776)

- Mathieu, P.-P., Sutton, R. T., Dong, B., & Collins, M.: Predictability of Winter Climate over the North Atlantic European Region during ENSO Events. *Journal of Climate*, 17(10), 1953–1974. [https://doi.org/10.1175/1520-0442\(2004\)017<1953:POWCOT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<1953:POWCOT>2.0.CO;2), 2004.
- Mori, M., Watanabe, M., Shiogama, H. *et al.*: Robust Arctic sea-ice influence on the frequent Eurasian cold winters in past decades. *Nature Geosci* 7, 869–873, <https://doi.org/10.1038/ngeo2277>, 2014.
- Mudryk, L. R. *et al.* : Characterization of Northern Hemisphere Snow Water Equivalent Datasets, 1981–2010, *Journal of Climate*. 28. 8037-8051. <https://doi.org/10.1175/JCLI-D-15-0229.1>, 2015.
- Mudryk, L. R. and C. Derksen: CanSISE Observation-Based Ensemble of Northern Hemisphere Terrestrial Snow Water Equivalent, Version 2. [Daily B5 SWE]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. doi: <https://doi.org/10.5067/96ltniikJ7vd>. [Accessed Aug. 16th 2019], 2017.
- 960 [Mudryk, L. R., Kushner, P. J., Derksen, C., and Thackeray, C. \(2017\), Snow cover response to temperature in observational and climate model ensembles, *Geophys. Res. Lett.*, 44, 919– 926, <https://doi.org/10.1002/2016GL071789>.](https://doi.org/10.1002/2016GL071789)
- Mudryk, L., Santolaria-Otín, M., Krinner, G., Ménégos, M., Derksen, C., Brutel-Vuilmet, C., Brady, M., & Essery, R.: Historical Northern Hemisphere snow cover trends and projected changes in the CMIP6 multi-model ensemble. *The Cryosphere*, 14(7), 2495–2514. <https://doi.org/10.5194/tc-14-2495-2020>, 2020.
- 970 [Muñoz-Sabater, J., \(2019\): ERA5-Land monthly averaged data from 1981 to present. Copernicus Climate Change Service \(C3S\) Climate Data Store \(CDS\). \(Accessed on 04-Apr-2020\), <https://doi.org/10.24381/cds.68d2bb3>](https://doi.org/10.24381/cds.68d2bb3)
- [Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C., and Thépaut, J.-N.: ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, *Earth Syst. Sci. Data*, 13, 4349–4383, <https://doi.org/10.5194/essd-13-4349-2021>, 2021.](https://doi.org/10.5194/essd-13-4349-2021)

- 975 Nakamura, T, Yamazaki, K, Iwamoto, K, Honda, M, Miyoshi, Y, Ogawa, Y, and Ukita, J.: A negative phase shift of the winter AO/NAO due to the recent Arctic sea-ice reduction in late autumn. *J. Geophys. Res. Atmos.*, 120, 3209– 3227. doi: [10.1002/2014JD022848](https://doi.org/10.1002/2014JD022848), 2015.
- Newman, M., Alexander, M. A., Ault, T. R., Cobb, K. M., Deser, C., Di Lorenzo, E., ... & Smith, C. A.: The Pacific decadal oscillation, revisited. *Journal of Climate*, 29(12), 4399-4427, 2016.
- 980 Ogawa, F., Keenlyside, N., Gao, Y., Koenigk, T., Yang, S., Suo, L., ... & Semenov, V.: Evaluating impacts of recent Arctic sea ice loss on the northern hemisphere winter climate change. *Geophysical Research Letters*, 45(7), 3255-3263, 2018.
- Orsolini, Y. J., Senan, R., Balsamo, G., Doblas-Reyes, F. J., Vitart, F., Weisheimer, A., Carrasco, A., & Benestad, R. E.: Impact of snow initialization on sub-seasonal forecasts. *Climate Dynamics*, 41(7), 985 1969–1982. <https://doi.org/10.1007/s00382-013-1782-0>, 2013.
- [Orsolini, Y. J., Senan, R., Vitart, F., Balsamo, G., Weisheimer, A., & Doblas-Reyes, F. J. \(2016\). Influence of the Eurasian snow on the negative North Atlantic Oscillation in subseasonal forecasts of the cold winter 2009/2010. *Climate Dynamics*, 47, 1325-1334. <https://doi.org/10.1007/s00382-015-2903-8>](https://doi.org/10.1007/s00382-013-1782-0)
- 990 Paik, S., & Min, S. K.: Quantifying the Anthropogenic Greenhouse Gas Contribution to the Observed Spring Snow-Cover Decline Using the CMIP6 Multimodel Ensemble. *Journal of Climate*, 33(21), 9261-9269, 2020.
- Peings, Y., & Magnusdottir, G.: Response of the wintertime Northern Hemisphere atmospheric circulation to current and projected Arctic sea ice decline: A numerical study with CAM5. *Journal of* 995 *Climate*, 27(1), 244-264, 2014.
- Peings, Y.: Ural blocking as a driver of early-winter stratospheric warmings. *Geophysical Research Letters*, 46(10), 5460-5468, 2019. <https://doi.org/10.1029/2019GL082097>
- Perlwitz, J., & Graf, H.-F.: The Statistical Connection between Tropospheric and Stratospheric Circulation of the Northern Hemisphere in Winter. *Journal of Climate*, 8(10), 2281–2295. 000 [https://doi.org/10.1175/1520-0442\(1995\)008<2281:TSCBTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(1995)008<2281:TSCBTA>2.0.CO;2), 1995.
- Pincus, R., Forster, P. M., & Stevens, B.: The Radiative forcing model intercomparison project (RFMIP): experimental protocol for CMIP6. *Geoscientific Model Development*, 9(9), 3447-3460, 2016.

- 005 [Plumb, R. A. \(1985\). On the Three-Dimensional Propagation of Stationary Waves, *Journal of Atmospheric Sciences*, 42\(3\), 217-229. \[https://doi.org/10.1175/1520-0469\\(1985\\)042<0217:OTTDPO>2.0.CO;2\]\(https://doi.org/10.1175/1520-0469\(1985\)042<0217:OTTDPO>2.0.CO;2\)](https://doi.org/10.1175/1520-0469(1985)042<0217:OTTDPO>2.0.CO;2)
- Pulliainen, J. et al.: Early snowmelt significantly enhances boreal springtime carbon uptake. *Proc. Natl Acad. Sci. USA* **114**, 11081–11086, 2017.
- Pulliainen, J., Luojuus, K., Derksen, C., Mudryk, L., Lemmetyinen, J., Salminen, M., Ikonen, J., Takala, M., Cohen, J., Smolander, T., & Norberg, J.: Patterns and trends of Northern Hemisphere snow mass
010 from 1980 to 2018. *Nature*, 581(7808), 294–298. <https://doi.org/10.1038/s41586-020-2258-0>, 2020.
- Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C.-J., Arsenault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D., & Toll, D.: The Global Land Data Assimilation System. *Bulletin of the American Meteorological Society*, 85(3), 381–394. <https://doi.org/10.1175/BAMS-85-3-381>, 2004.
- 015 Robinson, D. A. and T. W. Estilow: *Rutgers Northern Hemisphere 24 km Weekly Snow Cover Extent, September 1980 Onward, Version 1*. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi: <https://doi.org/10.7265/zzbm-2w05>. [accessed Jul. 22th 2019], 2021.
- Runge, J., Petoukhov, V., Donges, J. F., Hlinka, J., Jajcay, N., Vejmelka, M., ... & Kurths, J.: Identifying causal gateways and mediators in complex spatio-temporal systems. *Nature*
020 *communications*, 6(1), 1-10, 2015.
- Saito, K., & Cohen, J.: The potential role of snow cover in forcing interannual variability of the major Northern Hemisphere mode. *Geophysical Research Letters*, 30(6). <https://doi.org/10.1029/2002GL016341>, 2003.
- San Liang, X.: Unraveling the cause-effect relation between time series. *Physical Review E*, 90(5),
025 052150, 2014.
- Scaife, A. A., Arribas, A., Blockley, E., Brookshaw, A., Clark, R. T., Dunstone, N., Eade, R., Fereday, D., Folland, C. K., Gordon, M., Hermanson, L., Knight, J. R., Lea, D. J., MacLachlan, C., Maidens, A., Martin, M., Peterson, A. K., Smith, D., Vellinga, M., ... Williams, A.: Skillful long-range prediction of European and North American winters. *Geophysical Research Letters*, 41(7), 2514–2519.
030 <https://doi.org/10.1002/2014GL059637>, 2014.

Simon, A., Frankignoul, C., Gastineau, G., & Kwon, Y. O.: An observational estimate of the direct response of the cold-season atmospheric circulation to the Arctic Sea Ice Loss. *Journal of Climate*, 33(9), 3863-3882, 2020.

035 Smith, D. M., Eade, R., Andrews, M. B., Ayres, H., Clark, A., Chripko, S., ... & Walsh, A.: Robust but weak winter atmospheric circulation response to future Arctic sea ice loss. *Nature communications*, 13(1), 1-15, 2022.

[Sturm, M., Taras, B., Liston, G. E., Derksen, C., Jonas, T., & Lea, J. \(2010\). Estimating Snow Water Equivalent Using Snow Depth Data and Climate Classes, *Journal of Hydrometeorology*, 11\(6\), 1380-1394. <https://doi.org/10.1175/2010JHM1202.1>](https://doi.org/10.1175/2010JHM1202.1)

040 Von Storch, H., & Zwiers, F. W.: *Statistical analysis in climate research*. Cambridge university press, 1999.

Wallace, J. M., & Gutzler, D. S.: Teleconnections in the Geopotential Height Field during the Northern Hemisphere Winter. *Monthly Weather Review*, 109(4), 784–812. [https://doi.org/10.1175/1520-0493\(1981\)109<0784:TITGHF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1981)109<0784:TITGHF>2.0.CO;2), 1981.

045 Zhong, X., Zhang, T., Kang, S., & Wang, J. Snow depth trends from CMIP6 models conflict with observational evidence. *Journal of Climate*, 35(4), 1293-1307. <https://doi.org/10.1175/JCLI-D-21-0177.1>, 2022.

050

Table 1. Summary of the AGCMs used in this study. The short name is used in the text of this study instead of the full model name.

Model Name (short name)	#	Institution	Horizontal resolution (lat x lon)	# of vertical levels (top level)	# of members	Adjustment of SST/SIC	Snow cover available	CMIP6 External Forcing used	Reference
EC-Earth3 (EC-Earth3)	1	DMI	T255 (~80 km)	91 (0.01 hPa)	20	Yes	No	CMIP6	Döscher et al., (2022)
HadGEM3-GC3.1 (HadGEM3)	2	UoS	0.83° x 0.55° (~60 km)	85 (85 km)	10	No	No	HighResMIP	Walters et al. (2017)
ECHAM6.3 (ECHAM6)	3	MPI-M	T127 (~100km)	95 (0.01hPa)	10	Yes	No	CMIP6	Stevens et.al.(2013) Mueller et. al. (2018)
IAP4.1 (IAP4)	4	IAP	1.4° x1.4°	30 (2.2hPa)	15	Yes	No	1979-2005: CMIP5 historical 2006-2013: CMIP5 RCP8.5	Sun et al. (2012)
LMDZOR6 (LMDZOR6-LMDZ6)	5	LOCEAN-IPSL	1.26° x 2.5° (~150 km)	79 (0.01 hPa)	30	Yes	Yes	HighResMIP	Hourdin et al. (2020)
NorESM2-CAM6 (NorESM)	6	NERSC	0.94° x 1.25° (~100 km)	32 (3.4 hPa)	30	Yes	Yes	CMIP6	Bentsen et al. (2013) Seland et al. (2020)
CESM2-WACCM6 (CESM2)	7	WHOI-NCAR	0.94° x 1.25° (~100 km)	70 (4.5x10 ⁻⁶ hPa)	30	Yes	Yes	CMIP6	Gettelman et al. (2019)
CMCC-CM2-HR4 (CMCC)	8	CMCC	0.9° x 1.25° (~100 km)	30 (2 hPa)	10	No	Yes	HighResMIP	Cherchi et al. (2018)




Figure 1: (Top) Seasonal cycle of the area covered by snow, in 10^6 km^2 , in (a) North America and (b) Eurasia. (Bottom) Seasonal cycle of the Northern Hemisphere snow mass, in 10^{12} kg , in (c) North America and (d) Eurasia. Color curves show results from models. Thick black curves show observations. The grey shade provides the observational spread obtained from CanSISE.




Figure 2: (a) Mean January snow cover in CanSISE, in fraction. (b-i) ~~Mean~~ mean snow cover (color shade) bias in January, in fraction, calculated as the ALL ensemble mean minus CanSISE for each model. The contours indicate the mean snow cover fraction in CanSISE (contour interval 0.2) shown in (a).




Figure 3: Same as Fig. 2, but for the snow water equivalent, in m.




Figure 4: Time series of the anomalous area covered by snow in 10^6 km^2 (first row) and anomalous snow mass in 10^{12} kg (second row) in winter (from November to April), in (right) North America and (left) Eurasia in observation and the simulation ALL. (Third row) Same as first row, but for ALL minus NoSICvar. (Fourth row) Same as the second row, but for the difference ALL minus NoSICvar. Note the different scale in the y-axis for (left axis, black line) models and (right axis, blue line) ERA5-Land and CanSISE observations. The grey shading indicates the range between the minimum and the maximum values among the eight models. The black curve is the multi-model ensemble mean. The correlation between the multi-model mean and ERA5-Land is given in the bottom left corner of each panel, the star symbol indicating a p-value below 5%.





Figure 5: Trend in Northern Hemisphere (trend in a) snow mass over extent and (b) snow extent mass. Correlation between with ERA5-Land and simulated values for (c) snow mass over extent and (d) snow extent mass. Red (blue) symbol denotes the simulation ALL (-ALL minus NoSICvar/NoSIC) for 1979-2014. Grey symbol denotes the CanSISE observations for 1979-2010. Orange (blue sky) symbol is for the simulation ALL (ALL minus NoSICvar/NoSIC) when using linearly detrended fields. Northern Hemisphere trend in snow cover, in % per decade¹, for the multi-model mean of simulations (e-g) ALL and (h-j) the difference of (h-j) ALL minus NoSICvar/NoSIC; in (e) (h) November (f) (i) January and (g) (j) April. In (a-d), the size of the symbol is large (small) if the significance of the trends is below (above) 5%. In (e-j), black contours indicate where the p-values are lower than 5%.

55

60



65 Figure 6: Variance fraction (in %) of the interannual snow ~~coverwater equivalent~~ anomalies explained by SST, external forcing and sea ice concentration; for (a) November, (b) January and (c) April. The thick black curve indicates where the p-value of the ANOVA test is lower than 5% for at least 5 models out of 8. Variance fraction (in %) of the snow cover area over Eurasia (Eur) and North America (NAM) calculated for each model ~~separatelyseparaty~~ for (d) November, (e) January and (f) April. The number from 1 to 8 designates the model (see table 1 for correspondence with model name). The symbol star indicates when 30 members are available for both ALL and NoSICvar. The ~~effect of the~~ factor t , representing the influence of SST and external forcings, is ~~referredreferred~~ to as SST/ext. The ~~effect of the~~ factor e , representing the influence of SIC on the time mean snow, is ~~referredreferred~~ to as SIC. The ~~effect of the~~ factor $t:e$, representing the influence of SIC on the time-varying snow, is ~~referredreferred~~ to as SST/Ext:SIC. Numbers are given in bold font if the ANOVA test has a p-value below 5%.

70





Figure 7: (Left column) First and (center column) second EOFs of Northern Hemisphere snow cover anomalies in ERA5-Land during 1981-2014 for (first row) November, (second row) January and (third row) April. The variance explained by the EOFs is indicated. (Right column) Pattern correlation between the (blue bar) EOF1 and (orange bar) EOF2 of ERA5-Land and that found in other observational datasets MERRA2, NOAA-CDR and CanSISE.



85 Figure 8: (Left column) First EOF_{BC} of the snow cover anomalies associated with sea surface temperature and external forcing anomalies and (center column) first EOF_{SIC} associated with sea ice concentration anomalies, for (first row) November, (second row),
 90 The blue (orange) bars and the associated crosses provide the results when using PC1 (PC2) from observations. The dashed line provides provide the 5% -and level of statistical significance for the correlation when using ERA5-Land, MERRA2 or NOAA-CDR in 1981-2014 (32 degrees of freedom). The dotted line is the same as the dashed line but when using CanSISE in 1981-2010 (28 degrees of freedom).






Figure 9: (a) (b) (c) Regression of the SST, in $^{\circ}\text{C}$ (gray contours and color shade) on PC1_{BC} ; (d) (e) (f) Regression of the SST, in $^{\circ}\text{C}$ (color shade) and sea level pressure, in hPa (black contour, contour interval 2 hPa; dotted contours for negative values), on PC1_{BC} . (g) (h) (i) Regression of the SIC, in % (gray contours and color shade), and sea level pressure, in hPa (black contour, contour interval 2 hPa; dotted contours for negative values), on PC1_{SIC} . In all panels, color shade indicates p-value below 5% for the regression of the SST or SIC. Dots ~~indicate~~ ~~indicates~~ p-values below 5% for the SLP regression in panels (d)-(i). Left panels are for November. Center panels are for January. Right panels are for April.



105 **Figure 10: (First column from left) First EOF of the detrended Eurasian snow cover in ERA5-Land, shown as the regression of the**
 the snow cover, in %, onto the first PC. (Second column) Regression of the observed (color shade and grey contours) surface air
 110 temperature, in °C, and (black contour; contour interval 1 hPa) sea level pressure in hPa on PC1 at lag=0, and (third column) at lag
 = 1, i.e. for PC1 leading by one month the atmospheric fields. (Fourth column) Pattern correlation of the EOF1s and of the regressions
 of surface air temperature or sea level pressure on the Eurasian snow cover PC1 between ERA-Land and other snow cover datasets.
 T2M₀ and SLP₀ designate the pattern correlation of the regressions at lag = 0; T2M₁ and SLP₁ designate the pattern correlation at
 lag = 1. **The first** row is for November, **the** second row for January and **the** third row for April. In **the** second and third columns,
 color shadings indicate p-value below 5% for the **surface air temperature** regression.



115

120

Figure 11: (First row) First and (second row) second EOF_{int} of the Eurasian snow cover, in %, associated with internal atmospheric variability in the simulations ALL and NoSICvar~~NoSIC~~. The patterns are the average patterns of the eight models. The numbers displayed on top are the minimum and maximum variance explained by that EOF among the eight models. Dots indicate the locations where 7 models out of 8 have EOF_{int} anomalies with the same sign. (Last row) Spatial pattern correlation between the EOF_{int} obtained from each model and that of the model average. The blue (orange) bar shows the results when using the first (second) EOF. The numbers on the x-axis ~~designates~~ designates each model (see Table 1 for the corresponding model names). The symbol star indicates when 30 members are available for both ALL and NoSICvar~~NoSIC~~; for (left column) November, (center column) January and (right column) April.

125






Figure 12: Regression of the (color shade) 2 m air temperature (T2m), in °C, and (black contour, contour interval 0.5 hPa) SLP, in hPa, on (a) (b) (c) PC1_{Int} and (c) (d) (e) PC2_{Int}, for (left) November, (center) January and (right) April. All regressions are calculated at no lag (lag = 0 mth), and only use the T2m and SLP anomalies associated with the internal atmosphere-land variability, after the removal of the corresponding multi-model mean. The regression shows the multi-model mean regression map, with dots indicating when the sign of SLP anomalies is consistent in at least 7 models out of eight. The color shades are masked if the sign of T2m is not consistent in at least 7 models out of eight.

130



135


Figure 13: Same as Fig. 12a-c+2, but at (a) (b) (c) lag 1 and (d) (e) (f) lag 2, when the atmosphere follows by one month the (left) November, (center) January and (right) April snow cover index, as provided by PC1_{int}. **The last row shows the spatial pattern correlation north of 20°N between the SLP regression obtained from each model and that of the model average. The blue (green) bar shows the results at lag 1 (2). The numbers on the x-axis designate each model (see Table 1 for the corresponding model names). The symbol star indicates when 30 members are available for both ALL and NoSICvar; for (g) November, (h) January and (i) April.**



140

145


Figure 14: Same as Fig. 12a-c12, but using $PC2_{Int}$ instead of $PC1_{Int}$ at lag = 1 month, when the atmosphere follows by one month the (a) (d) November, (b) (e) January and (c) (f) April snow cover. The last row shows the spatial pattern correlation north of $20^{\circ}N$ between the SLP regression obtained from each model and that of the model average. The blue bar shows the results at lag 1. The numbers on the x-axis designate each model (see Table 1 for the corresponding model names). The symbol star indicates when 30 members are available for both ALL and NoSICvar; for (g) November, (h) January and (i) April.



150

155

Figure 15: (a) Regression of the daily air temperature anomalies over the polar cap (north of $60^{\circ}N$), in $^{\circ}C$, onto the First PC of the snow cover internal atmospheric variability, $PC1_{Int}$, in January and in LMDZOR6. The black line indicates the local statistical significance at the 5% level. The vertical dashed black line shows the days corresponding to January. (b) Explained variance fraction of the Arctic Oscillation defined as the first EOF of the sea level pressure north of $20^{\circ}N$, for a regression using $PC1_{Int}$, PCT_{50} and both PCT_{Int} and PCT_{50} as predictors. The explained variance fraction is significant at the 5% level in the three cases.



165 Figure 16: Regression of the (a,d) snow cover, in %, and sea level pressure (SLP; b,e) anomalies, in hPa, onto the January polar cap (north of 60°N) 50-hPa temperature anomalies in LMDZOR6. (c,f) Regression of the sea level pressure (SLP) anomalies, in hPa, onto the First PC of the snow cover internal atmospheric variability, $PC1_{Int}$, in January and LMDZOR6. The lag indicated is negative and given in month, indicating that the polar cap 50-hPa geopotential height or the $PC1_{Int}$ lags. The color shades are masked if the local statistical significance is above 5%.