	1	Responses <u>Reallocation</u> of elemental content and macromolecule of <u>in</u> the
	2	coccolithophore Emiliania huxleyi to reduced phosphorus availability and ocean
	3	acidification depend on light intensity acclimate to climate change
ļ	4	
	5	
	6	Yong Zhang ^{1,*} , Yong Zhang ^{1,§} , Shuai Ma ¹ , Hanbing Chen ² , Jiabing Li ¹ , Zhengke
	7	Li ³ , Kui Xu ⁴ , Ruiping Huang ⁵ , Hong Zhang ¹ , Yonghe Han ¹ , Jun Sun ⁶
	8	
	9	
	10	¹ College of Environmental and Resource Sciences, College of Carbon Neutral
	11	Modern Industry, Fujian Key Laboratory of Pollution Control and Resource
	12	Recycling, Fujian Normal University, Fuzhou, China
	13	² College of Life Science, Fujian Normal University, Fuzhou, China
	14	³ School of Food and Biological Engineering, Shanxi University of Science and
	15	Technology, Xi'an, China
	16	⁴ Hubei Key Laboratory of Edible Wild Plants Conservation and Utilization, Hubei
	17	Engineering Research Center of Special Wild Vegetables Breeding and
	18	Comprehensive Utilization Technology, College of Life Sciences, Hubei Normal
	19	University, Huangshi, China
	20	⁵ State Key Laboratory of Marine Environmental Science, College of Ocean and Earth
	21	Sciences, Xiamen University, Xiamen, China
	22	⁶ Institute for Advanced Marine Research, China University of Geosciences,
	23	Guangzhou, China
	24	

26	Running head: Physiology and biochemistry of E. huxleyi
27	
28	*Correspondence: Yong Zhang (yongzhang@fjnu.edu.cn)
29	
30	Keywords: Carbohydrate; CO ₂ ; coccolithophore; elemental content; light intensity;
31	phosphorus availability; protein.
32	
33	[§] Email: qsx20211022@student.fjnu.edu.cn
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	

51 Abstract

52 Global climate change leads to simultaneous changes in multiple environmental 53 drivers in the marine realm. Although physiological characterization of 54 coccolithophores have been studied under climate change, there is limited knowledge on the biochemical responses of this biogeochemically important phytoplankton 55 56 group to changing multiple environmental drivers. Here we investigate the interactive effects of reduced phosphorus availability (4 to 0.4 μ mol L⁻¹), elevated pCO₂ 57 concentrations (426 to 946 µatm) and increasing light intensity (40 to 300 µmol 58 photons $m^{-2} s^{-1}$) on elemental content and macromolecules of the cosmopolitan 59 60 coccolithophore Emiliania huxleyi. Reduced phosphorus availability reduces 61 particulate organic nitrogen and protein contents per cell under $40 \mu mol photons m^{-2}$ s^{-1} low light intensity, but not under 300 µmol photons m⁻² s⁻¹high light intensity. 62 63 Reduced phosphorus availability and <u>ocean acidification</u> elevated pCO_2 64 concentrations act synergistically to increase particulate organic carbon (POC) and 65 carbohydrate contents per cell under high light intensity 300 μ mol photons m⁻² s⁻¹ but not under low light intensity 40 μ mol photons m⁻² s⁻¹. Reduced phosphorus 66 67 availability, ocean acidification elevated pCO_2 concentrations and increasing light 68 intensity act synergistically to increase the allocation of POC to carbohydrates. Under 69 future ocean acidificationelevated pCO_2 concentrations and increasing light intensity, 70 enhanced carbon fixation could increase carbon storage in the phosphorus-limited 71 regions of the oceans where *E. huxleyi* dominates the phytoplankton assemblages. In 72 each light intensity, elemental carbon to phosphorus (C : P) and nitrogen to 73 phosphorus (N : P) ratios decrease with increasing growth rate. These results suggest 74 that coccolithophores could reallocate chemical elements and energy to synthesize 75 macromolecules efficiently, which allows them to regulate its elemental content and 76 growth rate to acclimate to changing environmental conditions.

77

78 **1 Introduction**

79 Continuous increase in atmospheric CO_2 level, as a consequence of anthropogenic 80 activities, leads to global and ocean warming, which in turn shoals the ocean upper 81 mixed layer (UML), hinders upward transport of nutrients from deeper ocean to the UML and increases light exposures to phytoplankton cells dwelling therein 82 83 (Steinacher et al., 2010; Wang et al., 2015). The dissolution of CO_2 in the oceans is 84 causing a significant chemical shift toward higher CO_2 and proton ([H⁺]) 85 concentrations, a process defined as ocean acidification (OA) (Caldeira and Wickett, 86 2003). These ocean changes expose phytoplankton cells within the UML to multiple 87 driversEnvironmental changes in the UML will expose phytoplankton cells to 88 physiological stress, and understanding the effects of changing multiple 89 environmental drivers on the physiology and biochemistry of marine phytoplankton is 90 important for projections of changes in the biogeochemical roles of phytoplankton in 91 the future ocean (Gao et al., 2019).

92 Coccolithophores take up carbon dioxide (CO₂) to produce particulate organic carbon (POC) via photosynthesis, and use bicarbonate (HCO_3^{-}) and calcium (Ca^{2+}) to 93 94 synthesize calcium carbonate plates (coccoliths, PIC) and release CO₂ via 95 calcification, and play a critical role in the marine carbon cycle (Rost and Riebesell, 96 2004). The cosmopolitan coccolithophore *Emiliania huxleyi* typically forms extensive 97 blooms that are easily detected by satellite remote sensing due to high light scattering 98 caused by coccoliths (Terrats et al., 2020; He et al., 2022). Within E. huxleyi blooms 99 in polar and subpolar oceans, dissolved nitrate and phosphate concentrations in surface seawater could be as lower as 0.95 μ mol L⁻¹ and 0.16 μ mol L⁻¹, respectively 100

(Townsend et al., 1994), light intensity are higher than 300 μ mol photons m⁻² s⁻¹ 101 (Tyrrell and Merico, 2004), and the mean concentrations of seawater CO₂ increased 102 103 by 21.0%–43.3% which weakens the oceanic CO₂ uptake from the atmosphere 104 (Kondrik et al., 2018). Emiliania huxleyi is also the dominant phytoplankton species 105 in the lower photic zone in the north-eastern Caribbean Sea (western Atlantic Ocean) 106 (Jordan and Winter, 2000) and in the South Pacific Gyre where dissolved nitrate and 107 phosphate concentration are about 1.0 μ mol L⁻¹ and 0.2 μ mol L⁻¹, respectively, and light intensity is lower than 20 μ mol photons m⁻² s⁻¹ (Beaufort et al., 2008; Perrin et 108 109 al., 2016). In the future ocean, numerous environmental factors will simultaneously 110 change and the extent of these changes may increase (Gao et al., 2019). To explore 111 how E. huxleyi acclimate to simultaneous changes in macronutrient concentration, 112 light intensity and CO₂ level, it is interesting to investigate their physiological and 113 biochemical processes, which can also help to project the effect of coccolithophores 114 on ocean carbon cycle and ecological systems.

115 For more than a decade, research has shown that *E. huxleyi* cells developed several 116 strategies to acclimate to reduced phosphorus availability, increasing light intensity 117 and ocean acidification (Leonardos and Geider, 2005; McKew et al., 2015; Wang et 118 al., 2022). Interactive effects of phosphorus availability and light intensity have 119 shown that under phosphorus limitation condition, cells increased expression and the 120 activity of alkaline phosphatase, and took up and used phosphorus efficiently under 121 high light intensity, whereas they lowered the phosphorus uptake rate under low light 122 intensity (Riegman et al., 2000; Perrin et al., 2016). In addition, the positive effect of 123 reduced phosphorus availability on cellular POC and PIC contents of E. huxleyi was 124 further enhanced by increasing light intensity due to high-light-induced increases in 125 CO₂ and HCO₃⁻ uptake rates under low phosphate availability (Leonardos and Geider,

126 2005). The negative effect of reduced phosphorus availability on cellular particulate 127 organic phosphorus (POP) content was partly compensated by increased PO_4^{3-} uptake 128 rate under increasing light intensity (Perrin et al., 2016). On the other hand, several 129 studies report that ocean acidification and reduced phosphorus availability acted 130 synergistically to increase the <u>cellular</u> POC content, especially at high light intensity, 131 and acted antagonistically to affect cellular PIC content of E. huxleyi (Leonardos and 132 Geider, 2005; Matthiessen et al., 2012; Zhang et al., 2020). In addition, ocean 133 acidification normally amplified the positive effect of increasing light intensity on 134 cellular POC content (Rokitta and Rost, 2012; Heidenreich et al., 2019). Due to high 135 proton concentration-induced reduction in HCO_3^- uptake rate, ocean acidification 136 could weaken or counteract the positive effect of increasing light intensity on cellular 137 PIC content (Rokitta and Rost, 2012; Kottmeier et al., 2016). Overall, while recent 138 studies have focused on physiological performance of E. huxleyi and their effects on 139 marine biogeochemical cycling of carbon, little information is available about the 140 biochemical response of E. huxlevi to reduced phosphorus availability, increasing 141 light intensity and ocean acidification.

142 The objective of this study is to investigate the combined effects of reduced 143 phosphorus availability, increasing light intensity and ocean acidification on cellular 144 elemental contents, the carbon (C) : nitrogen (N) : phosphorus (P) ratio and 145 macromolecules of E. huxleyi, and to analyze the effects of macromolecules on 146 elemental contents. Under reduced phosphorus availability, increasing light intensity 147 and ocean acidification, we hypothesize that increased <u>cellular</u> POC content is more 148 likely to be caused by increased carbohydrate content. In addition, we discuss the 149 potential mechanisms for changing <u>cellular</u> PIC content in response to changed levels 150 of phosphate, light, and CO₂, which is important for projections of changes in 151 coccolithophore biogeochemistry and ecology in the future ocean.

152

153 2 Materials and Methods

154 2.1 Experimental setup

Emiliania huxleyi strain RCC1266 (morphotype A) was originally isolated from shelf 155 156 waters around Ireland (49°30' N, 10°30' W) in 2007 and obtained from the Roscoff 157 algal culture collection (Fig. S1). Emiliania huxleyi was cultured under a 14 h : 10 h light : dark cycle (light period: 06:00 to 20:00 h) in a thermo-controlled incubator 158 159 (MGC-400H, Shanghai Yiheng Scientific Instrument) at 20°C in semicontinuous 160 cultures. The artifical seawater (ASW) media was prepared according to Berges et al. (2001) with a salinity of 33 psu, a boron concentration of 372 μ mol L⁻¹, and with the 161 addition of 2350 μ mol L⁻¹ bicarbonate to achieve the total alkalinity (TA) of 2350 162 μ mol L⁻¹, and enriched with 64 μ mol L⁻¹ NO₃⁻, f/8 concentrations for trace metals 163 164 and vitamins (Guillard and Ryther, 1962). The experiment was conducted in two parts (Fig. S24). The first part (Part 1) was performed at 40 μ mol photons m⁻² s⁻¹ (low light 165 intensity, LL) and the second one (Part 2) was at 300 μ mol photons m⁻² s⁻¹ (high light 166 167 intensity, HL). The LL intensity used here corresponds to the lower end of the 168 irradiance range of the UML, and the HL intensity represents to the irradiance in the 169 surface ocean (Jin et al., 2016; Perrin et al., 2016). For each part of the experiment, 170 dissolved inorganic phosphorus (DIP) concentration and ocean acidification were combined in a fully factorial design: high DIP concentration (4 μ mol L⁻¹)+low CO₂ 171 172 level (426 µatm, current CO₂ level) (HP+LC, treatment 1 in LL and treatment 5 in 173 HL), high DIP concentration (4 μ mol L⁻¹)+high CO₂ level (946 μ atm, future CO₂ 174 level) (HP+HC, treatment 2 in LL and treatment 6 in HL), low DIP concentration $(0.43 \mu mol L^{-1})$ + low CO₂ level (426 μatm) (LP+LC, treatment 3 in LL and treatment 175

7 in HL), and low DIP concentration (0.43 μ mol L⁻¹)+high CO₂ level (946 μ atm) 176 (LP+HC, treatment 4 in LL and treatment 8 in HL). High DIP concentration is replete 177 178 for physiological process of *E. huxleyi*, and low DIP concentration corresponds to the 179 upper end of the range of phosphate concentration in the coastal waters (Larsen et al., 180 2004) at the end of the incubation, low DIP concentration limits the growth of E. *huxleyi* (see below). There were eight treatments totally and four biological replicates 181 182 for each treatment (Fig. S24). In all cases, cell densities were lower than 78,000 cells mL^{-1} and the cells were acclimated to each treatment for at least 8 generations before 183 184 physiological and biochemical parameters were measured.

185 At LL intensity (Part 1), for the treatments of HP+LC and HP+HC, the ASW media were enriched with 4 μ mol L⁻¹ PO₄³⁻ and aerated for 24 h at 20°C with filter-sterilized 186 187 (PTFE filter, 0.22 µm pore size, Nantong) air pumped from the room. The pH_{Total} 188 (total scale) values of the media under both HP+LC and HP+HC treatments were 189 about 8.04. The dry air was humidified with Milli-Q water prior to the aeration to 190 minimize evaporation. Under the HP+HC treatment, the pH_{Total} values of the media 191 were adjusted to 7.74 by stepwise additions of CO₂-saturated seawater, and the ratio 192 was about 6.5 mL CO2-saturated seawater : 1000 mL ASW media. The CO2-saturated 193 seawater was achieved by bubbling pure CO₂ gas into 500 ml ASW media with a total 194 alkalinity of 2350 μ mol L⁻¹ for 2 h. For the treatments of LP+LC and LP+HC, the ASW media were enriched with 0.4 μ mol L⁻¹ PO₄³⁻ and aerated for 24 h at 20°C with 195 196 filtered room air. Under the LP+HC treatment, the pH_{Total} values of the media were also adjusted to 7.74 as described above. The HP+LC, HP+HC, LP+LC and LP+HC 197 198 seawater were then filtered (0.22 µm pore size, Polycap 75 AS, Whatman) and 199 carefully pumped into autoclaved 50 mL (for TA measurements), 600 mL (for 200 pre-experimental cultures) and 2350 mL (for experimental cultures) polycarbonate

201 bottles (Nalgene) with no headspace to minimize gas exchange. The volumes of 202 culture inoculum were calculated to match the volumes of media taken out from the 203 bottles prior to inoculation. The cells were inoculated to achieve an initial density of 5000 cells ml⁻¹ in the HP+LC and HP+HC conditions, respectively, and cultured for 2 204 205 days, then diluted to the initial density again. These processes were performed three times in 600 mL bottles for pre-experimental cultures at 40 μ mol photons m⁻² s⁻¹ (LL) 206 207 of photosynthetically active radiation (PAR; measured using a LI-190SA quantum 208 sensor, Beijing Ligaotai Technology Co. Ltd.). In the main experimental cultures-in the HP+LC and HP+HC conditions at LL intensity, the cells were, respectively, 209 210 transferred from 600 mL to 2350 mL bottles at the same time, and cultured for 211 another 2 days (Fig. S24b). Culture bottles were rotated 10 times until cells were 212 mixed at 09:00 h, 13:00 h and 19:00 h. Based on changes in cell densities during the 213 incubations, we calculated that at LL intensity, cells were acclimated to HP+LC and 214 HP+HC conditions for 10 generations. In the second day of the main experimental 215 cultures, subsamples were taken for measurements of cell densities, pH_{Total}, TA, 216 cellular contents of total particulate carbon (TPC), particulate organic carbon (POC), 217 nitrogen (PON) and phosphorus (POP), carbohydrate and protein. At the end of the 218 cultures under the previous conditions, cell samples with an initial density of 5000 cells ml⁻¹ were transferred from HP+LC condition (treatment 1) to LP+LC condition 219 220 (treatment 3), and from HP+HC condition (treatment 2) to LP+HC condition 221 (treatment 4) at LL intensity. The cells were acclimated to LP+LC and LP+HC 222 conditions for 8 generations before subsamples were taken for measurements, which 223 allows cells to have enough time periods to change growth rate against the low DIP 224 concentration.



were transferred from 40 (LL) to 300 μ mol photons m⁻² s⁻¹ (HL) of PAR with initial 226 227 cell density of 5000 cells ml⁻¹. The cells were cultured under the HP+LC and HP+HC 228 conditions for 2 days, respectively, and then diluted back to the initial cell density. 229 These processes were performed three times in 600 mL bottles at HL intensity, and 230 then the main experimental cultures were conducted in 2350 mL bottles. The cells 231 were, respectively, acclimated to HP+LC and HP+HC conditions for at least 8 232 generations at HL intensity. On the second day of the incubation, subsamples were 233 taken for measurements of the parameters. After that, cell samples with an initial density of 5000 cells ml⁻¹ were transferred from HP+LC condition (treatment 5) to 234 235 LP+LC condition (treatment 7), and from HP+HC condition (treatment 6) to LP+HC 236 condition (treatment 8). At HL intensity, cell samples were acclimated for at least 8 237 generations in LP+LC and LP+HC conditions, respectively, before subsamples were 238 taken for measurements.

239

240 2.2 Phosphate concentration and carbonate chemistry measurements

241 In the beginning and on the second day of the incubations, samples for determinations 242 of phosphate concentration (20 mL), pH_{Total} value (20 mL) and total alkalinity (TA) 243 (50 mL) were, respectively, filtered (PTFE filter, 0.22 µm pore size, Nantong) 7 h 244 after the onset of the light period (at 13:00). Dissolved inorganic phosphorus (DIP) 245 concentration was measured using a spectrophotometer (SP-722, Shanghai Spectrum 246 Instruments) following the phosphomolybdate method (Hansen and Koroleff, 1999). 247 The bottle for pH measurement was filled from bottom to top with overflow and 248 closed without a headspace. The pH_{Total} value was measured immediately at 20°C 249 using a pH meter which was corrected with a standard buffer of defined pH in 250 seawater (Dickson, 1993). TA samples were treated with 10 µL saturated HgCl₂ solution and stored in the dark at 4.0°C, and TA was measured at 20°C by potentiometric titration (AS-ALK1+, Apollo SciTech) according to Dickson et al. (2007). Carbonate chemistry parameters were estimated from TA and pH_{Total} using the CO2SYS program of Pierrot et al. (2006) with carbonic acid constants, K_1 and K_2 , taken from Roy et al. (1993).

256

257 2.3 Cell density and elemental content measurements

Twenty milliliter samples to monitor the cell density were taken daily at 13:30 h, and fresh media with the same DIP concentration and carbonate chemistry as in the initial treatment conditions were added as top-up. Cell densities were determined using a MultisizerTM3 Coulter Counter (Beckman Coulter). Growth rates were calculated for each replicate according to the equation: $\mu = (\ln N_t - \ln N_0) / d$, where N_t and N_0 refer to the cell densities on the second day and in the beginning of the main experiment, respectively, and *d* was the growth period in days.

265 After mixing, samples for determinations of TPC (300 mL), POC and PON (300 266 mL), and POP (300 mL) were obtained by filtering onto GF/F filters (precombusted at 267 450°C for 6 h) at the same time (14:00 h) in each treatment. For POC and PON 268 measurements, samples were fumed with HCl for 12 h to remove inorganic carbon. 269 TPC, POC and PON samples were dried at 60°C for 12 h and analyzed using an 270 Elementar CHNS analyzer (Vario EL cube, GmbH, Germany). Cellular particulate 271 inorganic carbon (PIC) content was calculated as the difference between cellular TPC and POC contents (Fabry and Balch, 2010). To remove dissolved inorganic 272 phosphorus from the GF/F filters, POP samples were rinsed three times with 0.17 mol 273 L^{-1} Na₂SO₄. After that, 2 mL 0.017 mol L^{-1} MgSO₄ solution was added onto filters, 274 and POP samples were dried at 90°C for 12 h, and then combusted at 500°C for 6 h to 275

276 remove POC and digested by 0.2 mol L^{-1} HCl (Solórzano and Sharp, 1980). 277 Phosphorus concentration was measured using a microplate reader (Thermo Fisher) 278 following the ammonium molybdate method (Chen et al., 1956) using 279 adenosine-5-triphosphate disodium trihydrate as a standard.

280

281 2.4 Protein and carbohydrate measurements

282 Samples for determinations of protein (600 mL) and carbohydrate (600 mL) were, 283 respectively, filtered onto polycarbonate filters (0.6 µm pore size, Nuclepore, 284 Whatman) and onto precombusted GF/F filters at 14:30 h. Protein samples were 285 extracted by bead milling (FastPrep Lysing Matrix D) in 0.5 mL 1× protein extraction 286 buffer (lithium dodecyl sulfate, ethylene diamine tetraacetic acid, Tris, glycerol and 287 4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride). Bead milling was performed four times for 1 min at 6.5 ms⁻¹, and samples were placed on ice for 2 min 288 289 between each round of bead milling to prevent degradation. The samples were then 290 centrifuged at 10,000 \times g for 5 min (Centrifuge 5418 R, Eppendorf, Germany), and 291 extracted protein in the supernatant was quantified using the BCA assay with bovine 292 gamma globulin as a standard using a microplate reader (Ni et al., 2016). Carbohydrate samples were hydrolyzed with 12.00 mol L^{-1} H₂SO₄ in the dark for 1 h 293 and diluted by Milli-O water to a final H_2SO_4 concentration of 1.20 mol L⁻¹. Then, 294 295 samples were sonicated for 5 min, vortexed for 30 s, and boiled at 90°C for 3 h 296 (Pakulski and Benner, 1992). The extracted carbohydrate was determined by phenol 297 sulfuric reaction with D-glucose as standard (Masuko et al., 2005).

298

299 **2.5 Data analysis**

BOD The percentages of carbon (C : carbohydrate is 40% and C : protein is 53%) and

301 nitrogen (N: protein is 16%) contributed by carbohydrate and protein were calculated 302 from the elemental composition of biochemical classes compiled by Geider and 303 LaRoche (2002). A three-way ANOVA was used to determine the main effects of DIP 304 concentration, light intensity and CO₂ level, and their interactions on each variable. A 305 Tukey post hoc test was performed to identify significant differences between two 306 DIP concentrations, two light intensities and two CO₂ levels. A Shapiro-Wilk test was conducted to analyze the normality of residuals and a Levene test was conducted 307 308 graphically to test for homogeneity of variances. The significant difference between 309 treatments was set at p < 0.05. All data analyses were conducted using the statistical 310 software *R* with the packages carData, lattice and nlme (R version 3.5.0).

311

312 3 Results

313 3.1 Dissolved inorganic phosphorus concentration and carbonate chemistry314 parameters

315 During the incubations, organismal activity significantly reduces dissolved inorganic 316 phosphorus (DIP) concentrations (Table 1). Under high phosphorus (HP) treatment, 317 DIP concentrations decrease by 20.32% in low light (LL) and low CO₂ (LC), by 318 22.32% in LL and high CO₂ (HC), by 27.66% in high light (HL) and LC, and by 31.58% in HL and HC. Under low phosphorus (LP) treatment, DIP concentrations 319 decrease from 0.43 μ mol L⁻¹ at the beginning of the experiment to be lower than 0.04 320 321 μ mol L⁻¹ (the detection limit) at the end of the incubation in LL and LC conditions, in 322 LL and HC conditions, in HL and LC conditions, and in HL and HC conditions. 323 During the incubations, at LL intensity, pH_T values increase by, on average, 0.02 in

HP+LC, by 0.03 in HP+HC, by 0.09 in LP+LC, and by 0.10 in LP+HC conditions

325 (Table 1). At HL intensity, pH_T values increase by 0.05 in HP+LC, by 0.06 in HP+HC,

by 0.12 in LP+LC, and by 0.09 in LP+HC conditions. Correspondingly, at LL
intensity, seawater CO₂ concentrations decrease by 5.53% in HP+LC, by 6.89% in
HP+HC, by 22.76% in LP+LC, and by 22.77% in LP+HC. At HL intensity, seawater
CO₂ concentrations decrease by 16.18% in HP+LC, by 16.41% in HP+HC, by
28.92% in LP+LC, and by 22.30% in LP+HC. Overall, organismal activity has larger
effects on carbonate chemistry under the LP treatment than under the HP treatment.

332

333 3.2 Growth rate

334 The effect of increasing light intensity on growth rate is positive which can be seen by 335 comparing growth rate in the HL regimes with their paired LL regimes (Fig. 1a, b; 336 Table 2), though the extent of increase in growth rate depends on CO₂ levels and 337 phosphate availability. Compared to LL intensity, growth rates at HL intensity 338 increased by 48.48% in HP+LC, by 50.87% in HP+HC, by 60.86% in LP+LC, and by 339 60.80% in LP+HC (Tukey post hoc test, all values of p < 0.01). The effect of 340 increasing CO₂ levels on growth rate depends on light intensity and phosphate 341 availability (Fig. 1a, b). Compared to LC level, growth rates in HC level decreased by 342 3.08% in LL and HP condition (p = 0.48), by 16.13% in LL and LP condition (p < 0.48) 343 0.01), by 1.50% in HL and HP condition (p = 0.68), and by 16.27% in HL and LP 344 condition (p < 0.01). The effect of reduced phosphorus availability on growth rate is 345 negative and the extent of reduction in growth rate depends on light intensity and CO₂ 346 levels (Fig. 1a,b). Compared to HP availability, growth rates in LP availability 347 decreased by 8.46% in LL and LC condition (p < 0.01), by 20.81% in LL and HC 348 condition (p < 0.01), by 0.76% in HL and LC condition (p = 0.99), and by 15.63% in 349 HL and HC condition (p < 0.01), and did not change significantly in HL and LC 350 condition (p = 0.99). These results show that high CO₂ levels and low phosphorus

availability acted synergistically to reduce growth rate of *E. huxleyi*, and increasing

light intensity could partly counteract this response (Table S1).

353

354 **3.3 <u>Cellular</u> POC, PON, POP and PIC contents**

355 The effect of increasing light intensity on <u>cellular</u> POC content is positive, which was 356 observed by comparing POC content in all the HL regimes with their paired LL regimes (Fig. 1bc,d). The extent of increase in POC content depends on CO₂ levels 357 358 and phosphate availability. Compared to LL intensity, cellular POC contents at HL 359 intensity increased by 27.15% in HP+LC, by 26.51% in HP+HC, by 43.24% in 360 LP+LC, and by 58.13% in LP+HC conditions (Tukey post hoc test, all values of p < p361 0.01). The effect of increasing CO₂ levels on POC content is light and phosphate 362 dependent and can be seen by comparing POC content in the HC regimes with their paired LC regimes (Fig. 1bc,d). At LL intensity, cellular POC contents are not 363 364 significantly different between HP+LC, HP+HC, LP+LC and LP+HC conditions (all 365 values of p > 0.1). At HL intensity, compared to LC level, cellular POC contents in HC level increased by 5.12% in HP condition (p = 0.74), and by 8.28% in LP 366 367 condition (p = 0.07). The effect of phosphate reduction on POC content is light and 368 CO₂ dependent, which can be seen by comparing POC content in the LP regimes with 369 those in their paired HP regimes (Fig. 1bc,d). At LL intensity, cellular POC contents 370 did not significantly differ between LP and HP availability. At HL intensity, 371 compared to HP availability, cellular POC contents in LP availability increased by 372 11.80% in LC condition (p = 0.02), and by 15.28% in HC condition (p < 0.01). These 373 results show that ocean acidification and reduced phosphorus availability acted 374 synergistically to increase POC contents in HL condition but not in LL condition 375 (Table S1).

376 The effect of increasing light intensity on <u>cellular</u> PON content depends on CO₂ 377 levels and phosphate availability (Fig. 1ce,f). Compared to LL intensity, cellular PON 378 contents at HL intensity increased by 12.03% in HP+LC condition (p = 0.27), by 379 19.54% in HP+HC condition (p < 0.01), by 22.68% in LP+LC condition (p < 0.01), 380 and by 30.90% in LP+HC condition (p < 0.01). The effect of increasing CO₂ levels on 381 PON content is light and phosphate dependent, which can be seen by comparing POC 382 content in the HC regimes with their paired LC regimes (Fig. 1ce,f). Compared to LC 383 level, cellular PON contents in HC level did not change significantly in LL and HP 384 condition, in LL and LP condition, in HL and LP condition, and increased by 14.68% 385 in HL and HP condition (p = 0.02). The effect of phosphate reduction on PON content 386 is CO₂ and light dependent, which can be seen by comparing PON content in the LP 387 regimes with those in their paired HP regimes (Fig. 1ce,f). Compared to HP 388 availability, cellular PON contents in LP availability decreased by 16.59% in LL and 389 LC condition (p = 0.01), by 24.03% in LL and HC condition (p < 0.01), by 8.35% in 390 HL and LC condition (p = 0.43), and by 17.32% in HL and HC condition (p < 0.01). 391 These results show that increasing light intensity compensated for the negative effect 392 of phosphate reduction on PON content (Table S1).

393 The effect of increasing light intensity on POP content is positive and can be seen 394 by comparing POP content in the HL regimes with their paired LL regimes, though 395 the extent of increase in POP content depends on CO₂ levels and phosphate 396 availability (Fig. 1dg,h). Compared to LL intensity, cellular POP contents at HL 397 intensity increased by 35.79% in HP+LC, by 41.70% in HP+HC, by 57.22% in 398 LP+LC, and by 56.44% in LP+HC conditions (Tukey post hoc test, all values of p < p399 0.01). Ocean acidification did not change the POP contents significantly in LL and 400 HP condition, in LL and LP condition, in HL and HP condition, and in HL and LP 401 condition under all treatments used here (all values of p > 0.53) (Fig. 1dg,h). Reduced 402 phosphorus availability significantly decreased the POP contents, which can be seen 403 by comparing POP content in the LP regimes with their paired HP regimes, though 404 the extent of reduction in POP content depends on light intensity and CO₂ levels (Fig. 405 1dg,h). Compared to HP availability, <u>cellular</u> POP contents in LP availability 406 decreased by 52.96% in LL and LC condition, by 54.03% in LL and HC condition, by 407 46.11% in HL and LC condition, and by 49.51% in HL and HC condition (all values 408 of p < 0.01). These results show that reduced phosphorus availability had a larger 409 effect on POP content than that of ocean acidification and increasing light intensity 410 (Table S1).

411 The effect of increasing light intensity on PIC content is positive, which can be 412 seen by comparing PIC content in the HL regimes with their paired LL regimes, 413 though the extent of increase in PIC content depends on CO₂ levels and phosphorus 414 availability (Fig. 1ei,j). Compared to LL intensity, cellular PIC contents at HL 415 intensity increased by 77.87% in HP+LC, by 70.28% in HP+HC, by 98.31% in 416 LP+LC, and by 90.68% in LP+HC conditions (Tukey post hoc test, all values of p <417 0.01). The effect of increasing CO_2 levels on PIC content is negative and can be seen 418 by comparing PIC content in the HC regimes with those in their paired LC regimes 419 (Fig. 1ei,j). The extent of reduction in PIC content depends on light intensity and 420 phosphorus availability. Compared to LC level, cellular PIC contents under ocean 421 acidification decreased by 31.43% in LL and HP condition (p = 0.09), by 16.00% in 422 LL and LP condition (p = 0.67), by 35.02% in HL and HP condition (p < 0.01), and 423 by 21.12% in HL and LP condition (p < 0.01). The effect of phosphate reduction on 424 PIC content is positive which can be seen by comparing PIC content in the LP 425 regimes with their paired HP regimes, though the extent of increase in PIC content depends on light intensity and CO₂ levels (Fig. 1 $\underline{ei,j}$). Compared to HP availability, cellular PIC contents in LP availability increased by 16.00% in LL and LC condition (p = 0.83), by 41.26% in LL and HC condition (p = 0.16), by 29.98% in HL and LC condition (p < 0.01), and by 60.44% in HL and HC condition (p < 0.01). These results show that high light intensity and low phosphorus availability acted synergistically to increase PIC content, which counteracts the negative effect of ocean acidification on PIC content (Table S1).

433

434 **3.4** Carbohydrate and protein contents

435 The effect of increasing light intensity on carbohydrate content is positive and can be 436 seen by comparing carbohydrate content in the HL regimes with their paired LL 437 regimes, though the extent of increase in carbohydrate depends on CO₂ levels and 438 phosphorus availability (Fig. 2a, b). Compared to LL intensity, cellular carbohydrate 439 contents at HL intensity increased by 148.81% in HP+LC condition, by 139.42% in 440 HP+HC condition, by 179.12% in LP+LC condition, and by 204.42% in LP+HC 441 condition (all values of p < 0.01). The effect of increasing CO₂ levels on carbohydrate 442 content is light and phosphate dependent which can be seen by comparing 443 carbohydrate content in the HC regimes with their paired LC regimes (Fig. 2a,b). 444 Compared to LC level, cellular carbohydrate contents under ocean acidification did 445 not change significantly under LL condition, and increased by 26.55% in LL and HP 446 condition (p = 0.58), by 8.91% in LL and LP condition (p = 0.99), by 21.32% in HL 447 and HP condition (p = 0.02), and by 18.45% in HL and LP condition (p < 0.01). The 448 effect of phosphate reduction on carbohydrate content is light and CO₂ dependent and 449 can be seen by comparing carbohydrate content in the LP regimes with their paired 450 HP regimes (Fig. 2a, b). Compared to HP availability, cellular carbohydrate contents 451 in LP availability did not change significantly in LL and LC condition, in LL and HC 452 condition (both p > 0.65) and increased by 40.13% in HL and LC condition (p < 0.01), 453 and by 36.00% in HL and HC condition (p < 0.01). These results show that increasing 454 light intensity dominantly increased carbohydrate content, and ocean acidification and 455 reduced phosphorus availability acted synergistically to increase carbohydrate 456 contents under high light intensity (Table S1).

457 The effect of increasing light intensity on <u>cellular</u> protein content is positive, which 458 can be seen by comparing protein content in the HL regimes with their paired LL 459 regimes, though the extent of increase in protein content depends on CO₂ level and 460 phosphorus availability (Fig. 2bc,d). Compared to LL intensity, cellular protein 461 contents at HL intensity increased by 24.76% in HP+LC condition, by 30.43% in 462 HP+HC condition, by 68.09% in LP+LC condition, and by 65.39% in LP+HC condition (all values of p < 0.01). The effect of increasing CO₂ levels on protein 463 464 content can be seen by comparing protein content in the HC regimes with their paired 465 LC regimes (Fig. 2be,d). Compared to LC level, cellular protein contents under ocean acidification did not change significantly in LL and HP condition, in LL and LP 466 condition, in HL and HP condition, and in HL and LP condition (all values of p > p467 0.09). The effect of phosphate reduction on protein content is light and CO₂ 468 469 dependent, which can be seen by comparing protein content in the LP regimes with 470 their paired HP regimes (Fig. 2bc,d). Compared to HP availability, cellular protein 471 content in LP availability decreased by 27.88% in LL and LC condition, by 28.80% in 472 LL and HC condition (both p < 0.01) and did not change significantly in HL and LC 473 condition, and in HL and HC condition (both p > 0.11). These results show that high 474 light intensity counteracted the negative effect of low phosphorus availability on 475 protein content, and ocean acidification had less effect on protein content (Table S1).

477 3.5 Percentage of POC allocated to carbohydrate (carbohydrate-C : POC) and 478 protein (protein-C : POC)

479 Increasing light intensity increased the percentage of POC allocated to carbohydrate 480 (carbohydrate–C : POC), which can be seen by comparing carbohydrate–C : POC in 481 the HL regimes with their paired LL regimes, though the extent of increase in 482 carbohydrate–C : POC depends on CO_2 levels and phosphorus availability (Fig. 2ce,f). 483 Compared to LL intensity, carbohydrate-C : POC at HL intensity increased by 484 95.60% in HP+LC condition, by 97.69% in HP+HC condition, by 95.05% in LP+LC 485 condition, and by 83.37% in LP+HC condition (all values of p < 0.01). The effect of 486 increasing CO₂ levels on carbohydrate–C : POC is light and phosphate dependent, and 487 can be seen by comparing carbohydrate-C : POC in the HC regimes with their paired 488 LC regimes (Fig. 2ce,f). Compared to LC level, carbohydrate–C : POC under ocean 489 acidification increased by 20.12% in LL and HP condition, by 11.42% in LL and LP 490 condition, by 20.36% in HL and HP condition, and by 4.40% in HL and LP condition 491 (all values of p > 0.08). The effect of phosphate reduction on carbohydrate–C : POC 492 is light and CO_2 dependent, which can be seen by comparing carbohydrate-C : POC 493 in the LP regimes with those in their paired HP regimes (Fig. 2<u>ce,f</u>). Compared to HP 494 availability, carbohydrate-C : POC in LP availability increased by 25.61% in LL and 495 LC condition (p = 0.16), by 17.37% in LL and HC condition (p = 0.47), by 25.81% in 496 HL and LC condition (p < 0.01), and by 8.11% in HL and HC condition (p < 0.05). 497 These results show that increasing light intensity, ocean acidification and reduced 498 phosphorus availability acted synergistically to increase the percentage of POC 499 allocated to carbohydrate (Table S1).

500 Increasing light intensity did not significantly change the percentage of POC

501 allocated to protein (protein-C : POC) in HP+LC, HP+HC, LP+LC and LP+HC 502 conditions under the phosphorus availability and CO_2 levels used here (Fig. 2dg,h). Ocean acidification did not significantly affect the protein-C : POC in LL and HP, in 503 504 LL and LP, in HL and HP, and in HL and LP conditions. The effect of phosphate 505 reduction on protein–C: POC is light and CO₂ dependent, which can be seen by 506 comparing the protein-C : POC in the LP regimes with their paired HP regimes (Fig. 507 $2d_{g,h}$). Compared to HP availability, protein-C : POC in LP availability decreased by 508 27.39% in LL and LC condition (p < 0.01), by 23.05% in LL and HC condition (p < 0.01) 0.01), by 12.81% in HL and LC condition (p = 0.09), and by 21.77% in HL and HC 509 510 condition (p < 0.01). These results show that reduced phosphorus availability 511 dominantly reduced the protein-C : POC, and increasing light intensity and ocean 512 acidification had less effects on protein-C : POC (Table S1). On the other hand, 513 increasing light intensity, reduced phosphorus availability and ocean acidification did 514 not significantly change the percentage of PON allocated to protein (protein–N : PON) 515 (Fig. 2e).

516

517 **3.6** Elemental stoichiometry and protein content as a function of growth rate

518 Reduced phosphorus availability increased the POC : PON ratio, and the extent of the 519 increase was larger under HL than LL intensity (Fig. S5a). At LL and HL intensities, both POC : POP ratio and PON : POP ratio were linearly and negatively correlated 520 521 with growth rates (Fig. 3a,b). In LL and HL conditions, POC : POP ratio decreased linearly with increasing growth rate ($R^2 = 0.71$, F = 32.08, p < 0.01 in LL condition; 522 $R^2 = 0.53$, F = 14.63, p < 0.01 in HL condition). Similarly, in LL and HL conditions, 523 PON : POP ratio decreased linearly with increasing growth rate ($R^2 = 0.69$, F = 29.23, 524 p < 0.01 in LL condition; $\mathbb{R}^2 = 0.50$, F = 13.31, p < 0.01 in HL condition). In all 525

treatments, protein content increased linearly with increasing growth rate ($R^2 = 0.76$, *F* = 151.14, *p* < 0.01) (Fig. 3c), and POC content increased linearly with increasing carbohydrate content ($R^2 = 0.94$, *F* = 435.10, *p* < 0.01) (Fig. 3d).

529

530 4 Discussion

531 Coccolithophores make an important contribution to marine biological carbon 532 pumpplay a complex role in the marine carbon cycle through production and export of 533 organic carbon to depth but also through the carbonate counter pump, and their 534 responses to global climate change could have significant consequences for marine 535 carbon cycling (Riebesell et al., 2017). The bloom-forming coccolithophore E. huxleyi, 536 dominating the assemblages in seawater under limited phosphorus condition, is likely 537 to be exposed to increasing light intensity and ocean acidification in the future ocean 538 (Kubryakova et al., 2021). In this study, we observed that increasing light intensity 539 compensates for the negative effects of low phosphorus availability on cellular protein 540 and nitrogen contents (Figs. 1 and 2). Reduced phosphorus availability, increasing 541 light intensity and ocean acidification act synergistically to increase cellular contents 542 of carbohydrate and POC, and the allocation of POC to carbohydrate. These 543 regulation mechanismschanges in E. huxleyi could provide vital information for 544 evaluating carbon cycle in marine ecosystems under global change.

Ribonucleic acid (RNA) is the main phosphorus-containing macromolecule within
the cell (Geider and La Roche, 2002). Therefore, the reduced phosphorus availability
dominantly reduces the RNA content (Fig. S<u>6</u>5), which contributes to low POP
contents (McKew et al., 2015) (Fig. 1<u>dg,h</u>). In eukaryotic cells, ribosomal RNA
(rRNA) constitutes about 80% of the total RNA and is mainly used to create ribosome
(Dyhrman, 2016). Thus, reduced RNA contents decrease the numbers of ribosome,

551 which has a potential to decrease protein synthesis (Dyhrman, 2016; Rokitta et al., 552 2016). On the other hand, the separation of the regression line between POC : POP 553 ratio (or PON : POP ratio) and growth rate under low and high light intensities 554 suggests different POP storage contents in E. huxleyi among different light intensities (Perrin et al., 2016). On the other hand, ILow light intensity down-regulates the 555 556 expression of genes related to nitrate reductase and nitrite reductase, and then reduces 557 the nitrate uptake and assimilation efficiency of E. huxleyi and other phytoplankton 558 species (Perrin et al., 2016; Lu et al., 2018), which exacerbates the negative effect of 559 low phosphorus availability on protein synthesis and PON contents (Figs. 1ce and 560 2be). Besides that, low light intensity significantly reduces the rates of RNA synthesis, 561 carbohydrate synthesis and cell division (Zhang et al., 2021), which adds to the 562 negative effect of low phosphorus availability on growth rate of E. huxleyi (Fig. 1a). 563 Under high light intensity and low CO₂ level, reduced phosphorus availability did not 564 change growth rate and protein content (Figs. 1ab and 2bd), which suggests that E. 565 huxleyi might compensate for low phosphate-induced decreases in ribosome content 566 by increasing protein synthesis efficiency under increasing light intensity (Reith and 567 Cattolico, 1985). Under high light intensity and ocean acidification, reduced 568 phosphorus availability did not significantly change cellular protein content while 569 reduced growth rate, which might indicate the lowered protein synthesis efficiency 570 (McKew et al., 2015).

Several studies report that reduced phosphorus availability (0.4–0.5 μmol L⁻¹) did
not change growth rates significantly during the short-time (2 or 3 days) incubations
under low CO₂ level and high light intensity (Rokitta et al., 2016; Zhang et al., 2020;
Wang et al., 2022) (Fig. S6). The reasons could be that *E. huxleyi* cells developed
high affinity for phosphate and increased the uptake rate of phosphate (Wang et al.,

576 2022) and could replace the phospholipid membrane with non-phosphorus membrane 577 during the short-time incubation of phosphorus limitation (Shemi et al., 2016). Our data showed that reduced phosphorus availability and ocean acidification acted 578 579 synergistically to reduce growth rate under both low and high light intensities (Fig. 580 1a,b). One of the reasons could be that low pH value under ocean acidification up 581 regulates the expressions of a series of genes involved in ribosome metabolism, such 582 as genes of large subunit ribosomal protein L3, L38E, L30E (RP-L3, RP-L38E, 583 RP-L30E), and small subunit ribosomal protein S3E, S5E, SAE (PR-S3E, PR-S5E, 584 RP-SAE) in E. huxleyi (Wilson and Doudna Cate, 2012) (Fig. S7). Under ocean 585 acidification, up regulation of expression of these genes has the potential to drive cells 586 to allocate more phosphorus to synthesize ribosome, and to reduce the allocation of 587 phosphorus to DNA replication (Rokitta et al., 2011), which exacerbates the limitation 588 of reduced phosphorus availability on the rate of cell division in E. huxleyi (Rouco et 589 al., 2013). Under phosphorus-replete condition, more phosphorus is reallocated to 590 ribosome metabolism under ocean acidification which could facilitate nitrogen 591 assimilation (Fig. 2bd). Overall, under high light intensity, ocean acidification is 592 likely to facilitate E. huxleyi cells to increase nitrogen content in phosphorus-replete 593 condition and to reduce growth rate in phosphorus-limitation.

In this study, we found that low light intensity dominantly limits carbon assimilation of *E. huxleyi* and reduces the effects of phosphate availability and ocean acidification on <u>cellular</u> carbohydrate and POC contents (Figs. 1<u>be</u> and 2a). However, under high light intensity, *E. huxleyi* had high carbohydrate and POC contents while low growth rate under reduced phosphorus availability and ocean acidification (Figs. 1<u>a.bb,d</u> and 2<u>a</u>b), which suggests that carbon assimilation rate did not change significantly while cell division rate decreased (Matthiessen et al., 2012; Perrin et al., 2016). Furthermore, carbohydrate is a carbon- and energy-storing macromolecule, and
protein is related to growth rate (Geider and La Roche, 2002). Under high light
intensity, reduced phosphorus availability and ocean acidification, *E. huxleyi* cells
could synthesize more carbohydrate to store carbon and energy but didn't increase
protein content, which contributes to the large percentage of POC allocated to
carbohydrate (Fig. 2<u>c</u>f).

607 The physiological reasons for reduced calcification rate under ocean acidification 608 could be due to high proton concentration-induced reduction in HCO_3^- uptake rate 609 (Meyer and Riebesell, 2015; Kottmeier et al., 2016). The molecular mechanisms for 610 low PIC content under ocean acidification may be due to down-regulation of a series 611 of genes potentially involved in ion transport and pH regulation, such as genes of 612 calcium/proton exchanger (CAX3), sodium/proton exchanger (NhaA2) and 613 membrane-associated proton pump (PATP) (Mackinder et al., 2011; Lohbeck et al., 614 2014). On the other hand, increasing light intensity up-regulates a series of genes 615 related to ion transport, such as gene of CAX3, gene of Cl⁻/HCO₃⁻ exchanger and 616 genes of various subunits of a vacuolar H⁺-ATPase (V-ATPase) and so on (Rokitta et 617 al., 2011). Up-regulation of these genes in high light intensity has the potential to facilitate cells to take up HCO_3^- and Ca^{2+} , and to pump proton outside the cells, and 618 619 then leads to large PIC content of E. huxleyi (Kottmeier et al., 2016). Our data suggest 620 that increasing light intensity counteracts the negative effect of ocean acidification on 621 cellular PIC content of *E. huxleyi* (Fig. 1ei,j). These results are consistent with the findings of Feng et al. (2020) who reported that combinations of increasing light 622 623 intensity and ocean acidification increase the expression of genes involved in 624 calcium-binding proteins (CAM and GPA), which has the potential to increase 625 calcium influx into cells and then compensate for the effect of reduced HCO₃⁻ uptake

626 rate on calcification. It is also suggested that increasing light intensity could cause 627 cells to remove H⁺ faster which neutralizes the effect of high proton concentration on 628 calcification (Jin et al., 2017). On the other hand, reduced phosphorus availability 629 extends the G1 phase of cell cycle where calcification occurs, which prolongs the 630 calcification time and then increases cellular PIC content (Müller et al., 2008). In addition, reduced phosphorus availability up-regulates expressions of genes of Ca2+ 631 632 uptake, proton removal and carbonic anhydrase, and then increases coccolith 633 production (Wang et al., 2022), which contribute to a larger PIC content and 634 counteract the negative effect of ocean acidification on PIC contents (Borchard et al., 635 2011) (Fig. 1eij). Furthermore, one of the reasons for larger PIC contents under 636 reduced phosphorus availability and increasing light intensity conditions are likely 637 due to larger and more numerous coccoliths (Gibbs et al., 2013; Perrin et al., 2016). 638 Overall, responses of calcification of E. huxleyi to ocean climate change are complex 639 than previously thought (Meyer and Riebesell, 2015), and it is worth exploring the 640 underlying mechanisms of calcification under changing multiple environmental 641 drivers (Mackinder et al., 2011; Feng et al., 2020).

642 Cellular POP content of E. huxleyi generally decreased, and POC : POP ratio and 643 PON : POP ratio increased with reducing phosphorus availability (Leonardo and 644 Geider, 2005; McKew et al., 2015). The negative correlations between growth rate 645 and POC : POP ratio or PON : POP ratio under each light intensity are consistent with 646 the growth rate hypothesis (Fig. 3), which proposes that growth rate increases with 647 increasing RNA : protein ratio. Phosphorus in RNA accounts for a high percentage of 648 total POP, whereas nitrogen in protein is the main form of PON (Zhang et al., 2021), 649 and the growth rate hypothesis suggests that growth rate could increase with decreasing POC : POP ratio or PON : POP ratio (Sterner and Elser, 2002). Our results 650

651	suggest that E. huxleyi could reallocate chemical elements and energy to synthesize
652	carbohydrate, protein and RNA efficiently, and then regulate its elemental
653	stoichiometry and growth rate to acclimate to reduced phosphorus availability, ocean
654	acidification and increasing light intensity (Moreno and Martiny, 2018).

In the future ocean, large carbohydrate and POC contents, POC : PON ratio, and POC : POP ratio of coccolithophores indicate increases in carbon export to the deep ocean that may affect the efficiency of the biological carbon pump and the marine biogeochemical cycle of carbon (Meyer and Riebesell, 2015). In addition, increased cellular PIC content under phosphorus limitation condition may have the potential to weaken CO₂ uptake of the oceans in phosphorus-limited marine environments. In summary, responses of coccolithophores to climate change is likely to affect the marine carbon cycle in the future (Riebesell et al., 2017).

676

678	Data availability. The data are available upon request to the corresponding author								
679	(yongzhang@fjnu.edu.cn).								
680									
681	Author contributions. YZ (yongzhang@fjnu.edu.cn), ZL and KX contributed to the								
682	design of the experiment. YZ (yongzhang@fjnu.edu.cn), YZ								
683	(qsx20211022@student.fjnu.edu.cn), SM, HC and RH performed this experiment and								
684	biochemical analyses. YZ (yongzhang@fjnu.edu.cn) wrote the first manuscript draft.								
685	All authors contributed to the data analyses and editing of the paper.								
686									
687	Competing interests. The authors declare that they have no conflict of interest.								
688									
689	Acknowledgments. We would like to thank Professor Zoe V. Finkel for providing the								
690	Emiliania huxleyi RCC1266, Dr. Vinitha Ebenezer for her helpful revision of the								
691	manuscript, and Dr. Peng Jin and two anonymous reviewers for their helpful								
692	suggestions which have help us to improve the manuscript. This work was supported								
693	by the National Natural Science Foundation of China (41806129 [YZ], 32001180								
694	[ZKL]).								
695									
696									
697									
698									
699									
700									

701 **References**

- Beaufort, L., Couapel, M., Buchet, N., Claustre, H., and Goyet, C.: Calcite production
 by coccolithophores in the southeast Pacific Ocean, Biogeosciences, 5, 1101–
 1117, doi:10.5194/bg-5-1101-2008, 2008.
- Berges, J. A., Franklin, D. J., and Harrison, P. J.: Evolution of an artificial seawater
 medium: improvements in enriched seawater, artificial water over the past two
 decades, J. Phycol., 37, 1138–1145, doi: 10.1046/j.1529-8817.2001.01052.x,
 2001.
- Borchard, C., Borges, A. V., Händel, N., and Engel, A.: Biogeochemical response of *Emiliania huxleyi* (PML B92/11) to elevated CO₂ and temperature under
 phosphorus limitation: A chemostat study, J. Exp. Mar. Biol. Ecol., 410, 61–71,
 doi: 10.1016/j.jembe.2011.10.004, 2011.
- Caldeira, K., and Wickett, M. E.: Oceanography: anthropogenic carbon and ocean pH,
 Nature, 425, 365, doi: 10.1038/425365a, 2003.
- Chen, P., Toribara, T. T., and Warner, H.: Microdetermination of phosphorus, Anal.
 Chem., 28, 1756–1758, doi: 10.1021/ac60068a036, 1956.
- Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practice for ocean
 CO₂ measurements, PICES Special Publication 3, pp. 191, 2007.
- Dickson, A. G.: pH buffers for seawater media based on the total hydrogen ion
 concentration scale, Deep-Sea Res., 40, 107–118, doi:
 10.1016/0967-0637(93)90055-8, 1993.
- 722 Dyhrman, S. T.: Nutrients and their acquisition: phosphorus physiology in microalgae,
- in: The physiology of microalgae, edited by: Borowitzka, M. A., Beardall, J. and
- 724 Raven, J. A., Springer, Heidelberg, 155–183, doi: 10.1007/978-3-319-24945-2,
- **725** 2016.

- Fabry, V. J., and Balch, W. M.: Direct measurements of calcification rates in
 planktonic organisms, in: Guide to best practices for ocean acidification research
 and data reporting, edited by: Riebesell, U., Fabry, V. J., Hansson, L. and
 Gattuso, J. P., Luxembourg, Publications Office of the European Union, p. 201–
 212, doi: 10.2777/66906, 2010.
- Feng, Y. Y., Roleda, M. Y., Armstrong, E., Summerfield, T. C., Law, C. S., Hurd, C.
 L., and Boyd, P. W.: Effects of multiple drivers of ocean global change on the
 physiology and functional gene expression of the coccolithophore *Emiliania huxleyi*, Glob. Chang. Biol., 26, 5630–5645, doi: 10.1111/GCB.15259, 2020.
- Gao, K., Beardall, J., Häder, D. P., Hall-Spencer, J. M., Gao, G., and Hutchins, D. A.:
 Effects of ocean acidification on marine photosynthetic organisms under the
 concurrent influences of warming, UV radiation, and deoxygenation, Front. Mar.
 Sci., 6, 322, https://doi.org/10.3389/fmars.2019.00322, 2019.
- Geider, R. J., and LaRoche, J.: Redfield revisited: variability of C:N:P in marine
 microalgae and its biochemical basis, Eur. J. Phycol., 37, 1–17, doi:
 10.1017/S0967026201003456, 2002.
- 742 Gibbs, S. J., Poulton, A. J., Brown, P. R., Daniels, C. J., Hopkins, J., Young, J. R.,
- Jones, H. L., Thiemann, G. J., O'Dea, S. A., and Newsam, C.: Species-specific
- growth response of coccolithophores to Palaeocene-Eocene environmental
- 745 change, Nat. Geosci., 6, 218–222, doi: 10.1038/ngeo1719, 2013.
- Guillard, R. R. L., and Ryther, J. H.: Studies of marine planktonic diatoms. I.
 Cyclotella nana Hustedt and Detonula confervacea Cleve, Can. J. Microbiol., 8,
 229–239, doi: 10.1139/m62-029, 1962.
- Hansen, H. P., and Koroleff, F.: Determination of nutrients, in: Methods of seawater
 analysis, edited by: Grasshoff, K., Kremling, K. and Ehrhardt, M., WILEY-VCH

- 751 Publishers, ISBN: 3-527-29589-5, 1999.
- He, S., Le, C., He, J., and Liu, N.: Empirical algorithm for detecting coccolithophore
 blooms through satellite observation in the Barents Sea, Remote Sens. Environ.,
 270, 112886, doi: 10.1016/j.res.2021.112886, 2022.
- Heidenreich, E., Wördenweber, R., Kirschhöfer, F., Nusser, M., Friedrich, F., Fahl, K., 755 756 Kruse, O., Rost, B., Franzreb, M., Brenner-Weiß, G., and Rokitta, S.: Ocean 757 acidification has little effect on the biochemical composition of the 758 coccolithophore Emiliania huxlevi. PLoS One, 14. e0218564, doi: 759 10.1371/journal.pone.0218564, 2019.
- Jin, P., Gao, G., Liu, X., Li, F., Tong, S., Ding, J., Zhong, Z., Liu,N., and Gao, K.:
 Contrasting photophysiological characteristics of phytoplankton assemblages in
 the Northern South China Sea, PLoS ONE, 11, e0153555, doi:
 10.1371/journal.pone.0153555, 2016.
- Jin, P., Ding, J. C., Xing, T., Riebesell, U., and Gao, K. S.: High levels of solar
 radiation offset impacts of ocean acidification on calcifying and non-calcifying
 strains of *Emiliania huxleyi*, Mar. Ecol. Prog. Ser., 568, 47–58, doi:
 10.3354/meps12042, 2017.
- Jordan, R. W., and Winter, A: Assemblages of coccolithophorids and other living
 microplankton off the coast of Puerto Rico during January–May 1995, Mar.
 Micropaleontol., 39, 113–130, doi: 10.1016/S0377-8398(00)00017-7, 2000.
- Kondrik, D. V., Pozdnyakov, D. V., and Johannessen, O. M.: Satellite evidence that *E*.
- *huxleyi* phytoplankton blooms weaken marine carbon sinks, Geophys. Res. Lett.,
- 773 45, e2017GL076240, doi: 10.1002/2017GL076240, 2018.
- Kottmeier, D. M., Rokitta, S. D., and Rost, B.: Acidification, not carbonation, is the
 major regulator of carbon fluxes in the coccolithophore *Emiliania huxleyi*, New

- 776 Phytol., 211, 126–137, doi: 10.1111/nph.13885, 2016.
- Kubryakova, E. A., Kubryakov, A. A., and Mikaelyan, A. S.: Winter coccolithophore
 blooms in the Black Sea: Interannual variability and driving factors, J. Mar. Syst.,
 213, 103461, doi: 10.1016/j.jmarsys.2020.103461, 2021.
- Larsen, A., Flaten, G. A. F., Sandaa, R., Castberg, T., Thyrhaug, R., Erga, S. R.,
 Jacquet, S., and Bratbak, G.: Spring phytoplankton bloom dynamics in
 Norwegian coastal waters: Microbial community succession and diversity,
 Limnol. Oceanogr., 49, 180–190, doi: 10.4319/lo.2004.49.1.0180, 2004.
- Leonardos, K., and Geider, R. J.: Elevated atmospheric carbon dioxide increases
 organic carbon fixation by *Emiliania huxleyi* (haptophyta), under nutrient-limited
 high-light conditions, J. Phycol., 41, 1196–1203, doi:
 10.1111/j.1529-8817.2005.00152.x, 2005.
- Lohbeck, K., Riebesell, U., and Reusch, T. B. H.: Gene expression changes in the
 coccolithophore *Emiliania huxleyi* after 500 generation of selection to ocean
 acidification, Proc. R. Soc. B., 281, 20140003, doi: 10.1098/rspb.2014.0003,
 2014.
- Lu, Y., Wen, Z., Shi, D., Chen, M., Zhang, Y., Bonnet, S., Li, Y., Tian, J., and Kao, S.
 J.: Effect of light on N₂ fixation and net nitrogen release of *Trichodesmium* in a field study, Biogeosciences, 15, 1–12, doi: 10.5194/bg-15-1-2018, 2018.
- Mackinder, L., Wheeler, G., Schroeder, D., von Dassow, P., Riebesell, U., and
 Brownlee, C.: Expression of biomineralization-related ion transport genes in *Emiliania huxleyi*, Env. Microbiol., 13, 3250–3565,
 doi:10.1111/j.1462-2920.2011.02561.x, 2011.
- Masuko, T., Minami, A., Iwasaki, N., Majima, T., Nishimura, S. I., and Lee, Y. C.:
 Carbohydrate analysis by a phenol-sulfuric acid method in microplate format,

- 801 Anal. Biochem., 339, 69–72, doi: 10.1016/j.ab.2004.12.001, 2005.
- Matthiessen, B., Eggers, S. L., and Krug, S. A.: High nitrate to phosphorus regime
 attenuates negative effects of rising pCO₂ on total population carbon
 accumulation, Biogeosciences, 9, 1195–1203, doi: 10.5194/bg-9-1195-2012,
 2012.
- McKew, B. A., Metodieva, G., Raines, C. A., Metodiev, M. V., and Geider, R. J.:
 Acclimation of *Emiliania huxleyi* (1516) to nutrient limitation involves precise
 modification of the proteome to scavenge alternative sources of N and P,
 Environ. Microbiol., 17, 4050–4062, doi: 10.1111/1462-2920.12957, 2015.
- Meyer, J., and Riebesell, U.: Reviews and syntheses: responses of coccolithophores to
 ocean acidification: a meta-analysis, Biogeosciences, 12, 1671–1682, doi:
 10.5194/bg-12-1671-2015, 2015.
- Moreno, A. R., and Martiny, A. C.: Ecological stoichiometry of ocean plankton, Ann.
 Rev. Mar. Sci., 10, 43–69, doi: 10.1146/annurev-marine-121916-063126, 2018.
- 815 Müller, M. N., Antia, A. N., and LaRoche, J.: Influence of cell cycle phase on
- 816 calcification in the coccolithophore *Emiliania huxleyi*, Limnol. Oceanogr., 53,
 817 506–512, doi: 10.4319/lo.2008.53.2.0506, 2008.
- Ni, G., Zimbalatti, G., Murphy, C. D., Barnett, A. B., Arsenault, C. M., Li, G.,
 Cockshutt, A. M., and Campbell, D. A.: Arctic *Micromonas* uses protein pools
 and non-photochemical quenching to cope with temperature restrictions on
 Photosystem II protein turnover, Photosynth. Res., 131, 203–220, doi:
 10.1007/s11120-016-0310-6, 2016.
- Pakulski, J. D., and Benner, R.: An improved method for the hydrolysis and MBTH
 analysis of dissolved and particulate carbohydrates in seawater, Mar. Chem., 40,
- 825 143–160, doi: 10.1016/0304-4203(92)90020-B, 1992.

- Perrin, L., Probert, I., Langer, G., and Aloisi, G.: Growth of the coccolithophore *Emiliania huxleyi* in light- and nutrient-limited batch reactors: relevance for the
 BIOSOPE deep ecological niche of coccolithophores, Biogeosciences, 13, 5983–
 6001, doi: 10.5194/bg-13-5983-2016, 2016.
- Pierrot, D., Lewis, E., and Wallace, D. W. R.: MS Excel program developed for CO₂
 system calculations, ORNL/CDIAC-105, Carbon Dioxide Information Analysis
 Centre, Oak Ridge National Laboratory, U.S., Department of Energy, doi:

833 10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a, 2006.

- 834Rversion3.5.0.:TheRfoundationforstatisticalcomputingplatform,835x86_64-w64-mingw32/x64.Availablefrom:
- 836 https://cran.r-project.org/bin/windows/ base/old/3.5.0, 2018.
- Reith, M. E., and Cattolico, R. A.: Chloroplast protein synthesis in the chromophytic
 alga *Olisthodiscus luteus*, Plant Physiol., 79, 231–236, doi: 10.1104/pp.79.1.231,
 1985.
- Riebesell, U., Bach, L. T., Bellerby, R. G. J., Rafael Bermúdez Monsalve, J.,
 Boxhammer, T., Czerny, J., Larsen, A., Ludwig, A., and Schulz, K. G.:
 Competitive fitness of a predominant pelagic calcifier impaired by ocean
 acidification, Nat. Geosci., 10, 19–24, dio: 10.1038/NGEO2854, 2017.
- Riegman, R., Stolte, W., Noordeloos, A. A. M., and Slezak, D.: Nutrient uptake and
 alkaline phosphatase (EC 3:1:3:1) activity of *Emiliania huxleyi*(Prymnesiophyceae) during growth under N and P limitation in continuous
 cultures, J. Phycol., 36, 87–96, 2000.
- Rokitta, S. D., von Dassow, P., Rost, B., and John, U.: P- and N-depletion trigger
 similar cellular responses to promote senescence in eukaryotic phytoplankton,
 Front. Mar. Sci., 3, 109, doi: 10.3389/fmars.2016.00109, 2016

- Rokitta, S. D., and Rost, B: Effects of CO₂ and their modulation by light in the
 life-cycle stages of the coccolithophore *Emiliania huxleyi*, Limnol. Oceanog., 57,
 607–618, doi: 10.4319/lo.2012.57.2.0607, 2012.
- 854 Rokitta, S. D., de Nooijer, L. J., Trimborn, S., de Vargas, C., Rost, B., and John, U.: Transcriptome analyses reveal differential gene expression patterns between the 855 856 life-cycle stages of Emiliania huxleyi (haptophyta) and reflect specialization to 857 different ecological niches, J. Phycol., 47, 829-838. doi: 858 10.1016/S0924-9338(02)00624-7, 2011.
- Rost, B., and Riebesell, U.: Coccolithophores and the biological pump: responses to
 environmental changes, in: Coccolithophores : from molecular biology to global
 impact, edited by: Thierstein, H. R. and Young, J. R., Springer, Berlin, 99–125,
 2004.
- Rouco, M., Branson, O., Lebrato, M., and Iglesias-Rodr guez, M.: The effect of
 nitrate and phosphate availability on *Emiliania huxleyi* (NZEH) physiology
 under different CO₂ scenarios, Front. Microbiol., 4, 1–11, doi:
 10.3389/fmicb.2013.00155, 2013.
- Roy, R. N., Roy, L. N., Vogel, K. M., Porter-Moore, C., Pearson, T., Good, C. E.,
 Millero, F. J., and Campbell, D. C.: Thermodynamics of the dissociation of boric
 acid in seawater at S 5 35 from 0 degrees C to 55 degrees C, Mar. Chem., 44,

870 243–248, doi: 10.1016/0304-4203(93)90206-4, 1993.

- 871 Shemi, A., Schatz, D., Fredricks, H. F., Van Mooy, B. A. S., Porat, Z., and Vardi, A.:
- 872 Phosphorus starvation induces membrane remodeling and recycling in *Emiliania*
- *huxleyi*, New Phytol., 211, 886–898, doi: 10.1111/nph.13940, 2016.

- Solórzano, L., and Sharp, J. H.: Determination of total dissolved phosphorus and
 particulate phosphorus in nature waters, Limnol. Oceanogr., 25, 754–758, doi:
 10.4319/lo.1980.25.4.0754, 1980.
- 877 Steinacher, M., Joos, F., Frölicher, T. L., Bopp, L., Cadule, P., Cocco, V., Doney, S.
- 878 C., Gehlen, M., Lindsay, K., Moore, J. K., Schneider, B., and Segschneider, J.:
 879 Projected 21st century decrease in marine productivity: a multi-model analysis,

880 Biogeosciences, 7, 979–1005, doi: 10.5194/bg-7-979-2010, 2010.

- 881 Sterner, R. W., and Elser, J. J.: Ecological Stoichiometry: The Biology of Elements
 882 from Molecules to the Biosphere, Princeton University Press, Princeton, 2002.
- Terrats, L., Claustre, H., Cornec, M., Mangin, A., and Neukermans, G.: Detection of
 coccolithophore blooms with BioGeoChemical-Argo floats, Geophys. Res. Lett.,
 47, e2020GL090559, doi: 10.1029/2020GL090559, 2020.
- Townsend, D. W., Keller, M. D., Holligan, P. M., Ackleson, S. G., and Balch, W. M.: 886 887 Blooms of the coccolithophore Emiliania huxleyi with respect to hydrography in 888 the Gulf of Maine, Cont. Shelf Res., 14, 979-1000, doi: 889 10.1016/0278-4343(94)90060-4, 1994.
- 890 Tyrrell, T., and Merico, A.: *Emiliania huxleyi*: bloom observations and the conditions
- that induce them, in: Coccolithophores : from molecular biology to global impact,
 edited by: Thierstein, H. R. and Young, J. R., Springer, Berlin, 75–97, 2004.
- Wang, C., Wang, J., Li, L., Wang, Y., and Lin, S.: P-limitation promotes carbon
 accumulation and sinking of *Emiliania huxleyi* through transcriptomic
 reprogramming, Front. Mar. Sci., 9, 860222, doi: 10.3389/fmars.2022.860222,
 2022.
- Wang, G., Xie, S. P., Huang, R. X., and Chen. C.: Robust warming pattern of global
 subtropical oceans and its mechanism, J. Climate, 28, 8574–8584, doi:

899 10.1175/jcli-d-14-00809.1, 2015.

- Wilson, D. N., and Doudna Cate J. H.: The structure and function of the eukaryotic
 ribosome, CSH Perspect. Biol., 4, a011536, doi: 10.1101/cshperspect.a011536,
 2012.
- 203 Zhang, Y., Collins, S., and Gao, K.: Reduced growth with increased quotas of
 particulate organic and inorganic carbon in the coccolithophore *Emiliania*pos *huxleyi* under future ocean climate change conditions, Biogeosciences, 17, 6357–
 po6 6375, doi: 10.5194/bg-17-6357-2020, 2020.
- 907 Zhang, Y., Li, Z. K., Schulz, K. G., Hu, Y., Irwin, A. J., and Finkel, Z. V.: 908 Growth-dependent changes in elemental stoichiometry and macromolecular 909 allocation in the coccolithophore Emiliania huxleyi under different 910 environmental conditions, Limnol. Oceanogr., 66, 2999-3009, doi: 911 10.1002/lno.11854, 2021.
- 912
- 913
- 914
- 915

- 917
- 918
- 919
- 920
- 921
- 922
- 923

924 Figure Legends

925 Figure 1. Growth rate (a), cellular contents of particulate organic carbon (POC, b), 926 nitrogen (PON, c) and phosphorus (POP, d), and particulate inorganic carbon (PIC, e) 927 of Emiliania huxleyi RCC1266 in the treatments of high phosphorus availability and 928 low CO₂ level (HP+LC), high phosphorus availability and high CO₂ level (HP+HC), 929 low phosphorus availability and low CO₂ level (LP+LC), and low phosphorus 930 availability and high CO₂ level (LP+HC) under low light (empty, 40 µmol photons m⁻ 2 s⁻¹) and high light (fill, 300 µmol photons m⁻² s⁻¹) intensities. Different letters 931 932 represent significant differences in each parameters between treatments (p < 0.05). 933 The data represents the means and standard deviation of four independent cultures.

934

935 Figure 2. Cellular contents of carbohydrate (a) and protein (b), and the percentages of 936 POC allocated to carbohydrate (c) and protein (d), and the percentage of PON allocated to protein (e) of *E. huxleyi* RCC1266 in the treatments of high phosphorus 937 938 availability and low CO₂ level (HP+LC), high phosphorus availability and high CO₂ 939 level (HP+HC), low phosphorus availability and low CO₂ level (LP+LC), and low 940 phosphorus availability and high CO₂ level (LP+HC) under low light (empty) and 941 high light (fill) intensities. Different letters represent significant differences in each 942 parameters between treatments (p < 0.05). The data represents the means and standard deviation of four independent cultures. Please see figure 1 for more detailed 943 944 information.

Figure 3. Cellular POC : POP ratio (a), PON : POP ratio (b), and protein content (c)
of *E. huxleyi* RCC1266 as a function of growth rate, and cellular POC content as a
function of carbohydrate (d) in the treatments of high phosphorus availability and low

949	CO ₂ level (HP+LC, \Box), high phosphorus availability and high CO ₂ level (HP+HC,
950	\bigcirc), low phosphorus availability and low CO ₂ level (LP+LC, \triangle), and low phosphorus
951	availability and high CO ₂ level (LP+HC, \diamond) under low light (LL, empty) and high
952	light (HL, fill) intensities. Each point indicates an individual replicate under each
953	treatment. Please see figure 1 for more detailed information.
954	
955	
956	
957	
958	
959	
960	
961	
962	
963	
964	
965	
966	
967	
968	
969	
970	
971	
973	
2.2	39

Table 1. Carbonate chemistry parameters and dissolved inorganic phosphorus (DIP) concentration at the end of the incubation. The values are means \pm standard deviation (sd) of four replicates. Respectively, LL and HL represent 40 and 300 µmol photons m^{-2} s⁻¹ of photosynthetically active radiation (PAR), and HP and LP represent 4 and 0.43 µmol L⁻¹ PO₄³⁻ at the beginning of the incubations.

			nCO_2	рН	ТА	DIC	HCO ₂ -	CO_{2}^{2-}	DIP
			(µatm)	(total scale)	$(\mu mol L^{-1})$				
LL	HP	LC	403±4	8.06±0.01	2346±23	2074±21	1861±18	200±2	3.20±0.03
		HC	881±20	7.77±0.01	2351±33	2216±32	2074±30	114±2	3.12±0.08
	LP	LC	329±4	8.13±0.01	2332±24	2024±22	1787±19	225±3	< 0.04
		HC	730±8	$7.84{\pm}0.01$	2349±24	2189±23	2033±22	132±2	< 0.04
HL	HP	LC	357±9	8.09±0.01	2235±41	1959±35	1749±30	199±6	2.86 ± 0.06
		HC	791±7	$7.80{\pm}0.01$	2296±20	2151±20	2007±18	118±1	$2.70{\pm}0.06$
	LP	LC	303±4	8.16±0.01	2354±13	2024±11	1773±9	241±3	< 0.04
		HC	735±19	7.83±0.01	2319±69	2162±65	2011±60	128±5	< 0.04
	HL	LL HP LP HL HP LP	LL HP LC HC LP LC HC HL HP LC HC LP LC HC	$\begin{array}{c cccc} & & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & $	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	$\begin{array}{c c c c c c c c c c c c c c c c c c c $

993	Table 2. Growth rate (d^{-1}) , cellular contents of POC, PON, POP, PIC, carbohydrate
994	(CHO) and protein (Pro) (pg cell ^{-1}), and the ratios of POC : PON, POC : POP, PON :
995	POP and PIC : POC, and the percentages of POC allocated to carbohydrate (CHO–C :
996	POC) and protein (Pro-C : POC), and the percentage of PON allocated to protein
997	(Pro–N : PON) (%). LC and HC represent low CO ₂ (426 μatm) and high CO ₂ (946
998	µatm) levels, respectively. Please see table 1 for more detailed information.

		Low ligh	t intensity		High light intensity					
	ł	ΙP	LP		H	ΗP	LP			
	LC	HC	LC HC		LC HC		LC	HC		
Growth rate	$0.91{\pm}0.03$	0.88 ± 0.01	$0.83{\pm}0.02$	$0.70{\pm}0.03$	1.35 ± 0.03	1.33 ± 0.02	1.34 ± 0.02	1.12 ± 0.02		
POC	8.34±0.57	8.73 ± 0.32	$8.20{\pm}0.36$	$8.04{\pm}0.24$	10.53 ± 0.70	11.04 ± 0.42	11.73±0.19	12.70±0.50		
PON	$1.49{\pm}0.16$	1.58 ± 0.12	1.23 ± 0.06	1.20 ± 0.10	1.65 ± 0.05	$1.89{\pm}0.07$	1.51 ± 0.07	1.56 ± 0.06		
РОР	0.16±0.01	0.15 ± 0.01	$0.08 {\pm} 0.01$	0.07 ± 0.01	0.22 ± 0.01	0.21 ± 0.01	0.12 ± 0.01	$0.10{\pm}0.01$		
PIC	2.12±0.22	1.45±0.17	2.44±0.11	2.06±0.37	3.74±0.22	2.41±0.41	4.83±0.34	3.79±0.49		
POC:PON	6.57±0.43	6.46±0.52	7.78 ± 0.46	7.87±0.54	7.45±0.28	6.83±0.32	9.09±0.44	9.50±0.11		
POC:POP	133.3±7.8	153.8±13.9	282.4±31.2	313.0±40.5	124.2±3.5	137.8±8.5	259.7±23.2	316.9±30.4		
PON:POP	20.40±2.53	23.98±3.37	36.30±3.54	40.10±7.42	16.68 ± 0.47	20.21±1.47	28.63±2.80	33.33±2.85		
PIC:POC	0.26±0.04	0.17 ± 0.02	$0.30{\pm}0.02$	0.26±0.04	0.36 ± 0.02	0.22 ± 0.04	0.41±0.03	$0.30{\pm}0.04$		
СНО	1.45±0.15	1.81 ± 0.16	1.79±0.16	1.94±0.16	3.58±0.41	4.30±0.17	4.96±0.24	5.85 ± 0.49		
Protein	5.23±0.55	5.37±0.39	3.73±0.27	3.80±0.15	6.45±0.36	6.97±0.22	6.25±0.29	6.28±0.30		
CHO-C:POC	6.95±0.41	8.31±0.85	8.71±0.70	9.64±0.58	13.62±1.43	15.60±0.98	16.92±1.04	18.39±0.96		
Pro-C:POC	33.26±3.24	32.58±1.98	24.15±2.52	25.07±1.69	32.49±1.69	33.51±1.41	28.23±1.35	26.21±1.27		
Pro-N:PON	56.84±8.96	54.55±6.51	48.41±2.46	51.07±5.40	62.62 ± 2.88	59.12±1.21	66.35±4.06	64.44±2.73		





1012 Figure 1



Figure 2



1024 Figure 3

Supplemental Information

Materials and Methods

<u>1 RNA Analyses</u>

Total RNA was extracted by using TRIzol reagent (Invitrogen) and the RNeasy Plus Mini Kit (Qiagen). Samples were resuspended in 1.2 mL TRIzol reagent (Thermo Fisher Scientific) in lysing matrix D tubes (MP Biomedicals), homogenized by a FastPrep-24 machine (MP Biomedicals, 3 cycles, 8.0 m s⁻¹, 30 s, 3 min ice-chilling at interval), followed by centrifugation at 12,000 \times g for 3 min at 4 °C (Eppendorf 5430R). RNA was extracted using a standard phenol-chloroform method (Chomczynski and Sacchi 1987). For more detail information, please see Zhang et al. (2021).

<u>2 Bioinformatics</u>

The quality of the raw reads was assessed by Fastqc v.0.11.8 (Andrews, 2010) and Fastq screen v.0.13.0 (Wingett and Andrews, 2018) and summarized using Multiqc (Ewels et al., 2016). Trimming of the raw reads was performed to remove low-quality bases and adapter sequences with Trimmomatic v.0.38 (Bolger et al., 2014). *De novo* transcriptome assembly was performed with the Trinity's version 2.11.0 (Grabherr et al., 2011), and the low-quality assembly were removed with CD-HIT (Li and Godzik, 2006). A preliminary assessment of *de novo* assembly quality was performed with Transrate (Smith-Unna et al., 2016) and Busco (Hara et al., 2015), and the completeness assessment yielded high scores for all assemblies. Open reading frames (ORFs) were then predicted using TransDecoder version v5.5.0 (Haas et al., 2013), and were then annotated by Blastx, Hmmpress, Signalp, Rnammer, PFam (Lagesen et al., 2007). All annotations were loaded and integrated with Trinotate v3.0.0 (Haas et al., 2007).

al., 2013). ORFs were further functionally annotated and assigned to the KEGG and GhostKOALA (Moreno-Santillán et al., 2019). Cleaned and trimmed reads of each sample were mapped to the assembled transcriptomes by salmon v0.9.1 (Patro et al., 2017). The differential expression was them calculated by DESeq2 v1.24.0 (Love et al., 2014) with a Benjamini-Hochberg adjusted *p*-value < 0.05. Data analysis and visualizations were made using R v.3.6.1 (Team, 2020), packages ggplot2 v.3.2.0 (Wickham, 2016) and Pheatmap v.1.0.12 (Kolde, 2015).

Table S1. Results of three-way ANOVAs of the effects of light intensity (L), dissolved inorganic phosphorus (PO_4^{3-}) concentration (P), pCO_2 level (C), and their interaction on growth rate, cellular contents of POC, PON, POP, PIC, carbohydrate and protein, and the ratios of POC : PON, POC : POP, PON : POP and PIC : POC, and the percentages of POC allocated to carbohydrate (carbohydrate–C : POC) and protein (protein–C : POC), and the percentage of PON allocated to protein (protein–N : PON). Please see figure 1 for more detailed information.

Parameter		L	Р	С	L×P	L×C	P×C	L×P×C
Growth rate	F	4588.5	198.3	225.7	6.7	14.7	100.9	8.6
	р	< 0.01	< 0.01	< 0.01	=0.02	< 0.01	< 0.01	< 0.01
POC	F	864.7	0.6	0.6	36.4	8.1	0.2	2.6
	р	< 0.01	=0.46	=0.46	< 0.01	< 0.01	=0.65	=0.12
PON	F	172.2	70.6	17.7	1.2	6.3	10.0	0.2
	р	< 0.01	< 0.01	< 0.01	=0.33	=0.02	< 0.01	=0.65
РОР	F	188.3	724.8	8.2	6.6	0.1	0.2	0.2
	р	< 0.01	< 0.01	< 0.01	=0.02	=0.89	=0.65	=0.65
PIC	F	329.4	109.1	47.2	17.7	8.9	1.9	0.1
	р	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	=0.17	=0.98
POC : PON	F	207.6	396.4	0.1	14.9	0.2	9.3	2.1
	р	< 0.01	< 0.01	=0.96	< 0.01	=0.65	< 0.01	=0.16
POC : POP	F	7.7	351.3	18.7	0.1	0.3	2.6	1.0
	р	=0.01	< 0.01	< 0.01	=0.85	=0.62	=0.12	=0.32
PON : POP	F	29.4	183.1	29.9	1.9	0.1	0.3	0.1
	р	< 0.01	< 0.01	< 0.01	=0.18	=0.97	=0.61	=0.94
PIC : POC	F	90.2	71.9	77.6	0.6	6.6	2.4	0.3
	р	< 0.01	< 0.01	< 0.01	=0.46	=0.02	=0.14	=0.61
Carbohydrate	F	925.2	37.7	23.6	40.9	7.9	0.9	0.9
	р	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	=0.34	=0.34
Protein	F	782.4	77.0	1.0	21.1	0.8	2.0	0.8
	р	< 0.01	< 0.01	=0.34	< 0.01	=0.38	=0.17	=0.37
Carbohydrate-C : POC	F	793.3	76.2	17.1	4.9	0.7	0.6	0.1
	р	< 0.01	< 0.01	< 0.01	=0.04	=0.41	=0.43	=0.95
Protein–C : POC	F	6.3	120.0	0.7	2.4	0.6	2.2	2.7
	р	=0.02	< 0.01	=0.42	=0.13	=0.45	=0.15	=0.11
Protein-N : PON	F	36.3	4.3	2.8	9.2	0.7	0.7	0.2
	р	< 0.01	=0.05	=0.10	< 0.01	=0.41	=0.41	=0.63



Figure S1. A scanning electron microscopy (SEM) picture of the coccosphere and coccolith of *E. huxleyi* RCC 1266.





Figure S24. Flow chart of the experimental processes (**a**) and flow chart of the pre-culture and experimental cultures in each treatment (**b**). At the beginning of the incubations, HP and LP represent 4 and 0.43 μ mol L⁻¹ PO₄^{3–}, and LC and HC represent low (about 426 μ atm) and high CO₂ (about 946 μ atm) concentrations, respectively.



Figure S32. Cell density of *E. huxleyi* RCC1266 in the treatments of HP+LC, HP+HC, LP+LC and LP+HC under low light (empty) and high light (fill) intensities during the experimental cultures. The cells were inoculated to achieve an initial density of about 5000 cell mL⁻¹, and cultured in each experimental condition for 2 days and then diluted to the initial cell density again. This process was repeated four times. The data represents the means and standard deviation of four independent cultures in the fourth incubation. Please see figure 1 for more detailed information.



Figure S43. In comparison to the HP+LC treatment, changes in growth rate (\mathbf{a}, \mathbf{b}) and cellular contents of POC (\mathbf{c}, \mathbf{d}) , PON (\mathbf{e}, \mathbf{f}) , POP (\mathbf{g}, \mathbf{h}) , PIC (\mathbf{i}, \mathbf{j}) , carbohydrate (\mathbf{k}, \mathbf{l}) and protein (\mathbf{m}, \mathbf{n}) of *E. huxleyi* RCC1266 in the treatments of HP+HC, LP+LC and LP+HC under low light (empty) and high light (fill) intensities. The data represents the means of four independent cultures. Please see figure 1 for more detailed information.



Figure S54. The ratios of POC : PON (**a**), POC : POP (**b**), PON : POP (**c**) and PIC : POC (**d**) of *E. huxleyi* RCC1266 in the treatments of HP+LC, HP+HC, LP+LC and LP+HC under low light (empty) and high light (fill) intensities. Different letters represent significant differences in each parameters between treatments (p < 0.05). The data represents the means and standard deviation of four independent cultures. Please see figure 1 for more detailed information.



Figure S<u>6</u>5. Cellular RNA content (a) and POC normalized RNA content (b) of *E. huxleyi* RCC1266 in the treatments of HP+LC and LP+LC under high light intensity. Different letters in each panel represent significant differences between treatments (p < 0.05). The data represents the means and standard deviation of four independent cultures. Please see figure 1 for more detailed information.



Figure S6. Cell density of *E. huxleyi* RCC1266 under different phosphorus concentrations (P) at 150 μ mol photons m⁻² s⁻¹ of photosynthetically active radiation under low CO₂-level.



Log2 (Fold change)

Figure S7. Heatmap of significant change in expression of genes linked to ribosome metabolism (Ribosome_ko03010) under the low pH treatment (ocean acidification) in comparison to the high pH treatment (present CO₂ level). Red indicates up-regulation of gene, and blue indicates down-regulation of gene. *RP-S* presents genes linked to small subunit ribosomal protein, and *RP-L* presents genes linked to large subunit ribosomal protein.

References

- Andrews, S.: FASTQC: a quality control tool for high throughput sequence data, Cambridge, 2010. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.
- Bolger, A. M., Lohse, M., and Usadel, B.: TRIMMOMATIC : a flexible trimmer for illumina sequence data, Bioinformatics, 30, 2114–2120, 2014.
- Chomczynski, P., and Sacchi, N.: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction, Anal. Biochem., 162, 156–159, doi: 10.1006/abio.1987.9999, 1987.
- Ewels, P., Magnusson, M., Lundin, S., and Käller, M.: MULTIQC: summarize analysis results for multiple tools and samples in a single report, Bioinformatics, 32, 3047–3048, 2016.
- Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, e., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N, and Regev, A.: Full-length transcriptome assembly from RNA-Seq data without a reference genome, Nat. Biotechnol., 29, 644–652, 2011.
- Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M. B., Eccles, D., Li, B., Lieber M., MacManes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R., Willian, T., Dewey, C. N., Henschet, R., LeDuc, R. D., Friedman, N., and Regev, A: *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis, Nat. Protoc., 8, 1494–1512, 2013.
- Hara, Y., Tatsumi, K., Yoshida, M., Kajikawa, E., Kiyonari, H., and Kuraku, S.: Optimizing and benchmarking *de novo* transcriptome sequencing: from library

preparation to assembly evaluation, BMC Genomics, 16, 977, 2015.

Kolde, R.: Package 'pheatmap'. R. Package, 1, 790, 2015.

- Lagesen, K., Hallin, P., Rodland, E. A., Staerfeldt, H. H., Rognes, T., and Ussery, D.W.: RNAmmer: Consistent and rapid annotation of ribosomal RNA genes, Nucleic Acids Res., 35, 3100–3108, 2007.
- Li, W., and Godzik, A.: Cd-hit: A fast program for clustering and comparing large sets of protein or nucleotide sequences, Bioinformatics, 22, 1658–1659, 2006.
- Love, M. I., Huber, W., and Anders, S.: Moderated estimation of fold change and dispersion for RNA-seq data with DES EQ 2, Genome Biol., 15, 550, 2014.
- Moreno-Santillán, D. D., Machain-Williams, C., Hernández-Montes, G., and Ortega, J.: *De novo* transcriptome assembly and functional annotation in five species of bats, Sci. Rep., 9, 6222, 2019.
- Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., and Kingsford, C.: Salmon provides fast and bias-aware quantification of transcript expression, Nat. Methods., 14, 417–419, 2017.
- R Development Core Team: R: a language and environment for statistical computing, v. 4.0.3. [WWW document] URL www.r-project.org [accessed 10 October 2020].
- Smith-Unna, R., Boursnell, C., Patro, R., Hibberd, J. M. and Kelly, S.: TransRate: Reference-free quality assessment of *de novo* transcriptome assemblies, Genome Res., 26, 1134–1144, 2016.
- Wickham, H.: ggplot2: elegant graphics for data analysis, R package v.3.3.2. [WWW document] URL. https://cran.r-project.org/web/packages/ggplot2/index.html [accessed 17 June 2020], 2016.
- Wingett, S. W and Andrews, S.: FASTQ screen: a tool for multi-genome mapping and quality control, F1000Research, 7, 1338, 2018.

Zhang, Y., Li, Z. K., Schulz, K. G., Hu, Y., Irwin, A. J., and Finkel, Z. V.: Growth-dependent changes in elemental stoichiometry and macromolecular allocation in the coccolithophore *Emiliania huxleyi* under different environmental conditions, Limnol. Oceanogr., 66, 2999–3009, doi: 10.1002/lno.11854, 2021.