

Signatures of the Madden-Julian Oscillation in Middle Atmosphere zonal mean Temperature: Triggering the Interhemispheric Coupling pattern

Christoph G. Hoffmann¹, Lena G. Buth^{1,2}, and Christian von Savigny¹

¹Institute of Physics, University of Greifswald, Felix-Hausdorff-Str. 6, 17489 Greifswald, Germany

²now at Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

Correspondence: Christoph G. Hoffmann (christoph.hoffmann@uni-greifswald.de)

Abstract. The influence of the Madden-Julian oscillation (MJO) ~~is the dominant mode of intraseasonal variability in the troposphere. The influence of the MJO~~ on the middle atmosphere (MA) and particularly on ~~its MA~~ temperature is of interest for both the understanding of MJO-induced teleconnections and research on the variability of the ~~middle atmosphere. However, only few studies dealing with the influence of the MJO on MA temperature are available.~~

5 MA. We analyze statistically the connection of the MJO and the MA zonal mean temperature based on observations by the MLS satellite instrument. We consider all eight MJO phases, different seasons and the state of the quasi-biennial oscillation (QBO). We show that the MA temperature is influenced by the state of MJO in large areas of the MA and under roughly all considered atmospheric conditions. The zonal mean temperature response ~~is characterized by~~ shows a particular spatial pattern, which we link to the interhemispheric coupling (IHC) mechanism, ~~a known dynamical feature of the MA as a major outcome of this study.~~ The strongest temperature deviations are on the order of $\pm 10\text{K}$ and are found in the polar winter MA during boreal winter ~~when the QBO is in the and the QBO~~ easterly phase. ~~Other atmospheric conditions also show temperature responses with the characteristic spatial pattern, but weaker and more noisy. The QBO turns out to have a relatively big influence during boreal winter but only a small influence during austral winter. We also discuss the role of sudden stratospheric warmings (SSWs), which have an ambivalent influence on our interpretation, because they introduce strong temperature variability in the polar winter MA themselves. In addition, SSWs are one possibility to explain the QBO influence during boreal winter. Furthermore, we also~~ We also analyze the change of the temperature response pattern while the MJO progresses from one phase to the next. We find a ~~largely systematic reaction of the MA to the phase changes, particularly a~~ gradual altitude shift of ~~the MA temperature parts of the IHC~~ response pattern, which can be seen more or less clearly depending on the atmospheric conditions.

10 ~~Overall, a major outcome of the present study is the finding that the tropospheric MJO can trigger the IHC mechanism, which affects many areas of the MA. It is therefore a noteworthy example for the complex couplings~~ The found statistical link between the MJO and the MA temperature highlights illustratively the far reaching connections across different atmospheric layers and geographical regions in the atmosphere. Additionally, it highlights close linkages of known dynamical features of the atmosphere, particularly the MJO, the IHC, the QBO, and SSWs.

15

20

25 Because of the wide coverage of atmospheric regions and included dynamical features, the results might help to further constrain the underlying dynamical mechanisms and could be used as a benchmark for the representation of atmospheric couplings on the intraseasonal timescale in atmospheric models.

Copyright statement. TEXT

1 Introduction

30 The **Madden-Julian Oscillation** (MJO), first documented by Madden and Julian (1972), is known as the dominant mode of intraseasonal variability in the tropical troposphere (Zhang, 2005). The major directly observable feature is a center of anomalously strong deep convection, flanked by regions of reduced convection. This structure appears periodically over the Indian Ocean and travels eastward over the maritime continent until it disappears over the Pacific Ocean. The MJO is a highly variable feature of the atmosphere and, e.g., its period varies strongly between about 30 and 90 days 35 (Zhang, 2005). Its appearance influences weather patterns in many equatorial regions, e.g., the monsoons in Asia and Australia (Zhang, 2005).

Although the obvious convective disturbance is limited to the Indo-Pacific region, dynamical anomalies, e.g., in terms of vorticity, are measured around the whole equator (e.g., Cassou, 2008). Moreover, the MJO has also influences around the whole globe and is itself influenced from extratropical regions making the MJO a component of global teleconnection patterns 40 (Lau and Waliser, 2012, Chapter 14). A prominent example in the northern hemisphere is a link between the MJO and the North Atlantic Oscillation (Cassou, 2008). Affected are different parameters of the polar climate system like the Arctic surface temperature (Yoo et al., 2012) or even the Arctic sea ice concentration (Henderson et al., 2014). For a review of the observed teleconnections between the tropics and the polar regions on different time scales see, e.g., Yuan et al. (2018).

Due to the intraseasonal time scale on which the MJO acts, a deeper knowledge on the processes that control it is expected to 45 improve the forecast skills of longer-term weather forecasts (e.g., Zhang, 2013). The understanding of the mechanisms of the teleconnections is important to improve the representation of the MJO in models but might in turn help to improve the forecasts skills also in the extratropics, e.g., in terms of the prediction of extreme weather events (Lau and Waliser, 2012, Chapter 14). Mechanisms for the teleconnections with the extratropics are reviewed by, e.g., Yuan et al. (2018). Inner-tropospheric interactions are based on the excitation of Rossby waves by the strong convection, which is associated with the MJO and 50 propagation of zonal mean zonal wind anomalies, which are triggered by the MJO.

A further mechanism for MJO-related teleconnections is proposed by Garfinkel et al. (2012) for the extended boreal winter season. It also considers the propagation of Rossby waves, but this time not only the horizontal poleward propagation but also the vertical propagation into the stratosphere. These waves could influence the polar vortex and modulate the appearance of midwinter major sudden stratospheric warmings (SSW), so that SSWs follow certain MJO phases. It was already known that 55 the occurrence of sudden stratospheric warmings (SSW) during Arctic winter can then influence the tropospheric state below

(e.g. Baldwin and Dunkerton, 2001). The complete chain can therefore lead to an influence of the MJO on the polar troposphere mediated by the stratosphere with a longer time scale (one to two months) than the inner-tropospheric connections.

There is also increasing evidence that the variability of the MJO itself is partly controlled by the stratosphere. A prominent example for this is the influence of the quasi-biennial oscillation (QBO; see Baldwin et al. (2001) for a review of the QBO) during boreal winter: Yoo and Son (2016) found that the MJO amplitude tends to be larger during the QBO easterly phase and smaller during the QBO westerly phase and several studies followed up on this (e.g., Son et al., 2017; Marshall et al., 2017; Zhang and Zhang, 2018; Densmore et al., 2019; Wang and Wang, 2021). The influence on the MJO from above is also covered by a more general recent review on the stratospheric influence on the tropical troposphere by Haynes et al. (2021).

Consequently, there is increasing awareness for the potential of considering troposphere-stratosphere couplings in both directions in weather prediction systems, as shown by, e.g., Domeisen et al. (2020). And at least from these examples, it becomes clear that it is of importance to consider at least the stratosphere to completely understand the functioning and interactions of the MJO in the climate system.

In addition to this troposphere-related motivation, there is a second major motivation to study the connection of MJO and the middle atmosphere (MA), which arises from the interest in the variability of the MA itself. One important aspect of MA research is the determination of a long-term trend in MA temperature, which is an important indicator of anthropogenic climate change (e.g., Randel et al., 2009; Santer et al., 2013; Maycock et al., 2018; Beig et al., 2003; Beig, 2011). Other aspects are, e.g., the understanding of the implications of solar variability on the climate system (e.g., Gray et al., 2010), the influence of large volcanic eruptions (e.g., Timmreck, 2012), and of course the monitoring of stratospheric ozone. All these aspects have in common that they are connected to temporal variability of different MA parameters, with temperature being an essential one. A major difficulty is the disentanglement of all the sources of temperature variability and the unambiguous attribution to the respective causes to understand the individual processes separately. In order to increase the robustness of analyses and interpretations, also the smaller sources of temperature variability should be known and quantified.

The MJO has, to our knowledge, gotten only little attention as a one possible independent source of temperature variability in the MA. In addition to the already mentioned analysis by Garfinkel et al. (2012), the studies by Yang et al. (2017, 2019) are in this context of relevance for our analysis. These publications cover the MJO influence on the MA temperature among other parameters for different geographical regions, seasons and atmospheric conditions. They are mostly based on modeled and reanalyzed data, whereas it is our aim to provide a purely observational perspective. Sun et al. (2021) analyze the effect of the MJO on the northern mesosphere during boreal winter. They also mostly rely on modeled data but use satellite observations as support, particularly, the same observational dataset as we use in the following (Sect. 2.1). There are more studies relevant in the broader context, but either they treat the influence on temperature not as the main point (e.g., Moss et al. (2016); Tsuchiya et al. (2016); Wang et al. (2018a), which focus on the MJO influence on wave activity) or are limited to different atmospheric regions (e.g., Yang et al. (2018); Kumari et al. (2020, 2021), which discuss the MJO influence on tides in the mesosphere / lower thermosphere region). The relationships to the relevant studies are discussed in more detail below (Sect. 4).

We note as an additional aspect for the motivation that the occurrence rates of individual MJO phases appear to be subject to climate change (Yoo et al., 2011). Therefore, a linkage of the MJO and the MA could represent an additional pathway of

anthropogenic climate change into the MA. Conversely, the characterization of the MJO-MA relationship might also be helpful to understand climate change related changes in the teleconnection patterns mentioned above.

All the above-mentioned aspects motivate the exploration of already existing observations with respect to the influence of the MJO on the MA, particularly on the MA key parameter temperature. However, the mentioned publications on the MJO influences on the MA temperature mostly deal with modeled or reanalyzed data, which has the advantage that respective mechanisms can be more easily identified in these more comprehensive datasets. Contrarily, this publication aims at contributing a global quantification of the influence of the MJO on MA temperatures based observed data, to gain additional independent and purely observationally determined information ~~as complement to modeling~~ to complement modelling efforts.

Most related studies first determine the MJO phase with the strongest response and concentrate in the following analysis mostly on this phase (e.g., Garfinkel et al., 2012; Yang et al., 2017). The temporal evolution of the atmosphere is then considered in time lags around the appearance of this phase. We consider instead all eight MJO phases, so that the transition of the MA temperature response from one MJO phase to the next becomes visible. Furthermore, we concentrate in this paper on the analysis of zonal mean temperatures for the sake of brevity. Nevertheless, the dataset provides also a wealth of three-dimensionally resolved information, which we intend to present in a later paper.

The present paper contains a description of the datasets and methodology in Sect. 2. The results, i.e. the zonal mean MA temperature responses to the individual MJO phases for different atmospheric conditions, are shown in Sect. 3. Section 4 contains an extensive discussion, in which the results are related to known dynamical features of the MA, before we conclude the paper in Sect. 5

2 Datasets and analysis approach

110 2.1 Datasets

We analyze the temperature data product measured by NASA's Microwave Limb Sounder (MLS) on the Aura satellite (Schwartz et al., 2020), version 5. A previous version of the dataset has been validated by Schwartz et al. (2008). Differences to the new version are mentioned in Livesey et al. (2020). Data screening and exclusion has been performed according to all suggestions in the MLS quality document (Livesey et al., 2020), including the screening for cloud effects based on the MLS ice water content (IWC) product. We analyze the complete reasonable vertical range from 261 hPa to 0.00046 hPa and also use the complete temporal coverage of the ongoing measurements of about 17 years at the time of the analysis (approximately August 2004 to September 2021). The data has a spatial resolution, which roughly decreases with height in a range from about 4 km vertical and 170 km horizontal resolution at 261 hPa to 13 km vertical and 316 km horizontal resolution at 0.00046 hPa (Livesey et al., 2020). The data has been analyzed on the original pressure grid. One has to keep in mind that the analysis grid is finer than the varying vertical resolution and that adjacent altitudes are not completely independent of each other. The original data follows the satellite track and has been averaged to fit onto temporally and horizontally regular grids. Particularly, for this paper we apply a zonal averaging with 10° resolution in the latitudinal direction.

For the characterization of the MJO, we use the OLR-based MJO Index (OMI; OLR stands for outgoing longwave radiation) introduced by Kiladis et al. (2014). We calculate the OMI values from OLR using the open source OMI calculation package 125 (Hoffmann et al., 2021; Hoffmann, 2021), version 1.2.2. According to Hoffmann et al. (2021) a high agreement between with the original calculation routine, which is close to being identical, is expected. For the calculation, interpolated OLR data according to Liebmann and Smith (1996) has been used. From the two OMI index values per time step, the phase and the strength of the MJO for each day can easily be calculated following the phase diagrams in Kiladis et al. (2014) and Wheeler and Hendon (2004).

130 We note that the MJO is subject to a seasonal variability (e.g., Zhang, 2005). The common MJO concept with the characteristic eastward propagation applies best for boreal winter. During boreal summer, the propagation direction includes a northward component (e.g., Wang et al., 2018b) and the feature is often given a distinct name, the boreal summer intraseasonal oscillation (BSISO). This has also implications for the definition of appropriate MJO indices and partly particular BSISO indices are used for analyses with a special focus on boreal summer (e.g., Kikuchi et al., 2012). However, since our analysis is not 135 restricted to boreal summer and Wang et al. (2018b) states that the MJO index OMI is also capable of reasonably tracking the BSISO, we apply OMI consistently for all seasons. This has the advantage of comparability of the results for different seasons. Nevertheless, we might check our boreal summer results with a special BSISO index in future.

140 For the characterization of the QBO, we use monthly mean zonal wind data above Singapore at 50 hPa (Naujokat, 1986). We apply the most simple discrimination approach: positive zonal wind values indicate a QBO westerly phase and negative values a QBO easterly phase.

2.2 Approach

Our analysis approach is a superposed epoch analysis (SEA) also known as composite analysis. The complete analysis chain consists, however, of several steps, which are computed for each geolocation of the previously mentioned regular spatial grid separately.

145 First, a temperature anomaly is computed by applying a boxcar running average with a window length of 90 d as a low-pass filter and calculating the difference between the original and the low-pass filtered time series. Afterwards the anomaly is smoothed with a 10-day running average. The latter is mainly done for reasons of comparability with other analyses and is strictly speaking disadvantageous since it introduces a statistical dependence of the previously independent individual days. Therefore, we have checked that, in practice, the results do not change significantly when the 10-day filter is not applied. The 150 resulting anomaly time series contains therefore due to the filtering only strongly damped variations on much longer (e.g., the seasonal cycle) and much shorter (e.g. weather fluctuations) time scales than the MJO, but still captures a broad range of frequencies in the intraseasonal time scale, relevant to the MJO.

Second, the data is selected according to the environmental conditions. We take generally only days into account, during 155 which the MJO strength was greater than 1. This is a common choice in many MJO related studies to make sure that there is actually an MJO signature apparent on all considered days. Depending on the particular experimental setups described below, the data is also selected with respect to particular seasons and with respect to the state of the QBO. Specifically, all days, which




Figure 1. Examples of the temperature responses to the eight MJO phases at two geolocations for the boreal winter / QBO easterly situation. The examples have been selected for demonstration purposes to represent a particularly strong response (upper half) and a weak response (lower half). The following description applies to both cases: The main results, the average temperature anomalies for each MJO phase, are shown as the black lines in the larger panels together with the eight standard errors of the mean shown as error bars. The blue line shows the fitted sine curve. Temperature responses, which are significant according to the MCIP method (see Sect. 2.2 for details on the MC quantification) are marked with red stars. Results according to the MCS significance estimation method, which estimates the significance of a systematic variation over all eight phases, are shown in the lower smaller panels on the right, respectively: The blue bars constitute a histogram of all sine amplitudes derived based on random data, whereas the sine amplitude of the real data is shown as a red arrow. The resulting percentage of random amplitudes lower than the real one is also given in the lower right text field. The upper smaller panels on the right simply indicate how many days of the data went into each of the eight temperature anomaly average values.

do not match the wanted setup criteria, are removed from the dataset prior to the SEA. We consider the seasons boreal winter / austral summer (December, January and February) and boreal summer / austral winter (June, July, August). [The complete set of selection criteria for a particular experiment is sometimes called "atmospheric conditions" or similar in the remainder of the manuscript.](#)

Third, the actual SEA is carried out: the selected temperature observations are grouped by the related MJO phases of each day and then averaged, so that one temperature mean value per MJO phase is calculated, complemented by the corresponding standard errors of the mean. The basic idea is that temperature fluctuations, which are not correlated with the MJO phase propagation will average out, while fluctuations correlated with the MJO phases will add up, so that remaining epoch averaged anomalies indicate a statistical connection between the MJO phase propagation and the MA temperature. Note that we apply a small correction after this basic step: a mean over the complete data subset at the respective geolocation (i.e., independent of the MJO phase) is subtracted from all eight average values. This overall mean value is usually close to 0 K, because the whole calculation is carried out on temperature anomalies, which are scattered around zero. However, the average of the selected subset of the data may slightly deviate from zero, which would complicate the comparison of the results for different 165 geolocations if not corrected for. Fig. 1 shows two examples of the SEA results, which represent a strong and a weak response. The number of days that go into the individual averages depends of course strongly on the selection criteria, but also somewhat 170 on the geolocation and the MJO phase itself. Rough numbers are about 400 days per MJO phase if only the MJO strength filtering is applied, 100 days with additionally the seasonal selection and 50 days with a combined seasonal and QBO selection.

As stated before (Sect. 1), we consider all eight phases of the MJO so that the transition of the MA temperature response 175 from one phase to the other becomes visible. At least in an ideal case, the response could vary like sine function over the course of the eight MJO phases. Following this notion, we fit a sine function to the eight mean values as the ~~forth~~^{fourth} analysis step to further characterize the behavior across the MJO phases. The parameters amplitude, phase and offset can freely be adjusted by the fit, whereas the period is fixed to 8 MJO phases. The fit results can be analyzed with different foci. However, in this study 180 only the resulting amplitudes are used for a significance estimation in the next analysis step. The two examples in Fig. 1 also show the fitted sine curves. It is seen that the strong response exhibits indeed a sine-like behavior and the amplitude roughly represents the strength of the deviations, whereas the appearance of the weak response is more noisy, which results in a even weaker sine amplitude.

The fifth and final step is a significance estimation using a Monte Carlo (MC) method. For this, the analysis is basically 185 repeated multiple times, each time with randomly modified input data. The resulting distribution of artificial results can be used to estimate how likely a particular temperature response can be the result of random fluctuations instead of physical reasons. As we have pointed out in Hoffmann and von Savigny (2019), there is a wide scope of individual decisions in the design of the MC calculations, which can influence the final significance estimation. In order to not distract the reader from the basic results, we concentrate here on only one version of the random data generation (as most other publications also do), which is well comparable to many previous publications. Particularly, we randomly redistribute the MJO index time series 1000 190 times, i.e. the attribution of MJO phases and strengths to the individual days of the time series is changed. The temperature data remains thereby untouched. The SEA calculation is repeated for each random data realization resulting in a distribution of possible temperature responses. Whereas we only present one kind of random data generation, we still show two different kinds of the final quantification of the significance. The first one, abbreviated as MCIP for "Monte Carlo Individual Phases" in the following, is also best comparable to previous studies, particularly if they are focused on individual MJO phases. It is 195 simply checked for each MJO phase separately if the absolute value of the real SEA anomaly result for a particular MJO phase

is greater than 95 % of the absolute values of the responses based on random data. Hence, this significance estimation checks if the derived SEA temperature anomaly is strong enough to be only very unlikely produced by random fluctuations for each MJO phase separately. Significant average values according to this method are marked with a red star in Fig. 1. An advantage is that this significance estimation exists separately for each MJO phase. However, this method has the disadvantage that it
200 may underestimate significance. E.g., if we consider the result for MJO phase 1 in the upper panel of Fig. 1, the response is relatively close to 0 K and consequently also likely reproduced by random data and therefore not marked as significant. However, the development of the response over all eight MJO phases indicates that the response of phase 1 could be a part of an overall systematic variation, albeit approximately at the zero-crossing of the variation. Therefore, we use a second quantification approach, abbreviated as MCS for "Monte Carlo Sine" in the following, in which it is checked that the real
205 amplitude of the fitted sine function is greater than 95 % of the fitted amplitudes based on the random data. Hence, this quantification approach evaluates the systematic behavior over all 8 phases and is therefore also based on a larger number of samples (all days that go into the eight averages instead only those, which go into one specific average). Examples of the distributions of sine amplitudes are also given in the lower smaller plots in Fig. 1.

We would like to point out that the overall approach aims at establishing a statistical connection between the MJO phase evolution and the MA temperature without probing a causal or physical connection. Strictly speaking, also the direction of the influence is not checked. However, on the one hand, describing a statistical connection first helps to clearly define the observational aspects, which the ultimately sought physical mechanism has to cover. This is the basic aim of the present manuscript. On the other hand, by considering previous publications, which overlap in particular aspects, it becomes expectable that a physical mechanism actually exists (i.e. that we do not describe a purely statistical artifact) as we will outline in Sect. 4.

215 3 Results: MJO influence on zonal mean MA temperature

This study is focused on the zonal mean response of MA temperature to the MJO. Nevertheless, a quick inspection of the full three-dimensional structure of the response exhibits also zonal structures, particularly in the lower stratosphere. These structures partly cancel out by the zonal averaging so that the zonal mean anomaly values shown in the following are likely lower than some of the individual local responses.

220 3.1 No filtering for environmental conditions

As will become obvious in the following, external conditions like the season and the QBO have a strong influence on the strength and spatial structure of the response. Hence, analyzed data that comprises different states of these conditions might also result in reduced responses due to interferences of the individual responses to these conditions. Therefore, we do not discuss the unfiltered analysis here in detail. ~~Instead, we start directly with the boreal winter situation in the following (Sect. 3.2).~~
225 Nevertheless, we show the results for the unfiltered data (only a filtering for MJO strength greater than 1 is applied) in the supplement (Fig. S1) and note here that the derived temperature anomalies connected to the MJO influence are in this case in the range of ± 2 K.




Figure 2. Meridional plane temperature responses for all eight MJO phases for the boreal winter / austral summer situation. Each panel contains the temperature anomaly response to a particular MJO phase for all geolocations in the zonal mean. All eight panels share the same color scale on the right. Insignificant values according to the MCIP method are marked with gray dots. Insignificant areas according to the MCS method are marked with black crossed lines. Note that the insignificance pattern according to the MCS method is identical for all 8 MJO phases, since the overall systematic behavior is evaluated (slight differences of the hatches between the different panels are due to numerical inaccuracies in the rendering of the image and do not indicate differences in the significance estimation results).

3.2 Boreal winter / austral summer

We start with the data restricted to boreal winter, since it is known that the strongest signal is likely produced by the MJO 230 during that season. Fig. 2 shows the meridional plane temperature responses for each of the eight MJO phases. The strongest response is seen for the MJO phases 5 and 6 in the northern polar region, i.e., in the winter hemisphere. This response is spread vertically and divided into two zones with opposite sign visible as strong red and blue areas in the figure. The departures from the mean state are as high as $\pm 6\text{ K}$ even in this zonal mean picture ~~and are therefore noticeably higher than for the unfiltered case mentioned above (Sect. 3.1).~~

235 A broader overall pattern in the meridional plane response can be recognized more or less for each MJO phase. However, it is also clearest for MJO phase 5, so that this phase serves as an example and is shown again with some visual guidance in Fig. 3 to support the following description: The two strong anomalies over the winter pole described before (indicated with




Figure 3. Repetition of the data shown for MJO phase 5 in Fig. 2, but with the five-zone-response highlighted to guide the reader better through the written description. The two dash-dotted circles indicate the two zones of the polar winter dipole in the temperature response. The two dashed circles mark the corresponding equatorial dipole, which has a reversed sign compared to the polar dipole. The solid circle marks the fifth anomaly zone located in the summer mesosphere, which has the same sign as the upper equatorial anomaly. See Sect. 3.2 for details.

dash-dotted circles in Fig. 3) constitute a vertical dipole with the positive anomaly around 3 hPa and the negative anomaly around about 0.03 hPa. This dipole is part of a remarkable global structure, which comprises three more zones of spatially 240 coherent anomalies, [which are described in the following](#). The departures from the mean state for these three zones are with less than roughly 1 K much weaker than the polar ones, but are still mostly significant, particularly when considering the MCS significance estimation. Two of these weaker anomalies constitute another vertical dipole in similar altitudes as the first one, but located around the equator (indicated with dashed circles in Fig. 3). This equatorial dipole has, however, opposite sign compared to the polar dipole, so that all four zones together look like quadrupole spanning over the complete winter MA. Put 245 in other words, areas of mutually opposing responses between the stratosphere and the mesosphere as well as the equatorial and the polar winter latitudes can be identified. Whereas these four zones are roughly found in the winter hemisphere, the fifth response zone is in the summer polar latitudes (indicated with a solid circle in Fig. 3). It is even higher up between 0.01 hPa

and 0.001 hPa and its extent is much smaller than that of the first four zones. It shows a positive anomaly and looks like a polar summer mesosphere extension of the positive equatorial anomaly around 0.1 hPa. As we will come back to this pattern 250 several times, we will call it temporarily "five-zone-response" for a clearer identification in the following. Its origin will also be discussed below in Sect. 4.1.

A comment on the appearance of the strongest response particularly in the northern winter middle atmosphere is due, since this is a region that is in any case characterized by strong variability. This strong variability is connected to the properties of the polar vortex and particularly the appearance of SSWs (e.g., Baldwin et al., 2021). Whereas the SEA method applied here 255 is designed to average out all variability that is not correlated with the MJO phase evolution, the success of this elimination depends on the length of the analyzed time series and the magnitude of the unrelated variability. Hence one might argue that the analyzed time span of 17 years is not enough for the elimination of the strong uncorrelated polar winter variability, so that the method produces false responses of temperature to the MJO in the polar winter region. While we can indeed not 260 totally exclude that uncorrelated variability still influences the results, we have several indications that the strong polar signal is overall truly correlated with the MJO phases. First, there are indications that the occurrence of SSWs themselves is correlated to the MJO (Garfinkel et al., 2012). Hence, the results presented here must not necessarily be interpreted in a way that they are compromised by the strong SSW variability, but they could also be interpreted as supportive for the idea of an interaction 265 between the MJO and SSWs. Second, as will be discussed But we note that this pattern has already been described in the context of a dynamical feature in the MA, the interhemispheric coupling, as we will discuss in Sect. 3.5, a similar pattern exists also for the southern hemisphere polar winter, which is dynamically much more quiescent. Third, although much weaker, also the polar summer hemisphere temperatures show partly responses to the MJO, see e.g., MJO phase 2 in Fig. 2 between 10 and 0.1. Hence, there seems to be a route for MJO variability into the polar middle atmosphere independent of vortex dynamics. As a forth argument, one could consider repeating the analysis only for boreal winters without SSW. We have quickly checked 270 this approach, which indicated that the signature is still clearly visible. However, the number of days that go into the analysis is quite low, when applying this additional filter so that the interpretation needs particular care and we have excluded this from the current analysis. All in all, we think that at least the baseline of the strong polar winter response is actually truly connected to the MJO. The full strength is then probably result of an amplification by an interaction of MJO and SSWs, although we cannot totally exclude that it is partly also caused by uncorrelated strong variability 4.1.

The five-zone-response appears more or less intense for each of the MJO phases while it has an opposite sign for some of 275 the phases (compare, e.g., the responses for the phases 6 and 2 and 6 in Fig. 2). Of interest could therefore also be the transition of the response from one MJO phase to the next to get an idea of possible temporal systematics. For some of the phases, a gradual systematic change of the response is indeed recognizable, e.g., a downward shift of the polar winter dipole from MJO phase 2 to phase 4. Furthermore, some of the opposite MJO phases show also temperature responses with an opposing sign, as expected for an ideal temperature oscillation during the course of one MJO cycle (e.g., the previously mentioned phases 2 and 280 6). However, the mapping of the responses of opposite phases is not perfectly symmetric. E.g., the phases 5 and 6 appear to be opposed by the phases 2 and 3 instead of 1 and 2 and the responses of the opposite phases 3 and 7 show roughly no sign switch at all. We will elaborate on the aspect of the phase transition in the context of the QBO influence in Sect. 3.3.

285 We note that the northern winter middle atmosphere is, of course, known to be subject to strong variability in any case, which is connected to the properties of the polar vortex and particularly the appearance of SSWs. Hence, one might argue that our observation that the strongest response appears particularly in this region suggests that it is a statistical artifact, i.e. that it is caused by strong uncorrelated fluctuations, which are not sufficiently averaged out. Indeed, it cannot be totally excluded that parts of the strong uncorrelated variability would remain in the averaged data, but we discuss some arguments supporting the existence of a real signal in Sect. 4.2. Particularly, SSWs could be themselves influenced by the MJO and therefore be part of the mechanism producing the temperature signals found here.

290 **3.3 Boreal winter and QBO easterly**

Fig. 4 shows a similar analysis but with data selected for boreal winter and QBO easterly conditions. Overall, the responses look similar to those for boreal winter data in Fig. 2 (note that the color scale has a different range). Particularly the five-zone-response is recognizable for each MJO phase, although the responses for the MJO phases 1 and 8 look somewhat disturbed (which will become part of the interpretation below). The significant part of the summer mesosphere extension appears generally to be higher up at around 0.001 hPa for the boreal winter and QBO easterly situation.

295 ~~Similar to Fig. 2 but for the boreal winter / austral summer and QBO easterly situation. Note that the color scale is different compared to Fig. 2.~~

300 It is obvious that the temperature anomalies are even stronger than for boreal winter filtering alone. The strongest response is again seen in MJO phase 6, here now on the order of ± 10 K. ~~This suggests that the state of the QBO has an important influence on the propagation of the MJO signal into the MA during boreal winter.~~

Also the transition of the response pattern from one MJO phase to the next becomes clearer when the data is additionally filtered for QBO easterly conditions and we will discuss the main aspects in the following. First, it appears that the responses of the MJO phases 2 to 4 look quite similar. And at least for the MJO phases 2 and 3 the

305 It is of interest to look again at the transitions of the response patterns between different MJO phases, which now emerge more clearly: At a simple level, it is seen that roughly two classes of the response pattern exist, which correspond to the one or the other sign of the response of their opposing phases 6 and 7 is switched, as might be expected for an ideal conception. These phases 6 and 7 are also part of three connected phases, which have the same sign of the anomaly pattern: the five-zone-response anomalies. Important is the ordered appearance of these two classes: one class is coherently observed during the first half of the MJO cycle (particularly during the phases 5 to 7). Hence, the response of the phases 2 to 4 seems to be opposed by the phases 5 to 7, which does not totally comply with the ideal case, in which phases 6 to 8 would be expected as opponents of the phases 2 to 4. It is also seen, that during which, e.g., the polar winter dipole response seems to descent from phase stratosphere is colder than normal), whereas as the other class is seen for the second half of the MJO cycle (particularly during the phases 5 to phase 7, which is not clearly seen for phases 2 to 4. It is remarkable that the only gradual variations between phases 2 to 4 on the one side and 5 to 7 on the other side are divided by an abrupt switch of the sign from phase 4 to phase 5, between which the response pattern roughly completely reverses. Hence, at first sight one interpretation of the phase transition scheme could be that there are two slowly varying states of MA temperature, between which an abrupt switch occurs. This suggests that there should be a




Figure 4. Similar to Fig. 2 but for the boreal winter / austral summer and QBO easterly situation. Note that the color scale is different compared to Fig. 2.

switch back around the MJO phases 8 and 1, but the responses of these phases are a bit more complicated to interpret, because a third pronounced and significant anomaly zone appears above the pole and extends the polar dipole. Nevertheless, if one assumes a descent of the negative (blue) polar temperature anomaly from phase 7 to 8 and extends this in a cyclic manner to

320 phase during which, e.g., the polar winter stratosphere is warmer than normal). This coherent behavior, with opposing MJO phases showing roughly temperature responses with also opposing signs, further supports that a real systematic response to the MJO is found here. The responses to the phases 1 and 2, a not-so-abrupt backward transition could be perceived here. However, if the idea of a continuously descending pattern is extended to all phases, a second interpretation appears more probable: It is 8 are somewhat more difficult to interpret since the polar winter anomaly dipole is extended by a third anomaly zone higher up.

325 However, these phases can be systematically integrated in a more detailed recognition of the response transitions between the MJO phases as detailed in the following.

Looking more closely, a gradual change of the response from one MJO phase to the next within the two classes is seen in terms of a descent of the response pattern. This is most easily seen by in the second half of the MJO cycle by starting with MJO phase 5 and following the negative polar winter anomaly with the eyes and starting with MJO phase 5 (blue area in the 330 upper right of the MJO phase 5 panel in Fig. 4): The negative anomaly is there in the highest altitudes and then descends in

the following MJO phases including 8 and 1 until it reaches the lowest altitude phase 8. Such a behavior can also be roughly identified during the first half of the MJO cycle, when starting in MJO phase 2. After phase 2 the transition comes slowly to an end, which is finally reached in MJO phases 3 and 1 with the upper positive polar winter anomaly (red area in the upper right of the MJO phase 1 panel in Fig. 4) and following its vertical position towards phase 4. Afterwards, Note that the summer mesosphere extension of the five-zone-response remains at its altitude during all eight MJO phases.

Moreover, one could also combine the gradual changes within both classes to a roughly continuous descent during the situation is abruptly reset during or after phase 4 and starts from the beginning with the reversed pattern in the new phase 5. In this picture course of a complete MJO cycle. This is most easily seen when starting again in phase 5, but this time with the positive stratospheric winter anomaly (red area on the right at about 5 hPa of the MJO phase 5 panel in Fig. 4). Although somewhat more irregular, this anomaly descends down to about 100 hPa in MJO phase 8. Now, the additionally appearing second polar positive anomaly areas area in the highest altitudes during the phases 8 and 1 in the highest altitudes could be seen as replacement for the vanishing positive anomaly in the lowest altitudes. Note that the summer mesosphere extension of the five-zone-response remains at its altitude during all eight MJO Then the descent could be continued in MJO phase 1 with the upper positive polar winter anomaly as before. Hence, the special responses of the phases 8 and 1 would realize the transition from the one class of the response to the other class.

The only pronounced discontinuity of the transition appears between the MJO phases 4 and 5, where the sign of the response is abruptly switched. Therefore, the response evolution could be interpreted in a way that it starts in phase 5, descends during the following MJO phases including 8 and 1 until the pattern looks reversed in its sign and is then abruptly reset into its original state after MJO phase 4. However, one could also speculate that the MJO phases 4 or 5 should also have a third polar anomaly zone, which is only by chance not very pronounced in this particular dataset. Such a third anomaly zone should this time be negative (opposed to the third zone of the phases 1 and 8), which would then indicate a smooth backward transition and, therefore, close the cycle of the MA temperature response during one MJO cycle. Although we have here no convincing indication for the existence of such a third zone, the speculation is motivated by the following discussion of the austral winter situation (Sect. 3.5).

We have used the boreal winter and QBO easterly situation to describe the two response pattern classes and the gradual transitions, since both are most clearly visible for this case. But at least the two response classes and in outlines also parts of the gradual transition can also be seen for other atmospheric conditions.

We would like to emphasize once more the statistical nature of the analysis. Particularly, we do not say that a particular MJO phase is the causal reason for the MA temperature pattern, which we obtain for that phase. Since propagation speeds have to be considered in a physical mechanism, it seems more likely that a pattern seen for one MJO phase has been triggered by previous MJO phases. This is, however, not problematic for our interpretation, since we only aim at a statistical description of the MA temperature anomaly at given states of the MJO independent of the physical mechanism. This description can still be used to constrain potential physical mechanisms, since a suitable mechanism should be able to reproduce also the statistical results presented here. We expect that particularly the coherent transitions of the responses between the MJO phases will be a helpful constraint for the temporal behavior of potential mechanisms.




Figure 5. Similar to Fig. 4 but for the boreal winter / austral summer and QBO westerly situation. Note that the color scale is different compared to Fig. 4.

3.4 Boreal winter and QBO westerly

We have also checked the MA temperature response for the boreal winter and QBO westerly situation and find that it is much less clear than in the boreal winter and QBO easterly case (Sect. 3.3). This meets the expectations as will be further discussed in Sect. 4.5. Nevertheless, the five-zone-response is also recognizable for some of the MJO phases (Fig. 5). Also other features

370 like the third polar anomaly zone above the polar dipole are visible again (MJO phase 5 in Fig. 5). The strongest response appears in MJO phase 5 and is on the order of $\pm 6\text{ K}$, which is the same order of magnitude as for the boreal winter situation without QBO filtering. Therefore, the MJO influence on MA temperature has not totally vanished and has probably also to be considered under QBO westerly conditions. However, due to the more noisy spatial pattern and the somewhat smaller amount of significant locations, it appears reasonable not to discuss Fig. 5 in detail. We note that the The corresponding structure might

375 become clearer in the future, when longer observational records are available.

Still, we would like to mention that the responses to the individual MJO phases have roughly the opposite sign compared to the respective MJO phase for QBO easterly filtering, although shifts in the altitudes of the response features make a 1-to-1 attribution difficult (compare, e.g., phases 2 and 7 in Fig. 4 and Fig. 5). The Furthermore, the MJO phase 5 appears to have the




Figure 6. Similar to Fig. 2 but for the austral winter / boreal summer situation. Note that the color scale is different compared to Fig. 2.

same special role as phases 1 and 8 for QBO easterly filtering. ~~Note that Hence,~~ the appearance of this feature is also shifted

380

by about half of an MJO cycle.

3.5 Austral winter / boreal summer

The analysis of the data restricted to austral winter (Fig. 6) shows generally that the previously described boreal winter responses can also be detected in the southern hemisphere, although they are weaker and more patchy. The strongest anomalies are again found in the polar winter regions, but have with ± 3 K only half of the magnitude compared to the boreal winter data,

385

which has not been filtered for the QBO (Fig. 2). The polar winter response shows generally again the pattern of a vertical dipole, ~~although individual exceptions like MJO phase 5 exist, in which while individual exceptions with a third zone appears as described before exist also here, e.g., for the MJO phase 5.~~

The equatorial vertical dipole is partly somewhat more difficult to identify but is still clearly visible for many MJO phases, at least for the phases 3, 4, 6, 7, and 8. Also the summer mesosphere extension, this time in the northern hemisphere of course, can be identified in the response of most MJO phases, e.g., for MJO

390

phase 6. However, the summer mesosphere response is more complex for some of the MJO phases: ~~it~~ It has a dipole structure itself (e.g., MJO phases 3 and 7). It is expected from the previous results that the summer mesospheric anomaly has the same sign as the upper part of the equatorial dipole, which is actually seen also for these phases. But in addition, we find another

anomaly zone above, which has the opposite sign, forming a summer mesospheric dipole. It is remarkable that this dipole also reverses the sign for the opposing MJO phases 3 and 7, which ~~suggest more a systematic behavior rather than a random effect~~ indicates a real systematic behavior.

The two significance estimation approaches provide quite different results: the MCIP method shows less and smaller significant areas, which is connected to the generally weaker response. The MCS suggests instead a significant systematic response over all eight phases in almost all places of the meridional plane. Hence, although the responses are weak in terms of the individual temperature anomalies, they appear to be clearly systematically correlated with the eight MJO phases. This might be connected to the fact that this season is dynamically less disturbed so that less interfering disturbances mask the pure MJO signal.

We note that it is not necessarily expected that the austral winter response is recognizable as a counterpart of the boreal winter response and its existence appears noteworthy on its own. This is also because the austral winter hemisphere is usually dynamically much more quiescent than the boreal winter hemisphere. In particular, ~~(major)~~ SSWs are very rare events in the southern hemisphere and only two such warmings have been observed since routine observations started some decades ago, namely in the austral winters of 2002 and 2019 (e.g., Jucker et al., 2021; Allen et al., 2003; von Savigny et al., 2005). While the first one is not included in the MLS period anyway, we have repeated our analysis with the MLS dataset restricted to the period before the end of 2018, so that also the second warming is not included (Fig. S2 in the supplement). It turns out that the pattern of the temperature response to the MJO phases remains basically unchanged when this warming is excluded. In contrast to the expectations, the strongest anomalies are even a bit stronger without 2019. This shows that the appearance of SSWs is not a precondition for the MJO-induced temperature response pattern. A fact that was unclear from analyzing the boreal winter response alone as discussed before (Sect. 3.2). However, due to the different magnitudes of the responses, the results still indicate that the dynamical disturbances of the boreal winter hemisphere are important for an amplification of the MJO signal.

When comparing the boreal and austral winter five-zone-response (Fig. 2 and Fig. 6), it becomes evident that the sign of the response is identical for most individual MJO phases ~~(e. g., for MJO phase 3 the polar negative anomaly is located below the positive anomaly in both Fig. 2 and Fig. 6 and so forth)~~. This is particularly true, when comparing the austral winter response to the boreal winter and QBO easterly situation (Fig. 4 and Fig. 6), which shows a higher level of structure as discussed before. Hence, the phasing of the response with respect to the MJO trigger appears to be generally similar for both hemispheres (whereas it was unclear if it is maybe changed by the QBO westerly phase during boreal winter, as discussed in Sect. 3.4)

Also the phase transition shows similarities with the boreal winter ~~situation and particularly the boreal winter and QBO easterly (and QBO easterly)~~ situation: Again the ~~responses of the phases 2 to 4 are quite similar and are opposed by the responses of the phases 6 to 8, which also show great similarities among each other. The phases 1 and 5 deviate somewhat: the pattern of response 1 is more noisy and not that regular. The response of phase 5 shows again 3 polar winter response zones instead of two, as it was seen for the phases 1 and 8 in the boreal winter case~~ two classes (different signs of the pattern) are seen and can again be attributed to the first and the second half of the MJO cycle, respectively. Overall, one might in outlines ~~see also here a gradual transition~~ also see here the gradual descent of the pattern from one MJO phase to the next ~~in a way that the~~

polar structure slowly descents. This time one could start in MJO phase 2 with tracking the polar positive anomaly. Again, one can start with MJO phase 5 and track the upper negative polar anomaly (blue region) at 0.01 hPa (on the left of the MJO phase 430 5 panel in Fig. 6). It descends down to about 0.1 hPa in phase 4 and further down to about 8 and then further across phase 1 down to 5 hPa in the phases 8 and 1. The abrupt change to the next cycle would then be between the phases 1 and 2 (it was between 4 and 5 for the phase 4).

But in contrast to the boreal winter / QBO easterly (ease) and the phase with the additional polar replacement area would be phase 5 (instead of 1) situation (Fig. 4), the reset of the pattern between the MJO phases 4 and 8 for boreal winter / QBO 435 easterly). This looks like these characteristic phases are shifted by about half an MJO cycle between the 2 cases, however, one has to consider that we tracked this time the positive anomaly instead of the negative one, which is also a change by half a cycle.

This raises for us the question if maybe the phases with the abrupt changes are only missing for some reason a third polar replacement area in the highest altitudes in both cases, boreal winter and austral winter. If these additional areas were there, 440 one would probably identify a completely smooth transition without any abrupt changes. To exemplify this, one could also start at phase 5 in Fig. 6 with following the upper negative anomaly around 0.01 with the eyes. This is a start identical to the description for the boreal winter / QBO easterly situation. One can now identify roughly a descend down to 0.1 in phase 8 and even a further descent from phase 8 does not appear to be so abrupt. Instead, a third anomaly zone (negative) appears here 445 in the responses of the phases 4 and 5, which suggests the interpretation as a smooth backward transition, because the lower vanishing negative anomaly can be seen as being replaced by the higher one (compare the respective discussion of the phases 1 to 4, where it is located between 1 and 10). To close the cycle continuously, one could attribute the lower negative anomaly of MJO phase 5 at 10 to the end of the descent and the upper one to the start of the new cycle. Hence, one could find a closed 450 cycle for the negative anomaly in the austral winter case. This was impossible (8 in Sect. 3.3). Considering that the third anomaly zone (positive) of phase 8, which was already discussed for the boreal winter / QBO easterly case because a second negative polar area was missing in the phases 4 and 5 in this case. Hence, this aspect should gain particular attention in future studies with other or longer datasets to clarify if the temperature response pattern maybe really completes a closed cycle in both hemispheres, also appears at least to some extent for the austral winter, one could suspect that the cycle of the temperature 455 responses during the course of the MJO is actually closed without an abrupt backward transition.

3.6 Austral winter / boreal summer and state of the QBO

455 We have also checked the QBO influence for the austral winter situation and find that the QBO has not such a big impact during this season in contrast to boreal winter situation described before. The results are shown in Fig. 7 for QBO easterly and Fig. 8 for QBO westerly. Although some differences between the MA temperature responses for QBO easterly and westerly are seen, they do not have a clear systematical structure so that these differences could also be random effects due to the different sampling periods. The only noteworthy difference is that the austral winter / QBO easterly case (Fig. 7) does not show a clear 460 descent of the pattern (nevertheless, the two classes/signs of the pattern corresponding to the first and second half of the MJO cycle are still clearly visible), whereas this descent can be seen in outlines in the austral winter / QBO westerly case (Fig. 8).




Figure 7. Similar to Fig. 6 but for the austral winter / boreal summer and QBO easterly situation. Note that the color scale is different compared to Fig. 6.

Together with the previous Sect. 3.5, we can conclude that the structure of the responses and the transitions between the phases for austral winter compare best to the boreal winter situation under QBO easterly conditions, while the austral winter response itself shows no clear QBO influence. Possible reasons for this are discussed in Sect. 4.5.

465 4 Discussion and relation of the results to previous research

4.1 Similarity with interhemispheric coupling

We have carved out a characteristic spatial pattern, the five-zone-response as we called it, of the MA temperature response to the MJO in the meridional plane, which we described in detail in Sect. 3.2 and the following subsections five-zone-response as we temporarily called it. It was a major aim of the present manuscript to communicate this broad overarching picture and 470 particularly to include the complete meridional plane, all MJO phases, different seasons, etc. To stay focused, the study has a purely statistical character and does not determine causalities or physical mechanisms. Another characteristic of the present study is that it is mostly based on purely observational data.




Figure 8. Similar to Fig. 6 but for the austral winter / boreal summer and QBO westerly situation. Note that the color scale is different compared to Fig. 6.

There are a number of previous studies in this area of research with partly overlapping aspects. These studies are typically less broad in the above sense, but cover instead also the physical explanation approaches of the found patterns.

475

This section aims at briefly relating our results to the previous publications to use synergies in both directions: The broader picture put up here may help to integrate and interrelate the more focused studies, whereas the more detailed considerations of the previous studies support the physical plausibility of the present statistical results and provide approaches to explanations.

It appears that this pattern is

4.1 Similarity with interhemispheric coupling (IHC)

480

The five-zone-response looks qualitatively similar to the temperature pattern of the a known dynamical feature of the MA, the interhemispheric coupling (IHC). IHC was first described by Becker et al. (2004) in the context of the anomalous southern hemisphere winter of 2002 and is basically a sequence chain of dynamical disturbances, which also result in temperatures anomalies. The spatial structure of these anomalies starts with a deviation of the planetary wave (PW) drag in the winter stratosphere and then spreads across the winter MA towards the summer mesopause region. These dynamical disturbances cause also temperatures anomalies with a spatial structure that is generally comparable to the one found here. A mechanism

for the IHC was already proposed in the first publications and then detailed by Körnich and Becker (2010). Some details are still subject of the scientific discussion and refinement as, e.g., in Yasui et al. (2021).

After its discovery, the existence of IHC was further supported by satellite observations of, e.g., noctilucent clouds (Karlsson et al., 2007, 2009b) and simulations with the Canadian Middle Atmosphere Model (CMAM, Karlsson et al., 2009a).

(Karlsson et al., 2009a). These simulations can be used for a ~~first quick~~ quantitative comparison of the spatial structure with our results. This comparison reveals that the pattern fits indeed also well in terms of the pressure levels of the strongest deviations (compare, e.g., Fig. 6 for time lag 15 d in Karlsson et al. (2009a) with our Fig. 2 for MJO phase 5, which are the periods of the strongest responses in both studies). We see, however, a much stronger magnitude of the effect with about ± 6 K for northern winter filtering and ± 10 K for the northern winter / QBO ~~east easterly~~ filtering instead of ± 3 K in Karlsson et al. (2009a).

~~Observational temperature anomalies in the context of IHC are reported by Yasui et al. (2021). Although they use the same dataset as in our study, their results may still serve as independent crosscheck because they do not link their results to the MJO at all, so that we will compare the results briefly in the following: Yasui et al. (2021) use a previous version of the MLS data and consider the data only up to 2014, which is, however, not expected to make an important difference. Their focus is also the boreal winter season, but they do not filter for the QBO. They also compute composites of the MLS temperature data, which are, however, not temporally triggered by the MJO, but by the appearance of SSWs (although more precisely, a correlated equatorial temperature anomaly is used as trigger). Their major result in the present context is the middle column of their Fig. 5, where they show the meridional plane temperature anomaly composites of the MLS data for different time lags around their trigger dates. Generally, they also find the Karlsson et al. (2009a) have further pointed out that the IHC temperature pattern can develop with both signs depending on the sign of the initial planetary wave drag disturbance. This fits to the two classes of response, which we have seen for the first and the second part of the MJO cycle, respectively. The first half fits roughly to the "five-zone-response weaker wave drag scenario" pattern, although the summer mesosphere zone is partly above the highest MLS observation altitude. They see a much stronger magnitude of in Karlsson et al. (2009a), whereas the second half fits to the anomalies with up to ± 30 K using the SSW trigger compared to the ± 6 K to ± 10 K we found for the MJO trigger. As the analysis by Yasui et al. (2021) is designed to extract warm polar winter stratosphere situations, they find the "stronger wave drag scenario".~~

As an interim conclusion, we state, that the five-zone-response is probably identical to the already known IHC temperature pattern. Hence, the MA part of the formation mechanism of the five-zone-response only with this sign and the opposite response is not available for comparison to our results. Nevertheless, the similarities of the covered SSW situation with our results of the MJO phases 5 and 6 (extended by MJO phase 7 and maybe 8 if filtering for QBO easterly conditions is applied) are obvious. The use is probably the known IHC mechanism.

One of the new aspects of the present study is that the emergence of the IHC pattern has now been linked statistically to the MJO phase evolution, which leads to the strong expectation that the MJO can indeed physically cause the initial PW drag deviation and with that trigger the IHC in both directions. So far, the IHC has been studied mainly in the context of SSW events. The MJO as a trigger is different from SSWs in such a way that, firstly, the MJO evolves continuously, periodically

and relatively smoothly in contrast to the event-like SSWs. Secondly, the MJO repeats itself on the shorter intraseasonal timescale. Hence, a connected new aspect of this study is that the MA temperature adapts in large parts of the meridional plane continuously to the PW drag changes on the MJO time scale via the IHC mechanism. A new question for the research on the IHC emerges from our observation that the pattern appears to descend during the course of the MJO. To our knowledge, an 525 altitude-time dependence of the IHC temperature pattern has not been explicitly described or explained in the literature so far.

We find some more aspects being broadly consistent with the IHC literature after quick ad hoc checks, which we mention in the following: First, also Yasui et al. (2021) show the SSW-triggered IHC temperature pattern (particularly their Fig. 5), which can be compared to our MJO-triggered results. They use a log-pressure coordinate ~~in Yasui et al. (2021)~~, which complicates the direct quantitative comparison of the vertical structure ~~with our pressure-resolved results a bit~~, but ad hoc conversions 530 ~~attributions~~ of the coordinates indicate that the altitudes of at least the stronger polar dipole fit quite well to ~~each other and that also the descent with time is comparable: The dipole descents between the time lags -8 to 0 and +8 to +16 about 15 to 20, which roughly fits to the descent from MJO phase 5 to 7 in Fig 4 (although we have applied the additional QBO filtering in this case)~~.

Karlsson et al. (2009a) point out that the same IHC pattern can develop with the opposite sign depending on the sign of the 535 initial dynamical disturbance (planetary wave drag; details on the mechanism follow below). We have seen clearest in Fig. 4 for the boreal winter and QBO easterly filtering that we get the opposite sign of the IHC pattern for MJO phases that are half an MJO cycle (i.e. 4 phases) apart. Specifically, the pattern of MJO phase 6 fits to the stronger wave drag scenario after a time lag of about 15 in Karlsson et al. (2009a), whereas MJO phase 2 fits to the weaker wave drag scenario with the same time lag (although phase 3 shows slightly a stronger response). The transition between these opposing structures is further discussed 540 below in Sect. ??.

our results. Note that also a descent of the pattern is implicitly shown in their data, but here related to real physical time instead to MJO phases like in our analysis. A rough conversion of time lags to MJO phases indicates, however, also an overall 545 similarity. We note that the ~~study by~~ ~~Yasui et al. (2021) contains two more details in their temperature anomalies and their schematic of the mechanisms (their Figs. 5 and 15)~~, which might be comparable to mentioned features of our results. We have not checked the quantitative similarity in terms of the vertical levels etc., but still we would like to briefly mention these possibly common features: first, strongest temperature anomalies found by Yasui et al. (2021) for the SSW trigger are with $\pm 30\text{ K}$ much stronger than the magnitudes found here for the MJO trigger. Second, Yasui et al. (2021) have not only considered a vertical dipole as the temperature response above the winter pole, but consider a third anomaly zone above the mesopause, which has the same sign as the stratospheric anomaly. We have ~~also repeatedly~~ seen this in the responses of individual MJO 550 phases, e.g., phase 8 for the boreal winter and QBO easterly situation (Fig. 4). ~~Second~~~~Third~~, Yasui et al. (2021) partly show that the summer mesospheric response of the IHC ~~may itself be being itself~~ a vertical dipole with a positive and a negative anomaly ~~instead of only one zone~~ for certain time lags, ~~which we have found at least~~. We have seen such a feature for the austral winter situation (Fig. 6).

~~Another~~~~Fourth, another~~ subtle but noteworthy detail appears when comparing our seasonally resolved results to Fig. 2 555 of Karlsson et al. (2007) although the quantities shown are not directly comparable: Without going into the details of their

analysis, one important indicator for the IHC is the blue area in the northern polar area in the top panel of their figure. The top panel resembles our boreal winter situation and it is seen that the blue area is directly located above the winter pole. When considering the austral summer situation (their middle panel), the blue area is ~~expected above the southern hemisphere polar latitudes. There is indeed a blue area, but it is~~ not located directly above the pole but instead shifted northwards to about 560 -50° latitude. We see a similar difference when comparing the polar temperature responses of our analysis. Whereas the polar response dipole is directly located above the northern polar region during boreal winter (Fig. 2), it is shifted somewhat more towards the equator for the austral winter situation (Fig. 6).

~~Overall, we conclude from these comparisons that the temperature response pattern found here in relation to the MJO resembles the one that is discussed as the characteristic structure of IHC.~~

565 4.2 The interconnection of IHC, SSWs and the MJO

~~A mechanism for the IHC was already proposed in the first publications and then detailed in Körnich and Beeker (2010). The major points are outlined here only briefly with the notice that details, which are not very important for our study, are still subject of the scientific discussion and refinement as, e.g., in Yasui et al. (2021). An overview is given in Fig. 1 of Körnich and Beeker (2010). The starting point is a deviation of the planetary wave (PW) drag in the winter stratosphere. The theory of IHC is at that point independent of the origin of this deviation. The changed PW drag changes the background dynamical conditions in the winter stratosphere, i.e. the strength of the westerlies and the Brewer-Dobson circulation. This changes also the adiabatic vertical motions and influences thereby the temperatures in the winter stratosphere above the winter pole and close to the equator. This explains the lower two temperature deviations of the quadrupole. The changed stratospheric circulation also changes the filtering of vertically propagating gravity waves, which influences the mesospheric meridional circulation, which has again a direct influence on temperature via adiabatic vertical motions. Altogether, this explains the upper two temperature deviations of the quadrupole. The propagation of the signal to~~ With the present findings, a statistical triangular relation between IHC, SSW and the MJO becomes obvious: First, the IHC pattern is inherently connected to SSWs, since SSWs may produce or amplify the polar winter part of the summer mesosphere, where the fifth temperature anomaly is located, is then explained by IHC temperature pattern. Second, the changed equator to summer pole temperature gradient in 575 the mesosphere and corresponding dynamical anomalies.

~~The results presented in Sect. 3 and discussed here now suggest that the MJO can be a source for the initial PW drag anomaly, which triggers the IHC anomaly pattern. On the one hand this might be expected, since the MJO is known to interact with PWs (e.g., Lau and Waliser, 2012, Chapter 14), which conveys its influence to the extratropics (e.g., Cassou, 2008) and occurrence of SSWs has been linked to the evolution of the stratosphere (e.g., Garfinkel et al., 2012; Yang et al., 2017, 2019). On the other hand it could also be expected that the signal is washed out on its way to the mesosphere considering that the MJO evolves simultaneously further in the troposphere and is probably involved in further interactions with PWs. MJO phases by Garfinkel et al. (2012). As a new aspect, we have now introduced the third side of the triangle, as we have statistically linked the IHC directly to the MJO. Overall, it should be expected that the connections between all the three features should be mutually consistent, which is briefly discussed in the following.~~

590 We note that we are not aware of any explicit description of a potential descent of the IHC pattern in the theory, as we have found it here. Intuitively, one could speculate on two possible initial reasons for such a descent: First, the descent could be an inherent feature of the IHC mechanism, so that an IHC pattern that is once triggered will always descend in the time after. But first of all, we would like to make an aside on SSWs being a potential disturbing factor for our analysis: The SEA method is designed to average out all variability that is not correlated with the MJO phase evolution, but the success of this 595 elimination can be reduced by unrelated variability with comparatively large magnitudes. We find the strongest responses to the MJO in the northern hemisphere winter, i.e., exactly where SSWs produce their strong variability. Hence, the objection might be raised that this strong polar winter variability is actually uncorrelated with the MJO and not sufficiently eliminated, so that the method produces false responses. While we can indeed not totally exclude that uncorrelated variability still influences the results, we have several indications that the strong polar signal is overall truly correlated with the MJO phases. The first one 600 is the fact that the occurrence of SSWs has been found to be itself correlated to the MJO as mentioned before and as further discussed below. Hence, the results presented here could also be interpreted as supportive for this idea instead of being an artifact. Second, it could be a specialty of the MJO-IHC interaction in a way that each MJO phase triggers the IHC pattern at a different altitude. In any case, having the decent of the pattern in mind could foster further research on IHC in terms of theory and observations: A possible decent should be considered in theoretical studies for a further refinement of the IHC mechanism 605 and some observational studies are based on evaluating data at only particular altitude levels, which might miss some of the effect due to the varying altitude levels of the pattern. we have found a similar pattern also for the southern hemisphere polar winter, which is dynamically much more quiescent and periods without SSWs can be analyzed (Sect. 3.5). Third, one could consider repeating the analysis also for boreal winters only without SSW periods. We have quickly checked this approach, which indicated that the signature is still clearly visible. However, the number of days that go into the analysis is quite low 610 when applying this additional filter so that the interpretation needs particular care and has not been included here.

4.3 Connection to SSWs and planetary wave forcing

The IHC pattern is connected to SSWs, since SSWs may produce or amplify the polar winter part of the IHC temperature pattern. Garfinkel et al. (2012) has statistically linked the occurrence of SSWs to the evolution of the MJO phases. We have now statistically linked the IHC to the MJO. Hence, statistical connections between all the three features have been made and 615 the linkages should be mutually consistent, at least in so far as the strong variability allows for rough comparisons. This appears to be the case, as we will briefly discuss in the following.

We have shown mentioned above (Sect. 4.13.3 and others) that the MA temperature responses to MJO phases 5 and 6 (extended by 7 and 8 in case of the QBO easterly filtering) during boreal winter fit to the IHC coupling pattern shown in Yasui et al. (2021). The situation analyzed in Yasui et al. (2021) is basically a SSW situation, so that roughly can be roughly 620 divided into two classes with opposite sign. The responses, which roughly belong to the second half of the MJO cycle (phases 5 to 8) can be attributed to an SSW-like pattern, whereas the first half of the cycle shows a the pattern with reversed sign.

The results of the second half of the MJO cycle show a warm polar winter stratosphere and can therefore be compared to those of Garfinkel et al. (2012), where the MJO-SSW connection is directly explored related to the SSW case. Garfinkel et al.

(2012) find MJO phase 7 as the dominating phase directly preceding SSWs (time lag of 1 to 12 days). This phase is also part of the second half of the MJO cycle, so that the results are at least consistent in this respect. Considering that the analyses differ in their setups (Garfinkel et al. (2012) consider the extended boreal winter, do not distinguish between the two QBO states, use a different MJO index, etc.) and that these analyses are always complicated by the inherent large variability, it might not be reasonable to discuss quantitative similarities and discrepancies in detail. However, an additional semi-quantitative observation can still be made: From phasing is broadly consistent with our results. However, from our analyses, one would have concluded that MJO phases 5 and 6 (instead of phase 7 highlighted by Garfinkel et al. (2012)) show the strongest SSW-like responses (compare Figs. 2 and 4). But one has to keep in mind that, which might look like a discrepancy. However, the pattern in our analysis descents from phase 5 to phase 8 at least for the boreal winter / QBO easterly case, so that we see a continuous evolution of the pattern. Garfinkel et al. (2012) use instead a sharp SSW trigger detection criterion, which is based on the atmospheric state at 10 hPa, which. This is at the lower bound of the polar positive temperature anomaly found here for MJO phases 5 and 6, but more central in the positive anomaly for phase 7. Hence, what we already identify as a SSW-like pattern in phases 5 and 6 might, with regard to the exact altitudes, be a precursor of the real SSW-like pattern considered by Garfinkel et al. (2012), which is then in terms of altitudes only reached in phase 7. This would also be roughly consistent with the result of Garfinkel et al. (2012) that MJO phase 6 is a precursor for SSWs with a longer time lag of 13 to 24 days. As said before, these comparisons should only be treated as We note that the comparison is only a rough qualitative cross checks and not as reliable quantitative results check since the analyses differ in their setups and are always complicated by the inherent large variability.

We note as a byproduct of the discussion that the descent of the IHC pattern, which has been shown in the present analysis, might be relevant also for other comparisons and considerations in this area of research similar to the example above.

Overall, we consider the general consistency with Garfinkel et al. (2012) as important support for the hypothesis that the statistical relationship between the MJO and the IHC brought up in this study is a real effect and not a statistical artifact.

4.3 MJO-IHC linkage: potential coupling by planetary wave forcing

The joint reason of SSW and IHC occurrences is known to be a deviation in the planetary wave PW forcing of the winter stratosphere. Whereas the forcing is always stronger than normal for SSW situations, the The sign of IHC coupling pattern has also been found for the opposite case. Karlsson et al. (2009a) call those two IHC modes depends on the sign of the PW drag deviation, called strong or weak planetary wave events forcing, respectively. As expected, the IHC pattern for the strong events resembles that of SSWs. The findings of the present study put now the MJO in the context of these deviations: The MJO acts probably as one source of such planetary wave disturbances. Moreover, it can both

Our findings now relate the MJO to such PW drag deviations: Although not proven by our statistical approach, the MJO appears to act as a source of the initial PW disturbances, which can weaken and strengthen the planetary wave activity and the previous discussion suggests that relevant PW activity in the winter stratosphere. More precisely, the first half of the MJO cycle (phases 1 to 4) tends to weaken the planetary wave activity, whereas the second half (phases 5 to 8) tends to strengthen it.

660 The It is therefore of interest to consider the study by Wang et al. (2018a), which explores the effect of MJO phase occurrences on wave activity in the northern hemisphere winter stratosphere is explored in more detail by Wang et al. (2018a). They, i.e. in the region where the IHC mechanism needs the initial PW drag deviation. Wang et al. (2018a) find that a higher occurrence frequency of MJO phase 4 is in line with weaker wave activity and a stronger polar vortex in the stratosphere, i.e. the opposite of a SSW situation. This is roughly consistent with our findings in the sense that MJO phase 4 belongs to the first half of the MJO cycle, for which we also found the opposite of the SSW situation. The reason find the weak PW drag case. Furthermore, Wang et al. (2018a) find higher wave activity for MJO phase 7, which is also roughly consistent with our 665 findings, since phase 7 belongs to the second half of the MJO cycle.

The reason for the PW drag modulation is according to Wang et al. (2018a) not that the MJO induced that MJO wave activity is generally weaker, but that MJO wave activity is in antiphase with the climatologically apparent waves in the respective region for MJO phase 4. Furthermore, Wang et al. (2018a) state that the opposite is true 4 and in phase for MJO phase 7, so that a higher occurrence frequency of phase 7 results in higher wave activity and a SSW-like situation. This is also roughly 670 consistent with our findings, since phase 7 belongs to the second half of the MJO cycle. However, 7. Hence, a mechanism for the relationship between the tropospheric MJO and the MA IHC pattern found in our study should probably consider such PW wave interferences as a link.

675 However, our results also raise new questions in this context: Wang et al. (2018a) state that all other MJO phases should not have a strong influence on the extratropical stratosphere, because there is no clear in-phase or antiphase relationship of the MJO related waves and the climatological waves. We find instead a response throughout the middle atmosphere for roughly all MJO phases. This would only be consistent if the clear MA responses found here for all MJO phases were could be consistent as long as the MA responses to all other MJO phases are considered as time-lagged results of the forcings of only the MJO 680 phases 4 and 7. In this case it would, however, be This is indeed not excluded by our results, instead the nature of the responses and the gradual transitions between them is an open question. However, it is then dubious that the main forcing occurs towards the end of those consecutive MJO phases, which lead to the same MA response according to our results show the appropriate sign of the response (phases 1 to 4 and 5 to 8, respectively). Hence, the explanation of Wang et al. (2018a) for all phases except 4 and 7 seems not to be in line with our findings and it appears. This could mean that the mechanism, which explains 685 our results, is more complex. But it appears also to be appropriate to question if Wang et al. (2018a) really describe the full mechanism. We note also here that the statistical approaches are not directly comparable. Whereas we make a composite analysis including data of all days with a certain MJO phase, Wang et al. (2018a) analyze changes that are correlated with changes in the occurrence frequency per winter of individual MJO phases and, hence, effects that are more relevant to the interannual time scale relationship of MJO and wave activity.

690 4.4 Timing of the MA response Comparison with respect to previous studies covering the phase transition relationship of the MJO and MA temperature

4.4.1 General remarks

We have described before that there are characteristic responses of the MA temperature for each MJO phase with systematic transitions between them. The coarsest classification of the eight individual responses would be the discrimination of the two cases "weak wave driving" and the SSW-like "strong wave driving". The two classes consist roughly of the subsequent MJO phases 1 to 4 (weak) and 5 to 8 (strong), particularly in the boreal winter and QBO easterly case (Sect. 3.3). Wang et al. (2018a) provide a possible explanation for such two classes, although it appears to us that it does not completely explain our results, as discussed in Sect. 4.2.

Particularly, the responses also evolve gradually within these two classes, which manifests itself in a descent of the spatial structure from phase to phase. This results in an almost continuous cyclic development of the MA temperature responses over the eight MJO phases (details and limitations of this generalized statement are extensively discussed in Sect. 3, particularly in the Sects. 3.3 and 3.5).

Indeed, it appears plausible that a periodically evolving trigger, like the MJO, should also produce roughly a systematically and periodically evolving response as we have found here. However, in terms of mechanisms it appears quite remarkable to us that the response evolves so systematically. This is not only because the MJO is only quasi-periodic with strong variability and hemispheric and seasonal asymmetries. Moreover, the MJO propagates itself zonally and changes its properties gradually while evolving in time. At each point in time, the current state of the MJO could in principle interact with other relevant parts of the atmosphere, causing disturbances, which travel horizontally and vertically into the MA, where they could in principle interfere with arbitrary other disturbances of the preceding MJO states. This could lead to a great variety of superimpositions of MJO responses in the MA, which could also, e.g., totally cancel out each other or look much noisier than the results found here. Instead, the overall net effect of the MJO seems to be more like a simpler periodic trigger that initiates the IHC pattern with the one or the other sign on a coarse scale and that causes, seen on a finer scale, gradual altitude shifts from phase to phase.

The clarification of the exact mechanism is not within the scope of this paper. Contrarily, in the light of many-faceted properties of the MJO-MA temperature interaction brought up in Sect. 3, we think that coming to a full explanation needs future research dedicated to this topic. There are some studies already dealing with the MJO-MA temperature relationship for particular MJO phases, which can be compared to our results.

However, independent of an explicit formulation of the mechanism, the statistical relationship between the MJO phases and the MA temperature, which is described in this study, might already serve as a benchmark for complex atmospheric models. Particularly those models, which are used for intraseasonal weather forecasting depend on having a good representation of the MJO and its teleconnection patterns in which the stratosphere is involved. We have carved out a characteristic and complex response pattern for the major parameter temperature, which connects the MJO with the complete MA in all latitudes and heights and we think that this pattern and its evolution with the MJO phases can be used as a challenge for atmospheric models, which implicitly concerns many aspects of atmospheric couplings. In order to support these kinds of evaluations, we briefly interrelate our results with previous studies in the following subsection 4.4.1. While the present study does not go as far into possible explanations of the relationship as some of these studies, it adds the overarching picture of the interhemispheric

725 coupling to this topic, which could help to combine the previous findings. Simultaneously, the previous results can further support the statistical findings of the present analysis.

4.4.1 Relation to previous studies

730 There are a few studies, which go beyond the specific case of the MJO-SSW connection discussed before (Sect. 4.2) and give more general information on the MJO-MA temperature relationship for particular MJO phases, which can be compared to our results. However, We note, however, that strict comparisons are difficult, not only because the of the partly high variability of MA temperature, different analysis setups and datasets differ different datasets, but also because most many studies analyze the state of the MA only after selected MJO phases while considering relatively long time lags. Hence, they study the temporal evolution after a certain state of the MJO. This is certainly useful to explore causalities of the propagation mechanisms of the disturbances and to examine lead times of potential predictions based on the MJO states. But it also bears ambiguities because 735 the MJO evolves itself during these as a function of relatively long time lags. E.g., a state of the MA that is found 30 days after MJO phase one, could also be attributed to MJO phase 4 or 5 with almost no time lag assuming one MJO phase lasts for about 7 days. We have intentionally decided to study the statistical relationships after only selected MJO phases. Our analysis works without time lags, but with respect to and considers instead all eight MJO phases as alternative an approach, although we might extend these analyses to a treatment of time lags in the future for comparison purposes. For now, comparisons have 740 to ad hoc to uncover the gradual changes of the responses between the individual MJO phases. This means that comparisons have ad hoc to assign the MJO phase evolution to specific time lags, which is of course difficult due to the high variability of the MJO. As an example, also Garfinkel et al. (2012) mention such an assignment, which considers a periodicity of 30 to 60 days for a complete MJO cycle and thus must remain a rough estimate. by using rough mean MJO periods. Hence, the reader should have in mind that the comparisons have a rough and partly qualitative character.

745 The influence of the MJO on mesospheric temperatures during boreal winter has recently been analyzed by Sun et al. (2021). They partly also use the MLS dataset employed here, but mostly rely on modeled data of the model SD-WACCM. A period of 100 days around SSWs is excluded, which further limits the comparability to our results. Sun et al. (2021) concentrate mainly on the temporal evolution after MJO phase 4 and on the northern hemisphere. As one main result they report a cooling of the northern hemisphere mesosphere 35 days after MJO phase 4. In qualitative agreement, we also see the mesospheric cooling 750 after MJO phase 4, namely in the phases 5 to 7 (Fig. 2). However, converted to time lags this appears to be earlier than the time lag of 35 days. Sun et al. (2021) also show the meridional plane temperature response of the MA for different time lags after MJO phase 4 (their Fig. 2). In consistency with our results, also the polar vertical dipole is seen with the change of the sign at about 1. Furthermore, they also show the temporal switch of the dipole's sign, here between time lag 25 and 30. They do not show much of the equatorial area, but a reversed equatorial dipole can be perceived to some extent. The order of 755 magnitude of the temperature anomaly MJO-related temperature anomaly shown in Sun et al. (2021) is with ± 5 K comparable to our analysis, as well as the fact that the strongest anomalies are observed, when the stratosphere shows the warm anomaly. The agreement of the orders of magnitude is particularly remarkable, since the periods of strongest variability are missing due to the exclusion of SSWs. Overall, we conclude that apart from the fact that differences are expected due to the different

approaches, the general response pattern is roughly consistent with our findings, while the timing is hard to compare, so that
760 consistency in this respect cannot be evaluated.

A study broadly comparable to Sun et al. (2021), but focused on the stratosphere and based on reanalysis data is presented
765 by Yang et al. (2019). They analyze the northern hemisphere stratospheric temperature response to partly all MJO phases, but
highlight a stratospheric warming after MJO phase 4 with a time lag of 30. This can be regarded as consistent with our results in
the same sense as discussed before in the context of Sun et al. (2021) for the mesosphere. Furthermore, the in Sun et al. (2021)
770 . Also the order of magnitude of the temperature response is in their analyses reported Yang et al. (2019) is comparable to our
results of the boreal winter situation without QBO filtering (Fig. 2), i.e., in the situation for which the analysis setups are closest
to each other. Their Fig. 1(a) shows the evolution of the response with the MJO phases at 10

The northern winter polar vertical dipole of the five-zone-response is, e.g., confirmed by Sun et al. (2021, Fig. 2). They also
770 show the temporal switch of the dipole's sign, here between time lag 25 for a polar cap average. Comparing their results for a d
and 30 d after MJO phase 4. A consistent sign change of the temperature anomaly is also found in Yang et al. (2019, Fig. 1(a))
for a time lag of 0 d with our polar results reveals a general consistency in the sense: It is seen that the stratospheric response is
negative for MJO phases 2 and 3 and positive for the phases 5 and 6, as in our study. Altitude-resolved results are also shown
in their Fig. 2 (left column) and should be visually compared by the readers themselves.

Yang et al. (2017) present a similar analysis based on reanalysis and modeled data, but with respect to the austral
775 winters and, thus, roughly comparable to our Fig. 6. In an overview of the southern polar cap temperature response to all
MJO phases at 10 timing, Sun et al. (2021) report a cooling of the northern hemisphere mesosphere 35 days after MJO phase
4. Yang et al. (2019) report a corresponding stratospheric warming after MJO phase 4 with a time lag of 30 (their Fig. 2 (A)
and (B)) they d. Although both might still fit to the appearance of this pattern in the second half of the MJO cycle in our data,
we would have expected it earlier in terms of time lags after phase 4.

The results for the austral winter can be compared to Yang et al. (2017, Fig. 2 (A) and (B)). They find cold anomalies at
780 10 hPa and a time lag of 0 d roughly for the MJO phases 1 to 3 and warm anomalies for the phases 5 to 7. This is roughly
consistent with our results; however, we see the more pronounced corresponding responses shifted by one MJO phase, namely
We see the corresponding responses slightly shifted to the MJO phases 2 to 4 for the cold anomaly and 6 to 8 for the warm
anomaly. Yang et al. (2017) discuss the responses to the MJO phases 1 and 5 in more detail and present data for, respectively,
785 and also somewhat different in altitude, but a rough consistency can be stated. A comparison of the complete meridional plane
for these phases. The temperature response for MJO phase 5(their Fig. 4) appears roughly consistent, which is included in
Yang et al. (2017, Fig. 4), also reveals a general consistency with our results. Particularly, the results for a moderate the time
lag of 10 d after phase 5 fit quite compare well to our results of following MJO phase 6. MJO phase 6 response. Also, our
magnitude of the anomalies, which is smaller than during boreal winter in the northern hemisphere, fits roughly to the results
790 of Yang et al. (2017). Note that this plot in Yang et al. (2017) also contains a summer mesosphere response in the northern
hemisphere, which also fits to our results. Our results for MJO phase 1 are quite noisy and hard to compare.

A We note that although the study by Yoo et al. (2012) is actually focused on the tropospheric connection between the
MJO and the Arctic, however, the upper vertical range of their analysis slightly overlaps with the lower edge of our analysis

around 200, so that a rough comparison is possible. Still, one should have in mind that the datasets could be affected by boundary effects etc. in these locations especially because these pressure levels are not in the focus of the respective analyses. Yoo et al. (2012) use reanalysis data and concentrate on the MJO phases 1 and 5 during boreal winter. Their Fig. 6 shows meridional plane temperature responses for different time lags, of which the upper parts can an inner-tropospheric MJO teleconnection, the lowest altitude levels of our analysis can still be compared to the lower parts of our Fig. 2. Their response for MJO phase 1 with time lag 0 shows a positive anomaly over the Arctic, which is consistent with the Arctic response at lower vertical levels in our analysis. A time lag of 8 might roughly be attributed to MJO phase 2 in our analysis and also here positive anomalies in this region are found in both studies. Also their response for phase 5 with time lag 0 shows a positive response in higher levels of the Arctic region, which is also roughly consistent with our analysis. Interestingly, the sign of the response changes in their study when looking at time lag 8 their highest levels. Without going into details, the tropospheric northern meridional plane response in Yoo et al. (2012, Fig. 6) at different time lags after MJO phase 1 and 5, which might be attributed roughly to MJO phase 6 in our study. Consistently, we also find a change to a negative anomaly in this region with the transition from phase 5 to phase 6. Also the order of magnitude of the response is with about ± 1 K consistent between the studies. As stated before, we are cautious with the interpretation of this small overlapping region. However, if the similarities were accepted as real, this could mean that the resembles roughly the evolution with the MJO phases found here. Therefore, the responses at the lower pressure altitude levels in our study could be are probably more a result of the tropospheric processes analyzed by Yoo et al. (2012) rather than a result of stratospheric processes discussed here so that they do not have to be explained within the dynamical framework of IHC. We also note that Furthermore, Yoo et al. (2012) report on stronger poleward Rossby wave propagation in connection with MJO phase 5 and weaker poleward wave propagation in connection with phase 1. They do not discuss a potential vertical propagation into the stratosphere, so that their findings are not necessarily applicable to our case. Nevertheless, Although this strictly applies only for the troposphere, we note that this attribution is roughly consistent with our finding that the MA response to the first half of the MJO cycle resembles the weak wave driving IHC pattern and the second half the discussion of the weak and strong wave driving IHC pattern (Sect. 4.1) and should therefore be considered when working on the questions raised in Sect. 4.3.

4.5 Influence of the QBO on the MA temperature response to the MJO

As shown above, filtering the data not only for the boreal winter but additionally for QBO easterly conditions, carves out the IHC pattern more clearly in terms of stronger anomalies and smoother transitions of the pattern between the MJO phases (For the influence of the QBO on the MJO-MA connection (e.g., Sect. 3.3). For this influence on the MJO-MA connection, at least two possible types of interference with the QBO have to be considered: The influence of the QBO on the MJO itself as well as the inner-stratospheric influence of the QBO on the polar stratospheric dynamics.

4.5.1 Influence of the QBO on the MJO

The influence of the QBO on the MJO is currently an active field of research. Yoo and Son (2016) found that the MJO amplitude indeed varies with the QBO phase during boreal winter in a way that the MJO is stronger during the QBO easterly

phase and weaker during the QBO westerly phase. Afterwards Zhang and Zhang (2018) argued that the MJO is not necessarily stronger during the QBO easterly phase, but that the number of active MJO days (days during which a strength threshold is exceeded) is increased. Wang and Wang (2021) state that both kinds of interpretations are actually connected and should not be treated independently. In any case, the fact that there is something like ~~and still debated~~ (e.g., Yoo and Son, 2016; Zhang and Zhang, 2018; ~~Yoo and Son, 2016; Son et al., 2017~~) ~~that the MJO effect on the MA temperature is most pronounced under these conditions~~. Note that, ~~combined with our results~~, this would be a net effect of the tropical lower stratosphere on the complete MA conveyed by the troposphere, where the MJO is active.

Regarding the seasonal differences, the QBO influence on the MJO has mainly been reported for boreal winter. More precisely, Yoo and Son (2016) and Son et al. (2017) state that they find no significant relationship of the QBO and the MJO for other seasons. However (e.g., Yoo and Son, 2016; Son et al., 2017). However, using two more sophisticated analysis approaches (QBO determination and BSISO index), Densmore et al. (2019) claim that there is also an QBO-MJO relationship during boreal summer, which is reversed compared to boreal winter. While the analysis by Densmore et al. (2019) is limited to the MJO phases 4 and 5, it also applies two more sophisticated analysis approaches, which might help to identify a potentially weaker signal also during boreal summer. First, the state of the QBO is determined more sophisticatedly and, second, an BSISO index is used to characterize the state of the MJO during boreal summer in addition the common MJO indices.

Our examination of the QBO influence on the MJO-MA connection during boreal summer (Sect. 3.6 and Figs. 7 and 8) showed no clear indication for a QBO influence during boreal summer. This can be seen as consistent with both previous notions: We cannot identify a strong impact of the QBO during boreal summer ~~like Yoo and Son (2016) and Son et al. (2017)~~, but we also ~~do did~~ not use the more sophisticated QBO identification approach and the specialized BSISO index as in Densmore et al. (2019), which would support the identification of weaker signals ~~approaches, which might be necessary to detect the signal during boreal summer~~.

4.5.2 Influence of the QBO on the polar stratosphere

The second possible type of interference is the inner-stratospheric influence of the equatorial QBO on the polar winter stratosphere (e.g., Holton and Tan, 1980), which results in the polar winter stratosphere tending to be colder during the QBO westerly phase and warmer during the QBO easterly phase (e.g., Labitzke and van Loon, 1992). This observation is connected to a known influence of the QBO on the occurrence of SSWs during boreal winter (e.g., Camp and Tung, 2007). Hence, ~~our study remains inconclusive with respect to the QBO influence during boreal summer, but a more sophisticated future analysis might be of interest, which could, as a side note, also benefit from an extension of the analysis by Densmore et al. (2019) to the other MJO phases~~ both the QBO and the MJO have an influence on the occurrence on SSWs, which have a large influence on the polar MA temperature. Therefore, one reason for the finding that the QBO apparently changes the MJO's influence on MA temperature could be that both influence the occurrence of the SSWs simultaneously with possible interactions or interferences of the effects.

860 We note that this QBO effect is so far implicitly restricted to the polar winter stratosphere. However, this region is particularly important as it acts as the starting point for the emergence of the IHC pattern (Sect. 4.1). Consistently, it has already been shown that the QBO influence on this region also influences other regions of the MA according to the IHC mechanism (Espy et al., 2011). Moreover, Murphy et al. (2012) show that some details of the IHC mechanism are themselves modified by the QBO phase. Hence, it is plausible that the temperature response to the MJO, which is conveyed by the IHC mechanism, 865 is subject to this QBO influence in the complete MA.

Overall, there

4.5.3 Concluding remarks on the QBO influence

Overall, we cannot discriminate the relative importance of the two possible types of the QBO influence on the MA temperature response to the MJO. Whereas a consistency of major aspects can be seen for the QBO influence on the MJO itself (Sect. 4.5.1), 870 we cannot draw any reliable conclusions on the importance of the inner-stratospheric influence (Sect. 4.5.2). This might be possible in the future, when longer datasets allow for a more detailed filtering with respect to other environmental parameters, particularly the exclusion of SSWs.

There is obviously a seasonal asymmetry of the QBO influence on the MA temperature response to the MJO (Sects. 3.3 and 3.6): The QBO easterly phase is needed to produce the clearest response pattern during boreal winter, whereas a comparable 875 (although weaker and more noisy) structure is produced during boreal summer in the southern hemisphere regardless of the QBO phase. One possible explanation for this apparent asymmetry could be the generally higher variability of the northern hemisphere boreal winter situation. This might demand for a particularly strong MJO signal to dominate the intraseasonal variability in the northern hemisphere. And the influence of the QBO easterly phase on the MJO during boreal winter could be a major factor to facilitate this strong MJO signal during this season. In contrast, a weaker MJO might already be able 880 to dominate the signal in the more quiescent southern hemisphere, so that a QBO support is, regardless of its still debated existence, not needed to produce the austral winter signal. Another explanation (which strictly speaking belongs to the next Sect. 4.5.2), could be the mentioned modification of the occurrence rates of SSWs by the QBO, which is roughly only relevant during boreal winter, because SSWs only rarely happen at all in the southern hemisphere.

4.5.4 Influence of the QBO on the polar stratosphere

885 The second possible type of interference is the inner-stratospheric influence of the equatorial QBO on the polar winter stratosphere. This connection was first described by Holton and Tan (1980) and implies in the current context of temperature that the polar winter stratosphere tends to be colder during QBO westerly phase and warmer during the QBO easterly phase (e.g., Labitzke and van Loon, 1992). We note that this observation is connected to a known influence of the QBO on the 890 occurrence of SSWs during boreal winter (e.g., Camp and Tung, 2007), so that this aspect is not independent from the discussion in Sect. 4.2. This could mean that, e.g., the strength with which the MJO can influence the polar stratospheric temperature by modulating SSWs is itself affected or limited by the QBO's influence on SSWs.

895 In any case, the QBO is overall known to have a similar effect on the polar stratosphere as was found for the MJO in this study, but on the biennial rather than on the intraseasonal time scale. One could expect intuitively that these two effects appear superimposed in the current analysis, so that particular combinations of MJO phases and QBO states interfere constructively or destructively and therefore lead to extreme values of the stratospheric temperature anomaly. As described before (Sect. 3.4), the QBO westerly response for boreal winter in MA temperature is not as systematic and less significant as the QBO easterly response for the same season, so that it is almost impossible to deduce a deeper systematic behavior and we cannot examine this intuitive expectation in detail. However, we found that the polar winter stratospheric responses for some MJO phases are actually reversed for the QBO easterly and westerly situation,

900 4.6 Limitations of the presented analysis

905 To obtain a complete picture, it will be necessary to study the MJO influence on the MA temperature also with respect to additional environmental conditions, at least the state of the El Niño–Southern Oscillation (ENSO) and the solar 11-year cycle. The length of the analyzed datasets currently limits the number of filters, which can be simultaneously applied, so that we have focused on those influences, which we expected to be most important based on previous research. However, particularly the discrimination of the QBO states based on comparatively only few years of data might be affected by, e.g., the cold polar stratospheric responses for the QBO westerly phase appear partly during the same MJO phases (e.g., 6 and 7) as the warm responses for the QBO easterly phase and not during the opposite MJO phases (Sect. 3.4 and Figs. 4 and 5). This does not seem to be consistent with this simple expectation and suggests a more complex connection.

910 We note that together with the QBO influence, also the solar underlying state of the 11-year cycle has been shown to be important for the polar winter stratospheric temperature (e.g., Labitzke and van Loon, 1992). This is worth mentioning because of two reasons: First, the solar 11-year cycle is another source of variability, which is not considered in this study and which could therefore disturb the pure signal of MJO and QBO in MA temperature. The influence of the solar variability should therefore be analyzed in a future study. Such a study would also be of interest, because an influence of solar variability directly on the MJO has also been indicated (Hood, 2017; Hoffmann and von Savigny, 2019), so that there are also at least 2 types of interference possible. Second, also the influences of QBO and solar 11-year cycle on the polar stratospheric temperature turned out to be not additive (e.g., Labitzke and van Loon, 1992; Camp and Tung, 2007), similar to what our analysis indicates here for the superimposed QBO and MJO influences. As written before, also a precise discrimination of boreal winters with and without SSWs is of interest but also demands for a longer dataset.

920 Overall, we cannot discriminate the relative importance of the two possible types of QBO interference with the MJO influence on MA temperature introduced here and possible further influences like the solar cycle. Whereas a consistency of major aspects can be seen for the QBO influence on the MJO itself (Sect. 4.5.1), we cannot draw any reliable conclusions on the importance of the inner stratospheric influence discussed in this section. This might be possible in the future, when longer datasets allow for a more detailed filtering with respect to other environmental parameters like the solar activity. Furthermore, we emphasize again the statistical nature of our analysis, which cannot examine the causality and possible physical mechanisms. The present publication aims at making the revealed statistical indications for a connection of the MJO

and the IHC already available to the community in order to foster research on the underlying mechanisms. However, to prevent the publication of possible statistical artifacts or spurious correlations, we have embedded our results into previous research, which shows the overall plausibility of a real effect, for which the physical reason can be established in the future.

The discussion of the inner-stratospheric QBO influence on the MA temperature response to the MJO in this section was 930 so far restricted to the polar winter stratospheric temperature. This region acts as the starting point for the emergence of the IHC pattern (Sect. 4.1) and it has indeed been shown that the QBO influence on this region also influences other regions of the MA according to the IHC mechanism (Espy et al., 2011). Moreover, Murphy et al. (2012) show that some details of IHC mechanism themselves are modified by the QBO phase. Hence, also the temperature response to We note that the statistical approach does also not determine the direction of a physical influence. From the previous literature, it appears likely that an 935 influence in the MJO in the complete MA, which is conveyed by the IHC mechanism, as we have indicated before, should be subject to this QBO influence. Therefore, the complete MA can and should be considered to further isolate the relevant effects in future studies direction from the tropospheric MJO to the MA temperature via an alteration of the wave driving exists. However, possible influences in the opposite direction or feedback loops are not excluded by our study.

5 Summary and conclusions

940 There is rising interest in MJO influences on the middle atmosphere: First, because the stratosphere is one route for MJO-related teleconnections, which is, e.g., of practical interest for weather forecasting efforts on the intraseasonal timescale. Second, it is of interest for MA research to identify all sources of variability in the MA, e.g., in order to cleanly isolate anthropogenic signals. And third it is naturally of interest for basic research on troposphere-stratosphere coupling. The MA temperature is 945 obviously a key parameter in this context. Contrarily, only few studies deal explicitly with the influence of the MJO on MA temperature. The available studies are typically mostly based on modeled or reanalyzed data and are limited to individual MJO phases or particular geographical regions.

We aimed at contributing to this field by statistically analyzing We have statistically analyzed the connection of the MJO and the MA temperature based purely on observations . We included, moreover, of MA temperature. We have included all MJO 950 phases in the analysis , in order to reveal also the transitions of the MA response when the MJO progresses from one phase to the next. As statistical approach, we have applied a superposed epoch analysis to the 17-year global temperature dataset of the MLS satellite instrument. For the MJO characterization, we have used the OMI MJO index. We have distinguished different environmental conditions, particularly different seasons and states of the QBO. For the sake of brevity, we have focused this paper on zonally averaged data.

We have indeed found that the MA temperature is influenced in a statistical sense by the state of the MJO in large areas of 955 the MA and under roughly all considered atmospheric conditions. Still, the strength of the influence varies considerably with both the atmospheric conditions and the region of the MA.

Aside from more occasional weaker spatial response areas Most notably, a pronounced characteristic response pattern, which we called here temporarily "five-zone-response", manifested itself under many atmospheric conditions. The first two zones

(compare, Fig. 3) are two anomaly areas with opposing sign in the polar winter stratosphere and mesosphere, together constituting a dipole response above the winter pole. These zones show mostly the strongest anomalies among all five zones. A second dipole, comprising the third and ~~forth~~ fourth zone, is found in the equatorial stratosphere and mesosphere. This dipole has a reversed sign compared to the polar winter dipole. The fifth zone is found above the summer pole in the mesopause region, so that the five-zone-response spans almost the complete meridional plane. The anomaly of the fifth zone has the same sign as the equatorial mesospheric zone. The overall sign of the complete five-zone-response is different for different MJO phases. It has been checked with two Monte Carlo approaches that the anomalies of at least these five zones are broadly significant for most analyzed cases.

A pattern like the five-zone-response ~~in~~ is known in MA research from ~~the interhemispheric coupling mechanism~~ a feature called interhemispheric coupling (IHC). This IHC mechanism basically describes, how an initial disturbance in planetary wave drag in the winter stratosphere propagates due to a chain of different dynamical effects through the MA ~~until it reaches the summer mesosphere. It~~ an thereby causes a similar pattern of temperature disturbances as we have seen here for the MJO influence. More detailed comparisons of the particular locations of the disturbances with the original IHC publications have confirmed the similarity. Moreover, the IHC mechanism works for weak and strong planetary wave activity, causing the one or the other sign of the temperature anomaly pattern as we found it here for different MJO phases.

Hence, one major outcome of the present ~~study is the finding observational and statistical analysis is the idea~~ that the MJO can lead to the initial planetary wave drag disturbance of the IHC mechanism. The MJO ~~can~~ could therefore systematically trigger the IHC mechanism ~~and it can do this~~ with the one or the other sign, i.e. ~~with a lead to the initial~~ strengthening or weakening of the planetary wave driving in the winter stratosphere. The present study therefore statistically connects an atmospheric feature mostly known in tropospheric research, the MJO, with one mostly known in MA research, the IHC. The MJO as a further potential trigger of the IHC brings new aspects into the research on MA dynamics, since its time scale and periodicity characteristics are quite different from SSWs, which have so far been considered as the starting point of IHC studies.

The analysis, applied individually to boreal or austral winter data, has revealed that the five-zone-response appears in both seasons with the response being roughly mirrored at the equator. The boreal winter response is, however, much stronger. Furthermore, the response during the boreal winter season shows also a strong dependence on the QBO phase, which we have not convincingly seen for the austral winter. Overall, the strongest anomalies and the clearest spatial pattern appear for boreal winter combined with the QBO easterly phase. In this case, we have found temperature anomalies in the range of $\pm 10\text{K}$ with the strongest anomalies being located over the winter pole. For ~~boreal winter and the QBO westerly phase the pattern is much less clear but the anomalies are still in the range of $\pm 6\text{K}$. For~~ austral winter the anomalies are with $\pm 3\text{K}$ much smaller, but nevertheless, the spatial pattern can easily be identified for most MJO phases. ~~When the austral winter data is separated according to the QBO phases, the anomalies are with $\pm 4\text{K}$ slightly higher, but the spatial pattern is by far not as sensitive to the QBO as during boreal winter.~~

~~One possible reason for the strong QBO influence during boreal winter is a potential alteration of the MJO strength by the QBO during this season. This QBO-MJO connection has been proposed in the literature in the last few years. This kind of~~

connection would indeed be consistent with our findings such that the easterly phase of the QBO strengthens the MJO during boreal winter, as we have discussed above in more detail. This would mean that the QBO influences the tropospheric MJO, which afterwards influences the MA temperature with a corresponding strength.

A dynamical feature in the MA closely related to the IHC are SSWs, which are mostly observed in the northern winter hemisphere apart from only two exceptions that occurred in the southern hemisphere so far. They essentially produce a strong positive temperature anomaly in the polar winter stratosphere, which is then part of the IHC temperature anomaly pattern.

During the last decade, an influence of the MJO on the occurrence of SSWs has been reported in the literature. Since we have in this study now linked the MJO also to IHC, we have checked and confirmed that there is generally a mutual consistency of these three features in terms of the MJO phases, for which a warming in the polar stratosphere is found. We note, however, that the strong atmospheric variability as well as different analysis setups and properties of the datasets only allow for rough consistency checks instead of precise comparisons. Therefore, the readers should pay particular attention to the detailed discussion above if more multifaceted information is needed.

Two more aspects related to SSWs are here of interest. First, the fact that SSWs occur mostly in the northern winter hemisphere, where they produce large temperature deviations in the polar regions, fits well to our finding that the largest temperature anomalies of our complete analysis occur in this area. This could be seen as support of the idea that the MJO and SSWs are interconnected and produce together the strongest responses correlated to the MJO phases. However, one could also argue that possible uncorrelated temperature deviations caused by the SSWs are so large that they are not completely eliminated by our statistical approach, so that the polar winter response is only (partly) an artifact. While we cannot totally rule out the latter, we have discussed several arguments, which indicate that the boreal winter polar temperature anomalies are really correlated with the MJO and probably amplified by the occurrence of SSWs. One of these arguments is that the same pattern of the temperature responses to the MJO can be seen in the southern hemisphere winter, even if the periods with the rare SSWs in this hemisphere are excluded from the analysis. The second aspect of interest in the context of SSWs is that a connection between the QBO phase and the occurrence of SSWs is known such that the QBO easterly phase supports the occurrence of SSWs. Together with the MJO-SSW connection, this could be a second reason for the strong QBO dependence of the MA temperature response to the MJO during boreal winter.

We also analyzed the change of the MA temperature responses while the MJO transitions through its eight phases. For most atmospheric conditions, we could identify a systematic behavior, which is again most clear for boreal winter and QBO easterly conditions. It can be described in two steps: First, the responses to the eight phases can mostly be attributed to either a "weak wave driving" or a "strong wave driving" class according to the sign of the five-zone-response. Moreover, responses that are attributed to the same class belong to subsequent MJO phases (e.g., for the boreal winter and QBO easterly situation the responses to the MJO phases 1 to 4 belong to the weak wave driving class whereas the responses to the phases 5 to 8 belong to the strong wave driving class). This supports the idea that we observe a real systematic connection between MJO and MA temperature with our statistical approach. Second, for many transitions from one MJO phase to the next a gradual shift of the five-zone-response to lower altitudes can be observed, suggesting the MA reacts systematically to each of the MJO phases.

1030 We are not aware of any explicit discussion of such a descent of a temperature anomaly pattern in the literature, which is related to the IHC or the MJO influence on MA temperature. Hence, this is likely a new aspect, for which a mechanism has still to be established. E.g., it appears not to be directly clear whether each phase of the MJO triggers the IHC pattern at a different altitude or if the IHC pattern inherently descends after it has ~~be-been~~ triggered once by a certain MJO phase or a combination of both.

1035 ~~Overall, we have presented statistical evidence for a connection of the tropospheric feature MJO and the MA feature IHC and have supported with additional arguments that a real causal connection likely exists. We think that this~~ The results presented here have a statistical character and do therefore not guarantee a real causal connection or uncover the physical mechanism. However, by embedding our results in the context of previous research, we have made plausible that a real causal connection as underlying reason is to be expected. This also contains a discussion of the interferences of the found MJO-IHC link with the QBO, SSWs, etc.

1040 We note that the present study and the mentioned previous research can benefit from each other. The previous studies provide partly more detailed analyses including physical mechanisms for particular aspects, which supports the plausibility of the current statistical findings. The present study now provides an overarching framework for previously found aspects by identifying the MA temperature response to the MJO as the IHC feature, which spans almost the complete MA.

1045 We have not attempted to present a physical mechanism together with the statistical results here. Despite the existence of mechanism approaches for individual aspects in the literature, establishing an overall mechanism appears to require more extensive future research. Such a mechanism should, in our view, cover at least the following aspects: The nature of the generation of the initial wave disturbance by all MJO phases, the propagation of the signal in both hemispheres as well as in summer and winter, the propagation in the winter hemisphere with and without SSWs, the interaction with the QBO, and the descent of the IHC temperature pattern while the MJO evolves. Of particular interest would be a detailed temporal view on the 1050 propagation of the disturbances in the MA that explains, which of the foregoing MJO phases are actually causally responsible for the MA temperature pattern that emerges statistically for a particular MJO phase in our results.

1055 Given that our statistical findings are physically substantiated in the future, we think that the described influence of the MJO on MA temperature is a noteworthy example for the complex couplings across different atmospheric layers and geographical regions in the atmosphere: in this case the source of disturbances is located at lower altitudes in the tropics (while it is probably itself influenced from above) and turns out to have observable implications up to 100 km altitude and from pole to pole. Moreover, the results further demonstrate the interconnections of different atmospheric features, particularly the MJO, the IHC, the QBO, and SSWs.

1060 Because of the wide coverage of atmospheric regions and included dynamical features, we think that the observational results presented here can ~~help to further constrain the underlying mechanisms. For the same reasons and because of the simplicity of the resulting temperature response pattern, the results presented here appear also to~~ be a good benchmark for atmospheric models covering the complete atmosphere. This applies particularly for those models, which depend especially on a good representation of troposphere-stratosphere couplings on the intraseasonal timescale, like, e.g., models that are used for intraseasonal weather predictions.

1065 *Code and data availability.* The releases of the OMI calculation package are freely available from a persistent storage referenced by Hoff-
mann (2021), while ongoing improvements are continuously found at <https://github.com/cghoffmann/mjoindices>, last access August 16,
2022. The MLS temperature data is available for download at <https://doi.org/10.5067/Aura/MLS/DATA2520>, last access September 15, 2022.
NOAA Interpolated Outgoing Longwave Radiation (OLR) data provided by the NOAA PSL, Boulder, Colorado, USA, is available from their
Web site <https://psl.noaa.gov/data/gridded/data.olrcdr.interp.html>, last access August 16, 2022. The zonal wind data for the QBO character-
ization is provided by Freie Universität Berlin at their website <https://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html>, last
1070 access August 16, 2022.

Author contributions. CGH has outlined the project, performed the analysis and written the manuscript. LGB has carried out a pilot study
under the supervision of CGH and CvS showing the potential of the analysis, CvS has extensively discussed the approach and the results
during all phases of the project. All authors have reviewed, discussed and improved the manuscript.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

1075 *Acknowledgements.* [We would like to thank the two anonymous reviewers for their valuable comments, which helped to improve the paper significantly.](#) This work was supported by the University of Greifswald. We are grateful to the teams of the NASA MLS instrument, of the
NOAA OLR dataset, and of the QBO dataset maintained by the Freie Universität Berlin for providing their valuable datasets. The analysis has
benefited from discussions on natural variability of the MA temperature and MA dynamics within the DFG (German Research Foundation)
research unit VolImpact, particularly within the subproject VolDyn.

Allen, D. R., Bevilacqua, R. M., Nedoluha, G. E., Randall, C. E., and Manney, G. L.: Unusual Stratospheric Transport and Mixing during the 2002 Antarctic Winter, *Geophys. Res. Lett.*, 30, 1599, <https://doi.org/10.1029/2003GL017117>, 2003.

Baldwin, M. P. and Dunkerton, T. J.: Stratospheric Harbingers of Anomalous Weather Regimes, *Science*, 294, 581–584, <https://doi.org/10.1126/science.1063315>, 2001.

Baldwin, M. P., Gray, L. J., Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton, J. R., Alexander, M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnersley, J. S., Marquardt, C., Sato, K., and Takahashi, M.: The Quasi-biennial Oscillation, *Rev. Geophys.*, 39, 179–229, <https://doi.org/10.1029/1999RG000073>, 2001.

Baldwin, M. P., Ayarzagüena, B., Birner, T., Butchart, N., Butler, A. H., Charlton-Perez, A. J., Domeisen, D. I. V., Garfinkel, C. I., Garny, H., Gerber, E. P., Hegglin, M. I., Langematz, U., and Pedatella, N. M.: Sudden Stratospheric Warmings, *Rev. Geophys.*, 59, e2020RG000708, <https://doi.org/10.1029/2020RG000708>, 2021.

Becker, E., Müllemann, A., Lübken, F.-J., Körnich, H., Hoffmann, P., and Rapp, M.: High Rossby-wave Activity in Austral Winter 2002: Modulation of the General Circulation of the MLT during the MaCWAVE/MIDAS Northern Summer Program, *Geophys. Res. Lett.*, 31, L24S03, <https://doi.org/10.1029/2004GL019615>, 2004.

Beig, G.: Long-Term Trends in the Temperature of the Mesosphere/Lower Thermosphere Region: 1. Anthropogenic Influences, *J. Geophys. Res. Space Phys.*, 116, A00H11, <https://doi.org/10.1029/2011JA016646>, 2011.

Beig, G., Keckhut, P., Lowe, R. P., Roble, R. G., Mlynczak, M. G., Scheer, J., Fomichev, V. I., Offermann, D., French, W. J. R., Shepherd, M. G., Semenov, A. I., Remsberg, E. E., She, C. Y., Lübken, F. J., Bremer, J., Clemesha, B. R., Stegman, J., Sigernes, F., and Fadnavis, S.: Review of Mesospheric Temperature Trends, *Rev. Geophys.*, 41, 1015, <https://doi.org/10.1029/2002RG000121>, 2003.

Camp, C. D. and Tung, K.-K.: The Influence of the Solar Cycle and QBO on the Late-Winter Stratospheric Polar Vortex, *J. Atmos. Sci.*, 64, 1267–1283, <https://doi.org/10.1175/JAS3883.1>, 2007.

Cassou, C.: Intraseasonal Interaction between the Madden–Julian Oscillation and the North Atlantic Oscillation, *Nature*, 455, 523–527, <https://doi.org/10.1038/nature07286>, 2008.

Densmore, C. R., Sanabia, E. R., and Barrett, B. S.: QBO Influence on MJO Amplitude over the Maritime Continent: Physical Mechanisms and Seasonality, *Mon. Wea. Rev.*, 147, 389–406, <https://doi.org/10.1175/MWR-D-18-0158.1>, 2019.

Domeisen, D. I. V., Butler, A. H., Charlton-Perez, A. J., Ayarzagüena, B., Baldwin, M. P., Dunn-Sigouin, E., Furtado, J. C., Garfinkel, C. I., Hitchcock, P., Karpechko, A. Y., Kim, H., Knight, J., Lang, A. L., Lim, E.-P., Marshall, A., Roff, G., Schwartz, C., Simpson, I. R., Son, S.-W., and Taguchi, M.: The Role of the Stratosphere in Subseasonal to Seasonal Prediction: 2. Predictability Arising From Stratosphere-Troposphere Coupling, *J. Geophys. Res. Atmos.*, 125, e2019JD030923, <https://doi.org/10.1029/2019JD030923>, 2020.

Espy, P. J., Ochoa Fernández, S., Forkman, P., Murtagh, D., and Stegman, J.: The Role of the QBO in the Inter-Hemispheric Coupling of Summer Mesospheric Temperatures, *Atmos. Chem. Phys.*, 11, 495–502, <https://doi.org/10.5194/acp-11-495-2011>, 2011.

Garfinkel, C. I., Feldstein, S. B., Waugh, D. W., Yoo, C., and Lee, S.: Observed Connection between Stratospheric Sudden Warmings and the Madden-Julian Oscillation, *Geophys. Res. Lett.*, 39, L18 807, <https://doi.org/10.1029/2012GL053144>, 2012.

Gray, L. J., Beer, J., Geller M., Haigh J. D., Lockwood M., Matthes K., Cubasch U., Fleitmann D., Harrison G., Hood L., Luterbacher J., Meehl G. A., Shindell D., van Geel B., and White W.: Solar Influences on Climate, *Rev. Geophys.*, 48, RG4001, <https://doi.org/10.1029/2009RG000282>, 2010.

Haynes, P., Hitchcock, P., Hitchman, M., Yoden, S., Hendon, H., Kiladis, G., Kodera, K., and Simpson, I.: The Influence of the Stratosphere on the Tropical Troposphere, *J. Meteorol. Soc. Jpn.*, 99, 803–845, <https://doi.org/10.2151/jmsj.2021-040>, 2021.

Henderson, G. R., Barrett, B. S., and M. Lafleur, D.: Arctic Sea Ice and the Madden–Julian Oscillation (MJO), *Clim. Dynam.*, 43, 2185–2196, <https://doi.org/10.1007/s00382-013-2043-y>, 2014.

1120 Hoffmann, C. G.: Mjoindices: Complete Implementation of the OMI MJO Index Algorithm in Python 3, Zenodo, <https://doi.org/10.5281/zenodo.4852196>, 2021.

Hoffmann, C. G. and von Savigny, C.: Indications for a Potential Synchronization between the Phase Evolution of the Madden–Julian Oscillation and the Solar 27-Day Cycle, *Atmos. Chem. Phys.*, 19, 4235–4256, <https://doi.org/10.5194/acp-19-4235-2019>, 2019.

Hoffmann, C. G., Kiladis, G. N., Gehne, M., and von Savigny, C.: A Python Package to Calculate the OLR-Based Index of the Madden–Julian-Oscillation (OMI) in Climate Science and Weather Forecasting, *J. Open Res. Softw.*, 9, 9, <https://doi.org/10.5334/jors.331>, 2021.

1125 Holton, J. R. and Tan, H.-C.: The Influence of the Equatorial Quasi-Biennial Oscillation on the Global Circulation at 50 {mb}, *J. Atmos. Sci.*, 37, 2200–2208, [https://doi.org/10.1175/1520-0469\(1980\)037<2200:TIOTEQ>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2200:TIOTEQ>2.0.CO;2), 1980.

Hood, L. L.: QBO/Solar Modulation of the Boreal Winter Madden-Julian Oscillation: A Prediction for the Coming Solar Minimum, *Geophys. Res. Lett.*, 44, 3849–3857, <https://doi.org/10.1002/2017GL072832>, 2017.

1130 Jucker, M., Reichler, T., and Waugh, D. W.: How Frequent Are Antarctic Sudden Stratospheric Warnings in Present and Future Climate?, *Geophys. Res. Lett.*, 48, e2021GL093215, <https://doi.org/10.1029/2021GL093215>, 2021.

Karlsson, B., Körnich, H., and Gumbel, J.: Evidence for Interhemispheric Stratosphere-Mesosphere Coupling Derived from Noctilucent Cloud Properties, *Geophys. Res. Lett.*, 34, L16 806, <https://doi.org/10.1029/2007GL030282>, 2007.

Karlsson, B., McLandress, C., and Shepherd, T. G.: Inter-Hemispheric Mesospheric Coupling in a Comprehensive Middle Atmosphere Model, *J. Atmos. Sol.-Terr. Phys.*, 71, 518–530, <https://doi.org/10.1016/j.jastp.2008.08.006>, 2009a.

1135 Karlsson, B., Randall, C. E., Benze, S., Mills, M., Harvey, V. L., Bailey, S. M., and Russell III, J. M.: Intra-Seasonal Variability of Polar Mesospheric Clouds Due to Inter-Hemispheric Coupling, *Geophys. Res. Lett.*, 36, L16 806, <https://doi.org/10.1029/2009GL040348>, 2009b.

Kikuchi, K., Wang, B., and Kajikawa, Y.: Bimodal Representation of the Tropical Intraseasonal Oscillation, *Clim. Dynam.*, 38, 1989–2000, <https://doi.org/10.1007/s00382-011-1159-1>, 2012.

1140 Kiladis, G. N., Dias, J., Straub, K. H., Wheeler, M. C., Tulich, S. N., Kikuchi, K., Weickmann, K. M., and Ventrice, M. J.: A Comparison of OLR and Circulation-Based Indices for Tracking the MJO, *Mon. Wea. Rev.*, 142, 1697–1715, <https://doi.org/10.1175/MWR-D-13-00301.1>, 2014.

Körnich, H. and Becker, E.: A Simple Model for the Interhemispheric Coupling of the Middle Atmosphere Circulation, *Adv. Space. Res.*, 45, 661–668, <https://doi.org/10.1016/j.asr.2009.11.001>, 2010.

1145 Kumari, K., Oberheide, J., and Lu, X.: The Tidal Response in the Mesosphere/Lower Thermosphere to the Madden-Julian Oscillation Observed by SABER, *Geophys. Res. Lett.*, 47, e2020GL089172, <https://doi.org/10.1029/2020GL089172>, 2020.

Kumari, K., Wu, H., Long, A., Lu, X., and Oberheide, J.: Mechanism Studies of Madden-Julian Oscillation Coupling Into the Mesosphere/Lower Thermosphere Tides Using SABER, MERRA-2, and SD-WACCMX, *J. Geophys. Res. Atmos.*, 126, e2021JD034595, <https://doi.org/10.1029/2021JD034595>, 2021.

1150 Labitzke, K. and van Loon, H.: On the Association between the QBO and the Extratropical Stratosphere, *J. Atmos. Sol.-Terr. Phys.*, 54, 1453–1463, [https://doi.org/10.1016/0021-9169\(92\)90152-B](https://doi.org/10.1016/0021-9169(92)90152-B), 1992.

Lau, W. K.-M. and Waliser, D. E.: Intraseasonal Variability in the Atmosphere-Ocean Climate System, Springer, Berlin, Heidelberg, 2012.

1155 Liebmann, B. and Smith, C. A.: Description of a Complete (Interpolated) Outgoing Longwave Radiation Dataset, *Bull. Amer. Meteor. Soc.*, 77, 1275–1277, 1996.

Livesey, N. J., Read, W. G., Wagner, P. A., Froidevaux, L., Santee, M. L., Schwartz, M. J., Lambert, A., Millán Valle, L. F., Pumphrey, H. C., Manney, G. L., Fuller, R. A., Jarnot, R. F., Knosp, B. W., and Lay, R. R.: *Aura Microwave Limb Sounder (MLS) Version 5.0x Level 2 and 3 Data Quality and Description Document (JPL D-105336 Rev. A)*, 2020.

Madden, R. A. and Julian, P. R.: Description of Global-Scale Circulation Cells in the Tropics with a 40–50 Day Period, *J. Atmos. Sci.*, 29, 1109–1123, [https://doi.org/10.1175/1520-0469\(1972\)029<1109:DOGS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1972)029<1109:DOGS>2.0.CO;2), 1972.

Marshall, A. G., Hendon, H. H., Son, S.-W., and Lim, Y.: Impact of the Quasi-Biennial Oscillation on Predictability of the Madden–Julian Oscillation, *Clim. Dynam.*, 49, 1365–1377, <https://doi.org/10.1007/s00382-016-3392-0>, 2017.

Maycock, A. C., Randel, W. J., Steiner, A. K., Karpechko, A. Y., Christy, J., Saunders, R., Thompson, D. W. J., Zou, C.-Z., Chrysanthou, A., Luke Abraham, N., Akiyoshi, H., Archibald, A. T., Butchart, N., Chipperfield, M., Dameris, M., Deushi, M., Dhomse, S., Di Genova, G., Jöckel, P., Kinnison, D. E., Kirner, O., Ladstädter, F., Michou, M., Morgenstern, O., O’Connor, F., Oman, L., Pitari, G., Plummer, D. A., Revell, L. E., Rozanov, E., Stenke, A., Visioni, D., Yamashita, Y., and Zeng, G.: Revisiting the Mystery of Recent Stratospheric Temperature Trends, *Geophys. Res. Lett.*, 45, 9919–9933, <https://doi.org/10.1029/2018GL078035>, 2018.

Moss, A. C., Wright, C. J., and Mitchell, N. J.: Does the Madden–Julian Oscillation Modulate Stratospheric Gravity Waves?, *Geophys. Res. Lett.*, 43, 3973–3981, <https://doi.org/10.1002/2016GL068498>, 2016.

1170 Murphy, D. J., Alexander, S. P., and Vincent, R. A.: Interhemispheric Dynamical Coupling to the Southern Mesosphere and Lower Thermosphere, *J. Geophys. Res. Atmos.*, 117, D08 114, <https://doi.org/10.1029/2011JD016865>, 2012.

Naujokat, B.: An Update of the Observed Quasi-Biennial Oscillation of the Stratospheric Winds over the Tropics, *J. Atmos. Sci.*, 43, 1873–1877, [https://doi.org/10.1175/1520-0469\(1986\)043<1873:AUOTOQ>2.0.CO;2](https://doi.org/10.1175/1520-0469(1986)043<1873:AUOTOQ>2.0.CO;2), 1986.

Randel, W. J., Shine, K. P., Austin, J., Barnett, J., Claud, C., Gillett, N. P., Keckhut, P., Langematz, U., Lin, R., Long, C., Mears, C., Miller, A., Nash, J., Seidel, D. J., Thompson, D. W. J., Wu, F., and Yoden, S.: An Update of Observed Stratospheric Temperature Trends, *J. Geophys. Res. Atmos.*, 114, D02 107, <https://doi.org/10.1029/2008JD010421>, 2009.

1175 Santer, B. D., Painter, J. F., Bonfils, C., Mears, C. A., Solomon, S., Wigley, T. M. L., Gleckler, P. J., Schmidt, G. A., Doutriaux, C., Gillett, N. P., Taylor, K. E., Thorne, P. W., and Wentz, F. J.: Human and Natural Influences on the Changing Thermal Structure of the Atmosphere, *Proc Natl Acad Sci USA*, 110, 17 235, <https://doi.org/10.1073/pnas.1305332110>, 2013.

1180 Schwartz, M., Livesey, N., and Read, W.: MLS/Aura Level 2 Temperature V005, Goddard Earth Sciences Data and Information Services Center (GES DISC), <https://doi.org/10.5067/Aura/MLS/DATA2520>, 2020.

Schwartz, M. J., Lambert, A., Manney, G. L., Read, W. G., Livesey, N. J., Froidevaux, L., Ao, C. O., Bernath, P. F., Boone, C. D., Cofield, R. E., Daffer, W. H., Drouin, B. J., Fetzer, E. J., Fuller, R. A., Jarnot, R. F., Jiang, J. H., Jiang, Y. B., Knosp, B. W., Krüger, K., Li, J.-L. F., Mlynczak, M. G., Pawson, S., Russell, J. M., Santee, M. L., Snyder, W. V., Stek, P. C., Thurstans, R. P., Tompkins, A. M., Wagner, P. A., Walker, K. A., Waters, J. W., and Wu, D. L.: Validation of the Aura Microwave Limb Sounder Temperature and Geopotential Height Measurements, *J. Geophys. Res. Atmos.*, 113, D15S11, <https://doi.org/10.1029/2007JD008783>, 2008.

Son, S.-W., Lim, Y., Yoo, C., Hendon, H. H., and Kim, J.: Stratospheric Control of the Madden–Julian Oscillation, *J. Climate*, 30, 1909–1922, <https://doi.org/10.1175/JCLI-D-16-0620.1>, 2017.

Sun, C., Yang, C., and Li, T.: Dynamical Influence of the Madden–Julian Oscillation on the Northern Hemisphere Mesosphere during the 1190 Boreal Winter, *Sci. China Earth Sci.*, 64, 1254–1266, <https://doi.org/10.1007/s11430-020-9779-2>, 2021.

Timmreck, C.: Modeling the Climatic Effects of Large Explosive Volcanic Eruptions, *WIREs Clim Change*, 3, 545–564, <https://doi.org/10.1002/wcc.192>, 2012.

Tsuchiya, C., Sato, K., Alexander, M. J., and Hoffmann, L.: MJO-related Intraseasonal Variation of Gravity Waves in the Southern Hemisphere Tropical Stratosphere Revealed by High-Resolution AIRS Observations, *J. Geophys. Res. Atmos.*, 121, 7641–7651, <https://doi.org/10.1002/2015JD024463>, 2016.

1195 von Savigny, C., Rozanov, A., Bovensmann, H., Eichmann, K.-U., Noël, S., Rozanov, V., Sinnhuber, B.-M., Weber, M., Burrows, J. P., and Kaiser, J. W.: The Ozone Hole Breakup in September 2002 as Seen by SCIAMACHY on ENVISAT, *J. Atmos. Sci.*, 62, 721–734, <https://doi.org/10.1175/JAS-3328.1>, 2005.

Wang, F. and Wang, L.: An Exploration of the Connection between Quasi-Biennial Oscillation and Madden-Julian Oscillation, *Environ. Res. Lett.*, 16, 114 021, <https://doi.org/10.1088/1748-9326/ac3031>, 2021.

1200 Wang, F., Tian, W., Xie, F., Zhang, J., and Han, Y.: Effect of Madden-Julian Oscillation Occurrence Frequency on the Interannual Variability of Northern Hemisphere Stratospheric Wave Activity in Winter, *J. Climate*, 31, 5031–5049, <https://doi.org/10.1175/JCLI-D-17-0476.1>, 2018a.

Wang, S., Ma, D., Sobel, A. H., and Tippett, M. K.: Propagation Characteristics of BSISO Indices, *Geophys. Res. Lett.*, 45, 9934–9943, <https://doi.org/10.1029/2018GL078321>, 2018b.

1205 Wheeler, M. C. and Hendon, H. H.: An All-Season Real-Time Multivariate MJO Index: Development of an Index for Monitoring and Prediction, *Mon. Wea. Rev.*, 132, 1917–1932, [https://doi.org/10.1175/1520-0493\(2004\)132<1917:AARMMI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<1917:AARMMI>2.0.CO;2), 2004.

Yang, C., Li, T., Smith, A. K., and Dou, X.: The Response of the Southern Hemisphere Middle Atmosphere to the Madden-Julian Oscillation during Austral Winter Using the Specified-Dynamics Whole Atmosphere Community Climate Model, *J. Climate*, 30, 8317–8333, <https://doi.org/10.1175/JCLI-D-17-0063.1>, 2017.

1210 Yang, C., Smith, A. K., Li, T., and Dou, X.: The Effect of the Madden-Julian Oscillation on the Mesospheric Migrating Diurnal Tide: A Study Using SD-WACCM, *Geophys. Res. Lett.*, 45, 5105–5114, <https://doi.org/10.1029/2018GL077956>, 2018.

Yang, C., Li, T., Xue, X., Gu, S.-y., Yu, C., and Dou, X.: Response of the Northern Stratosphere to the Madden-Julian Oscillation During Boreal Winter, *J. Geophys. Res. Atmos.*, 124, 5314–5331, <https://doi.org/10.1029/2018JD029883>, 2019.

1215 Yasui, R., Sato, K., and Miyoshi, Y.: Roles of Rossby Waves, Rossby–Gravity Waves, and Gravity Waves Generated in the Middle Atmosphere for Interhemispheric Coupling, *J. Atmos. Sci.*, 78, 3867–3888, <https://doi.org/10.1175/JAS-D-21-0045.1>, 2021.

Yoo, C. and Son, S.-W.: Modulation of the Boreal Wintertime Madden-Julian Oscillation by the Stratospheric Quasi-Biennial Oscillation, *Geophys. Res. Lett.*, 43, 1392–1398, <https://doi.org/10.1002/2016GL067762>, 2016.

1220 Yoo, C., Feldstein, S., and Lee, S.: The Impact of the Madden-Julian Oscillation Trend on the Arctic Amplification of Surface Air Temperature during the 1979–2008 Boreal Winter, *Geophys. Res. Lett.*, 38, L24 804, <https://doi.org/10.1029/2011GL049881>, 2011.

Yoo, C., Lee, S., and Feldstein, S. B.: Mechanisms of Arctic Surface Air Temperature Change in Response to the Madden–Julian Oscillation, *J. Climate*, 25, 5777–5790, <https://doi.org/10.1175/JCLI-D-11-00566.1>, 2012.

Yuan, X., Kaplan, M. R., and Cane, M. A.: The Interconnected Global Climate System—A Review of Tropical–Polar Teleconnections, *J. Climate*, 31, 5765–5792, <https://doi.org/10.1175/JCLI-D-16-0637.1>, 2018.

1225 Zhang, C.: Madden-Julian Oscillation, *Rev. Geophys.*, 43, RG2003, <https://doi.org/10.1029/2004RG000158>, 2005.

Zhang, C.: Madden–Julian Oscillation: Bridging Weather and Climate, *Bull. Amer. Meteor. Soc.*, 94, 1849–1870, <https://doi.org/10.1175/BAMS-D-12-00026.1>, 2013.

Zhang, C. and Zhang, B.: QBO-MJO Connection, *J. Geophys. Res. Atmos.*, 123, 2957–2967, <https://doi.org/10.1002/2017JD028171>, 2018.