



An overview of sedimentary volcanism on Mars

Petr Brož¹, Dorothy Oehler², Adriano Mazzini³, Ernst Hauber⁴, Goro Komatsu⁵, Giuseppe Etiope⁶,
Vojtěch Cuřín⁷

¹Institute of Geophysics of the Czech Academy of Sciences, Boční II/1401, 141 31 Prague, Czech Republic

5 ²Planetary Science Institute, Tucson, AZ, United States

³Centre for Earth Evolution and Dynamics, University of Oslo, Norway

⁴Institute of Planetary Research, DLR, Rutherfordstr. 2, 12489, Berlin, Germany

⁵International Research School of Planetary Sciences, Università d'Annunzio, Viale Pindaro 42, 65127 Pescara, Italy

⁶Istituto Nazionale di Geofisica e Vulcanologia, Sezione Roma 2, Italy; Faculty of Environmental Science and Engineering,

10 Babes Bolyai University, Cluj-Napoca, Romania

⁷Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Czech Republic

Correspondence to: Petr Brož (petr.broz@ig.cas.cz)



Abstract

15 Extensive fields of sub-kilometre-to kilometre-scale mounds, cones, domes, shields, and flow-like edifices cover large parts
of the martian lowlands. These features have been compared to structures on Earth produced by sedimentary volcanism – a
process that involves subsurface sediment/fluid mobilization and commonly releases methane to the atmosphere. It was
proposed that such processes might help to explain the presence of methane in martian atmosphere and may also have produced
habitable, subsurface settings of potential astrobiological relevance. However, it remains unclear whether sedimentary
20 volcanism on Earth and Mars share genetic similarities; hence whether methane, or other gases were released on Mars during
this process. The aim of this review is to summarize the current knowledge about mud-volcano-like structures on Mars, address
the critical aspects of this process, identify key open questions, and point to areas where further research is needed to
understand this phenomenon and its importance for the Red Planet’s geological evolution. We show here that after several
decades of exploration, the amount of evidence supporting martian sedimentary volcanism has increased significantly, but as
25 critical ground truth is still lacking, alternative explanations cannot always be ruled out. We also highlight that the lower
gravity and temperatures on Mars compared to Earth control the dynamics of clastic eruptions as well as surface emplacement
mechanism and resulting morphologies of erupted material. This implies that shapes and triggering mechanisms of mud-
volcano-like structures may be different from those observed on Earth. Therefore, comparative studies should be done with
caution. To provide a better understanding of the significance of these abundant features on Mars, we argue for follow-up
30 studies targeting putative sedimentary volcanic features identified on the planet’s surface and, if possible, for in situ
investigations by landed missions such as that currently in progress by the Zhurong rover.

1 Introduction

The buoyant ascent of liquefied, fluid-rich and fine-grained sediments through a lithologic succession and its subsequent
intrusion or extrusion (hereafter referred to as mud volcanism; Kopf, 2002) is a common phenomenon on Earth. Also known
35 as subsurface sediment and fluid mobilization (van Rensbergen et al., 2003), it is observed in sedimentary basins typically
characterised by the rapid accumulation of fine-grained and organic-rich deposits (Mazzini and Etiope, 2017 and references
therein). Images of the martian surface acquired in the early 1970’s by the Mariner 9 and Viking Orbiter missions revealed the
existence of large outflow channels likely incised by flooding events capable of transporting large amount of sediments (e.g.,
Baker and Milton, 1974; Komatsu and Baker, 1997) that were subsequently deposited in giant impact basins acting as local
40 depocenters (Lucchitta et al., 1986). Despite these observations, mud volcanism was never considered as a significant process
that could have shaped the surface of Mars. Only some early works hypothesised this type of activity based on low-resolution
imagery at specific localities (e.g., Davis and Tanaka, 1995; Tanaka, 1997; Ori et al., 2000, 2001). A renewed interest for
subsurface sediment mobilization processes emerged when higher-resolution images became available through the Mars
Global Surveyor (MGS), Mars Express (MEX), and Mars Reconnaissance Orbiter (MRO) missions and, importantly, some
45 later studies reporting the detection of methane in the martian atmosphere (Krasnopolsky et al., 2004; Formisano et al., 2004;



Mumma et al., 2009; Webster et al., 2018). On Earth mud volcanism is an important source of methane to the atmosphere (e.g., Dimitrov, 2003; Milkov et al., 2003; Etiope and Milkov, 2004). By analogy it was then hypothesised that gas released at mud-volcano-like features could be responsible for the atmospheric methane observed on Mars (e.g., Skinner and Mazzini, 2009; Komatsu et al., 2011; Etiope et al., 2011; Oehler and Etiope, 2017). Whereas today the existence of methane on Mars is
50 debated (e.g., Oehler and Etiope, 2021; Grenfell et al., 2022) after the ExoMars Trace Gas Orbiter (TGO) failed to detect it (Korablev et al., 2019; Knutsen et al., 2021), the interest in mud volcano-like structures on Mars remains high. This geological phenomenon may have an important role in atmosphere gas emissions. Additionally, the eruption of deeply buried deposits at putative clastic eruptions sites provides a window into the subsurface and the sedimentary history of Mars which is otherwise
55 only provided by impact crater excavation (e.g., crater central uplifts; Cockell and Barlow, 2002; Mustard et al., 2009; Quantin et al., 2012;). The source region of remobilised sediments and fluids may have been located in habitable deep environments (e.g., Michalski et al., 2013; Stamenković et al., 2021; see the review by Cockell [2014] for limitations of subsurface habitat niches). Therefore, erupted materials represent prime targets for in-situ investigations and the search for biosignatures (Westall et al., 2015).

The first hints at possible mud volcano-like structures on Mars were provided by Viking Orbiter images which revealed the
60 presence of dozens to thousands of small pitted cones on the floors of large ancient impact basins (e.g., Allen, 1979; Frey et al., 1979). After several decades of Mars exploration, the amount of evidence supporting sedimentary volcanism has significantly increased (Fig. 1). The aim of this review is to summarize the current knowledge, identify key open questions, and point to areas where further research is needed to understand this phenomenon and its importance for Mars' geological evolution. Section 2 provides the definition of mud volcanism and review its main characteristics observed on Earth. Here, we
65 also identify some key requirements needed to generate terrestrial mud volcanism and, by analogy, some key questions that should be addressed when searching for evidences to validate the presence of possible subsurface sediment mobilization on the Red Planet. In Section 3, we describe the morphological and morphometrical properties of potential mud volcano-like surface structures and show some typical examples for the different morphological classes, highlighting the similarities and dissimilarities to terrestrial mud volcanoes. Section 4 defines the specific martian environmental conditions and their influence
70 on potential sedimentary volcanic processes. Section 5 assess the prerequisites and the possible timing for the occurrence of sedimentary volcanism during the martian history. In Section 6 we discuss putative sedimentary volcanism in the context of Mars' geologic evolution and its implications for habitability. Finally, in Section 7 we also suggest measurements by present and future missions (including promising sampling locations) as well as laboratory experiments and modelling activities to test the sedimentary volcanism hypothesis on Mars.
75 Finally we note that the authors of the paper have different opinions and views about the subject of this paper. So several aspects addressed in the work are not unanimously agreed, and therefore the paper represents an arena of debate, rather than a consensual overview.



2 Surface mud-fluid manifestations on Earth: definitions, origin, and variants

In this section, we briefly illustrate the main features of surface sedimentary mud-fluid manifestations observed on Earth, which include sedimentary volcanoes (or mud volcanoes) and some variants. We also emphasize on the issue of the potentially ambiguous identification of the various surface manifestations and terminology reported in the literature. A correct definition of the process and related surface manifestations is essential for studying and understanding the putative sedimentary volcanism and mud volcano-like structures identified on Mars. Once the minimum requirements, conditions and factors, necessary to generate terrestrial mud volcanoes are defined, by analogy the same prerequisite can be inferred for the martian structures discussed in this work, assuming these are truly a product of sedimentary volcanism. More specifically, we will try to answer two important questions: (a) is the surface mud always stemming from a deep shale mobilised (diapir) along a fault? and (b) is the presence of gas necessary in sedimentary volcanism? Is that gas mostly methane, like on Earth? If the answers are yes, the martian mud-volcano-like structures are very likely sites where biomarkers can be better preserved and are, or have been, sites of methane release to the martian atmosphere.

2.1 Sedimentary volcanism: definitions and genetic process

Terrestrial sedimentary volcanoes are widely studied from geological, geophysical and geochemical points of view; for the details on their geographic distribution, inventories, characteristics, and impact on the environment, atmosphere and energy resource exploration, the reader may primarily refer to the main review works and references therein (e.g., Dimitrov, 2002; Kopf, 2002; Mazzini and Etiope, 2017).

However, it is crucial to clarify that not all muddy-fluid manifestations are mud volcanoes. The term “mud volcano” should not be used for any gas manifestation resembling a mud pool or where extrusive mud gives rise to small conic edifices. Many CO₂-vents, related to geothermal or hydrothermal environments, may show such characteristics. It is not only a problem of semantics, because the attribution of “mud volcano” to a surface gas manifestation implies the existence of a series of specific geologic processes and features.

The definition of mud volcano is strictly linked to the genetic mechanisms that are essential to generate this specific geological phenomenon. The processes are schematically illustrated in Fig. 2. Basically, a mud volcano is characterised by the surface release of mostly deep-seated shale deposits that are brecciated and transported along a fracture system or fault (the shale is for a mud volcano what magma is for an igneous volcano). The shale and other rock types are fractured and mobilised, in a diapir style, due to the combination of three factors: (a) lateral tectonic compression; (b) gravitational instability of shale, i.e., typically when shale is a low density sediment underlying another sedimentary rock with higher density; (c) overpressure of gas and water from an underlying source. This source is generally a gas reservoir, which provides the necessary pressure, likely the factor triggering the mobilization of the shale (Mazzini and Etiope, 2017). The erupted sediments are defined with the term “mud breccia” and consist of a mix of ground-up and brecciated sediments and rocks that originate from various units and formations intersected by the conduit. Two components are typically identified in the mud breccia a) a fine-grained matrix



110 (consisting mostly of clay and fine sediments in large part originating from the low density shales) and b) a set of rock clasts
with sizes ranging from a few cm to cubic metres stripped from the various sedimentary formations that have been pierced by
the diapir. The depth of the reservoirs (which can be found by seismic images and/or drillings) and the origin of the clasts are
fundamental to assess the minimum depth of the mud volcano system. Mud volcanoes are located almost exclusively in
convergent basins but they can occur along any type of fault, with normal, reverse or strike-slip kinematics (Ciotoli et al.
115 2020).

Sedimentary volcanism on Earth is exclusively observed in petroleum-bearing sedimentary basins, i.e., in areas with gas-oil
systems (source rocks, reservoirs, generally in faulted anticlines). Therefore, the erupted gas is mainly methane, associated
with other hydrocarbon gases in trace amounts (ethane, propane, butane) and non-hydrocarbon gases typically occurring in
sedimentary reservoirs (CO₂, N₂, minor amounts of He, and H₂S). Methane is mostly of deep thermogenic origin, but often
120 including shallower components of secondary microbial methane (Etiope et al., 2009). A few mud volcanoes in petroliferous
basins may release more N₂, due to differential solubility in uplifted basins, which also induces higher helium concentrations
(Etiope et al., 2011). Water is typically salty and enriched in various elements, largely sourced from the brines accumulated in
the gas-oil reservoir (fossil water) in addition to connate water (trapped in sediments), which may also derive from illitization
of clayey minerals (smectite-illite transformation; Mazzini and Etiope, 2017). Meteoric water can also mix with the fossil
125 water during its upwelling.

2.2 Variants

The mud volcano system depicted in Fig. 2 may be more complex and a few variants of this classic scheme may exist. In some
cases, the gas may stem from multilayer reservoirs and the mobilised shales may originate from different sedimentary units
and, likewise, multiple source rocks may be involved in the plumbing system (e.g. Guliyev and Feizullayev, 1997; Inan et al.,
130 1997; Cooper et al., 2001). In addition, a set of other surface sedimentary mud-fluid manifestations exists on Earth that may
have sizes and morphologies that resemble those of mud volcanoes. For this reason, the attribution of sedimentary volcanism
for these variants may be uncertain, ambiguous, or susceptible to subjective and/or misleading interpretations. However, more
detailed studies reveal that these structures have genetic mechanisms, and accordingly names, that are different from those of
sedimentary volcanism. For example:

135 (a) *Sediment-Hosted Geothermal Systems*. Some sedimentary basins, featuring petroleum systems, developed above deep
geothermal systems (e.g. along grabens), which can be associated with igneous intrusions, deep magmatic chambers, or lateral
migration of hydrothermal fluids. These systems can be characterised by elevated CO₂ and H₂O pressures. These settings are
referred as “Sediment-Hosted Geothermal Systems” (SHGSs), where CO₂-dominated fluids migrate upward and mix with the
gaseous hydrocarbons hosted in the shallower sedimentary formations (i.e., Procesi et al., 2019). SHGSs are essentially hybrid
140 systems and the relative fraction of the two endmembers (sedimentary CH₄ and geothermal CO₂, with related waters) may
vary greatly. By definition, however, a SHGS releases on the surface a mixture with CO₂ concentrations >50 vol % (Procesi



et al., 2019) and it may have the features of a hydrothermal system, with large amounts of water vapour (as in the case of the Lusi system in Indonesia; Mazzini et al., 2012).

145 (b) *Artesian systems*. At mud spring sites, overpressured water displaces fine-grained sediments during upwelling forming localised pools. These are generally shallow systems (tens or a few hundreds of metres deep), with surface manifestations within the range of some metres and mud flows whose extension may vary depending on the terrain morphology. These springs do not necessarily contain gas (e.g. Bristow et al., 2000). Dewatering structures like sand volcanoes may also display conical shaped features reaching > 10 m in size. These are shallow-rooted structures resulting from the remobilization of shallow non-consolidated sediments (Gill et al., 1957).

150 (c) *Sedimentary diapirism*. Here shales are purely driven by gravitative instability (similar to salt diapirism), without the need of gas overpressure; the movement is continuous but extremely slow and does not produce fluid eruptions or episodic activity on the surface (e.g. Bouriak et al., 2000; Henry et al., 2022; Bulanova et al., 2018).

155 (d) *Injected sands* are rapid phenomena mostly triggered by sand fluidization due to earthquakes, do not require gas overpressure and do not show presence of mud breccia. They may also be called sand injectites, they occur at depth and require the fracturing of overburden sediments through which fluidised sands migrate eventually reaching the surface (i.e. extrudites) (Jolly and Lonergan, 2002; Hurst et al., 2006; Polteau et al., 2008).

160 (e) *Pingoes* are morphological deformation of the ground, typically associated with frozen aquifers (so they exclusively exist on Earth in high latitude regions), as well as associated with the presence of permafrost or gas hydrates; they are quite shallow and do not imply relevant transport of sediments but rather only surficial deformation. Their morphology may resemble that of small mud volcanoes. Although gas is not necessary to form a pingo (which is essentially a hydrological phenomenon), methane can be released when pingoes are emplaced along faults or at pockmarks sites (Andreassen et al. 2017; Hodson et al. 2020).

2.3 Morphologies, rheology and mud flows in sedimentary volcanism

165 Mazzini and Etiope (2017) provide an overview of the dimensions (from metre scale up to 12 km in diameter) and morphologies observed at numerous mud volcano sites worldwide and propose a classification identifying the main processes that control the ultimate shape of the structures and the extensions of the mud flows. These processes are directly connected with the mechanisms of eruption/erosion that characterize each site and can be applied also to the other surface mud-fluid manifestations described herein. Dynamic factors include the eruption frequency and the vigour (i.e. more explosive events vs less destructive events). The amount of gas/water/sediments-rocks released during the eruptions will affect the rheology of the mud and, accordingly, the ultimate morphology of the flows. The local topography pre-existing the eruptions will affect the shape of the surface manifestations. Mechanical factors include a) the interaction between the erupted media (more or less viscous) and the surface (more or less erodible) hosting the flows, b) the type of erosion (operated by e.g. submarine currents, wind, rain) that is different depending of the geographical setting, or c) the type of e.g. subsidence associated with subsurface



dynamics. Different albedo observed at the mud flows can be used to infer the erosive efficiency of the erupted media (e.g. 175 Mazzini et al. 2021), or the chronology of the eruptive phases (Mazzini et al. 2009).

2.4 Answering the key questions

With the clarifications given above, and based on the numerous observations and studies on mud volcanoes worldwide, we will now try to answer to the two key questions about sedimentary volcanism:

(a) is the surface mud always stemming from a deep shale mobilised (diapir) along a fault?

180 (b) Is the presence of gas necessary?

(a) On Earth, at least for the relatively large (tens to thousands of metres in diameter) structures, it has been documented that the main mud component typically originates from deep shale units, upwelling even from depths of several kilometres, as demonstrated by studies conducted on mud breccia deposits (e.g., Cita et al. 1981; Inan et al. 1997, Akhmanov et al. 2003). 185 Mud breccia and faults appear essential components of sedimentary volcanism. For small structures, a few centimetres or metres wide, the mud can be shallower and the process may be more similar to an artesian system described above. Accordingly, the answer would be “yes” for traditional and relatively large sedimentary volcanoes, with mud breccia containing clasts and biomarkers transported from great depths.

(b) All non-ambiguous (and relatively large) mud volcanoes on Earth systematically release methane (and other gaseous 190 hydrocarbons), because they develop within petroleum systems. However, as mentioned above, some manifestations may also release substantial amounts of CO₂ (hybrid SHGS) or N₂ (uplifted basins). Therefore, the answer to the second question is also “yes”. On Earth, sedimentary volcanism is always associated with gas emissions, as the gas is fundamental for the rapid and episodic shale mobilization (Fig. 2). Gas may not be necessary for those peculiar structures, such as sedimentary diapirism and injected sands or other dewatering structures, described above, whose genetic mechanisms are different. Fundamentally, they 195 do not have the “eruptive and fluid discharge” character that is typical of sedimentary volcanism.

In conclusion, it is evident that the size of the emission structure, the presence of gas, mud breccia and faults are relevant to understand the genetic process driving mud-fluid manifestations. Larger (tens-hundreds metres wide) structures imply deeper roots, and their formation needs relevant fluid pressures and mobilizations of large volumes of sediments. Smaller structures (<a few metre wide) may represent shallower processes, possibly involving local aquifers and gas in the shallow subsoil. The 200 “size”, “mud breccia” and “fault” factors, observable in the available images from Mars, shall be carefully considered in the interpretation of the mud-volcano-like structures on the Red Planet, as discussed below. The release of gas can only be detected by on-site ground measurements (rovers) or, in case of substantial gas plumes, by orbiters (Oehler and Etiope, 2021).



3 Observations

In this section, we briefly summarize the current knowledge about the morphology of mud-volcano-like structures on Mars. Four individual sub-sections will focus on a) sub-kilometre to kilometre-scale circular mounds widely spread across the northern lowlands, b) the kilometre-scale topographically positive features of various shapes and often associated with flow-like edifices, c) kilometre-scale flows, and d) hundreds of kilometre-long flows and deposits. These divisions are made in order to group features that bear similar morphological, morphometrical and spatial similarities, although overlapping characteristics in some parameters often exist among these groups.

3.1 Morphologic expression of putative sedimentary volcanism on Mars

3.1.1. Sub-kilometre to kilometre-scale circular mounds

Many tens of thousands of bright, sub-kilometre- to kilometre-scale, circular mounds occur in the northern plains of Mars (Fig. 3). The largest abundances are in Acidalia and Utopia Planitiae, but additional examples occur in Chryse and Arcadia Planitiae, Cydonia Mensae, the Isidis-Utopia overlap, Isidis Planitia, and possibly in the Scandia region (*e.g.*, Davis and Tanaka, 1995; Tanaka, 1997, 2005; Tanaka et al., 2000, 2003, 2008; Farrand et al., 2005; Kite et al., 2007; Rodríguez et al., 2007; Skinner and Tanaka, 2007; Allen et al., 2009, 2013; McGowan, 2009; Oehler and Allen, 2009, 2010, 2011, 2012 a, b; Skinner and Mazzini, 2009; Komatsu, 2010; McGowan and McGill, 2010; Allen et al., 2013; Ivanov et al., 2014; Orgel et al., 2015, 2019; Komatsu et al., 2011, 2016; Hemmi and Miyamoto, 2018; De Toffoli et al., 2019, 2021). While the great majority of these mounds occur in the northern lowlands, possible examples have been reported from several localities in the highlands, including craters in Arabia Terra (Pondrelli et al., 2011; Franchi et al., 2014), and a flat-floored depression in Terra Sirenum (Hemmi and Miyamoto, 2017).

The mounds have typically been compared to a variety of terrestrial analogues (*e.g.*, rootless cones, scoria cones, tuff cones, pingos, erosional remnants, clathrate degasification structures, and mud volcanoes), and in many cases, a mud volcano interpretation has been deemed most consistent with the observed morphologies, geologic setting, associated flow structures, and evidence for a fine-grained (low thermal inertia) sediment size.

The mounds in Acidalia Planitia (Fig. 3) were described in detail by Oehler and Allen (2010). In that study, the total area of their occurrence was outlined, and in about half that area, 18,000+ examples were mapped. Based on that, ~40,000 such features were estimated for southern Acidalia Planitia. Hemmi and Miyamoto (2018) studied 1,300 mounds in southern Acidalia, using 40 digital elevation models (DEMs) to measure mound heights and basal diameters. These two studies provide the following observations: The mounds in Acidalia (Fig.3) are circular to subcircular in plan view, with diameters ranging from ~0.3 to 2.2 km (average ~0.8 km). The heights of the examples measured by Hemmi and Miyamoto (2018) average 15.2 m, with a range of 1.1 to 69.5 m. In cross-sectional profile, many of the mounds appear as domes (Fig. 3a), commonly displaying a central depression (Figs. 3c-d), or steep-sided cones. Many are surrounded by moats (Fig. 3d). Some bright



235 circular mounds may appear to be nearly flat and more irregular in plan view, and these can sometimes enclose large, boulder-
like knobs (Fig. 3e). Farrand et al. (2005) first recognised and termed these morphologies as domes, cones, and splotches. Most
have a high albedo and smooth surfaces relative to the surrounding plains. Some have apron-like extensions of the smooth,
high albedo material onto the plains. Similar smooth-textured, high-albedo material forms occasional, lobate flow-like
structures that emanate from some of the mounds (Figs. 3c,f). Thermal Emission Imaging System (THEMIS) Nighttime
Infrared (IR) images show the mounds to be dark in Nighttime IR (compare Figs. 3a-b), implying that their thermal inertia is
240 lower than that of the plains and likely reflects a finer grain size than that of the plains (Oehler and Allen, 2010). Farrand et al.
(2005) concluded that "dried, loosely cemented, mud deposits would be a good match to both the albedo and thermal inertia
of these mounds". The mineralogy of the bright mounds in Acidalia has been investigated with data from the Compact
Reconnaissance Imaging Spectrometer for Mars (CRISM), but results have not been definitive, possibly due to enhanced
coatings of poorly crystalline materials (Oehler and Allen, 2010).

245 **3.1.2 Kilometre-scale cones, domes and shields**

While the circular mounds discussed in previous subsection show relative uniformity in their appearance within individual
fields, this is not the case for members of the kilometre-scale feature type. Members of this group have been reported from
Valles Marineris in Candor and Coprates Chasmata (Harrison and Chapman, 2008; Chan et al., 2010; Okubo, 2016; Kumar et
al., 2019; Wheatley et al., 2019), in the southern part of Chryse Planitia (Komatsu, 2010; Komatsu et al., 2011, 2016; Brož et
250 al., 2019), in the Amenthes/Nepentes region (Skinner and Tanaka, 2007), or in Hydraotes Chaos (Meresse et al., 2008).
Edifices these regions display a wide range of shapes and sizes (Fig. 4), both among individual fields as well as when compared
between them. Some edifices have a conical morphology including well-developed summit craters (Fig. 4a), others have no
such craters (Harrison and Chapman, 2008; Chan et al., 2010; Okubo, 2016; Kumar et al., 2019; Wheatley et al., 2019). Other
structures have the cross-sectional shape of domes with steep sides and flat summit areas that may feature small central knobs
255 (Fig. 4b). Additional morphologies include shield- or pie-like edifices with one or multiple summit craters and well-developed
lobate margins (Fig. 4c; Komatsu et al., 2016; Brož et al., 2019). The central craters associated with kilometre-scale cones and
shield-like edifices might be either enclosed (Fig. 4a) or breached (Fig. 4e). In several regions, some of these edifices are
associated with flow-like units over which they might be superposed (Fig. 4d; Okubo, 2016; Brož et al., 2017), or embayed by
(Fig. 4e; Skinner and Tanaka, 2007; Brož and Hauber, 2013).

260 Edifices of this group have broad variations in their sizes, both among features in individual fields as well as among various
fields. For example, the field of ~130 conical features within Coprates Chasma range in sizes from 0.2 km to 2 km in diameter,
with average of 0.8 km (based on 59 edifices; Brož et al., 2017). Edifices are from 75 to 250 m high and their flanks are
between 15 and 25° (based on 6 edifices; Brož et al., 2015a). On the other hand, ~170 pitted cones in Amenthes/Nepentes
region are much wider than the edifices in Coprates Chasma and range from 3 to 15 km in diameter (average 7.8 km, based on
265 measurements of 92 edifices). They are also higher, from 30 to 370 m (average ~120 m, based on measurements of 53 edifices)



and their slopes are typically below 10°. Features among this field also have much wider craters as compared to those in Coprates Chasma, and the craters are often breached. The crater floors can be below the level of the surrounding plains (Brož and Hauber, 2013). The field displaying the largest known variation among individual edifices is situated in the southern part of Chryse Planitia. Here, more than 1,300+ structures can be classified into five different types – cratered cones, shield-like
270 edifices, domes with a small central knob on their summit craters, irregular pies without significant topographical expression and kilometre-scale flows (Brož et al., 2019). These edifices can range in size from 0.2 km up to 20 km.

Because they have a similar variety of shapes and are commonly associated with flows, it is common for terrestrial km-scale mud volcanoes to be proposed as potential analogue (e.g., Jakubov et al., 1971; Aliyev et al., 2015; Mazzini and Etiope, 2017). However, the debate whether some of these kilometre-scale features were formed by sedimentary volcanism is not yet settled
275 as igneous volcanism has been invoked as an alternative mechanism for their formation in several cases (Lucchitta, 1990; Meresse et al., 2008; Brož and Hauber, 2013; Brož et al., 2015a; 2015b; 2017). If this is indeed the case, some of these features might represent martian analogues to terrestrial scoria cones, lava domes, tuff rings and/or tuff cones.

The spectral data from CRISM have been used to gain insight into their formation mechanism, however, the data did not unambiguously show the presence of phyllosilicates, carbonates, or sulphates in association with these edifices (Dapremont
280 and Wray, 2021). Hence, their sedimentary origin is not supported or refuted by mineralogical information. Additionally, Brož et al. (2017) reported the presence of partially dehydrated opaline silica, polyhydrated and monohydrated sulphate, high-calcium pyroxene, likely combined with olivine, and hydrous silica in the summit area of one cone, and mafic minerals associated with a flow-like unit in Coprates Chasma. These authors proposed that igneous volcanism might explain these observations, hence questioning a sedimentary origin of the field in Coprates Chasma (Okubo, 2016). To date, spectral
285 observations do not support a sedimentary origin over an igneous one, or vice versa.

3.1.3 Kilometre-scale flows

Kilometre-scale flows (KSFs) have been identified on several martian low plains; Chryse (Komatsu et al. 2011, 2016; Brož et al., 2019; 2022a), Acidalia (Ivanov et al., 2015; Ivanov and Hiesinger, 2020), Utopia (Ivanov et al., 2014, 2015) and Elysium Planitiae (Wilson and Mougini-Mark, 2014), as well as at the highland-lowland boundary (HLB) between southern Utopia
290 Planitia and Terra Cimmeria (Skinner et al., 2007, 2009). They represent a morphologically diverse group of landforms characterised by a) spatially extensive spreading controlled by local topography, and b) the presence of lobate margins indicative of flow processes. In some martian volcanic provinces, like Tharsis or Elysium, such landforms have been interpreted as lava flows (e.g., Hartmann & Berman, 2000; Bleacher et al., 2007; Keszthelyi et al., 2008; Hauber et al., 2009, 2011). Based on remote sensing data, however, subsurface sediment mobilization has been proposed as the most likely
295 formation mechanism for some of these features (e.g. Skinner and Tanaka, 2007; Ivanov et al., 2014, 2015; Wilson & Mougini-Mark, 2014; Komatsu et al. 2011, 2016; Okubo, 2016; Brož et al., 2019).



KSFs are often elongated in plan-map view as the material released from their source areas flowed along the local topographic gradient and/or filled nearby terrain depression(s). However, in flat areas of Chryse and Utopia Planitiae their length-to-width ratio is comparatively lower. KSFs vary greatly in areal extent and may feature individual or overlapping flow-like structures (e.g., Komatsu et al., 2016; Brož et al., 2022). The largest examples are situated within Utopia Planitia. Here many overlapping flows spread over areas ranging in size from 10^2 to 10^3 km² (Cuřín et al., 2023). However, similar extents have been found for individual KSF such as, e.g., Zephyria Fluctus which has an area of 153 km² and an average thickness of 3.8 m (Wilson and Mouginiš-Mark, 2014). This unique flow has been interpreted to be formed by the emplacement of mud by Wilson and Mouginiš-Mark (2014), however, its igneous origin was not definitely ruled out. Individual flows in Utopia Planitia typically cover an area of ~35 km² and have cross-sectional shapes of plateaus standing above surrounding terrains to heights of 20-30 metres (Cuřín et al., 2023). These are comparable to the examples found in Acidalia Planitia and Utopia-Cimmeria HLB (Skinner et al., 2007, 2009). The flows present in Chryse Planitia are smaller (few 10s of km²; Brož et al., 2019).

Most KSFs comprise central and marginal units, each with specific morphologies depending on the specific region on Mars. In Utopia and Acidalia, KSFs are characterised by smooth central flat parts with randomly distributed rimless depressions (Fig. 5e; Ivanov et al., 2015) and marginal units consisting of coalescing pits, distributory branching, and lobate hummocks (Fig. 5e,f). The KSFs of the Utopia-Cimmeria HLB consist of lobate material with varying thickness (Skinner et al., 2009), but the difference between their central and marginal units is not as pronounced. Zephyria Fluctus' central laminar flow unit has a unique polygonal surface texture with steep-sided margins (Fig. 5h,i; Wilson and Mouginiš-Mark, 2014). The channelised flow-like features of Chryse Planitia are more complex and consist of three morphological elements: a central depression, leveed channels, and a distal portion of the fading channel(s) (Fig. 5a; Komatsu et al., 2016; Brož et al., 2019; 2022a). Topographically low mounds inside the central depressions have been interpreted to mark the positions of feeder vents (Brož et al., 2022a). Similarly, Zephyria Fluctus seems to be sourced from a circular pond-like depression (Fig. 5, Wilson and Mouginiš-Mark, 2014).

In Acidalia and Chryse Planitiae, KSFs are often associated with circular mounds. A similar spatial association is observed in Utopia Planitia, where plateau-like outflows are often emanating from elongated ridges (Cuřín et al., 2023). In Utopia and Acidalia Planitiae, KSFs appear in proximity of craters with pancake-like ejecta, ghost craters (Ivanov et al., 2015) and giant polygons (Ivanov et al., 2014; Cuřín et al., 2023).

KSFs are discernible in THEMIS Nighttime IR imagery. Across all the mentioned locations, the central units are darker than the marginal one, but their contrast to the surrounding terrain varies. In Chryse and Utopia Planitiae, they appear darker than the surroundings, while KSFs in Acidalia Planitia as well as Zephyria Fluctus appear brighter than the surrounding landforms. Using THEMIS Nighttime IR data, Komatsu et al. (2011) examined the material forming one of the KSFs within Chryse Planitia and the associated conical and dome-shaped edifices. They found that these features had a lower thermal inertia (i.e. potentially finer grainsize) than the surrounding units, making a composition by compact igneous rocks unlikely. However, it should be noted that if these flows were formed of unconsolidated fine-grained pyroclastic material, they would show a lower thermal inertia as well.



Similar flows as KSFs can be also found in association in some regions of Mars with large valleys systems, e.g., like flows associated with Granicus and Tinjar Valles that are situated west of the Elysium Mons volcano. These flows have been interpreted to be mud flows, however, they likely formed due to the permafrost melting caused by the volcanic activity (Russell et al., 2003; Pedersen, 2013), not due to sedimentary volcanism. It remains unclear at the moment, if the mud was produced in
335 the subsurface during melting, or formed on the surface by mixing of the released water with surface sediments.

3.2 Distribution and geologic setting

The martian mud-volcano-like surface features have been mainly observed in various parts of the northern lowlands of Mars, especially in Utopia, Chryse and Acidalia Planitiae. The circular mounds in Acidalia commonly occur at elevations below -
4000 m in locations associated with giant polygons, which also are almost exclusively found in the northern plains of Mars
340 (Oehler and Allen, 2012a; Allen et al., 2013; Moscardelli et al., 2012). The mounds occur individually, in pairs, in irregular clusters, and in chains. The chains may be relatively rectilinear in plan view, overlying troughs forming giant polygons (Gallagher et al., 2018), or they may be curvilinear, aligned along arcuate ridges in a type of landform termed “thumbprint terrain” (Guest et al., 1977). Facies analysis incorporating catchment areas and transport distances of sediment deposition from the Noachian to Hesperian fluvial systems and late Hesperian outflow floods predicts that southern Acidalia would be the
345 depocenter for fine-grained, distal-facies muds (Rice and Edgett, 1997; Allen et al., 2013; Oehler and Allen, 2010, 2012a, b, 2021). The long transport distances from Noachian catchments in the highlands to depocenters in southern Acidalia would promote excellent grain size separation, such that distal facies sediments in southern Acidalia would be expected to contain thick accumulations of mud. This prediction is supported by Salvatore and Christensen (2014a, b) who used high resolution data sets to investigate morphology and spectral signatures and concluded that southern Acidalia is a region of extensive, fine-
350 grained and water-saturated sedimentation, and Ivanov and Hiesinger (2020) who conducted a photogeological study along with crater-size-frequency distributions and concluded that a volatile-saturated, mud-rich unit was deposited in the southern Acidalia plains. Thumbprint terrain in two regions of Acidalia has been interpreted as the result of impact-generated tsunamis (Rodríguez, et al., 2016; Costard et al., 2017), supporting the concept of a northern ocean and water-saturated sediments in the Acidalia depocenters. The mounds that occur in curvilinear chains in this terrain have been interpreted as mud-volcano-like
355 structures formed by rapid compaction in high energy flows caused by the tsunami-producing impacts (Di Pietro et al., 2021). After deposition of Noachian to Hesperian fluvial units within and below a) the region now covered by the Vastitas Borealis Interior and Marginal units of Tanaka et al., 2005 or b) the late Hesperian lowlands unit (IH1) of Tanaka et al., 2014, the late Hesperian, circum-Chryse outflow floods would have injected enormous quantities of sediments into the same area (Lucchitta et al., 1986; Baker et al., 2015; Alemanno et al., 2018). Both the earliest outflow sediments and the underlying, older strata
360 would have been rapidly buried. On Earth, rapid burial of a volatile- and mud-rich section is ideal for development of subsurface overpressure and initiation of mud volcanism (Kopf, 2002). The circum-Chryse outflow sediments could have done



the same on Mars (Oehler and Allen, 2010). This process might additionally explain the approximate co-location of bright mounds and giant polygons in Acidalia (Oehler and Allen, 2012a; Allen et al., 2013; Orgel et al., 2019).

365 The Chryse outflow floods and sediment emplacement might be also responsible for the formation of kilometre-scale cones, domes and shields situated in the southern part of Chryse Planitia, in a region near the termini of several large outflow channels, namely in the Simud, Ares and Tiu Valles (Komatsu et al., 2011, 2016; Pajola et al., 2016). Here, more than 1,300+ edifices, classified in five different types (including kilometre-sized cones, domes, shields and KSF), spread over the area of 700,000 km² (Brož et al., 2019). The distribution of these features shows that they are clustered and anticorrelated to the erosional remnants of ancient highlands, suggesting a genetic link between their distribution and the sedimentary deposits over
370 which they are superposed. Their distribution also shows that different types of features occur preferentially at specific latitudes (Fig. 5 in Brož et al., 2019), although the area of their extension often overlaps with regions populated also by other types. Such distribution patterns could be related to a model of sandar facies as suggested for this region by Rice and Edgett (1997). These authors identified three facies types (proximal, midfan, and distal facies) in a lateral sequence progressing from south to north. Most of the known features were mapped in the zone of the midfan facies, however, no clear correlation between
375 feature type and distance to outflow channel termini was found (Brož et al., 2019).

Another distinct distribution pattern was found for a field of ~170 pitted cones and 80+ smaller mounds (Skinner and Tanaka, 2007; Brož and Hauber, 2013) in the Amenthes/Nephtes region situated close to the dichotomy boundary, between the cratered highlands of Tyrrhena Terra in the south and smoother plains of Utopia Planitia in the north. Here, the edifices are aligned, from west to east, in a NW-SE and then W-E direction parallel to the southern margin of Utopia Planitia. To explain
380 such distribution, Skinner and Tanaka (2007) proposed the existence of annular ring basins in an impact tectonics scenario that would have acted as locations for sediment accumulation in southern Utopia Planitia and hence a source reservoir for sedimentary volcanism. Also field of ~130 pitted cones reported from Valles Marineris in Coprates Chasmata (Harrison and Chapman, 2008; Chan et al., 2010; Okubo, 2016; Brož et al., 2017; Kumar et al., 2019; Wheatley et al., 2019) show aligned distribution in NW-SE direction. Brož et al. (2017) proposed that such distribution is controlled by structures oriented roughly
385 parallel to the long axis of the Coprates Chasma tectonic graben.

The southwestern part of Utopia Planitia, in the region of Adamas Labyrinthus, also displays evidence for possible sedimentary volcanism in the form of a field of more than 300 of KSFs. They have been firstly described by Ivanov et al. (2014) who referred to them as «etched flows». An additional mapping campaign performed by Cuřin et al. (2023) categorised these KSFs into 4 classes ('hills', 'ridges', 'plateaus', and 'complexly layered units'). Several KSFs can be also found throughout Acidalia
390 Planitia southward of Acidalia Mensae (Ivanov et al., 2015; Ivanov and Hiesinger 2020), as well as around the Utopia-Cimmeria HLB (Skinner et al., 2007, 2009); although no comprehensive inventory of their presence in these regions exists. In Elysium Planitia, the single flow of Zephyria Fluctus with a supposed sedimentary origin is present within the lower unit of the Medusae Fossae Formation (Fig. 5g,h,i; Wilson and Mouginis-Mark, 2014).

It remains unclear at which depth the source reservoir for the hypothesised sedimentary volcanoes is located. Hemmi and
395 Miyamoto (2018) estimated source depths for the mounds in Acidalia to be 110 – 850 m (if the mounds were formed



subaerially) and 30 – 450 m (if the mounds were formed sub-aqueously). Their work was based on bulk densities, fractures associated with co-located giant polygons, and an isostatic compensation model where the depth of the mud source was estimated from mound heights. De Toffoli et al. (2019) estimated source depths for mounds in Arcadia Planitia of 16 – 18 km. Their work was based on fractal analysis of the mounds to assess whether their spatial distributions were consistent with control by underlying fractures and then on the assumption that upper cut offs determined by the fractal analysis reflect the depths of the fluid source. The orders-of-magnitude difference in estimated depths to source reservoirs of these two studies highlights uncertainties in both the approaches utilised for these assessments as well as the understanding of the origin of the bright mounds in the northern plains of Mars.

3.3 Ages

With the exception of really extensive flows, the age of martian mud-volcano-like structures is difficult to determine as they do not represent units of sufficient size for crater counting (e.g., Warner et al., 2015). Moreover, many of them have a relatively rugged topography with steep slopes. Hence, it is typically only possible to date spatially larger units with a known relative stratigraphic position with respect to the hypothesised mud volcanoes (i.e. either the edifices are superposed on these units or are partly buried/embayed by them) (e.g., Brož and Hauber, 2013; Brož et al., 2019). This approach enables bracketing their ages by maximum and minimum ages, however, this approach is commonly fraught with large uncertainties.

This can be illustrated on the example of circular mounds in Acidalia. In most areas, these mounds have erupted onto the Vastitas Borealis Formation (VBF), a Late Hesperian to Early Amazonian (~3.2 to 1.75 Ga) unit interpreted as either a paleo-ocean deposit (Kreslavsky and Head, 2002) or a mixture of Noachian to Hesperian materials and local outflow-channel sediments (Tanaka et al., 2003). Since the majority of the mounds overlie the VBF, they must be younger than its Late Hesperian to Early Amazonian age. Nevertheless, the minimum age has not been established. Some studies interpret the mounds as generally ancient features that formed on early Mars while fluids were still abundant in the shallow subsurface of the northern plains and perhaps while an ocean existed (e.g., Oehler and Allen, 2010; Oehler and Allen, 2012 a; Allen et al., 2013). This interpretation would be consistent with the interpretations of «thumbprint» terrain as a product of tsunamis. However, this view has been challenged by Rodríguez et al. (2019) who suggest a later stage of sedimentary volcanism that postdates a possible ocean. Supporting a more recent origin of the features, De Toffoli et al. (2019) proposed that relatively small mounds (0.3 to 0.5 km in diameter) associated with thumbprint terrain in Arcadia Planitia may be young, with a “last occurrence” of ~370 Ma. That age is based on craters < 1 km in diameter, which because of their small size, could reflect a resurfacing age rather than a formation age (see Warner et al., 2015 for discussion of crater sizes and age interpretations). Similar uncertainties are associated with the ages of kilometre-sized cones, domes, shields and KSFs. Specifically, the features within the southern part of Chryse Planitia are spread over a unit that was significantly resurfaced 3.2 Ga years ago (Brož et al., 2019) and that experienced at least another two resurfacing events. As the features seems to be formed after these resurfacing events and they are partly covered by secondaries formed during the formation of Mojave crater, which has been



dated to 4.7 Ma (Werner et al., 2014), this gives a range in their age between 880 and 5 Ma (Brož et al., 2019). Similarly, cones within Coprates Chasma were emplaced on top of sedimentary deposits of Hesperian age (Okubo, 2016), but again the unit
430 has been later resurfaced and the cones formed after this resurfacing event. So they are likely Middle to Late Amazonian in age since some are superposed by a young landslide (Brož et al., 2017). However, some known edifices might be older, like pitted cones in Amenthes/Nephtes region that are more than ~2.4 Ga old (Brož and Hauber, 2013).

The ages of KSFs in Utopia and Acidalia are similarly uncertain. Ivanov et al. (2015) tied their formation to the later stages of VBF emplacement, hence their ages should be 3.57 Ga and 3.61, respectively. Zephyria Fluctus, situated within the Late
435 Hesperia transition unit (Tanaka et al., 2005; 2014) has the visual appearance of a very recent flow (Mouginis-Mark, 2013), but its precise age has not yet been determined.

4 Effect of the environment on sedimentary volcanism

Mars is a planet with very different environmental properties as compared to Earth. Both the surface gravitational acceleration and the current atmospheric pressure are lower, at 3.71 m/s² vs. 9.81 m/s² and ~600–1000 Pa vs. ~10⁵ Pa, respectively. Today,
440 the range of atmospheric pressure at the surface of Mars can reach likely up to 12.4 mbar at the deepest point on the bottom of the ancient impact basin, Hellas, (Haberle et al., 2001, Wray, 2021) and can drop to 0.7 mbar on the top of the highest mountain, Olympus Mons. The surface temperature is on average -60°C, but it can range from -143°C at the poles up to +35°C in equatorial regions. Although average temperatures are far below the freezing point of water, locally higher temperatures at favourable seasons might be reached, theoretically enabling liquid water to be present on the surface today (Wallace and Sagan,
445 1979; Brass, 1980; Carr, 1983; Haberle et al., 2001; Hecht, 2002; Möhlmann, 2004; Kossacki et al., 2006; Bargery et al., 2010). However, such water would be likely very limited in time as the low atmospheric pressure would trigger boiling, freezing, and eventually evaporation into the atmosphere (e.g., Hecht, 2002; Wray, 2021). These processes might have inhibited the ability of water to propagate over the martian surface during most of its history.

Initial studies of mud behaviour at martian surface pressure were performed by Wilson and Mouginis-Mark (2014). The authors
450 proposed that the water present in the mud would be unstable and evaporate from the mud flow, ultimately removing the latent heat from the mixture. This implies that the residual water present in the mud mixture would freeze relatively quickly, in range of hours to days. Additional insight came from experimental work of Brož et al. (2020a,b), where the behaviour of low viscosity mud was experimentally studied in a low-pressure chamber partly simulating the Mars environment. Their results showed that low viscosity mud flows could actually propagate over cold (<273 K) and warm (>273 K) surfaces at martian atmospheric
455 pressure, but the mechanism of such propagation would be different from that observed on Earth. On Mars, mud propagating over cold surfaces would rapidly freeze on the surface of the flow due to evaporative cooling (Bargery et al., 2010) forming an icy-crust leading to propagation in a similar manner to pahoehoe lava flows observed on Earth (Hon et al., 1994). Once such an icy crust has formed, the interior of the mud flow is protected from additional evaporative cooling. As a consequence, mud remains liquid inside the crust for prolonged periods of time and propagates via mud tubes (analogous to lava tubes;



460 Calvari and Pinkerton, 1999). In contrast, low viscosity mud propagating over a warm surface boils and levitates above the
surface (Brož et al., 2020b). As the water content within martian mud flows might have varied, mud flows may have had
different viscosities. Brož et al. (2022b) experimentally revealed that the exposure of high viscosity mud to low atmospheric
pressure also leads to the formation of an icy crust, but also to a volume increase. This phenomenon occurs since the low
atmospheric pressure causes an instability of the water present in the mud mixture, leading to the formation of expanding
465 bubbles, which cannot escape from the high viscosity mud and increases the volume of the mud by up to 15%. Low-pressure
experiments hence demonstrate that the propagation and behaviour of mud on Mars indeed differs distinctly from that on Earth.
Sedimentary edifices built by mud flows on Mars are therefore expected to differ in shape and morphology from their terrestrial
counterparts (Brož et al., 2019; 2020a; 2022a).

Useful insights to understand the mechanisms of mud flows at low atmospheric pressures might come from theoretical
470 considerations of igneous volcanism on Mars, for which the roles of different gravity and atmospheric pressure have been
intensively studied (e.g., Dehn and Sheridan, 1990; Wilson and Head, 1994; Wilson and Head, 2004; Parfitt and Wilson, 2008;
Brož et al., 2021 and references therein). These investigations showed that a low surface pressure environment is capable of
affecting the ascent of magma in the feeding conduit (e.g., Wilson and Head, 1994; Parfitt and Wilson, 2008). In fact, the low
atmospheric pressure would favour the formation of bubbles and their growth within the ascending magma and hence reduce
475 its density. Such decrease would cause a larger density contrast to the surrounding rocks, hence in buoyancy, and would
ultimately increase the speed of the magmas ascent. Similarly, the rapid decompression of the mud during its ascent together
with rapid degassing might also favour its release via low energy explosive eruptions. In this case, muddy eruptions similar in
nature to Hawaiian and/or Strombolian igneous eruptions, might be more frequent on Mars than on Earth because significant
gas expansion within the final phase of its ascent might increase the ejection velocities of the expelled mixture (Wilson and
480 Head, 1994).

As surface atmospheric pressure has varied on all timescales (for a recent review see Jakosky, 2021) and temperatures were
likely higher in the past (Haberle et al., 2017), martian mud flows might have been emplaced at times where the importance
of evaporative cooling may have been significantly reduced. It is reasonable to expect that the shapes of martian sedimentary
volcanoes may then vary significantly among various fields due to local environmental properties (Brož et al., 2022a; Cuřín et
485 al., 2023). To our knowledge, no dedicated studies exist that investigated such morphological variabilities. To perform such
study is, however, particularly challenging since the ages of possible martian sedimentary volcanoes as well as the exact
paleopressures in Mars' history are only poorly constrained (e.g., Kite et al., 2014).

In addition to low atmospheric pressure, Mars also has a lower gravity as compared to Earth, and it remains unknown how this
may affect sedimentary eruptions. Once again, a comparison to igneous volcanism can provide valuable hints. It has been
490 shown that the lower gravity on Mars tends to reduce the speed of lava flows and increase their thickness, as the liquid would
spread laterally to a lesser degree (Wilson and Head, 1994; Rowland et al, 2004). As a consequence, this would change the
heat loss rate of the lava flow. In a similar manner, mud flows could form thicker accumulations without losing heat. Moreover,
the low temperatures at the martian surface could also impede the ability of water to infiltrate into the subsurface (Conway et



al., 2011; McCauley et al., 2002; Pfeffer and Humphrey, 1998), limiting water loss from the mud mixture and thus maintaining
495 the viscosity of the flow (i.e. it does not become more viscous by losing water). Finally, the low atmospheric pressure would
also reduce the importance of cooling the mud flows by convection of the overlying atmosphere.

The lower gravity on Mars could also hinder the occurrence of sedimentary volcanism, as the lithostatic pressure within the
crust has a different gradient to that on Earth. In other words, in order to achieve the same pressure as at a certain depth on
Earth, it is necessary to be approximately three times deeper on Mars (i.e. the sedimentary basins need to be three times
500 thicker). Therefore, from the theoretical point of view, it might be more challenging to have suitable depths of strata for
sedimentary volcanism to arise on Mars compared to on Earth.

Taken together, there is currently insufficient understanding of the processes that might lead to subsurface sediment
mobilization on Mars. There are also no analytical or numerical models to explain such processes, despite significant progress
in analogue modelling over the last years. The main challenge is determining how the historical variations in atmospheric
505 surface pressure and the low gravity on Mars might affect this process. Namely, for mud-volcano like edifices formed under
atmospheric conditions inhibiting the presence of liquid water on the surface for prolonged period of time, it is unknown how
evaporative cooling and the subsequent formation of an icy-crust on mud flows affects their rheology and their ability to spread
across the surface. Also we do not know what the pressure gradient driving the flow of the mud up to the surface is, how a
turbulent flow regime would affect the freezing rate of the mud, or when these mud volcano-like features formed and what the
510 conditions were like on the surface of Mars at the time. In summary, these gaps complicate the development of numerical
models that would be capable of reconstructing the formation of martian sedimentary volcanoes and would enable us to study
the role of individual parameters in sedimentary volcanic processes on Mars. In turn, these shortcomings limit our ability to
predict the surface morphology of the resulting sedimentary edifices, and hence what to search for.

5 Ideal prerequisites to form sedimentary volcanism on Mars?

515 From a general point of view, the initiation of sediment mobilization on Mars requires several conditions to be met: (a) the
relatively quick accumulation of sufficiently thick sedimentary deposits, (b) the presence of at least some strata of fine-grained
sediments and liquid water within these deposits, and (c) overpressurization of the sediments or gravitational instability as a
trigger of fluid expulsion (and likely gas expulsion as well). In other words, sedimentary volcanism on Mars cannot occur
anywhere, but should be restricted to specific locations where favourable conditions were present in the past.

520 As a first prerequisite for sediment mobilization and mud volcanism, sizable depocenters must have been available that could
be filled over time with sufficiently thick volatile-rich sedimentary strata. On Earth, such areas of sediment accumulation are
often linked to plate boundaries (e.g., Dickinson, 1974; Miall, 1984), whereas Mars is a one-plate planet (Solomon, 1978) and
does not show evidence for plate tectonics (Tosi and Padovan, 2021). Therefore, other suitable locations with sufficient
sedimentary accumulation have to be selected. The ideal candidate locations displaying such characteristics were identified by
525 many studies and include a) ancient, large impact basins that would have acted as sinks for water and sediments that were



transported in giant flood events through outflow channels (e.g., McGill, 1989; Frey et al., 2002; Ivanov et al., 2014; Jones et al., 2016), and b) large troughs created by extensional tectonics and/or collapse (notably Valles Marineris), some of which may have once hosted lakes (e.g., Harrison and Grimm, 2004; Harrison and Chapman, 2008; Warner et al., 2013; Okubo, 2016). For example, Tewelde and Zuber (2013) proposed that 2–4 km of sediment fill accumulated in Acidalia Planitia and up to 5 km of sediments were deposited in Utopia Planitia over time. However, at the moment, the exact sediment thicknesses and sedimentation rates in the basins remain unknown.

The second prerequisite is the infilling of such depocenters with fine-grained sediments that could be potentially mobilised. Several mechanisms have been proposed for the transport and deposition of such fine-grained sediments. For example, Okubo (2016) proposed that eolian deposition within the Hesperian epoch might have partly infilled the Candor Chaos and Coprates Chasma regions and that these sediments might have been later buried by Middle Amazonian sediments and mobilised into large muddy laccoliths. Alternatively, Ivanov et al. (2014) proposed that fine-grained sediments might have originated from an ancient muddy ocean fed by outflow channels floods (see also Jöns, 1985). Similar outflow channel floods were also proposed as the source for fine-grained sediments within Chryse Planitia, as analogous to terrestrial outflow events (Rice and Edgett, 1997). An additional mechanism is the deposition of fine-grained volcanic ash originating from explosive volcanic eruptions (Ivanov et al., 2012) resulting in thick and extensive pyroclastic deposits deposited over volatile-rich units (see Brož et al., 2021 and references therein). These pyroclastic deposits could then be subsequently buried by younger material.

The third prerequisite is the generation of overpressure or gravitational instability required to mobilize sediment, water and/or other volatiles. Hypotheses include a) top-down freezing of water-bearing (i.e. muddy) sediment bodies, i.e. the gradual thickening of a cryosphere (Clifford et al., 2010; Ivanov et al., 2014; Ivanov and Hiesinger, 2020) and b) rapid burial of sedimentary strata by mass-wasting processes like lahars, landslides, impacts, etc. (e.g., Tanaka, 1999; Skinner and Tanaka, 2007; Skinner and Mazzini, 2009). Overpressure resulting from these types of processes may be enhanced by gas released if clathrates are destabilised (as could occur by uplift, sublimation of a frozen ocean, temperature changes, etc. (e.g., Oehler and Allen, 2010; Oehler and Etiope, 2017), d) seismic activity (e.g., Skinner and Tanaka, 2007), e) rapid changes in the local tectonic regime (e.g., Hemmi and Miyamoto, 2017) or f) combinations of these mechanisms. In addition, one must consider the possibility that large quantity of clathrates may have been destabilised, if, for example, a frozen ocean sublimated away. If that occurred, then clathrate destabilization might be a major source of overpressure and mud volcanism in some areas. However, currently, there is neither ground truth nor numerical modelling to assess such scenarios.

Despite this uncertainty it is reasonable to expect that one or more of these mechanisms were likely active in Mars` history. As individual martian sedimentary depocenters were subject to different geological settings and conditions, and the climate and aqueous activity on Mars changed over time (cf. previous Section 4), possible sediment mobilizations did not necessarily happen at the same epochs. This could apply to both the time when the sediments were emplaced and when they were subsequently mobilised in the subsurface. It is most likely that water-rich sediments were emplaced when liquid water was more widespread on Mars than in the Middle to Late Amazonian. It seems reasonable to assume that the sediments were therefore accumulated when the valley networks and, perhaps more importantly, the outflow channels were formed (Late



560 Noachian-Early Hesperian, and Late Hesperian-Early Amazonian, respectively). As for the mobilization, it is also likely that most potential triggers for mobilization would have been more active in the Noachian and Hesperian than in the Middle to Late Amazonian (e.g., large impacts: Hartmann and Neukum, 2001; seismicity: Knapmeyer et al., 2008). The exception is the thickening of the cryosphere (top-down freezing), which is probably an ongoing process. As a result of such changes and the associated effects of the martian environment on mud behaviour (see previous Section 4), martian putative sedimentary
565 volcanoes and mud flows might show a large morphological and chronological variability. In fact, distinct depocenters would host differently-sized mud reservoirs, with sediment of diverse lithologies and geochemical provenance, contain variable amounts of available water, and experienced diverse conditions (e.g., overpressurization, density stratification) that may have triggered sediment liquefaction and mobilization. Variations within this parameter space could have controlled the effusion rates during mud expulsion as well as the water content within the mud, and in turn the viscosity of the ascending mixture
570 might show a broad variability (e.g., Brož et al., 2019) resulting in a potentially large morphological diversity of sedimentary landforms (see previous section).

6 Addressing the key questions about sedimentary volcanism for Mars

This review aims to outline the fundamental concepts and definitions of mud volcanism on Earth and to identify and describe martian examples of features that could be associated with a similar formation mechanism. The fundamental question is: Are
575 these martian features analogous to terrestrial mud volcanoes? Or, in contrast: Is it impossible to provide a definitive answer based on our current knowledge? Throughout this review we have applied the term «mud volcano-like» for structures that may have had a similar formation process to mud volcanoes on Earth. Considering the parameters that are commonly used to define mud volcanism-related phenomena on Earth, there are several aspects that need to be resolved and further investigated before we can apply Earth-based definitions to martian structures. Whereas the surface morphology of the investigated features is
580 relatively well characterised at least at the meter to kilometre scale (Section 3), the subsurface mechanisms leading to sediment mobilization as well as the link to tectonic settings remain unclear. We still cannot constrain the composition and origin of the mobilised sediments emplaced on the surface, the associated amount of water and gases, nor the triggering mechanism that would initiate sedimentary volcanism on Mars. Section 2.4 underlined key questions that should be addressed to better assess the potential for mud volcanism on Mars.

585 *Sedimentary deposits and structural discontinuities.* Most of the largest terrestrial mud volcanoes are formed in areas of major compressional stress at plate boundaries, and the source areas are at kilometre-scale depths (Mazzini and Etiope, 2017). It is known that thick accumulations of layered sediments exist on Mars, and they can locally reach thicknesses of several kilometres (e.g., Malin and Edgett, 2001; Milliken et al., 2010). Although these deposits (e.g., in Valles Marineris or in Gale crater) are exposed at the surface and are most likely not a viable source for deep-seated mud volcanism, several ancient
590 depocenters especially in large, ancient impact basins are believed to host kilometres-thick depositional units at depth (e.g., Lucchitta et al., 1986; Goldspiel and Squyres, 1991) that could act as significant reservoirs of mobile and potentially buoyant



595 sediments which could be ultimately erupted at the surface. At the termination of outflow channels, voluminous sediments must have been deposited by the flooding events and reach minimum thicknesses of more than several hundred metres (e.g., Tanaka, 1997). Direct observational evidence of such deeply buried sediments is lacking, however, and multispectral analyses of materials excavated by impact crater formation in the northern lowlands do not strongly support thick deposits of sediments in the basins with the most numerous mud volcano candidates (Chryse, Acidalia, Utopia) (Pan et al., 2017). Geophysical results are inconclusive, too. Although the average bulk density of the martian crust is lower than expected than for a mafic (basaltic) crust (Goossens et al., 2017) and lateral variations may exist, the average density seems to be lower in the southern highlands than in the northern lowlands (Wieczorek et al., 2022), which appears inconsistent with large volumes of low-density sediments in the lowlands (although it should be noted that the highest crustal densities in the lowlands correspond to volcanic provinces, not the large impact basins; Belleguic et al., 2005; Goossens et al., 2017). Nevertheless, recent seismic observations obtained through measurements by the InSight lander indicate lower shear-wave velocities in the lowlands as compared to the highlands, which could be due to thick accumulations of sediments (Li et al., 2022). New orbiters with powerful radar instruments and high-resolution gravity and measurements (Genova, 2020; Oberst et al., 2022) would be essential to identify and characterize any voluminous wet sediments in the deep martian subsurface.

600 The importance of faults and other structural discontinuities to facilitate the sediment/fluid flow on Earth has been highlighted in Section 2. Mars has lacked plate tectonics for most (or possibly all) of its history (e.g., Grott et al., 2013, Smrekar et al., 2019 and references therein), and although poorly constrained, the martian subsurface is considered to be extensively faulted and fractured after a long history of impacts, the formation and evolution of the dichotomy, and the uplift of Tharsis (Golombek and Phillips, 2010). The northern lowland areas, where most of the candidate mud volcanoes are observed, are structurally dominated by wrinkle ridges, considered to be evidence for contractional deformation of a layered substrate (see review of Mueller and Golombek, 2004). Therefore, there are certainly tectonic discontinuities in the areas of possible sedimentary volcanism. However, they are mostly indicative of compressional stress, and the dip angle of the thrust faults associated with wrinkle ridges is relatively shallow (average $\sim 30^\circ$, according to many models; e.g., Karagoz et al., 2022). Although there may be a large range of possible dips (see discussion by Andrews-Hanna, 2020), it is not clear whether these faults may have acted as preferential pathways for liquefied sediment.

605 *Hydrocarbons.* As explained in Section 2, mud volcanism on Earth is exclusively observed in hydrocarbon-bearing sedimentary basins and commonly associated with the release of significant volumes of methane (Mazzini and Etiope, 2017), a gas that significantly contributes to the greenhouse effect. At the moment, we do not know what type of gases (if any) may have been released if subsurface sediment mobilization was ever active on Mars. A summary of current observations on that regard have been provided by Oehler and Etiope (2021). The authors discuss the potential for methane on Mars (with both abiotic and potential biotic sources assessed) as well as the discrepancies between the methane measurements by Curiosity (Webster et al., 2018) and the non-detections by the ExoMars Trace Gas Orbiter (Korablev et al., 2019). Oehler and Etiope (2021) conclude that there is a strong case for the production of abiotic methane in the subsurface of Mars. Much of that could have been produced in the early history of the planet and may have already seeped to the surface. If so, remaining quantities



are likely to be trapped by the planet wide cryosphere, resulting in minor and episodic releases – potentially below detection limits of TGO. Nevertheless, methane may have been present in the martian subsurface in substantial amounts early in the planet's history. In addition, since the martian atmosphere is CO₂-rich, both methane and CO₂ are potential gases that could have been released with sedimentary volcanism on Mars. If we assume that martian sedimentary volcanoes were indeed
630 accompanied by large releases of methane (or other greenhouse gases), this would imply that those gases could have contributed to transitory climate warming in the past as well as the potential for the existence of methane reservoirs at depth (Oehler and Etiope, 2021). However, as no signs of active sedimentary volcanism on Mars have been discovered so far, and the ages of putative sedimentary edifices determined by crater counting suggest an activity at least dozens or hundreds of millions of years ago (Brož and Hauber, 2012; Brož et al., 2017; 2019), it is therefore highly unlikely that ancient sedimentary
635 volcanism is the source for martian atmospheric methane detected today (e.g., Giuranna et al., 2019; Oehler and Etiope, 2021 and references therein). Lefèvre and Forget (2009) showed that methane should have a relatively short lifetime in the current atmosphere of Mars – around 300 years in the upper atmosphere, but only 200 days or less when close to the surface. Therefore, recent methane detections are likely released by different mechanisms, e.g., by seepage from partly sealed subsurface reservoirs. And as such, it remains unclear at the moment if a genetic link between martian mud volcano-like structures and
640 methane releases exists at all.

Triggering mechanisms: On Earth, gravitational instability of shales and gas overpressure as well as water present at buried deposits are crucial factors to promote sediment mobilization (Mazzini and Etiope, 2017). It is extremely unlikely that all these mechanisms operated (or even operate) in the subsurface of Mars (e.g., Oehler and Allen, 2010) due to the likely absence of comparable quantities of hydrocarbons and plate tectonics. Other mechanisms (summarised in Section 5) have therefore been
645 proposed. For example, it was proposed by Oehler and Etiope (2021) that clathrate destabilization might account for the majority of the bright mounds in Acidalia where shallow-rooted conduits are suggested following a model similar to that recorded from, e.g., Lake Baikal (Khlystov et al., 2019). If such scenario is proven to be correct, the vast number of these mounds would reveal the extent to which clathrates are buried across the martian northern lowlands.

However, as none of these scenarios are supported by ground truth evidence nor by physical models nor by numerical
650 modelling results, this limits our ability to understand the way the sedimentary volcanism and the associated possible release of greenhouse gasses would affect the evolution of Mars. And hence, if sedimentary volcanism could help to alter the atmosphere of early Mars or not (Wordsworth, 2016).

7 Ground truthing and tests of the sedimentary volcano hypothesis

Currently available remote sensing data, despite their diversity and high spatial resolution, provide only limited insights into
655 sedimentary volcanism, and, in many cases, the surface morphologies can have alternative interpretations (e.g., Beven, 1996). Spectral data obtained from orbit are inconclusive (e.g., Komatsu et al., 2016; Dapremont and Wray, 2021). A definitive proof of sedimentary volcanism on Mars is hampered by the lack of *in-situ* data (ground truth) that should provide unambiguous



evidence. Until now, in situ observations supporting the presence of subsurface sediment mobilization are very limited. In this section, we describe the most promising examples that have been, or can be, visited on the martian surface, and suggest tests
660 of the sedimentary volcano hypothesis.

7.1. Sedimentary pipes in Gale crater

The Mars Science Laboratory mission with its rover, Curiosity, has traversed fluvio-lacustrine and aeolian sedimentary rocks that were deposited in Gale crater ~3.6 to 3.2 Gy ago (Rubin et al., 2017). Structures interpreted to be pipes formed by vertical movement of fluidised sediment were observed at several locations (Fig. 6a-d) (Rubin et al., 2017). Circular rings of erosion-
665 resistant material with diameters of 7 to 70 cm rise a few centimetres above their surroundings and display cementation and concentric internal layering. They are associated with other potential fluidised sediment features such as sedimentary dikes (Fig. 6e; Grotzinger et al., 2013) and deformational structures and may be analogues to clastic pipes on the Colorado Plateau (Ormö et al., 2004; Mahaney et al., 2004; Wheatley et al., 2016, 2019). Clastic pipes are injection features that vertically crosscut bedding with sharp contacts (Figs. 7). They display cylindrical morphologies, massive or radially graded interiors,
670 and raised outer rims. Increased grain size and subsequent cementation along the more porous edges makes the rims more resistant to weathering. Pipes have crosscutting relationships with other pipes due to multiple formation events or migrating eruption centres, and they are associated with other soft-sediment deformation features. Terrestrial clastic pipes form via liquefaction and fluidization, which require a near-surface groundwater system to initiate.

Another potentially interesting and relevant analogue is identified in western Japan along the Kii Peninsula coast where ancient
675 mud volcanism is preserved and exposed in sedimentary sequences (Komatsu et al., 2019). Coarse-grained shallow marine sediment sequences of the Miocene are intruded by the underlying fine-grained sediment. The intruding mudstone deposits exhibit diverse types of stratigraphic features, including mud dikes intruding into overlying layers or diapirs in contact with surrounding strata (Fig. 7). Like many pipes on Earth, the structures in Gale crater are more resistant to erosion than the host rock; they form near other pipes, dikes, or deformed sediment; and some contain internal concentric or eccentric layering.
680 These structures provide new evidence of the importance of subsurface aqueous processes in shaping the near-surface geology of Mars (Rubin et al., 2017).

7.2. Zhurong rover study of one putative martian sedimentary volcano

The Chinese Zhurong rover onboard Tianwen-1 landed on southern Utopia Planitia (25.066° N, 109.925° E) on 15 May 2021. In the vicinity of the landing site, orbital imagery revealed the presence of a field of cone-shaped edifices (Liu et al., 2021; Ye
685 et al., 2021; Zhao et al., 2021). In the field, there is one pitted cone with a height of 80 m and a basal diameter of 800 m located about 16 km southeast of the landing site, and it has attracted the attention of the mission science team (Liu et al., 2021). Alternative interpretations, including cinder cones, sedimentary volcanoes or pingos, have been proposed to explain the origin of this structure. Sedimentary volcanism appears to be the preferred origin as reported in recent studies (Ye et al., 2021; Huang et al., 2022), although other small mounds in the region have been interpreted as lava domes (Lin et al., 2023). In-situ study



690 of the closest cone to the landing site would provide a great opportunity for ground-truthing of one example of putative
sedimentary volcanoes on Mars. The identification of clay minerals, like smectite, or illite (Mazzini and Etiope, 2017 and
references therein), as a main bulk component of the pitted cones would be strong evidence for a mud volcano origin.

7.3. Future in-situ investigation of hypothesised sedimentary volcanism

Some proposals have been made regarding the importance of in-situ investigation of purported sedimentary volcanism
695 (Komatsu et al., 2014), and some candidate landing sites can be listed at some areas of hypothesised sedimentary volcanism
such as those in Chryse Planitia (Rodriguez et al., 2007; Komatsu et al., 2011, 2016; Brož et al., 2019; Komatsu and Brož,
2021).

The conceivable in-situ investigations at a future landing site may include those for a) geology, b) geo-biochemistry, and c)
biology. First of all, the origin of the edifices must be investigated. In-situ lithological and mineralogical examination of
700 edifices' surface or sub-surface would be essential to distinguish the formation processes. A confirmed sedimentary volcanism
scenario would bring insights about the water occurrence both on the surface and in the subsurface of Mars. Further, an analysis
of the erupted mud breccia clasts would provide unprecedented information regarding the subsurface geology at these sites.

On Earth mud volcanoes host a large variety of microbial communities that thrive particularly at seepage sites (e.g., Wrede et
al., 2012; Kokoscha et al., 2015; Tu et al., 2017; Lee et al., 2021; Miyake et al., in press). It has even been suggested that fluids
705 ascending from deeply subducted slabs may have lead to low-temperature alteration environments in the conduits of
serpentinite mud volcanoes that provided suitable niches for early life (e.g., Pons et al., 2011; Fryer et al., 2012, 2020). The
study of putative mud volcano-like structures has therefore a great potential for astrobiology investigations and to collect
fossilised microorganisms (e.g., Komatsu and Ori, 2000). Such investigation of martian sedimentary volcanoes should focus
on localities where mud eruptions occurred, including summit craters and small mud mounds (called gryphons) where
710 emissions of mud and gas might continue even after the major eruptions for prolonged period of time. However, young fresh-
looking mudflows emanating away from the summit craters are also promising candidate targets. It is recommended to conduct
drilling ≥ 2 m into the mud in order to sample materials less exposed to the harsh surface environment and protected from
surface radiation (Pavlov et al., 2022). The only currently existing rover equipped to drill to that depth is the ExoMars rover
(Vago et al., 2017), which is planned to land in the Oxia Planum region. While the main targets are phyllosilicate-bearing
715 layered deposits, there are some km-sized mounds in the area that form part of a regional population of mounds in the southern
and eastern marginal regions of the Chryse impact basin (McNeil et al., 2021). Some of these mounds have been interpreted
to be products of sedimentary volcanism (Adler et al., 2022), and if the ExoMars rover lands near one of these mounds, it is
recommended that it investigates it in situ.



8 Conclusions

720 After several decades of Mars exploration, there is a growing consensus that a phenomenon similar to sedimentary volcanism on Earth may have been active on the Red Planet. An improved understanding of martian geological settings and higher quality remote sensing images have helped in assessing the origin (sedimentary versus igneous) of numerous enigmatic features observed on the surface. This manuscript reviews the martian regions where mud-volcano like features occur and provides detailed descriptions of the observed morphologies and potential formation mechanisms.

725 The greatest abundance of mud volcano-like features occurs in the northern lowlands of Mars, commonly north of the latitudinal maximum limit that has governed the landing-site selection for previous landed missions. The largest abundances are in Acidalia and Utopia Planitiae, where a hundred thousand of such structures occur, but examples occur also in Chryse, Arcadia and Isidis Planitiae. Outside the northern lowlands, only a few fields of such mounds have been identified, for example in Arabia Terra, Valles Marineris or Terra Sirenum. These numerous mounds clearly reflect major events in the history of the
730 martian lowlands - events that could have initiated warming episodes and may even have global climatic relevance. In addition to these numerous, near-circular edifices, larger and morphologically/morphometrically more diverse features of similar origin have been also discovered. They are represented by kilometre-scale edifices with cross-sectional shapes of domes, shields, or flow-like features with a length of up to dozens of kilometres.

As outlined in this overview, there should be favourable conditions in the subsurface of Mars that enable the process of
735 sediment mobilization and hence sedimentary volcanism. This is because thick accumulations of layered sediments exist on this planet acting as significant reservoirs of mobile and potentially buoyant sediments and at the same time these layers might be extensively faulted and fractured enabling mobilised sediment to propagate to the surface. However, it is currently unknown what the exact mechanism responsible for such mobilization would be, because we do not know what type of gases (if any) may have been released. Therefore, whether the process of subsurface sediment mobilization on Mars is an exact analogue of
740 the process of terrestrial mud volcanism which is governed by gas emissions.

If the described mud-volcano like structures on Mars share a formation mechanism similar to that observed on Earth, they would represent excellent targets for future landed missions, as they could archive unaltered biosignatures, if life ever developed on Mars (Oehler and Etiope, 2021). On Earth, mud volcanoes are localities where methane is continuously released and represent ideal niches for habitability (e.g., Knittel and Boetius, 2009). They also provide a window into deep biosphere
745 potential sedimentary strata (e.g., Plümper et al., 2017), as mud volcanoes bring sediments from metres to kilometres of depth to the surface via a relatively low temperature and pressure processes (i.e. unlike impact ejecta that are associated with high pressure and temperatures). The study of similar features on Mars could, thus, provide insightful information about currently inaccessible buried stratigraphy.

Currently, new insights might come from the Chinese rover, Zhurong that is near a potential mud-volcano-like mound in
750 Utopia Planitia. Observations from this rover might provide important observations about the origin of the material(s) forming the edifice. Continuing work on analogues, using laboratory simulations of mud flows under Mars-like conditions, and



associated numerical models should help to assess the significance of some of the various edifices found on Mars. And hence this might help to predict the surface morphology of the resulting sedimentary edifices, and what to search for.

755 Future exploration, using drones or helicopters, may allow sample analysis or even sample collection from some of the potential sedimentary volcanoes in northern latitudes. This would then help to address many of our current knowledge gaps and provide direct insight about the mechanism of mud dynamics on Mars. The next decades of Mars exploration should therefore involve efforts to better explore these enigmatic features, which could be excellent candidates for future missions aimed at biosignature detection (Oehler and Etiope, 2021).

References

- 760 Adler, J.B., Bell, J.F., Warner, N.H., Dobrea, E.N., and Harrison, T.N.: Regional geology of the Hypanis Valles system, Mars. *Journal of Geophysical Research: Planets*, 127, e2021JE006994, doi:10.1029/2021JE006994, 2022.
- Akhmanov, G. G., Premoli Silva, I., Erba, E., and Cita, M. B.: Sedimentary succession and evolution of the Mediterranean Ridge western sector as derived from lithology of mud breccia clasts, *Marine Geology*, 195 (1–4), 277–299, doi:10.1016/S0025-3227(02)00693-X, 2003.
- 765 Alemanno, G., Orofino, V., Mancarella, F.: Global map of Martian fluvial systems: Age and total eroded volume estimations. *Earth and Space Science*, 5 (10), 560–577, doi.org/10.1029/2018EA000362, 2018.
- Allen, C.C.: Volcano/ice interactions on Mars, *J. Geophys. Res.*, 84, 8048–8059, doi:10.1029/JB084iB14p08048, 1979.
- Allen, C.C., Oehler, D.Z., and Baker, D.M.: Mud volcanoes – A new class of sites for geological and astrobiological exploration of Mars. 40th Lunar and Planetary Science Conference, Abs. #1749, 2009.
- 770 Allen, C.C., Oehler, D., Etiope, G., Van Rensbergen, P., Baciu, C., Feyzullayev, A., Martinelli, G., Tanaka, K., and Van Rooij, D.: Fluid expulsion in terrestrial sedimentary basins: a process providing potential analogs for giant polygons and mounds in the martian lowlands. *Icarus* 224:424–432, doi:10.1016/j.icarus.2012.09.018, 2013.
- Aliyev, A., Guliyev, F., Dadashev, F. G., and Rahmannov, R. R.: *Atlas of the world mud volcanoes*, Nafta-Press, ISBN 978-9952-437-60-7, 2015.
- 775 Andreassen, K., Hubbard, A., Winsborrow, M., Patton, H., Vadakkepuliambatta, S., Plaza-Faverola, A., Gudlaugsson, E., Serov, P., Deryabin, A., Mattingsdal, R., Mienert, J., and Büinz, S.: Massive blow-out craters formed by hydrate-controlled methane expulsion from the Arctic seafloor, *Science* 356, 6341, 948–953, doi:10.1126/science.aal4500 , 2017.
- Baker, V.R. and Milton, D. J.; Erosion by catastrophic floods on Mars and Earth, *Icarus*, 23, 27–41, doi:10.1016/0019-780 1035(74)90101-8, 1974.
- Baker, V.R., Hamilton, C.W., Burr, D.M., Gulick, V., Komatsu, G., Luo, W., Rice, Jr., J.W., and Rodríguez, J.A.P.: Fluvial geomorphology on Earth-like planetary surfaces: a review. *Geomorphology* 245, 149–182, doi:10.1016/j.geomorph.2015.05.002, 2015.



- 785 Belleguic, V., Lognonné, P., and Wieczorek, M.: Constraints on the Martian lithosphere from gravity and topography data, *J. Geophys. Res.* 110, E11005, doi:10.1029/2005JE002437, 2005.
- Beven, K.: Equifinality and uncertainty in geomorphological modelling. In *The Scientific Nature of Geomorphology*, Proc. 27th Binghampton Symp. Geomorphol., edited by B. L. Rhoads and C. E. Thorn, pp. 289–313, Wiley, Chichester, 1996.
- 790 Bouriak, S., Vanneste, M., and Saoutkine, A.: Inferred gas hydrates and clay diapirs near the Storegga Slide on the southern edge of the Vøring Plateau, offshore Norway, *Marine Geology*, 163, 1–4, 125–148, doi:10.1016/S0025-3227(99)00115-2, 2000.
- Brand, U., Blamey, N., Barbelli, C., Griesshaber, E., Posenato, R., Angiolini, L., Azmy, K., Farabegoli, E., and Came, R.: Methane hydrate: Killer cause of Earth's greatest mass extinction. *Palaeoworld* 25, 496–507. doi:10.1016/j.palwor.2016.06.002, 2016.
- 795 Brass, G. W.: Stability of brines on Mars, *Icarus* 42, 20–28, doi:10.1016/0019-1035(80)90237-7, 1980.
- Bristow, C. R., Gale, I. N., Fellman, E., Cox, B. M., Wilkinson, I. P., and Riding, J. B.: The lithostratigraphy, biostratigraphy and hydrogeological significance of the mud springs at Templars Firs, Wootton Bassett, Wiltshire: Proceedings of the Geologists' Association, v. 111, no. 3, p. 231–245, 2000.
- Brož, P., and Hauber, E.: Hydrovolcanic tuff rings and cones as indicators for phreatomagmatic explosive eruptions on Mars, 800 *J. Geophysical Res.* 118, 1656–1675, doi:10.1002/jgre.20120, 2013.
- Brož, P., Čadek, O., Hauber, E., and Rossi, A.P.: Scoria cones on Mars: detailed investigation of morphometry based on high-resolution digital elevation models, *J. Geophysical Res.* 120, 1512–1527, doi:10.1002/2015JE004873, 2015a.
- Brož, P., Hauber, E., Platz, T., and Balme, M.R.: Evidence for Amazonian highly viscous lavas in the southern highlands on Mars, *Earth Planet. Sci. Lett.* 415, 200–212, doi:10.1016/j.epsl.2015.01.033, 2015b.
- 805 Brož, P., Hauber, E., Wray, J.J., and Michael, G.: Amazonian volcanism inside Valles Marineris on Mars, *Earth Planet. Sci. Lett.* 473, 122–130, doi:10.1016/j.epsl.2017.06.003, 2017.
- Brož, P., Hauber, E., van de Burgt, I., Špillar, V., and Michael, G.: Subsurface sediment mobilization in the southern Chryse Planitia on Mars, *Journal of Geophysical Research*, 124, 703–720, doi:10.1029/2018JE005868, 2019.
- Brož, P., Krýza, O., Wilson, L., Conway, S.J., Hauber, E., Mazzini, A., Raack, J., Patel, M. R., Balme, M.R., and Sylvest, 810 M.E.: Experimental evidence for lava-like mud flows under Martian surface conditions, *Nat. Geosci.* 13, 403–407, doi:10.1038/s41561-020-0577-2, 2020a.
- Brož, P., Krýza, O., Conway, S.J., Mueller, N.T., Hauber, E., Mazzini, A., Raack, J., Patel, M.R., Balme, M.R., and Sylvest, M.E.: Mud flow levitation on Mars: insights from laboratory simulations. *Earth and Planetary Science Letters* 545, doi:10.1016/j.epsl.2020.116406, 2020b.
- 815 Brož, P., Bernhardt, H., Conway, S. J., Parekh, R.: An overview of explosive volcanism on Mars. *Journal of Volcanology and Geothermal Research*. 409, doi:10.1016/j.jvolgeores.2020.107125, 2021.



- Brož, P., Hauber, E., Conway, S.J., Luzzi, E., Mazzini, A., Noblet, A., Jaroš, J., Fawdon, P., and Markonis, Y.: New Evidence for Sedimentary Volcanism on Chryse Planitia, Mars, *Icarus* 382, 115038, doi:10.1016/j.icarus.2022.115038, 2022a.
- Brož, P., Krýza, O., Conway, S.J., Mazzini, A., Hauber, E., Sylvest, M.E., and Patel, M.R.: Volumetric changes of mud on Mars: evidence from laboratory simulations. In: 53rd lunar and Planetary science conference, #1337, 2022b.
- 820
- Bulanova, I. A., Solovyeva, M. A., Akhmanov, G. G., Khlystov, O. M., and Starovoytov, A. V.: Results of geological and geophysical studies of the Elovsky (Lake Baikal): Proceedings of VII International conference "Marine Research and Education (MARESEDU-2018)" Moscow, 19-22 November 2018, v. 2, no. LLC "PolyPRESS", Tver', 2019, p. 153–154, 2018.
- 825
- Burr, D.M., Brunom B.C., Lanagan, P.D., Glaze, L.S., Jaeger, W.L., Soare, R.J., Wan Bun Tseung, J.-M., Skinner, J.A., Baloga, S.M.: Mesoscale raised rim depressions (MRRDs) on Earth: A review of the characteristics, processes, and spatial distributions of analogs for Mars. *Planetary and Space Science*, 57, 579-596, doi:10.1016/j.pss.2008.11.011, 2009.
- Calvari, S., and Pinkerton, H.: Lava tube morphology on Etna and evidence for lava flow emplacement mechanisms, *Journal of Volcanology and Geothermal Research*, 90, 263-280, doi:10.1016/S0377-0273(99)00024-4, 1999.
- 830
- Carr, M. H.: Stability of streams and lakes on Mars. *Icarus* 56, 476–495, doi:10.1016/0019-1035(83)90168-9, 1983.
- Chan, M.A., Ormo, J., Murchie, S., Okubo, C.H., Komatsu, G., Wray, J.J., McGuire, P., and McGovern, J.A.: Geomorphic knobs of Candor Chasms, Mars: new Mars Reconnaissance Orbiter data and comparisons to terrestrial analogs. *Icarus* 205:138–153, doi:10.1016/j.icarus.2009.04.006, 2010.
- 835
- Ciotoli G. Procesi M., Etiopie G., Fracassi U., and Ventura G.: Influence of tectonics on global scale distribution of geological methane emissions, *Nature Comm.* 11, 2305, doi:10.1038/s41467-020-16229-1, 2020.
- Cita, M. B., Ryan, W. B. F., and Paggi, L.: Prometheus mudbreccia: An example of shale diapirism in the Western Mediterranean Ridge, *Ann. Geol. Pays Hellen.*, 30, 543–570, 1981.
- Clifford, S. M.: A model for the hydrologic and climatic behavior of water on Mars, *J. Geophys. Res.* 98, 10973-1016, doi:10.1029/93JE00225, 1993.
- 840
- Clifford, S. M., Lasue, J., Heggy, E., Boisson, J., McGovern, P., and Max, M. D.: Depth of the Martian cryosphere: Revised estimates and implications for the existence and detection of subpermafrost groundwater, *J. Geophys. Res.*, 115, E07001, doi:10.1029/2009JE003462, 2010.
- Cockell, C.S.: Trajectories of Martian Habitability, *Astrobiology*, 14, 182-203, doi:10.1089/ast.2013.1106, 2014.
- 845
- Conway, S.J., Lamb, M.P., Balme, M.R., Towner, M.C., and Murray, J.B.: Enhanced runout and erosion by overland flow at low pressure and subfreezing conditions: Experiments and application to Mars, *Icarus* 211 (1), 443–457. doi:10.1016/j.icarus.2010.08.0266, 2011.
- Cooper, C.: Mud volcanoes of Azerbaijan visualized using 3D seismic depth cubes: the importance of overpressured fluid and gas instead of non extant diapirs: Abstract Vol. Subsurface Sediment Mobilization Conf, 10-13 September, Ghent, Belgium, p. 71, 2001.
- 850



- Costard, F., Séjourné, A., Jelloun, K., Clifford, S., Lavigne, F., Di Pietro, I., and Souley, S.: Modeling tsunami propagation and the emplacement of thumbprint terrain in an early Mars ocean, *J. Geophys. Res. Planets* 122, doi:10.1002/2016JE005230, 2017.
- 855 Cuřin, V., Brož, P., Hauber, E., and Markonis, Y.: Mud flows in Southwestern Utopia Planitia, Mars, *Icarus*, doi:10.1016/j.icarus.2022.115266, 2023.
- Dapremont, A. M., and Wray, J.J.: Insights into Mars mud volcanism using visible and near-infrared spectroscopy, *Icarus* 359, 114299, doi:10.1016/j.icarus.2020.114299, 2021.
- Davis, P.A., and Tanaka, K.L.: Curvilinear ridges in Isidis Planitia, Mars – The result of mud volcanism? 24th Lunar and Planetary Science, 321–322, 1995.
- 860 Dehn, J., and Sheridan, M.F.: Cinder cones on the Earth, Moon, Mars, and Venus: A computer model. 21st Lunar and Planetary Science Conference, #270, 1990.
- De Toffoli, B., Pozzobon, R., Massironi, M., Mazzarini, F., Conway, S., and Cremonese, G.: Surface expressions of subsurface sediment mobilization rooted into a gas hydrate-rich cryosphere on Mars, *Scientific Reports* 9, 8603, doi:10.1038/s41598-019-45057-7, 2019.
- 865 De Toffoli, B., Massironi, M., Mazzarini, F., and Bistacchi, A.: Rheological and mechanical layering of the crust underneath thumbprint terrains in Arcadia Planitia, Mars. *J. Geophys. Research: Planets* 126 (11), doi:10.1029/2021JE007007, 2021.
- Dickinson, W.R.: *Tectonics and Sedimentation*. SEPM Special Publication, Vol. 22, SEPM Society for Sedimentary Geology, doi:10.2110/pec.74.22, 1974.
- 870 Dimitrov, L.I.: Mud volcanoes—the most important pathway for degassing deeply buried sediments. *Earth Sci. Rev.* 59, 49–76, doi:10.1016/S0012-8252(02)00069-7, 2002.
- Dimitrov, L.I.: Mud volcanoes—a significant source of atmospheric methane. *Geo-Mar Lett* 23, 155–161, doi:10.1007/s00367-003-0140-3, 2003.
- Di Pietro, I., Séjourné, A., Costard, F., Ciazela, J., and Rodríguez, A.P.: Evidence of mud volcanism due to the rapid 875 compaction of martian tsunami deposits in southeastern Acidalia Planitia, Mars, *Icarus* 354, 114096, doi:10.1016/j.icarus.2020.114096, 2021.
- Etiopie, G., Feyzullayev, A., and Baciu, C.L.: Terrestrial methane seeps and mud volcanoes: a global perspective of gas origin. *Mar. Pet. Geol.* 26, 333–344, doi:10.1016/j.marpetgeo.2008.03.001, 2009.
- Etiopie, G., Baciu, C., and Schoell, M.: Extreme methane deuterium, nitrogen and helium enrichment in natural gas from the 880 Homorod seep (Romania), *Chemical Geology*, 280, 89–96, doi:10.1016/j.chemgeo.2010.10.019, 2011.
- Farrand, W.H., Gaddis, L.R., and Keszthlyi, L.: Pitted cones and domes on Mars: Observations in Acidalia Planitia and Cydonia Mensae using MOC, THEMIS, and TES data, *J. Geophys. Res.* 110. doi:10.1029/2004JE002297, 2005.
- Feyzullayev, A. A.: Mud volcanoes in the South Caspian basin: Nature and estimated depth of its products, *Natural Science* 4(7), doi:10.4236/ns.2012.47060, 2012.



- 885 Formisano, V., Atreya, S., Encrenaz, T., Ignatiev, N., Giuranna, M.: Detection of Methane in the Atmosphere of Mars, *Science*, 306, 1758-1761, doi:10.1126/science.1101732, 2004.
- Franchi, F., Rossi, A.P., Pondrelli, M., and Cavalazzi, B.: Geometry, stratigraphy and evidences for fluid expulsion within Crommelin crater deposits, Arabia Terra, Mars. *Planetary and Space Science*, 92, 34-48, doi:10.1016/j.pss.2013.12.013, 2014.
- 890 Frey, H.M., Lowry, B.L., and Chase, S.A.: Pseudocraters on Mars, *J. Geophys. Res.* 84, 8075–8086, doi:10.1029/JB084iB14p08075, 1979.
- Frey, H.V., Roark, J.H., Shockey, K.M., Frey, E.L., and Sakimoto, S.E.H.: Ancient lowlands on Mars, *Geophysical Research Letters*, 29, 1384. doi:10.1029/2001GL013832, 2002.
- Fryer, P.: Serpentinite Mud Volcanism: Observations, Processes, and Implications. *Annual Review of Marine Science*, 4, 345-373, doi:10.1146/annurev-marine-120710-100922, 2012.
- 895 Fryer et al.: Mariana serpentinite mud volcanism exhumes subducted seamount materials: implications for the origin of life. *Phil. Trans. R. Soc. A378*: 20180425, doi:10.1098/rsta.2018.0425, 2020.
- Gallagher, C., Balme, M., Soare, R., Conway, S.J.: Formation and degradation of chaotic terrain in the Galaxias regions of Mars: implications for near-surface storage of ice, *Icarus* 309, 69-83, doi: 10.1016/j.icarus.2018.03.002. 2018.
- 900 Genova, A.: ORACLE: A mission concept to study Mars' climate, surface and interior, *Acta Astronautica* 166, 317–329, doi: 10.1016/j.actaastro.2019.10.006, 2020.
- Gill, W. D., and Kuenen, P. H.: Sand volcanoes on slumps in the Carboniferous of County Clare, Ireland, *Quarterly Journal of the Geological Society*, 113, 1-4, 441-460, doi:10.1144/GSL.JGS.1957.113.01-04.19, 1957.
- Giuranna, M., Viscardy, S., Daerden, F., Neary, L., Etiope, G., Oehler, D., Formisano, V., Aronica, A., Wolkenberg, P., Aoki, S., Cardesin-Moinelo, A., Marin-Yaseli de la Parra, J., Merritt, D., Amoroso, M.: Independent confirmation of a methane spike on Mars and a source region east of Gale Crater. *Nat. Geosci.* 12, 326–332, doi:10.1038/s41561-019-0331-9, 2019.
- 905 Goldspiel, J.M., and Squyres, S.W.: Ancient aqueous sedimentation on Mars. *Icarus*, 89, 392-410, doi:10.1016/0019-1035(91)90186-W, 1991.
- 910 Guest, J. E., Butterworth, P. S., and Greeley, R.: Geological observations in the Cydonia Region of Mars from Viking, *J. Geophys. Res.*, 82 (28), 4111–4120, doi:10.1029/JS082i028p04111, 1977.
- Golombek, M. P. and Phillips, R. J.: Mars Tectonics. Chapter 5, *in* *Planetary Tectonics*, (eds. T.R. Watters and R.A. Schultz), Cambridge University Press, Cambridge, UK, pp. 183-232.
- Goossens, S., Sabaka, T. J., Genova, A., Mazarico, E., Nicholas, J. B., and Neumann, G. A.: Evidence for a low bulk crustal density for Mars from gravity and topography, *Geophys. Res. Lett.* 44, 7686–7694, doi:10.1002/2017GL074172, 2017.
- Graue, K.: Mud volcanoes in deepwater Nigeria, *Mar. Petrol. Geol.* 17, 959–974, doi:10.1016/S0264-8172(00)00016-7, 2000.
- Grenfell, J.L., Wunderlich, F., Sinnhuber, M., Herbst, K., Lehmann, R., Scheucher, M., Gebauer, S., Arnold, G., and Rauer H.: Atmospheric processes affecting methane on Mars, *Icarus*, 382, 114940, doi:10.1016/j.icarus.2022.114940, 2022.



- Grotzinger, J.P., et al.: A habitable fluvio-lacustrine environment at Yellowknife Bay, Gale Crater, Mars, *Science* 343, 1242777, doi:10.1126/science.1242777, 2013.
- Grott, M., Baratoux, D., Hauber, E., Sautter, V., Mustard, J., Gasnault, O., Ruff, S.W., Karato, S.-I., Debaille, V., Knapmeyer, M., Sohl, F., Van Hoolst, T., Breuer, D., Morschhauser, A., Toplis, M.J.: Long-term evolution of the Martian crust-mantle system, *Space Sci. Rev.*, doi:10.1007/s11214-012-9948-3, 2013.
- Guliyev, I. S., and Feizullayev, A. A.: All about Mud volcanoes, Nafta Press, Baku, 52 p, 1997.
- 925 Haberle, R.M., McKay, C.P., Schaeffer, J., Cabrol, N.A., Grin, E.A., Zent, A.P., and Quinn, R.: On the possibility of liquid water on present-day Mars, *J. Geophys. Res.*, 106(E10), 23317– 23326, doi:10.1029/2000JE001360, 2001.
- Haberle, R.M., Catling, D.C., Carr, M.H., and Zahnle, K. J.: The Early Mars Climate System. In: Haberle, R. M. et al. (Eds.) *The Atmosphere and Climate of Mars*, pp. 526–568. Cambridge University Press, doi:doi:10.1017/9781139060172.017, 2017.
- 930 Harrison, K.P., and Grimm, R.E.: Tharsis recharge: a source of groundwater for Martian outflow channels. *Geophysical Research Letters* 31(14), L14703, doi:10.1029/2004GL020502, 2004.
- Harrison, K.P., and Chapman, M.G.: Evidence for ponding and catastrophic floods in central Valles Marineris, Mars, *Icarus* 198, 351–364. doi:10.1016/j.icarus.2008.08.003, 2008.
- Hartmann, W. K., and Neukum, G.: Cratering chronology and the evolution of Mars. *Space Science Reviews*, 96(1/4), 165–
935 194, doi:10.1023/A:1011945222010, 2001.
- Hemmi, R., and Miyamoto, H.: Distribution, morphology, and morphometry of circular mounds in the elongated basin of northern Tera Sirenum, Mars., *Progress in Earth and Planetary Science* 4, 26, doi:10.1186/s40645-017-0141-x, 2017.
- Hemmi, R., and Miyamoto, H.: High-resolution topographic analyses of mounds in southern Acidalia Planitia, Mars: Implications for possible mud volcanism in submarine and subaerial environments, *Geosciences* 8, 152; 940 doi:10.3390/geosciences8050152. 2018.
- Hodson, A. J., Nowak, A., Hornum, M. T., Senger, K., Redeker, K., Christiansen, H. H., and Marca, A.: Sub-permafrost methane seepage from open-system pingos in Svalbard, *The Cryosphere*, 14, 3829–3842, doi:10.5194/tc-14-3829-2020, 2020.
- Hon, K., Kauahikaua, J., Denlinger, R., and Mackay, K.: Emplacement and inflation of pahoehoe sheet flows: observation and measurements of active lava flows on Kilauea Volcano, Hawaii. *Geol. Soc. Am. Bull.* 106, 351–370, doi:10.1130/0016-7606(1994)106%3C0351:EAIOPS%3E2.3.CO;2, 1994.
- Huang, H.; Liu, J.; Wang, X.; Chen, Y.; Zhang, Q.; Liu, D.; Yan, W.; Ren, X.: The Analysis of Cones within the Tianwen-1 Landing Area: *Remote Sens.*, 14, 2590, doi:10.3390/rs14112590, 2022.
- Hurst, A., Cartwright, J. A., Huuse, M., and Duranti, D.: Extrusive sandstones (extrudites): a new class of stratigraphic trap? 950 *Geological Society, London, Special Publications*, 254, 1, p. 289, 2006.



- Inan, S., Yalcin, M. N., Guliyev, I. S., Kuliev, K., and Feyzullayev, A. A.: Deep Petroleum occurrences in the Lower Kura Depression, South Caspian Basin, Azerbaijan: an organic geochemical and basing modeling study, *Marine & Petroleum Geology* 14, 731–762, doi:10.1016/s0264-8172(97)00058-5, 1997.
- Ivanov, M.A., Hiesinger, H., Erkeling, G., Hielscher, F., and Reiss, D.: Major episodes of geologic history of Isidis Planitia on Mars, *Icarus* 218 (1), 24–46, doi:10.1016/j.icarus.2011.11.029, 2012.
- Ivanov, M.A., Hiesinger, H., Erkeling, G., and Reiss, D.: Mud volcanism and morphology of impact craters in Utopia Planitia on Mars: Evidence for the ancient ocean, *Icarus* 228, 121–140, doi:10.1016/j.icarus.2013.09.018, 2014.
- Ivanov, M.A., and Hiesinger, H.: The Acidalia Mensa region on Mars: A key element to test the Mars ocean hypothesis, *Icarus* 349, 113874, doi:10.1016/j.icarus.2020.113874, 2020.
- Jakosky, B.M.: Atmospheric Loss to Space and the History of Water on Mars, *Annual Review of Earth and Planetary Sciences*, 49:1, 71-93, doi:10.1146/annurev-earth-062420-052845, 2021.
- Jakubov, A. A., AliZade, A. A., and Zeinalov, M. M.: Mud volcanoes of the Azerbaijan SSR: Atlas (in Russian), Azerbaijan Academy of Sciences, Baku, 1971.
- Jolly, R. J. H., and Lonergan, L.: Mechanism and control on the formation of sand intrusion, *Journal of the Geological Society* 159, 5, 605–617, doi:10.1144/0016-764902-025, 2002.
- Jones, E., Caprarelli, G., and Osinski, G. R.: Insights into complex layered ejecta emplacement and subsurface stratigraphy in Chryse Planitia, Mars, through an analysis of THEMIS brightness temperature data. *Journal of Geophysical Research: Planets*, 121, 986–1015. doi:10.1002/2015JE004879, 2016.
- Jöns, H.-P.: Late sedimentation and late sediments in the northern lowlands on Mars, *Lunar and Planetary Science XVI*, 414–415 (Abstract), 1985.
- Karagoz, O., Kenkmann, T., and Wulf, G.: Insights into the subsurface structure of wrinkle ridges on Mars. *Earth and Planetary Science Letters*, 595, 117759doi: 10.1016/j.epsl.2022.117759, 2022.
- Kargel, J.S., Tanaka, K.L., Baker, V.R., Komatsu, G., and MacAyeal, D.R.: Formation and dissociation of clathrate hydrates on Mars: Polar caps, Northern plains, and Highlands. 31st LPSC, Abs. # 1891, 2000.
- Khlystov, O., Poort, J., Mazzini, A., Akhmanov, G.G., Minami, H., Hachikubo, A., Khabuev, A., Kazakov., A.V., De Batist, M., Naudts, L., Chensky, A.G., Vorobeva, S.: Shallow-rooted mud volcanism in Lake Baikal. *Marine and Petroleum Geology* 102, doi:10.1016/j.marpetgeo.2019.01.005, 580-589, 2019.
- Kite, E. S.; Hovius, N., Hillier, J. K., and Besserer, J.: Candidate Mud Volcanoes in the Northern Plains of Mars, *American Geophysical Union, Fall Meeting 2007*, abstract id.V13B-1346, 2007.
- Kite, E. S., Williams, J.-P., Lucas, A., and Aharonson, O.: Low palaeopressure of the martian atmosphere estimated from the size distribution of ancient craters, *Nat. Geosci.* 7, 335–339, doi:10.1038/ngeo2137, 2014.
- Knapmeyer, M., Oberst, J., Hauber, E., Wählisch, M., Deuchler, C., and Wagner, R.: Working models for spatial distribution and level of Mars' seismicity, *J. Geophys. Res.*, 111, E11006, doi:10.1029/2006JE002708, 2006.



- Knittel, K., and Boetius, A.: Anaerobic Oxidation of Methane: Progress with an Unknown Process. *Annual Review of Microbiology*, 63, 311-334, doi:10.1146/annurev.micro.61.080706.093130, 2009.
- Knutsen, E.W., Villanueva, G.L., Liuzzi, G., Crismani, M.M.J., Mumma, M.J., Smith, M.D., Vandaele, A.C., Aoki, S., Thomas, I. R., Daerden, F., Viscardy, S., Erwin, J.T., Trompet, L., Neary, L., Ristic, B., Lopez-Valverde, M.A., Lopez-Moreno, J.J., Patel, M.R., Karatekin, O., and Bellucci, G.: Comprehensive investigation of Mars methane and organics with ExoMars/NOMAD, *Icarus*, 357, 114266, doi:10.1016/j.icarus.2020.114266, 2021.
- 990 Kokoschka, S., Dreier, A., Romoth, K., Taviani, M., Schäfer, N., Reitner, J. and Hoppert, M.: Isolation of Anaerobic Bacteria from Terrestrial Mud Volcanoes (Salse di Nirano, Northern Apennines, Italy), *Geomicrobiology Journal*, 32:3-4, 355-364, doi:10.1080/01490451.2014.940632, 2015.
- Komatsu, G. A possible mud volcano field in Chryse Planitia, Mars, Abstract, EPSC2010-131, European Planetary Science Congress, Rome, 2010.
- 995 Komatsu, G., and Baker V.R.: Paleohydrology and flood geomorphology of Ares Vallis. *J. Geophys. Res.*, 102, 4151-4160, doi.org/10.1029/96JE02564, 1997.
- Komatsu, G., and Ori, G.G.: Exobiological implications of potential sedimentary deposits on Mars, *Planet. Space Sci.* 48/11, 1043–1052, doi:10.1016/S0032-0633(00)00078-7, 2000.
- Komatsu, G., Ori, G.G., Cardinale, M., Dohm, J.M., Baker, V.R., Vaz, D.A., Ishimaru, R., Namiki, N., and Matsui, T., Roles of methane and carbon dioxide in geological processes on Mars. *Planet. Space Sci.* 59, 169–181, doi:10.1016/j.pss.2010.07.002, 2011.
- 1000 Komatsu, G., Ishimaru, R., Miyake, N., Ohno, S., and Matsui, T.: Astrobiological Potential of Mud Volcanism on Mars. #1085, *Lunar and Planet. Sci. Conf.* 45th, 2014.
- Komatsu, G., Okubo, C.H., Wray, J.J., Ojha, L., Cardinale, M., Murana, A., Orosei, R., Chan, M.A., Ormö, J., and Gallagher, R.: Small edifice features in Chryse Planitia, Mars: Assessment of a mud volcano hypothesis. *Icarus* 268, 56–75, doi:10.1016/j.icarus.2015.12.032, 2016.
- 1005 Komatsu, G., Ishimaru, R., Kawai, K., Miyake, N., Kobayashi, M., Sakuma, H., and Matsui, T.: Sedimentary records of ancient mud volcanism: How do we identify mud volcanoes in the stratigraphy of Mars? *Lunar and Planet. Sci. Conf.* 50th, 2019.
- 1010 Komatsu, G., and Brož, P.: Southern Chryse Planitia on Mars as a potential landing site: investigation of hypothesized sedimentary volcanism. #1164, *Lunar and Planet. Sci. Conf.* 52th, 2021.
- Kopf, A.J.: Significance of mud volcanism. *Rev. Geophys.* 40 (2), 2–52, 2002.
- Korablev, O., Vandaele, A.C., Montmessin, F. et al. No detection of methane on Mars from early ExoMars Trace Gas Orbiter observations, *Nature* 568, 517–520, doi:10.1038/s41586-019-1096-4, 2019.
- 1015 Kossacki, K. J., Markiewicz, W. J., Smith, M. D., Page, D., and Murray, J.: Possible remnants of a frozen mud lake in southern Elysium, Mars. *Icarus* 181, 363–374, 2006.



- Krasnopolsky, V.A., Maillard, J.P., and Owen, T.O.: Detection of methane in the martian atmosphere: evidence for life?, *Icarus*, 172, 537-547, doi:10.1016/j.icarus.2004.07.004, 2004.
- 1020 Kumar, P.S., Krishna, N., Prasanna Lakshmi, K.J., Raghukanth, S.T.G., Dhabu, A., and Platz, T.: Recent seismicity in Valles Marineris, Mars: Insights from young faults, landslides, boulder falls and possible mud volcanoes. *Earth Planet. Sci. Lett.* 505, 51–64. doi:10.1016/j.epsl.2018.10.008, 2019.
- Lee, D.-H., Kim, J.-H., Lee, Y.M., Kim, J.-H., Jin, Y.K., Paull, C., Ryu, J.-S. and Shin, K.-H.: Geochemical and Microbial Signatures of Siboglinid Tubeworm Habitats at an Active Mud Volcano in the Canadian Beaufort Sea. *Front. Mar. Sci.* 8:656171, doi:10.3389/fmars.2021.656171, 2021.
- 1025 Lefèvre, F. and Forget F.: Observed variations of methane on Mars unexplained by known atmospheric chemistry and physics. *Nature* 460:720–72, doi:10.1038/nature08228, 2009.
- Li, J., Beghein, C., Lognonné, P., McLennan, S. M., Wiczeorek, M., Panning, M., et al.: Different Martian Crustal Seismic Velocities across the Dichotomy Boundary from Multi-Orbiting Surface Waves. *Geophysical Research Letters* 49, e2022GL101243, doi:10.1029/2022GL101243, 2022.
- 1030 Lin, Y. (林雨), Zhao, J. (赵健楠), Wang, L. (王乐), Huang, J. (黄俊), Zhang, L. (张良), Xiao, L. (肖龙): Evaluation of small-sized mounds formation mechanisms in China's Zhurong landing region, *Icarus*, 389, 115256, doi:10.1016/j.icarus.2022.115256, 2023.
- Liu, J.; Li, C.; Zhang, R.; Rao, W.; Cui, X.; Geng, Y.; Jia, Y.; Huang, H.; Ren, X.; Yan, W.; et al.: Geomorphic Contexts and Science Focus of the Zhurong Landing Site on Mars: *Nat. Astron.* 6, 65–71, doi:10.1038/s41550-021-01519-5, 2021.
- 1035 Lucchitta, B.K.: Young volcanic deposits in the Valles Marineris, Mars? *Icarus* 86, 476–509, doi:10.1016/0019-1035(90)90230-7, 1990.
- Lucchitta, B. K., Ferguson, H. M., and Summers, C.: Sedimentary deposits in the Northern Lowland Plains, Mars, *J. Geophys. Res.*, 91 (B13), E166– E174, doi:10.1029/JB091iB13p0E166, 1986.
- Mahaney, W.C., Milner, M.W., Netoff, D.I., Malloch, D., Dohm, J.M., Baker, V.R., Miyamoto, H., Hare, T.M., and Komatsu, G., Ancient wet aeolian environments on Earth: clues to presence of fossil/live microorganisms on Mars. *Icarus*, 171, 39-53, 2004.
- 1040 Malin, M. C. and Edgett, K.S.: Sedimentary rocks of early Mars, *Science* 290, 1927–1937, doi:10.1126/science.290.5498.1927, 2000.
- Max, M.D., and Clifford, S.M.: Initiation of Martian outflow channels: related to the dissociation of gas hydrate? *Geol. Res. Lett.* 28 (9), 1787–1790, 2001.
- 1045 Max, M.D., Clifford, S.M., and Johnson, A.H.: Hydrocarbon system analysis for methane hydrate exploration on Mars. *Am. Assoc. Petrol. Geol. Memoir* 101, 99–114. doi:10.1306/13361573M1013546, 2013.
- Mazzini, A., Svensen, H., Planke, S., Guliyev, I., Akhmanov, G. G., Fallik, T., and Banks, D.: When mud volcanoes sleep: Insight from seep geochemistry at the Dashgil mud volcano, Azerbaijan: *Marine and Petroleum Geology* 26, 9, p. 1704–1715, 2009.
- 1050



- Mazzini A., Etiope, G., and Svensen H.: A new hydrothermal scenario for the 2006 Lusi eruption, Indonesia. Insights from gas geochemistry. *Earth Plan. Sci. Lett.*, 317–318, 305–318, 2012.
- Mazzini A., and Etiope, G., Mud volcanism: an updated review. *Earth Sci. Rev.*, 168, 81–112. doi:10.1016/j.earscirev.2017.03.001, 2017.
- 1055 Mazzini, A., Akhmanov, G., Manga, M., Sciarra, A., Huseynova, A., Huseynov, A., and Guliyev, I.: Explosive mud volcano eruptions and rafting of mud breccia blocks: *Earth and Planetary Science Letters* 555, 116699, 2021.
- McCauley, C. A., White, D. M., Lilly, M. R., and Nyman, D. M.: A comparison of hydraulic conductivities, permeabilities and infiltration rates in frozen and unfrozen soils. *Cold Regions Science and Technology*, 34(2), 117–125. doi:10.1016/S0165-232X(01)00064-7, 2002.
- 1060 McGill, G.E., Buried topography of Utopia, Mars: Persistence of a giant impact depression. *Journal of Geophysical Research – Solid Earth* 94, B3. doi:10.1029/JB094iB03p02753, 1989.
- McGowan, E.: Spatial distribution of putative water related features in Southern Acidalia/Cydonia Mensae, Mars, *Icarus* 202:78–89, 2009.
- McGowan, E.M. and McGill, G.E.: The Utopia/Isidis overlap; possible conduit for mud volcanism. 41st Lunar and Planetary Science Conference, Abs. # 1070. 2010.
- 1065 McNeil, J.D., Fawdon, P., Balme, M.R., and Coe, A.L.: Morphology, morphometry and distribution of isolated landforms in southern Chryse Planitia, Mars, *Journal of Geophysical Research: Planets*, 126, e2020JE006775, doi:10.1029/2020JE006775, 2021.
- Meresse, S., Costard, F., Mangold, N., Masson, P., Neukum, G., and the HRSC Co-I Team, Formation and evolution of the chaotic terrains by subsidence and magmatism: *Hydraotes Chaos, Mars, Icarus* 194, 487–500. doi:10.1016/j.icarus.2007.10.023, 2008.
- 1070 Miall, A.D.: *Principles of Sedimentary Basin Analysis*. Springer, New York, 1984.
- Michalski, J.R., Cuadros, J., Niles, P.B., Parnell, J., Rogers, A.D., and Wright, S.P.: Groundwater activity on Mars and implications for a deep biosphere, *Nature Geoscience* 6:133–138, doi:10.1038/ngeo1706, 2013.
- 1075 Milliken, R. E., Grotzinger, J. P., and Thomson, B. J.: Paleoclimate of Mars as captured by the stratigraphic record in Gale Crater, *Geophys. Res. Lett.* 37, L04201, doi:10.1029/2009GL041870, 2010.
- Milkov, A. V., Sassen, R., Apanasovich, T. V., and Dadashev, F. G.: Global gas flux from mud volcanoes: A significant source of fossil methane in the atmosphere and the ocean. *Geophys. Res. Lett.*, 30, 1037, doi:10.1029/2002GL016358, 2003.
- 1080 Miyake, N., Ishimaru, R., Komatsu, G., Matsui, T.: Characterization of archaeal and bacterial communities thriving in methane-seeping on-land mud volcanoes, Niigata, Japan. *International Microbiology*, <https://doi.org/10.1007/s10123-022-00288-z>, in press.
- Moscardelli, L., Dooley, T., Dunlap, D., Jackson, M., and Wood, L.: Deep-water polygonal fault systems as terrestrial analogs for large-scale Martian polygonal terrains, *GSA Today* volume 22, no. 8, 4-9, doi:10.1130/GSATG147A.1, 2012.



- Möhlmann, D.T.F.: Water in the upper martian surface at mid- and low-latitudes: presence, state, and consequences. *Icarus*, 168, 318-323, doi:10.1016/j.icarus.2003.11.008, 2004.
- Mueller, K., and Golombek, M.: Compressional Structures on Mars. *Annual Review of Earth and Planetary Sciences*, 32, 435-464, doi:10.1146/annurev.earth.32.101802.120553, 2004.
- Mumma, M. J., Villanueva, G. L., Novak, R. E., Hewagama, T., Bonev, B. P., DiSanti, M. A., Mandell, A. M., Smith, M. D.: Strong Release of Methane on Mars in Northern Summer 2003. *Science*, 323, 1041-1045, doi:10.1126/science.1165243, 2009.
- Mustard, J.F., Ehlmann, B.L., Murchie, S.L., Poulet, F., Mangold, N., Head, J.W., Bibring, J.-P., and Roach, L.H.: Composition, morphology and stratigraphy of Noachian crust around the Isidis Basin, *J Geophys Res* 114, doi:10.1029/2009JE003349, 2009.
- Oberst, J., Wickhusen, K., Gwinner, K., Hauber, E., et al.: Planetary polar explorer – the case for a next-generation remote sensing mission to low Mars orbit, *Exp. Astron.*, doi:10.1007/s10686-021-09820-x, 2022.
- Oehler, D.Z. and Allen, C.C.: Mud volcanoes in the martian lowlands: potential windows to fluid-rich samples from depth. 40th Lunar and Planetary Science Conference, Abs. # 1034, 2009.
- Oehler, D.Z. and Allen, C.C.: Evidence for pervasive mud volcanism in Acidalia Planitia, Mars. *Icarus* 208, 636–657, 2010.
- Oehler, D.Z. and Allen, C.C.: Habitability of a large ghost crater in Chryse Planitia, Mars. In *International Conference: Exploring Mars Habitability*, Lisbon, Portugal, 2011.
- Oehler, D.Z., and Allen, C.C.: Giant polygons and mounds in the lowlands of Mars: Signatures of an ancient ocean? *Astrobiology* 12 (6), doi:10.1089/ast.2011.0803, 2012a.
- Oehler, D.Z., and Allen, C.C.: Focusing the search for biosignatures on Mars: Facies prediction with an example from Acidalia Planitia, *SEPM Special Publ. No. 102*, 183–194, in *Sedimentary Geology of Mars*, Society for Sedimentary Geology, 2012b.
- Oehler D., and Etiope G.: Methane Seepage on Mars: Where to Look and Why, *Astrobiology* 17(12), 1233-1264, doi: 10.1089/ast.2017.1657, 2017.
- Oehler D., and Etiope G.: Methane on Mars: subsurface sourcing and conflicting atmospheric measurements. In: Soare R., Conway S., Williams J.P., Oehler D. (Eds.), *Mars Geological Enigmas: From the Late Noachian Epoch to the Present Day*, 1st Ed., pp. 149-174. Elsevier, 2021.
- Oehler, D.Z., Salvatore, M., Etiope, G., and Allen, C.C.: Focusing the search for organic biosignatures on Mars. 52nd Lunar and Planetary Science Conference, Abs. # 1353, 2021.
- Okubo, C.H.: Bedrock Geologic and Structural Map through the Western Candor Colles Region of Mars, U.S. Geological Survey Scientific Investigations Map 3309, 2014.
- Okubo, C.H.: Morphologic evidence of subsurface sediment mobilization and mud volcanism in Candor and Coprates Chasmata, Valles Marineris, Mars, *Icarus* 269, 23–37, doi:10.1016/j.icarus.2015.12.051, 2016.



- Orgel, C., Hauber, E., Skinner J.A. Jr., van Gasselt, S., Ransdale, J., Balme, M., Séjourné, A., and Keresz-Turi, A.: Distribution, origin and evolution of hypothesized mud volcanoes, thumbprint terrain and giant polygons in Acidalia, Utopia and Arcadia Planitiae: Implications for sedimentary processes in the northern lowlands of Mars. 46th Lunar and Planetary Science Conference, Abs. # 1862, 2015.
- 1120
- Orgel, C., Hauber, E., Van Gasselt, S., Reiss, D., Johnsson, A., Ramsdale, J.D., Smith, I., Swirad, Z.M., Séjourné, A., Wilson, J.T., Balme, M.R., Conway, S.J., Costard, F., Eke, V.R., Gallagher, C., Kereszturi, A., Losiak, A., Massey, R.J., Platz, T., Skinner, J.A., and Teodoro, L.R.F., Grid mapping the northern plains of Mars: A new overview of recent water- and ice-related landforms in Acidalia Planitia, *J. Geophys. Research: Planets* 124 (2), 454–482, doi:10.1029/2018JE005664, 2019.
- 1125
- Ori, G.G., Marinangeli, L., Komatsu, G., 2000. Gas (methane?)-related features on the surface of Mars and subsurface reservoirs. In: Proceedings of the Lunar Planetary Science Conference XXXI, #1550, abstract [CD-ROM].
- Ori, G.G., Komatsu, G., Ormö, J., Marinangeli, L., 2001. Subsurface models for the formation of mound-like morphologies on Mars. In: Proceedings of the Lunar Planetary Science Conference XXXII, #1539, abstract [CD-ROM].
- 1130
- Ormö, J., Komatsu, G., Chan, M.A., Beitler, B., and Parry, W.T.: Geological features indicative of processes related to the hematite formation in Meridiani Planum and Aram Chaos, Mars: A comparison with diagenetic hematite deposits in southern Utah, USA, *Icarus* 171, 295–316, doi:10.1016/j.icarus.2004.06.001, 2004.
- Page, D.P.: A candidate methane-clathrate destabilization event on Mars: a model for sub-millennial-scale climatic change on Earth, *Gondwana Res.* 59, 43–56, doi:10.1016/j.gr.2018.03.010, 2018.
- 1135
- Pajola, M., Rossato, S., Baratti, E., Mangili, C., Mancarella, F., McBride, K., and Coradini, M.: The Simud–Tiu Valles hydrologic system: A multidisciplinary study of a possible site for future Mars on-site exploration, *Icarus*, 268, 355–381, doi:10.1016/j.icarus.2015.12.049, 2016.
- Pan, L., Ehlmann, B. L., Carter, J., and Ernst, C. M.: The stratigraphy and history of Mars' northern lowlands through mineralogy of impact craters: A comprehensive survey, *J. Geophys. Res. Planets* 122, 1824–1854, doi:10.1002/2017JE005276, 2017.
- 1140
- Parfitt, E.A., and Wilson, L.: *Fundamentals of Physical Volcanology*, 256 pp. Blackwell, Oxford, U. K., 2008.
- Pavlov, A.A., McLain, H.L., Glavin, D.P., Roussel, A., Dworkin, J.P., Elsila, J.E., and Yocum, K.M.: Rapid Radiolytic Degradation of Amino Acids in the Martian Shallow Subsurface: Implications for the Search for Extinct Life, *Astrobiology* 1099–1115, doi:10.1089/ast.2021.0166, 2022.
- 1145
- Pedersen, G. B. M.: Frozen Martian lahars? Evaluation of morphology, degradation and geologic development in the Utopia–Elysium transition zone, *Planetary and Space Science* 85, 59–77, doi:10.1016/j.pss.2013.05.020, 2013.
- Pfeffer, W., and Humphrey, N.: Formation of ice layers by infiltration and refreezing of meltwater. *Annals of Glaciology*, 26, 83–91. doi:10.3189/1998AoG26-1-83-91, 1998.



- 1150 Plümper, O., King, H.E., Geisler, T., Liu, Y., Pabst, S., Savov, I.P., Rost, D., and Zack, T.: Subduction zone forearc serpentinites as incubators for deep microbial life. *Proceedings of the National Academy of Sciences*, 114, 4324–4329, doi:10.1073/pnas.1612147114, 2017.
- Polteau, S., Mazzini, A., Galland, O., Planke, S., and Malthe-Sorensen, A.: Saucer-shaped intrusions: Occurrences, emplacement and implications, *Earth and Planetary Science Letters* 266, 1–2, 195–204, doi:10.1016/j.epsl.2007.11.015, 2008.
- 1155 Pondrelli, M., Rossi, A.P., Ori, G.G., Van Gasselt, S., Praeg, D., and Ceramicola, S.: Mud volcanoes in the geologic record of Mars: the case of Firsoff Crater, *Earth Planet Sci Lett* 304:511–519, doi:10.1016/j.epsl.2011.02.027, 2011.
- Pons, M.-L., Quitté, G., Fujii, T., Rosing, M.T., Reynard, B., Moynier, F., Douchet, C., and Albarède, F.: Early Archean serpentine mud volcanoes at Isua, Greenland, as a niche for early life. *Proc.Natl Acad. Sci. USA* 108, 17 639–17 643, doi:10.1073/pnas.1108061108, 2011.
- 1160 Procesi, M., Ciotoli, G., Mazzini, A., and Etiope, G.: Sediment-Hosted Geothermal Systems: review and first global mapping. *Earth Sci. Rev.*, 192, 529–544, doi:10.1016/j.earscirev.2019.03.020, 2019.
- Quantin, C., Flahut, J., Clenet, H., Allemand, P., and Thomas, P.: Composition and structures of the subsurface in the vicinity of Valles Marineris as revealed by central uplifts of impact craters, *Icarus* 221:436–452, doi:10.1016/j.icarus.2012.07.031 2012.
- 1165 Rampe, E.B., Blake, D.F., Bristow, T.F., Ming, D.W., Vaniman, D.T., Morris, R.V., Achilles, C.N., Chipera, S.J., Morrison, S.M., Tu, V.M., Yen, A.S., Castle, N., Downs, G.W., Downs, R.T., Grotzinger, J.P., Hazen, R.M., Treiman, A.H., Peretyazhko, T.S., Des Marais, D.J., Walroth, R.C., Craig, P.I., Crisp, J.A., Lafuente, B., Morookian, J.M., Sarrazin, P.C., Thorpe, M.T., Bridges, J.C., Edgar, L.A., Fedo, C.M., Freissinet, C., Gellert, R., Mahaffy, P.R., Newsom, H.E., Johnson, J.R., Kah, L.C., Siebach, K.L., Schieber, J., Sun, V.Z., Vasavada, A.R., Wellington, D., and Wiens, R.C.:
- 1170 Mineralogy and geochemistry of sedimentary rocks and eolian sediments in Gale crater, Mars: A review after six Earth years of exploration with Curiosity, *Geochemistry*, 80, 125605, doi:10.1016/j.chemer.2020.125605, 2020.
- Rice, J.W., Jr., and Edgett, K.S.: Catastrophic flood sediments in Chryse basin, Mars, and Quincy basin, Washington: application of sandar facies model, *J Geophys Res* 102:4185–4200, doi:10.1029/96JE02824, 1997.
- Rodríguez, J.A.P., Tanaka, K.L., Kargel, J.S., Dohm, J.M., Kuzmin, R., Fairén, A.G., Sasaki, S., Komatsu, G., Schulze-
- 1175 Makuch, D., and Jianguo, Y.: Formation and disruption of aquifers in southwestern Chryse Planitia, Mars, *Icarus* 191 (2), 545–567, doi:10.1016/j.icarus.2007.05.021, 2007.
- Rodríguez, J. A. P., et al.: Tsunami waves extensively resurfaced the shorelines of a receding, early Martian ocean, *Nature Scientific Reports* 6, 25106, doi:10.1038/srep25106, 2016.
- Rodríguez, J.A.P., Kargel, J.S., Oehler, D.Z., Crown, D.A., Baker, V.R., and Komatsu, G.: Potential cryospheric mud
- 1180 volcanism in the northern plains of Mars, Geologic and astrobiological implications. 50th Lunar and Planetary Science Conference, Abs. # 2580, 2019.



- Rowland, S. K., Harris, A., L., and Garbeil, H.: Effects of Martian conditions on numerically modeled, cooling-limited, channelized lava flows. *Journal of Geophysical Research – Planets* 109, Issue E10, doi:10.1029/2004JE002288, 2004.
- 1185 Rubin, D.M., Faïren, A., Martínez-Frías, J., Frydenvang, J., Gasnault, O., Galfenbaum, G., Goetz, W., Grotzinger, J.P., Le Mouélic, S., Mangold, N., Newsom, H., Oehler, D.Z., Rapin, W., Schieber, J., and Weins, R.C.: Fluidized-sediment pipes in Gale Crater, Mars, and possible Earth analogs, *Geology* 45, 7–10, doi:10.1130/G38339.1, 2016.
- Russell, P. S.: Elysium-Utopia flows as mega-lahars: A model of dike intrusion, cryosphere cracking, and water-sediment release, *Journal of Geophysical Research* 108 (E6), doi:10.1029/2002je001995, 2003
- 1190 Salvatore, M.R. and Christensen, P.R.: Evidence for widespread aqueous sedimentation in the northern plains of Mars, *Geology* 42, 423–426, doi:10.1130/G35319.1, 2014a.
- Salvatore, M.R. and Christensen, P.R.: On the origin of the Vastitas Borealis Formation in Chryse and Acidalia Planitiae, Mars, *J Geophys Res: Planets* 119, 2437–2456, doi:10.1002/2014JE004682, 2014b.
- Skinner Jr., J.A., and Tanaka, K.L.: Evidence for and implications of sedimentary diapirism and mud volcanism in the southern Utopia highland–lowland boundary plain, Mars. *Icarus* 186 (1), 41–59, doi:10.1016/j.icarus.2006.08.013, 2007.
- 1195 Skinner, J.A., Jr., and Mazzini, A.: Martian mud volcanism: terrestrial analogs and implications for formational scenarios, *Marine and Petroleum Geology* 26:1866–1878, doi:10.1016/j.marpetgeo.2009.02.006, 2009.
- Smrekar, S.E., Lognonné, P., Spohn, T. et al.: Pre-mission InSights on the Interior of Mars. *Space Sci Rev* 215, 3, doi:10.1007/s11214-018-0563-9, 2019.
- Solomon, S.C.: On volcanism and thermal tectonics on one-plate planets. *Geophys. Res. Lett.*, 5: 461–464, doi:10.1029/GL005i006p00461, 1978.
- 1200 Stamenković, V., and 98 co-authors: Deep Trek: Science of Subsurface Habitability & Life on Mars, *Bulletin of the AAS*, 53(4), doi:10.3847/25c2cfcb.dc18f731, 2021.
- Tanaka, K.L.: Sedimentary history and mass flow structures of Chryse and Acidalia Planitiae, Mars, *Journal of Geophysical Research* 102, 4131–4150, doi:10.1029/96JE02862, 1997.
- 1205 Tanaka, K.L.: Debris-flow origin for the Simud/Tiu deposit on Mars, *Journal of Geophysical Research: Planets* 104, 8637–8652, doi:10.1029/98JE02552, 1999.
- Tanaka, K.L., Joyal, T., and Wenker, A.: The Isidis Plains Unit, Mars: Possible catastrophic origin, tectonic tilting, and sediment loading. 31st Lunar and Planetary Science Conference, Abs. # 2023. 2000.
- Tanaka, K.L., Skinner Jr., J.A., Hare, T.M., Joyal, T., and Wenker, A.: Resurfacing history of the Northern Plains of Mars based on geologic mapping of Mars Global Surveyor data, *J. Geophys. Res.* 108 (E4), GDS 24-1–GDS 24-32. Doi:10.1029/2002JE001908, 2003.
- 1210 Tanaka, K.L.: Geology and insolation-driven climatic history of Amazonian north polar materials on Mars, *Nature* 437, 991–994, doi:10.1038/nature04065, 2005.
- Tanaka, K.L., Skinner, J.A., Jr., and Hare, T.M.: Geologic Map of the Northern Plains of Mars, U.S. Geological Survey Science Investigations Map 2888, scale 1:15,000,000. Available online at <https://pubs.usgs.gov/sim/2005/2888>, 2005.
- 1215



- Tanaka, K.L., Rodríguez, J.A.P., Skinner Jr., J.A., Mourke, M.C., Fortezzo, C.M., Herkenhoff, K.E., Kolb, E.J., and Okubo, C.H.: North polar region of Mars: Advances in stratigraphy, structure, and erosional modification, *Icarus* 196 (2), 318–358, doi:10.1016/j.icarus.2008.01.021, 2008.
- 1220 Tanaka, K.L., Skinner, J.A. Jr., Dohm, J.M., Irwin, R.P. III, Kolb, E.J., Fortezzo, C.M., Platz, T., Michael, G.G., and Hare T.M.: Geologic Map of Mars, U.S. Geological Survey Scientific Investigations Map 3292, scale 1:20,000,000, <http://dx.doi.org/10.3133/sim3292>, 2014.
- Tewelde, Y., Zuber, M.T.: Determining the fill thickness and densities of Mars' northern lowlands, 44th LPSC, Abs. #2151, 2013.
- 1225 Tosi, N., and Padovan, S.: Mercury, Moon, Mars. In *Mantle Convection and Surface Expressions* (eds H. Marquardt, M. Ballmer, S. Cottaar and J. Konter), doi:10.1002/9781119528609.ch17, 2021.
- Tu, T.-H., Wu, L.-W., Lin, Y.-S., Imachi, H., Lin, L.-H. and Wang, P.-L.: Microbial Community Composition and Functional Capacity in a Terrestrial Ferruginous, Sulfate-Depleted Mud Volcano. *Front. Microbiol.* 8:2137, doi:10.3389/fmicb.2017.02137, 2017.
- 1230 Tu, V.M., Rampe, E.B., Bristow, T.F., Thorpe, M.T., Clark, J.V., Castle, N., Fraeman, A.A., Edgar, L.A., McAdam, A., Bedford, C., Achilles, C.N., Blake, D., Chipera, S.J., Craig, P.I., Des Marais, D.J., Downs, G.W., Downs, R.T., Fox, V., Grotzinger, J.P., Hazen, R.M., Ming, D.W., Morris, R.V., Morrison, S.M., Pavri, B., Eigenbrode, J., Peretyazhko, T.S., Sarrazin, P.C., Sutter, B., Treiman, A.H., Vaniman, D.T., Vasavada, A.R., Yen, A.S., and Bridges, J.C.: A Review of the Phyllosilicates in Gale Crater as Detected by the CheMin Instrument on the Mars Science Laboratory, Curiosity Rover. *Minerals* 11, 847, doi:10.3390/min11080847, 2021.
- 1235 van Rensbergen, P., Hillis, R. R., Maltman, A. J. and Morley, C. K.: Subsurface sediment mobilization: Introduction. Geological Society, London, Special Publications, 216(1), 1–8, doi:10.1144/GSL.SP.2003.216.01.01, 2003.
- Vago, J. L., Westall, F., Coates, A. J., Jaumann, R., Korablev, O., Ciarletti, V., Mitrofanov, I., Josset, J. L., De Sanctis, M. C., Bibring, J. P., et al.: Habitability on Early Mars and the Search for Biosignatures with the Exomars Rover, *Astrobiology* 2017, 17, 471–510, doi:10.1089/ast.2016.1533, 2017.
- 1240 Wallace, D. and Sagan, C.: Evaporation of ice in planetary atmospheres: ice-covered rivers on Mars, *Icarus* 39, 385–400, doi:10.1016/0019-1035(79)90148-9, 1979.
- Warner, N. H., Sowe, M., Gupta, S., Dumke, A., and Goddard, K.: Fill and spill of giant lakes in the eastern Valles Marineris region of Mars, *Geology* 41 (6), 675–678, doi:10.1130/G34172.1, 2013.
- 1245 Warner, N. H., Gupta, S., Calef, F. Grindrod, P., Boll, N., and Goddard, K.: Minimum effective area for high resolution crater counting of martian terrains, *Icarus* 245, 198–240, doi:10.1016/j.icarus.2014.09.024, 2015.
- Webster, C. R., et al.: Background levels of methane in Mars' atmosphere show strong seasonal variations, *Science*, 360, 1093–1096, doi:10.1126/science.aaq0131, 2018.
- Werner, S. C., Ody, A., Poulet, F.: The Source Crater of Martian Shergottite Meteorites, *Science*, 343, 1343–1346, doi:10.1126/science.1247282, 2014.



- 1250 Westall, F., Foucher, F., Bost, N., Bertrand, M., Loizeau, D., Vago, J.L., Kminek, G., Gaboyer, F., Campbell, K.A., Bréhéret, J.G., Gautret, P., and Cockell, C.S.: Biosignatures on Mars: What, Where, and How? Implications for the Search for Martian Life, *Astrobiology* (11):998-1029, doi:10.1089/ast.2015.1374, 2015.
- Wheatley, D.F., Chan, M.A., and Sprinkel, D.A.: Clastic pipe characteristics and distributions throughout the Colorado Plateau: Implications for paleoenvironment and paleoseismic controls, *Sedimentary Geology*, 344, 20–33. doi:10.1016/j.sedgeo.2016.03.027, 2016.
- 1255 Wheatley, D. F., Chan, M. A., and Okubo, C. H.: Clastic pipes and mud volcanism across Mars: Terrestrial analog evidence of past martian groundwater and subsurface fluid mobilization, *Icarus* 328, 141–151. doi:10.1016/j.icarus.2019.02.002, 2019.
- Wieczorek, M. A., Broquet, A., McLennan, S. M., Rivoldini, A., Golombek, M., Antonangeli, D., et al.: InSight constraints on the global character of the Martian crust, *Journal of Geophysical Research: Planets* 127, e2022JE007298, doi:10.1029/2022JE007298, 2022.
- 1260 Wilson, L., and Head, J.W.: Review and analysis of volcanic eruption theory and relationships to observed landforms, *Rev. Geophys.* 32, 221–263, doi:10.1029/94RG01113, 1994.
- Wilson, L., and Head, J.W.: Evidence for a massive phreatomagmatic eruption in the initial stages of formation of the Mangala Valles outflow channel, Mars, *Geophys. Res. Lett.* 31, L15701, doi:10.1029/2004GL020322, 2004.
- 1265 Wordsworth, R.D.: The Climate of Early Mars. *Annual Review of Earth and Planetary Sciences*, 44, 381-408, doi:10.1146/annurev-earth-060115-012355, 2016.
- Wray, J.J.: Contemporary Liquid Water on Mars? *Annual Review of Earth and Planetary Sciences* 2021 49:1, 141-171, doi:10.1146/annurev-earth-072420-071823, 2021.
- 1270 Wrede, C., Brady, S., Rockstroh, S., Dreier, A., Kokoschka, S., Heinzemann, S.M., and Heller, C.: Aerobic and anaerobic methane oxidation in terrestrial mud volcanoes in the Northern Apennines, *Sedimentary Geology*, 263 pp. 210-219, 2012.
- Wu. X. et al.: Geological characteristics of China’s Tianwen-1 landing site at Utopia Planitia, Mars, *Icarus* 370, 114657, doi:10.1016/j.icarus.2021.114657, 2021.
- 1275 Ye, B.; Qian, Y.; Xiao, L.; Michalski, J.R.; Li, Y.; Wu, B.; Qiao, L.: Geomorphologic exploration targets at the Zhurong landing site in the southern Utopia Planitia of Mars, *Earth Planet. Sci. Lett.*, 576, 117199, doi:10.1016/j.epsl.2021.117199, 2021.
- Zhao, J.; Xiao, Z.; Huang, J.; Head, J.W.; Wang, J.; Shi, Y.; Wu, B.; Wang, L.: Geological characteristics and targets of high scientific interest in the Zhurong landing region on Mars, *Geophys. Res. Lett.*, 48, e2021GL094903, doi:10.1029/2021GL094903, 2021.
- 1280



Figures






Figure 1: Global MOLA topographic map showing the positions of the main fields of Martian mud-volcano-like structures. Letters refer to local evidence of sedimentary volcanism in: a) Simud & Tiu Valles, b) Hydraotes Chaos, c) Firsoff crater, d) Medusae Fossae Formation, e) unnamed flat-floored basin in Terra Sirenum, f) Scandia Colles; and roman numerals refer to regional evidence in: I) Chryse and Acidalia Planitiae, II) Isidis Planitia, III) Adamas Labyrinthus, IV) Amenthes/Nephtes region, V) Arcadia Planitia, and VI) Candor and Coprates Chasmata. See the main text for full description of the evidence. MOLA Science Team.



1290


Figure 2: A conceptual drawing illustrating the main elements associated with mud volcanism (MV) on Earth both on the surface and in the subsurface. Red dashed line represents fault zone and different colours for arrows mark various sources for liquids. Figure is adapted from Mazzini and Etiope (2017).



1295 **Figure 3: Sub-kilometre- to Kilometre-scale, circular mounds in Acidalia Planitia. Panels (a), (c) and (e) are from Mars Reconnaissance Orbiter (MRO) Context Camera (CTX) mosaics (© Google Earth/Mars). Panel (b) is from Mars Odyssey THEMIS-Nighttime infrared (IR) global mosaic, v. 14. (Arizona State University/USGS). Panel (d) is an MRO HiRISE image. Panel (f) is an MRO CTX image. Panel (a), A variety of types of bright mounds. Centred 41.12°N, 26.34°W. Rectangles show locations of Panels (c) - (e). Panel (b), Same area as Panel (a) showing dark responses of the bright mounds and surrounding materials in Nighttime IR. Arrows as in Panel (a). Compare Figs. 3a and b to see the different responses in Nighttime IR of mounds and impact craters. Panel (c), Pitted cones showing central depressions and material on flanks apparently superposed on the darker substrate of the plains. Panel (d), HiRISE image (ESP_026732_2215_RED) showing a bright mound with a central depression and surrounding moat. Panel (e), splotch-like, bright mound with entrained, boulder-sized angular knobs. Panel (f), CTX image (J03_045945_2201_XN_40N027W) showing flow-like features extending from a bright mound (top arrow) to darker apparent flows (lower two arrows). Centred 40.25°N, 27.13°W; location ~62 km SSW of Panel (a). North is up in all. HiRISE imagery: NASA/JPL/University of Arizona, THEMIS imagery NASA/JPL-Caltech/Arizona State University and CTX imagery NASA/JPL/Malin Space Science Systems.**

1300

1305



1310

1315

1320

Figure 4: Examples of kilometre-scale features from various regions of Mars. Panel (a) shows a conical feature with steep flanks and wide central crater (HiRISE ESP_022025_2000, centred 19.73°N, 322.44°E), (b) a domical feature with small central knob in the summit area (HiRISE ESP_025137_1995, centred 19.04°N, 322.64°E), (c) a shield-like or pie-like feature with central breached crater (HiRISE ESP_025704_2005, centred 20.186°N, 321.259°E), (d) a small cluster of conical and domical edifices associated with flow-like unit, and (e) a cluster of wide conical edifices with wide breached central craters surrounded by flow-like unit. Edifices on panels (a,b,c) are situated in the southern part of Chryse Planitia. The fields of cones on panel (d) and (e) are situated on the floor of Coprates Chasma (CTX P13_006269_1670_XN_13S062W, centred 12.711°S, 297.67°E), and in Nephtentes/Amenthes region (CTX G01_018499_1961_XN_16N252W, centred 16.194°N, 107.373°E), respectively. Except the panel (e), north is up. HiRISE imagery: NASA/JPL/University of Arizona and CTX imagery NASA/JPL/Malin Space Science Systems.





Figure 5: An example of kilometre-sized flows (KSFs) from three regions of Mars showing large variability in their shapes and general appearance. While the KSFs within Chryse Planitia captured on panels (a,b,c) consist of three different parts – source area, open channel and leaves – and are often associated with bright smooth units, the KSFs in Adamas Labyrinthus (d,e,f) do not show such an appearance. Instead they have the shape of a plateau. Panels (d,e,f) show plateau-shaped KSFs from Adamas Labyrinthus that are standing above the surrounding terrain to a height of 20-30 metres and they do not show clear source area from which the material originated. A unique KSF is the flow of Zephyria Fluctus which shows well-developed large source area (g), as well as former flow pattern on its surface (h) and well-developed steep-sided margins and lobes (i). Panel (a) based on CTX image P17_007639_1997_XN_19N034W (centred 19.85°N, 326.02°E), (b) CTX F01_036121_2011_XN_21N034W (centred 20.41°N, 325.69°E), (c) CTX B19_016856_1990_XI_19N035W (centred 20.228°N, 324.05°E), (d) CTX G21_026424_2175_XN_37N257W (centred 102.2°E, 37.52°N, 102.2°E), (e) CTX P17_007779_2181_XN_38N259W (centred 39.14°N, 100.92°E), (f) CTX P17_007502_2195_XI_39N256W (centred 38.63°N, 104.2°E), (g) HiRISE ESP_037169_1805 (centred 0.59°N, 155.29°E), (h) HiRISE ESP_027464_1805 (centred 0.65°N, 155.39°E), and (i) HiRISE ESP_028941_1810 (centred 0.795°N, 155.592°E) respectively. HiRISE imagery: NASA/JPL/University of Arizona and CTX imagery NASA/JPL/Malin Space Science Systems.



1340 **Figure 6: Examples of outcrop features observed by the Curiosity rover and hypothesised to have formed by upward**
injection of sediment through the Martian crust. Gale Crater. (a – d). Various pipes observed at Dingo Gap and Marias
Pass. Figures are adapted from Rubin et al. (2017). (e). Dike observed at Yellowknife Bay (arrows). Image numbers for
(a) 0528MR0020830010303294E01_DXXX, (b) MR002083, (c) NLB_445620806EDR_F0261274NCAM00354M, (d)
1051ML0046250040306086E01_DXXX, and (e) 20170206PIA17595-16. NASA/JPL-Caltech/MSSS.

1345




Figure 7: Terrestrial analogues for the martian outcrop features hypothesised to have formed by upward injection of sediment. (a – d) Clastic pipes widely observed within the Colorado Plateau. (e, f) Sedimentary dikes exposed horizontally (e, the dike width is about 30 cm) and vertically (f) along the western coast of Kii Peninsula. These dikes resulted from an ancient (as old as Miocene) process of subsurface sediment mobilization. Photos in panels (a-d) by David Wheatley (all rights reserved) and (e-f) by Goro Komatsu.

1350

Acknowledgment

We are thankful to David Wheatley for providing us images of clastic pipes in Figure 7 and Dave Rubin for helping us to locate martian images used in Figure 6. AM acknowledge the support of the Research Council of Norway (NFR) through the HOTMUD project number 288299 and its centres of Excellence funding scheme, project number 223272 (CEED). VČ acknowledge the support of IGA Faculty of Environmental Sciences CZU Prague through the grant "Kilometre-Sized Flows on Mars: Controlling Factors and Geomorphology – No. 2022B0038".

1355