

---

1 **Response of soil nutrients and erodibility to slope aspect in the northern**  
2 **agro-pastoral ecotone, China**

3 Yuxin Wu<sup>a,b,c</sup>, Guodong Jia<sup>a,b,c,\*</sup>, Xinxiao Yu<sup>a,b,c,\*</sup>, Honghong Rao<sup>d</sup>, Xiuwen Peng<sup>e</sup>,  
4 Yusong Wang<sup>a,b,c</sup>, **Yushi Wang<sup>a,b,c</sup>**

5 <sup>a</sup>Key Laboratory of State Forestry and Grassland Administration on Soil and Water  
6 Conservation, Beijing Forestry University, Beijing 100083, PR China

7 <sup>b</sup>The Metropolitan Area Forest Ecosystem Research Station, School of Soil and Water  
8 Conservation, Beijing Forestry University, Beijing 100083, PR China

9 <sup>c</sup>The Metropolitan Area Field Scientific Observation Research Station, School of Soil  
10 and Water Conservation, Beijing Forestry University, Beijing 100083, PR China

11 <sup>d</sup>**School of Science, East China University of Technology, Nanchang 330013, PR**  
12 **China**

13 <sup>e</sup>Shanghai Investigation, Design & Research Institute Co., Ltd, Shanghai 200126, PR  
14 China

15 \* Corresponding author. Address: No.35 Tsinghua East Road, Haidian District,  
16 Beijing Forestry University, 100083 Beijing, China.

17 Email address: jiaguodong1111@163.com(G.Jia). yuxinxiao1111@163.com(X.Yu).

18 **Abstract**

19 Soil erosion, considered a major environmental and social problem, leads to the  
20 loss of soil nutrients and the degradation of soil structure, **impacts** plant growth.  
21 However, data on the effects of land use changes caused by vegetation restoration on  
22 soil nutrients and erodibility at different slope aspects is limited. This study was  
23 conducted to detect the response of soil nutrients and erodibility of different slope  
24 aspects in a typical watershed of the northern agro-pastoral ecotone in China. The  
25 following indexes were used to determine the improvement of soil nutrients and  
26 erodibility through a weighted summation method: comprehensive soil nutrient index  
27 and comprehensive soil erodibility index. The results showed that the vegetation types  
28 with the highest comprehensive soil quality index (CSQI) on the western, northern,  
29 southern, and eastern slopes were *Pinus sylvestris* and *Astragalus melilotoides* (1.45),  
30 *Caragana korshinskii* and *Capillipedium parviflorum* (2.35), *Astragalus melilotoides*  
31 (4.78), and *Caragana korshinskii* and *Lespedeza bicolor* (5.00), respectively. Slope  
32 aspect had a significant effect on understory vegetation characteristics, soil nutrients,  
33 and soil erodibility. Understory vegetation and soil characteristics ~~could~~ **explained**  
34 50.86–74.56% of the total variance in soil nutrients and erodibility of slope aspect.

---

35 Mean weight diameter and total phosphorus were the main factors affecting CSQI on  
36 different slope aspects. Our study suggested the combinations of species, such as *C.*  
37 *korshinskii* and *L. bicolor*, were the best species to include on any slope aspect in  
38 regards to improving soil nutrients and soil erodibility.

39 **Keywords:** Slope aspect; Soil nutrients; Soil erodibility; Soil erosion; Vegetation  
40 restoration; Land use

## 41 1. Introduction

42 Soil erosion, considered a major environmental and social problem, leads to the  
43 loss of soil nutrients and the degradation of soil structure, influences the functional  
44 capacity of soils on a global scale (Singh and Panda, 2017; Wen et al., 2021).  
45 Vegetation restoration is an important method of ecological restoration that aims to  
46 control soil erosion and prevent soil degradation (Schmiedel et al., 2017; Zhang et al.,  
47 2021). Vegetation restoration can improve the soil structure and nutrients, which in  
48 turn promotes the restoration of soil quality and function (Guo et al., 2021; Li et al.,  
49 2017). Changes in land use due to vegetation restoration play an important role in  
50 improving the environment and ecosystem function, as well as improving soil quality  
51 and soil nutrient cycling (Akiyama and Kawamura, 2007; Singh and Gupta, 2018).

53 Previous studies have shown that the plants selected for vegetation restoration  
54 projects drive land use change and alter soil properties, thus affecting soil erodibility  
55 (Wang et al., 2019b, a; Zhang et al., 2019). Many studies have also elucidated the  
56 influences of land use change on soil nutrients and have confirmed that revegetation is  
57 an effective way to enhance soil nutrients (Huang et al., 2020; Li et al., 2020; Yang et  
58 al., 2021; Zhu et al., 2020). Most studies have only focused on one aspect; thus, they  
59 lack comprehensive consideration and evaluation of the impact of land use changes  
60 caused by vegetation restoration on soil nutrients and erodibility. However, it is not  
61 clear which plants selected for restoration are the most effective in enhancing soil  
62 nutrients and reducing soil erodibility. The lack of a comprehensive understanding  
63 prevents us from gaining the best ecological benefits from vegetation restoration.  
64 Therefore, studies must be conducted on the response of soil nutrients and erodibility  
65 to different vegetation restoration types.

66 Soil erodibility is the sensitivity of the soil surface to erosion processes (Batista  
67 et al., 2023; Bryan et al., 1989). It is a necessary parameter for establishing soil loss  
68 equations and erosion models. There is currently no soil erosion model that can  
69 accurately predict soil erosion, although there are many related models (de Vente et al.,

---

70 2013, 2008). At present, the soil erodibility K-factor, as defined in the general soil  
71 loss equation (USLE), is the most widely used measure (Wischmeier and Smith,  
72 1978). In addition to K, other soil indexes have been adopted, including saturated  
73 hydraulic conductivity (SHC), soil disintegration rate (SDR), mean weight diameter  
74 (MWD), soil structural stability index (SSSI), clay ratio (CR), and soil organic carbon  
75 cementing agent index (SCAI), to quantify soil erodibility (Dong et al., 2022a; Guo et  
76 al., 2021; Wang et al., 2018; Zhang et al., 2019). Soil organic carbon, nitrogen, and  
77 phosphorus as well as their stoichiometry is also essential for assessing soil quality as  
78 well as ecosystem productivity and functionality (Borchard et al., 2017; Li et al., 2020;  
79 Masciandaro and Ceccanti, 1999; Schloter et al., 2003). A single index cannot fully  
80 reflect all soil properties; therefore, it is necessary to develop a comprehensive soil  
81 index using several related indicators.

82 In addition to soil properties, topographic factors also significantly affect soil  
83 nutrients and erodibility (Bangroo et al., 2017; Nabiollahi et al., 2018; Qin et al., 2016;  
84 Zhang et al., 2018). Slope aspect can affect the growth of plants due to a combination  
85 of factors, such as light, temperature, wind speed, and precipitation, which can cause  
86 significant changes in the ecological relationship between plants and the environment  
87 (Li et al., 2018; Tamene et al., 2020; Zhang et al., 2020). This is especially true for  
88 harsh climates such as cold, dry alpine regions in the north, in which plants are more  
89 sensitive to environmental changes. However, the optimal vegetation restoration type  
90 has primarily been studied by slope gradient and slope position (Dong et al., 2022a;  
91 Guo et al., 2021; Wen et al., 2021). There is a lack of systematic evaluation of the  
92 effects of land use changes caused by vegetation restoration on soil nutrients and  
93 erodibility on different slope aspects. Therefore, the classification of slope aspect  
94 needs to be further refined to elucidate the response of different slope aspects to  
95 changes in soil nutrients and erodibility caused by revegetation.

96 The ecologically fragile northern agro-pastoral zone in China is located in an  
97 erosion zone affected by both wind and water; soil erosion in this zone is considered  
98 very serious (Guo et al., 2019). Recently, the Chinese government has planned and  
99 carried out a series of ecological restoration projects in this region, including the  
100 Beijing-Tianjin Wind and Sand Source Control Project, the Beijing-Hebei Water  
101 Protection Forest Project, and the Sebei Forest Plantation Afforestation Project. These  
102 ecological restoration projects have effectively reduced land erosion and  
103 desertification, and have significantly delayed the onslaught of wind and sand (Wang

---

104 et al., 2021b; Zeng et al., 2014; Zhang et al., 2017). However, the method used for  
105 afforestation, which mainly consists of plantations, is affected by differences in water,  
106 heat, wind, and sand in the different habitats, making it difficult to achieve vegetation  
107 restoration in some ecologically fragile areas, and the selection of suitable tree species  
108 is still equivocal.

109 Based on the abovementioned scientific gaps, we hypothesize that both slope  
110 aspect and land use types can significantly alter soil structure and properties to  
111 influence soil nutrients and erodibility under vegetation restoration. We further  
112 hypothesize that the western slope may have the lowest comprehensive soil quality  
113 index compared to other slope aspects. Therefore, we selected four slope aspects  
114 (west, north, south, and east) that have four different land use types (degraded land,  
115 grasslands, shrublands and woodlands) in a typical watershed of the northern  
116 agro-pastoral ecotone with three specific purposes: 1) to determine the impact of  
117 different vegetation types on different slope aspects on soil nutrient improvement and  
118 soil erodibility enhancement; 2) to determine the key influencing factors affecting soil  
119 nutrients and erodibility of the four slope aspects; and 3) to provide optimal  
120 revegetation models for improving soil nutrients and reducing soil erodibility on  
121 different slope aspects.

## 122 **2. Materials and Methods**

### 123 **2.1. Study area**

124 This study was conducted in the Yangcaogou Watershed (41°4'~41°8' N,  
125 114°58'~115°2' E; Fig.1), Chongli District, Zhangjiakou City, Hebei Province, China.  
126 The watershed is located in a typical ecological transition zone of the agro-pastoral  
127 ecotone in northern China (Wu et al., 2023). The study site spans an area of 10.6 km<sup>2</sup>  
128 with an altitude ranging from 1084 to 1575 m. It belongs to a typical temperate  
129 continental monsoon semi-arid climate with an annual average temperature of 3.5 °C.  
130 The average annual rainfall is 401.6 mm. The rainy season occurs from June to  
131 September (Chang et al., 2021; Guo et al., 2019). The main soil type is classified as  
132 chestnut soil in both the Chinese Soil Taxonomy and the World Reference Base for  
133 Soil Resources (Schad, 2017). Most of the study area consists of Proterozoic soil rock  
134 formations. Owing to irrational human reclamation and grazing, there is very serious  
135 soil and gully erosion. Over the past decade, due to the implementation of the  
136 Beijing–Tianjin Sandstorm Source Control Project, soil erosion and desertification  
137 has been effectively mitigated (Wang et al., 2020b). However, native plant

---

138 populations have been diminished and instead the area is planted with trees, shrubs,  
139 and herbs.

## 140 **2.2. Selection of sites and determination of slope aspect**

141 The study was conducted during the 2021 growing season. A comprehensive  
142 field survey was conducted on the dominant plant species and soil properties of each  
143 of the following land use types: degraded land, grasslands, shrublands, and woodlands  
144 in the Yangcaogou watershed. Grasslands, shrublands, and woodlands were restored  
145 from degraded land over the past 12 years. The degraded land (loss of soil material  
146 from wind and water erosion, degradation of physical, chemical and biological  
147 properties of soil) was previously degraded cropland. All land use types were  
148 vegetated and restored in the form of engineering measures such as fish scale pits  
149 (Wang et al., 2014b) and parallel ditches (Barua and Alam, 2013).

150 In addition to the degraded land, the other three land use types were all sampled  
151 along complete slope aspects at the E, W, N, and S slopes. It includes 28 sample sites  
152 (20 m × 20 m) of a degraded land, two grasslands, two shrublands and two woodlands  
153 on each slope aspect. Three sampling quadrats (1 m × 1 m) were set up in each  
154 sample site to investigate and record the species, height, richness, coverage,  
155 aboveground biomass, belowground biomass, and litter biomass of herbs. Height was  
156 measured as the average height of herbs in the sample. Biomass coverage was  
157 determined following the visual method (Proulx and Mazumder, 1998). Richness was  
158 calculated by measuring the number of individuals of each herb in the quadrat and  
159 calculating the percentage of its occurrence (Dou et al., 2023). Belowground biomass  
160 and soil samples were collected with a 9 cm diameter soil drill. The measured land  
161 use types, major plant species, and understory vegetation characteristics at each  
162 selected field site are listed in Table S1.

163 Following the methods described by (Yimer et al., 2006), study sites were  
164 selected that included the four land use types on each of the four slope aspects: east,  
165 west, north, and south. Eastern, western, northern, and southern slopes are also known  
166 as semi-sunny, semi-shady, shady, and sunny slopes (Che et al., 2022; Chen et al.,  
167 2021b). ~~In this region, it is difficult to find degraded land because the vast majority of  
168 the degraded land had been converted to artificial forest and grass vegetation.~~ In this  
169 region, four unrestored degraded land were selected as representatives from the  
170 western slope. The slope gradients and positions were similar for all selected sample

---

171 sites (Fig. 1).

### 172 **2.3. Soil sampling and analysis**

173 Three quadrants were selected at each site to investigate vegetation and collect  
174 soil samples. For each sampling point, a steel cutting ring (100 cm<sup>3</sup>) was used to  
175 obtain 75 soil samples (25 sites × three sampling points). The saturated hydraulic  
176 conductivity of the soil were evaluated using the constant head permeability test  
177 (Chandler and Chappell, 2008). The mean weight diameter was measured by screens  
178 with different pore sizes (0.25, 0.50, 1.00, 2.50 and 5.00 mm) (Campo et al., 2008).  
179 After air-drying via dry screening, 50 g of the soil samples were placed on the sieve of  
180 a soil aggregate analyzer (TTF-100 model, China), then completely immersed in  
181 water, and shaken up and down 30 times for 1 minute (Wang et al., 2014a). After  
182 shaking, samples were removed from the settling cylinder, and the remaining  
183 aggregates on each sieve were put into an aluminum box for drying. Finally, the  
184 samples were weighed and the dried aggregates were recorded.

185 Soil characteristics of different vegetation types at different slope aspects are  
186 listed in Table S2. Topsoil samples were collected from 0–10 cm using a cutting ring.  
187 Samples were brought back to the lab to oven-dried at 105°C for 24 hours. Then, the  
188 soil bulk density (SBD) (Lardy et al., 2022; Moreira et al., 2020) and soil capillary  
189 porosity (SCP) (Singh and Pollard, 1958) were measured. In addition, 225 mixed soil  
190 samples (25 sites × three quadrats/site × three samples/quadrat) were collected as soil  
191 samples. Among them, the particle size distribution of clay content (Cl), silt content  
192 (Si), sand content (Sa) was determined by a Microtrac S3500 laser particle sizer  
193 (Malvern 3000, UK). Total nitrogen (TN) and total phosphorus (TP) were determined  
194 by the dichromate oxidation (Bremner, 1996) and HClO<sub>4</sub>-H<sub>2</sub>SO<sub>4</sub> methods (Kisand,  
195 2005), respectively. Soil pH (Cornfield, 1954) was determined using a pH meter at a  
196 2.5 soil:1 water ratio.

### 197 **2.4 Calculation of soil indexes**

198 Saturated hydraulic conductivity of the soil (K<sub>s</sub>) (Campo et al., 2008), mean  
199 weight diameter (MWD) (Ortas and Lal, 2012), soil disintegration rate (SDR) (Guo et  
200 al., 2021), soil structure stability index (SSSI) (Nichols and Toro, 2011), soil organic  
201 carbon cementing agent index (SCAI) (Dong et al., 2022a) and K factor (Jiang et al.,  
202 2020; Li et al., 2012) were used to express the soil erodibility. These indexes were  
203 calculated using equations (1) - (5):

$$K_S = \frac{QL}{Aht} \quad (1)$$

where Q is the outflow volume (ml), A is the soil column section (mm<sup>2</sup>), t is the time (min), h is the head difference (mm), and L is the height of the soil column (mm).

$$MWD = \sum_{i=1}^n (w_i/m_t)d_i \quad (2)$$

Where  $w_i$  is the mass of the i-th level of aggregates or other soil material (g),  $m_t$  is the sample mass, and  $d_i$  is the mean diameter of the i-th level of aggregates or other soil material (mm).

$$SDR = \frac{M_1 - M_2}{t_2 - t_1} \times 100\% \quad (3)$$

Where  $M_1$  and  $M_2$  are the weight of the soil before ( $t_1$ ) and after ( $t_2$ ) disintegration, respectively.

$$SSSI = 100\% \times \frac{SOMC}{Cl+Si} \quad (4)$$

$$K = \left\{ 0.2 + 0.3 \exp \left[ -0.0256Sa \left( 1 - \frac{Si}{100} \right) \right] \right\} \left( \frac{Si}{Cl+Si} \right)^{0.3} \times \left( 1 - \frac{0.25C}{C + \exp(3.72 - 2.95C)} \right) \left( 1.0 - \frac{0.7SN1}{SN1 + \exp(-5.51 + 22.9SN1)} \right) \quad (5)$$

Where SOMC is the content of soil organic matter (Kar et al., 2023),  $C = 0.583 \times SOMC$ ; Cl and Si represent the clay and silt content (%), respectively;  $SN1 = 1 - Sa/100$ ; K represents the soil loss rate per unit area under rainfall erosivity conditions for a specified soil on a standard plot (Jiang et al., 2020; Renard et al., 1997). A previous study indicates the rationality and validity of estimating K in the Zhangjiakou region using this model (Wang et al., 2020a).

In order to further evaluate soil nutrients and erodibility, comprehensive soil nutrient and erodibility index were calculated using equations 6 and 7, respectively:

$$CSNI = \sum_i^n K_{ni} \cdot C_{ni} \quad (6)$$

$$CSEI = \sum_i^n K_{ei} \cdot C_{ei} \quad (7)$$

Where  $K_{ni}$  and  $C_{ni}$  are the weight and score of soil nutrient index respectively,  $K_{ei}$  and  $C_{ei}$  are the weight and score of soil erodibility index respectively, and n is the number of indexes.

The weight of each soil nutrient index and soil erodibility index was determined using a principal component analysis (PCA) (Pandey et al., 2021; Wang et al., 2018). The scores of SHC, MWD, SSSI, SOC, TN, and TP scores were calculated using a

---

233 "reverse S" function, which was calculated using equations 8.

$$234 \quad f(x) = \begin{cases} 1 & , x \geq b \\ \frac{x-a}{b-a} & , a < x < b \\ 0 & , x \leq a \end{cases} \quad (8)$$

235 The SDR and K factor scores were calculated by "S" function, as shown in  
236 equations 9.

$$237 \quad f(x) = \begin{cases} 1 & , x \leq b \\ \frac{x-a}{b-a} & , a > x > b \\ 0 & , x \geq a \end{cases} \quad (9)$$

238 Comprehensive soil quality index (CSQI) is used to express soil quality, which  
239 takes into account both soil nutrients and erodibility (De Laurentiis et al., 2019; Dong  
240 et al., 2022b). The CSQI was calculated as follows (Eq. 10):

$$241 \quad CSQI = \frac{CSNI}{CSEI} \quad (10)$$

242 where CSQI ( $> 0$ ), CSNI (0-1) and CSEI (0-1) are the comprehensive soil quality,  
243 nutrient, and erodibility indexes, respectively.

## 244 2.5. Statistical analysis

245 ~~Excel 2016~~ and SPSS Ver. 20 software were used for data processing and  
246 statistical analysis, and ArcGIS 10.4.1 and Origin 2021 were used for graphing. A  
247 one-way analysis of variance (ANOVA) was used to compare soil nutrient and  
248 erodibility indexes of different slope aspects and different land use types. The effects  
249 of land use types, slope aspects and their interaction on soil nutrients and erodibility  
250 indexes were tested using a two-way ANOVA. Pearson's correlation coefficient was  
251 used to determine the correlation between soil nutrient, erodibility, and quality  
252 indexes and their influencing factors. The contributions of understory vegetation and  
253 soil characteristics to total variance in soil nutrients and erodibility indicators were  
254 determined using a redundancy analysis (RDA) (Capblancq et al., 2018; Peres-Neto et  
255 al., 2006). A random forest algorithm based on R software was used to analyze the  
256 importance of impact factors from different slope aspects (Schonlau and Zou, 2020;  
257 Vincenzi et al., 2011). The importance index was determined as the average accuracy  
258 reduction. When the importance index is higher, it means that the corresponding  
259 factor holds more weight (Chen et al., 2021a; Hao et al., 2015).

## 260 3. Results



---

### 261 **3.1. Changes in the characteristics of understory vegetation on different slope** 262 **aspects**

263 Slope aspect significantly influenced some of the characteristics of understory  
264 vegetation such as aboveground biomass (AGB) and belowground biomass (BGB).  
265 All measured characteristics of understory vegetation on the western slope were lower  
266 than that of other three slope aspects. AGB and BGB was significantly lower for the  
267 western slope than the eastern slope (Fig. 2). AGB and BGB on the eastern slope were  
268 significantly higher than those on the western slope by 63.40% and 78.40%,  
269 respectively (Fig. 2d, e). The measured plant characteristics from the eastern and  
270 western slopes were not significantly different from those on the northern and  
271 southern slopes. There were significant differences among the four land use types for  
272 all characteristics measured for the western slope (Table S1). BH, R, and AGB of  
273 understory vegetation were significantly higher for the woodland than for the other  
274 three land use types (Fig. 2). Overall, shrubland had the highest litter biomass on each  
275 slope aspect, while degraded land on the western slope had the lowest.

### 276 **3.2. Changes in soil nutrients on different slope aspects**

277 Slope aspect significantly affected soil nutrients. Soil organic carbon (SOC),  
278 total nitrogen (TN), and total phosphorus (TP) were significantly lower in soil  
279 collected from the western slope than the eastern slope (Fig. 2). SOC of the eastern  
280 slope was 0.96–1.38 times greater than that of other slopes, respectively (Fig. 2g). TN  
281 was highest on the eastern slope and was 0.39 g kg<sup>-1</sup> and 0.28 g kg<sup>-1</sup> greater than that  
282 on the western and northern slopes, respectively (Fig. 2h). Similarly, the TP of the  
283 eastern slope was significantly greater than that of the southern and eastern slopes by  
284 59.60% and 17.37%, respectively (Fig. 2i). When all slope aspects were considered,  
285 comprehensive soil nutrient index (CSNI) was significantly lower on the western  
286 slope than on the other three slope aspects. The highest CSNI was found for both  
287 southern slope (0.81) and eastern slope (0.86) (Fig. 3). For a given slope aspect, land  
288 use types also significantly influenced soil nutrients (Fig. S1). For example, on the  
289 western slope, the SOC of forested land was significantly higher than other restored  
290 land uses by 11.81–150.84% depending on the comparison. SOC, TN, and TP of  
291 degraded land were significantly lower than that of other land use types. CSNI was  
292 influenced by land use type, slope aspect, and their interactions (Table 1). Compared  
293 to degraded land, CSNI was significantly higher for all three land uses, with the  
294 greatest increase in CSNI for shrubland (0.75), followed by woodland and grassland

---

295 (Fig. 4).

### 296 **3.3. Changes in soil erodibility under vegetation restoration**

297 The effect of slope aspect on soil erodibility indicators was significant (Table 1  
298 and 2). Among the four slope aspects, SHC of the soil collected from the eastern slope  
299 was the greatest, and was significantly greater than that of the western and northern  
300 slopes by 311.16% and 187.10%, respectively. MWD was highest on the eastern slope  
301 (3.65 mm), followed by the southern and northern slopes. MWD among the four  
302 slopes was significantly different. SSSI of the western slope was the lowest (0.41 g  
303 kg<sup>-1</sup>), and it was significantly lower than the other three slope aspects. In contrast, the  
304 highest SCAI was found on the western slope, and it was significantly higher than the  
305 other slope aspects by 46.10%–59.70%, respectively. When all slope aspects were  
306 considered, the southern (0.26) and eastern (0.20) slopes had the highest  
307 comprehensive soil erodibility index (CSEI) reduction capacity (Fig. 3). For any  
308 given slope aspect, land use types also greatly influenced soil erodibility indicators  
309 (Table 2). On the western slope, MWD was significantly increased by 0.67 mm–1.59  
310 mm. On the northern slope, the SHC of woodland was significantly higher than that of  
311 shrubland (by 117.67%) and grassland (by 94.24%), respectively. On the southern  
312 slope, the K in the grassland land use type was significantly lower than that in  
313 woodland and shrubland. On the eastern slope, soil disintegration rates of the three  
314 restored land uses were significantly different, with the highest SDR in the woodlands.  
315 CSEI was influenced by land use type, slope aspect, and their interactions (Table 1).  
316 The CSEI of all three restored land uses was significantly lower by (63.01–64.70%)  
317 compared to the degraded land (Fig. 4).

### 318 **3.4. Changes in comprehensive soil quality index under vegetation restoration**

319 When all slope aspects are considered, there were significant differences in  
320 comprehensive soil quality index (CSQI), with the eastern slope (2.46) having the  
321 greatest capacity to increase CSQI (Fig. 3). Compared to degraded land, the CSQI of  
322 grassland, shrubland and woodland increased significantly by 2.51, 2.65, and 2.44,  
323 respectively (Fig. 4). CSQI was influenced by land use type, slope aspect, and their  
324 interactions (Table 1).

325 The differences in CSQI of different vegetation types were compared to  
326 determine the optimal vegetation restoration type for different slope aspects. On the  
327 western slope, the WGCP grassland (*Capillipedium parviflorum*) and WWPS  
328 woodland (*Pinus sylvestris* and *Astragalus melilotoides*) had relatively high CSQIs.

---

329 They were significantly higher than that of other vegetation types (Fig. 5a). Therefore,  
330 these two plant communities may be selected for restoration practices on the western  
331 slope. On the northern slope, the CSQI of the shrubland (NSCK) was significantly  
332 higher and second highest in grassland (NGBI). The combination of *Caragana*  
333 *korshinskii* and *Capillipedium parviflorum* (NSCK) could also be selected as taxa for  
334 restoration vegetation (Fig. 5b). On the southern slope, the CSQI of grassland (SGAM)  
335 was significantly higher than that of other vegetation types (Fig. 5c). The SGAM was  
336 dominated by the herb *Astragalus melilotoides*, which had the highest CSQI. *A.*  
337 *melilotoides* could be selected for improving soil quality on the southern slope. On the  
338 eastern slope, the CSQI of the shrubland (ESCK) was relatively higher than that of  
339 other sites (Fig. 5d). The ESCK was dominated by *Caragana korshinskii* and  
340 *Lespedeza bicolor*, which had the highest CSQI. Therefore, these species should be  
341 selected for improving soil quality on the eastern slope.

### 342 **3.5. Key factors and their contributions on different slope aspects**

343 The RDA followed by Monte Carlo permutation tests revealed that the variations  
344 in the nine measured soil quality indicators were significantly influenced by  
345 understory vegetation and soil characteristics on the four slope aspects ( $P < 0.01$ , Fig.  
346 6). On the western slope, 62.7% of the total variance can be explained by understory  
347 vegetation and soil characteristics (Fig. 6a), with understory vegetation and soil  
348 characteristics explaining 43.11% and 19.59% of the total variance, respectively. For  
349 the northern slope, the understory vegetation and soil characteristics contributed  
350 50.86% of the total variance of soil quality (Fig. 6b), of which understory vegetation  
351 and soil characteristics accounted 33.28% and 17.58% of the total variance,  
352 respectively. On the southern slope, the total variance in soil quality of 54.23% could  
353 be explained by understory vegetation and soil characteristics, of which the  
354 combination of soil and roots contributed 44.56% and 9.67% of total variance,  
355 respectively (Fig. 6c). However, on the eastern slope, the understory vegetation and  
356 soil characteristics contributed 74.56% of the total variance of soil quality (Fig. 6d),  
357 of which understory vegetation and soil characteristics accounted for 56.81% and  
358 17.59% of the total variance, respectively.

359 The random forest analysis highlighted the importance of 21 modeling factors to  
360 determine the restoration characteristics of understory vegetation and the physical and  
361 chemical characteristics of topsoil on different slope aspects. MWD, TP, saturated  
362 hydraulic conductivity (SHC), and soil disintegration rate (SDR) were the main

---

363 factors influencing understory vegetation and soil properties on different slope aspects.  
364 The mean accuracy reduction was calculated using the random forest method. Using  
365 this calculation, we obtained an MWD of 13.40, TP of 13.30, SHC of 12.60, and SDR  
366 of 8.20 (Fig. S2).

#### 367 **4. Discussion**

##### 368 **4.1. Effects of slope aspect on understory vegetation characteristics**

369 ~~Slope aspect, one of the most important topographic factors, may impacts~~  
370 ~~vegetation characteristics due to differences in sunlight, moisture, temperature, and~~  
371 ~~soil (Fig. 2). Soil is the material basis for plant growth, and there is an important~~  
372 ~~relationship between plant growth, development, and distribution and the soil~~  
373 ~~characteristics of different slope aspects (Gao, 2017; Zhou et al., 2020). There is a~~  
374 ~~synergistic evolutionary and adaptive relationship between plant growth and survival~~  
375 ~~in the environment. Moreover, plants grow differently on different slope aspects,~~  
376 ~~showing plastic responses depending on their habitat (Che et al., 2022; Sharma et al.,~~  
377 ~~2010).~~

378 Although slope aspect, one of the most important topographic factors, may  
379 impacts vegetation characteristics due to differences in sunlight, moisture,  
380 temperature, and soil, our results showed that most of the characteristics of understory  
381 vegetation had no significant differences based on the different slope aspects. This  
382 may be due to the fact that the understory plants were shaded by the taller trees and  
383 shrubs (Niinemets, 2010). Aboveground biomass was greater on the eastern and  
384 southern slopes than on the northern and western slopes. Vegetation density was  
385 lowest on the western slope. These findings indicated that aboveground biomass is  
386 closely related to sunshine hours. Sunshine hours affect the balance of heat and water  
387 (Chen et al., 2021b; Shi et al., 2021). This contributed to the low aboveground  
388 biomass of the western slope. Similarly, belowground biomass declined from the  
389 eastern, southern, northern, and western slopes. This may be due to the difference in  
390 the aboveground biomass of the four slope aspects. Aboveground biomass impacts  
391 belowground biomass (Sun et al., 2022), and the belowground biomass was  
392 significantly lower on the western slope than on the eastern slope.

393 ~~In view of the influence of slope aspect on the establishment of restored~~  
394 ~~vegetation in the study area, the number of seedlings on the western and northern~~  
395 ~~(shaded) slopes should be increased at the early stage of vegetation restoration in the~~  
396 ~~northern agro-pastoral ecotone. In addition, timely replanting and follow-up~~

---

397 ~~application of nitrogen fertilizer during the restoration process will help to reduce the~~  
398 ~~differences in vegetation growth caused by the inherent differences among the slope~~  
399 ~~aspects.~~

#### 400 **4.2. Effects of slope aspect on soil nutrients**

401 ~~Soil nutrients play an important role in the maintenance and improvement of soil~~  
402 ~~quality. Soil nutrients are an important reflection of the ecological effects of~~  
403 ~~vegetation restoration~~ (Salekin et al., 2021; Wang et al., 2012; Yüksek and Yüksek,  
404 2021). Our results show that the conditions related to slope aspect have significant  
405 effects on single soil nutrient indicators and the comprehensive soil nutrient index  
406 (Figs. 2, 5). In the same area, soil nutrients can vary depending on the slope aspect (Li  
407 et al., 2021; Sharma et al., 2010). On different slope aspects, TN, TP, and the  
408 comprehensive soil nutrient index of surface soil were highest on the eastern and  
409 southern slopes, while the soil organic carbon content was highest on the northern  
410 slope. Plants need to absorb a large amount of fast-acting nitrogen and phosphorus  
411 during vegetative growth, and the nutrients required for plant growth are converted  
412 from organic matter in the soil. The lowest SOC, TN, TP, and the comprehensive soil  
413 nutrient index on the western slope are due to the fact that it was located in the  
414 wind-water erosion zone of the northern agro-pastoral ecotone, and the topsoil has  
415 been lost due to long-term wind erosion.

416 The effect of different slope aspect conditions on soil pH was limited. This is  
417 because plant root systems and sediments were not abundant in the case of vegetation  
418 restoration of just 12a (Bai et al., 2020). The organic acid content was low when  
419 combined with organic matter during decomposition and vegetation restoration;  
420 therefore, it was insufficient to lower the pH of the surface soil (Seddaiu et al., 2013).  
421 ~~Because controlling wind speed is the key to soil nutrient enhancement, future~~  
422 ~~restoration projects that take place in dry alpine areas (i.e., the western and northern~~  
423 ~~slopes) should prioritize the use of thickened non-woven fabric of at least 50 g m<sup>2</sup> for~~  
424 ~~better insulation and to block wind, which is conducive to seed germination and~~  
425 ~~seedling growth.~~

#### 426 **4.3. Effects of slope aspect on soil erodibility**

427 ~~Soil erodibility is commonly used to characterize the susceptibility of soils to~~  
428 ~~water erosion and is influenced by vegetation and soil characteristics.~~ Our results  
429 show that slope aspect has a significant effect on single soil erodibility indexes as well  
430 as comprehensive soil erodibility index. In general, soil erodibility decreases from the

---

431 western slope to the eastern slope (Table 2), a pattern that may be related to the  
432 geographical location, altitude, temperature, and semi-arid climate of the region. Due  
433 to **special location**, the western and northern slopes are susceptible to year-round gales  
434 from the northwestern interior and Siberia, resulting in varying environmental  
435 conditions ~~on the different slope aspects~~. However, the soil water content of the  
436 northern slope (shaded slope) is higher than that of the western slope, which may be  
437 more favorable for vegetation restoration on the northern slope (Liu et al., 2020); the  
438 western slope may be more vulnerable to erosion. Wind speed and soil moisture are  
439 key factors controlling the process of vegetation restoration (Hupet and Vanclooster,  
440 2002; Meng et al., 2018), and these factors further influence soil erodibility (Sun et al.,  
441 2016). ~~Therefore, future studies should investigate methods to enhance vegetation  
442 restoration while utilizing soil water resources available on the different slope aspects  
443 and reducing soil erodibility.~~

#### 444 **4.4. Relationship between soil nutrients and soil erodibility**

445 The comprehensive soil nutrient index was significantly positively correlated  
446 with saturated hydraulic conductivity, mean weight diameter, and soil structure  
447 stability index (Fig. 7), while the comprehensive soil nutrient index was highly  
448 significantly negatively correlated with the comprehensive soil erodibility index, with  
449 an explanation of 88% (Table S1). Many previous studies have reported similar  
450 results (Dong et al., 2022a; Zhu et al., 2018). In this study, higher saturated hydraulic  
451 conductivity, mean weight diameter, and soil structure stability index and lower soil  
452 disintegration rate, K, and SOC cementing agent index indicate better soil structure  
453 and lower soil erodibility. These characteristics can significantly reduce runoff and  
454 sediment loss, which can result in soil nutrient accumulation (Pan and Shangguan,  
455 2006; Sun et al., 2015; Zheng et al., 2021). Therefore, revegetation increases soil  
456 nutrients and reduces soil erodibility, which further change vegetation and soil  
457 characteristics. In addition, these factors could reduce soil nutrient loss and further  
458 promote soil nutrient accumulation by reducing soil erodibility.

459 The comprehensive soil erodibility index was highly significantly negatively  
460 correlated with SOC, TN, and TP (Fig. 7). Previous studies have shown that soil  
461 organic matter and SOC are closely related to soil erodibility (Wang et al., 2019b).  
462 SOC acts as a cement for soil aggregation, which improves soil structural stability  
463 through the formation of aggregates, thus reducing soil erodibility. Soil nitrogen  
464 indirectly affects soil erodibility by promoting plant growth and development,

---

465 increasing the accumulation of SOC in plants. In addition, nitrogen enrichment  
466 increased soil macroparticles and mean weight diameter, which directly affected soil  
467 erodibility. Similar to nitrogen, phosphorus is one of the essential elements for plant  
468 growth and development, and the phosphorus content of soil determines the  
469 development of soil microorganisms and root systems, which will further influence  
470 the input of soil organic carbon and the formation of soil aggregates.

#### 471 **4.5. Key factors impacting soil and vegetation related to slope aspect**

472 ~~The interaction between soil and vegetation in the study area is complex.~~  
473 ~~Because in the early stages of vegetation recovery, soil factors are unstable and~~  
474 ~~vegetation is in the adaptation stage (Peng et al., 2009):~~ The results derived from the  
475 random forest method showed **that mean weight diameter and TP were the main**  
476 **influencing factors** ~~influencing the surface soil indicators.~~ The main adhesion agents  
477 for the formation of aggregates included clay content, SOC and cementation. The  
478 mean weight diameter was significantly and positively correlated with soil organic  
479 carbon and clay content. The magnitude of mean weight diameter affects soil  
480 structural stability and root establishment, which varies due to environmental factors  
481 on different slope aspects. Soil phosphorus is an important element necessary for  
482 plant growth and development, and rapid growth requires more soil phosphorus, so  
483 there were some differences between different land use types on different slope  
484 aspects. The difference in TP between slope aspect affected the amount of inorganic  
485 phosphorus available for uptake by plants, and the lower phosphorus content limited  
486 plant growth. ~~Saturated hydraulic conductivity reflects the permeability of soil and is~~  
487 ~~an important indicator of soil erodibility.~~ ~~Differences in aboveground and~~  
488 ~~belowground biomass of different slope aspects lead to different soil root traits, which~~  
489 ~~affect the magnitude of saturated hydraulic conductivity.~~ ~~The soil disintegration rate~~  
490 ~~was significantly negatively correlated with soil organic carbon, clay content, and~~  
491 ~~mean weight diameter, and differences in soil microbial, nutrient, and root~~  
492 ~~characteristics between slope aspects resulted in significant variations in the soil~~  
493 ~~disintegration rate.~~ By analyzing the main factors influencing surface soil quality in  
494 different slope aspects, timely application of phosphorus fertilizer in vegetation  
495 restoration projects could help accelerate the process of afforestation.

#### 496 **4.6. Optimal land use type and plant species based on slope aspect**

497 Our study has shown that vegetation restoration can be an effective measure to  
498 improve soil nutrients and reduce soil erodibility. Moreover, the restored land use

---

499 types and plant species to improve soil quality differed significantly depending on the  
500 slope aspect. Therefore, according to the differences in water, heat, wind, and sand on  
501 different slope aspects ~~in the northern agro-pastoral ecotone of China~~, the selection of  
502 land use and its corresponding vegetation types should be carefully considered ~~when~~  
503 ~~planning restoration projects to improve soil quality. The comprehensive soil nutrient,~~  
504 ~~erodibility, and quality indexes were established with a comprehensive investigation~~  
505 ~~of various soil nutrient and erodibility indexes. The optimal types of vegetation~~  
506 ~~restoration for different slope aspects was clarified.~~ Our findings both agree with and  
507 differ from previous studies (Colgan et al., 2010; Dong et al., 2022a; Wang et al.,  
508 2021a). Studies that found contrasting results are likely due to the environmental  
509 conditions (e.g. climate, rainfall, topographic conditions, seed bank, soil texture) of  
510 the different slopes aspects. It is noteworthy that herbaceous vegetation on the  
511 western slope is prone to severe shallow nutrient loss and soil erosion because of  
512 strong wind conditions and sandy soil (Guo et al., 2020). Therefore, the use of  
513 herbaceous vegetation should be carefully considered as the primary restoration  
514 vegetation species. Fortunately, our proposal (*Caragana korshinskii* and *Lespedeza*  
515 *bicolor*) satisfied this requirement. In addition, wind also contributes to soil erosion in  
516 this region; however, limited research has been conducted on wind erosion and  
517 combined erosion by wind and water. Future studies should be conducted on  
518 combined erosion by wind and water study to better characterize soil erosion.

## 519 **5. Conclusions**

520 We found that some understory vegetation characteristics and soil properties  
521 varied significantly with slope aspect. Soil nutrients and erodibility reflected by soil  
522 organic carbon, total nitrogen, total phosphorus, saturated hydraulic conductivity, soil  
523 disintegration rate, mean weight diameter, soil structure stability index, soil erodibility  
524 factor, and soil organic carbon cementing agent index, respectively, were also  
525 influenced by slope aspect and land use. Furthermore, comprehensive soil nutrient,  
526 erodibility, and quality indexes also varied significantly with slope aspect, land use,  
527 and predominant plant species. Slope aspect strongly modified the relationship  
528 between comprehensive soil nutrient, erodibility, and quality indexes as well as  
529 understory vegetation characteristics and soil properties. Our study found that  
530 *Caragana korshinskii* and *Lespedeza bicolor* were the best taxa to include on any  
531 slope aspect to improve soil nutrients and prevent soil erosion. This study provides  
532 insight into the rational planning of vegetation restoration measures on all slope



---

533 aspects in the northern agro-pastoral ecotone in semi-arid areas. **Future work will**  
534 **focus on land degradation associated with soil erosion from water and storms in the**  
535 **region.**

#### 536 **Date Availability**

537 Data will be made available on request.

#### 538 **Author contributions.**

539 Yuxin Wu: Writing-original draft. Guodong Jia: Project administration, Funding  
540 acquisition, Writing-review and editing. Xinxiao Yu: Project administration, Funding  
541 acquisition, Writing-review and editing. Honghong Rao: Methodology and Formal  
542 analysis. Xiuwen Peng: Investigation. Yusong Wang **and Yushi Wang**: Investigation.

#### 543 **Competing interests.**

544 The author declares that the publication of this scientific paper has no conflict of  
545 interest.

#### 546 **Acknowledgements.**

547 We are grateful for the grants from the National Key Research and Development  
548 Program of China (2022YFF1302502-03) (China) and the National Natural Science  
549 Foundation of China (42230714).

#### 550 **References**

551 Akiyama, T. and Kawamura, K.: Grassland degradation in China: Methods of  
552 monitoring, management and restoration, *Grassland Science*, 53, 1–17,  
553 <https://doi.org/10.1111/j.1744-697X.2007.00073.x>, 2007.

554 Bai, Y., Zha, X., and Chen, S.: Effects of the vegetation restoration years on soil  
555 microbial community composition and biomass in degraded lands in Changting  
556 County, China, *J. For. Res.*, 31, 1295–1308,  
557 <https://doi.org/10.1007/s11676-019-00879-z>, 2020.

558 Bangroo, S. A., Najjar, G. R., and Rasool, A.: Effect of altitude and aspect on soil  
559 organic carbon and nitrogen stocks in the Himalayan Mawer Forest Range, *CATENA*,  
560 158, 63–68, <https://doi.org/10.1016/j.catena.2017.06.017>, 2017.

561 Barua, G. and Alam, W.: An analytical solution for predicting transient seepage  
562 into ditch drains from a ponded field, *Advances in Water Resources*, 52, 78–92,  
563 <https://doi.org/10.1016/j.advwatres.2012.09.002>, 2013.

564 Batista, P. V. G., Evans, D. L., Cândido, B. M., and Fiener, P.: Does soil thinning  
565 change soil erodibility? An exploration of long-term erosion feedback systems, *SOIL*,  
566 9, 71–88, <https://doi.org/10.5194/soil-9-71-2023>, 2023.

567 Borchard, N., Adolphs, T., Beulshausen, F., Ladd, B., Gießelmann, U. C.,  
568 Hegenberg, D., Mösel, B. M., and Amelung, W.: Carbon accrual rates, vegetation  
569 and nutrient dynamics in a regularly burned coppice woodland in Germany, *GCB*  
570 *Bioenergy*, 9, 1140–1150, <https://doi.org/10.1111/gcbb.12408>, 2017.

571 Bremner, J.: *Methods of Soil Analysis Part 3, Chemical Methods*, Chemical  
572 *Methods (Methodsofsoilan3)*, 1996.

573 Bryan, R. B., Govers, G., and Poesen, J.: The concept of soil erodibility and

---

574 some problems of assessment and application, *CATENA*, 16, 393–412,  
575 [https://doi.org/10.1016/0341-8162\(89\)90023-4](https://doi.org/10.1016/0341-8162(89)90023-4), 1989.

576 Campo, J., Gimeno-García, E., Andreu, V., González-Pelayo, O., and Rubio, J. L.:  
577 Aggregation of under canopy and bare soils in a Mediterranean environment affected  
578 by different fire intensities, *CATENA*, 74, 212–218,  
579 <https://doi.org/10.1016/j.catena.2008.05.002>, 2008.

580 Capblancq, T., Luu, K., Blum, M. G. B., and Bazin, E.: Evaluation of  
581 redundancy analysis to identify signatures of local adaptation, *Mol Ecol Resour*, 18,  
582 1223–1233, <https://doi.org/10.1111/1755-0998.12906>, 2018.

583 Chandler, K. R. and Chappell, N. A.: Influence of individual oak (*Quercus robur*)  
584 trees on saturated hydraulic conductivity, *Forest Ecology and Management*, 256,  
585 1222–1229, <https://doi.org/10.1016/j.foreco.2008.06.033>, 2008.

586 Chang, X., Sun, L., Yu, X., Liu, Z., Jia, G., Wang, Y., and Zhu, X.: Windbreak  
587 efficiency in controlling wind erosion and particulate matter concentrations from  
588 farmlands, *Agriculture, Ecosystems & Environment*, 308, 107269,  
589 <https://doi.org/10.1016/j.agee.2020.107269>, 2021.

590 Che, C., Xiao, S., Ding, A., Peng, X., and Su, J.: Growth response of plantations  
591 *Hippophae rhamnoides* Linn. on different slope aspects and natural *Caragana opulens*  
592 Kom. to climate and implications for plantations management, *Ecological Indicators*,  
593 138, 108833, <https://doi.org/10.1016/j.ecolind.2022.108833>, 2022.

594 Chen, Y., Zheng, W., Li, W., and Huang, Y.: Large group activity security risk  
595 assessment and risk early warning based on random forest algorithm, *Pattern*  
596 *Recognition Letters*, 144, 1–5, <https://doi.org/10.1016/j.patrec.2021.01.008>, 2021a.

597 Chen, Z., Wang, G., Pan, Y., Yang, X., and Shen, Y.: Water use patterns differed  
598 notably with season and slope aspect for *Caragana korshinskii* on the Loess Plateau of  
599 China, *CATENA*, 198, 105028, <https://doi.org/10.1016/j.catena.2020.105028>, 2021b.

600 Colgan, R., Atkinson, C. J., Paul, M., Hassan, S., Drake, P. M. W., Sexton, A. L.,  
601 Santa-Cruz, S., James, D., Hamp, K., Gutteridge, C., and Ma, J. K.-C.: Optimisation  
602 of contained *Nicotiana tabacum* cultivation for the production of recombinant protein  
603 pharmaceuticals, *Transgenic Res*, 19, 241–256,  
604 <https://doi.org/10.1007/s11248-009-9303-y>, 2010.

605 Cornfield, A. H.: The phosphate status of garden soils in relation to soil pH, *Plant*  
606 *Soil*, 5, 243–245, <https://doi.org/10.1007/BF01395899>, 1954.

607 De Laurentiis, V., Secchi, M., Bos, U., Horn, R., Laurent, A., and Sala, S.: Soil  
608 quality index: Exploring options for a comprehensive assessment of land use impacts  
609 in LCA, *Journal of Cleaner Production*, 215, 63–74,  
610 <https://doi.org/10.1016/j.jclepro.2018.12.238>, 2019.

611 Dong, L., Li, J., Zhang, Y., Bing, M., Liu, Y., Wu, J., Hai, X., Li, A., Wang, K.,  
612 Wu, P., Shangguan, Z., and Deng, L.: Effects of vegetation restoration types on soil  
613 nutrients and soil erodibility regulated by slope positions on the Loess Plateau,  
614 *Journal of Environmental Management*, 302, 113985,  
615 <https://doi.org/10.1016/j.jenvman.2021.113985>, 2022a.

616 Dong, L., Li, J., Zhang, Y., Bing, M., Liu, Y., Wu, J., Hai, X., Li, A., Wang, K.,  
617 Wu, P., Shangguan, Z., and Deng, L.: Effects of vegetation restoration types on soil  
618 nutrients and soil erodibility regulated by slope positions on the Loess Plateau,  
619 *Journal of Environmental Management*, 302, 113985,  
620 <https://doi.org/10.1016/j.jenvman.2021.113985>, 2022b.

621 Dou, P., Miao, Z., Wang, J., Huang, J., Gao, Q., Wang, K., and Wang, K.: The  
622 key to temperate savanna restoration is to increase plant species richness reasonably,  
623 *Front. Environ. Sci.*, 11, 1112779, <https://doi.org/10.3389/fenvs.2023.1112779>, 2023.

624 Gao, S.: The Impact of Different Aspects to Vegetation Characteristics and  
625 Composition in *Stipa krylovii* Steppe in Gacha Area—A Case of Alatantaogaotu  
626 Gacha, Abaga County, GSER, 06, 58–64, <https://doi.org/10.12677/GSER.2017.62007>,  
627 2017.

628 Guo, M., Chen, Z., Wang, W., Wang, T., Wang, W., and Cui, Z.: Revegetation  
629 induced change in soil erodibility as influenced by slope situation on the Loess  
630 Plateau, *Science of The Total Environment*, 772, 145540,  
631 <https://doi.org/10.1016/j.scitotenv.2021.145540>, 2021.

632 Guo, Q., Cheng, C., Jiang, H., Liu, B., and Wang, Y.: Comparative rates of wind  
633 and water erosion on typical farmland at the northern end of the Loess Plateau, China,  
634 *Geoderma*, 352, 104–115, <https://doi.org/10.1016/j.geoderma.2019.05.011>, 2019.

635 Guo, W.-Z., Chen, Z.-X., Wang, W.-L., Gao, W.-W., Guo, M.-M., Kang, H.-L., Li,  
636 P.-F., Wang, W.-X., and Zhao, M.: Telling a different story: The promote role of  
637 vegetation in the initiation of shallow landslides during rainfall on the Chinese Loess  
638 Plateau, *Geomorphology*, 350, 106879,  
639 <https://doi.org/10.1016/j.geomorph.2019.106879>, 2020.

640 Hao, P., Zhan, Y., Wang, L., Niu, Z., and Shakir, M.: Feature selection of time  
641 series MODIS data for early crop classification using random forest: A case study in  
642 Kansas, USA, *Remote Sensing*, 7, 5347–5369, <https://doi.org/10.3390/rs70505347>,  
643 2015.

644 Huang, C., Zeng, Y., Wang, L., and Wang, S.: Responses of soil nutrients to  
645 vegetation restoration in China, *Reg Environ Change*, 20, 82,  
646 <https://doi.org/10.1007/s10113-020-01679-6>, 2020.

647 Hupet, F. and Vanclooster, M.: Intraseasonal dynamics of soil moisture  
648 variability within a small agricultural maize cropped field, *Journal of Hydrology*, 261,  
649 86–101, [https://doi.org/10.1016/S0022-1694\(02\)00016-1](https://doi.org/10.1016/S0022-1694(02)00016-1), 2002.

650 Jiang, Q., Zhou, P., Liao, C., Liu, Y., and Liu, F.: Spatial pattern of soil  
651 erodibility factor (K) as affected by ecological restoration in a typical degraded  
652 watershed of central China, *Science of The Total Environment*, 749, 141609,  
653 <https://doi.org/10.1016/j.scitotenv.2020.141609>, 2020.

654 Kar, S. K., Singh, R. M., Patra, S., Sankar, M., Kumar, S., and Singh, A.:  
655 Implication of land use shifting on land degradation and restoration potential of  
656 conservation agriculture in India's North-West Himalayan region, *Geoderma Regional*,  
657 32, e00616, <https://doi.org/10.1016/j.geodrs.2023.e00616>, 2023.

658 Kisand, A.: Distribution of sediment phosphorus fractions in hypertrophic  
659 strongly stratified Lake Verevi, in: *Lake Verevi, Estonia — A Highly Stratified  
660 Hypertrophic Lake*, vol. 182, edited by: Ott, I. and Kõiv, T., Springer-Verlag,  
661 Berlin/Heidelberg, 33–39, [https://doi.org/10.1007/1-4020-4363-5\\_3](https://doi.org/10.1007/1-4020-4363-5_3), 2005.

662 Lardy, J. M., DeSutter, T. M., Daigh, A. L. M., Meehan, M. A., and Staricka, J.  
663 A.: Effects of soil bulk density and water content on penetration resistance,  
664 *Agricultural & Env Letters*, 7, <https://doi.org/10.1002/ael2.20096>, 2022.

665 Li, H., Zhu, H., Qiu, L., Wei, X., Liu, B., and Shao, M.: Response of soil OC, N  
666 and P to land-use change and erosion in the black soil region of the Northeast China,  
667 *Agriculture, Ecosystems & Environment*, 302, 107081,  
668 <https://doi.org/10.1016/j.agee.2020.107081>, 2020.

669 Li, R., Zhang, W., Yang, S., Zhu, M., Kan, S., Chen, J., Ai, X., and Ai, Y.:  
670 Topographic aspect affects the vegetation restoration and artificial soil quality of  
671 rock-cut slopes restored by external-soil spray seeding, *Sci Rep*, 8, 12109,  
672 <https://doi.org/10.1038/s41598-018-30651-y>, 2018.

673 Li, T., Zeng, J., He, B., and Chen, Z.: Changes in Soil C, N, and P

---

674 Concentrations and Stoichiometry in Karst Trough Valley Area under Ecological  
675 Restoration: The Role of Slope Aspect, Land Use, and Soil Depth, *Forests*, 12, 144,  
676 <https://doi.org/10.3390/f12020144>, 2021.

677 Li, W., Yan, M., Qingfeng, Z., and Zhikaun, J.: Effects of Vegetation Restoration  
678 on Soil Physical Properties in the Wind-Water Erosion Region of the Northern Loess  
679 Plateau of China, *Clean Soil Air Water*, 40, 7–15,  
680 <https://doi.org/10.1002/clen.201100367>, 2012.

681 Li, Z., Liu, C., Dong, Y., Chang, X., Nie, X., Liu, L., Xiao, H., Lu, Y., and Zeng,  
682 G.: Response of soil organic carbon and nitrogen stocks to soil erosion and land use  
683 types in the Loess hilly–gully region of China, *Soil and Tillage Research*, 166, 1–9,  
684 <https://doi.org/10.1016/j.still.2016.10.004>, 2017.

685 Liu, L., Gudmundsson, L., Hauser, M., Qin, D., Li, S., and Seneviratne, S. I.:  
686 Soil moisture dominates dryness stress on ecosystem production globally, *Nat*  
687 *Commun*, 11, 4892, <https://doi.org/10.1038/s41467-020-18631-1>, 2020.

688 Masciandaro, G. and Ceccanti, B.: Assessing soil quality in different  
689 agro-ecosystems through biochemical and chemico-structural properties of humic  
690 substances, *Soil and Tillage Research*, 51, 129–137,  
691 [https://doi.org/10.1016/S0167-1987\(99\)00056-2](https://doi.org/10.1016/S0167-1987(99)00056-2), 1999.

692 Meng, Z., Dang, X., Gao, Y., Ren, X., Ding, Y., and Wang, M.: Interactive effects  
693 of wind speed, vegetation coverage and soil moisture in controlling wind erosion in a  
694 temperate desert steppe, Inner Mongolia of China, *J. Arid Land*, 10, 534–547,  
695 <https://doi.org/10.1007/s40333-018-0059-1>, 2018.

696 Moreira, W. H., Tormena, C. A., de Lima, R. P., Anghinoni, G., and Imhoff, S.:  
697 The influence of sowing furrow opening and wetting and drying cycles on soil  
698 physical quality under no-tillage in Southern Brazil, *Soil and Tillage Research*, 204,  
699 104711, <https://doi.org/10.1016/j.still.2020.104711>, 2020.

700 Nabiollahi, K., Golmohamadi, F., Taghizadeh-Mehrjardi, R., Kerry, R., and  
701 Davari, M.: Assessing the effects of slope gradient and land use change on soil quality  
702 degradation through digital mapping of soil quality indices and soil loss rate,  
703 *Geoderma*, 318, 16–28, <https://doi.org/10.1016/j.geoderma.2017.12.024>, 2018.

704 Nichols, K. A. and Toro, M.: A whole soil stability index (WSSI) for evaluating  
705 soil aggregation, *Soil and Tillage Research*, 111, 99–104,  
706 <https://doi.org/10.1016/j.still.2010.08.014>, 2011.

707 Niinemets, Ü.: A review of light interception in plant stands from leaf to canopy  
708 in different plant functional types and in species with varying shade tolerance, *Ecol*  
709 *Res*, 25, 693–714, <https://doi.org/10.1007/s11284-010-0712-4>, 2010.

710 Ortas, I. and Lal, R.: Long-Term Phosphorus Application Impacts on  
711 Aggregate-Associated Carbon and Nitrogen Sequestration in a Vertisol in the  
712 Mediterranean Turkey, *Soil Science*, 177, 241–250,  
713 <https://doi.org/10.1097/SS.0b013e318245d11c>, 2012.

714 Pan, C. and Shangguan, Z.: Runoff hydraulic characteristics and sediment  
715 generation in sloped grassplots under simulated rainfall conditions, *Journal of*  
716 *Hydrology*, 331, 178–185, <https://doi.org/10.1016/j.jhydrol.2006.05.011>, 2006.

717 Pandey, S., Kumar, P., Zlatic, M., Nautiyal, R., and Panwar, V. P.: Recent  
718 advances in assessment of soil erosion vulnerability in a watershed, *International Soil*  
719 *and Water Conservation Research*, 9, 305–318,  
720 <https://doi.org/10.1016/j.iswcr.2021.03.001>, 2021.

721 Peng, S.-L., Hou, Y.-P., and Chen, B.-M.: Vegetation Restoration and Its Effects  
722 on Carbon Balance in Guangdong Province, China, *Restoration Ecology*, 17, 487–494,  
723 <https://doi.org/10.1111/j.1526-100X.2008.00399.x>, 2009.

724 Peres-Neto, P. R., Legendre, P., Dray, S., and Borcard, D.: Variation partitioning  
725 of species data matrices: estimation and comparison of fractions, *Ecology*, 87,  
726 2614–2625, [https://doi.org/10.1890/0012-9658\(2006\)87\[2614:VPOSDM\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[2614:VPOSDM]2.0.CO;2),  
727 2006.

728 Proulx, M. and Mazumder, A.: Reversal of grazing impact on plant species  
729 richness in nutrient-poor vs. nutrient-rich ecosystems, *Ecology*, 79, 2581–2592,  
730 [https://doi.org/10.1890/0012-9658\(1998\)079\[2581:ROGIOP\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1998)079[2581:ROGIOP]2.0.CO;2), 1998.

731 Qin, Y., Feng, Q., Holden, N. M., and Cao, J.: Variation in soil organic carbon by  
732 slope aspect in the middle of the Qilian Mountains in the upper Heihe River Basin,  
733 China, *CATENA*, 147, 308–314, <https://doi.org/10.1016/j.catena.2016.07.025>, 2016.

734 Renard, K. G., Foster, G. R., Weesies, G. A., McCool, D. K., and Yoder, D. C.  
735 (Eds.): *Predicting soil erosion by water: a guide to conservation planning with the  
736 revised universal soil loss equation (RUSLE)*, Washington, D. C, 384 pp., 1997.

737 Salekin, S., Bloomberg, M., Morgenroth, J., Meason, D. F., and Mason, E. G.:  
738 Within-site drivers for soil nutrient variability in plantation forests: A case study from  
739 dry sub-humid New Zealand, *CATENA*, 200, 105149,  
740 <https://doi.org/10.1016/j.catena.2021.105149>, 2021.

741 Schad, P.: World Reference Base for Soil Resources, in: Reference Module in  
742 Earth Systems and Environmental Sciences, Elsevier, B9780124095489106000,  
743 <https://doi.org/10.1016/B978-0-12-409548-9.10496-8>, 2017.

744 Schloter, M., Dilly, O., and Munch, J. C.: Indicators for evaluating soil quality,  
745 *Agriculture, Ecosystems & Environment*, 98, 255–262,  
746 [https://doi.org/10.1016/S0167-8809\(03\)00085-9](https://doi.org/10.1016/S0167-8809(03)00085-9), 2003.

747 Schmiedel, U., Kruspe, M., Kayser, L., and Oettlé, N.: The ecological and  
748 financial impact of soil erosion and its control – a case study from the semiarid  
749 northern cape province, south africa, *Land Degrad. Develop.*, 28, 74–82,  
750 <https://doi.org/10.1002/ldr.2513>, 2017.

751 Schonlau, M. and Zou, R. Y.: The random forest algorithm for statistical learning,  
752 *The Stata Journal*, 20, 3–29, <https://doi.org/10.1177/1536867X20909688>, 2020.

753 Seddaiu, G., Porcu, G., Ledda, L., Roggero, P. P., Agnelli, A., and Corti, G.: Soil  
754 organic matter content and composition as influenced by soil management in a  
755 semi-arid Mediterranean agro-silvo-pastoral system, *Agriculture, Ecosystems &  
756 Environment*, 167, 1–11, <https://doi.org/10.1016/j.agee.2013.01.002>, 2013.

757 Sharma, C. M., Baduni, N. P., Gairola, S., Ghildiyal, S. K., and Suyal, S.: Effects  
758 of slope aspects on forest compositions, community structures and soil properties in  
759 natural temperate forests of Garhwal Himalaya, *Journal of Forestry Research*, 21,  
760 331–337, <https://doi.org/10.1007/s11676-010-0079-y>, 2010.

761 Shi, X., Du, C., Guo, X., and Shi, W.: Heterogeneity of water-retention capacity  
762 of forest and its influencing factors based on meta-analysis in the  
763 Beijing-Tianjin-Hebei region, *J. Geogr. Sci.*, 31, 69–90,  
764 <https://doi.org/10.1007/s11442-021-1833-0>, 2021.

765 Singh, G. and Panda, R. K.: Grid-cell based assessment of soil erosion potential  
766 for identification of critical erosion prone areas using USLE, GIS and remote sensing:  
767 A case study in the Kapgari watershed, India, *International Soil and Water  
768 Conservation Research*, 5, 202–211, <https://doi.org/10.1016/j.iswcr.2017.05.006>,  
769 2017.

770 Singh, J. S. and Gupta, V. K.: Soil microbial biomass: A key soil driver in  
771 management of ecosystem functioning, *Science of The Total Environment*, 634,  
772 497–500, <https://doi.org/10.1016/j.scitotenv.2018.03.373>, 2018.

773 Singh, K. and Pollard, A. G.: Relationship between soil structure, soil cultivation,

---

774 nitrogen uptake and crop growth. III.—Effects of cultivation on the porosity of soil  
775 and its compactness and on crop development and yields, *J. Sci. Food Agric.*, 9,  
776 454–462, <https://doi.org/10.1002/jsfa.2740090712>, 1958.

777 Sun, J., YU, X., Fan, D., Liang, H., Chang, Y., and Li, H.: Impact of vegetation  
778 cover on surface runoff hydraulic characteristics with simulated rainfall, *Acta*  
779 *Ecologica Sinica*, 35, <https://doi.org/10.5846/stxb201310302620>, 2015.

780 Sun, L., Zhang, G., Luan, L., and Liu, F.: Temporal variation in soil resistance to  
781 flowing water erosion for soil incorporated with plant litters in the Loess Plateau of  
782 China, *CATENA*, 145, 239–245, <https://doi.org/10.1016/j.catena.2016.06.016>, 2016.

783 Sun, Y., Wang, Y., Yan, Z., He, L., Ma, S., Feng, Y., Su, H., Chen, G., Feng, Y., Ji,  
784 C., Shen, H., and Fang, J.: Above- and belowground biomass allocation and its  
785 regulation by plant density in six common grassland species in China, *J Plant Res*,  
786 135, 41–53, <https://doi.org/10.1007/s10265-021-01353-w>, 2022.

787 Tamene, G. M., Adiss, H. K., and Alemu, M. Y.: Effect of Slope Aspect and Land  
788 Use Types on Selected Soil Physicochemical Properties in North Western Ethiopian  
789 Highlands, *Applied and Environmental Soil Science*, 2020, 1–8,  
790 <https://doi.org/10.1155/2020/8463259>, 2020.

791 de Vente, J., Poesen, J., Verstraeten, G., Van Rompaey, A., and Govers, G.:  
792 Spatially distributed modelling of soil erosion and sediment yield at regional scales in  
793 Spain, *Global and Planetary Change*, 60, 393–415,  
794 <https://doi.org/10.1016/j.gloplacha.2007.05.002>, 2008.

795 de Vente, J., Poesen, J., Verstraeten, G., Govers, G., Vanmaercke, M., Van  
796 Rompaey, A., Arabkhedri, M., and Boix-Fayos, C.: Predicting soil erosion and  
797 sediment yield at regional scales: Where do we stand?, *Earth-Science Reviews*, 127,  
798 16–29, <https://doi.org/10.1016/j.earscirev.2013.08.014>, 2013.

799 Vincenzi, S., Zucchetta, M., Franzoi, P., Pellizzato, M., Pranovi, F., De Leo, G.  
800 A., and Torricelli, P.: Application of a Random Forest algorithm to predict spatial  
801 distribution of the potential yield of *Ruditapes philippinarum* in the Venice lagoon,  
802 Italy, *Ecological Modelling*, 222, 1471–1478,  
803 <https://doi.org/10.1016/j.ecolmodel.2011.02.007>, 2011.

804 Wang, B., Xue, S., Liu, G. B., Zhang, G. H., Li, G., and Ren, Z. P.: Changes in  
805 soil nutrient and enzyme activities under different vegetations in the Loess Plateau  
806 area, Northwest China, *CATENA*, 92, 186–195,  
807 <https://doi.org/10.1016/j.catena.2011.12.004>, 2012.

808 Wang, B., Zhang, G.-H., Shi, Y.-Y., and Zhang, X. C.: Soil detachment by  
809 overland flow under different vegetation restoration models in the Loess Plateau of  
810 China, *CATENA*, 116, 51–59, <https://doi.org/10.1016/j.catena.2013.12.010>, 2014a.

811 Wang, H., Zhang, G., Li, N., Zhang, B., and Yang, H.: Soil erodibility influenced  
812 by natural restoration time of abandoned farmland on the Loess Plateau of China,  
813 *Geoderma*, 325, 18–27, <https://doi.org/10.1016/j.geoderma.2018.03.037>, 2018.

814 Wang, H., Zhang, G., Li, N., Zhang, B., and Yang, H.: Soil erodibility as  
815 impacted by vegetation restoration strategies on the Loess Plateau of China: Effect of  
816 vegetation restoration on soil erodibility, *Earth Surf. Process. Landforms*, 44, 796–807,  
817 <https://doi.org/10.1002/esp.4531>, 2019a.

818 Wang, H., Zhang, G., Li, N., Zhang, B., and Yang, H.: Variation in soil  
819 erodibility under five typical land uses in a small watershed on the Loess Plateau,  
820 China, *CATENA*, 174, 24–35, <https://doi.org/10.1016/j.catena.2018.11.003>, 2019b.

821 Wang, H., Sun, B., Yu, X., Xin, Z., and Jia, G.: The driver-pattern-effect  
822 connection of vegetation dynamics in the transition area between semi-arid and  
823 semi-humid northern China, *CATENA*, 194, 104713,

---

824 <https://doi.org/10.1016/j.catena.2020.104713>, 2020a.

825 Wang, H., Wang, J., and Zhang, G.: Impact of landscape positions on soil  
826 erodibility indices in typical vegetation-restored slope-gully systems on the Loess  
827 Plateau of China, *CATENA*, 201, 105235,  
828 <https://doi.org/10.1016/j.catena.2021.105235>, 2021a.

829 Wang, S., Zhang, B., Xie, G., Zhai, X., and Sun, H.: Vegetation cover changes  
830 and sand-fixing service responses in the Beijing–Tianjin sandstorm source control  
831 project area, *Environmental Development*, 34, 100455,  
832 <https://doi.org/10.1016/j.envdev.2019.08.002>, 2020b.

833 Wang, S., Zhang, B., Wang, S., and Xie, G.: Dynamic changes in water  
834 conservation in the Beijing–Tianjin Sandstorm Source Control Project Area: A case  
835 study of Xilin Gol League in China, *Journal of Cleaner Production*, 293, 126054,  
836 <https://doi.org/10.1016/j.jclepro.2021.126054>, 2021b.

837 Wang, Z.-J., Jiao, J.-Y., Su, Y., and Chen, Y.: The efficiency of large-scale  
838 afforestation with fish-scale pits for revegetation and soil erosion control in the steppe  
839 zone on the hill-gully Loess Plateau, *CATENA*, 115, 159–167,  
840 <https://doi.org/10.1016/j.catena.2013.11.012>, 2014b.

841 Wen, H., Ni, S., Wang, J., and Cai, C.: Changes of soil quality induced by  
842 different vegetation restoration in the collapsing gully erosion areas of southern China,  
843 *International Soil and Water Conservation Research*, 9, 195–206,  
844 <https://doi.org/10.1016/j.iswcr.2020.09.006>, 2021.

845 Wischmeier, W. H. and Smith, D. D.: Predicting rainfall erosion losses,  
846 *Agricultural Handbook*, 1978.

847 Wu, Y., Yu, X., and Jia, G.: Seasonal Variation of Soil Erodibility Under  
848 Vegetation Restoration in the Agro-pastoral Ecotone of Northern China, *J Soil Sci*  
849 *Plant Nutr*, <https://doi.org/10.1007/s42729-023-01183-w>, 2023.

850 Yang, X., Shao, M., Li, T., Zhang, Q., Gan, M., Chen, M., and Bai, X.:  
851 Distribution of soil nutrients under typical artificial vegetation in the desert–loess  
852 transition zone, *CATENA*, 200, 105165, <https://doi.org/10.1016/j.catena.2021.105165>,  
853 2021.

854 Yimer, F., Ledin, S., and Abdelkadir, A.: Soil organic carbon and total nitrogen  
855 stocks as affected by topographic aspect and vegetation in the Bale Mountains,  
856 Ethiopia, *Geoderma*, 135, 335–344, <https://doi.org/10.1016/j.geoderma.2006.01.005>,  
857 2006.

858 Yükses, T. and Yükses, F.: Effects of altitude, aspect, and soil depth on carbon  
859 stocks and properties of soils in a tea plantation in the humid Black Sea region, *Land*  
860 *Degrad Dev*, 32, 4267–4276, <https://doi.org/10.1002/ldr.4033>, 2021.

861 Zeng, X., Zhang, W., Cao, J., Liu, X., Shen, H., and Zhao, X.: Changes in soil  
862 organic carbon, nitrogen, phosphorus, and bulk density after afforestation of the  
863 “Beijing–Tianjin Sandstorm Source Control” program in China, *CATENA*, 118,  
864 186–194, <https://doi.org/10.1016/j.catena.2014.01.005>, 2014.

865 Zhang, B., Zhang, G., Zhu, P., and Yang, H.: Temporal variations in soil  
866 erodibility indicators of vegetation-restored steep gully slopes on the Loess Plateau of  
867 China, *Agriculture, Ecosystems & Environment*, 286, 106661,  
868 <https://doi.org/10.1016/j.agee.2019.106661>, 2019.

869 Zhang, J., Chen, H., Fu, Z., and Wang, K.: Effects of vegetation restoration on  
870 soil properties along an elevation gradient in the karst region of southwest China,  
871 *Agriculture, Ecosystems & Environment*, 320, 107572,  
872 <https://doi.org/10.1016/j.agee.2021.107572>, 2021.

873 Zhang, L., Cao, W., and Fan, J.: Soil organic carbon dynamics in Xilingol

874 grassland of northern China induced by the Beijing-Tianjin Sand Source Control  
875 Program, *Front. Earth Sci.*, 11, 407–415, <https://doi.org/10.1007/s11707-016-0589-9>,  
876 2017.

877 Zhang, X., Hu, M., Guo, X., Yang, H., Zhang, Z., and Zhang, K.: Effects of  
878 topographic factors on runoff and soil loss in Southwest China, *CATENA*, 160,  
879 394–402, <https://doi.org/10.1016/j.catena.2017.10.013>, 2018.


880 Zhang, X., Adamowski, J. F., Liu, C., Zhou, J., Zhu, G., Dong, X., Cao, J., and  
881 Feng, Q.: Which slope aspect and gradient provides the best afforestation-driven soil  
882 carbon sequestration on the China's Loess Plateau?, *Ecological Engineering*, 147,  
883 105782, <https://doi.org/10.1016/j.ecoleng.2020.105782>, 2020.

884 Zheng, J. Y., Zhao, J. S., Shi, Z. H., and Wang, L.: Soil aggregates are key factors  
885 that regulate erosion-related carbon loss in citrus orchards of southern China: Bare  
886 land vs. grass-covered land, *Agriculture, Ecosystems & Environment*, 309, 107254,  
887 <https://doi.org/10.1016/j.agee.2020.107254>, 2021.

888 Zhou, X., Ke, T., Li, S., Deng, S., An, X., Ma, X., De Philippis, R., and Chen, L.:  
889 Induced biological soil crusts and soil properties varied between slope aspect, slope  
890 gradient and plant canopy in the Hobq desert of China, *CATENA*, 190, 104559,  
891 <https://doi.org/10.1016/j.catena.2020.104559>, 2020.

892 Zhu, G., Deng, L., and Shangguan, Z.: Effects of soil aggregate stability on soil  
893 N following land use changes under erodible environment, *Agriculture, Ecosystems &  
894 Environment*, 262, 18–28, <https://doi.org/10.1016/j.agee.2018.04.012>, 2018.


895 Zhu, M., Yang, S., Ai, S., Ai, X., Jiang, X., Chen, J., Li, R., and Ai, Y.: Artificial  
896 soil nutrient, aggregate stability and soil quality index of restored cut slopes along  
897 altitude gradient in southwest China, *Chemosphere*, 246, 125687,  
898 <https://doi.org/10.1016/j.chemosphere.2019.125687>, 2020.



899

900 **Fig. 1.** Location map of the sampling points in the study area. The first letter: D, G, S  
901 and W represent degraded land, grassland, shrubland and woodland. The sampling  
902 sites from west to east were: DAA, degraded land; GAS, *Artemisia sacrorum*; GAM,  
903 *Astragalus melilotoides*; WPS, *Pinus sylvestris*; WLG, *Larix gmelinii*; SHR,  
904 *Hippophae rhamnoides*; SCK, *Caragana korshinskii*.






906


907 **Fig. 2.** Variation of understory vegetation characteristics and soil nutrients with slope  
 908 aspects. BH, biomass height; R, richness; BC, biomass coverage; AGB, aboveground  
 909 biomass; BGB, belowground biomass; LB, litter biomass; SOC, soil organic carbon;  
 910 TN, total nitrogen; TP, total phosphorus; W, west; N, north; S, south; E, east. Different  
 911 letters indicate significant differences among different seasons at  $P < 0.05$  level.

912



913


914 **Fig.3.** Variation of comprehensive soil nutrient, erodibility and quality index with  
 915 slope aspects. CSNI, comprehensive soil nutrient index; CSEI, comprehensive soil  
 916 erodibility index; CSQI, comprehensive soil quality index. Different letters indicate  
 917 significant differences among different slope aspects at  $P < 0.05$  level.



919

920 **Fig. 4.** Variation of comprehensive soil nutrient, erodibility and quality index with  
 921 land use. Different letters indicate significant differences among different land use  
 922 types at  $P < 0.05$  level.


923




924

925 **Fig. 5.** Variation in comprehensive soil quality index with vegetation types along  
 926 slope aspects. WDAA, *Artemisia annua*; WGAM, NGAM and SGAM, *Astragalus*  
 927 *melilotoides*; NGBI, *Bothriochloa ischaemum*; EGSM, *Artemisia sacrorum*,  
 928 *Astragalus melilotoides*; WGCP, NGCP and SGCP, *Capillipedium parviflorum*;  
 929 WSHR, NSHR, SSHR and ESHR, *Hippophae rhamnoides*; WSCK, NSCK, SSCK

930 and ESCK, *Caragana korshinskii*; WWLG, NSWG, SSWG and ESGW, *Larix*  
 931 *gmelinii*; WWPS, NWPS, SWPS and EWPS, *Pinus sylvestris*. Different letters  
 932 indicate significant differences among different seasons at  $P < 0.05$  level.



933  
 934 **Fig. 6.** Results of redundancy analysis (RDA) among soil quality parameters and  
 935 characteristics of vegetation and soil on four slope aspects. BH: biome height; R:  
 936 richness; BC: biome coverage; AGB: aboveground biomass; BGB: belowground  
 937 biomass; LB: litter biomass; Sand: sand content; Silt: silt content; Clay: clay content;  
 938 SWC: soil water content; SBD: soil bulk density; SOC: soil organic carbon; TN: total  
 939 nitrogen; TP: total phosphorus; SHC, saturated hydraulic conductivity; SDR, soil  
 940 disintegration rate; MWD, mean weight diameter; K, soil erodibility factor; SSSI, soil  
 941 structure stability index; SCAI, SOC cementing agent index.



942  
943

944 **Fig. 7.** Correlation analysis of CSNI, CSEI and CSQI with vegetation and soil  
945 characteristics. Red indicates a positive correlation, blue indicates a negative  
946 correlation, and the color depth indicates Pearson coefficients \* $p < 0.05$ , \*\* $p < 0.01$   
947 and \*\*\* $p < 0.001$ ,  $n = 84$ . CSNI, comprehensive soil nutrient index; CSEI,  
948 comprehensive soil erodibility index; CSQI, comprehensive soil quality index.

949

950 **Table 1**

951 The two-way ANOVA result for soil nutrient and erodibility. SOC: soil organic carbon;  
952 TN: total nitrogen; TP: total phosphorus; CSNI: comprehensive soil nutrient index;  
953 SHC: saturated hydraulic conductivity; SDR: soil disintegration rate; MWD: mean  
954 weight diameter; K: soil erodibility factor; SSSI: soil structure stability index; SCAI:  
955 SOC cementing agent index; CSEI: comprehensive soil erodibility index; CSQI:  
956 comprehensive soil quality index.

soil variables	Land use type		Slope aspect		Land use ×Slope aspect	
	F	P	F	P	F	P
Soil nutrient						
SOC	1200.37	0.000	50.985	0.000	5.818	0.000

TN	520.016	0.000	79.681	0.000	24.354	0.000
TP	382.353	0.000	6.718	0.000	6.764	0.000
CSNI	832.059	0.000	46.447	0.000	6.851	0.000
Soil erodibility						
SHC	824.538	0.000	54.173	0.000	52.672	0.000
SDR	799.513	0.000	6.632	0.001	3.956	0.000
MWD	1667.15	0.000	180.654	0.000	10.673	0.001
K	859.009	0.000	14.423	0.000	23.822	0.000
SSSI	517.098	0.000	41.05	0.000	26.717	0.000
SCAI	693.653	0.000	15.553	0.000	6.623	0.000
CSEI	1120.468	0.000	38.983	0.000	6.369	0.000
Soil quality						
CSQI	642.05	0.000	103.399	0.000	35.679	0.000

957

958 **Table 2**

959 Soil erodibility indicators of different land use types at different slope aspect (mean  
960  $\pm$  SD). SHC, saturated hydraulic conductivity; SDR, soil disintegration rate; MWD,  
961 mean weight diameter; K, soil erodibility factor; SSSI, soil structure stability index;  
962 SCAI, SOC cementing agent index. Different capital letters indicate significant  
963 differences between slope aspects ( $p < 0.05$ ), different lowercase letters indicate  
964 significant differences between the land use types ( $p < 0.05$ ).

Slope aspect	Land use	SHC mm min <sup>-1</sup>	SDR g min <sup>-1</sup>	MWD mm	K t·hm <sup>2</sup> ·h·hm <sup>-2</sup> MJ <sup>-1</sup> ·mm <sup>-1</sup>	SSSI g kg <sup>-1</sup>	SCAI mm kg <sup>-1</sup> g <sup>-1</sup>
	Degraded land	0.13±0.02cC	1.64±0.19aA	0.79±0.02dD	0.33±0.01aA	0.25±0.01dB	20.23±0.81aA
W	grassland	0.28±0.04bC	0.29±0.04cA	1.83±0.06bD	0.26±0.01dA	0.51±0.06bB	9.09±0.97bA
	shrubland	0.32±0.07bC	0.82±0.53bA	2.38±0.32aD	0.32±0.01bA	0.46±0.04cB	9.03±0.80bA
	Woodland	0.53±0.06aC	1.58±0.07aA	1.46±0.15cD	0.27±0.01cA	0.61±0.05aB	7.53±0.70cA
N	grassland	0.28±0.03bB	0.26±0.02cB	2.32±0.47bC	0.31±0.01aAB	0.50±0.06aA	8.30±0.94aB

---

	shrubland	0.31±0.04bB	0.73±0.44bB	2.84±0.12aC	0.29±0.04aAB	0.58±0.08aA	8.14±0.95aB
	Woodland	0.60±0.07aB	1.26±0.17aB	1.76±0.29cC	0.29±0.01aAB	0.57±0.03aA	7.90±0.39aB
	grassland	0.93±0.11bA	0.24±0.01cBC	3.28±0.04aB	0.25±0.01cB	0.51±0.10bA	9.16±1.74aB
S	shrubland	1.31±0.20aA	0.40±0.11bBC	3.32±0.06aB	0.31±0.01aB	0.53±0.03bA	8.27±0.40abB
	Woodland	1.45±0.14aA	1.17±0.06aBC	3.25±0.07aB	0.28±0.01bB	0.67±0.10aA	6.94±1.00bB
	grassland	1.55±0.18aA	0.24±0.01cC	4.06±0.14aA	0.29±0.01aB	0.59±0.02bA	7.28±0.29bB
E	shrubland	1.71±0.06aA	0.31±0.07bC	3.46±0.09bA	0.26±0.02bB	0.61±0.05bA	8.18±0.89aB
	Woodland	1.73±0.12aA	0.38±0.03aC	3.42±0.10bA	0.28±0.01bB	0.71±0.05aA	6.41±0.44cB

---

965