

Offshore seismicity clusters in the West Iberian Margin illustrated by two decades of events.

Gabriela Fernandez-Viejo¹, Carlos Lopez-Fernandez¹, Patricia Cadenas²

Con formato: Portugués (Portugal)

¹Department of Geology, University of Oviedo, Jesús Arias de Velasco s/n 33005 Oviedo, Spain

²Memorial University of Newfoundland, Dept. Earth Sciences

Correspondence to: Gabriela Fernandez-Viejo (fernandezgabriela@uniovi.es)

Con formato: Inglés (Estados Unidos)

Abstract. An analysis of two decades (2003-2022) of seismicity recorded by the Spanish and Portuguese seismic networks along the West Iberian passive margin has resulted in a better understanding of the distribution of moderate seismic activity in this intraplate submarine area. The study provides a precise trend of specific alignments inferred from the density maps of seismicity, giving an accurate depiction of event distribution along two wide stripes that extend ~~for~~ 700 km ~~long~~ through the ocean floor in a WNW-ESE direction. These bands are parallel to the Africa-Eurasia plate boundary but are distinctly separated from its related seismicity by approximately 300 km and 700 km, respectively. This is a sufficient distance to be considered as intraplate activity. When trying to relate this seismicity to structural and geophysical features, a conclusive picture doesn't emerge. The earthquakes occur indiscriminately across thinned continental, hyperextended, and exhumed mantle rift domains. They fade out in the proximity of undisputed oceanic crust, but some events extend beyond. The hypocentral depths signal a considerable amount of events nucleating in the upper mantle. The focal mechanisms, although scarce, are predominantly strike-slip. Considering these observations, hypothesis ranging from subduction initiation, development of strained corridors or local structures of the margin, ~~have been are~~ discussed in order to explain this relatively anomalous seismicity. However, some of them ~~do not portray lack~~ convincing arguments, while others are too ~~unspecifie vague~~. None of them are flawless, suggesting that several factors may be at play. Despite being one of the most probed passive margins in the world, the present geodynamical status of the West Iberian Margin manifested in its modern seismicity remains unknown. Interpreting this data within a global tectonic plate framework, together with the potential addition of seafloor seismometers, may provide the key to understanding this activity along one of the most archetypical margins of the Atlantic Ocean.

25 1 Introduction

1.1 Physiographic aspects and brief geological history of the West Iberian Margin

The West Iberian Margin (WIM) is an 850 km long passive margin parallel to the N-S trending rectilinear coast of western Iberia that resulted from a tectonic evolution marked by a multiphase rifting history (Pinheiro et al., 1996; Péron-Pinvidic et al., 2007; Tucholke et al., 2007; Alves et al., 2009; Pereira et al., 2016; Granado et al., 2021). Currently, the dynamics of the margin are influenced by two significant plate limits. To the west lies the mid-Atlantic ridge (MAR) constructive plate boundary. Meanwhile, to the south, the margin is impacted by the complex NW Africa-SW Eurasia plate limit, known as the Azores-Gibraltar Fault Zone (AGFZ) (Fig. 1). To the north, the West Iberian Margin (WIM) adjoins the E-W trending southern Biscay passive margin, which ~~locates is located~~ to the south of the extinct Bay of Biscay ridge.

Along the WIM, the 30-85 km wide continental platform is noticeably intersected by deep E-W/NE-SW trending canyons, namely Nazaré (NZc), Setubal (STc), and San Vicente (SVC) (Fig. 1). Northwards of NZc, 200 km from the coast and separated from the platform by the Galicia Interior Sedimentary Basin (GIB), the Galicia Bank (GB) stands out, which is an elevated area with a relief of over 3,500 m, along with other minor topographic seamounts. Westwards of the GB, the Iberian Abyssal Plain (IAP) emerges as an expansive and flat ocean floor that reaches a depth of 5,300 m. As we move towards the MAR, the

surface becomes considerably rough, indicative of oceanic crust. To the west of the IAP, the prominent Kings Through (KT) 40 is a 450 km long and WNW-ESE trending scar in the oceanic floor (Fig. 1) (Srivastava and Roest, 1992), which consists of a central depression flanked by ridges. It is proposed that this feature formed from 56 to 25 Myr during the-Mesozoic rifting along the crest of a hotspot, resulting in an aseismic ridge, which represented the plate boundary between Iberia and Eurasia. This boundary shifted and was active along the Bay of Biscay between 49-36 Myr, extending eastwards along the Pyrenees (Roest and Srivastava, 1991) and finally moved to its present location along the AGFZ at Chron 13 (Macchiavelli et al., 2017). 45 Between NZc and STc, the Extremadura Spur (Es) separates the IAP from the Tagus Abyssal Plain (TAP), a 200 km wide flat area and situated at 5,000 m depth. The TAP is limited to the west by the Madeira Tore Rise and to the southeast by the Gorringe Bank, one of the principal seismogenic sources in the region. The presence of ~~these reliefs~~areas of relief has been associated with N-S normal faults and ~~NE-SWW-E~~ thrusts (Boillot et al., 1988a, Pinheiro et al., 1996; [Álvarez-Marrón](#), et al., 1997, Vazquez et al., 2008; Duarte et al., 2013; Sallarès et al., 2013; [Martínez-Lorient et al., 2013](#)), 50 which in turn can be related to ~~the~~margin reactivation during the Mesozoic and the Paleogene (e.g., Jiménez-Munt y Negredo, 2013). Furthermore, the Extremadura Spur and related reliefs are connected to extensive Late Cretaceous alkaline magmatic occurrences (Pereira et al., 2016; Escada et al., 2022).

The mid-Atlantic ridge (MAR), a constructive plate boundary situated about 1,500 km west of the WIM, separates the North American plate from the Eurasian plate. With a slow-spreading rate, it accretes at a rate of 24-26 mm/y at 43°N (Somoza et 55 al., 2019). Unlike other areas of the North Atlantic, the MAR is remarkably devoid of significant transform faults, spanning over 1400 km without any such faults between 40.6°N and 43.6°N and between 43.6°N and 47.6°N, except for short ones with negligible displacement. This unique characteristic of the ridge in this section of the North Atlantic contrasts with its displacement by relevant transform faults both to the south and north of the WIM ([fig. 1](#)).

The AGFZ plate boundary that limits the WIM southwards is structurally much more complex. The Southwest Iberian Margin 60 (SWIM) faults (Zitellini et al., 2009; Martínez-Lorient et al., 2013) connects the Gloria Fault with the eastern part of the Africa-Europe plate boundary (Fig. 1). [To the East of the Strait of Gibraltar, the plate boundary is defined by a right-lateral transpressive shear one \(Morel and Meghraoui, 1996\). And in a The AGFZ is sometimes described as a diffuse convergent plate boundary to the East of Gibraltar due to the ample area of deformation. But a recent seismic study interprets Gomez de la Peña et al., 2002 interpret](#) long and active faults accommodating most of the deformation in the Western Mediterranean [\(Gomez de la Peña et al., 2022\)](#)

[The Mediterranean. The kinematic evolution of the Iberian margin during the Atlantic opening—along the WIM abyssal plain](#) is well determined by magnetic anomalies after the Cretaceous normal polarity superchron (Roest and Srivastava, 1991). However, the early rift to drift seafloor evolution from 200 Myr to 83 Myr, remains controversial, mainly due to the dual assignment of the J anomaly either to a real isochron (e.g., Barnett-Moore et al., 1996) or to a polygenic magmatic event right 70 before oceanic inception (e.g., Nirrengarten et al., 2017).

1.2 Crustal structure and domains of the WIM

The WIM fits into the category of a non-volcanic passive margin with a narrow platform (Lymer and Reston, 2022). It developed conjugate to the Newfoundland margin during the Mesozoic rifting (Péron-Pinvidic and Manatschal, 2009). Its crustal structure, although debated in [minor aspects](#)detail, is formed by a series of structural domains from East to West (e.g., 75 Péron-Pinvidic and Manatschal, 2009; [Ramos et al., 2016](#); Druet et al., 2019; Merino et al., 2021) and includes a Variscan basement (Schulmann et al., 2022). The segmentations observed within the WIM resulted from the northward migration of extension and successive late Jurassic to early Cretaceous extensional phases (e.g., Malod and Mauffret, 1990; Péron-Pinvidic et al., 2007; Tucholke et al., 2007; Merino et al., 2021).

Following the onset of breakup within the Pangea supercontinent 200 Myr ago, the pre-Jurassic template of the North Atlantic 80 was subjected to three main phases of rifting that took place from Late Triassic to Early Jurassic, Late Jurassic to Early

Cretaceous, and during the Mid Cretaceous. The different rifting stages led to a system of tilted blocks limited by westward-dipping extensional detachment faults (Lymer and Reston, 2022), which show a N-S orientation in the North and variable trends in the South. At the end of rift-related deformation, during the Late Cretaceous, the WIM experienced continental break-up (Soares et al., 2012; Welford and King, 2021), leading to irregular regions of exhumed and serpentized mantle (Afilhado et al., 2008; Merino et al., 2021; Grevemeyer et al., 2022). The onset of seafloor spreading occurred at least in two pulses, the first one at the end of the Jurassic between the Grand Banks and the TAP, and a second one during the early Cretaceous between the IAP and the Grand Banks, following the temporal northward opening of the Atlantic Ocean.

85 As a consequence of the later Alpine orogeny that formed the actual reliefs on land, there was an important compressive deformation, which produced the reactivation of some extensional faults as thrusts, or as transpressive strike slip faults (e.g., Pereira et al., 2011; Martínez-Loriente et al., 2013; Ramos et al., 2016; Terrinha et al., 2019). The Cenozoic deformation abruptly terminates against the Ocean-Continent transition (OCT).

90 Published works investigating the Mesozoic rift domains in the WIM distinguish two sectors. In the northern sector, the continental hyperextended crust stretches for more than 300 km and the exhumed mantle domain extends westwards of the GB (Druet et al., 2019). In the southern sector, contrastingly, the hyperextended crust extends for 200 km before reaching the 95 same transition (Pedrera et al., 2017). The OCT extends between 12°10'W and 12°30'W in the IAP (Whitsmarsh et al., 1990) and it would be interpreted to extend N°10 for 130 km until Extremadura Spur. According to Grevemeyer et al. (2022), the OCT would run between 12.5°W and 13°W and in the TAP, while Merino et al. (2021) locates the OCT between 13-13.5°W.

100 Crustal thickness based on refraction profiles north of the Extremadura Spur, report thicknesses that shallow from almost 30 km close to the coast to 15 km in the ocean side with a quick thinning happening in 50 km extension (Dean et al., 2000). Based on seismic-derived gravity modelling, Amigo Marx et al. (2022), differentiate an 18 km thick proximal domain, a <18 km thick extended domain, and a < 7km thick hyperextended domain. The exhumed mantle domain has been drilled at a serpentinite ridge (Boillot et al., 1988b). Westwards of the Extremadura Spur, a thin lower crust has been interpreted beneath a dense upper crust. The thickness of the crust decreases quickly towards the West, while it does so gradually towards the 105 North, creating two distinct domains. The first one is located near the continent and characterized by a thin and light crust, while the second one is further away and has a denser and thicker crust (Amigo-Marx et al., 2022).

110 Using seismic, wells and gravity data, Granado et al. (2021) modelled a more complex 3D crustal scale structure of the WIM with two necking domains along the GIB and the DGMDeep Galicia basin in the west of the Galicia Bank, suggesting two rift directions. The GIB encompasses localized regions of significantly thinned crust that are segmented by NE-SW trending transfer zones and it has been interpreted as a failed rift (Murillas et al., 1990; King et al., 2020). The GB is a 15-20 km thick block of continental crust with relatively low thinning (Péron-Pinvidic and Manatschal, 2010; King et al., 2020). The North has a narrower exhumed mantle domain than the southern WIM. The outer necking zone merges southwards with the eastern one following a NW-SE orientation. The southern sector has a narrow continental domain, with a broader exhumed mantle domain.

115 The limit between the North and South-Central WIM, follows the NW-SE orientation that is observed by Granado et al., 2021 in the Bouguer gradient anomaly maps THD-(total horizontal derivative) and dZ(vertical derivative). According to these authors, Granado et al. (2021) it relates to the Variscan fabric onshore, suggesting unrecognized transfer structures to explain the transition. A similar transfer zone SW of the GB was suggested earlier (Pinheiro et al., 1992; Whitsmarsh et al., 1993) based on the analysis of magnetic anomalies and named as the Figueira Fracture zone, which would represent an original 120 fracture zone born at the time of initiation of seafloor spreading.

So, within the general dichotomy North WIM-South WIMvariations of the WIM, we can consider in our study area two of the three WIM segments from north to south: the Galicia margin that includes the GB and IAP, as the northern segment, and the southern segmentcentral segment, that includes the Extremadura Spur and the TAP. The seismicity related to the third and

southernmost segment of the WIM, the so-called Southwest Iberian Margin is out of the scope of this study, as it is considered
125 related to the AGFZ plate boundary (Buorn et al., 1988).

1.2 Stress state, seismicity and kinematics in the WIM

Seismicity is the expression of active stresses within plates, and these forces are intimately linked to kinematics. The so-called Iberian plate takes its name from periods where effectively, the Iberian Peninsula and part of its margins acted as an independent minor tectonic plate between the surrounding and larger European and African plates, for instance from Late

130 Eocene (Chron 18.42 Ma) to early Miocene time (chron 6c, 24 ma)—(e.g., (Roest and Srivastava, 1991). At earlier periods, Iberia was attached to Africa, having a plate boundary running in the northern part through the Bay of Biscay and the actual Pyrenees, and subsequently, moved in solidarity with Europe. The presence of thick post-orogenic and post-tectonic units offshore sealing the Biscay accretionary wedge (Álvarez-Marrón et al., 1997; Fernández-Viejo et al., 2012) among other things confirm that Iberia is now part of the Eurasian plate without discussion. Nevertheless, Iberian kinematics have

135 been for a long time a cause of study and controversy. The M sequence of magnetic anomalies along the Newfoundland margin and WIM forms one of the primary constraints for plate kinematic reconstructions characterising the motion of Iberia during the Mesozoic (Vissers and Mejer, 2012; Sibuet et al., 2004, Barnett-Moore et al., 2016 and references therein). However, the validity of these anomalies has been questioned (Bronner et al., 2011). The difficulty in distinguishing different crustal types only adds uncertainty (Granado et al., 2021, and references therein). The seafloor spreading history of the western and northern

140 margins of Iberia has been described involving successive jumps of the plate boundary between Iberia, Europe, and Africa from north (Bay of Biscay axis, and KT) to south (AGFZ) (Roest and Srivastava, 1991; Macchiavelli et al., 2017).

Complexity in kinematics is normally inversely proportional to the size of the plate. Therefore, numerous models of Iberia whereabouts-kinematics have been published in the last decades, the most modern ones approaching the problem through deformable plate tectonic models (King and Welford, 2022). These have provided insight into the kinematic role of continental

145 blocks and their interplay with large and micro-tectonic plates during the formation of the north Atlantic rifted margins (e.g., Nirrengarten et al., 2018; Peace et al., 2019; King and Welford, 2022).

Macchiavelli et al. (2017) indicates a spreading rate of 23 mm/yr north of 40°W latitude and 20.5 mm year south of 38 °W and 2 mm/yr of N-S convergence between Iberia and Africa since the Miocene. The earlier partitioning of convergence between

150 north and south became totally accommodated by the southern boundary from early Miocene onwards. The total convergence oscillates between 3-6 mm/y (Fernandes et al., 2003; Nocquet, 2012; Serpelloni, et al., 2004). It is characterized around the

Azores archipelago by an NNE-SW extension that passes to the East to a dextral strike-slip border along the Gloria fault. Along the Gulf of Cádiz, the boundary becomes an obliquely convergent boundary, N45°W (Ribeiro et al., 1996; Stich et al., 2006; Pedrera et al., 2011) with a velocity of 4.5 mm/year (Nocquet and Calais, 2004; Stich et al., 2006), that accommodates

155 intense deformation along shear zones (Terrinha et al., 2009), and through the reactivation of thrusts in the SWIM (Martinez-

Loriente et al., 2013; Ramos et al., 2016). A neotectonic model across the southern plate boundary indicates that friction coefficients between 0.06 and 0.1 are the best solutions and that the maximum slip rates predicted outside the ridge occur along

160 the Gloria fault, Although the western segment of the Gloria fault which is almost devoid of seismicity (Jimenez-Munt and Negredo, 2003), which could indicate an expression of fault locking or slow creeping (Hensen et al, 2019), the central and eastern segments have a record of high magnitude earthquakes, the M8.4 event of 1941 or the M7.1 event in 1931. The deformation here is attributed to a wide zone area (Lay, 2019) or to the existence of hydrated mantle permitting water circulation along fractures down to 12 km below the sea floor (Batista et al. 2016, 2017)

Geodetic velocities (Garate et al., 2015, Palano et al., 2015) also reveal an important deformation in the southern sector of Iberia, while in the interior of the Iberian Peninsula the crustal deformation is locally accommodated, with rates of less than

15 nanostrain/yr (15 mm/yr/1000 km). The stations situated in central and northern Portugal are displaced to the north around

165 1 mm/yr (Palano et al., 2015). Along the NW margin, the geodetic data provide evidence of E-W contraction, with rates of

Con formato: Francés (Francia)

55 nanostrain/yr (55 mm/yr/1000 km). In the WIM, there is movements to the NW reaching 3 mm/year. In the active stress map of the Iberia, the transition from a compressive to an extensivonal regime occurs just around the Lisboa-Nazaré latitude (Fig.—1). A peculiarity worthy of note is the large amount of compression that the map shows exists in this area and the radical change to extension north of this transition.

170 Within the global picture, the present-day NNW-SSE compressional stress, has been practically constant since the Miocene (Andeweg et al., 1999; de Vicente et al., 2008 and references therein), and oriented at high angles to both north and west coasts of Iberia, being subparallel to the tensile stresses induced by the lateral density variations along these Iberian margins (Andeweg et al., 1999).

175 These complex systems of forces and movements outlined above is are the engine producing the intraplate earthquakes in the WIM. Seismicity in the southernmost WIM is overwhelmingly clustered, as expected on both plate boundaries, along the AGFZ, with significantly complex patterns, and along the MAR. However, within mainland Western Iberia and along adjacent offshore domains, away from both boundaries, seismicity has been repeatedly classified as diffuse. There is a general consensus on noting that the magnitude, and the number of events are difficult to reconcile with a typical intraplate location (Custodio et al., 2015; Kruger et al., 2020; Veludo et al., 2017; Duarte et al., 2013; Ribeiro, 2002).

180 The seismicity at the WIM hasn't been directly approached in published studies, and the absence of Ocean Bottom Seismometers within the central and northern WIM, except for some local studies in neighbouring areas. (Kruger et al., 2020) has discouraged an in depth profound study of their characteristics. In an extended study including mainland Portugal, Custodio et al. (2015) indicated that the marine earthquakes do collapse into well-defined clusters while cumulative seismic moment and epicentre density decrease from South to North. The analysis of some focal mechanisms corroborates confirms

185 that Portugal is under horizontal pressure in the NNW-SSE direction, with a great proportion of strike-slip solution and some reverse oblique mechanisms onshore (Borges et al., 2001).

2 Gathering seismicity in the WIM

190 The data gathered for this study belong to the permanent seismic networks of Portugal and Spain. In the Portuguese network, operated by the Instituto Português do Mar e da Atmosfera (IPMA) there is a network of 50 land stations, 5 in the Madeira archipelago and another 35 in the Azores. The Spanish network, operated by the Instituto Geográfico Nacional de España (IGN), maintains 60 stations on land and 56 in the Canary Islands. We have retrieved data from the available catalogues since January 2003 until December 2022 in the area between 38°N and 45°N and 8°W and 20° W, for earthquakes showing magnitudes > 2.5.

195 In total, 278 events were collected from the IPMA network and 649 from the IGN network. Integrating both catalogues, and selecting, in case of coincidence, those with the lesser error in location, we elaborated compiled a final working catalogue of 708 events. From these, we established several requisites of quality required to be included in this study (RMS < 1, Smax < 25, Smin < 25 and Err < 75), resulting in 606 events, 352 from the Spanish network and 254 from the Portuguese network. (Table 1).

200 Likewise, we have included the existing historical seismicity data in the IGN catalogues and collected included the work of Ferrao et al. (2016), which contains information on 10 historical events that occurred within the study area. The 9 focal mechanisms considered in this work from events occurring in the marine domain have been obtained from 14 focal mechanisms included in the CMT Catalogue (Dziewonski et al., 1981; Ekström et al., 2012) and from the IGN Seismic Moment Tensor Database (Rueda and Mezcua, 2005).

205 We should be aware of some of the data limitations and reliability. The seismotectonic interpretations in continental margins under the sea present an added difficulty due to the inconclusive mapping of structures in the sea floor. Despite being well probed, the WIM contains many highs, roughness, and bathymetric heterogeneities that make it difficult to map tectonic

features, specifically faults that could act as seismogenic sources. The structural variability observed along the margin added uncertainty in the data processing when introducing velocity models in the location programs of the events. Additionally, crustal depths in the margin vary, but a considerable number of events show hypocentres below the interpreted Moho depth.

210 The acquisition mode poses the second limitation as the land stations are located to the east, leading to a biased location error. This results in uncertainty in some locations and the lack of full azimuth to relocate with confidence. To minimize the impact of these shortcomings, only the events that comply with the requirements were considered reliable and retrieved:

- a) Magnitude should be over > 2.5.
- b) Events with location uncertainty of < 25 km in plan with average values < 10 km, and 75 km in depth. It is precisely 215 this last parameter the one that poses the main problem, although the average error estimated has been of 4.9 km.
- c) The average GAP (maximum angle separating two seismic stations, measured from the epicentre of an earthquake) < 183, for all the events used in this study being the ones with higher GAP discarded for this purpose.

220 The information collected was integrated through a geographic information system with (i) bathymetric information provided by the European Marine Observation and Data Network (EMODnet Bathymetry Consortium, 2020), (ii) geological structures mapped in the margin-continental platform (Somoza et al., 2021), (iii) gravity and magnetic data (Granado et al., 2021, GeoMapAPP-Ryan et al., 2009), and (iv) data from crustal Moho Depth (Granado et al., 2021; Whitmarsh, 1990; GeoMapApp - Ryan et al., 2009). The ArcMap v.10.3.1 (ESRI) program was used.

3 Results, ~~elusters~~clusters, and trends

225 3.1 Distribution, ~~magnitudes~~magnitudes, and focal mechanisms of the events in the WIM

Figure 2 displays the concentration of activity over the last 20 years, which extends from the coast to 198°W. The figure shows the distribution of events collected and processed, represented on a marine bathymetric chart (Fig. 2a), along with a density map of events (Fig. 2b) that has been calculated.

According to those results, events are more abundant at the coast, blending in with the onshore, more dispersed seismicity.

230 Going-seawards, we start to recognise a subtle clustering, which seems to define two separated diffuse bands that extend up to 18°W longitude and become narrower with distance. Further ~~up~~ towards the MAR, there ~~is-are~~ some isolated events. The orientation of these bands is about N80°W. Although these blurred alignments have been inferred before (Custodio et al., 2015) the discrimination of events made in this study ~~permits to get rid~~allows removal of badly located earthquakes that add to fuzziness ~~giving place to~~resulting in a sharper image of these two seismic alignments. However, the separation in two bands 235 is not that simple. The density maps in figure 2b also show that the northern band is composed of two distinct clusters that while following the same rough NW-SE trend, they are separated for more than 100 km band space were seismicity is very scarce.

Figure 3 represents the depth distribution of events, superimposed ~~to-on~~ crustal ~~domain~~ cross-sections representative of the latitudes of the alignments (Granado et al., 2021). The bands observed are named in this study as the Galicia (N) ~~for the~~ 240 Northern segment of the WIM and the Lisbon one~~band~~(S) ~~for the~~ Central segment of the WIM. Specifically, they can be described by particular characteristics:

245 a) Events offshore Galicia, Northern WIM (North). This density band of events is oriented N75°W from the coast until 17.5°W. It spreads over 670 km and ends in the proximity of the bottom topographic expression of the Azores-Biscay rise (Fig. 2a). It seems there is a clustered seismicity in two main spots along the band with a ~~middle~~-gap in the middle over 100 km extension.

Offshore in this sector during the studied period, we registered a total of 165 events over 2.5, (8 events/yr), and 21 over 4, (1/yr). Most of them are over 20 km depth, and progressively shallower to the west. The depths of events include mantle depths

below the geophysical inferred Moho depth, which rises from 22 km ~~in~~ at the coast up to 11 km oceanwards, at the end of the lineation. The β -value is 0.87.

250 In profile 1 (Fig.-3), which represents the crustal cross-section type for the Galicia Margin, there is an abundance of events in the thickest crustal areas, corresponding to the GB or to the proximal domain. The number of events is larger at the transition between ~~hyperextended crust and exhumed mantle~~ the westernmost necking and hyperthinned domains interpreted by Granado et al. (2021). There is ~~an arguably but~~ noticeable ~~50 km wide~~ gap in event distribution west of the Galicia Bank, followed by a second cluster in the distal margin, which coincides with an area of topographic roughness in the ocean bottom.

255 The figures also show that there is a non-negligible percentage of events inside the uppermost mantle, especially within the transition between the ~~hyperextended westernmost necking~~ and ~~exhumed mantle~~ hyperthinned domains. In a N-S crustal profile within the Galicia Bank cluster, ~~a~~ particular set of south-dipping earthquakes can be observed ~~in~~ (profile 5a, Fig.-3). This is interpreted as the expression of the thrusts, resulting from the limited Alpine convergence ~~and incipient subduction in the Bay of Biscay~~ that led to the partial closure of the Bay ~~of Biscay~~ (Alvarez Álvarez-Marreón et al., 1997; Ayarza et al., 2004;

260 Fernández-Viejo et al., 2012). Some of the focal mechanism in this area give a component of ~~in~~ reverse fault, which would agree with this interpretation.

b) ~~Events offshore Lisbon, Central WIM (South)~~. This second band runs ~~along for~~ 670 km between the coast of Portugal and 16.8°W. We can ~~subdivide define~~ two ~~subclusters~~ ~~subtrends~~, the first one oriented N80°W from the coast until 12.5°W. 320 km ~~away~~ part, a small bend occurs ~~adopting a position to an orientation~~ N75°E along a further 160 km until 16.8W.

265 ~~This band presents more events than the Galicia one~~ In the Central WIM, seismicity is much more abundant than in the Northern WIM, although magnitudes are similar. In this sector, for the 20-year period analysed, we ~~got recognised~~ a total of 295 events with magnitude over 2.5, (15 events/yr.), 20 over 4-4.9, (1/yr.) and 3 over 5, (2 in the last decade). Most of them are concentrated in the first 30 km of depth, being progressively more superficial, ~~as crust thickness diminishes oceanwards~~. The β -value is 0.96.

270 The focal mechanisms of the ~~events further away than far away events than~~ 14°W, present a significant component of strike slip. The depth ~~of the events, which relates to an unknown error~~ ~~solutions reveal earthquakes nucleating below, is more than~~ the Moho depth, which in this area decreases from 23 km in the coast to 12 km oceanwards, at the end of the lineation.

Again, in the depth distribution profiles (Fig. 3, profile 2) it seems that there ~~is~~ are more events up to the transition between hyperextended crust and exhumed mantle and ~~a lack of them~~ ~~less number of events beneath the~~ in the oceanic crust until the 275 western termination ~~of the profiles~~ where there is an increase in the amount of seismicity and at greater depths, ~~in the oceanic domain. This decrease cannot be considered such an evident gap as in the northern WIM at the same longitudes.~~

There is a puzzling vertical alignment (Fig. 3, profile 5b) below the Extremadura Spur, indicating some type of structure northwards and probably a second one ~~within to the north of~~ it. There may be a manifestation of a splay fault delimiting this topographic high. ~~Both vertical alignments occur within the exhumed mantle section.~~

280 A detail worthy ~~of to~~ note is that the highest magnitude earthquakes in this area occur in the subcrustal mantle ~~and below two seamounts~~ (Fig. 3, profile 4). ~~The northernmost event locates ~10 km westwards of a seamount~~ that may ~~be correspond related to volcanism, to a volcanic edifices as there is~~ are some volcanic edifices in the surface (Escada et al., 2022). ~~The referred vertical alignment would be consistent with a volcanic origin for those particular events.~~

285 Although the graphics show how the depth of the earthquakes is ~~accordingly in general~~, diminishing towards the ocean along the whole WIM, there is a considerable amount of earthquakes nucleating in the uppermost mantle, and especially in the transition areas to exhumed mantle domain and towards the oceanic crust at the end of the alignments. This suggest that seismicity occurs in the whole margin ~~crust~~, down to lithospheric levels. Within the ~~Lisbon (S) alignment~~ Central WIM, the hypocenters are slightly deeper than in the ~~Galicia (N) alignment~~ Northern WIM clusters, even when crustal depths are similar in both areas. The dispersion of events is greater at around longitude 14°W, maybe indicating the presence of some type of 290 seismic boundary or discontinuity in the N-S direction that traverses the whole WIM. ~~Profiles 3 and 4 in figure 3, show the~~

Con formato: Español (España)

Con formato: Español (España)

Con formato: Español (España)

distribution at depth within the oceanic domains of the margin revealing the separation of the both both bands. Also Also, they illustrate how hypocenters get shallower towards the west. In general, earthquakes along both alignments seem to cluster around marine reliefs, of either tectonic origin (Galicia Bank) or with volcanic additions (Extremadura Spur).

Earthquakes located below the Moho may indicate a strong, seismogenic upper mantle that can sustain large stresses, later released during brittle rupture. This type of “ductile earthquakes” have been proposed for Wyoming lithospheric mantle (Prieto et al., 2017), or in Newport-Inglewood fault attributed to a system of seismic asperities in ductile fault zones. Once a ductile mylonitic structure has developed a shear zone, subsequent cataclastic deformation is consistently localized in a narrow zone (Takahashi et al., 2017). The role of temperature in the rheology of the oceanic lower crust and lithospheric mantle is not well understood. Earthquakes occur throughout the oceanic crust and upper mantle, the later with olivine-dominated rheology. The brittle ductile transition is associated with a threshold temperature of about 600°C, which represents a transition from velocity weakening to velocity strengthening, consistent with the focal depth of earthquakes in the oceanic lithosphere elsewhere (Boettcher et al., 2007).

In the area of study, but in the most southern segment of the WIM, Geissler et al. (2012) provided evidence that most of the earthquakes within the Gulf of Cadiz have hypocentral depths between 40 and 55 km. In the SWIM, And Grevemeyer et al. (2016) also in the SWIM, provide evidence for earthquakes rupturing at 30-50 km depth, hypothesizing that they are caused by the proximity to the AGFZ plate boundary and by an elastic behaviour at the continent-ocean transition zone, which is further supported by low heat flow and the amount of regional stresses caused by the Africa and Eurasia² collision. Being an old passive margin, shortening and deformation of such a rigid lithosphere, may cause intense deformation at mantle depths. Mantle nucleation of events may additionally be encouraged by the presence of fluids or fluid migration in the mantle, either by reducing the effective normal stress or promoting strain localization along the shear zones. The velocity gradients observed on the edges of rift domains would make them preferred sites for reactivation due to inherited strength contrasts down to mantle depths.

Regarding the focal mechanisms in the study area, corresponding to events of magnitude 4.1 to 5.9 (Table 2), indicate mostly strike-slip movement.

315 3.2 WIM seismicity and geophysical data

The superposition of seismicity onto different types of geophysical data can be used to infer relationships that may give hints ~~of~~ to the seismicity sources. In the absence of confidently mapped submarine faults in the area, figure 4 depicts the event distribution obtained in this study overlaid to a different set of geophysical data for assessment. In the first case, the events are overlaid onto the Moho depth map. As expected, depth of events follows crustal thickness decrease seawards. However, as many events seem to nucleate in the uppermost mantle, we must infer that the crust and the upper mantle in the WIM must be strong, and that neither the Moho discontinuity nor the transition upper-lower crust are the rheological fundamental boundaries concerning seismicity.

When the seismicity is overlaid on the calculated gravity (Fig.–4b), the main relation ~~arise~~ arises from the fact that seismicity indeed does align where the basement is uplifted (i.e. Galicia Bank or Extremadura Spur), and therefore, follows the gravimetric expression of thickened crustal blocks, something that could be expected given that ~~gradients of velocity~~ velocity gradients have a notable influence on crustal strength. Regarding the magnetic map (Fig.–4c), it is difficult to establish a relationship, attending due to the main N-S disposition of the oceanic magnetic anomalies. The complexity in the magnetic imprint of the continental margin reveals some relation of several events to particular magnetic structures. Finally, figure 4d shows the seismicity overlaid onto the WIM rift domains map, which shows that seismicity does not nucleate preferentially in any domain. Although events are somewhat more abundant in the proximal domain, they do appear on exhumed mantle and hyper-thinned crust, indicating that the weakness of the domains is adequate for producing earthquakes anywhere. Seismicity

almost abruptly stops around the area of undisputed oceanic crust marked by anomaly M3 (figure 4c). Nonetheless, there is still a few events westward toward the MAR, and they follow the N80°W direction too.

Moreover, it also shows that the south alignment Central WIM seismicity happens south of the boundary between Northern and southern WIM, if we consider this boundary the location of the change in width of the exhumed domain, therefore ignoring rift domains transitions altogether.

4 Causes of the intraplate seismicity along the West Iberian margin

The concentration of seismicity at the northern and western coasts is explained in general terms as the internal deformation being the prime mode to accommodate strain induced by the convergence between Africa and Eurasia and the impossibility of propagating west of the Iberian peninsulaIberian Peninsula due to the Atlantic ridge push.

Pioreviously to discussing the possible sources of the WIM seismicity, a summary of the characteristics of the studied events that are reliable and therefore treated as strong constraints are:

- a) Seismic events are concentrated along two distinct WNW-ESE trends parallel to the active AGFZ Eurasia-Africa plate boundary. The interval distance between bands is similar, around 300 km (Fig.-21).
- 345 b) Magnitude is low to moderate, and hypocentral depths include the whole crust and upper mantle.
- c) The bands do not follow inherited tectonic trends. The events cut different rift domains and are prone to cluster along transitional areas and topographic highs. There is not an evident link to gravity or magnetic anomalies.
- d) They do show an association with topographical highs, especially with the western border of the Galicia Bank and the northern part of the Extremadura Spur.
- 350 e) The few focal mechanism available indicate overwhelmingly strike slip types.

A note on reliability of observations and data limitations

The text highlights the limitations of seismotectonic interpretations due to the inconclusive mapping of structures in the sea floor of the WIM. Despite being well probed, the margin contains many highs, roughness, and bathymetric heterogeneities that make it difficult to map tectonic features, specifically faults that could act as seismogenic sources. The structural variability observed along the margin added uncertainty in the data processing when introducing velocity models in the location programs of the events. Additionally, crustal depths in the margin vary, but a considerable number of events show hypocentres below the interpreted Moho depth.

The acquisition mode poses the second limitation as the land stations are located to the east, leading to a biased location error. This results in uncertainty in some locations and the lack of full azimuth to relocate with confidence. To minimize the impact of these shortcomings, only the events that comply with the requirements were considered reliable and retrieved:

- a) Magnitude should be over > 2.5.
- b) Events with location uncertainty of < 25 km in plan with average values < 10 km, and 75 km in depth. It is precisely this last parameter the one that poses the main problem, although the average error estimated has been of 4.9 km.
- 360 c) The average GAP < 183, for all the events used in this study being the ones with higher GAP discarded for this purpose.

Based on the aforementioned observations, we may have a walk throughreview the various hypotheses that have been proposed to explain seismic activity in the WIM plus a few new ideas. We can evaluate the validity of these hypotheses against the data collected over the past two decades.

4.1 Intraplate seismicity generated by unmapped faults. Strike-slip corridors, shear zones, transform faults.

370 Some causes for local clusters of events within the WIM have been put forward in the literature: for instance, Vazquez et al. (2008) indicated that the GB resulted from the reactivation of two major faults, forming an ample shear zone. Borges et al. (2001) related the central WIM seismicity to strike-slip movements in certain structures. Recently, Somoza et al. (2021)

Con formato: Fuente: Negrita

375 indicated speculate that to the north of Nazaré fault, seismicity responds to a reactivation and inversion of thrusts situated in the ocean seamount, that even connect to faults in the continent. The Nazaré fault would be prolonged-extended up to the MAR, linking with the Kurchatov transform fault (Fig. 1). However, there is not enough data to support this interpretation.

In any case, local clusters do not explain the overall distribution in the WIM. We have observed also local clusters such as the southward directed cluster in the GB related to the arrested subduction in the Bay of Biscay (figure Fig. 3, profile 5a) or the vertical alignment beneath some volcanic edifices along the Extremadura Spur that may relate to either some magmatic activity or to an splay fault quasy vertical (Figure 3, profile 5b).

380 For the bigger picture and an inclusive theory for the WIM seismicity, one of the recurring hypothesis indicated by some authors was the presence of shear corridors, (Custodio et al., 2015; Whitsmarsh et al., 1990) due to the particular stress state of the margin and the Iberian Peninsula.

Their oblivious indifference to structures of the crust or rheological boundaries of the rifted margin, all support the view that the events we are studying respond to the actual stress state, drawing the incipient silhouette of rectilinear lithospheric fractures or zones of strain parallel to the south and north of well-formed transform faults, one of them a plate boundary. These events may signal the formation of fracture zones that mimic oceanic transform faults. Contrastingly In contrast to the ones formed during the ocean opening, these ones are forming in the opposite direction, from land to sea. These faults may correspond to delayed accommodation structures of the extension and opening of the Atlantic in these latitudes, where oblique extension and the jump of plate boundaries (Srivastava et al., 1990), might have impeded the formation of classic oceanic transform faults at the time.

390 This process would be framed within the peripatetic behaviours of the Iberian microplate. The topic of oblique spreading along the WIM has not been deeply studied approached in depth. However, there is some evidence that the opening of the Atlantic, and especially during the Bay of Biscay rift, Iberia, may have been “bouncing” from its perpendicular to the extension direction. Favouring this option is the fact that oblique spreading is very common in slow spreading ridges, as lithosphere is 395 relatively cold and crustal growth is decelerated (Peyve, 2009). Another argument in agreement with this hypothesis is that mid-ocean ridges with oblique spreading are not dissected by transform faults. Shallow tectonics dictates their segmentation rather than mantle convection.

In the other hand, the presence of these heterogeneities stand out from the rift domains map, where the width of domains changes at the Lisbon lineation and north of the Galicia lineation (Fig. 3).

400 Published interpretation relying on the study of seismic profiles that cross the WIM seismicity bands, do not show evident crustal strike-slip structures. However, it is important to note that sub-vertical strike-slip faults that promote horizontal displacements are difficult to image in near vertical reflection data, and have a limited expression on the floor surface, even more if they are relatively recent.

405 The claim for these NW-SE oriented transform corridors to facilitate extension along the MAR, working as “pseudo transform faults” (Fig. 5a), reinforces the need to accommodate differential stresses from South to North along the WIM, which have resulted from the rotations and subtle drift of the Iberian plate during the past 30 Myr, together with the current NW-SE oriented compression. In this case, the seismicity alignments may correspond to those shallow accommodation zones, similar to shear zones that act as nascent oceanic transform faults, without dissecting the ridge (yet).

410 This compartmentalization of a passive margin into strike slip narrow belts between the MAR and a deformable plate interior, is more consistent with soft plate idealizations-concepts than with perfectly rigid plates and narrow boundaries (Storti et al., 2007), which takes us to the next hypothesis to try to fit this seismicity into the plate tectonics theory.

4.2 A diffuse and ~1000 km wide plate boundary

Plate rigidity and narrow plate boundaries are central assumptions of the original plate tectonics theory. But in the last decades 415 evidences have has been piling up to suggest that in many cases, plate boundaries can be very wide and plates may not be as

Con formato: Fuente: Negrita

rigid as assumed (Ribeiro, 2002). The diffuse plate boundary hypothesis tries to reconcile observations related to intraplate activity by proposing that large plates may be composed of a few smaller subplates and crustal blocks, so more rigid portions are in relative motion with each other separated by diffuse boundaries with clustered deformation (Gordon, 1998). The diffuse plate boundaries are much broader than traditional boundaries such as mid-ocean ridges, trenches and/or oceanic transform

420 faults (Zatman et al., 2000).

In this way, the passive interior of tectonic plates can be dissected by strike slip deformation belts propagating towards MAR segments that separate adjacent, partially coupled lithospheric slivers with differential translations, and deformation rates. These internal differential displacements are then accommodated by distributed deformation within the plate interior, so that localized deformation is ultimately dissipated into a non-rigid plate.

425 Applying this model to WIM is challenging because

published studies proposing soft plate deformation and the formation of weakened zones involve the spatial extension of transform faults through oceanic fracture zones, possibly weakened by serpentinization, that are used to transfer deformation from mid ocean ridges towards neighbouring plate interiors. In the WIM case, as pointed out earlier, the Galicia (N) and Lisbon (S) lineationseismicity bands are discontinuous, do not intersect the MAR and specifically stop in the vicinity of the 430 oceanic crust before reaching the MAR, so a full lithospheric detachment is not occurring. Therefore, it is more likely that these lineations may be inherited from the rifting period, or from post rift accommodation and thermal subsidence rather than to the sea floor spreading stage of the Atlantic.

Along the southern Biscay margin, the Ventaniella seismicity line (Fernandez-Viejo et al., 2014; López-Fernández et al., 2018) shows a similar NW-SE trend and includes earthquakes of similar magnitude, distribution and focal mechanism as the Galicia 435 (N) and Lisbon (S) lineation identified in the WIM. Contrastingly, it appears in a continental margin whose direction is perpendicular to the orientation of the WIM direction (the Bay of Biscay margin). If the three seismicity bands are expressions of shear corridors inside an exceptionally wide plate boundary, this hypothesis would deserve its credit, and the fact that they appear in two completely perpendicular trending passive margins, even crossing oceanic and continental parts of the Iberian plate, would agree with support it.

440 Asti et al. (2022) proposed a 400 km wide, diffuse Iberia-Eurasia plate boundary transecting Iberia that would have been active during the Mesozoic, and extended from the Iberian Rift system to the Armorican shelf, which was associated with the contemporaneous opening of the North Atlantic and Bay of Biscay. The deforming domain between these two boundaries was a fuzzy region, where the exact locations of different tectonic structures and rift basins is still debated. If a similar diffuse plate boundary may exists today, it would be almost double in width and would encompass the entire Peninsula except for its

445 northeastern third (Pyrenees and related foreland basin).

One advantage of the diffuse plate boundary theory (Fig. 5b) is that some of the inconsistencies between the deformations inferred from plate kinematic models and geological observations (which are extraordinarily abundant in Iberian kinematics-see; Barnett-Moore et al., (2016), and references therein) may be then reconciled. The same amount of total displacement is partitioned between a series of subparallel strike slip corridors, each of them accommodating a portion of the total displacement 450 between the stable parts of the African and European plates. This would significantly broaden the range of geodynamical models in a region with such as slow lithospheric deformation. If deformation between Africa and-Eurasia is distributed inside Iberia, these internal corridors of deformation can accommodate the convergence of both macroplates, while assisting synchronically the E-W Atlantic extension.

Nevertheless, if a great part of Iberia belongs to a “diffuse plate boundary”, we still need to explain why the corridors are 455 produced at those particular separation of about 300 km.—Does that separation depends on the width of the subplates related to the “rigid microplate”? Does it depend on the velocity rate of the ridge? This needs further investigations out of the scope of this study, but it points to some kind of relationship to be able to form measurable strike-slip corridors between areas with contrasting crustal lithospheric thickness, plate width and undergoing different stress fields. In any case, a complete radical

hypothesis has been advanced in the literature, which approaches the dynamics of the WIM ~~in~~~~from~~ a very different perspective:
460 the closing of the Atlantic Basin.

4.3 Subduction initiation and oceanic dynamics that may be, or have been, at play in the WIM

Several authors have suggested the possibility of a developing subduction zone in the SWIM, which could explain the moderate seismicity and the large magnitude of the 1755 Lisbon earthquake in an intraplate setting (Vilanova et al., 2003; Martinez-

465 Loriente et al., 2021). Ribeiro et al. (1996) proposed that ~~the WIM is the site of~~ an incipient northward propagating subduction zone ~~may be~~ nucleating at the Gorringe Bank (figure 5c). The ongoing terminal stage of collision between Africa and Eurasia produces compressive stresses along the SWIM, which could trigger passive margin reactivation. [Gutscher et al., \(2012\) show evidences of subduction towards the East in the Gibraltar arc and in the West within the southern Iberian margin](#) NE-SW thrust systems extending 300 km along the SWIM accommodate the ~~are orthogonal~~-convergence ([Gutscher et al., 2012](#)) ~~and while~~ 470 younger thrust faults are nucleating along the ~~west Portuguese passive margin~~ SWIM ([Zitellini et al., 2009; Terrinha et al., 2019; Martínez-Loriente et al., 2013; 2021](#)) or in the Tagus Abyssal plain ([Neves et al., 2009](#)). Duarte et al. (2013) ~~suggest speculate~~ that these structures may be indicative of the onset of margin tectonic inversion and the nucleation of a new subduction, ~~which develops due to the westward propagation of the Gibraltar Arc and related subduction~~ ([Gutscher et al., 2012](#)).

Comentado [p1]: Gutscher et al., 2002 (Evidence for active subduction beneath Gibraltar), y Mediadea et al., 2004 ([Structure and evolution of the "Olistostrome" complex of the Gibraltar Arc in the Gulf of Cádiz \(eastern Central Atlantic\): evidence from two long seismic cross ...](#)), hay que revisar que muestran NE-SW thrust systems in the SWIM. Revisar también Terrinha et al. 2019

475 However, [focusing on the central and northern WIM, away from the complexity of the plate boundary](#), the buoyancy of old and cold oceanic lithosphere makes the spontaneous initiation of subduction in ~~a~~ passive margins difficult, and significant weakening mechanisms are required to fail in subduction (e.g., Stern and Gerya, 2018). The Azores, Madeira plume, or the peridotite ridge in the Galicia margin may serve as potential weakening mechanisms. A 100 my old oceanic crust ~~that~~ would need a significant compressive stress to fail (Zhong et al., 2019), which is not observed at the WIM.

480 Thermo-mechanical models suggest that magma-poor margins with hyperextended and exhumed mantle domains are favourable sites for subduction initiation, as the serpentized mantle facilitates strain localization and the progressive development of a major shear zone that ultimately links up with a zone of high strain the continental lower crust along the necking zone (Auzemery et al., 2021). Shearing then propagates into the mantle or initiate a subduction plate boundary. The exhumation and hydration of the mantle in the footwall of major detachments induces weak decollement layers (Brun and

485 Beslier, 1996) prone to tectonic reactivation, which would diminish the required—compressive stress to initiate subduction (Hirt et al., 2013). [However, the seismicity observed as we have seen it is not restricted to the exhumed mantle domains— and the focal mechanisms either suggest an inverse fault](#). Subduction could be easier in the southern WIM as a relevant transform fault, the Gloria fault, puts in contact two different lithospheric age crusts making for additional stresses and weakened lithosphere. [Although the initiation of subduction by passive margin collapse is extremely unrealistic, i.e. \(Stern and Gerya., 490 2018\), the westward migration and propagation of the Gibraltar Arc along the SWIM may be responsible for induced subduction initiation \(Duarte et al., 2013\).— The majority of focal mechanisms along the WIM events outside of the plate boundary are strike-slip, which does not support the lithospheric collapse in a subduction interface propagating northwards along the WIM.— Although some type of underthrusting ~~cannot be ruled out~~ and convergence is evidenced in the south \(Gutscher et al., 2012, Martínez-Loriente et al., 2021\).— seismicity distribution and focal mechanisms, which are predominantly strike-](#)

Con formato: Fuente: Times New Roman

495 [slip, does not support unequivocally the presence of a subduction zone the northward propagation of a subduction interface.](#)

Con formato: Fuente: Times New Roman

Con formato: Fuente: Times New Roman

Con formato: Fuente: Times New Roman

4.4 Oceanic complexity, inheritance, formation of strained corridors in the Atlantic ocean

Within the other big oceanic basin of the Earth, in the Pacific, crustal studies evidence complex tectonic settings. In triple junctions, triangular microplates without any bounding continental margins grow, as exemplified by the Galapagos microplate.

500 Researchers have found strong evidence that indeed, one ~~of~~ such microplates grew to become the Pacific plate, ~~for instance~~ (Boschman and Hinsberger, 2016). Schouten et al. (2008), studying distributed deformation at oceanic triple junction showed that lithospheric plates undergo significant internal deformation as their boundaries rapidly evolve, as exemplified by the ~~triple~~

junction (RRR) Cocos-Nazca. The Cocos Nazca rift tip does not meet the East ~~P~~acific ~~R~~ise. Instead, two secondary rifts form the link. The active incipient rift is just the latest of a sequence of southeast trending fractures that progressively stepped ~~sideways~~ during the last 5 Myr, successively accommodating minor N-S extension of the generated oceanic crust (Smith, 505 2011).

The roughness of the abyssal plain around the King's ~~T~~rough may represent the incipient and ~~left-alone~~~~abandoned~~ rotation of a potential mini Galapagos-like microplate. The current scars ~~related~~~~associated~~ to the triple junction on the King's ~~T~~rough area with seafloor expression relate to inherited lithospheric fractures, which developed during the evolution of the triple point and the plate boundary migration. However, even when considering the inheritance of such lithospheric discontinuities, the 510 seismicity in the WIM does not appear connected to it in particular, more than following the general NNW-SSE trend. And as figure 2 shows, seismicity is mainly within the continental rifted crust and not so abundant within the oceanic domains.

Palano et al. (2015) propose a current large-scale clockwise rotation that makes Iberia ~~to~~ act as a microplate with a southern limit in the AGFZ Nubia-Eurasia convergent zone. Somoza et al. (2021) indicate that the main weakened zones along ~~the~~-N and NW Iberia are the old inherited limits between the oceanic domains of Eurasia and Iberia (King's Trough, Azores-Biscay 515 rise, Jean-Charcot Seamount). They question the clockwise rotation in the oceanic domain, which would be dominated by the stress derived from the ridge propagation in the segment between Azores and King's Trough, with a \sim N120° propagation direction. They also suggest a left-lateral shear zone resulting from the northward movement of the continental Iberia and the SE movement of oceanic Iberia along the OCT.

According to the possible hypothesis above mentioned, the one that presents less problems in this study is that the pre-existing 520 Mesozoic rift structure, together with the oblique oceanic spreading, when the plate boundary was shifting southwards towards the AGFZ, may have left weakened areas along the WIM. These inherited zones may be prone to reactivation and release seismicity under the ~~actual~~~~prevailing~~ stress regime. Inherited features revealed by the seismicity are for instance, the old proto-subduction interface in the north of Galicia Bank (figure 3, profile 5) or the associated volcanic edifices around the Extremadura Spur. On the contrary, ~~the~~ the remarkable reduction of exhumed domains towards the central parts seem not to 525 have any expression in the current seismicity distribution.

5 Conclusions

Analysis of two decades of seismicity data recorded by the Spanish and Portuguese seismic networks along the West Iberian 530 passive margin (WIM), covering an area between 38° and 45°N latitude, shows that two NNW-SSE bands concentrate 98% of the intraplate activity. The northern band is located offshore Galicia, while the southern band is in offshore Lisbon. The observed seismicity is higher than expected for a passive margin, and coincides with the area where the mid-Atlantic ridge lacks conspicuous transform faults. Focal mechanisms indicate strike-slip deformation, and hypocentral depths suggest crustal and upper mantle sources.

Various hypotheses have been proposed to explain this seismicity, including the formation of shear corridors/transform faults, 535 a diffuse plate boundary, initiation of subduction, and other less understood mechanisms associated with weakened zones inherited from the Mesozoic rifting, complex ocean opening, and/or triple point migration in the Iberia microplate. Based on the parameters obtained in this study, a conservative approach to explain the WIM seismicity would be the presence of strained shear corridors that nucleate preferentially on topographical heterogeneities of the seafloor. These corridors form in response to the accommodation of two perpendicular sources of stress: the E-W extension associated with the MAR, and the N-S compression from the Azores-Gibraltar Fracture Zone. The drastic crustal thickness variations from east to west and the 540 nucleation of small events along multiple faults that make up the block morphology of the Mesozoic rifted margin conform the occurrence of areas prone to strain release. The presence of sub-Moho earthquakes also indicates that the upper mantle in the WIM acts as a high-strength layer, particularly beneath the hyperextended and exhumed mantle transition areas.

Several plate-scale processes may be acting simultaneously to produce the observed seismicity, such as the possible initiation of convergence-related deformation in the SWIM or the inherited compartmentalization of the lithosphere in the northern part 545 of the margin due to triple junction migration. Further study of seismicity in the WIM and neighbouring oceanic areas, together with the arrival of seafloor stations, may provide further insights to better understand this marine intraplate clustered activity.

Data and resources

Seismicity data used in this study were collected from the IGN Spanish seismic network at www.ign.es (last accessed December 2022) and from IPMA Portuguese seismic network at www.ipma.pt/en/ (last accessed December 2022). Gravity, 550 magnetic and crustal data were obtained from GeoMapApp at <https://www.geomapapp.org/> (last accessed January 2023).

Acknowledgements

The research leading to these results received funding from the Project MCIU-22-PID2021-123116NB-I00 by the Spanish Science and Innovation Ministry. P. Cadenas hold a Marie Skłodowska-Curie Actions, SUBIMAP, grant agreement ID: 895895. Chat GPT was used to improve the English writing and clarity in some paragraphs.

555

References

Álvarez-Marrón, J., Rubio, E., and Torne, M.: Subduction-related structures in the North Iberian Margin, *J. Geophys. Res.* 102, 22497-22511, doi:10.1029/97JB01425, 1997.

Ayarza, P., Mrtinez-Catalán, J.R., [Álvarez-Marrón](#), J., Zeyen, H., and Juhlin, C.: Geophysical constraints on 560 the deep structure of a limited ocean-continent subduction zone at the North iberian margin, *Tectonics*, 23, TC101, doi:1.1029/2002TC001487, 2004.

Afilhado, A., Matias, L., Shiobara, H., Hirn, A., Mendes-Victor, L., and Shimamura, H.: From unthinned continent to ocean: The deep structure of the West Iberia passive continental margin at 38N, *Tectonophysics*, 458, 9-50, doi:10.1016/j.tecto.2008.03.002, 2008

565 Alves, T., Moita, C., Cunha, T., Ullnaess, M., Myklebust, R., Monteiro, J.H., and Manuppella, G.: Diachronous evolution of Late Jurassic-Cretaceous continental rifting in the northwest Atlantic (west Iberian margin), *Tectonics*, 28, TC4003, doi:10.1029/2008TC002337, 2009

Amigo Marx, B., Fernandez, O., Olaiz, A., Poblet, J., and Zamora, G.: On the influence of Variscan inheritance on rifting of the Western Iberian margin, *Terra Nova*, 34, 424-432, doi:10.1111/ter.12595, 2022.

570 Andeweg, B., De Vicente, G., Cloetingh, S.A.P.L., Giner, J., and Muñoz-Martin, A.: Local stress fields and intraplate deformation of Iberia: variations in spatial and temporal interplay of regional stress sources, *Tectonophysics*, 305, 153-164, doi:10.1016/S0040-1951(99)00004-9, 1999.

Asti, R., Sasipiturry, N., and Angrand, P.: The Mesozoic Iberia-Eurasia diffuse plate boundary: A wide domain of distributed transtensional deformation progressively focusing along the North Pyrenean Zone, *Earth. Sci. Rev.*, 230, 104040, 575 doi:10.1016/j.earscirev.2022.104040, 2022.

Auzemery, A., Willingshofer, E., Sokoutis, D., Brun, J.P., and Cloetingh, S.A.: Passive margin inversion controlled by stability of the mantle lithosphere, *Tectonophysics*, 817, 229042, doi:10.1016/j.tecto.2021.229042, 2021.

Barnett-Moore, N., M. Hosseinpour, and Maus, S.: Assessing discrepancies between previous plate kinematic models of 580 Mesozoic Iberia and their constraints, *Tectonics*, 35, 1843-1862, doi:10.1002/2015TC004019, 2016.

580 Boettcher, M. S., Hirth, G., and Evans, B.: Olivine friction at the base of oceanic seismogenic zones. *J. Geophys. Res.*, 112, B01205, [doi:10.1029/2006JB004301](https://doi.org/10.1029/2006JB004301), 2007.

Boillot, G., Girardeau, J., and Kornprobst, J.: Rifting of the Galicia margin: crustal thinning and emplacement of mantle rocks on the seafloor. In Boillot, G., Winterer, E.L., et al., *Proc. ODP, Sci. Results*, 103: College Station, TX (Ocean Drilling Program), 741-756, [doi:10.2973/odp.proc.sr.103.179.1988](https://doi.org/10.2973/odp.proc.sr.103.179.1988), 1988a

585 Boillot, G., Keith W. Dunham, Philip A. Meyers, and Eileen, S.H.: *Proceedings of the Ocean Drilling Program: Scientific Results*, Ocean Drill. Program, College Station, Texas, 1-858, 1988b

Borges, J.F., Fitas, A.J.S., Bezzeghoud, M., and Teves-Costa , P.: Seismotectonics of Portugal and its adjacent Atlantic Area, *Tectonophysics*, 337, 373-387, [doi:10.1016/S0040-1951\(00\)00291-2](https://doi.org/10.1016/S0040-1951(00)00291-2), 2001.

Boschman, L.M., and van Hinsbergen , D.J.J.: On the enigmatic birth of the Pacific Plate within the Panthalassa Ocean, *Sci. 590 Adv.* 2(7) e1600022, [doi:10.1126/sciadv.1600022](https://doi.org/10.1126/sciadv.1600022), 2016.

Bronner, A., Sauter, D., Manatschal, G., Péron-Pinvidic, G. and Munschy, M.: Magmatic breakup as an explanation for magnetic anomalies at magma-poor rifted margins, *Nature Geosci.* 4, 549-553, [doi:10.1038/ngeo1201](https://doi.org/10.1038/ngeo1201), 2011.

Brun, J.P., and Beslier, M.O.: Mantle exhumation at passive margins, *Earth. Planet. Sci. Lett.*, 142, 161-173, [doi:10.1016/0012-821X\(96\)00080-5](https://doi.org/10.1016/0012-821X(96)00080-5), 1996.

595 Buñor E, Udías A, Colombás MA: Seismicity, source mechanisms and seismotectonics of the Azores-Gibraltar plate boundary, *Tectonophysics*, 152, 89-118, 1988.

Custódio, S., Dias, N. A., Carrilho, F., Góngora, E., Rio, I., Marreiros, C., Morais, I., Alves, P. and Matias, L.: Earthquakes in western Iberia: Improving the understanding of lithospheric deformation in a slowly deforming region, *Geophys. J. Int.* 203(1), 127-145, [doi:10.1093/gji/ggv285](https://doi.org/10.1093/gji/ggv285), 2015.

600 Dean, S.M., Minshull, T.A., Whitmarsh, R.B., Louden, K.E.: Deep structure of the ocean-continent transition in the southern Iberia abyssal Plain from seismic refraction profiles: The IAM-9 transect at 40°20'N, *Journal of Geophysical Res.*, 105: 5859-5885, 2000.

De Vicente, G., Cloetingh, S., Muñoz-Martín, A., Olaiz, A., Stich, D., Vegas, R., Galindo-Zaldívar, J., and Fernández-Lozano, J.: Inversion of moment tensor focal mechanisms for active stresses around the microcontinent Iberia: Tectonic implications, *605 Tectonics*, 27. TC1009, [doi:org/10.1029/2006TC002093](https://doi.org/10.1029/2006TC002093), 2008.

Druet, M., Muñoz-Martín, A., Granja-Bruña, J.L., Carbó-Gorosabel, A., Llanes, P., Catalán, M., Maestro, A., Bohoyo, F., and Martín-Dávila , J.: Bouguer anomalies of the NW Iberian continental margin and the adjacent abyssal plains, *J. Map. Geogr. Libr.*, 15, 635-641, [doi:10.1080/17445647.2019.1644214](https://doi.org/10.1080/17445647.2019.1644214), 2019.

Duarte, J.C., Rosas, F.M., Terrinha, P., Schellart, W., Boutelier, D., Gutscher, M.A. and Ribeiro , A.: Are subduction zones 610 invading the Atlantic? Evidence from the southwest Iberia Margin, *Geology*, 41, 839-842, [doi:10.1130\(G34100.1](https://doi.org/10.1130(G34100.1), 2013

Dziewonski, A.M., T.A. Chou, and Woodhouse, J.H.: Determination of earthquake source parameters from waveform data for studies of global and regional seismicity, *J. Geophys. Res.*, 86, 2825-2852, doi:10.1029/JB086iB04p02825, 1981.

EMODnet Bathymetry Consortium (2020). EMODnet Digital Bathymetry (DTM 2020), [doi:10.12770/bb6a87dd-e579-4036-abe1-e649cea9881a](https://doi.org/10.12770/bb6a87dd-e579-4036-abe1-e649cea9881a), 2020.

615 Ekström, G., M. Nettles, and Dziewonski, A. M.: The global CMT project 2004-2010: Centroid-moment tensors for 13,017 earthquakes, *Phys. Earth Planet. Inter.*, 200-201, 1-9, [doi:10.1016/j.pepi.2012.04.002](https://doi.org/10.1016/j.pepi.2012.04.002), 2012.

Escada, C., Represas, P., Santos, F., Pereira, R., Mata, J., and Rosas, F.M.: New evidence of Late Cretaceous magmatism on the offshore central West Iberian margin (Extremadura Spur) from potential field data, *Tectonophysics*, 832, 229354, [doi:10.1016/j.tecto.2022.229354](https://doi.org/10.1016/j.tecto.2022.229354), 2022.

620 Fernandes, R. M. S., Ambrosius, B. A. C., Noomen, R., Bastos, L., Wortel, M. J. R., Spakman, W., and Govers, R.: The relative motion between Africa and Eurasia as derived from ITRF2000 and GPS data, *Geophys. Res. Lett.*, 30, 1828, [doi:10.1029/2003GL017089](https://doi.org/10.1029/2003GL017089), 2003.

Fernández-Viejo, G., López-Fernández, C., Domínguez-Cuesta, M.J., and Cadenas, P.: How much confidence can be conferred on tectonic maps of continental shelves? The Cantabrian-Fault case, *Sci. Rep.*, 4, 3661, [doi:10.1038/srep03661](https://doi.org/10.1038/srep03661), 2014.

Con formato: Inglés (Estados Unidos)

625 Fernandez-Viejo, G., Pulgar, J.A., Gallastegui, J. and Quintana, L.: The fossil accretionary wedge of the Bay of Biscay: critical wedge analysis on depth-migrated seismic sections and geodynamical implications, *J Geol.*, 120, 1-18. [doi:10.1086/664789](https://doi.org/10.1086/664789), 2012.

Con formato: Inglés (Estados Unidos)

Con formato: Inglés (Estados Unidos)

Ferrão, C., Bezzeghoud, M., Caldeira, B., and Borges, J.F.: The Seismicity of Portugal and Its Adjacent Atlantic Region from 1300 to 2014: Maximum Observed Intensity (MOI) Map, *Seismol. Res. Lett.*, 87, 743-750, [doi:10.1785/0220150217](https://doi.org/10.1785/0220150217), 2016.

630 Garate, J., Martin-Davila, J., Khazaradze, G., Echeverria, A., Asensio, E., Gil, A.J., de Lacy, M.C., Armenteros, J.A., Ruiz, A.M., Gallastegui, J., [AlvarezÁlvarez](#)-Lobato, F., Ayala, C., Rodríguez-Caderot, G., Galindo-Zaldívar, J., Rimi, A. and Harnafi, M.: Topo-Iberia project: CGPS crustal velocity field in the Iberian Peninsula and Morocco, *GPS Solut.*, 19, 287-295, [doi:10.1007/s10291-014-0387-3](https://doi.org/10.1007/s10291-014-0387-3), 2015.

635 Geissler, W.H., Matias, L., Stich, D., Carrilho, F., Jokat, W., Monna, S., Ibenbrahim, A., Mancilla, F., Gutscher, M.A., Sallares, V., and Zitellini, N.: Focal mechanisms for sub-crustal earthquakes in the Gulf of Cadiz from a dense OBS deployment, *Geophys. Res. Lett.*, 37, L18309, [doi: 10.1029/2010GL044289](https://doi.org/10.1029/2010GL044289), 2010.

Gomez de la Peña, L., Ranero, C.R., Gràcia, E., Booth-Rea, G., Azañón, J.M., Tinivella, U., and Yelles-Chaouche, A.: Evidence for a developing plate boundary in the western Mediterranean, *Nat. Commun.*, 13, 4786. [doi:10.1038/s41467-022-31895-z](https://doi.org/10.1038/s41467-022-31895-z), 2022.

640 Gonzalez, A., Cordoba, D. and Vales, D.: Seismic crustal structure of Galicia continental margin, NW Iberian Peninsula, *Geophys. Res. Lett.*, 26, 1061-1064, 1999.

Gordon R.G.: The plate tectonic approximation: Plate nonrigidity, diffuse plate boundaries, and global plate reconstructions, *Ann. Rev. Earth Planet. Sci.*, 26, 615-642, 1998

645 Granado, C., Muñoz-Martín, A., Olaiz, A.J., Fernández, O., and Druet, M.: 3D crustal-scale structure of the West Iberia margin: a novel approach to integrated structural characterization of passive margins, *Mar. Geophys. Res.*, 42, 10, [doi:10.1007/s11001-021-09432-2](https://doi.org/10.1007/s11001-021-09432-2), 2021.

Grevemeyer, I., Matias, L., and Silva, S.: Mantle earthquakes beneath the South Iberia continental margin and Gulf of Cádiz - constraints from an onshore-offshore seismological network, *J. Geodyn.*, 99, 39-50, [doi:10.1016/j.jog.2016.06.001](https://doi.org/10.1016/j.jog.2016.06.001), 2016.

650 Grevemeyer, I., Ranero, C.R., Papenberg, C., Sallares, V., Bartolomé, R., Prada, M., Batista, L., Neres, M.: The continent-to-ocean transition in the Iberia Abyssal Plain, *Geology*, 50, 615-619, 2022.

Gutscher, M.A., Dominguez, S., Westbrook, G.K., Le Roy, P., Rosas, F., Duarte, J.C., Terrinha, P., Miranda, J.M., Graindorge, D., Gailler, A., Sallares, V., and Bartolome R.: The Gibraltar subduction: a decade of new geophysical data. *Tectonophysics*, 574, 72-91, [doi:10.1016/j.tecto.2012.08.038](https://doi.org/10.1016/j.tecto.2012.08.038), 2012.

655 Hirt, G., and Guillot, S.: Rheology and tectonic significance of serpentine, *Elements*, 9, 107-113, [doi:10.2113/gselements.9.2.107](https://doi.org/10.2113/gselements.9.2.107), 2013.

Jimenez-Munt, I., and Negredo, A.: Neotectonic modeling of the western part of the Africa-Eurasia plate boundary: from the Mid-atlantic ridge to Algeria, *Earth Planet. Sci. Lett.*, 205, 257-271, [doi:10.1016/S0012-821X\(02\)01045-2](https://doi.org/10.1016/S0012-821X(02)01045-2), 2003.

King, M.T., Welford, J.K., and Peace, A.L.: Investigating the role of the Galicia Bank on the formation of the North West Iberian margin using deformable plate tectonic models, *Tectonophysics*, 789, 228537, [doi:10.1016/j.tecto.2020.228537](https://doi.org/10.1016/j.tecto.2020.228537), 2020.

660 King, M.T., and Welford, J.K.: Advances in deformable plate tectonic models: 2. Reconstructing the southern North Atlantic back through time, *G-cubed*, 23, e2022GC010373, [doi:10.1029/2022GC010373](https://doi.org/10.1029/2022GC010373), 2022.

Krüger, F., Dahm, T., and Hannemann, K.: Mapping of Eastern North Atlantic Ocean seismicity from Po/So observations at a mid-aperture seismological broad-band deep sea array, *Geophys. J. Int.*, 221, 1055-1080. [doi:10.1093/gji/gja054](https://doi.org/10.1093/gji/gja054), 2020.

665 Laske, G., Masters, G., Ma, Z. and Pasyanos, M.: Update on CRUST1.0 - A 1-degree Global Model of Earth's Crust, *Geophys. Res. Abstr.* 15 EGU2013-2658, <https://ui.adsabs.harvard.edu/abs/2013EGUGA..15.2658L>, 2013.

Código de campo cambiado

Con formato: Inglés (Reino Unido)

Con formato: Inglés (Reino Unido)

López-Fernández, C., Fernández-Viejo, G., Olona, J., and Llana-Fúnez, S.: Intraplate seismicity in Northwest Iberia along the trace of the Ventaniella fault: a case for fault intersection at depth, *Bull. Seism. Soc. Am.*, 108, 604-618, [doi:10.1785/0120170215](https://doi.org/10.1785/0120170215), 2018.

670 Lymer, G., and Reston, T.J.: The West Iberian margin: past and current research concepts and future challenges. *Continental rifted margins 2. Case examples*. Eds: Wiley and Sons, 256 p., 2022

Macchiavelli, C., Vergés, J., Schettino, A., Fernández, M., Turco, E., Casciello, E., Torne, M., Pierantoni, P.P. and Tunini, L.: A new southern North Atlantic isochron map: Insights into the drift of the Iberian Peninsula since the Late Cretaceous, *J. Geophys. Res. B*, 122, 9603-9626, [doi:10.1002/2017JB014769](https://doi.org/10.1002/2017JB014769), 2017.

675 Malod, J.A. and Mauffret, M.: Iberian plate motions during the Mesozoic, *Tectonophysics*, 184, 261-278, The Geological Society of London, London, 1990.

Martínez-Loriente, S., Gracia, E., Bartolome, R., Sallarès, V., Connors, C., Perea, H., Lo Iacono, C., Klaeschen, D., Terrinha, P., Danobeitia, J.J. and Zitellini, N.: Active deformation in old oceanic lithosphere and significance for earthquake hazard: Seismic imaging of the Coral Patch Ridge area and neighboring abyssal plains (SW Iberian Margin), *G-cubed*, 14, 2206-2231, 2013.

680 Martinez-Loriente, S., Sallares, V., and Gracia, E.: The Horseshoe abyssal plain thrust could be the source of the 1755 Lisbon earthquake and tsunami. *Commun. Earth. Environ.*, 2, 145, [doi:10.1038/s43247-021-00216-5](https://doi.org/10.1038/s43247-021-00216-5), 2021.

Merino, I., Ranero, C.R., Prada, M., Sallares, V., and Grevemeyer, I.: The Rift and Continent-Ocean Transition Structure under the Tagus Abyssal Plain West of the Iberia, *J. Geophys. Res. B. Solid Earth Planets*, 126, e2021JB022629, [doi:10.1029/2021JB022629](https://doi.org/10.1029/2021JB022629), 2021.

685 Meyer, B., Saltus, R. and Chulliat, A.: EMAG2v3: Earth Magnetic Anomaly Grid (2-arc-minute resolution). Version 3. NOAA National Centers for Environmental Information, doi:10.7289/V5H70CVX, Accessed [December 2023], 2017

Murillas, J., Mougenot, D., Boulot, G., Comas, M.C., Banda, E. and Mauffret, A.: Structure and evolution of the Galicia Interior Basin (Atlantic western Iberian continental margin), *Tectonophysics* 184 297-319, [doi:10.1016/0040-1951\(90\)90445-E](https://doi.org/10.1016/0040-1951(90)90445-E), 1990.

690 Neves, M.C., Terrinha, P., Alfilhado, A., Moulin, M., Matias, L., Rosas, F. (2009). [Response of a multi-domain continental margin to compression: Study from seismic reflection-refraction and numerical modelling in the Tagus Abyssal plain. Tectonophysics](https://doi.org/10.1016/j.tecto.2008.05.008), 468 (1-4), 113-130, <https://doi.org/10.1016/j.tecto.2008.05.008>.

Con formato: Portugués (Portugal)

Con formato: Inglés (Canadá)

Con formato: Inglés (Canadá)

Con formato: Inglés (Canadá)

Nirrengarten, M., Manatschal, G., Tugend, J., Kusznir, N.J. and Sauter, D.: Nature and origin of the J-magnetic anomaly offshore Iberia-Newfoundland: implications for plate reconstructions, *Terra Nova*, 29, 20-28, [doi:10.1111/ter.12240](https://doi.org/10.1111/ter.12240), 2017.

695 Nirrengarten, M., Manatschal, G., Tugend, J., Kusznir, N. J. and Sauter, D.: Kinematic Evolution of the Southern North Atlantic: Implications for the Formation of Hyperextended Rift Systems, *Tectonics*, 37, 89-118, [doi:10.1002/2017TC004495](https://doi.org/10.1002/2017TC004495), 2018.

Nocquet, J.M.: Present-day kinematics of the Mediterranean: A comprehensive overview of GPS results, *Tectonophysics*, 579, 220-242, doi:10.1016/j.tecto.2012.03.037, 2012.

700 Nocquet, J.M., and Calais, E.: Geodetic measurements of crustal deformation in the western Mediterranean and Europe, *Pure Appl. Geophys.*, 161, 661-681. [doi:10.1007/s00224-003-2468-z](https://doi.org/10.1007/s00224-003-2468-z), 2004.

Palano, M., Gonzalez, P.J., and Fernandez, J.: The diffuse plate boundary of Nubia and Iberia in the western Mediterranean: crustal deformation evidence for viscous coupling and fragmented lithosphere, *Earth Planet. Sci. Lett.*, 430, 439-447, [doi:10.1016/j.epsl.2015.08.040](https://doi.org/10.1016/j.epsl.2015.08.040), 2015.

705 Peace, A.L., Welford, J.K., Ball, P.J., and Nirrengarten, M.: Deformable plate tectonic models of the southern North Atlantic, *J. Geodyn.*, 128, 11-37, [doi:10.1016/j.jog.2019.05.005](https://doi.org/10.1016/j.jog.2019.05.005), 2019.

Pedrera, A., Ruiz-Constán, A., Galindo-Zaldívar, J., Chalouan, A., Sanz de Galdeano, C., Marín-echado, C., Ruano, P., Benmakhlof, M., Akil, M., López-Garrido, A.C., Chabli, A., Ahmamou, M., and González-Castillo, L.: Is there an active

subduction beneath the Gibraltar orogenic arc. Constraints from Pliocene to Present-day stress field, *J. Geodyn.*, 52, 83-96, 710 [doi:10.1016/j.jog.2010.12.003](https://doi.org/10.1016/j.jog.2010.12.003), 2011.

Pedrera, A., García-Senz, J., Ayala, C., Ruiz-Constán, A., Rodriguez-Fernández, L. R., Robador, A. and González Menéndez, L.: Reconstruction of the exhumed mantle across the North Iberian Margin by crustal-scale 3-D gravity inversion and geological cross section, *Tectonics*, 413, 150-154, [doi:10.1002/2017TC004716](https://doi.org/10.1002/2017TC004716), 2017.

Péron-Pinvidic, G., Manatschal, G., Minshull, T.A. and Sawyer, D.S.: Tectonosedimentary evolution of the deep Iberia-715 Newfoundland margins: evidence for a complex breakup history, *Tectonics*, 26, TC2011, [doi:10.1029/2006TC0019070](https://doi.org/10.1029/2006TC0019070), 2007.

Péron-Pinvidic, G., Manatschal, G.: The final rifting evolution at deep magma-poor passive margins from Iberia-Newfoundland: a new point of view, *Int. J. Earth Sci.*, 98, 1581-1597, [doi:10.1007/s00531-008-0337-9](https://doi.org/10.1007/s00531-008-0337-9), 2009.

Pereira, R., Alves, T. and Cartwright, J.: Post rift compression of the SW Iberian margin (eastern North Atlantic): a case for prolonged inversion in the ocean-continent transition zone. *J. Geol. Soc.*, 168, 1249-1263, doi:10.1144/0016-76492010-151, 720 2011.

Pereira, R., Alves, T. and Mata, J.: Alternating crustal architecture in West Iberia: a review of its significance in the context of NE Atlantic rifting, *J. Geol. Soc.*, 174, 522-540, [doi:10.1144/jgs2016-050](https://doi.org/10.1144/jgs2016-050), 2016.

Pinheiro, L.M., Whitmarsh, R.B. and Miles, P. R.: The ocean-continent boundary off the western continental margin of Iberia-II. Crustal structure in the Tagus Abyssal Plain, *Geophys. J. Int.*, 109, 106-124, [doi:10.1111/j.1365-246X.1992.tb00082.x](https://doi.org/10.1111/j.1365-246X.1992.tb00082.x), 725 1992.

Pinheiro, L.M., Wilson, R.C.L., Pena dos Reis, R., Whitmarsh, R. B., and Ribeiro, A.: The western Iberia margin: A geophysical and geological overview, *Proc. Ocean Drill. Prog. Sci. Results*, 149 3.1996

Ramos, A., Fernández, O., Terrinha, P. and Muñoz, J.A.: Extension and inversion structures in the Tethys-Atlantic linkage zone, Algarve Basin, Portugal, *Int. J. Earth Sci.*, 105, 1663-1679, [doi:10.1007/s00531-015-1280-1](https://doi.org/10.1007/s00531-015-1280-1), 2016

730 Ribeiro, A., Cabral, J., Baptista, R. and Matias, L.: Stress pattern in Portugal mainland and the adjacent Atlantic region, West Iberia, *Tectonics*, 15, 641-659, [doi:10.1029/95TC03683](https://doi.org/10.1029/95TC03683), 1996.

Ribeiro, A.: *Soft Plate and Impact Tectonics*, Springer Verlag, Heidelberg, 1-324, 2002.

Roest, W.R. and Srivastava, S.P.: Kinematics of the plate boundaries between Eurasia, Iberia and Africa in the North-Atlantic from the Late Cretaceous to the present, *Geology*, 19, 613-616, [doi:10.1130/0091-7613](https://doi.org/10.1130/0091-7613), 1991.

735 Rueda, J. and Mezcua, J.: Near-real-time Seismic Moment-tensor Determination in Spain, *Seismol. Res. Lett.*, 276, 455-465, [doi:10.1785/gssrl.76.4.455](https://doi.org/10.1785/gssrl.76.4.455), 2005.

Ryan, W.B.F., Carbotte, S.M., Coplan, J., O'Hara, S., Melkonian, A., Arko, R., Weissel, R.A., Ferrini, V., Goodwillie, A., Nitsche, F., Bonczkowski, J. and Zemsky, R.: Global Multi-Resolution Topography (GMRT) synthesis data set, *Geochem. Geophys. Geosyst.*, 10, Q03014, [doi:10.1029/2008GC002332](https://doi.org/10.1029/2008GC002332), 2009.

740 Sallarès, V., Martínez-Loriente, S., Prada, M., Gràcia, E., Ranero, C., Gutscher, M.A., Bartolome, R., Gailler, A., Dañobeitia, J.J. and Zitellini, N.: Seismic evidence of exhumed mantle rock basement at the Gorringe Bank and the adjacent Horseshoe and Tagus abyssal plains (SW Iberia), *Earth. Planet. Sci.*, 365, 120-131, [doi:10.1016/j.epsl.2013.01.021](https://doi.org/10.1016/j.epsl.2013.01.021), 2013.

Sandwell, D.T., Müller, R.D., Smith, W.H.F., Garcia, E. and Francis, R.: New global marine gravity model from CryoSat-2 and Jason-1 reveals buried tectonic structure, *Science* 346, 6205, 65-67, [doi:10.1126/science.1258213](https://doi.org/10.1126/science.1258213), 2014.

745 Schouten, H., Smith, D.K., Montési, L.G.J., Zhu, W. and Klein, E.M.: Cracking of lithosphere north of the Galapagos triple junction, *Geology*, 36, 339-342, [doi:10.1130/G24431A.1](https://doi.org/10.1130/G24431A.1), 2008.

Schulmann, K., Edel, J.B., Martínez Catalán, J.R., Mazur, S., Guy, A., Lardeaux, J.M., Ayarza, P., Palomeras, I.: Tectonic evolution and global crustal architecture of the European Variscan belt constrained by geophysical data, *Earth. Sci. Rev.*, 234, 104195, [doi:10.1016/j.earscirev.2022.104195](https://doi.org/10.1016/j.earscirev.2022.104195), 2022.

750 Serpelloni, E., Vannucci, G., Pondrelli, S., Argnani, A., Casula, G., Anzidei, M., Baldi, P. and Gasperini, P.: Kinematics of the Western Africa-Eurasia plate boundary from focal mechanisms and GPS data, *Geophys. J. Int.*, 169, 1180-1200, doi:10.1111/j.1365-246X.2007.03367.x, 2004.

Sibuet, J., Srivastava, S.P. and Spakman, W.: Pyrenean orogeny and plate kinematics, *J. Geophys. Res.*, 109, B08104, doi:10.1029/2003JB002514, 2004.

755 Smith, D.K.: Grid(s) of multibeam bathymetry at Galapagos Triple Junction, Palisades, New York: *Integr. Earth Data Appl.* Doi:10.1594/IEDA/100005, 2011.

Srivastava, S.P., Schouten, H., Roest, W.R., Klitgord, K., Kovacs, L.C., Verhoef, J., and Macnab, R.: Iberian plate kinematics: a jumping plate boundary between Eurasia and Africa, *Nature*, 344, 756-759, 1990.

760 Stich, D., Serpelloni, E., Mancilla, F.L., and Morales, J. Morales: Kinematics of the Iberia-Maghreb plate contact from seismic moment tensors and GPS observations, *Tectonophysics*, 426, 295-317, doi:10.1016/j.tecto.2006.08.004, 2006.

Soares, D.M., Alves, T.A., and Terrinha, P.: The breakup sequence and associated lithospheric breakup surface: Their significance in the context of rifted continental margins (West Iberia and Newfoundland margins, North Atlantic). *Earth. Planet. Sci. Lett.*, 355-356, 311-326, 2012.

Somoza, L., Medialdea, T., González, F. J., León, R., Palomino, D., Rengel, J., Fernández-Salas, L.M. and Vázquez, J.: 765 Morphostructure of the Galicia continental Margin and Adjacent Deep Ocean Floor: From Hyperextended Rifted to Convergent Margin Styles. *Mar. Geology*, 407, 299-315, doi:10.1016/j.margeo.2018.11.011, 2019.

Somoza, L., Medialdea, T., González, F. J., Machancoses, S., Candón, J. A., Cid, C., Calado, A., Afonso, A., Pinto Ribeiro, L., Blasco, I., Albuquerque, M., Asensio-Ramos, M., Bettencourt, R., De Ignacio, C., López-Pamo, E., Ramos, B., Rincón-Tomás, B., Santofimia, E., Souto, M., Tojeira, I., Viegas, C. and Madureira, P.: High-Resolution Multibeam Bathymetry of 770 the Northern Mid-Atlantic Ridge at 45-46°N: the Moytirra Hydrothermal Field, *J. Maps*, 17, 184-196. doi:10.1080/17445647.2021.1898485, 2021.

Stern, R.J., and Gerya, T.: Subduction initiation in nature and models: A review. *Tectonophysics*, 746, 173-198, doi:10.1016/j.tecto.2017.10.014, 2018.

Srivastava, S.P. and Roest, W.R.: King's Trough: reactivated pseudo-fault of a propagating rift, *Geophys. J. Int.*, 108, 143-150, 775 doi:10.1111/j.1365-246X.1992.tb00845.x, 1992.

Storti, F., Soto Marin, R., Rossetti, R. and Casas Sainz, A.M.: Evolution of experimental thrust wedges accreted from along-strike tapered, silicone-floored multilayers, *J. Geol. Soc.*, 164, 73-85, doi:10.1144/0016-76492005-186, 2007.

Takahashi, M., van den Ende, M.P.A., Niemeijer, A.R. and Spiers, C.J.: Shear localization in a mature mylonitic rock analog during fast slip, *Geochem. Geophys. Geosyst.*, 18, 513-530, doi:10.1002/2016GC006687, 2007.

780 Terrinha, P., Matias, L., Vicente, J., Duarte, J., Luís, J., Pinheiro, L., Lourenço, N., Diez, S., Rosas, F., Magalhães, V., Valadares, V., Zitellini, N., Roque, C., and Mendes Víctor, L.: Morphotectonics and Strain Partitioning at the Iberia-Africa Plate Boundary from Multibeam and Seismic Reflection Data, *Mar. Geology*, 267, 156-174, doi:10.1016/j.margeo.2009.09.012, 2009.

Terrinha, P., Ramos, A., Neres, M., Valadares, V. Duarte, J. et al.: The alpine orogeny in the West and Southwest Iberia 785 margins. In Terrinha, P. et al. (2019). *The Alpine Orogeny in the West and Southwest Iberia Margins*. In: Quesada, C., Oliveira, J. (eds) *The Geology of Iberia: A Geodynamic Approach. Regional Geology Reviews*. Springer, Cham. 487-505, doi:10.1007/978-3-030-11295-0_11, 2019.

Tucholke, B.E., Sawyer, D. and Sibuet, J. C.: Breakup of the Newfoundland-Iberia rift, In: G. D. Karner, G. Manatschal and L. M. Pinheiro (Eds.), *Imaging, Mapping and Modelling Continental Lithosphere Extension and Breakup*, *Geol. Soc. London Spec. Publ.*, 282, 9-46, 2007.

Vázquez, J.T., Medialdea, T., Ercilla, G., Somoza, L., Estrada, F., Fernández-Puga, M.C., Gallart, J., Gràcia, E., Maestro, A., and Sayago, M.: Cenozoic deformational structures on the Galicia Bank Region (NW Iberian continental margin), *Mar. Geology*, 249, 128-149, 2008.

Veludo, I., Dias, N. A., Fonseca, P. E., Matias, L., Carrilho, F., Haberland, C. and Villaseñor, A.: Crustal seismic structure beneath Portugal and southern Galicia (Western Iberia) and the role of Variscan inheritance, *Tectonophysics*, 717, 645-664, [doi:10.1016/j.tecto.2017.08.018](https://doi.org/10.1016/j.tecto.2017.08.018), 2017.

Vilanova, S.P., Nunes, C.F., and Fonseca, J.F.B.D.: Lisbon 1755: A Case of Triggered Onshore Rupture? *Bull. Seismol. Soc. Am.*, 93, 2056-2068, [doi:10.1785/0120020245](https://doi.org/10.1785/0120020245), 2003.

Vilanova, S.P., Nemser, E.S., Besana-Ostman, G.M., Bezzeghoud, M., Borges, J.F., Brum da Silveira, A., Cabral, J., Carvalho, J., Cunha, P.P., Dias, R.P., Madeira, J., Lopes, F.C., Oliveira, C.S., Pereira, H., García-Mayordomo, J., Wong, I., Arvidsson, R., and Fonseca, J.F.B.D.: Incorporating Descriptive Metadata into Seismic Source Zone Models for Seismic-Hazard Assessment: A Case Study of the Azores-West Iberian Region, *Bull. Seismol. Soc. Am.*, 104, 1212-1229, [doi:10.1785/0120130210](https://doi.org/10.1785/0120130210), 2014.

Vissers, R., and P.T. Meijer: Mesozoic rotation of Iberia: Subduction in the Pyrenees?, *Earth Sci. Rev.*, 110, 93-110, [doi:10.1016/j.earscirev.2011.11.001](https://doi.org/10.1016/j.earscirev.2011.11.001), 2012.

Whitmarsh, R.B., Miles, P.R., and Mauffret, A.: The ocean-continent boundary off the western continental margin of Iberia-I. Crustal structure at 40°30'N. *Geophys. J. Int.*, 103, 509-531, [doi:10.1111/j.1365-246X.1990.tb01788.x](https://doi.org/10.1111/j.1365-246X.1990.tb01788.x), 1990.

Whitmarsh, R.B., Pinheiro, L.M., Miles, P.R., Recq, M., and Sibuet, J.C.: Thin crust at the Western Iberia Ocean-Continent Transition and ophiolites. *Tectonics*, 12, 1230-1239, doi:10.1029/93TC00059, 1993.

Zatman, S.: On steady rate coupling between an elastic upper crust and a viscous interior, *Geophys. Res. Lett.*, 27, 2421-2424, doi:10.1029/2000GL011592, 2000.

Zitellini, N., Gracia, E., Matias, L., Terrinha, P., Abreu, M.A., Dealteris, G., Henriet, J.P., Dañobeitia, J.J., Masson, D.G., Mulder, T., Ramella, R., Somoza, L., and S. Diez: The quest for the Africa-Eurasia plate boundary west of the Strait of Gibraltar. *Earth Planet. Sci. Lett.*, 280, 13-50, [doi:10.1016/j.epsl.2008.12.005](https://doi.org/10.1016/j.epsl.2008.12.005), 2009.

Zhong, T., Gordon, R. G., and C. Wang: Oblique seafloor spreading across intermediate and superfast spreading centers, *Earth Planet. Sci. Lett.*, 495, 146-156, [doi:10.1016/j.epsl.2018.05.001](https://doi.org/10.1016/j.epsl.2018.05.001), 2019.

820

825

FIGURES





Figure 1. Study area. The WIM and its main physiographic features including their West (mid Atlantic ridge (MAR), Eurasia- America) and South (Azores-Gibraltar fault zone AGFZ, Eurasia-Africa) plate boundaries. GB: Galicia Bank, IAP: Iberian Abyssal Plain, GIB: Galician interior Basin, VGs: Vasco de Gama seamount, Vs: Vigo seamount, Ps: Porto seamount, NZc: Nazare canyon, Es: Extremadura Spur, STc: Setubal canyon, TAP: Tagus abyssal plane, SVC: San Vicente canyon, FZ: fracture Zone (Kurtachov). Cs: Charcot seamount, As: Atlantic seamount.






Figure 2. a) Seismicity studied in this work showing the events according to magnitude. b) Density map of events studied in this work highlighting the alignments in purple discontinuous lines 1) Galicia, north and 2), Lisbon, south (see text) some focal mechanisms are shown. Seismicity catalogue in the WIM in the period 2003-2022 (magnitude > 2.5, RMS < 1, Smaj < 25, Smin < 25 and Err < 75), according to the parameters described in the methodology section. Topographic base from EMODnet Bathymetry Consortium (2020). Focal mechanism from CMT (blue) Catalog and IGN (red) Seismic Moment Tensor Database.

P1 to P5 white solid lines show the position of the profiles shown in figure 4. Red circle is the 637-drill location cited in Boillot et al. (1988a)

Con formato: Color de fuente: Automático




Figure 3. Representation of cross-sections of seismicity at depth overlaid on crustal cross-sections modified from Granado Amigo Marx et al. (2021-2022), showing the crustal domains. The location of the profiles is shown in figure 2a and the seismicity included in each profile comprises 100 km band along that direction. **Profile 1)** Depth profile of seismicity along alignment North (Galicia) b) Profile along alignment south c), d) North south cross-sections along different longitudes; oceanic domain, and e) close up in cross-section of the seismicity clusters around Galicia Bank and Extremadura Spur.







Figure 4. Map of the WIM seismicity located in this study overlaid on: a) crustal depth of the WIM (Laske et al., 2013); b) gravity map (Sandwell et al., 2014); c) magnetic map (Meyer et al., 2017); d) rift domains of the WIM (Granado et al., 2021).



855

Figure 5. Schematic cartoons to illustrate the different hypothesis on the origin of seismicity of the WIM. a) shear corridors, inverse transform faults b) diffuse plate boundary c) initiation of subduction.

860 TABLES

Table 1. Synthesis of the statistical parameters of the seismicity catalogue prepared from the data of the IGN and IPMA seismic networks.

	RMS	Smaj	Smin	Err	GAP
Minimum	0.10	1.5	0.8	0.1	4
Maximum	0.99	24.8	22.9	74	335
Average	0.57	9.48	5.56	4.98	182.44
Standard deviation	0.21	5.59	4.09	11.46	87.29

Table 2. Focal mechanisms collected for the study area from CMT Catalog and IGN Seismic Moment Tensor Database.

LONGITUDE	LATITUDE	DEPTH (km)	Mw	DATE	DATABASE
-14.4400	39.4800	32	5.9	24/01/1983	CMT
-09.1204	43.7570	27	4.3	23/04/2006	IGN
-14.2965	39.9127	3	4.3	18/08/2007	IGN
-10.6200	40.1800	19	4.8	04/09/2018	CMT
-10.5665	40.2162	29	4.9	04/09/2018	IGN
-14.7700	40.1800	12	4.7	01/11/2018	CMT
-14.9012	40.2458	25	4.7	01/11/2018	IGN
-07.7683	44.1478	21	4.2	29/10/2021	IGN
-09.4919	42.5781	12	4.1	27/01/2022	IGN

← **Tabla con formato**

865