



Measurement Report: The Palau Atmospheric Observatory and its Ozonesonde Record - Continuous Monitoring of Tropospheric Composition and Dynamics in the Tropical West Pacific

Katrin Müller¹, Jordis S. Tradowsky^{2,3}, Peter von der Gathen¹, Christoph Ritter¹, Sharon Patris⁴, Justus Notholt⁵, and Markus Rex¹

¹Alfred Wegener Institut, Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

²Norwegian Meteorological Institute, Oslo, Norway

³Bodeker Scientific, Alexandra, New Zealand

⁴Coral Reef Research Foundation, Palau

⁵University of Bremen, Bremen, Germany

Correspondence: Katrin Müller (katrin.mueller@awi.de)

Abstract. The Tropical West Pacific is recognized as an important region for stratosphere-troposphere exchange, but has been a measurement gap in the global ozone sounding network. The Palau Atmospheric Observatory (PAO) was established to study the atmospheric composition above the remote Tropical West Pacific with a comprehensive instrumental setup. Since 2016, two laboratory containers in Palau host an Fourier-transform infrared spectrometer, a lidar (micro lidar until 2016, cloud and aerosol lidar from 2018), a Pandora 2S photometer and laboratory space for weather balloon soundings with ozone-, water-vapor-, aerosol- and radiosondes. In this analysis, we focus on the continuous, fortnightly ozone sounding program with Electrochemical Concentration Cell (ECC) ozone sondes.

The aim of this study is to introduce the PAO and its research potential, present the first observation of the typical seasonal cycle of tropospheric ozone in the Tropical West Pacific based on a multiannual record of in situ observations, and investigate major drivers of variability and seasonal variation from 01/2016 until 12/2021 related to the large scale atmospheric circulation. We present the PAO ozone (O_3) volume mixing ratios (VMR) and relative humidity (RH) time series complemented by other observations.

The site is exposed to year-round high convective activity reflected in dominating low O_3 VMR and high RH. In 2016, the impact of the strong El Niño is evident as a particularly dry, ozone-rich episode. The main modulator of annual tropospheric O_3 variability is identified as the movement of the Intertropical Convergence Zone (ITCZ), with lowest O_3 VMR in the free troposphere during the ITCZ position north of Palau. An analysis of the relation of O_3 and RH for the PAO and selected sites from the Southern Hemispheric Additional OZonesondes (SHADOZ) network reveals three different regimes. Palau's O_3 /RH distribution resembles the one in Fiji, Java and American Samoa, but is unique in its seasonality and its comparably narrow Gaussian distribution around low O_3 VMR and the evenly distributed RH. A previously found bimodal distribution of O_3 VMR and RH could not be seen in the Palau record.

Due to its unique remote location, Palau is an ideal atmospheric background site to detect changes in air dynamics imprinted on the chemical composition of the tropospheric column. The efforts to establish, run and maintain the PAO have succeeded to



fill an observational gap in the remote Tropical West Pacific and give good prospects for ongoing operations. The ECC sonde record will be integrated into the SHADOZ database in the near future.

25 1 Introduction

The Tropical West Pacific (TWP) is the source region of stratospheric air in boreal winter (Newell and Gould-Stewart, 1981; Fueglistaler et al., 2004; Krüger et al., 2008). Rex et al. (2014) showed that air masses entering the stratosphere in the TWP, also originate in and transit through the local troposphere. Thus, the tropospheric chemical composition above the TWP influences the stratospheric composition on a global scale. The tropospheric composition above the Pacific island state of Palau (7.3°
30 N, 134.5° E), located in the remote TWP, is determined by the unique local air chemistry within the global tropospheric O₃ minimum and the complex interplay of general circulation patterns characterizing the region, such as the Hadley, Walker and Monsoon Circulations (Fig.1). The remote TWP is far from industrial human activities and essentially devoid of major air pollution, particularly from NO_x, leading to the globally lowest tropospheric O₃ columns year-round (e.g. Thompson et al., 2003b; Rex et al., 2014; Müller et al., 2023). A corresponding OH minimum prolongs lifetimes of various chemical species
35 and thus enables their enhanced transport into the stratosphere (Rex et al., 2014). Locally dominant marine convection and vertical mixing as well as transport of ozone-precursor-free air masses with the trade winds from the East, i.e. from the vast Pacific Ocean, foster the clean background state of the troposphere throughout the year. However, air mass transport from remote, polluted locations to the TWP tied to dynamical variations introduces disturbances in the otherwise humid, ozone-poor tropospheric column (Anderson et al., 2016; Müller et al., 2023). Hence, long-term monitoring of various chemical
40 components above Palau can provide insights into dynamical changes in this region with O₃ and water vapor (H₂O) being particularly valuable tracers.

The Palau Atmospheric Observatory (PAO) was established on the grounds of the Palau Community College (PCC) in downtown Koror in 2015 to observe and study the atmosphere above the TWP. It is equipped with a comprehensive instrumental setup which was originally supplied through the EU-funded StratoClim (Stratospheric and upper tropospheric processes
45 for better Climate predictions, 2015-2019) project. Since 2019 the PAO continues operations as a measurement site run by the Alfred-Wegener-Institut (AWI) with further contributions from the Institute of Environmental Physics (IUP), University of Bremen. Two laboratory containers host a Fourier-transform infrared (FTIR) spectrometer, a lidar (micro lidar until 2016, cloud and aerosol lidar from 2018 onwards), a Pandora 2S photometer and laboratory space for weather balloon soundings with ozone- (O₃), water vapor- (H₂O), aerosol- and radiosondes (Fig. 2). In particular, the ozone sounding program with
50 Electrochemical Concentration Cell (ECC) ozonesondes has been successfully running fortnightly soundings with only minor interruptions since the beginning of 2016. The timelines of all measurements are summarized in Fig. 3. Convective mixing and wash-out are dominant modulators of the local air composition due to Palau's location in the global warm pool characterized by the highest sea surface temperatures (SST) globally (see Fig. 1). High-energy convection accompanies the upwelling branches of global atmospheric circulation patterns, which are following the seasonal movement of the Intertropical Convergence Zone (ITCZ). Rainfall rates are highest during the Western Pacific Monsoon between July and October and lowest during
55



boreal spring, but compared to other tropical locations the dry season is less pronounced (Australian Bureau of Meteorology and CSIRO, 2014; Miles et al., 2020). Palau's climate is, thus, specifically affected by the position of the ITCZ, the South Pacific Convergence Zone (SPCZ), the Western Pacific Monsoon and the trade winds. Relevant subseasonal and interannual variations are the Madden-Julian Oscillation (MJO), the El Niño Southern Oscillation (ENSO) and the Quasi-Biennial Oscillation (QBO). The importance of the TWP for global ozone chemistry lies within the interplay of circulation patterns in the upper troposphere/ lower stratosphere (UTLS), i.e. the tropospheric circulation causing strong upward motion of air and the stratospheric Brewer-Dobson circulation which lifts the air further into the stratosphere where it is then transported polewards (Holton et al., 1995). Around 16 km altitude, in the Tropical Tropopause Layer (TTL), mass fluxes in the Brewer Dobson circulation become comparable to those in the Hadley circulation (e.g. Folkins, 2002; Pan et al., 2014). The Brewer Dobson circulation lifts and cools the thermal tropopause when it is strongest during boreal winter (Fueglistaler et al., 2009). O₃ VMR in the TTL are controlled by several overlapping processes, i.e. the dominating, slow radiatively driven ascent, the onset of stratospheric photo-chemical in situ production and the meridional mixing of air from the extra-tropical lower stratosphere. The TTL, as defined by Fueglistaler et al. (2009) has its globally lowest temperatures in this geographic region. It controls final dehydration of air masses and thus stratospheric H₂O, a driver of important O₃ loss cycles in the stratosphere (e.g., Randel and Jensen, 2013). A dynamically driven seasonal cycle of O₃ in the TTL, in turn, influences TTL temperatures (Randel et al., 2007). The coupling between ENSO and the Quasi-Biennial Oscillation (QBO) is a main driver of interannual variability of both O₃ and H₂O in the TTL (e.g. Garfinkel and Hartmann, 2007; Taguchi, 2010; Diallo et al., 2018).

Balloon-borne measurements with Electrochemical Concentration Cell (ECC) ozonesondes are the most practical way to observe O₃ in situ, especially at more remote sites (e.g. Thompson et al., 2019). The highly vertically resolved ozonesonde measurements provide a-priori profiles for many remote sensing techniques including microwave and FTIR instruments which are either ground-based or on satellites (e.g., Vigouroux et al., 2008; Hubert et al., 2016). The ECC sonde measurement technique itself is being continuously validated to yield an overall accuracy of 5 %-15 % in the troposphere, if standard operating procedures (SOP) (Smit et al., 2007; Thompson et al., 2019; Smit and Thompson, 2021) are followed carefully. The latter is especially important in the Tropics, where the instrumental uncertainty is highest in the TTL and critically dependent on the so-called background current (see also Smit, 2014; Rex et al., 2014; Vömel et al., 2020). Reports of near-zero O₃ concentrations in the upper troposphere (UT) of the TWP (Kley et al., 1996; Rex et al., 2014) depend on the details of the background current correction in the data set which limits the detection limit of the sondes under these conditions (Voemel and Diaz, 2010; Rex et al., 2014; Newton et al., 2016).

The latest large campaign conducted in the TWP assessing the tropospheric O₃ column took place during the CONTRAST/-CAST (Convective Transport of Active Species in the Tropics/ Coordinated Airborne Studies in the Tropics) project in January/February 2014 (Pan et al., 2017). Besides ground-based operations, CONTRAST research aircraft were deployed from an airbase in Guam (see Fig. 1), providing a variety of in situ measurements. Apart from various major research campaigns on air chemistry in the TWP since the 1980s (e.g. overview in Müller, 2020), mainly the SHADOZ (Southern Hemispheric Additional OZonesondes) network has been providing regular high-quality monitoring of O₃ in the tropics since 1998 (Thompson et al., 2003a, b, 2007, 2011, 2012, 2017; Witte et al., 2017, 2018).



As there is no long term site located in the vicinity of Palau, prior to the establishment of the PAO, the TWP constituted a gap in the international O₃ sounding network (see also <https://woudc.org/data/stations/>). However, a long-term data record of tropospheric composition in the TWP is required to assess annual and inter-annual variability. To fill this gap, we present the growing record of O₃ profiles, and additional atmospheric variables, measured at PAO.

95 Our overall objective is to survey the potential of measurements obtained at the PAO to explain the air composition and driving meteorological and chemical processes in the TWP. We introduce the PAO and its measurement record from 01/2016 until 12/2021. After an overview of the instrumental setup we focus on the tropospheric O₃ profiles measured by ECC ozone sondes in order to assess dynamical influences on the TWP air composition and their seasonality. We investigate the wind patterns during the timeframe of presented observations from PAO by including additional ground station data. By comparing
100 the Palau tropospheric O₃ time series with observations from selected SHADOZ stations and from the CONTRAST campaign, we examine the regional importance of the PAO to monitor the variability of the tropospheric circulation in the TWP.

2 Overview of the measurements

2.1 Balloon-borne observations: ECC ozone and radio sondes

We present the PAO data record of ECC ozonesondes from 21.01.2016 until 31.12.2021 with 198 fortnightly balloon soundings.
105 At least one intensive campaign with several launches a week took place every year except 2020 and 2021, when this was not feasible due to the Corona pandemic. Most soundings were conducted during daytime, typically around mid-day, however, not at a fixed hour. 17 night-time launches during campaigns were performed with additional use of Cryogenic Frostpoint Hygrometers (CFHs) and Compact Optical Backscatter and Aerosol Lidar Detector (COBALD) sondes and/or coinciding measurements with the Compact Cloud and Aerosol Lidar (ComCAL) (Fig. 2, 3). In this study, no differentiation between day-
110 and night-time profiles was made, as we assume no significant diurnal cycle for O₃ and RH within the tropospheric column.

The quality of observations has been monitored closely throughout the full measurement period. Personnel from the local Coral Reef Research Foundation (CRRF) has been trained to perform the fortnightly soundings, ensuring a high standard. In anticipation of the seasonality of the tropospheric O₃ minimum, intensive campaigns were planned and conducted in fall and spring. The interannual persistence and year-round existence of the tropospheric O₃ minimum in the TWP was already
115 suggested by satellite observations (see Rex et al., 2014) and is now confirmed using in situ measurements collected at the PAO.

Developed in the 1960s (Komhyr, 1969), the basic electrochemical measurement principle of ECC ozonesondes and the related equations are still in use. Ambient air is pumped through an aqueous potassium iodide (KI) sensing solution (cathode) in a Teflon cell by a small gas sampling pump. In a redox reaction all sampled O₃ is converted to iodine, which itself is reduced
120 back to iodide in contact with a platinum electrode. The cathode cell is attached, via an ion bridge, to an anode cell filled with



a saturated KI solution. The generated electrical current I_m [μA] in the external circuit connecting the cells, corrected for a residual background current I_b , is directly proportional to the partial pressure of O_3 (P_{O_3}), calculated by

$$P_{O_3} = \frac{R}{2F\eta_C} \cdot (I_m - I_b) \cdot T_{box} \cdot \frac{c_{pumpcorr}}{f_{pump}}, \quad (1)$$

with R as the universal gas constant, F as the Faraday constant (factor 2 accounts for the stoichiometry, i.e. $2 e^-$ per O_3 molecule), $\eta_C \approx 1$ as a factor for the conversion efficiency, T_{box} [K] as the so-called “box” or pump temperature, f_{pump} [$cm^3 s^{-1}$] as the gas volume flow rate and finally, $c_{pumpcorr}$ as an empirical correction factor for the pump efficiency depending on pressure (see Tab.A1 for details on corrections).

Experiments conducted at the PAO led to the decision to use the pressure dependent background current correction for the calculation of O_3 VMR in this study and the companion study by Müller et al. (2023). Müller (2020) presents the development, status and preliminary results of a device attachable to regular ECC sondes which measures the background current in-flight. The device enables time-controlled intermissions by switching from a measurement of ozone in the ambient air to measurements using ozone-free air. The ozone-free measurement periods in modified ECC sondes launches in Palau and Ny Ålesund support the application of a pressure-dependent background current correction to the raw tropospheric signal.

The assessment of potential near-zero O_3 measurements in the TWP and the scientific discussion on that topic (e.g. Smit, 2014; Rex et al., 2014; Voemel and Diaz, 2010; Newton et al., 2016) demanded a particular focus on ensuring the high-quality of the equipment and attention to detail during all steps required for generating the PAO data record, including the pre-flight preparations of the sonde and the post-processing of the raw signal. The Assessment of Standard Operating Procedures for OzoneSondes (ASOPOS), an international consortium of ozonesonde experts established under the umbrella of the WMO’s Global Atmosphere Watch (GAW) program has put together a reference for standard operating procedures (SOP) based on extensive assessments (GAW Report 201, Smit (2014), updated in Smit and Thompson (2021)). The GAW recommendations for our sonde model, type 6A manufactured by Science Pump Corporation (SPC), have been carefully followed with the exception of the background current correction method and are summarized in Table A1.

Preparation of the ECC sondes according to SOP is done in two steps, the first step three to thirty days before launch, and a second preparation within 24 hours before the launch, usually on the launch day. In Palau, the preparation and storage in between take place inside an air-conditioned and dehumidified lab container, providing a stable environment with temperatures around $25^\circ C$ and RH around 40 %. All parameters measured during preparations are documented and available as metadata. Raw pump current measurements during flight are available to readers upon request in addition to the Vaisala software output files and sounding data in NASA Ames format. Data presented in here have been calculated from raw data with specifically developed software (using Eq.1 and corrections summarized in Tab.A1).

Vaisala radiosondes were used for data transmission as well as to measure pressure, temperature and humidity and to provide the GPS coordinates. The RS92-SGP model and its interface to the ECC sonde was used until October 2017 (see Dirksen et al. (2014) for technical details and quality assessment). Thereafter, the ground receiving unit and software were switched to facilitate use of Vaisala’s new standard radiosonde RS41-SGP (Vaisala, 2017; Sun et al., 2019). In support of international



efforts to assess differences between the old and new radiosonde (for example by the Global Climate Observing System
155 Reference Upper-Air Network, GRUAN) four dual soundings with both radiosonde types, RS92 and RS41, were conducted in
spring 2018 (Dirksen et al., 2019). We expect no major impact due to this change regarding the tropospheric O₃ monitoring.
However, the RH measurements show significant differences, particularly in the upper troposphere, which will be investigated
in the future.

The new RS41 radiosondes facilitate monitoring of the ECC sonde voltage and motor current. First measurements with the
160 new RS41 revealed suspected performance issues of ECC sonde wet cell batteries, which were then replaced with dry batteries.
Comparison with satellite observations from Aura MLS (Microwave Limb Sounder, Froidevaux et al. (2008), not shown here)
reveal discrepancies in stratospheric O₃ observations for around 20 % of soundings in the data set. At pressure levels below 20
hPa (altitudes above the 20 hPa level), the affected ozonesondes measured 1-2 ppm less O₃ (10-20 %) than the average of 5
satellite measurements of the nearest overpass. An early drop of the wet cell battery voltage would be a possible cause, but as
165 the battery voltage was not monitored with RS92 radiosondes, clarification of this effect is not possible. We assume no impact
on the tropospheric profiles below 20 km and advise to be cautious of this effect when using the time series for either total
column or stratospheric O₃ assessment.

2.2 Balloon-borne observations: water vapor and aerosol sondes

As stratospheric water vapor controls the radiative budget of the stratosphere and has great impact on global warming (e.g.
170 Riese et al., 2012; Solomon et al., 2010), the growing PAO record of CFH soundings can be relevant in monitoring and
assessing the variability of this important quantity (doi in preparation). 17 measurements with CFH sondes by Environmental
Science (EN-SCI) were performed from 2016-2021 (see Sect. 2.1 and Fig. 3) in combination with ECC and COBALD sondes.
The CFH instruments use a chilled-mirror principle to detect water vapor up to 28 km altitude with an uncertainty below 10
% (Vömel et al., 2007). The COBALD technique by MyLab, Switzerland, enables measurement of aerosol backscatter at two
175 optical wavelengths, 455 nm (visible blue) and 940 nm (infrared), and thus detection of cirrus clouds and aerosol layers in
the TTL (e.g. Brunamonti et al., 2018). COBALD requires launches after dark, which determined the combined ECC-CFH-
COBALD launches to be at night-time. COBALD deployment has been in cooperation with the ETH Zurich, where the sonde
was developed based on an original setup by Rosen and Kjöme (1991). The combined soundings of CFH and COBALD are an
excellent in situ validation for the ComCAL lidar measurements, which will be subject of upcoming studies.

180 2.3 Fourier-transform infrared (FTIR) spectrometer

The solar absorption spectrometry has been established as a powerful tool to study the composition of the atmosphere. Using
the sun as an external light source allows to measure the concentrations of about 30 trace gases in the atmosphere. Besides
the total column the analysis of the spectral line shape allows retrieval of the concentration profiles in about 2-4 atmospheric
layers up to 30 km. The PAO is equipped with the Bruker 120 M mobile high resolution Fourier-transform infrared (FTIR)
185 spectrometer. The solar tracker is mounted on the top of the hosting container. If weather conditions permit (no clouds, sunny),
the dome opens, the solar tracker follows the sun, and spectra are recorded. O₃ is measured around 2000 cm⁻¹ at a resolution of



0.005 cm⁻¹. Measurement times for individual scans are about 2-5 min. The retrieval is performed using SFIT4, a code based on the optimal estimation method. O₃ can be retrieved at about four independent layers between the surface and about 30 km.

In addition, many other trace gases show absorption features in the measured spectra, like CH₄, CO, OCS, CH₂O, HCN, C₂H₆ and N₂O. Currently the instrument is modified to cover also the long-wave infrared region, allowing a better retrieval of O₃ and studying other trace gases, like HNO₃.

2.4 Compact cloud aerosol lidar (ComCAL)

The „Compact Cloud Aerosol Lidar“ (ComCAL) deployed at the PAO since 2018 has originally been designed in 2006 (Immler et al., 2006) and is hosted in its own laboratory container. As a multi-wavelength lidar system capable of observing aerosol and cloud particle backscatter, extinction and depolarisation, it superseded the previously installed micro Lidar installed in the first laboratory container, measuring only in spring 2016 (see Cairo et al. (2021) and Fig. 3). The ComCAL consists in its current configuration of flash lamp pumped Nd:YAG laser from Quantel Brilliant with 20 Hz and an energy of 120 mJ / 180 mJ and 65 mJ for the three colors of 355 nm, 532 nm and 1064 nm, respectively. The recording telescope is a 40 cm Newton with 1200 mm focal length and 1mm field stop (0.83 mrad field of view). The signals are detected by Hamamatsu photomultipliers for all colors, except for 1064, for which a Licel APD is employed. The Transient Recorders are also from Licel (12bit, 20 MHz) and collect the signals in both photo counting and analog mode. The depolarization is measured at 355 nm and 532 nm by using a Glen-Taylor prism in the detection branch that is coupled to the laser. Hence, the lidar profiles parallel (“p”) and perpendicular (“s”) to the laser are recorded alternately by the same photomultipliers. This configuration facilitates the calibration of the depolarization, however, as the “s” signals perpendicular to the laser polarization are generally weak, the signal to noise ratio is lower for the “s” polarization, compared to the “p” signals.

Using a monostatic design (same optical axis for laser and recording telescope) valid data is obtained above 500 m altitude. Lidar data are stored with a resolution of around 2 minutes (1200 laser shots) and 7.5 m. The evaluation is done according to (Klett, 1985) and (Ansmann et al., 1992), typically with 10 min and 60 m resolution. The particle extinction is measured by the inelastic Raman scattering of N₂ at 387 nm. With a resolution of 2h / 60m during night-time typically a signal to noise ratio of about 10 is found for this Raman channel in approximately 6 km altitude. In summary, the ComCAL lidar is capable to track aerosol layers even in the UTLS region, as well as subvisible or optically thin cirrus, at least concerning the backscatter coefficient and the depolarization. Hence, analysis of cloud properties, as outlined by Cairo et al. (2021) are feasible. Moreover, the system shall be used in conjunction with the balloon-borne soundings at the PAO to gain more information about the TTL, e.g. to what extent clouds and aerosol layers may be related to features in temperature, RH or O₃ concentration.

2.5 Auxiliary Data (Palau weather station, SHADOZ)

Located 800 m South-West of the PAO on a slope on the same island, the US National weather service and NOAA operate a general weather station (PTRO 91408) with twice daily radio soundings. In August 2018, the launch site was switched to a location further away (8km) in Airai, next to the Palau International Airport.



As a tropical station the PAO shares technical challenges as well as certain atmospheric characteristics with O₃ measurement stations assembled in the SHADOZ network (e.g. Thompson et al., 2003a, b, 2007, 2011, 2019). Since 1998, SHADOZ fills observational gaps in the ozonesonde record of the Southern Hemisphere and the tropics by providing consistent data from remote tropical and subtropical sites. To embed our O₃ profile observations in a larger geographical context, we include data from selected SHADOZ stations in our analysis (see Fig.1 and Tab.1). The data version 6 is used in this study and incorporates a reprocessing of all data according to state-of-the-art procedures (see Smit et al., 2012; Deshler et al., 2017; Witte et al., 2017; Thompson et al., 2017; Witte et al., 2018).

The latest JOSIE campaign (Juelich Ozone Sonde Intercomparison Experiment) was dedicated to tropical stations and their particular challenges and confirmed the high quality of methods and operating procedures used within SHADOZ (Thompson et al., 2019). In terms of common characteristics in free-tropospheric and TTL ozone, the SHADOZ stations can be divided into several groups: the Western Pacific and eastern Indian Ocean (Kuala Lumpur, Java, Fiji, American Samoa) the equatorial Americas (San Cristobal, Costa Rica, Paramaribo), the Subtropics (Hanoi, Hilo, Irene, Réunion) and the Atlantic and Africa (Natal, Ascension, Cotonou, Nairobi) (Thompson et al., 2012). We focus our comparison on the four Western Pacific stations. In addition, Hanoi, being the second closest SHADOZ station, Costa Rica, located at a similar latitude, and Hilo, as another Pacific station North of the equator, are considered (compare Table 1).

3 Results and Discussion

We present the PAO time series of ECC ozone and radiosoundings from 2016 until 2021 as well as accompanying measurements of other parameters. First, the local tropospheric variability is discussed in the context of global circulation patterns. Then, the PAO data are compared to measurements from selected SHADOZ sites and from the CONTRAST campaign to highlight the regional relevance.

3.1 Local tropospheric O₃ variability

Figure 4 shows time series of ozone and humidity measured above Palau from 2016 until 2021 as time-height cross-sections in the geopotential height range from 0-20 km. To provide context, the Multivariate ENSO Index (MEI, Wolter and Timlin, 2011) is shown in panel c). Arrows at the top of panels a) and b) indicate the launch times of ozonesondes and radiosondes at PAO. For the plot the data was time interpolated between the next measurements. If no sonde launch was conducted for more than 20 days, a gap is visible in the time series plot. The tropospheric ozone VMRs are lowest from July to October (< 30 ppb) and peak in northern hemispheric spring (20-100 ppb). The boundary layer (0-2 km) is generally very humid and ozone-poor, mostly > 80% RH and < 20 ppb O₃ VMR. In the mid-troposphere (2-14 km) greater differences throughout the year occur for both O₃ and RH (from < 10% to > 90%). The pronounced anti-correlation of O₃ and RH in at these altitudes can be used to determine the origin of air masses (Müller et al., 2023).

Figure 5 shows in more detail the O₃ VMR, relative humidity and additionally the temperature in the geopotential height range from 14-20 km in the UTLS. The evolution of the humidity profile at these altitudes clearly shows the change of ra-



diosonde instrumentation which took place in 2017 when the Vaisala RS92 was replaced with its successor, the RS41, which tends to report higher relative humidity values in the UTLS. Generally, the TTL composition is modulated by the seasonality of the Brewer Dobson circulation and the changing position of the ITCZ, which is shifting the Hadley circulation north- and southwards over the course of the year. O₃ VMR range from roughly 20 - 150 ppb at approx. 16 km, which can be related to
255 both variability in tropopause height and uplift of ozone-poor air masses in active convection. RH varies from 0 to > 50 %. Day-to-day-variability seems to play a strong role in TTL RH variability, particularly below around 16 km.

The overall variability of O₃ VMR in the tropospheric column above Palau seen in Figure 4 can be largely explained by annual variations of the large-scale circulation (compare Müller et al., 2023). Besides the overall high convective activity in the TWP warm pool, Palau's climate is governed by trade winds, the position of the ITCZ and the Western Pacific Monsoon.

260 Interannual variations of the chemical air composition above the TWP are tied to ENSO (e.g. Shiotani, 1992; Gettelman and Forster, 2002; Smith et al., 2012). The PAO data record starts during the strong El Niño event in 2016 (e.g. Huang et al., 2016; Diallo et al., 2018), which can be seen from the Multivariate ENSO index in Fig. 4. The index shows a return to neutral conditions in May/June of 2016. The years 2017 and 2018 were governed by moderate La Niña events. A weak El Niño in 2018/2019 was followed by a longer and stronger La Niña phase that is still ongoing in the beginning of 2023. In the following,
265 we focus on the prominent El Niño event of 2016. A longer time series will eventually allow investigation of the full ENSO variability.

In analogy to the impact of the strong 1997/8 El Niño, Palau experienced a drought during spring 2016 with critically low water reserves on the island (Di Liberto, 2016; Polhemus, 2017). During El Niño, the warm pool shifts towards the central Pacific and the TWP experiences negative SST anomalies and positive anomalies of outgoing longwave radiation along
270 with a weakening of the trade winds, leading to suppressed convection and decreased cloud coverage and precipitation. Our measured time series reveals clear corresponding characteristics during this time period, i.e. an unusual dry and ozone-rich mid-troposphere (Fig. 4) and a weaker vertical gradient in O₃ VMR over the whole TTL, with enhanced O₃ levels in the upper troposphere (UT) below the cold point tropopause and decreased O₃ in the lower stratosphere (LS) above 18 km (Fig. 4).

Less up-lift of ozone-poor boundary layer air and less wash-out of O₃ precursors due to the strong El Niño event can explain
275 the high mid-tropospheric and UT O₃ VMR between March and July 2016 above Palau. Individual observations of higher than average O₃ in the UT are made every year during this season and are visible as tongue-like features in Fig. 4 (a). Decreased LS O₃ VMR, as can be seen in Fig. 5 (a) at the start of the time series, have been documented as a zonal El Niño response by different studies and are attributed to enhanced tropical upwelling (e.g. Randel et al., 2009; Konopka et al., 2016; Diallo et al., 2018).

280 Figures 5 (b-c) show the ozonopause, defined as the level of 90 ppb O₃ VMR following Prather et al. (2011), as a black solid line relating the TTL O₃ seasonality to photo-chemical processes. The ozonopause generally follows the temperature cycle with a maximum in winter, when the thermal tropopause is coldest and reaches its maximum altitude. In spring/early summer 2016, it differs clearly from the unusual cold thermal tropopause (Fig. 5 (c)). This TTL temperature anomaly has been associated with warm ENSO events by various studies (e.g. Kiladis et al., 2001; Randel and Thompson, 2011; Paulik and
285 Birner, 2012). LS water vapor levels are higher during this period compared to the following years. This will be assessed more



thoroughly in a future study deploying the PAO data record from CFH soundings, which do not include the inhomogeneity in the RH data record shown here caused by the change in radiosonde models. In general, global lower stratospheric water vapor is tied to the variability of TTL temperatures, since the final dehydration of air masses takes place in the TTL (e.g. (Gettelman and Forster, 2002; Schoeberl and Dessler, 2011; Fueglistaler, 2012; Randel and Jensen, 2013)).

290 Figure 6 shows monthly mean O_3 VMRs (c) and altitude cross sections with individual O_3 VMR measurements averaged in 300 m bins around 8 km (d) and 18 km (e). O_3 measurements are complemented in the Fig. with monthly means of meridional (a) and zonal (b) wind for the time series alongside the monthly latitudinal position of the ITCZ derived from TRMM data averaged from 1998 to 2009 for a region between 125-175° E and 0-20° N (after Shonk et al., 2018) (f). The O_3 column has two dominant annual features, i.e. a mid-tropospheric cycle with a clear minimum from July to October (10-30 ppb, Fig. 6(d)), and
295 a reverse cycle of greater amplitude in the TTL with a maximum from June to September (Fig. 6(e)). In the mid-troposphere, individual measurements outside the minimum show a greater spread compared to the persistent signal in the TTL, inter- as well as intra-annually, as can be inferred from Figure 6 (d) and (e). Around 8 km altitude, VMR as low as during the mid-tropospheric minimum period are occasionally also measured between January and June in years other than the El Niño year 2016. It should be noted that during the timeframe in which O_3 in the mid-troposphere is expected to reach its minimum,
300 the sounding frequency was increased in most years, to increase the robustness of particularly low O_3 observations. The TTL seasonal cycle is a known zonal phenomenon, which is mostly controlled by the Brewer-Dobson circulation (Thompson et al., 2003a, b; Randel et al., 2007).

The annual movement of the ITCZ can explain the periodicity of the annual O_3 cycle in the mid-troposphere. The ITCZ crosses Palau's latitude in June and October (for the definition of ITCZ latitude see caption of Fig. 6 and Shonk et al. (2018)).
305 On average, the ITCZ is located 6° N of the equator, but moves towards the warming hemisphere throughout the year. Therefore, Palau is generally close to the ITCZ, but closest during its northernmost position from July until October. The dominant influence of the Northern Hadley cell and thus transport of air masses to Palau with northeasterly trade winds is only interrupted during these few months, when the Western Pacific Monsoon reaches Palau from the South-West below 5 km altitude. Above 5 km though, the influence of the monsoon circulation vanishes and easterly winds dominate throughout the remaining
310 tropospheric column. Additional moisture transported to Palau in this wettest season of the year corresponds well to the lowest O_3 measurements. But measurements of low O_3 (see Fig. 6 (d)) paired with high RH occur year-round and are interpreted as the result of convective activity, which transports boundary layer air upwards, which is humid and ozone poor. To some extent this convection persists year-round due to the overall close proximity of the ITCZ. The wind components measured above Palau (Fig. 6 (a,b)) show a seasonality corresponding to the ITCZ movement with particularly opposite regimes in the
315 meridional wind for June to October compared to November until May. The shift in regimes corresponds well with the annual movement of the ITCZ shown in Fig. 6 (f). The tropospheric O_3 seasonality and in particular the transport processes driving this pattern are discussed in more detail by Müller et al. (2023). The average zonal wind in the UTLS has to be treated with caution due to the irregular shifting between westerlies and easterlies with the QBO.



3.2 Regional tropospheric O₃ variability

320 3.2.1 Comparison with SHADOZ

As O₃ and H₂O are valuable tracers of dynamical transport in the TWP, we compared the O₃-RH relation measured at the PAO to other selected SHADOZ stations to assess their interdependency and variability in the region. Given its position far away from major anthropogenic emissions sources the PAO provides an atmospheric background site well-suited to study the impact of global climate change on seasonal air composition and dynamics. In Figure 7, 2D histograms for O₃ VMR
325 (≤ 100 ppb) versus RH at selected SHADOZ stations and at the PAO are shown for the free troposphere (3-14 km). Data is essentially unsmoothed, i.e. no spatial or temporal averaging beyond raw data processing has been applied, in order to preserve the simultaneously measured combination of the two tracers. This is an important advantage over averaging of data into mean seasonal or annual profiles with altitude, where the effects of layered structures in certain altitudes might be canceled out by an equally dominant background (compare Pan et al., 2015; Müller et al., 2023). The sampling frequencies are not regular
330 and for different stations data has been accumulated over different time periods (compare Tab.1). The resulting distributions, therefore, cannot be compared in a quantitative climatological manner, but we can assume a good representation of interannual variability for SHADOZ data (time series of 13 to 21 years). Histogram values in Fig. 7 are normalized to the total number of observations in the free troposphere for each station. Thus, they can be interpreted as a statistical probability distribution for a certain O₃/RH combination to occur.

335 In this sense, the given stations can be classified into three groups based on the distribution of O₃ versus RH:

1. a fairly narrow, almost Gaussian distribution of O₃ VMR with evenly distributed RH (Costa Rica, Kuala Lumpur),
2. predominantly dry air spread over a wider range of O₃ VMR (Hanoi, Hilo), and
3. a mixture of the two previous categories: an “L”-shaped distribution with a dominant mode of low O₃ over the whole RH range and a tail towards higher O₃ VMR corresponding to low RH values (Palau, Fiji, Java, American Samoa).

340 With a reference to these categories, the statistical O₃ box plots for the eight stations are arranged as a longitudinal cross-section over the Pacific Ocean in Fig. 8 (compare Thompson et al., 2012, 2017). The most frequently observed O₃ VMR (maxima in the marginal 1D histograms of Fig. 7) are superimposed as orange circles on the boxplots. In Figure 8, we see the separation of the Pacific island stations (turquoise boxes) from the other groups in terms of O₃ VMR (lowest median, mean values and maximum frequency). The similarity between the Western Pacific stations can be explained by their tropical
345 marine climate and location far from industrial centers. Without NO_x emissions, O₃ loss in a marine, humid environment will dominate over O₃ production.

With O₃ as an indicator for its precursor NO_x (Crawford et al., 1997; Gao et al., 2014; Rex et al., 2014) we would expect a decrease from East to West, while crossing the remote Pacific and following the trade winds within the ITCZ (Thompson et al., 2012), as more and more NO_x is lost by conversion to HNO₃ and subsequent washout in convection is outweighing NO_x
350 production (Graedel et al., 1994). But this lateral ozone gradient is not apparent in any of our statistical measures. Instead,



according to the statistics for the free troposphere, Palau sets the lowest boundaries for all selected stations, except for the minimum range value (excluding outliers).

The Palau data further stand out due to the exceptionally narrow O_3 distribution around ~ 22 ppb (compare marginal 1-D histogram in Fig. 7). The fairly even distribution of RH rather resembles histograms classified into group 1. This consistency within the near-equatorial Northern hemispheric stations, all located within $\pm 5^\circ$ latitude of Palau, is caused by their close vicinity to the average position of the ITCZ around 6° North and thus overall high rainfall rates (e.g. Schneider et al., 2014). The most unique attribute of the distribution based on PAO data is the essential absence of a dry season in terms of tropospheric humidity values. The driest air masses paired with enhanced O_3 VMR (horizontal part of the “L”-shape) are mostly limited to the time period of November until April (for more details on seasonal distributions, see Müller (2020)). But while moist air masses above 40 % are rare during the dry seasons for other stations in group 3, these are frequently observed year-round in Palau, also during the drier period, with the exception of the strong El Niño in 2016. The RH distributions of Fiji, America Samoa and Java are skewed towards lower values during their pronounced dry seasons. Weak overall anti-correlation of O_3 and RH occurs for Fiji and American Samoa data (Fiji: $R=-0.46$, American Samoa: $R=-0.42$). Kuala Lumpur and Costa Rica (group 1) constitute the boundaries of the longitudinal Pacific cross-section (brown boxes in Fig. 8) and are continuously influenced by continental pollution and deep convection, with most frequent O_3 VMR observations at around 30 and 40 ppb, respectively, and evenly distributed RH. That means O_3 VMR are indeed enhanced compared to the most frequent O_3 measurements in group 3, but there are hardly any observations greater than 60 ppb, possibly due to local mixing processes and O_3 loss in the humid free troposphere. Kuala Lumpur is almost completely lacking observations below 10 % RH (see Fig. 7 (h)).

Both subtropical stations, Hanoi and Hilo, in group 2 (purple boxes in Fig. 8) are similarly affected by pollution with a significantly wider spread in their O_3 distributions compared to all other stations, especially towards higher values (Fig. 7 (e, f)). Their free-tropospheric O_3 to RH ratio is therefore especially different from the one at Palau. The tendency towards lower RH values is tied to their location close to the subsiding branch of the Hadley circulation. The less humid troposphere chemically acts in favor of higher O_3 VMR. The semi-permanent North Pacific High above Hawaii, together with the cold SST in the East Pacific, is responsible for a stable trade wind inversion which suppresses deep convective activity above Hilo and enables long-range transport of potentially polluted air masses from both Asia and North America (e.g., Oltmans et al., 2004). Hanoi, in turn, is affected by Asian outflow within shorter distances to the sources (Ogino et al., 2013).

For this study, we have neglected variations in altitude despite the inhomogeneous distribution of O_3 values in the free troposphere. We particularly excluded the boundary layer and TTL, but included the level of deep convective outflow (10-12 km altitude). O_3 VMR are typically very low at this level in the TWP (e.g., Folkins, 2002; Gettelman and Forster, 2002; Pan et al., 2014). At 200 hPa, approx. 25 % of the O_3 observations in Palau are below 20 ppb (not shown here, c.f. Müller, 2020). For American Samoa, Fiji and Java, Solomon et al. (2005) found even higher percentages between 30% and 45 %. However, at lower altitudes, Palau experiences higher fractions of low O_3 VMR compared to their study and the overall statistics for the free troposphere (Fig. 8) attest to the exceptional dominance of the ozone-poor tropospheric column compared to other SHADOZ sites.



385 3.2.2 Comparison with CONTRAST

A detailed analysis of the relation between tropospheric O₃ VMR and RH in the TWP has been performed by Pan et al. (2015) for data from the CONTRAST campaign in 2014 (see Pan et al. (2017) and Fig. 1). They found a bi-modal distribution of free-tropospheric O₃ VMR within the altitude profile. The humid, ozone-poor background or “primary mode” was isolated by simply removing all “dry” data with RH less than 45 %, which indicated an entirely convective control of the primary mode.

390 The study thus still refrains from resolving the individual vertical structure of layers, but proposes the RH threshold as an overall free-tropospheric criterion for the primary mode.

For Palau a similar analysis does not reveal a bi-modal distribution for the full data set (Fig. 9 (c)), and a threshold of 45 % does not separate all higher O₃ observations from the primary low O₃ mode (Fig. 9 (d)). Figure 9 (a) depicts the layer-normalized density distribution for all O₃ observations at the PAO from the ground to 15 km, with data binning according to Pan et al. (2015). Using the same method of analysis, the panel on the right, Fig. 9 (b), shows only “wet” data, with RH greater than 45%. The primary mode, i.e. the layer maximum, is highlighted in dark colors and obvious within both distributions. The “wet” data feature less frequent occurrences of O₃ measurements with more than 45 ppb. However, the highest frequency of O₃ observations in the mid-tropospheric layer of expected highest occurrence of enhanced O₃ air masses (from 320 to 340 K potential temperature) is approximately the same for both the full and the “wet” data set at around 20 ppb (Fig. 9 (c,d)). The 400 tail in the “wet” distribution is reduced, but not completely absent, as it was shown for the CONTRAST data set (see Pan et al., 2015, Fig.3f). The Palau data does not show a clear separation into two modes in the first place (compare Fig. 9 (c) with Fig. 3c in Pan et al. (2015)).

The differences with the aircraft measurements performed in the TWP from the Guam air base can be understood by looking at the uniqueness and seasonality of the free-tropospheric O₃ to RH relation at Palau (Fig. 7. In the PAO time series, air masses 405 with higher O₃ content (> 40 ppb) also occur under wet conditions (> 45 %). Likewise, dry and ozone-poor air is frequently observed, presumably measured in higher altitudes and explained by the temperature-dependent vertical RH-gradient (e.g., Mapes, 2001). The central role of local convection in homogenizing air masses, lifting ozone-poor air from the ground and further depleting O₃ in the tropospheric column, is inevitable. But RH is not sufficient as a stand-alone indicator for a convective and thus local profile in the TWP.

410 Müller et al. (2023) show a different approach to separate humid, ozone-poor atmospheric background air masses from the Palau time series and, with this, contribute to the debate on the origin of frequent filaments of high O₃ and low RH interrupting the tropospheric background in the wider TWP (e.g. Anderson et al., 2016).

4 Conclusions

The PAO with its comprehensive instrumental setup monitors different atmospheric constituents since 2016. In particular, the 415 O₃ time series from regular ECC ozone sonde measurements fills an observational gap in the international ozone sounding network. The PAO, being a highly valuable candidate for SHADOZ, will be included in the network in the near future (private conversation with SHADOZ’s principal investigator Ryan Stauffer and Anne Thompson).



First analysis of the PAO O₃ and RH records (2016-2021) highlights the suitability of Palau as a site for studies of the atmospheric background enabling a comprehensive analysis of the influence of climate change on dynamics and air chemistry
420 in the TWP. The regional tropospheric O₃ minimum is confirmed by the in situ measurements showing that lowest O₃ values accompanied by mostly high RH are observed in the mid-troposphere from July until October. The regional TTL seasonality of O₃ is confirmed and first signals of interannual variability due the ENSO are evident. Overall, the seasonal tropospheric O₃ variability is governed by the movement of the ITCZ, which is close to Palau year-round. In the UTLS, the Brewer-Dobson circulation and its interplay with the uprising branch of the Hadley circulation take over control. As the ITCZ acts as a
425 boundary for interhemispheric transport (Sun et al., 2022), its position North of Palau from July until October inhibits transport of potentially polluted northern hemispheric air masses to the region, which further promotes the seasonal mid-tropospheric O₃ minimum and is further explored in back trajectory analysis by Müller et al. (2023).

The comparison between data from the PAO and selected SHADOZ stations for the free troposphere in the O₃/RH space emphasizes the uniqueness of the atmospheric composition above Palau. Low O₃ values attributed to a background atmosphere
430 occur year-round in dry or humid conditions within the free troposphere, but show a seasonality. Dry air masses of enhanced O₃ VMR, the horizontal part of an “L”-shape in the O₃/RH distribution, are mostly observed from November until April.

Comparison of the PAO data set with data from the CONTRAST aircraft campaign caution against the generalization of conclusions from a regionally and timely limited snapshot in the TWP. The aircraft campaign took place during the time of the year with greatest occurrence of enhanced O₃ VMR. Pan et al. (2015) proposed a “fundamental bimodal distribution
435 of tropical tropospheric ozone” for the whole TWP based on the analysis of CONTRAST data, which separates ozone-poor humid air masses from ozone-rich dry air. The application of their method on the full PAO time series, however, did not confirm the general existence of such a separation, which occurs only during individual years or seasons. Müller et al. (2023) successfully used a different, statistical approach on the PAO time series to separate humid ozone-poor background air masses from anomalously dry and ozone-rich air.

While this paper provides a first overview of the O₃ data record and associated meteorological variables, we encourage the scientific community to use the PAO datasets to study relevant phenomena in this undermonitored region of the Pacific Ocean. Analysis of the water vapor measurements by CFH sondes as well as aerosol and cloud observations by ComCAL will give further insights on UTLS processes in this key region of global stratosphere-troposphere exchange. In the stratosphere, water vapor drives important gas-phase O₃ loss cycles as the primary source of HO_x and is involved in the heterogeneous chemical
445 processes leading to the formation of polar stratospheric clouds, thus eventually promoting chlorine activation and resulting in polar ozone loss (e.g. Solomon et al., 1986; Manney et al., 1994; Crutzen et al., 1995, see Diallo et al., 2018 and references therein).

Code and data availability. All code used to produce the data and results is available upon request. The ozonesonde dataset is available under
450 <https://doi.org/10.5281/zenodo.6920648> and will be included in the SHADOZ database in the future. SHADOZ data can be accessed via the archive at <https://tropo.gsfc.nasa.gov/shadoz/Archive.html>. The Palau weather station data (station code: PTRO 91408) was accessed via the



upper-air sounding database provided by the University of Wyoming (weather.uwyo.edu/upperair/sounding.html). ECMWF ERA5 data used for SST in Fig. 1 was accessed via the Copernicus Climate Change Service Climate Data Store (CDS): <https://cds.climate.copernicus.eu/>. The ENSO Index data is provided by NOAA (<https://psl.noaa.gov/enso/mei/>).

Appendix A

455 All O₃ VMR profiles measured at the PAO from 2016 to 2021 are shown in Fig. A1 sorted by month and alongside some statistical measures. Details on the statistical evaluation are given in Müller (2020). Table A1 provides specifications on the operating procedures of ECC soundings established at the PAO, please note the deviation from international SOP regarding the background current correction.



460 *Author contributions.* KM prepared the first draft and worked closely with JT on the text and figures. PG and SP were involved in Sect. 2.1 and 2.5. JN provided input for Sect. 2.3, CR for Sect. 2.4. PV and MR supported the analysis and provided effective and constructive comments to improve the manuscript. KM, SP, JT and others performed the measurements. All authors contributed to writing the paper.

Competing interests. None of the authors has any financial or personal relationships that could inappropriately influence or bias the content of the paper.

465 *Acknowledgements.* The setup of the PAO and this study was mostly funded by the European Union under the 7th Framework Program as part of the StratoClim project (www.stratoclim.org), grant number 603557. The authors want to thank Dr. Patrick Tellei, President of the Palau Community College, for provision of space, German Honorary Consul Thomas Schubert, for overall support, and various people and institutions for operations at the PAO: Pat and Lori Colin (CRRF), Gerda Ucharm (CRRF), Ingo Beninga (impres GmbH), Wilfried Ruhe(impres GmbH), Winfried Markert (Uni Bremen), Tine Weinzierl (Uni Bremen) and Jürgen "Egon" Graeser (AWI). The authors want to further thank Jonathan Shonk for kindly providing data for Figure 6 (f).



470 References

- Anderson, D. C., Nicely, J. M., Salawitch, R. J., Canty, T. P., Dickerson, R. R., Hanisco, T. F., Wolfe, G. M., Apel, E. C., Atlas, E., Bannan, T., Bauguitte, S., Blake, N. J., Bresch, J. F., Campos, T. L., Carpenter, L. J., Cohen, M. D., Evans, M., Fernandez, R. P., Kahn, B. H., Kinnison, D. E., Hall, S. R., Harris, N. R., Hornbrook, R. S., Lamarque, J.-F., Le Breton, M., Lee, J. D., Percival, C., Pfister, L., Pierce, R. B., Riemer, D. D., Saiz-Lopez, A., Stunder, B. J., Thompson, A. M., Ullmann, K., Vaughan, A., and Weinheimer, A. J.: A Pervasive Role for Biomass
475 Burning in Tropical High Ozone/Low Water Structures, *Nature Communications*, 7, 10 267, <https://doi.org/10.1038/ncomms10267>, 2016.
- Ansmann, A., Wandinger, U., Riebesell, M., Weitkamp, C., and Michaelis, W.: Independent Measurement of Extinction and Backscatter Profiles in Cirrus Clouds by Using a Combined Raman Elastic-Backscatter Lidar, *Applied Optics*, 31, 7113, <https://doi.org/10.1364/AO.31.007113>, 1992.
- Australian Bureau of Meteorology and CSIRO: Climate Variability, Extremes and Change in the Western Tropical Pacific: New Science and
480 Updated Country Reports 2014, Centre for Australian Weather and Climate Research, Collingwood, Vic., 2014.
- Brunamonti, S., Jorge, T., Oelsner, P., Hanumanthu, S., Singh, B. B., Kumar, K. R., Sonbawne, S., Meier, S., Singh, D., Wienhold, F. G., Luo, B. P., Boettcher, M., Poltera, Y., Jauhiainen, H., Kayastha, R., Karmacharya, J., Dirksen, R., Naja, M., Rex, M., Fadnavis, S., and Peter, T.: Balloon-Borne Measurements of Temperature, Water Vapor, Ozone and Aerosol Backscatter on the Southern Slopes of the Himalayas during StratoClim 2016–2017, *Atmospheric Chemistry and Physics*, 18, 15 937–15 957, [https://doi.org/10.5194/acp-18-15937-](https://doi.org/10.5194/acp-18-15937-2018)
485 2018, 2018.
- Cairo, F., De Muro, M., Snels, M., Di Liberto, L., Bucci, S., Legras, B., Kottayil, A., Scoccione, A., and Ghisu, S.: Lidar Observations of Cirrus Clouds in Palau (7°33' N, 134°48' E), *Atmospheric Chemistry and Physics*, 21, 7947–7961, [https://doi.org/10.5194/acp-21-7947-](https://doi.org/10.5194/acp-21-7947-2021)
2021, 2021.
- Crawford, J. H., Davis, D. D., Chen, G., Bradshaw, J., Sandholm, S., Kondo, Y., Merrill, J., Liu, S., Browell, E., Gregory, G., Anderson, B.,
490 Sachse, G., Barrick, J., Blake, D., Talbot, R., and Poeschel, R.: Implications of Large Scale Shifts in Tropospheric NO_x Levels in the Remote Tropical Pacific, *Journal of Geophysical Research: Atmospheres*, 102, 28 447–28 468, <https://doi.org/10.1029/97JD00011>, 1997.
- Crutzen, P. J., Grooss, J. U., Bruhl, C., Muller, R., and Russell, J. M.: A Reevaluation of the Ozone Budget with HALOE UARS Data: No Evidence for the Ozone Deficit, *Science*, 268, 705–708, <https://doi.org/10.1126/science.268.5211.705>, 1995.
- Deshler, T., Stübi, R., Schmidlin, F. J., Mercer, J. L., Smit, H. G. J., Johnson, B. J., Kivi, R., and Nardi, B.: Methods to Homogenize
495 Electrochemical Concentration Cell (ECC) Ozonesonde Measurements across Changes in Sensing Solution Concentration or Ozonesonde Manufacturer, *Atmospheric Measurement Techniques*, 10, 2021–2043, <https://doi.org/10.5194/amt-10-2021-2017>, 2017.
- Di Liberto, T.: Drought in the Pacific (NOAA), <https://www.climate.gov/news-features/event-tracker/drought-pacific>, 2016.
- Diallo, M., Riese, M., Birner, T., Konopka, P., Müller, R., Hegglin, M. I., Santee, M. L., Baldwin, M., Legras, B., and Ploeger, F.: Response of Stratospheric Water Vapor and Ozone to the Unusual Timing of El Niño and the QBO Disruption in 2015–2016, *Atmospheric Chemistry and Physics*, 18, 13 055–13 073, <https://doi.org/10.5194/acp-18-13055-2018>, 2018.
500
- Dirksen, R. J., Sommer, M., Immler, F. J., Hurst, D. F., Kivi, R., and Vömel, H.: Reference Quality Upper-Air Measurements: GRUAN Data Processing for the Vaisala RS92 Radiosonde, *Atmospheric Measurement Techniques*, 7, 4463–4490, [https://doi.org/10.5194/amt-7-4463-](https://doi.org/10.5194/amt-7-4463-2014)
2014, 2014.
- Dirksen, R. J., Bodeker, G. E., Thorne, P. W., Merlone, A., Reale, T., Wang, J., Hurst, D. F., Demoz, B. B., Gardiner, T. D., Ingleby, B.,
505 Sommer, M., von Rohden, C., and Leblanc, T.: Progress in Managing the Transition from the RS92 to the Vaisala RS41 as the Operational Radiosonde within the GCOS Reference Upper-Air Network, 2019.



- Folkins, I.: Tropical Ozone as an Indicator of Deep Convection, *Journal of Geophysical Research*, 107, 4184, <https://doi.org/10.1029/2001JD001178>, 2002.
- Froidevaux, L., Jiang, Y. B., Lambert, A., Livesey, N. J., Read, W. G., Waters, J. W., Browell, E. V., Hair, J. W., Avery, M. A., McGee, T. J., Twigg, L. W., Sunnicht, G. K., Jucks, K. W., Margitan, J. J., Sen, B., Stachnik, R. A., Toon, G. C., Bernath, P. F., Boone, C. D., Walker, K. A., Filipiak, M. J., Harwood, R. S., Fuller, R. A., Manney, G. L., Schwartz, M. J., Daffer, W. H., Drouin, B. J., Cofield, R. E., Cuddy, D. T., Jarnot, R. F., Knosp, B. W., Perun, V. S., Snyder, W. V., Stek, P. C., Thurstans, R. P., and Wagner, P. A.: Validation of Aura Microwave Limb Sounder Stratospheric Ozone Measurements, *Journal of Geophysical Research*, 113, D15S20, <https://doi.org/10.1029/2007JD008771>, 2008.
- 510
- Fueglistaler, S.: Stepwise Changes in Stratospheric Water Vapor?, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2012JD017582>, 2012.
- 515
- Fueglistaler, S., Wernli, H., and Peter, T.: Tropical Troposphere-to-Stratosphere Transport Inferred from Trajectory Calculations, *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004069>, 2004.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical Tropopause Layer, *Reviews of Geophysics*, 47, RG1004, <https://doi.org/10.1029/2008RG000267>, 2009.
- 520
- Gao, R. S., Rosenlof, K. H., Fahey, D. W., Wennberg, P. O., Hints, E. J., and Hanisco, T. F.: OH in the Tropical Upper Troposphere and Its Relationships to Solar Radiation and Reactive Nitrogen, *Journal of Atmospheric Chemistry*, 71, 55–64, <https://doi.org/10.1007/s10874-014-9280-2>, 2014.
- Garfinkel, C. I. and Hartmann, D. L.: Effects of the El Niño–Southern Oscillation and the Quasi-Biennial Oscillation on Polar Temperatures in the Stratosphere, *Journal of Geophysical Research*, 112, D19 112, <https://doi.org/10.1029/2007JD008481>, 2007.
- 525
- Gottelman, A. and Forster, P. d. F.: A Climatology of the Tropical Tropopause Layer., *Journal of the Meteorological Society of Japan*, 80, 911–924, <https://doi.org/10.2151/jmsj.80.911>, 2002.
- Graedel, T., Crutzen, P. J., Kläger, F., and Brühl, C.: *Chemie der Atmosphäre: Bedeutung für Klima und Umwelt*, Spektrum-Lehrbuch, Spektrum Akademischer Verlag, Heidelberg, 1994.
- 530
- Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-Troposphere Exchange, *Reviews of Geophysics*, 33, 403–439, 1995.
- Huang, B., L'Heureux, M., Hu, Z.-Z., and Zhang, H.-M.: Ranking the Strongest ENSO Events While Incorporating SST Uncertainty, *Geophysical Research Letters*, 43, 9165–9172, 2016.
- Hubert, D., Lambert, J.-C., Verhoelst, T., Granville, J., Keppens, A., Baray, J.-L., Bourassa, A. E., Cortesi, U., Degenstein, D. A., Froidevaux, L., Godin-Beekmann, S., Hoppel, K. W., Johnson, B. J., Kyrölä, E., Leblanc, T., Lichtenberg, G., Marchand, M., McElroy, C. T., Murtagh, D., Nakane, H., Portafaix, T., Querel, R., Russell III, J. M., Salvador, J., Smit, H. G. J., Stebel, K., Steinbrecht, W., Strawbridge, K. B., Stübi, R., Swart, D. P. J., Taha, G., Tarasick, D. W., Thompson, A. M., Urban, J., van Gijssel, J. A. E., Van Malderen, R., von der Gathen, P., Walker, K. A., Wolfram, E., and Zawodny, J. M.: Ground-Based Assessment of the Bias and Long-Term Stability of 14 Limb and Occultation Ozone Profile Data Records, *Atmospheric Measurement Techniques*, 9, 2497–2534, [https://doi.org/10.5194/amt-9-2497-](https://doi.org/10.5194/amt-9-2497-2016)
- 540
- 2016, 2016.
- Immler, F., Beninga, I., Ruhe, W., Stein, B., Mielke, B., Rutz, S., Terli, O., and Schrems, O.: A New LIDAR System for the Detection of Cloud and Aerosol Backscatter, Depolarization, Extinction, and Fluorescence, Reviewed and revised papers presented at the 23rd International Laser Radar Conference, 24-28 July 2006 Nara Japan, Editors: Chikao Nagasawa, Nobuo Sugimoto, I, pp. 35–38, 2006.



- Kiladis, G. N., Straub, K. H., Reid, G. C., and Gage, K. S.: Aspects of Interannual and Intraseasonal Variability of the Tropopause and Lower Stratosphere, *Quarterly Journal of the Royal Meteorological Society*, 127, 1961–1983, <https://doi.org/10.1002/qj.49712757606>, 2001.
- 545 Klett, J. D.: Lidar Inversion with Variable Backscatter/Extinction Ratios, *Applied optics*, 24, 1638–1643, 1985.
- Kley, D., Crutzen, P. J., Smit, H. G. J., Vomel, H., Oltmans, S. J., Grassl, H., and Ramanathan, V.: Observations of Near-Zero Ozone Concentrations Over the Convective Pacific: Effects on Air Chemistry, *Science*, 274, 230–233, <https://doi.org/10.1126/science.274.5285.230>, 1996.
- 550 Komhyr, W. D.: Electrical Concentration Cells for Gas Analysis, *Annales Geophysicae*, 25, 203–210, 1969.
- Komhyr, W. D.: Operations Handbook - Ozone Measurements to 40-Km Altitude with Model 4A Electrochemical Concentration Cell (ECC) Ozonesondes (Used with 1680-MHz Radiosondes), Technical Memorandum ERL ARL-149, National Oceanic and Atmospheric Administration (NOAA), Silver Spring, MD, United States, 1986.
- Konopka, P., Ploeger, F., Tao, M., and Riese, M.: Zonally Resolved Impact of ENSO on the Stratospheric Circulation and Water Vapor Entry Values, *Journal of Geophysical Research: Atmospheres*, 121, 11,486–11,501, <https://doi.org/10.1002/2015JD024698>, 2016.
- 555 Krüger, K., Tegtmeier, S., and Rex, M.: Long-Term Climatology of Air Mass Transport through the Tropical Tropopause Layer (TTL) during NH Winter, *Atmospheric Chemistry and Physics*, 8, 813–823, <https://doi.org/10.5194/acp-8-813-2008>, 2008.
- Manney, G. L., Froidevaux, L., Waters, J. W., Zurek, R. W., Read, W. G., Elson, L. S., Kumer, J. B., Mergenthaler, J. L., Roche, A. E., O'Neill, A., Harwood, R. S., Mackenzie, I., and Swinbank, R.: Chemical Depletion of Ozone in the Arctic Lower Stratosphere during Winter 1992–93, *Nature*, 370, 429–434, <https://doi.org/10.1038/370429a0>, 1994.
- 560 Mapes, B. E.: Water's Two Height Scales: The Moist Adiabatic and the Radiative Troposphere, *Quarterly Journal of the Royal Meteorological Society*, 127, 2353–2366, <https://doi.org/10.1002/qj.49712757708>, 2001.
- Miles, W., Grecni, Z., Matsutaro, E. X., Colin, P., Keener, V., and Golbuu, Y.: Climate Change in Palau: Indicators and Considerations for Key Sectors, Tech. rep., Zenodo, <https://doi.org/10.5281/ZENODO.4124259>, 2020.
- 565 Müller, K.: Characterization of Ozone and the Oxidizing Capacity of the Tropical West Pacific Troposphere, Ph.D. thesis, Universität Bremen, <https://doi.org/10.26092/ELIB/463>, 2020.
- Müller, K., Wohltmann, I., von der Gathen, P., and Rex, M.: Air Mass Transport to the Tropical West Pacific Troposphere Inferred from Ozone and Relative Humidity Balloon Observations above Palau, to be submitted, 2023.
- Newell, R. E. and Gould-Stewart, S.: A Stratospheric Fountain?, *Journal of the Atmospheric Sciences*, 38, 2789–2796, 1981.
- 570 Newton, R., Vaughan, G., Ricketts, H. M. A., Pan, L. L., Weinheimer, A. J., and Chemel, C.: Ozonesonde Profiles from the West Pacific Warm Pool: Measurements and Validation, *Atmospheric Chemistry and Physics*, 16, 619–634, <https://doi.org/10.5194/acp-16-619-2016>, 2016.
- Ogino, S.-Y., Fujiwara, M., Shiotani, M., Hasebe, F., Matsumoto, J., T. Hoang, T. H., and T. Nguyen, T. T.: Ozone Variations over the Northern Subtropical Region Revealed by Ozonesonde Observations in Hanoi, *Journal of Geophysical Research: Atmospheres*, 118, 3245–3257, <https://doi.org/10.1002/jgrd.50348>, 2013.
- 575 Oltmans, S.J., Johnson, B.J., Harris, J.M., Thompson, A. M., Liu, H.Y., Chan, C.Y., Vömel, H., Fujimoto, T., Brackett, V.G., Chang, W.L., et al.: Tropospheric Ozone over the North Pacific from Ozonesonde Observations, *Journal of Geophysical Research*, 109, <https://doi.org/10.1029/2003JD003466>, 2004.
- Pan, L. L., Paulik, L. C., Honomichl, S. B., Munchak, L. A., Bian, J., Selkirk, H. B., and Vömel, H.: Identification of the Tropical Tropopause Transition Layer Using the Ozone-Water Vapor Relationship, *Journal of Geophysical Research: Atmospheres*, 119, 3586–3599, <https://doi.org/10.1002/2013JD020558>, 2014.
- 580



- Pan, L. L., Honomichl, S. B., Randel, W. J., Apel, E. C., Atlas, E. L., Beaton, S. P., Bresch, J. F., Hornbrook, R., Kinnison, D. E., Lamarque, J.-F., Saiz-Lopez, A., Salawitch, R. J., and Weinheimer, A. J.: Bimodal Distribution of Free Tropospheric Ozone over the Tropical Western Pacific Revealed by Airborne Observations, *Geophysical Research Letters*, 42, 7844–7851, <https://doi.org/10.1002/2015GL065562>, 2015.
- 585 Pan, L. L., Atlas, E. L., Salawitch, R. J., Honomichl, S. B., Bresch, J. F., Randel, W. J., Apel, E. C., Hornbrook, R. S., Weinheimer, A. J., Anderson, D. C., Andrews, S. J., Baidar, S., Beaton, S. P., Campos, T. L., Carpenter, L. J., Chen, D., Dix, B., Donets, V., Hall, S. R., Hanisco, T. F., Homeyer, C. R., Huey, L. G., Jensen, J. B., Kaser, L., Kinnison, D. E., Koenig, T. K., Lamarque, J.-F., Liu, C., Luo, J., Luo, Z. J., Montzka, D. D., Nicely, J. M., Pierce, R. B., Riemer, D. D., Robinson, T., Romashkin, P., Saiz-Lopez, A., Schauffler, S., Shieh, O., Stell, M. H., Ullmann, K., Vaughan, G., Volkamer, R., and Wolfe, G.: The Convective Transport of Active Species in the Tropics (CONTRAST) Experiment, *Bulletin of the American Meteorological Society*, 98, 106–128, <https://doi.org/10.1175/BAMS-D-14-00272.1>, 2017.
- 590 Paulik, L. C. and Birner, T.: Quantifying the Deep Convective Temperature Signal within the Tropical Tropopause Layer (TTL), *Atmospheric Chemistry and Physics*, 12, 12 183–12 195, <https://doi.org/10.5194/acp-12-12183-2012>, 2012.
- Polhemus, D. A.: Drought in the U.S.-Affiliated Pacific Islands: A Multi-level Assessment, <https://doi.org/10.21429/C9ZS74>, 2017.
- 595 Prather, M. J., Zhu, X., Tang, Q., Hsu, J., and Neu, J. L.: An Atmospheric Chemist in Search of the Tropopause, *Journal of Geophysical Research*, 116, D04 306, <https://doi.org/10.1029/2010JD014939>, 2011.
- Randel, W. J. and Jensen, E. J.: Physical Processes in the Tropical Tropopause Layer and Their Roles in a Changing Climate, *Nature Geoscience*, 6, 169–176, <https://doi.org/10.1038/ngeo1733>, 2013.
- Randel, W. J. and Thompson, A. M.: Interannual Variability and Trends in Tropical Ozone Derived from SAGE II Satellite Data and SHADOZ Ozonesondes, *Journal of Geophysical Research*, 116, D07 303, <https://doi.org/10.1029/2010JD015195>, 2011.
- 600 Randel, W. J., Park, M., Wu, F., and Livesey, N.: A Large Annual Cycle in Ozone above the Tropical Tropopause Linked to the Brewer–Dobson Circulation, *Journal of the Atmospheric Sciences*, 64, 4479–4488, <https://doi.org/10.1175/2007JAS2409.1>, 2007.
- Randel, W. J., Garcia, R. R., Calvo, N., and Marsh, D.: ENSO Influence on Zonal Mean Temperature and Ozone in the Tropical Lower Stratosphere, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL039343>, 2009.
- 605 Rex, M., Wohltmann, I., Ridder, T., Lehmann, R., Rosenlof, K., Wennberg, P., Weisenstein, D., Notholt, J., Krüger, K., Mohr, V., and Tegtmeier, S.: A Tropical West Pacific OH Minimum and Implications for Stratospheric Composition, *Atmospheric Chemistry and Physics*, 14, 4827–4841, <https://doi.org/10.5194/acp-14-4827-2014>, 2014.
- Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact of Uncertainties in Atmospheric Mixing on Simulated UTLS Composition and Related Radiative Effects: IMPACT OF MIXING ON RADIATIVE EFFECTS, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2012JD017751>, 2012.
- 610 Rosen, J. M. and Kjome, N. T.: Backscattersonde: A New Instrument for Atmospheric Aerosol Research, *Applied Optics*, 30, 1552–1561, 1991.
- Schneider, T., Bischoff, T., and Haug, G. H.: Migrations and Dynamics of the Intertropical Convergence Zone, *Nature*, 513, 45–53, <https://doi.org/10.1038/nature13636>, 2014.
- 615 Schoeberl, M. R. and Dessler, A. E.: Dehydration of the Stratosphere, *Atmospheric Chemistry and Physics*, 11, 8433–8446, <https://doi.org/10.5194/acp-11-8433-2011>, 2011.
- Shiotani, M.: Annual, Quasi-Biennial, and El Niño-Southern Oscillation (ENSO) Time-Scale Variations in Equatorial Total Ozone, *Journal of Geophysical Research*, 97, 7625, <https://doi.org/10.1029/92JD00530>, 1992.



- Shonk, J. K. P., Guilyardi, E., Toniazzo, T., Woolnough, S. J., and Stockdale, T.: Identifying Causes of Western Pacific ITCZ Drift in ECMWF
620 System 4 Hindcasts, *Climate Dynamics*, 50, 939–954, <https://doi.org/10.1007/s00382-017-3650-9>, 2018.
- Smit, H. and Thompson, A.: *Ozonesonde Measurement Principles and Best Operational Practices: ASOPOS 2.0 (Assessment of Standard Operating Procedures for Ozonesondes)*, WMO, World Meteorological Organization, GAW Report, p. 173, 2021.
- Smit, H. G. J.: *Quality Assurance and Quality Control for Ozonesonde Measurements in GAW*, GAW Report No. 201, WMO, 2014.
- Smit, H. G. J., Sträter, W., Johnson, B. J., Oltmans, S. J., Davies, J., Tarasick, D. W., Hoegger, B., Stubi, R., Schmidlin, F. J., Northam,
625 T., Thompson, A. M., Witte, J. C., Boyd, I., and Posny, F.: Assessment of the Performance of ECC-ozonesondes under Quasi-Flight Conditions in the Environmental Simulation Chamber: Insights from the Juelich Ozone Sonde Intercomparison Experiment (JOSIE), *Journal of Geophysical Research*, 112, D19 306, <https://doi.org/10.1029/2006JD007308>, 2007.
- Smit, H. G. J., Oltmans, S., Deshler, T., Tarasick, D. W., Johnson, B. J., Schmidlin, F. J., Stuebi, R., and Davies, J.: *SI2N/O3S-DQA Activity: Guide Lines for Homogenization of Ozone Sonde Data*, SPARC-IGACO-IOC Report, 2012.
- 630 Smith, I. N., Moise, A. F., and Colman, R. A.: Large-Scale Circulation Features in the Tropical Western Pacific and Their Representation in Climate Models, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016667>, 2012.
- Solomon, S., Garcia, R. R., Rowland, F. S., and Wuebbles, D. J.: On the Depletion of Antarctic Ozone, *Nature*, 321, 755–758, <https://doi.org/10.1038/321755a0>, 1986.
- Solomon, S., Thompson, D. W. J., Portmann, R. W., Oltmans, S. J., and Thompson, A. M.: On the Distribution and Variability of Ozone in
635 the Tropical Upper Troposphere: Implications for Tropical Deep Convection and Chemical-Dynamical Coupling, *Geophysical Research Letters*, 32, L23 813, <https://doi.org/10.1029/2005GL024323>, 2005.
- Solomon, S., Rosenlof, K. H., Portmann, R. W., Daniel, J. S., Davis, S. M., Sanford, T. J., and Plattner, G.-K.: Contributions of Stratospheric Water Vapor to Decadal Changes in the Rate of Global Warming, *Science*, 327, 1219–1223, <https://doi.org/10.1126/science.1182488>, 2010.
- 640 Sun, B., Reale, T., Schroeder, S., Pettey, M., and Smith, R.: On the Accuracy of Vaisala RS41 versus RS92 Upper-Air Temperature Observations, *Journal of Atmospheric and Oceanic Technology*, 36, 635–653, <https://doi.org/10.1175/JTECH-D-18-0081.1>, 2019.
- Sun, X., Palm, M., Notholt, J., and Müller, K.: Determination of the Chemical Equator from GEOS-Chem Model Simulation: A Focus on the Tropical Western Pacific Region, Preprint, *Dynamics/Atmospheric Modelling/Troposphere/Physics (physical properties and processes)*, <https://doi.org/10.5194/acp-2022-653>, 2022.
- 645 Taguchi, M.: Observed Connection of the Stratospheric Quasi-Biennial Oscillation with El Niño–Southern Oscillation in Radiosonde Data, *Journal of Geophysical Research*, 115, D18 120, <https://doi.org/10.1029/2010JD014325>, 2010.
- Thompson, A. M., Witte, J. C., McPeters, R. D., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V. W. J. H., Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami, S., Toshihiro, O., Johnson, B. J., Vömel, H., and Labow, G. J.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 Tropical Ozone Climatology 1. Comparison with Total Ozone Mapping Spectrometer (TOMS) and
650 Ground-Based Measurements, *Journal of Geophysical Research*, 108, 8238, <https://doi.org/10.1029/2001JD000967>, 2003a.
- Thompson, A. M., Witte, J. C., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V. W. J. H., Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami, S., Ogawa, T., Fortuin, J. P. F., and Kelder, H. M.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 Tropical Ozone Climatology 2. Tropospheric Variability and the Zonal Wave-One, *Journal of Geophysical Research*, 108, 8241, <https://doi.org/10.1029/2002JD002241>, 2003b.
- 655 Thompson, A. M., Witte, J. C., Smit, H. G. J., Oltmans, S. J., Johnson, B. J., Kirchhoff, V. W. J. H., and Schmidlin, F. J.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2004 Tropical Ozone Climatology: 3. Instrumentation, Station-to-Station Variability,



- and Evaluation with Simulated Flight Profiles, *Journal of Geophysical Research*, 112, D03 304, <https://doi.org/10.1029/2005JD007042>, 2007.
- 660 Thompson, A. M., Allen, A. L., Lee, S., Miller, S. K., and Witte, J. C.: Gravity and Rossby Wave Signatures in the Tropical Troposphere and Lower Stratosphere Based on Southern Hemisphere Additional Ozonesondes (SHADOZ), 1998–2007, *Journal of Geophysical Research*, 116, D05 302, <https://doi.org/10.1029/2009JD013429>, 2011.
- 665 Thompson, A. M., Miller, S. K., Tilmes, S., Kollonige, D. W., Witte, J. C., Oltmans, S. J., Johnson, B. J., Fujiwara, M., Schmidlin, F. J., Coetzee, G. J. R., Komala, N., Maata, M., bt Mohamad, M., Nguyo, J., Mutai, C., Ogino, S.-Y., Da Silva, F. R., Leme, N. M. P., Posny, F., Scheele, R., Selkirk, H. B., Shiotani, M., Stübi, R., Levrat, G., Calpini, B., Thouret, V., Tsuruta, H., Canossa, J. V., Vömel, H., Yonemura, S., Diaz, J. A., Tan Thanh, N. T., and Thuy Ha, H. T.: Southern Hemisphere Additional Ozonesondes (SHADOZ) Ozone Climatology (2005–2009): Tropospheric and Tropical Tropopause Layer (TTL) Profiles with Comparisons to OMI-based Ozone Products: SHADOZ OZONE DURING AURA, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2011JD016911>, 2012.
- 670 Thompson, A. M., Witte, J. C., Sterling, C., Jordan, A., Johnson, B. J., Oltmans, S. J., Fujiwara, M., Vömel, H., Allaart, M., PETERS, A., Coetzee, G. J. R., Posny, F., Corrales, E., Diaz, J. A., Félix, C., Komala, N., Lai, N., Ahn Nguyen, H. T., Maata, M., Mani, F., Zainal, Z., Ogino, S.-y., Paredes, F., Penha, T. L. B., Silva, F. R., Sallons-Mitro, S., Selkirk, H. B., Schmidlin, F. J., Stübi, R., and Thiongo, K.: First Reprocessing of Southern Hemisphere Additional Ozonesondes (SHADOZ) Ozone Profiles (1998–2016): 2. Comparisons With Satellites and Ground-Based Instruments, *Journal of Geophysical Research: Atmospheres*, 122, <https://doi.org/10.1002/2017JD027406>, 2017.
- 675 Thompson, A. M., Smit, H. G. J., Witte, J. C., Stauffer, R. M., Johnson, B. J., Morris, G., von der Gathen, P., Van Malderen, R., Davies, J., PETERS, A., Allaart, M., Posny, F., Kivi, R., Cullis, P., Hoang Anh, N. T., Corrales, E., Machinini, T., da Silva, F. R., Paiman, G., Thiong’o, K., Zainal, Z., Brothers, G. B., Wolff, K. R., Nakano, T., Stübi, R., Romanens, G., Coetzee, G. J. R., Diaz, J. A., Mitro, S., Mohamad, M., and Ogino, S.-Y.: Ozone-sonde Quality Assurance: The JOSIE–SHADOZ (2017) Experience, *Bulletin of the American Meteorological Society*, 100, 155–171, <https://doi.org/10.1175/BAMS-D-17-0311.1>, 2019.
- Vaisala: Vaisala Radiosonde RS41 Measurement Performance, Tech. rep., 2017.
- 680 Vigouroux, C., De Mazière, M., Demoulin, P., Servais, C., Hase, F., Blumenstock, T., Kramer, I., Schneider, M., Mellqvist, J., Strandberg, A., Velasco, V., Notholt, J., Sussmann, R., Stremme, W., Rockmann, A., Gardiner, T., Coleman, M., and Woods, P.: Evaluation of Tropospheric and Stratospheric Ozone Trends over Western Europe from Ground-Based FTIR Network Observations, *Atmospheric Chemistry and Physics*, 8, 6865–6886, <https://doi.org/10.5194/acp-8-6865-2008>, 2008.
- Voemel, H. and Diaz, K.: Ozone Sonde Cell Current Measurements and Implications for Observations of Near-Zero Ozone Concentrations in the Tropical Upper Troposphere, *Atmospheric Measurement Techniques*, 3, 495–505, <https://doi.org/10.5194/amt-3-495-2010>, 2010.
- 685 Vömel, H., David, DE., and Smith, K.: Accuracy of Tropospheric and Stratospheric Water Vapor Measurements by the Cryogenic Frost Point Hygrometer: Instrumental Details and Observations, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Vömel, H., Smit, H. G. J., Tarasick, D., Johnson, B., Oltmans, S. J., Selkirk, H., Thompson, A. M., Stauffer, R. M., Witte, J. C., Davies, J., Van Malderen, R., Morris, G. A., Nakano, T., and Stübi, R.: A New Method to Correct the Electrochemical Concentration Cell (ECC) Ozone-sonde Time Response and Its Implications for “Background Current” and Pump Efficiency, *Atmospheric Measurement Techniques*, 690 13, 5667–5680, <https://doi.org/10.5194/amt-13-5667-2020>, 2020.
- Witte, J. C., Thompson, A. M., Smit, H. G. J., Fujiwara, M., Posny, F., Coetzee, G. J. R., Northam, E. T., Johnson, B. J., Sterling, C. W., Mohamad, M., Ogino, S.-Y., Jordan, A., and da Silva, F. R.: First Reprocessing of Southern Hemisphere Additional Ozonesondes (SHADOZ) Profile Records (1998–2015): 1. Methodology and Evaluation: SHADOZ Reprocessed Ozone-sonde Profiles, *Journal of Geophysical Research: Atmospheres*, 122, 6611–6636, <https://doi.org/10.1002/2016JD026403>, 2017.

<https://doi.org/10.5194/egusphere-2023-1023>

Preprint. Discussion started: 12 June 2023

© Author(s) 2023. CC BY 4.0 License.



- 695 Witte, J. C., Thompson, A. M., Smit, H. G. J., Vömel, H., Posny, F., and Stübi, R.: First Reprocessing of Southern Hemisphere ADditional OZonesondes Profile Records: 3. Uncertainty in Ozone Profile and Total Column, *Journal of Geophysical Research: Atmospheres*, 123, 3243–3268, <https://doi.org/10.1002/2017JD027791>, 2018.
- Wolter, K. and Timlin, M. S.: El Niño Southern Oscillation Behaviour since 1871 as Diagnosed in an Extended Multivariate ENSO Index (MEI.Ext), *International Journal of Climatology*, 31, 1074–1087, <https://doi.org/10.1002/joc.2336>, 2011.




Figure 1. Location of Palau (black star), selected SHADOZ sites (black circle, see Tab.1) and CONTRAST campaign domain around the airbase in Guam (grey circle). Longterm Mean Sea Surface Temperatures (SST) in °C derived from monthly ERA5 data from 1959-2021 show the location of the global warm pool area.




Figure 2. The current instrumental setup at the Palau Atmospheric Observatory in two adjacent laboratory containers. The MaxDOAS instrument Pandora2S is not discussed here. The micro-Lidar deployed in 2016 is not shown and was mounted on top of the front container.




Figure 3. Timeline of the measurements performed in Palau using different instruments. Electrochemical Concentration Cell (ECC) ozone sondes have been launched bi-weekly or more frequently during campaigns. The accompanying radiosonde model was switched from Vaisala RS92 to RS41 in October 2017. Cryogenic Frostpoint Hygrometers (CFHs) and Compact Optical Backscatter and Aerosol Lidar Detector (COBALD) sondes were launched together with ECC sondes 13 times during campaigns. The Fourier-transform infrared (FTIR) spectrometer has been deployed with some interruptions. The Italian micro Lidar ran only in spring 2016, while the Compact Cloud Aerosol Lidar (ComCAL) system was installed in 2018.




Figure 4. Time height cross sections of (a) ozone (O_3) volume mixing ratio and (b) relative humidity (RH), (c) Multivariate ENSO index (MEI) from 2016 to 2021. The O_3 and RH measurements have been performed at the PAO. The ENSO Index data is provided by NOAA (<https://psl.noaa.gov/enso/mei/>), (Wolter and Timlin, 2011).




Figure 5. As panels (a), (b) and (c) of Figure 4, but only for the geopotential height range from 14–20 km. The black solid line in (b) and (c) denotes the ozonepause defined as O_3 VMR observations of 90 ppb (linearly interpolated) after Prather et al. (2011). The black horizontal dashed line in (a) at 18 km highlights the a prominent layer of the annual cycle as visualized in Figure 6 (e).




Figure 6. Monthly mean zonal (a) and meridional (b) wind from daily radio soundings from the operational NOAA-associated observing site at the Palau airport (WMO station ID PTRO 91408) averaged from 2016-2019 as well as O₃ VMR (c) (compare Müller et al., 2023), averaged from 08/2016-2021 (i.e. excluding El Niño 2016) as time-height cross-sections, alongside mean O₃ VMR of individual soundings of all years (different markers and colors) in 300 m altitude layers centered around 8 km (d) and 18 km (e). Panel (f) shows the monthly mean latitudinal movement of the ITCZ from TRMM averaged from 1998-2009 for 125-175° E, 0-20° N after (Shonk et al., 2018), whereby the ITCZ location is defined in terms of the zonal mean rainfall rate across this region using a threshold of 50% of the peak zonal-mean rainfall intensity. Palau’s latitude is marked by the horizontal turquoise line.




Figure 7. Free-tropospheric (3–14 km) relation between O_3 and RH in a 2D (hexagonally binned) histogram, normalized to the total count of data points per station, using all measured data pairs with ≤ 100 ppb O_3 VMR for Palau (a) and selected SHADOZ stations (b–h); color shading indicates percentage of total count per gridpoint and marginal plots give individual 1D histograms, normalized to unity.




Figure 8. Boxplots for free-tropospheric (3-14 km) O_3 VMR ≤ 100 ppb for Palau and selected SHADOZ stations, colored by the group categories derived from O_3 /RH distributions (compare Fig. 7): brown for group 1, purple for group 2, turquoise for group 3; outliers are not shown and whiskers are a function of the interquartile range ($IQR = Q3 - Q1$, i.e. $Q3$ or $Q1 +$ or $- 1.5 \cdot IQR$ respectively); blue squares are mean values; orange circles refer to the most frequent O_3 VMR as illustrated in the marginal 1-D histograms in Fig. 7; arrangement of boxplots on the horizontal axis is an approximation to longitudinal positions.




Figure 9. Relative frequency distribution (normalized to the layer maximum) of tropospheric (0-15 km) O₃ VMR with altitude (100 m bins) for all observations (a) and “humid” observations with RH greater 45 % (b) for Palau from 2016-2021, (c) and (d) histograms show relative frequency distributions of O₃ VMR for a layer between 320 and 340 K potential temperature for all (c) and only “wet” (d) observations, respectively; black horizontal lines in (a, b) indicate the approximate location of the boundaries for the potential temperature based selection of the data in (c, d) (compare Fig. 3 in Pan et al., 2015).



Site	Location [°lat, lon]	Time Record mm/yyyy	Distance to Palau [km]
Java	7.5S, 112.6E	01/1998-10/2013	3100
Hanoi	21N, 106E	09/2004-12/2020	3400
Kuala Lumpur	2.7N, 102E	01/1998-12/2021	3700
Fiji	18S, 178 E	01/1998-12/2021	5600
American Samoa	14S, 171W	01/1998-12/2021	6500
Hilo	19N, 157 W	01/1998-12/2021	7400
Costa Rica	10N, 84W	07/2005-12/2021	15400

Table 1. SHADOZ stations and respective time record used in this paper, with distance as the crow flies to Palau; see Fig. 1, compare e.g. Thompson et al., 2012.




Figure A1. All measured individual O₃ VMR profiles sorted by month and year (colored markers), the monthly mean (solid black line) excluding El Niño 2016, i.e. starting 08/2016, and monthly statistics (dashed line for mean between 5 and 10 km, dotted line for mean of minimum values per sounding between 5 and 10 km); number (#) of profiles per year given in brackets.



Specification	PAO	Comments
Ozonsonde Type	SPC, model 6A	
Radiosonde Type	Vaisala RS92-SGP and RS41-SGP	change in October 2017
Sensing Solution Type	1.0 % KI, full buffer	
Cathode Solution Volume	3ml	
Background Current	all recorded	I_{b2} used for correction
Temperature Pump Location	internal	in Teflon block of pump
Pumpflow Measurement	bubble flow meter	
Source of Zero Ozone	Vaisala chemical destruction filter or self-built charcoal/ desiccant filter	dry storage ensured
Laboratory Conditions	pressure, temperature and relative humidity recorded	air-conditioned and de-humidified laboratory space
Software Data Reduction	retrieval with commercial Vaisala software package	own calculations from raw cell current used in post-processing and data analysis
Corrections applied for		
Background Current	yes	pressure-dependent, see Sect.2.1
Pump Efficiency	yes	according to Komhyr (1986)
Humidity Effect in Flow Rate	no	contribution was found to be -1 to -2 % or -1 ppb in the free troposphere
Total Column Ozone	no	

Table A1. Major specifications and Standard Operating Procedures (SOP) of O₃ soundings at the Palau Atmospheric Observatory (PAO), in accordance with recommendations from Smit et al. (2012) and Smit (2014), except for the applied pressure-dependent background current correction.