



Detecting nitrogen oxide emissions in Qatar and quantifying emission factors of gas-fired power plants - A four-years study

Anthony Rey-Pommier^{1,2}, Frédéric Chevallier¹, Philippe Ciaï^{1,2}, Jonilda Kushta², Theodoros Christoudias²,
I. Safak Bayram³ and Jean Sciare².

¹ Laboratoire des Sciences du Climat et de l'Environnement, LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, 91198 Gif-sur-Yvette, France

² The Cyprus Institute, Climate and Atmosphere Research Center, 2121 Nicosia, Cyprus

³ Department of Electronic and Electrical Engineering, University of Strathclyde, Glasgow G1 1XW, United Kingdom

Correspondence: Anthony Rey-Pommier (anthony.rey-pommier@lsce.ipsl.fr)

Abstract. Nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$), produced in urban areas and industrial facilities (particularly in fossil fuel-fired power plants), are major sources of air pollutants, with implications for human health, leading local and national authorities to estimate their emissions using inventories. In Qatar, these inventories are not systematically updated, while the country is experiencing fast economic growth. Here, we use spaceborne retrievals of nitrogen dioxide (NO_2) columns at high spatial resolution from the TROPOMI instrument to estimate NO_x emissions in Qatar from 2019 to 2022 with a flux-divergence scheme, according to which emissions are calculated as the sum of a transport term and a sink term representing the three-body reaction comprising NO_2 and hydroxyl radical (OH). Our results highlight emissions from gas power plants in the north-east of the country, and from the urban area of the capital Doha. The emissions from cement plants in the west and different industrial facilities in the south-east are under-estimated, due to frequent low-quality measurements of NO_2 columns in these areas. Our top-down model estimates a weekly cycle with lower emissions on Fridays compared to the rest of the week, which is consistent with social norms in the country, and an annual cycle with mean emissions of 9.56 kt per month for the four-year period. These monthly emissions differ from CAMS-GLOB-ANT_v5.3 and EDGARv6.1 global inventories, for which the annual cycle is less marked and the average emissions are respectively 1.44 and 1.68 times higher. Our emission estimates are correlated with local electricity generation, and allow to infer a mean NO_x emission factor of $0.557 \text{ t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$ for the three gas power plants in the Ras Laffan area.

1 Introduction

Nitrogen oxides are reactive trace gases that can be converted into other chemical species, including ozone and fine particulate matter. Emissions of NO_x can originate from natural sources (from fires, lightning and soils), but the majority originates from anthropogenic sources, such as vehicle engines and heavy industrial facilities like power plants, steel mills and cement kilns (Vuuren, 2011). High levels of NO_x in the troposphere contribute to the formation of acid rain and smog. They also have a significant effect on human health by causing various respiratory diseases (Bovensmann et al., 1999; Burnett et al., 2004; EPA, 2016; He et al., 2020). To limit those impacts, national and regional governments generally enact a series of air pollution control strategies, which generally take the form of proscriptions on certain polluting technologies, with the aim of reducing the concentration of pollutants at the local level to targets that must be reached within a given timeframe. In the Middle East region, such mitigation strategies are quite recent. The region is thus experiencing increasing levels of NO_x pollution from anthropogenic sources (Osipov et al., 2022), with levels that remain high by international standards (Lelieveld et al., 2015). During the last two decades, Qatar experienced a rapid development based on oil and gas, leading to a degradation of air quality (Mansouri Daneshvar, Hussein Abadi, 2017). However, no official report concerning emissions of pollutants has been publicly available in the last 12 years.

Spectrally resolved satellite measurements of solar backscattered UV-Visible radiation enable the quantification of NO_2 in the atmosphere. Such measurements have been providing information on the spatial distribution of tropospheric NO_2 for more than 20 years, allowing the identification of many NO_x sources globally (Richter, Burrows, 2002; Celarier et al., 2008). In October 2017, the Sentinel-5 Precursor satellite was launched. Its main instrument is the TROPospheric Monitoring



45 Instrument (TROPOMI), which provides daily tropospheric NO_2 column densities at high spatial resolution with a large swath width. Columns images can then be used to retrieve NO_x emissions with the use of the continuity equation in steady state. This scheme, known as flux-divergence, requires the use of other physical quantities. In this study, we use this method to quantify the emissions in Qatar based on retrievals from 2019 to 2022 and overcome the absence of recent reporting.

This article is organised as follows: Section 2 provides a description of Qatar and its main emission sources. Section 3 provides a description of the datasets used in this study. Section 4 presents the flux-divergence method to infer NO_x emissions at the scale of the country. Section 5 presents the spatial distribution of NO_x emissions and their seasonal variations. It also presents the limitation of our method to infer emissions for areas above which TROPOMI measurements are often of low quality. Section 6 confronts our estimated emissions to local electricity generation and existing global inventories to provide a sectoral approach of NO_x emissions. Section 7 presents the limits and the uncertainties of the model, and Section 8 our 55 concluding remarks. For the purposes of our study, NO_x emissions are expressed as NO_2 throughout this article.

2 General features of Qatar and overview of reported NO_x emissions in 2007

Qatar is a country with an area of $11,600 \text{ km}^2$, located on the northeast coast of the Arabian Peninsula. The country shares its sole terrestrial border with Saudi Arabia in the south, and has maritime boundaries with Bahrain in the west, Iran in the north and the United Arab Emirates in the east. Qatar has the third largest proven natural gas reserves in the world and non-negligible oil reserves (EIA, 2021). Since 1973, oil and gas revenues increased dramatically, making the country the third largest exporter of natural gas (OPEC, 2020). The resulting economic growth raised Qatar to one of the countries with the highest per capita incomes in the world. Between 2001 and 2019, the average GDP growth rate of the country was 9.1 % (World Bank, 2022), driven by the exploitation of the oil and gas fields, which account for 85 % of its exports and over 60 % of its gross domestic product. Because the incomplete combustion of hydrocarbons produces NO_x , the exploitation of such 65 oil and gas resources is a source of air pollution: as a consequence, the transport sector is a source of emissions, as well as the power sector, which is dominated in Qatar by gas power plants. Other sectors, such as cement production, also contribute to the total NO_x budget. Little information is available on the country's emissions and the share of these different sectors. The last official communication dates back to 2011 for emissions in 2007, and uses the IPCC Common Reporting Format sector classification (UNFCCC, 2011). It estimates total emissions of 163 kt (assumed to be expressed as NO_2), with the following 70 shares:

- **Power generation:** There are several gas-fired power plants in Qatar which, until the end of 2022, provided all the electricity in the country, with a generation that increased from 4.5 TWh in 1990 to 48 TWh in 2021 (EIA, 2022). According to the report for emissions in 2007, they produced 32.03 kt of NO_x , which corresponded to 19.7 % of the country's total emissions. The report indicates an emission factor equal to $0.109 \text{ t}_{\text{NO}_x} \cdot \text{TJ}^{-1}$ or $0.392 \text{ t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$, which is a low but realistic emission factor for an electric mix dominated by gas. More recent data on NO_x emissions from the power sector are not available.
- **Cement production:** The country has three cement production sites. According to the report for emissions in 2007, cement production was responsible for 23.4 % of the country's total NO_x emissions. It should be noted that due to the high temperatures in summer, outdoor activities at the productions sites are reduced, which can result in lower activity 80 between June and September.
- **Road transport:** According to the report for emissions in 2007, the road transport sector accounted for 22.7 % of the country's total NO_x emissions (27.6 % for the total transport sector). Fuel sales do not seem to have any annual seasonality (Al-Attiyah Foundation, 2018): the corresponding NO_x emissions are therefore not variable with seasons. However, they may change gradually with the nature of the vehicle fleet, which includes a growing number of diesel 85 vehicles.
- **Manufacture of Solid Fuels and Other Energy Industries:** This sector includes upstream oil and gas activities and processing operations. Most of historical upstream operations took place in the Dukhan field, located in the west of the country, but the majority of the extraction currently takes place offshore in the North Field, which is one of the largest non-associated gas fields in the world. According to the report for emissions in 2007, these activities accounted for 25.7 % of the country's NO_x emissions. 90

Figure 1 shows the locations of the industrial facilities in Qatar which are likely to emit significant amounts of NO_x . Four regions can be distinguished. The west of the country comprises two large-capacity cement plants ($7.0 \text{ Mt} \cdot \text{yr}^{-1}$ and $5.0 \text{ Mt} \cdot \text{yr}^{-1}$). The south-east includes a gas power plant, a cement plant and an aluminium smelter, while the north-east comprises the industrial area of Ras Laffan, including three gas power plants. Finally, the urban area of Doha, in the 95 centre-east of the country, concentrates the majority of the population, as well as five gas power plants.




Figure 1: Location of the main industrial facilities (cement kilns, power plants and aluminium smelters) in Qatar. Urban areas are displayed in gray and Doha's city centre is denoted with a star. Industrial facilities and cities in other countries are not displayed. Map tiles by Stamen Design under CC BY 3.0. Data © OpenStreetMap contributors.

3 Instruments and data

3.1 TROPOMI NO₂ retrievals

NO₂ can be observed from space with satellite instruments based on its strong absorption features in the 400–465 nm wavelength region (Vandaele et al., 1998). By comparing observed spectra with a reference spectrum, the amount of NO₂ in a portion of the atmosphere between the instrument and the surface can be derived. The TROPOspheric Monitoring Instrument (TROPOMI), onboard the European Space Agency's (ESA) Sentinel-5 Precursor (S-5P) satellite, is one of those instruments, providing daily measurements of NO₂ around 13:30 local time (LT). Its high spatial resolution (originally 3.5×7 km² at nadir, improved to 3.5×5.5 km² since 6 August 2019) allows to observe some of the fine-scale structure of NO₂ pollution, such as within cities (Beirle et al., 2019; Demetillo et al., 2020), or near power plants and industrial facilities (Shikwambana et al., 2020; Saw et al., 2021). Tropospheric vertical column densities (VCDs, or simply "columns") are provided by the algorithm after measurement by the instrument, which represents the vertically integrated number of NO₂ molecules per surface unit between the surface and the tropopause. An algorithm also provides an air mass factor (AMF), which is used to convert slant column densities into vertical column densities. This factor depends on many parameters, including the albedo of the viewed surface, the vertical distribution of the absorber and the viewing geometry. It is a source of structural uncertainty in NO₂ measurements (Boersma et al., 2004; Lorente et al., 2019), which becomes non-negligible in polluted environments. The large swath width of the instrument (~2600 km) makes it possible to construct NO₂ images of VCDs on large spatial scales. We use TROPOMI NO₂ retrievals from 2019 to 2022 (S5P-PAL reprocessed data with processor version 2.3.1 from January 2019 to October 2021, OFFL stream with processor version 2.3.1 from November 2021 to October 2022, and OFFL stream with processor version 2.4.0 from November 2022 to December 2022) over Qatar. The arid climate of the country, which offers a large number of clear-sky days throughout the year, enables the calculation of monthly averages based on multiple observations. Its intensive anthropogenic activity, concentrated on a small number of areas, allows to observe high NO₂ concentration patterns. Due to its short lifetime (about 1 to 10 hours), background NO₂ levels can be orders of magnitude lower than levels near polluted areas. Pollution patterns are therefore characterised by large signal-to-noise ratios above main emitters. Finally, TROPOMI products provide a quality assurance value q_a , which ranges from 0 (no data) to 1 (high-quality data). For our analysis of concentrations, we selected NO₂ retrievals with q_a values greater than 0.75, which systematically correspond to clear-sky conditions (Eskes et al., 2022), and gridded these retrievals at a spatial resolution of 0.0625°×0.0625°. From August 2019, this resolution is lower than that of the instrument, thus providing a grid for which NO₂ VCDs correspond to one or more measurements. The observed plumes remain correctly resolved.




Figure 2: TROPOMI observation of NO_2 column densities above Qatar on 12 December 2019. White pixels correspond to areas with low-quality data ($q_a < 0.75$) or no data.

The domain of interest extends between the meridians 50°E and 52°E and between the parallels 24°N and 27°N . TROPOMI tropospheric columns show polluted areas, such as the Ras Laffan area, but also the large urban area of Doha. Observed VCDs are high (5 to 30×10^{15} molecules. cm^{-2}), which facilitates the observation of NO_2 plumes. An example of a TROPOMI image is provided in Figure 2. Within the domain, the meridional component of the wind is generally oriented towards the south. The zonal component varies throughout the year. Strong wind events can cause the pollution from Bahrain to reach the coasts of Qatar. Finally, several areas in the west of the country are not visible by TROPOMI most of the time, because the quality of the corresponding measurements is too low. This phenomenon, which is also present in the east of the country (slightly south of downtown Doha) but with a smaller spatial extent, is studied in details in Section 5.3.

3.2 ERA5

The horizontal wind vector field $\mathbf{w} = (u, v)$ is taken from the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 data archive (fifth generation of atmospheric reanalyses) at a horizontal resolution of $0.25^\circ \times 0.25^\circ$ on 37 pressure levels (Hersbach et al., 2020). The hourly values have been linearly interpolated to the TROPOMI orbit timestamp and re-gridded to a $0.0625^\circ \times 0.0625^\circ$ resolution.

3.3 CAMS real-time fields

The Copernicus Atmospheric Monitoring Service (CAMS) global near-real-time service provides analyses and forecasts for reactive gases, greenhouse gases and aerosols. Data is gridded on 25 vertical pressure levels with a horizontal resolution of $0.4^\circ \times 0.4^\circ$ and a temporal resolution of 3 hours (Huijnen et al., 2019). Here, CAMS concentration fields of hydroxyl radical (OH) are used to calculate NO_2 sinks from TROPOMI observations. We also use CAMS temperature field T to account for the reaction rate for HNO_3 production through the calculation of the NO_2 lifetime according to Burkholder et al. (2020). The hourly values are also linearly interpolated to the TROPOMI orbit timestamp and re-gridded to a $0.0625^\circ \times 0.0625^\circ$ resolution.

3.4 Inventories

The Emissions Database for Global Atmospheric Research (EDGARv6.1) and the CAMS global anthropogenic emissions (CAMS-GLOB-ANT_v5.3) are global inventories that provide $0.1^\circ \times 0.1^\circ$ gridded emissions for different sectors on a monthly basis. EDGARv6.1 emissions are based on activity data (population, energy production, fossil fuel extraction, industrial processes, agricultural statistics, etc.) derived from the International Energy Agency (IEA) and the Food and Agriculture Organization (FAO), corresponding emission factors, national and regional information on technology mix data and end-of-pipe measurements. The inventory covers the years 1970-2018. CAMS-GLOB-ANT_v5.3 is developed within the framework of the Copernicus Atmospheric Monitoring Service (Granier et al., 2019). For this inventory, NO_x emissions are based on



various sectors in the EDGARv5.0 emissions up to 2015 which are extrapolated to 2021 using sectorial trends from the Community Emissions Data System (CEDS) inventory (Hoesly et al., 2018) up to 2019. From one inventory to another, the names and definitions of the sectors may differ. In EDGARv6.1 and CAMS-GLOB-ANT_v5.3, the emissions for a given country are derived from the type of technologies used, the dependence of emission factors on fuel type, combustion conditions, as well as activity data and low-resolution emission factors (Janssens-Maenhout et al., 2019).

3.5 Electricity consumption and production data

As the power sector is one of the main drivers of NO_x , we use electricity production and consumption data at several time scales, detailed in Section 5.5. Daily load profiles from February 2016 to January 2017 are taken from Bayram et al. (2018) and used to calculate monthly ratios between the average power demand and the power demand during the overpass of TROPOMI. These ratios are assumed to be valid from 2019 to 2022. We also use monthly electricity generation time series from 2019 to 2022, which are provided by the Qatar Ministry of Development Planning and Statistics (PSA, 2023). From 2019 to 2021, these time series can be completed with monthly reports from Kahramaa, which transmits and distributes the electricity generated by each of the country's power plants with great accuracy (Kahramaa, 2023). The corresponding data for 2022 is not available yet.

4 Method

4.1 Mask and background removal

In satellite retrievals, the NO_2 signal from a sparsely populated area or a small industrial facility may be covered by noise or by the signal generated by natural NO_x emissions. As a consequence, detecting traces of non-natural emissions in TROPOMI NO_2 images is not a straightforward process. In the absence of anthropogenic sources, the NO_2 columns that are observed constitute a tropospheric background that varies between 0.2 and 1.0×10^{15} molecules. cm^{-2} . At the global scale, this background is mostly due to soil emissions in the lower troposphere (Yienger, Levy, 1995; Hoelzemann et al., 2004). In the upper troposphere, NO_2 sources include lightning, convective injection and downwelling from the stratosphere (Ehhalt et al., 1992), but the factors controlling the resulting concentrations are poorly understood. According to state-of-art estimates, anthropogenic NO_x accounts for most of the emissions at the global scale, whereas natural emissions from fires, soils and lightning are smaller (Jaeglé et al., 2005; Müller, Stavrakou, 2005; Lin, 2012). In Qatar, the desert climate limits lightning and fire, and the share of the corresponding emissions within the background is thus expected to be low. As a first step, we estimate this background by excluding the part of the domain which is outside the territory of Qatar, as well as other regions with human activity, which include Bahrain and the neighbouring part of Saudi Arabia. The remaining pixels constitute an "external mask" within which we calculate the 5th percentile of the TROPOMI columns. The corresponding value is defined as the tropospheric background. Figure 3 displays the external mask, as well as an "internal mask" that gathers only pixels above Qatar's territory. This external mask is used in Section 5 and Section 6.




Figure 3: Masks used for NO_x emissions estimates: Green cells are used to estimate the NO_2 tropospheric background that is removed from vertical column densities before calculation of emissions (external mask). Blue cells are used to count NO_x emissions attributable to Qatar (internal mask). Orange cells, which cover urban and industrial areas from other countries, are not considered.



4.2 Flux-divergence method

185 As a second step, we derive top-down NO₂ production maps with the flux-divergence method, which consists in applying the continuity equation in steady state as:

$$e_{\text{NO}_2} = \text{div}(\Omega_{\text{NO}_2} \mathbf{w}) + \Omega_{\text{NO}_2} / \tau \quad (1)$$

The previous equation highlights a transport term $D = \text{div}(\Omega_{\text{NO}_2} \mathbf{w})$, obtained by multiplying NO₂ VCDs with horizontal wind speeds and the NO₂ sinks $S = \Omega_{\text{NO}_2} / \tau$. As the local overpass time of TROPOMI is close to 13:30 LT, sinks are dominated by the chemical loss due to the reactions of NO₂ with OH (leading to formation of HNO₃), which can be described
190 by a first-order time constant τ as:

$$\tau = \frac{1}{k_{\text{mean}}(T, [M]) \cdot [\text{OH}]} \quad (2)$$

Here k_{mean} is the reaction rate that characterises the different reactions between NO₂ and OH. Burkholder et al. (2020) provide a general expression of this reaction rate with respect to atmospheric conditions (temperature T and total air concentration $[M]$).

In the atmosphere, OH is a dominant oxidant species. It is mainly produced during daylight hours by interaction between
195 water and atomic oxygen produced by ozone dissociation (Levy, 1971), and its concentration is therefore strongly correlated with solar ultraviolet radiation (Rohrer, Berresheim, 2006). The direct measurement of OH is possible using spectroscopic methods but the spatial representativeness of the data is limited due to its short lifetime. Most global analyses thus estimate OH budgets from other variable species (Li et al., 2018; Wolfe et al., 2019) with high associated uncertainties (Huijnen et al., 2019). In polluted air, another mechanism for OH production is the reaction between NO and HO₂. This reaction, referred
200 to as the NO_x recycling mechanism, illustrates the nonlinear dependence of the OH concentration on NO₂ (Valin et al., 2011; Lelieveld et al., 2016). Here, considering OH as the only sink assumes other sinks are negligible. We consider such an approximation as valid (Rey-Pommier et al., 2022). This hypothesis can be validated a posteriori by analysing the calculated NO_x emission maps and verifying the absence of highly negative emissions (which would correspond, all other things being equal, to an underestimation of the sink term) or emissions having the shape of a plume (which would correspond, all other
205 things being equal, to an overestimation of the sink term).

Finally, it should be noted that anthropogenic combustion activities produce mainly NO, which is transformed into NO₂ by reaction with ozone O₃. NO₂ is then photolyzed during the day, reforming NO (Seinfeld, 1989). This photochemical equilibrium between NO and NO₂ can be highlighted with the concentration ratio $L = [\text{NO}_x] / [\text{NO}_2]$. NO_x emissions are therefore obtained by multiplying NO₂ production e_{NO_2} by L . As diurnal NO concentrations in urban areas are generally
210 above 20 ppb, the characteristic stabilization time of this ratio never exceeds a few minutes (Graedel et al., 1976; Seinfeld and Pandis, 2006). This time being lower than the order of magnitude of the inter-mesh transport time (about 30 min considering the resolution used and the mean wind module in the region), we can reasonably neglect the effect of the stabilization time of the conversion factor on the total composition of the emissions and treat each cell of the grid independently from its direct environment. In Rey-Pommier et al. (2022), this ratio was estimated using the CAMS NO and NO₂ concentration fields in
215 Egypt. For Qatar, CAMS data show many outliers for these concentration fields: for some pixels, NO concentrations can be equal to zero. They can be also two times higher than NO₂ concentrations in places without any significant feature that would explain it. We therefore do not use these fields and choose a fixed value of 1.32 for this ratio, as used by Beirle et al. (2019). After removal of outliers in the statistics, CAMS values for L suggest that the average value for this ratio ranges between 1.17 and 1.60 with small spatial variations.

220 5 Results

5.1 Selection of TROPOMI images and averages

We apply Equation 1 to daily TROPOMI images and average the resulting emissions to obtain monthly daytime emissions. This process can be hindered due to the presence of poor quality measurements that prevent the calculation of the divergence in the transport term each day. This happens when a significant part of the country is covered by clouds. Furthermore, the
225 resolution used to grid data (0.0625° × 0.0625°, i.e. about 6.3 × 6.9 km² in this region) is lower than the initial resolution of TROPOMI (3.5 × 7 km² until 6 August 2019). For the first eight months of this study, our gridding therefore results in maps having many pixels without data. In the process of estimating mean monthly emissions, we consider these two situations and do not take into account in the averaging the days for which corresponding TROPOMI images have more than 70 % of the territory of Qatar (defined by the internal mask) without data or with poor quality data ($q_a < 0.75$). Furthermore, Bahrain is located 30 km away from Qatar, and the centres of the capitals of the two countries are separated by about 130 km.
230



Pollution from Bahrain, which is usually transported to Qatar, can reach Doha during strong wind events. In such situations, the errors in ERA5 can alter the estimation of the transport term and thus the NO_x estimates. As a consequence, we also remove days with high wind speeds in the Bahrain/Qatar direction, i.e. days for which the average wind over Bahrain and the marine area between the two countries has a speed higher than 30 km.h^{-1} and an angle between -15° ($E\frac{1}{4}SE$) and -75° ($S\frac{1}{4}SE$). Figure 4 shows the amount of days considered in the calculation of the average emissions of a given month between 2019 and 2022. It also shows the number of days that have been discarded and the reasons for the corresponding discards.




Figure 4: Number of days involved in the mean monthly estimates of NO_x emissions (blue) and days that have been discarded due to strong winds blowing from Bahrain to the south-east (brown), large areas within the internal mask with low-quality data (yellow), or both (red).

Here, monthly average emissions are calculated with between 8 and 25 days of TROPOMI images, with low values for the seven first months of the study due to the lower resolution of TROPOMI during this period. Excluding these months, averages are calculated with 17.9 days. Cases for which too many pixels are not observed by TROPOMI in the internal mask occur more frequently than cases with strong winds from Bahrain. Overall, we consider most of the inferred monthly averages to be robust. However, at the small scale, there are a small number of pixels with a constant absence of reliable VCD estimations due to low-quality measurements, resulting with pixels having estimations of NO_x with a poor reliability. The treatment of such pixels is precised in Section 5.3.

5.2 Spatial distribution of NO_x emissions

We use the flux-divergence model to obtain NO_x emission maps within our domain. For each applicable day (as described in Section 5.1), a background is calculated using the external mask and daily emissions are estimated. These daily emissions are then averaged to obtain a representation of the emissions at 13:30 LT on a monthly scale. For October 2022, a maximum of 25 daily emissions have been average to calculate the corresponding mean monthly emissions map, which is displayed in Figure 5. When the transport term is integrated over large spatial scales, it cancels out due to the mass balance in the continuity equation between NO_2 sources and NO_2 sinks. Although the sink term is responsible for most of the total NO_x budget within the domain, the transport term can reach high values at small scale, highlighting hotspots where emissions are concentrated. Such hotspots comprise gas power plants in the north-eastern part of the country and industrial facilities in the south-east. High emissions are also observed in Doha, but in that case, the sink term accounts for most of the NO_x budget due to the spread of sources in the urban area. The west part of the country, where two cement plants are located, also shows significant emissions. High emissions outside of Qatar are observed in the urban areas of Manama (Bahrain) and Dammam (Saudi Arabia). In the south-eastern corner of the domain, an area of high emissions also seems to stand out from the desert. This area corresponds to a cross-border centre between Saudi Arabia and the United Arab Emirates. It is possible that an important road traffic there is responsible for such emissions, but we have no information to support this hypothesis. Finally, we observe high emissions at sea, up to about 30 km from the east coast of the country. These emissions, of about $2.0 \times 10^{15} \text{ molecules.cm}^{-2}.\text{h}^{-1}$, cannot be solely attributed to shipping activity (Rey-Pommier et al., 2022). Because in this region, the centre of the $40 \text{ km} \times 44 \text{ km}$ CAMS pixel is close to the coast, the NO_2 lifetime is probably under-estimated. The rest of the domain has relatively low emissions, which is consistent with the absence of major sources of NO_x .




Figure 5: Mean NO_x emissions above Qatar (13:30 LT): transport term (a), sink term (b), and resulting emissions (c) for October 2022. Power plants are denoted with dots, cement plants with squares, and Doha's city centre with a star.




Figure 6: Mean NO_x emissions above Qatar (13:30 LT): transport term (a), sink term (b), and resulting emissions (c) for the 2019-2022 period. Power plants are denoted with dots, cement plants with squares, and Doha's city centre with a star.

Nevertheless, it can be noted that inferred emissions in the domain remain relatively noisy. From monthly maps, we calculate average emissions at 13:30 for the period 2019-2022, which are displayed on Figure 6. The hotspots identified earlier remain visible with a improved signal-to-noise ratio. In particular, emissions in the north-east region are superior to 3.0×10^{15} molecules. cm^{-2} for only 7 pixels, which are among the pixels closest to the power plants in the region. In the eastern part of the country, NO_x emissions reflect the main directions of urban expansion in Doha (from the coast towards the west and the southwest), with emissions ranging from 2×10^{15} molecules. $\text{cm}^{-2} \cdot \text{h}^{-1}$ to 10×10^{15} molecules. $\text{cm}^{-2} \cdot \text{h}^{-1}$. The desert areas display very low emissions (about 0.2×10^{15} molecules. $\text{cm}^{-2} \cdot \text{h}^{-1}$), indicating a slight overestimation of the average sink term or an underestimation of the NO_2 background. Emissions at sea remain abnormally high on the east coast, but the effect is lower than what is observed on Figure 5. However, unlike for October 2022, for which the emissions from cement plants could be identified, we observe low emissions in the western part of the country.

5.3 Availability of TROPOMI data above industrial sites

High-quality measurements by TROPOMI are infrequent above the western and south-eastern part of Qatar, with pixels regularly corresponding to measurements with $q_a < 0.75$. For these pixels, the inferred emissions are only calculated with an



average over a very small number of measurements. In many cases, no measurements are available, and emissions can not be attributed to the pixels. Figure 7 shows the mean TROPOMI coverage over the domain for year 2020 counted as fraction of days with $q_a < 0.75$ within the year. In addition to the west and south-east (corresponding to the southern part of Doha) already identified, the north-east is also affected, but to a lesser extent. This situation is the same for years 2019 and 2021, and the identified areas have a fraction of observable days lower than 40 %. The situation is different for year 2022, during which the west has a fraction of about 70 % and the south-east 50 % (see Supplementary Material). The effect on total inferred emissions is not negligible, since pixels that are concerned include intensive emitters, such as an industrial area in the south-east and the two large cement plants in the west. For instance, during the 36 months of the period 2019-2021, inferred emissions above these cement plants are lower than 1.5×10^{15} molecules.cm⁻².h⁻¹ for 8 months and greater than this value for 8 months. Emissions could not be inferred for at least one of the plants for the remaining 20 months. Conversely, emissions could be inferred for 10 months out of 12 in 2022, and they were higher than 1.5×10^{15} molecules.cm⁻².h⁻¹ for 6 of those months. In the TROPOMI products, the value of q_a includes automated quality assurance parameters related to different algorithms concerning the presence of clouds, aerosol particles, pressure levels and other physical quantities. For situations with $q_a < 0.75$, retrievals are only sensitive to the NO₂ concentrations above the clouds and will depend on model assumptions for the missing part. They still contain useful information, but this information should be carefully interpreted. Here, all areas that are affected by these low values are located near the coasts. These areas are not characterised by persistent clouds, but they are all located within or close to urban or industrial centres, in which aerosol emissions can be high. However, other regions in Qatar and the neighbouring countries are characterised by high emissions of aerosols without q_a values being particularly high there. For instance, many studies have been providing estimates of NO_x emissions in Riyadh (Beirle et al., 2019) without indicating anything similar. We observe that these high frequencies of low q_a values are found in several urban and industrial areas on both coasts of the Persian Gulf (especially in Saudi Arabia, United Arab Emirates and Iran). This suggests that the identified persistent low values for q_a come from the calculation of one or several quality assurance parameters. Because the areas displaying low values usually have a rectangular shape, it also suggests a low resolution for these parameters.




Figure 7: TROPOMI observation density for 2020.

Here, we observe that the use of the OFFL stream (version 2.3.1) from November 2021 to October 2022 generates less pixels for which the value of q_a falls below 0.75 than the use of the S5P-PAL data from January 2019 to October 2021. Furthermore, the last two months of 2022, calculated with the version 2.4.0 of the NO₂ product (operational since July 2022), do not show any area with particularly low quality flag values (see Supplementary Material). The latest version of the user manual indicates that in this last version, the surface albedo climatology (SAC) used in the NO₂ fitting window and derived from OMI and GOME-2 was replaced by a SAC derived from TROPOMI observations (Eskes et al., 2022). This new TROPOMI SAC is consistently applied in the cloud fraction, the cloud pressure retrievals, and in the air-mass factor calculation. In the low- q_a areas identified on Figure 7, it is therefore possible that high levels of aerosol-generating industrial activities, in a small number of pixels in the grid, results in a value of q_a below 0.75 in those pixels, but also in all the surrounding pixels due to the low resolution of the parameter responsible for exceeding the threshold, without these pixels being located over particularly emissive zones. This effect would be absent in the version 2.4.0 of the TROPOMI product. For the first 34 months of our study, the calculated monthly emissions for these areas would be then underestimated, because they would correspond to the days with low industrial activity.



To evaluate the impact of this effect, we apply the same model using a threshold value of q_a corresponding to 0.7. This includes many pixels that would be rejected with a threshold of 0.75. As the TROPOMI manual recommends not to use this threshold level, we do not consider these emissions to be representative of human activity in Qatar, and they are not treated as emissions estimates. Nevertheless, it allows us to evaluate the effect of non-detection of cement plants and power plants in the south-east of the domain. The use of a threshold value of $q_a = 0.7$ increases the inferred fluxes by 43.8 % above cement plants in the west, and by 33.4 % above the industrial area in the south-east. Fluxes for the Ras Laffan area in the north and central Doha, which are less affected by the effect described here, are increased by 2.6 % and 3.0 % respectively. It is difficult to determine whether these increases are realistic because they were obtained using columns where NO_2 might have been estimated above aerosol and cloud layers. However, they indicate that our model, ran with TROPOMI data before November 2021, leads to a systematic underestimation of NO_x emissions in two of the four most NO_x -intensive areas of the country. We note however that on average, these increases are compensated: with a threshold of $q_a = 0.7$, total fluxes within the internal mask are reduced by 2.1 % because fluxes outside the main emissive areas, which comprises most of the mask, are reduced by about 17 %.

5.4 Weekly cycle

In Qatar, the official rest day is Friday, and the economic activity of the country is lower during this day than during the other days of the week. We therefore try to characterise this feature, by evaluating the weekly cycle of NO_x emissions. We use the TROPOMI-inferred emissions to obtain averages per day of the week. We use the flux-divergence method to calculate the emissions within the internal mask. We remove the days for which most of the area contains poor quality measurements and those for which the wind from Bahrain blows to the south-east, in the same way as it was conducted in Section 5.1.




Figure 8: Mean weekly profiles of anthropogenic NO_x emissions at 13:30 LT in Qatar using TROPOMI observations for 2019 to 2022. The average emissions for the four-year period is displayed with the black line.

For a given day, the empty pixels (i.e. those to which it was not possible to assign emissions) are filled with the average of the emissions obtained for the remaining pixels. This choice leads to an underestimation of the emissions when the main emitting areas are covered by clouds or aerosols, and to an overestimation in the opposite case. To limit this effect, we remove from our estimate of annual averages the emissions below the 5th percentile and above the 95th percentile. Figure 8 shows the resulting daily emissions for the four-year period. Each year, a Friday minimum is observed, defining a weekly cycle. This trend is also observed for mean NO_2 column densities. Fridays have average emissions of 9.04 t.h^{-1} , which is lower than average emissions for the rest of the week, which reach 14.15 t.h^{-1} . In Middle-Eastern countries, this "week-end" effect had already been inferred from satellite measurements by Stavrakou et al. (2020) and Rey-Pommier et al. (2022), and from ground measurements by Butenhoff et al. (2015) on NO_2 concentrations, but with a lower difference between Fridays and the other days.

5.5 Scaling of emissions and annual cycle

In this study, we estimate the emissions at around 13:30 LT, which corresponds to the moment when TROPOMI overpasses the country. However, anthropogenic activity is not uniform throughout the day, and the emissions inferred from parameters calculated at 13:30 do not correspond to the average emissions of the country. Using the power consumption data from Bayram et al. (2018), Figure 9 shows the mean load curve for the country on April 2016, July 2016, October 2016 and



January 2017. These curves show that the average power injected into the grid is slightly lower than the power injected at 13:30. The ratio between the two powers is minimal in June and reaches 0.911, and maximal in February where it reaches 0.976 (see Supplementary Material). Moreover, according to the traffic congestion index in Doha (Tomtom, 2023), the road traffic around 13:30 shows a congestion peak, slightly lower than the other peaks that occur at the beginning and the end of daytime. This suggests that the average emissions from the road transport sector are also close to those at 13:30, with a similar ratio to that of the power sector.




Figure 9: Monthly mean power demand in Qatar for four different months in 2016-2017.

We therefore re-scale the inferred NO_x emissions at the scale of the country by multiplying the monthly estimate for 13:30 with the power ratio to estimate mean monthly emissions. We acknowledge this is a simplification: the power sector has a variable emission factor due to the different technologies in the power plants that are used to respond to the electricity demand over the year, and the road congestion level does not necessarily reflect traffic emissions. Moreover, other sectors, such as cement production, whose seasonality is unknown to us, might behave differently.

Within the internal mask, Figure 10 shows total NO_x emissions in Qatar using our method. Since the NO_2 background has been removed from VCD calculations, the total NO_x budget calculated for each month corresponds to the NO_x production by human activities. The annual variability is marked, with higher emissions in summer and lower emissions in winter. On average, NO_x emissions reach 9.56 kt per month. With mean emissions of 9.29 kt and 10.08 kt, 2019 and 2022 appear to be the years with lowest and highest mean emissions respectively. No seasonal cycle seems to appear for VCDs, which suggests more intense oxidation in summer, which is captured by higher OH values for summer months in CAMS data. The uncertainties, detailed in Section 7, are higher for the months for which emissions could not be attributed for the western and south-eastern parts of the territory. This hinders year-on-year comparisons. Consequently, although emissions in the March/April/May (MAM) 2020 period are 96 % and 87 % of the level of emissions in MAM 2019 and MAM 2021 respectively, we can not assert whether this effect can be attributed to the restrictions put in place from March to May 2020 to tackle the Covid-19 pandemic. This caution is all the more necessary as these restrictions have been less stringent than in other countries (Hale et al., 2021). Similarly, higher emissions during 2022 could be due to the different values reached by q_a in the 2.4.0 version of the TROPOMI product with respect to the 2.3.1 version. However, it can not be excluded that these higher emissions were due to a more intense activity in 2022 in the context of organising the FIFA World Cup.




Figure 10: Total TROPOMI-derived anthropogenic NO_x emissions in Qatar from 2019 to 2022.



6 Comparison with air pollution inventories and electricity generation data

As Qatar is a small country with an undiversified economy, the annual cycle of the TROPOMI-inferred emissions can be interpreted. Reported emissions reached 163 kt in 2007, but a significant part did not take place within the terrestrial
 375 borders of the country. To compare such reported emissions with those calculated within the internal mask, emissions from upstream oil and gas operations (41.86 kt), which mostly take place offshore, must be removed from the previous budget. We also remove emissions from navigation (4.35 kt), civil aviation (3 kt) and fugitive emissions (4.28 kt). Reported terrestrial emissions therefore reach the approximate value of 109.51 kt for 2007, i.e. an average of 9.13 kt per month, with the main NO_x emitting sectors being power generation, transport and cement. Total emissions, as well as and their share in the total
 380 NO_x budget, may have varied since then. This section confronts our estimated emissions with local electricity generation and existing global inventories in order to provide a sectoral approach of NO_x emissions and their annual variability.

6.1 Comparison with electricity generation data

In Qatar, all electricity production comes from gas-fired power plants, with three plants located in the north-east (Ras Laffan area). Other gas power plants are located in the urban area of Doha and in the south-east of the country. Figure 11 shows the
 385 total electricity production and the share of the power plants in Ras Laffan between 2019 and 2021 as provided by Kahramaa. Data regarding the electricity generation by power plant in 2022 is not yet available. The electricity generation is 2 to 2.5 times higher during summer months than during winter months. This increased generation is mostly due to the use of air conditioning (Gastli et al., 2013), which is also captured on Figure 9.




Figure 11: Electricity generation in Qatar from 2019 to 2021 according to the Planning and Statistics Authority reports. Generation from the three Ras Laffan power plants, in the north-eastern part of the country, are displayed in yellow, orange and brown, while generation from other power plants are displayed in light blue.

The inferred emissions have a behaviour that is similar to electricity generation, with higher emissions in the warmer
 390 months. The comparison of the time series of NO_x emissions and the electricity generation data for the four-years period provides a correlation coefficient of $R^2 = 0.400$. This value is relatively low, but it should be noted that monthly emission maps can be very noisy due to averaging over too few days. For instance, on April 2019, the spatial distribution of emissions for this month shows non-negligible emissions in the central part of the country where no significant emitter is located. We can therefore consider that the average is not calculated with enough data to limit noise effects. When we only keep in the analysis
 395 the mean monthly emissions calculated on the basis of more than 18 days (the average over the 4-year period being 17.1), 19 points out of 48 are retained and the correlation is improved, with a coefficient which then reaches $R^2 = 0.657$, with a slope of $1.773 \text{ t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$. This value would be equal to the emission factor of the power sector, provided that it was the only source of variable NO_x emissions during the months that are considered. This value must therefore be considered as an upper limit estimate. Figure 12 shows the comparison between electricity generation and emissions in the two cases. The points
 400 that are retained in the second case mostly correspond to months with high emissions, notably in autumn. Among these 19 points, only 2 correspond to winter or spring months (December to May) whereas 7 correspond to summer months (June to August) and 10 to autumn months (September to November). The improved correlation must be interpreted with caution, but it might be possible that other NO_x-producing sectors have a seasonality which is the opposite to that of the power sector. If the transport sector has no particular seasonal cycle, it is possible that industry has a higher production during winter and
 405 spring months, leading to higher industrial NO_x emissions during this period. As mentioned before, mid-day temperatures in summer are high in Qatar, preventing labor in outdoor activities, and this interpretation could be investigated. With the exception of the two cement plants in the west of the country, most of the industries are located in the urban area of Doha or Ras Laffan, where emissions from several sectors overlap over distances smaller than the resolution of TROPOMI.




Figure 12: Comparison between monthly TROPOMI-derived NO_x emissions above Qatar and electricity generation according to Planning and Statistics Authority reports. A linear regression between two datasets is displayed with a green dashed line. (a) All months in the period 2019-2022 are included. (b) Months that are included are those for which inferred emissions have been obtained with an averaging of 18 daily emissions or more. The color scale represents the number of days involved in the calculation of points.

The method can also be applied to a smaller area of the domain. The north-eastern part of the country is interesting in this respect: it is very sparsely populated and the human activity there is mainly associated with oil and gas activities and the production of electricity. In this region, three power plants (Ras Laffan A, B and C) have a total capacity of 3.35 GW, and are responsible of about 45 % of the total electricity generation throughout the year, as shown on Figure 11. These power plants are independent water and power plants (IWPPs), serving as both desalination plants and power plants. Other than these plants, the region also has gas liquefaction facilities, as Qatar is a major producer of LNG. Refineries are also present. In the life cycle of gas, only extraction and combustion are high NO_x-emitting processes: emissions from midstream (liquefaction and transport) and downstream (refining) activities are 20 to 30 times lower per unit of fuel (Marais et al., 2022). As the oil and gas extraction processes are located outside this area (mostly offshore), it can be considered that the majority of the NO_x emitted in this region come from the combustion of gas in the power plants. This is not the case for other power plants in the country, which are located in urban areas where emissions from different sectors overlap. We focus on this area and calculate the corresponding monthly emissions. The three power plants are located within the same pixel, and the difficulty for TROPOMI to visualise this region with good quality measurements is less pronounced than for other regions identified in Section 5.3. Because Ras Laffan plants are not the only ones participating in the country's electricity generation, the use of load curves is not longer valid to infer total emissions on a given period, and only emissions at 13:30 can be considered. Assuming a correct calculation of emissions, the observed emissions display a Gaussian distribution around the power plants. Using a zonal cross-section of obtained emissions in a 28-km band centered around the mean latitude of the plants, we estimate the mean emissions by fitting a Gaussian curve $e_{\text{NO}_x}(\lambda) = B + \frac{E_0}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(\lambda-\lambda_0)^2}{2\sigma^2}\right)$ on the profile, with B the observed emissions above desert areas and seas, λ_0 the mean longitude of the power plants, σ the standard deviation (measured as an angle) and E_0 the total emissions. λ is the zonal angle. The best approximation is displayed on Figure 13.




Figure 13: Zonal cross sections of 13:30 NO_x emissions at the latitude of Ras Laffan power plants. The average profile for 2019-2022 is displayed in red, and the corresponding Gaussian fit in black. Individual monthly profiles are displayed in light grey. The positions of the three power plants are marked with vertical blue lines.



Monthly profiles peak around the location of the power plants whose height vary without displaying any particular seasonality. The fit of the averaged profile leads to a low positive background of 67.8 kg.h^{-1} per unit of band surface, i.e. about $0.390 \text{ mg.m}^{-2}.\text{h}^{-1}$, and average emissions of 1.86 t.h^{-1} for the three power plants. Two lower peaks are also identified in the west, corresponding to emissions from southern Bahrain and Dammam (Saudi Arabia).

Here, only a small part of the domain is considered, for which the power sector is by far the major contributor to NO_x emissions. It is therefore not necessary to consider the seasonality of the other sectors in the estimation of an emission factor, as it was done earlier when the emissions of the whole domain were considered. However, as the scaling assumption can no longer be used, other assumptions must be made. One assumption would be that these plants serve as intermediate-load power plants: 13:30 being a peak of consumption at any time of the year, the Ras Laffan plants would be running very close to their maximum capacity, and the annual variability of power generation by the three plants observed in Figure 11 would be due to the decrease of power generation during low demand hours. This would be consistent with the lack of seasonality detected in the monthly profiles in Figure 13. In this case, comparing the emissions at 13:30 with the total capacity of the plants provides an emission factor of $0.557 \text{ t}_{\text{NO}_x}.\text{GWh}^{-1}$. As the total capacity is used in the calculation, this value must be considered as a lower limit estimate.

Another possible assumption is that the three plants are used as peak-load power plants: in this case, they would be used at 13:30 to respond to the daily peak. This use would increase with the height of the peak, and therefore with the temperature. In this case, the annual variability of power generation by the three plants would be due to their variable use at 13:30. The absence of a clear annual variability in our TROPOMI-inferred emissions would be explained by the large uncertainties in the calculation of our emissions, which concern only 6 pixels of our domain. Here, the large share of the three power plants in the electricity production seems to rule out a peak-load functioning. However, it is also possible that some of the plants correspond to different functions within the national grid.

6.2 Comparison with emission inventories

We compare TROPOMI-derived NO_x emissions to the Emissions Database for Global Atmospheric Research (EDGARv6.1) for 2018 and the CAMS global anthropogenic emissions (CAMS-GLOB-ANT_v5.3) from 2019 to 2022. The two inventories provide $0.1^\circ \times 0.1^\circ$ sectorial gridded emissions on a monthly basis. After aggregating the different sectors of activity, CAMS-GLOB-ANT and EDGAR inventories directly provide the anthropogenic NO_x emissions over the same domain. Both inventories display NO_x emissions that are significantly higher than our estimates. According to CAMS-GLOB-ANT, the annual emissions for 2019, 2020, 2021 and 2022 are 164.8, 164.6, 165.3 and 166.4 kt respectively, whereas we only estimate emissions of 111.5, 111.8, 114.4 and 121.0 kt for these same years. According to EDGAR, NO_x emissions are even higher, reaching 193.8 kt in 2018. It should be noted that the same version of EDGAR estimates emissions of 111.1 kt in 2007, which is similar to the reported terrestrial emissions for the same year, as well as our estimated emissions for 2019-2022.

The sectoral shares of emissions estimated by EDGAR and CAMS-GLOB-ANT differ. In EDGAR, the transport sector accounts for 42 to 54 % of the emissions, whereas it has a nearly constant share of 58 % in CAMS-GLOB-ANT. In the remaining emissions, industry accounts for about 2.8 kt per month in both inventories, which constitutes a lower share in EDGAR (between 13 and 23% of terrestrial emissions) than in CAMS-GLOB-ANT (about 18% of terrestrial emissions). Such shares are lower than reported shares for 2007, where the cement sector alone accounted for 23.4% of total emissions. Finally, power sector emissions in both inventories account for a significant share of total emissions, but with a higher value for EDGAR (35 % of the total budget for 2018) than for CAMS-GLOB-ANT (20 % of total emissions in 2019-2022). The two values are respectively higher and lower than the reported share for power emissions, which was estimated at 28.1 % for terrestrial emissions in 2007. Another major difference concerns the seasonal cycle of those power emissions. In 2018, EDGARv6.1 shows a pronounced variability, with extremes of 7.0 kt emitted in May and 3.5 kt in December (see Supplementary Material). The ratio between these two values remains lower than the ratio between the extremes of the monthly electricity generations, which has varied between 2.2 and 2.8 over the last four years, but it highlights a difference with the previous version of EDGAR (EDGARv5.0) which showed almost no seasonality in power emissions. This lack of seasonality is also found in the emissions of CAMS-GLOB-ANT_v5.3, which uses EDGARv5.0 as a basis. We also note that while these seasonality differences between TROPOMI-inferred emissions and inventories were already highlighted for the previous versions of EDGAR and CAMS-GLOB-ANT in Egypt, the difference between mean annual NO_x emissions was much lower (Rey-Pommier et al., 2022).

6.3 Emission factor for the power sector

The report for emissions in 2007 indicates an emission factor for the power sector of $0.392 \text{ t}_{\text{NO}_x}.\text{GWh}^{-1}$, and the corresponding power plants are still operating today. In the previous sections, we estimated an emission factor of $1.773 \text{ t}_{\text{NO}_x}.\text{GWh}^{-1}$ using all monthly emissions in the internal mask and assuming the power sector was the only source of variable NO_x emissions. We



also estimated a value of $0.557 \text{ t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$ for Ras Laffan power plants using mean emissions at 13:30 under the hypothesis of a intermediate-load power plant functioning. To infer emission factors from inventory estimates, power emissions in EDGAR and CAMS-GLOB-ANT and total electricity generation provided by the Planning and Statistics Authority (PSA) between 2018 and 2022 can also be used to infer emission factors of $1.406 \text{ t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$ and $0.647 \text{ t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$ respectively. Table 1 summarises all the different emission factors that have been calculated. Even though EDGAR’s emission factor is particularly high compared to other estimates, the five listed values are plausible, as the majority of gas-fired power plants have a NO_x emission factor value between 0.1 and $10 \text{ t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$ (Miller, 2004).

Year(s)	Source	Data for NO_x emissions	Data for power	Method	Emission factor ($\text{t}_{\text{NO}_x} \cdot \text{GWh}^{-1}$)
2007	All PPs	Reported emission factor (Initial National Communication to the UNFCCC)			0.392
2018	All PPs	EDGARv6.1 inventory NO_x emissions	Annual power generation (PSA)	Ratio between total power emissions and electricity generation	1.406
2019-2022	All PPs	CAMS-GLOB-ANT_v5.3 inventory NO_x emissions	Monthly power generation (PSA)	Ratio between total power emissions and electricity generation	0.674
2019-2022	Internal mask	Monthly TROPOMI-inferred NO_x emissions	Monthly power generation (PSA)	Linear fit between monthly power emissions and electricity generation	1.773
2019-2022	Ras Laffan power plants	13:30 TROPOMI-inferred NO_x emissions	Power plant capacities (PSA)	Gaussian fit of zonal cross-section for Ras Laffan power plants	0.557

Table 1: NO_x emission factors for electricity production inferred from different sources and methods. The value is calculated from monthly emissions and electricity generation, fluxes at 13:30 and capacities or directly reported by Qatar authorities.

The comparison between these different emission factors should be made with caution. The factor reported by the report for 2007 emissions takes into account all of the power plants in the country, the oldest having been built in 1980. Moreover, some units of one of the Ras Laffan plants were built between 2010 and 2011, i.e. after the publication of the report. It is also possible that other power plants have been upgraded in the last years. Finally, the reported emission factor for 2007 accounts for the public electricity production and the water production, but the monthly reports from Kahramaa do not specify whether the electricity generation from the Ras Laffan power plants account for the large amount of electricity that is used in desalination processes (Kahramaa, 2023). Monthly reports show that the water production from the IWPPs in the country does not have a clear annual cycle, the production generally ranging between 50 and 60 millions of cubic meters per month in the entire country. The amount of electricity needed for this water production varies with the technologies used in the IWPPs, and accounts for about 20 % of the consumption in total electricity (Okonkwo et al., 2021). Consequently, if the electricity used for water production is not accounted for in the electricity generation for monthly reports, then all inferred emission factors are over-estimated.

7 Uncertainties and assessment of results

The long time series of average NO_2 concentrations in cities in several countries with similar economic and industrial developments (Dubai, Manama, Kuwait City) show a stagnation of NO_x levels over the period 2005-2014 (Lelieveld et al., 2015). It is therefore not clear whether the NO_x emissions of the country in 2019-2022 have increased from their value in 2007, as suggested by CAMS and EDGAR, or decreased, as our TROPOMI-based estimates seem to suggest. We don’t know exactly how parameters in EDGARv6.1 and CAMS-GLOB-ANT_v5.3 are estimated in order to calculate NO_x emissions, which means we cannot conduct a discussion on such bottom-up estimates accordingly. However, we can identify several factors that could influence of our top-down NO_x emissions. The following aspects can be considered:

- As discussed in Section 5.3, a large portion of the territory, which includes some of the most emissive facilities in the country, is not visible by TROPOMI with high-quality data, and emissions from very emissive pixels might be underestimated. As this situation only occurs from 2019 to 2021, the magnitude of the induced underestimation is the difference between inferred emissions in 2022 and inferred emissions in 2019-2021, i.e. about 6 %.

- TROPOMI retrievals are sometimes biased due to poor estimation of the air mass factor or local effects under particular vertical distribution conditions (Griffin et al., 2019; Lorente et al., 2019; Judd et al., 2020; Wang et al., 2021). It should be noted that the latest versions of TROPOMI (v2.x) have tropospheric VCDs that are between 10 % and 40 % larger than the first versions (v1.x), depending on the level of pollution and season (Van Geffen et al., 2022). The chemical transport model TM5, which is used in the retrieval of the operational TROPOMI data, has been shown to underestimate surface level pollution while overestimating NO_2 at higher levels above the sea (Latsch et al., 2023; Rieß et al., 2023). Such a bias is probably not negligible and probably accounts for significant part of the difference between our estimates and inventories, but also within our estimates, as surface albedo values differ between desert areas and urban zones.

- The vertical levels on which parameters in Equation 1 are estimated might be incorrect. Most studies use values



averaged within the PBL, which is higher in summer (Lorente et al., 2019; Lama et al., 2020), or averaged within a fixed vertical layer (Beirle et al., 2019). Here, we interpolated the fields of w , T and $[OH]$ using the first pressure levels in ERA5 and CAMS at a mean pressure of 987.5 hPa. As Qatar is a flat country, this level corresponds to an altitude between 220 and 250 m. Because vertical transport of NO_x , which is emitted mainly from combustion engines and industrial stacks, is generally minor compared to horizontal transport (Sun, 2022), NO_x is confined to these first hundred metres above ground level. In Egypt, Rey-Pommier et al. (2022) have shown that the consideration of higher levels decrease mean temperature which tends to increase emissions, balanced by the decrease of mean OH concentrations with altitude which tends to decrease the sink. The net effect would be a decrease of emissions of a few percents during most of the year.

- The NO_x -to- NO_2 ratio might locally be under-estimated. The conversion of NO to NO_2 by the reaction with O_3 is balanced by the photolysis of NO_2 which reforms NO, leading to a stabilisation of the ratio a few kilometers downwind the source. The stationary regime can also be displaced by the presence of volatile organic compounds. Near important sources of emissions, which mainly produce NO, a high NO_x -to- NO_2 ratio should be expected. We can therefore consider that for pixels containing hotspots close to their boundaries, such as power plants, it should be much higher than the chosen value of 1.32 (Hanrahan, 1999; Goldberg et al., 2022), leading to an underestimation of our emissions locally. With power plants concentrating a large share of the NO_x emissions in our study, this effect is probably not negligible and probably accounts for a large part of the difference between our estimates and inventories.

- The sink term might be under-estimated or over-estimated. If the transport term is assumed to be estimated correctly, an underestimation of the sink term would lead to significant negative emissions in many parts of the domain, while an overestimation would lead to non-emitting areas with NO_x emissions that are higher than noise. None of these effects seem to appear for averaged emissions for the 2019-2022 period, but monthly maps show significant portions of the territory with negative emissions in winter and spring months, while summer months display abnormally high emissions over deserts and seas. Those effects are not visible for most of the autumn months. We can therefore assume that the sink term is slightly under-estimated in winter and spring months and over-estimated in summer months, due to errors in the estimation of OH in CAMS and/or the presence of additional sinks in summer. Accounting for this effect would not change the averaged emissions for the period 2019-2022, but would reduce the seasonality observed on Figure 6. The emission factor calculated from emissions in the internal mask would also be reduced.

These effects must be taken into account for understanding the differences between our model, reported emissions and inventory emissions. The uncertainties used here must also be considered. NO_x emissions, as estimated with the flux-divergence method, are calculated with the use of several quantities. The uncertainty ranges shown in Figures 10 and 12 are calculated from uncertainty statistics whose references are presented in this section, assuming that they correspond to standard deviations. The uncertainty of tropospheric NO_2 columns under polluted conditions is dominated by the sensitivity of satellite observations to air masses in the lower troposphere, expressed by the AMF, and the corresponding relative uncertainty is of the order of 30 % (Boersma et al., 2004), and is likely related to the a priori profiles used within the operational retrieval that do not reflect well the concentration peak of NO_2 near the ground. For the Middle East region, the impact of the a priori profile is less critical, as surface albedo is generally high and cloud fractions are generally low. Thus, we do not expect a bias of this importance, and consider a relative uncertainty of 30 % for the tropospheric column as reasonable. For wind module, we assume an uncertainty of $3 \text{ m}\cdot\text{s}^{-1}$ for both zonal and meridional wind components. For OH concentrations, the analysis of different methods conducted by Huijnen et al. (2019) showed smaller differences for low latitudes than for extratropics, but still significant. We thus take a relative uncertainty of 30 % for OH concentration. For the reaction rate k_{mean} , we use the value of the corresponding relative uncertainty, as estimated by Burkholder et al. (2020).

8 Conclusion

In this study, we investigated the potential of a top-down model of NO_x emissions based on TROPOMI retrievals and a flux-divergence scheme applied at high resolution over Qatar for the last four years (2019-2022). This scheme requires different parameters to be calculated and consists in the calculation of a transport term that uses horizontal wind, and the calculation of a sink term that requires temperature data and OH concentration to illustrate the chemical loss of NO_x . Results illustrate the difference between localised and diffuse sources of NO_x . For diffuse sources such as the Doha urban area, the transport term and the sink term similarly contribute to the total NO_x budget, whereas the transport term is higher than the sink term for localised sources such as the gas power plants located in the north-east of the country. The emissions from other hotspots, such as cement plants in the west part of the country, could not be correctly estimated from 2019 to 2021 due to unavailability of high-quality TROPOMI retrievals. Our estimated NO_x emissions show a weekly variability which is consistent with the social norms of the country and an annual variability which is consistent with its electricity generation. In this study, estimated emissions are similar to reported emissions in 2007, but they are 1.44 times lower than emissions in the CAMS-GLOB-ANT_v5.3 inventory for 2019-2022 and 1.68 times lower than emissions in EDGARv6.1 for 2018. They have an annual cycle whose relative amplitude is higher than those two inventories. These notable differences may be subject



575 to further discussion regarding sectoral activity data and emission factors used in global inventories. Finally, this study is
also an attempt to retrieve the NO_x emission factor of the power sector. This top-down estimation is made possible by the
desert features of Qatar, which allow to consider only one chemical sink, and also by the low diversity of its economy in which
the power sector is the major source of variable NO_x . An emission factor is also estimated for a group of isolated gas power
plants. The emission factors are 1.42 and 4.52 times higher than the value reported for 2007. Moreover, the emission factor of
580 the entire power sector is higher than that of the three isolated gas power plants, indicating higher emission factors for other
power plants in Doha. Although several limitations exist in the estimation of these results, we believe that such a calculation
can be reproduced in other countries and for power plants that share similar characteristics. Overall, this study highlights the
potential of TROPOMI to compensate for non-existent, inaccurate or outdated inventories by providing low-latency emissions
estimates. The development of similar applications is likely to provide a better monitoring of global anthropogenic emissions,
585 therefore helping countries to report their emissions of air pollutants and greenhouse gases as part of their strategies and
obligations to tackle air pollution issues and climate change.

Code availability. Code will be made available on request.

Data availability. The TROPOMI Sentinel 5P Product Algorithm Laboratory (S5P-PAL) reprocessed data (pro-
cessor version 2.3.1) from January 2019 to October 2020 has been used. The OFFL stream has been used af-
590 terwise, using processor versions 2.3.1 and 2.4.0 from November 2022. The TROPOMI NO_2 product is pub-
licly available on the TROPOMI Open hub (<http://www.tropomi.eu/data-products/data-access>, TROPOMI Data
Hub, 2022) while the S5P-PAL reprocessed data can be found on the S5P-PAL data portal ([https://data-
portal.s5p-pal.com](https://data-portal.s5p-pal.com), ESA, 2022). CAMS data can be downloaded from the Copernicus Climate Data Store
(<https://ads.atmosphere.copernicus.eu/cdsapp#!/dataset/cams-global-atmospheric-composition-forecasts>, ECMWF, 2022a).
595 The European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 reanalysis be downloaded from the Coperni-
cus Climate Data Store (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels-monthly-means>,
ECMWF, 2022b). Electricity generation data can be downloaded from The Planning and Statistics Authority Por-
tal (<https://www.psa.gov.qa/en/statistics1/Pages/default.aspx>, last access: 31 January 2023), and from Kahramaa Por-
tal (<https://www.km.qa/MediaCenter/Pages/Publications.aspx>, last access: 24 January 2023). Emissions reported for
600 year 2007 is available on the UNFCCC website (<https://unfccc.int/resource/docs/natc/qatnc1.pdf>, last access: 26 August
2022). EDGARv6.1 emissions are provided by https://edgar.jrc.ec.europa.eu/emissions_data_and_maps. CAMS-GLOB-
ANT_v5.3 emissions are available at <https://eccad3.sedoo.fr>.

Author contributions. AR analysed the data, prepared the main software code and wrote the paper. FC provided the
TROPOMI NO_2 data product and corresponding gridded maps. ISB provided time series of electricity consumption. PC,
605 TC, JK, and JS contributed to the improvement of the method and the interpretation of the results. All authors read and
agreed on the published version of the paper.

Financial support. This study has been funded by the European Union's Horizon 2020 research and innovation programme
under grant agreement N°856612 (EMME-CARE).

Competing interests. The authors declare that they have no conflict of interest.

610 References

- Al-Attiyah Foundation*. Gear change: Vehicle fuel efficiency in the GCC, [https://www.abhafoundation.org/media-
uploads/rpapers/Research%20series-17-Date-01-2018-Fuel%20Efficiency%20GCC.pdf](https://www.abhafoundation.org/media-uploads/rpapers/Research%20series-17-Date-01-2018-Fuel%20Efficiency%20GCC.pdf). 2018.
- Bayram Islam Safak, Saffouri Faraj, Koc Muammer*. Generation, analysis, and applications of high resolution electricity load
profiles in Qatar // *Journal of Cleaner Production*. 2018. 183. 527–543.
- 615 *Beirle S, Borger C, Dörner S, Li A, Hu Z, Liu F, Wang Y, Wagner T*. Pinpointing nitrogen oxide emissions from space, *Sci.*
Adv., 5, eaax9800. 2019.
- Boersma KF, Eskes HJ, Brinksma EJ*. Error analysis for tropospheric NO_2 retrieval from space // *Journal of Geophysical*
Research: Atmospheres. 2004. 109, D4.



- 620 *Bovensmann Heinrich, Burrows JP, Buchwitz M, Frerick Johannes, Noel Suresh, Rozanov VV, Chance KV, Goede APH.*
SCIAMACHY: Mission objectives and measurement modes // *Journal of the atmospheric sciences.* 1999. 56, 2. 127–150.
- Burkholder JB, Sander SP, Abbatt JPD, Barker JR, Cappa C, Crounse JD, Dibble TS, Huie RE, Kolb CE, Kurylo MJ, others .* Chemical kinetics and photochemical data for use in atmospheric studies; evaluation number 19. 2020.
- Burnett Richard T, Stieb Dave, Brook Jeffrey R, Cakmak Sabit, Dales Robert, Raizenne Mark, Vincent Renaud, Dann Tom.*
625 Associations between short-term changes in nitrogen dioxide and mortality in Canadian cities // *Archives of Environmental Health: An International Journal.* 2004. 59, 5. 228–236.
- Butenhoff Christopher L, Khalil M Aslam K, Porter William C, Al-Sahafi Mohammed Saleh, Almazroui Mansour, Al-Khalaf Abdulrahman.* Evaluation of ozone, nitrogen dioxide, and carbon monoxide at nine sites in Saudi Arabia during 2007 // *Journal of the Air & Waste Management Association.* 2015. 65, 7. 871–886.
- 630 *Celarie EA, Brinksma EJ, Gleason JF, Veefkind JP, Cede A, Herman JR, Ionov Dimitri, Goutail Florence, Pommereau J-P, Lambert J-C, others .* Validation of Ozone Monitoring Instrument nitrogen dioxide columns // *Journal of Geophysical Research: Atmospheres.* 2008. 113, D15.
- Demetillo Mary Angelique G., Navarro Aracely, Knowles Katherine K., Fields Kimberly P, Geddes Jeffrey A., Nowlan Caroline R., Janz Scott J, Judd Laura M, Al-Saadi Jassim, Sun Kang, McDonald Brian C., Diskin Glenn S., Pusede Sally E.* Observing nitrogen dioxide air pollution inequality using high-spatial-resolution remote sensing measurements in
635 Houston, Texas // *Environmental Science & Technology.* 2020. 54, 16. 9882–9895.
- EIA . Natural gas, <https://www.eia.gov/international/data/world/natural-gas/dry-natural-gas-reserves>. 2021.
- EIA . Analysis - Energy Sector Highlights, <https://www.eia.gov/international/data/country/QAT>. 2022.
- EPA US. Integrated science assessment for oxides of nitrogen–health criteria // US Environmental Protection Agency, Washington, DC. 2016.
- 640 *Ehhalt Dieter H, Rohrer Franz, Wahner Andrea.* Sources and distribution of NO_x in the upper troposphere at northern mid-latitudes // *Journal of Geophysical Research: Atmospheres.* 1992. 97, D4. 3725–3738.
- Eskes H, Geffen J van, Boersma F, Eichmann KU, Apituley A, Pedergnana M, Sneep M, Veefkind JP, Loyola D.* Sentinel-5 Precursor/TROPOMI Level 2 Product User Manual Nitrogen Dioxide. 2022.
- Gastli Adel, Charabi Yassine, Alammari Rashid A, Al-Ali Ali M.* Correlation between climate data and maximum electricity
645 demand in Qatar. 2013. 565–570.
- Goldberg Daniel L, Harkey Monica, Foy Benjamin de, Judd Laura, Johnson Jeremiah, Yarwood Greg, Holloway Tracey.* Evaluating NO_x emissions and their effect on O₃ production in Texas using TROPOMI NO₂ and HCHO // *Atmospheric Chemistry and Physics.* 2022. 22, 16. 10875–10900.
- Graedel TE, Farrow LA, Weber TA.* Kinetic studies of the photochemistry of the urban troposphere // *Atmospheric Environment (1967).* 1976. 10, 12. 1095–1116.
650
- Granier Claire, Darras Sabine, Gon Hugo Denier van der, Jana Doubalova, Elguindi Nellie, Bo Galle, Michael Gauss, Marc Guevara, Jalkanen Jukka-Pekka, Kuenen Jeroen, others .* The Copernicus atmosphere monitoring service global and regional emissions (April 2019 version). 2019.
- Griffin Debra, Zhao Xiaoyi, McLinden Chris A, Boersma Folkert, Bourassa Adam, Dammers Enrico, Degenstein Doug, Eskes Henk, Fehr Lukas, Fioletov Vitali, others .* High-resolution mapping of nitrogen dioxide with TROPOMI: First
655 results and validation over the Canadian oil sands // *Geophysical Research Letters.* 2019. 46, 2. 1049–1060.
- Hale Thomas, Angrist Noam, Goldszmidt Rafael, Kira Beatriz, Petherick Anna, Phillips Toby, Webster Samuel, Cameron-Blake Emily, Hallas Laura, Majumdar Saptarshi, others .* A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker) // *Nature Human Behaviour.* 2021. 5, 4. 529–538.
- 660 *Hanrahan Patrick L.* The plume volume molar ratio method for determining NO₂/NO_x ratios in modeling—Part II: Evaluation studies // *Journal of the Air & Waste Management Association.* 1999. 49, 11. 1332–1338.
- He Mike Z, Kinney Patrick L, Li Tiantian, Chen Chen, Sun Qinghua, Ban Jie, Wang Jiaonan, Liu Siliang, Goldsmith Jeff, Kioumourtzoglou Marianthi-Anna.* Short-and intermediate-term exposure to NO₂ and mortality: a multi-county analysis in China // *Environmental Pollution.* 2020. 261. 114165.



- 665 *Hersbach Hans, Bell Bill, Berrisford Paul, Hirahara Shoji, Horányi András, Muñoz-Sabater Joaquín, Nicolas Julien, Peubey Carole, Radu Raluca, Schepers Dinand, others* . The ERA5 global reanalysis // Quarterly Journal of the Royal Meteorological Society. 2020. 146, 730. 1999–2049.
- Hoelzemann Judith J, Schultz Martin G, Brasseur Guy P, Granier Claire, Simon Muriel* . Global Wildland Fire Emission Model (GWEM): Evaluating the use of global area burnt satellite data // Journal of Geophysical Research: Atmospheres. 670 2004. 109, D14.
- Hoesly Rachel M, Smith Steven J, Feng Leyang, Klimont Zbigniew, Janssens-Maenhout Greet, Pitkanen Tyler, Seibert Jonathan J, Vu Linh, Andres Robert J, Bolt Ryan M, others* . Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS) // Geoscientific Model Development. 2018. 11, 1. 369–408.
- 675 *Huijnen Vincent, Pozzer Andrea, Arteta Joaquim, Brasseur Guy, Bouarar Idir, Chabrillat Simon, Christophe Yves, Doumbia Thierno, Flemming Johannes, Guth Jonathan, others* . Quantifying uncertainties due to chemistry modelling–evaluation of tropospheric composition simulations in the CAMS model (cycle 43R1) // Geoscientific Model Development. 2019. 12, 4. 1725–1752.
- Jaeglé Lyatt, Steinberger Linda, Martin Randall V, Chance Kelly* . Global partitioning of NO_x sources using satellite observations: Relative roles of fossil fuel combustion, biomass burning and soil emissions // Faraday discussions. 2005. 130. 680 407–423.
- Janssens-Maenhout Greet, Crippa Monica, Guizzardi Diego, Muntean Marilena, Schaaf Edwin, Dentener Frank, Bergamaschi Peter, Pagliari Valerio, Olivier Jos GJ, Peters Jeroen AHW, others* . EDGAR v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012 // Earth System Science Data. 2019. 11, 3. 959–1002.
- 685 *Judd Laura M, Al-Saadi Jassim A, Szykman James J, Valin Lukas C, Janz Scott J, Kowalewski Matthew G, Eskes Henk J, Veeffkind J Pepijn, Cede Alexander, Mueller Moritz, others* . Evaluating Sentinel-5P TROPOMI tropospheric NO₂ column densities with airborne and Pandora spectrometers near New York City and Long Island Sound // Atmospheric measurement techniques. 2020. 13, 11. 6113–6140.
- Kahramaa* . Qatar General Electricity and Water Corporation, <https://www.km.qa/MediaCenter/Pages/Publications.aspx>. 690 2023.
- Lama Srijana, Houweling Sander, Boersma K Folkert, Eskes Henk, Aben Ilse, Gon Hugo AC Denier van der, Krol Maarten C, Dolman Han, Borsdorff Tobias, Lorente Alba* . Quantifying burning efficiency in megacities using the NO₂/CO ratio from the Tropospheric Monitoring Instrument (TROPOMI) // Atmospheric Chemistry and Physics. 2020. 20, 17. 10295–10310.
- Latsch Miriam, Richter Andreas, Burrows John P* . Improving the detection of global NO_x emissions from shipping in S5P/TROPOMI data. 695 2023.
- Lelieveld Jos, Beirle Steffen, Hörmann Christoph, Stenchikov Georgiy, Wagner Thomas* . Abrupt recent trend changes in atmospheric nitrogen dioxide over the Middle East // Science advances. 2015. 1, 7. e1500498.
- Lelieveld Jos, Gromov Sergey, Pozzer Andrea, Taraborrelli Domenico* . Global tropospheric hydroxyl distribution, budget and reactivity // Atmospheric Chemistry and Physics. 2016. 16, 19. 12477–12493.
- 700 *Levy H* . Normal atmosphere: Large radical and formaldehyde concentrations predicted // Science. 1971. 173, 3992. 141–143.
- Li Mengze, Karu Einar, Brenninkmeijer Carl, Fischer Horst, Lelieveld Jos, Williams Jonathan* . Tropospheric OH and stratospheric OH and Cl concentrations determined from CH₄, CH₃Cl, and SF₆ measurements // NPJ Climate and Atmospheric Science. 2018. 1, 1. 1–7.
- Lin J-T* . Satellite constraint for emissions of nitrogen oxides from anthropogenic, lightning and soil sources over East China on a high-resolution grid // Atmospheric Chemistry and Physics. 2012. 12, 6. 2881–2898. 705
- Lorente A, Boersma KF, Eskes HJ, Veeffkind JP, Van Geffen JHGM, De Zeeuw MB, Denier Van Der Gon HAC, Beirle Steffen, Krol MC* . Quantification of nitrogen oxides emissions from build-up of pollution over Paris with TROPOMI // Scientific reports. 2019. 9, 1. 1–10.
- Mansouri Daneshvar Mohammad Reza, Hussein Abadi Nasrin* . Spatial and temporal variation of nitrogen dioxide measurement in the Middle East within 2005–2014 // Modeling Earth Systems and Environment. 2017. 3, 1. 1–9. 710



- Marais Eloise A, Akker Orianna, Wiedinmyer Christine.* Greenhouse gas and air pollutant emissions from power barges (powerships) // *Environmental Science: Advances.* 2022. 1, 2. 164–169.
- Miller Paul Joseph.* North American power plant air emissions. 2004.
- Müller J-F, Stavrou T.* Inversion of CO and NO_x emissions using the adjoint of the IMAGES model // *Atmospheric Chemistry and Physics.* 2005. 5, 5. 1157–1186.
- 715 *OPEC ASB.* OPEC Annual Statistical Bulletin 2019. 2020.
- Okonkwo Eric C, Abdullatif Yasser M, Tareq Al-Ansari.* A nanomaterial integrated technology approach to enhance the energy-water-food nexus // *Renewable and Sustainable Energy Reviews.* 2021. 145. 111118.
- Osipov Sergey, Chowdhury Sourangsu, Crowley John N, Tadic Ivan, Drewnick Frank, Borrmann Stephan, Eger Philipp, Fachinger Friederike, Fischer Horst, Predybaylo Evgeniya, others .* Severe atmospheric pollution in the Middle East is attributable to anthropogenic sources // *Communications Earth & Environment.* 2022. 3, 1. 1–10.
- 720 *PSA .* Planning and Statistics Authority, Qatar Monthly Statistics, <https://www.psa.gov.qa/en/statistics1/Pages/default.aspx>. 2023.
- Rey-Pommier Anthony, Chevallier Frédéric, Ciais Philippe, Broquet Grégoire, Christoudias Theodoros, Kushta Jonilda, Hauglustaine Didier, Sciare Jean.* Quantifying NO_x emissions in Egypt using TROPOMI observations // *Atmospheric Chemistry and Physics Discussions.* 2022. 1–21.
- 725 *Richter A, Burrows JP.* Tropospheric NO₂ from GOME measurements // *Advances in Space Research.* 2002. 29, 11. 1673–1683.
- Rieß T Christoph VW, Vliet Jasper van, Van Roy Ward, Laat Jos de, Dammers Enrico, Boersma Folkert.* Aircraft validation reveals a 20% low bias in TROPOMI NO₂ over sea caused by TM5 a priori profiles. 2023.
- 730 *Rohrer Franz, Berresheim Harald.* Strong correlation between levels of tropospheric hydroxyl radicals and solar ultraviolet radiation // *Nature.* 2006. 442, 7099. 184–187.
- Saw Gautam Kumar, Dey Sagnik, Kaushal Hemant, Lal Kanhaiya.* Tracking NO₂ emission from thermal power plants in North India using TROPOMI data // *Atmospheric Environment.* 2021. 259. 118514.
- 735 *Seinfeld John H.* Urban air pollution: state of the science // *Science.* 1989. 243, 4892. 745–752.
- Seinfeld and Pandis John H and Spyros N.* Atmospheric chemistry and physics from air pollution to climate change. 2006.
- Shikwambana Lerato, Mhangara Paidamwoyo, Mbatha Nkanyiso.* Trend analysis and first time observations of sulphur dioxide and nitrogen dioxide in South Africa using TROPOMI/Sentinel-5P data // *International Journal of Applied Earth Observation and Geoinformation.* 2020. 91. 102130.
- 740 *Stavrou T, Müller J-F, Bauwens M, Boersma KF, Geffen J van.* Satellite evidence for changes in the NO₂ weekly cycle over large cities // *Scientific reports.* 2020. 10, 1. 1–9.
- Sun Kang.* Derivation of Emissions From Satellite-Observed Column Amounts and Its Application to TROPOMI NO₂ and CO Observations // *Geophysical Research Letters.* 2022. 49, 23. e2022GL101102.
- Tomtom .* TomTom, Doha traffic, Qatar, https://www.tomtom.com/en_gb/traffic-index/doha-traffic/. 2023.
- 745 *UNFCCC .* UNFCCC Nationally Determined Contributions Registry, <https://unfccc.int/resource/docs/natc/qatnc1.pdf>. 2011.
- Valin LC, Russell AR, Hudman RC, Cohen RC.* Effects of model resolution on the interpretation of satellite NO₂ observations // *Atmospheric Chemistry and Physics.* 2011. 11, 22. 11647–11655.
- Van Geffen Jos, Eskes Henk, Compernelle Steven, Pinardi Gaia, Verhoelst Tijn, Lambert Jean-Christopher, Sneep Maarten, Ter Linden Mark, Ludewig Antje, Boersma K Folkert, others .* Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-based data // *Atmospheric Measurement Techniques.* 2022. 15, 7. 2037–2060.
- 750



Vandaele Ann Carine, Hermans Christian, Simon Paul C, Carleer Michel, Colin Réginald, Fally Sophie, Merienne Marie-France, Jenouvrier Alain, Coquart Bernard. Measurements of the NO₂ absorption cross-section from 42 000 cm⁻¹ to 10 000 cm⁻¹ (238–1000 nm) at 220 K and 294 K // *Journal of Quantitative Spectroscopy and Radiative Transfer*. 1998. 59, 3-5. 171–184.

Vuuren LF; Smith SJ; Dentener F. DP; Bouwman. Global Projections for Anthropogenic Reactive Nitrogen Emissions to the Atmosphere: An Assessment of Scenarios in the Scientific Literature // *Current Opinion in Environmental Sustainability*. 2011. 3. 359–369.

Wang Yuan, Yuan Qiangqiang, Li Tongwen, Zhu Liye, Zhang Liangpei. Estimating daily full-coverage near surface O₃, CO, and NO₂ concentrations at a high spatial resolution over China based on S5P-TROPOMI and GEOS-FP // *ISPRS Journal of Photogrammetry and Remote Sensing*. 2021. 175. 311–325.

Wolfe Glenn M, Nicely Julie M, St. Clair Jason M, Hanisco Thomas F, Liao Jin, Oman Luke D, Brune William B, Miller David, Thames Alexander, González Abad Gonzalo, others . Mapping hydroxyl variability throughout the global remote troposphere via synthesis of airborne and satellite formaldehyde observations // *Proceedings of the National Academy of Sciences*. 2019. 116, 23. 11171–11180.

World Bank . World Bank Indicators, <https://data.worldbank.org/indicator>. 2022.

Yienger JJ, Levy H. Empirical model of global soil-biogenic NO_x emissions // *Journal of Geophysical Research: Atmospheres*. 1995. 100, D6. 11447–11464.

770