

Evaluation of vertical transport in ERA5 and ERA-Interim reanalysis using high-altitude aircraft measurements in the Asian summer monsoon 2017

Bärbel Vogel¹, C. Michael Volk², Johannes Wintel^{2,*}, Valentin Lauther², Jan Clemens^{1,3}, Jens-Uwe Grooß¹, Gebhard Günther¹, Lars Hoffmann³, Johannes C. Laube¹, Rolf Müller¹, Felix Ploeger¹, and Fred Stroh¹

¹Institute of Energy and Climate Research (IEK-7), Forschungszentrum Jülich, Jülich, Germany

²Institute for Atmospheric and Environmental Research, University of Wuppertal, Wuppertal, Germany

³Jülich Supercomputing Centre, Forschungszentrum Jülich, Jülich, Germany

* now at: Curt-Engelhorn-Centre of Archaeometry gGmbH, Mannheim, Germany

Correspondence: B. Vogel (b.vogel@fz-juelich.de)

Abstract.

During the Asian monsoon season, greenhouse gases as well as pollution emitted near the ground are rapidly uplifted by convection up to an altitude of ~ 13 km, with slower ascent and mixing with the stratospheric background above. Here, we address the robustness of the representation of these transport processes in different reanalysis data sets using ERA5, ERA-Interim and ERA5 $1^\circ \times 1^\circ$. This transport assessment includes mean age of air from global 3-dimensional simulations by the Lagrangian transport model CLaMS as well as different trajectory-based transport times and associated ascent rates compared with observation-based age of air and ascent rates of long-lived trace gases from airborne measurements during the Asian summer monsoon 2017 in Nepal.

Our findings confirm that the ERA5 reanalysis yields a better representation of convection than ERA-Interim resulting in different transport times and air mass origins at the Earth's surface. In the Asian monsoon region above 430 K, mean age of air driven by ERA-Interim is too young, whereas mean age of air from ERA5 $1^\circ \times 1^\circ$ is too old, but somewhat closer to the observations. The mean effective ascent rates derived from ERA5 and ERA5 $1^\circ \times 1^\circ$ back-trajectories are in good agreement with the observation-based mean ascent rates unlike ERA-Interim, which is much faster above 430 K. Although a reliable CO_2 reconstruction is a challenge for model simulations, we show that up to 410 K, the CO_2 reconstruction using ERA5 agrees best with high-resolution in situ aircraft CO_2 measurements, indicating a better representation of Asian monsoon transport in the newest ECMWF reanalysis product ERA5.

1 Introduction

20 The global amount of greenhouse gases (GHGs) in the atmosphere has increased because of worldwide anthropogenic emissions. In particular, the rapid increase of anthropogenic CO₂ emissions in South Asia contributes strongly to the acceleration of the CO₂ growth rate, e.g. the anthropogenic CO₂ emission rate from India was the fourth highest worldwide in 2017 (behind China, the USA and the European Union) (Friedlingstein et al., 2019, 2022). In addition to GHGs also pollution, water vapour, aerosol particles and their precursors as well as some ozone-destroying substances have high emission rates in Asia
25 and can be transported very fast into the lower stratosphere during the Asian summer monsoon season (e.g. Brunamonti et al., 2018; Hanumanthu et al., 2020; Adcock et al., 2021; Appel et al., 2022; Vogel et al., 2023). Subsequently these trace gases as well as aerosol particles can be distributed into the global northern lower stratosphere over a period of several weeks (e.g. Ploeger et al., 2013; Müller et al., 2016; Vogel et al., 2016; Yu et al., 2017; Rolf et al., 2018; Lauther et al., 2022). To better understand the impact of anthropogenic emissions in Asia on the atmosphere, it is important to evaluate the transport of air in
30 the Asian summer monsoon region into the lower stratosphere represented in meteorological reanalyses in combination with unique high-resolution aircraft measurements obtained over the northern Indian subcontinent during the StratoClim aircraft campaign in summer 2017 (Stroh and StratoClim-Team, 2023).

From about June to September, the Asian summer monsoon constitutes a seasonally persistent zonally restricted circulation pattern transporting climate-relevant emissions rapidly from surface sources to higher altitudes, i.e. to the lower stratosphere
35 (e.g. Mason and Anderson, 1963; Randel and Park, 2006; Park et al., 2007; Pan et al., 2016; Vogel et al., 2015; Ploeger et al., 2017; Vogel et al., 2023). The Asian summer monsoon is associated with deep convection over the Indian subcontinent and an anticyclonic flow in the upper troposphere and lower stratosphere (UTLS) over the Asian monsoon region spanning from northeast Africa to the Pacific (e.g. Park et al., 2007). Air parcels are uplifted quickly by convection followed by slow diabatic uplift in the UTLS superimposed on the anticyclonic flow (e.g. Brunamonti et al., 2018; Vogel et al., 2019; Legras and Bucci,
40 2020; von Hobe et al., 2021), while in other regions within the tropical transition layer the heating rates are in general smaller during boreal summer (Vogel et al., 2019). The higher the air parcels are located above the level of maximum convective outflow (~ 360 K, ~ 13 km), the larger the contribution of air masses is from outside the Asian monsoon anticyclone (i.e. from the stratospheric background) to the upward spiralling flow (Vogel et al., 2019, 2023).

In state-of-the-art chemistry transport models, the transport of air parcels differs, because different methods (Eulerian, La-
45 grangian), different vertical velocities (kinematic, diabatic) and different meteorological reanalyses are used to drive the models (e.g. Stenke et al., 2009; Bergman et al., 2013; Brinkop and Jöckel, 2019; Tao et al., 2019; Ploeger et al., 2019; Bucci et al., 2020; Legras and Bucci, 2020; Tegtmeier et al., 2020; Clemens et al., 2023). Further, the implementation of convection and irreversible mixing differs from model to model (e.g. Brinkop and Jöckel, 2019; Konopka et al., 2019; Wohltmann et al., 2019; Hoffmann et al., 2023). The aim of this study is to infer differences of vertical transport in the Asian monsoon region using
50 three data sets: two reanalyses provided by the European Centre for Medium-Range Weather Forecasts (ECMWF), namely, ERA-Interim and its successor ERA5 as well as its down-scaled version ERA5 $1^\circ \times 1^\circ$, a computing-time-saving alternative to the full resolution ERA5 data.

In general, differences between ERA-interim and ERA5 are attributed to the better spatial and temporal resolution of the ERA5 reanalysis, which allows for a better representation of convective updrafts, gravity waves and tropical cyclones (e.g. Hoffmann et al., 2019; Li et al., 2020; Legras and Bucci, 2020; Malakar et al., 2020). Consequently, ERA5 provides a more accurate representation of the lapse rate tropopause height than ERA-Interim (e.g. Hoffmann et al., 2022; Tegtmeier and Krüger, 2022).

In the Asian monsoon anticyclone slow diabatic uplift in the range of 1-1.5K per day occurs above the level of maximum convective outflow using Lagrangian transport simulations driven by ERA-Interim (Vogel et al., 2019). However, it was found consistently in several previous studies that in general the vertical velocities in ERA-Interim are 30-50% too fast in the tropics (Dee et al., 2011; Ploeger et al., 2012; Schoeberl et al., 2012). Tegtmeier and Krüger (2022) summarise that diabatic vertical ascent appears to be faster in ERA-Interim, which produces a residence time (between 370 K and 400 K) of ~ 2 months in the tropical tropopause layer, in contrast to residence times of ~ 3 months or longer based on other reanalyses (e.g. MERRA, MERRA-2, or CFSR, however ERA5 was not included here). This bias seems to be corrected in ERA5 manifesting in weaker diabatic heating rates in the tropics resulting in a greater age of air (i.e., larger mean stratospheric transit times) and thus a significantly slower Brewer–Dobson circulation in ERA5 compared to ERA-Interim (Ploeger et al., 2021). Different residence times in the UTLS would change the chemical composition at these altitudes and even small changes of radiatively active trace gases such as O₃, H₂O or aerosol particles could have important local radiative impacts (e.g. Riese et al., 2012; Vernier et al., 2015; Fadnavis et al., 2019; Bian et al., 2020).

For the StratoClim aircraft campaign during the Asian summer monsoon 2017, in general, a higher consistency with observations and a better reproducibility of pollution features could be found using ERA5 compared to ERA-Interim using diabatic trajectory calculations back to cloud top altitudes (Bucci et al., 2020). ERA5 improves in general the transport in the Asian monsoon 2017, however upward transport in the region of the Tibetan Plateau should be considered with caution (Legras and Bucci, 2020). Considering the transport of air masses contributing to the Asian Tropopause Aerosol Layer (ATAL) measured in the region of the Asian monsoon anticyclone during August 2016, ERA5 shows in general faster transport of air from the ground to ATAL altitudes (up to ≈ 410 K) due to a better representation of convection. In addition, more continental source regions contributing to the ATAL are found in ERA5, whereas in ERA-Interim more marine sources are attributed to air at ATAL altitudes (Clemens et al., 2023).

Using ERA-Interim compared to ERA5 reanalysis yield different vertical velocities or ascent rates in the region of the Asian monsoon anticyclone that has consequences for global transport simulations. To assess such global simulations it is essential to understand the strengths and weaknesses of the newest ECMWF product ERA5 in particular in the Asian monsoon region. In this work, differences in the transport of air in the regions of the Asian summer monsoon 2017 will be inferred using the Chemical Lagrangian Model of the Stratosphere (CLaMS) driven by the three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$). Model results will be assessed using unique airborne measurements up to ~ 20 km during the Asian summer monsoon 2017 conducted during the StratoClim aircraft campaign in Nepal (Stroh and StratoClim-Team, 2023). Trajectory-based transport times, origin of air at the Earth's surface, mean effective ascent rates, transport time distribution as well as mean age of air from 3-dimensional CLaMS simulations are compared using the three data sets (ERA-Interim, ERA5 and




Figure 1. Regional map of the aircraft measurements over the Indian subcontinent. The flight paths of the eight local scientific flights (F01-F08) by the high altitude research aircraft Geophysica are shown. The scientific flights were carried out every second day from Kathmandu (Nepal) between 27 July and 10 August 2017. In addition, the locations of the measurement sites for Greenhouse gases in Nainital (NTL, India) and Comilla (CLA, Bangladesh) used for CO₂ reconstruction are indicated (figure adapted from Vogel et al. (2023)).

ERA5 1° × 1°). In addition, simulated mean age of air is compared to observation-based mean age of air inferred from long-lived trace gases such as C₂F₆, HFC-125, SF₆ and N₂O. In addition, C₂F₆ and HFC-125 are used to compare observation-based mean ascent rates with those from different reanalyses.

Further, a unique set of CO₂ aircraft measurements featuring high temporal and vertical resolution up to ~20 km was obtained during the StratoClim aircraft campaign 2017 (Stroh and StratoClim-Team, 2023). Measured CO₂ profiles were successfully reconstructed using ground-based measurements of CO₂ mainly from Nainital (northern India) by Lagrangian model simulations using ERA5 reanalysis leading to an improved understanding of the vertical structure of CO₂ in the monsoon region (Vogel et al., 2023). Here we use the same CO₂ reconstruction method as used in Vogel et al. (2023), but focus on the differences between the three data sets (ERA-Interim, ERA5 and ERA5 1° × 1°). In general, our results show that using ERA5 reanalysis yield a better agreement with aircraft measurements conducted over the Indian subcontinent in summer 2017 as ERA-Interim.

2 Measurements during the Asian summer monsoon 2017

In the frame of the StratoClim project funded by the European Commission, a measurement campaign using the Russian Geophysica high altitude research aircraft was conducted in Kathmandu (Nepal) in summer 2017 (see Fig. 1) to measure a

variety of trace gases and aerosol characteristics for the first time in the Asian monsoon anticyclone up to 20 km altitude (corresponding to ~ 55 hPa or ~ 475 K potential temperature) (Stroh and StratoClim-Team, 2023). The StratoClim measurements constitute a unique data set to characterise major processes which dominate particle and trace gas transport from the northern Indian subcontinent, one of the most polluted regions of the world, into the lower stratosphere.

CO_2 and N_2O were detected using the multi-tracer in situ instrument HAGAR (Volk et al., 2000; Homan et al., 2010) operated by the University of Wuppertal. Apart from CO_2 and N_2O , it also provides simultaneous in situ measurements of CH_4 , CFC-12, CFC-11, H-1211, SF_6 and H_2 . Except for CO_2 , which is measured at high time resolution (3 to 5 s) by non-dispersive infrared absorption (NDIR), all the other species were measured by gas chromatography with electron capture detection (GC/ECD) every 90 s. The instrument is calibrated every 7.5 min during flight with either of two standard gases, which are inter-calibrated in the laboratory with standards provided by NOAA GML. For StratoClim the accuracy of CO_2 was estimated to be about 0.2 ppm and about 2 ppb for N_2O (more details can be found in Vogel et al., 2023).

The long-lived trace gases C_2F_6 , HFC-125 and SF_6 were collected with the whole air sampler of Utrecht University operated on board the Geophysica research aircraft (e.g. Laube et al., 2010a). Ambient air was compressed into evacuated stainless-steel canisters (2 L) using a metal bellows pump that has been previously shown to not impact trace gas mixing ratios (e.g. Kaiser et al., 2006). The samples were transported to the University of East Anglia (UEA) for analysis on a high sensitivity gas chromatograph-trisector mass spectrometer system (Laube et al., 2010b). More details on the whole air sampler measurements during StratoClim and the used analytical technique can be found in Adcock et al. (2021).

120

3 Lagrangian transport simulations

3.1 CLaMS trajectory calculations

Back-trajectory calculations were performed using the trajectory module of the Chemical Lagrangian Model of the Stratosphere (CLaMS) (McKenna et al., 2002b, a; Pommrich et al., 2014, and references therein) which was developed with the aim to study transport and chemical processes in the atmosphere in the presence of strong tracer gradients. Here, CLaMS diabatic backward trajectories were started along the entire flight paths (every 1 second) of all StratoClim Geophysica flights conducted over the northeastern part of the Indian subcontinent. Overall ~ 110000 back-trajectories are calculated, between 9000 and 16000 per flight depending on the length of the flights.

For comparison, the back-trajectory calculations are driven by three data sets, two reanalyses in different resolutions provided by the European Centre for Medium-Range Weather Forecasts (ECMWF): ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$. The new ERA5 reanalysis (Hersbach et al., 2020) is a high-resolution atmospheric data set with 137 vertical levels up to 0.01 hPa, a horizontal resolution of ~ 31 km (T_L 639) and a hourly time resolution. We retrieved the data on a $0.3^\circ \times 0.3^\circ$ horizontal grid. The ECMWF's prior reanalysis ERA-Interim (Dee et al., 2011) has 60 vertical levels up to 0.01 hPa, a horizontal resolution of ~ 79 km (T_L 255) (corresponding to $1^\circ \times 1^\circ$ horizontal grid) and a 6-hourly time resolution.

135 Further, we use a version of ERA5 with lower resolution referred to as ‘ERA5 $1^\circ \times 1^\circ$ ’ (similar to Ploeger et al., 2021; Konopka et al., 2022; Clemens et al., 2023). ERA5 $1^\circ \times 1^\circ$ data are directly provided by the ECMWF on a $1^\circ \times 1^\circ$ horizontal grid after down-scaling the original data by truncation of the spherical harmonics representation to a $1^\circ \times 1^\circ$ horizontal grid (corresponding to T_L 255). In addition, the time resolution is down-sampled to every 6 hours, for better comparability with ERA-Interim. However, the vertical resolution is not changed and is the same as in the original ERA5 reanalysis. ERA5 $1^\circ \times 1^\circ$ data are a computing-time-saving alternative to the full resolution ERA5 data and are particularly suited for 3-dimensional global multi-annual CLaMS simulations.

In the CLaMS model, potential temperature is used as the vertical coordinate when the pressure is less than about 300 hPa, (i.e. in the upper troposphere and in the stratosphere); when the pressure is greater than about 300 hPa (more accurately, for pressure p exceeding a reference level of $p/p_{\text{surface}} = 0.3$), a pressure-based orography-following hybrid coordinate (in units of K) is used (Pommrich et al., 2014). In potential temperature levels above about 300 hPa, the vertical velocity (i.e. $d\Theta/dt$) is determined solely by the total heating rate (Pommrich et al., 2014; Ploeger et al., 2021). Total diabatic heating rates including clear-sky radiative heating, cloud radiation, latent heat release, as well as turbulent and diffusive heat transport for the upper troposphere and stratosphere are deduced from ECMWF reanalyses. However, also in the first layers below the reference level the vertical coordinate is still close to diabatic, as the transition from potential temperature to an orography-following vertical coordinate occurs rather slow (e.g. Pommrich et al., 2014). Therefore, the vertical velocity includes information on convective transport as resolved in the reanalysis vertical wind and total diabatic heating rate (see Pommrich et al., 2014).

Trajectories are considered ending in the model boundary layer referred to as ‘model BL’, when they are located for the first time below about 2–3 km above surface considering orography (i.e., the vertical hybrid pressure–potential–temperature coordinate (ζ) fulfils $\zeta \leq 120$ K) (details see e.g. Vogel et al., 2015, 2019).

155 The upward transport and convection in CLaMS (in both trajectory calculations as well as in three-dimensional simulations) depends on the employed reanalysis data (ERA-Interim, ERA5, ERA5 $1^\circ \times 1^\circ$) which differ strongly in the representation of convection (e.g. Hoffmann et al., 2019; Li et al., 2020; Clemens et al., 2023). The differences between ERA5 and ERA-Interim is attributed among other issues to the better spatial and temporal resolution of the ERA5 reanalysis, which allows for a better representation of convective updrafts. Therefore, in ERA-Interim convection over Asia is underestimated compared to ERA5. In our study no additional parametrisation for convection is used for the CLaMS simulations, only the convection already included in the reanalysis is considered.

3.2 Method for CO₂ reconstruction

Vogel et al. (2023) demonstrated that high-resolution CO₂ profiles measured in situ during the StratoClim campaign in summer 2017 reflect the seasonal variability of CO₂ at ground level. In addition, CO₂ is chemically inert in the troposphere and stratosphere and can be used as an age tracer considering time periods of several months (e.g. Boering et al., 1996; Andrews et al., 2001; Ray et al., 2022). Therefore a reasonable reconstruction of vertical CO₂ was conducted successfully using CLaMS back-trajectories driven by ERA5 reanalysis using ground-based CO₂ measurement (Vogel et al., 2023). Following the approach by Vogel et al. (2023), here we apply the same method for CO₂ reconstruction, however the differences between ERA5

compared to ERA-Interim and ERA5 $1^\circ \times 1^\circ$ will be analysed to infer possible differences in transport of air masses between
170 the three data sets.

The method for CO₂ reconstruction used in Vogel et al. (2023) is briefly summarised hereafter. CO₂ mixing ratios from
ground-based observations on the Indian subcontinent (Fig. 2) measured during the time when the CLaMS back-trajectories
reach the model BL are used for CO₂ reconstruction. For that purpose different CO₂ ground-based observations (all shown in
Fig. 3) available on different time scales (monthly, weekly or daily) were interpolated in time on a common daily grid to get a
175 CO₂ mixing ratio for every day from each used measurement site for the CO₂ reconstruction. These calculated CO₂ mixing
ratios define CO₂ in the model boundary layer and are transported passively along the trajectory to the location and time of the
Geophysica flight path, i.e. CO₂ is treated as chemically inert over the time of the back-trajectory calculation.

As a second step, a regional mask was developed where CO₂ is prescribed in the model BL depending on different geo-
graphical regions (see Fig. 3). In each of these geographical regions referred to as ‘BL region’ CO₂ is prescribed using one
180 specific measurement site, e.g. trajectories ending in the BL region marked in green and dark-red (roughly Indian Subconti-
nent and Tibetan Plateau) are prescribed using ground-based measurements from Nainital and the BL region marked in yellow
(roughly Bangladesh) is prescribed using ground-based measurements from Comilla. Unfortunately the coverage of ground-
based measurements of CO₂ over the Indian subcontinent in 2016 to 2017 is sparse, therefore only data from Nainital and
Comilla are available. Additional CO₂ ground-based time series from other geographical regions influencing the Geophysica
185 measurements provided by measurement sites for Greenhouse gases in Mt. Waliguan (WLG, China), Bukit Kototabang (BKT,
Indonesia), Mauna Loa (MLO, Hawaii) and Samoa (SMO, Cape Matatula) are used for CO₂ reconstruction (more details about
the used ground-based observations can be found in Vogel et al., 2023).

Ground-based CO₂ (provided by the World Data Centre for Greenhouse Gases (WDCGG), <https://gaw.kishou.go.jp>) mea-
sured in Mouna Loa (Hawaii) and in Cape Matatula (Samoa) (Thoning et al., 2021, <http://doi.org/10.7289/V5X0659V>) as well
190 as their average (black dashed-dotted line) are also shown in Fig. 2 as reference for the tropical background (e.g. Boering
et al., 1996; Andrews et al., 1999). The comparison of the different seasonal cycles of the ground-based CO₂ measurements
demonstrates that the seasonal CO₂ maximum over the Indian subcontinent during pre-monsoon is much larger than the CO₂
maximum of ground-based CO₂ of the tropical background.

The definition of the different model boundary layer regions are adjusted according to the available measurement sites. Case
195 studies with different regional masks defining the model boundary layer regions were performed and the regional mask was
developed according to the best agreement of reconstructed and measured vertical CO₂ profiles. Further, the local air mass
transport influencing Nainital is taken into account as explained in Vogel et al. (2023).

The seasonal variability of CO₂ over the northern Indian subcontinent (mean value between 20–30°N and 75–95°E) of the
lowest model level at 975 hPa of the GOSAT-L4B product (Matsunaga, T. and Maksyutov, S. (eds.), 2018) is shown in Fig. 2 for
200 comparison to ground-based CO₂ measurements. The GOSAT-L4B product is a model simulation using CO₂ surface fluxes
inferred from column-averaged satellite measurements (Maksyutov et al., 2013). The lowest model level of GOSAT-L4B is
closest to the inferred CO₂ surface fluxes and is not strongly influenced by the tracer transport of the underlying transport
model. GOSAT-L4B CO₂ over the northern Indian subcontinent has the same seasonality as ground-based CO₂ over Nainital,




Figure 2. Temporal variability of ground-based CO₂. The variability of ground-based CO₂ is shown at Nainital and Comilla (geographical positions see Fig. 1). In addition, the seasonal variability of CO₂ over the northern Indian subcontinent (mean value between 20–30°N and 75–95°E) of the lowest model level at 975 hPa of the GOSAT-L4B product for comparison to ground-based CO₂ measurements is shown. Further, ground-based CO₂ measured in Mouna Loa (Hawaii) and in Cape Matatula (Samoa) as well as their average (black dashed-dotted line) as reference for the tropical background are given. The pre-monsoon period (March–May) when a seasonal CO₂ maximum is expected is high-lighted (light-grey) as well as the period of the StratoClim aircraft campaign during monsoon 2017 (dark-grey).

however the minimum and maximum values differ strongly, highlighting the need of ground-based CO₂ measurements over
 205 the Indian subcontinent in addition to satellite-based estimation of CO₂ surface fluxes (a more detailed discussion can be found
 in Vogel et al., 2023).

3.3 Mean age of air from 3-dimensional CLaMS simulations

Trajectory-based transport times are compared to mean age of air from global 3-dimensional CLaMS chemistry transport model
 210 simulations performed over a time period of several decades to consider in addition the transport time of aged air, that is not
 considered in our back-trajectory calculations ending on 1 June 2016. Global 3-dimensional CLaMS simulations are based on
 3-dimensional forward trajectories and a parametrisation of small-scale mixing depending on the shear in the large-scale flow
 (e.g. Pommrich et al., 2014). The model simulations are driven with either ERA5 1° × 1° or ERA-Interim reanalysis winds and
 diabatic heating rates, as described in more detail by Ploeger et al. (2021). Similar as for the CLaMS trajectory calculations,
 215 convection resolved in the reanalysis vertical winds and total diabatic heating rates are used for the 3-dimensional CLaMS
 simulation (see Sect. 3.1). Apart from small-scale mixing, the vertical transport in CLaMS trajectory calculations and in the
 3-dimensional CLaMS simulations is treated in the same way.




Figure 3. Regional mask for CO₂ reconstruction using CO₂ ground-based measurements at different sites in Asia and the Pacific. In each model boundary layer (BL) region (marked by different colours) CO₂ is prescribed from one specific measurement site: tropical southern hemisphere (tSH) by Samoa (SMO), Indian subcontinent (India) by Nainital (NTL), Bangladesh (BGD) by Comilla (CLA), Tibetan Plateau (TIB) by Nainital (NTL), marine northern hemisphere (mNH) by Mauna Loa (MLO), continental northern hemisphere (cNH) by Mt. Waliguan (WLG) and Warm Pool region (Wpool) by Bukit Kototabang (BKT) (figure adapted from Vogel et al. (2023)).

Ploeger et al. (2021) performed global 3-dimensional CLaMS simulations to calculate the age of air spectrum, the distribution of transit times through the stratosphere, at each location in the stratosphere based on chemically inert pulse tracers. In our study, the globally calculated mean age of air by Ploeger et al. (2021) is interpolated along all Geophysica flights paths (F01-F08). Thus a direct comparison to the trajectory-based transport time is possible. In Ploeger et al. (2021), 60 different tracer pulses are released at the tropical surface (30°S-30°N), more specifically by a mixing ratio boundary condition in the lowest model layer. The chosen pulse frequency of 2 months allows a 2-month resolution of the age spectrum along the transit time axis for 10 years of transit time (more details in Ploeger et al., 2021). Mean age of air is calculated in three different ways to enable assessing uncertainties arising from the method. First, mean age is calculated as the first moment (mean) of the age spectrum, Second, this age spectrum-based mean age has been corrected for its finite tail (truncated to 10 years), by applying an exponential correction fit. Third, mean age has also been calculated from a clock-tracer with a linear increase in the entire lowest model layer. Due to the methodological differences, the age spectrum-based mean age is expected to yield the youngest estimate, the spectrum-based mean age including the tail correction yields the oldest estimate, and the clock-tracer mean age values lie in between. The range between these three different mean age estimates can be interpreted as an estimate for methodological uncertainties arising from the mean age calculation. Compared to the trajectory-based transport times, all three mean age estimates are expected to result in higher values, as they include the effects of mixing and recirculation of old stratospheric air into the tropics, which is absent in the pure trajectory calculations ending on 1 June 2016.

season	time period	start time	age of air
monsoon 2017	June–September 2017	1 June 2017	~ 2 months
pre-monsoon 2017	March–May 2017	1 March 2017	~ 2–5 months
winter 16/17	December 2016 – February 2017	1 Dec 2016	~ 5–8 months
post-monsoon 2016	October–November 2016	1 Oct 2016	~ 8–10 months
monsoon 2016	June–September 2016	1 June 2016	~ 10–14 months
aged air	older than 1 June 2016		> 14 months

Table 1. Time periods and trajectory-based age of air of considered seasons on Indian subcontinent. The analysis of CLaMS back-trajectories is performed back until the start time of each season. For each season, air parcels that were released at the model boundary layer (BL) are analysed. The longest simulation time is back until 1 June 2016 (~ one year). Air parcels that are located in the free atmosphere on 1 June 2016 are considered as aged air.

235 4 Results

CLaMS diabatic backward trajectories driven by three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$) were started along all Geophysica flight tracks (F01-F08) to infer a trajectory-based transport time from the location of the measurement back to the time when the back-trajectory reached the model BL. The trajectories are calculated back to 1 June 2016 and are analysed within different time periods to identify the source regions at the model BL depending on season (see Tab. 1). However, most
240 back-trajectories reach the model BL much later than 1 June 2017, which implies that air parcels probed during the Geophysica flights were released at the model BL much later than 1 June 2017, e.g. 64% (63%) of all air parcels are from the monsoon season 2017 using ERA5 (ERA-Interim) reanalysis.

The higher the sampled air parcels are located the longer are their simulated trajectory-based transport times which is to be expected. However, there is also a strong variability of transport times between individual air parcels at the same level of
245 potential temperature indicating mixing of air masses of different transport times (or different ages) as well as from different origins.

4.1 Transport times and mean age of air

The mean fraction of air from the model BL and the trajectory-based mean transport time are calculated from all backward trajectories started along the Geophysica flight tracks depending on flight height. Fig. 4 shows the trajectory-based mean
250 transport time (averaged as median in 2 K intervals) and the mean fraction of air from the model BL depending on potential temperature and accumulated back to the start times of monsoon 2017 and monsoon 2016 using the three data sets. To calculate the trajectory-based mean transport time only the fraction of trajectories from the model BL is considered, thus older air masses are neglected as first approximation.

All simulations show that considering a trajectory length back to the start time of monsoon 2017 yields a boundary layer
255 fraction between 80% and 100% below 370K; above the boundary layer fraction is decreasing rapidly and reaches 0% around




Figure 4. Mean fraction of air from the model BL (top) and the trajectory-based mean transport time (bottom) as calculated from all backward trajectories started along the Geophysica flight tracks averaged as median in 2 K intervals and accumulated back to the start times of monsoon 2017 (1 June 2017, dotted lines) and monsoon 2016 (1 June 2016, solid lines). The trajectory calculations are driven by three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$) indicated by different colours. The differences (Δ) between ERA-Interim and ERA5 and ERA5 $1^\circ \times 1^\circ$, respectively, are shown in the right panels.




Figure 5. Trajectory-based mean transport time back to monsoon 2016 (same as in Fig. 4a) and the clock-tracer mean age of air inferred from 3-dimensional CLaMS simulations both averaged as median in 2 K intervals. The trajectory calculations are driven by three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$). Mean age of air is only available for 3-dimensional CLaMS simulations driven by ERA-Interim and ERA5 $1^\circ \times 1^\circ$. The time differences (Δ Time) between clock-tracer mean age of air and trajectory-based mean transport time are shown in the right panel. Methodological differences in calculating mean age of air (Sect. 3.3) are indicated as shading (in light-blue and light-grey). The left envelope represents the age spectrum-based mean age and the right envelope the spectrum-based mean age including the tail correction (below 400 K the difference between the three methods is minor and therefore not shown).

420 K (Fig. 4a). The transition at ~ 370 K corresponds to the crossover level near 364 K found in the Asian summer monsoon 2017 by Legras and Bucci (2020). The crossover level marks the separation of descending and ascending motion and thus confirms that convection as represented in the reanalysis data is included in CLaMS backward trajectories. Our results show that a trajectory length of about two months is too short for a comprising simulation of the chemical composition of the Asian monsoon anticyclone because only very young air masses are considered.

Using a trajectory length back to the start time of monsoon 2016 (≈ 10 -14 months), a boundary layer fraction of almost 100% is reached up to 410 K. Above, the boundary layer fraction is slowly decreasing and depends strongly on the used ECMWF reanalysis and implicates different mean transport times at these altitudes (Fig. 4b).

Below 410 K convection in ERA5 yields faster transport times (up to ≈ 20 days) and higher model boundary layer (BL) fractions than in ERA-Interim. Vice versa, above 420 K air masses have faster transport times up to two months (≈ 60 days) from the model BL to the UTLS in ERA-Interim compared to ERA5 corresponding to a 20% higher fraction from the model BL. Transport times inferred from ERA5 $1^\circ \times 1^\circ$ trajectories have in principle a similar behaviour as transport times based on ERA5, however the variability is somewhat different, caused by the different temporal and horizontal resolution.

Considering in addition aged air (older than 1 June 2016) the mean age of air from a 3-dimensional CLaMS simulation (Sect. 3.3) interpolated along the flight tracks (and averaged as median in 2 K intervals) is compared to the trajectory-based

mean transport times calculated from pure back-trajectory calculations (Fig. 5). Mean age of air is only available for 3-dimensional CLaMS simulations driven by ERA-Interim and ERA5 $1^\circ \times 1^\circ$. The trajectory-based mean transport times of ERA5 and ERA5 $1^\circ \times 1^\circ$ in the lower stratosphere are very similar as shown in Fig. 4b, therefore we assume that also the mean age of air from 3-dimensional CLaMS simulations driven by ERA5 and ERA5 $1^\circ \times 1^\circ$ would be similar.

275 N_2O profiles measured during the StratoClim campaign indicate strong mixing with older stratospheric air above ~ 400 K (Fig. 6a; a more detailed discussion can be found in (Vogel et al., 2023)). Halon-1211 (which has a shorter live-time than N_2O) measurements aboard Geophysica (see Fig. 2 in Adcock et al., 2021) indicate that just below ~ 400 K a minor impact of older air is found. This is in agreement with the CLaMS backward-trajectory calculations, that results in a BL fraction of a few percent below 100% at these levels of potential temperature (Fig. 4a). Mixing with older stratospheric air above ~ 400 K
280 is evident in both the trajectory-based mean transport time as well as the mean age of air inferred from 3-dimensional CLaMS simulations.

However, there is a strong difference between the used ECMWF reanalyses as already found in trajectory-based mean transport times at potential temperatures higher than 410 K. ERA-Interim results in a mean age of about 2 years, while using ERA5 $1^\circ \times 1^\circ$ yields a mean age of more than 3 years at 470 K (Fig. 5). The differences from using ERA-Interim and ERA5 $1^\circ \times$
285 1° are much larger than from methodological differences to calculate the mean age of air (the clock-tracer mean age, age spectrum-based mean age and spectrum-based mean age including the tail correction; see Sect. 3.3).

Below 400 K potential temperature there is a difference between the trajectory-based mean transport time and the mean age of air of ≈ 80 days and ≈ 130 days using ERA-Interim or ERA5 $1^\circ \times 1^\circ$, respectively. Moreover, trajectory-based transport times are restricted to below about 1 year (1 June 2016), which basically excludes influence of downward transport from the
290 stratosphere. However due to the calculation of the mean value of the transport times of many single trajectories, this statistical treatment represents mixing between different air masses.

The 3-dimensional CLaMS simulations (see Sect. 3.3) used here to calculate the mean age of air also include parameterised small-scale mixing (dependent on the deformation rate in the large-scale flow) which causes an additional ageing of air compared to the pure trajectory calculations (e.g. Konopka et al., 2019). But also differences in the treatment of the lower model
295 boundary could cause the difference between trajectory-based mean transport time and the mean age of air from global CLaMS simulations below 400 K (see Fig. 5). The age of air tracer is released in the lowest model layer, whereas the trajectory-based mean transport time is related to the top of the model boundary layer ($\zeta=120$ K ~ 2 -3 km above surface following orography).

To validate clock-tracer mean age of air as well as trajectory-based transport times from CLaMS we use N_2O measured by
300 the HAGAR instrument during the StratoClim research flights. We compute mean age of air (Γ) from measured N_2O using $\Gamma - \text{N}_2\text{O}$ correlations by Andrews et al. (2001) and Engel et al. (2002) based on aircraft and balloon measurements. We use Eq. 3 by Andrews et al. (2001) derived for N_2O mixing ratios of the year 1997:

$$\Gamma = 0.0566 \times (313 - \text{N}_2\text{O}[1997]) - 0.000195 \times (313 - \text{N}_2\text{O}[1997])^2. \quad (1)$$

This $\Gamma - \text{N}_2\text{O}$ correlation is adapted to N_2O mixing ratios (in ppb) for the year 2017 as follows:

$$305 \quad \text{N}_2\text{O}[1997] = \text{N}_2\text{O}[2017] \times (313/335). \quad (2)$$

In addition, the mean age of air is calculated using a correlation by Engel et al. (2002) which is based on measurements from 1997 and 2000 and is also adapted to N_2O mixing ratios for the year 2017.

$$\Gamma = 6.03 - 0.0136 \times \text{N}_2\text{O}[1997] + 8.5892 \times 10^{-5} \times \text{N}_2\text{O}[1997]^2 - 3.376968 \times 10^{-7} \times \text{N}_2\text{O}[1997]^3 \quad (3)$$

Figure 6b shows the $\Gamma - \text{N}_2\text{O}$ correlations (valid above 375 K) from Andrews et al. (2001) and Engel et al. (2002) compared
310 to clock-tracer mean age of air derived from global 3-dimensional CLaMS simulations driven by ERA-Interim and ERA5 $1^\circ \times 1^\circ$
 1° reanalysis. In the Asian monsoon region, clock-tracer mean age of air based on ERA-Interim is lower than observation-based
estimates while mean age of air based on ERA5 $1^\circ \times 1^\circ$ is somewhat older, but a little closer to the observations. For N_2O
larger than ~ 310 ppb (between 380 K and 410 K) simulated mean age of air for both ERA5 $1^\circ \times 1^\circ$ as well as ERA-Interim is
somewhat older than the observation-based mean age of air, likely related to an underestimation of subgrid-scale convective
315 transport processes in the model (see Konopka et al., 2019, and discussion above).

Measured N_2O profiles indicate strong mixing with older stratospheric air only above ~ 400 K (Fig. 6a), therefore we can
also compare trajectory-based transport times with observation-based mean age of air below 400 K. In Fig. 6b, trajectory-based
mean transport times (back to 1 June 2016) for potential temperature levels between 375 K and 400 are added. At these alti-
tudes, a very good agreement between observation-based mean age of air and trajectory-based transport times is found using
320 both ERA-Interim as well as ERA5 to drive the trajectories. Therefore, CLaMS back-trajectories are very well suited for CO_2
reconstruction in particular below 400 K (Sect. 4.4). CLaMS mean age of air above 400 K will be further used for comparison
to observation-based mean age from HFC-125 and C_2F_6 which are used to derive ascent rates (Sect. 4.3) .

4.2 Air mass origin and its vertical propagation

325 For a better source attribution of the StratoClim aircraft measurements it is important to identify the source regions at the
model BL. During monsoon 2017 most air parcels were released in the northern part of the Indian subcontinent, the Tibetan
Plateau, Bay of Bengal and eastern China (Fig. 7), however the details differ between the three data sets. Using ERA-Interim
in general more marine sources are found in the western Pacific compared to ERA5 for monsoon 2017. A cluster of air parcels
at the model BL is found over the western Pacific caused by typhoon activity at $\sim 20^\circ\text{N}$ 125°E influencing research flight F08
330 (Fig. A1) using ERA5 reanalysis (details see (Stroh and StratoClim-Team, 2023)), whereas this typhoon signature is not found
using ERA-Interim and is only very weakly represented in ERA5 $1^\circ \times 1^\circ$. Due to the better representation of convection, a
slightly higher fraction of air is transported during monsoon 2017 from the model BL using ERA5 (64%) compared to ERA-
Interim (63%) and ERA5 $1^\circ \times 1^\circ$ (63%). The frequency distributions for each research flight (F01-F08) for monsoon 2017
using ERA5 are shown in the supplementary Fig. A1.

335 During pre-monsoon 2017 the origins are shifted towards the tropics to the northern Inter-Tropical Convergence Zone (ITCZ)
e.g. over the Indian Ocean and the western Pacific (see supplementary Fig. A2). For winter 16/17, the origins move further




Figure 6. Airborne N₂O measurements from the StratoClim campaign in Kathmandu (Nepal) during July and August 2017 (left). In addition, the mean WMO tropopause using ERA5 (Hoffmann and Spang, 2022) as well as the lowest and highest tropopause (grey dashed lines) over Kathmandu during the flight days are shown. Mean age versus N₂O from Andrews et al. (2001) and Engel et al. (2002) adapted to the year 2017 compared to clock-tracer mean age of air derived from global 3-dimensional CLaMS simulations driven by ERA-Interim and ERA5 1° × 1° reanalysis (right). Only N₂O measurements from the HAGAR instrument above 375 K potential temperature are shown. Further, trajectory-based transport times using ERA5 and ERA-Interim (back to 1 June 2016) are added for potential temperature levels between 375 K and 400 K.

to the south to the southern Inter-Tropical Convergence Zone (ITCZ) mostly over the Warm Pool region, northern Australia and western Pacific. The contributions from post-monsoon 2016 and monsoon 2016 are minor. Differences between ERA5 and ERA-Interim are obvious in transport time e.g. during pre-monsoon 2017 and winter 16/17, 25% of air is from model BL using
 340 ERA-Interim and only 20% using ERA5. Thus faster ERA-Interim vertical velocities in the UTLS (as already shown in Fig. 4) have an impact on the spatial distribution of the air mass origin in the model BL and yield differences between ERA-Interim and ERA5. During pre-monsoon 2017 the highest frequency distributions using ERA5 are found in the Bay of Bengal, whereas in ERA-Interim high fractions are found in continental Asia, Bay of Bengal and the tropical western Pacific.

Due to the different vertical velocities in ERA-Interim, ERA5 and ERA5 1° × 1° (Fig. 4), the propagation of air masses from
 345 different model BL regions into the lower stratosphere varies in the region of the Asian monsoon. To infer these differences the regional mask (Fig. 3) introduced in Sect. 3.2 is applied. Fig. 8 shows the fraction of air from the model BL, split into the BL regions (Fig. 3) as well as the fractions of the free atmosphere using three data sets. The fractions of air are accumulated back to starting times of different seasons: monsoon 2017 (a), pre-monsoon 2017 (b), winter 16/17 (c), post-monsoon 2016 (d), monsoon 2016 (e). The longer the trajectories the higher are the contributions from the model BL and the lower are the
 350 fractions from the free atmosphere. The quality of the reconstruction of CO₂ depends on the trajectory length, therefore it is

important to know the contributions from the model BL in each altitude. The trajectory length can be too short (and thus miss contributions from the model BL) or too long (resulting in higher uncertainties) (for a detailed discussion on this issue see Vogel et al., 2023).

355 Below 380 K, in general contribution from continental regions (Indian Subcontinent, Bangladesh, Tibetan Plateau and the continental northern hemisphere) are higher using ERA5 compared to ERA-Interim (Fig. 9). While using ERA-Interim, higher contributions from the marine northern hemisphere and the Warm Pool region are found in these levels of potential temperature. Between 380 K and 420 K, the contributions are vice versa and more marine sources (marine Northern Hemisphere, Warm Pool region) are found using ERA5 than using ERA-Interim. In particular above 430 K the impact of the tropical southern hemisphere is much stronger in ERA-Interim compared to ERA5 (Fig. 9).

360 Using ERA5 $1^\circ \times 1^\circ$ data to drive the CLaMS trajectories yields comparable results using ERA5 reanalysis (Figs. 8 and 9) and differences between ERA5 and ERA5 $1^\circ \times 1^\circ$ are in general much lower than 10 percentage points in contrast to differences between ERA5 and ERA-Interim (~ 20 percentage points).

Using cloud top altitudes from geostationary satellites to identify convection which occurred 30 days before the StratoClim measurements, Bucci et al. (2020) (see Fig. 10 therein) found that up to an altitude of 17 km (~ 400 K) convective sources 365 contribute more than 95% to the composition of the air probed during all flights. However, they calculate back-trajectories only back to cloud top altitudes. Nevertheless, in CLaMS trajectories only the convection inherent in the ERA5 and ERA-Interim reanalyses is included and therefore, small scale convection could be underestimated. Further, in Bucci et al. (2020) only very young air masses (younger than 30 d) are considered, therefore contributions from pre-monsoon 2017 and winter 16/17 are not covered. A more detailed comparison between the approach used in Bucci et al. (2020) and our analysis can be found elsewhere 370 (Stroh and StratoClim-Team, 2023).

4.3 Effective ascent rates and transport time distribution

From CLaMS trajectories effective ascent rates are calculated as difference in potential temperature along the backward trajectories for a time interval of 1 day and 20 days before the time of the aircraft measurements. The effective ascent rate is an integrated quantity, depends on time and is not an instantaneous ascent rate at a specific location in the atmosphere. The effective 375 ascent rates are calculated for the three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$; see Fig 10). Negative effective ascent rates reflect the descent of air masses just before the flight such as descending air from the lower stratosphere mixing into the air of the Asian monsoon anticyclone.

The effective ascent rates calculated over 24 h just before the aircraft measurements (1 day) reflect the short term evolution of the sampled air mass and can be impacted by recent convective events (e.g. Fig. 10b at 390 K) or stratospheric intrusions 380 i.e. mixing with older stratospheric air (Fig. 10b at ~ 415 K and ~ 435 K). Therefore, strong differences between the three data sets are found (Fig. 10a-c). Less impact of convection as well as of stratospheric intrusions are found in ERA-Interim and in ERA5 $1^\circ \times 1^\circ$. Below 360 K effective ascent rates over 1 day in ERA5 up to ~ 50 K/d, in ERA5 $1^\circ \times 1^\circ$ up to ~ 30 K/d and in ERA-Interim only up to ~ 20 K/d are found (not shown here).




Figure 7. Frequency distribution (number of trajectories normalised by the total number of trajectories started along the flight path) of the locations where air parcels were traced back to the model BL. Trajectories driven by ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$ reanalysis were started along the entire flight paths (every 1 second) of all eight Geophysica flights. The frequency distributions are shown for monsoon 2017 (a zoom of Asia marked as grey box is shown right beside). The frequency distribution is calculated in longitude-latitude bins of $2.0^\circ \times 1.5^\circ$. The percentages indicate the fraction of air parcels released at the model BL within monsoon 2017. In summary, 90% (93%) using ERA5 (ERA-Interim) of the air parcels were released at the model BL after 1 June 2016, the other 10% (7%) originates from aged air. The detailed pattern of the frequency distribution depends on the used reanalyses.




Figure 8. The fraction of air from the model boundary layer (BL) and the free atmosphere. The fraction from the model BL and from the free atmosphere is calculated from all backward trajectories started along the Geophysica flight tracks averaged in 2 K intervals and accumulated back to the start times of different seasons (rows), namely monsoon 2017, pre-monsoon 2017, winter 16/17, post-monsoon 2016, monsoon 2016 (detailed start times are listed in Tab. 1) and for three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$; columns). The fraction of air for monsoon 2016 (last row) referred to as the free atmosphere corresponds to the fraction of ‘aged’ defined in Tab. 1. The fraction of air from the model BL is divided in the different BL regions as shown in Fig. 3.




Figure 9. The difference of the fraction of air from the model boundary layer (BL) between back-trajectories driven by ERA5 and ERA-Interim (left) as well as between back-trajectories driven by ERA5 and ERA5 $1^\circ \times 1^\circ$ (right) depending on potential temperature. The difference is accumulated back to monsoon 2016 (1 June 2016) and fractions are averaged in 4 K intervals in contrast to Fig. 8 (last row; where 2 K intervals are used) to highlight the main impact of the BL sources (however the values of percentage points depends on the used averaging interval). The model boundary layer regions (Fig. 3) are summarised into three regions: continental (India, BGD, TIB, cNH) and marine regions (mNH, Wpool) mainly from the northern Hemisphere as well as the tropical Southern Hemisphere (tSH).

The mean effective ascent rates over 20 days (Fig. 10d-e) reflect a time-averaged ascent rate which is impacted by (vertical
 385 as well as horizontal) mixing of air masses of different origin and age. In general, the mean effective ascent rates inferred from ERA-Interim are higher compared to ERA5 and ERA5 $1^\circ \times 1^\circ$. To evaluate the mean effective ascent rates calculated from CLaMS back-trajectories, mean ascent rates from air samples collected with the whole air sampler (WAS) of Utrecht University are estimated using measurements of the long-lived trace gases HFC-125 and C_2F_6 . Both trace gases are chemically inert in the troposphere and stratosphere and have been demonstrated to be suitable to derive observation-based mean age ages of air
 390 as both have very long atmospheric lifetimes (HFC-125 >800 years (Leedham Elvidge et al., 2018) and $C_2F_6 \approx 10,000$ years (Worton et al., 2007)). As the concept of age of air inferred from measurements only works in the stratosphere, a reference level of 390 K (corresponding to a mean age of 0 as determined via polynomial fit functions, Figure 11) is used.

The observation-based mean age of air based on HFC-125 and C_2F_6 at 470 K is about ~ 2 -2.5 years (Fig. 11), clock-tracer mean age of air inferred from 3-dimensional CLaMS simulations driven by ERA-Interim is younger than 2 years and ~ 2 -
 395 3 years using ERA5 $1^\circ \times 1^\circ$ in this altitude (Fig. 11). Observation-based mean age of air inferred from HFC-125 and C_2F_6 is based on a reference level of 390 K, while clock-tracer mean age of air is based on Earth's surface. From trajectory-based transport times, a time lag of about 2-3 months between Earth's surface and 390 K can be estimated. Taken this time lag into account, mean age of air driven by ERA-Interim is too young at this altitude, whereas mean age of air from ERA5 $1^\circ \times 1^\circ$ is somewhat too old at 470 K. Further, observation-based mean age of air based on SF_6 is compared to observation-based mean
 400 age of air based on HFC-125 and C_2F_6 (Fig. 11), however observation-based mean age of air based on SF_6 is about half a year




Figure 10. Effective ascent rates calculated as difference in potential temperature along backward trajectories driven by three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$) to the time of the StratoClim measurements for a time interval of 1 day (top) and 20 days (bottom) and their mean values in 2 K intervals. The ascent rates are calculated for all trajectories calculated for research flights F01-F08. Negative effective ascent rates reflect the descent of air masses.




Figure 11. Observation-based mean age of air (left) and observation-based mean ascent rates above 390 K (right) derived from trace gas measurements of air samples collected with the whole air sampler (WAS) of Utrecht University during the eight StratoClim research flights over the Indian subcontinent in summer 2017. Note that negative observation-based mean age of air (< -0.1 year) found below 390 K are not shown. In addition, clock-tracer mean age of air for each air sample is shown derived from global 3-dimensional CLaMS simulations driven by the ERA-Interim and ERA5 $1^\circ \times 1^\circ$ reanalysis (right).

older at 470 K than from HFC-125 and C₂F₆ caused by SF₆ sources in Asia (Adcock et al., 2021). Therefore, SF₆ is a rather unsuitable chemical age tracer for the Asian monsoon region.

The mean ascent rate for each air sample is simply derived from dividing the potential temperature difference to the reference level of 390 K by the age of air derived from the two tracers (Figure 11). The mean age of air reflects an integrated three-dimensional transit time, impacted by vertical and horizontal transport as well as mixing processes. These processes are likely to be increasingly influential the further away the air is from the 390 K reference surface, with a clear tendency to increase the observation-based mean age of air because of the horizontal transport (in-mixing) of aged stratospheric air. In the trajectory-based mean effective ascent for 20 days vertical and horizontal transport as well as mixing processes are included and therefore comparable with observation-based ascent rates.

Between 390 and 430 K, there is a variability of the mean effective ascent rates derived from HFC-125 and C₂F₆ from 0.2 up to 2.3 K/d. Above 430 K, the observation-based ascent rate converges to ~ 0.2 K/d. Here, the mean effective ascent rate derived from ERA5 (as well as ERA5 $1^\circ \times 1^\circ$) back-trajectories over a time interval of 20 days (~ 0.2 - 0.3 K/d) is in good agreement with observation-based mean ascent rates derived from air samples collected by the whole air sampler. Mean effective ascent rates derived from ERA-Interim back-trajectories are much faster (≈ 0.5 K/d) above 430 K.

Our analysis agrees with previous studies, that found consistently that in general the vertical velocities in ERA-Interim are too fast in the tropics (Dee et al., 2011; Ploeger et al., 2012; Schoeberl et al., 2012). In addition, our findings show that the mean effective ascent rates of ~ 0.2 - 0.3 K/d derived from ERA5 in the region of the Asian monsoon in the lower stratosphere (430 K-480 K) agree very well with observation-based mean ascent rates derived from long-lived trace gases.




Figure 12. Normalised frequency distribution of the transport time from 360 K (\approx the level of maximum convective outflow) to the location of the aircraft measurement along the CLaMS backward trajectories (referred to as ‘transport time distribution’) using the three data sets. The transport time distribution is shown for different levels of potential temperature (for 2 K intervals) for a time resolution of 5 and 10 days. In the legend, the transport time to the maximum peak for each level of potential temperature is given.

We show that the ascent rates along CLaMS backward trajectories depend on the used ECMWF reanalyses. However they
 420 further depend on the considered altitude range. Thus, in general ERA-Interim in the UTLS is faster than ERA5. However, due to a better representation of convection, air masses can be uplifted faster as well as up to higher levels of potential temperature by convection when using the ERA5 reanalysis compared to ERA-Interim. These differences have an impact on the frequency distribution of the transport time of backward trajectories from the main convective outflow to the sample region (referred to as ‘transport time distribution’) on different levels of potential temperature using the three data sets.

425 Transport time distributions with a time resolution of 5 and 10 days using the three data sets (Fig. 12) reflect the faster vertical velocities found in the UTLS using ERA-Interim, thus the maximum peak of the age spectrum is in general at shorter transport times compared to ERA5 and ERA5 $1^\circ \times 1^\circ$. However, at lower potential temperatures the impact of convection which is different in the used reanalyses has to be taken into account. Thus a better representation of convection (visible in the two peaks at 380 K for a time resolution of 5 days; black line) and slower vertical velocities found in ERA5 in the UTLS can result in a
 430 spectrum peak at similar transit times to the maximum peak as a coarser resolution of convection and faster vertical velocities




Figure 13. Airborne CO₂ measurements from the StratoClim campaign in Kathmandu (Nepal) during July and August 2017. Each air parcel is coloured by the trajectory-based transport time from the model boundary layer (BL) to the time of measurements inferred by Lagrangian back-trajectory calculations driven by ERA-Interim (a). Air parcels located in the model BL are not shown. Aged air (air located in the free atmosphere on 1 June 2016) is marked in black. Further, the differences of the trajectory-based transport time between ERA-Interim and ERA5 (b) are shown from back-trajectories reaching the model BL. In addition, the mean WMO tropopause (Hoffmann and Spang, 2022) as well as the lowest and highest tropopause (grey dashed lines) over Kathmandu during the aircraft campaign (27 July - 10 August 2017) are added using ERA-Interim (a) and ERA5 (b) reanalysis.

found in ERA-Interim. Therefore, the maximum peaks for 390 K and 400 K using ERA-Interim are at similar transport times.

4.4 Reconstruction of CO₂ from airborne measurements

Airborne CO₂ measurements from the StratoClim campaign in Kathmandu (Nepal) during July and August 2017 are shown in
 435 Fig. 13. Each air parcel is coloured by the trajectory-based transport time from the model BL to the time of measurements inferred by Lagrangian back-trajectory calculations driven by ERA-Interim (Fig. 13a). Trajectory-based transport times increase with the altitude of sampled air parcels (as already shown in Fig. 4). However, there is also a strong variability of transport times between individual air parcels at the same level of potential temperature indicating mixing of air masses of different transport times or of different age (Fig. 13a). Moreover, differences in transport times of individual air parcels are found using
 440 instead of ERA-Interim ERA5 (Fig. 13b) reanalyses. In the stratosphere ERA-Interim has the tendency to be faster (shorter

transport times) than ERA5 (bluish data points). In the UTLS, certain air masses are found experienced faster upward transport by convection using ERA5 than using ERA-Interim (reddish points).

In addition, differences in the tropopause height (in particular for the local minimum and maximum) exist using ERA5 and ERA-Interim. Hoffmann and Spang (2022) found that the standard deviation of the tropical tropopause height are \approx 30-50% higher in ERA5 (T_L 639; high-resolution version) compared to ERA-Interim, mostly related to explicitly resolved gravity waves in ERA5, which are absent in ERA-Interim due to its coarser spatial resolution. Tegtmeier et al. (2020) attributed tropopause shifts between ERA-Interim and ERA5 to higher vertical resolution of ERA5, having 3 times more levels in the tropical tropopause than ERA-Interim.

For a reliable reconstruction of measured vertical CO_2 profiles over the entire altitude range, both accurate back-trajectory calculations as well as precise CO_2 concentrations at the ground are required. For the latter purpose, a regional mask was developed where CO_2 is prescribed in the model BL depending on different BL regions (Fig. 3).

To reconstruct vertical profiles of trace gases in the region of the Asian monsoon up to 410 K potential temperature the fraction of air from the model BL has to be $\sim 100\%$, otherwise mixing with aged air has to be taken into account. In Sect. 4.1, it was shown that using a trajectory length back to the start time of monsoon 2016 (≈ 10 -14 months), a boundary layer fraction of about 100% is reached up to 410 K. Above 410 K mixing with older air masses successively occurred and the fraction from the model BL rapidly decreases.

Figure 14 (top) shows reconstructed CO_2 using three data sets for back-trajectory calculations until 1 June 2016 neglecting the contributions from the free atmosphere (aged air). The comparison with measured in situ CO_2 profiles shows a good overall agreement from the model BL up to ~ 410 K for ERA5 trajectories. A reconstruction using ERA-Interim shows a stronger dispersion between 390 K and 420 K, because here ERA-Interim vertical velocities are faster than ERA5. Further, below 370 K the measured variability of CO_2 caused by convection (low CO_2 at ≈ 360 K) is better reproduced using ERA5 due to the better representation of convection. A CO_2 reconstruction using trajectories driven by ERA5 $1^\circ \times 1^\circ$ is somewhere between ERA-Interim and ERA5.

Above ~ 410 K, aged air has to be taken into account as discussed in Sect. 4.1. Figure 14 (bottom) shows reconstructed CO_2 , but using in addition GOSAT-L4B CO_2 data for the fraction of aged air. For back-trajectories ending in the free atmosphere CO_2 is reconstructed from GOSAT-L4B data that are providing CO_2 values up to 10 hPa (for details see Vogel et al., 2023). Here, for each 1 K interval the median of all air parcels considering both the fraction from the model BL as well as from the aged air is calculated. This approach allows the mixing of air at the top of the Asian monsoon anticyclone between air from the boundary layer and air from (stratospheric) background to be considered.

Caused by too fast vertical velocity in the UTLS in ERA-Interim (according to effective ascent rates inferred from whole air sampler measurements) higher CO_2 from the model BL is found in the lower stratosphere. Even including the contribution of aged air from GOSAT-L4B data yields slightly higher reconstructed CO_2 above 420 K using ERA-Interim compared to the measurements. Using ERA5 as well as ERA5 $1^\circ \times 1^\circ$ yield slightly lower reconstructed CO_2 above 420 K. However, above 410 K the quality of the GOSAT-L4B needs also to be taken into account for an assessment of the quality of CO_2 reconstruction. GOSAT-L4B data depend on CO_2 fluxes at the Earth's surface (GOSAT-L4A data), on model resolution as well as on vertical

transport in the used atmospheric transport model, which could have a too fast transport in the lower stratosphere, similar as ERA-Interim.

In summary, there are differences in CO₂ reconstruction using the three data sets, whereby the statistical variability of the CO₂ reconstruction is in the range of the measurements. A CO₂ reconstruction using ERA5 agrees best with the measured
480 vertical CO₂ profile up to 410 K, although there is only a slight difference when using ERA-Interim reanalysis. It should be noted that the used CO₂ reconstruction technique has limitations because of the very low number of sites measuring ground-based CO₂ over the Indian subcontinent in 2016-2107. The UTLS is a very sensitive region regarding the interplay between deep convection and vertical velocities in the lower stratosphere influencing the vertical transport of CO₂ and thus the CO₂ reconstruction using trajectory calculations.

485

5 Conclusions

It was reported previously that because of a better spatial and temporal resolution, the ERA5 reanalysis yields a better representation of convection than the predecessor ERA-Interim (e.g. Hoffmann et al., 2019; Li et al., 2020; Legras and Bucci, 2020; Malakar et al., 2020). Further it was shown that the vertical transport in ERA-Interim is too fast in the tropical UTLS
490 (Dee et al., 2011; Ploeger et al., 2012; Schoeberl et al., 2012; Tegtmeier and Krüger, 2022). At higher northern-hemispheric stratospheric levels above the tropical tropopause layer, it was reported that ERA5 transport is likely too slow (Ploeger et al., 2021). In general, our findings confirm these results, however in our study, we focus in detail on the Asian summer monsoon region.

Differences in transport of air in the region of the Asian summer monsoon 2017 were inferred using the Chemical Lagrangian
495 Model of the Stratosphere (CLaMS) driven by three data sets, namely two ECMWF reanalyses in different resolution (ERA-Interim, ERA5 and ERA5 1° × 1°). The model results were assessed using unique airborne measurements up to ~20 km (~475 K) during the Asian summer monsoon 2017 conducted with the Geophysica aircraft during the StratoClim campaign in Nepal (Stroh and StratoClim-Team, 2023). CLaMS diabatic backward trajectories were calculated for all Geophysica research flights (F01-F08) performed over the Indian subcontinent. Trajectory-based transport times, origin of air at the Earth's surface,
500 mean effective ascent rates, transport time distributions as well as mean age of air from 3-dimensional CLaMS simulations were compared using the three data sets.

Below 410 K convection as represented in ERA5 yields faster trajectory-based upward transport (up to ≈ 20 days) than ERA-Interim. On the other hand, air masses above 420 K show up to two months (≈ 60 days) shorter trajectory-based transport times from the model BL to the UTLS in ERA-Interim compared to ERA5. A better representation of convection and slower vertical
505 velocities above the convection found in ERA5 can yield similar transport times to the UTLS (~380 K–390 K) as a coarser resolution of convection and faster vertical velocities as found in ERA-Interim. Therefore, the frequency distribution of the transport time from the level of maximum convective outflow (~360 K) to different levels of the aircraft measurement based on back-trajectories (denoted as transport time distribution) are very sensitive to the three data sets.




Figure 14. Reconstructed CO_2 using back-trajectory calculations until 1 June 2016 driven by three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$) compared to HAGAR CO_2 airborne-measurements. Reconstructed CO_2 is shown using the regional mask shown in Fig. 3 for the fraction of trajectories ending in the model BL driven by the three data sets (top). Reconstructed CO_2 is shown as median calculated from all trajectories until 1 June 2016 in 1 K intervals for research flights F01-F08. Reconstructed CO_2 is shown using in addition GOSAT-L4B CO_2 data for the fraction of trajectories ending in the free atmosphere, mainly from stratospheric background (bottom). Bars indicate the range between the 25 and 75 percentile.

Below 380 K, contributions from continental regions (Indian Subcontinent, Bangladesh, Tibetan Plateau and the continental northern hemisphere) to air masses along the flight paths of all 8 local research flights (F01–F08) are higher using ERA5 compared to ERA-Interim while in ERA-Interim higher contributions from the marine northern hemisphere and the Warm Pool region are found. Although using ERA-Interim for back-trajectory calculations in general more marine sources at these altitudes are identified, the signal from typhoon activity in the western Pacific (in particular during research flight F08 on 10 August 2017) is not visible when using ERA-Interim. Above 380 K, it is the other way round and more marine sources are found using ERA5 and a stronger impact of the tropical southern hemisphere is found using ERA-Interim.

Above 430 K, the mean effective ascent rates derived from ERA5 back-trajectories over a time interval of 20 days ($\approx 0.2\text{--}0.3$ K/day) are in good agreement with the observation-based mean ascent rates inferred from long-lived trace gases such as C_2F_6 and HFC-125 derived from air samples collected by the whole air sampler aboard Geophysica. Mean effective ascent rates derived from ERA-Interim back-trajectories are much faster ≈ 0.5 K/day than observation-based mean ascent rates at these altitudes. Caused by the difference in the mean effective ascent rates when using two ECMWF reanalyses a different mean age of air is calculated at higher altitudes. At 470 K, a mean age of air of younger than 2 years is calculated using 3-dimensional CLaMS simulations driven by ERA-Interim, while using ERA5 $1^\circ \times 1^\circ$ a mean age between 2 and 3 years is calculated (a 3-dimensional CLaMS simulation driven by ERA5 is not yet available). At these altitudes, observation-based age of air from C_2F_6 and HFC-125 is up to $\sim 2\text{--}2.5$ years. In the monsoon region above 430K, the mean age of air using ERA-Interim mean age is in general too young, while ERA5 $1^\circ \times 1^\circ$ is somewhat too old, but closer to observation-based age of air derived from N_2O than using ERA-Interim. Thus uncertainties regarding the correct simulation of mean age of air in the lower stratosphere (430 K – 480 K) still remain.

Trajectory-based transport times inferred from ERA5 $1^\circ \times 1^\circ$ trajectories have in principle a similar behaviour as the times obtained using ERA5, however the variability is somewhat different, caused by the reduced temporal and horizontal resolution in ERA5 $1^\circ \times 1^\circ$. Details in simulated transport e.g. air mass origin, impact of tropical cyclones, transport time distribution and the vertical dispersion are different when employing ERA5 and ERA5 $1^\circ \times 1^\circ$. For long-term simulations over several years or decades ERA5 $1^\circ \times 1^\circ$ appear to be an acceptable (computing-time-saving) data set, however for detailed transport calculations in the Asian monsoon region (e.g analysing aircraft or balloon measurements) the full resolution ERA5 reanalysis resolve more small-scale features and variability.

Further, high-resolution CO_2 profiles measured aboard Geophysica were reconstructed using ground-based measurements of CO_2 mainly from Nainital (northern India) by Lagrangian model simulations using three data sets (ERA-Interim, ERA5 and ERA5 $1^\circ \times 1^\circ$) leading to an improved understanding of the vertical structure of CO_2 in the monsoon region. A reliable reconstruction (simulation) of vertical CO_2 profiles during the Asian monsoon is a challenge for model simulations because the seasonal variability of CO_2 at the ground, mixing with aged stratospheric air as well as the vertical velocities (including convection as well as vertical ascent caused by diabatic heating in the UTLS) have to be simulated accurately. Our analysis shows that by using the ERA5 reanalysis for CO_2 reconstruction a slightly better agreement with high-resolution in situ aircraft CO_2 measurements is obtained compared to ERA-Interim. However at higher altitudes (above 410 K), uncertainties remain in the used reconstruction approach, mainly caused by the limitations of the GOSAT-L4B CO_2 data used for characterising aged

stratospheric air, demonstrating the need for better global CO₂ simulations. Further, a sufficiently dense coverage of continuous
545 quality-controlled ground-based monitoring of CO₂ over the Indian subcontinent is a prerequisite for a reliable simulation of
vertical CO₂ profiles in the region of the Asian summer monsoon.

Code and data availability. The StratoClim data can be downloaded from the HALO database at <https://halo-db.pa.op.dlr.de/mission/101>.
For more details on the measurements please contact C. Michael Volk (M.Volk@uni-wuppertal.de) for HAGAR N₂O and CO₂ and Jo-
hannes Laube (j.laube@fz-juelich.de) for C₂F₆ and HFC-125 whole air sampler measurements. Ground-based CO₂ from Nainital and
550 Comilla were provided through National Institute for Environmental Research (NIES) available under Terao et al. (2022a, b). Ground-
based CO₂ measurements from other sites can be downloaded from the World Data Centre for Greenhouse Gases (WDCGG) (<https://gaw.kishou.go.jp>) and GOSAT-L4B CO₂ data under (https://data2.gosat.nies.go.jp/index_en.html). The ERA-Interim and ERA5 tropopause
data are available under Hoffmann, Lars; Spang, Reinhold, 2021, "Reanalysis Tropopause Data Repository", <https://datapub.fz-juelich.de/slcs/tropopause>. The CLaMS trajectory code is available on a GitLab server at <https://jugit.fz-juelich.de/clams/CLaMS>.




Figure A1. Frequency distribution (fd) of the air mass origins at the model boundary layer (BL) for each research flight F01-F08 using ERA5 reanalysis for monsoon season 2017.

555 Appendix A: Supplement: Air mass origin and trajectory-based transport time

Author contributions. FS was leading the organisation and coordination of the StratoClim aircraft campaign. CMV, JW and VL were responsible for the measurements and analysis of airborne CO₂ profiles. JL was responsible for the observation-based mean age as well as ascent rates and FP for age of air from 3-dimensional CLaMS simulations. LH provided tropopause altitudes. FP, GG, JC, JG and LH helped with provisioning of ECMWF reanalyses. CLaMS trajectory calculations and CO₂ reconstruction were performed by BV. The study was conceived by BV, CMV and RM, whereby the results were discussed by all co-authors. The paper was written by BV with contributions from all co-authors.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

Acknowledgements. The authors are indebted to many local institutions, authorities, as well as individuals for making the StratoClim aircraft field campaign a success. We are especially grateful to the Nepalese, Indian, and Bangladeshi authorities for granting clearances as




Figure A2. Frequency distribution (fd) of the air mass origins at the model boundary layer (BL) similar as Fig. 7, but for pre-monsoon 2017, winter 16/17, post-monsoon 2016 and monsoon 2016. The percentages indicate the fraction of air parcels released at the model BL within a certain season. The detailed patterns of the frequency distribution depend strongly on the considered season as well as on used three data sets (ERA-Interim, ERA5 and ERA5 1° × 1°).




Figure A3. Airborne CO₂ measurements from the StratoClim campaign in Kathmandu (Nepal) and trajectory-based transport time for each research flight F01–F08 using ERA5 reanalysis (similar as Fig. 13, but for single flights). Each air parcel is coloured by the trajectory-based transport time from the model boundary layer (BL) to the time of measurements. Air parcels located in the model BL are not shown. Aged air (air located in the free atmosphere on 1 June 2016) is marked in black. In addition, the mean WMO tropopause (Hoffmann and Spang, 2022) as well as the lowest and highest tropopause (grey dashed lines) over Kathmandu on the day of the measurement are shown inferred from ERA5.

565 well as the Kathmandu airport authorities for their local support. Strong support by several local science partners is highly appreciated. We
thank the Geophysica aircraft crews and pilots. The European Commission has granted and funded the StratoClim project within Framework
Programme 7 under ENV.2013.6.1-2, Grant agreement No. 603557. The HAGAR operations and data analysis was supported by Thorben
Beckert from University Wuppertal and were partly funded by the German Helmholtz Association within the Helmholtz-CAS Joint Re-
search Group No. 307. The Nainital and Comilla measurements were performed by Manish Naja from Aryabhata Research Institute of
570 Observational Sciences, Md. Kawser Ahmed from University of Dhaka, and Shohei Nomura, Toshinobu Machida, Motoki Sasakawa Hi-
toshi Mukai from NIES, and supported by the Environment Research and Technology Development Fund (grant nos. JPMEERF20152002,
20182002 and 21S20800) of the Environmental Restoration and Conservation Agency of Japan. Further, the authors gratefully acknowledge
the World Data Centre for Greenhouse Gases (WDCGG) for providing CO₂ ground-based measurements in particular Yong Zhang from
China Meteorological Administration, Beijing, China; Kirk Thoning, Pieter Tans, Ed Dlugokencky and Xin Lan from Earth System Research
575 Laboratory (NOAA), Boulder, US. Further, we would like to thank the Japan Aerospace Exploration Agency (JAXA), the National Institute
for Environmental Studies (NIES), and the Ministry of the Environment (MOE) for providing the GOSAT L4B-data product in particular
Shamil Maksyutov. We thank the European Centre for Medium-Range Weather Forecasts (ECMWF) for providing the ERA-Interim and the
ERA5 reanalyses and the Jülich Supercomputing Centre (JSC; Research Centre Jülich, Germany) for computing time on the supercomputer
JUWELS (project CLaMS-ESM) and for storage resources on the meteocloud data archive. JCL received funding from the ERC (EXC3ITE-
580 678904-ERC-2015-STG). Finally, we acknowledge our colleagues from IEK-7 (Research Centre Jülich) Mohamadou Diallo, Paul Konopka,
Nicole Spelten and Nicole Thomas for support and discussion.

References

- Adcock, K. E., Fraser, P. J., Hall, B. D., Langenfelds, R. L., Lee, G., Montzka, S. A., Oram, D. E., Röckmann, T., Stroh, F., Sturges, W. T., Vogel, B., and Laube, J. C.: Aircraft-Based Observations of Ozone-Depleting Substances in the Upper Troposphere and Lower Stratosphere in and Above the Asian Summer Monsoon, *J. Geophys. Res.*, 126, e2020JD033 137, <https://doi.org/https://doi.org/10.1029/2020JD033137>, 2021.
- Andrews, A. E., Boering, K. A., Daube, B. C., Wofsy, S. C., Hints, E. J., Weinstock, E. M., and Bui, T. B.: Empirical age spectra for the lower tropical stratosphere from in situ observations of CO₂: Implications for stratospheric transport, *J. Geophys. Res.*, 104, 26.581–26.595, 1999.
- Andrews, A. E., Boering, K. A., Daube, B. C., Wofsy, S. C., Loewenstein, M., H., Podolske, J. R., Webster, C. R., Herman, R. L., Scott, D. C., Flesch, G. J., Moyer, E. J., Elkins, J. W., Dutton, G. S., Hurst, D. F., Moore, F. L., Ray, E. A., Romashkin, P. A., and Strahan, S. E.: Mean age of stratospheric air derived from in situ observations of CO₂, CH₄ and N₂O, *J. Geophys. Res.*, 106, 32 295–32 314, 2001.
- Appel, O., Köllner, F., Dragoneas, A., Hünig, A., Molleker, S., Schlager, H., Mahnke, C., Weigel, R., Port, M., Schulz, C., Drewnick, F., Vogel, B., Stroh, F., and Borrmann, S.: Chemical analysis of the Asian tropopause aerosol layer (ATAL) with emphasis on secondary aerosol particles using aircraft-based in situ aerosol mass spectrometry, *Atmos. Chem. Phys.*, 22, 13 607–13 630, <https://doi.org/10.5194/acp-22-13607-2022>, 2022.
- Bergman, J. W., Fierli, F., Jensen, E. J., Honomichl, S., and Pan, L. L.: Boundary layer sources for the Asian anticyclone: Regional contributions to a vertical conduit, *J. Geophys. Res.*, 118, 2560–2575, <https://doi.org/10.1002/jgrd.50142>, 2013.
- Bian, J., Li, D., Bai, Z., Li, Q., Lyu, D., and Zhou, X.: Transport of Asian surface pollutants to the global stratosphere from the Tibetan Plateau region during the Asian summer monsoon, *Natl. Sci. Rev.*, 7, 516–533, <https://doi.org/10.1093/nsr/nwaa005>, 2020.
- Boering, K. A., Wofsy, S. C., Daube, B. C., Schneider, H. R., Loewenstein, M., Podolske, J. R., and Conway, T. J.: Stratospheric Mean Ages and transport rates from observations of carbon dioxide and nitrous oxide, *Science*, 274, 1340–1343, 1996.
- Brinkop, S. and Jöckel, P.: ATTILA 4.0: Lagrangian advective and convective transport of passive tracers within the ECHAM5/MESy (2.53.0) chemistry–climate model, *Geosci. Model Dev.*, 12, 1991–2008, <https://doi.org/10.5194/gmd-12-1991-2019>, 2019.
- Brunamonti, S., Jorge, T., Oelsner, P., Hanumanthu, S., Singh, B. B., Kumar, K. R., Sonbawne, S., Meier, S., Singh, D., Wienhold, F. G., Luo, B. P., Boettcher, M., Poltera, Y., Jauhiainen, H., Kayastha, R., Karmacharya, J., Dirksen, R., Naja, M., Rex, M., Fadnavis, S., and Peter, T.: Balloon-borne measurements of temperature, water vapor, ozone and aerosol backscatter on the southern slopes of the Himalayas during StratoClim 2016–2017, *Atmos. Chem. Phys.*, 18, 15 937–15 957, <https://doi.org/10.5194/acp-18-15937-2018>, 2018.
- Bucci, S., Legras, B., Sellitto, P., D’Amato, F., Viciani, S., Montori, A., Chiarugi, A., Ravegnani, F., Ulanovsky, A., Cairo, F., and Stroh, F.: Deep-convective influence on the upper troposphere–lower stratosphere composition in the Asian monsoon anticyclone region: 2017 StratoClim campaign results, *Atmos. Chem. Phys.*, 20, 12 193–12 210, <https://doi.org/10.5194/acp-20-12193-2020>, 2020.
- Clemens, J., Vogel, B., Hoffmann, L., Griessbach, S., Thomas, N., Fadnavis, S., Müller, R., Peter, T., and Ploeger, F.: Identification of source regions of the Asian Tropopause Aerosol Layer on the Indian subcontinent in August 2016, *EGU sphere*, 2023, 1–39, <https://doi.org/10.5194/egusphere-2022-1462>, accepted, 2023.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Mor-

- crette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 620 Engel, A., Strunk, M., Müller, M., Haase, H., Poss, C., Levin, I., and Schmidt, U.: Temporal development of total chlorine in the high-latitude stratosphere based on reference distributions of mean age derived from CO_2 and SF_6 , *J. Geophys. Res.*, 107, <https://doi.org/10.1029/2001JD000584>, 2002.
- Fadnavis, S., Müller, R., Kalita, G., Rowlinson, M., Rap, A., Li, J.-L. F., Gasparini, B., and Laakso, A.: The impact of recent changes in Asian anthropogenic emissions of SO_2 on sulfate loading in the upper troposphere and lower stratosphere and the associated radiative changes, *Atmos. Chem. Phys.*, 19, 9989–10 008, <https://doi.org/10.5194/acp-19-9989-2019>, 2019.
- 625 Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Hauck, J., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Bakker, D. C. E., Canadell, J. G., Ciais, P., Jackson, R. B., Anthoni, P., Barbero, L., Bastos, A., Bastrikov, V., Becker, M., Bopp, L., Buitenhuis, E., Chandra, N., Chevallier, F., Chini, L. P., Currie, K. I., Feely, R. A., Gehlen, M., Gilfillan, D., Gkritzalis, T., Goll, D. S., Gruber, N., Gutekunst, S., Harris, I., Haverd, V., Houghton, R. A., Hurtt, G., Ilyina, T., Jain, A. K., Joetzjer, E., Kaplan, J. O., Kato, E., Klein Goldewijk, K., Korsbakken, J. I., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Marland, G., McGuire, P. C., Melton, J. R., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Neill, C., Omar, A. M., Ono, T., Peregón, A., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Séférian, R., Schwinger, J., Smith, N., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F. N., van der Werf, G. R., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2019, *Earth System Science Data*, 11, 1783–1838, <https://doi.org/10.5194/essd-11-1783-2019>, 2019.
- 630 Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, O., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Schwingshackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth System Science Data*, 14, 1917–2005, <https://doi.org/10.5194/essd-14-1917-2022>, 2022.
- 645 Hanumanthu, S., Vogel, B., Müller, R., Brunamonti, S., Fadnavis, S., Li, D., Ölsner, P., Naja, M., Singh, B. B., Kumar, K. R., Sonbawne, S., Jauhiainen, H., Vömel, H., Luo, B., Jorge, T., Wienhold, F. G., Dirksen, R., and Peter, T.: Strong day-to-day variability of the Asian Tropopause Aerosol Layer (ATAL) in August 2016 at the Himalayan foothills, *Atmos. Chem. Phys.*, 20, 14 273–14 302, <https://doi.org/10.5194/acp-20-14273-2020>, 2020.
- 650 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.

- 655 Hoffmann, L. and Spang, R.: An assessment of tropopause characteristics of the ERA5 and ERA-Interim meteorological reanalyses, *Atmos. Chem. Phys.*, 22, 4019–4046, <https://doi.org/10.5194/acp-22-4019-2022>, 2022.
- Hoffmann, L., Günther, G., Li, D., Stein, O., Wu, X., Griessbach, S., Heng, Y., Konopka, P., Müller, R., Vogel, B., and Wright, J. S.: From ERA-Interim to ERA5: the considerable impact of ECMWF's next-generation reanalysis on Lagrangian transport simulations, *Atmos. Chem. Phys.*, 19, 3097–3124, <https://doi.org/10.5194/acp-19-3097-2019>, 2019.
- 660 Hoffmann, L., Baumeister, P. F., Cai, Z., Clemens, J., Griessbach, S., Günther, G., Heng, Y., Liu, M., Haghighi Mood, K., Stein, O., Thomas, N., Vogel, B., Wu, X., and Zou, L.: Massive-Parallel Trajectory Calculations version 2.2 (MPTRAC-2.2): Lagrangian transport simulations on graphics processing units (GPUs), *Geosci. Model Dev.*, 15, 2731–2762, <https://doi.org/10.5194/gmd-15-2731-2022>, 2022.
- Hoffmann, L., Konopka, P., Clemens, J., and Vogel, B.: Lagrangian transport simulations using the extreme convection parameterization: an assessment for the ECMWF reanalyses, *Atmos. Chem. Phys.*, 23, 7589–7609, <https://doi.org/10.5194/acp-23-7589-2023>, 2023.
- 665 Homan, C. D., Volk, C. M., Kuhn, A. C., W., A., Baehr, J., Viciani, S., Ulanovski, A., and Ravegnani, F.: Tracer measurements in the tropical tropopause layer during the AMMA/SCOUT-O3 aircraft campaign, *Atmos. Chem. Phys.*, 10, 3615–3627, <https://doi.org/10.5194/acp-10-3615-2010>, 2010.
- Kaiser, J., Engel, A., Borchers, R., and Roeckmann, T.: Probing stratospheric transport and chemistry with new balloon and aircraft observations of the meridional and vertical N₂O isotope distribution, *Atmos. Chem. Phys.*, 6, 3535–3556, [https://doi.org/10.5194/acp-6-3535-](https://doi.org/10.5194/acp-6-3535-2006)
670 2006, 2006.
- Konopka, P., Tao, M., Ploeger, F., Diallo, M., and Riese, M.: Tropospheric mixing and parametrization of unresolved convective updrafts as implemented in the Chemical Lagrangian Model of the Stratosphere (CLaMS v2.0), *Geosci. Model Dev.*, 12, 2441–2462, <https://doi.org/10.5194/gmd-12-2441-2019>, 2019.
- Konopka, P., Tao, M., von Hobe, M., Hoffmann, L., Kloss, C., Ravegnani, F., Volk, C. M., Lauther, V., Zahn, A., Hoor, P., and Ploeger, F.:
675 Tropospheric transport and unresolved convection: numerical experiments with CLaMS-2.0/MESSy, *Geosci. Model Dev.*, 15, 7471–7487, <https://doi.org/10.5194/gmd-15-7471-2022>, 2022.
- Laube, J. C., Engel, a., Bönisch, H., Möbius, T., Sturges, W. T., Braß, M., and Röckmann, T.: Fractional release factors of long-lived halogenated organic compounds in the tropical stratosphere, *Atmos. Chem. Phys.*, 10, 1093–1103, [https://doi.org/10.5194/acp-10-1093-](https://doi.org/10.5194/acp-10-1093-2010)
2010, 2010a.
- 680 Laube, J. C., Martinerie, P., Witrant, E., Blunier, T., Schwander, J., Brenninkmeijer, C. A. M., Schuck, T. J., Bolder, M., Röckmann, T., van der Veen, C., Bönisch, H., Engel, A., Mills, G. P., Newland, M. J., Oram, D. E., Reeves, C. E., and Sturges, W. T.: Accelerating growth of HFC-227ea (1,1,1,2,3,3,3-heptafluoropropane) in the atmosphere, *Atmos. Chem. Phys.*, 10, 5903–5910, <https://doi.org/10.5194/acp-10-5903-2010>, 2010b.
- Lauther, V., Vogel, B., Wintel, J., Rau, A., Hoor, P., Bense, V., Müller, R., and Volk, C. M.: In situ observations of CH₂Cl₂ and CHCl₃
685 show efficient transport pathways for very short-lived species into the lower stratosphere via the Asian and the North American summer monsoon, *Atmos. Chem. Phys.*, 22, 2049–2077, <https://doi.org/10.5194/acp-22-2049-2022>, 2022.
- Leedham Elvidge, E., Bönisch, H., Brenninkmeijer, C. A. M., Engel, A., Fraser, P. J., Gallacher, E., Langenfelds, R., Mühle, J., Oram, D. E., Ray, E. A., Ridley, A. R., Röckmann, T., Sturges, W. T., Weiss, R. F., and Laube, J. C.: Evaluation of stratospheric age of air from CF₄, C₂F₆, C₃F₈, CHF₃, HFC-125, HFC-227ea and SF₆; implications for the calculations
690 of halocarbon lifetimes, fractional release factors and ozone depletion potentials, *Atmos. Chem. Phys.*, 18, 3369–3385, <https://doi.org/10.5194/acp-18-3369-2018>, 2018.

- Legras, B. and Bucci, S.: Confinement of air in the Asian monsoon anticyclone and pathways of convective air to the stratosphere during the summer season, *Atmos. Chem. Phys.*, 20, 11 045–11 064, <https://doi.org/10.5194/acp-20-11045-2020>, 2020.
- 695 Li, D., Vogel, B., Müller, R., Bian, J., Günther, G., Ploeger, F., Li, Q., Zhang, J., Bai, Z., Vömel, H., and Riese, M.: Dehydration and low ozone in the tropopause layer over the Asian monsoon caused by tropical cyclones: Lagrangian transport calculations using ERA-Interim and ERA5 reanalysis data, *Atmos. Chem. Phys.*, 20, 4133–4152, <https://doi.org/10.5194/acp-20-4133-2020>, 2020.
- Maksyutov, S., Takagi, H., Valsala, V. K., Saito, M., Oda, T., Saeki, T., Belikov, D. A., Saito, R., Ito, A., Yoshida, Y., Morino, I., Uchino, O., Andres, R. J., and Yokota, T.: Regional CO₂ flux estimates for 2009–2010 based on GOSAT and ground-based CO₂ observations, *Atmos. Chem. Phys.*, 13, 9351–9373, <https://doi.org/10.5194/acp-13-9351-2013>, 2013.
- 700 Malakar, P., Kesarkar, A., Bhate, J., Singh, V., and Deshamukhya, A.: Comparison of Reanalysis Data Sets to Comprehend the Evolution of Tropical Cyclones Over North Indian Ocean, *Earth and Space Science*, 7, e2019EA000978, <https://doi.org/https://doi.org/10.1029/2019EA000978>, 2020.
- Mason, R. B. and Anderson, C. E.: The development and decay of the 100-mb. summertime anticyclone over southern Asia, *Monthly Weather Review*, 91, 3–12, [https://doi.org/10.1175/1520-0493\(1963\)0912.3.CO;2](https://doi.org/10.1175/1520-0493(1963)0912.3.CO;2), 1963.
- 705 Matsunaga, T. and Maksyutov, S. (eds.): A Guidebook on the Use of Satellite Greenhouse Gases Observation Data to Evaluate and Improve Greenhouse Gas Emission Inventories, Satellite Observation Center, National Institute for Environmental Studies, Japan, 2018.
- McKenna, D. S., Groß, J.-U., Günther, G., Konopka, P., Müller, R., Carver, G., and Sasano, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 2. Formulation of chemistry scheme and initialization, *J. Geophys. Res.*, 107, 4256, <https://doi.org/10.1029/2000JD000113>, 2002a.
- 710 McKenna, D. S., Konopka, P., Groß, J.-U., Günther, G., Müller, R., Spang, R., Offermann, D., and Orsolini, Y.: A new Chemical Lagrangian Model of the Stratosphere (CLaMS): 1. Formulation of advection and mixing, *J. Geophys. Res.*, 107, 4309, <https://doi.org/10.1029/2000JD000114>, 2002b.
- Müller, S., Hoor, P., Bozem, H., Gute, E., Vogel, B., Zahn, A., Bönisch, H., Keber, T., Krämer, M., Rolf, C., Riese, M., Schlager, H., and Engel, A.: Impact of the Asian monsoon on the extratropical lower stratosphere: trace gas observations during TACTS over Europe 2012, *Atmos. Chem. Phys.*, 16, 10 573–10 589, <https://doi.org/10.5194/acp-16-10573-2016>, 2016.
- 715 Pan, L. L., Honomichl, S. B., Kinnison, D. E., Abalos, M., Randel, W. J., Bergman, J. W., and Bian, J.: Transport of chemical tracers from the boundary layer to stratosphere associated with the dynamics of the Asian summer monsoon, *Journal of Geophysical Research: Atmospheres*, 121, 14,159–14,174, <https://doi.org/10.1002/2016JD025616>, 2016.
- Park, M., Randel, W. J., Gettleman, A., Massie, S. T., and Jiang, J. H.: Transport above the Asian summer monsoon anticyclone inferred from Aura Microwave Limb Sounder tracers, *J. Geophys. Res.*, 112, D16309, <https://doi.org/10.1029/2006JD008294>, 2007.
- 720 Ploeger, F., Konopka, P., Müller, R., Fueglistaler, S., Schmidt, T., Manners, J. C., Groß, J.-U., Günther, G., Forster, P. M., and Riese, M.: Horizontal transport affecting trace gas seasonality in the Tropical Tropopause Layer (TTL), *J. Geophys. Res.*, 117, D09303, <https://doi.org/10.1029/2011JD017267>, 2012.
- Ploeger, F., Günther, G., Konopka, P., Fueglistaler, S., Müller, R., Hoppe, C., Kunz, A., Spang, R., Groß, J.-U., and Riese, M.: Horizontal water vapor transport in the lower stratosphere from subtropics to high latitudes during boreal summer, *J. Geophys. Res.*, 118, 8111–8127, <https://doi.org/10.1002/jgrd.50636>, 2013.
- 725 Ploeger, F., Konopka, P., Walker, K., and Riese, M.: Quantifying pollution transport from the Asian monsoon anticyclone into the lower stratosphere, *Atmos. Chem. Phys.*, 17, 7055–7066, <https://doi.org/10.5194/acp-17-7055-2017>, 2017.

- Ploeger, F., Legras, B., Charlesworth, E., Yan, X., Diallo, M., Konopka, P., Birner, T., Tao, M., Engel, A., and Riese, M.: How robust are stratospheric age of air trends from different reanalyses?, *Atmos. Chem. Phys.*, 19, 6085–6105, <https://doi.org/10.5194/gmd-12-2441-2019>, 2019.
- Ploeger, F., Diallo, M., Charlesworth, E., Konopka, P., Legras, B., Laube, J. C., Grooß, J.-U., Günther, G., Engel, A., and Riese, M.: The stratospheric Brewer–Dobson circulation inferred from age of air in the ERA5 reanalysis, *Atmos. Chem. Phys.*, 21, 8393–8412, <https://doi.org/10.5194/acp-21-8393-2021>, 2021.
- Pommrich, R., Müller, R., Grooß, J.-U., Konopka, P., Ploeger, F., Vogel, B., Tao, M., Hoppe, C. M., Günther, G., Spelten, N., Hoffmann, L., Pumphrey, H.-C., Viciani, S., D’Amato, F., Volk, C. M., Hoor, P., Schlager, H., and Riese, M.: Tropical troposphere to stratosphere transport of carbon monoxide and long-lived trace species in the Chemical Lagrangian Model of the Stratosphere (CLaMS), *Geosci. Model Dev.*, 7, 2895–2916, <https://doi.org/10.5194/gmd-7-2895-2014>, 2014.
- Randel, W. J. and Park, M.: Deep convective influence on the Asian summer monsoon anticyclone and associated tracer variability observed with Atmospheric Infrared Sounder (AIRS), *J. Geophys. Res.*, 111, D12314, <https://doi.org/10.1029/2005JD006490>, 2006.
- Ray, E. A., Atlas, E. L., Schauffler, S., Chelpon, S., Pan, L., Bönisch, H., and Rosenlof, K. H.: Age spectra and other transport diagnostics in the North American monsoon UTLS from SEAC⁴RS in situ trace gas measurements, *Atmos. Chem. Phys.*, 22, 6539–6558, <https://doi.org/10.5194/acp-22-6539-2022>, 2022.
- Riese, M., Ploeger, F., Rap, A., Vogel, B., Konopka, P., Dameris, M., and Forster, P.: Impact of uncertainties in atmospheric mixing on simulated UTLS composition and related radiative effects, *J. Geophys. Res.*, 117, D16305, <https://doi.org/10.1029/2012JD017751>, 2012.
- Rolf, C., Vogel, B., Hoor, P., Afchine, A., Günther, G., Krämer, M., Müller, R., Müller, S., Spelten, N., and Riese, M.: Water vapor increase in the lower stratosphere of the Northern Hemisphere due to the Asian monsoon anticyclone observed during the TACTS/ESMVal campaigns, *Atmospheric Chemistry and Physics*, 18, 2973–2983, <https://doi.org/10.5194/acp-18-2973-2018>, 2018.
- Schoeberl, M. R., Dessler, A. E., and Wang, T.: Simulation of stratospheric water vapor and trends using three reanalyses, *Atmos. Chem. Phys.*, 12, 6475–6487, <https://doi.org/10.5194/acp-12-6475-2012>, 2012.
- Stenke, A., Dameris, M., Grewe, V., and Garny, H.: Implications of Lagrangian transport for simulations with a coupled chemistry-climate model, *Atmos. Chem. Phys.*, 9, 5489–5504, <http://www.atmos-chem-phys.net/9/5489/2009/>, 2009.
- Stroh, F. and StratoClim-Team: First detailed airborne and balloon measurements of microphysical, dynamical, and chemical processes in the Asian Summer Monsoon Anticyclone: overview and selected results of the 2016/2017 StratoClim field campaigns, *Atmos. Chem. Phys.*, to be submitted, 2023.
- Tao, M., Konopka, P., Ploeger, F., Yan, X., Wright, J. S., Diallo, M., Fueglistaler, S., and Riese, M.: Multitimescale variations in modeled stratospheric water vapor derived from three modern reanalysis products, *Atmos. Chem. Phys.*, 19, 6509–6534, <https://doi.org/10.5194/acp-19-6509-2019>, 2019.
- Tegtmeier, S. and Krüger, K.: Tropical Tropopause Layer, in: SPARC, 2022: SPARC Reanalysis Intercomparison Project (S-RIP) Final Report, edited by Fujiwara, M., Manney, G. L., Gray, L. J., and Wright, J. S., SPARC Report No. 10, chap. 08, WCRP-6/2021, <https://doi.org/https://doi.org/10.17874/800dee57d13>, 2022.
- Tegtmeier, S., Anstey, J., Davis, S., Dragani, R., Harada, Y., Ivanciu, I., Pilch Kedzierski, R., Krüger, K., Legras, B., Long, C., Wang, J. S., Wargan, K., and Wright, J. S.: Temperature and tropopause characteristics from reanalyses data in the tropical tropopause layer, *Atmos. Chem. Phys.*, 20, 753–770, <https://doi.org/10.5194/acp-20-753-2020>, 2020.
- Terao, Y., Nomura, S., Mukai, H., Machida, T., Sasakawa, M., Ahmed, M. K., and Patra, P. K.: Atmospheric carbon dioxide dry air mole fraction at Comilla, Bangladesh, NIES, <https://doi.org/10.17595/20220301.002>, ver.2022.0 (Last updated: 2022/03/01), 2022a.

- Terao, Y., Nomura, S., Mukai, H., Machida, T., Sasakawa, M., and Naja, M.: Atmospheric carbon dioxide dry air mole fraction at Nainital, India, NIES, <https://doi.org/10.17595/20220301.001>, ver.2022.0 (Last updated: 2022/03/01), 2022b.
- 770 Thoning, K., Crotwell, A., and Mund, J.: Atmospheric Carbon Dioxide Dry Air Mole Fractions from continuous measurements at Mauna Loa, Hawaii, Barrow, Alaska, American Samoa and South Pole. 1973-2020, Tech. rep., Version 2021-08-09 National Oceanic and Atmospheric Administration (NOAA), Global Monitoring Laboratory (GML), Boulder, Colorado, USA, <https://doi.org/10.15138/yaf1-bk21>, 2021.
- Vernier, J. P., Fairlie, T. D., Natarajan, M., Wienhold, F. G., Bian, J., Martinsson, B. G., Crumeyrolle, S., Thomason, L. W., and Bedka, K. M.: Increase in upper tropospheric and lower stratospheric aerosol levels and its potential connection with Asian pollution, *J. Geophys. Res.*, 120, 1608–1619, <https://doi.org/10.1002/2014JD022372>, 2015.
- 775 Vogel, B., Günther, G., Müller, R., Groß, J.-U., and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and of the extratropical lowermost stratosphere, *Atmos. Chem. Phys.*, 15, 13 699–13 716, <https://doi.org/10.5194/acp-15-13699-2015>, 2015.
- Vogel, B., Günther, G., Müller, R., Groß, J.-U., Afchine, A., Bozem, H., Hoor, P., Krämer, M., Müller, S., Riese, M., Rolf, C., Spelten, N., Stiller, G. P., Ungermann, J., and Zahn, A.: Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012, *Atmos. Chem. Phys.*, 16, 15 301–15 325, <https://doi.org/10.5194/acp-16-15301-2016>, 2016.
- 780 Vogel, B., Müller, R., Günther, G., Spang, R., Hanumanthu, S., Li, D., Riese, M., and Stiller, G. P.: Lagrangian simulations of the transport of young air masses to the top of the Asian monsoon anticyclone and into the tropical pipe, *Atmos. Chem. Phys.*, 19, 6007–6034, <https://doi.org/10.5194/acp-19-6007-2019>, 2019.
- 785 Vogel, B., Volk, C. M., Wintel, J., Lauther, V., Müller, R., Patra, P. K., Riese, M., Terao, Y., and Stroh, F.: Reconstructing high-resolution in-situ vertical carbon dioxide profiles in the sparsely monitored Asian monsoon region, *Commun Earth Environ*, 4, <https://doi.org/10.1038/s43247-023-00725-5>, 2023.
- Volk, C., Riediger, O., Strunk, M., Schmidt, U., Ravegnani, F., Ulanovsky, A., and Rudakovand, V.: In situ tracer measurements in the tropical tropopause region during APE-THESEO, in: *Eur. Comm. Air Pollut. Res. Report*, vol. 73, pp. 661–664, 2000.
- 790 von Hobe, M., Ploeger, F., Konopka, P., Kloss, C., Ulanowski, A., Yushkov, V., Ravegnani, F., Volk, C. M., Pan, L. L., Honomichl, S. B., Tilmes, S., Kinnison, D. E., Garcia, R. R., and Wright, J. S.: Upward transport into and within the Asian monsoon anticyclone as inferred from StratoClim trace gas observations, *Atmos. Chem. Phys.*, 21, 1267–1285, <https://doi.org/10.5194/acp-21-1267-2021>, 2021.
- Wohltmann, I., Lehmann, R., Gottwald, G. A., Peters, K., Protat, A., Louf, V., Williams, C., Feng, W., and Rex, M.: A Lagrangian convective transport scheme including a simulation of the time air parcels spend in updrafts (LaConTra v1.0), *Geosci. Model Dev.*, 12, 4387–4407, <https://doi.org/10.5194/gmd-12-4387-2019>, 2019.
- 795 Worton, D. R., Sturges, W. T., Gohar, L. K., Shine, K. P., Martinerie, P., Oram, D. E., Humphrey, S. P., Begley, P., Gunn, L., Barnola, J.-M., Schwander, J., and Mulvaney, R.: Atmospheric Trends and Radiative Forcings of CF₄ and C₂F₆ Inferred from Firn Air, *Environmental Science & Technology*, 41, 2184–2189, <https://doi.org/10.1021/es061710t>, 2007.
- Yu, P., Rosenlof, K. H., Liu, S., Telg, H., Thornberry, T. D., Rollins, A. W., Portmann, R. W., Bai, Z., Ray, E. A., Duan, Y., Pan, L. L., 800 Toon, O. B., Bian, J., and Gao, R.-S.: Efficient transport of tropospheric aerosol into the stratosphere via the Asian summer monsoon anticyclone, *Proc. Natl. Acad. Sci.*, 114, 6972–6977, <https://doi.org/10.1073/pnas.1701170114>, 2017.