



Empirical upscaling of OzFlux eddy covariance for high-resolution monitoring of terrestrial carbon uptake in Australia.

Chad. A. Burton¹, Luigi. J. Renzullo^{2,1}, Sami. W. Rifai³, and Albert. I.J.M. Van Dijk¹

¹ Fenner School of Environment & Society, Australian National University, Canberra, ACT, Australia.

² Bureau of Meteorology, Canberra, ACT, Australia

³ School of Biological Sciences, The University of Adelaide, Adelaide SA, Australia.

Correspondence to: Chad Burton (chad.burton@anu.edu.au)

Abstract. We develop high resolution (1 km) estimates of Gross Primary Productivity (GPP), Ecosystem Respiration (ER) and Net Ecosystem Exchange (NEE) over the Australian continent for the period January 2003 to June 2022 by empirical upscaling of flux tower measurements. We compare our estimates with nine other products that cover the three broad categories that define current methods for estimating the terrestrial carbon cycle and assess if consistencies between datasets can point to the correct dynamics of Australia's carbon cycle. Our results indicate that regional empirical upscaling greatly improves upon the existing global empirical upscaling efforts, outperforms process-based models, and agrees much better with the dynamics of CO₂ flux over Australia as estimated by two regional atmospheric inversions. Our nearly 20-year estimates of terrestrial carbon fluxes revealed Australia is a strong net carbon sink of -0.44 (IQR=0.42) PgC/year on-average, with an inter-annual variability of 0.18 PgC/year and an average seasonal amplitude of 0.85 PgC/yr. Annual mean carbon uptake estimated from other methods ranged considerably, while carbon flux anomalies showed much better agreement between methods. NEE anomalies were predominately driven by cumulative rainfall deficits and surpluses, resulting in larger anomalous responses from GPP over ER. In contrast, we show that the long-term average seasonal cycle is dictated more by the variability in ER than GPP, resulting in peak carbon uptake typically occurring during the cooler, drier Austral autumn, and winter months. This new estimate of Australia's terrestrial carbon cycle provides a benchmark for assessment against Land Surface Model simulations, and a means for monitoring of Australia's terrestrial carbon cycle at an unprecedented high-resolution. We call this new estimate of Australia's terrestrial carbon cycle, "AusEFlux" (Australian Empirical Fluxes).

1. Introduction

The global terrestrial biosphere has acted as a net carbon sink, absorbing approximately 29 % of anthropogenic CO₂ emissions each year and thereby mitigating impacts from global warming (Friedlingstein et al., 2022). Australia's vast semi-arid ecosystems play a large and critical role in controlling the inter-annual variability (IAV) of the global terrestrial carbon sink, and are therefore of crucial importance to understand if we are to make reliable predictions about the fate of the global



carbon cycle under a warming climate (Ahlström et al., 2015; Chen et al., 2017; Ma et al., 2016; Poulter et al., 2014; Metz et al., 2023). However, uncertainties in the methods used for quantifying components of the terrestrial biosphere preclude
35 definitive inferences about the magnitude of Australia's terrestrial carbon sink, the seasonal and inter-annual oscillations, and the drivers of change in carbon flux variability.

Several methods exist to quantify the spatio-temporal dynamics of the terrestrial carbon cycle. Dynamic Global Vegetation Models (DGVMs) and Land Surface Models (LSMs) simulate responses of vegetation to changes in climate by parameterising ecological processes but are limited by several uncertainties that relate to their parametrisations and limited
40 inclusion of key ecological processes (Kowalczyk et al., 2006; Li et al., 2021; Quillet et al., 2010). Uncertainties in these models can lead to large differences in land carbon flux estimates, even where similar models are used (Teckentrup et al., 2021). For example, over a 19-year period from 2003 to 2021, the Community Atmosphere Biosphere Land Exchange (CABLE) model extracted from TRENDY v10 estimates Australia's annual mean GPP to be 3.01 PgC/yr (Friedlingstein et al., 2022) while a regionally forced CABLE run using a similar model configuration estimates GPP to be greater than 50 %
45 higher at 4.58 PgC/yr (Villalobos et al., 2022).

Atmospheric inversion methods, which rely upon atmospheric CO₂ measurements and an atmospheric transport model, provide a semi-empirical method for quantifying aspects of the carbon cycle, but their capacity to spatially resolve CO₂ fluxes is severely constrained by the sparse observational network of measuring sites (51 sites globally, with only four locations in Australia) (Rödenbeck et al., 2018). Satellite-based remote sensing of atmospheric CO₂ has become possible
50 using the Greenhouse Gas Observing Satellite (GOSAT) and the Orbiting Carbon Observatory (OCO-2 and OCO-3) satellites (Basu et al., 2013; Eldering et al., 2017). This allows for spatially comprehensive monitoring of CO₂ sources and sinks over continental to global scales. Several global inversion studies have incorporated these datasets, but results over Australia have been contradictory (Basu et al., 2013; Chevallier et al., 2014; Detmers et al., 2015). Villalobos et al. (2022) conducted a regional atmospheric inversion over Australia assimilated with OCO-2 data to infer a gridded estimate (~81 km
55 cells) of NEE for 2015-2019. They found Australia was a strong annual carbon sink (-0.47 PgC/yr) on average, and that peak carbon uptake occurred during the cooler, drier months of the austral winter. Similarly, using an atmospheric inversion of GOSAT satellite measurements, Metz et al. (2023) found that Australia's seasonal CO₂ flux variability coincided with the onset of rainfall after the dry season, leading to CO₂ flux releases during the October-December period, and carbon uptake occurring during the drier March-September period. These studies provided valuable insight into the dynamics of
60 Australia's terrestrial carbon cycle, but their very coarse spatial resolution prevents these approaches from resolving spatially detailed estimates of Australia's carbon cycle.

A third approach relies on data-driven machine learning (ML) methods to upscale eddy covariance (EC) micrometeorological tower data from global networks of long-term carbon and water flux measurement sites. This approach has the advantage of relying on a denser network of empirical observations than the atmospheric inversion approaches (for
65 example, the popular FLUXNET2015 dataset contains 206 sites (Pastorello et al., 2020)). Another advantage of data-driven ML approaches is their ability to accurately model highly nonlinear relationships to explanatory variables, as is common in

complex environmental systems. Nevertheless, the results of global empirical upscaling products, most notably FLUXCOM (Jung et al., 2020; Tramontana et al., 2016), are prone to several limitations, including: significantly underestimating the magnitude of the IAV of carbon fluxes, an inability to resolve carbon flux trends (e.g. from CO₂ fertilisation), and overestimating the size of the tropical carbon sink (Jung et al., 2020). Over Australia, two FLUXCOM products: ‘FLUXCOM-Met’ and FLUXCOM-RS’, show substantially different mean annual NEE fluxes of -0.23 and -0.05 PgC/yr, respectively (averaged over the period 2003-2015). Furthermore, the annual mean GPP and ER components show a > 60 % difference in magnitude between the two products. IAV of NEE, as estimated by one standard deviation in the fluxes, is also subdued compared with estimates from LSMs and atmospheric inversions.

This lack of agreement between the different approaches to quantifying Australia’s land carbon sinks and sources calls into question how well constrained the magnitudes, IAV, temporal trends, and spatial allocations of Australia’s land carbon fluxes are. Here we explore the potential for empirical upscaling of the regional “OzFlux” eddy covariance network (Isaac et al., 2017) to better characterise Australia’s terrestrial carbon cycle. Models built on global datasets (and with a strong northern hemisphere bias) will necessarily need to generalise across vastly different climates, ecosystem types, and plant functional traits, limiting their ability to accurately represent ecosystem dynamics in regions where ecosystem responses do not conform to the dominant dynamics in the global dataset. This may especially be the case in Australia where extreme climate variability and evolutionary isolation have created sclerophyllous, evergreen, woody species that do not fit into standard globally predominant plant functional types used by LSMs (Beringer et al., 2016; Beringer et al., 2022; Williams and Woinarski, 1997). Furthermore, Australia’s data record of EC flux tower measurements has grown substantially in the intervening years since the inception of the commonly used FLUXNET2015 training dataset. For example, the FLUXCOM product included data from only four EC flux towers over Australia (~43 site-years of data), and the current FLUXNET2015 dataset contains 23 sites equating to ~115 site years of Australian data. Contrast this with the full OzFlux dataset over Australia which, as of January 2022, contains 33 sites and 238 site-years of data. These later years of EC flux tower measurements since 2015 are especially valuable given they have recorded a period of extreme climate variability in Australia such as the historic drought from 2017-2019 (Devanand et al., 2023 in review), culminating in the Black Summer bushfires (Byrne et al., 2021), and the subsequent triple La Niña with record breaking rainfall in eastern Australia from 2020-2023. A further advantage of upscaling fluxes at a regional scale is the ability to take advantage of higher-resolution input datasets than is tractable at the global scale, both due to the unavailability and uncertainty of global high-resolution datasets and the computational constraints that attend global upscaling.

Our objectives for this study are as follows:

- Develop an accurate, high-resolution (~1 km) empirical upscaling of Net Ecosystem Exchange (NEE), Ecosystem Respiration (ER), and Gross Primary Productivity (GPP) for Australia covering the period January 2003 to June 2022.
- Evaluate our empirical upscaling of Australian flux data in comparison with LSM, inversion-derived, and global empirical upscaling estimates of the carbon cycle with the aim of identifying consistencies between datasets that may point to the correct dynamics of Australia’s terrestrial carbon cycle.



- Assess if the upscaling approach can offer new insights into Australia’s carbon cycle, and/or affirm if the upscaling can replicate known biogeochemical controls on the carbon cycle.

2. Data & Methods

105 2.1 Data

2.1.1 CO₂ flux tower data

We used monthly fluxes of NEE, GPP, and ER produced by the OzFlux regional network of eddy covariance flux towers. These data are processed to Level 6 and are freely accessible through the Terrestrial Ecosystem Research Network (TERN 2023). Twenty-nine of the 33 freely available sites were selected. The four sites that were excluded showed strong
110 landscape heterogeneity within the flux tower footprint, insufficient temporal duration, or non-representative landcover (e.g., almond farms). A summary of the selected sites and their locations is shown in Figure A1. The Level 6 OzFlux data used in this study provides two separate estimates of constituent carbon fluxes derived from two methods for partitioning NEE into its component fluxes of GPP and ER. This study uses the ‘SOLO’ data version which is calculated using a data-driven nocturnal respiration approach for partitioning where respiration is modelled using an artificial neural network driven by air
115 and soil temperature, soil water content, and satellite-observed greenness (MODIS EVI) data (Isaac et al., 2017). We trained ML models with the flux data at a monthly temporal resolution using 2,825 monthly observations, equating to 235 site-years.

2.1.2 Gridded explanatory variables

The variables in Table 1 were selected for inclusion in the modelling framework as they were considered to cover most of the expected climate and landscape controls on the terrestrial carbon cycle in Australia. All datasets were temporally
120 resampled to monthly resolution and reprojected onto a common 1-km x 1-km geographic grid for prediction; the training procedure uses data extracted from the same 1-km gridded data (using the pixel located over the EC tower).

2.1.3 Comparison datasets

Datasets included for comparative purposes cover the three current categories of methods for estimating the exchange of terrestrial carbon with the atmosphere: process-based models, empirical upscaling of eddy covariance data, and atmospheric
125 inversions. Observation-based GPP products derived from light-use-efficiency methods and solar-induced fluorescence are also included for completeness. Where possible, datasets are processed and plotted in their native resolutions to avoid introducing errors from spatially resampling finer-resolution datasets to very coarse resolutions (or vice-versa). The exceptions to this are the MODIS-GPP and DIFFUSE-GPP products (described below) which were resampled to 1 km resolutions to match the resolutions of our ML upscaling product.

130



Table 1: Gridded feature layers used in the modelling framework to train and predict terrestrial carbon fluxes over Australia.

Explanatory Variable (abbreviation)	Description	Data Source & Reference
Land Surface Temp. (LST), Normalised Difference Water Index (NDWI), Kernel Normalised Vegetation Index (kNDVI),	A suite of MODIS derived products characterising the land surface responses to climate. In addition, fractional anomalies are calculated for the kNDVI variable to account for disturbances from fire or land-use change. Fractional anomalies are calculated against a long-term climatological mean from 2003-2021.	MODIS Collections (version 6.1) downloaded from Google Earth Engine: https://developers.google.com/earth-engine/datasets/catalog/modis
Average Air Temp. (Tavg), Vapour Pressure Deficit (VPD), Incoming Shortwave Radiation (srad), Total Precipitation (rain)	~1 km resolution gridded climate products based on topographically conditional spatial interpolation of Australia's extensive network of weather stations. In addition, fractional anomalies are also calculated for all variables except VPD. In addition to monthly fractional rainfall anomalies, three-, six-, and twelve-month cumulative fractional rainfall anomalies are added to help characterise memory and lag in the carbon response to water deficit.	ANUClimate: https://dapds00.nci.org.au/thredds/catalogs/gh70/catalog.html (Hutchison et al., 2014)
LST minus Tavg (LST-Tair)	The subtraction of air temperature from land surface temperature is indicative of vegetation canopy moisture stress	Derived from MODIS LST and ANUClimate Tavg
Fraction Trees (trees), Fraction C4 grass (C4_grass), Fraction Grass (grass), Fraction Bare (bare),	Per-pixel fractions of trees, grass, and bare derived from temporal decompositions of MODIS NDVI into persistent and recurrent fractions. An estimate of the proportion of C4 grass is also included. These variables are static and represent conditions in 2020.	Correspondence (Donohue, 2021)
Vegetation Height (VegH)	A per-pixel estimate of vegetation height in metres. This variable is static and represents the average vegetation height from 2007-2010.	Accessible from https://dapds00.nci.org.au/thredds/catalog/ub8/au/LandCover/OzWALD_L_C/catalog.html (Liao et al., 2020)



2.1.3.1 Process-model simulations

135 We compared our results with two runs of the CABLE model. The first was a regional, fine resolution (0.25°) offline run
forced by Australian regional climate drivers that follows the protocol from Haverd et al. (2018) but with land use remaining
static at the year 2000 (hereafter referred to as CABLE-BIOS3). CABLE-BIOS3 net biosphere production (NBP) includes
GPP and autotrophic and heterotrophic respiration, but does not include fire disturbances, harvest, erosion or export of
carbon through rivers (a fuller description of the set-up is outlined in Villalobos et al. (2022)). A second CABLE run was
140 extracted from the TRENDY v10 ensemble (Friedlingstein et al., 2022), hereafter referred to as CABLE-POP. This dataset
has a spatial resolution of 1° , is forced by global climate data and NBP includes additional fluxes from fire emissions and
land use change.

2.1.3.2 FLUXCOM

Our regional ML upscaling product is compared with the well-known global ML upscaling product, FLUXCOM (Jung et al.,
145 2020; Tramontana et al., 2016). FLUXCOM is built using similar machine learning methods to those used in this study,
though trained on the global FLUXNET2015 dataset. Two products are available, FLUXCOM-RS was trained exclusively
on MODIS remote sensing data, and FLUXCOM-RS+METEO (FLUXCOM-Met hereafter) which is trained on climate
reanalysis data and climatological remote sensing data (Jung et al., 2020). For FLUXCOM-Met, we use the multi-model
mean of the ERA5-based product. Both RS-METEO and RS products are assessed here and were downloaded at monthly
150 temporal resolution from the Max Planck Institute for Biogeochemistry ([https://www.bgc-
jena.mpg.de/geodb/projects/Home.php](https://www.bgc-jena.mpg.de/geodb/projects/Home.php), last access 13/01/2023).

2.1.3.3 Atmospheric Inversions

A regional inverse modelling product, produced by Villalobos et al. (2022) was included for comparison as it provides a
wholly independent measure of NEE. This regional inversion estimates carbon fluxes over the Australian continent from
155 2015-2019 by assimilating of carbon-dioxide measurements from the Orbiting Carbon Observatory-2 (OCO-2) satellite. The
product is provided at ~ 81 km spatial resolution and monthly temporal resolution (available for download from
<https://zenodo.org/record/6649768>). NEE in this dataset includes fire emissions and fossil fuel emissions, so to facilitate
better comparisons fossil fuel emissions were subtracted from the NEE time-series. A second regional satellite-assimilated
atmospheric inversion from Metz et al. (2023) is also included. This timeseries represents the spatially averaged net flux of
160 CO_2 over the Australian TRANSCOM region (which includes New Zealand). Therefore, the time-series is only shown where
Australian-wide spatially averaged time-series are plotted, and some differences between time-series may be attributable to
the inclusion of the New Zealand land mass in the estimate.



2.1.3.4 Observation-based GPP products

We compare our GPP estimates with a suite of observation-based GPP products: the MODIS Terra GPP product
165 (MOD17A2H), based on a per-biome light-use efficiency approach (Running et al., 2015); the GOSIF GPP product,
generated through a data-driven approach based on OCO-2 SIF soundings, MODIS remote sensing data, and meteorological
reanalysis data (Li and Xiao, 2019); and DIFFUSE GPP which is based on total and diffuse irradiance and the fraction of
shortwave irradiance absorbed by foliage (Donohue et al., 2014). All datasets are averaged to monthly temporal resolution,
and MODIS-GPP and DIFFUSE-GPP are spatial resampled to 1-km grid cells by averaging the 250 m pixels within each 1
170 km pixel grid.

2.1.4 Fire emissions

Fire emissions were added to our estimates of NEE from the Global Fire Assimilation System version 12 (GFASv12) (Kaiser
et al., 2012). Daily fire emissions are temporally resampled to monthly totals by summing daily values.

2.1.5 Bioclimatic regions

175 Bioclimatic regions used for separating fluxes into specific ecosystems were identical to those defined in Haverd et al.
(2013) and include six bioclimatic classes: tropics, savanna, warm temperate, cool temperate, Mediterranean, and desert
(Fig. 9a).

2.2 Methods

180 2.2.1 Empirical ML upscaling

The most common ML models implemented in the literature on empirical upscaling of EC data are random forest regression,
support vector regression, model tree ensembles, piecewise regression models, and artificial neural networks (Verrelst et al.,
2015). Random forest (RF) regression has proven itself to be the go-to model for many remote sensing-based studies owing
to its high accuracy, robustness to over-fitting, scalability, and easy to configure hyperparameters (Belgiu and Drăguț, 2016).
185 In recent years, gradient-boosting decision tree (GBDT) learning algorithms have also proven to be highly accurate and
robust to overfitting (Chen and Guestrin, 2016; Wei et al., 2019). Here, rather than rely on any one ML method, we rely on
both RF and GBDT methods to develop an ensemble of predictions.

Beyond the ML algorithm used, there are numerous other sources of uncertainty associated with the empirical
upscaling of EC flux tower data. Epistemic uncertainties arise from the limitations of the training data (e.g., biases in the
190 locations sampled), and uncertainties in the features used for training, as well as the hyperparameters used during model
optimisation. In addition to these reducible (or at least quantifiable) epistemic uncertainties, aleatoric uncertainties arise
from the uncertainties of the eddy covariance measurements themselves (Isaac et al., 2017), along with the non-deterministic



dependencies between variables (Hüllermeier and Waegeman, 2021). Here we attempt to account for a portion of the empirical uncertainty by iterating the training data and the models used for fitting. During model fitting, two randomly selected EC sites are removed from the training data and both a GBDT model (from the python package LightGBM (Ke et al., 2017) and a RF model are fit on the remaining data (hyperparameter optimization is conducted on every fit). This procedure is repeated 15 times to increase the likelihood of every site being removed from the training dataset, resulting in 30 unique models. These 30 models are used to generate 30 gridded estimates for each of the variables modelled (GPP, ER, and NEE). In the results that follow, we report the interquartile range of these 30 predictions as our envelope of uncertainty, and the ‘best-estimate’ as the median of the ensemble predictions.

The overall modelling framework is summarised in Figure 1. Each flux is independently modelled, and therefore there is no inherent exact mass balance between GPP-ER and NEE. The same predictor variables were used for modelling each flux, so the resulting products originate from a consistent set of drivers.




Figure 1: A flow chart showing the modelling framework for creating gridded estimates of GPP, ER, and NEE for the Australian continent.

2.2.2 Model evaluation

The accuracy of each ML model in the ensemble was assessed using a nested, time-series-split cross-validation approach (Fig. 2). This approach ensured minimal data leakage between training and testing sets, while still allowing the algorithm to ‘see’ all the sites during training; a desirable feature in the cross-validation technique due to the relatively limited number of sites (n=29), with some ecosystems sampled by only one or two flux towers (e.g., alpine regions, cereal cropping). Five outer cross-validation splits are performed, with each split containing 20 % test data from every site (as a discrete length of time equal to 20% of the total length of the dataset), while the remaining 80 % of the data is used for training. Five ‘inner’ cross-validation splits were conducted to optimise the hyperparameter selection for the outer loop. Across the five outer cross-validation splits, all samples in the dataset were tested. Mean Absolute Error (MAE) and the coefficient of



determination (R^2) are reported to assess the accuracy of the fit for each of the variables modelled. The cross-validation scores reported in the results section summarise the train-test splits of all 30 model fits. Throughout the remainder we use the terms ‘observed’ and ‘predicted’ to refer to in-situ measurements from EC towers and the predictions, respectively. We also use the convention of negative NEE values referring to net carbon uptake by the land surface.




Figure 2: A schematic representation of a single cross-validation split using a nested, time-series cross-validation procedure. Five outer splits and five inner splits were conducted per model iteration. For each split, models are trained on data from every site included in that model iteration (i.e., 80 % of every site) and tested on a continuous period for every site (i.e., 20 % of each site). For each subsequent split, the test period is moved forward in time.

In addition to evaluating the overall predictive capacity using temporal cross-validation, we also perform an intercomparison between the results of this study and similar products covering Australia. This is performed through scatter plots of modelled vs observed fluxes for several products (statistics for comparison are MAE and r^2 – the square of Pearson’s correlation), through comparison of the mean seasonal cycles disaggregated by bioclimatic region, and through the assessment of annual anomalies. It is important to note that NEE calculated through empirical upscaling of EC flux tower data is conceptually distinct from inversion-based NEE and process-model NBP. The addition of fire-emissions to our estimates of NEE narrows the conceptual distance between the estimates, and where a conceptual difference still applies, we contend that fluxes from other sources are unlikely to be large enough to warrant the additional complexity of their inclusion.

3. Results

3.1 Cross-validation performance

Temporal cross-validation results revealed a comparatively high degree of agreement between observations and predictions (Fig. 3). As for other regional and global upscaling products, GPP and ER were predicted with better skill than NEE. GPP scored a $R^2 = 0.91$ and MAE = 19.4 gC/m²/month. For ER, $R^2 = 0.89$ and MAE = 15.8 gC/m²/month, while for NEE, $R^2 = 0.68$ and MAE = 17.9 gC/m²/month. To understand how well the predictions reproduce annual mean fluxes, and the per-biome predictability of fluxes, we produced scatter plots comparing the annual mean fluxes of the EC flux tower sites with the annual mean fluxes of the median of the prediction ensemble (Fig. 3d-f). Regardless of biome, annual mean fluxes were



exceptionally well reproduced by the median of the ensemble with the ‘all-data’ fit closely matching the one-to-one line. The climatological seasonal cycles of NEE at each of the EC sites were also very well reproduced (Fig. A2).

245




Figure 3. Pooled temporal cross-validation results for EC flux tower sites: scatter plots of observed and predicted monthly (a) GPP, (b) ER and (c) NEE, with heat colours indicating data density; and scatter plots of observed and predicted annual mean (d) GPP, (e) ER, and (f) NEE, with colour coding indicating bioclimatic regions, as shown in Figure 9a.

250

Scatter plots showing the trend and strength of the relationships between EC flux tower observations and modelled values for other products can be found in the Appendices (Fig. A3). The EC flux tower values are compared with the nearest pixel in each product, and the products have been kept in their native resolution. This means there may be some bias to products with a higher spatial resolution, but errors introduced from resampling are avoided. Only those products with a reasonably high spatial resolution have been compared with the flux tower (i.e., 1° resolution CABLE-POP and the OCO-2 inversion have been excluded), and comparisons were only made for periods where all included products have data. Most products perform reasonably well at predicting GPP (Fig. A3a-f). Typically, products show an overestimation of small GPP and ER values, and an underestimation of large values, except for CABLE-BIOS3’s which overestimates GPP and ER across the distribution. CABLE-BIOS3’s estimates of NEE showed almost no correlation with EC flux tower observations,


255



260 recording a r^2 of 0.04 (Fig. A3j). The FLUXCOM NEE products performed considerably worse than the cross-validation scores reported in this study (Fig. A3k-l).

3.2 Feature importance

To understand which explanatory variables most impacted flux predictions, feature importance plots were produced using the Shapley Additive Explanations (SHAP) Python library (Lundberg & Lee, 2017). Shapley values represents the average marginal contribution of a feature value across all possible coalitions (Lundberg et al., 2020). The feature importance bar plots of Figure 4 show the top five ranked features for each modelled flux, ranked in descending order with the most important variables at the top. These plots were derived by calculating the mean absolute SHAP values for each feature in each model iteration, and subsequently averaging those values across all the models in the ensemble. Flux predictions were strongly influenced by the remote sensing variables of kNDVI and NDWI which respond to canopy density, health, and water status. Solar radiation and average air temperature were the most important climate variable across the fluxes. The land cover variables of vegetation height and fraction of trees also proved important for flux predictions.



275 **Figure 4. Shapley additive explanation (SHAP) feature importance plots. (a) GPP, (b) ER, (c) NEE. The plots summarise feature importance across all models in the ensemble by first calculating mean absolute SHAP values for each feature in each model, and then averaging those values across all the models in the ensemble. The error bars show the 95 % confidence interval.**

SHAP dependence plots for kNDVI along with the four principal climate drivers in the model (temperature, rainfall, solar radiation, and VPD) aid in the interpretation of feature importance (Fig. 5; these plots were created using a single optimised GDBT model fit on all the training data). In these plots, the feature values are plotted against their corresponding SHAP values, and the dots are coloured by, in the case of the climate drivers, kNDVI, and in the case of kNDVI, by the values of the feature that has the strongest interaction effect with kNDVI. A strong interaction between two variables produces a distinct vertical colour gradient. The dependency plots for the climate features are coloured by kNDVI as it aids in approximately disaggregating the influence of climate on carbon fluxes between the wetter, cooler, and high kNDVI coastal fringe regions of the Australian continent from the drier, warmer, lower kNDVI regions of Australia's (semi) arid interior. In the dependence plot for kNDVI (Fig. 5a), solar radiation shows a clear interaction effect. Where kNDVI is low (<



290

~0.2), increasing solar radiation produces predictions of GPP that are relatively lower than in regions with higher kNDVI. Solar radiation was the third most important feature in the prediction of GPP (Fig. 4a), and high kNDVI regions had a greater light sensitivity than low kNDVI regions (Fig. 5b).




Figure 5. SHAP dependency scatter plots for kNDVI, along with the four principal climate features (solar radiation, air temperature, rainfall and VPD). In the case of (a,f,k) the SHAP values are coloured by the feature with the largest interaction effect, while the climate variable SHAP values are coloured by their interaction with kNDVI. Note that the y-axis scale is different for each sub plot.

295

Solar radiation and kNDVI were also key predictors for ER, following similar relationships as GPP, but the overall amplitude of increase is less (Fig. 5f & 5g). ER also sees a greater influence from air temperature (Fig. 5h) and rainfall (Fig. 5i) than GPP, where higher values of these variables increased predicted rates of ER. In the case of air temperature, in areas of high kNDVI the rate of ER increase was greater than in low kNDVI regions. Rates of ER respiration increase sharply with increased rainfall, but for low kNDVI, predictions of ER increase at a more rapid rate than for high kNDVI (Fig. 5i).

300

Relationships between features and NEE predictions are more difficult to interpret given the likelihood of complex interaction effects when modelling the carbon balance (NEE) *versus* modelling only ER or GPP. The most important features for the NEE predictions are kNDVI and NDWI, average air temperature, and solar radiation (Fig. 4c). Increasing solar radiation typically resulted in more negative NEE predictions (greater uptake of carbon) (Fig. 5l). The rate of increase

305




in carbon uptake under increasing solar radiation is lower where kNDVI is low, while regions of high kNDVI see a much greater sensitivity to increases in solar radiation. Increasing air temperature tends to result in more positive NEE predictions (Fig. 5m), though the relationship does not follow a simple trajectory. For high kNDVI, temperature increases at the highest end of the distribution ($>25^{\circ}\text{C}$) result in a strong positive rate of change in NEE predictions (i.e., greater release of carbon).
310 For very low kNDVI, temperature changes have a much more modest impact on NEE.

3.3 Prediction uncertainties

The coefficient of variation between the thirty ensemble members provides a spatial indication of uncertainty in CO_2 flux predictions (Fig. 6). We use a non-standard definition of the coefficient of variation where the median absolute deviation between the long-term annual means of each ensemble member were divided by the median of the ensemble annual means,
315 expressed as an absolute value. Both GPP and ER show comparatively low variability across predictions, where the greatest coefficient of variation values is found in the arid interior (Fig. 6a and 6b). NEE shows stronger variation between ensemble members in some of the arid regions of the north-west, the savannah regions of western Queensland, and the agricultural regions of the Western Australian wheat belt and the Murray-Darling Basin (MDB) (Fig. 6c and 6d). In the case of the arid and savanna regions, the uncertainty coincides with areas where annual mean NEE is close to zero, so small deviations in
320 predictions can result in high relative uncertainty (refer to the annual mean flux map in Figure 8g). However, in parts of the aforementioned agricultural regions, uncertainty is both high in relative and absolute terms (again refer to Figure 8g). This may be due several factors such as: the dearth of OzFlux towers in agricultural regions, the heterogeneity of crop types and agricultural practices that are likely not well represented by our feature layers, and the potential for large amounts of carbon to be laterally exported out of these regions through harvest of agricultural commodities.

325




330 **Figure 6.** Prediction uncertainty estimated from iterating EC flux tower data and model type: (a-c) displays the absolute coefficient of variation for (a) GPP, (b) ER, and (c) NEE, defined as the median absolute deviation between all ensemble members, and the median of the ensembles, expressed as an absolute value. (d) shows the fraction of ensemble members where the sign of annual mean NEE (positive or negative) agrees, i.e., if all ensemble members agree on the sign of NEE then the values is one, and if positive and negative estimates are each produced by half of the members, then value is 0.5

3.4 Upscaling results and comparison with other products

3.4.1 Annual mean and IAV of carbon fluxes across Australia

335 We adopted the model ensemble median as our best estimate, and the interquartile range (IQR) of estimates as a measure of uncertainty. During 2003 to 2022 Australia's terrestrial ecosystems were a strong net carbon sink on an annual mean basis



340 **Figure 7. Monthly carbon fluxes summed across Australia from 2003 to 2021. (a) GPP, (b) ER, (c) NEE. Shading around time-series shows the interquartile range of the prediction ensembles, and the solid blue line shows the median of the ensemble predictions. Orange lines shows the 12-month running mean of the median model. Box plots are based on the median model prediction and show the long-term mean (green triangle), median (line within box) and interquartile ranges (boxes) averaged over the entire time series. c) also shows NEE after adding fire emissions (green line), as estimated by the GFASv12 product.**

345 of -0.44 (IQR= 0.42) PgC/year (Fig. 7c) (including fire emissions). IAV defined as one standard deviation of the annual mean timeseries is 0.18 PgC/year and the average seasonal range of NEE is 0.85 PgC/year. The annual mean estimates of NEE from this study show a greater terrestrial carbon uptake than any of the LSMs or FLUXCOM products, while the regional atmospheric inversion (which also includes fire emissions) predicts a very similar annual mean carbon uptake of -



0.47 PgC/year (though this is assessed over a much shorter period than the other products). IAV of NEE for the other products ranges from 0.06 PgC/yr for FLUXCOM-Met, to 0.26 PgC/year for the OCO-2 inversion. The GOSAT-inversion
350 conducted by Metz et al. (2023) estimated IAV of 0.207 PgC/yr across the Australia TRANSCOM region. CABLE-BIOS3
also shows comparatively high IAV of 0.23 PgC/year (Fig. 9f). The per-pixel plots of Figure 8g-i show how annual NEE
fluxes are spatially allocated. The strongest carbon sinks are seen along the forested coastal regions of the eastern seaboard
from western Tasmania to northern New South Wales, the south-west corner of Western Australia including the southern
part of the Great Western Woodlands, and the tropical part of the Northern Territory. The regions of strongest IAV in NEE
355 are in the savanna regions of northern Australia, the intensive agricultural regions of the MDB, and the Channel Country of
south-west Queensland and into South Australia where episodic river basins such as the Coopers and Meullers Creek
periodically fill during anomalously large rainfall events (Fig. 8h). The climatological ‘month-of-maximum’ NEE plot in
Figure 8i shows the month during which NEE typically achieves its most negative value (greatest carbon uptake), and the
plot shows clear delineations along bioclimatic regions.

360 Annual mean GPP across Australia averaged 4.25 (0.91) PgC/year, with an IAV of 0.50 PgC/year and an average
seasonal range of 1.47 PgC/yr (Fig. 7a). Averaged over Australia, our estimate of GPP closely approximates that of GOSIF
and MODIS, with the uncertainty envelope encompassing these two products. In contrast, DIFFUSE, FLUXCOM, and
CABLE-POP report lower estimates (Fig. 9a). The IAV between products varies substantially with both FLUXCOM
products showing the lowest IAV in GPP (FLUXCOM-Met: 0.13 PgC/year, FLUXCOM-RS: 0.23 PgC/year), while this
365 study and CABLE-BIOS3 (0.78 PgC/year) display the strongest IAV.

ER averaged 3.64 (1.01) PgC/year (Fig. 7b), with an IAV of 0.34 PgC/year and an average seasonal range of 1.56
PgC/year, notably higher than GPP. Agreement between products is generally poor, though the long-term mean of
FLUXCOM-Met and this study agree (Fig. 9b). CABLE-BIOS3 show the most IAV in ER (0.56 PgC/year), while the two
FLUXCOM products record very low IAV, with FLUXCOM-RS equal to 0.07 PgC/year, and FLUXCOM-Met 0.09
370 PgC/year.





Figure 8. Per pixel summaries derived from the median of the prediction ensemble. Annual means fluxes of GPP (a), ER (d), and NEE (g), Standard deviation in annual mean fluxes of GPP (b), ER (e), and NEE. Climatological month of maximum flux, GPP (c), ER (f), and NEE (i).



375


380

Figure 9. Twelve-month rolling mean terrestrial carbon fluxes from a suite a products covering Australia, compared with this study. Right-side plots (d-f) show the anomalies of the left-side plots (a-c), where the monthly anomalies are calculated using a climatology that starts in 2003 and ends at the maximum length of the available time-series for each product. The numbers in the left-side plots show the long-term annual mean flux for a given product. Blue shading around ‘This Study’ shows the interquartile range from prediction ensembles.

3.4.2 Climatological carbon fluxes

385

Figure 10e-g shows the climatological seasonal cycles of the component terrestrial fluxes summed across Australia (climatologies were calculated starting in 2003 and extending over the full remaining length of the time-series for each product). The seasonal cycle of this study’s NEE differs substantially from those of the LSMs and FLUXCOM-Met (Fig. 10g). According to our results, a climatological peak in terrestrial carbon uptake occurs for Australia during the cooler, drier months of March-September. Examination of the equivalent plots for GPP (Fig. 10e) and ER (Fig. 10f), shows that concomitant increases in ER during periods of peak GPP mean that the time of greatest primary production across Australia (December - March) are not coincident with peak carbon uptake. This result contrasts with the findings of the LSMs and FLUXCOM-Met which show peak carbon coinciding with peak GPP in Austral summer (Fig. 10g). Despite displaying a



390

Figure 10. Climatological seasonal cycles. (a) Map of bioclimatic regions. (b-d) Bioregion specific annual climatological seasonal cycles for GPP, ER, and NEE, respectively. (e-f) Annual climatological seasonal cycles averaged across Australia.



greater amplitude of seasonal variability, the NEE seasonal cycle of the regional OCO-2 inversion largely matches our estimate. The GOSAT-Inversion also displays similarities with this study and the OCO-2 inversion. However, the GOSAT-
395 inversion shows a second peak in July – it is unclear from the dataset provided if this might be due to the inclusion of New Zealand in the analysis area.

Breaking the fluxes down into bioclimatic zones (Fig. 10a-d), we can observe two processes that predominately dictate the typical seasonal pattern of NEE in Australia. Firstly, seasonal variations in ER in the desert region (peak-to-peak amplitude = 0.66 PgC/yr) exceed GPP variations (amplitude = 0.46 PgC/yr). Beginning in March and extending through the
400 autumn and winter period, ER declines more rapidly than GPP resulting in enhanced carbon uptake during this period. Secondly, in the savanna region we observe a sharp response in ER following the end-of-dry-season rainfall events that exceed the response from GPP, resulting in a net carbon pulse to the atmosphere in the Oct-Dec period (fluxes from these regions are re-plotted in Figure A5 to enhance interpretability). The interaction between these two processes likely explains most of the seasonal variation in Australia's terrestrial carbon cycle, and is responsible for peak carbon uptake in Australia
405 occurring in the autumn-winter months, while the carbon sink tends to be weakest during the Oct-Dec period.

We found that the largest discrepancies between products also occurs in the desert region (Fig. 10a). The LSMs, FLUXCOM-Met, GOSIF, and this study all report GPP peaking in February-March, with the nadir of GPP occurring during the May-Sept period (Fig. 9b). On the other hand, MODIS-GPP and FLUXCOM-RS show an inverted climatology to the other products that are unlikely to be accurate given the monsoonal climate drivers in the region with >70 % of the typical
410 annual median rainfall falling between November and April (Bowman et al., 2010). The CABLE-POP model appeared to significantly underestimate both GPP and ER in desert regions (Fig. 10b-c). This may explain why the Australia-wide seasonal NEE curve from CABLE-POP (Fig. 10g) does not align with the results of this study despite a similar spatial pattern in the month-of-maximum NEE flux plot (Fig. A4). The desert and savanna regions typically contribute the most to annual fluxes in other products, but CABLE-POP's NEE fluxes are comparatively more influenced by the savanna and
415 tropical regions. This is most likely due to CABLE-POP's representation of vegetation cover fractions over inland Australia which show the desert region as entirely bare (Teckentrup et al., 2021). FLUXCOM-RS follows a similar trajectory in the Australia-wide NEE to that of our estimate, though with considerably less seasonal amplitude (Fig. 10g). Examining the bioclimatic zones, we see that this is mainly due to an incorrect GPP seasonal cycle in the desert region, combined with a very low amplitude in the seasonal cycle of ER in the desert (Fig. 10c). The seasonal cycle of FLUXCOM-Met is markedly
420 different from FLUXCOM-RS. The per-biome fluxes from FLUXCOM-Met appear more realistic than those FLUXCOM-RS, but produce an inverted Australia-wide NEE seasonal cycle to our estimate (Fig. 10g). This is due to greater amplitude declines in seasonal GPP compared with ER, especially in the warm and cool temperate regions.

3.5 Drivers of carbon flux anomalies

As a simple means for interpreting the drivers of carbon flux anomalies, temporal Pearson correlations between carbon flux
425 anomalies and climate anomalies (respective to 2003-2021 averages) for each bioclimatic zone were conducted (Table 2).



Correlations were calculated per-pixel and then averaged over the bioclimatic zone. Caution in interpreting the results is warranted as the terrestrial carbon cycle is intrinsically complex, and nonlinear. With that caveat: for GPP, ER, and NEE, cumulative rainfall anomalies almost universally correlate most strongly with carbon flux anomalies. In the case of NEE, across all bioclimatic regions monthly rainfall anomalies were insignificantly correlated. Yet, the cumulative rainfall anomalies proved to be the strongest correlate (where a cumulative rainfall surplus resulted in negative NEE anomalies i.e., greater carbon uptake). In the case of the desert region, correlations of monthly rainfall anomalies jumped from a statistically insignificant r-value of -0.08 to a strong significant correlation of -0.50 for six-month cumulative rainfall anomalies (Table 2), similar scores were found for the savanna region. Correlations for non-lagged monthly rainfall anomalies in the savanna and desert regions were both much higher for ER than for GPP, suggesting ER responds more quickly to wetting than GPP in the arid and semi-arid regions of Australia.

4. Discussion

Our estimate of the long-term annual mean carbon sink over Australia (-0.44 PgC/yr) is higher than those reported by any study besides the regional OCO-2 inversion (-0.47 PgC/yr). We found evidence that Australia is, on average, a stronger annual carbon sink than previous CABLE LSM and FLUXCOM estimates have concluded: the consistency between our estimate of the mean sink and the OCO-2 inversion; the fact that 25 out of the 29 OzFlux EC sites used here report strong annual mean carbon sinks (Figure A7), and the theoretical argument that ML predictions tend to produce good estimates of the mean. Although upscaling uncertainties are present, our efforts to characterise a significant portion of the empirical uncertainty indicate that Australia's status as a comparatively strong net carbon sink holds up.

Carbon flux anomalies show better agreement between diverse methods, with our estimate, CABLE-BIOS3, and the GOSAT Inversion all largely agreeing on the timing and magnitude of NEE anomalies. The largest annual anomaly, the 2010-11 La Niña anomaly of -0.70 PgC/year reported here (based on a 12-month rolling mean) also aligns well with the -0.77 PgC reported by Ma et al. (2016) and the -0.79 PgC anomaly reported by Poulter et al. (2014). The OCO-2 Inversion, our study, CABLE-BIOS3, and the GOSAT-Inversion also converge on a NEE IAV of ~0.2 PgC/yr (the range among these products is 0.18 to 0.26 PgC/year). Cross-validation showed that our predictions generally underestimate large NEE fluxes (both positive and negative fluxes, Fig. 3). Thus, it is fair to assume that the inter-annual (and seasonal) variability in NEE should be larger than the estimate reported by this study, and perhaps the larger variability of the inversions is closer to the truth. This study is consistent with other studies in showing NEE anomalies in Australia are driven by a greater response of GPP over ER to anomalous rainfall periods (Ahlström et al., 2015; Ma et al., 2016; Poulter et al., 2014; Haverd et al., 2016; Trudinger et al., 2016; Teckentrup et al., 2021; Fig. 9). This is especially the case where rainfall anomalies are cumulative.



460

Table 2. Temporal Pearson correlations between carbon flux anomalies, climate anomalies, and kNDVI anomalies. Every flux and climate variable anomaly are based on a 2003-2021 baseline. The highest correlation for each flux and bioclimatic zone is shown in bold (for the climate variables only, kNDVI correlations are ignored), and the colour scheme indicates the strength of the correlations ranging from -1 (blue) to 1 (red).

Flux	Variable	Bioclimatic region					
		Tropics	Savanna	Warm Temperate	Cool Temperate	Mediterranean	Desert
GPP	Rainfall	0.17	0.27	0.21	0.15	0.25	0.39
	Rainfall Cml-3	0.28	0.46	0.51	0.41	0.48	0.66
	Rainfall Cml-6	0.33	0.54	0.57	0.47	0.57	0.78
	Rainfall Cml-12	0.26	0.59	0.50	0.44	0.52	0.74
	Air Temp.	-0.01	-0.36	-0.25	-0.11	-0.23	-0.36
	Solar Rad.	-0.23	-0.43	-0.28	-0.16	-0.29	-0.45
	kNDVI	0.86	0.88	0.88	0.81	0.84	0.80
ER	Rainfall	0.45	0.49	0.54	0.45	0.60	0.55
	Rainfall Cml-3	0.39	0.54	0.67	0.58	0.69	0.68
	Rainfall Cml-6	0.38	0.62	0.67	0.59	0.72	0.78
	Rainfall Cml-12	0.22	0.63	0.60	0.56	0.68	0.79
	Air Temp.	0.07	-0.31	-0.17	0.06	-0.14	-0.28
	Solar Rad.	-0.52	-0.59	-0.56	-0.38	-0.55	-0.54
	kNDVI	0.68	0.78	0.81	0.69	0.71	0.75
NEE	Rainfall	0.12	-0.02	0.09	0.05	0.07	-0.08
	Rainfall Cml-3	-0.13	-0.31	-0.30	-0.23	-0.22	-0.42
	Rainfall Cml-6	-0.25	-0.40	-0.41	-0.33	-0.32	-0.50
	Rainfall Cml-12	-0.34	-0.49	-0.36	-0.29	-0.30	-0.41
	Air Temp.	0.15	0.35	0.31	0.25	0.28	0.44
	Solar Rad.	-0.09	0.15	0.00	0.00	0.01	0.20
	kNDVI	-0.69	-0.79	-0.70	-0.67	-0.68	-0.57

The strong correlations between cumulative rainfall anomalies and NEE anomalies provides some additional support to the study of Cranko Page et al. (2022), who showed that the inclusion of rainfall lags increased the predictability of site-level NEE in Australia. Australia contributes substantially to the IAV of the global terrestrial carbon sink, an important advantage of our high-resolution dataset is that it allows us to identify and monitor fine-resolution hotspots of IAV (maps showing greater detail are shown in Figure A8). Through future updating of this dataset, the ecosystems that play an outsized role in



controlling Australia's mean carbon sink and contribute substantially to its IAV can begin to be systematically monitored for change.

470 We have shown that climatological peak terrestrial carbon uptake in Australia occurs in the Austral autumn and winter months owing mostly to more rapid declines in rates of ER compared with GPP over the arid regions of Australia. Concomitant increases in ER during times of high GPP mean that periods of peak primary production do not necessarily coincide with peak carbon uptake on a seasonal basis. This finding agrees with Renchon et al. (2018) at the Cumberland Plains EC flux tower site, where the forest was a CO₂ sink in winter and a source in summer due to larger seasonal
475 amplitudes in ER. Similarly, Metz et al. (2023) found that seasonal rainfall in semi-arid regions after the dry season drives pulses of heterotrophic respiration that precede the GPP response, leading to net carbon uptake not beginning until March. Our results confirm that ER over the savanna region responds quickly to seasonal rainfall events at the end of the dry-season, while GPP responds more slowly resulting in carbon pulses to the atmosphere during the Oct-Dec period. Correspondingly, we also find non-lagged correlations between monthly rainfall climatologies and climatological ER stronger than those for
480 GPP over the semi-arid regions of Australia (Fig. A6).

While our estimate shows substantial improvement over the existing empirical upscaling products covering Australia, the dataset can be improved in several ways. More EC flux tower data can be incorporated into the training dataset as new sites come online and/or existing sites continue to retrieve data. New or improved feature layers can be incorporated as they become available (e.g., time-varying estimates of the percentages of trees, grass and bare). Finally, the
485 dataset may be extended further back in time through the inclusion of other satellite missions such as Landsat and AVHRR. Regardless, the dataset provides a step-forward in our means for assessing the complex, seasonal, and interannual dynamics of Australia's carbon cycle.


5. Conclusions

Despite the sparseness of Australia's comparatively small network of EC towers we show that regional empirical upscaling
490 can improve considerably upon existing global upscaling products, outperform existing LSMs, perform similar to or better than other empirical GPP products, and replicate the dynamics of CO₂ flux over Australia as estimated by two regional atmospheric inversions. Our estimate suggests Australia was a strong carbon sink (2003-2022 average) with an annual mean uptake of -0.44 (0.42) PgC/year, has an IAV of 0.18 PgC/year, and an average seasonal amplitude of 0.85 PgC/yr. Estimates of the annual mean carbon uptake from other methods varied considerably, and only our study and the OCO-2 inversion
495 agreed. However, carbon flux anomalies showed much better agreement between methods. NEE anomalies were predominately driven by cumulative rainfall deficits and surpluses, resulting in larger anomalous responses from GPP over ER. In contrast, the long-term average seasonal cycle is dictated more by the variability in ER than GPP, resulting in peak carbon uptake typically occurring during the cooler, drier Austral autumn and winter months. Our new estimates of Australia's terrestrial carbon cycle fluxes improve upon our understanding of the magnitudes, seasonal cycles, and processes




500 governing Australia’s terrestrial carbon cycle and provides a new benchmark for assessment against future LSM developments, and a means for high-resolution monitoring of Australia’s terrestrial carbon cycle.

Appendix A: Figures



505 **Figure A1.** Locations of ‘OzFlux’ eddy covariance flux tower sites used in this study. The table on the right lists some key climate features of each site, such as the annual mean rainfall total, annual mean temperature, its elevation above sea level, and the average monthly kNDVI of the site. The ‘stamen’ basemap is provided by OpenStreetMap.



510

Figure A2. Climatological seasonal cycles of NEE for each EC flux tower site used in this study, plotted along with the seasonal cycle of the predictions from the nearest pixel to the tower.

515

520






Figure A3. Scatter plots of modelled vs EC flux tower monthly carbon fluxes for a suite of products. The EC tower flux values are compared with the nearest pixel in each product, and the products have been kept in their native resolution. Only those products with a reasonably high spatial resolution have been compared with the flux tower (i.e., 1-degree resolution CABLE-POP and the OCO-2 Inversion have been excluded from these plots).


525



530 **Figure A4.** Climatological month of maximum flux plots. In the case of NEE (b), the pixels show the month of the most negative value (i.e. largest carbon sink). Climatologies are calculated from 2003 and extend to the full length of the available time-series for each product, indicated in the subtitle of each plot.



535 **Figure A5.** (a) Flux climatologies for the Savanna and Desert region, showing the same results as those in Figure 10, but shown on single plot to enhance interpretability. (b) NEE per bioclimatic region calculated by subtracting GPP from ER (i.e., not directly modelled), presented here to show how the fluxes interact to produce NEE.



540 **Figure A6.** Per pixel temporal Pearson correlations between ER climatologies and rainfall climatologies.





Figure A7. Boxplots of annual cumulative NEE for each of the sites used in the empirical upscaling.



545

Figure A8. Maps of annual mean NEE and standard deviation in annual mean NEE zoomed in on three regions to show the landscape features resolved by a high-resolution (1km) dataset of NEE. The top three panels show a region in central Queensland that extends from the episodic rivers in the south-east (e.g., Coopers Creek), to Townsville in the north west. Panel (c) shows a true colour satellite image (sourced from Esri World Imagery), panel (a) shows the long-term annual mean, and (b) shows the standard deviation in the annual means. Panels d-f show the same but for south-east Australia extending from Adelaide in the west to Mallacoota in the east. Panels g-f show the same but for south-west Western Australia.

550



Code Availability.

The code used to conduct all analysis shown in this manuscript is available on the open-source repository:

https://github.com/cbur24/NEE_modelling

555 **Data Availability.**

The surface gridded carbon fluxes are available from the Zenodo repository at: <https://doi.org/10.5281/zenodo.7947265>.

These fluxes have been resampled to a 5 km grid to facilitate easier uploading and sharing. Full resolution datasets can be provided on request.

Author Contributions.

560 CB and LR conceived the study, CB performed all analysis and drafted the manuscript. SR, AVD, and LR provided extensive intellectual input and provided extensive edits to the manuscript.

Competing interests.

The authors declare that they have no conflict of interest.

Acknowledgements

565 The authors would like to thank Dr Yohanna Villalobos for providing the CABLE-BIOS3 and the OCO-2 Inversion datasets used in the intercomparison. We recognise the efforts of Dr Randall Donohue who provided access to several datasets and valuable intellectual discussion. We would also like to thank the efforts of the Terrestrial Ecosystem Research Network (TERN) infrastructure, which is enabled by the Australian Government's National Collaborative Research Infrastructure Strategy (NCRIS). Lastly, we thank the National Computing Infrastructure (NCI) which provides a research compute
570 environment without which this work would not be possible.

Financial Support

The first author is supported by a research scholarship provided by Geoscience Australia, funded by the Australian Government.

575



References

- Ahlström, A., Raupach, M. R., Schurgers, G., Smith, B., Arneeth, A., Jung, M., Reichstein, M., Canadell, J. G., Friedlingstein, P., and Jain, A. K.: The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink, *Science*, 348, 895-899, 2015.
- 580 Basu, S., Guerlet, S., Butz, A., Houweling, S., Hasekamp, O., Aben, I., Krummel, P., Steele, P., Langenfelds, R., and Torn, M.: Global CO₂ fluxes estimated from GOSAT retrievals of total column CO₂, *Atmospheric Chemistry and Physics*, 13, 8695-8717, 2013.
- Belgiu, M. and Drăguț, L.: Random forest in remote sensing: A review of applications and future directions, *ISPRS journal of photogrammetry and remote sensing*, 114, 24-31, 2016.
- Beringer, J., Moore, C. E., Cleverly, J., Campbell, D. I., Cleugh, H., De Kauwe, M. G., Kirschbaum, M. U., Griebel, A., Grover, S., and Huete, A.: Bridge to the future: Important lessons from 20 years of ecosystem observations made by the OzFlux network, *Global Change*
585 *Biology*, 2022.
- Beringer, J., Hutley, L. B., McHugh, I., Arndt, S. K., Campbell, D., Cleugh, H. A., Cleverly, J., Resco de Dios, V., Eamus, D., Evans, B., Ewenz, C., Grace, P., Griebel, A., Haverd, V., Hinko-Najera, N., Huete, A., Isaac, P., Kanniah, K., Leuning, R., Liddell, M. J., Macfarlane, C., Meyer, W., Moore, C., Pendall, E., Phillips, A., Phillips, R. L., Prober, S. M., Restrepo-Coupe, N., Rutledge, S., Schroder, I., Silberstein, R., Southall, P., Yee, M. S., Tapper, N. J., van Gorsel, E., Vote, C., Walker, J., and Wardlaw, T.: An introduction to the
590 Australian and New Zealand flux tower network – OzFlux, *Biogeosciences*, 13, 5895-5916, 10.5194/bg-13-5895-2016, 2016.
- Byrne, B., Liu, J., Lee, M., Yin, Y., Bowman, K. W., Miyazaki, K., Norton, A. J., Joiner, J., Pollard, D. F., and Griffith, D. W.: The carbon cycle of southeast Australia during 2019–2020: Drought, fires, and subsequent recovery, *AGU Advances*, 2, e2021AV000469, 2021.
- Chen and Guestrin: Xgboost: A scalable tree boosting system, *Proceedings of the 22nd acm sigkdd international conference on knowledge discovery and data mining*, 785-794,
- 595 Chen, M., Rafique, R., Asrar, G. R., Bond-Lamberty, B., Ciais, P., Zhao, F., Reyer, C. P., Ostberg, S., Chang, J., and Ito, A.: Regional contribution to variability and trends of global gross primary productivity, *Environmental Research Letters*, 12, 105005, 2017.
- Chevallier, F., Palmer, P. I., Feng, L., Boesch, H., O'Dell, C. W., and Bousquet, P.: Toward robust and consistent regional CO₂ flux estimates from in situ and spaceborne measurements of atmospheric CO₂, *Geophysical Research Letters*, 41, 1065-1070, 2014.
- Cranko Page, J., De Kauwe, M. G., Abramowitz, G., Cleverly, J., Hinko-Najera, N., Hovenden, M. J., Liu, Y., Pitman, A. J., and Ogle, K.:
600 Examining the role of environmental memory in the predictability of carbon and water fluxes across Australian ecosystems, *Biogeosciences*, 19, 1913-1932, 10.5194/bg-19-1913-2022, 2022.
- Detmers, R., Hasekamp, O., Aben, I., Houweling, S., Van Leeuwen, T., Butz, A., Landgraf, J., Köhler, P., Guanter, L., and Poulter, B.: Anomalous carbon uptake in Australia as seen by GOSAT, *Geophysical Research Letters*, 42, 8177-8184, 2015.
- Devanand, A., Falster, G. M., Gillett, Z. E., Hobeichi, S., Holgate Chiara M., Jin, C., Mu, M., Parker, T., Rifai, S. W., Rome, K. S.,
605 Stojanovic, M., Vogel, E., Abram, N. J., Abramowitz, G., Coats, S., Evans, J. P., Gallant, A. J. E., Pitman, A. J., Power, S. B., Rauniyar, S. P., Taschetto, A. S., and Ukkola, A. M.: The Tinderbox Drought in southeast Australia, 2017-2019: characteristics, causes, and implications for the future submitted to *Science Advances*, 2023.
- Donohue: Testing the accuracy of Gramina, a model for monitoring Australia's pasture and grassland biomass, CSIRO, Canberra, Australia., CSIRO, Canberra, Australia., 2021.
- 610 Donohue, Hume, I., Roderick, M. L., McVicar, T. R., Beringer, J., Hutley, L. B., Gallant, J. C., Austin, J. M., Van Gorsel, E., and Cleverly, J. R.: Evaluation of the remote-sensing-based DIFFUSE model for estimating photosynthesis of vegetation, *Remote Sensing of Environment*, 155, 349-365, 2014.
- Eldering, A., O'Dell, C. W., Wennberg, P. O., Crisp, D., Gunson, M. R., Viatte, C., Avis, C., Braverman, A., Castano, R., Chang, A., Chapsky, L., Cheng, C., Connor, B., Dang, L., Doran, G., Fisher, B., Frankenberg, C., Fu, D., Granat, R., Hobbs, J., Lee, R. A. M.,
615 Mandrake, L., McDuffie, J., Miller, C. E., Myers, V., Natraj, V., O'Brien, D., Osterman, G. B., Oyafuso, F., Payne, V. H., Pollock, H. R., Polonsky, I., Roehl, C. M., Rosenberg, R., Schwandner, F., Smyth, M., Tang, V., Taylor, T. E., To, C., Wunch, D., and Yoshimizu, J.: The Orbiting Carbon Observatory-2: first 18 months of science data products, *Atmos. Meas. Tech.*, 10, 549-563, 10.5194/amt-10-549-2017, 2017.
- Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Bakker, D. C., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., and Pongratz, J.: Global carbon budget 2021, *Earth System Science Data*, 14, 1917-2005, 2022.
- Haverd, V., Smith, B., and Trudinger, C.: Dryland vegetation response to wet episode, not inherent shift in sensitivity to rainfall, behind Australia's role in 2011 global carbon sink anomaly, *Global Change Biol*, 22, 2315-2316, 2016.
- Haverd, V., Raupach, M. R., Briggs, P. R., Canadell, J. G., Isaac, P., Pickett-Heaps, C., Roxburgh, S. H., van Gorsel, E., Viscarra Rossel, R. A., and Wang, Z.: Multiple observation types reduce uncertainty in Australia's terrestrial carbon and water cycles, *Biogeosciences*, 10,
625 2011-2040, 10.5194/bg-10-2011-2013, 2013.
- Hüllermeier, E. and Waegeman, W.: Aleatoric and epistemic uncertainty in machine learning: An introduction to concepts and methods, *Machine Learning*, 110, 457-506, 2021.
- Hutchison, M., Kesteven, J., and Xu, T.: ANUClimate collection [dataset], 2014.
- 630 Isaac, P., Cleverly, J., McHugh, I., Van Gorsel, E., Ewenz, C., and Beringer, J.: OzFlux data: network integration from collection to curation, *Biogeosciences*, 14, 2903-2928, 2017.



- Jung, M., Schwalm, C., Migliavacca, M., Walther, S., Camps-Valls, G., Koirala, S., Anthoni, P., Besnard, S., Bodesheim, P., and Carvalhais, N.: Scaling carbon fluxes from eddy covariance sites to globe: synthesis and evaluation of the FLUXCOM approach, *Biogeosciences*, 17, 1343-1365, 2020.
- 635 Ke, G., Meng, Q., Finley, T., Wang, T., Chen, W., Ma, W., Ye, Q., and Liu, T.-Y.: Lightgbm: A highly efficient gradient boosting decision tree, *Advances in neural information processing systems*, 30, 2017.
- Kowalczyk, E., Wang, Y., Law, R., Davies, H., McGregor, J., and Abramowitz, G.: The CSIRO Atmosphere Biosphere Land Exchange (CABLE) model for use in climate models and as an offline model, *CSIRO Marine and Atmospheric Research Paper*, 13, 42, 2006.
- Li and Xiao: A global, 0.05-degree product of solar-induced chlorophyll fluorescence derived from OCO-2, MODIS, and reanalysis data, *Remote Sensing*, 11, 517, 2019.
- 640 Li, X., Ma, H., Ran, Y., Wang, X., Zhu, G., Liu, F., He, H., Zhang, Z., and Huang, C.: Terrestrial carbon cycle model-data fusion: Progress and challenges, *Science China Earth Sciences*, 64, 1645-1657, 2021.
- Liao, Z., Van Dijk, A. I., He, B., Larraondo, P. R., and Scarth, P. F.: Woody vegetation cover, height and biomass at 25-m resolution across Australia derived from multiple site, airborne and satellite observations, *International Journal of Applied Earth Observation and Geoinformation*, 93, 102209, 2020.
- 645 Ma, X., Huete, A., Cleverly, J., Eamus, D., Chevallier, F., Joiner, J., Poulter, B., Zhang, Y., Guanter, L., and Meyer, W.: Drought rapidly diminishes the large net CO₂ uptake in 2011 over semi-arid Australia, *Scientific Reports*, 6, 1-9, 2016.
- Metz, E.-M., Vardag, S. N., Basu, S., Jung, M., Ahrens, B., El-Madany, T., Sitch, S., Arora, V. K., Briggs, P. R., and Friedlingstein, P.: Soil respiration-driven CO₂ pulses dominate Australia's flux variability, *Science*, 379, 1332-1335, 2023.
- 650 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., and Humphrey, M.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, *Scientific data*, 7, 1-27, 2020.
- Poulter, B., Frank, D., Ciais, P., Myneni, R. B., Andela, N., Bi, J., Broquet, G., Canadell, J. G., Chevallier, F., and Liu, Y. Y.: Contribution of semi-arid ecosystems to interannual variability of the global carbon cycle, *Nature*, 509, 600-603, 2014.
- Quillet, A., Peng, C., and Garneau, M.: Toward dynamic global vegetation models for simulating vegetation-climate interactions and feedbacks: recent developments, limitations, and future challenges, *Environmental Reviews*, 18, 333-353, 2010.
- 655 Renchon, A. A., Griebel, A., Metzen, D., Williams, C. A., Medlyn, B., Duursma, R. A., Barton, C. V., Maier, C., Boer, M. M., and Isaac, P.: Upside-down fluxes Down Under: CO₂ net sink in winter and net source in summer in a temperate evergreen broadleaf forest, *Biogeosciences*, 15, 3703-3716, 2018.
- Rödenbeck, C., Zaehle, S., Keeling, R., and Heimann, M.: How does the terrestrial carbon exchange respond to inter-annual climatic variations? A quantification based on atmospheric CO₂ data, *Biogeosciences*, 15, 2481-2498, 10.5194/bg-15-2481-2018, 2018.
- 660 Teckentrup, L., De Kauwe, M. G., Pitman, A. J., Goll, D. S., Haverd, V., Jain, A. K., Joetzer, E., Kato, E., Lienert, S., Lombardozzi, D., McGuire, P. C., Melton, J. R., Nabel, J. E. M. S., Pongratz, J., Sitch, S., Walker, A. P., and Zaehle, S.: Assessing the representation of the Australian carbon cycle in global vegetation models, *Biogeosciences*, 18, 5639-5668, 10.5194/bg-18-5639-2021, 2021.
- Terrestrial Ecosystem Research Network (TERN): <https://www.tern.org.au/>, last
- 665 Tramontana, G., Jung, M., Schwalm, C. R., Ichii, K., Camps-Valls, G., Ráduly, B., Reichstein, M., Arain, M. A., Cescatti, A., and Kiely, G.: Predicting carbon dioxide and energy fluxes across global FLUXNET sites with regression algorithms, *Biogeosciences*, 13, 4291-4313, 2016.
- Trudinger, C. M., Haverd, V., Briggs, P. R., and Canadell, J. G.: Interannual variability in Australia's terrestrial carbon cycle constrained by multiple observation types, *Biogeosciences*, 13, 6363-6383, 10.5194/bg-13-6363-2016, 2016.
- 670 Verrelst, J., Camps-Valls, G., Muñoz-Mari, J., Rivera, J. P., Veroustraete, F., Clevers, J. G., and Moreno, J.: Optical remote sensing and the retrieval of terrestrial vegetation bio-geophysical properties—A review, *ISPRS Journal of Photogrammetry and Remote Sensing*, 108, 273-290, 2015.
- Villalobos, Y., Rayner, P. J., Silver, J. D., Thomas, S., Haverd, V., Knauer, J., Loh, Z. M., Deutscher, N. M., Griffith, D. W. T., and Pollard, D. F.: Interannual variability in the Australian carbon cycle over 2015–2019, based on assimilation of Orbiting Carbon Observatory-2 (OCO-2) satellite data, *Atmos. Chem. Phys.*, 22, 8897-8934, 10.5194/acp-22-8897-2022, 2022.
- 675 Wei, Z., Meng, Y., Zhang, W., Peng, J., and Meng, L.: Downscaling SMAP soil moisture estimation with gradient boosting decision tree regression over the Tibetan Plateau, *Remote Sensing of Environment*, 225, 30-44, 2019.
- Williams, J. and Woinarski, J.: *Eucalypt ecology: individuals to ecosystems*, Cambridge University Press 1997.