Supplementary Information

Distinct Photochemistry in Glycine Particles Mixed with Different Atmospheric Nitrate Salts

Zhancong Liang^{1,2}, Zhihao Cheng¹, Ruifeng Zhang^{1,2}, Chak K. Chan^{1, 3*}

¹School of Energy and Environment, City University of Hong Kong, Hong Kong, China.

²City University of Hong Kong Shenzhen Research Institute, Shenzhen, China

³Division of Physical Science and Engineering, King Abdullah University of Science and

Technology, Thuwal, 23955-6900, Saudi Arabia

Correspondence to: Chak K. Chan (chak.k.chan@cityu.edu.hk; chak.chan@kaust.edu.sa)

Content

Figure S1. The schematic of the aerosol flow-cell system.

Figure S2. The schematic of the aqueous reactor system.

Figure S3. The Full width at half maxima of nitrate peak and glycine peak in AN+GC particles at different RHs.

Figure S4. The Raman characteristic peak of GSN.

Figure S5. Ionic chromatography of Fresh and UV-aged AN+GC and SN+GC particles extract.

Figure S6. Raman spectrum of ammonium acetate solution.

Figure S7. The decay kinetics of nitrate in AN+GC and SN+GC particles at 80% RH.

Figure S8. Raman spectrum of AN+Ala and SN+Ala mixed particles after 0 and 8 h irradiation at 80% RH.



Figure S1. The schematic of the aerosol flow-cell system.



Figure S2. The schematic of the aqueous reactor system.



Figure S3. The Full width at half maxima of nitrate peak (at ~1050 cm⁻¹) and glycine peak (~900 cm⁻¹) in AN+GC particles at different RHs.



Figure S4. The Raman characteristic peak of GSN.



Figure S5. Ionic chromatography of Fresh and UV-aged (8 h) AN+GC and SN+GC particles extract. The particles were aged at 80%RH.



Figure S6. Raman spectrum of ammonium acetate solution.



Figure S7. The decay kinetics of nitrate in AN+GC and SN+GC particles at 80% RH.



Figure S8. Raman spectrum of AN+Ala and SN+Ala mixed particles after 0 and 8 h irradiation at 80% RH.