

# A model of the within-population variability of budburst in forest trees

Jianhong Lin<sup>1, \*</sup>, Daniel Berveiller<sup>1</sup>, Christophe François<sup>1</sup>, Heikki Hänninen<sup>2, 3</sup>, Alexandre Morfin<sup>1</sup>, Gaëlle Vincent<sup>1</sup>, Rui Zhang<sup>2, 3</sup>, Cyrille Rathgeber<sup>4</sup>, Nicolas Delpierre<sup>1, 5, \*</sup>

<sup>1</sup> Université Paris-Saclay, CNRS, AgroParisTech, Ecologie Systématique et Evolution, 91190, Gif-sur-Yvette, France.

<sup>2</sup> State Key Laboratory of Subtropical Silviculture, Zhejiang A&F University, Hangzhou, China

<sup>3</sup> SFGA Research Center for *Torreya grandis*, Zhejiang A&F University, Hangzhou, China

<sup>4</sup> INRAE, SILVA, Université de Lorraine, AgroParisTech, Nancy, France

<sup>5</sup> Institut Universitaire de France (IUF)

\* Correspondence to: jianhong.lin@universite-paris-saclay.fr, nicolas.delpierre@universite-paris-saclay.fr

**Abstract.** Spring phenology is a key indicator of temperate and boreal ecosystems' response to climate change. To date, most phenological studies have analyzed the mean date of budburst in tree populations while overlooking the large variability of budburst among individual trees. The consequences of neglecting the within-population variability (WPV) of budburst when projecting the dynamics of tree communities are unknown. Here, we develop the first model designed to simulate the WPV of budburst in tree populations. We calibrated and evaluated the model on 48,442 budburst observations collected between 2000 and 2022 in three major temperate deciduous trees, namely, hornbeam (*Carpinus betulus*), oak (*Quercus petraea*) and chestnut (*Castanea sativa*). The WPV model received support for all three species, with a root mean square error of  $5.6 \pm 0.3$  days. Retrospective simulations over 1961-2022 indicated earlier budburst as a consequence of ongoing climate warming. However, simulations revealed no significant change for the duration of budburst (DurBB, i.e., the time interval from BP20 to BP80, which respectively represent the date when 20% and 80% of trees in a population have reached budburst), due to a lack of significant temperature increase during DurBB in the past. This work can serve as a basis for the development of models targeting intra-population variability of other functional traits, which is of increasing interest in the context of climate change.

Keywords: budburst variability; model; temperate trees; climate warming; budburst duration; population.

## 1. Introduction

Phenology, as the study of recurrent biological events such as budburst in spring, has attracted increasing attention due to climate warming (Piao et al., 2019). The timing of leaf phenology in spring is a major indicator of climate warming (Parmesan and Yohe, 2003) and is mainly modulated by temperature (Menzel et al., 2006; Zhang et al., 2022; Zhang et al., 2021; Chen et al., 2018; Vitasse et al., 2009a), photoperiod (Delpierre et al., 2016; Fu et al., 2019; Vitasse and Basler, 2013; Meng et al., 2021) and soil moisture (Liu et al., 2022; Luo et al., 2021). In the northern hemisphere, it is well established that spring phenological events have been advanced by climate warming (Walther et al., 2002; Menzel et al., 2006), although this advancement is currently slowing down (Fu et al., 2015; Chen et al.,

36 2019). To date, massive efforts have been made to study the spatiotemporal variability of leaf phenology among tree  
37 populations and across years (Delpierre et al., 2016; Fu et al., 2015; Meng et al., 2021; Chen et al., 2018). However,  
38 the variability of leaf phenology within populations has received little attention to date (Scotti et al., 2016; Delpierre  
39 et al., 2017), which is in line with the general focus of ecological studies on average traits (Violle et al., 2012). This  
40 is intriguing, since the within-population (i.e., tree-to-tree) variability of phenological events is vast and can even be  
41 equivalent to that observed among populations (Delpierre et al., 2017; Vitasse et al., 2009a; Rathgeber et al., 2011).  
42 It typically takes 1 to 4 weeks from the first to the last tree to burst buds in a population (Denechere et al., 2021),  
43 with an average of 19 days (Delpierre et al., 2017). Furthermore, the duration from the first to last tree to burst buds  
44 in a given population varies annually (Denechere et al., 2021).

45 The large within-population variability (WPV) of budburst observed in natural tree populations is considered to  
46 result from their exposure to a large range of fluctuating environmental (e.g., frost) and biotic (e.g., herbivores and  
47 pathogens) selection pressures, which alternatively favor trees that burst buds early or late (Alberto et al., 2011).  
48 From an evolutionary point of view, this phenotypic diversity has an adaptive value at the population scale, because  
49 the environment is likely to change across the lifetime of trees (Petit and Hampe, 2006; Morente-Lopez et al., 2022;  
50 Blanquart et al., 2013). For instance, if a local climate becomes suitable in early spring under climate warming, trees  
51 that burst buds early will benefit from an extended growing season, thus maximizing their carbon assimilation and  
52 possibly their biomass production (Zohner et al., 2020; Delpierre et al., 2009; Richardson et al., 2010), which will  
53 allow them to gradually occupy a dominant position in the population. Moreover, early budburst enables trees to  
54 escape pathogens (e.g., for oak, see Dantec et al., 2015). On the contrary, if freezing events occur frequently in early  
55 spring with the advance of budburst, late trees can grow better by avoiding freezing injury (Delpierre et al., 2017;  
56 Zohner et al., 2020; Puchalka et al., 2016). Moreover, the WPV also affects interactions with competing plants and  
57 herbivores (Hart et al., 2016; Renner and Zohner, 2018).

58 **The internal mechanism of the WPV of budburst is probably underpinned by genetic diversity, as evidenced by the**  
59 **variability of phenological traits among individual trees that experience similar environmental conditions (Bontemps**  
60 **et al., 2016; Delpierre et al., 2017). This genetic determinism is further reflected in the year-to-year repeatability of**  
61 **the phenological ranking of individuals within tree populations (Delpierre et al., 2017). In addition to this genetic**  
62 **determinism, the WPV is also likely influenced by micro-environmental variations such as the unbalanced**  
63 **distribution of soil-water content within populations, edaphic conditions, or microtopography (Delpierre et al., 2017;**  
64 **Denechere et al., 2021; Scotti et al., 2016). To the best of our knowledge, the question of whether and to what extent**  
65 **would the WPV of budburst be modified in the current context of climate change has not been addressed so far.**

66 Phenological research has made extensive use of modeling to study the response of the spatiotemporal variability of  
67 budburst to climate warming (Zhang et al., 2022; Meng et al., 2021; Delpierre et al., 2009; Chuine and Regniere,  
68 2017). The models postulate that temperature and photoperiod are the main environmental cues that trigger budburst  
69 in boreal and temperate (Delpierre et al., 2009; Kramer, 1994; Hänninen and Kramer, 2007), subtropical (Zhang et  
70 al., 2022; Du et al., 2019), and tropical trees (Chen et al., 2017). **In process-based models for spring phenology, the**  
71 **effects of environmental factors (mainly air temperature) on budburst are quantified (Zhang et al., 2022; Hänninen,**

2016; Jewaria et al., 2021). Firstly, dormancy state of buds reached in the previous autumn is released due to exposure to low temperature, that is, removing the growth-arresting physiological factors in the bud (the chilling requirement of dormancy release). Secondly, when dormancy is relieved to a certain extent, high temperatures drive the process of ontogenetic development, that is, visible bud elongation and swelling that results in budburst (forcing requirement of ontogenetic growth). Meanwhile, there is an interaction between these two stages in the models, namely, ontogenetic growth is influenced by dormancy release (Hänninen, 2016; Hänninen and Kramer, 2007; Vegis, 1964). Lundell et al. (2020) further proved that this interaction can be affected by prevailing temperatures. One important point is that these models do not pay attention to the WPV of phenological traits. They have been parameterized and applied to predict the mean or median date of budburst in a given tree population (Lundell et al., 2020; Kramer, 1994; Zhang et al., 2022). In other words, these models simulate the timing of budburst as a discrete event in the population without considering the WPV of leaf phenology. To the best of our knowledge, only two studies to date, notably (Rousi and Heinonen, 2007) in Birch (*Betula pendula*) and (Langvall et al., 2001) in Norway spruce (*Picea abies* (L.) Karst.), have attempted to establish a link between WPV and environmental conditions through the temperature sum required for the opening of buds at the scale of individual trees. At the scale of tree populations, a distribution of temperature sums to budburst was also used in the so-called physio-demo-genetic (PDG) models (Kramer et al., 2008; Oddou-Muratorio and Davi, 2014) to simulate the adaptive potential of tree populations. However, a systematic model for the WPV of budburst is still lacking.

Here we developed a model that simulates the WPV of budburst in temperate deciduous trees. We calibrated and validated the model over an extensive budburst dataset acquired from five tree populations at the individual tree scale over 23 years (representing 48,442 observations). Specially, we aim to 1) develop the WPV model and validate its ability for predicting the progress of budburst in tree populations, 2) use the model to in a retrospective simulation exercise testing whether the duration of budburst period in the population changed with climate warming in the recent decades.

## 2. Materials and Methods

### 2.1 Study sites

We used budburst data collected from two forests located near Paris (France): Barbeau (48.476° N, 2.780° E, 95 m asl) and Orsay (48.705° N, 2.167° E, 105 m asl). At these sites, the progress of budburst was observed at the individual scale in populations of three major temperate deciduous tree species, namely, hornbeam (*Carpinus betulus* L.), oak (*Quercus petraea* (Matt.) Liebl) and chestnut (*Castanea sativa* Mill.). Hornbeam is an early leafing tree species, chestnut is a late species while oak is intermediate. Hornbeam and oak are present in both forests, while chestnut is present in Orsay only (Table 1). For each species, we focused on healthy and dominant trees, except for hornbeam (an understory species). We collected budburst observations from 2000 to 2022, which yielded a dataset comprising 5 populations and 103 population-years. In each population, we observed between 28 and 309 individuals (mean 90) (Table 1).

106

107

108 **Table 1. Description of the phenological and meteorological datasets.**

Phenology Site	Coordinate	Meteorological station	Coordinate	Species	Number of year	Number of data	Number of trees (min / max / average)	Observation years
Orsay	48.705° N, 2.165° E	Gometz-le- Châtel	48.677° N, 2.136° E	<i>Quercus</i>	23	153	29/190/85	2000-2022
				<i>Carpinus</i>	20	124	29/146/50	2002-2006, 2008- 2022
				<i>Castanea</i>	21	112	29/192/80	2000-2007, 2010- 2022
Barbeau	48.476° N, 2.780° E	Châtelet-en- Brie	48.491° N, 2.802° E	<i>Quercus</i>	20	87	29/309/154	2003-2022
				<i>Carpinus</i>	19	64	28/241/114	2004-2022

109

## 110 2.2 Phenology dataset

111 A team of eight local observers (including most of the authors of this paper) conducted the observations of  
 112 developing buds in the tree crowns throughout spring. The observers used binoculars and occasionally received  
 113 training in order to reduce observer bias (Liu et al., 2021). The interval between phenological observations was of 4  
 114 days on average (from 2 to 7 days). A tree was considered to have burst its buds when at least 50% of the buds in the  
 115 upper third of the crown presented leaves that extended beyond the tip of the scales, which corresponded to stage  
 116 BBCH 9 (Meier, 1997). At each observation date, we calculated the percentage of trees that had reached budburst in  
 117 the tree population, dividing the number of trees at BBCH 9 by the total number of trees observed on that date and  
 118 multiplying the result by 100.

## 119 2.3 Temperature data

120 We obtained the mean daily temperature data from the meteorological station nearest to the study sites (Table 1).  
 121 However, there were missing values in the temperature data collected from the stations, especially before 1970. To  
 122 fill these gaps and predict the missing data in order simulate budburst in previous years, we used the SAFRAN  
 123 reanalysis data (grid-resolution of 8\*8 km<sup>2</sup>) (Vidal et al., 2010), which we de-biased by establishing a linear  
 124 regression between the local and corresponding SAFRAN temperature data from September of previous year to June.

## 125 2.4 Model description

126 We introduce a novel model, named the within-population variability (WPV) model, which was constructed to  
 127 predict the progress of budburst in tree populations (i.e., percentage of trees having burst buds at a given date in a  
 128 tree population). We hypothesized that the difference between individuals in the population was reflected in the  
 129 difference of the forcing accumulation requirement ( $F^*$ ).

130 We built the WPV model by modifying a state-of-the-art process-based model that simulated a discrete budburst  
 131 event (i.e., budburst of an individual plant or mean budburst date in a tree population) (Lundell et al., 2020). In short,  
 132 the model represents the release of endodormancy through the accumulation of “chilling” temperatures and simulates  
 133 the ontogenetic growth of buds through the accumulation of “forcing” temperatures. One particularity of the model is  
 134 that ontogenetic growth is regulated by the state of rest break and the prevailing temperature (Lundell et al., 2020;  
 135 Hänninen, 1990; Hänninen and Kramer, 2007; Vegis, 1964). The ontogenetic competence,  $Co$  (a dimensionless [0, 1]

136 multiplier), is applied to represent this regulation (Lundell et al., 2020; Hänninen and Kramer, 2007; Hänninen,  
 137 2016). In the model, budburst is considered to occur at date  $t$  when a given sum of the forcing temperature is reached  
 138 such that  $F(t) \geq F^*$ . In the WPV model, we assumed that  $F^*$  followed a normal distribution at the level of the tree  
 139 population (see Fig. S1 for a flow chart of the model). At each date  $t$ , the model simulates the proportion of the  
 140 population (BP, for **budburst percent**) that has fulfilled the forcing accumulation requirement:

141 
$$F^* = (\mu, \sigma^2) \quad \text{eq.1}$$

142 
$$BP(t) = 0.5 \times (1 + \text{erf}(\frac{F(t)-\mu}{\sigma \times \text{sqrt}(2)})) \times 100 \quad \text{eq.2}$$

143  
 144 Where  $F(t)$  is the forcing degree-day accumulation reached on day  $t$ ,  $\mu$  is the mean of normal distribution,  $\sigma$  is the  
 145 standard deviation of normal distribution, and  $\text{erf}$  is the Gaussian error function.  
 146

147 The forcing accumulation  $F(t)$  is calculated as the integral of a “forcing rate” as follows:

148 
$$F(t) = \sum_{d=270}^t Rf_{act} \quad \text{eq.3}$$

149 Where  $d$  is the start date of forcing accumulation ( $d = \text{DoY } 270$  in the previous year). In this model, the stage of  
 150 dormancy release and the stage of ontogenetic growth can occur simultaneously (i.e., the model belongs to the  
 151 “parallel” model category) (Hänninen, 2016; Chuine and Regnier, 2017). However, the forcing rate  $Rf_{act}$ , which is  
 152 the actual rate of ontogenetic growth, is affected by both temperatures and ontogenetic competence ( $Co$ ). It is  
 153 calculated as follows:

154 
$$Rf_{act}(t) = Rf(t) * Co(t) \quad \text{eq. 4}$$

155 Where  $Rf(t)$  is the potential rate of ontogenetic growth on day  $t$ , and  $Co$  is the ontogenetic competence on day  $t$ ;  
 156 these two variables are calculated as follows:

157 
$$Rf(t) = \begin{cases} 0, & T(t) < T_b \\ T(t) - T_b, & T(t) \geq T_b \end{cases} \quad \text{eq. 5}$$

158 Where  $T(t)$  is the daily mean air temperature on day  $t$  and  $T_b$  is the temperature threshold ( $^{\circ}\text{C}$ ) above which forcing  
 159 accumulation occurs.

160 The ontogenetic competence  $Co$  varies over time and is simulated as:

161 
$$Co(t) = \max(0; \min(1; g \times T(t) + h + \frac{Sr(t)}{100} * (1 - h))) \quad \text{eq.6}$$

162 Where  $Co(t)$  is the ontogenetic competence on day  $t$ , in the range  $[0, 1]$ , which modulates the effect of the state of  
 163 rest break on the rate of ontogenetic growth (see Fig. S2). When  $Co=0$ , ontogenetic growth is stopped. The ability of  
 164 ontogenetic growth is restored between  $Co=0$  and  $Co=1$  with rest breaking. Finally,  $g$  and  $h$  are parameters (Lundell  
 165 et al., 2020), and  $Sr(t)$  is the state of rest break on day  $t$ , which is calculated as follows:

166 
$$Sr(t) = C_{tot}/C_{cri} \quad \text{eq.7}$$

167 Where  $C_{cri}$  is the chilling requirement for rest completion, and  $C_{tot}$  is the actual accumulation of chilling temperature,  
 168 quantified as the number of chilling units (in chill units C.U.) and calculated from DoY=270 of the previous year up  
 169 to day  $t$  as follows:

170 
$$C_{tot} = \sum_{d=270}^t Rc \quad \text{eq.8}$$

171 Where the daily rate of chilling accumulation ( $Rc$ ) is calculated as follows:

172 
$$Rc = \begin{cases} 1, & T(t) < T_c \\ 0, & T(t) \geq T_c \end{cases} \quad \text{eq.9}$$

173 Where  $T_c$  is the temperature threshold ( $^{\circ}\text{C}$ ) below which chilling accumulation occurs.

## 174 2.5 Parameter estimation

175 We calibrated the model using budburst data obtained during the period 2000-2016 in Orsay (all three species:  
 176 hornbeam, oak, chestnut) and then validated it using data from 2017-2022 in Orsay (three species) and from 2000-  
 177 2022 in Barbeau (two species: hornbeam and oak). The model was therefore calibrated over 17 years for the three  
 178 species (Orsay populations, representing 52, 71 and 50 observation dates for hornbeam, oak and chestnut,  
 179 respectively) and validated over 29 site-years for hornbeam and oak (representing 89 and 114 observation dates,  
 180 resp.), and 6 years (29 observation dates) for chestnut. A previous study (Vitasse et al., 2009b) provided evidence of  
 181 similar apparent phenological responses to temperature among populations of the same species located as far as 650  
 182 km apart, which also suggests the low differentiation of phenological traits across populations. Orsay and Barbeau  
 183 populations are separated by a distance of 50 km and experience a similar climate. This is why we used the Barbeau  
 184 data as a validation counterpart to the Orsay data used for calibration. The model predicts the percentage of budburst  
 185 in the population (from 0% to 100% budburst) along with the corresponding date. Thus, we calculated the root mean  
 186 square error (RMSE) over two dimensions (Fig. S3). First, we calculated RMSE over the percentage of budburst in  
 187 the tree population (i.e., comparing the difference between the observed and predicted budburst percent occurring on  
 188 the same day of the year, DoY).

189 
$$RMSE_{BP} = \sqrt{\frac{\sum_{i=1}^n (\sqrt{num} \times (BP_{obs,i} - BP_{pred,i})^2)}{\sum_{i=1}^n \sqrt{num}}} \quad \text{eq.10}$$

190 Where  $RMSE_{BP}$  is the root mean square error for budburst percent (expressed in percent),  $num$  is the number of trees  
 191 observed on a given day of the year,  $BP_{obs,i}$  is the observed percentage of budburst of datum  $i$ ,  $BP_{pred,i}$  is the predicted  
 192 percentage of budburst of same datum, and  $n$  is the total number of data (e.g.,  $n=50$  in a hypothetic case where the  
 193 percentage of budburst has been observed five times per year on average over 10 years in a given population). We  
 194 used  $\sqrt{num}$  as a weight in the calculation of squared errors to compensate for the fact that a very large number of  
 195 trees (i.e.,  $>300$  trees) were observed at some dates: these observations are more representative of the actual  
 196 percentage of budburst in the population (as compared to observations established for a smaller number of trees),  
 197 although they also tend to overrepresent them in the calculation of errors.

198 We then calculated the RMSE of dates (i.e., comparing the difference, in number of days, between the observations  
 199 and predictions for the same percentage of budburst; Fig. S3).

$$200 \quad RMSE_{DoY} = \sqrt{\frac{\sum_{i=1}^n (\sqrt{num} \times (DoY_{obs,i} - DoY_{pred,i}))^2}{\sum_{i=1}^n \sqrt{num}}} \quad \text{eq.11}$$

201 Where  $RMSE_{DoY}$  is the root mean square error for the budburst date (in days),  $num$  is the number of trees observed,  
 202  $DoY_{obs,i}$  is the observed date of budburst of datum  $i$  (e.g., the date when we observed 24% budburst for the  
 203 population of interest in a given year),  $DoY_{pred,i}$  is the predicted date of budburst of the same datum (e.g., the date  
 204 when the model predicted 24% budburst in the same tree population and year), and  $n$  is the total number of data.

205 Finally, we calculated the total RMSE as follows:

$$206 \quad RMSE_{tot} = RMSE_{BP} + RMSE_{DoY} \quad \text{eq.12}$$

207 We used  $RMSE_{tot}$  as an aggregate, multi-objective cost function (similar to, e.g., Keenan et al., 2011) during the  
 208 calibration procedure. We used the function *optim* to calibrate the model parameters with R statistical software  
 209 v.4.0.3 (R Development Core Team, 2020). In order to ensure that the *optim* algorithm reached the global minimum  
 210 of the cost function, we ran it 768 times for each calibration, starting from different, random combinations of initial  
 211 parameters, which appear in Table S1, and retained the parameter set yielding the overall lowest  $RMSE_{tot}$ . One  
 212 possible issue with aggregate multi-objective cost functions such as eq. 12 is that the same minimum  $RMSE_{tot}$  can be  
 213 achieved with multiple combinations of  $RMSE_{BP}$  and  $RMSE_{DoY}$ . In order to evaluate this, we produced a figure to  
 214 show the relation between  $RMSE_{tot}$  and  $RMSE_{BP}$  or  $RMSE_{DoY}$  (Fig. S4). Using  $RMSE_{tot}$  instead of  $RMSE_{BP}$  or  
 215  $RMSE_{DoY}$  yielded very similar results, with differences of less than 1% or 1 day when using  $RMSE_{tot}$  instead of  
 216  $RMSE_{BP}$  or  $RMSE_{DoY}$ , respectively (Table S2). In addition to RMSE, we used mean bias error and the correlation  
 217 coefficient ( $r$ ) and  $p$  value to evaluate the model forecast accuracy (in terms of budburst percentage or days), which  
 218 are calculated as follows:

$$219 \quad \text{mean bias} = \frac{1}{N} \sum_{i=1}^N (obs_i - pred_i) \quad \text{eq.13}$$

220 Where  $obs_i$  and  $pred_i$  are the  $i^{\text{th}}$  observation and prediction, respectively,  $N$  is the number of observations.

$$221 \quad r = \frac{\sum_{i=1}^N (obs_i - obs_{mean})(pred_i - pred_{mean})}{\sqrt{\sum_{i=1}^N (obs_i - obs_{mean})^2} \sqrt{\sum_{i=1}^N (pred_i - pred_{mean})^2}}$$

222 Where  $obs_{mean}$  and  $pred_{mean}$  are the mean of observation and prediction, respectively.

## 223 2.6 Evaluating the modelled $F^*$ distributions

224 To validate the modelled  $F^*$  distribution, we simulated the distribution of the forcing accumulation at the date of  
 225 each BP observation. Because there are different observed BP in each year. We binned the observed BP data into 11  
 226 groups (i.e., BP0, BP10, BP20...BP100, e.g., we regard the data between BP5 (date at which 5% trees burst buds) to  
 227 BP15 (date at which 15% trees burst buds) as group ‘‘BP10’’; note that BP0 refers to dates at which 5% or less trees



228 have burst buds, and BP100 refers to dates at which 95% or more trees have burst buds). Then we used a sigmoid  
229 function to simulate the relation between BP and averaged corresponding forcing accumulation across all the years.  
230 We also calculated their first derivatives (i.e., the increasing of BP per unit of forcing accumulation). Moreover, we  
231 calculated the distribution of observed BP across all the years.

## 232 **2.7 Evaluating the response of the within-population variability of budburst to climate warming**


233 We used our model to predict budburst in the past (1961-2022) using historical daily mean temperature data and gap-  
234 filled data using debiased SAFRAN reanalysis of temperatures (see above).

235 As explained earlier, our model simulates the percentage of budburst in a tree population at a given date. To evaluate  
236 the response of the WPV of budburst to climate warming, we focused on the particular dates at which 20% and 80%  
237 of trees in a given population had reached budburst (termed BP20 and BP80, respectively) and the duration between  
238 these two dates ( $DurBB = BP80 - BP20$ ), which we consider to represent the variability of budburst within the  
239 population for a given year. BP20 represents the “beginning” of budburst in the tree population, whereas BP80  
240 represents its “end.” We chose these quantiles instead of more extreme quantiles of distribution (e.g., 5% and 95%),  
241 because they are well represented in our dataset (Fig. 1), thus implying higher model accuracy. For sake of model  
242 evaluation, we calculated the DurBB in observed phenology data. Specifically, we selected years which had records  
243 before BP20 and after BP80. Then the date of BP20 or BP80 were calculated by using the nearest two data (one is  
244 below BP20 or BP80, another is above BP20 or BP80) through interpolation (e.g., 15 % budburst percent is on DoY  
245 80 and 25 % budburst percent is on DoY 84. We can obtain the date of BP20 by interpolation, that is DoY 82).

## 246 **2.8 Statistical analyses**

247 For each population, we quantified by linear regression the sensitivity of budburst date (BP20 and BP80) and the  
248 DurBB to time ( $\text{days year}^{-1}$ ) and to Jan-May temperature ( $\text{days } ^\circ\text{C}^{-1}$ ). Analysis of Variance (ANOVA) was used to  
249 analysis whether the significance of the regression slopes ( $\alpha=0.05$ ). All simulations and statistical analyses were  
250 carried out with R statistical software v.4.0.3 (R Development Core Team, 2020).





251  
 252 **Fig. 1. Observed percentage of budburst in five tree populations during the period 2000-2022. The size of the points is**  
 253 **scaled with the square root of the number of trees observed. The lines connect the dates of the same year.**

254 **3. Results**


255 **3.1 Phenological observations**

256 Figure 1 shows the observed percentages of budburst in the five tree populations monitored from 2000 to 2022.  
 257 These percentage data were established based on 48,442 observations of budburst collected from individual trees.

258 Among the species, hornbeam was the earliest to reach budburst, typically over DoY 70-100, followed by oak over  
259 DoY 90-110, and finally, chestnut over DoY 100-130. The budburst dates of the oak and hornbeam populations at  
260 Barbeau and Orsay were very close, with average differences of 2 and 1 days (Table S3). The duration of budburst in  
261 the population (DurBB) (i.e., time interval, in days, during which the proportion of trees having reached budburst  
262 increases from 20% to 80%) differs for each species depending on the site and year, with a mean of 8 days over the  
263 whole dataset and ranging from 3 days for hornbeam at Orsay in 2018 and 2021 to 21 days for oak at Orsay in 2012  
264 (Fig.1).


### 265 **3.2 Model performance**

266 For all the populations considered here, the WPV model predicted with good accuracy the progress of budburst in  
267 tree populations during spring as well as the interannual variability of budburst (Fig. 2, Fig. 3; see Fig. S5 for a  
268 comparison of observed and simulated time series). **The model predicted the percentage of budburst in tree**  
269 **populations with an error ( $RMSE_{BP}$ ) of  $16\% \pm 0.2\%$  for the calibration dataset (correlation coefficient of predictions**  
270 **vs. observations:  $0.81 \pm 0.02$ ,  $P < 0.001$ ) and  $20\% \pm 2.1\%$  for the validation dataset (correlation:  $0.76 \pm 0.07$ ,  $P < 0.001$ ).**  
271 **This corresponded to prediction errors for the date of budburst ( $RMSE_{DOY}$ ) of  $3.9 \pm 0.6$  days for the calibration**  
272 **dataset (correlation:  $0.89 \pm 0.04$ ,  $P < 0.001$ ) and  $5.6 \pm 0.3$  days for the validation dataset (correlation:  $0.81 \pm 0.10$ ,**  
273  **$P < 0.001$ ).** This compared well to the time resolution of the phenological observations (3-7 days). The mean bias was  
274 less than 1 day (Fig. 3).



275

276 **Fig. 2. Evaluation of the within-population variability (WPV) model predicting the budburst percentage over calibration**  
 277 **(red points) and validation (blue points) data. The points of circle are observed in Orsay and of triangle are observed in**  
 278 **Barbeau. The points establish the correspondence between the observed and predicted percentage of budburst on an**  
 279 **observation day in the population of interest. The one-to-one relation is shown as the black line. RMSE which is the root**  
 280 **mean square error for the budburst percentage, mean bias and correlation coefficient (r) are shown. There are 52, 71 and**  
 281 **50 points (i.e., observation dates) for calibration and 89, 114, 29 points for validation for hornbeam, oak and chestnut,**  
 282 **respectively. P-values of the correlation coefficients appear as (\*: P<0.05, \*\*: P<0.01, \*\*\*: P<0.001).**



283


284 **Fig. 3. Evaluation of the within-population variability (WPV) model predicting budburst dates over calibration (red points)**  
 285 **and validation (blue points) data. The points of circle are observed in Orsay and of triangle are observed in Barbeau. The**  
 286 **points establish the correspondence between the observed and predicted budburst date on one observation day in the**  
 287 **population of interest. The one-to-one relation is shown as the black line. RMSE which is root mean square error for the**  
 288 **budburst percentage, mean bias and correlation coefficient (r) are shown. There are 52, 71 and 50 points (i.e., observation**  
 289 **dates) for calibration and 89, 114, 29 points for validation for hornbeam, oak and chestnut, respectively. P-values of the**  
 290 **correlation coefficients appear as (\*: P<0.05, \*\*: P<0.01, \*\*\*: P<0.001).**

291 **3.3 Parameter variations across species**

292 As mentioned earlier, we assumed that the forcing requirement ( $F^*$ ) followed a normal distribution. The calibration  
 293 procedure yielded a set of distribution curves that differed across species (Fig. 4). We observed that the distribution  
 294 of  $F^*$  had a highest mean and standard deviation for oak compared with hornbeam and chestnut (Fig. 4, Table 2).  
 295 The distributions of  $F^*$  compared well to the actual distribution of forcing accumulation established from  
 296 observations (Fig. 5b, e, h), validating the choice of the normal distribution. However, the modelled distribution did  
 297 not overlap exactly the distribution established from observed data, because the distribution of observations along the  
 298 BP scale was uneven (Fig. 5c, f, i). The temperature threshold for chilling accumulation ( $T_c$ ) ranged from 9.7°C for  
 299 chestnut to 10.5°C for hornbeam and oak (Table 2). The temperature threshold for forcing accumulation ( $T_f$ ) ranged  
 300 from 3.9°C for hornbeam to 7.7°C for chestnut (Table 2, Fig. S2). In all species, buds could not begin ontogenetic  
 301 growth until the accumulation of chilling to a certain extent (i.e., parameter h was negative for all populations, Table  
 302 2). We found that the threshold of chilling accumulation necessary for the onset of forcing accumulation (i.e., value  
 303 of Sr(t) from which Co becomes positive) was very high for early species and decreased for late species (e.g., value  
 304 of h increased approximately from -0.98 in hornbeam to 0 in chestnut; see Table 2 and Fig. S2). Prevailing  
 305 temperatures could compensate for the lack of chilling accumulation (positive parameter g; Table 2) in hornbeam  
 306 and oak, but not in chestnut (g=0).


307  
 308 **Table 2. Parameter values of the WPV model for three populations.  $\mu$  (°C-days) and  $\sigma$  (°C-days) are the mean and**  
 309 **standard deviation of the distribution of  $F^*$ , respectively (Eqn. 1).  $T_b$  and  $T_c$  (°C) are the threshold temperatures for the**  
 310 **accumulation of forcing and chilling temperatures, respectively (Eqns. 5 and 9).  $g$  (°C<sup>-1</sup>) and  $h$  (dimensionless) are the**  
 311 **parameters determining the interactive effect of the state of rest break and the prevailing air temperature on the**  
 312 **ontogenetic competence (Eqn. 6).  $C_{cri}$  (number of days) is the chilling requirement of rest completion.**

Species	Site	$\mu$	$\sigma$	$T_b$	$T_c$	$g$	$h$	$C_{cri}$
<i>Carpinus</i>	Orsay	138.4	29.6	3.9	10.5	0.0080	-0.98	155.5
<i>Quercus</i>	Orsay	150.4	37.2	5.3	10.5	0.0032	-0.89	153.0
<i>Castanea</i>	Orsay	131.2	30.7	7.7	9.7	0.0000	-0.02	144.9



313

314 **Fig. 4.** Normal distribution of the forcing requirement ( $F^*$ ) for three tree species.




315

316 **Fig. 5.** Evaluating the modelled  $F^*$  distributions. Subplots (a, d and g) represent the relation between budburst percentage  
 317 (BP) and forcing accumulation. The black points and error bars represent the forcing accumulation required to reach a

318 given budburst percentage in observed data (average across years  $\pm$  one standard deviation). The red curves represent a  
 319 sigmoid function fitted to the black dots (a, d, g), and its first derivative (b, e, h). The blue curve represents predictions  
 320 based on the parameters in Table 2. Subplots (b, e and h) represent the increasing of BP per unit of forcing accumulation.  
 321 Subplots (c, f and i) show the distribution of observed data points in the budburst dataset.

### 322 3.4 Retrospective analysis for within-population variability of budburst


323 Over the past six decades (1961-2022), spring average temperature increased by  $+1.9^{\circ}\text{C}$  in Orsay and  $+1.4^{\circ}\text{C}$  in  
 324 Barbeau (Fig. S6). Over this time period, our retrospective simulations suggest that the beginning (20%, BP20) and  
 325 end (80%, BP80) of budburst in tree populations has advanced significantly for all the species (Fig. 6), with  
 326 respectively  $1.7 \pm 0.6$  days decade $^{-1}$  (mean  $\pm$  SD across species) and  $1.8 \pm 0.4$  days decade $^{-1}$  and apparent  
 327 temperature sensitivities of  $5.8 \pm 0.4$  days  $^{\circ}\text{C}^{-1}$  and  $5.6 \pm 0.4$  days  $^{\circ}\text{C}^{-1}$ . These similar trends regarding the beginning  
 328 and end of budburst result in an unchanged duration of the budburst period (DurBB in the considered populations  
 329 over the past 62 years (no trend in DurBB is significantly different from zero in Fig. 7,  $P > 0.05$ ). **Meanwhile, the**  
 330 **results about temperature sensitivity were similar which were negative for BP20 and BP80 for all three species based**  
 331 **on pre-season temperature preceding budburst (Table S4).** Notably, the interannual variability of DurBB was large  
 332 (Fig.6), and fairly simulated by our model (RMSE of  $3.4 \pm 1.8$  days).



333



334 Fig. 6. Simulated occurrence of the beginning (20%, BP20 in blue) and end (80%, BP80 in red) of budburst using the  
 335 WPV model for three tree species during the period 1961-2022. The fitted lines highlight the trends over the past 62 years.  
 336 Text in blue (red) shows the sensitivity of BP20 (BP80) to time and mean spring temperature (from January to May),  
 337 respectively. **The trends in days/decade and days/°C are displayed on the figure.** The sensitivity values are tested by linear  
 338 regression analyses (\*:  $P < 0.05$ , \*\*:  $P < 0.01$ , \*\*\*:  $P < 0.001$ ) and adjusted coefficient of determination ( $R^2_{adj}$ ) is shown.



339

340 Fig. 7. Simulated duration of budburst in the population (DurBB) using the WPV model for three tree species during the  
 341 period 1961-2022. The fitted line depicts the change in DurBB over the past 62 years. The sensitivity of DurBB to time and  
 342 mean spring temperature (from January to May) are tested by linear regression analyses (ns:  $P > 0.05$ ) and adjusted  
 343 coefficient of determination ( $R^2_{adj}$ ) is shown. **The trends in days/decade and days/°C are displayed on the figure.** The  
 344 black points are the actual durations of budburst observed in the data (i.e., restricted to years when both BP20 and BP80  
 345 are available in a population).

#### 346 4. Discussion

347 To the best of our knowledge, this paper presents the first model simulating the within-population variability of  
 348 budburst in tree populations. An added value of this model is that it can simulate the duration of budburst in tree  
 349 populations. The central hypothesis of the model is that  $F^*$ , the amount of accumulated forcing temperature required  
 350 for trees to budburst, follows a normal distribution in tree populations. The ability of the model to simulate the

351 dynamics of budburst over the calibration and validation data, as well as the good agreement between the observed  
352 and the simulated  $F^*$  distributions (Fig. 5), lend support to this hypothesis for all the species and populations  
353 considered. Our model yielded RMSE for the validation data (5.4 to 5.9 days), which are close to the temporal  
354 resolution of the spring phenology observation (from 2-7 days) and similar to the typical prediction accuracy of  
355 models simulating discrete (i.e., population average) budburst dates (e.g., Basler, 2016).

356 The variability in the timing of budburst among individuals in tree populations is considered to be mainly determined  
357 by genetic diversity (Bontemps et al., 2016; Delpierre et al., 2017; Jarvinen et al., 2003; Rousi and Heinonen, 2007;  
358 Rusanen et al., 2003) followed by the influence of the microenvironment (Delpierre et al., 2017; Rousi and Heinonen,  
359 2007). The phenological ranking of individuals is largely conserved in tree populations (Delpierre et al., 2017),  
360 leading to the identification of “early”, “intermediate” and “late” trees (Malyshev et al., 2022). Further, the  
361 distribution of budburst categories is not uniform in natural tree populations, with numerous “intermediate”  
362 individuals and comparatively fewer “early” and “late” trees (Malyshev et al., 2022; Chesnoiu et al., 2009; Zohner et  
363 al., 2018; Caradonna et al., 2014), which lends further support to a unimodal distribution such as the normal law. **Our**  
364 **data further show the same trees are always early/late within the population with corresponding low/high forcing**  
365 **accumulation requirements (Fig. S7).** Our model reproduces this phenomenon, with categories of “early,”  
366 “intermediate,” and “late” trees corresponding to increasing values of  $F^*$ . This core assumption of the model is  
367 supported by previous empirical studies, which observe that the variability of  $F^*$  could represent the variability of  
368 budburst among trees (Langvall et al., 2001; Rousi and Heinonen, 2007). Nevertheless, we could have chosen to  
369 assign the variance among individuals to one or several other parameters of the model, related to the fact that genetic  
370 variations may affect any of the plant traits determining the modelled parameters. For instance, Gauzere et al. (2019)  
371 found that the temperature yielding mid-forcing during ecodormancy ( $T_{50}$ ) was more sensitive than  $F^*$  in the  
372 UniChill model, which suggests that this parameter is another good candidate for identifying the phenological  
373 behavior of individual trees in a population. Thus, we constructed a model assuming that the threshold for forcing  
374 temperature ( $T_b$ , i.e., parameter of our model analogous to  $T_{50}$ ) followed a normal distribution, whereas  $F^*$  was fitted  
375 as a constant parameter for the population. This model fitted the data less effectively in both the calibration and  
376 validation steps (see Fig. S8 and S9 compared with Fig. 2 and 3), which further supports our decision to assign the  
377 among-individual variance to  $F^*$ . **We further tested to assign the among-individual variance to the parameters for**  
378 **phase of dormancy release (e.g., chilling requirement of rest completion ( $C_{cri}$ ) and the threshold temperatures for the**  
379 **accumulation of chilling temperatures ( $T_c$ )), also using a normal distribution. However, the model fitted the data even**  
380 **worse than in our attempt of fitting a normal distribution of  $T_b$ .** Questions remain regarding the actual shape of the  
381  $F^*$  distribution. Indeed, natural selection can lead to traits that are not normally distributed (Caradonna et al., 2014),  
382 and uneven distribution of observations may contribute to the non-perfect overlapping of observed and simulated  $F^*$   
383 distributions (Fig. 5). However, earlier results (Vallet, 2020) showed that the form of the distribution had little  
384 influence on the prediction accuracy.

385 We built the WPV model based on a two-phase parallel model framework, which describes the cumulative effect of  
386 chilling and forcing temperatures on the endodormancy and ecodormancy phases, respectively (Hänninen, 2016;  
387 Hänninen and Kramer, 2007; Lundell et al., 2020; Chuine and Regniere, 2017). This model structure is in line with

388 our current understanding of the physiological and molecular basis of dormancy in which the dynamics of the  
389 dormancy mechanism are emphasized as opposed to a strict classification between the dormancy stages (Lundell et  
390 al., 2020; Cooke et al., 2012). In this study, the threshold of chilling accumulation is up to 10.5°C for oak and  
391 hornbeam. It is consistent with the experimental results in Baumgarten et al. (2021) which challenge the common  
392 assumption that optimal chilling temperatures range ca. 4–6°C, showing 10°C is also effective for chilling  
393 accumulation in six dominant temperate European tree species including oak. Furthermore, the model uses the  
394 concept of ontogenetic competence ( $Co$ ) to simulate the process of regulation for the rate of ontogenetic growth by  
395 the state of rest break, a phenomenon that has found support in phenological experiments (Lundell et al., 2020;  
396 Zhang et al., 2022). Our results demonstrate that in the investigated species,  $Co$  is 0 until dormancy is released to a  
397 certain extent (Fig. S2), that is, ontogenetic growth cannot start before a certain amount of chilling accumulation has  
398 been reached, which is consistent with previous findings (Lundell et al., 2020; Zhang et al., 2022). According to the  
399 calibrated parameter values, ontogenetic competence is also influenced by the prevailing temperature, although the  
400 effect is minimal. Indeed, parameter  $g$ , which is related to the effect of the prevailing temperature, ranges from 0 to  
401 0.0080 (Table 2), which is comparable to values found in a previous study (Lundell et al., 2020). To some extent in  
402 this model, one consequence is that the effect of the prevailing temperature can compensate for the deficiency of  
403 chilling accumulation.

404 Beyond introducing a model to describe the WPV of budburst in tree populations, our study aimed to quantify the  
405 response of the duration of budburst (DurBB) to climate warming. We used temperature data to simulate the  
406 occurrence of 20% (BP20) and 80% (BP80) budburst, and DurBB over the past decades. Our results suggest that the  
407 start and end of budburst in tree populations have advanced over the past 62 years with climate warming (Fig. 6),  
408 which is consistent with previous results showing advances in the population average dates of budburst (Wenden et  
409 al., 2020; Menzel et al., 2006; Fu et al., 2015). In addition, our model simulates sensitivities of budburst to time and  
410 temperature that are comparable to values reported earlier (Vitasse et al., 2009b, see Table S5). Our results point to  
411 significant sensitivities to both time and temperature for oak as well as significant sensitivity to temperature for  
412 hornbeam, which is consistent with the results of Vitasse et al. (2009b). **The advancement of budburst would**  
413 **increase the possibility of spring frost damage (Liu et al., 2018; Vitasse et al., 2018), influencing tree physiology and**  
414 **growth with possible impacts on the productivity of forests (Vitasse et al., 2019) or even the distribution of tree**  
415 **species (Chuine, 2010).**

416 Our retrospective simulations suggest that there was not trend in the duration of budburst in tree populations, DurBB,  
417 over the past 62 years (Fig. 7), in spite of climate warming (Fig. S6). Since both BP20 and BP80 advanced at a  
418 similar rate, DurBB did not evolve over time over the 1961-2022 period. Interestingly, the analysis of temperature  
419 data revealed no significant warming in the period of time from BP20 to BP80 over the past decades ( $P>0.05$ , Fig.  
420 S10). This could explain why DurBB (time interval between BP20 and BP80) did not change over time, in spite of  
421 the strong trends in both BP20 and BP80, caused by climate warming. **However, interannual variability of DurBB**  
422 **was large, which was reproduced by the WPV model (Fig. 7).** Moreover, our study sites are located in the temperate  
423 zone, at the heart (for oak and hornbeam) and at the north (chestnut) of our study species distribution areas (Caudullo  
424 et al., 2017). At those sites, trees can accumulate enough chilling, or at least, chilling accumulation is not a limitation

425 for ontogenetic growth in nature so far, meaning that budburst is still advancing (Wenden et al., 2020; Piao et al.,  
426 2019). Thus, the phenomenon by which DurBB increased with insufficient chilling accumulation in a given  
427 population (see Zhang et al., 2021, their Fig. 2, 3, 4 for evidence in subtropical trees) did not appear in our  
428 retrospective simulations. **However, we can infer that if chilling accumulation can't be fulfilled under future,  
429 continuous climate warming, it will take more time to fulfill the forcing requirement for late trees with a high forcing  
430 requirement, leading to the prolonging of DurBB. A longer duration of budburst would increase the possibility of  
431 damage (i.e., freezing, insect damage).**

432 We acknowledge that the projections of the WPV of budburst produced by the model are uncertain, first and  
433 foremost because the parameter values were inferred from observation data collected in natural conditions as  
434 opposed to controlled experiments (Hanninen et al., 2019). Another cause of uncertainty is the ability of the  
435 phenological response of plants to acclimatize to the changing climate (Bennie et al., 2010). Under the hypothesis of  
436 plant acclimatization, the parameters of the WPV model could have changed over the past decades, and would  
437 further change with ongoing climate warming. Consequently, related experiments are urgently needed to improve  
438 our understanding of the WPV of budburst to infer more reliable parameters and analyze the behavior of phenology  
439 models in different climates (Hanninen et al., 2019). However, because our model addresses for the first time  
440 explicitly the within-population variation of the physiological traits affecting phenology, it can contribute as a  
441 framework for future experimental studies. In our study, we only considered the effect of temperature on budburst.  
442 **However, other environmental factors may also affect budburst (e.g., photoperiod, soil moisture and the interaction  
443 between factors). Previous studies showed that photoperiod is expected to modulate the timing of budburst in late-  
444 successional species such as oak and chestnut, but not in early-successional species such as hornbeam (Basler and  
445 Korner, 2012), but see a counter-example on oak in Malyshev et al. (2018). Moreover, photoperiod may have a more  
446 complex interaction mechanism with temperature in terms of regulating the time of budburst (Meng et al., 2021).  
447 And negative correlations between spring soil moisture and the start of the growing season were found in the  
448 Mongolian Plateau (Luo et al., 2021).** We envision that improved versions of the WPV of budburst could be  
449 proposed based on a more comprehensive understanding of the potential mechanism between phenology and  
450 environmental factors in the future.

## 451 **5. Conclusion**

452 In conclusion, our work presents a novel model, simulating the continuity of budburst in tree populations in spring.  
453 This phenological model can be adapted to the study of other stages of the tree phenological cycle, which are all of  
454 continuous nature in tree populations (e.g., leaf senescence, wood formation etc.). We found budburst was advanced  
455 in the past 62 years due to climate warming. However, the duration of budburst period of population was not affected  
456 by increasing temperature. This is the first model simulating the within population variability of budburst in the  
457 population. It provides a basis for implementation of a module in models directly interested in the within-population  
458 variability of phenological and other functional traits (e.g., physio-demo-genetic models). It can also be used as a  
459 stand-alone, to study the dynamics of phenological traits from the scale of individuals to the population and  
460 community in the context of climate change.

461 **Code and data availability**

462 The related phenology data and R code for the phenological model are openly accessible under  
463 <https://doi.org/10.5281/zenodo.7962840> and <https://doi.org/10.5281/zenodo.10020474>, respectively.

464 **Authors' contributions**

465 ND and JL designed the research. ND, JL, AM, GV, DB collected phenological data. JL and ND performed the  
466 research. JL wrote the manuscript with substantial inputs from all co-authors.

467 **Competing interests**

468 The authors declare that they have no conflict of interest.

469 **Acknowledgements**

470 We acknowledge Eric Dufrêne for setting up the phenological surveys. We are also grateful to Eric Dufrêne and  
471 Jean-Yves Pontalier for their invaluable contributions regarding the collection of phenological data. **We are also**  
472 **grateful for the constructive comments provided by Marc Peaucelle and Yongshuo H. Fu.** This work was supported  
473 by the China Scholarship Council (202008330320) and the ANR (FOREPRO project, grant number ANR-19-CE32-  
474 0008).

475 **References**

- 476 Alberto, F., Bouffier, L., Louvet, J. M., Lamy, J. B., Delzon, S., and Kremer, A.: Adaptive responses for seed and  
477 leaf phenology in natural populations of sessile oak along an altitudinal gradient, *J Evol Biol*, 24, 1442-1454,  
478 10.1111/j.1420-9101.2011.02277.x, 2011.
- 479 Basler, D.: Evaluating phenological models for the prediction of leaf-out dates in six temperate tree species across  
480 central Europe, *Agricultural and Forest Meteorology*, 217, 10-21, 10.1016/j.agrformet.2015.11.007, 2016.
- 481 Basler, D. and Korner, C.: Photoperiod sensitivity of bud burst in 14 temperate forest tree species, *Agricultural and*  
482 *Forest Meteorology*, 165, 73-81, 10.1016/j.agrformet.2012.06.001, 2012.
- 483 Baumgarten, F., Zohner, C. M., Gessler, A., and Vitasse, Y.: Chilled to be forced: the best dose to wake up buds  
484 from winter dormancy, *New Phytol*, 230, 1366-1377, 10.1111/nph.17270, 2021.
- 485 Bennie, J., Kubin, E., Wiltshire, A., Huntley, B., and Baxter, R.: Predicting spatial and temporal patterns of bud-burst  
486 and spring frost risk in north-west Europe: the implications of local adaptation to climate, *Global Change Biology*,  
487 16, 1503-1514, 10.1111/j.1365-2486.2009.02095.x, 2010.
- 488 Blanquart, F., Kaltz, O., Nuismer, S. L., and Gandon, S.: A practical guide to measuring local adaptation, *Ecol Lett*,  
489 16, 1195-1205, 10.1111/ele.12150, 2013.
- 490 Bontemps, A., Lefevre, F., Davi, H., and Oddou-Muratorio, S.: In situ marker-based assessment of leaf trait  
491 evolutionary potential in a marginal European beech population, *J Evol Biol*, 29, 514-527, 10.1111/jeb.12801, 2016.

492 CaraDonna, P. J., Iler, A. M., and Inouye, D. W.: Shifts in flowering phenology reshape a subalpine plant community,  
493 Proc Natl Acad Sci U S A, 111, 4916-4921, 10.1073/pnas.1323073111, 2014.

494 Caudullo, G., Welk, E., and San-Miguel-Ayanz, J.: Chorological maps for the main European woody species, Data  
495 Brief, 12, 662-666, 10.1016/j.dib.2017.05.007, 2017.

496 Chen, L., Huang, J. G., Ma, Q., Hanninen, H., Tremblay, F., and Bergeron, Y.: Long-term changes in the impacts of  
497 global warming on leaf phenology of four temperate tree species, Global Change Biology, 25, 997-1004,  
498 10.1111/gcb.14496, 2019.

499 Chen, L., Huang, J. G., Ma, Q., Hanninen, H., Rossi, S., Piao, S., and Bergeron, Y.: Spring phenology at different  
500 altitudes is becoming more uniform under global warming in Europe, Global Change Biology, 24, 3969-3975,  
501 10.1111/gcb.14288, 2018.

502 Chen, X. Q., Wang, L. X., and Inouye, D.: Delayed response of spring phenology to global warming in subtropics  
503 and tropics, Agricultural and Forest Meteorology, 234, 222-235, 10.1016/j.agrformet.2017.01.002, 2017.

504 Chesnoiu, E. N., Șofletea, N., Curtu, A. L., Toader, A., Radu, R., and Enescu, M.: Bud burst and flowering  
505 phenology in a mixed oak forest from Eastern Romania, Annals of Forest Research, 52, 199-206,  
506 doi:10.15287/afr.2009.136, 2009.

507 Chuine, I. and Regniere, J.: Process-Based Models of Phenology for Plants and Animals, Annual Review of Ecology,  
508 Evolution, and Systematics, 48, 159-182, 10.1146/annurev-ecolsys-110316-022706, 2017.

509 Chuine, I.: Why does phenology drive species distribution?, Philosophical Transactions of the Royal Society B-  
510 Biological Sciences, 365, 3149-3160, 10.1098/rstb.2010.0142, 2010.

511 Cooke, J. E., Eriksson, M. E., and Junttila, O.: The dynamic nature of bud dormancy in trees: environmental control  
512 and molecular mechanisms, Plant Cell Environ, 35, 1707-1728, 10.1111/j.1365-3040.2012.02552.x, 2012.

513 Dantec, C. F., Ducasse, H., Capdevielle, X., Fabreguettes, O., Delzon, S., and Desprez-Loustau, M. L.: Escape of  
514 spring frost and disease through phenological variations in oak populations along elevation gradients, Journal of  
515 Ecology, 103, 1044-1056, 10.1111/1365-2745.12403, 2015.

516 Delpierre, N., Guillemot, J., Dufrene, E., Cecchini, S., and Nicolas, M.: Tree phenological ranks repeat from year to  
517 year and correlate with growth in temperate deciduous forests, Agricultural and Forest Meteorology, 234, 1-10,  
518 10.1016/j.agrformet.2016.12.008, 2017.

519 Delpierre, N., Dufrene, E., Soudani, K., Ulrich, E., Cecchini, S., Boe, J., and Francois, C.: Modelling interannual and  
520 spatial variability of leaf senescence for three deciduous tree species in France, Agricultural and Forest Meteorology,  
521 149, 938-948, 10.1016/j.agrformet.2008.11.014, 2009.

522 Delpierre, N., Vitasse, Y., Chuine, I., Guillemot, J., Bazot, S., Rutishauser, T., and Rathgeber, C. B. K.: Temperate  
523 and boreal forest tree phenology: from organ-scale processes to terrestrial ecosystem models, *Annals of Forest  
524 Science*, 73, 5-25, 10.1007/s13595-015-0477-6, 2016.

525 Denechere, R., Delpierre, N., Apostol, E. N., Berveiller, D., Bonne, F., Cole, E., Delzon, S., Dufrene, E., Gressler, E.,  
526 Jean, F., Lebourgeois, F., Liu, G., Louvet, J. M., Parmentier, J., Soudani, K., and Vincent, G.: The within-population  
527 variability of leaf spring and autumn phenology is influenced by temperature in temperate deciduous trees, *Int J  
528 Biometeorol*, 65, 369-379, 10.1007/s00484-019-01762-6, 2021.

529 Du, Y. J., Pan, Y. Q., and Ma, K. P.: Moderate chilling requirement controls budburst for subtropical species in  
530 China, *Agricultural and Forest Meteorology*, 278, 107693, ARTN 107693, 10.1016/j.agrformet.2019.107693, 2019.

531 Fu, Y. H., Zhang, X., Piao, S., Hao, F., Geng, X., Vitasse, Y., Zohner, C., Penuelas, J., and Janssens, I. A.: Daylength  
532 helps temperate deciduous trees to leaf-out at the optimal time, *Global Change Biology*, 25, 2410-2418,  
533 10.1111/gcb.14633, 2019.

534 Fu, Y. H., Zhao, H., Piao, S., Peaucelle, M., Peng, S., Zhou, G., Ciais, P., Huang, M., Menzel, A., Penuelas, J., Song,  
535 Y., Vitasse, Y., Zeng, Z., and Janssens, I. A.: Declining global warming effects on the phenology of spring leaf  
536 unfolding, *Nature*, 526, 104-107, 10.1038/nature15402, 2015.

537 Gauzere, J., Lucas, C., Ronce, O., Davi, H., and Chuine, I.: Sensitivity analysis of tree phenology models reveals  
538 increasing sensitivity of their predictions to winter chilling temperature and photoperiod with warming climate,  
539 *Ecological Modelling*, 411, ARTN 108805, 10.1016/j.ecolmodel.2019.108805, 2019.

540 Gauzere, J., Delzon, S., Davi, H., Bonhomme, M., Garcia de Cortazar-Atauri, I., and Chuine, I.: Integrating  
541 interactive effects of chilling and photoperiod in phenological process-based models. A case study with two  
542 European tree species: *Fagus sylvatica* and *Quercus petraea*, *Agricultural and Forest Meteorology*, 244-245, 9-20,  
543 10.1016/j.agrformet.2017.05.011, 2017.

544 Hanninen, H., Kramer, K., Tanino, K., Zhang, R., Wu, J., and Fu, Y. H.: Experiments Are Necessary in Process-  
545 Based Tree Phenology Modelling, *Trends Plant Sci*, 24, 199-209, 10.1016/j.tplants.2018.11.006, 2019.

546 Hänninen, H.: Modelling bud dormancy release in trees from cool and temperate regions, *Acta Forestalia Fennica*,  
547 10.14214/aff.7660, 1990.

548 Hänninen, H.: Boreal and temperate trees in a changing climate: Modelling the ecophysiology of seasonality,  
549 Dordrecht: Springer Science +Business Media, 2016.

550 Hänninen, H. and Kramer, K.: A framework for modelling the annual cycle of trees in boreal and temperate regions,  
551 *Silva Fennica*, 41, 167-205, 2007.

552 Hart, S. P., Schreiber, S. J., and Levine, J. M.: How variation between individuals affects species coexistence, *Ecol  
553 Lett*, 19, 825-838, 10.1111/ele.12618, 2016.



554 Jarvinen, P., Lemmetyinen, J., Savolainen, O., and Sopanen, T.: DNA sequence variation in BpMADS2 gene in two  
555 populations of *Betula pendula*, *Mol Ecol*, 12, 369-384, 10.1046/j.1365-294x.2003.01740.x, 2003.

556 Jewaria, P. K., Hanninen, H., Li, X., Bhalerao, R. P., and Zhang, R.: A hundred years after: endodormancy and the  
557 chilling requirement in subtropical trees, *New Phytol*, 231, 565-570, 10.1111/nph.17382, 2021.

558 Keenan, T. F., Carbone, M. S., Reichstein, M., and Richardson, A. D.: The model–data fusion pitfall: assuming  
559 certainty in an uncertain world, *Oecologia*, 167, 587-597, 10.1007/s00442-011-2106-x, 2011.

560 Kramer, K.: A Modeling Analysis of the Effects of Climatic Warming on the Probability of Spring Frost Damage To  
561 Tree Species in the Netherlands and Germany, *Plant Cell Environ*, 17, 367-377, 10.1111/j.1365-  
562 3040.1994.tb00305.x, 1994.

563 Kramer, K., Buiteveld, J., Forstreuter, M., Geburek, T., Leonardi, S., Menozzi, P., Povillon, F., Schelhaas, M., du  
564 Cros, E. T., Vendramin, G. G., and van der Werf, D. C.: Bridging the gap between ecophysiological and genetic  
565 knowledge to assess the adaptive potential of European beech, *Ecological Modelling*, 216, 333-353,  
566 10.1016/j.ecolmodel.2008.05.004, 2008.

567 Langvall, O., Nilsson, U., and Orlander, G.: Frost damage to planted Norway spruce seedlings - influence of site  
568 preparation and seedling type, *Forest Ecology and Management*, 141, 223-235, 10.1016/S0378-1127(00)00331-5,  
569 2001.

570 Liu, G. H., Chuine, I., Denechere, R., Jean, F., Dufrene, E., Vincent, G., Berveiller, D., and Delpierre, N.: Higher  
571 sample sizes and observer inter-calibration are needed for reliable scoring of leaf phenology in trees, *Journal of*  
572 *Ecology*, 109, 2461-2474, 10.1111/1365-2745.13656, 2021.

573 Liu, Q., Piao, S. L., Janssens, I. A., Fu, Y. S., Peng, S. S., Lian, X., Ciais, P., Myneni, R. B., Penuelas, J., and Wang,  
574 T.: Extension of the growing season increases vegetation exposure to frost, *Nature Communications*, 9, 426, ARTN  
575 426, 10.1038/s41467-017-02690-y, 2018.

576 Liu, Z., Fu, Y. H., Shi, X., Lock, T. R., Kallenbach, R. L., and Yuan, Z.: Soil moisture determines the effects of  
577 climate warming on spring phenology in grasslands, *Agricultural and Forest Meteorology*, 323, 109039,  
578 <https://doi.org/10.1016/j.agrformet.2022.109039>, 2022.

579 Lundell, R., Hanninen, H., Saarinen, T., Astrom, H., and Zhang, R.: Beyond rest and quiescence (endodormancy and  
580 ecodormancy): A novel model for quantifying plant-environment interaction in bud dormancy release, *Plant Cell*  
581 *Environ*, 43, 40-54, 10.1111/pce.13650, 2020.

582 Luo, M., Meng, F., Sa, C., Duan, Y., Bao, Y., Liu, T., and De Maeyer, P.: Response of vegetation phenology to soil  
583 moisture dynamics in the Mongolian Plateau, *CATENA*, 206, 105505, <https://doi.org/10.1016/j.catena.2021.105505>,  
584 2021.

585 Malyshev, A. V., Henry, H. A. L., Bolte, A., Khan, M. A. S. A., and Kreyling, J.: Temporal photoperiod sensitivity  
586 and forcing requirements for budburst in temperate tree seedlings, *Agricultural and Forest Meteorology*, 248, 82-90,  
587 10.1016/j.agrformet.2017.09.011, 2018.

588 Malyshev, A. V., van der Maaten, E., Garthen, A., Mass, D., Schwabe, M., and Kreyling, J.: Inter-Individual  
589 Budburst Variation in *Fagus sylvatica* Is Driven by Warming Rate, *Front Plant Sci*, 13, 853521,  
590 10.3389/fpls.2022.853521, 2022.

591 Meier, U.: Growth stages of mono- and dicotyledonous plants., *BBCH Monograph*, Blackwell Wissenschafts-Verlag  
592 Berlin Wien, 1997.

593 Meng, L., Zhou, Y., Gu, L., Richardson, A. D., Penuelas, J., Fu, Y., Wang, Y., Asrar, G. R., De Boeck, H. J., Mao, J.,  
594 Zhang, Y., and Wang, Z.: Photoperiod decelerates the advance of spring phenology of six deciduous tree species  
595 under climate warming, *Global Change Biology*, 27, 2914-2927, 10.1111/gcb.15575, 2021.

596 Menzel, A., Sparks, T. H., Estrella, N., Koch, E., Aasa, A., Ahas, R., Alm-Kubler, K., Bissolli, P., Braslavská, O.,  
597 Briede, A., Chmielewski, F. M., Crepinsek, Z., Curnel, Y., Dahl, A., Defila, C., Donnelly, A., Filella, Y., Jatcza, K.,  
598 Mage, F., Mestre, A., Nordli, O., Penuelas, J., Pirinen, P., Remisova, V., Scheifinger, H., Striz, M., Susnik, A., Van  
599 Vliet, A. J. H., Wielgolaski, F. E., Zach, S., and Züst, A.: European phenological response to climate change matches  
600 the warming pattern, *Global Change Biology*, 12, 1969-1976, 10.1111/j.1365-2486.2006.01193.x, 2006.

601 Morente-Lopez, J., Kass, J. M., Lara-Romero, C., Serra-Diaz, J. M., Soto-Correa, J. C., Anderson, R. P., and Iriondo,  
602 J. M.: Linking ecological niche models and common garden experiments to predict phenotypic differentiation in  
603 stressful environments: Assessing the adaptive value of marginal populations in an alpine plant, *Global Change*  
604 *Biology*, 28, 4143-4162, 10.1111/gcb.16181, 2022.

605 Oddou-Muratorio, S. and Davi, H.: Simulating local adaptation to climate of forest trees with a Physio-Demo-  
606 Genetics model, *Evol Appl*, 7, 453-467, 10.1111/eva.12143, 2014.

607 Parmesan, C. and Yohe, G.: A globally coherent fingerprint of climate change impacts across natural systems, *Nature*,  
608 421, 37-42, 10.1038/nature01286, 2003.

609 Petit, R. J. and Hampe, A.: Some evolutionary consequences of being a tree, *Annual Review of Ecology Evolution*  
610 *and Systematics*, 37, 187-214, 10.1146/annurev.ecolsys.37.091305.110215, 2006.

611 Piao, S., Liu, Q., Chen, A., Janssens, I. A., Fu, Y., Dai, J., Liu, L., Lian, X., Shen, M., and Zhu, X.: Plant phenology  
612 and global climate change: Current progresses and challenges, *Global Change Biology*, 25, 1922-1940,  
613 10.1111/gcb.14619, 2019.

614 Puchalka, R., Koprowski, M., Przybylak, J., Przybylak, R., and Dabrowski, H. P.: Did the late spring frost in 2007  
615 and 2011 affect tree-ring width and earlywood vessel size in Pedunculate oak (*Quercus robur*) in northern Poland?,  
616 *Int J Biometeorol*, 60, 1143-1150, 10.1007/s00484-015-1107-6, 2016.

617 R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing,  
618 Vienna, Austria. URL <https://www.R-project.org/>, 2020

619 Rathgeber, C. B., Rossi, S., and Bontemps, J. D.: Cambial activity related to tree size in a mature silver-fir plantation,  
620 *Ann Bot*, 108, 429-438, 10.1093/aob/mcr168, 2011.

621 Renner, S. S. and Zohner, C. M.: Climate Change and Phenological Mismatch in Trophic Interactions Among Plants,  
622 Insects, and Vertebrates, *Annu Rev Ecol Evol S*, 49, 165-182, 10.1146/annurev-ecolsys-110617-062535, 2018.

623 Richardson, A. D., Black, T. A., Ciais, P., Delbart, N., Friedl, M. A., Gobron, N., Hollinger, D. Y., Kutsch, W. L.,  
624 Longdoz, B., Luysaert, S., Migliavacca, M., Montagnani, L., Munger, J. W., Moors, E., Piao, S., Rebmann, C.,  
625 Reichstein, M., Saigusa, N., Tomelleri, E., Vargas, R., and Varlagin, A.: Influence of spring and autumn  
626 phenological transitions on forest ecosystem productivity, *Philos Trans R Soc Lond B Biol Sci*, 365, 3227-3246,  
627 10.1098/rstb.2010.0102, 2010.

628 Rousi, M. and Heinonen, J.: Temperature sum accumulation effects on within-population variation and long-term  
629 trends in date of bud burst of European white birch (*Betula pendula*), *Tree Physiol*, 27, 1019-1025,  
630 10.1093/treephys/27.7.1019, 2007.

631 Rusanen, M., Vakkari, P., and Blom, A.: Genetic structure of *Acer platanoides* and *Betula pendula* in northern  
632 Europe, *Can J Forest Res*, 33, 1110-1115, 10.1139/X03-025, 2003.

633 Scotti, I., González-Martínez, S. C., Budde, K. B., and Lalagüe, H.: Fifty years of genetic studies: what to make of  
634 the large amounts of variation found within populations?, *Annals of Forest Science*, 73, 69-75, 10.1007/s13595-015-  
635 0471-z, 2016.

636 Vallet, L.: Modélisation de la dynamique intra-populationnelle du débourrement en Ile-de-France, MSc report,  
637 Université Paris-Saclay, Orsay, France, 2020.

638 Vegis, A.: Dormancy in Higher Plants, *Annual Review of Plant Physiology*, 15, 185-+,  
639 10.1146/annurev.pp.15.060164.001153, 1964.

640 Vidal, J. P., Martin, E., Franchisteguy, L., Baillon, M., and Soubeyroux, J. M.: A 50-year high-resolution  
641 atmospheric reanalysis over France with the Safran system, *International Journal of Climatology*, 30, 1627-1644,  
642 10.1002/joc.2003, 2010.

643 Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., Jung, V., and Messier, J.: The return of  
644 the variance: intraspecific variability in community ecology, *Trends Ecol Evol*, 27, 244-252,  
645 10.1016/j.tree.2011.11.014, 2012.

646 Vitasse, Y. and Basler, D.: What role for photoperiod in the bud burst phenology of European beech, *European*  
647 *Journal of Forest Research*, 132, 1-8, 10.1007/s10342-012-0661-2, 2013.

648 Vitasse, Y., Porte, A. J., Kremer, A., Michalet, R., and Delzon, S.: Responses of canopy duration to temperature  
649 changes in four temperate tree species: relative contributions of spring and autumn leaf phenology, *Oecologia*, 161,  
650 187-198, 10.1007/s00442-009-1363-4, 2009a.

651 Vitasse, Y., Delzon, S., Dufrêne, E., Pontailier, J.-Y., Louvet, J.-M., Kremer, A., and Michalet, R.: Leaf phenology  
652 sensitivity to temperature in European trees: Do within-species populations exhibit similar responses?, *Agricultural  
653 and Forest Meteorology*, 149, 735-744, 10.1016/j.agrformet.2008.10.019, 2009b.

654 Vitasse, Y., Schneider, L., Rixen, C., Christen, D., and Rebetez, M.: Increase in the risk of exposure of forest and  
655 fruit trees to spring frosts at higher elevations in Switzerland over the last four decades, *Agricultural and Forest  
656 Meteorology*, 248, 60-69, 10.1016/j.agrformet.2017.09.005, 2018.

657 Walther, G. R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J., Fromentin, J. M., Hoegh-Guldberg, O.,  
658 and Bairlein, F.: Ecological responses to recent climate change, *Nature*, 416, 389-395, 10.1038/416389a, 2002.

659 Wenden, B., Mariadassou, M., Chmielewski, F. M., and Vitasse, Y.: Shifts in the temperature-sensitive periods for  
660 spring phenology in European beech and pedunculate oak clones across latitudes and over recent decades, *Global  
661 Change Biology*, 26, 1808-1819, 10.1111/gcb.14918, 2020.

662 Zhang, R., Lin, J. H., Wang, F. C., Delpierre, N., Kramer, K., Hanninen, H., and Wu, J. S.: Spring phenology in  
663 subtropical trees: Developing process-based models on an experimental basis, *Agricultural and Forest Meteorology*,  
664 314, ARTN 108802, 10.1016/j.agrformet.2021.108802, 2022.

665 Zhang, R., Lin, J. H., Wang, F. C., Shen, S. T., Wang, X. B., Rao, Y., Wu, J. S., and Hanninen, H.: The chilling  
666 requirement of subtropical trees is fulfilled by high temperatures: A generalized hypothesis for tree endodormancy  
667 release and a method for testing it, *Agricultural and Forest Meteorology*, 298, ARTN 108296,  
668 10.1016/j.agrformet.2020.108296, 2021.

669 Zohner, C. M., Mo, L., and Renner, S. S.: Global warming reduces leaf-out and flowering synchrony among  
670 individuals, *Elife*, 7, 10.7554/eLife.40214, 2018.

671 Zohner, C. M., Mo, L., Sebald, V., Renner, S. S., and Dornelas, M.: Leaf-out in northern ecotypes of wide-ranging  
672 trees requires less spring warming, enhancing the risk of spring frost damage at cold range limits, *Global Ecology  
673 and Biogeography*, 29, 1065-1072, 10.1111/geb.13088, 2020.

674