

# Amplified surface warming in the Southwest Pacific during the mid-Pliocene (3.3–3.0 Ma) and future implications

Georgia R. Grant<sup>1</sup>, Jonny H.T. Williams<sup>2</sup>, Sebastian Naeher<sup>1</sup>, Osamu Seki<sup>3</sup>, Erin L. McClymont<sup>4</sup>, Molly O. Patterson<sup>5</sup>, Alan M. Haywood<sup>6</sup>, Erik Behrens<sup>2</sup>, Masanobu Yamamoto<sup>3</sup>, Katelyn Johnson<sup>1</sup>

<sup>1</sup>GNS Science, Lower Hutt, New Zealand

<sup>2</sup>NIWA, Wellington, New Zealand

<sup>3</sup>Hokkaido University, Sapporo, Hokkaido, Japan

<sup>4</sup>Durham University, Durham, United Kingdom

<sup>5</sup>Binghamton University, SUNY, New York, America

<sup>6</sup>University of Leeds, Leeds, United Kingdom

*Correspondence to:* Georgia R. Grant (G.Grant@gns.cri.nz)

## Abstract

Based on Nationally-Determined Contributions concurrent with Shared Socio-economic Pathway (SSP) 2-4.5, the IPCC predicts global warming between 2.1–3.5°C (very likely range 10<sup>th</sup>-90<sup>th</sup> percentile) by 2100 AD. However, global average temperature is a poor indicator of regional warming and Global Climate Models (GCMs) require validation with instrumental or proxy data from geological archives to assess their ability to simulate regional ocean and atmospheric circulation, and thus, to evaluate their performance for regional climate projections. The Southwest Pacific is a region that performs poorly when GCMs are evaluated against instrumental observations. The New Zealand Earth System Model (NZESM) was developed from the United Kingdom Earth System Model (UKESM) to better understand Southwest Pacific response to global change, by including a nested ocean grid in the Southwest Pacific with 80% greater horizontal resolution than the global-scale host.

Here, we reconstruct regional Southwest Pacific sea surface temperature (SST) for the mid-Pliocene Warm Period (mPWP; 3.3–3.0 Ma), which has been widely considered a past analogue with an equilibrium surface temperature response of +3°C to an atmospheric CO<sub>2</sub> concentration of ~350–400ppm, to assess the warming distribution in the Southwest Pacific. This study presents proxy SSTs from seven deep sea sediment cores distributed across the Southwest Pacific. Our reconstructed SSTs are derived from molecular biomarkers preserved in the sediment - alkenones (i.e.,  $U_{37}^{K'}$  index) and isoprenoid glycerol dialkyl glycerol tetraethers (i.e., TEX<sub>86</sub> index) and are compared with SSTs reconstructed from the Last Interglacial (125 ka), Pliocene Model Intercomparison Project (PlioMIP) outputs and transient climate model projections (NZESM and UKESM) of low to high range SSPs for 2090-2099 AD.

Mean interglacial equilibrium SSTs during the mPWP for the Southwest Pacific sites, were on average, 4.2°C (1.8–6.1°C likely range) above pre-industrial and show good agreement with model outputs from NZESM and UKESM under mid-range SSP 2-4.6 conditions. These results highlight that not only is the mPWP an appropriate analogue when considering future temperature change in the centuries to come, but also demonstrate that the Southwest Pacific region will experience warming that exceeds that of the global mean if atmospheric CO<sub>2</sub> remains above 350 ppm.

## 1 Introduction


The latest IPCC climate projections to 2100 AD project average global surface warming of between 1.4-4.4°C depending on the emissions pathway (IPCC, 2022). While limiting global warming to 1.5°C urgently requires policies and actions to bring about steep emission reductions this decade, global warming could be stabilised at 2.0°C, if the latest Nationally Determined Contributions are achieved (Meinshausen, 2022). Despite stabilising at 2.0°C, heat taken up by the ocean and the polar ice sheets would ensure global sea-level would continue to rise for centuries to come (IPCC, 2022). Warming above 2.0°C may trigger rapid unstoppable collapse of the marine-based sectors of the Antarctic Ice Sheets, with one model for a high-emissions scenario suggesting global mean sea-level rise of up to 2 m by 2100 AD and 13 m by 2300 AD (DeConto *et al.*, 2021; IPCC, 2022). Notwithstanding the high-end scenarios, a stability threshold for Antarctic ice shelves is crossed above +2.0°C that commits the planet to multi-metre, multi-century sea-level rise (DeConto and Pollard, 2016; Golledge *et al.*, 2019; Lowry *et al.*, 2021). Additionally, the regional expression of global warming can differ significantly from global averages, as is evident from most land regions currently recording warming which exceeds the global average (Hoegh-Guldberg *et al.*, 2018; Sutton and Bowen, 2018; Doblas-Reyes *et al.*, 2021). Regionally focussed climate models are necessary for island nations with oceanic influence and dramatic topography such as New Zealand, since these parameters are unresolvable at the spatial resolutions used by climate models with a low, uniform resolution (Doblas-Reyes *et al.*, 2021).

Here, we consider the regional climate of the Southwest Pacific and Southern Ocean, which is often misrepresented due to coarse resolution and biases introduced in global climate models (Behrens *et al.*, 2020, 2022; Williams *et al.*, 2023). Steep regional gradients in SST, salinity and nutrients, characterise water masses spanning the Southwest Pacific and New Zealand continent (Zealandia - Te Riu-a-Māui) (Ridgway, 2007; Chiswell *et al.*, 2015; Chiswell, 2021), which represents a key location for southward heat transport balanced by northward flow of deep western boundary currents (Carter *et al.*, 2004). Subtropical waters are transported southward through surface eddies and the East Australian Current and Tasman Front (Fig 1; Behrens *et al.*, 2019). Zealandia is situated at the confluence of relatively cool, fresh, nutrient-rich Subantarctic Waters and warm, salty, nutrient-poor Subtropical Waters, defining the Subtropical Front (e.g., Chiswell *et al.*, 2015; Fig. 1). The NZESM was developed from its parent model, the UKESM, to address the need for higher spatial resolution in models across Zealandia (Williams *et al.*, 2016). An increased horizontal grid resolution from 1° to 0.2° better simulates boundary currents and surface eddies, and result in an increased meridional heat transport from the equator to higher southern latitudes (Behrens *et al.*, 2019) and is in better agreement with historical observations compared to the UKESM (Behrens *et al.*, 2020).

Past climate data allow the reconstruction of the equilibrium climate states in response to both fast and slow Earth system feedbacks involving the cryosphere, ocean and atmospheric circulation and the carbon cycle. Data from these geological archives for times representing higher-than-present CO<sub>2</sub> worlds have been widely used in climate model-intercomparison projects (CMIPs) to assess the performance of transient GCMs run to equilibrium (e.g. Haywood *et al.*, 2019; Masson-Delmotte *et al.*, 2013). While most CMIPs reconcile global mean temperatures, they poorly reconcile regional climatic patterns such as polar amplification (Naish & Zwartz, 2012; Haywood *et al.*, 2019; Masson-Delmotte *et al.*, 2013; Fischer *et al.*, 2018). This is in part due to the incomplete spatial coverage

of the geological data, accuracy and quality of the data, the resolution of GCM grids and their treatment of mid- to high-latitude polar processes. Equilibrium Climate Sensitivity (ECS) (model warming associated with a doubling of CO<sub>2</sub> once the energy balance has reached equilibrium) is one important measure of how models perform on longer timescales. An increase in ECS from CMIP Phase 5 to CMIP Phase 6 ensemble has been linked to shortwave cloud feedbacks, which has significant impact over the Southern Ocean (Zelinka *et al.*, 2020; Zhu *et al.*, 2021). Higher ECS is more consistent with estimates of warmer-than-present paleo climate sensitivity (Forster *et al.*, 2021).

We assess the magnitude and distribution of warming for the Southwest Pacific for various emissions scenarios and discuss the differences between the global climate models and paleoclimate reconstructions and consider the implications for interpreting projections of future warming in the SW Pacific.



**Figure 1: Location map for sites used in the Southwest Pacific Sea Surface Temperature (SST) reconstruction (North top of page). Sites in black have been previously published and sites in red were analysed in this study. Present day surface ocean circulation and fronts referenced in text are displayed. Note ODP 806 (0.3° N 159.4° E) is not displayed in this projection. Bathymetry is plotted using ‘ggOceanMaps’ (Vihtakari, 2022) and bathymetry data are sourced from Amante and Eakins (2009).**

### 1.1 Paleoclimate analogues

*Mid-Pliocene Warm Period (3.3 – 3.0 Ma)*

Global temperatures (+2–3°C) last experienced during the mPWP (3.3–3.0 Ma) may be reached by 2100 AD if  
105 emissions are abated in line with the SSP2-4.5 scenario, which is the pathway aligned to current policy (not the  
aspirational 1.5°C Paris-target) (Burke *et al.*, 2018). The mPWP spans a 300 kyr period when atmospheric CO<sub>2</sub>  
was comparable to present day (mean 390 ppm; Chalk *et al.*, 2017; De La Vega *et al.*, 2020). During this period  
interglacial global temperatures were 2–3°C warmer (Dowsett *et al.*, 2013; Masson-Delmotte *et al.*, 2013), and  
the amplitude of glacial-interglacial sea-level change was likely between 6 and 17 m (16<sup>th</sup>-84<sup>th</sup> percentile) (Grant  
110 *et al.*, 2019; Grant and Naish, 2021). Such a rise in global sea-level implies melting of the Greenland Ice Sheet  
(Koenig *et al.*, 2015; Batchelor *et al.*, 2019), West Antarctic Ice Sheet (Naish *et al.*, 2009; McKay *et al.*, 2012)  
and parts of marine-based East Antarctic Ice Sheet (Cook *et al.*, 2013; Patterson *et al.*, 2014; Bertram *et al.*, 2018).  
Therefore, the interglacial periods of mPWP are considered to be the most accessible and suitable past analogue,  
or window, into the future equilibrium response of the Earth system to warming in line with SSP2-4.5 (Naish &  
115 Zwart, 2012; Dowsett *et al.*, 2013; Haywood *et al.*, 2019).

The mPWP has been the focus of several major international research initiatives. The Pliocene Research,  
Interpretation and Synoptic Mapping (PRISM) project (Dowsett *et al.*, 2013; 2016) undertook a global  
compilation of paleoclimate data, primarily surface temperature reconstructions. The Pliocene Modelling  
120 Intercomparison Project (PlioMIP) presents a multi-model ensemble with various ECS run for mPWP conditions  
(Haywood *et al.*, 2011; 2016; 2020). The recent IPCC summary of ECS ranges from median values of 2.5–3.7°C  
(Forster *et al.*, 2021). This ECS summary does not include model-based estimates, but does include emergent  
constraints (Hargreaves and Annan, 2016; Renoult *et al.*, 2020) utilising PlioMIP (Haywood *et al.*, 2020) and  
proxy temperature and CO<sub>2</sub> reconstructions (Martinez-Boti *et al.*, 2015; Sherwood *et al.*, 2020).  
125 Marine Isotope Stage (MIS) KM5c (3.2 Ma) interglacial became a focus for reconstructing warming within mPWP  
as insolation values and the orbital configuration were most similar to the Holocene interglacial (Haywood *et al.*,  
2020; McClymont *et al.*, 2020). While, based on less data points, this approach has better agreement between  
models and observations and revealed a higher ECS of 2.6–4.8°C for conditions of MIS KM5c from the PlioMIP  
Phase 2 ensemble (PlioMIP2; Haywood *et al.*, 2020). A recent review of SSTs in the mPWP for MIS KM5c by  
130 the PlioVAR working group (Pliocene climate variability on glacial-interglacial timescales; McClymont *et al.*,  
2020) used alkenones to reconstruct an average global SST warming of 3.2–3.4°C above pre-industrial SST. This  
is slightly warmer than PlioMIP2 simulations, where global surface air temperature over oceans were ~2.8°C  
above pre-industrial. However, differences are suggested to be due to regional ocean circulation and proxy signals  
(McClymont *et al.*, 2020).

135 While interglacial minima and glacial maxima in the benthic δ<sup>18</sup>O stack (MISs) have been the primary means of  
reconstructing the timing and magnitude of global sea-level variations over the last 5 Ma (Lisiecki and Raymo,  
2005), for some time intervals (i.e., mPWP) global sea-level is known to fluctuate at a higher frequency than can  
be assessed in the benthic δ<sup>18</sup>O stack (Grant *et al.*, 2019). This is also the case for other proxies with variable  
140 sampling resolution such as SST that have not been tuned to the δ<sup>18</sup>O stack (e.g. Herbert *et al.*, 2010; Fig. 2). The  
reliance on orbitally tuned timescales in deep ocean paleoclimate records has potentially led to the  
misinterpretations of the timing, frequency and amplitude of glacial-interglacial climate change. This is  
particularly the case in the Pliocene and Early Pleistocene where there are less globally distributed δ<sup>18</sup>O records

and many are of coarse sampling resolution (Lisiecki and Raymo, 2005). In a number of studies (Lisiecki and Raymo, 2005; Miller *et al.*, 2012; Grant *et al.*, 2019), average glacial climate conditions (global surface temperature and sea-level) during the mPWP have been considered similar to those of the Holocene.

#### *Last interglacial (125 ka)*

Finally, we briefly compare these results to the Last Interglacial MIS 5e (~125 ka) as many of the sites investigated here were also used by Cortese *et al.* (2013) in a proxy SST study. Peak interglacial SSTs were reconstructed from core-top planktonic foraminiferal assemblages, calibrated to modern SSTs and then applied to paleo assemblages (Cortese *et al.*, 2013). The Southwest Pacific study presented warming focused in Tasmania and western New Zealand and proposed a strengthened East Australian Current bathing Tasmania with warmer water (Cortese *et al.*, 2013). MIS 5e represents a lower global average temperature increase of 1–2°C above pre-industrial in response to changing orbital configurations on radiative forcing (rather than CO<sub>2</sub>), associated with 6–9 m of sea-level rise, which together with the mPWP analogue discussed above implies extreme sensitivity of the polar ice sheets to relatively small changes in global mean surface temperature (Dutton *et al.*, 2013).




Figure 2: mid-Pliocene Warm Period (mPWP) climate context showing reconstructions of a) a combined signal of global sea-level and ocean temperature from deep sea benthic  $\delta^{18}\text{O}$  data (Lisiecki and Raymo, 2005) spanning mPWP to present, b) daily insolation at  $65^\circ\text{N}$  (21<sup>st</sup> June: Laskar *et al.*, 2004), c) global relative sea-level change from the PlioSeaNZ record, Whanganui Basin, New Zealand (Grant *et al.*, 2019), d) a combined signal of global sea-level and ocean temperature from deep sea benthic  $\delta^{18}\text{O}$  data mPWP (3.3–3.0 Ma) of (Lisiecki and Raymo, 2005), e) atmospheric CO<sub>2</sub> from  $\delta^{11}\text{B}$ -pH proxy (De La Vega *et al.*, 2020) and f) tropical Sea Surface Temperatures (SSTs) from alkenone paleothermometry (Herbert *et al.*, 2010). Pre-industrial (PI) estimates are also shown.

## 1.2 Future scenarios

Here we display model results of future projections from NZESM and UKESM. These are previously published and thus introduced here, while the comparisons to data presented in this study are discussed in detail in Section 3.


The NZESM (Williams *et al.*, 2016, Behrens *et al.*, 2020) is based on the UKESM (Sellar *et al.*, 2019; Senior *et al.*, 2020), a CMIP6 earth system model (ESM) containing a dynamic atmosphere, ocean, prognostic sea ice, complex atmospheric chemistry and ocean biogeochemistry. Via a two-way nesting scheme, ocean physical parameters were dynamically downscaled from 1° to 0.2° in the NZESM to better simulate boundary currents and mesoscale variability, instrumental for southward heat transport (Behrens *et al.*, 2019). This nesting improves the steady state simulated sea surface properties (Behrens *et al.*, 2020; 2022). With the exception of a solar-cycle-dependence of the ozone photolysis scheme included in the NZESM (Dennison *et al.*, 2019), the atmospheric physics is identical to the UKESM in all other respects. Globally averaged SSTs are marginally warmer than the UKESM in all pathways up to 2100 AD, but that difference is reduced as the magnitude of warming increases under higher-emission scenarios (Fig. 3). Indeed, for 2090–2099 AD in SSP3-7.0, the mean difference between the two models is essentially zero for higher greenhouse gas levels. This global signal is, dominated by the southern hemisphere warming induced by increased southward heat transport from the tropics in the NZESM.

The latest climate projections are grouped according to primary Shared Socioeconomic Pathways (SSPs; Lee *et al.*, 2021) forced by various greenhouse gas emissions and other radiative forcings and simulated by the CMIP6 (Eyring *et al.*, 2016). These pathways are differentiated by degrees of very likely warming by 2100 AD, i.e. 1.3–2.4°C (SSP1 - sustainability), 2.1–3.5°C (SSP2 - middle of the road), 2.5–4.6°C (SSP3 - regional rivalry) (Chen *et al.*, 2021; O'Neill *et al.*, 2016). NZESM and UKESM were both run for SSP1-2.6, SSP2-4.5 and SSP3-7.0 and broadly correspond to low, medium and high emissions scenarios and were run out to 2100 AD. While UKESM was run for other SSPs, NZESM was not, so we have restricted comparison to these scenarios.

The UKESM (a CMIP ensemble member) and NZESM have an ECS of 5.4°C (Sellar *et al.*, 2019) which is higher than the likely range (high confidence) for ECS as 2.5–4°C (Zelinka *et al.*, 2020). This is most clearly seen in the degree of global warming (Fig. 3a) compared to the regional warming of the high-resolution NZESM ocean-grid area (Fig. 3b). Climate scenarios by the 2090-2099 AD period generated warming of i) ~3°C globally and ~2°C regionally for SSP1-2.6, ii) 4°C globally and 3°C regionally for SSP2-4.5, and iii) 6°C globally and 4.5°C regionally for SSP3-7.0 (Fig. 3). This differs (close to half) from mean global warming from the CMIP6 model ensemble of ~1.8°C, 2.7°C and 3.6°C for SSP1-2.6, 2-4.5 and 3-7.0 respectively by 2100 AD (IPCC, 2022). Annual mean SSTs were extracted for all sites and are reported here. Sites in the tropics (ODP 806) and Southern Ocean (ANDRILL) were excluded as they are outside of the NZESM high-resolution ocean-grid region.

As a reference or pre-industrial control, the results generated from the Hadley Centre Global Sea Ice and Sea Surface Temperature (HadISST) model were used from 1870–1879 AD (NCAR, 2022; Rayner *et al.*, 2003). Best practise of model assessment is to present anomalies with reference to pre-industrial runs from the same model. As a pre-industrial run is unavailable for the NZESM, we have used the single reference of HadISST for all model

and proxy anomaly assessment. HadISST was selected as it is the most complete reanalysis product nearest to pre-industrial conditions.



210 **Figure 3: Mean Surface Air Temperature from NZESM and UKESM simulations of low- to high-range emission Shared Socio-economic Pathways (SSPs) for a) the global region and b) the area covered by the high-resolution ocean grid of NZESM. Results generated from the UKESM (Sellar *et al.*, 2019) and NZESM (Williams *et al.*, 2016, Behrens *et al.*, 2022) projections used in this study are extracted for all SSPs for 2090–2099 AD.**

## 215 **2 Methods**

To enable comparison of past SSTs with future projections, we assess the full duration and glacial to interglacial amplitude of the mPWP, for sites across the Southwest Pacific region. In this approach, glacial and interglacial modal means are determined statistically to ensure the pattern and magnitude of warming is more representative of mPWP interglacial climate conditions as opposed to the single peak interglacial conditions that have been the focus of previous climate reconstructions (Dowsett *et al.*, 2016, Haywood *et al.*, 2020, McClymont *et al.*, 2020).

We have applied the  $U_{37}^{K'}$  index (unsaturated ketone index; Prahl and Wakeham, 1987) to reconstruct SSTs at four new sites (DSDP 590, ODP 1123, ODP 1168 and ODP 1172) which complement three sites with previously published  $U_{37}^{K'}$ -derived SSTs (DSDP 593; McClymont *et al.*, 2016; DSDP 594 and ODP 1125; Caballero-Gill *et al.*, 2019). We have also applied TEX<sub>86</sub> (TetraEther index of tetraether consisting of 86 carbon atoms) most commonly correlated with SST or shallow subsurface (50–200 m) temperatures (Tierney and Tingley, 2015) at two of the sites (DSDP 590 and ODP 1172). Two additional sites, ODP 806 (Eastern Equatorial Pacific: Medina-Elizalde and Lea, 2010) and ANDRILL (Ross Sea, Antarctica; McKay *et al.*, 2012) are located outside of the Southwest Pacific and provide a meridional climate context.

We extract site-specific simulated SSTs from PlioMIP and future UKESM and NZESM to compare the reconstructed pattern of warming in the Southwest Pacific during the mPWP.

## 2.1 mid-Pliocene Warm Period records

Sea surface temperature records from nine sites are presented in this study, including published SST data from five sites and new SST data from four sites to improve the geographical resolution across the Southwest Pacific and surrounding water masses. Inclusion of tropical site ODP 806 and Antarctic ANDRILL site in the Ross Sea allows us to present a latitudinal transect from 0.3° N to 77° S, within longitudes 155° E to 165° W (Fig. 1; Table 1). Sites were selected from cores that were available through International Ocean Drilling Program (IODP) and predecessor drilling programmes. Sampling of new sites was evenly distributed across the mPWP (Table S1 and S2), with age models selected from the most up to date publications (Table 1). The age models used in previously published SST records are retained here (Table 1). Published age models by Karas *et al.* (2011), Patterson *et al.* (2016) and McClymont *et al.* (2016) are calibrated to the deep sea  $\delta^{18}\text{O}$  benthic stack (Lisiecki and Raymo, 2005). In the case of sites ODP 1172 and ODP 1168, we use the integrated shipboard age models for the mPWP (Exon *et al.*, 2001). Linear interpolation of magnetostratigraphy provided by Exon *et al.* (2001) was used in absence of high-resolution  $\delta^{18}\text{O}$  records for site ODP 1168 and 1172 that could be correlated to the deep sea  $\delta^{18}\text{O}$  benthic stack.

Sediment samples obtained from four sites (ODP 1168, ODP 1172, ODP 1123, DSDP 590) were analysed for alkenone-based SST reconstructions using the  $U_{37}^{K'}$  index (e.g., Prahl and Wakeman, 1987; Section 2.2) at a target temporal resolution of less than 10 kyr (Table 1). For two sites (DSDP 590 and ODP 1172), additional analysis of glycerol dialkyl glycerol tetraethers (GDGTs) were undertaken to derive for  $\text{TEX}_{86}$ -based SST estimates (e.g., Schouten *et al.*, 2002).  $U_{37}^{K'}$  derived SSTs were reported previously for DSDP 594 and ODP 1125 (Caballero-Gill *et al.*, 2019; McClymont *et al.*, 2020), and DSDP593 McClymont *et al.*, 2016). Temperature reconstructions for the ANDRILL core were based on the  $\text{TEX}_{86}$  index (McKay *et al.*, 2012) and ODP 806 was analysed for Mg/Ca of planktic foraminifera *Globigerinoides sacculifer* (Medina-Elizalde and Lea, 2010) renamed *Trilobatus sacculifer* (Spezzaferri *et al.*, 2015). ANDRILL sediment samples represent interglacial periods as organic material at this location is not preserved during glacial intervals. These sediments are poorly constrained to specific interglacial periods and are not assigned specific ages (McKay *et al.*, 2012). Reported SST results exclude sites ANDRILL and ODP 806 as HadISST, NZESM and UKESM cannot be produced for the ANDRILL site (presently covered by the Ross Ice Shelf) and the NZESM high resolution model does not cover the region in which ANDRILL and ODP806 are located, thus we cannot provide comparisons. Furthermore, they are not alkenone-derived SST estimates.

265

270



**Table 1. Mid-Pliocene Warm Period (mPWP) site identification and location with associated surface water mass, sampling period and resolution (italicised in parenthesis), and source references for previously published data or age models used in association with new analyses.**

Site	Latitude	Longitude	Surface Water Mass	Period ( <i>sampling</i> )	Reference
ANDRILL1B	-77.889	167.089	Antarctic Shelf Water	Interglacials during mPWP (3000-3300 ka)	McKay <i>et al.</i> , 2012
DSDP594	-45.524	174.948	Subtropical Frontal Zone	3000-3299 ka (3 kyr)	Caballero-Gill <i>et al.</i> , 2019; McClymont <i>et al.</i> , 2020
ODP1172	-43.960	149.928	Subtropical Frontal Zone	3000-3301 ka (8 kyr)	Age Model: Exon <i>et al.</i> , 2001; Data – this study
ODP1168	-42.610	144.413	Subtropical Frontal Zone	3008-3290 ka (7 kyr)	Age model: Exon <i>et al.</i> , 2001; Data – this study
ODP1125	-42.550	-178.166	Rekohu Eddy (extension of Tasman Front)	3000-3299 ka (2 kyr)	Caballero-Gill <i>et al.</i> , 2019; McClymont <i>et al.</i> , 2020
ODP1123	-41.786	-171.499	Subtropical Water	3004-3300 ka (10 kyr)	Age Model: Patterson <i>et al.</i> , 2018; Data – this study
DSDP593	-40.508	167.675	Subtropical Water (Tasman Sea)	3025-3295 ka (10 kyr)	McClymont <i>et al.</i> , 2016; McClymont <i>et al.</i> , 2020
DSDP590	-31.167	163.3595	Subtropical Water	3017-3300 ka (15 kyr)	Age model: Karas <i>et al.</i> , 2011; Data – this study
ODP806	0.3185	159.361	Western Pacific Warm Pool	3000-3086 ka (2 kyr)	Medina-Elizalde and Lea, 2010

## 2.2 Biomarker ( $U_{37}^{K'}$ and TEX<sub>86</sub>) sea surface temperature reconstructions

Organic biomarkers preserved in marine sediments are important proxies for past water temperatures (e.g., de Bar *et al.*, 2019; Herbert *et al.*, 2010; Hollis *et al.*, 2019). The  $U_{37}^{K'}$  index has been applied successfully to reconstruct SSTs in marine settings worldwide from low to high latitudes (e.g., Herbert, 2014). Although this proxy is calibrated to annual average SST using linear regressions based on sediment core top data between 60°N and 60°S (Müller *et al.*, 1998; Conte *et al.*, 2006; Rosell-Melé and Prah, 2013), reconstructed SSTs can be biased towards higher temperatures due to peak alkenone production during the bloom period, which is commonly spring or early summer (Conte *et al.*, 2006; Prah *et al.*, 2010). However, other studies used a combination of measurements and modelling to show that the maximum seasonality variations can be up to ~2.5°C at high latitudes (Conte *et al.*, 2006; Prah *et al.*, 2010; Max *et al.*, 2020, McClymont *et al.*, 2020). To address the decreased response of  $U_{37}^{K'}$  at high temperatures (>24°C), Tierney and Tingley (2018) developed a Bayesian B-spline regression model (BAYSPLINE). Previous studies, including some utilised here (e.g., McClymont *et al.*, 2020), applied the linear core top calibration of Müller *et al.* (1998). However, because site DSDP 590 produces SST more than 24°C and there is little difference between the calibrations at mid-latitudes (maximum of 0.7°C), we have used the BAYSPLINE calibration and applied this to all sites (Appendix A). This results in slightly cooler temperatures (maximum <0.7°C; Table S1) but the difference remains within the calibration uncertainties (1.4°C below 24°C; Tierney and Tingley, 2018).

Additionally, for comparison with alkenone-based SST reconstructions, two sites (DSDP 590 and ODP 1172) were analysed for isoprenoid glycerol dialkyl glycerol tetraethers (isoGDGTs), which are produced by marine Thaumarchaeota (Schouten *et al.*, 2002; 2013) and used, to reconstruct TEX<sub>86</sub>-derived SSTs. Because only a limited number of samples for two sites were analysed for TEX<sub>86</sub> (n=27) within the mPWP, the results are not used in analysis to determine reported means, but are discussed in Appendix A.

### 2.3 Data analysis

Probability distributions of the mPWP proxy SSTs, grouped by site, are displayed using ‘vioplot’ which graphically normalises the distribution for ease of comparison (Fig. 4). The plots often show a bimodal distribution curve which we infer to represent two normal distributions centred around mean interglacial and mean glacial SSTs. Single mode distributions may reflect lower variability between glacial–interglacial conditions (e.g., low-latitude tropical sites), lower sample resolution that does not capture glacial–interglacial cyclicity, or sampling that favours either glacial or interglacial conditions (as is the case for ANDRILL, which is biased to interglacial ice retreat facies). Interglacials are typically identified through benthic  $\delta^{18}\text{O}$  record cyclicity and tuning these records to the global benthic  $\delta^{18}\text{O}$  stack. However, glacial-interglacial cyclicity can be quite variable between different members in the stack during the mPWP (Lisiecki and Raymo, 2004), and this also occurs between records from the Southwest Pacific sites (e.g., McClymont *et al.*, 2020). Furthermore, a number of these sites do not have  $\delta^{18}\text{O}$  records and the SST records are not consistently cyclical or high-enough resolution to determine glacial and interglacials values. For that reason, we have employed a statistical package which identifies two modes that are considered to represent average glacial and interglacial means, and thus, places more emphasis on values that record interglacial and deglacial transitions with less emphasis on glacial or interglacial extremes.

The temperature distributions for each site (excluding ANDRILL as interglacial values only) were assessed for bi-modal distribution to identify mean glacial and interglacial modes using the ‘noramlmixEM2comp’ function in package ‘mixtools’ (Benaglia *et al.*, 2010). This employs an expectation-maximization (EM) algorithm to fit an equal two-component mixture model, assuming normal distributions. This is an automated process (samples are not identified as glacial or interglacial) assuming equal two-part mixture and normal distributions of these mixtures. While the accuracy of these results is dependent on the assumptions applied here - that the glacial and interglacials present a normal distribution and have an equal bi-modal component - it is a systematic approach that applies statistical analysis to objectively identify the variance within the data and attribute that to glacial and interglacial conditions recorded in the data (Fig. B2). We acknowledge that this is an imperfect approach. However, we consider that this reduces bias introduced when visually selecting interglacial or glacial samples reliant on discrete values or temporal constraints (the latter are age model dependent). Secondly, this reduces emphasis on extreme warming during some interglacials of the mPWP and varying responses of the sites so we can be more confident that the SSTs are reflective of the broader climate conditions of the mPWP. These interglacial modes are used for plotted and tabulated comparisons to the UKESM and NZESM projections presented in the results below. Uncertainty ( $1\sigma$ ) associated with the  $U_{37}^{K'}$ -BAYSPLINE calibration is  $\pm 1.4^\circ\text{C}$  below  $\sim 23.4^\circ$  and non-linear above (Tierney and Tingley, 2018). Therefore, the higher SSTs of DSDP 590 have

335 a higher uncertainty (average of  $\pm 2.4^{\circ}\text{C}$ )(Table 2). The uncertainties for all proxy SSTs are taken as the mean of  
all sites ( $\pm 1.5^{\circ}\text{C}$ ) for absolute SST and when referenced to pre-industrial HadISST (Table 2).

### 3 Results


#### 3.1 mid-Pliocene Warm Period Sea Surface Temperature signature

340 With respect to pre-industrial (HadISST), mean site SSTs for the mid-latitudes (45 to 30°S) range from 0.8–6.6°C  
with a likely (16<sup>th</sup> – 84<sup>th</sup> percentile range) of 2–4.7°C (3.4°C average for all sites). Minimum SST ranges for the  
sites are -3.5–1.7°C (-0.3°C average for all sites) and maximum SSTs range from 3.5 to 7.5°C (5.8°C average of  
all sites) (Table 2). Interglacial modal means, used in this study as moderate warm conditions, range between 1.3–  
5.4°C (average 4.2°C) warmer than HadISST for the Southwest Pacific mid-latitude sites.

345 The sites presented in this study are sampled over glacial–interglacial cycles for which the total glacial–interglacial  
amplitude of SSTs range from ~4.4–7.5°C (Fig. 4; Table 2) (excluding ANDRILL and ODP 806). Interglacial and  
glacial modal means determined by the bimodal statistical analysis (Section 2.3) are generally comparable to the  
16<sup>th</sup> and 84<sup>th</sup> percentile (within ~1°C), highlighting that these modes are reflective of the likely range rather than  
accounting for extreme values representing the tails of the p-distribution used to estimate the total glacial–  
350 interglacial amplitude range, which has a mean of 6.1°C (Table 2). The difference between glacial and interglacial  
modal means is approximately half that of the total glacial-interglacial amplitude (~3°C; Table 2). The meridional  
gradients for mean glacials or interglacials do not differ significantly but do show a flattened gradient for  
interglacial modal means between site DSDP 590 and ODP 806 (30–0° S; Fig. 4b) due to the low SST distribution  
of site ODP 806.

355 The sites warm (~0 – 20°C) from the pole to site ODP 1125 (north Chatham Rise), before a reduction in SSTs are  
seen at sites ODP 1123 (offshore Chatham Rise) and DSDP 593 (eastern Tasman Sea), then returning to high  
temperatures >25°C at sites north of 32°S (DSDP 590 and ODP 806) which show comparable peak temperatures  
(Fig. 4). While latitude is generally correlated with SST, surface water mass and regional currents alter this  
360 relationship. Site DSDP 594 south of the Subtropical Front in surface Subantarctic Water is noticeably colder than  
sites situated either within (ODP 1172), or just north of (ODP 1168, ODP 1125, ODP 1123), the Subtropical Front.  
However, current proximity to the Subtropical Front doesn't appear to be a main driver either.

DSDP 593 and DSDP 594 (north and south of the Subtropical Front) show the least warming above pre-industrial,  
365 but interglacial modal means still warm 1–2 °C. Sites that show significant interglacial modal mean warming  
above the global mPWP average are offshore Tasmania (ODP 1168 and ODP 1172) and site ODP 1125 (northern  
Chatham Rise) which all display warming between 4.8–5.4°C, and DSDP 590 (north Tasman Sea) presents  
extreme warming of 6.7°C (Table 2).



370

375

**Figure 4: Time series and SST distribution for mid-Pliocene Warm Period (3.3 – 3.0 Ma) records. a)  $U_{37}^{K'}$ -SST calibrations of BAYSPLINE (solid; Tierney and Tingley, 2018) with the difference to Müller *et al.* (1998) shaded are plotted for all timeseries, with ODP 806 (Mg/Ca SST derived) and excluding ANDRILL (not plotted due to poor age control). b) Probability distributions ('violin plots') of the SST timeseries. Interglacial (red) and glacial (blue) modal means are also shown (Table 2; Fig. B1). Data for this plot are provided in Table S1 and Table S2.**

380

**Table 2. Statistical distribution of mid-Pliocene Warm Period Sea Surface Temperature (SST) anomalies relative to HadISST (1870-1879 AD) using  $U_{37}^{K'}$  BAYSPLINE derived SSTs. Reported minimums, 16<sup>th</sup> percentile, mean (50<sup>th</sup> percentile), 84<sup>th</sup> percentile and maximum SST anomalies are shown with  $U_{37}^{K'}$  BAYSPLINE reported 1 $\sigma$  uncertainty. Glacial and interglacial modal means are calculated as described in methods and the total range is calculated as the difference between maximum and minimum temperature.**

Site	Min. (°C)	16 <sup>th</sup> (°C)	Mean (°C)	84 <sup>th</sup> (°C)	Max. (°C)	$U_{37}^{K'} \pm 1\sigma$ (°C)	Glacial Modal-mean (°C)	Interglacial Modal-mean (°C)	Total range (°C)
DSDP594	-3.5	-0.9	0.8	2.2	3.5	1.4	-1.5	1.3	7.0
ODP1172	-1.1	2.4	3.7	5.0	6.4	1.4	3.5	4.8	7.5
ODP1168	0.8	2.6	4.0	5.3	6.9	1.4	3.2	4.9	6.1
ODP1125	1.4	2.9	4.8	6.3	7.2	1.4	2.8	5.4	5.8
ODP1123	0.7	2.0	2.9	4.5	5.1	1.4	2.3	4.2	4.4
DSDP593	-2.2	-0.8	1.0	2.4	3.8	1.4	-0.6	1.9	6.0
DSDP590	1.7	5.8	6.6	7.2	7.5	2.4	3.2	6.7	5.8
Mean	-0.3	2.0	3.4	4.7	5.8	1.5	1.8	4.2	6.1
Variance	5.2	6.7	5.8	5	4	1	5	5.4	3.1

385

## 3.2 Global Climate Models


### 3.2.1 PlioMIP

Standardised boundary conditions used by all 16 models participating in PlioMIP, is termed the PlioCore experiment (Haywood *et al.*, 2016; 2020), based on the latest PRISM4 climate reconstruction for MIS KM5c (Dowsett *et al.*, 2016). Data presented here is the multi-model mean of PlioCore (Haywood *et al.*, 2020) with site specific SSTs extracted and referenced to the HadISST pre-industrial reanalysis (NCAR, 2022) for comparison to the mPWP proxy SSTs (Table 3; Fig. 5). Due to the poor spatial distribution of this study (although considerably higher than previous studies in the region), we are unable to sum the temperature distribution over latitudinal ranges of 30° for comparison to meridional gradients reported elsewhere (e.g., Haywood *et al.*, 2020; McClymont *et al.*, 2020). However, we provide a comparison of mPWP site data to PlioMIP latitudinal averages (1° resolution) between longitudes 140°E – 160°W, alongside site-specific SST from the PlioMIP experiment (Haywood *et al.*, 2020).

Specific site warming for PlioMIP does not vary significantly from the meridional gradient except for ODP 1123 (Fig. 5b). Sites ODP 1123, DSDP 593 and DSDP 594 all present SST anomalies within 1°C for PlioMIP and mPWP (Table 3). However, on average for the sites, PlioMIP SST anomaly is 2.4°C, while mPWP is 4.2°C (Table 3).

**Table 3. Summary site mean Sea Surface Temperature (°C) for HadISST (1870-1879 AD), PlioMIP (multi-model mean) and interglacial modal-mean mPWP  $U_{37}^{K'}$  BAYSPLINE derived (this study), and SST Anomaly (relative to HadISST) for PlioMIP and mPWP.**

	HadISST	PlioMIP		mPWP	
	SST (°C)	SST (°C)	SST anomaly (°C)	SST (°C)	SST anomaly (°C)
DSDP594	10.8	12.7	1.9	12.1	1.3
ODP1172	12.4	14.9	2.5	17.2	4.8
ODP1168	12.6	16	3.4	17.5	4.9
ODP1125	13.6	16.2	2.6	19	5.4
ODP1123	14.3	17.6	3.3	18.5	4.2
DSDP593	15	16.1	1.1	16.9	1.9
DSDP590	20.7	22.6	1.9	27.4	6.7
Mean	14.2	16.6	2.4	18.4	4.2
Variance	9.9	9.9	2.3	15.3	5.4



410 **Figure 5: Ensemble mean regional Sea Surface Temperature (SST) from core PlioMIP experiments with mid-Pliocene**  
**Warm Period (mPWP) site mean interglacial SST plotted using the same temperature scale. b) absolute SST between**  
**31–46°S for PlioMIP latitudinal mean (blue dotted), PlioMIP site specific (blue solid) with transparent blue shading**  
**for minima-maxima and heavier blue shading for  $\pm 1$  standard deviation of the PlioMIP ensemble and mPWP site**  
**specific (red dashed) with mean-maxima shaded in red.**

415

### 3.2.2 Future Earth System Model simulations

On average, NZESM simulations show higher warming than the coarser resolution UKESM in all scenarios presented here (Table 4; Fig. 6). However, for the sites investigated, UKESM simulations shows more variability between sites due to minimal warming at ODP 594 and extreme warming at site ODP 1172 (Table 4; Fig. 6). As for proxy data in the previous section, statistical summaries refer to the mid-latitude sites (excluding ANDRILL and ODP 806).

420

Projections for 2090–2099 AD for SSP1-2.6, SSP2-4.5 and SSP3-7.0 show a stable pattern of warming for both models (Table 4; Fig. 6). However, warming at sites ODP 1172, ODP 1168, ODP 1123 and ODP 1125 in UKESM simulations increases above NZESM with higher emission scenarios, while DSDP 594 and 593 remain significantly higher in NZESM simulations over UKESM (Table 4). NZESM and UKESM simulations for SSP3-7.0 have similar mean warming (+4.5°C and +4.4°C respectively) to the mPWP (+4.2°C), with the means strongly biased by differences in DSDP 594, 593 and 590 (Table 4). The intense warming recorded at site DSDP 590 during the mPWP is particularly visible in latitudinal gradient comparisons (Fig. 6c,f,i) and highlights the importance of comparing site specific data.

425


430

435

**Table 4. Site annual mean Sea Surface Temperature anomalies (°C) for UKESM and NZESM with respect to HadISST (1870-1879 AD) for SSP1-2.6, SSP2-4.5, SSP3-7.0 at 2095 AD (2090–2099 AD). Global SST for the mPWP refers to the compilation by McClymont *et al.*, 2020.**

2090-2099 AD							
Site	UKESM			NZESM			mPWP SST
	SSP 1 (°C)	SSP 2 (°C)	SSP 3 (°C)	SSP 1 (°C)	SSP 2 (°C)	SSP 3 (°C)	
DSDP594	0.1	1	2.3	2.3	3.3	4.9	1.3
ODP1172	2.2	5.7	7.8	3.2	4.8	6.7	4.8
ODP1168	0.6	3.1	5.4	1.4	2.7	3.9	4.9
ODP1125	2.2	3.4	4.4	2.2	3.3	4.6	5.4
ODP1123	2	3.4	4.3	1.1	2.2	3.5	4.2
DSDP593	0.5	1.9	3.3	1.4	3	4	1.9
DSDP590	1.3	2.4	3.4	1.8	2.8	3.9	6.7
Mean	1.3	3	4.4	1.9	3.2	4.5	4.2
Variance	2.1	4.7	5.5	2.1	2.6	3.2	3.2
Global	3	4	6	3	4	6	3.2

440



**Figure 6: Regional Sea Surface Temperature (SST) anomalies to HadISST (1870 – 1879 AD) for SSP1-2.6 (a–c), SSP2-4.5 (d–f), SSP3-7.0 (g–i) in 2090–2099 AD compared to mid-Pliocene Warm Period (mPWP) site mean interglacial SST anomalies (filled circles using same colour scale as map). Left panels are NZESM, middle panels are UKESM, and right panels are site SST anomalies between 31–46°S for mPWP (red dotted), UKESM (purple dashed) and NZESM (black solid).**

445

## 4 Discussion

### 4.1 Pliocene analogue


The mPWP encompasses several glacial cycles throughout the 300 kyr, with variable climate conditions (insolation, CO<sub>2</sub> and tropical SST; Fig. 2). Many studies have therefore focused on a single interglacial (MIS KM5c) as insolation values are near identical to today (Laskar *et al.*, 2004; Haywood *et al.*, 2020; McClymont *et al.*, 2020). However, this requires confidence in age models and ultimately tuning of records. The approach taken here, of assessing glacial and interglacial characteristics spanning the whole mPWP interval, aims to smooth glacial and interglacial extremes and represent the more “likely” climate conditions for equilibrium glacial and interglacial states across the region. Interglacial SST site modal means (this study) between 30° S and 45° S average at +4.2°C (Table 2) for global SST estimates of 3.2–3.4°C (McClymont *et al.*, 2020). In comparison, for the same sites, PlioMIP SSTs average 2.4±2.1°C (MIS KM5c ; Haywood *et al.*, 2020; Table 3), with global multi-mode median of 3.0°C (10<sup>th</sup>-90<sup>th</sup> percentiles: 2.1-4.8 °C) (Haywood *et al.*, 2020). Thus, this study demonstrates an amplified warming signal in the Southwest Pacific relative to the global mean temperature that is not displayed in the PlioMIP simulations. Likewise, the mean of glacial modes is +2°C with reference to HadISST (Table 2), where glacial conditions of the mPWP are often considered comparable to pre-industrial (Lisiecki and Raymo, 2005), which is however, poorly studied. The comparable SSTs for sites with previously published values for MIS KM5c provides confidence in the approach of representing interglacial modal means used in this study and highlights the importance of regional variability in site selection to determine regional response (Fig. 7a). While we acknowledge the sites provided in this study are still spatially limited, they provide a significant increase to the resolution of sampling in this area for the mPWP.

Site DSDP 590 (northern Tasman Sea) presents the highest SSTs, which is currently north of the Tasman Front outlet of EAC. The location of the Tasman Front is controlled by the northern tip of New Zealand’s North Island, which was at a slightly lower latitude in the Pliocene (Strogen *et al.*, 2022) which may have allowed for a more northern Tasman Front, directing warmer waters across site DSDP 590. Alternatively, the warming at DSDP 590 may be explained by a broadening and invigoration of the Tasman Front, which may be at the expense of flow to the EAC-extension (Hill *et al.*, 2011). While a strengthening of the EAC is expected, the magnitude and distribution of that strengthening is argued (Hill *et al.*, 2011). This circulation shift could also account for a lower degree of warming observed in the mPWP at site ODP 1172, situated in the southern extent of the EAC. Furthermore, redirected flow through the Tasman Front, which ultimately bathes the Chatham Rise, may account for the high degree of warming displayed by sites ODP 1123 and ODP 1125 (Table 2).

Furthermore, previous studies for the Last Interglacial (MIS 5e; 125 ka) suggest a warming of southern and eastern New Zealand (specifically based on data from site ODP 1123) may be a result of an increased and extended flow of the EAC becoming entrained in the Subtropical Front that would bathe the Chatham Rise sites (Fig 7a; Cortese *et al.*, 2013). The results presented here support a strengthening of the EAC and outlets relative to pre-industrial and modern, which is consistent with paleo studies for Late Pleistocene interglacials (Bostock *et al.*, 2015; Cortese *et al.*, 2013) and suggest these currents may have multiple ways of operating under warmer climates. Indeed,



485 modern EAC transport and outlets are underestimated by most models (Chiswell *et al.*, 2015; Sen Gupta *et al.*, 2016, 2021).



490 **Figure 7: Absolute Sea Surface Temperatures as a latitudinal transect of the Southwest Pacific with a) HadISST (NCAR, 2022), mid-Pliocene Warm Period interglacial (solid red with the red ribbon showing mean to maximum SSTs), PlioMIP multi-model mean (solid blue) and Marine Isotope Stage (MIS) 5e (~125 ka) (dashed yellow; Cortese *et al.*, 2013), and b) HadISST (dotted grey; NCAR, 2022), mid-Pliocene Warm Period interglacial modal mean (solid red), NZESM (dashed black) and UKESM (dashed purple) for SSP2-4.5 2090–2099 AD (Williams *et al.*, 2016; Sellar *et al.*, 2019).**

495 **4.1.1 Paleo – model comparison**

Warming during the interglacial modal means of the mPWP can be simplified as  $>4^{\circ}\text{C}$  above pre-industrial in five of the seven mid-latitude sites across the region, with two sites (DSDP 593 and DSDP 594) showing moderate warming ( $<2^{\circ}\text{C}$ ) (Table 2). This pattern is broadly reflected in both UKESM and NZESM projected scenarios explored here, with closer fit under middle of the road emission scenario (Fig. 6 & 7b; Table 4). NZESM and UKESM show a general trend (for the seven mid-latitude sites) of closer correlation to mPWP at lower temperature sites with increasing underestimation at sites with higher SSTs for all scenarios except for SSP3-7.0 (Fig. 6). These low temperature sites (DSDP 593 & DSDP 594) are also the two sites where UKESM provides systematically warmer values than NZESM (Fig. 6; Table 4).

505 The key differences between the UKESM and NZESM can be summarised by more distributed region-wide warming in the NZESM, with reduced warming along the EAC and offshore eastern New Zealand (Fig. 6). The pattern of NZESM SST field reflects local oceanographic grid refinement, which improves the fidelity of complex regional current transport and the representation of ocean fronts (Behrens *et al.*, 2020). The concentrated regionally-limited warming of the UKESM, is less consistent with the mPWP signature of warming across the region (Fig. 6). Specifically, NZESM present lower SSTs for the EAC relative to the UKESM, which is more

consistent with SSTs at site ODP 1172 during the mPWP (Fig. 6). This also corroborates the apparent intense warming observed at site DSDP 590 in the northern Tasman Sea during the mPWP, because of increased flow eastward to the Tasman Front at the expense of an invigorated EAC but may also reflect the paleogeographic positioning of site DSDP 590 and the Tasman Front (Strogen *et al.*, 2022). Significantly, while the NZESM produces a Subtropical Front further south than the UKESM, bathing DSDP 594 in warmer waters, this does not extend to the eastern Chatham Rise sites, which is not consistent with mPWP observations (Fig. 6g–i). Lastly, we note that warming at site ODP 1168 (southwest Tasmania) is comparable to site ODP 1172 (southeast Tasmania; EAC) during the mPWP, which is inconsistent with both NZESM and UKESM (Fig. 6).

Modelled SSTs at the sites show increased variability under higher-emission scenarios (Fig. 6) more comparable to the range and magnitude of mPWP observations, however, this is driven by closer values at DSDP 590 and tends to overestimate SSTs at the other sites (Fig. 6; Table 4 & S5). Rather, for both UKESM and NZESM, SSP2-4.5 for 2090–2099 AD show the least deviation to all sites reconstructed for the mPWP (Fig. 6d–f; & 7b). While the magnitude of warming changes significantly with SSP projections, we consider both UKESM and NZESM produce a pattern of warming consistent with site observations of the mPWP.

Considering regional SSTs in the meridional context, we have compared HadISST, mPWP and PlioMIP with Last Interglacial MIS 5e (Fig. 7a) and ESM scenarios for SSP2-4.5 by 2090–2099 AD (Fig. 7b). The glacial and interglacial gradients of the mPWP are relatively consistent and show a much steeper gradient in comparison to the interglacial MIS 5e (+1–2°C) and pre-industrial HadISST (Fig. 7a). While future ESM projections for SSP2-4.5 2090–2099 AD (as the closest scenario to mPWP) shows much more comparable distribution with the mPWP (Fig. 7b).

#### 4.2 Global and regional warming

In comparing our interglacial modal mean mPWP SSTs, reconstructed using organic biomarker proxies, in conjunction with data sampled for the same sites from transient ESM simulations to run to 2100 AD, we acknowledge that the ESM values do not reflect the future equilibrium temperature responses for these Southwest Pacific sites. However, in most cases, average regional temperatures at the sites studied are expected to increase beyond 2100 AD as longer duration feedbacks in the Earth climate system play out. To consider the difference between equilibrium and transient climates we discuss the variance of global and regional SSTs between paleo and future scenarios.

Global SSTs during mPWP interglacials are ~3°C above pre-industrial (Masson-Delmotte *et al.*, 2013; Dowsett *et al.*, 2016; McClymont *et al.*, 2020; Haywood *et al.*, 2020), comparable to expected warming (2.1–3.5°C) for SSP2-4.5 by 2100 AD (IPCC, 2022). The pattern and magnitude of regional warming is similar between the mPWP and ESM simulations under SSP2-4.5 (Table 4; Fig 6–7b), however the global warming generated by the ESMs under SSP2-4.5 is 4°C. Thus, while these mPWP proxy SSTs present a higher degree of warming than global, the same degree of warming from UKESM and NZESM requires 1°C higher global temperature increase.

The ECS of the UKESM (and NZESM) is 5.4°C (Sellar *et al.*, 2019; Senior *et al.*, 2020), which exceeds that of estimates for the mPWP of 2.6–4.8°C (MIS KM5c; Haywood *et al.*, 2020), and far exceeds that of the likely range (2.5–4°C) for the CMIP6 ensemble ((Forster *et al.*, 2021). This is of importance because the Southern Ocean has long been identified as having significant deviation from models to observations and it is uncertain whether high ECS models (linked to shortwave cloud feedbacks; Zelinka *et al.*, 2020) act to better estimate observations (Schuddeboom and McDonald, 2021). Here, we show the high ECS simulations of NZESM and UKESM, present a comparable warming signature seen during the mPWP in the Southwest Pacific, as opposed to the lower ECS PlioMIP simulations (Fig. 7b). These results demonstrate that while higher ECS models do produce more extreme regional temperature response under transient climates and ~100 year-timescales, they require a higher degree of global warming, suggesting longer-term feedbacks including ice dynamics may play a significant role in accurately determining committed warming, particularly for this region in proximity to Antarctica and the Southern Ocean. Furthermore, the use of lower ECS models (e.g. majority of the CMIP6 ensemble) for regional downscaling in the Southwest Pacific may be underestimating the amplified warming signal we see in the mPWP and ESM SSP2-4.5 scenarios (Appendix C).

## 5 Conclusions

The regional expression of warming differs from the global average on a variety of timescales and has significant implications for the frequency and extent of climate induced hazards related to weather, sea-level rise and socio-economic factors. Our mPWP proxy SST reconstructions for interglacial modal means show warming at sites across the Southwest Pacific averaged at 4.2°C, that is 1-2°C above global warming (Masson-Delmotte *et al.*, 2013). This mPWP SST signature contains significant regional variability that is not seen in PlioMIP multi-model mean and exceeds the Southwest Pacific PlioMIP average of 2.4°C (Haywood *et al.*, 2020), but do replicate warming at the three sites used in the PRISM climate reconstruction (Dowsett *et al.*, 2016; McClymont *et al.*, 2020).

The NZESM and UKESM show relatively consistent warming under low- and high-emission pathway simulations, but the NZESM presents slightly warmer site averages in all scenarios. The most comparable warming to mPWP by the ESMs is for 2090-2099 AD under the SSP2-4.5 scenario that is expected to reach 2.1-3.5°C globally by 2100 AD (IPCC, 2022). However, the global warming for these ESMs under this pathway is ~4°C, which relates to the high ECS of the models. This suggests, that high ECS models better replicate the regional warming signature in the Southwest Pacific, and that low ECS models in the CMIP6 ensemble may underestimate warming in the Southwest Pacific. Ultimately, testing of longer-term scenarios using NZESM, to accommodate for long feedbacks, for instance, potentially including a quantitative ice-sheet model (Smith *et al.*, 2021), would provide insight into impacts of warming on ocean currents in the Southwest Pacific and determine the effect of transient and equilibrium climate responses.

Paleoclimate reconstructions, such as those presented in this study, act as the only available evidence of equilibrium climate response to conditions predicted for the near future. While equilibrium climate states are not directly comparable to the transient future projections, expected sustained warming may result in comparable conditions.

590 **Appendix A**

Lipid biomarkers were analysed in the Organic Geochemistry Laboratory at GNS Science as reported in Naeher *et al.* (2012, 2014) and Ohkouchi *et al.* (2005) with modifications. In brief, freeze-dried and homogenized sediment samples (10–17 g) were extracted four times with dichloromethane (DCM)/ methanol (MeOH) (3:1, v:v) by ultrasonication for 20 min each time. Elemental sulfur was removed by activated copper. The total lipid  
595 extracts (TLEs) were divided into three fractions via liquid chromatography over silica columns using *n*-hexane (F1), *n*-hexane/DCM (1:2, v:v; F2) and DCM/MeOH (1:1, v:v; F3), respectively.

The F2 fractions containing alkenones were analysed using gas chromatography mass spectrometry (GC-MS) on an Agilent 7890A GC System, equipped with an Agilent J&W HP-1ms capillary column [60 m × 0.32 mm inner  
600 diameter (i.d.) × 0.25 μm film thickness (f.t.)], and connected through a splitter to an Agilent 5975C inert MSD mass spectrometer and flame ionisation detector (FID). The oven was heated from 70°C to 280°C at 20°C min<sup>-1</sup>, then at 4°C min<sup>-1</sup> to 320°C and held isothermal for 20 minutes with a total run time of 40.5 minutes. Helium was used as carrier gas with a constant flow of 1.0 mL min<sup>-1</sup>. Samples (1 μL) were injected splitless at an inlet temperature of 320°C. The MS was operated in electron impact ionisation mode at 70 eV using a source  
605 temperature of 230°C. For alkenone identification, the MS was operated in simultaneous full scan and single ion monitoring (SIM) mode at *m/z* 55, 58, 97, 109.1, 526.5, 528.5 and 530.5. Alkenones were quantified using FID. Glycerol dialkyl glycerol tetraethers (GDGTs), present in the F3 fractions, were dissolved in *n*-hexane/isopropanol (99:1, v:v) and filtered with 0.45 μm PTFE filters prior to liquid chromatography mass spectrometry (LC-MS) analysis at the University of Hokkaido, Japan. GDGTs were analysed on an Agilent 1260 HPLC system coupled  
610 to an Agilent 6130 Series quadrupole MS. Separation was achieved using a Prevail Cyano column (2.1 × 150 mm, 3 μm; Grace Discovery Science, USA) maintained at 30°C following the method of Hopmans *et al.* (2000) and Schouten *et al.* (2007). Conditions were: flow rate 0.2 ml/min, isocratic with 99% *n*-hexane and 1% isopropanol for the first 5 min followed by a linear gradient to 1.8% isopropanol over 45 min. Ionization was achieved using atmospheric pressure, positive ion chemical ionization. The spectrometer was run in selected ion monitoring mode  
615 (*m/z* 743.8, 1018, 1020, 1022, 1032, 1034, 1036, 1046, 1048, 1050, 1292.3, 1296.3, 1298.3, 1300.3, and 1302.3). Compounds were identified by comparing mass spectra and retention times with those in the literature (Hopmans *et al.*, 2000).

The  $U_{37}^{K'}$  index is defined based on the relative abundance of the C<sub>37:2</sub> and C<sub>37:3</sub> alkenones according to Prahl and  
620 Wakeham (1987) as follows:

$$U_{37}^{K'} = [C_{37:2}] / ([C_{37:2}] + [C_{37:3}]) \quad (1)$$

We used the calibration of Müller *et al.* (1998) and BAYSPLINE (Tierney and Tingley, 2018) to reconstruct SSTs  
625 from the  $U_{37}^{K'}$  index.

The  $TEX_{86}$  index is based on the relative distribution of isoprenoidal glycerol dialkyl glycerol tetraethers (isoGDGTs) in marine sediments, originally defined by Schouten *et al.* (2002):

$$TEX_{86} = \frac{[GDGT-2]+[GDGT-3]+[GDGT-4']}{[GDGT-1]+[GDGT-2]+[GDGT-3]+[GDGT-4']} \quad (2)$$

630 where GDGT-1, GDGT-2 and GDGT-3 are characterized by one, two and three cyclopentane moieties and cren' is the regioisomer of crenarchaeol. This index derived from core top samples was calibrated to SSTs using linear regressions as proposed by Schouten *et al.* (2002) and Kim *et al.* (2008).

To test the reliability of reconstructed SSTs and to increase confidence in the choice of the applied calibrations, we have compared  $U_{37}^{K'}$  and  $TEX_{86}$  SST at two sites. While  $U_{37}^{K'}$  SSTs using the BAYSPLINE (Tierney and Tingley, 2018) does yield slightly cooler temperatures (up to 0.7 °C) at higher-latitude sites than the calibration of Müller *et al.* (1998), the  $TEX_{86}$  SSTs differ by +6.4 to -16.9°C dependent on the calibration used (Figs. A1, A2). This proxy may be compromised at sites with high soil organic matter inputs (Hopmans *et al.*, 2004) and high contributions of sedimentary GDGTs (Pancost *et al.*, 2001; Zhang *et al.*, 2011) which is considered negligible in open-marine environments. Other non-temperature controls such as oxygen concentrations, growth phases, nutrient cycling may be introduced in upwelling zones but are not able to be addressed here due to limited understanding of these effects (Elling *et al.*, 2014; Qin *et al.*, 2015; Hollis *et al.*, 2019). Non-linear calibrations such as the  $TEX_{86}^H$  index (Kim *et al.*, 2010) were developed to extend the calibrated SST range of the previous calibrations, however this may underestimate SSTs in ancient greenhouse climates (Tierney and Tingley, 2015; O'Brien *et al.*, 2017, Hollis *et al.*, 2019) and a non-linear relationship contradicts available experimental evidence suggesting a linear relationship with SST (Pitcher *et al.*, 2010; Schouten *et al.*, 2013; Elling *et al.*, 2014). Therefore, a Bayesian approach (BAYSPAR; Tierney and Tingley, 2015) was developed to consider spatially varying uncertainty derived from modern SST distribution is widely used. Additionally, a new machine-learning approach (OPTIMAL: Optimised Palaeothermometry from Tetraethers via MACHine Learning) aims to address uncertainty in the method application to paleo SST and determine SST beyond the modern range (>30°C) (Dunkley Jones *et al.*, 2020). In comparing the two independent biomarker proxies of derived SST using  $U_{37}^{K'}$  - BAYSPLINE with  $TEX_{86}$  calibrations of Schouten *et al.* (2002), Kim *et al.* (2010), Tierney and Tingley (2015), and Dunkley Jones *et al.* (2020), we find all calibrations are comparable ( $\sim\pm 5^\circ\text{C}$ ) but the BAYSPAR approach of Tierney and Tingley (2015) displays the closest values to  $U_{37}^{K'}$ -BAYSPLINE (Fig. A1 & A2; Table A1). Notably, the calibrations of Schouten *et al.* (2002), Kim *et al.* (2010) and BAYSPAR (Tierney and Tingley, 2015) show less scatter at higher temperatures (>25°C; Fig. A1), while the OPTIMAL calibration (Dunkley Jones *et al.*, 2020) presents offsets of up to -15°C (Fig. A2) in comparison to  $U_{37}^{K'}$ -BAYSPLINE.

650 The  $TEX_{86}$  calibration of Tierney and Tingley (2015) (BAYSPAR) shows the closest values to the  $U_{37}^{K'}$  - SST BAYSPLINE and lowest scatter (Figs. A1, A2; Table A1), and therefore are selected for display in Fig. 4. In contrast, the calibration of Schouten *et al.* (2002) shows larger scatter in reconstructed SSTs than BAYSPAR (Figs. A1, A2). The calibration of Kim *et al.* (2010) yields similar SST estimates as Schouten *et al.* (2002) and BAYSPAR, but seems to overestimate SSTs at lower temperatures. In contrast, OPTIMAL (Dunkley-Jones *et al.*, 2020) appears to underestimate SSTs at higher temperatures. Importantly, the general agreement in SST reconstruction from two independent biomarker provides higher confidence in the results.




Figure A1: Comparison between  $U_{37}^{K'}$  derived SST using BAYSPLINE with  $TEX_{86}$  Index SST calibrations of Schouten *et al.* (2002), Kim *et al.*, (2010), Dunkley Jones *et al.* (2020) and Tierney and Tingley (2015).





Figure A2: Comparison between  $U_{37}^{K'}$  derived SST using BAYSPLINE with  $TEX_{86}$  Index SST calibrations of Schouten *et al.* (2002), Kim *et al.*, (2010), OPTIMAL (Dunkley Jones *et al.*, 2020) and BAYSPAR (Tierney and Tingley, 2015).

**Table A1: Comparison between  $U_{37}^{K'}$  derived SST using BAYSPLINE with TEX<sub>86</sub> Index SST calibrations of Schouten *et al.* (2002), Kim *et al.* (2010), OPTIMAL (Dunkley Jones *et al.*, 2020) and BAYSPAR (Tierney and Tingley, 2015).**

TEX <sub>86</sub> calibrations	Average Difference (°C) of TEX <sub>86</sub> SST –relative to BAYSPLINE SST reconstructions
BAYSPAR 2015	0.1
Kim2010	0.8
OPTIMAL2020	-2.3
Schouten2002	-0.6


**Appendix B**



680 **Figure B1. Bimodal analysis for each site after Benaglia *et al.*, (2010), excluding ANDRILL as it only represents interglacial conditions, displaying density curves with calculated bimodal distributions interpreted as glacial distributions (red) and interglacial (green). Code is available.**



## Appendix C



685 **Figure C1. Extended version of Figure 6 to include a second high ECS model (CESM2: ECS 5.2-5.6°C; Danabasoglu et al., 2020) and the lowest ECS model in CMIP6 (INM: ECS 1.8°C; Volodin et al., 2018). Regional Sea Surface Temperature (SST) anomalies to HadISST (1870 – 1879 AD) for SSP1-2.6 (a–c), SSP2-4.5 (d–f), SSP3-7.0 (g–i) in 2090–2099 AD compared to mid-Pliocene Warm Period (mPWP) site mean interglacial SST anomalies (filled circles using same colour scale as map). Panels (a-c) are NZESM, panels (d-f) are UKESM, panels (g-i) are CESM2, panels (j-l) are INM and the far-right panels (m-o) are site SST anomalies between 31–46°S for mPWP (red dotted), UKESM (purple dashed) and NZESM (black solid), CESM2 (dark green solid) and INM (light green dashed). The INM low ECS model shows a significantly different pattern of warming to the high ECS models of NZESM, UKESM and CESM2.**

690

## Code and Data availability

Data tables and supplementary tables: DOI 10.5281/zenodo.7109199

Script and necessary data files: <https://github.com/GRG-GNS/Pliocene-SST-Southwest-Pacific>

695 Table S1. All site sea surface temperature (SST; °C) data used in results with index and calibrations of Müller98 (Müller *et al.*, 1998) and BAYSPLINE (Tierney and Tingley, 2018). The proxy type and references are also provided.

Table S2. Site sample data for analyses undertaken this study, including all and TEX<sub>86</sub> index calculations and calibrations. References for calibrations are contained within column headers.

700 Table S3. Seasonal and annual mean sea surface temperature (SST; °C) model outputs of HadISST (NCAR, 2022), UKESM (Sellar *et al.*, 2019), NZESM (Williams *et al.*, 2016) at the seven Southwest Pacific sites (DSDP 594, ODP 1172, ODP 1168, ODP 1125, ODP 1123, DSDP 593, DSDP 590) for SSP2 2040 AD (2036-2045 AD), and SSP1, 2, and 3 2095 AD (2090-2099 AD). Including UKESM and NZESM with respect to HadISST.

705 Table S4. Site sea surface temperature (SST; °C) annual means and seasonal range for UKESM and NZESM SSP2-4.5 2036-2045 AD, with MPWP interglacial modal means and total glacial range (maximum to minimum SST).

Table S5. Compiled sea surface temperature (SST; °C) interglacial means for MIS 5e (125kyr; Cortese *et al.*, 2013) and mPWP (3.3-3.0 Ma) and model annual means for HadISST (1870-1879 AD), and SSP2-4.5 2090-2099 AD for UKESM, NZESM.

## 710 Sample availability

Samples were obtained from the International Ocean Discovery Program, Texas A&M University.

## Author Contribution

GRG, JHTW and SN designed the project. SN, OS and MY measured and analysed the data. JHTW and AMH provided climate model simulations. GRG prepared the manuscript with the contribution of all authors.

## 715 Competing interests

The authors declare that they have no conflict of interest.

## Acknowledgements

720 We thank the International Ocean Discovery Program (IODP), which provided the samples, and the Australia-New Zealand IODP Consortium (ANZIC), which provided Legacy Analytical Funding for this study. ANZIC is supported by the Australian Government through the Australian Research Council's LIEF funding scheme [LE160100067] and the ANZIC of universities and government agencies. We acknowledge support from GNS Science and the New Zealand Ministry of Business, Innovation and Employment through the Global Change Through Time research program and Organic Geochemistry Laboratory (contract C05X1702). JW obtained funding and support through the Ministry of Business Innovation and Employment Deep South National Science 725 Challenge project number C01X1412. The development of the UKESM, was supported by the Met Office Hadley

Centre Climate Programme funded by BEIS and Defra (GA01101) and by the Natural Environment Research Council (NERC) national capability grant for the UK Earth System Modelling project, grant number NE/N017951/1. JW wishes to acknowledge the use of New Zealand eScience Infrastructure (NeSI) high performance computing facilities, consulting support and training services as part of this research. New Zealand's national facilities are provided by NeSI and funded jointly by NeSI's collaborator institutions and through the Ministry of Business, Innovation & Employment's Research Infrastructure programme, [www.nesi.org.nz](http://www.nesi.org.nz). OS acknowledges the support by Japan Society of Promotion of Science funded by Ministry of Education, Culture, Sports, Science and Technology, Japan (KAKENHI 26287129 and 17H06318). We also wish to acknowledge the invaluable sea surface temperature compilations and assessment by PlioVar (Pliocene Climate Variability), a Past Global Changes (PAGES) working group and the contributing authors.

## References

- Ahn, S., Khider, D., Lisiecki, L. E., Lawrence, C. E.: A probabilistic Pliocene-Pleistocene stack of benthic  $\delta^{18}\text{O}$  using a profile hidden Markov model, *Dynamics and Statistics of the Climate System*, 2:1. <https://doi.org/10.1093/climsys/dzx002>, 2017.
- Amante, C. and B.W. Eakins.: ETOPO1 1 Arc-Minute Global Relief Model: Procedures, Data Sources and Analysis. NOAA Technical Memorandum NESDIS NGDC-24. National Geophysical Data Center, NOAA. ,2009.
- Batchelor, C.L., Margold, M., Krapp, M., Murton, D.K., Dalton, A.S., Gibbard, P.L., Stokes, C.R., Murton, J.B. and Manica, A.: The configuration of Northern Hemisphere ice sheets through the Quaternary. *Nature communications*, 10(1), pp.1-10. <https://doi.org/10.1038/s41467-019-11601-2>, 2019.
- Behrens, E., Fernandez, D., and Sutton, P: Meridional oceanic heat transport influences marine heatwaves in the Tasman Sea on interannual to decadal timescales. *Front. Mar. Sci.* 6, 228. <https://doi.org/10.3389/fmars.2019.00228>, 2019.
- Behrens, E., Williams, J., Morgenstern, O., Sutton, P., Rickard, G. and Williams, M.J.: Local grid refinement in New Zealand's earth system model: Tasman Sea ocean circulation improvements and super-gyre circulation implications. *Journal of Advances in Modeling Earth Systems*, 12(7), <https://doi.org/10.1029/2019MS001996>, 2019.
- Behrens, E., Rickard, G., Rosier, S., Williams, J., Morgenstern, O. and Stone, D.: Projections of Future Marine Heatwaves for the Oceans Around New Zealand Using New Zealand's Earth System Model. *Frontiers in Climate*, p.19. <https://doi.org/10.3389/fclim.2022.798287>, 2022.
- Benaglia, T., Chauveau, D., Hunter, D.R. and Young, D.S.: mixtools: an R package for analyzing mixture models. *Journal of statistical software*, 32, pp.1–29. <https://github.com/dsy109/mixtools> , 2010.
- Bertram, R.A., Wilson, D.J., van de Flierdt, T., McKay, R.M., Patterson, M.O., Jimenez-Espejo, F.J., Escutia, C., Duke, G.C., Taylor-Silva, B.I. and Riesselman, C.R.: Pliocene deglacial event timelines and the biogeochemical response offshore Wilkes Subglacial Basin, East Antarctica. *Earth and Planetary Science Letters*, 494, pp.109-116. <https://doi.org/10.1038/s41586-018-0501-8>, 2018.
- Bostock, H.C., Hayward, B.W., Neil, H.L., Sabaa, A.T. and Scott, G.H.: Changes in the position of the Subtropical Front south of New Zealand since the last glacial period. *Paleoceanography*, 30(7), pp.824–844. <https://doi.org/10.1002/2014PA002652>, 2015.

- Burke, K.D., Williams, J.W., Chandler, M.A., Haywood, A.M., Lunt, D.J. and Otto-Bliesner, B.L.: Pliocene and Eocene provide best analogs for near-future climates. *Proceedings of the National Academy of Sciences*, 115(52), pp.13288–13293. <https://doi.org/10.1073/pnas.1809600115>, 2018.
- Caballero-Gill, R.P., Herbert, T.D. and Dowsett, H.J.: 100-kyr paced climate change in the Pliocene warm period, Southwest Pacific. *Paleoceanography and Paleoclimatology*, 34(4), pp.524–545. <https://doi.org/10.1029/2018PA003496>, 2019.
- Carter, R.M., McCave, I.N., and Carter, L.: Leg 181 synthesis: fronts, flows, drifts, volcanoes, and the evolution of the southwestern gateway to the Pacific Ocean, eastern New Zealand. In Richter, C. (Ed.), Proc. ODP, Sci. Results, 181: College Station, TX (Ocean Drilling Program), 1–111. doi:10.2973/odp.proc.sr.181.210.2004, 2004.
- Chalk, T.B., Hain, M.P., Foster, G.L., Rohling, E.J., Sexton, P.F., Badger, M.P., Cherry, S.G., Hasenfratz, A.P., Haug, G.H., Jaccard, S.L. and Martínez-García, A.: Causes of ice age intensification across the Mid-Pleistocene Transition. *Proceedings of the National Academy of Sciences*, 114(50), pp.13114–13119. [10.1073/pnas.1702143114](https://doi.org/10.1073/pnas.1702143114), 2017.
- Chen, L., Cao, L., Zhou, Z., Zhang, D. and Liao, J.: A New Globally Reconstructed Sea Surface Temperature Analysis Dataset since 1900. *Journal of Meteorological Research*, 35(6), pp.911-925. <https://doi.org/10.1007/s13351-021-1098-7>, 2021.
- Chiswell, S.M., Bostock, H.C., Sutton, P.J. and Williams, M.J.: Physical oceanography of the deep seas around New Zealand: a review. *New Zealand Journal of Marine and Freshwater Research*, 49(2), pp.286–317. <https://doi.org/10.1080/00288330.2014.992918>, 2015.
- Chiswell, S.M.: Atmospheric wavenumber-4 driven South Pacific marine heat waves and marine cool spells. *Nature Communications*, 12(1), pp.1-8. <https://doi.org/10.1038/s41467-021-25160-y>, 2021.
- Conte, M.H., Sicre, M.A., Rühlemann, C., Weber, J.C., Schulte, S., Schulz-Bull, D. and Blanz, T.: Global temperature calibration of the alkenone unsaturation index (UK'37) in surface waters and comparison with surface sediments. *Geochemistry, Geophysics, Geosystems*, 7(2). <https://doi.org/10.1029/2005GC001054>, 2006.
- Cook, C.P., Van De Fliedrt, T., Williams, T., Hemming, S.R., Iwai, M., Kobayashi, M., Jimenez-Espejo, F.J., Escutia, C., González, J.J., Khim, B.K. and McKay, R.M.: Dynamic behaviour of the East Antarctic ice sheet during Pliocene warmth. *Nature Geoscience*, 6(9), pp.765-769. <https://doi.org/10.1038/ngeo1889>, 2013.
- Cortese, G., Dunbar, G.B., Carter, L., Scott, G., Bostock, H., Bowen, M., Crundwell, M., Hayward, B.W., Hargreaves, J. C. and Annan, J. D.: Could the Pliocene constrain the equilibrium climate sensitivity?, *Clim. Past*, 12, 1591–1599, <https://doi.org/10.5194/cp-12-1591-2016>, 2016.
- Howard, W., Martínez, J.I. and Moy, A.: Southwest Pacific Ocean response to a warmer world: insights from Marine Isotope Stage 5e. *Paleoceanography*, 28(3), pp.585–598. <https://doi.org/10.1002/palo.20052>, 2013.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., et al. The Community Earth System Model Version 2 (CESM2). *Journal of Advances in Modeling Earth Systems*, 12. <https://doi.org/10.1029/2019MS001916>, 2020.
- DeConto, R.M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise. *Nature*, 531(7596), pp.591-597. <https://doi.org/10.1038/nature17145>, 2016.
- DeConto, R.M., Pollard, D., Alley, R.B., Velicogna, I., Gasson, E., Gomez, N., Sadai, S., Condrón, A., Gilford, D.M., Ashe, E.L. and Kopp, R.E.: The Paris Climate Agreement and future sea-level rise from Antarctica. *Nature*, 593(7857), pp.83-89. <https://doi.org/10.1038/s41586-021-03427-0>, 2021.

- De La Vega, E., Chalk, T.B., Wilson, P.A., Bysani, R.P. and Foster, G.L.: Atmospheric CO<sub>2</sub> during the Mid–Piacenzian Warm Period and the M2 glaciation. *Scientific reports*, 10(1), pp.1–8. 10.1038/s41598-020-67154-8, 2020.
- De Bar, M.W., Rampen, S.W., Hopmans, E.C., Damsté, J.S.S. and Schouten, S.: Constraining the applicability of organic paleotemperature proxies for the last 90 Myrs. *Organic Geochemistry*, 128, pp.122–136. <https://doi.org/10.1016/j.orggeochem.2018.12.005>, 2019.
- 810 Dennison, F., Keeble, J., Morgenstern, O., Zeng, G., Abraham, N.L. and Yang, X.: Improvements to stratospheric chemistry scheme in the um-ukca (v10. 7) model: Solar cycle and heterogeneous reactions. *Geoscientific Model Development*, 12(3), pp.1227-1239. <https://doi.org/10.5194/gmd-12-1227-2019>, 2019.
- Doblas–Reyes, F.J., A.A. Sörensson, M. Almazroui, A. Dosio, W.J. Gutowski, R. Haarsma, R. Hamdi, B. Hewitson, W.–T. Kwon, B.L. Lamptey, D. Maraun, T.S. Stephenson, I. Takayabu, L. Terray, A. Turner, and Z. 815 Zuo.: Linking Global to Regional Climate Change. In Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change Masson–Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.). Cambridge University Press. In Press. Doi:10.1017/9781009157896.012., 820 2021.
- Dowsett, H.J., Robinson, M.M., Stoll, D.K., Foley, K.M., Johnson, A.L., Williams, M. and Riesselman, C.R.: The PRISM (Pliocene palaeoclimate) reconstruction: time for a paradigm shift. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 371(2001), p.20120524. [doi.org/10.1098/rsta.2012.0524](https://doi.org/10.1098/rsta.2012.0524), 2013.
- 825 Dowsett, H., Dolan, A., Rowley, D., Moucha, R., Forte, A.M., Mitrovica, J.X., Pound, M., Salzmann, U., Robinson, M., Chandler, M. and Foley, K.: The PRISM4 (mid–Piacenzian) paleoenvironmental reconstruction. *Climate of the Past*, 12(7), pp.1519–1538. <https://doi.org/10.5194/cp-12-1519-2016>, 2016.
- Dunkley Jones, T., Eley, Y.L., Thomson, W., Greene, S.E., Mandel, I., Edgar, K. and Bendle, J.A.: OPTiMAL: A new machine learning approach for GDGT–based palaeothermometry. *Climate of the Past*, 16(6), pp.2599– 830 2617. <https://doi.org/10.5194/cp-16-2599-2020>, 2020.
- Dutton, A., Carlson, A.E., Long, A.J., Milne, G.A., Clark, P.U., DeConto, R., Horton, B.P., Rahmstorf, S. and Raymo, M.E.: Sea-level rise due to polar ice-sheet mass loss during past warm periods. *Science*, 349(6244), [DOI: 10.1126/science.aaa4019](https://doi.org/10.1126/science.aaa4019), 2015.
- Elling, F.J., Könneke, M., Lipp, J.S., Becker, K.W., Gagen, E.J. and Hinrichs, K.U.: Effects of growth phase on 835 the membrane lipid composition of the thaumarchaeon *Nitrosopumilus maritimus* and their implications for archaeal lipid distributions in the marine environment. *Geochimica et Cosmochimica Acta*, 141, pp.579–597. [Doi.org/10.1016/j.gca.2014.07.005](https://doi.org/10.1016/j.gca.2014.07.005), 2014.
- Eyring, V., Bony, S., Meehl, G.A., Senior, C.A., Stevens, B., Stouffer, R.J. and Taylor, K.E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, 9(5), pp.1937–1958. <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- 840 Exon, N.F., Kennett, J.P., Malone, M.J., *et al.*: Proc. ODP, Init. Repts., 189: College Station, TX (Ocean Drilling Program). [Doi.org/10.2973/odp.proc.ir.189.2001](https://doi.org/10.2973/odp.proc.ir.189.2001), 2001.



- Fischer, H., Meissner, K.J., Mix, A.C. *et al.*: Palaeoclimate constraints on the impact of 2 °C anthropogenic warming and beyond. *Nature Geosci* **11**, 474–485. <https://doi.org/10.1038/s41561-018-0146-0>, 2018.
- 845 Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen, M.D. Palmer, M. Watanabe, M. Wild, and H. Zhang,: The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, 850 J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 923–1054, doi: 10.1017/9781009157896.009, 2021.
- Golledge, N.R., Keller, E.D., Gomez, N., Naughten, K.A., Bernales, J., Trusel, L.D. and Edwards, T.L.: Global environmental consequences of twenty-first-century ice-sheet melt. *Nature*, *566*(7742), pp.65–72. 855 <https://doi.org/10.1038/s41586-019-0889-9> , 2019.
- Grant, G.R., Naish, T.R., Dunbar, G.B., Stocchi, P., Kominz, M.A., Kamp, P.J., Tapia, C.A., McKay, R.M., Levy, R.H. and Patterson, M.O.: The amplitude and origin of sea-level variability during the Pliocene epoch. *Nature*, *574*(7777), pp.237–241. <https://doi.org/10.1038/s41586-019-1619-z>, 2019.
- Grant, G. and Naish, T.: Pliocene sea-level revisited: is there more than meets the eye? *PAGES Magazine* *29*, 860 <https://doi.org/10.22498/pages.29.1.34>, 2021.
- Haywood, A.M., H.J. Dowsett, M.M. Robinson, D.K. Stoll, A.M. Dolan, D.J. Lunt, B. Otto-Bliesner, and M.A. Chandler.: Pliocene Model Intercomparison Project (PlioMIP): Experimental design and boundary conditions (Experiment 2). *Geosci. Model Dev.*, *4*, 571–577, doi:10.5194/gmd-4-571-2011. 2011.
- Haywood, A.M., Dowsett, H.J., Dolan, A.M., Rowley, D., Abe-Ouchi, A., Otto-Bliesner, B., Chandler, M.A., 865 Hunter, S.J., Lunt, D.J., Pound, M. and Salzmann, U.: The Pliocene model intercomparison project (PlioMIP) phase 2: scientific objectives and experimental design. *Climate of the Past*, *12*(3), pp.663–675. <https://doi.org/10.5194/cp-12-663-2016>, 2016.
- Haywood, A.M., Valdes, P.J., Aze, T., Barlow, N., Burke, A., Dolan, A.M., Von Der Heydt, A.S., Hill, D.J., Jamieson, S.S.R., Otto-Bliesner, B.L. and Salzmann, U.: What can Palaeoclimate Modelling do for you?. *Earth 870 Systems and Environment*, *3*(1), pp.1–18. <https://doi.org/10.1007/s41748-019-00093-1> , 2019.
- Haywood, A.M., Tindall, J.C., Dowsett, H.J., Dolan, A.M., Foley, K.M., Hunter, S.J., Hill, D.J., Chan, W.L., Abe-Ouchi, A., Stepanek, C. and Lohmann, G.: The Pliocene Model Intercomparison Project Phase 2: large-scale climate features and climate sensitivity. *Climate of the Past*, *16*(6), pp.2095–2123. <https://doi.org/10.5194/cp-16-2095-2020>, 2020.
- 875 Herbert, T.D., Peterson, L.C., Lawrence, K.T. and Liu, Z.: Tropical ocean temperatures over the past 3.5 million years. *Science*, *328*(5985), pp.1530–1534. /doi/10.1126/science.1233137, 2010.
- Herbert, T.D.: 8.15 Alkenone Paleotemperature Determinations. *Treatise on Geochemistry*, edited by: Holland, HD and Turekian, KK, Elsevier, Oxford, pp.361–378. <https://doi.org/10.1016/B978-0-08-095975-7.00615-X>, 2014.
- 880 Hermanson, L., Smith, D., Seabrook, M., Bilbao, R., Doblus-Reyes, F., Tourigny, E., Lapin, V., Kharin, V.V., Merryfield, W.J., Sospedra-Alfonso, R. and Athanasiadis, P.: WMO global annual to decadal climate update: a

- prediction for 2021–25. *Bulletin of the American Meteorological Society*, 103(4), pp.E1117-E1129. 10.1175/BAMS-D-20-0311.1, 2022.
- 885 Hill, K.L., Rintoul, S.R., Ridgway, K.R. and Oke, P.R.: Decadal changes in the South Pacific western boundary current system revealed in observations and ocean state estimates. *Journal of Geophysical Research: Oceans*, 116(C1). <https://doi.org/10.1029/2009JC005926> , 2011.
- Hoegh–Guldberg, O., Jacob, D., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante, R., Ebi, K., Engelbrecht, F., Guiot, J. and Hijioaka, Y.: Impacts of 1.5 C global warming on natural and human systems. Global warming of 1.5°C. <https://www.ipcc.ch/sr15/chapter/chapter-3/> , 2018.
- 890 Hollis, C.J., Dunkley Jones, T., Anagnostou, E., Bijl, P.K., Cramwinckel, M.J., Cui, Y., Dickens, G.R., Edgar, K.M., Eley, Y., Evans, D. and Foster, G.L.: The DeepMIP contribution to PMIP4: methodologies for selection, compilation and analysis of latest Paleocene and early Eocene climate proxy data, incorporating version 0.1 of the DeepMIP database. *Geoscientific Model Development*, 12(7), pp.3149–3206. <https://doi.org/10.5194/gmd-12-3149-2019> , 2019.
- 895 Hopmans, E.C., Weijers, J.W., Schefuß, E., Herfort, L., Damsté, J.S.S. and Schouten, S., A novel proxy for terrestrial organic matter in sediments based on branched and isoprenoid tetraether lipids. *Earth and Planetary Science Letters*, 224(1–2), pp.107–116. [Doi.org/10.1016/j.epsl.2004.05.012](https://doi.org/10.1016/j.epsl.2004.05.012) , 2004.
- IPCC: Summary for Policymakers [H.-O. Pörtner, D.C. Roberts, E.S. Poloczanska, K. Mintenbeck, M. Tignor, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem (eds.)]. In: *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 3–33, [doi:10.1017/9781009325844.001](https://doi.org/10.1017/9781009325844.001), 2022.
- 900 Kim, J.H., Van der Meer, J., Schouten, S., Helmke, P., Willmott, V., Sangiorgi, F., Koç, N., Hopmans, E.C. and Damsté, J.S.S.: New indices and calibrations derived from the distribution of crenarchaeal isoprenoid tetraether lipids: Implications for past sea surface temperature reconstructions. *Geochimica et Cosmochimica Acta*, 74(16), pp.4639–4654. [Doi.org/10.1016/j.gca.2010.05.027](https://doi.org/10.1016/j.gca.2010.05.027), 2010.
- Karas, C., Nürnberg, D., Tiedemann, R. and Garbe-Schönberg, D.: Pliocene climate change of the Southwest Pacific and the impact of ocean gateways. *Earth and Planetary Science Letters*, 301(1-2), pp.117-124. [Doi.org/10.1016/j.epsl.2010.10.028](https://doi.org/10.1016/j.epsl.2010.10.028) , 2011.
- 910 Koenig, S.J., Dolan, A.M., De Boer, B., Stone, E.J., Hill, D.J., DeConto, R.M., Abe-Ouchi, A., Lunt, D.J., Pollard, D., Quiquet, A. and Saito, F.: Ice sheet model dependency of the simulated Greenland Ice Sheet in the mid-Pliocene. *Climate of the Past*, 11(3), pp.369-381. <https://doi.org/10.5194/cp-11-369-2015> , 2015.
- Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A.C.M. and Levrard, B.: A long-term numerical solution for the insolation quantities of the Earth. *Astronomy & Astrophysics*, 428(1), pp.261-285. <https://doi.org/10.1051/0004-6361:20041335> , 2004.
- 915 Lee, J.–Y., J. Marotzke, G. Bala, L. Cao, S. Corti, J.P. Dunne, F. Engelbrecht, E. Fischer, J.C. Fyfe, C. Jones, A. Maycock, J. Mutemi, O. Ndiaye, S. Panickal, and T. Zhou: Future Global Climate: Scenario–Based Projections and Near–Term Information. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson–Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang,

- K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press. In Press. Doi:10.1017/9781009157896.006, 2021.
- 925 Lisiecki, L.E. and Raymo, M.E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records. *Paleoceanography*, 20(1). <https://doi.org/10.1029/2004PA001071>, 2005.
- Lowry, D.P., Krapp, M., Golledge, N.R. and Alevropoulos–Borrill, A.: The influence of emissions scenarios on future Antarctic ice loss is unlikely to emerge this century. *Communications Earth and Environment*, 2(1), pp.1–14. Doi.org/10.1038/s43247-021-00289-2, 2021.
- 930 Max, L., Lembke–Jene, L., Zou, J., Shi, X. and Tiedemann, R.: Evaluation of reconstructed sea surface temperatures based on U37k' from sediment surface samples of the North Pacific. *Quaternary Science Reviews*, 243, p.106496. <https://doi.org/10.1016/j.quascirev.2020.106496>, 2020.
- Masson–Delmotte, V., Schulz, M., Abe–Ouchi, A., Beer, J., Ganopolski, A., Rouco, J.G., Jansen, E., Lambeck, K., Luterbacher, J., Naish, T. and Osborn, T.: Information from paleoclimate archives. In *Climate change 2013: the physical science basis: Contribution of Working Group I to the Fifth Assessment Report of the*
- 935 *Intergovernmental Panel on Climate Change* (pp. 383–464). Cambridge University Press. Doi:10.1017/CBO9781107415324.013, 2013.
- Martínez-Botí, M., Foster, G., Chalk, T. et al. Plio-Pleistocene climate sensitivity evaluated using high-resolution CO<sub>2</sub> records. *Nature* 518, 49–54. <https://doi.org/10.1038/nature14145>, 2015.
- 940 McClymont, E.L., Elmore, A.C., Kender, S., Leng, M.J., Greaves, M. and Elderfield, H.: Pliocene-Pleistocene evolution of sea surface and intermediate water temperatures from the Southwest Pacific. *Paleoceanography*, 31(6), pp.895–913. <https://doi.org/10.1002/2016PA002954>, 2016.
- McClymont, E.L., Ford, H.L., Ho, S.L., Tindall, J.C., Haywood, A.M., Alonso–Garcia, M., Bailey, I., Berke, M.A., Littler, K., Patterson, M.O. and Petrick, B.: Lessons from a high–CO<sub>2</sub> world: an ocean view from ~ 3 million years ago. *Climate of the Past*, 16(4), pp.1599–1615. <https://doi.org/10.5194/cp-16-1599-2020>, 2020.
- 945 McKay, R., Naish, T., Carter, L., Riesselman, C., Dunbar, R., Sjunneskog, C., Winter, D., Sangiorgi, F., Warren, C., Pagani, M. and Schouten, S.: Antarctic and Southern Ocean influences on Late Pliocene global cooling. *Proceedings of the National Academy of Sciences*, 109(17), pp.6423–6428. <https://doi.org/10.1073/pnas.1112248109>, 2012.
- Medina-Elizalde, M. and Lea, D.W.: Late Pliocene equatorial Pacific. *Paleoceanography*, 25(2). 950 <https://doi.org/10.1029/2009PA001780>, 2010.
- Meinshausen, M., Lewis, J., McGlade, C., Gütschow, J., Nicholls, Z., Burdon, R., Cozzi, L. and Hackmann, B.: Realization of Paris Agreement pledges may limit warming just below 2° C. *Nature*, 604(7905), pp.304–309. 10.1038/s41586-022-04553-z, 2022.
- 955 Miller, K.G., Wright, J.D., Browning, J.V., Kulpecz, A., Kominz, M., Naish, T.R., Cramer, B.S., Rosenthal, Y., Peltier, W.R. and Sostdian, S.: High tide of the warm Pliocene: Implications of global sea level for Antarctic deglaciation. *Geology*, 40(5), pp.407–410. Doi.org/10.1130/G32869.1, 2012.
- Müller, P.J., Kirst, G., Ruhland, G., Von Storch, I. and Rosell-Melé, A.: Calibration of the alkenone paleotemperature index U37K' based on core-tops from the eastern South Atlantic and the global ocean (60 N–60 S). *Geochimica et Cosmochimica Acta*, 62(10), pp.1757–1772. Doi.org/10.1016/S0016-7037(98)00097-0, 1998.



- 960 Naehrer, S., Smittenberg, R.H., Gilli, A., Kirilova, E.P., Lotter, A.F. and Schubert, C.J.: Impact of recent lake eutrophication on microbial community changes as revealed by high resolution lipid biomarkers in Rotsee (Switzerland). *Organic geochemistry*, 49, pp.86-95. [10.1016/j.orggeochem.2012.05.014](https://doi.org/10.1016/j.orggeochem.2012.05.014), 2012.
- Naehrer, S., Niemann, H., Peterse, F., Smittenberg, R.H., Ziegler, P.K. and Schubert, C.J.: Tracing the methane cycle with lipid biomarkers in Lake Rotsee (Switzerland). *Organic geochemistry*, 66, pp.174-181. <https://doi.org/10.1016/j.orggeochem.2013.11.002>, 2014.
- 965 Naish, T., Powell, R., Levy, R., Wilson, G., Scherer, R., Talarico, F., Krissek, L., Niessen, F., Pompilio, M., Wilson, T. and Carter, L.: Obliquity-paced Pliocene West Antarctic ice sheet oscillations. *Nature*, 458(7236), pp.322-328. [Doi.org/10.1038/nature07867](https://doi.org/10.1038/nature07867), 2009.
- Naish, T. and Zwartz, D.: Looking back to the future. *Nature Climate Change*, 2(5), pp.317-318. <https://doi.org/10.1038/nclimate1504>, 2012.
- 970 NCAR: National Center for Atmospheric Research Staff (Eds). "The Climate Data Guide: SST data: HadiSST v1.1." Retrieved from <https://climatedataguide.ucar.edu/climate-data/sst-data-hadisst-v11>, Last modified 19 Jun 2022.
- Ohkouchi, N., Xu, L., Reddy, C.M., Montluçon, D. and Eglinton, T.I.: Radiocarbon dating of alkenones from marine sediments: I. Isolation protocol. *Radiocarbon*, 47(3), pp.401-412. [Doi.org/10.1017/S0033822200035189](https://doi.org/10.1017/S0033822200035189), 2005.
- 975 O'Brien, C.L., Robinson, S.A., Pancost, R.D., Damsté, J.S.S., Schouten, S., Lunt, D.J., Alsenz, H., Bornemann, A., Bottini, C., Brassell, S.C. and Farnsworth, A.: Cretaceous sea-surface temperature evolution: Constraints from TEX86 and planktonic foraminiferal oxygen isotopes. *Earth-Science Reviews*, 172, pp.224-247. [10.1016/j.earscirev.2017.07.012](https://doi.org/10.1016/j.earscirev.2017.07.012), 2017.
- O'Neill, B.C., Tebaldi, C., Van Vuuren, D.P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.F., Lowe, J. and Meehl, G.A.: The scenario model intercomparison project (ScenarioMIP) for CMIP6. *Geoscientific Model Development*, 9(9), pp.3461-3482. <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.
- 985 Pancost, R.D., Bouloubassi, I., Aloisi, G. and Damsté, J.S.S.: Three series of non-isoprenoidal dialkyl glycerol diethers in cold-seep carbonate crusts. *Organic geochemistry*, 32(5), pp.695-707. [https://doi.org/10.1016/S0146-6380\(01\)00015-8](https://doi.org/10.1016/S0146-6380(01)00015-8), 2001.
- Patterson, M.O., McKay, R., Naish, T., Escutia, C., Jimenez-Espejo, F.J., Raymo, M.E., Meyers, S.R., Tauxe, L. and Brinkhuis, H.: Orbital forcing of the East Antarctic ice sheet during the Pliocene and Early Pleistocene. *Nature Geoscience*, 7(11), pp.841-847. <https://doi.org/10.1038/ngeo2273>, 2014.
- 990 Patterson, M.O., McKay, R., Naish, T., Bostock, H.C., Dunbar, R., Ohneiser, C., Woodard, S.C., Wilson, G. and Caballero-Gill, R.: A Southwest Pacific perspective on long-term global trends in Pliocene-Pleistocene stable isotope records. *Paleoceanography and Paleoclimatology*, 33(7), pp.825-839. <https://doi.org/10.1029/2017PA003269>, 2018.
- Pitcher, A., Rychlik, N., Hopmans, E.C., Spieck, E., Rijpstra, W.I.C., Ossebaar, J., Schouten, S., Wagner, M. and Sinninghe Damsté, J.S.: Crenarchaeol dominates the membrane lipids of *Candidatus Nitrososphaera gargensis*, a thermophilic Group I. 1b Archaeon. *The ISME Journal*, 4(4), pp.542-552. [Doi.org/10.1038/ismej.2009.138](https://doi.org/10.1038/ismej.2009.138), 2010.
- Prahl, F.G. and Wakeham, S.G.: Calibration of unsaturation patterns in long-chain ketone compositions for palaeotemperature assessment. *Nature*, 330(6146), pp.367-369. <https://doi.org/10.1038/330367a0>, 1987.

- 1000 Prah, F.G., Rontani, J.F., Zabeti, N., Walinsky, S.E. and Sparrow, M.A.: Systematic pattern in U37K'–  
Temperature residuals for surface sediments from high latitude and other oceanographic settings. *Geochimica et  
Cosmochimica Acta*, 74(1), pp.131-143. Doi.org/10.1016/j.gca.2009.09.027, 2010.
- Qin, W., Carlson, L.T., Armbrust, E.V., Devol, A.H., Moffett, J.W., Stahl, D.A. and Ingalls, A.E.: Confounding  
effects of oxygen and temperature on the TEX86 signature of marine Thaumarchaeota. *Proceedings of the  
1005 National Academy of Sciences*, 112(35), pp.10979–10984. Doi.org/10.1073/pnas.1501568112, 2015.
- R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing,  
Vienna, Austria. URL <https://www.R-project.org, 2022>.
- Rayner, N. A.; Parker, D. E.; Horton, E. B.; Folland, C. K.; Alexander, L. V.; Rowell, D. P.; Kent, E. C.; Kaplan,  
A.: Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth  
1010 century. *J. Geophys. Res.*108 (D14), 4407, <https://doi.org/10.1029/2002JD002670>, 2003.
- Ridgway, K.R.: Long-term trend and decadal variability of the southward penetration of the East Australian  
Current. *Geophysical Research Letters*, 34(13). <https://doi.org/10.1029/2007GL030393> , 2007.
- Renoult, M., Annan, J. D., Hargreaves, J. C., Sagoo, N., Flynn, C., Kapsch, M.-L., Li, Q., Lohmann, G.,  
Mikolajewicz, U., Ohgaito, R., Shi, X., Zhang, Q., and Mauritzen, T.: A Bayesian framework for emergent  
1015 constraints: case studies of climate sensitivity with PMIP, *Clim. Past*, 16, 1715–1735, <https://doi.org/10.5194/cp-16-1715-2020>, 2020.
- Rosell-Melé, A. and Prah, F.G.: Seasonality of UK' 37 temperature estimates as inferred from sediment trap data.  
*Quaternary Science Reviews*, 72, pp.128-136. 10.1016/j.quascirev.2013.04.017, 2013.
- Schouten, S., Hopmans, E.C., Schefuß, E. and Damste, J.S.S.: Distributional variations in marine crenarchaeotal  
1020 membrane lipids: a new tool for reconstructing ancient sea water temperatures?. *Earth and Planetary Science  
Letters*, 204(1–2), pp.265–274. 10.1016/S0012-821X(02)00979-2 , 2002.
- Schouten, S., Hopmans, E.C. and Damsté, J.S.S.: The organic geochemistry of glycerol dialkyl glycerol tetraether  
lipids: A review. *Organic geochemistry*, 54, pp.19–61. 10.1016/j.orggeochem.2012.09.006 , 2013.
- Schuddeboom, A.J. and McDonald, A.J.: The Southern Ocean Radiative Bias, Cloud Compensating Errors, and  
1025 Equilibrium Climate Sensitivity in CMIP6 Models. *Journal of Geophysical Research: Atmospheres*, 126(22),  
p.e2021JD035310. <https://doi.org/10.1029/2021JD035310>, 2021.
- Sellar, A.A., Jones, C.G., Mulcahy, J.P., Tang, Y., Yool, A., Wiltshire, A., O'connor, F.M., Stringer, M., Hill, R.,  
Palmieri, J. and Woodward, S.: UKESM1: Description and evaluation of the UK Earth System Model. *Journal of  
Advances in Modeling Earth Systems*, 11(12), pp.4513–4558. <https://doi.org/10.1029/2019MS001739>, 2019.
- 1030 Sen Gupta, A., McGregor, S., Van Sebille, E., Ganachaud, A., Brown, J.N. and Santoso, A.: Future changes to  
the Indonesian Throughflow and Pacific circulation: The differing role of wind and deep circulation  
changes. *Geophysical Research Letters*, 43(4), pp.1669–1678. <https://doi.org/10.1002/2016GL067757> , 2016.
- Sen Gupta, A., Stellingma, A., Pontes, G.M., Taschetto, A.S., Vergés, A. and Rossi, V.: Future changes to the upper  
ocean Western Boundary Currents across two generations of climate models. *Scientific reports*, 11(1), pp.1–12.  
1035 <https://doi.org/10.1038/s41598-021-88934-w> , 2021.
- Senior, C.A., Jones, C.G., Wood, R.A., Sellar, A., Belcher, S., Klein-Tank, A., Sutton, R., Walton, J., Lawrence,  
B., Andrews, T. and Mulcahy, J.P.: UK community Earth system modelling for CMIP6. *Journal of Advances in  
Modeling Earth Systems*, 12(9), p.e2019MS002004. <https://doi.org/10.1029/2019MS002004> , 2020.

- 1040 Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., et al. An assessment of Earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics*, 58. <https://doi.org/10.1029/2019RG000678>, 2020.
- Spezzaferri, S., Kucera, M., Pearson, P.N., Wade, B.S., Rappo, S., Poole, C.R., Morard, R. and Stalder, C.: Fossil and genetic evidence for the polyphyletic nature of the planktonic foraminifera "Globigerinoides", and description of the new genus *Trilobatus*. *PLoS One*, 10(5), p.e0128108. <https://doi.org/10.1371/journal.pone.0259924> , 2015.
- 1045 Smith, R.S., Mathiot, P., Siahhaan, A., Lee, V., Cornford, S.L., Gregory, J.M., Payne, A.J., Jenkins, A., Holland, P.R., Ridley, J.K. and Jones, C.G.: Coupling the UK Earth System Model to dynamic models of the Greenland and Antarctic ice sheets. *Journal of Advances in Modeling Earth Systems*, 13(10), p.e2021MS002520. <https://doi.org/10.1029/2021MS002520> , 2021.
- Strogen, D.P., Seebeck, H., Hines, B.R., Bland, K.J. and Crampton, J.S.: Palaeogeographic evolution of Zealandia: mid-Cretaceous to present. *New Zealand Journal of Geology and Geophysics*, pp.1-30. <https://doi.org/10.1080/00288306.2022.2115520> , 2022.
- 1050 Sutton, P.J. and Bowen, M.: Ocean temperature change around New Zealand over the last 36 years. *New Zealand Journal of Marine and Freshwater Research*, 53(3), pp.305–326. <https://doi.org/10.1080/00288330.2018.1562945>, 2019.
- 1055 Tierney, J.E. and Tingley, M.P.: A TEX86 surface sediment database and extended Bayesian calibration. *Scientific data*, 2(1), pp.1-10. <https://doi.org/10.1038/sdata.2015.29> , 2015.
- Tierney, J. E., and Tingley, M. P.: BAYSPLINE: A new calibration for the alkenone paleothermometer. *Paleoceanography and Paleoclimatology*, 33, 281– 301. <https://doi.org/10.1002/2017PA003201>, 2018.
- Vihtakari M.: ggOceanMaps: Plot Data on Oceanographic Maps using 'ggplot2'. <https://mikkovihtakari.github.io/ggOceanMaps/> , 2022.
- 1060 Volodin, Evgenii M., Mortikov, Evgeny V., Kostykin, Sergey V., Galin, Vener Ya., Lykossov, Vasily N., Gritsun, Andrey S., Diansky, Nikolay A., Gusev, Anatoly V., Iakovlev, Nikolay G., Shestakova, Anna A. and Emelina, Svetlana V.: Simulation of the modern climate using the INM-CM48 climate model. *Russian Journal of Numerical Analysis and Mathematical Modelling*, 33, 6, pp. 367-374. <https://doi.org/10.1515/rnam-2018-0032>, 2018.
- 1065 Williams, J., Morgenstern, O., Varma, V., Behrens, E., Hayek, W., Oliver, H., Dean, S., Mullan, B. and Frame, D.: Development of the New Zealand Earth System Model. *Weather and Climate*, 36, pp.25–44. 2016.
- Williams, J., Behrens, E., Morgenstern, O., Teixeira, J.C., Varma, V., and Hayek, W.: Regional ocean grid refinement and its effect on simulated atmospheric climate. ESS Open Archive . Submitted to Weather and Climate. DOI: 10.22541/essoar.167642236.61101960/v1, 2023.
- 1070 Zelinka, M.D., Myers, T.A., McCoy, D.T., Po-Chedley, S., Caldwell, P.M., Ceppi, P., Klein, S.A. and Taylor, K.E.: Causes of higher climate sensitivity in CMIP6 models. *Geophysical Research Letters*, 47(1), p.e2019GL085782. doi:10.1029/2019GL085782, 2020.
- Zhang, Y.G., Zhang, C.L., Liu, X.L., Li, L., Hinrichs, K.U. and Noakes, J.E.: Methane Index: A tetraether archaeal lipid biomarker indicator for detecting the instability of marine gas hydrates. *Earth and Planetary Science Letters*, 307(3–4), pp.525–534. DOI: 10.1016/j.epsl.2011.05.031, 2011.
- 1075 Zhu, J., Otto-Bliesner, B.L., Brady, E.C., Poulsen, C.J., Tierney, J.E., Lofverstrom, M. and DiNezio, P.: Assessment of equilibrium climate sensitivity of the Community Earth System Model version 2 through

simulation of the Last Glacial Maximum. *Geophysical Research Letters*, 48(3), p.e2020GL091220.  
1080 <https://doi.org/10.1029/2020GL091220>, 2021.