

# Cloud properties and their projected changes in CMIP models with low/medium/high climate sensitivity

Lisa Bock<sup>1</sup> and Axel Lauer<sup>1</sup>

<sup>1</sup>Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

**Correspondence:** L. Bock (lisa.bock@dlr.de)

**Abstract.** Since the release of the first CMIP6 simulations one of the most discussed topics is the higher effective climate sensitivity (ECS) of some of the models resulting in an increased range of ECS values in CMIP6 compared to previous CMIP phases. An important contribution to ECS is the cloud climate feedback. Although climate models have continuously been developed and improved over the last decades, a realistic representation of clouds remains challenging. Clouds contribute to the large uncertainties in modeled ECS, as projected changes in cloud properties and cloud feedbacks also depend on the simulated present-day fields.

In this study we investigate the representation of both cloud physical and radiative properties from a total of 51 CMIP5 and CMIP6 models. ECS is used as a simple metric to group the models as the sensitivity of the physical cloud properties to warming is closely related to cloud feedbacks, which in turn are known to have a large contribution to ECS. Projected changes of cloud properties in future scenario simulations are analyzed by ECS group. In order to help interpreting the projected changes, model results from historical simulations are also analyzed.

The results show that differences in the net cloud radiative effect as a reaction to warming among the three model groups are driven by changes in a range of cloud regimes rather than individual regions. In polar regions, high ECS models show a weaker increase in the net cooling effect of clouds due to warming than the low ECS models. At the same time, high ECS models show a decrease in the net cooling effect of clouds over the tropical ocean and the subtropical stratocumulus regions whereas low ECS models show either little change or even an increase in the cooling effect. Over the Southern Ocean, the low ECS models show a higher sensitivity of the net cloud radiative effect to warming than the high ECS models.

## 1 Introduction

Climate models are an essential tool for projecting future climate. Within the context of the Coupled Model Intercomparison Project (CMIP, <https://www.wcrp-climate.org/wgcm-cmip>), a World Climate Research Programme (WCRP) initiative, several modeling groups worldwide provide a set of coordinated simulations with different Earth system models (ESMs) of the past (historical) time period and different future scenarios. The main objective of CMIP is to better understand past, present, and future climate, its variability and future change arising from both natural, unforced variability and in response to changes in radiative forcing in a multi-model context.

25 Across the different CMIP phases, several improvements e.g. in the climatological large-scale patterns of temperature, water  
vapor, and zonal wind speed were found with the latest phase models (CMIP6, Eyring et al. (2016)) typically performing  
slightly better than their CMIP3 and CMIP5 predecessors when compared to observations (Bock et al., 2020). While this is  
also the case for some cloud properties and selected regions such as the Southern Ocean, clouds remain challenging for global  
climate models with many known biases remaining in CMIP6 (Lauer et al., 2023). As such, clouds continue to play a significant  
30 role in uncertainties of climate models and climate projections (Bony et al., 2015).

One of the extensively discussed topics in analyses of the CMIP6 ensemble is the higher effective climate sensitivity (ECS)  
of some models and therefore the increased range in ECS now between 1.8 and 5.6 K compared to 2.1 and 4.7 in the CMIP5  
ensemble (Meehl et al., 2020; Bock et al., 2020; Schlund et al., 2020). ECS provides a single number, defined as the change  
in global mean near-surface air temperature resulting from a doubling of the atmospheric CO<sub>2</sub> concentration compared to  
35 preindustrial conditions, once the climate has reached a new equilibrium (Gregory et al., 2004). A possible reason for the  
increase of ECS in some models is improvements in cloud representation in these models. Zelinka et al. (2020) show that the  
increased range of ECS in the CMIP6 models could be explained by an increased range in cloud feedbacks. Studies using single  
models concluded that the increased climate sensitivity found in these models is largely determined by cloud microphysical  
processes (Zhu et al., 2022; Frey and Kay, 2018; Gettelman et al., 2019; Bodas-Salcedo et al., 2019). They also point out that  
40 the simulated present-day mean state of cloud properties is related to the simulated cloud feedback but could also be connected  
to other coupled feedbacks (Andrews et al., 2019).

As future changes in cloud properties are closely connected to cloud feedbacks and cloud feedbacks are known to be strongly  
correlated with ECS (see Sect. 3.1), we use ECS as a simple proxy to group the ensemble of CMIP5 and CMIP6 models for  
this analysis. This facilitates the analysis and allows for obtaining more general conclusions beyond individual models that  
45 can vary widely in their sensitivity to the prescribed forcings. A particular focus of this study is whether there are systematic  
differences in cloud-related quantities between the different ECS groups. The sensitivity of the physical properties to warming  
is analyzed, as this gives some insight into the uncertainty of the projected cloud properties and their potential contribution to  
cloud feedbacks and ECS.

A comparison of the present-day climatologies of key cloud properties from the different ECS model groups is done to help  
50 interpreting simulated future changes in cloud properties. The present-day performance of CMIP models has been investigated  
for example by Kuma et al. (2023), who applied an artificial neural network to derive cloud types from radiation fields. They  
found that results from models with a high ECS agree on average better with observations than from models with a low ECS.  
Jiang et al. (2021) found that the models' ECS is positively correlated with the integrated cloud water content and water  
vapor performance scores for both CMIP6 and CMIP5 models. In contrast, Brunner et al. (2020) showed that some CMIP6  
55 models with a high future warming compared to other models receive systematically lower performance weights when using  
anomaly, variance, and trend of surface air temperature, and anomaly and variance of sea level pressure to assess the models'  
performance.

A number of different feedbacks are relevant to ECS with cloud feedbacks being an important contribution. In order to  
assess whether there are systematic differences in simulated cloud properties among models with different ECS, we compare

60 the simulated cloud properties from three groups of models sorted by their ECS values and quantify how the projected changes  
in cloud properties and cloud radiative effects differ. In Section 2 we briefly introduce the models and observations used as  
well as the software tool applied to compare the models. The representation of cloud radiative effects for all three groups is  
also compared with observational data in Section 3 followed by an analysis of the projected future changes in cloud properties  
and radiative effects. Section 4 summarizes the discussion and conclusions.

## 65 **2 Data**

### **2.1 Models**

In this study we use model simulations from the CMIP Phases 5 (Taylor et al., 2012) and 6 (Eyring et al., 2016). The individual  
models are detailed in Table 1. All model data are freely available via the Earth System Grid Federation (ESGF), which is an  
international collaboration that manages the decentralized database of CMIP output.

70 For the analysis presented here, we use historical simulations over the time period 1985–2004 (Table 1) and the scenario  
simulations of the Representative Concentration Pathway (RCP) 8.5 from CMIP5 and the Shared Socioeconomic Pathways  
(SSP) 5-8.5 simulations from CMIP6 for the years 2081-2100. The historical simulations use prescribed natural and anthro-  
pogenic climate forcings such as concentrations of greenhouse gases and aerosols. We only consider one ensemble member  
per model, typically the first member "r1i1p1" (CMIP5) and "r1i1p1f1" (CMIP6). As the intermodel spread is typically much  
75 larger than the interensemble spread we do not expect our results to change significantly when using more ensemble members  
for each model. For further details on the model simulations, we refer to Taylor et al. (2012) and Eyring et al. (2016).

ECS and cloud feedbacks are calculated using the simulations forced by an abrupt quadrupling of CO<sub>2</sub> (abrupt-4×CO<sub>2</sub>) and  
the preindustrial control simulations (piControl) following the method described in Andrews et al. (2012) and Schlund et al.  
(2020).

80 In total, the CMIP ensemble investigated here consists of 24 CMIP5 and 27 CMIP6 models that provide the output needed  
for this analysis. We grouped them into the three groups "low", "medium" and "high" by their ECS values (see Table 1). The  
thresholds for the three groups are chosen in a way that each of the three groups contains the same number of models. Multi-  
model group means are calculated as 20-year means over all models in the high, medium and low ECS group applying equal  
weights to each model.

### 85 **2.2 Observations**

The Clouds and Earth's Radiant Energy Systems (CERES) Energy Balanced and Filled (EBAF) Ed4.2 dataset (Loeb et al.,  
2018; Kato et al., 2018) provides global monthly mean top of atmosphere (TOA) longwave (LW), shortwave (SW), and net  
radiative fluxes under clear-sky and all-sky conditions, which are used as a reference dataset to calculate the cloud radiative  
effects and the TOA outgoing radiation. CERES instruments are flown on NASA's Terra and Aqua satellites.

**Table 1.** List of CMIP5 and CMIP6 models grouped by ECS value into three roughly equally sized groups "high" (ECS > 4.0 K), "medium" (2.87 K < ECS < 4.0 K) and "low" (ECS < 2.87 K).

Number	CMIP5 model	CMIP6 model	ECS (K)	Citation
1		CanESM5	5.62	Swart et al. (2019)
2		HadGEM3-GC31-LL	5.55	Williams et al. (2018); Kuhlbrodt et al. (2018)
3		HadGEM3-GC31-MM	5.42	Williams et al. (2018); Kuhlbrodt et al. (2018)
4		UKESM1-0-LL	5.34	Sellar et al. (2019)
5		CESM2	5.16	Danabasoglu et al. (2020)
6		CNRM-CM6-1	4.83	Volodine et al. (2019)
7		KACE-1-0-G	4.77	Lee et al. (2020a)
8		CNRM-ESM2-1	4.76	S��ferian et al. (2019)
9		CESM2-WACCM	4.75	Danabasoglu et al. (2020)
10		NESM3	4.72	Cao et al. (2018)
11	MIROC-ESM		4.67	Watanabe et al. (2011)
12	HadGEM2-ES		4.61	Collins et al. (2011)
13		IPSL-CM6A-LR	4.56	Boucher et al. (2020)
14		TaiESM1	4.31	Lee et al. (2020b)
15	IPSL-CM5A-LR		4.13	Dufresne et al. (2013)
16	IPSL-CM5A-MR		4.12	Dufresne et al. (2013)
17	CSIRO-Mk3-6-0		4.08	Rotstayn et al. (2010)
1	GFDL-CM3		3.97	Donner et al. (2011)
2	BNU-ESM		3.92	Ji et al. (2014)
3	ACCESS1-0		3.83	Bi et al. (2013)
4	CanESM2		3.69	Arora et al. (2011)
5	MPI-ESM-LR		3.63	Giorgetta et al. (2013); Stevens et al. (2013)
6		CMCC-ESM2	3.58	Cherchi et al. (2019)
7	ACCESS1-3		3.53	Bi et al. (2013)
8		CMCC-CM2-SR5	3.52	Cherchi et al. (2019)
9	MPI-ESM-MR		3.46	Giorgetta et al. (2013); Stevens et al. (2013)
10	FGOALS-g2		3.38	Li et al. (2013)
11		MRI-ESM2-0	3.15	Yukimoto et al. (2019); Mizuta et al. (2012)
12		GISS-E2-1-H	3.11	Kelley et al. (2020)
13		BCC-CSM2-MR	3.04	Wu et al. (2019)
14		FGOALS-f3-L	3.00	He et al. (2020)
15		MPI-ESM1-2-LR	3.00	Mauritsen et al. (2019)
16		MPI-ESM1-2-HR	2.98	Muller et al. (2018)
17	CCSM4		2.94	Gent et al. (2011)
18		FGOALS-g3	2.88	Li et al. (2020)
1	bcc-csm1-1-m		2.86	Wu et al. (2010); Wu (2012)
2	bcc-csm1-1		2.83	Wu et al. (2010); Wu (2012)
3	NorESM1-M		2.80	Bentsen et al. (2013)
4		GISS-E2-1-G	2.72	Kelley et al. (2020)
5	MIROC5		2.72	Watanabe et al. (2010)
6		MIROC-ES2L	2.68	Hajima et al. (2020)
7		MIROC6	2.61	Tatebe et al. (2019)
8	IPSL-CM5B-LR		2.60	Hourdin et al. (2013)
9	MRI-CGCM3		2.60	Yukimoto et al. (2012)
10		NorESM2-LM	2.54	Seland et al. (2020)
11		NorESM2-MM	2.50	Seland et al. (2020)
12	GFDL-ESM2M		2.44	Donner et al. (2011)
13	GFDL-ESM2G		2.39	Donner et al. (2011)
14	GISS-E2-H		2.31	Schmidt et al. (2006)
15		CAMS-CSM1-0	3.29	Rong et al. (2018)
16	GISS-E2-R		2.11	Schmidt et al. (2006)
17	inmcm4		2.08	Volodin et al. (2010)

90 The dataset covers the whole years of the time period 2001-2022. We would like to note that the time period from the models used for comparison with the CERES-EBAF dataset (Sect. 2.1) does not match exactly the observed years. Given that ESMs are not expected to reproduce the exact observed phase of climate modes, which largely control present-day variability of clouds but rather their statistical properties, it is probably not surprising, however, that this difference in the time periods has very little impact on the multi-year group averages when comparing multi-year climatologies of cloud parameters.

95 Starting with CERES-EBAF Ed4.1, the dataset provides adjusted clear-sky fluxes which are now defined in a manner that is more in line with how clear-sky fluxes are represented in climate models. The uncertainty estimates for  $1^\circ \times 1^\circ$  regional monthly net cloud radiative effects are about  $7 \text{ W m}^{-2}$  (CERES-EBAF, 2021).

## 2.3 ESMValTool


100 All analyses in this study are performed with the open-source community diagnostics and performance metrics tool for evaluation of ESMs “Earth System Model Evaluation Tool” (ESMValTool) version 2 (Eyring et al., 2020; Lauer et al., 2020; Righi et al., 2020; Weigel et al., 2021; Schlund et al., 2023). All figures from this paper can be reproduced by running the ESMValTool "recipe" (configuration script defining all datasets, processing steps and diagnostics) recipe\_bock23acp.yml (see also the Code and Data Availability Section).

## 3 Analysis

### 105 3.1 ECS and cloud feedback

The large spread in ECS of CMIP6 models could be mainly explained by uncertainties in the simulated net cloud feedback. The net cloud feedback is defined as change in the sum of shortwave and longwave cloud radiative effects at the top of the atmosphere (TOA) per degree of surface warming (2-m temperature) calculated as the difference between abrupt-4xCO<sub>2</sub> simulations and the corresponding piControl simulations. The TOA shortwave and longwave cloud radiative effects are calculated as the differences between the respective TOA all-sky and clear-sky radiative fluxes. While this method is commonly used to calculate cloud feedbacks, we would like to note that the results using this method are not exactly the same as those calculated with a more accurate offline radiative transfer method (Soden et al., 2004). Particularly the shortwave CRE can be affected in regions with high surface albedos such as polar latitudes if the surface albedo changes between the two model simulations e.g. because of melting sea ice (Shell et al., 2008). The net cloud feedback is typically dominated by the shortwave component (Zelinka et al., 2020).

115 The relation between ECS and simulated cloud feedbacks is illustrated in Figure 1, which shows the correlation between net, shortwave and longwave cloud feedbacks and ECS in the CMIP5 and CMIP6 models (Table 1). The relation between net cloud feedback and ECS is dominated by the shortwave cloud feedback, which shows a strong correlation with ECS ( $r = 0.66$  and a small p value of  $p = 3.6e-9$ ). For the longwave cloud feedback there is only a weak (negative) correlation with ECS ( $p =$   
120  $0.05$ ).




**Figure 1.** Scatterplot of the global mean a) net, b) shortwave and c) longwave cloud feedback (x-axis) and ECS (y-axis) of the CMIP models (Table 1) with regression line including the confidence interval of the regression of 95%. Horizontal dashed lines indicate separations of the three ECS groups (see Table 1).

As the representation of clouds and their sensitivity to climate change have a strong impact on the ECS (Zelinka et al., 2020; Bjordal et al., 2020; Bony et al., 2015) and because the range of ECS obtained from the ensemble of CMIP6 models is larger than the one from the previous model generations (Meehl et al., 2020), this motivated us to look into the differences in present-day performance and future projections of physical cloud parameters from models with low/medium/high ECS.

125 Figure 2 shows the geographical distributions of the net, shortwave and longwave cloud feedbacks averaged over all models within each group. The pattern of the net cloud feedback is dominated by the geographical distribution of the shortwave cloud feedback. On global average, the high ECS group has the largest net cloud feedback of  $0.41 \text{ W m}^{-2}$ , followed by the medium ECS group ( $0.01 \text{ W m}^{-2}$ ) and the low ECS group ( $-0.20 \text{ W m}^{-2}$ ). The group mean net cloud feedback changes sign at around  $60^\circ\text{S}$  and  $80^\circ\text{N}$  in all three groups. The sign change at around  $60^\circ\text{S}$  in the shortwave cloud feedback corresponds to the latitude region where a change from clouds with an ice component in the piControl simulations to clouds consisting almost entirely of liquid droplets in the abrupt-4xCO<sub>2</sub> experiment (cloud phase feedback) starts to contribute significantly to the total cloud feedback (Ceppi et al., 2017). With increasing latitude there is an increasing ice fraction in the model clouds that supports a negative shortwave feedback as cloud particles can change phase with warming. Particularly over the Arctic and the tropical Pacific, the (negative) shortwave cloud feedback is partly or fully compensated by a (positive) longwave cloud feedback resulting in rather small net cloud feedback values.


135 The high ECS models show a more positive net cloud feedback in the Tropics and midlatitudes, especially over the Southern Ocean, than the other two groups. The group mean of the low ECS models shows a distinct negative net cloud feedback in the Tropics, particularly in the tropical Pacific. This signal is much weaker in the other two groups. The reason is a more pronounced negative shortwave cloud feedback particularly over the Pacific Intertropical Convergence Zone (ITCZ) and South Pacific Convergence Zone (SPCZ) in the group mean of the low ECS models.



**Figure 2.** Geographical maps of net (a,b,c), shortwave (d,e,f) and longwave (g,h,i) cloud feedback for high (left), medium (middle) and low (right) ECS groups.

### 3.2 Present-day cloud fields

The representation of cloud properties in ESMs is related to the simulated cloud feedback (Zelinka et al., 2020, 2022). In order to help interpreting differences in simulated future changes in cloud properties due to warming among the three ECS groups, we compare the geographical distributions of the climatologies from the individual groups with each other. Here, we focus on the most climate-relevant parameters, which are total cloud fraction, liquid water and ice water path (Figure 3) and longwave, shortwave and net cloud radiative effects (Figure 4). For a direct comparison with satellite observations, output of cloud-related quantities from satellite simulators would be needed. Such output is, however, only available in very limited form or not at all from the models. Comparisons of the model results with observations are therefore restricted to cloud radiative effects, for



**Figure 3.** Geographical map of the multi-year annual mean total cloud fraction (a-c), ice water path (d-f) and liquid water path (g-i) for group means of historical CMIP simulations from all three ECS groups.

which data are available that could be compared directly with climate model results without using satellite simulators or other  
 150 sampling strategies (Loeb and Staff, 2022, see Section 2.2 for more details on CERES-EBAF).

### Total cloud fraction

The annual mean total cloud fraction from all ECS groups (Figure 3a-c) shows the known geographical patterns: maxima  
 155 over land in the Tropics due to strong convection, minima in the subtropics because of descending air with local maxima in  
 stratocumulus regions off the west coasts of the continents (Africa, North and South America), maxima in the midlatitudes  
 over the ocean especially over the Southern Ocean and minima over polar regions where the air is very cold and dry.



**Table 2.** Mean values of each group together with the 25% and 75% quantiles (in parentheses) calculated by bootstrapping (1000 times, sample size = number of models in the group). The second line gives the 25% and 75% quantiles calculated from all individual models.

Variable	Mean		
	high ECS	med ECS	low ECS
Total Cloud Fraction (%)	64.1 (63.3, 65.0) (61.9, 68.8)	59.8 (58.9, 60.0) (56.7, 62.5)	59.5 (59.0, 59.8) (57.8, 61.9)
Ice Water Path ( $\text{g m}^{-2}$ )	37.0 (34.3, 40.1) (19.1, 51.9)	34.6 (30.3, 38.6) (17.6, 40.6)	40.7 (35.5, 45.2) (14.9, 42.3)
Liquid Water Path ( $\text{g m}^{-2}$ )	65.0 (61.0, 68.5) (55.3, 68.4)	72.1 (67.1, 76.8) (54.6, 86.1)	83.2 (78.5, 87.9) (60.4, 105.5)
Net Cloud Radiative Effect ( $\text{W m}^{-2}$ )	-22.8 (-23.4, -22.3) (-24.7, -20.7)	-23.2 (-23.6, -22.8) (-25.0, -21.9)	-25.8 (-26.3, -25.3) (-28.2, -23.6)

The group mean of the high ECS models (Figure 3a) shows the largest global mean of 64.1% in total cloud cover compared to 59.8% and 59.5% for the medium and low ECS groups, respectively. Compared with the inter-model spread given by the quantiles (see Table 2), this difference is a robust signal. Especially the maxima in total cloud cover over the Southern Ocean and the northern Atlantic and Pacific (Figure 3a) are more pronounced (differences up to 10%) in the group mean of the high ECS group, which leads to a slight reduction in the known bias of total cloud cover from CMIP models compared with the CMIP5/6 multi-model mean (Lauer et al., 2023). The minima polewards of about  $75^\circ$  are more pronounced in the low and medium ECS models (Figure 3b,c).

### Ice water path

The global distribution of ice water path (Figure 3d-f) from all ECS group means shows the maximum in the ITCZ due to frequent convection of up to  $0.16 \text{ kg m}^{-2}$ . The absolute minima of ice water path are found in the subtropics in the subsidence regions west of continents. High amounts of cloud ice are also found along the stormtracks in midlatitudes, with values decreasing towards the poles.

The inter-model spread in the global mean ice water path (Table 2) is large resulting in a large overlap between the different ECS groups and no statistically significant difference in the global mean ice water path from the three ECS groups. There are, however, some differences in regional features of the ice water path distribution. The maximum of the ice water path values in the Tropics related to the ITCZ are highest in the low ECS group. In contrast, the maxima in midlatitudes, especially over the Southern Ocean, are most pronounced in the high ECS models (Figure 3d).

**Table 3.** Root mean square difference (RMSD) and pattern correlation of each group mean together with the 25% and 75% quantiles (in parentheses) calculated by bootstrapping (1000 times, sample size = number of models in the group). The second line gives the 25% and 75% quantiles calculated from all individual models. The RMSD values and the correlation are calculated in comparison to the corresponding reference dataset CERES-EBAF (see Section 2.2).

Variable	RMSD ( $\text{W m}^{-2}$ )			Correlation		
	high ECS	med ECS	low ECS	high ECS	med ECS	low ECS
Net Cloud Radiative Effect	9.3 (9.0, 9.9) (9.5, 14.1)	9.2 (9.0, 9.7) (10.5, 13.0)	12.3 (11.7, 13.0) (11.9, 18.3)	0.86 (0.84, 0.86) (0.74, 0.84)	0.86 (0.85, 0.87) (0.70, 0.82)	0.79 (0.75, 0.80) (0.55, 0.77)
Shortwave Cloud Radiative Effect	8.4 (8.2, 9.1) (9.2, 13.2)	7.8 (7.8, 8.4) (10.6, 12.9)	11.1 (10.7, 12.0) (12.0, 18.5)	0.92 (0.90, 0.92) (0.79, 0.91)	0.93 (0.91, 0.93) (0.81, 0.87)	0.87 (0.85, 0.88) (0.73, 0.83)
Longwave Cloud Radiative Effect	5.7 (5.6, 6.1) (7.0, 8.6)	6.1 (6.0, 6.4) (7.1, 8.3)	5.9 (5.7, 6.4) (6.3, 9.5)	0.92 (0.91, 0.92) (0.82, 0.87)	0.93 (0.92, 0.93) (0.83, 0.88)	0.92 (0.91, 0.92) (0.78, 0.89)

### Liquid water path

The ECS group means of cloud liquid water path (Figure 3g-i) show local maxima in the ITCZ but the largest values of liquid water path are found in the extratropics in the stormtrack regions, mainly over the Southern Ocean and the northern Atlantic. There are no local maxima in the stratocumulus regions seen in all three group means which is related to a known bias of underestimating stratocumulus clouds in the CMIP models (e.g., Jian et al., 2020). In addition to the higher cloud cover in the high ECS group in these regions the clouds seem to be less bright in comparison the two other groups. This indicates an improvement of the representation of stratocumulus clouds in the high ECS group, which is consistent with the findings of Cesana et al. (2023).


The high ECS group mean shows the lowest global mean ( $65.0 \text{ g m}^{-2}$ ) followed by the medium ECS group and the low ECS group with  $72.1 \text{ g m}^{-2}$  and  $83.2 \text{ g m}^{-2}$ , respectively. This negative correlation between the group-averaged global mean liquid water path and ECS seems quite robust regarding the relatively small inter-model spread in these two variables within each group. The relative differences between the ECS groups are uniformly distributed with no region being particularly pronounced.

### Cloud radiative effects

The cloud radiative effects are calculated as the differences in top of the atmosphere clear-sky and all-sky radiative fluxes (for details see Section 3.1). The CERES-EBAF data show a global mean net cooling due to clouds of about  $-18 \text{ W m}^{-2}$  (Figure 4a). Clouds have a warming radiative effect in particular over regions with a high surface albedo like ice-covered regions in Greenland and Antarctica and the desert regions in North Africa. A large negative (cooling) net radiative effect of clouds is found over the stratocumulus regions in the subtropics and in the midlatitude stormtrack regions. In the ITCZ there is a partly compensating effect between the shortwave and longwave radiative effects leading to smaller absolute net values than in the stratocumulus and stormtrack regions.

Compared with CERES-EBAF, the amplitude of the global mean net cloud radiative effect is slightly overestimated in the models with the largest bias in the low ECS group (mean bias =  $-4.8 \text{ W m}^{-2}$ , RMSD =  $12.3 \text{ W m}^{-2}$ ) and the smallest bias in

## TOA Net Cloud Radiative Effect



**Figure 4.** Geographical map of the multi-year annual mean net cloud radiative effect from (a) CERES-EBAF Ed4.2 (OBS) and (b,c,d) group means of historical CMIP simulations from all three ECS groups.

195 the high ECS group (mean bias =  $-1.8 \text{ W m}^{-2}$ , RMSD =  $9.3 \text{ W m}^{-2}$ ) (see also Table 3). While the global mean biases of the group means are within the observational uncertainty range, the RMSD values are larger than the ones of different individual observational datasets when compared to a reference dataset consisting of an average over different products (Lauer et al., 2023).

Biases in simulated sea surface temperatures (SSTs) can affect simulated cloud properties. We therefore also analyzed some  
 200 results from AMIP simulations that use the atmosphere components of the CMIP models and for which SSTs and sea ice concentrations from observations are prescribed (not shown). Similar to previous studies (e.g., Lauer and Hamilton, 2013; Lauer et al., 2023), we found rather little differences in the multi-year climatologies of cloud properties from the models between the historical and AMIP runs analyzed here. For the net cloud radiative effect, differences between the annual mean climatologies from the historical and AMIP simulations are below  $5 \text{ W m}^{-2}$  throughout most of the globe but differences in

205 the ITCZ and tropical Atlantic can reach up to about  $10 \text{ W m}^{-2}$ . Globally averaged, the mean bias for the three group averages ranges between 0.3 and  $0.6 \text{ W m}^{-2}$ , RMSE between 2.7 and  $3.3 \text{ W m}^{-2}$  and pattern correlations between 0.97 and 0.98.

The geographical patterns of the three model groups agree well with the CERES-EBAF observations (Figure 4). The linear pattern correlations of the annual average net cloud radiative effect from the high and the medium ECS group means with observations are slightly higher (0.86) than that of the low (0.79) ECS group. This is also reflected in the range of correlation  
210 values from the individual models in each group given by the 25% and 75% quantiles. These range lies between 0.55 and 0.77 in the low ECS group, between 0.70 and 0.82 in the medium ECS group and between 0.74 and 0.84 in the high ECS group. For comparison, the range of correlation coefficients of different observational datasets is 0.98-0.99 (Lauer et al., 2023). The correlation values of the shortwave and longwave cloud radiative effects are larger for all ECS groups.

The peaks of positive cloud forcing over land over Greenland, North Africa and the west coast of North and South America  
215 are underestimated in all three groups. In these regions, however, observational uncertainties are expected to be large because of high surface albedo, topography or very low cloud cover. The largest positive bias for all groups is found over the stratocumulus regions with up to  $46 \text{ W m}^{-2}$  locally. Apart from this, the low ECS group shows particularly between  $30^\circ\text{S}$  and  $30^\circ\text{N}$  (Figure 4d), a too strong net cloud radiative effect resulting mainly from a too strong shortwave cooling of the clouds in this latitude belt (Figure 6e) seemingly caused by the largest cloud water path values of all three ECS groups (Figure 6b,c).

### 220 3.3 Differences in projected future cloud properties


In order to investigate the sensitivity of cloud parameters simulated by the three ECS groups to future warming, we compare the changes in selected cloud properties and cloud radiative effects in future simulations from each group. For CMIP6 we calculate the changes as differences between data from SSP5-8.5 and for CMIP5 from RCP8.5 and results for the respective historical simulations.

225 The zonally averaged group means (Figure 6a-f, upper panels) show the results from the historical and the scenario simulations for the investigated cloud properties (total cloud fraction, ice and liquid water path and cloud radiative effects) for the different ECS groups. Projected zonal mean changes per degree warming (near-surface temperature increase) are displayed in the panels below (Figure 6a-f, lower panels). Additionally, we show the sensitivity of cloud parameters from each ECS group over the ocean for selected regions. The relative changes (calculated as the differences between the scenario value and the  
230 historical value divided by the historical value) in cloud parameters per degree warming averaged over selected regions (Figure 5) are shown in Figure 7: 1) Arctic, 2) Southern Ocean, 3) tropical ocean, 4) Pacific ITCZ and the stratocumulus regions 5a) South East Pacific, 5b) South East Atlantic and 5c) North East Pacific.

In the following, we discuss the differences in projected future cloud properties for each cloud parameter.

#### **Total cloud cover**

235 For zonal mean cloud cover (Figure 6a), the comparison of the historical runs with the scenario simulations shows an increase in the zonal mean cloud cover in particular over the polar regions north and south of about  $70^\circ$ . This positive sensitivity to




**Figure 5.** Maps of selected regions: 1) Arctic (70-90°N), 2) Southern Ocean (30-65°S), 3) Tropical Ocean (30°N-30°S), 4) Pacific ITCZ (0-12°N, 135°O-85°W) and the three stratocumulus regions 5a) South East Pacific (10-30°S, 75-95°W), 5b) South East Atlantic (10-30°S, 10°W-10°O) and 5c) North East Pacific (15-35°N, 120-140°W).

warming shows maximum values ranging between about  $0.5\% \text{ K}^{-1}$  for the high, about  $1\% \text{ K}^{-1}$  for the medium and  $1.4\% \text{ K}^{-1}$  for the low ECS groups.


Particularly in the Tropics and in SH mid- and high latitudes, the sensitivity of simulated cloud cover to warming is quite different among the high ECS group and the two other groups. While the low and medium ECS groups show a mostly positive sensitivity in the Tropics, the high ECS group shows a negative sensitivity of cloud cover to warming of about 0.5 to  $-1.5\% \text{ K}^{-1}$ . Averaged over the tropical ocean (Figure 7c), the behavior of the high ECS models is significantly different than that of the two other groups. All high ECS models show a decrease in total cloud cover over the tropical ocean while the individual models in the two other groups do not agree on the sign of the change.

In all three subtropical stratocumulus regions investigated (North East Pacific, South East Pacific and South East Atlantic), the high ECS group shows a decrease in total cloud cover (Figure 7e). In contrast, the low and medium ECS groups show particularly in the Southern Hemisphere stratocumulus regions an increase in total cloud cover that is most pronounced in the low ECS group.

In general, there is a decrease in cloud fraction in midlatitudes which is most pronounced in the high ECS group and becomes weaker towards the poles. In SH mid- and high latitudes south of  $45^\circ\text{S}$ , the low ECS group shows a strong positive sensitivity of up to more than  $1\% \text{ K}^{-1}$  while the high ECS group shows a negative sensitivity of about  $-1\% \text{ K}^{-1}$  at  $45^\circ\text{S}$ . South of  $55^\circ\text{S}$ , the high ECS group also shows a positive sensitivity of total cloud cover. The medium ECS group lies in between the low and high ECS groups but is in general closer to the low ECS group. Averaged over the Southern Ocean (latitude belt  $30\text{-}65^\circ\text{S}$ ), the high ECS models mostly show a negative sensitivity while the individual models in the two other groups show positive and negative sensitivities.



**Figure 6.** Upper panels: zonally averaged group means of (a) total cloud fraction, (b) liquid water path, (c) ice water path and (d) net, (e) shortwave and (f) longwave cloud radiative effect from historical simulations (solid lines) and RCP8.5 / SSP5-8.5 scenarios (dashed lines) for the three different ECS groups. The reference dataset CERES-EBAF Ed4.2 is shown as solid black lines in panels d-f. Lower panels: corresponding relative differences of all zonally averaged group means between the RCP8.5 / SSP5-8.5 scenarios and the corresponding historical simulations. Shading indicates the 5% and 95% quantiles of the single model results.



**Figure 7.** Relative change (calculated as the difference between the scenario value and the historical value divided by the historical value) of total cloud fraction (clt), ice water path (iwp), liquid water path (lwp) and net cloud radiative effect (netcre) per degree warming averaged over selected regions over the ocean: (a) Arctic (70-90°N), (b) Southern Ocean (30-65°S), (c) tropical ocean (30°N-30°S), (d) Pacific ITCZ (0-12°N, 135°O-85°W) and e) the three stratocumulus regions South East Pacific (10-30°S, 75-95°W), South East Atlantic (10-30°S, 10°W-10°O) and North East Pacific (15-35°N, 120-140°W) (see also Figure 5). In the box plot, each box indicates the range from the first quartile to the third quartile, the vertical line shows the median and the whiskers the minimum and maximum values excluding the outliers. Outliers are defined as being outside 1.5 times the interquartile range.

## Cloud liquid and ice water path

In the Tropics between about 10°S and 10°N, the cloud ice water path shows a strong sensitivity to warming of up to 9% K<sup>-1</sup> and 10% K<sup>-1</sup> in all three ECS groups (Figure 6b). The zonally averaged ice water path increases also in all three groups north and south of about 60°N/S with the high ECS group showing the strongest sensitivity to warming. Particularly in the Arctic  
260 north of 80°N, the sensitivity of the simulated ice water path to warming is about twice as high in the high ECS group ( 4% K<sup>-1</sup>) than in the medium and low ECS groups ( 2% K<sup>-1</sup>). In midlatitudes, all groups show a negative sensitivity to warming with the high ECS group typically showing the strongest sensitivity in the Northern Hemisphere among the three ECS groups.

Similarly to the ice water path, also the zonally averaged liquid water path increases with temperature in all three groups in the polar regions (Figure 6c). This is consistent with the findings of Lelli et al. (2023) who report an observed trend to brighter  
265 and more liquid clouds in satellite measurements over the Arctic. In contrast to the ice water path, the lowest ECS group shows the highest sensitivity in the Arctic latitude belt. Averaged over the whole Arctic, however, there are no significant differences in ice and liquid water path over the ocean between the different ECS groups (Figure 7a).

The amplitude of the decrease in ice water path per degree warming is peaking at about 35°S and N and is about twice as large in the Southern Hemisphere than in the Northern Hemisphere. Beyond about 60°N and S, there is an increase in ice  
270 water path that is getting more pronounced towards the poles. This increase in ice water path with warming is even stronger for the liquid water path with no significant differences between the ECS groups. This increase in liquid water path can be partly explained by a phase change from cloud ice to liquid at higher temperatures.

In the stratocumulus regions (Figure 7e), liquid water path increases in the low ECS model group while it decreases in the high ECS group. The medium ECS group lies in between the two with many of the individual models disagreeing on the sign  
275 of the change. This behavior is consistent with the sensitivity of the changes in total cloud cover in these regions. We would like to note that ice water path values are typically very small in the stratocumulus regions. Relative changes can therefore be large without being physically relevant.

Over the Southern Ocean, the decrease in ice water path and the increase in liquid water path with warming is also not statistically significantly different among the three ECS groups. Averaged over the whole Southern Ocean (Figure 7b), all high  
280 ECS models show a decrease in cloud ice water path whereas about half of the low ECS models show an increase.

## Cloud radiative effects

Over the northern polar region the cooling effect of the net cloud radiative effect increases significantly for all three ECS groups (Figure 6def). Averaged over the whole Arctic (Figure 7a), the low ECS group shows the strongest increase in cooling among the three ECS groups. The increase in net cloud radiative effect is dominated by a stronger shortwave cloud radiative effect  
285 that is only partly compensated by a larger longwave cloud radiative effect. This is driven particularly by an increase in cloud liquid water path and only to a smaller extent to an increase in cloud ice water path and total cloud cover (Figures 6abc).

North of about 50°N and south of about 50°S, all three ECS groups show stronger shortwave cloud radiative effects, i.e. stronger cooling, in the future scenarios than in the historical simulations. In contrast, the shortwave cloud radiative effect is



reduced in the projections in mid- and low latitudes. Here, the low ECS group shows the smallest changes, while the reductions  
290 in shortwave cloud radiative effect per degree of warming are strongest in the high ECS group. This is mainly driven by a  
reduction in total cloud cover alongside a reduction in liquid water path that can only be compensated within about  $\pm 10^\circ$   
around the Equator by an increase in cloud ice water path (Figure 6abc).

On average, there is a small decrease in the amplitude of the net radiative effect between about 1 and 3%  $\text{K}^{-1}$  for high ECS  
models in the latitude belt  $50^\circ\text{S}$  to  $50^\circ\text{N}$ . For the two other groups there is a small increase in the amplitude. Beyond  $50^\circ\text{N}$  and  
295  $50^\circ\text{S}$  the amplitude of the net cloud radiative effect increases (i.e. more negative) per degree temperature change with a peak  
at about  $65^\circ\text{S}$  and  $80^\circ\text{N}$  of about 25% and 30%, respectively, per degree temperature increase. Ceppi et al. (2016) shows that  
this cloud response results from an increasing cloud optical depth with temperature which is in agreement with the increased  
liquid water path in Figure 6c.

In the Tropics, the high ECS group shows the strongest weakening of the net cloud radiative effect. This is caused by a  
300 reduced shortwave cooling (Figure 6e) connected to the decrease in total cloud fraction. In contrast, the medium and low ECS  
groups show a stronger net cloud radiative effect (i.e. more negative) with warming in the future projections. This different  
behavior can also be seen in Figure 7c.

Driven mostly by the changes in total cloud cover and liquid water path, the cooling effect of the net cloud radiative effect in  
the stratocumulus regions amplifies with warming in the low ECS group while it gets weaker in the high ECS group (Figure 7d).  
305 Again, the medium ECS group is in between the two other groups with many individual models within this group disagreeing  
on the sign of the change in the net cloud radiative effect with warming.

#### 4 Summary and conclusions

The uncertainty in the representation of clouds and their response to climate change is one of the main contributors to the  
overall uncertainty in effective climate sensitivity and thus projections of future climate. The increased range of ECS obtained  
310 from the ensemble of CMIP6 models compared to previous CMIP phases motivated us to look into the differences in present-  
day and future projections of cloud parameters. Of particular interest was whether there are systematic differences in projected  
cloud properties among the models contributing to differences in ECS. We therefore sorted 51 CMIP5 and CMIP6 models  
providing the required output in three equally sized ECS groups. Models with an ECS higher than 4.0 K belong to the "high"  
ECS group, with an ECS between 2.87 K and 4.0 K to the "medium" and with an ECS lower than 2.87 K to the "low" ECS  
315 group. Furthermore, historical simulations of the models were compared to each other to obtain a qualitative overview on the  
differences among the three model groups in simulating observed cloud patterns and properties.

We found higher total cloud cover values in the high ECS group mean with especially the maxima over the Southern Ocean  
and the northern Atlantic and Pacific being more pronounced. The high ECS group mean also shows the largest values of ice  
water path over the Southern Ocean among the three groups whereas the expected maxima of the ice water path in the Tropics  
320 related to the ITCZ are more pronounced in the low ECS group mean. The liquid water path is lowest in the high ECS group  
mean over the whole globe.

When comparing the group mean net cloud radiative effects to the observationally based CERES-EBAF dataset, we found the bias, RMSD and correlation to be significant worse for the low ECS group than for the two other groups. The low ECS group shows the highest overestimation of the net cloud radiative effect in the Tropics and at the same time the highest ice  
325 water path in this region among the three model groups.

In order to investigate the sensitivity of cloud parameters to future warming simulated by the three ECS groups, we compared results from historical simulations with the ones from RCP8.5 and SSP5-8.5 runs from each group. We found that in polar regions, the increase in cloud cover per degree of warming is strongest in the low ECS models, which is about a factor of 2-3 higher than in the high ECS models. Together with an increase in cloud ice and liquid water path, the cooling effect of the net  
330 cloud radiative effect increases significantly for all three ECS groups particularly in the northern polar region. These simulated future changes in all three groups in polar regions are consistent with satellite observations showing an increase in the observed brightness of Arctic clouds in recent years (Lelli et al., 2023). Averaged over the whole Arctic, the low ECS group shows the strongest increase in the cooling effect of the shortwave cloud radiative effect among the three ECS groups.

In midlatitudes and in the Tropics, the three model groups do not agree on the sign of the sensitivity of cloud cover to  
335 warming. While the high ECS models show a decrease in cloud fraction particularly in SH mid- and high latitudes south of  $45^{\circ}\text{S}$ , the low ECS group shows a strong positive sensitivity of up to more than  $1\% \text{K}^{-1}$ . Over the tropical ocean, all high ECS models show a decrease in total cloud cover while the individual models in the two other groups do not agree on the sign of the change. The shortwave cloud radiative effect is reduced in the projections in mid- and low latitudes with the low ECS group showing the smallest changes, while the reductions in shortwave cloud radiative effect per degree of warming are strongest in  
340 the high ECS group. This is mainly driven by a reduction in total cloud cover alongside a reduction in liquid water path that can only be compensated within about  $\pm 10^{\circ}$  around the Equator by an increase in cloud ice water path. Between about  $10^{\circ}\text{S}$  and  $10^{\circ}\text{N}$  all three ECS groups show a strong sensitivity of the cloud ice water path to warming of up to  $9\% \text{K}^{-1}$  and  $10\% \text{K}^{-1}$ . This increase in cloud ice water path is expected to be related to stronger and/or more frequent deep convection as the main increase in the vertical distribution of cloud ice occurs in the upper troposphere around 300 hPa and higher (not shown).

345 Similarly, the behavior of the three ECS groups is different in the subtropical stratocumulus regions. The high ECS group shows a decrease in total cloud cover with warming, the low and medium ECS groups show particularly in the SH stratocumulus regions an increase in total cloud cover. Together with changes in liquid water path following changes in cloud cover, the cooling effect of the net cloud radiative effect in the stratocumulus regions amplifies with warming in the low ECS group while it gets weaker in the high ECS group.

350 Over the Southern Ocean, we found a decrease in ice water path and an increase in liquid water path with warming. These changes, however, are not statistically significantly different among the three ECS groups. Averaged over the whole Southern Ocean (latitude belt  $30\text{-}65^{\circ}\text{S}$ ), all high ECS models agree in a future decrease in cloud ice water path whereas about half of the low ECS models show a positive and half of the models a negative change in cloud ice. This might be connected to the higher ice water path over the Southern Ocean of the high ECS group mean in today's climate.

355 Our results suggest that the differences in the net cloud radiative effect as a response to warming and thus differences in ECS among the CMIP models are not solely driven by an individual region but rather by changes in a range of cloud regimes

leading to differences in the net cloud radiative effects. Contributors are changes in all different global cloud regimes, in polar regions, in tropical and subtropical regions and in midlatitudes. In polar regions, high ECS models show a significantly weaker increase in the net cooling effect of clouds due to warming than the low ECS models. At the same time, high ECS models show a decrease in the net cooling effect of clouds over the tropical ocean and the subtropical stratocumulus regions. In both regions low ECS models show either little change or even an increase in the cooling effect as a consequence of warming. The differences among the ECS groups in the Southern Ocean fit consistently into this picture, showing a higher sensitivity of the net cloud radiative effect to warming in the low ECS models than in the high ECS models. We thus conclude that changes in all three regions contribute to the amplitude of simulated ECS.

*Code and data availability.* All model simulations used for this paper are publicly available on ESGF. The observational dataset CERES-EBAF Ed4.2 (see Section 2.2) is not distributed with the ESMValTool that is restricted to the code as open source software, but the ESMValTool provides a script with exact downloading and processing instructions to recreate the dataset used in this publication. All diagnostics used for this paper will be made available in the ESMValTool after acceptance of this publication. ESMValTool v2 is released under the Apache License, version 2.0. The latest release of ESMValTool v2 is publicly available on Zenodo at <https://doi.org/10.5281/zenodo.3401363>. The source code of the ESMValCore package, which is installed as a dependency of the ESMValTool v2, is also publicly available on Zenodo at <https://doi.org/10.5281/zenodo.3387139>. ESMValTool and ESMValCore are developed on the GitHub repositories available at <https://github.com/ESMValGroup> with contributions from the community very welcome. For more information, we refer to the ESMValTool website (<https://www.esmvaltool.org>). All figures from this paper can be reproduced with the ESMValTool "recipe" (configuration script defining all datasets, processing steps and diagnostics to be applied) recipe\_bock23acp.yml.

*Author contributions.* LB performed the analysis, prepared all figures, and lead the writing of the manuscript. AL contributed to the scientific interpretation of the results and the writing of the manuscript.

*Competing interests.* No competing interests.

*Acknowledgements.* This project has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement 101003536 (ESM2025—Earth System Models for the Future) and from the ESA Climate Change Initiative Climate Modelling User Group (ESA CCI CMUG).

We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP6. We thank the climate modeling groups for producing and making available their model output, the Earth System Grid Federation (ESGF) for archiving the data and providing access, and the multiple funding agencies that support CMIP6 and ESGF. This work used resources of the Deutsches Klimarechenzentrum (DKRZ) granted by its Scientific Steering Committee (WLA) under project ID bd0854.

The CERES-EBAF data were obtained from the NASA Langley Research Center Atmospheric Science Data Center.

We thank Mattia Righi (DLR) and three anonymous reviewers for their helpful comments on the manuscript.

## References

- Andrews, T., Gregory, J. M., Webb, M. J., and Taylor, K. E.: Forcing, feedbacks and climate sensitivity in CMIP5 coupled atmosphere-ocean climate models, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL051607>, 2012.
- 390 Andrews, T., Andrews, M. B., Bodas-Salcedo, A., Jones, G. S., Kulhbrodt, T., Manners, J., Menary, M. B., Ridley, J., Ringer, M. A., and Sellar, A. A.: Forcings, feedbacks and climate sensitivity in HadGEM3-GC3. 1 and UKESM1, *Journal of Advances in Modeling Earth Systems*, 2019.
- Arora, V. K., Scinocca, J. F., Boer, G. J., Christian, J. R., Denman, K. L., Flato, G. M., Kharin, V. V., Lee, W. G., and Merryfield, W. J.: Carbon emission limits required to satisfy future representative concentration pathways of greenhouse gases, *Geophys Res Lett*, 38, <https://doi.org/10.1029/2010gl046270>, 2011.
- 395 Bentsen, M., Bethke, I., Debernard, J. B., Iversen, T., Kirkevag, A., Seland, O., Drange, H., Roelandt, C., Seierstad, I. A., Hoose, C., and Kristjansson, J. E.: The Norwegian Earth System Model, NorESM1-M - Part 1: Description and basic evaluation of the physical climate, *Geosci Model Dev*, 6, 687–720, <https://doi.org/10.5194/gmd-6-687-2013>, 2013.
- Bi, D. H., Dix, M., Marsland, S. J., O'Farrell, S., Rashid, H. A., Uotila, P., Hirst, A. C., Kowalczyk, E., Golebiewski, M., Sullivan, A., Yan, H. L., Hannah, N., Franklin, C., Sun, Z. A., Vohralik, P., Watterson, I., Zhou, X. B., Fiedler, R., Collier, M., Ma, Y. M., Noonan, J., Stevens, L., Uhe, P., Zhu, H. Y., Griffies, S. M., Hill, R., Harris, C., and Puri, K.: The ACCESS coupled model: description, control climate and evaluation, *Aust Meteorol Ocean Aust Meteorol Ocean*, 63, 41–64, 2013.
- 400 Bjordal, J., Storelvmo, T., Alterskjær, K., and Carlsen, T.: Equilibrium climate sensitivity above 5°C plausible due to state-dependent cloud feedback, *Nature Geoscience*, 13, 718–721, <https://doi.org/10.1038/s41561-020-00649-1>, 2020.
- 405 Bock, L., Lauer, A., Schlund, M., Barreiro, M., Bellouin, N., Jones, C., Meehl, G. A., Predoi, V., Roberts, M. J., and Eyring, V.: Quantifying Progress Across Different CMIP Phases With the ESMValTool, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD032321, <https://doi.org/10.1029/2019JD032321>, 2020.
- Bodas-Salcedo, A., Mulcahy, J. P., Andrews, T., Williams, K. D., Ringer, M. A., Field, P. R., and Elsaesser, G. S.: Strong dependence of atmospheric feedbacks on mixed-phase microphysics and aerosol-cloud interactions in HadGEM3, *Journal of Advances in Modeling Earth Systems*, 2019.
- 410 Bony, S., Stevens, B., Frierson, D. M. W., Jakob, C., Kageyama, M., Pincus, R., Shepherd, T. G., Sherwood, S. C., Siebesma, A. P., Sobel, A. H., Watanabe, M., and Webb, M. J.: Clouds, circulation and climate sensitivity, *Nature Geoscience*, 8, 261–268, <https://doi.org/10.1038/ngeo2398>, 2015.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D'Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, E., L., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Otlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002010, <https://doi.org/10.1029/2019ms002010>, 2020.
- 420

- Brunner, L., Pendergrass, A. G., Lehner, F., Merrifield, A. L., Lorenz, R., and Knutti, R.: Reduced global warming from CMIP6 projections when weighting models by performance and independence, *Earth System Dynamics*, 11, 995–1012, <https://doi.org/10.5194/esd-11-995-2020>, 2020.
- 425
- Cao, J., Wang, B., Young-Min, Y., Ma, L., Li, J., Sun, B., Bao, Y., He, J., Zhou, X., and Wu, L.: The NUIST Earth System Model (NESM) version 3: description and preliminary evaluation, *Geoscientific Model Development*, 11, 2975–2993, 2018.
- Ceppi, P., McCoy, D. T., and Hartmann, D. L.: Observational evidence for a negative shortwave cloud feedback in middle to high latitudes, *Geophysical Research Letters*, 43, 1331–1339, <https://doi.org/10.1002/2015GL067499>, [\\_eprint: https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015GL067499](https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015GL067499), 2016.
- 430
- Ceppi, P., Brient, F., Zelinka, M. D., and Hartmann, D. L.: Cloud feedback mechanisms and their representation in global climate models, *WIREs Climate Change*, 8, e465, <https://doi.org/10.1002/wcc.465>, publisher: John Wiley & Sons, Ltd, 2017.
- CERES-EBAF: CERES-EBAF Ed4.1 Data Quality Summary V3, [https://ceres.larc.nasa.gov/documents/DQ\\_summaries/CERES\\_EBAF\\_Ed4.1\\_DQS.pdf](https://ceres.larc.nasa.gov/documents/DQ_summaries/CERES_EBAF_Ed4.1_DQS.pdf), (last access on 2023-11-23), 2021.
- 435
- Cesana, G., Ackerman, A., Črnivec, N., Pincus, R., and Chepfer, H.: An observation-based method to assess tropical stratocumulus and shallow cumulus clouds and feedbacks in CMIP6 and CMIP5 models, *Environmental Research Communications*, 5, 045 001, 2023.
- Cherchi, A., Fogli, P. G., Lovato, T., Peano, D., Iovino, D., Gualdi, S., Masina, S., Scoccimarro, E., Materia, S., and Bellucci, A.: Global Mean Climate and Main Patterns of Variability in the CMCC-CM2 Coupled Model, *Journal of Advances in Modeling Earth Systems*, 11, 185–209, 2019.
- 440
- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S.: Development and evaluation of an Earth-System model-HadGEM2, *Geosci Model Dev*, 4, 1051–1075, <https://doi.org/10.5194/gmd-4-1051-2011>, 2011.
- Danabasoglu, G., Lamarque, J. F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, 445 W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *J Adv Model Earth Sy*, 12, 2020.
- Donner, L. J., Wyman, B. L., Hemler, R. S., Horowitz, L. W., Ming, Y., Zhao, M., Golaz, J. C., Ginoux, P., Lin, S. J., Schwarzkopf, M. D., 450 Austin, J., Alaka, G., Cooke, W. F., Delworth, T. L., Freidenreich, S. M., Gordon, C. T., Griffies, S. M., Held, I. M., Hurlin, W. J., Klein, S. A., Knutson, T. R., Langenhorst, A. R., Lee, H. C., Lin, Y. L., Magi, B. I., Malyshev, S. L., Milly, P. C. D., Naik, V., Nath, M. J., Pincus, R., Ploshay, J. J., Ramaswamy, V., Seman, C. J., Shevliakova, E., Sirutis, J. J., Stern, W. F., Stouffer, R. J., Wilson, R. J., Winton, M., Wittenberg, A. T., and Zeng, F. R.: The Dynamical Core, Physical Parameterizations, and Basic Simulation Characteristics of the Atmospheric Component AM3 of the GFDL Global Coupled Model CM3, *J Climate*, 24, 3484–3519, <https://doi.org/10.1175/2011jcli3955.1>, 455 2011.
- Dufresne, J. L., Foujols, M. A., Denvil, S., Caubel, A., Marti, O., Aumont, O., Balkanski, Y., Bekki, S., Bellenger, H., Benschila, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., de Noblet, N., Duvel, J. P., Ethe, C., Fairhead, L., Fichet, T., Flavoni, S., Friedlingstein, P., Grandpeix, J. Y., Guez, L., Guilyardi, E., Hauglustaine, D., Hourdin, F., Idelkadi, A., Ghattas, J., Joussaume, S., Kageyama, M., Krinner, G., Labetoulle, S., Lahellec, A., Lefebvre, M. P., Lefebvre, F., Levy, C., Li, Z. X., 460 Lloyd, J., Lott, F., Madec, G., Mancip, M., Marchand, M., Masson, S., Meurdesoif, Y., Mignot, J., Musat, I., Parouty, S., Polcher, J.,

- Rio, C., Schulz, M., Swingedouw, D., Szopa, S., Talandier, C., Terray, P., Viovy, N., and Vuichard, N.: Climate change projections using the IPSL-CM5 Earth System Model: from CMIP3 to CMIP5, *Clim Dynam*, 40, 2123–2165, <https://doi.org/10.1007/s00382-012-1636-1>, 2013.
- 465 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geosci Model Dev*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Eyring, V., Bock, L., Lauer, A., Righi, M., Schlund, M., Andela, B., Arnone, E., Bellprat, O., Brotz, B., Caron, L. P., Carvalhais, N., Cionni, I., Cortesi, N., Crezee, B., Davin, E. L., Davini, P., Debeire, K., de Mora, L., Deser, C., Docquier, D., Earnshaw, P., Ehbrecht, C., Gier, B. K., Gonzalez-Reviriego, N., Goodman, P., Hagemann, S., Hardiman, S., Hassler, B., Hunter, A., Kadow, C., Kindermann, S., Koirala, S., Koldunov, N., Lejeune, Q., Lembo, V., Lovato, T., Lucarini, V., Massonnet, F., Muller, B., Pandde, A., Perez-Zanon, N., Phillips, A., Predoi, V., Russell, J., Sellar, A., Serva, F., Stacke, T., Swaminathan, R., Torralba, V., Vegas-Regidor, J., von Hardenberg, J., Weigel, K., and Zimmermann, K.: Earth System Model Evaluation Tool (ESMValTool) v2.0-an extended set of large-scale diagnostics for quasi-operational and comprehensive evaluation of Earth system models in CMIP, *Geosci Model Dev*, 13, 3383–3438, <https://doi.org/10.5194/gmd-13-3383-2020>, 2020.
- 470
- 475 Frey, W. R. and Kay, J. E.: The influence of extratropical cloud phase and amount feedbacks on climate sensitivity, *Climate Dynamics*, 50, 3097–3116, <https://doi.org/10.1007/s00382-017-3796-5>, 2018.
- Gent, P. R., Danabasoglu, G., Donner, L. J., Holland, M. M., Hunke, E. C., Jayne, S. R., Lawrence, D. M., Neale, R. B., Rasch, P. J., Vertenstein, M., Worley, P. H., Yang, Z. L., and Zhang, M. H.: The Community Climate System Model Version 4, *J Climate*, 24, 4973–4991, <https://doi.org/10.1175/2011jcli4083.1>, 2011.
- 480 Gattelman, A., Hannay, C., Bacmeister, J. T., Neale, R. B., Pendergrass, A. G., Danabasoglu, G., Lamarque, J. F., Fasullo, J. T., Bailey, D. A., Lawrence, D. M., and Mills, M. J.: High Climate Sensitivity in the Community Earth System Model Version 2 (CESM2), *Geophys Res Lett*, 46, 8329–8337, <https://doi.org/10.1029/2019gl083978>, 2019.
- Giorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Bottinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H. D., Ilyina, T., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Mueller, W., Notz, D., Pithan, F., Raddatz, T., Rast, S., Redler, R., Roeckner, E., Schmidt, H., Schnur, R., Segschneider, J., Six, K. D., Stockhause, M., Timmreck, C., Wegner, J., Widmann, H., Wieners, K. H., Claussen, M., Marotzke, J., and Stevens, B.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *J Adv Model Earth Sy*, 5, 572–597, <https://doi.org/10.1002/Jame.20038>, 2013.
- 485
- Gregory, J. M., Ingram, W., Palmer, M., Jones, G., Stott, P., Thorpe, R., Lowe, J., Johns, T., and Williams, K. J. G. R. L.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophysical research letters*, 31, 2004.
- 490 Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geosci Model Dev*, 13, 2197–2244, 2020.
- He, B., YU, Y., Bao, Q., Lin, P., Liu, H., Li, J., Wang, L., Liu, Y., Wu, G., Chen, K., Guo, Y., Zhao, S., Zhang, X., Song, M., and Xie, J.: CAS FGOALS-f3-L model dataset descriptions for CMIP6 DECK experiments, *Atmospheric and Oceanic Science Letters*, 13, 582–588, <https://doi.org/10.1080/16742834.2020.1778419>, 2020.
- 495

- Hourdin, F., Grandpeix, J. Y., Rio, C., Bony, S., Jam, A., Cheruy, F., Rochetin, N., Fairhead, L., Idelkadi, A., Musat, I., Dufresne, J. L., Lahellec, A., Lefebvre, M. P., and Roehrig, R.: LMDZ5B: the atmospheric component of the IPSL climate model with revisited parameterizations for clouds and convection, *Clim Dynam*, 40, 2193–2222, <https://doi.org/10.1007/s00382-012-1343-y>, 2013.
- 500 Ji, D., Wang, L., Feng, J., Wu, Q., Cheng, H., Zhang, Q., Yang, J., Dong, W., Dai, Y., Gong, D., Zhang, R. H., Wang, X., Liu, J., Moore, J. C., Chen, D., and Zhou, M.: Description and basic evaluation of Beijing Normal University Earth System Model (BNU-ESM) version 1, *Geosci Model Dev*, 7, 2039–2064, <https://doi.org/10.5194/gmd-7-2039-2014>, 2014.
- Jian, B., Li, J., Zhao, Y., He, Y., Wang, J., and Huang, J.: Evaluation of the CMIP6 planetary albedo climatology using satellite observations, *Climate Dynamics*, 54, 5145–5161, <https://doi.org/10.1007/s00382-020-05277-4>, 2020.
- 505 Jiang, J. H., Su, H., Wu, L., Zhai, C., and Schiro, K. A.: Improvements in Cloud and Water Vapor Simulations Over the Tropical Oceans in CMIP6 Compared to CMIP5, *Earth and Space Science*, 8, e2020EA001 520, <https://doi.org/10.1029/2020EA001520>, 2021.
- Kato, S., Rose, F. G., Rutan, D. A., Thorsen, T. J., Loeb, N. G., Doelling, D. R., Huang, X., Smith, W. L., Su, W., and Ham, S.-H.: Surface Irradiances of Edition 4.0 Clouds and the Earth’s Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Data Product, *Journal of Climate*, 31, 4501 – 4527, <https://doi.org/10.1175/JCLI-D-17-0523.1>, 2018.
- 510 Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., and others: GISS-E2. 1: Configurations and climatology, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 025, publisher: Wiley Online Library, 2020.
- Kuhlbrot, T., Jones, C. G., Sellar, A., Storkey, D., Blockley, E., Stringer, M., Hill, R., Graham, T., Ridley, J., and Blaker, A.: The Low-Resolution Version of HadGEM3 GC3. 1: Development and Evaluation for Global Climate, *Journal of advances in modeling earth systems*, 515 10, 2865–2888, 2018.
- Kuma, P., Bender, F. A.-M., Schuddeboom, A., McDonald, A. J., and Seland, O.: Machine learning of cloud types in satellite observations and climate models, *Atmospheric Chemistry and Physics*, 23, 523–549, <https://doi.org/10.5194/acp-23-523-2023>, 2023.
- Lauer, A. and Hamilton, K.: Simulating Clouds with Global Climate Models: A Comparison of CMIP5 Results with CMIP3 and Satellite Data, *Journal of Climate*, 26, 3823 – 3845, <https://doi.org/10.1175/JCLI-D-12-00451.1>, place: Boston MA, USA Publisher: American Meteorological Society, 2013.
- 520 Lauer, A., Eyring, V., Bellprat, O., Bock, L., Gier, B. K., Hunter, A., Lorenz, R., Pérez-Zanón, N., Righi, M., Schlund, M., Senftleben, D., Weigel, K., and Zechlau, S.: Earth System Model Evaluation Tool (ESMValTool) v2.0 – diagnostics for emergent constraints and future projections from Earth system models in CMIP, *Geosci. Model Dev.*, 13, 4205–4228, <https://doi.org/10.5194/gmd-13-4205-2020>, 2020.
- Lauer, A., Bock, L., Hassler, B., Schröder, M., and Stengel, M.: Cloud Climatologies from Global Climate Models – A Comparison of CMIP5 and CMIP6 Models with Satellite Data, *Journal of Climate*, 36, 281–311, <https://doi.org/10.1175/JCLI-D-22-0181.1>, 2023.
- 525 Lee, J., Kim, J., Sun, M.-A., Kim, B.-H., Moon, H., Sung, H. M., Kim, J., and Byun, Y.-H.: Evaluation of the Korea Meteorological Administration Advanced Community Earth-System model (K-ACE), *Asia-Pacific Journal of Atmospheric Sciences*, 56, 381–395, <https://doi.org/10.1007/s13143-019-00144-7>, section: 381, 2020a.
- Lee, W. L., Wang, Y. C., Shiu, C. J., Tsai, I. C., Tu, C. Y., Lan, Y. Y., Chen, J. P., Pan, H. L., and Hsu, H. H.: Taiwan Earth System Model 530 Version 1: description and evaluation of mean state, *Geosci Model Dev* *Geosci Model Dev*, 13, 3887–3904, 2020b.
- Lelli, L., Vountas, M., Khosravi, N., and Burrows, J. P.: Satellite remote sensing of regional and seasonal Arctic cooling showing a multi-decadal trend towards brighter and more liquid clouds, *Atmospheric Chemistry and Physics*, 23, 2579–2611, <https://doi.org/10.5194/acp-23-2579-2023>, 2023.



- Li, L., Lin, P., Yu, Y., Wang, B., Zhou, T., Liu, L., Liu, J., Bao, Q., Xu, S., Huang, W., Xia, K., Pu, Y., Dong, L., Shen, S., Liu, Y., Hu, N.,  
535 Liu, M., Sun, W., Shi, X., Zheng, W., Wu, B., Song, M.-R., Liu, H., Zhang, X., Wu, G., Xue, W., Huang, X., Yang, G., Song, Z., and Qiao,  
F.: The Flexible Global Ocean-Atmosphere-Land System Model version g2, *Adv. Atmos. Sci.*, in press, <https://doi.org/10.1007/s00376-012-2140-6>, 2013.
- Li, L. J., Yu, Y. Q., Tang, Y. L., Lin, P. F., Xie, J. B., Song, M. R., Dong, L., Zhou, T. J., Liu, L., Wang, L., Pu, Y., Chen, X. L., Chen, L., Xie,  
540 Z. H., Liu, H. B., Zhang, L. X., Huang, X., Feng, T., Zheng, W. P., Xia, K., Liu, H. L., Liu, J. P., Wang, Y., Wang, L. H., Jia, B. H., Xie,  
F., Wang, B., Zhao, S. W., Yu, Z. P., Zhao, B. W., and Wei, J. L.: The Flexible Global Ocean-Atmosphere-Land System Model Grid-Point  
Version 3 (FGOALS-g3): Description and Evaluation, *J Adv Model Earth Sy*, 12, 2020.
- Loeb, N. and Staff, N.: The Climate Data Guide: CERES EBAF: Clouds and Earth's Radiant Energy Systems (CERES) Energy Balanced and Filled (EBAF), <https://climatedataguide.ucar.edu/climate-data/ceres-ebaf-clouds-and-earths-radiant-energy-systems-ceres-energy-balanced-and-filled>, (last access on 2023-11-23), 2022.
- 545 Loeb, N. G., Doelling, D. R., Wang, H., Su, W., Nguyen, C., Corbett, J. G., Liang, L., Mitrescu, C., Rose, F. G., and Kato, S.: Clouds and  
the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Top-of-Atmosphere (TOA) Edition-4.0 Data Product,  
*Journal of Climate*, 31, 895 – 918, <https://doi.org/10.1175/JCLI-D-17-0208.1>, 2018.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler,  
550 S., Flaeschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohengger, C., Ilyina, T., Jahns,  
T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornblueh, L.,  
Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Mobis, B., Muller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D.,  
Nyawira, S. S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H.,  
Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch,  
555 J. S., Tian, F. X., Voigt, A., Vrese, P., Wieners, K. H., Wilkenskield, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M  
Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO<sub>2</sub>, *J Adv Model Earth Sy J Adv Model Earth Sy*, 11,  
998–1038, 2019.
- Meehl, G. A., Senior, C. A., Eyring, V., Flato, G., Lamarque, J. F., Stouffer, R. J., Taylor, K. E., and Schlund, M.: Context for inter-  
preting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth system models, *Sci Adv*, 6, eaba1981,  
<https://doi.org/10.1126/sciadv.aba1981>, edition: 2020/07/09, 2020.
- 560 Mizuta, R., Yoshimura, H., Murakami, H., Matsueda, M., Endo, H., Ose, T., Kamiguchi, K., Hosaka, M., Sugi, M., Yukimoto, S., and  
others: Climate simulations using MRI-AGCM3. 2 with 20-km grid, *Journal of the Meteorological Society of Japan. Ser. II*, 90, 233–258,  
publisher: Meteorological Society of Japan, 2012.
- Muller, W. A., Jungclaus, J. H., Mauritsen, T., Baehr, J., Bittner, M., Budich, R., Bunzel, F., Esch, M., Ghosh, R., Haak, H., Ilyina, T.,  
Kleine, T., Kornblueh, L., Li, H., Modali, K., Notz, D., Pohlmann, H., Roeckner, E., Stemmler, I., Tian, F., and Marotzke, J.: A Higher-  
565 resolution Version of the Max Planck Institute Earth System Model (MPI-ESM1.2-HR), *J Adv Model Earth Sy J Adv Model Earth Sy*, 10,  
1383–1413, 2018.
- Righi, M., Andela, B., Eyring, V., Lauer, A., Predoi, V., Schlund, M., Vegas-Regidor, J., Bock, L., Brötz, B., de Mora, L., Diblen, F.,  
Dreyer, L., Drost, N., Earnshaw, P., Hassler, B., Koldunov, N., Little, B., Loosveldt Tomas, S., and Zimmermann, K.: Earth System Model  
Evaluation Tool (ESMValTool) v2.0 – technical overview, *Geosci. Model Dev.*, 13, 1179–1199, [https://doi.org/10.5194/gmd-13-1179-](https://doi.org/10.5194/gmd-13-1179-2020)  
570 2020, 2020.

- Rong, X. Y., Li, J., Chen, H. M., Xin, Y. F., Su, J. Z., Hua, L. J., Zhou, T. J., Qi, Y. J., Zhang, Z. Q., Zhang, G., and Li, J. D.: The CAMS Climate System Model and a Basic Evaluation of Its Climatology and Climate Variability Simulation, *J Meteorol Res-Prc J Meteorol Res-Prc*, 32, 839–861, 2018.
- 575 Rotstayn, L. D., Collier, M. A., Dix, M. R., Feng, Y., Gordon, H. B., O’Farrell, S. P., Smith, I. N., and Syktus, J.: Improved simulation of Australian climate and ENSO-related rainfall variability in a global climate model with an interactive aerosol treatment, *Int J Climatol*, 30, 1067–1088, <https://doi.org/10.1002/joc.1952>, 2010.
- Schlund, M., Lauer, A., Gentine, P., Sherwood, S. C., and Eyring, V.: Emergent constraints on Equilibrium Climate Sensitivity in CMIP5: do they hold for CMIP6?, *Earth Syst. Dynam. Discuss.*, 2020, 1–40, <https://doi.org/10.5194/esd-2020-49>, 2020.
- 580 Schlund, M., Hassler, B., Lauer, A., Andela, B., Jöckel, P., Kazeroni, R., Loosveldt Tomas, S., Medeiros, B., Predoi, V., Sénési, S., Servonnat, J., Stacke, T., Vegas-Regidor, J., Zimmermann, K., and Eyring, V.: Evaluation of native Earth system model output with ESMValTool v2.6.0, *Geoscientific Model Development*, 16, 315–333, <https://doi.org/10.5194/gmd-16-315-2023>, 2023.
- Schmidt, G. A., Ruedy, R., Hansen, J. E., Aleinov, I., Bell, N., Bauer, M., Bauer, S., Cairns, B., Canuto, V., Cheng, Y., Del Genio, A., Faluvegi, G., Friend, A. D., Hall, T. M., Hu, Y. Y., Kelley, M., Kiang, N. Y., Koch, D., Lacis, A. A., Lerner, J., Lo, K. K., Miller, R. L., Nazarenko, L., Oinas, V., Perlwitz, J., Perlwitz, J., Rind, D., Romanou, A., Russell, G. L., Sato, M., Shindell, D. T., Stone, P. H., Sun, S., 585 Tausnev, N., Thresher, D., and Yao, M. S.: Present-day atmospheric simulations using GISS ModelE: Comparison to in situ, satellite, and reanalysis data, *J Climate*, 19, 153–192, <https://doi.org/10.1175/Jcli3612.1>, 2006.
- Seland, O., Bentsen, M., Olivie, D., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y. C., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y. C., Griesfeller, J., Grini, A., Guo, C. C., Ilicak, M., Karset, I. H. H., Landgren, O., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z. S., Heinze, C., Iversen, T., and Schulz, M.: Overview of 590 the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario simulations, *Geosci Model Dev Geosci Model Dev*, 13, 6165–6200, 2020.
- Sellar, A. A., Jones, C. G., Mulcahy, J., Tang, Y., Yool, A., Wiltshire, A., O’connor, F. M., Stringer, M., Hill, R., and Palmieri, J.: UKESM1: Description and evaluation of the UK Earth System Model, *Journal of Advances in Modeling Earth Systems*, 2019.
- Shell, K. M., Kiehl, J. T., and Shields, C. A.: Using the Radiative Kernel Technique to Calculate Climate Feedbacks in NCAR’s Community 595 Atmospheric Model, *Journal of Climate*, 21, 2269 – 2282, <https://doi.org/10.1175/2007JCLI2044.1>, 2008.
- Soden, B. J., Broccoli, A. J., and Hemler, R. S.: On the Use of Cloud Forcing to Estimate Cloud Feedback, *Journal of Climate*, 17, 3661 – 3665, [https://doi.org/10.1175/1520-0442\(2004\)017<3661:OTUOCF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<3661:OTUOCF>2.0.CO;2), 2004.
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornbluh, L., Lohmann, U., Pincus, R., Reichler, T., and Roeckner, E.: Atmospheric component of the MPI-M Earth 600 System Model: ECHAM6, *J Adv Model Earth Sy*, 5, 146–172, <https://doi.org/10.1002/jame.20015>, 2013.
- Swart, N. C., Cole, J. N., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., and Hanna, S.: The Canadian Earth System Model version 5 (CanESM5. 0.3), *Geoscientific Model Development*, 12, 4823–4873, 2019.
- Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O., Guérémy, J.-F., Moine, M.-P., Msadek, R., Ribes, A., Rocher, M., 605 Roehrig, R., Salas-y Mélia, D., Sanchez, E., Terray, L., Valcke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and Madec, G.: Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and Future Climate, *Journal of Advances in Modeling Earth Systems*, n/a, <https://doi.org/10.1029/2019ms001791>, 2019.

- Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura, K., Takata, K., O'ishi, R., Yamazaki, D., Suzuki, T., Kurogi, M., Kataoka, T., Watanabe, M., and Kimoto, M.: Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geosci Model Dev Geosci Model Dev*, 12, 2727–2765, 2019.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485–498, 2012.
- Volodire, A., Saint-Martin, D., Senesi, S., Decharme, B., Alias, A., Chevallier, M., Colin, J., Gueremy, J. F., Michou, M., Moine, M. P., Nabat, P., Roehrig, R., Melia, D. S. Y., Seferian, R., Valcke, S., Beau, I., Belamari, S., Berthet, S., Cassou, C., Cattiaux, J., Deshayes, J., Douville, H., Ethe, C., Franchisteguy, L., Geoffroy, O., Levy, C., Madec, G., Meurdesoif, Y., Msadek, R., Ribes, A., Sanchez-Gomez, E., Terray, L., and Waldman, R.: Evaluation of CMIP6 DECK Experiments With CNRM-CM6-1, *J Adv Model Earth Sy J Adv Model Earth Sy*, 11, 2177–2213, 2019.
- Volodin, E. M., Dianskii, N. A., and Gusev, A. V.: Simulating present-day climate with the INMCM4.0 coupled model of the atmospheric and oceanic general circulations, *Izv Atmos Ocean Phy+*, 46, 414–431, <https://doi.org/10.1134/S000143381004002x>, 2010.
- Watanabe, M., Suzuki, T., O'ishi, R., Komuro, Y., Watanabe, S., Emori, S., Takemura, T., Chikira, M., Ogura, T., Sekiguchi, M., Takata, K., Yamazaki, D., Yokohata, T., Nozawa, T., Hasumi, H., Tatebe, H., and Kimoto, M.: Improved Climate Simulation by MIROC5. Mean States, Variability, and Climate Sensitivity, *J Climate*, 23, 6312–6335, <https://doi.org/10.1175/2010JCLI3679.1>, 2010.
- Watanabe, S., Hajima, T., Sudo, K., Nagashima, T., Takemura, T., Okajima, H., Nozawa, T., Kawase, H., Abe, M., Yokohata, T., Ise, T., Sato, H., Kato, E., Takata, K., Emori, S., and Kawamiya, M.: MIROC-ESM 2010: model description and basic results of CMIP5-20c3m experiments, *Geosci Model Dev*, 4, 845–872, <https://doi.org/10.5194/gmd-4-845-2011>, 2011.
- Weigel, K., Bock, L., Gier, B. K., Lauer, A., Righi, M., Schlund, M., Adeniyi, K., Andela, B., Arnone, E., Berg, P., Caron, L.-P., Cionni, I., Corti, S., Drost, N., Hunter, A., Lledó, L., Mohr, C. W., Paçal, A., Pérez-Zanón, N., Predoi, V., Sandstad, M., Sillmann, J., Sterl, A., Vegas-Regidor, J., von Hardenberg, J., and Eyring, V.: Earth System Model Evaluation Tool (ESMValTool) v2.0 – diagnostics for extreme events, regional and impact evaluation, and analysis of Earth system models in CMIP, *Geoscientific Model Development*, 14, 3159–3184, <https://doi.org/10.5194/gmd-14-3159-2021>, 2021.
- Williams, K. D., Copsey, D., Blockley, E. W., Bodas-Salcedo, A., Calvert, D., Comer, R., Davis, P., Graham, T., Hewitt, H. T., Hill, R., Hyder, P., Ineson, S., Johns, T. C., Keen, A. B., Lee, R. W., Megann, A., Milton, S. F., Rae, J. G. L., Roberts, M. J., Scaife, A. A., Schiemann, R., Storkey, D., Thorpe, L., Watterson, I. G., Walters, D. N., West, A., Wood, R. A., Woollings, T., and Xavier, P. K.: The Met Office Global Coupled Model 3.0 and 3.1 (GC3.0 and GC3.1) Configurations, *Journal of Advances in Modeling Earth Systems*, 10, 357–380, <https://doi.org/10.1002/2017ms001115>, 2018.
- Wu, T. W.: A mass-flux cumulus parameterization scheme for large-scale models: description and test with observations, *Clim Dynam*, 38, 725–744, <https://doi.org/10.1007/s00382-011-0995-3>, 2012.
- Wu, T. W., Yu, R. C., Zhang, F., Wang, Z. Z., Dong, M., Wang, L. N., Jin, X., Chen, D. L., and Li, L.: The Beijing Climate Center atmospheric general circulation model: description and its performance for the present-day climate, *Clim Dynam*, 34, 123–147, <https://doi.org/10.1007/s00382-008-0487-2>, 2010.
- Wu, T. W., Lu, Y. X., Fang, Y. J., Xin, X. G., Li, L., Li, W. P., Jie, W. H., Zhang, J., Liu, Y. M., Zhang, L., Zhang, F., Zhang, Y. W., Wu, F. H., Li, J. L., Chu, M., Wang, Z. Z., Shi, X. L., Liu, X. W., Wei, M., Huang, A. N., Zhang, Y. C., and Liu, X. H.: The Beijing Climate Center Climate System Model (BCC-CSM): the main progress from CMIP5 to CMIP6, *Geosci Model Dev Geosci Model Dev*, 12, 1573–1600, 2019.

- Yukimoto, S., Adachi, Y., Hosaka, M., Sakami, T., Yoshimura, H., Hirabara, M., Tanaka, T. Y., Shindo, E., Tsujino, H., Deushi, M., Mizuta, R., Yabu, S., Obata, A., Nakano, H., Koshiro, T., Ose, T., and Kitoh, A.: A New Global Climate Model of the Meteorological Research Institute: MRI-CGCM3-Model Description and Basic Performance, *J Meteorol Soc Jpn*, 90a, 23–64, <https://doi.org/10.2151/jmsj.2012-A02>, 2012.
- 650 Yukimoto, S., Kawai, H., Koshiro, T., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yabu, S., Yoshimura, H., Shindo, E., Mizuta, R., Obata, A., Adachi, Y., and Ishii, M.: The Meteorological Research Institute Earth System Model Version 2.0, MRI-ESM2.0: Description and Basic Evaluation of the Physical Component, *J Meteorol Soc Jpn* *J Meteorol Soc Jpn*, 97, 931–965, 2019.
- Zelinka, M. D., Myers, T. A., Mccoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., and Taylor, K. E.: Causes of Higher  
655 Climate Sensitivity in CMIP6 Models, *Geophys Res Lett*, 47, <https://doi.org/10.1029/2019GL085782>, 2020.
- Zelinka, M. D., Klein, S. A., Qin, Y., and Myers, T. A.: Evaluating Climate Models’ Cloud Feedbacks Against Expert Judgment, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035 198, <https://doi.org/10.1029/2021JD035198>, 2022.
- Zhu, J., Otto-Bliesner, B. L., Brady, E. C., Gettelman, A., Bacmeister, J. T., Neale, R. B., Poulsen, C. J., Shaw, J. K., McGraw, Z. S., and  
660 Kay, J. E.: LGM Paleoclimate Constraints Inform Cloud Parameterizations and Equilibrium Climate Sensitivity in CESM2, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002 776, <https://doi.org/10.1029/2021MS002776>, 2022.