

Weather persistence on sub-seasonal to seasonal timescales: a methodological review

Alexandre Tuel¹ and Olivia Martius^{1,2}

¹Institute of Geography and Oeschger Centre for Climate Change Research, University of Bern, Switzerland

²Mobilair Lab for Natural Risks, University of Bern, Switzerland

Correspondence: Alexandre Tuel (alexandre.tuel@giub.unibe.ch)

Abstract. Persistence is an important concept in meteorology. It refers to surface weather or the atmospheric circulation either remaining in approximately the same state (stationarity) or repeatedly occupying the same state (recurrence) over some prolonged period of time. Persistence can be found at many different timescales; however, the sub-seasonal to seasonal (S2S) timescale is especially relevant in terms of impacts and atmospheric predictability. For these reasons, S2S persistence has been attracting increasing attention by the scientific community. The dynamics responsible for persistence and their potential evolution under climate change are a notable focus of active research. However, one important challenge facing the community is how to define persistence, from both a qualitative and quantitative perspective. Despite a general agreement on the concept, many different definitions and perspectives have been proposed over the years, among which it is not always easy to find one's way. The purpose of this review is to present and discuss existing concepts of weather persistence, associated methodologies and physical interpretations. In particular, we call attention to the fact that persistence can be defined as a global or as a local property of a system, with important implications in terms of methods but also impacts. We also highlight the importance of timescale and similarity metric selection, and illustrate some of the concepts using the example of summertime atmospheric circulation over Western Europe.

1 Introduction

Surface weather persistence at sub-seasonal to seasonal (S2S) timescales can have severe impacts on human and natural systems. Long-lasting dry conditions, for instance, can lead to droughts and wildfires and can affect agriculture and energy production. Long-lasting wet spells may cause severe flooding and crop loss. Persistent surface weather can result either from quasi-stationary, long-lived atmospheric circulation conditions (stationarity) or from repeated, shorter-lived circulation features (recurrence). Recurrence refers to the repeated occurrence of similar large-scale circulation types or weather systems within some (S2S) time interval, usually with brief interruptions. Many recent high-impact weather and climate events were linked to persistent quasi-stationary or recurrent weather conditions. An example for recurrence are the Western European floods in July 2021 that occurred at the end of an extreme wet spell in Western Europe. The wet spell resulted from repeated atmospheric blocks and Rossby wave breaking episodes (Tuel et al., 2022b). Other examples of recurrence include the floods in the UK during winter 2013-2014 and in Queensland (Australia) in February-April 2022 both were caused by sequences of cyclones

25 (Huntingford et al., 2014) (Wikipedia, 2022; Floodlist, 2022). An example for quasi-stationarity is the catastrophic flooding
in Pakistan in summer 2022 that resulted (in part) from long-lasting and particularly heavy monsoon rains (Mallapaty, 2022).
Intense heatwaves and associated atmospheric circulations also tend to be persistent (often arising from a combination of
recurrence and quasi-stationarity) (Lorenz et al., 2010), as in Western Europe in 2003 (Black et al., 2004; García-Herrera
et al., 2010), Western Russia in 2010 (Drouard and Woollings, 2018; Di Capua et al., 2021) or China (WMO, 2022) and India
30 (Bloomberg, 2022) in 2022.

S2S weather persistence offers the potential for improved predictability at the S2S timescale (Franzke et al., 2011), which is
highly relevant for risk preparedness, and is attracting increased attention from the research community (e.g., Vitart et al., 2017;
Meehl et al., 2021; Domeisen et al., 2022). However, a persistent state is not necessarily well predictable, and persistent states
with low predictability can cause large errors in sub-seasonal weather forecasts (Quandt et al., 2017).
35 Persistence is also an important aspect of climate model evaluation and climate projections. Whether global climate models
are able to correctly simulate persistence is key to the robustness of long-term projections, especially of high-impact weather
events – all the more so as climate projections suggest enhanced persistence (Li and Thompson, 2021; Hoffmann et al., 2021;
Tuel and Martius, 2021a) in the future.

Characterising weather persistence is therefore key to our understanding of the atmospheric circulation and its predictability,
40 and the associated hazards. Previous studies have focused on weather persistence from varied perspectives. Some assessed the
persistence of specific weather systems or features, like atmospheric blocking (e.g. Liu, 1994), Rossby waves (Röthlisberger
et al., 2019), or teleconnection patterns (Barnes and Hartmann, 2010). Others analysed specific episodes of particularly persistent
weather conditions (e.g. Black et al., 2004; Di Capua et al., 2021; Tuel et al., 2022b), while others yet characterised the overall
tendency of the atmospheric circulation and surface weather to exhibit persistence (e.g. MacDonald, 1992; Li and Thompson,
45 2021; Hoffmann et al., 2021). However, while previous studies generally agree on what persistence means conceptually, past
work on this topic has involved many different definitions, often causing confusion and leading to different interpretations of
persistence. Many case studies have also described observed situations as persistent based on subjective analyses rather than
quantitative metrics. A further source of confusion is that "recurrence" is used in the literature to refer not only to successive
occurrences of the same weather pattern at close intervals – what we will focus on in this review – but also to the states of the
50 atmosphere with the highest probability of occurrence (Michelangeli et al., 1995).

It is difficult to give a unique definition of weather persistence. Besides, it may not even be desirable as different interpretations
are possible and useful, depending on the system and timescale of analysis, and on the motivations and goals of the study. Our
goal here is therefore to review existing concepts of weather persistence, associated methodologies and physical interpretations.
We present and structure a wide variety of approaches, definitions, techniques and metrics that have been used to analyse these
55 concepts and that allow answering one or several of the following questions:

- Is there persistence in the data?
- What are the persistent timescales in the data?
- In which specific periods does persistence occur?




Figure 1. Overview of the persistence methods discussed in this paper. Section numbers relative to each methods are indicated in bold between brackets.

– What are the persistent locations in the state space?

60 While persistence occurs at many different timescales, we specifically focus on the S2S timescale – often highly impact-relevant, and certainly important for predictability – but most methods and arguments apply in principle also to longer timescales. An overview of the methods is given in Figure 1. We begin in Section 2 by introducing the two aspects of persistence: stationarity and recurrence. Section 3 then discusses several different perspectives on persistence to consider when choosing an analysis methodology. Finally, we present a detailed list of methods to detect or quantify persistence in Sections 4 and 5. We illustrate
 65 most of the methods with examples taken from the literature or with our own analyses of summertime atmospheric circulation over Europe (details about the data we use are given in the appendix). We keep the interpretation of the results to a minimum, the point being to illustrate the methods and not to analyse European summer circulation persistence in detail.

2 Persistence: stationarity or recurrence?

Persistence in a dynamical system (climate variable, atmospheric circulation field, etc.) arises from the repeated occurrence of
 70 the same value(s) or pattern(s) over a period of time. Successive occurrences can follow each other continuously – a situation we refer to as "stationary" – or in an interrupted sequence – in which case we speak of "recurrence". Persistence therefore comes in two flavours, stationarity and recurrence, which we illustrate with the example of extreme warm conditions in a region north of the Black Sea in Figure 2. Here, extreme warmth is defined as a daily-mean temperature exceeding 1.5 standard deviations above

its annual cycle. In summer 2010, the region experienced persistent (stationary) extreme warmth, with temperatures remaining continuously above the threshold for 41 days straight (Figure 2-a). By contrast, in spring 1979, extreme warm temperatures occurred frequently throughout May, but the three extreme warm episodes within one month were separated by at least 5 days with near-average temperatures (Figure 2-c). The corresponding evolution of the atmospheric circulation can be conceptually described as a slow-moving trajectory for stationary conditions (Figure 2-b) and a more rapidly-evolving trajectory that revisits the same point repetitively (Figure 2-d).

The literature often uses "persistence" as a synonym for "stationarity", (e.g., Dole and Gordon, 1983; Barnston and Livezey, 1987; Franzke et al., 2011; Di Lorenzo and Mantua, 2016; Fereday, 2017; Liu et al., 2018; Du et al., 2019; Francis et al., 2020; Hoffmann et al., 2021; Li and Thompson, 2021), i.e., persistence is associated with long-lived flow anomalies and little change in atmospheric circulation and surface weather. Recurrence has by contrast attracted less scientific attention. Most studies on recurrence have focused on extreme/impactful events (e.g., Mailier et al., 2006; Barton et al., 2016; Dacre and Pinto, 2020; Tuel and Martius, 2022a) and Rossby waves (e.g., Röthlisberger et al., 2019; Ali et al., 2021). Yet, from a physical perspective, stationarity and recurrence are intimately related. Recurring weather systems typically result from stationary favourable large-scale conditions, like sea-surface temperature anomalies or the location of extratropical jets (e.g., Tuel and Martius, 2022b). Temporal dependence, or memory, in a system can thus translate in practice into both stationarity and recurrence (Franzke, 2013). Additionally, from the impacts perspective, it makes sense to look at stationarity and recurrence together, since both can cause prolonged, impactful surface weather. Recurrent Rossby waves, for instance, modulate the persistence of surface temperature and precipitation anomalies (Röthlisberger et al., 2019; Ali et al., 2021) while long-lived SST anomaly patterns like ENSO can remotely trigger recurrent extreme weather (Gershunov and Barnett, 1998).

Note that "recurrence", as we define it here, is sometimes referred to as "temporal" or "serial clustering" (Franzke, 2013), for instance in the case of recurrent cyclones (Mailier et al., 2006) or heavy precipitation events (Barton et al., 2016). Note also that "recurrence" is also frequently used in meteorology to refer to preferred patterns that repeatedly occur in a time series, but not necessarily at close intervals (e.g., Vigaud et al., 2018; Kornhuber et al., 2019; Son et al., 2021) – in the case of atmospheric circulation patterns, one speaks of circulation or weather "regimes" (Michelangeli et al., 1995). This is different from our definition, in which recurrence specifically relates to the repeated occurrence of the same patterns over S2S timescales. Such patterns may be rare in the full dataset and would therefore not be considered as "recurrent" in the regime approach, but can be highly relevant from an impacts perspective. Hannachi et al. (2017) gave a comprehensive overview of the weather regime approach and associated methodologies. We also discuss the relevance of the weather regime perspective for stationarity analysis in section 4.2.1.

Because stationarity and recurrence are two faces of the same coin, distinguishing one from the other may not be evident nor necessarily relevant.

First, the distinction often depends on the variable of interest. Recurrent weather systems can indeed result in stationary surface weather anomalies, and vice-versa. For example, the long-lived (stationary) heatwaves of the 2010 and 2021 summers in Western Russia (Fig. 2-a) and the Baltic were linked to recurrent atmospheric blocks (Drouard and Woollings, 2018; Tuel et al., 2022b). Likewise, prolonged droughts or wet spells may result from recurrent Rossby wave activity (Röthlisberger et al., 2019; Ali




Figure 2. Illustrating stationarity and recurrence on S2S timescales. (a,c) Example of daily mean 2-meter temperature series averaged over the 33° - 43° E/ 50° - 57° N region illustrating (a) stationarity (summer 2010) and (c) recurrence (spring 1979) in extreme warm temperatures on S2S timescales (black: observations; gray: mean annual cycle; light gray: +1.5 standard deviation from the mean). Data is from the ERA5 reanalysis (Hersbach et al., 2020). (b,d) Idealised system trajectories (thick black lines) in the phase space corresponding to stationarity (b) and recurrence (d). The background PDF is shown in light contours. Panels (b,d) reproduced with permission from Hannachi et al. (2017).

et al., 2021, 2022). Stationary surface warm and humid conditions can also trigger recurrent thunderstorm activity (Mohr et al., 2020). Conversely, recurrent extreme precipitation events (e.g., Barton et al., 2022) or extratropical cyclones (e.g., Dacre and Pinto, 2020) are often linked to stationary jet states that last for much longer than the lifetime of individual weather systems. Second, the longer the timescale of analysis, the less obvious the difference between stationarity and recurrence. Impacts for example often depend on anomalies of surface temperature or precipitation averaged or accumulated over several weeks to months (e.g., droughts). Weekly or monthly values may thus sometimes be preferred to daily ones, in which case synoptic-scale stationarity and recurrence would both result in large weekly or monthly anomalies that would result from simple "persistence".

Nevertheless, the distinction between stationarity and recurrence remains highly relevant for several reasons: from a methodological perspective (sections 4 and 5), but also for forecasting and process understanding at the synoptic timescale, as well

as for some stakeholders, like insurers (for whom it matters whether impacts resulted from a single or multiple events). For
120 process understanding, taking a weather systems perspective is often very relevant. In such a case, distinct weather systems
(like cyclones) can in principle be separated from one another, and long-lived single systems be distinguished from multiple
short-lived systems occurring in close succession. It therefore matters whether persistence is driven by recurrence or stationarity.
The distinction is also important to assess whether numerical models simulate persistence for the right physical reasons.

3 Considerations and methodological approaches for persistence analysis in weather and climate data

125 Before reviewing how persistence can be apprehended and quantified, let us begin with some basic notations and definitions.
In the following, we denote by $\mathbf{x}(t) \in \mathbb{R}^m$ the dynamical system under analysis.

$\mathbf{x}(t)$ could for example be the weather evolution in a region, the time evolution of one variable at one location, or the general
circulation. $(\mathbf{x}(t))_t$ evolves within a state space $\mathcal{S}$ that consists of the set of all possible *system states* $\{s_i\}_i$ that the system can
occupy. To each time t corresponds a single *system value*, $\mathbf{x}(t)$, and a single system state. The same system state can be attained
130 at multiple different time steps. The series of successive system values constitutes the system *trajectory*. In practice, only a
finite number of observations are available $(\mathbf{x}(t_k))_{1 \leq k \leq N}$ to characterize the persistence of a system.

Characterising a system as "persistent" can mean different things. Consequently, it is important to always specify the perspective
that is taken to avoid confusion. First, there are different flavours to persistence (global, state or episodic persistence: section
3.1). Second, persistence can be studied from a Lagrangian or a Eulerian perspective (section 3.2). Finally, persistence is linked
135 to a similarity metric (section 3.3) and a notion of timescale (section 3.4), for which different approaches are possible.

3.1 Global, state and episodic persistence

Persistence (whether stationarity or recurrence) is a broad concept that covers different kinds of behaviour in dynamical systems.
We might say, for instance, that temperature is more persistent than precipitation because temperature evolves, on average,
over longer timescales than precipitation. In this sense, persistence quantifies the system's inertia. However, if we qualify last
140 summer's weather as particularly persistent, we mean something different: namely, that last summer's weather varied much less
than what one could have reasonably expected in a normal summer. Here, persistence refers to some unusual behaviour of the
system over a particular period. And saying that zonal jets or atmospheric blocks are persistent means again something else:
that these particular states of the circulation tend to be more long-lived or recurrent than other states.

This leads us to make the distinction between three types of persistence: global, state and episodic persistence.

- 145 1. *Global persistence* characterises the tendency to stationarity or recurrence across the whole trajectory of the dynamical
system. We choose this term because persistence in this sense is a "global" property of the system: it is not restricted to
any particular system state or time period. Global stationarity translates into the tendency for the system to change little at
small timescales (successive values being close to each other). In mathematical terms, this translates into $\langle \left| \frac{d\mathbf{x}(t)}{dt} \right| \rangle \ll \frac{\sigma_x}{T}$
where $\langle \cdot \rangle$ is a time average, σ_x is the standard deviation of the series $x(t)$ and T its typical timescale of evolution.
150 Global persistence can sometimes be impact-relevant (e.g., trends in global persistence under climate change can be

important for impacts). However, it does not make distinctions between the different system states. Consequently, global persistence is not suited to characterise persistent system states or specific time intervals with persistent system behaviour and is generally not the best-suited approach for risk assessment. Global persistence is, however, strongly related to intrinsic system predictability, since present values of the system largely constrain its future values. It can thus yield important information for numerical forecasting, including at the S2S timescale. Most global persistence methods focus on stationarity – like the autocorrelation coefficient (section 4.1.1) or the Hurst exponent (section 4.1.2) – but some exist for recurrence as well (section 5.2.2). Many studies look at global persistence: we may cite, for instance, MacDonald (1992); Pfleiderer and Coumou (2018); Pfleiderer et al. (2019) and Li and Thompson (2021) who analyse the persistence of temperature series and Hoffmann et al. (2021) who consider 10-day atmospheric flow persistence.

2. *State persistence* relates to the persistent behaviour of specific system states. State space persistence analysis consists in either identifying persistent system states – e.g. with the quasi-stationary (section 4.2.2) or the optimally persistent pattern (section 4.2.3) methods – or characterising the persistence of given states – with methods like residence times (section 4.2.5) for stationarity or Ripley’s K for recurrence (section 5.1.3). State persistent is highly relevant from the impacts, the forecasting and the process understanding perspectives. Knowing which states are persistent, and to what extent, is useful i) to make the link to surface impacts and ii) to know which weather patterns or sequences may be more predictable than others. It also helps to determine the physical processes that support or are responsible for the persistence. The state approach to persistence has been used to shed light on stationary states of the North Atlantic circulation and their predictability (Faranda et al., 2017b), to characterise the stationarity of continental-scale weather patterns (Francis et al., 2018, 2020) or drought (Ford and Labosier, 2014), or to quantify recurrence in extra-tropical cyclones (Mailier et al., 2006) and extreme precipitation (Tuel and Martius, 2021a).
3. *Episodic persistence* is tied to specific time intervals (or "episodes") during which the system exhibits stationary or recurrent behaviour. It is in that sense a purely local property that characterises the anomalous behaviour of the system over a limited time period. Importantly, episodic persistence can occur simply by chance in any dynamical system, even in systems that exhibit no global persistence. Episodic persistence is therefore not necessarily relevant for predictability and process understanding. Still, persistent periods can always be analysed to look for potential drivers that discriminate pure "statistical flukes" from possibly predictable events. Episodic persistence is however well-suited for impacts analysis, because it can make a direct link between periods of persistent weather and impacts. Relevant methods include running window techniques (section 4.3) for stationarity and window counts (section 5.1.1) for recurrence. Hoffmann et al. (2021) investigated for example stationarity in 10-day sequences of atmospheric circulation, while Bevacqua et al. (2020) and Kopp et al. (2021) looked at sub-seasonal periods with recurrent cyclones and extreme precipitation events.

While it helps to capture the various interpretations of persistence, this classification is not perfect, and there is some overlap between categories. In practice, the state or episodic perspectives can also be used to quantify global persistence (by averaging persistence metrics across systems states or time intervals) and global persistence metrics can be computed on subsets of the data to identify persistent periods. Some methods, like recurrence plots (section 5.2.2), can even deal with all three types of

185 persistence.

The three types of persistence are also not independent from one another. Global persistence, for instance, can emerge from repeated occurrences in one or a handful of system states only, while the rest of the trajectory, if analysed separately, may not be qualified as persistent. There are also strong relationships between state and episodic persistence. The most common system states to occur during persistent periods are indeed likely to be persistent states. Correspondingly, persistent states, when they occur, are likely to be associated with persistent periods. Persistent states can thus be uncovered from the knowledge of persistent periods, for instance with pattern recognition or clustering algorithms applied to system values during persistent periods, or simply by averaging system values during persistent periods (e.g., Faranda et al., 2017b; Hoffmann et al., 2021). However, state persistence only characterises the average behaviour of system states – it does not imply that all occurrences of a persistent state will necessarily be persistent. Consequently, there is no one-to-one relationship between persistent states and persistent periods. Some system states can behave in a persistent way under certain conditions but not under others. The lifetime and travel speed of atmospheric blocks, for instance, is affected by land-atmosphere feedbacks or upstream latent heating (Steinfeld et al., 2020). Similarly, extratropical cyclones may occur in sequences but also as single events (Dacre and Pinto, 2020). Additionally, persistent periods may be associated with a variety of system states. While we expect persistent states to be the most frequent ones during persistent periods, some states which on average are not persistent can still at times by pure chance behave persistently. Still, this classification is useful to illustrate the different methodological ways that weather persistence can be tackled, and we rely on it to structure the description of methods in sections 4 and 5.

3.2 Lagrangian and Eulerian perspectives

Weather persistence is most often analysed from a Eulerian perspective, i.e., persistence of the same weather pattern or conditions at a fixed location in space. By contrast, in the Lagrangian perspective the focus is on the persistence of a given weather pattern in time. In the Eulerian framework, the system $\mathbf{x}(t)$ represents a time series over a domain fixed with time, whereas in the Lagrangian one, $\mathbf{x}(t)$ follows individual weather patterns or the background atmospheric flow. The Eulerian stationarity of a quantity $\psi(\mathbf{x}, t)$ translates into $\partial\psi/\partial t = 0$ while the Lagrangian stationarity implies $\partial\psi/\partial t + \mathbf{u} \cdot \nabla\psi = 0$ where $\mathbf{u}$ is the background flow. For example, temperature anomalies can be tracked over time Kornhuber and Tamarin-Brodsky (e.g., 2021) or analysed at a fixed location Pfeleiderer and Coumou (e.g., 2018) and Li and Thompson (2021). Similarly, weather systems such as blocking, cyclones or vortices (Bray and Cavallo, 2022) can be analysed at a fixed location or following the weather systems. For example, Økland and Lejenäs (1987) contrast the persistence of blocking at fixed longitudes against the persistence of individual blocking episodes. Another example is Kossin (2018), who characterise the persistence in tropical cyclones by their translation speed.

Both the Eulerian and the Lagrangian perspectives are relevant for impact and risk assessment. The former links persistence to impacts at a given location, and the latter highlights impacts along the trajectory of a weather pattern (system) during its lifetime. Indeed, the same weather system can produce hazardous weather over large areas, putting strain on the resources of insurance companies or governments. The two perspectives can be brought together by considering the translation speed

and lifetime of the tracked weather systems. Systems with long lifetimes and low translation speed lead to both Eulerian and Lagrangian persistence. By contrast, long-lived systems that travel fast are persistent from a Lagrangian perspective only. Likewise, slow-moving but short-lived systems are not Lagrangian-persistent, but Eulerian persistence can still be detected if several such systems occur over the same area in close succession.

3.3 Quantifying similarity

Assessing persistence typically requires quantifying the self-similarity of system values $\mathbf{x}(t)$ with a metric (e.g., Wharton et al., 2008; Zerzucha and Walczak, 2012; Ali et al., 2020; Ontañón, 2020). By self-similarity, we refer to the tendency for successive values of $\mathbf{x}(t)$ to be similar to each other, according to some metric. This is not to be confused with the concept of geometric self-similarity in fractal geometry. Metric selection is an important step that should be done with care, because it conditions how persistence is quantified and some metrics are not suited to certain kinds of data (e.g., heavily skewed). There are two main classes of similarity metrics:

1. *categorical metrics*: with categorical metrics, system values are either similar (if they belong to the same category) or not (if they don't).
2. *continuous metrics*: continuous metrics measure the degree of similarity between two system values in a continuous way.

Categorical metrics focus on specific features of the system, like the presence of a given weather pattern or the occurrence of a specific event. They require the set of all $\mathbf{x}(t)$ values to be classified into distinct categories (usually from 2 to a few dozen). Categories can represent pre-defined system states of interest (e.g., a warm/cold anomaly, a given phase of a teleconnection mode, or the occurrence of a specific weather pattern, like a block) (e.g., Pinto et al., 2014; Drouard and Woollings, 2018; Pfleiderer and Coumou, 2018; Ali et al., 2021; Kopp et al., 2021), but can also be obtained objectively with dimension reduction/pattern recognition methods. Examples include Principal Component Analysis (or Empirical Orthogonal Functions, EOF) (Fereday, 2017, e.g.), Optimally Interpolated Patterns (OPP) (Hannachi, 2008), Self-Organising Maps (SOM) (e.g., Francis et al., 2018; Weiland et al., 2021; Rousi et al., 2022b) or clustering algorithms (probabilistic, hierarchical or non-hierarchical) (e.g., Demuzere et al., 2011; Hannachi et al., 2012; Grams et al., 2017). When $\mathbf{x}(t)$ represents 2D circulation data (like sea-level pressure or geopotential height), the ensemble of system states is often referred to as "weather regimes" (Michelangeli et al., 1995; Grams et al., 2017; Francis et al., 2018) (see section 4.2.1). In EOF analysis, the distance metric is the L^2 norm (Euclidean distance), but most clustering methods can work with any distance metric. The challenge with dimension reduction methods is choosing the number of categories to retain (EOFs, clusters, SOM nodes, etc.). A high number can capture rare states of the system, but at the cost of making persistence more difficult to assess (since sequences when the system remains in the exact same category will become less frequent).

Continuous metrics measure the degree of similarity between any pair of system values in a continuous way. Common examples of continuous metrics include the Euclidean distance (e.g., Faranda et al., 2017b), pattern correlation (e.g., Mo and Ghil, 1987), or more complex similarity indices like the Teweles–Wobus score (e.g., Horton et al., 2017; Blanchet et al., 2018) or the image structural similarity index (SSIM) (e.g., Hoffmann et al., 2021).

In comparison to categorical metrics, continuous metrics offer the advantage that they do not require specifying features/events of interest beforehand. They are also more flexible insofar as similarity can be quantified with respect to any system value as reference, and not just representative values for each category. Since they do not require simplifying the state space, continuous metrics may also be able to pick up rare persistent patterns that are missed by dimension reduction methods that focus on the most common patterns in a series. These advantages come at a cost: working with continuous metrics, especially complex ones, can be more computationally intensive, and sometimes more difficult to interpret physically.

3.4 Persistence timescales

Persistence is linked to a notion of timescale during which the system continuously remains in the same state (for stationarity) or occupies that same state repeatedly (for recurrence). There are three common ways to approach persistence timescales.

The first option is to choose a single, fixed timescale for analysis. This choice can be guided by impact and forecasting considerations, by physical knowledge of the underlying system, or by observations of persistent events (e.g., Huntingford et al., 2014; Lawrence et al., 2020; Overland and Wang, 2021; Rakovec et al., 2022; Rousi et al., 2022a). This is the most common approach to analyse episodic persistence, but it is also applicable to global persistence. Stationarity can be quantified by the average similarity between n successive states (for continuous similarity metrics Kolstad et al. (see e.g., 2017); Hoffmann et al. (see e.g., 2021)) or by the variety of system states during an n -step window (for categorical metrics Fereday (see e.g., 2017); Richardson et al. (see e.g., 2019)). Stationarity can also be inferred from extreme anomalies of circulation, temperature or precipitation during n -step windows. For instance, Gálfi et al. (2019) and Tuel and Martius (2023) identify persistent warm and cold spells by averaging temperature anomalies over 1-3 weeks.

Recurrence can similarly be assessed by calculating the number of times that a particular system state or event occurs during n -step windows (e.g., Mailier et al., 2006; Pinto et al., 2014; Kopp et al., 2021; Tuel and Martius, 2022a). A fixed timescale can also be used as a threshold to separate stationary from non-stationary events, by requiring stationary events to last at least n steps. For instance, Francis et al. (2018) and Francis et al. (2020) define persistent periods by requiring the circulation pattern in a region to remain in the same state for at least four consecutive days. Mann et al. (2018) similarly define persistent resonant wave events as those lasting at least 10 days. Dole and Gordon (1983) also take this approach for the persistence of point-wise geopotential anomalies. Note that this fixed timescale can also consist of a single time step. In this case, "persistence" characterises how much past system values determine future ones. Li and Thompson (2021), for instance, characterise persistence with the lag-1 autocorrelation coefficient, Röthlisberger and Martius (2019) look at 1-day transition probabilities between different system states, and Vautard (1990); Michelangeli et al. (1995) and Hannachi et al. (2017) calculate time derivatives of geopotential fields to identify quasi-stationary states.

The second option is to select an analysis method that explores a range of timescales and pinpoints the relevant persistence timescales, sometimes accompanied by some notion of statistical significance. This approach only works for global persistence. Autocorrelation analysis, for instance, detects the timescales at which the system exhibits significant lagged memory (section 4.1.1). Spectral analysis can likewise highlight important timescales of variability that can be linked to stationarity (section 4.1.2). For recurrence, methods like Ripley's K function indicate the timescales at which recurrence is statistically significant

(section 5.1.3).

Finally, the third option is to characterise persistence not by a single timescale, but by a distribution of timescales. To assess stationarity, one can typically work with the distribution of persistent event durations. Persistent events are periods during which the system verifies a persistence criterion: for stationarity successive system values must be similar; and for recurrence the same event must occur multiple times, each occurrence being separated by at most n time steps from the previous one. For recurrence, it is also possible to consider the distribution of inter-event times (e.g., Altmann and Kantz, 2005). The distribution of event lengths can be further summarised by considering the average or maximum event length (Faranda et al., 2017b; Rousi et al., 2022b), or by modeling it with an exponential or power-law distribution (section 4.2.5). This approach has been applied to numerous cases: heatwaves (Lorenz et al., 2010), droughts (Meng et al., 2017; Moon et al., 2018), wet spells (Ali et al., 2021), atmospheric blocks (Liu, 1994), circulation patterns (Huguenin et al., 2020), and mid-latitude cyclone clustering (Bevacqua et al., 2020).

4 Stationarity

The diversity of perspectives on persistence translates into a wide range of methods, of which we give an overview in the following two sections dedicated respectively to stationarity and recurrence. Following the distinctions introduced in section 3, we separate methods that quantify global, state, and episodic persistence (though some methods can be used for more than one). We also specify whether methods can only be used with a single timescale, or whether they quantify persistence across timescales. An overview of methods is shown in Table 1.

4.1 Global stationarity

We begin with several methods that quantify global stationarity in one-dimensional time series. They characterise stationarity in the series as a whole, but are generally unable to identify stationary states or periods. However, they often allow the user to characterise the timescales of variability and persistence in the data, and are hence relevant for system predictability and process understanding.

4.1.1 Autocorrelation

Autocorrelation is a frequently used measure of stationarity in weather and climate science. If X_t is a continuous, one-dimensional process of mean μ and variance σ^2 , its Pearson autocorrelation coefficient at lag k is defined as:

$$\rho(k) = \frac{\mathbb{E}[(X_t - \mu)(X_{t+k} - \mu)]}{\sigma^2} \quad (1)$$

where $\mathbb{E}$ denotes the expectancy with respect to the distribution of X_t . Potential cycles and long-term trends should be removed from the data before analysis (Weiss and Weiss, 1999). In equation (1), X_t can also be replaced by its rank, which yields the alternative Spearman autocorrelation, more robust to non-linear behaviour in the data. The definition of equation (1) can also be extended to third-order statistics to capture interactions coming from non-linear correlation, yielding the bicorrelation

$$\rho(k_1, k_2) = \frac{\mathbb{E}[(X_t - \mu)(X_{t+k_1} - \mu)(X_{t+k_2} - \mu)]}{\sigma^3} \text{ (Pires and Hannachi, 2021).}$$

Several summary metrics for autocorrelation exist, like the autocorrelation timescale:

$$T_\rho^e = \min_{k>0} \{\rho(k) \leq \exp(1)\} \quad (2)$$

the decorrelation time (Hannachi, 2021)

$$320 \quad T_\rho^d = 1 + 2 \sum_{k=1}^N \rho(k) \quad (3)$$

where N is the number of timesteps (if ρ is integrable), or the characteristic time T_ρ^c (Trenberth, 1984) (Figure 3-a):

$$T_\rho^c = 1 + \sum_{k=1}^N \left(1 - \frac{k}{N}\right) \rho(k) \quad (4)$$

Both T_ρ^d and T_ρ^c roughly approximate the average time between successive independent values. The lag-1 autocorrelation $\rho(1)$ can also be used as a stationarity metric (Li and Thompson, 2021), which is equivalent to fitting a linear regression
 325 model between successive system values: $\mathbf{x}(t+1) = \beta \mathbf{x}(t) + \epsilon(t)$, where $\epsilon(t)$ is a white-noise process. If $\mathbf{x}(t)$ is normalised, the regression slope β is equal to $\rho(1)$. Note that more complex models are possible: linear models for continuous series can be extended by the large class of Autoregressive-Moving Average (ARMA) models.

Autocorrelation has many advantages: a simple definition, ease of interpretation ($\rho(k)$ is related to the linear regression coefficient of X_{t+k} against X_t), and high flexibility; it is already implemented in common programming languages, it requires
 330 no subjective threshold choice, and results can be easily reproduced. By varying the lag k , it can measure stationarity at all timescales. It also comes with a notion of statistical significance: given a confidence level, it is possible to say whether the obtained autocorrelation is significant (indicating a link at lag k) or not, pointing to the relevant stationarity timescales in the series. On the downside, autocorrelation only works for one-dimensional data and requires a large number of values as input. Therefore it is best suited to measuring stationarity globally. One can compute autocorrelation on a subset of the
 335 data only (summer values or a specific time interval, for instance) if the number of data points is large enough, in which case autocorrelation may be used to characterise stationarity locally in time. Additionally, autocorrelation only highlights linear relationships (though bicorrelation can help capture non-linearity). Finally, autocorrelation measures the strength of the connection between lagged system values, but not how far apart they might be in the state space.

Example applications include Horel (1985a) and Barnston and Livezey (1987) who calculated lag autocorrelation on principal
 340 component time series of monthly Northern Hemisphere geopotential fields. MacDonald (1992) used autocorrelation to detect stationarity in monthly temperature series, as did Weiss and Weiss (1999) to assess stationarity in ENSO. Weatherhead et al. (2010) characterised daily temperature stationarity in the Arctic based on lag-1 autocorrelation. Degenhardt and Ólafsson (2019) and Kolstad et al. (2015) calculated lag-1 autocorrelation to highlight intra-seasonal stationarity of monthly-mean temperatures in Iceland and Europe, respectively, and Li and Thompson (2021) applied autocorrelation to daily temperature series and found
 345 it was strongly related to the average length of warm and cold episodes across the world. Kolstad et al. (2017) even used autocorrelation analysis in a causal discovery framework by regressing temperature values against previous ones and including




Figure 3. (a) Autocorrelation characteristic time (T_p^C ; equation (4)) of JJA daily rescaled Z500. (b) Long-range dependence parameter d of unfiltered surface wind speeds. Only those values that are significant at the 5% level are displayed. Reproduced with permission from Franzke (2013).

potential covariates.

We show on Figure 3-a the characteristic time for daily rescaled 500 hPa geopotential height over Europe during summer. While the results say nothing about the role of individual weather systems, the larger values over the British Isles and Western Russia
 350 seem consistent with more frequent persistent atmospheric blocks over these regions.

4.1.2 Asymptotic methods to characterise variability across timescales

Persistence in a system is associated with memory effects that lead to variability being concentrated at long, rather than short timescales. How variability in the series is distributed across timescales is therefore an important indicator of global persistence and can point to relevant persistence timescales. Specifically, persistence often translates into scaling laws: in the time series
 355 variability as a function of frequency or timescale (for stationarity) (Bunde et al., 2002), but also in the distribution of inter-event times (for recurrence, see section 5.1.4). The slope of these laws indicates the degree of persistence. Time series also often exhibit different scaling laws over different frequency intervals, highlighting how persistence may differ across timescales in the data.

360 A first scaling law can be obtained by considering how autocorrelation decreases as a function of time lag. The more stationary a series, the less rapidly its autocorrelation should decrease. Conceptually, time series can be broadly divided between short-range and long-range dependence series. For a short-range dependent series (like autoregressive models), the autocorrelation decreases rapidly with the time lag, eventually reaching 0 after a certain lag or decaying exponentially:

$$\rho(k) \sim \alpha^k \quad (5)$$

365 as $k \rightarrow \infty$, and with $0 < |\alpha| < 1$ (with $\alpha < 0$ corresponding to anti-persistence). By contrast, in a long-range dependent series, the autocorrelation decays following a power-law:

$$\rho(k) \sim k^{2d-1} \quad (6)$$

where d is called the dependence parameter ($0 < d < \frac{1}{2}$) (Beran, 2017; Franzke, 2013). A lower d is associated with a slower decay of the autocorrelation function, hence with more stationary systems. White noise has $d = 0$. Witt and Malamud (2013) and Franzke (2013) discuss several ways how d can be estimated.

Like autocorrelation α and d (the dependence parameter) measure global stationarity in continuous time series because they are based on asymptotic relationships. However, they cannot highlight specific stationarity timescales in a series, as autocorrelation does. Applications to atmospheric time-series show that temperature exhibits long-term dependence (e.g., Yuan et al., 2010; Koscielny-Bunde et al., 1998; Eichner et al., 2003), while precipitation exhibits either short- or long-term dependence (Potter, 1979; Hannachi, 2014; Yang and Fu, 2019). Franzke (2013) computed d and found long-range dependence in North Atlantic winds (Figure 3-b).

The Hurst coefficient (or exponent) H (Hurst, 1951) is a common measure of memory in a time series which is obtained from a second scaling law. Specifically, H characterises how a time series $(X_t)_t$ fluctuates relative to its mean. Noting

$$380 \quad Z_i^n = \sum_{t=(i-1)n+1}^{in} (X_t - \mathbb{E}[X_t]) \quad (7)$$

the cumulative fluctuations of $(X_t)_t$ relative to its mean, calculated over intervals of size n , Hurst argued, empirically from geophysical time series, that these fluctuations could exhibit scale-invariant properties. In other words, cumulative fluctuations over two different timescales n and m are related through

$$Z_i^n \stackrel{d}{=} \left(\frac{n}{m}\right)^H Z_j^m \quad \forall (i, j, k, l) > 0 \quad (8)$$

385 where $\stackrel{d}{=}$ stands for equality in distribution. H ranges from 0 to 1. $H < 1/2$ indicates anti-persistent behaviour, such that successive increments of X_t relative to its mean $(X_t - \mathbb{E}[X_t])$ tend to be negatively correlated, and the time series fluctuates substantially at short timescales. By contrast, $H > 1/2$ points to persistent behaviour, in which successive increments are positively correlated, and variability is concentrated at long timescales. $H = 1/2$ corresponds to white noise (no temporal correlation). Thus, the higher H is, the smoother the time series. H is also theoretically related to the dependency parameter (as $H = 2d + 1$) and the power spectrum exponent (see below) (Franzke et al., 2020).

H can be estimated in many ways (see e.g., Koutsoyiannis, 2003; De las Nieves López García and Requena, 2019; Franzke et al., 2020). In the literature, H has mainly been used to characterise long-term dependence/stationarity (Mandelbrot and Wallis, 1969), for instance in series of monthly- or annual-mean temperature (MacDonald, 1992; Kumar et al., 2013), precipitation (Bunde et al., 2013) or drought indices (Tatli, 2015). However, some studies also calculated it for daily time series (e.g., Rehman and Siddiqi, 2009; Velásquez Valle et al., 2013).

Like the Hurst exponent, spectral analysis also characterises how a time series' variability is distributed across timescales. The power spectrum is commonly defined as the Fourier transform of the autocorrelation function ρ :

$$S(f) = \int_{-\infty}^{\infty} \rho(k) e^{-2i\pi f k} dk \quad (9)$$

where f is the frequency and k the time lag. In a pure white noise series, the variability is distributed equally across frequencies. Hence, the power spectral density is a constant. If temporal dependence is present, however, the power spectrum typically exhibits a power-law decrease with frequency f :

$$S(f) \sim f^{-\beta} \quad (10)$$

where β , called the power spectrum exponent, indicates the degree of stationarity in the time series ($0 < \beta < 1$). For statistically stationary series, one can also show that $\beta = 2H - 1$ where H is the Hurst coefficient (Parzen, 1986).

The larger β is, the more variance is concentrated at low frequencies. This implies a memory effect at low frequencies that relates to stationarity in the time series. Such scaling behaviour is very common in climatological series, and the spectrum is often divided into distinct scaling regimes corresponding to specific frequency intervals. Fraedrich and Larnder (1993) discuss the example of precipitation in continental Europe, and relate the different regimes to specific physical processes and timescales (from individual storms at high frequencies to climate fluctuations at low frequencies) (Figure 4). Yang and Fu (2019) obtain similar results using hourly and daily precipitation data for the United States. Pelletier and Turcotte (1997) also used power spectra to quantify stationarity in various monthly climatic series, as did Ault et al. (2014) for drought stationarity and MacDonald (1992) for temperature stationarity. Telesca et al. (2016) analysed stationary regimes in 10-minute wind series across Switzerland.

Note that, as with the autocorrelation, the definition of the power spectrum can be extended to third-order statistics to yield the bispectrum (the Fourier transform of the bicornelation function) (Pires and Hannachi, 2021). While the power spectrum characterises how the signal's variance is distributed across timescales, the bispectrum provides the contribution of each pair of frequency to the signal's skewness. Pires and Hannachi (2021) for instance calculated the bispectrum of the 3-monthly ENSO series to better understand its predictability.

4.2 State stationarity

We now turn to methods that focus on stationarity of specific system states. Unlike global methods, which take one-dimensional time series as input, several of the following methods are directly applicable to multidimensional data. We begin with methods that identify stationary states from the system trajectory (sections 4.2.1-4.2.4), before discussing methods that quantify the average stationarity of a system state (sections 4.2.5-4.2.6).

4.2.1 Weather regimes

We begin with an identification method for stationary states based on the concept of "weather regimes" (Michelangeli et al., 1995). This concept emerges from the realisation that the extra-tropical atmospheric circulation evolves mainly as a succession




Figure 4. Schematic diagram of the scaling regimes of continental European rainfall obtained by spectral analysis, along with the hypothesised meteorological interpretations of the various regimes. Reproduced under the terms of the Creative Commons CC BY license from Fraedrich and Larnder (1993).

of a handful of large-scale circulation patterns (Hannachi et al., 2017). These preferred sub-seasonal flow patterns, or weather regimes, tend to be quasi-stationary over timescales of a few days to a few weeks and are therefore strongly related to weather persistence. They account for much of the low-frequency atmospheric variability at intra-seasonal timescales (Pandolfo, 1993; Hannachi et al., 2017). The existence of weather regimes in the mid-latitudes has long been recognised, such as the concept of *Grosswetterlagen* (Baur, 1951) or atmospheric blocking (e.g., Namias, 1964). They have attracted considerable attention, in particular because of the potential long-range predictability they offer (e.g., Ghil and Robertson, 2002; Büeler et al., 2021) and their link to surface impacts (e.g., Grams et al., 2017).

There are many ways to calculate weather regimes (see Huth et al. (2008) and Hannachi et al. (2017) for detailed overviews). The most common methods include pattern recognition and dimensionality reduction techniques, applied to a proxy variable for the atmospheric circulation like 500 hPa geopotential height or sea-level pressure. Examples include orthogonal pattern decomposition techniques like Principal Component Analysis (in the temporal or spectral domain) (Fereday, 2017; Grams et al., 2017) or Optimally Interpolated Patterns (OPP) (Hannachi, 2008), self-organizing maps (Huth et al., 2008; Francis et al., 2020; Weiland et al., 2021), archetypal analysis (Hannachi and Trendafilov, 2017; Chapman et al., 2022), and clustering algorithms (Figure 5). The latter can be probabilistic (e.g., Gaussian mixture models; Woollings et al., 2010), hierarchical (e.g., Ward clustering; Hannachi et al., 2012) and non-hierarchical (e.g., k-means or Partitioning Around Medoids; Grams et al., 2017). Various statistics (gap statistic, silhouette coefficient, etc.) can help objectively select an optimal number of clusters, many

of which are available from the R package `clusterCrit` (Desgraupes, 2018). Note that in practice, the input data should be normalised to remove long-term and seasonal trends to focus exclusively on intra-seasonal variability (Grams et al., 2017). Regimes can also be identified as local maxima in the (multidimensional) PDF of the target field, obtained empirically through e.g., kernel smoothing (Kimoto and Ghil, 1993; Woollings et al., 2010), or with more complex tools of network theory (Mukhin et al., 2022) and topology (Strommen et al., 2022). Finally, Franzke et al. (2011) identify regimes in the North Atlantic jet position with a Hidden Markov Model (HMM). HMMs are a powerful tool that brings together Markov models and Gaussian mixture models. Given N unknown (hidden) states, the HMM models the distribution of the observed series conditionally on each hidden state, with the sequence of hidden states following a first-order Markov process (Franzke et al., 2008). The transition matrix, hidden states and conditional distributions can be estimated simultaneously.

The main drawback of the weather regime approach is that it is biased toward preferred, or frequent, states. Stationary but rare flow patterns that fall outside the range of the major regimes may therefore be missed. Additionally, certain patterns may fall in between regimes and are not or mis-classified. Still, weather regimes are very useful because they transform complex multidimensional systems into categorised, one-dimensional series (according to which regime the system is closest to at each time step).

4.2.2 Quasi-stationary states

One major disadvantage of weather regimes is that they strongly simplify the state space. They also focus on the patterns that account for most of the variability in the data, regardless of their persistence. They may thus miss rare yet impact-relevant persistent patterns. Several other techniques exist to directly extract stationary patterns from the system trajectory. Stationary circulation patterns can be seen as mathematically quasi-stationary solutions of the atmosphere's equations of evolution (Mo and Ghil, 1987). Such solutions are characterized by average system time derivatives close to zero, meaning that the system tends to remain in their vicinity longer than elsewhere in the state space – hence their link to stationary states. Stationary states can therefore be directly identified from the system's dynamics by looking for states whose time derivative is close to zero. If we know the system's exact evolution equations (in the case of simplified models, for example), strictly stationary solutions can be directly computed (e.g., Charney and DeVore, 1979; Legras and Ghil, 1985; Mo and Ghil, 1987). In practice, however, this is rarely the case, and quasi-stationary (or "metastable") states are defined in a statistical sense only, as those for which system tendencies (i.e., time derivatives) approach zero (Vautard, 1990):

$$\{\mathbf{x}^* \text{ such that } \mathcal{T}(\mathbf{x}^*) \approx 0\} \quad (11)$$

$\mathcal{T}(\mathbf{x}^*)$ is the composite tendency at $\mathbf{x}^*$, defined as the average (or area-average for multidimensional fields) of instantaneous tendencies at all occurrences of $\mathbf{x}^*$ – in practice at all times when the system is in the neighbourhood of $\mathbf{x}^*$:

$$\mathcal{T}(\mathbf{x}^*) = \sum_i \frac{d\mathbf{x}}{dt} \bigg|_{\mathbf{x}(t)} \mathbb{1}\{d(\mathbf{x}^*, \mathbf{x}(t)) \leq d_0\} \quad (12)$$

where $d(\cdot, \cdot)$ is a similarity metric and d_0 some small threshold. It is also possible to weigh the terms in equation (12) according to $d(\mathbf{x}^*, \mathbf{x}(t))$ (Vautard, 1990; Michelangeli et al., 1995). When dealing with complex multidimensional systems, it is simpler




Figure 5. Summer weather regimes computed by k-means clustering from daily Z500 fields over the North Atlantic–European sector from 1950 to 2003 (data from NCEP-NCAR reanalysis). To eliminate transient and ambiguous episodes, only sequences of 5 days or more occupied by the same regime are retained (hence total regime frequency summing to $\approx 70\%$). Reproduced with permission from Cassou et al. (2005).

to calculate tendencies on the time series of the system’s leading principal components. Equation (12) can then more easily be solved by minimising $|\mathcal{T}(\mathbf{x})^2|$ in $\mathbf{x}$, with distinct solutions corresponding to different quasi-stationary states. In practice, instantaneous tendencies at any $\mathbf{x}^*$ can exhibit a large variance, and time series are commonly low-pass filtered to remove the short-term noise that would complicate solving for equation (12). Additionally, $|\mathcal{T}(\mathbf{x})^2|$ can be difficult to minimise as it is
480 piecewise constant (due to the finite number of observed $\mathbf{x}$ values). Only approximate solutions are achievable. It may thus be difficult to know precisely how many zeros of $|\mathcal{T}(\mathbf{x})^2|$ exist and whether two approximate solutions correspond to the same minimum. Vautard (1990) presents a method to select relevant solutions.

It is important to note that even if $T_i(\mathbf{x}^*) \approx 0$, instantaneous tendencies $\left. \frac{d\mathbf{x}}{dt} \right|_{\mathbf{x}(t)}$ in equation (12) may be far from zero (Vautard, 1990). Indeed, the time derivative of $\mathbf{x}(t)$ usually depends not only on $\mathbf{x}$ but also on other time-dependent variables that may
485 evolve independently from $\mathbf{x}$.

Haines and Hannachi (1995) and Hannachi (1997) estimated quasi-stationary states over the North Pacific in the output from a global climate forced by perpetual January conditions, by projecting simplified dynamics (e.g., quasi-geostrophy) onto the leading modes of variability of the GCM simulation. Vautard (1990) found four main quasi-stationary patterns in wintertime




Figure 6. Five quasi-stationary states of JJA 5-day averaged rescaled Z500.

daily 700 hPa geopotential height fields over the Atlantic, and analysed their stationarity and onset/break characteristics. Michelangeli et al. (1995) also looked at 700 hPa geopotential fields and compared quasi-stationary states with the leading EOFs over the Atlantic and Pacific Oceans during winter. Mo and Ghil (1987) did a similar comparison for the Southern Hemisphere circulation during the austral winter.

Figure 6 shows five quasi-stationary Z500 anomaly patterns over Europe in summer. We obtained them from the 10 leading EOFs of the 5-day averaged Z500 fields, following Vautard (1990). The patterns were obtained by clustering the resulting approximate solutions with the highest number of clusters for which the resulting patterns were subjectively different. The solutions include blocking-type patterns over Scandinavia/Western Russia (Figure 6-a,b) and zonal patterns (Figure 6-c,d) that are similar to the results of Figures 5 and 11, and to the canonical patterns of variability over the Euro-Atlantic sector (Grams et al., 2017).

4.2.3 Optimally persistent patterns

The technique of optimally persistent patterns (OPPs) was introduced by DelSole (2001). OPP analysis is conceptually similar to principal component analysis; however, instead of patterns that maximise the variance in the observed series, OPPs are defined as the patterns whose time components (i.e., the projection of the observed series onto the OPP) are the most persistent. Persistence is here measured by the decorrelation time (equation (3)) or, alternatively, by the squared decorrelation time $T_2 = 1 + 2 \sum_{k=1}^N \rho(k)^2$ DelSole (2001) (with $\rho(k)$ the autocorrelation function and N the number of observations). If the dynamical system $\mathbf{x}(t)$ is




Figure 7. The pattern associated with the leading eigenvector maximising the squared decorrelation time for daily anomaly fields of 500 hPa geopotential height for the 1950–1999 period (data from NCEP-NCAR reanalysis; units in m). Adapted from DelSole (2001).

505 m-dimensional, then for an m-dimensional pattern $\mathbf{u}$, the time component of $\mathbf{u}$ is defined as $y(t) = \mathbf{u}^T \mathbf{x}(t)$. If $\rho_y(k)$ is its associated autocorrelation function, finding OPPs then consists in maximising $1 + 2 \sum_{k=1}^N \rho_y(k)$ or $1 + 2 \sum_{k=1}^N \rho_y^2(k)$. In the first case (maximising the decorrelation time), the optimisation reduces to an eigenvalue problem, details of which can be found in Hannachi (2021). The leading eigenvector maximises the decorrelation time. OPPs can then be obtained by projecting the observed data $\mathbf{x}(t)$ onto the time series associated with each eigenvector. The second case (maximising the squared decorrelation

510 time) leads to a more complex non-linear optimisation problem that can be solved iteratively (see details in DelSole (2001)). Note that as in section 4.2.2, the system should first be embedded into a lower-dimensional space, e.g., by projecting it onto the set of leading EOFs. Figure 7 shows the leading OPP obtained by DelSole (2001) by maximising the squared decorrelation time for daily Northern Hemisphere Z500 fields.

OPP are especially relevant for forecasting since they correspond to the patterns with the most low-frequency variability.

515 However, like EOFs and quasi-stationary states, OPPs are mathematical objects that do not necessarily correspond to "real" patterns ever attained by the system.

Other methods have been proposed to identify persistent patterns in a spatio-temporal field. One consists in minimising the one-step-ahead forecast error, where the forecast is obtained by projecting the observed field $\mathbf{x}(t)$ onto a set of temporal patterns. These patterns, named "Predictive Oscillation Patterns" by Kooperberg and O'sullivan (1996), can thus be considered

520 as the "most predictable" (i.e., persistent) patterns of a spatio-temporal field. Another possibility is to minimise the interpolation error variance between $\mathbf{x}(t)$ and its estimate $\hat{\mathbf{x}}(t)$ obtained from all $(\mathbf{x}(t'))_{t' \neq t}$ values. In this case one speaks of Optimally Interpolated Patterns (Hannachi, 2008). Further details for both methods can be found in Hannachi (2021).

4.2.4 Extreme values and dynamical systems theory

The quasi-stationary state method only focuses on the most stationary states of a system, and OPPs characterise stationarity for a handful of possible patterns only. By contrast, dynamical systems theory provides a convenient framework to describe the stationarity of any system state (Lucarini et al., 2016). In this framework, stationarity is defined for any point $\mathbf{x}_0$ of the state space as the inverse of the average residence time of trajectories around $\mathbf{x}_0$. The residence time is calculated based on the distance between successive system values, with two successive values deemed similar if their distance is below some small threshold. For any state $\mathbf{x}_0$, the probability that $\mathbf{x}(t)$ will remain in a close neighbourhood around $\mathbf{x}_0$ (a ball of radius ϵ) can be approximated by (Faranda et al., 2017a, b)

$$\mathbb{P}(d(\mathbf{x}(t), \mathbf{x}_0) \leq \epsilon) \simeq \exp \left\{ -\theta(\mathbf{x}_0) \frac{d(\mathbf{x}(t), \mathbf{x}_0) - \mu(\mathbf{x}_0)}{\sigma(\mathbf{x}_0)} \right\} \quad (13)$$

where $d(\cdot, \cdot)$ is a distance function (Euclidean distance in Faranda et al. (2017a, b)) and θ is called the extremal index. ϵ is usually chosen to be the 2nd percentile of the $d(\mathbf{x}(t), \mathbf{x}_0)$ values. θ can be estimated in various ways (Hamidieh et al., 2009; Holešovský and Fusek, 2022), two common ones being the intervals estimator of Ferro and Segers (2003) and the gap estimator of Süveges (Faranda et al., 2017a).

In extreme value statistics, θ is the inverse average duration of consecutive sequences of extreme events, and is used to cluster events (Ferro and Segers, 2003). A large θ therefore indicates that event occurrences tend to be isolated, while a low θ indicates that events occur as part of a larger group. In dynamical systems theory, stationarity is then defined as $1/\theta$ (some studies directly use θ as a measure of stationarity (e.g., Franzke, 2013)). A stationary point of the system (where $\frac{d\mathbf{x}}{dt} = 0$) has infinite stationarity. By contrast, if trajectories immediately leave the neighbourhood of $\mathbf{x}_0$ then the stationarity is equal to 1. In practice, long trajectories of $\mathbf{x}(t)$ are required to explore all possible states of the state space (also called the set of attractor points, or simply the attractor), and to best approximate sequences of states on the attractor. Note that the choice of ϵ constrains the values that θ can reach. In particular, very small values (necessary for equation (13) to hold) usually lead to $1/\theta$ being around 1-2 days (e.g., Faranda et al., 2017b, a; Holmberg et al., 2022), so that the resulting persistence metric is highly local in time and not necessarily relevant for S2S timescales.

The advantage of the method is that it is grounded in mathematical theory and, unlike approaches based on e.g., transition probabilities, it does not require categorising the data. It also provides an easily interpretable index of stationarity for each system state (the average residence time of the system's trajectory around that state). Holmberg et al. (2022) have for instance taken this approach to analyse the link between atmospheric circulation persistence and warm spells in Europe. Faranda et al. (2019) averaged θ values across all time steps to quantify global persistence in the North Atlantic flow. Going back to our example of the European summer circulation, the three major persistent Z500 anomaly patterns are obtained by clustering the top 3% (115) most persistent daily patterns (Figure 8). We chose 5 cluster centers empirically, after testing for 2-15 centers and selecting the highest number of clusters for which the resulting patterns were subjectively different. As in Figure 6, persistent states include blocking regimes (over Scandinavia, Western Russia and the North Sea; Figure 8-a,b,c), a zonal regime (Figure 8-d) and a European ridge regime (Figure 8-e).




Figure 8. Most persistent Z500 anomaly patterns during JJA, obtained by clustering the top 3% (115) daily patterns with the highest stationarity index $1/\theta$. The clustering algorithm is Partitioning Around Medoids (PAM). Also shown are the associated 200 hPa zonal wind anomalies (contours; same levels as Z500). The frequency of each pattern among the 115 most persistent ones is indicated in the top left-hand corner of each panel.

4.2.5 Residence times

A common way to quantify stationarity in time series relies on the concept of "residence time". Residence times extend the dynamical systems approach of the previous section. For a continuous series, the residence time R at time t is defined as the time during which the system remains similar to its value in t :

$$R(t) = \max_{k \geq 0} \{d(\mathbf{x}(t), \mathbf{x}(t+k')) \leq \epsilon \forall 0 \leq k' \leq k\} \quad (14)$$

where $d(\cdot, \cdot)$ is a similarity metric and ϵ is a small threshold. For a categorical series, R is similarly defined as the time during which the system remains in its state $\mathbf{x}(t)$ before transitioning to another state:

$$R(t) = \max_{k \geq 0} \{\mathbf{x}(t+k') = \mathbf{x}(t) \forall 0 \leq k' \leq k\} \quad (15)$$

The definition can be relaxed to allow for brief interruptions in a sequence of similar states (one "average" day between two sequences of warm or wet days, for instance) (e.g., Ali et al., 2021). This definition can be extended to sets $\tilde{\mathcal{S}} \subset \mathcal{S}$ of several system states: $R(t) = \max_{k \geq 0} \{\mathbf{x}(t+k') \in \tilde{\mathcal{S}} \forall 0 \leq k' \leq k\}$ (Richardson et al., 2019). Note that equations (14)-(15) take a Eulerian perspective; their parallel in the Lagrangian perspective is the concept of "survival time", i.e., the duration of a specific weather system or pattern along its trajectory (Liu et al., 2018; von Lindheim et al., 2021).

The residence time approach can characterise episodic, state, and also global stationarity, and is particularly useful for pre-
 570 dictability and risk assessment (De Luca et al., 2019; Francis et al., 2020; Berkovic and Raveh-Rubin, 2022). From the time
 perspective, large values of $R(t)$ indicate the most stationary periods. A minimum threshold is often defined to separate sta-
 tionary from non-stationary periods: for instance, 2 days for weather regimes (De Luca et al., 2019; Francis et al., 2020) and
 extreme precipitation (Du et al., 2022), 5-6 days for warm spells (Berkovic and Raveh-Rubin, 2022; Rousi et al., 2022b), 5-25
 days for geopotential anomalies (Dole and Gordon, 1983), or 2 seasons for droughts (Ford and Labosier, 2014).

575 In the state space, the stationarity of any state $\mathbf{x}_0$ can be described from the distribution of its residence times $R_{\mathbf{x}_0} \sim$
 $\mathbb{P}\{R(t) \text{ s.t. } \mathbf{x}(t) = \mathbf{x}_0 \text{ and } \mathbf{x}(t-1) \neq \mathbf{x}_0\}$ (Figure 9). The easiest is to calculate the mean (Kysely and Domonkos, 2006; Kučerová
 et al., 2017; Richardson et al., 2019), maximum (Rousi et al., 2022b), or some extreme percentile (Zolina et al., 2013) of
 $R_{\mathbf{x}_0}$. Like autocorrelation, it is also possible to characterise stationarity by the dependence of $\mathbb{P}(R_{\mathbf{x}_0} = n)$ on n (Sharma and
 Panu, 2014). Light distribution tails indicate an exponential decay and short-term memory (see section 4.2.6), while heavy tails
 580 (e.g., power-law (Bunde et al., 2013) or q-exponential (Weber et al., 2019) distributions) point to long-range dependence in the
 system. For example, Pfeleiderer and Coumou (2018) fitted exponential models to the distribution of consecutive warm and cold
 spells to quantify temperature persistence in the Northern Hemisphere, and Huguenin et al. (2020) did the same for weather
 types over Central Europe. Many other types of distribution can also be fitted (e.g., Deni et al., 2010; Zolina et al., 2013).
 Residence times can also be modeled as functions of covariates. For instance, Röthlisberger et al. (2019) and Ali et al. (2021)
 585 apply a Weibull regression model to relate the duration of dry and wet spells to Rossby wave activity in the mid-latitudes.
 Finally, averaging $R(t)$ over time t , or $R_{\mathbf{x}_0}$ across all possible system states, provides a convenient and easily interpretable global
 stationarity metric (e.g., Pfeleiderer et al., 2019).

4.2.6 Transition probabilities

Residence times characterise how long the system remains in the same state $\mathbf{x}_0$. By contrast, transition probabilities P describe
 590 the likelihood that the state of the system does not change at the next time step:

$$P(\mathbf{x}_0) = \mathbb{P}(\mathbf{x}(t+1) = \mathbf{x}_0 \mid \mathbf{x}(t) = \mathbf{x}_0) \quad (16)$$

Transition probabilities can be further divided by marginal probabilities to yield probability ratios (e.g., Kolstad et al., 2015):

$$PR(\mathbf{x}_0) = \frac{P(\mathbf{x}_0)}{\mathbb{P}(\mathbf{x}(t) = \mathbf{x}_0)} \quad (17)$$

A high transition probability $P(\mathbf{x}_0)$ or $PR(\mathbf{x}_0)$ means that the system likely remains in $\mathbf{x}_0$, which indicates persistence (Figure
 595 10-a). Like residence times, transition probabilities can be computed on specific system states only, or averaged across all system
 states (according to their frequency) to yield a measure of global stationarity. Their statistical significance can be estimated
 by comparing observed transition probabilities to those obtained from a large set of randomly shuffled series. However, unlike
 residence times, transition probabilities cannot identify stationary periods: they only work in the state space.

Økland and Lejenäs (1987) applied this method to blocking stationarity; Ford and Labosier (2014), Wilby et al. (2016) and Moon
 600 et al. (2018) to droughts; Fereday (2017) to large-scale sea-level pressure patterns; Guilbert et al. (2015) to daily precipitation;
 and Röthlisberger and Martius (2019) to dry and warm periods (Figure 10-a).




Figure 9. PDF of the North Atlantic jet latitude index (solid) together with the weighted Gaussian PDFs from the HMM: the southern (dashed), northern (dotted), and central (dashed-dotted) regimes. Regime duration curves (southern (black), northern (red), and central (blue)), expressed as the frequency of occurrences lasting at least n days. Dashed curves show the corresponding 2.5% and 97.5% confidence levels. Reproduced with permission from Franzke et al. (2011).

It is often convenient to assume that the distribution of system states at $t + 1$ only depends on the system state at t , in other words to approximate the system with a first-order Markov chain model (Sericola, 2013). Under this assumption, transition probabilities are directly related to the distribution of residence times. Indeed, if the probability of remaining in state $\mathbf{x}_0$ does not depend on how long the system has previously been in state $\mathbf{x}_0$: $\mathbb{P}(\mathbf{x}(t+1) = \mathbf{x}_0 | \mathbf{x}(t) = \mathbf{x}_0) = \alpha$, then the probability that the residence time of $\mathbf{x}_0$, $R_{\mathbf{x}_0}$, exceeds n is equal to the probability of finding a sequence of at least n consecutive $\mathbf{x}_0$ values, specifically:

$$\mathbb{P}(R_{\mathbf{x}_0} \geq n) = \mathbb{P}_t(\mathbf{x}(t-1) \neq \mathbf{x}_0; \mathbf{x}(t) = \mathbf{x}_0; \mathbf{x}(t+1) = \mathbf{x}_0; \dots; \mathbf{x}(t+n-1) = \mathbf{x}_0) \quad (18)$$

where the subscript in $\mathbb{P}_t$ specifies that the probability is taken with respect to the time variable, whereas in $\mathbb{P}(T_{\mathbf{x}_0} \geq n)$ it is taken with respect to the ensemble of residence times for $\mathbf{x}_0$. Thanks to the first-order Markov assumption, equation (18) factors as

$$\mathbb{P}(R_{\mathbf{x}_0} \geq n) = \mathbb{P}_t(\mathbf{x}(t-1) \neq \mathbf{x}_0; \mathbf{x}(t) = \mathbf{x}_0) \times \prod_{k=0}^{n-2} \mathbb{P}_t(\mathbf{x}(t+k+1) = \mathbf{x}_0 | \mathbf{x}(t+k) = \mathbf{x}_0) \quad (19)$$

Thus,

$$\mathbb{P}(R_{\mathbf{x}_0} \geq n) = \beta \times \alpha^{n-1} \quad (20)$$




Figure 10. (a) Climatological May–October (MJJASO) hot spell survival probabilities (shaded) in ERA-Interim reanalysis data. Red contours show MJJASO cyclone frequencies of 30% and 40%, respectively. Reproduced with permission from Röthlisberger and Martius (2019). (b) Change in the odds ratio of drought occurrence in spring given a 1-unit increase in winter standardised precipitation index, estimated from logistic regression. Thick black contours indicate statistical significance with 95% confidence. Adapted with permission from Ford and Labosier (2014).

615 where $\beta = \mathbb{P}_t(\mathbf{x}(t-1) \neq \mathbf{x}_0; \mathbf{x}(t) = \mathbf{x}_0)$ does not depend on t for a stationary system, and

$$\mathbb{P}(R_{\mathbf{x}_0} = n) = \beta \alpha^{n-1} (1 - \alpha) \quad (21)$$

The logarithm of the distribution of residence times is therefore a linear function of n , with the slope ($\log(\alpha)$) equal to the transition probability. The Markov assumption is thus consistent with a short-term dependent system, in which residence time probabilities decay exponentially with n (see section 4.2.5 and Bunde et al. (2013)). For instance, Huguenin et al. (2020) fit
620 such exponential laws and use their slope as measures of weather regime stationarity over Central Europe in current and future climates. This approach is well-suited to comparing two different series (e.g., two different periods or locations) because it separates changes in the marginal frequency of state $\mathbf{x}_0$ (only affecting β) from changes in stationarity (only affecting α).

4.2.7 Time series modeling

Another possibility to investigate state stationarity is to fit a statistical model to the data that links successive system values with each other:

$$\mathbf{x}(t+1) \stackrel{d}{=} \mathcal{F}(\{\mathbf{x}(k)\}_{k=1..t}, \{\theta(t)\}_{k=1..t}) \quad (22)$$

where $\stackrel{d}{=}$ stands for equality in distribution and $\{\theta(t)\}_{k=1..t}$ are covariates. Transition probabilities (section 4.2.6), for example, are equivalent to modeling a system's evolution by a transition matrix between all possible system states. Hidden Markov Models (section 4.2.1) likewise estimate transition matrices between hidden states. Logistic regression can be used to assess stationarity in a given state. Several studies used it to quantify stationarity in droughts, using previous precipitation or soil moisture anomalies as predictor variables ($\theta(t)$ in equation (22)) (Ford and Labosier, 2014; Meng et al., 2017) (Figure 10-b). The link to drought occurrence at the previous time step is implicitly taken into account with previous precipitation or soil moisture anomalies. If previous system values are explicitly included as covariates to the model, one speaks of autologistic regression (Wolters, 2017).

4.3 Running window methods for episodic persistence

The running window approach identifies persistent periods of a given fixed length in continuous or categorical data. It assesses the degree of stationarity over fixed time intervals by means of a "similarity index". Given a time interval $\mathcal{T} = \{t_1, \dots, t_n\}$ of length n and a similarity metric $d(\cdot, \cdot)$, the similarity index $S_n(\mathcal{T})$ is defined as the average similarity between system values in $\mathcal{T}$. This similarity can be computed with reference to one time step $\mathbf{x}(t_k)$ in particular (typically the first one):

$$S_n(\mathcal{T}) = \frac{1}{n-1} \sum_{\substack{1 \leq k' \leq n \\ k' \neq k}} d(\mathbf{x}(t_{k'}), \mathbf{x}(t_k)), \quad (23)$$

or it can measure the average similarity across all pairs of values:

$$S_n(\mathcal{T}) = \frac{2}{n(n-1)} \sum_{\substack{1 \leq k, k' \leq n \\ k \neq k'}} d(\mathbf{x}(t_k), \mathbf{x}(t_{k'})) \quad (24)$$

Different time intervals can then be ranked according to their degree of stationarity. Mo and Ghil (1987), for instance, use pattern correlation as similarity metric and set a threshold of 0.5 to separate stationary from non-stationary periods. Another possibility is to keep the N periods with the largest $S_n(\mathcal{T})$ (i.e., to use a percentile-based threshold). Stationary states can then be detected with e.g., pattern recognition techniques applied to the most stationary periods (Mo and Ghil, 1987). We show an example for summertime European circulation in Figure 11. The most common are blocking-type patterns over Northern Europe/Western Russia (Figure 11-a,b,c), but we also find two pronounced zonal patterns with a southward/northward shifted jet (Figure 11-d,e).

Averaging $S_n(\mathcal{T})$ values across multiple periods of length n can additionally provide for a measure of global stationarity, as did Hoffmann et al. (2021) to analyse 10-day persistence in summer atmospheric circulation. When working with categorical data,

similarity indices can be averaged for each system state to highlight the most stationary ones (Horel, 1985b).

The running window approach is well-suited to compute temporal trends in stationarity, since the similarity index can be defined at all time steps (Hoffmann et al., 2021). One limitation, however, is that results for the same system but calculated for different
655 period lengths are not directly comparable since the marginal distribution of $S_n(\mathcal{T})$ depends on n . Furthermore, $S_n(\mathcal{T})$ only measures the average similarity between successive values. A high $S_n(\mathcal{T})$ does not guarantee high stationarity; there could be breaks in between sequences of similar values. This is especially relevant for long periods during which strict persistence is unlikely to occur.

While we try to give a comprehensive view of common methods used in stationarity analysis, we could not include all the
660 methods that exist in the literature. Kornhuber and Tamarin-Brodsky (2021), for instance, define stationarity in a Lagrangian context as the zonal velocity of individual weather systems (in their case, localised temperature anomalies), while Hoskins and Woollings (2015) use Rossby wave phase speed as a proxy for weather stationarity in the mid-latitudes. Finally, in the framework of dynamical systems theory, stationarity can also be equated to the system remaining for some period of time in a small subset of the state space. During that period, the system may explore different configurations, but fewer ones than if it had been able
665 to evolve freely across the whole state space (Fereday, 2017). In this sense, persistent states are states from which the system takes (statistically) a long time to reach the rest of its state space. This perspective relates to the intransitivity theory of Lorenz (1990) which postulates that the evolution of the atmospheric circulation on S2S timescales can, under certain conditions, be governed by a few well-separated attractors, each with their own preferred states (Weiland et al., 2021).

Method	Section	Type	Definition	Features	Main limitations
Autocorrelation	4.1.1	Global	Measure of strength of linear dependence between successive system values; equation (1)	Lag-1 autocorrelation frequently used, but can also highlight relevant timescales of self-dependence in a series	Requires 1D data; characterises the strength of the dependence, not "similarity" per se
Dependence parameter	4.1.2	Global	Asymptotic characterisation of how autocorrelation evolves with time lag	Discriminates short- and long-range dependence	Requires 1D data; based on autocorrelation which characterises the strength of the dependence, not "similarity" per se
Hurst exponent	4.1.2	Global	Asymptotic characterisation of how cumulative fluctuations of a series around its mean evolve with timescale	Simple metric to identify stationary/anti-stationary/noise-like behaviour in time series	Requires 1D data
Power spectrum exponent	4.1.2	Global	Asymptotic characterisation of the distribution of time series variability across timescales	Relies on the power spectrum, for which many methods are available	Requires 1D data; power spectrum linked to autocorrelation which characterises the strength of the dependence, not "similarity" per se
Weather regimes	4.2.1	State	Common method to identify patterns accounting for high variability in multi-dimensional series	Many techniques available; unsupervised method; often good physical interpretation	Requires choosing a number of regimes; strong simplification of the state space, i.e. rare yet relevant system states may be missed; indirect link to persistence, which requires further analysis

Quasi-stationary states	4.2.2	State	Identifies stationary system states (time derivative ≈ 0)	Unsupervised method; best applied to low-dimension systems (dimension reduction required)	Challenging implementation; true number of states can be difficult to assess; stationarity constraint is strong (zero tendency) and highly local in time; no guarantee that resulting states are "physical"
Optimally persistent patterns	4.2.3	State	Identifies most autocorrelated patterns in multi-dimensional series and associated timescales	Unsupervised method; relevant for forecasting	Implementation can be challenging; limited number of resulting states and no guarantee that they are "physical"
Dynamical systems approach	4.2.4	State	Defines a persistence index for every point of a system's trajectory based on local residence times	Unsupervised method, well-grounded in mathematical theory; persistence index directly interpretable; works with continuous multi-dimensional data	Based on extreme value approximations; the metric is consequently highly local in time, and may not be relevant at S2S timescales
Residence times	4.2.5	Global/ State/ Episodic	Defined as the period during which a system remains similar to its initial value	Versatile and easy to interpret approach; residence times can be calculated across time or system states, on continuous or categorical data	Potentially difficult to summarise the distribution of residence times for a given system state
Transition probabilities	4.2.6	State	Probability that a system remains in the same state at the next time step	Easy to calculate and interpret; directly related to residence times under the Markov assumption	Requires categorical data; limited to one-step-ahead stationarity
Time series modeling	4.2.7	State	Statistical modeling of successive system values	Potentially very versatile approach that can include the effect of external covariates	Models may capture "dependence" rather than "similarity", and can quickly become complex and difficult to interpret

Running window methods	4.3	Episodic	Defines a stationarity value for all time steps at a given timescale	Results can be used to identify stationary states	Relative stationarity metric only; difficult to compare dif- ferent systems
---------------------------	-----	----------	--	--	---

Table 1: Overview of stationarity methods: method name; corresponding section; type of persistence it applies to; brief definition; main features; main limitations.




Figure 11. (a-e) Most persistent 10-day averaged Z500 anomaly patterns during JJA, obtained by clustering the top 10% (40) 10-day patterns with the highest similarity index (equation (24)). Similarity is measured with the SSIM index. The clustering algorithm is Partitioning Around Medoids (PAM). The associated 200 hPa zonal wind anomalies are shown by contour lines (same levels as Z500). The frequency of each pattern is also indicated. (f-j) 10-day averaged surface temperature anomalies associated with each persistent Z500 pattern.

5 Recurrence

We now discuss methods to capture recurrence in atmospheric data (see Table 2 for an overview). In contrast to stationarity, most methods here only quantify recurrence in a specific system state. They typically require binary time series as input (where "1" represents an occurrence of the system state or event of interest) and count event occurrences over time. Recurrence is therefore always defined for a given state or event, usually with the potential for large impacts like tropical (Mumby et al., 2011) and extra-tropical cyclones (Mailier et al., 2006; Vitolo et al., 2009), windstorms (Khare et al., 2015), or extreme precipitation episodes (Kopp et al., 2021). For this reason, we did not explicitly divide this section into global, state and episodic methods. Common methods are available to characterise episodic persistence (section 5.1.1) and the persistence of given states (sections 5.1.2-5.2.1). Few definitions of "global recurrence" exist, however. Additionally, no study has, to our knowledge, tried to analyse S2S recurrence in a "non-supervised" way, i.e. that would allow to objectively identify recurrent states. In section 5.2.2, we discuss the potential of recurrence plots to fill that gap.

Because most recurrence methods work directly with binary time series, point processes are a convenient theoretical tool in recurrence analysis. Point processes are a class of probability models for the random occurrence of points in a space (one- or multi-dimensional). The most simple hypothesis than can be made is that of complete serial randomness, i.e., points occur completely independently of one another. In this case, recurrence occurs by chance only. The homogeneous Poisson point process is a simple model without memory for binary series in which events occur without correlation and with a constant intensity. In a homogeneous Poisson process, the number of points in individual time intervals are independent, and the number of points in an interval of length τ is Poisson distributed with rate $\lambda\tau$, where λ is the process intensity rate. This model is useful to simulate binary series with complete serial randomness, and therefore provides an empirical basis to assess the statistical significance of various recurrence metrics in observations.

Separating distinct occurrences of a state, pattern or event is critical to distinguish recurrence from stationarity. Over short periods, like a season, it is in principle possible to visually separate multiple systems occurring in close succession, for instance with Hovmoeller diagrams (e.g., Tuel et al., 2022a; Rousi et al., 2022a). Most of the time, however, separation cannot be done by hand, and objective criteria are required. One option are physically-based detection/tracking algorithms – e.g., for cyclones (Hodges, 1995; Wernli and Schwierz, 2006) or atmospheric blocks (Schwierz et al., 2004; Steinfeld, 2021) – that can separate distinct weather systems. A second option is to use a minimum duration between events to classify them as distinct, with the choice depending on the underlying physical system or on impact considerations. Barton et al. (2016), Kopp et al. (2021) and Tuel and Martius (2021a), for instance, require a minimum of 2 days with non-extreme precipitation between extreme precipitation episodes (a typical timescale for extratropical cyclones). In Pfleiderer et al. (2019), a single day is used to separate distinct dry or warm periods.




Figure 12. (a) Example of a time series of extreme precipitation events (black dots) and 21-day periods containing more than 2 events (gray shading). Adapted from Kopp et al. (2021) under the terms of the Creative Commons Attribution 4.0 License. (b) Map of dispersion statistic ψ (equation (25)) of DJF cyclones over 1975–2004, averaged across an ensemble of CMIP5 models. Reproduced from Bevacqua et al. (2020) under the terms of the Creative Commons Attribution 4.0 license.

5.1 Diagnostic methods

5.1.1 Window counts

A simple way to identify recurrent periods is to look for periods with high event counts. One possibility is to set a time window and require the number of event counts during this window to exceed some threshold (2 or higher; Figure 12-a). Kopp et al. (2021) and Tuel and Martius (2022a) take this approach to analyse recurrent extreme precipitation episodes at the 2- and 3-week timescales. Pinto et al. (2014) likewise identify cyclone clusters over the British Isles as periods with 4 or more consecutive
 705 cyclones within 7 days. Another possibility is to look for clusters of events in which each event is separated by at most n time steps from the preceding one. For instance, Bevacqua et al. (2020) define cyclone clusters as sequences of $N \geq 2$ consecutive cyclones transiting within a given area and separated by 24 hours or less. This method is largely impact-driven and does not aim at modeling the binary series, assessing the significance of the clustering or calculating return periods for large event counts (Dacre and Pinto, 2020). For this, further steps must be taken.

5.1.2 Dispersion metrics

Dispersion metrics characterise the distribution of event occurrences in a time series. In a series with no tendency to clustering, high event counts over a given duration should be much less frequent than in a series where clustering is prevalent. If events occur with the same average probability in the two series, then in the clustered series, sequences with no events should mechanically be more frequent, and the variance in event counts will be higher. In a homogeneous Poisson process (see above), $N_k(\tau)$, event
 715 counts in disjoint intervals of length τ (indexed by k), follow a Poisson distribution with parameter $\lambda\tau$. Hence, the expected

value $\mu(\tau)$ and variance $\sigma(\tau)^2$ of $N_k(\tau)$ are both equal to $\lambda\tau$. Two common statistical dispersion metrics quantify deviations from this homogeneous behaviour. The dispersion statistic is a measure of temporal correlation over different timescales. It is defined as (Mailier et al., 2006)

$$\psi(\tau) = \frac{\sigma(\tau)^2}{\mu(\tau)} - 1 \quad (25)$$

720 ψ is related to the Fano Factor ($FF(\tau) = \psi(\tau) + 1$) (Telesca, 2007), also called index of dispersion (Mailier et al., 2006). A positive $\psi(\tau)$ ($\sigma(\tau)^2 > \mu(\tau)$) indicates over-dispersion: events tend to occur in clusters. A negative $\psi(\tau)$ points to under-dispersion: events occur more regularly than in a random process. Near-zero values are consistent with a homogeneous Poisson process (Figure 12-b).

Similarly, the Allan Factor (AF) is defined as the variance of successive event counts over an interval of length τ divided by
725 twice the average event count in τ steps (Telesca, 2007):

$$AF(\tau) = \frac{\mathbb{E}[(N_{k+1}(\tau) - N_k(\tau))^2]}{2\mu(\tau)} \quad (26)$$

For a homogeneous Poisson process, $AF(\tau) = 1$, while $AF(\tau) > 1$ indicates clustering behaviour. ψ and AF can be evaluated for a single window τ^* , but assessing how they evolve with τ helps better detect recurrent behaviour in the series (Serinaldi and Kilsby, 2013). In the absence of clustering, $\psi(\tau)$ and $AF(\tau)$ remain constant with τ . In clustered processes, however, both
730 metrics scale with τ , often with a power-law behaviour over some range $[\tau_1, \tau_2]$ which can be described by an exponent (Telesca, 2007), similar to the spectral analysis of persistence (section 4.1.2). One advantage of ψ and AF is that they are insensitive to the underlying event frequency, which makes them convenient to compare different locations or event identification algorithms with each other (Dacre and Pinto, 2020). However, the corresponding null hypothesis – that the absence of recurrence is equivalent to exponentially distributed event return times under the Poisson assumption – is not always relevant, and other approaches may
735 be better suited to detect deviations from memoryless processes (Blender et al., 2015).

The statistical significance of $\psi(\tau)$ and $AF(\tau)$ can be assessed empirically through Monte-Carlo sampling of random series without memory, or analytically. Mailier et al. (2006) discuss a chi-square test for $\psi(\tau)$ (for a fixed window τ), while Serinaldi and Kilsby (2013) introduce the sampling distribution of the empirical AF estimator. Note that seasonality in event counts can artificially inflate the dispersion statistic, so care must be taken either to remove the seasonality (e.g., Tuel and Martius, 2021a),
740 or to simulate random processes with the same seasonality as the observed signal (λ function of the month, for instance). Note also that $\psi(\tau)$ and $AF(\tau)$ both involve $\mu(\tau)$ in the denominator; in other words, these metrics are normalised by the marginal event frequency. The same $\psi(\tau)$ can be obtained for series with different event frequencies, which makes dispersion metrics less relevant for impacts than other methods (e.g, window counts; section 5.1.1).

Since its introduction by Mailier et al. (2006), the dispersion statistic has been frequently applied in recurrence analysis, namely
745 to detect temporal/serial clustering in cyclones, both tropical (Mumby et al., 2011; Wolff et al., 2016) and extratropical (Vitolo et al., 2009; Pinto et al., 2013, 2014; Economou et al., 2015; Pinto et al., 2016). The Allan factor has been applied to daily precipitation series (Serinaldi and Kilsby, 2013), to wave storms (Besio et al., 2017) and to extreme wind speeds in Switzerland (Telesca et al., 2020).

Alternative metrics that characterise the distribution of event counts have been proposed in the literature. For example, Kopp et al. (2021), working with extreme precipitation counts over 2- to 4- week timescales, proposed a new metric, $S_{cl}(\tau)$ – which they named the "clustering metric" – based on a weighted sum of window event counts. Like the dispersion statistic and Allan factor, $S_{cl}(\tau)$ requires choosing a window size τ . Clustering periods of length τ are then identified and ranked by applying a moving window sum to the binary series. The period with the most events is selected first; the corresponding events are removed from the series (to avoid any intersection between clustering periods); and the procedure continues until a minimum number of events is obtained, or until no more clustering periods (with at least two events) are found. This yields a set of K periods with decreasing event counts $n_k(\tau)$: $n_1(\tau) \geq n_2(\tau) \geq \dots \geq n_K(\tau)$. Kopp et al. (2021) then define their clustering metric by:

$$S_{cl}(\tau) = \sum_{1 \leq k \leq K} n_k(\tau) q_k \quad (27)$$

where q_k are weights that can be defined in various ways, as long as they verify (i) that q_k decreases with k (periods with fewer events are given less weight) and (ii) that $q(i) - q(i+1) > q(i+1) - q(i+2)$. In the end, this approach is equivalent to a scoring system for clustering periods, and the resulting metric $S_{cl}(\tau)$ correlates well with the dispersion statistic.

5.1.3 Ripley's K function

Given a system state $\mathbf{x}_0$, Ripley's K function measures the average number of occurrences of $\mathbf{x}_0$ within a given neighbourhood around a random occurrence of $\mathbf{x}_0$. It was originally developed for spatial data, but it can be applied in a temporal setting to characterise recurrence in $\mathbf{x}_0$. Let the binary variable $Y_t = \mathbb{1}\{X_t \text{ is in state } \mathbf{x}_0\}$; Ripley's K function for a neighbourhood of size n is then equal to (Ripley, 1981; Dixon, 2014):

$$K(n) = \lambda^{-1} \left(\mathbb{E} \left[\sum_{k=-n}^n Y_{t+k} \mid Y_t = 1 \right] - 1 \right) \quad (28)$$

where $\lambda = \mathbb{E}[Y_t]$ (frequency of $\mathbf{x}_0$ in X_t). The statistical significance of $K(n)$ can then be assessed by comparing it to values obtained for a homogeneous Poisson process of same length and with the same average rate of occurrence as Y_t (Figure 13). In equation (28), λ^{-1} can also be discarded, so that $K(n)$ represents an average event count and is more directly related to impacts. Ripley's K was used in meteorology to characterise S2S recurrence in extreme precipitation events by Barton et al. (2016), Tuel and Martius (2021a) and Tuel and Martius (2021b). Stephenson et al. (2004) and Hannachi (2010) also used Ripley's K to characterise the clustering of climate modes in the state space.

5.1.4 Distribution of inter-event times

Stationarity in a given system state $\mathbf{x}_0$ can be described by the corresponding distribution of residence times (section 4.2.5). Similarly, the distribution of inter-event times (also called "recurrence times" or "waiting times") characterises the recurrence in $\mathbf{x}_0$ from a global perspective (Altmann and Kantz, 2005). The recurrence time T_i is the duration between the i -th and $(i+1)$ -th occurrences of state $\mathbf{x}_0$ in a time series. We can easily see that both short and long recurrence times will be more frequent in a series that exhibits clustering than in a series where events occur independently of one another. The presence of a heavy tail in




Figure 13. (a) Assessing the significance of clustering with Ripley’s K function: conceptual example of Ripley’s K values on an observed series (black crosses) and the corresponding 95% uncertainty range obtained from random simulations of series with no clustering (blue shading). In this case, observations show significant clustering at the 10-20 day timescale. Reproduced from Tuel and Martius (2021a). (b) Map of 20-day Ripley’s K values for catchment-averaged extreme precipitation during fall in Switzerland. Hatching indicates statistical significance at the 95% level. Adapted from Tuel and Martius (2021b) under the terms of the Creative Commons Attribution 4.0 license.

the distribution of T_i therefore points to a tendency for temporal clustering in the series. If $\mathbf{x}_0$ is rare enough, the distribution of T_i may converge to well-known distributions like the exponential (a sign of short-term dependence, as with homogeneous Poisson processes), or the stretched exponential or power-law distribution (Eichner et al., 2007; Corral, 2015). When events correspond to extremes above a given threshold, the exponents that characterise these distribution are even related to the autocorrelation coefficient of the original series (Santhanam and Kantz, 2008; Kalra and Santhanam, 2021).

5.2 Stochastic modeling of recurrence

5.2.1 Event series

We have so far discussed diagnostic methods that characterise recurrence in binary time series. A different approach consists in fitting point processes to the series. A point process is a random process representing the occurrence times of specific events. It can be characterised by event times $T(n)$ (time of the n -th event in the series) or by the distribution of event counts $N(t_1, t_2)$ for any interval $[t_1, t_2]$. We can define its *intensity* or *hazard* function $\lambda(t)$ by

$$\lambda(t) = \lim_{\Delta t \rightarrow 0^+} \frac{1}{\Delta t} \mathbb{P}(N(t, t + \Delta t) = 1) \quad (29)$$

and, for any $t_1 < t_2$, its *intensity measure* $\Lambda(t_1, t_2)$ as

$$\Lambda(t_1, t_2) = \int_{t_1}^{t_2} \lambda(t) dt \quad (30)$$

Many different point processes have been described in the statistics literature (e.g., Cox and Isham, 1980), but clustering analyses in weather/climate studies have mainly relied on Poisson processes. In a Poisson process, $\lambda(t)$ is a deterministic function of time. If $\lambda(t)$ is constant, the process is said to be homogeneous; otherwise, the process said to be inhomogeneous. For any $t_1 < t_2$, $N(t_1, t_2)$ follows a Poisson distribution with rate $\Lambda(t_1, t_2)$:

$$N(t_1, t_2) \stackrel{d}{=} \text{Poisson}(\Lambda(t_1, t_2)) \quad (31)$$

and event numbers in disjoint intervals are independent from each other.

The point process approach has mainly been used to relate recurrence in extreme events to atmospheric or climate variability. To that end, $\lambda(t)$ is assumed to depend on given time covariates $X(t)$. A linear dependence corresponds to Poisson Generalised Linear Models (GLMs):

$$\lambda(t) = \beta \cdot X(t) \quad (32)$$

For instance, Villarini et al. (2011), Tuel and Martius (2022a) and Tuel and Martius (2022b) used Poisson GLMs to relate temporal clustering in heavy precipitation to large-scale modes of climate and atmospheric variability. Mailier et al. (2006) did the same for extratropical cyclones in the North Atlantic. The linear constraint can be lifted by using General Additive Models, or GAMs, in which the relationship between $\lambda(t)$ and the covariates is specified by flexible smooth functions (Hastie, 1992). Villarini et al. (2013) used GAMs to relate flood recurrence in the US Midwest to modes of climate variability like ENSO, and Barton et al. (2022) used GAMs to link temporal clustering in extreme precipitation across Europe to weather regimes in the North Atlantic. Another extension of the Poisson GLM is the Cox regression model (Smith and Karr, 1986), in which

$$\lambda(t) = \lambda_0(t) \exp(\beta \cdot X(t)) \quad (33)$$

where $\lambda_0(t)$ is a baseline intensity/hazard function and $\exp(\beta \cdot X(t))$ accounts for the time-varying effects of the selected covariates. Examples include Villarini et al. (2013), Mallakpour et al. (2017) and Yang and Villarini (2019), who applied Cox regression to recurrence in floods and extreme precipitation. More complex models are also possible; for instance, Khare et al. (2015) introduced a Poisson mixture model for windstorm clustering, in which $\lambda(t)$ is expressed as a stochastic function of time modulated by a gamma distribution (Khare et al., 2015). Note that models can be fitted directly by maximising the likelihood function (Yang and Villarini, 2019) or indirectly by working with window event counts (Mailier et al., 2006; Tuel and Martius, 2022a; Barton et al., 2022).

5.2.2 Recurrence plots

All previously discussed methods share the same limitation: they require binary time series as input, meaning that they are designed to characterise recurrence only in a given system state. It is not possible with such methods to explore recurrence




Figure 14. Effect of Euro-Atlantic zonal regime occurrence (corresponding 500 hPa geopotential mean and anomalies shown on left panel) on the temporal clustering of extreme precipitation at the three-week timescale during winter (measured as a probability multiplicative factor). Reproduced from Barton et al. (2022) under the terms of the Creative Commons Attribution 4.0 license.

in an "unsupervised" way, i.e., to automatically detect recurrent system states or sequences of states, and associated recurrent periods. Recurrence plots (RPs) may offer a solution to this problem.

An RP is a graphical representation of a system's self-similarity with time (Marwan et al., 2007). Mathematically, it is defined as a two-dimensional binary matrix $\mathbf{R} \in \{0, 1\}^{N \times N}$, with time on both axes, that consists of pairwise comparisons of system

825 values:

$$\mathbf{R}(i, j) = \Theta(\epsilon - d(\mathbf{x}(t_i), \mathbf{x}(t_j))) \quad (34)$$

with $d(\cdot, \cdot)$ a distance or similarity metric (by convention, $d(\mathbf{x}(t_i), \mathbf{x}(t_i)) = 0$), Θ the Heaviside step function, and ϵ a similarity threshold. If $d(\mathbf{x}(t_i), \mathbf{x}(t_j)) < \epsilon$, then $\mathbf{x}(t_i)$ and $\mathbf{x}(t_j)$ are similar and $\mathbf{R}(i, j) = 1$ represents a recurrence of the system. By definition, $\mathbf{R}$ is symmetric and $\forall 1 \leq i \leq N$, $\mathbf{R}(i, i) = 1$. A conceptual example is shown on Figure 15-a.

830 Before calculating $\mathbf{R}$, the system $\mathbf{x}$ can first be reduced to a lower-dimensional space (as in section 4.2.2), by e.g., projecting $\mathbf{x}$ onto a set of principal components (Marwan et al., 2007; Mukhin et al., 2022). Note that $\mathbf{R}$ depends on the selected similarity metric and on ϵ . For a given system $\mathbf{x}$, $\mathbf{R}$ is thus not uniquely defined. Choosing a value for ϵ should be done carefully. ϵ should be small enough to account for "real" recurrences in the data, but not too small to have enough recurrences to analyse. Several rules of thumb have been proposed in the literature (see Marwan et al. (2007) and references therein). For instance, ϵ can be

835 selected by requiring an average recurrence rate of 1-10%. For weather and climate data, ϵ could also be selected based on physical considerations and expert judgment.

RPs are a convenient way to visualise a system's trajectory, especially for multi-dimensional systems (like circulation fields). RPs capture persistent behaviour: vertical lines indicate stationarity and diagonal lines (outside of the main diagonal) indicate recurrence. RPs can thus highlight periods when the trajectory of a system roughly visits the same sequence of states or parts

840 of the state space.

Several measures have been proposed to quantify the presence of specific patterns in RPs. They are known collectively as

"recurrence quantification analysis" (Marwan, 2008). Common ones include the recurrence rate:

$$RR = \frac{1}{N^2} \sum_{1 \leq i, j \leq N} \mathbf{R}(i, j) \quad (35)$$

the determinism, which quantifies the fraction of recurrence points forming diagonal lines of minimum length l_{min} :

$$845 \quad DET = \frac{\sum_{l \geq l_{min}} l D(l)}{\sum_{l \geq 1} l D(l)} \quad (36)$$

where $D(l)$ is the number of diagonal lines of length l , and the laminarity, which quantifies the fraction of recurrence points forming vertical lines of minimum length l_{min} :

$$LAM = \frac{\sum_{l \geq l_{min}} l V(l)}{\sum_{l \geq 1} l V(l)} \quad (37)$$

where $V(l)$ is the number of vertical lines of length l . All three can be used as global measures of persistence (e.g., Ramirez-
850 Amaro and Figueroa-Nazuno, 2006). However, these indices do not a priori discriminate between stationarity and persistence (high values of RR , for instance, could be due to either). To do so would require making sure that similar time steps are separated by "0"s in the RP (Figure 15-b).

RPs can also detect local recurrence in time. With RPs, one can extend the counting approach introduced in section 5.1.1 to
855 multiple-day sequences of complex multi-dimensional data. Specifically, to search for sequences of p steps that recur over periods of q steps ($0 < p < q$), we can simply define a recurrence index $RI_{p,q}(t)$ as the number of diagonal lines of length p in the sub-matrix of $\mathbf{R}$ indexed by $[t, t+p] \times [t, t+q]$ (Figure 15-b). To make sure that distinct occurrences of the same sequence are separated, we can require diagonals to be separated by a minimum number of days (2 in Figure 15-b). It is also possible to allow for short breaks (e.g., 1 step) in the diagonals.

RPs have recently gained attention in a range of disciplines that deal with complex systems (Goswami, 2019), though only
860 rarely in environmental sciences. Ramirez-Amaro and Figueroa-Nazuno (2006) investigated the recurrence properties of major teleconnection patterns with RPs. Yiou et al. (2018) indirectly relied on RPs to identify the most recurrent states of North Atlantic circulation at the intra-seasonal timescale. Adeniji et al. (2018) used RPs to analyse the recurrence characteristics of hourly wind speed in Nigeria and Ray et al. (2019) to investigate daily temperature and humidity data across India. Recently, Mukhin et al. (2022) used RPs to detect weather regimes in the Northern Hemisphere. On Figure 16, we show results obtained
865 for 3-day sequences of similar upper-level circulation patterns over Europe in summer. We calculate $RI_{p,q}(t)$ with $p = 3$ days and $q = 21$ days and identify the two most common sequences among the ≈ 30 21-day windows with $RI_{p,q}(t) \geq 2$. The method identifies recurring blocks and cyclones over Scandinavia as the most frequent recurring sequences of circulation patterns. The results appear physically meaningful since counts of individual blocks and cyclones during the corresponding windows range from 1-2 (for blocks) and 4-11 (for cyclones). More accurate patterns could possibly be obtained by choosing more than 2
870 clusters, but the results of Figure 16 are only meant to illustrate the potential of the method.




Figure 15. Detecting recurrence with recurrence plots. (a) Conceptual example of a recurrence plot. The red rectangle highlights a period during which the system exhibits recurrence with 3 successive similar 4-day sequences occurring over a 12-day period (highlighted in red). (b) Zoom over the recurrence period with the three separate 4-day sequences highlighted by green dashed lines.




Figure 16. Two major 3-day recurring sequences of atmospheric circulation (Z500) over Europe during summer obtained by clustering the ≈ 30 21-day windows with $RI_{3,21}(t) \geq 2$ with the PAM algorithm. Stippling (respectively hatching) indicates atmospheric blocking (respectively cyclone) frequency anomalies larger than 30%. Similarity is measured with the SSIM index, with a similarity threshold ϵ in equation (34) of 0.25 (corresponding to a recurrence rate of $\approx 5\%$). Note that successive panels (a, b, c, and d, e, f) are 1-day apart.

Method	Section	Type	Definition	Features	Limitations
Window counts	5.1.1	State	Absolute event counts over windows of length τ	Simple, impact-driven approach; requires a fixed timescale	Basic characterisation of recurrence (e.g., no statistical significance)
Dispersion metrics	5.1.2	State	Characterise the degree of deviation in the distribution of event counts at a fixed timescale from homogeneous Poisson series	Statistical approach to recurrence	Not related to marginal event frequency; seasonality can make statistical significance assessment difficult
Ripley's K	5.1.3	State	Counts the average number of events in the neighbourhood of a given event	Statistical approach to recurrence; simple visualisation of recurrence across timescales; direct link to event counts (impacts)	Seasonality can make statistical significance assessment difficult
Inter-event times	5.1.4	State	Asymptotic characterisation of the distribution of inter-event times	Statistical approach to recurrence; discriminates short- and heavy tailed distributions where short-tailed ones are consistent with no-memory processes (recurrence occurs by chance)	Physical interpretation may be complex
Event count modeling	5.2.1	State	Fit a statistical model to $N(\tau)$ (event counts over windows of length τ), e.g. Poisson model	Requires a fixed timescale τ ; allows to model the effects of covariates on recurrence	Requires statistical assumptions on the distribution of event counts
Recurrence plots	5.2.2	Global/ State/ Episodic	2D binary matrix $\mathbf{R}$ describing system self-similarity at all time steps: $\mathbf{R}(i, j) = 1$ iff $d(\mathbf{x}(t_i), \mathbf{x}(t_j)) \leq \epsilon$; several global or local metrics can be defined	Requires a similarity threshold (ϵ); very versatile approach that can be applied to any kind of data	Can be complex to implement and may require custom metrics depending on the application

Table 2. As Table 1, but for recurrence methods.

Summary and outlook

Weather persistence on S2S timescales has been a topic of research since the early days of meteorology. Stationary or recurrent behaviour are common features of weather dynamics, and are strongly related to fundamental physical processes, weather predictability and surface weather impacts. Studying weather persistence is therefore important for theoretical as well as practical reasons. One challenge is that persistence remains a very broad concept that relates to different behaviours in dynamical systems. We propose a typology / structure for the broad concepts related to persistence. Namely that persistence is used to describe the average behaviour of a system, across its whole trajectory (global persistence); sometimes to refer to specific segments of this trajectory (episodic persistence); and sometimes to qualify the behaviour of particular system states (state persistence).

A wide range of methods have been introduced in the literature to describe persistence in weather and climate series, and several exist for each type of persistence. They offer many distinct and often complementary perspectives on persistence. Some methods quantify persistence in a statistical framework, which can be very relevant for weather forecasting. Others focus instead on persistent periods, including "statistical flukes", a useful approach for risk assessment but which says nothing about the overall behaviour of the series. Other methods yet aim at identifying which weather patterns tend to be more persistent than others.

The diversity of existing methods presented in this review reflects the fact that "persistence" is a multi-faceted concept. While we can agree on a general definition of the concept, many options exist when it comes to actually quantify persistence in real-world data. Though different methods may be related, each sheds light on a particular aspect of persistence. What is meant by "persistence" can not be dissociated from the metric used to quantify it. The choice of method should be guided by the end goal, whether process understanding, risk/impact assessment or predictability. Future research should nevertheless perhaps consider testing more systematically the robustness of results to the choice of persistence metric. This could matter particularly to better characterise potential trends in persistence under climate change and their associated impacts. As a final note, while we centred our review on S2S persistence, most of the methods we discussed apply in principle to other timescales, be they sub-daily or multi-decadal. What really differs is how to define the system to analyse (daily time steps may not be relevant for inter-annual variability, for instance) and how to interpret persistence.

Code availability. R code implementing several of the methods presented in this paper is available at <https://github.com/Quiriosity129/persistence>. Code to calculate the dynamical system metrics (section 4.2.4) is available for R at <https://github.com/thaos/dtheta> or from the R package *CSTools* (Perez-Zanon and coauthors, 2022), and for Python at <https://github.com/yrobink/CDSK>. The Structural Similarity Index (SSIM) can be calculated with the Python package *scikit-image* (van der Walt et al., 2014).

Appendix A: Data

We illustrate some of the methods (Figures 3, 6, 8, 11 and 16) with data from the ERA5 reanalysis (Hersbach et al. (2020); available from <https://dx.doi.org/10.24381/cds.bd0915c6>). We use geopotential height at 500 hPa and 2-meter temperature

for the months of June, July and August, and over the 1979-2020 period. We remove their seasonality and long-term trends by normalising the data with a moving 30-day, 7-year window (as in Pfeiderer and Coumou (2018)). The calculation of the blocking and cyclone indices in Figure 16 is described in Tuel and Martius (2022a).

905 *Author contributions.* AT: Conceptualisation; Formal analysis; Visualisation; Software; Writing - original draft.

OM: Conceptualisation; Funding acquisition; Supervision; Writing - review & editing.

Competing interests. O.M. serves as editor of Earth System Dynamics. The authors declare no further competing interests.

Acknowledgements. O.M. acknowledges support from the Swiss Science Foundation (SNSF) grant number 178751.

References

- Adeniji, A. E., Olusola, O. I., and Njah, A. N.: Comparative study of chaotic features in hourly wind speed using recurrence quantification analysis, *AIP Advances*, 8, 025 102, <https://doi.org/10.1063/1.4998674>, 2018.
- Ali, S. K., Aydam, Z. M., and Rashed, B. M.: Similarity metrics for classification: A Review, *IOP Conference Series: Materials Science and Engineering*, 928, 032 052, <https://doi.org/10.1088/1757-899X/928/3/032052>, 2020.
- Ali, S. M., Martius, O., and Röthlisberger, M.: Recurrent Rossby Wave Packets Modulate the Persistence of Dry and Wet Spells Across the
915 Globe, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL091452>, 2021.
- Ali, S. M., Röthlisberger, M., Parker, T., Kornhuber, K., and Martius, O.: Recurrent Rossby waves during Southeast Australian heat-waves and links to quasi-resonant amplification and atmospheric blocks, *Weather and Climate Dynamics Discussions*, 2022, 1–41, <https://doi.org/10.5194/wcd-2022-1>, 2022.
- Altmann, E. G. and Kantz, H.: Recurrence time analysis, long-term correlations, and extreme events, *Phys. Rev. E*, 71, 056 106,
920 <https://doi.org/10.1103/PhysRevE.71.056106>, 2005.
- Ault, T. R., Cole, J. E., Overpeck, J. T., Pederson, G. T., and Meko, D. M.: Assessing the Risk of Persistent Drought Using Climate Model Simulations and Paleoclimate Data, *Journal of Climate*, 27, 7529 – 7549, <https://doi.org/10.1175/JCLI-D-12-00282.1>, 2014.
- Barnes, E. A. and Hartmann, D. L.: Dynamical Feedbacks and the Persistence of the NAO, *Journal of the Atmospheric Sciences*, 67, 851–865, <https://doi.org/10.1175/2009JAS3193.1>, 2010.
- 925 Barnston, A. G. and Livezey, R. E.: Classification, Seasonality and Persistence of Low-Frequency Atmospheric Circulation Patterns, *Monthly Weather Review*, 115, 1083–1126, [https://doi.org/10.1175/1520-0493\(1987\)115<1083:CSAPOL>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<1083:CSAPOL>2.0.CO;2), 1987.
- Barton, Y., Giannakaki, P., von Waldow, H., Chevalier, C., Pfahl, S., and Martius, O.: Clustering of Regional-Scale Extreme Precipitation Events in Southern Switzerland, *Monthly Weather Review*, 144, 347–369, <https://doi.org/10.1175/MWR-D-15-0205.1>, 2016.
- Barton, Y., Rivoire, P., Koh, J., Ali, M. S., Kopp, J., and Martius, O.: On the temporal clustering of European extreme precipitation
930 events and its relationship to persistent and transient large-scale atmospheric drivers, *Weather and Climate Extremes*, 38, 100 518, <https://doi.org/10.1016/j.wace.2022.100518>, 2022.
- Baur, F.: Extended-Range Weather Forecasting, pp. 814–833, American Meteorological Society, Boston, MA, https://doi.org/10.1007/978-1-940033-70-9_66, 1951.
- Beran, J.: *Statistics for Long-Memory Processes*, Routledge, <https://doi.org/10.1201/9780203738481>, 2017.
- 935 Berkovic, S. and Raveh-Rubin, S.: Persistent warm and dry extremes over the eastern Mediterranean during winter: The role of North Atlantic blocking and central Mediterranean cyclones, *Quarterly Journal of the Royal Meteorological Society*, 148, 2384–2409, <https://doi.org/10.1002/qj.4308>, 2022.
- Besio, G., Briganti, R., Romano, A., Mentaschi, L., and De Girolamo, P.: Time clustering of wave storms in the Mediterranean Sea, *Natural Hazards and Earth System Sciences*, 17, 505–514, <https://doi.org/10.5194/nhess-17-505-2017>, 2017.
- 940 Bevacqua, E., Zappa, G., and Shepherd, T. G.: Shorter cyclone clusters modulate changes in European wintertime precipitation extremes, *Environmental Research Letters*, 15, 124 005, <https://doi.org/10.1088/1748-9326/abbde7>, 2020.
- Black, E., Blackburn, M., Harrison, G., Hoskins, B., and Methven, J.: Factors contributing to the summer 2003 European heatwave, *Weather*, 59, 217–223, <https://doi.org/10.1256/wea.74.04>, 2004.
- Blanchet, J., Stalla, S., and Creutin, J.-D.: Analogy of multiday sequences of atmospheric circulation favoring large rainfall accumulation
945 over the French Alps, *Atmospheric Science Letters*, 19, e809, <https://doi.org/10.1002/asl.809>, 2018.

- Blender, R., Raible, C. C., and Lunkeit, F.: Non-exponential return time distributions for vorticity extremes explained by fractional Poisson processes, *Quarterly Journal of the Royal Meteorological Society*, 141, 249–257, <https://doi.org/10.1002/qj.2354>, 2015.
- Bloomberg: Extreme Heat to Persist in India for Third Straight Month, <https://www.bloomberg.com/news/articles/2022-04-30/extreme-heat-to-persist-in-india-for-third-straight-month>, 2022.
- 950 Bray, M. T. and Cavallo, S. M.: Characteristics of long-track tropopause polar vortices, *Weather and Climate Dynamics*, 3, 251–278, <https://doi.org/10.5194/wcd-3-251-2022>, 2022.
- Bunde, A., Havlin, S., Koscielny-Bunde, E., and Schellnhuber, H. J.: Atmospheric Persistence Analysis: Novel Approaches and Applications, pp. 170–191, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-642-56257-0_5, 2002.
- Bunde, A., Büntgen, U., Ludescher, J., Luterbacher, J., and von Storch, H.: Is there memory in precipitation?, *Nature Climate Change*, 3, 174–175, <https://doi.org/10.1038/nclimate1830>, 2013.
- 955 Büeler, D., Ferranti, L., Magnusson, L., Quinting, J. F., and Grams, C. M.: Year-round sub-seasonal forecast skill for Atlantic–European weather regimes, *Quarterly Journal of the Royal Meteorological Society*, 147, 4283–4309, <https://doi.org/10.1002/qj.4178>, 2021.
- Cassou, C., Terray, L., and Phillips, A. S.: Tropical Atlantic Influence on European Heat Waves, *Journal of Climate*, 18, 2805 – 2811, <https://doi.org/10.1175/JCLI3506.1>, 2005.
- 960 Chapman, C. C., Monselesan, D. P., Risbey, J. S., Feng, M., and Sloyan, B. M.: A large-scale view of marine heatwaves revealed by archetype analysis, *Nature Communications*, 13, 7843, <https://doi.org/10.1038/s41467-022-35493-x>, 2022.
- Charney, J. G. and DeVore, J. G.: Multiple Flow Equilibria in the Atmosphere and Blocking, *Journal of Atmospheric Sciences*, 36, 1205–1216, [https://doi.org/10.1175/1520-0469\(1979\)036<1205:MFEITA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1979)036<1205:MFEITA>2.0.CO;2), 1979.
- Corral, A.: Scaling in the timing of extreme events, *Chaos, Solitons Fractals*, 74, 99–112, <https://doi.org/10.1016/j.chaos.2015.01.011>, 2015.
- 965 extreme Events and its Applications, 2015.
- Cox, D. and Isham, V.: *Point Processes*, Routledge, <https://doi.org/10.1201/9780203743034>, 1980.
- Dacre, H. F. and Pinto, J. G.: Serial clustering of extratropical cyclones: a review of where, when and why it occurs, *npj Climate and Atmospheric Science*, 3, 48, <https://doi.org/10.1038/s41612-020-00152-9>, 2020.
- De las Nieves López García, M. and Requena, J. P. R.: Different methodologies and uses of the Hurst exponent in econophysics, *Estudios de Economía Aplicada*, 37, 96–108, 2019.
- 970 De Luca, P., Harpham, C., Wilby, R. L., Hillier, J. K., Franzke, C. L. E., and Leckebusch, G. C.: Past and Projected Weather Pattern Persistence with Associated Multi-Hazards in the British Isles, *Atmosphere*, 10, 577, <https://doi.org/10.3390/atmos10100577>, 2019.
- Degenhardt, L. and Ólafsson, H.: Persistence of observed air temperatures in Iceland, *International Journal of Climatology*, 39, 1262–1275, <https://doi.org/10.1002/joc.5875>, 2019.
- 975 DelSole, T.: Optimally Persistent Patterns in Time-Varying Fields, *Journal of the Atmospheric Sciences*, 58, 1341 – 1356, [https://doi.org/10.1175/1520-0469\(2001\)058<1341:OPPITV>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<1341:OPPITV>2.0.CO;2), 2001.
- Demuzere, M., Kassomenos, P., and Philipp, A.: The COST733 circulation type classification software: an example for surface ozone concentrations in Central Europe, *Theoretical and Applied Climatology*, 105, 143–166, <https://doi.org/10.1007/s00704-010-0378-4>, 2011.
- Deni, S. M., Jemain, A. A., and Ibrahim, K.: The best probability models for dry and wet spells in Peninsular Malaysia during monsoon seasons, *International Journal of Climatology*, 30, 1194–1205, <https://doi.org/https://doi.org/10.1002/joc.1972>, 2010.
- 980 Desgraupes, B.: clusterCrit: An R Package for Computing Clustering Quality Indices, <https://CRAN.R-project.org/package=clusterCrit>, R package version 1.2.8, 2018.

- Di Capua, G., Sparrow, S., Kornhuber, K., Rousi, E., Osprey, S., Wallom, D., van den Hurk, B., and Coumou, D.: Drivers behind the summer 2010 wave train leading to Russian heatwave and Pakistan flooding, *npj Climate and Atmospheric Science*, 4, 55, <https://doi.org/10.1038/s41612-021-00211-9>, 2021.
- Di Lorenzo, E. and Mantua, N.: Multi-year persistence of the 2014/15 North Pacific marine heatwave, *Nature Climate Change*, 6, 1042–1047, <https://doi.org/10.1038/nclimate3082>, 2016.
- Dixon, P. M.: Ripley's K Function, in: *Wiley StatsRef: Statistics Reference Online*, John Wiley & Sons, Ltd, Chichester, UK, <https://doi.org/10.1002/9781118445112.stat07751>, 2014.
- 990 Dole, R. M. and Gordon, N. D.: Persistent Anomalies of the Extratropical Northern Hemisphere Wintertime Circulation: Geographical Distribution and Regional Persistence Characteristics, *Monthly Weather Review*, 111, 1567–1586, [https://doi.org/10.1175/1520-0493\(1983\)111<1567:PAOTEN>2.0.CO;2](https://doi.org/10.1175/1520-0493(1983)111<1567:PAOTEN>2.0.CO;2), 1983.
- Domeisen, D. I., White, C. J., Afargan-Gerstman, H., Muñoz, Á. G., Janiga, M. A., Vitart, F., Wulff, C. O., Antoine, S., Ardilouze, C., Batté, L., Bloomfield, H. C., Brayshaw, D. J., Camargo, S. J., Charlton-Pérez, A., Collins, D., Cowan, T., del Mar Chaves, M., Ferranti, L., Gómez, R., González, P. L., González Romero, C., Infanti, J. M., Karozis, S., Kim, H., Kolstad, E. W., LaJoie, E., Lledó, L., Magnusson, L., Malguzzi, P., Manrique-Suñén, A., Mastrangelo, D., Materia, S., Medina, H., Palma, L., Pineda, L. E., Sfetsos, A., Son, S.-W., Soret, A., Strazzo, S., and Tian, D.: Advances in the subseasonal prediction of extreme events: Relevant case studies across the globe, *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/BAMS-D-20-0221.1>, 2022.
- 995 Drouard, M. and Woollings, T.: Contrasting Mechanisms of Summer Blocking Over Western Eurasia, *Geophysical Research Letters*, 45, 12,040–12,048, <https://doi.org/10.1029/2018GL079894>, 2018.
- 1000 Du, H., Alexander, L. V., Donat, M. G., Lippmann, T., Srivastava, A., Salinger, J., Kruger, A., Choi, G., He, H. S., Fujibe, F., Rusticucci, M., Nandintsetseg, B., Manzanar, R., Rehman, S., Abbas, F., Zhai, P., Yabi, I., Stambaugh, M. C., Wang, S., Batbold, A., Oliveira, P. T., Adrees, M., Hou, W., Zong, S., Santos e Silva, C. M., Lucio, P. S., and Wu, Z.: Precipitation From Persistent Extremes is Increasing in Most Regions and Globally, *Geophysical Research Letters*, 46, 6041–6049, <https://doi.org/10.1029/2019GL081898>, 2019.
- 1005 Du, H., Donat, M. G., Zong, S., Alexander, L. V., Manzanar, R., Kruger, A., Choi, G., Salinger, J., He, H. S., Li, M.-H., Fujibe, F., Nandintsetseg, B., Rehman, S., Abbas, F., Rusticucci, M., Srivastava, A., Zhai, P., Lippmann, T., Yabi, I., Stambaugh, M. C., Wang, S., Batbold, A., de Oliveira, P. T., Adrees, M., Hou, W., e Silva, C. M. S., Lucio, P. S., and Wu, Z.: Extreme Precipitation on Consecutive Days Occurs More Often in a Warming Climate, *Bulletin of the American Meteorological Society*, 103, E1130 – E1145, <https://doi.org/10.1175/BAMS-D-21-0140.1>, 2022.
- 1010 Economou, T., Stephenson, D. B., Pinto, J. G., Shaffrey, L. C., and Zappa, G.: Serial clustering of extratropical cyclones in a multi-model ensemble of historical and future simulations, *Quarterly Journal of the Royal Meteorological Society*, 141, 3076–3087, <https://doi.org/10.1002/qj.2591>, 2015.
- Eichner, J. F., Koscielny-Bunde, E., Bunde, A., Havlin, S., and Schellnhuber, H.-J.: Power-law persistence and trends in the atmosphere: A detailed study of long temperature records, *Phys. Rev. E*, 68, 046 133, <https://doi.org/10.1103/PhysRevE.68.046133>, 2003.
- 1015 Eichner, J. F., Kantelhardt, J. W., Bunde, A., and Havlin, S.: Statistics of return intervals in long-term correlated records, *Phys. Rev. E*, 75, 011 128, <https://doi.org/10.1103/PhysRevE.75.011128>, 2007.
- Faranda, D., Messori, G., Alvarez-Castro, M. C., and Yiou, P.: Dynamical properties and extremes of Northern Hemisphere climate fields over the past 60 years, *Nonlinear Processes in Geophysics*, 24, 713–725, <https://doi.org/10.5194/npg-24-713-2017>, 2017a.
- Faranda, D., Messori, G., and Yiou, P.: Dynamical proxies of North Atlantic predictability and extremes, *Scientific Reports*, 7, 41 278, <https://doi.org/10.1038/srep41278>, 2017b.
- 1020

- Faranda, D., Alvarez-Castro, M. C., Messori, G., Rodrigues, D., and Yiou, P.: The hammam effect or how a warm ocean enhances large scale atmospheric predictability, *Nature Communications*, 10, 1316, <https://doi.org/10.1038/s41467-019-09305-8>, 2019.
- Fereday, D.: How Persistent Are North Atlantic-European Sector Weather Regimes?, *Journal of Climate*, 30, 2381–2394, <https://doi.org/10.1175/JCLI-D-16-0328.1>, 2017.
- 1025 Ferro, C. A. T. and Segers, J.: Inference for clusters of extreme values, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 65, 545–556, <https://doi.org/10.1111/1467-9868.00401>, 2003.
- Floodlist: Australia ? More Floods in Queensland After Widespread Heavy Rainfall, <https://floodlist.com/australia/queensland-floods-may-2022>, 2022.
- Ford, T. and Labosier, C. F.: Spatial patterns of drought persistence in the Southeastern United States, *International Journal of Climatology*, 34, 2229–2240, <https://doi.org/10.1002/joc.3833>, 2014.
- 1030 Fraedrich, K. and Larnder, C.: Scaling regimes of composite rainfall time series, *Tellus A: Dynamic Meteorology and Oceanography*, 45, 289 – 298, <https://doi.org/10.3402/tellusa.v45i4.14893>, 1993.
- Francis, J. A., Skific, N., and Vavrus, S. J.: North American Weather Regimes Are Becoming More Persistent: Is Arctic Amplification a Factor?, *Geophysical Research Letters*, 45, <https://doi.org/10.1029/2018GL080252>, 2018.
- 1035 Francis, J. A., Skific, N., and Vavrus, S. J.: Increased persistence of large-scale circulation regimes over Asia in the era of amplified Arctic warming, past and future, *Scientific Reports*, 10, 14 953, <https://doi.org/10.1038/s41598-020-71945-4>, 2020.
- Franzke, C., Crommelin, D., Fischer, A., and Majda, A. J.: A Hidden Markov Model Perspective on Regimes and Metastability in Atmospheric Flows, *Journal of Climate*, 21, 1740–1757, <https://doi.org/10.1175/2007JCLI1751.1>, 2008.
- Franzke, C., Woollings, T., and Martius, O.: Persistent Circulation Regimes and Preferred Regime Transitions in the North Atlantic, *Journal of the Atmospheric Sciences*, 68, 2809–2825, <https://doi.org/10.1175/JAS-D-11-046.1>, 2011.
- 1040 Franzke, C. L. E.: Persistent regimes and extreme events of the North Atlantic atmospheric circulation, *Phil. Trans. R. Soc. A*, 371, 20110 471, <https://doi.org/10.1098/rsta.2011.0471>, 2013.
- Franzke, C. L. E., Barbosa, S., Blender, R., Fredriksen, H.-B., Laepple, T., Lambert, F., Nilsen, T., Rypdal, K., Rypdal, M., Scotto, M. G., Vannitsem, S., Watkins, N. W., Yang, L., and Yuan, N.: The Structure of Climate Variability Across Scales, *Reviews of Geophysics*, 58, <https://doi.org/10.1029/2019RG000657>, 2020.
- 1045 Gálfi, V. M., Lucarini, V., and Wouters, J.: A large deviation theory-based analysis of heat waves and cold spells in a simplified model of the general circulation of the atmosphere, *Journal of Statistical Mechanics: Theory and Experiment*, 2019, 033 404, <https://doi.org/10.1088/1742-5468/ab02e8>, 2019.
- García-Herrera, R., Díaz, J., Trigo, R. M., Luterbacher, J., and Fischer, E. M.: A Review of the European Summer Heat Wave of 2003, *Critical Reviews in Environmental Science and Technology*, 40, 267–306, <https://doi.org/10.1080/10643380802238137>, 2010.
- 1050 Gershunov, A. and Barnett, T. P.: ENSO Influence on Intraseasonal Extreme Rainfall and Temperature Frequencies in the Contiguous United States: Observations and Model Results, *Journal of Climate*, 11, 1575 – 1586, [https://doi.org/10.1175/1520-0442\(1998\)011<1575:EIOIER>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<1575:EIOIER>2.0.CO;2), 1998.
- Ghil, M. and Robertson, A. W.: "Waves" vs. "particles" in the atmosphere's phase space: A pathway to long-range forecasting?, *Proceedings of the National Academy of Sciences*, 99, 2493–2500, <https://doi.org/10.1073/pnas.012580899>, 2002.
- 1055 Goswami, B.: A Brief Introduction to Nonlinear Time Series Analysis and Recurrence Plots, *Vibration*, 2, 332–368, <https://doi.org/10.3390/vibration2040021>, 2019.

- Grams, C. M., Beerli, R., Pfenninger, S., Staffell, I., and Wernli, H.: Balancing Europe's wind-power output through spatial deployment informed by weather regimes, *Nature Climate Change*, 7, 557–562, <https://doi.org/10.1038/nclimate3338>, 2017.
- 1060 Guilbert, J., Betts, A. K., Rizzo, D. M., Beckage, B., and Bombles, A.: Characterization of increased persistence and intensity of precipitation in the northeastern United States, *Geophysical Research Letters*, 42, 1888–1893, <https://doi.org/10.1002/2015GL063124>, 2015.
- Haines, K. and Hannachi, A.: Weather Regimes in the Pacific from a GCM, *Journal of Atmospheric Sciences*, 52, 2444 – 2462, [https://doi.org/10.1175/1520-0469\(1995\)052<2444:WRITPF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<2444:WRITPF>2.0.CO;2), 1995.
- Hamidieh, K., Stoev, S., and Michailidis, G.: On the Estimation of the Extremal Index Based on Scaling and Resampling, *Journal of*
 1065 *Computational and Graphical Statistics*, 18, 731–755, <https://doi.org/10.1198/jcgs.2009.08065>, 2009.
- Hannachi, A.: Low-Frequency Variability in a GCM: Three-Dimensional Flow Regimes and Their Dynamics, *Journal of Climate*, 10, 1357 – 1379, [https://doi.org/10.1175/1520-0442\(1997\)010<1357:LFVIAG>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<1357:LFVIAG>2.0.CO;2), 1997.
- Hannachi, A.: A New Set of Orthogonal Patterns in Weather and Climate: Optimally Interpolated Patterns, *Journal of Climate*, 21, 6724 – 6738, <https://doi.org/10.1175/2008JCLI2328.1>, 2008.
- 1070 Hannachi, A.: On the Origin of Planetary-Scale Extratropical Winter Circulation Regimes, *Journal of the Atmospheric Sciences*, 67, 1382 – 1401, <https://doi.org/10.1175/2009JAS3296.1>, 2010.
- Hannachi, A.: Intermittency, autoregression and censoring: a first-order AR model for daily precipitation, *Meteorological Applications*, 21, 384–397, <https://doi.org/10.1002/met.1353>, 2014.
- Hannachi, A.: Patterns Identification and Data Mining in Weather and Climate, Springer Nature, <https://doi.org/10.1007/978-3-030-67073-3>,
 1075 2021.
- Hannachi, A. and Trendafilov, N.: Archetypal Analysis: Mining Weather and Climate Extremes, *Journal of Climate*, 30, 6927 – 6944, <https://doi.org/10.1175/JCLI-D-16-0798.1>, 2017.
- Hannachi, A., Woollings, T., and Fraedrich, K.: The North Atlantic jet stream: a look at preferred positions, paths and transitions, *Quarterly Journal of the Royal Meteorological Society*, 138, 862–877, <https://doi.org/10.1002/qj.959>, 2012.
- 1080 Hannachi, A., Straus, D. M., Franzke, C. L. E., Corti, S., and Woollings, T.: Low-frequency nonlinearity and regime behavior in the Northern Hemisphere extratropical atmosphere, *Reviews of Geophysics*, 55, 199–234, <https://doi.org/10.1002/2015RG000509>, 2017.
- Hastie, T. J.: *Generalized Additive Models*, p. 59, Routledge, <https://doi.org/10.1201/9780203738535-7>, 1992.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D.,
 1085 Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hodges, K. I.: Feature Tracking on the Unit Sphere, *Monthly Weather Review*, 123, 3458 – 3465, [https://doi.org/10.1175/1520-0493\(1995\)123<3458:FTOTUS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1995)123<3458:FTOTUS>2.0.CO;2), 1995.
- 1090 Hoffmann, P., Lehmann, J., Fallah, B., and Hattermann, F. F.: Atmosphere similarity patterns in boreal summer show an increase of persistent weather conditions connected to hydro-climatic risks, *Scientific Reports*, 11, 22 893, <https://doi.org/10.1038/s41598-021-01808-z>, 2021.
- Holešovský, J. and Fusek, M.: Improved interexceedance-times-based estimator of the extremal index using truncated distribution, *Extremes*, <https://doi.org/10.1007/s10687-022-00444-8>, 2022.
- Holmberg, E., Messori, G., Caballero, R., and Faranda, D.: The counter-intuitive link between European heatwaves and atmospheric
 1095 persistence, *Earth System Dynamics Discussions*, 2022, 1–26, <https://doi.org/10.5194/esd-2022-50>, 2022.

- Horel, J. D.: Persistence of the 500 mb Height Field during Northern Hemisphere Winter, *Monthly Weather Review*, 113, 2030 – 2042, [https://doi.org/10.1175/1520-0493\(1985\)113<2030:POTMHF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1985)113<2030:POTMHF>2.0.CO;2), 1985a.
- Horel, J. D.: Persistence of Wintertime 500 mb Height Anomalies over the Central Pacific, *Monthly Weather Review*, 113, 2043 – 2048, [https://doi.org/10.1175/1520-0493\(1985\)113<2043:POWMHA>2.0.CO;2](https://doi.org/10.1175/1520-0493(1985)113<2043:POWMHA>2.0.CO;2), 1985b.
- 1100 Horton, P., Jaboyedoff, M., and Obled, C.: Global Optimization of an Analog Method by Means of Genetic Algorithms, *Monthly Weather Review*, 145, 1275 – 1294, <https://doi.org/10.1175/MWR-D-16-0093.1>, 2017.
- Hoskins, B. and Woollings, T.: Persistent Extratropical Regimes and Climate Extremes, *Current Climate Change Reports*, 1, 115–124, <https://doi.org/10.1007/s40641-015-0020-8>, 2015.
- Huguenin, M. F., Fischer, E. M., Kotlarski, S., Scherrer, S. C., Schwierz, C., and Knutti, R.: Lack of Change in the Projected Frequency and Persistence of Atmospheric Circulation Types Over Central Europe, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019GL086132>, 2020.
- 1105 Huntingford, C., Marsh, T., Scaife, A. A., Kendon, E. J., Hannaford, J., Kay, A. L., Lockwood, M., Prudhomme, C., Reynard, N. S., Parry, S., Lowe, J. A., Screen, J. A., Ward, H. C., Roberts, M., Stott, P. A., Bell, V. A., Bailey, M., Jenkins, A., Legg, T., Otto, F. E. L., Massey, N., Schaller, N., Slingo, J., and Allen, M. R.: Potential influences on the United Kingdom’s floods of winter 2013/14, *Nature Climate Change*, 4, 769–777, <https://doi.org/10.1038/nclimate2314>, 2014.
- 1110 Hurst, H. E.: Long-Term Storage Capacity of Reservoirs, *Transactions of the American Society of Civil Engineers*, 116, 770–799, <https://doi.org/10.1061/TACEAT.0006518>, 1951.
- Huth, R., Beck, C., Philipp, A., Demuzere, M., Ustrnul, Z., Cahynová, M., Kyselý, J., and Tveito, O. E.: Classifications of Atmospheric Circulation Patterns, *Annals of the New York Academy of Sciences*, 1146, 105–152, <https://doi.org/10.1196/annals.1446.019>, 2008.
- 1115 Kalra, D. S. and Santhanam, M. S.: Inferring long memory using extreme events, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 31, 113 131, <https://doi.org/10.1063/5.0064432>, 2021.
- Khare, S., Bonazzi, A., Mitas, C., and Jewson, S.: Modelling clustering of natural hazard phenomena and the effect on re/insurance loss perspectives, *Natural Hazards and Earth System Sciences*, 15, 1357–1370, <https://doi.org/10.5194/nhess-15-1357-2015>, 2015.
- Kimoto, M. and Ghil, M.: Multiple Flow Regimes in the Northern Hemisphere Winter. Part I: Methodology and Hemispheric Regimes, *Journal of Atmospheric Sciences*, 50, 2625 – 2644, [https://doi.org/10.1175/1520-0469\(1993\)050<2625:MFRITN>2.0.CO;2](https://doi.org/10.1175/1520-0469(1993)050<2625:MFRITN>2.0.CO;2), 1993.
- 1120 Kolstad, E. W., Sobolowski, S. P., and Scaife, A. A.: Intraseasonal Persistence of European Surface Temperatures, *Journal of Climate*, 28, 5365–5374, <https://doi.org/10.1175/JCLI-D-15-0053.1>, 2015.
- Kolstad, E. W., Barnes, E. A., and Sobolowski, S. P.: Quantifying the role of land-atmosphere feedbacks in mediating near-surface temperature persistence, *Quarterly Journal of the Royal Meteorological Society*, 143, 1620–1631, <https://doi.org/10.1002/qj.3033>, 2017.
- 1125 Kooperberg, C. and O’sullivan, F.: Predictive Oscillation Patterns: A Synthesis of Methods for Spatial-Temporal Decomposition of Random Fields, *Journal of the American Statistical Association*, 91, 1485–1496, <https://doi.org/10.1080/01621459.1996.10476716>, 1996.
- Kopp, J., Rivoire, P., Ali, S. M., Barton, Y., and Martius, O.: A novel method to identify sub-seasonal clustering episodes of extreme precipitation events and their contributions to large accumulation periods, *Hydrology and Earth System Sciences*, 25, 5153–5174, <https://doi.org/10.5194/hess-25-5153-2021>, 2021.
- 1130 Kornhuber, K. and Tamarin-Brodsky, T.: Future Changes in Northern Hemisphere Summer Weather Persistence Linked to Projected Arctic Warming, *Geophysical Research Letters*, 48, <https://doi.org/10.1029/2020GL091603>, 2021.

- Kornhuber, K., Osprey, S., Coumou, D., Petri, S., Petoukhov, V., Rahmstorf, S., and Gray, L.: Extreme weather events in early summer 2018 connected by a recurrent hemispheric wave-7 pattern, *Environmental Research Letters*, 14, 054 002, <https://doi.org/10.1088/1748-9326/ab13bf>, 2019.
- 1135 Koscielny-Bunde, E., Bunde, A., Havlin, S., Roman, H. E., Goldreich, Y., and Schellnhuber, H.-J.: Indication of a Universal Persistence Law Governing Atmospheric Variability, *Physical Review Letters*, 81, 729–732, <https://doi.org/10.1103/PhysRevLett.81.729>, 1998.
- Kossin, J. P.: A global slowdown of tropical-cyclone translation speed, *Nature*, 558, 104–107, <https://doi.org/10.1038/s41586-018-0158-3>, 2018.
- Koutsoyiannis, D.: Climate change, the Hurst phenomenon, and hydrological statistics, *Hydrological Sciences Journal*, 48, 3–24, <https://doi.org/10.1623/hysj.48.1.3.43481>, 2003.
- 1140 Kučerová, M., Beck, C., Philipp, A., and Huth, R.: Trends in frequency and persistence of atmospheric circulation types over Europe derived from a multitude of classifications, *International Journal of Climatology*, 37, 2502–2521, <https://doi.org/10.1002/joc.4861>, 2017.
- Kumar, S., Merwade, V., Kinter, J. L., and Niyogi, D.: Evaluation of Temperature and Precipitation Trends and Long-Term Persistence in CMIP5 Twentieth-Century Climate Simulations, *Journal of Climate*, 26, 4168 – 4185, <https://doi.org/10.1175/JCLI-D-12-00259.1>, 2013.
- 1145 Kysely, J. and Domonkos, P.: Recent increase in persistence of atmospheric circulation over Europe: comparison with long-term variations since 1881, *International Journal of Climatology*, 26, 461–483, <https://doi.org/10.1002/joc.1265>, 2006.
- Lawrence, Z. D., Perlwitz, J., Butler, A. H., Manney, G. L., Newman, P. A., Lee, S. H., and Nash, E. R.: The Remarkably Strong Arctic Stratospheric Polar Vortex of Winter 2020: Links to Record-Breaking Arctic Oscillation and Ozone Loss, *Journal of Geophysical Research: Atmospheres*, 125, <https://doi.org/10.1029/2020JD033271>, 2020.
- 1150 Legras, B. and Ghil, M.: Persistent Anomalies, Blocking and Variations in Atmospheric Predictability, *Journal of Atmospheric Sciences*, 42, 433 – 471, [https://doi.org/10.1175/1520-0469\(1985\)042<0433:PABAVI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1985)042<0433:PABAVI>2.0.CO;2), 1985.
- Li, J. and Thompson, D. W. J.: Widespread changes in surface temperature persistence under climate change, *Nature*, 599, 425–430, <https://doi.org/10.1038/s41586-021-03943-z>, 2021.
- Liu, P., Zhu, Y., Zhang, Q., Gottschalck, J., Zhang, M., Melhauser, C., Li, W., Guan, H., Zhou, X., Hou, D., Peña, M., Wu, G., Liu, Y., Zhou, L., He, B., Hu, W., and Sukhdeo, R.: Climatology of tracked persistent maxima of 500-hPa geopotential height, *Climate Dynamics*, 51, 701 – 717, <https://doi.org/10.1007/s00382-017-3950-0>, 2018.
- 1155 Liu, Q.: On the definition and persistence of blocking, *Tellus A*, 46, 286–298, <https://doi.org/10.1034/j.1600-0870.1994.t01-2-00004.x>, 1994.
- Lorenz, E. N.: Can chaos and intransitivity lead to interannual variability?, *Tellus A: Dynamic Meteorology and Oceanography*, 42, 378–389, <https://doi.org/10.3402/tellusa.v42i3.11884>, 1990.
- 1160 Lorenz, R., Jaeger, E. B., and Seneviratne, S. I.: Persistence of heat waves and its link to soil moisture memory, *Geophysical Research Letters*, 37, L09 703, <https://doi.org/10.1029/2010GL042764>, 2010.
- Lucarini, V., Faranda, D., Freitas, A. C. G. M. M., de Freitas, J. M. M., Holland, M., Kuna, T., Nicol, M., Todd, M., and Vaienti, S.: Extremes and Recurrence in Dynamical Systems, John Wiley & Sons, Inc, Hoboken, NJ, USA, <https://doi.org/10.1002/9781118632321>, 2016.
- MacDonald, G.: Persistence in Climate, Tech. rep., <https://irp.fas.org/agency/dod/jason/persistence.pdf>, 1992.
- 1165 Mailier, P. J., Stephenson, D. B., Ferro, C. A. T., and Hodges, K. I.: Serial Clustering of Extratropical Cyclones, *Monthly Weather Review*, 134, 2224–2240, <https://doi.org/10.1175/MWR3160.1>, 2006.
- Mallakpour, I., Villarini, G., Jones, M. P., and Smith, J. A.: On the use of Cox regression to examine the temporal clustering of flooding and heavy precipitation across the central United States, *Global and Planetary Change*, 155, 98–108, <https://doi.org/10.1016/j.gloplacha.2017.07.001>, 2017.

- 1170 Mallapaty, S.: Why are Pakistan's floods so extreme this year?, <https://www.nature.com/articles/d41586-022-02813-6>, 2022.
- Mandelbrot, B. B. and Wallis, J. R.: Some long-run properties of geophysical records, *Water Resources Research*, 5, 321–340, <https://doi.org/10.1029/WR005i002p00321>, 1969.
- Mann, M. E., Rahmstorf, S., Kornhuber, K., Steinman, B. A., Miller, S. K., Petri, S., and Coumou, D.: Projected changes in persistent extreme summer weather events: The role of quasi-resonant amplification, *Science Advances*, 4, eaat3272, <https://doi.org/10.1126/sciadv.aat3272>, 1175 2018.
- Marwan, N.: A historical review of recurrence plots, *The European Physical Journal Special Topics*, 164, 3–12, <https://doi.org/10.1140/epjst/e2008-00829-1>, 2008.
- Marwan, N., Carmen Romano, M., Thiel, M., and Kurths, J.: Recurrence plots for the analysis of complex systems, *Physics Reports*, 438, 237–329, <https://doi.org/10.1016/j.physrep.2006.11.001>, 2007.
- 1180 Meehl, G. A., Richter, J. H., Teng, H., Capotondi, A., Cobb, K., Doblas-Reyes, F., Donat, M. G., England, M. H., Fyfe, J. C., Han, W., Kim, H., Kirtman, B. P., Kushnir, Y., Lovenduski, N. S., Mann, M. E., Merryfield, W. J., Nieves, V., Pegion, K., Rosenbloom, N., Sanchez, S. C., Scaife, A. A., Smith, D., Subramanian, A. C., Sun, L., Thompson, D., Ummenhofer, C. C., and Xie, S.-P.: Initialized Earth System prediction from subseasonal to decadal timescales, *Nature Reviews Earth & Environment*, 2, 340–357, <https://doi.org/10.1038/s43017-021-00155-x>, 2021.
- 1185 Meng, L., Ford, T., and Guo, Y.: Logistic regression analysis of drought persistence in East China, *International Journal of Climatology*, 37, 1444–1455, <https://doi.org/10.1002/joc.4789>, 2017.
- Michelangeli, P.-A., Vautard, R., and Legras, B.: Weather Regimes: Recurrence and Quasi Stationarity, *Journal of the Atmospheric Sciences*, 52, 1237–1256, [https://doi.org/10.1175/1520-0469\(1995\)052<1237:WRRAS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<1237:WRRAS>2.0.CO;2), 1995.
- Mo, K. C. and Ghil, M.: Statistics and Dynamics of Persistent Anomalies, *Journal of Atmospheric Sciences*, 44, 877 – 902, [https://doi.org/10.1175/1520-0469\(1987\)044<0877:SADOPA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1987)044<0877:SADOPA>2.0.CO;2), 1190 1987.
- Mohr, S., Wilhelm, J., Wandel, J., Kunz, M., Portmann, R., Punge, H. J., Schmidberger, M., Quinting, J. F., and Grams, C. M.: The role of large-scale dynamics in an exceptional sequence of severe thunderstorms in Europe May-June 2018, *Weather and Climate Dynamics*, 1, 325–348, <https://doi.org/10.5194/wcd-1-325-2020>, 2020.
- Moon, H., Gudmundsson, L., and Seneviratne, S. I.: Drought Persistence Errors in Global Climate Models, *Journal of Geophysical Research: Atmospheres*, 123, 3483–3496, <https://doi.org/10.1002/2017JD027577>, 1195 2018.
- Mukhin, D., Hannachi, A., Braun, T., and Marwan, N.: Revealing recurrent regimes of mid-latitude atmospheric variability using novel machine learning method, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 32, 113 105, <https://doi.org/10.1063/5.0109889>, 2022.
- Mumby, P. J., Vitolo, R., and Stephenson, D. B.: Temporal clustering of tropical cyclones and its ecosystem impacts, *Proceedings of the National Academy of Sciences*, 108, 17 626–17 630, <https://doi.org/10.1073/pnas.1100436108>, 1200 2011.
- Namias, J.: Seasonal persistence and recurrence of European blocking during 1958–1960, *Tellus*, 16, 394–407, <https://doi.org/10.1111/j.2153-3490.1964.tb00176.x>, 1964.
- Økland, H. and Lejenäs, H.: Blocking and persistence, *Tellus A*, 39A, 33–38, <https://doi.org/10.1111/j.1600-0870.1987.tb00286.x>, 1987.
- Ontañón, S.: An overview of distance and similarity functions for structured data, *Artificial Intelligence Review*, 53, 5309–5351, <https://doi.org/10.1007/s10462-020-09821-w>, 1205 2020.
- Overland, J. E. and Wang, M.: The 2020 Siberian heat wave, *International Journal of Climatology*, 41, <https://doi.org/10.1002/joc.6850>, 2021.

- Pandolfo, L.: Observational Aspects of the Low-Frequency Intraseasonal Variability of the Atmosphere in Middle Latitudes, vol. 34 of *Advances in Geophysics*, pp. 93–174, Elsevier, [https://doi.org/10.1016/S0065-2687\(08\)60435-5](https://doi.org/10.1016/S0065-2687(08)60435-5), 1993.
- Parzen, E.: Quantile spectral analysis and long-memory time series, *Journal of Applied Probability*, 23, 41–54, <https://doi.org/10.2307/3214341>, 1986.
- Pelletier, J. D. and Turcotte, D. L.: Long-range persistence in climatological and hydrological time series: analysis, modeling and application to drought hazard assessment, *Journal of Hydrology*, 203, 198–208, [https://doi.org/10.1016/S0022-1694\(97\)00102-9](https://doi.org/10.1016/S0022-1694(97)00102-9), 1997.
- Perez-Zanon, N. and coauthors: CSTools: Assessing Skill of Climate Forecasts on Seasonal-to-Decadal Timescales, <https://CRAN.R-project.org/package=CSTools>, R package version 4.1.1, 2022.
- Pfleiderer, P. and Coumou, D.: Quantification of temperature persistence over the Northern Hemisphere land-area, *Climate Dynamics*, 51, 627–637, <https://doi.org/10.1007/s00382-017-3945-x>, 2018.
- Pfleiderer, P., Schleussner, C.-F., Kornhuber, K., and Coumou, D.: Summer weather becomes more persistent in a 2 degree C world, *Nature Climate Change*, 9, 666–671, <https://doi.org/10.1038/s41558-019-0555-0>, 2019.
- Pinto, J. G., Bellenbaum, N., Karremann, M. K., and Della-Marta, P. M.: Serial clustering of extratropical cyclones over the North Atlantic and Europe under recent and future climate conditions, *Journal of Geophysical Research: Atmospheres*, 118, 12,476–12,485, <https://doi.org/10.1002/2013JD020564>, 2013.
- Pinto, J. G., Gómar, I., Masato, G., Dacre, H. F., Woollings, T., and Caballero, R.: Large-scale dynamics associated with clustering of extratropical cyclones affecting Western Europe, *Journal of Geophysical Research: Atmospheres*, 119, 13,704–13,719, <https://doi.org/10.1002/2014JD022305>, 2014.
- Pinto, J. G., Ulbrich, S., Economou, T., Stephenson, D. B., Karremann, M. K., and Shaffrey, L. C.: Robustness of serial clustering of extratropical cyclones to the choice of tracking method, *Tellus A: Dynamic Meteorology and Oceanography*, 68, 32 204, <https://doi.org/10.3402/tellusa.v68.32204>, 2016.
- Pires, C. A. L. and Hannachi, A.: Bispectral analysis of nonlinear interaction, predictability and stochastic modelling with application to ENSO, *Tellus A: Dynamic Meteorology and Oceanography*, 73, 1–30, <https://doi.org/10.1080/16000870.2020.1866393>, 2021.
- Potter, K. W.: Annual precipitation in the northeast United States: Long memory, short memory, or no memory?, *Water Resources Research*, 15, 340–346, <https://doi.org/10.1029/WR015i002p00340>, 1979.
- Quandt, L.-A., Keller, J. H., Martius, O., and Jones, S. C.: Forecast Variability of the Blocking System over Russia in Summer 2010 and Its Impact on Surface Conditions, *Weather and Forecasting*, 32, 61 – 82, <https://doi.org/10.1175/WAF-D-16-0065.1>, 2017.
- Rakovec, O., Samaniego, L., Hari, V., Markonis, Y., Moravec, V., Thober, S., Hanel, M., and Kumar, R.: The 2018-2020 Multi-Year Drought Sets a New Benchmark in Europe, *Earth’s Future*, 10, <https://doi.org/10.1029/2021EF002394>, 2022.
- Ramirez-Amaro, K. and Figueroa-Nazuno, J.: Recurrence Plot Analysis and its Application to Teleconnection Patterns, in: 2006 15th International Conference on Computing, pp. 65–72, <https://doi.org/10.1109/CIC.2006.59>, 2006.
- Ray, R., Khondekar, M. H., Ghosh, K., and Bhattacharjee, A. K.: Complexity and periodicity of daily mean temperature and dew-point across India, *Journal of Earth System Science*, 128, 143, <https://doi.org/10.1007/s12040-019-1174-x>, 2019.
- Rehman, S. and Siddiqi, A.: Wavelet-based Hurst exponent and fractal dimensional analysis of Saudi climatic dynamics, *Chaos, Solitons & Fractals*, 40, 1081–1090, <https://doi.org/10.1016/j.chaos.2007.08.063>, 2009.
- Richardson, D., Kilsby, C. G., Fowler, H. J., and Bárdossy, A.: Weekly to multi-month persistence in sets of daily weather patterns over Europe and the North Atlantic Ocean, *International Journal of Climatology*, 39, 2041–2056, <https://doi.org/10.1002/joc.5932>, 2019.

- Ripley, B. D.: Spatial Statistics, Wiley Series in Probability and Statistics, John Wiley & Sons, Inc., Hoboken, NJ, USA, <https://doi.org/10.1002/0471725218>, 1981.
- Röthlisberger, M. and Martius, O.: Quantifying the Local Effect of Northern Hemisphere Atmospheric Blocks on the Persistence of Summer Hot and Dry Spells, *Geophysical Research Letters*, 46, 10 101–10 111, <https://doi.org/10.1029/2019GL083745>, 2019.
- Röthlisberger, M., Frossard, L., Bosart, L. F., Keyser, D., and Martius, O.: Recurrent Synoptic-Scale Rossby Wave Patterns and Their Effect on the Persistence of Cold and Hot Spells, *Journal of Climate*, 32, 3207–3226, <https://doi.org/10.1175/JCLI-D-18-0664.1>, 2019.
- Rousi, E., Fink, A. H., Andersen, L. S., Becker, F. N., Beobide-Arsuaga, G., Breil, M., Cozzi, G., Heinke, J., Jach, L., Niermann, D., Petrovic, D., Richling, A., Riebold, J., Steidl, S., Suarez-Gutierrez, L., Tradowsky, J., Coumou, D., Düsterhus, A., Ellsäßer, F., Fragakoulidis, G., Gliksman, D., Handorf, D., Haustein, K., Kornhuber, K., Kunstmann, H., Pinto, J. G., Warrach-Sagi, K., and Xoplaki, E.: The extremely hot and dry 2018 summer in central and northern Europe from a multi-faceted weather and climate perspective, *EGUsphere*, 2022, 1–37, <https://doi.org/10.5194/egusphere-2022-813>, 2022a.
- Rousi, E., Kornhuber, K., Beobide-Arsuaga, G., Luo, F., and Coumou, D.: Accelerated western European heatwave trends linked to more-persistent double jets over Eurasia, *Nature Communications*, 13, 3851, <https://doi.org/10.1038/s41467-022-31432-y>, 2022b.
- Santhanam, M. S. and Kantz, H.: Return interval distribution of extreme events and long-term memory, *Phys. Rev. E*, 78, 051 113, <https://doi.org/10.1103/PhysRevE.78.051113>, 2008.
- Schwierz, C., Croci-Maspoli, M., and Davies, H. C.: Perspicacious indicators of atmospheric blocking, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2003GL019341>, 2004.
- Sericola, B.: Markov Chains, John Wiley & Sons, Inc., Hoboken, NJ USA, <https://doi.org/10.1002/9781118731543>, 2013.
- Serinaldi, F. and Kilsby, C. G.: On the sampling distribution of Allan factor estimator for a homogeneous Poisson process and its use to test inhomogeneities at multiple scales, *Physica A: Statistical Mechanics and its Applications*, 392, 1080–1089, <https://doi.org/10.1016/j.physa.2012.11.015>, 2013.
- Sharma, T. C. and Panu, U. S.: Modeling of hydrological drought durations and magnitudes: Experiences on Canadian streamflows, *Journal of Hydrology: Regional Studies*, 1, 92–106, <https://doi.org/10.1016/j.ejrh.2014.06.006>, 2014.
- Smith, J. A. and Karr, A. F.: Flood Frequency Analysis Using the Cox Regression Model, *Water Resources Research*, 22, 890–896, <https://doi.org/10.1029/WR022i006p00890>, 1986.
- Son, R., Wang, S.-Y. S., Kim, S. H., Kim, H., Jeong, J.-H., and Yoon, J.-H.: Recurrent pattern of extreme fire weather in California, *Environmental Research Letters*, 16, 094 031, <https://doi.org/10.1088/1748-9326/ac1f44>, 2021.
- Steinfeld, D.: ConTrack - Contour Tracking of circulation anomalies in weather and climate data, <https://doi.org/10.5281/ZENODO.4765560>, 2021.
- Steinfeld, D., Boettcher, M., Forbes, R., and Pfahl, S.: The sensitivity of atmospheric blocking to upstream latent heating – numerical experiments, *Weather and Climate Dynamics*, 1, 405–426, <https://doi.org/10.5194/wcd-1-405-2020>, 2020.
- Stephenson, D. B., Hannachi, A., and O’Neill, A.: On the existence of multiple climate regimes, *Quarterly Journal of the Royal Meteorological Society*, 130, 583–605, <https://doi.org/10.1256/qj.02.146>, 2004.
- Strommen, K., Chantry, M., Dorrington, J., and Otter, N.: A topological perspective on weather regimes, *Climate Dynamics*, <https://doi.org/10.1007/s00382-022-06395-x>, 2022.
- Tatli, H.: Detecting persistence of meteorological drought via the Hurst exponent, *Meteorological Applications*, 22, 763–769, <https://doi.org/10.1002/met.1519>, 2015.
- Telesca, L.: Time-clustering of natural hazards, *Natural Hazards*, 6, 593–601, <https://doi.org/10.1007/s11069-006-9023-z>, 2007.

- Telesca, L., Lovallo, M., and Kanevski, M.: Power spectrum and multifractal detrended fluctuation analysis of high-frequency wind measurements in mountainous regions, *Applied Energy*, 162, 1052–1061, <https://doi.org/10.1016/j.apenergy.2015.10.187>, 2016.
- 1285 Telesca, L., Guignard, F., Laib, M., and Kanevski, M.: Analysis of temporal properties of extremes of wind measurements from 132 stations over Switzerland, *Renewable Energy*, 145, 1091–1103, <https://doi.org/10.1016/j.renene.2019.06.089>, 2020.
- Trenberth, K. E.: Some Effects of Finite Sample Size and Persistence on Meteorological Statistics. Part II: Potential Predictability, *Monthly Weather Review*, 112, 2369 – 2379, [https://doi.org/10.1175/1520-0493\(1984\)112<2369:SEOFSS>2.0.CO;2](https://doi.org/10.1175/1520-0493(1984)112<2369:SEOFSS>2.0.CO;2), 1984.
- Tuel, A. and Martius, O.: A global perspective on the sub-seasonal clustering of precipitation extremes, *Weather and Climate Extremes*, 33, 100 348, <https://doi.org/10.1016/j.wace.2021.100348>, 2021a.
- 1290 Tuel, A. and Martius, O.: A climatology of sub-seasonal temporal clustering of extreme precipitation in Switzerland and its links to extreme discharge, *Natural Hazards and Earth System Sciences*, 21, 2949–2972, <https://doi.org/10.5194/nhess-21-2949-2021>, 2021b.
- Tuel, A. and Martius, O.: Subseasonal Temporal Clustering of Extreme Precipitation in the Northern Hemisphere: Regionalization and Physical Drivers, *Journal of Climate*, 35, 3537–3555, <https://doi.org/10.1175/JCLI-D-21-0562.1>, 2022a.
- Tuel, A. and Martius, O.: The influence of modes of climate variability on the sub-seasonal temporal clustering of extreme precipitation, *iScience*, 25, 103 855, <https://doi.org/10.1016/j.isci.2022.103855>, 2022b.
- 1295 Tuel, A. and Martius, O.: On the persistence of warm and cold spells in the Northern Hemisphere extratropics: regionalisation, synoptic-scale dynamics, and temperature budget, *EGU sphere*, 2023, 1–36, <https://doi.org/10.5194/egusphere-2022-1478>, 2023.
- Tuel, A., Schaeffli, B., Zscheischler, J., and Martius, O.: On the links between sub-seasonal clustering of extreme precipitation and high discharge in Switzerland and Europe, *Hydrology and Earth System Sciences*, 26, 2649–2669, <https://doi.org/10.5194/hess-26-2649-2022>, 2022a.
- 1300 Tuel, A., Steinfeld, D., Ali, S. M., Sprenger, M., and Martius, O.: Large-scale drivers of persistent extreme weather during early summer 2021 in Europe, *Geophysical Research Letters*, 49, e2022GL099 624, <https://doi.org/10.1029/2022GL099624>, 2022b.
- van der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D., Yager, N., Gouillart, E., Yu, T., and the scikit-image contributors: scikit-image: image processing in Python, *PeerJ*, 2, e453, <https://doi.org/10.7717/peerj.453>, 2014.
- 1305 Vautard, R.: Multiple Weather Regimes over the North Atlantic: Analysis of Precursors and Successors, *Monthly Weather Review*, 118, 2056 – 2081, [https://doi.org/10.1175/1520-0493\(1990\)118<2056:MWROTN>2.0.CO;2](https://doi.org/10.1175/1520-0493(1990)118<2056:MWROTN>2.0.CO;2), 1990.
- Velásquez Valle, M. A., Medina García, G., Sánchez Cohen, I., Klaudia Oleschko, L., Ruiz Corral, J. A., and Korvin, G.: Spatial Variability of the Hurst Exponent for the Daily Scale Rainfall Series in the State of Zacatecas, Mexico, *Journal of Applied Meteorology and Climatology*, 52, 2771 – 2780, <https://doi.org/10.1175/JAMC-D-13-0136.1>, 2013.
- 1310 Vigaud, N., Robertson, A., and Tippet, M. K.: Predictability of Recurrent Weather Regimes over North America during Winter from Submonthly Reforecasts, *Monthly Weather Review*, 146, 2559–2577, <https://doi.org/10.1175/MWR-D-18-0058.1>, 2018.
- Villarini, G., Smith, J. A., Baeck, M. L., Vitolo, R., Stephenson, D. B., and Krajewski, W. F.: On the frequency of heavy rainfall for the Midwest of the United States, *Journal of Hydrology*, 400, 103–120, <https://doi.org/10.1016/j.jhydrol.2011.01.027>, 2011.
- Villarini, G., Smith, J. A., Vitolo, R., and Stephenson, D. B.: On the temporal clustering of US floods and its relationship to climate teleconnection patterns, *International Journal of Climatology*, 33, 629–640, <https://doi.org/10.1002/joc.3458>, 2013.
- 1315 Vitart, F., Ardilouze, C., Bonet, A., Brookshaw, A., Chen, M., Codorean, C., Déqué, M., Ferranti, L., Fucile, E., Fuentes, M., Hendon, H., Hodgson, J., Kang, H.-S., Kumar, A., Lin, H., Liu, G., Liu, X., Malguzzi, P., Mallas, I., Manoussakis, M., Mastrangelo, D., MacLachlan, C., McLean, P., Minami, A., Mladek, R., Nakazawa, T., Najm, S., Nie, Y., Rixen, M., Robertson, A. W., Ruti, P., Sun, C., Takaya, Y., Tolstykh, M., Venuti, F., Waliser, D., Woolnough, S., Wu, T., Won, D.-J., Xiao, H., Zaripov, R., and Zhang, L.: The Subseasonal to Seasonal (S2S)

- 1320 Prediction Project Database, *Bulletin of the American Meteorological Society*, 98, 163–173, <https://doi.org/10.1175/BAMS-D-16-0017.1>, 2017.
- Vitolo, R., Stephenson, D. B., Cook, I. M., and Mitchell-Wallace, K.: Serial clustering of intense European storms, *Meteorologische Zeitschrift*, 18, 411–424, <https://doi.org/10.1127/0941-2948/2009/0393>, 2009.
- von Lindheim, J., Harikrishnan, A., Dörffel, T., Klein, R., Koltai, P., Mikula, N., Müller, A., Névir, P., Pacey, G., Polzin, R., and Vercauteren, N.: Definition, detection, and tracking of persistent structures in atmospheric flows, <https://doi.org/10.48550/ARXIV.2111.13645>, 2021.
- 1325 Weatherhead, E., Gearheard, S., and Barry, R.: Changes in weather persistence: Insight from Inuit knowledge, *Global Environmental Change*, 20, 523–528, <https://doi.org/10.1016/j.gloenvcha.2010.02.002>, governance, Complexity and Resilience, 2010.
- Weber, J., Meyers, M., Beck, C., Timme, M., Pinto, J. G., Withaut, D., and Schäfer, B.: Wind Power Persistence Characterized by Superstatistics, *Scientific Reports*, 9, 19971, <https://doi.org/10.1038/s41598-019-56286-1>, 2019.
- 1330 Weiland, R. S., van der Wiel, K., Selten, F., and Coumou, D.: Intransitive Atmosphere Dynamics Leading to Persistent Hot-Dry or Cold-Wet European Summers, *Journal of Climate*, 34, 6303–6317, <https://doi.org/10.1175/JCLI-D-20-0943.1>, 2021.
- Weiss, J. P. and Weiss, J. B.: Quantifying Persistence in ENSO, *Journal of the Atmospheric Sciences*, 56, 2737–2760, [https://doi.org/10.1175/1520-0469\(1999\)056<2737:QPIE>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<2737:QPIE>2.0.CO;2), 1999.
- Wernli, H. and Schwerz, C.: Surface Cyclones in the ERA-40 Dataset (1958–2001). Part I: Novel Identification Method and Global Climatology, *Journal of the Atmospheric Sciences*, 63, 2486 – 2507, <https://doi.org/10.1175/JAS3766.1>, 2006.
- 1335 Wharton, E., Panetta, K., and Agaian, S.: Human visual system based similarity metrics, in: 2008 IEEE International Conference on Systems, Man and Cybernetics, pp. 685–690, <https://doi.org/10.1109/ICSMC.2008.4811357>, 2008.
- Wikipedia: 2022 Eastern Australia floods, https://en.wikipedia.org/wiki/2022_Eastern_Australia_floods, 2022.
- Wilby, R. L., Noone, S., Murphy, C., Matthews, T., Harrigan, S., and Broderick, C.: An evaluation of persistent meteorological drought using a homogeneous Island of Ireland precipitation network, *International Journal of Climatology*, 36, 2854–2865, <https://doi.org/10.1002/joc.4523>, 2016.
- 1340 Witt, A. and Malamud, B. D.: Quantification of Long-Range Persistence in Geophysical Time Series: Conventional and Benchmark-Based Improvement Techniques, *Surveys in Geophysics*, 34, 541–651, <https://doi.org/10.1007/s10712-012-9217-8>, 2013.
- Wolff, N. H., Wong, A., Vitolo, R., Stolberg, K., Anthony, K. R. N., and Mumby, P. J.: Temporal clustering of tropical cyclones on the Great Barrier Reef and its ecological importance, *Coral Reefs*, 35, 613–623, <https://doi.org/10.1007/s00338-016-1400-9>, 2016.
- 1345 Wolters, M. A.: Better Autologistic Regression, *Frontiers in Applied Mathematics and Statistics*, 3, <https://doi.org/10.3389/fams.2017.00024>, 2017.
- Woollings, T., Hannachi, A., and Hoskins, B.: Variability of the North Atlantic eddy-driven jet stream, *Quarterly Journal of the Royal Meteorological Society*, 136, 856–868, <https://doi.org/10.1002/qj.625>, 2010.
- 1350 World Meteorological Organization: Extreme weather in China highlights climate change impacts and need for early warnings, <https://public.wmo.int/en/media/news/extreme-weather-china-highlights-climate-change-impacts-and-need-early-warnings>, 2022.
- Yang, L. and Fu, Z.: Process-dependent persistence in precipitation records, *Physica A: Statistical Mechanics and its Applications*, 527, 121 459, <https://doi.org/10.1016/j.physa.2019.121459>, 2019.
- Yang, Z. and Villarini, G.: Examining the capability of reanalyses in capturing the temporal clustering of heavy precipitation across Europe, *Climate Dynamics*, 53, 1845–1857, <https://doi.org/10.1007/s00382-019-04742-z>, 2019.
- 1355 Yiou, P., Cattiaux, J., Ribes, A., Vautard, R., and Vrac, M.: Recent Trends in the Recurrence of North Atlantic Atmospheric Circulation Patterns, *Complexity*, p. 3140915, <https://doi.org/10.1155/2018/3140915>, 2018.

- Yuan, N., Fu, Z., and Mao, J.: Different scaling behaviors in daily temperature records over China, *Physica A: Statistical Mechanics and its Applications*, 389, 4087–4095, <https://doi.org/10.1016/j.physa.2010.05.026>, 2010.
- 1360 Zerzucha, P. and Walczak, B.: Concept of (dis)similarity in data analysis, *TrAC Trends in Analytical Chemistry*, 38, 116–128, <https://doi.org/10.1016/j.trac.2012.05.005>, 2012.
- Zolina, O., Simmer, C., Belyaev, K., Gulev, S. K., and Koltermann, P.: Changes in the Duration of European Wet and Dry Spells during the Last 60 Years, *Journal of Climate*, 26, 2022–2047, <https://doi.org/10.1175/JCLI-D-11-00498.1>, 2013.