The crustal structure of the Lomgmenshan fault zone and its implications for seismogenesis: New insight from aeromagnetic and gravity data

Hai Yang^{1,2,3}, Shengqing Xiong^{1,3}, Qiankun Liu¹, Fang Li¹, Zhiye Jia¹, Xue Yang¹, Haofei Yan¹, Zhaoliang Li¹

5

10

¹China Aero Geophysical Survey and Remote Sensing Center for Natural Resources, Beijing, 100083 ²State Key Laboratory of Lithospheric Evolution, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, 100029

³Key Laboratory of Airborne Geophysics and Remote Sensing Geology, Ministry of Natural Resources, Beijing 100083, China

Correspondence to: Shengqing Xiong (xsqagrs@126.com); Hai Yang (yanghai@mail.cgs.gov.cn)

Abstract. Although many geophysical models have been proposed in the Longmenshan fault zone (LFZ) and its surrounding areas, the deep structure of the seismic gap and its constraint of the Wenchuan and Lushan earthquakes remain uncertain. Based on the compiled aeromagnetic data and Bouguer gravity

- 15 data, we have tried to create a more detailed and reasonable magnetic and density model using 2D forward modeling and 3D inversion and made the deep structure of the LFZ visible. The research shows that structure is heterogenous across the LFZ. The earthquake epicenters are located in region with high magnetic anomalies and gravity gradients that are associated with rigid blocks where apt to accumulate stress. However, the seismic gap shows low magnetic anomalies and transition of gravity anomalies
- 20 related to a weak zone. The Sichuan Basin has two NE-trending banded high magnetic blocks extending beneath the LFZ that firmly support the crust of the Sichuan Basin was downward subduction toward the LFZ. More importantly, the basement subducts to approximately 33 km west of the Wenchuan-Maoxian fault with a low dip angle beneath the middle segment of the LFZ, whereas the distance decreases to approximately 17 and 19 km under the southern segment. Thus, the crust of the Sichuan Basin beneath
- 25 the middle segment extends farther than that beneath the southern segment with the seismic gap as the transition zone. Therefore, we propose that the structural heterogeneity of the basement on the western margin of the Sichuan Basin may be the main reason for the different focal mechanisms and geodynamics of the Wenchuan and Lushan earthquakes.

1 Introduction

- 30 In 2008 and 2013, the devastating Wenchuan (M_s 8.0) and Lushan (M_s 7.0) earthquakes successively struck the LFZ in the eastern margin of the Tibetan Plateau. The two earthquakes caused great losses of human lives and properties in China. Many researchers focus on the mechanism of two earthquakes because they occurred at a close distance with distinct fault geometry, surface rupture, coulomb stress, and deep structure (Chen et al., 2013; Li et al., 2013, 2014; Lei and Zhao, 2010; Pei et
- 35 al., 2010; Shan et al., 2013; Wang et al., 2009, 2014a, 2015, 2017a; Wu et al., 2013; Zhan et al., 2013; Zhao et al., 2013). For instance, the Wenchuan earthquake occurred on the Yingxiu-Beichuan fault in the middle segment of the LFZ, while the Lushan earthquake occurred on a blind reverse fault in the southern segment (Chen et al., 2013; Li et al., 2014; Wang et al., 2014a). The Wenchuan earthquake occurred on a thrust fault associated with dextral strike-slip movement and had a surface rupture extending more than
- 40 300 km toward the NE. However, the Lushan earthquake was dominated by concealed thrust fault with the rupture zone restricted to 30 km underground and no obvious surface rupture (Chen et al., 2013; Zhao et al., 2013). It is noted that there is a 40~60 km gap void of aftershocks for both earthquakes along the fault zone between the Wenchuan and Lushan earthquakes. (Du et al., 2013; Liu et al., 2018; Pei et al., 2014; Wang et al., 2015; Zhu et al., 2016).
- 45 Two different earthquakes happened in the LFZ within a short time and the risk of the seismic gap has challenged Earth scientists. In the past decade, numerous studies have been carried out on the high-resolution deep structure of the LFZ. It is generally accepted that the deep structure of the 2013 Lushan hypocenter is characterized by a high velocity (Vp, Vs), low Poisson's ratio, and high resistivity, whereas high velocity (Vp, Vs), high Poisson's ratio, and high resistivity were imaged at the 2008 Wenchuan hypocenter (Li et al., 2013; Lei and Zhao, 2010; Pei et al., 2010; Wang et al., 2009, 2015; Zhan et al., 2013). The gap area between two earthquakes is characterized by low velocities (Vp, Vs), high Poisson's ratios, and high conductivity (Liu et al., 2018; Pei et al., 2014; Wang et al., 2014c, 2015, 2017b; Zhan et al., 2013). Several models recently proposed by different research groups to explain the formation of a low-velocity seismic gap, (1) a weak and ductile deformation area formed by the strong compression between the Tibetan Plateau and the Sichuan Basin (Pei et al., 2014), (2) a fault zone caused mantle upwelling and partial melting (He et al., 2017; Liang et al., 2018), (3) fluid-bearing ductile flow from the

mid-lower crust of Tibet (Wang et al., 2014c). Therefore, the formation of the low velocity and highconductivity gap area beneath the seismic gap still cannot reach an agreement, because the large-scale crustal shortening resulted in complex deep structure under the LFZ.

- 60 Previous interpretation of the magnetic and gravity data trying to describe the magnetic and density differences along the LFZ. 1) the Wenchuan and Lushan earthquakes were separated by NNW-trending (near EW) high magnetic material, the Wenchuan earthquake distributed in the northeast segment of the negative magnetic anomaly belt (Yan et al., 2016), 2) the LFZ is characterized by large negative magnetic anomaly area caused by some crystalline complexes with reversal magnetization in the crust (Yan et al.,
- 65 2016; Zhang et al., 2010), 3) the LFZ lies in the gravity gradient zone that is related to the density discontinuity in deep (Chen et al., 2013; Zhang et al., 2010). However, the lack of detailed magnetic and gravity structure makes it difficult to understand the deep structure of the LFZ. To create an integrated geophysical model beneath the middle and southern segments of the LFZ, in this study we conducted a quantitative analysis using 2D forward modeling under the constraint of velocity model and 3D inversion.
- 70 Then, we discuss the structure geometry and physical property under the LFZ and adjacent two blocks, which is essential to advance our understanding on the genesis of the seismic gap and geodynamics of two earthquakes.

2. Tectonic setting

The LFZ lies in a convergent area between the Songpan–Ganzi block (SGB) and the Sichuan Basin at the eastern margin of the Tibet Plateau (Fig. 1a). It is characterized by a steep topographic gradient, complex strike-slip, and strong thrust motions with an average elevation rising 4000m above the Sichuan Basin over a distance of less than 100 km (Burchfiel et al., 2008; Chen and Wilson, 2008; Guo et al., 2013; Parsons et al., 2008; Royden et al., 2008). The fault zone is bounded by the Wenchuan - Maowen fault in the west, and the Guanxian-Anxian fault in the east. The Mian-Lue suture formed the northeast

80 boundary of the fault zone and connected with the Kangdian tectonic belt to the southwest. The SGB is covered by a 10 km Middle to Upper Triassic flysch formation in the northwest of the LFZ. Triassic syntectonic adakitic-type granitoids are widely distributed in the SGB, which are likely sourced from the partial melting of an underlying Proterozoic Yangtze-type crystalline basement (Dai et al., 2011; Hu et al., 2005; Roger et al.,2010; Zhang et al.,2006; Zhao et al., 2007a, b) (Fig.1b). The Sichuan Basin has a
Precambrian crystalline basement overlain by late Proterozoic to Cenozoic sedimentary cover.
Geological and geophysical studies suggest the crust of the Sichuan basin probably extends beneath the
LFZ (Burchfiel et al., 1995; Guo et al., 2013; Roger et al., 2010; Wang et al., 2015; Zhang et al., 2006;
Zhu et al., 2008).



Figure 1. Geological background of the LFZ. (a) Tectonic map of the LFZ and adjacent area. (b) Geological map of the LFZ, showing the location of modeling profiles as blue lines (modified after Li et al. (2008) and Yan (2011)). F1: Wenchuan-Maoxian fault, F2: Yinxiu-Beichuan fault, F3: Guanxian-Anxian fault, F4: Guangyuan-Dayi concealed fault, F5: Pingwu-Qingchuan fault, F6: Longquanshan fault; PG: Pengguan complex; BX: Baoxing complex; KD: Kangding complex; JZD: Jiaoziding complex. Red stars show the 2013 Lushan and 2008 Wenchuan earthquakes, respectively. Pink dots indicate the 1204 earthquakes (earthquake magnitude≥4) from 1970 to 2014. Blue line is the seismic profile from He et al. (2017). The pink squares are magnetic susceptibility measurement points.

There are four reversed thrust and strike-slip faults from northwest to southeast, including the Wenchuan-Maowen fault (F₁: rear fault), the Yingxiu-Beichuan fault (F₂: central fault), the Guanxian-Anxian fault (F₃: fore fault), and the Guangyuan-Dayi fault (F₄: range front blind fault) (Fig. 1b). The Wenchuan earthquake occurred in the Yingxiu-Beichuan fault. The Lushan earthquake occurred in a blind reverse fault between the Guanxian-Anxian fault and the Guangyuan-Dayi fault in the southern segment of the LFZ (Chen et al., 2013; Li et al., 2013; Wang et al., 2014c). Due to the distinct deformation style, evolution history, and foreland sediment, the LFZ can be divided into three segments along the strike (Fig. 1b). The northern segment has an imbricated thrust fault system in front of the Jiaoziding (JZD) complex and the Tangwangzhai syncline. The middle segment outcrops the Pengguan (PG) complex with

Proterozoic to Triassic klippes developing in front of it. The southern segment outcrops the Baoxing (BX)

complex with Silurian to Triassic klippes distributing in front of it (Li et al., 2008). Unlike the PG complex is fault contact with later formations, the BX complex is covered by a successive formation showing parallel unconformity or conformity. From the northeast to southwest of the LFZ, the deformation age is getting younger, the deformation is more brittle and extensive, and the Cenozoic activity is more intense (Li et al., 2008). The Wenchuan earthquake was located in the middle segment

of the LFZ, while the Lushan earthquake occurred in the southern segment. The seismic gap lies in the transition zone between the middle and southern segments.

3 Data and Method

110

115 **3.1 Rock magnetic susceptibility measurement**

The Proterozoic to Cenozoic strata are successively distributed in the LFZ and surrounding area. Measuring the magnetic susceptibility of these formations is important for the interpretation of aeromagnetic anomalies. Therefore, a field survey of rock magnetic susceptibility was conducted in the LFZ, especially the high aeromagnetic anomaly area. Each type of lithology test 30 points for reliable

- 120 statistic results using a KM-7 Magnetic Susceptibility kappameter (Fig. 1 and Table 1). The results show that Proterozoic serpentine has the highest magnetism with an average magnetic susceptibility value of 0.0765 SI. Proterozoic quartz diorite has a moderate magnetic susceptibility of 0.0238 0.0487 SI with an average value of 0.0377 SI, while Proterozoic granite has values of 0.0002 0.0247 SI with an average value of 0.0068 SI. The Triassic and Jurassic granites are widely distributed in the west of the LFZ, and
- 125 their magnetic susceptibility ranges from 0.0118 0.0201 SI, and their average value is 0.0167 SI. The magnetic susceptibility of the Siguniangshan granite ranges from 0.0077 0.0161 SI with an average value of 0.0123 SI. Most sedimentary formations are usually nonmagnetic. In general, the outcropping Proterozoic intermediate-felsic intrusive rocks and Triassic-Jurassic granite commonly have high magnetic susceptibility.

130

Formation/Period	lithologic	min	max	average	Point No.
KD	quartz diorite	0.0226	0.0562	0.0407	D04
KD	quartz diorite	0.0016	0.0542	0.0238	D09
KD	quartz diorite	0.0144	0.0868	0.0487	D10
KD	granite	0.0001	0.0029	0.0005	D05
KD	granite	0.0013	0.0082	0.0041	D06
KD	granite	0.0009	0.0048	0.0032	D07
KD	granite	0.0135	0.0359	0.0247	D08
KD	granite	0.0015	0.0224	0.0082	D11
PG	granite	0.0001	0.0008	0.0002	D15
PG	Serpentine (Not in-suit)	0.0490	0.1203	0.0765	D15-1
Devonian	Mica schist	0.0071	0.0840	0.0250	D12
Permian	Basalt (Not in-suit)	0.0062	0.0710	0.0361	D16
Triassic	granite	0.0460	0.0239	0.0118	D02
Triassic	granite	0.0105	0.0259	0.0182	D03
Triassic	Syenite (Not in-suit)	0.0122	0.0243	0.0201	D13
Jurassic	granite	0.0077	0.0161	0.0123	D14
Cretaceous	Mudstone	0.0003	0.0018	0.0011	D01

Table 1. Statistics of rock magnetic susceptibility in the LFZ

3.2 Aeromagnetic and gravity data

135 The total field magnetic anomaly data used in this study are based on a compilation provided by the China Aero Geophysical Survey and Remote Sensing Center for Natural Resources (Xiong et al., 2013, 2016a, b), covering the LFZ and surrounding area. These magnetic data were derived from 47 different airborne surveys from 1958 to 2013 with three kinds of scales. The line spacings of 1:500, 000-1:1000,000 aeromagnetic datasets are 5 and 10 km respectively. The sampling interval is 10-15m. The 140 line spacings of 1:100, 000-1:200,000 aeromagnetic datasets are 1 and 2 km respectively. The sampling interval is 10-15m. The line spacing of 1:50,000 aeromagnetic datasets is 500m. The sampling interval is 5-10m. All data were adjusted to 1km above Earth's surface and sutured to be a continuous, merged dataset with a 1km grid (Fig. 2a). A detailed description of these datasets, surveys, and data processing methodology can be found in the previous literature (Xiong et al., 2013). Such a merged dataset has been 145 corrected by the International Geomagnetic reference field model to eliminate the influence of the various latitude by differential reduction to pole method (RTP) in the Oasis Motaj software (https:// www. seequent. com/ products-solutions/ geosoft-oasis-montaj). As a result, the processed anomalies show a slight northward shift in the RTP anomaly map (Fig. 2b). To remove the effects of shallow magnetic

bodies, the aeromagnetic data were continued upward to 20 km above the Earth's surface (Fig. 3a). In

150 contrast, the first vertical derivative has been conducted to extract the anomalies derived from shallow magnetic bodies (Fig. 3b). The Bouguer gravity data used in this study are based on a compilation provided by Development Research Center of China Geological Survey (Zhang et al., 2011), covering the Longmenshan area with 1km grid (Fig. 4a). Such a dataset has been removed the regional gravity field by first vertical derivative method to enhance the density information from shallow crust.

155 **3.3 2D forward modeling**

The modeling of gravity and aeromagnetic data under the constraint of the velocity model produced an integrated and less ambiguous result. He et al. (2017) inversed three S-wave velocity models using the receiver function method based on the seismic data collected from 15 temporary seismological stations and 42 permanent seismological stations. These models crossed the earthquake epicenters and seismic gap show detailed velocity structure underneath the middle and southern segment of the LFZ. To create a reliable structure model accommodating three types of geophysical data, three S-wave velocity models were used to constrain the 2D magnetic and density modeling. The velocity model could provide information of Moho depth. Then, the physical properties of different rocks were modeled, the magnetic susceptibility contrast for the aeromagnetic data and the density for the gravity data. Meanwhile, this method easily adds geological and structural information from previous studies and the understanding of researchers during the modeling process.

The models were created using a 2D gravity and magnetic modeling package running on the Oasis Montaj Program. Sequential gravity-magnetic modeling was performed by defining the density surface in the crust according to the seismic models. The density values of the initial model referred to the previous studies (Wang et al., 2014b; Zhang et al., 2014). The initial magnetic susceptibility used in the modeling is a reference value from field observation to improve the interpretability of the model. For example, the magnetic susceptibilities of basement rocks in the Sichuan Basin refer to the values of outcropped Proterozoic intrusive rocks in the LFZ. Finally, the position, shape, dimensions, and contrast of the physical properties of the rocks were adjusted to obtain the best fit between the observed and calculated data.

3.4 3D inversion

180

To study the deep structure of the magnetic sources, we converted the aeromagnetic anomaly grid data into a subsurface susceptibility model. In this study, we applied 3D inversion of the aeromagnetic data (Li and Oldenburg, 1996), which has been widely used in geophysical exploration to create quantitative models of ore-related magnetic sources and deep structure (Aitken and Betts, 2009; Fullagar et al., 2004; Li et al., 1996; Lü et al., 2013; Oldenburg et al., 1997; Roy and Clowes, 2000; Silva et al., 2001; Wang et al., 2020a, b).

This inversion method assumes that the magnetic anomaly is caused by induced magnetization and there is no remnant magnetization. It discretizes the subsurface space into many rectangular cells with unknown susceptibilities. The inversion problem is formulated as an optimization problem. This method minimizes a trade-off between data misfit and a model norm subject to a positivity constraint. The objective function of this method is

$$\begin{aligned} & \text{Minimize } \phi = \parallel W_d(d-Gm) \parallel_2^2 + \parallel W_m(m-m_0) \parallel_2^2 \\ & \text{Subject to } m \geq 0 \end{aligned}$$

190 where μ is a regularization parameter, **d** is the observed data, **m** is the model, **G** is the sensitivity matrix, W_d is a diagonal data weighting matrix whose diagonal elements are reciprocals of estimated noise standard deviations, and W_m is a model weighting matrix that consists of a weighted sum of zerothand first-order finite difference matrixes. **m**₀ is a reference model.

The regularization parameter (μ) is following the traditional method. In 3D magnetic inversion,
there are two fitting processes including data fitting and model fitting. The regularization parameter is the ratio of the maximum value of data fitting to the maximum value of model fitting. In addition, The Generalized Cross Validation (GCV) method is also used to calculate the regularization parameters. The inversion results are basically consistent with the traditional method. The weighting matrix uses Depth weighting. The default setting for each iteration is stopping the iteration if the error is less than 0.001.

200 4. Results

4.1 Major features of aeromagnetic data

The Sichuan Basin is mainly characterized by NE-trending positive and negative banded magnetic anomalies of a large scale and high intensity (Fig. 2a). The positive anomalies are mostly 200-400 nT, and the negative anomaly is -300 nT. These features are different with the SGB and Yajiang Basin displaying low magnetic anomalies with intensities of 10-60 nT. The LFZ is characterized by a gradient zone of positive and negative anomalies on the aeromagnetic ΔT image that is different from the feature of Xianshuihe fault and sutures in this area (Fig. 2a). After reduction to the pole, the negative anomalies disappeared because of the elimination of oblique magnetization, and the LFZ moved above the banded positive anomalies (Fig. 2b). After 20 km of upward continuation, the regional magnetic anomaly is separated into two parts. The Sichuan Basin shows a large-scale high magnetic block in contrast to the broad and low magnetic in the SGB (Fig. 3a).

As a transition zone between the Sichuan Basin and SGB, the LFZ is characterized by a boundary of two magnetic anomaly areas, but it is mainly located in the magnetic anomaly area of the Sichuan Basin. This result strongly supports that the LFZ thrusts above the basement of the Sichuan Basin, and

- the magnetic boundary between the Sichuan Basin and the SGB is probably located northwest of the Wenchuan-Maoxian fault (F₁). More importantly, the fault zone can be divided into three segments according to the magnetic anomaly feature along the strike with the boundary of the Siguniangshan-Dayi and Gucheng-Wulian (I and II in Fig. 2a). The middle and southern segments are characterized by positive magnetic anomalies and associated negative anomalies on the aeromagnetic ΔT image, but the magnetic anomaly features are discontinuity separated by the boundary of the Siguniangshan-Dayi. The
- northern segment shows a small-scale linear magnetic anomaly zone on the negative background. The division of magnetic anomaly features is consistent with the segmentation based on the geological observation (Li et al., 2008). The seismic gap is located in the discontinuous area between the middle and southern segments. The Wenchuan earthquake and the Lushan earthquake occurred on the edge of
- two banded magnetic anomaly belts belonging to the middle and southern segments respectively.



Figure 2. Aeromagnetic anomaly feature of the LFZ and adjacent area. (a) Aeromagnetic ΔT anomaly image. (b) Reduction to the pole (RTP) image of the aeromagnetic ΔT data. I: boundary of Siguniangshan-Dayi; II: boundary of Gucheng-Wulian.



Figure 3. Potential field transformations of RTP aeromagnetic ΔT anomaly data. (a) 20 km upward continuation of RTP aeromagnetic ΔT data. (b) First vertical derivative of RTP aeromagnetic ΔT data.

In addition, there is an obvious magnetic discontinuous distributed from Daofu and Danba to

Chengdu, which separates two distinct magnetic anomaly areas (Fig. 2a). It is inferred that there is a

concealed fault named Daofu-Chengdu fault. The fault is divided into two segments by the Longmenshan fault zone. The western segment is distributed along Daofu - Qiaoqi and merges into the Xianshuihe fault in the northwest. The eastern segment started at Xiling, passed through Chengdu, and ended by the Longquanshan fault, which is characterized by an EW-trending linear discontinuity of magnetic anomalies extending approximately 90-100 km (Fig. 3b). It shows a small magnetic anomaly change after first vertical derivative calculation because the anomalies of the shallow structure were highlighted in the magnetic field. The Longquanshan fault (F₆) is a foreland uplift zone in the central Sichuan Basin. The fault doesn't show a magnetic gradient zone or discontinuous on the aeromagnetic anomaly image, which suggests that it is a shallow structure without cutting through the magnetic basement (Fig.2 and Fig.3).

245 4.2 Major features of gravity data

According to the Bouguer gravity anomaly image (Fig. 4a), the Sichuan Basin has a high Bouguer gravity anomaly with values ranging from -185 to -90 mgal. The SGB has a relatively low Bouguer gravity anomaly with values ranging from -435 to -250 mgal. The feature illustrates the crust of the Sichuan Basin has a higher density than that in the SGB. As a transition zone between two blocks, The

- 250 LFZ is characterized by a gradient zone with Bouguer gravity values of -290 to -185 mgal. From the first vertical derivate of the Bouguer gravity anomaly map, a NE-trending high gravity anomaly belt extends along the LFZ forming the western boundary of the Sichuan basin, which is probably caused by the high-density rocks in the upper crust.
- However, the gravity anomaly belt is discontinuous along the LFZ, which could be divided into two
 segments with the boundary of Siguniangshan-Dayi (I in Fig. 4a and 4b). The Bouguer gravity value of the middle segment is -250~-185 mgal, while that of the southern segment is -290~-215 mgal. Moreover, the first vertical derivate of Bouguer gravity anomaly in the middle segment is wider than that in the southern segment. The density differences along the LFZ might be caused by structural heterogeneities in the crust. The epicenter of the Wenchuan earthquake was located in a relatively high gravity anomaly area with Bouguer gravity values ranging from -215 to -225 mgal, while the Lushan earthquake was located in an area with low values of -250 to -260 mgal. The seismic gap is a transition zone between the

middle and southern segments. The Longquanshan fault doesn't show an obvious high-low gravity boundary in Figure 4a. However, it is characterized by an anomaly belt on the first vertical derivate image, which indicate the fault exist in the shallow crust.



265

Figure 4. (a) Bouguer gravity anomaly image of the LFZ and surrounding areas. (b) first vertical derivate of Bouguer gravity anomalies in the LFZ and adjacent areas. I: boundary of Siguniangshan-Dayi. The blue square is seismic gap.

4.3 Magnetic and gravity modeling

4.3.1 Profile AB

270 Profile AB passes through the epicenter of the Wenchuan earthquake (Fig. 5). The gravity anomaly feature show obvious change on both sides of the LFZ, and the value is slightly increased in the Longmenshan area caused by the uplift of a high-density geological body. The model shows high magnetic susceptibility and density in the crust of the Sichuan Basin, ranging in 0.0189 - 0.0440 SI and 2.65 - 2.75 g/cm³, respectively. The depth to the top of the magnetic basement is 5 - 7 km, and the thickness is approximately 15 - 20 km. The thickness of the magnetic basement is stable in the Sichuan Basin, but it gradually thins and disappears beneath the northwest of LFZ. In contrast, the crust of SGB shows low magnetic susceptibility and density, ranging in 0.0013 - 0.0019 SI and 2.68 - 2.70 g/cm³, respectively. The depth to the top of the magnetic basement is approximately 4 - 10 km, and the thickness is 12 - 18 km. There is a nonmagnetic area between the two basements with low density, which is a low-

280 velocity zone extending downward to the crustal low-velocity zone in the Vs model (He et al., 2017). This zone is probably a weak and brittle area formed by the collision of the Sichuan Basin and the SGB.

The model indicates that the crust of the Sichuan Basin has two high magnetic blocks (HMB-1 and HMB-2 in Fig. 5). The HMB-1 lies in the central Sichuan Basin with an average magnetic susceptibility of 0.0314 - 0.0440 SI, which corresponds to the NE-trending positive magnetic anomaly belt. The HMB-

1 dips to the southeast. The HMB-2 has an average magnetic susceptibility of 0.0189 SI and contact with the overlying high magnetic block in the southeast. Compared with the location of the Wenchuan-Maoxian fault on the Earth's surface, the magnetic basement of the Sichuan Basin obviously extends to the west of the Wenchuan-Maoxian fault with a distance of approximately 33 km. This result is consistent with the resistivity model inversed from magnetotelluric data and the velocity model from deep seismic reflection profile (Guo et al., 2013; Zhu et al., 2008). The Wenchuan earthquake and its aftershocks were distributed inside the magnetic basement of the Sichuan Basin.



Figure 5. Two-dimensional gravity and magnetic model and interpreted crustal structure along profile AB. ρ marks the density in units of g/cm³, and κ marks the magnetic susceptibility in units of 10⁻⁵SI. The black line is the boundary of the density block, and the blue line is the boundary of the magnetic block.

295

14

4.3.2. Profile CD

315

Profile CD passes through the epicenter of the Lushan earthquake (Fig. 6). In contrast to profile AB, the Bouguer gravity anomaly doesn't show obvious changes on both sides of the LFZ. The values only slightly increase in the Longmenshan area, which is caused by the upward thrusting of high-density geological bodies. However, the magnetic anomaly values show large changes on both sides of the LFZ. The values are high in the Sichuan Basin and decrease rapidly in the LFZ. Therefore, the model shows that the basement of the Sichuan Basin has high magnetic susceptibilities of 0.0250 - 0.0440 SI. The density is 2.65 - 2.70 g/cm³. The magnetic layer dips to the northwest. The depth to the top of the magnetic basement is approximately 5 - 11 km, and the thickness is approximately 17 - 23 km. In the west of the Wenchuan – Maoxian fault (F₁), there is a magnetic body with a moderate magnetic susceptibility of 0.0126 SI. The depth to the top of the magnetic body is 3 - 6 km, and the thickness is 16 - 20 km. The magnetic body is inferred to be intermediate-felsic intrusive rock because there are many Triassic and Jurassic granite, syenite, and granodiorite plutons outcropping on the surface with high magnetic susceptibility. The basement of the SGB has a low magnetic susceptibility of 0.0013-0.0038

310 SI. The depth to the top of the magnetic basement is approximately 4 - 7 km, and the thickness is approximately 13 - 16 km.

In this model, the magnetic basement of the Sichuan Basin extends approximately 19 km west of the Wenchuan – Miaoxian fault. And, the magnetic basement subducts beneath the LFZ at a high angle rather than a low angle in profile AB. The top of the magnetic basement shows large fluctuations, which suggests that the crystalline basement is highly thrusted and deformed in the southern segment of the LFZ. Moreover, there is no double-layer magnetic structure in this profile. The Lushan earthquake and its aftershocks are distributed in the rigid magnetic basement of the Sichuan Basin.



Figure 6. Two-dimensional gravity and magnetic model and interpreted crustal structure along profile CD. ρ marks the density in units of g/cm³, and κ marks the magnetic susceptibility in units of 10⁻⁵SI. The black line is the boundary of the density block, and the blue line is the boundary of the magnetic block.

4.3.3. Profile EF

Profile EF passes through the seismic gap (Fig. 7). The Bouguer gravity anomaly shows the same
pattern as profile AB, which displays obvious changes with the boundary of the LFZ. The basement of the Sichuan Basin has two high magnetic blocks as in profile AB. The HMB-1 is distributed in the central Sichuan Basin with a magnetic susceptibility of 0.0151 - 0.0377 SI. The depth to the top of the magnetic basement is 4 - 10 km, and the thickness is approximately 15 - 19 km. The HMB-2 is locally distributed beneath the Longmenshan area with a magnetic susceptibility of 0.0126 SI. In this model, the HMB-1 doesn't directly in contact with the HMB-2 and dips to the northwest. A large area of nonmagnetic sedimentary cover or basement is distributed between the two high magnetic blocks. A high magnetic anomaly peak in the northwest of LFZ, consistent with the outcropping location of the Siguniangshan granite, is modeled by a pluton with a magnetic susceptibility of 0.0126 SI. The pluton extends downward to the middle-upper crust from the surface, and the thickness is approximately 22 km. Particularly, unlike

335 the previous two profiles, the magnetic basement under the Longmenshan area does not have a complex thrust and nappe structure. And, the magnetic basement subducts approximately 17 km west of the Wenchuan-Miaoxian fault. The low-density layer in the middle and upper crust does not extend below the magnetic basement in the Sichuan Basin.



Figure 7. Two-dimensional gravity and magnetic model and interpreted crustal structure along profile EF. ρ marks the density in units of g/cm³, and κ marks the magnetic susceptibility in units of 10⁻⁵SI. The black line is the boundary of the density block, and the blue line is the boundary of the magnetic block.

4.4 3D inversion of aeromagnetic data

345 To make the deep structure of the LFZ visible, the subsurface space was divided into 280 × 256 × 200 cells forming a cube volume of 840 km× 768 km× 100 km. After 3D inversion, a susceptibility value for each cell was obtained. Cells with high susceptibility provide a good representation of magnetic sources for the magnetic highs. To observe the magnetic sources visually, Fig. 8 displays the inverted susceptibility model by an isosurface with a magnetic susceptibility value of 0.015 SI. We set the bottom 350 depth of the susceptibility model as 50km because the curie depth of this area is up to 45-50km (Xiong et al., 2016c). The inversion model shows that the isosurface delineates two high magnetic blocks under

the Sichuan basin (HMB1 and HMB2 in Fig. 8). The Wenchuan and Lushan earthquake lies on the different magnetic block. The seismic gap is characterized by a low magnetic area between two high magnetic blocks. It is worth noting that the magnetic block is uplifted to shallow surface beneath the LFZ which is well-matched with the outcropping of the Baoxing and Pengguan complex on the ground. The susceptibility model proved the strong crust shortening and basement deformation in western Sichuan basin.

355

The results of the inversion are shown by depth slices in the range of 10–40 km to provide physical property of rocks at different depth (Fig. 9). Figures 9a and 9b display that the high magnetic rocks are distributed in a small area beneath the LFZ. When the depth increases to 30 and 40km, two NE-trending high magnetic areas show a large scale in the Sichuan basin. Two earthquakes lie in two areas with local high magnetic susceptibility at 10 and 20km depth (Fig. 9a, b). The deep of the area show two high magnetic blocks at 30 and 40 km depth (Fig. 9c, d). It suggests that the magnetic bodies beneath the LFZ are shallow buried and extended to deep with large scale. The seismic gap is absent of high magnetic soft the inversion model along profiles AB, CD, and EF are shown in Fig. 10. The Wenchuan and Lushan earthquakes are near or in the high magnetic blocks, but the seismic gap shows a weak magnetic area which is the southwest end of the HMB2(Fig. 10a, b, c).



370 Figure 8. Three-dimensional susceptibility model of Longmenshan fault zone. HMB-1 and HMB-2: two high magnetic blocks in Sichuan basin.



Figure 9. Horizontal slices of the inverted susceptibility model at different depths from 10–40 km every 10 km.



Susceptibility(SI)

Figure 10. Vertical slice of the inversion model along line AB, CD and EF. Its location is marked in Fig. 2a.

375

5. Discussion

5.1 The source of aeromagnetic anomaly

Hu et al., 2005; Zhao et al., 2007a, b).

To understand the deep structure of the LFZ, geophysical data should be explained 380 comprehensively including aeromagnetic and gravity anomalies. Regionally, the LFZ shows similar magnetic and gravity anomaly feature to the Sichuan basin. The inversion models indicate that the basement of the Sichuan Basin has extended to the west of the Wenchuan-Maoxian fault and reached the deep of SGB. The results agreed with the previous conclusion that the LFZ has been thrust above the Sichuan Basin during the Late Indosinian-Early Yanshanian (Guo et al., 2013; Wang et al., 2015; Xiong et al., 2016a; Zhang et al., 2010; Zhu et al., 2008). The convergent process caused partial melting of the crust and formed a series of intermediate-felsic intrusive rocks in the SGB. The isotopic and chronology data show that these rocks have Proterozoic clastic zircon cores and Nd model ages (T_{DM}), which indicate that there is Proterozoic Yangtze-type continental crust beneath the Songpan-Ganzi area (Dai et al., 2011;

- 390 To find the source of the aeromagnetic anomaly, we are trying to analyze outcropped rock assemblage in the western margin of the Sichuan Basin. The outcropped Precambrian complexes, such as Kangding, Baoxing, and Pengguan, are well-matched with the high magnetic anomalies (Fig. 11). According to field observations of magnetic susceptibility, the quartz diorite, granite from Kangding Complex, and the Serpentine from Pengguan Complex usually have high magnetic susceptibility. These 395 Precambrian rock assemblages could produce large-scale and high magnetic anomalies. The Triassic-Jurassic Syenite and granite have moderate magnetic susceptibility in the SGB producing local magnetic anomalies. Based on the understanding of previous geological surveys, the basement is mainly composed of Neoarchean-Paleoproterozoic high-grade metamorphic rocks that are able to produce high magnetic anomaly in the Sichuan Basin, such as Kangding Group (SBGMR, 1991; Xiong et al., 2016a).
- 400 Recently, a large amount of geochronological and geochemical evidence has shown that the Kangding complex has arc signatures, representing metamorphic products of Neoproterozoic arc-related acidic plutons rather than Neoarchean and Paleoproterozoic crystalline basement (Chen et al., 2005; Du et al., 2007; Geng et al., 2007; Kang et al., 2017; Lai et al., 2015; Liu et al., 2009; Zhou et al., 2002). Meanwhile, zircon U–Pb data have shown that the Huangshuihe and Yanjing Groups also formed in the

- 405 Neoproterozoic, and the E'bian Group formed in the late Mesoproterozoic (Chen et al., 2018; Du et al., 2005; Ren et al., 2013). Moreover, granite and rhyodacite collected from the drill core of hydrocarbon exploration in the central Sichuan Basin have isotopic age of 701.5-794 Ma (Gu et al., 2014; Luo et al., 1986). These widespread Meso-Neoproterozoic rift-related magmatism and sedimentation records imply that Sichuan Basin probably played a key role in the assembly and breakup of the Rodinia supercontinent
- 410 (Cui et al., 2015). Therefore, the banded positive magnetic anomaly is probably related to Meso-Neoproterozoic magmatic events rather than the presence of rigid Neoarchean and Paleoproterozoic crystalline basements in the Sichuan Basin. The 2D and 3D models show two high magnetic blocks indicating there are probably two concealed Meso-Neoproterozoic igneous rock belts across the Sichuan Basin. These large-scale igneous rock belts formed the rigid blocks of the Sichuan Basin which subducted beneath the SGB and apt to accumulate stress.



Figure 11. RTP aeromagnetic ΔT anomaly contours and outcrops of Precambrian basement and intrusive rocks in 420 the western margin of the Sichuan Basin (modified after Du et al. (2014) and Geng et al. (2007)).

5.2 The crustal structure of the LFZ

445

The gravity and magnetic anomalies show obvious lateral changes along the strike of the LFZ, which can be divided into southern, middle, and northern segments (Fig. 2a and 4). The southern segment is characterized by the magnetic anomaly gradient zone and the low Bouguer gravity anomaly, while the middle segment has a magnetic anomaly gradient zone and high Bouguer gravity anomaly. The Wenchuan and Lushan earthquakes were distributed in the middle and southern segments respectively. The northern segment is characterized by a negative magnetic field with some of the linear magnetic anomaly zone superimposed. The division of aeromagnetic and gravity anomaly features is the same as the surface deformation segmentation of the LFZ (Li et al., 2008). Therefore, the deep structure reflected 430 by aeromagnetic and gravity anomaly probably controlled the evolution and deformation of the LFZ.

The 2D and 3D inversion of multi-sourced geophysical data suggests that the basement of the Sichuan Basin is heterogeneous in magnetic and density (Fig. 8, 9, 10, and 12). There are two NE-trending high magnetic blocks with different scales and magnetic anomaly intensities. The Wenchuan earthquake lies on the HMB-2 with a small scale and low intensity. The Lushan earthquake lies on the HMB-1 with a large scale and high intensity. The seismic gap is characterized by a low or non-magnetic area between two blocks. According to the 2D models, the magnetic basement subducts beneath the LFZ with a distance of approximately 33 km west of the Wenchuan-Maoxian fault in profile AB. However, the distances are 17 km and 19 km in profiles CD and EF, respectively. Therefore, the subducted distance of the basement has a large lateral change along the western margin of the Sichuan Basin. The basement beneath the middle segment of the LFZ wedges farther than that under the southern segment, forming a "stair-shaped" along the LFZ (Fig. 12).

The 2D models indicate that the two disastrous earthquakes and their aftershocks were mainly distributed in the magnetic basement of the Sichuan Basin (Fig. 9). Moreover, the 3D magnetic susceptibility model shows an obvious uplift beneath the LFZ suggesting the magnetic basement has undergone a strong deformation (Fig. 8 and 10). The feature is also proved by the highly shortened crystalline crust of the Yangtze block shown by the deep seismic reflection profile (Guo et al., 2013). Meanwhile, the thickness of the magnetic basement decreases gradually beneath the LFZ when the Yangtze's crust wedges into the eastern Tibetan Plateau (Fig. 5, 6, and 7). These characteristics indicate

that the crust of the Sichuan Basin has undergone strong elastic shortening. The seismic images show

450 that the crustal low-velocity zone extends beneath the epicenters of the Wenchuan and Lushan earthquakes (He et al., 2017). Therefore, the occurrence of the two earthquakes may be closely related to the destruction of the rigid magnetic basement through the detached upper crust of the SGB collided with the crust of the Sichuan Basin.



Figure 12. Western margin of the magnetic basement in the Sichuan Basin.

5.2 Seismogenesis mechanism of the LFZ

455

The Wenchuan earthquake occurred in the Yingxiu-Beichuan fault, which is part of the centralfront range fault system in the middle segment of the LFZ. The Lushan earthquake occurred in a blind reverse fault east of the Shuangshi-Dachuan fault, which is part of the front range fault system in the southern segment of the LFZ. Although the two segments are separated by seismic gaps with small distances, the geological deformation is quite different. The front-range structure in the southern segment is much more complicated than that in the middle segment. The range of the latest structural deformation increases from 30 km in the middle segment to 150 km in the southern segment (Xu et al., 2013). The

- 465 focal mechanism shows that the Lushan earthquake was a pure thrust event without obvious ruptures on both sides (Chen et al., 2013). The axis of the maximum horizontal stress is oriented NW-SE (Luo et al., 2015). However, the Wenchuan earthquake was dominated by thrusts with a dextral strike-slip component. The surface rupture mainly extends northeast with a distance of approximately 300 km. The axis of the maximum horizontal stress occurs in several different directions (Luo et al., 2015). Moreover,
- 470 high-resolution geodetic data show that the surface deformation has obvious changes on both sides of the seismic gap (Wang et al., 2011). These features indicate that the deformation mechanisms of the middle and southern segments are different. Therefore, several velocity and resistivity models were created to explain the genesis of the seismic gap and the deformation mechanism. The researches commonly indicate the presence of ductile zones or partial melting under the gap area (Liang et al., 2018; Pei et al.,
- 475 2014; Wang et al., 2014c). In this study, integrated 2D and 3D models suggest that the downward subducted basement of the Sichuan Basin is irregular in shape and heterogeneous in magnetic and density. The different focal mechanisms of the two earthquakes and the genesis of the seismic gap may be closely related to the differential thrusting mechanism caused by basement heterogeneity in the western margin of the Sichuan Basin.
- 480 Two earthquakes and their aftershocks occurred in the rigid magnetic basement of the Sichuan Basin, which is characterized by high-velocity areas in seismic images (Wang et al., 2015). There is a seismic gap with low Vp and Vs, high Poisson's ratio, and high conductivity between the two earthquakes. It is inferred to be a fluid-rich ductile crust extending to the middle and lower crust (Pei et al., 2014; Wang et al., 2015; Zhan et al., 2013). Consequently, compared with the magnetic basement under the earthquake epicenter, there is no high magnetic basement beneath the seismic gap. The inferred Daofu-Chengdu fault crosses the seismic gap. The fault could be divided into two segments. The western segment may be related to a tear zone and accompanied by the intrusion of intermediate-felsic rocks. The S-wave velocity models indicate that there is an NW-trending Moho uplift from the seismic gap to the

northwest, which is a fault zone that caused mantle upwelling and partial melting (He et al., 2017; Liang

et al., 2018). The eastern segment of the Daofu-Chengdu fault cuts the basement of the Sichuan Basin and extends to Longquanshan Mountain. The sedimentary covers are commonly nonmagnetic in the Sichuan Basin and cannot cause changes in magnetic anomalies. The magnetic anomaly decays obviously after 20 km upward continuation, which suggests that the displacement is small on both sides of the fault. The fault may be closely related to the early thrust in the southern segment of the LFZ and the uplift of Longquanshan Mountain.

This study proposes a schematic model (Fig. 13). The rigid basement of the Sichuan Basin subducts beneath the Songpan-Ganzi fold belt during the Late Indochina-early Yanshanian. The basement beneath the middle and northern segments of the LFZ extends further than under the southern segment. The heterogeneity of the basement caused the Songpan-Ganzi fold belt to tear into two pieces with different geodynamic systems (Slab 1 and Slab 2 in Fig. 13). Different geodynamic systems lead to different activities of the two slabs that form a tear zone for the emplacement of intermediate-acid intrusive rocks. This kind of mechanism is also found in the southern Tibetan Plateau which is characterized by the tearing and dischronal subduction of the Indian continental slab (Hou et al., 2006, 2011). With the continuous uplift of the Tibetan Plateau, the compression stress is increasing in the western margin of the Sichuan Basin. In 2008 and 2013, two earthquakes with different focal mechanisms occurred

505 Sichuan Basin. In 2008 and 2013, two earthquakes with different focal mechanisms occurred successively in the middle and southern segments of the LFZ. These two earthquakes were produced by two slabs thrust above the "stair-shaped" magnetic basement. Therefore, the Wenchuan earthquake occurred in the middle and northern segments of the LFZ, and the Lushan earthquake was restricted to the southern segment. Apparently, the central Sichuan Basin may be involved in the early thrust process 510 of the southern segment of the LFZ, which is represented by the displacement of the basement and formation of the Longquanshan fault (F₆).

The geodynamic process of the two earthquakes is summarized as follows:

(1) With the uplift of the Tibetan Plateau, the Songpan-Ganzi block continued to move toward the southeast and collided with the basement of the Sichuan Basin. Compression stresses were prior accumulation in the middle-northern segment because the basement beneath the middle and northern segment wedges farther than under the southern segment. Finally, the Wenchuan Ms 8.0 earthquake was

triggered in the middle segment of the LFZ. Due to different geodynamics between the middle and southern segments of the LFZ and the constraints of the tear zone and the Siguniangshan pluton, no large-scale surface rupture occurred to the southwest of the Wenchuan earthquake. However, the rupture extended approximately 340 km toward the northeast.

520

525

(2) The compression stress was completely released in the middle and northern segments of the LFZ after the happening of Wenchuan earthquake. The stress shifted to the southern segment and finally triggered the Lushan Ms7.0 earthquake. Due to the constraints of the heterogenous basement and the Xianshuihe fault on both sides, the Lushan earthquake was a pure thrust event without obvious rupture on both sides.



Figure 13. Schematic deep structure model of the Longmenshan fault zone and adjacent areas.

6. Conclusions

The aeromagnetic and gravity anomaly revealed significant variations in the physical property of 530 the crust along the LFZ. The anomaly feature of the LFZ could be divided into three segments that are consistent with the surface deformation. 2D forward modeling and 3D inversion obtained a comprehensive magnetic and density model in the LFZ and adjacent area. The result suggests that two
high magnetic blocks of the Sichuan Basin are subducted beneath the LFZ, which is inferred as MesoNeoproterozoic igneous rock belt apt to accumulate stress. The Wenchuan earthquake lies on a small
scale and moderate magnetic block while the Lushan earthquake lies on a large scale and high magnetic
block. There is a low or non-magnetic area that is void aftershocks for both earthquakes. The lateral
change of magnetic and gravity anomaly indicates structural heterogeneity along the LFZ. Moreover, the

magnetic basement beneath the middle segment wedges further and has a lower dip angle than under the

southern segment. The magnetic basement involves more intense deformation beneath the epicenter of the two earthquakes than under the seismic gap. Due to the irregular shape of the basement and the constraints of the tear zone and intermediate-acid intrusive rocks, the thrust mechanism is different in the middle and southern segments of the LFZ. It provides an essential tectonic framework for the genesis of the two earthquakes with different focal mechanisms. This study provides new geophysical evidence for mapping the deep structure of the LFZ.

545 Data availability

535

Data will be made available on request. The use of the Oasis Montaj software is authorized by Beijing Maiqin Nengyuan Jishufuwu Co., Ltd. (https://www.seequent.com/products-solutions/geosoft-oasis-montaj/).

Author contribution

550 Hai Yang was responsible for the idea and methodology of the study, field investigation, compilation and analysis of data, and writing the paper. Shengqing Xiong was responsible for the acquisition and management of financial support, contribution to experimental design. Qiankun Liu and Fang Li investigated and revised the ideas of the article. Zhiye Jia and Xue Yang processed the aeromagnetic and gravity data. Haofei Yan and Zhaoliang Li was responsible for the 3D inversion. All authors contributed to the review of the manuscript.

Competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Professor Chuntao Liang and Fujun He from Chengdu University of Technology for providing seismic imaging results and enthusiastic help. Thanks to all of our colleagues for their hard work. This research was supported by the National Natural Science Foundation of China grant (U2244220) and China Geological Survey Project grant (DD20190551, DD20230351). Thank you to the anonymous reviewers for their hard work and constructive comments.

565 References

Aitken, A. R. A. and Betts, P. G.: Multi - scale integrated structural and aeromagnetic analysis to guide tectonic models: An example from the eastern Musgrave Province, Central Australia, Tectonophysics, 476(3–4), 418–435, doi: 10.1016/j.tecto.2009.07.007, 2009.

Burchfiel, B.C., Chen, Z., Liu, Y., and Royden, L.H.: Tectonic of the Longmenshan and adjacent regions,
Central China, Int. Geol. Rev. 37, 661–735, doi: 10.1080/00206819509465424, 1995.

Burchfiel, B.C., Royden, L.H., van der Hilst, R.D., Hager, B.H., Chen, Z., King, R.W., Li, C., Lu, J., Yao, H., and Kirby, E.: A geological and geophysical context for the Wenchuan earthquake of 12 May 2008, Sichuan, People's Republic of China, GSA Today 18, 4–11, doi:10.1130/GSATG18A.1, 2008.

Chen, F.L., Xie, Y., Cui, X.Z., Sun, Z.M., Ren, G.M., and Deng, Q.: Geochronology, geochemistry and
tectonic implications of basalts from the Ebian Group in the western Yangtze block, South China, J.
Mineral Petrol., 38, 76-86, 2018. (in Chinese with English abstract)

Chen, L.C., Ran, Y.K., Wang, H., Li, Y.B., and Ma, X.Q.: The Lushan Ms7.0 earthquake and activity of the southern segment of the Longmenshan fault zone, Chin. Sci. Bull., 58, 3475–3482, doi: 10.1007/s11434-013-6009-6, 2013.

580 Chen, S. and Wilson, C.J.L.: Emplacement of the Longmen Shan Thrust–Nappe Belt along the eastern margin of the Tibetan Plateau, J. Struct. Geol. 18, 413–430, doi: 10.1016/0191-8141(95)00096-V, 2008.

Chen, Y.L., Luo, Z.H., Zhao, J.X., Li, Z.H., Zhang, H.F., and Song, B.: Petrogenesis and dating of the Kangding complex, Sichuan province, Sci. China Ser. D, 48, 622–634, doi: 10.1360/03yd0312, 2005.

Cui, X.Z.; Jiang, X.S.; Wang, J.; Wang, X.C.; Zhuo, J.W.; Deng, Q.; Liao, S.Y.; Wu, H.; Jiang, Z.F.; and
Wei, Y.N.: Mid-Neoproterozoic diabase dykes from Xide in the western Yangtze block, south china: new

evidence for continental rifting related to the breakup of Rodinia supercontinent, Precambrian Res., 268, 339-356, doi: 10.1016/j.precamres.2015.07.017, 2015.

Dai, Z.M., Sun, C.M., Zhang, K.Z., and Li, Z.J.: U-Pb dating of zircons from the Four-Girl Mountain Pluton in the Songpan-Ganze Terrane, and the relationship between the pluton and the Wenchuan Ms8.0 earthquake of 2008, Geology in China, 38, 623-636, doi: 10.3969/j.issn.1000-3657.2011.03.011, 2011. (in Chinese with English abstract)

Du, F., Long, F., Ruan, X., Yi, G.X., Gong, Y., Zhao, M., Zhang, Z.W., Qiao, H.Z., Wang, Z., and Wu, J.: The M7.0 Lushan earthquake and the relationship with the M8.0 Wenchuan earthquake in Sichuan, China, Chinese J. of Geophys., 56, 1772-1783, doi: 10.6038/cjg20130535, 2013. (in Chinese with English abstract)

595

590

605

615

Du, L.L., Geng, Y.S., Yang, C.H., Wang, X.S., Ren, L.D., and Zhou, X.W.: Geochemistry and SHRIMP U-Pb zircon chronology of basalts from the Yanbian Group in the Western Yangtze Block, Acta Geol. Sin., 79, 805–813, doi: 10.3321/j.issn:0001-5717.2005.06.009, 2005. (in Chinese with English abstract).

Du, L.L., Geng, Y.S., Yang, C.H., Wang, X.S., Zhou, X.W., and Ren, L.D.: New understanding on 600 Kangding Group on western margin of Yangtze Block: evidence from geochemistry and chronology, Acta Geol. Sin., 81, 1562–1577, doi: 10.3321/j.issn:0001-5717.2007.11.011, 2007. (in Chinese with English abstract).

Du, L.L., Guo, J.H., Nutman, A.P., Wymand, D., Geng, Y.S., and Yang, C.H.: Implications for Rodinia reconstructions for the initiation of Neoproterozoic subduction at ~860 Ma on the western margin of the Yangtze Block: Evidence from the Guandaoshan Pluton, Lithos, 196-197, 67-82, doi: 10.1016/j.lithos.2014.03.002, 2014.

Fullagar, P. K., Pears, G., Hutton, D., and Thompson, A. D.: 3D gravity and aeromagnetic inversion for MVT lead - zinc exploration at Pillara, Western Australia, Explor. Geophys., 35(2), 142-146, doi: 10.1071/EG04142, 2004.

610 Geng, Y.S., Yang, C.H., Wang, X.S., Ren, L.D., Du, L.L., and Zhou, X.W.: Age of crystalline basement in western margin of Yangtze Block, Geological Journal of China Universities, 13, 429-441, 2007. (in Chinese with English abstract).

Gu, Z.D., Zhang, W., and Yuan, M.: Zircon SHRIMP U-Pb dating of basal granite and its geological significance in Weiyuan area of Sichuan Basin, Chinese J. of Geol., 49, 202-213, doi: 10.3969/j.issn.0563-5020.2014.01.015, 2014. (in Chinese with English abstract)

Guo, X.Y., Gao, R., Randy Keller, G., Xu, X., Wang, H.Y., and Li, W.H.: Imaging the crustal structure beneath the eastern Tibetan Plateau and implications for the uplift of the Longmen Shan range, Earth Planet. Sci. Lett., 379, 72-80, doi: 10.1016/j.epsl.2013.08.005, 2013.

He, F.J., Liang, C.T., Yang, Y.H., Fang, L.H., and Su, J.R.: The crust structure of the unruptured segment 620 between Wenchuan and Lushan earthquake revealed by receiver functions, Chinese J. Geophys., 60, 2130-2146, doi: 10.6038/cjg20170609, 2017. (in Chinese with English abstract)

Hou, Z.Q., Zhao, Z.D., Gao, Y.F., Yang, Z.M., and Jiang, W.: Tearing and dischronal subduction of the

Indian continental slab: Evidence from Cenozoic Gangdese volcano-magmatic rocks in south Tibet, Acta Petrol. Sin., 22, 761-774, doi: 10.3969/j.issn.1000-0569.2006.04.001, 2006. (in Chinese with English abstract)

625 abstrac

650

Hou, Z.Q., Zhang, H.R., Pan, X.F., and Yang, Z.M.: Porphyry Cu (-Mo-Au) systems in non-arc settings: Examples from the Tibetan-Himalyan orogens and the Yangtze block, Ore Geology Reviews, 39, 21–45, doi: 10.1016/j.oregeorev.2010.09.002, 2011.

Hu, J.M., Meng, Q.R., Shi, Y.R., and Qu, H.J.: SHRIMP U-Pb Dating of zircons from granitoid bodies
in the Songpan-Ganzi terrane and its implication, Acta Petrol. Sin., 21, 867-880, doi: 10.3321/j.issn:1000-0569.2005.03.026, 2005. (in Chinese with English abstract)

Kang H., Li D.P., Chen Y.L., Hu G.Q., and Deng W.B.: Origin and tectonic implications of Kangding intrusive complexes in Sichuan Province: Evidence from zircon Hf isotope, Geol. in China, 44(6): 1175-1189, doi: 10.12029/gc20170610, 2017. (in Chinese with English abstract).

- 635 Lai, S.C., Qin, J.F., Zhu, R.Z, and Zhao, S.W.: Neoproterozoic quartz monzodiorite–granodiorite association from the Luding–Kangding area: Implications for the interpretation of an active continental margin along the Yangtze Block (South China Block), Precambrian Res., 267, 196-208, doi: 10.1016/j.precamres.2015.06.016, 2015.
- Lei, J., and Zhao D.: Structural heterogeneity of the Longmenshan fault zone and the mechanism of the
 2008 Wenchuan earthquake (Ms 8.0), Geochem. Geophys. Geosyst., 10(10), 1-17.
 doi:10.1029/2009GC002590, 2010.

Li, Y., and Oldenburg, D. W.: 3-D inversion of magnetic data. Geophysics, 61, 394-408, doi: 10.1190/1.1443968, 1996.

Li, Y.Q., Jia, D., Wang, M.M., Shaw, J.H., He, J.K., and Lin, A.M.: Structural geometry of the source
region for the 2013 Mw 6.6 Lushan earthquake: Implication for earthquake hazard assessment along the
Longmen Shan, Earth Planet. Sci. Lett., 390, 275–286, doi: 10.1016/j.epsl.2014.01.018, 2014.

Li, Z.W., Liu, S.G., Chen, H.D., Liu, S., Guo, B., and Tian, X.B.: Structural segmentation and zonation and differential deformation across and along the Lomgmen thrust belt, West Sichuan, China, Journal of Chengdu University of Technology (Science & Technology Edition), 35, 440-454, doi: 10.3969/j.issn.1671-9727.2008.04.014, 2008. (in Chinese with English abstract)

Li, Z.W., Tian, B.F., Sha, L., and Yang, J.S.: Asperity of the 2013 Lushan earthquake in the eastern margin of Tibetan Plateau from seismic tomography and aftershock relocation, Geophys. J. Int., 195, 2016-2022, doi: 10.1093/gji/ggt370, 2013.

Liang, C.T., Huang, Y.L., Wang, C.L., Liu, Z.Q., Yang, Y.H., Wu, J., and He, F.J.: Progress in the studies
of the seismic gap between the 2008 Wenchuan and 2013 Lushan earthquakes, Chinese J. of Geophy.,
61, 1996-2010, doi: 10.6038/cjg2018M0254, 2018. (in Chinese with English abstract)

Liu, S.W., Yan, Q.R., Li, Q.G., and Wang, Z.Q.: Petrogenesis of granitoid rocks in the Kangding Complex, western margin of the Yangtze Craton and its tectonic significance, Acta Petrol. Sin., 25, 1883–1896, doi: 10.2110/palo. 2009. p09-040r, 2009. (in Chinese with English abstract).

660 Liu, Z.Q., Liang, C.T., Hua, Q., Li, Y., Yang, Y.H., He, F.J., and Fang, L.H.: The seismic potential in the seismic gap between the Wenchuan and Lushan earthquakes revealed by the joint inversion of receiver functions and ambient noise data, Tectonics, 37, 4226-4238, doi: 10.1029/2018TC005151, 2018.

Lü, Q., Qi, G., and Yan, J.: 3D geologic model of Shizishan ore field constrained by gravity and magnetic interactive modeling: A case history, Geophysics, 78(1), 25–35, doi: 10.1190/GEO2012–0126.1, 2013.

665 Luo, Y., Zhao, L., Zeng, X.F., and Gao, Y.: Focal mechanisms of the Lushan earthquake sequence and spatial variation of the stress field, Sci. China Earth Sci., 58(7),1148-1158, doi: doi: 10.1007/s11430-014-5017-y, 2015.

Luo, Z.L.: Is there a paleo continental nucleus in central Sichuan? Journal of Chengdu College of Geology, 13, 65-73, 1986. (in Chinese with English abstract)

670 Oldenburg, D. W., Li, Y., and Ellis, R. G. Inversion of geophysical data over a copper gold porphyry deposit: A case history for Mt. Milligan, Geophysics, 62(5), 1419–1431, doi: 10.1190/1.1444246, 1997.

Parsons, T., Ji, C., and Kirby, E.: Stress changes from the 2008 Wenchuan earthquake and increased hazard in the Sichuan basin, Nature, 1–2. doi:10.1038/nature07177, 2008.

Pei, S., Su, J., Zhang, H., Sun, Y., Toksoz, M., Wang, Z., and Gao, X.: Three-dimensional seismic velocity
structure across the 2008 Wenchuan Ms 8.0 earthquake, Sichuan, China, Tectonophysics, 491, 211–217, doi: 10.1016/j.tecto.2009.08.039, 2010.

Pei, S.P., Zhang, H.J., Su, J.R., and Cui, Z.X.: Ductile Gap between the Wenchuan and Lushan Earthquakes Revealed from the Two-dimensional Pg Seismic Tomography, Sci. Rep., 4, 6489, doi: 10.1038/srep06489, 2014.

- 680 Ren, G.M., Pang, W.H., Sun, Z.M., and Yin, F.G.: Zircon SHRIMP U-Pb dating of basalt from Huangshuihe Group on the western margin of the Yangtze block and its geological significance, Geol. in China, 40, 1007-1015, doi: 10.3969/j.issn.1000-3657.2013.04.002, 2013. (in Chinese with English abstract)
- Roger, F., Jolivet, M., and Malavieille, J.: The tectonic evolution of the Songpan Garzê (North Tibet) and
 adjacent areas from Proterozoic to Present: a synthesis, J. Asian Earth Sci. 39, 254–269, doi: 10.1016/j.jseaes.2010.03.008, 2010.

Roy, B. and Clowes, R. M.: Seismic and potential - field imaging of the Guichon Creek batholith, British Columbia, Canada, to delineate structures hosting porphyry copper deposits, Geophysics, 65(5), 1418–1434, doi: 10.1190/1.1444831, 2000.

690 Royden, L.H., Burchfiel, B.C., and van der Hilst, R.D.: The geological evolution of the Tibetan Plateau, Science, 321, 1054–1958, doi: 10.1126/science.1155371, 2008.

Shan, B., Xiong, X., Zheng, Y., Jin, B.K., Liu, C.L., Xie, Z.J., and Hsu, H.T.: Stress changes on major faults caused by 2013 Lushan earthquake and its relationship with 2008 Wenchuan earthquake, Sci. China Earth Sci., 56, 1169–1176, doi10.1007/s11430-013-4642-1, 2013

695 Sichuan Bureau of Geology and Mineral Resources (SBGMR).: Regional Geology of Sichuan Province,

Beijing: Geological Publishing House, 1991. (in Chinese).

705

725

Silva, J. B. C., Medeiros, W. E., and Barbosa, V. C. F.: Potential - field inversion: Choosing the appropriate technique to solve a geologic problem, Geophysics, 66(2), 511, doi: 10.1190/1.1444941, 2001.

Wang, J., Yao, C., and Li, Z.: Aeromagnetic anomalies in central Yarlung - Zangbo suture zone (Southern Tibet) and their geological origins, J. Geophys. Res.: Sol. Ea., 125, e2019JB017351, doi: 10.1029/2019JB017351, 2020a.

Wang, J., Yao, C.L., Li, Z.L., Zheng, Y.M., Shen, X.H., Zeren, Z.M., and Liu, W.L.: 3D inversion of the Sichuan Basin magnetic anomaly in South China and its geological significance, Earth, Planets and Space, 72:40, doi: 10.1186/s40623-020-01167-5, 2020b

Wang, J.J., Xu, C.J., Freymueller, J. T., and Li, Z.H.: Probing Coulomb stress triggering effects for a Mw N 6.0 earthquake sequence from 1997 to 2014 along the periphery of the Bayan Har block on the Tibetan Plateau, Tectonophysics, 694, 249–267, doi: 10.1016/j.tecto.2016.11.009, 2017a.

Wang, M., Jia, D., Shaw, J.H., Hubbard, J., Plesch, A., Li, Y., and Liu, B.: The 2013 Lushan earthquake:
710 Implications for seismic hazards posed by the Range front blind thrust in the Sichuan Basin China, Geology, 42, 915–918, doi: 10.1130/G35809.1, 2014a.

Wang, P., Zhang, Z.J., Zhang, X., Han, Y.Y., Wang, M.L., Hou, J., and Xu, T.: Crustal density structure of the central Longmenshan and adjacent area and its geodynamic implications, Acta Petrol. Sin., 30, 1179-1187, 2014b. (in Chinese with English abstract)

715 Wang, Q., Qiao, X.J., Lan, Q.G., Jeffrey, F., Yang, S.M., Xu, C.J., et al.: Rupture of deep faults in the 2008Wenchuan earthquake and uplift of the Longmen Shan, Nat. Geosci., 4, 630-640, doi: 10.1038/ngeo1210, 2011.

Wang, Z., Huang, R.Q., and Pei, S.P.: Crustal defrormation along the Longmen-Shan fault zone and its implications for seismogenesis, Tectonophysics, 610, 128-137, doi: 10.1016/j.tecto.2013.11.004, 2014c.

Wang, Z., Su, J.R., Liu, C.X., and Cai, X.L.: New insights into the generation of the 2013 Lushan Earthquake (Ms 7.0), China, J. Geophys. Res. Sol. Ea., 120, 3507–3526, doi: 10.1002/2014JB011692, 2015.

Wang, Z., Wang, X.B., Huang, R.Q., and Liu, G.N.: Deep structure imaging of multi-geophysical parameters and seismogenesis in the Longmenshan fault zone, Chinese J. Geophys., 60, 2068-2079, doi: 10.6038/cjg20170604, 2017b. (in Chinese with English abstract)

Wang, Z., Y. Fukao, and Pei, S.: Structural control of rupturing of the Mw7.9 2008 Wenchuan Earthquake China, Earth Planet. Sci. Lett., 279, 131–138, doi: 10.1016/j.epsl.2008.12.038, 2009.

Wu, Y.Q., Jiang, Z.S., Wang, M. Chen, S., Liao, H., and Li, Q.: Preliminary results pertaining to coseismic displacement and preseismic strain accumulation of the Lushan Ms 7.0 earthquake as reflected by GPS
surveying, Chinese Sci. Bull., 58, 3460–3466, doi: 10.1007/s11434-013-5998-5, 2013.

Xiong, S.Q.: Chinese continent aeromagnetic and geological structure feature, Beijing: Geological

Publishing House, 1-370, 2016a. (in Chinese)

750

Xiong, S.Q., Fan, Z.G., Zhang, H.R., Guo, Z.H., Huang, X.Z., and Ding, Y.Y.: Chinese continent aeromagnetic map and its introduction, Beijing: Geological Publishing House, pp. 19, 2013. (in Chinese)

Xiong, S.Q., Yang, H., Ding, Y.Y., Li, Z.K., and Li, W.: Distribution of igneous rocks in China revealed by aeromagnetic data, J. Asian Earth Sci., 129, 231-242, doi: 10.1016/j.jseaes.2016.08.016, 2016b.

Xiong, S.Q., Yang, H., Ding, Y.Y., Li, Z.K.: Characteristics of Chinese continent curie point isotherm, Chinese J. Geophys., 59, 643-657, doi: 10.6038/cjg20161008, 2016c.

Xu, X.W., Chen, G.H., Yu, G.H., Cheng, J., Tan, X.B., Zhu, A.L., and Wen, X.Z.: Seismogenic structure
of Lushan earthquake and its relationship with Wenchuan earthquake, Earth Science Frontiers, 20, 11-20, 2013. (in Chinese with English abstract)

Yan, L.: Study on the active tectonics in Longmen Mountain area and surface rupture of Wenchuan earthquake, Chengdu: Chengdu university of Technology, PhD. Thesis, 1-114, 2011. (in Chinese with English abstract)

Yan, Y.F., Teng, J.W., Ruan, X.M., and Hu, G.Z.: Aeromagnetic field characteristics and the Wenchuan earthquake in the Longmenshan Mountains and adjacent areas, Chinese J. Geophys., 59, 197-214, doi: 10.6038/cjg20160117, 2016. (in Chinese with English abstract)

Zhan, Y., Zhao, G.Z., Unsworth, M., Wang, L.F., Chen, X.B., and Li, T.: Deep structure beneath the southwestern section of the Longmenshan fault zone and seimogenetic context of the 4.20 Lushan Ms7.0 earthquake, Chinese Sci. Bull., 58, 3467–3474, 2013.

Zhang, H.F., Zhang, L., Harris, N., Jin, L.L., and Yuan, H.: U-Pb zircon ages, geochemical and isotopic compositions of granitoids in Songpan-Garze fold belt, eastern Tibetan Plateau: constraints on petrogenesis and tectonic evolution of the basement, Contrib. Mineral Petrol. 152, 75–88, doi: 10.1007/s00410-006-0095-2, 2006.

755 Zhang, J.S., Gao, R., Zeng, L.S., Li, Q.S., Guan, Y., and He, R.Z.: Relationship between characteristics of gravity and magnetic anomalies and the earthquakes in the Longmenshan range and adjacent areas, Tectonophysics, 491, 218-229, doi: 10.1016/j.tecto.2009.12.004, 2010.

Zhang, M.H., He, H., and Wang, C.X.: The launch of a large regional gravity information system in China, Applied Geophysics, 8, 170-175, doi: 10.1007/s11770-011-0287-z, 2011.

Zhang, Y.Q., Teng, J.W., Wang, Q.S., and Hu, G.Z.: Density structure and isostatic state of the crust in the Longmenshan and adjacent areas, Tectonophysics, 619–620, 51–57, doi: 10.1016/j.tecto.2013.08.018, 2014.

Zhao, C.P., Zhou, L.Q., and Chen, Z.L.: Source rupture process of Lushan Ms7.0 earthquake, Sichuan, China and its tectonic implication, Chin Sci. Bull., 58, 1894-1900, 2013.

765 Zhao, Y.J., Yuan, C., Zhou, M.F., Yan, D.P., Long, X.P., and Li, J.L.: Geochemistry and petrogenesis of Laojungou and Mengtonggou granites in western Sichuan, China: constraints on the nature of Songpan-Ganzi basement, Acta Petrologica Sinica, 23, 995-1006, doi: 10.3321/j.issn:1000-0569.2007.05.013, 2007a. (in Chinese with English abstract)

775

780

Zhao, Y.J., Yuan, C., Zhou, M.F., Yan, D.P., Long, X.P., and Cai, K.D.: Post-orogenic extension of
Songpan-Garze orogen in Early Jurassic: Constraints from Niuxingou monzodiorite and Siguniangshan
A-type granite of western Sichuan, China, Geochimica, 36, 139-152, doi: 10.3321/j.issn:03791726.2007.02.003, 2007b. (in Chinese with English abstract)

Zhou, M.F., Yan, D.P., Kennedy, A.K., Li, Y.Q., and Ding J.: SHRIMP U-Pb zircon geochronological and geochemical evidence for Neoproterozoic arc-magmatism along the western margin of the Yangtze Block, South China, Earth and Planet. Sci. Lett., 196, 51-67, doi: 10.1016/S0012-821X(01)00595-7. 2002.

Zhu, S.B.: Is the 2013 Lushan earthquake (Mw = 6.6) a strong aftershock of the2008 Wenchuan, China mainshock (Mw = 7.9)? J. Geodyn., 99, 16-26, doi: 10.1016/j.jog.2016.05.002, 2016.

Zhu, Y.T., Wang, X.B., Yu, N., Gao, S.Q., Li, K., and Shi, Y.J.: Deep structure of magnetotelluric profile on Longmen Mts. and its relation to the Ms8.0 Wenchuan earthquake, Acta Geol. Sin., 82, 1769-1777, 2008. (in Chinese with English abstract)