≥ Draftable

Draftable Comparison Export

This document is an exported comparison with limited functionality, generated by Draftable Desktop. To access full functionality, use Draftable's powerful comparison viewer in any of our products.

Left document: cirrus_defi7_comp.pdf Right document: cirrus stratoclim revised.pdf

Kight document. chitus_stratochin_fevised.

What is this document?

This is a comparison of two documents. The two documents are interleaved such that the left document is displayed on even pages and the right document is displayed on odd pages.

Is there a specific way I should view this file?

This document is intended to be viewed in Two Page Continuous mode (or sometimes called 'Two Page Scrolling'). It should open in this mode by default when using Adobe Acrobat and most popular PDF readers.

If the document opens in a different view, you can often change this in the settings. In Adobe Acrobat, go to View > Page Display > Two Page Scrolling.

Why are there blank pages?

Blank pages are inserted to keep both documents as aligned as much as possible.

How do I read the changes?

Text deleted from the left document and, hence, not in right document is highlighted red. Text added to the right document and, hence, not in left document is highlighted green.

Tip for printing

When printing this document, we recommend printing double-sided and include this first page. This will result in the matching text being displayed on different pages and easily readable, much like a book.

For more information

Draftable offers powerful document comparison solutions for all use-cases. To view our products, please visit our website: <u>draftable.com</u>.

A Comparative Analysis of In-Situ Measurements of High Altitude Cirrus in the Tropics

Francesco Cairo¹, Martina Krämer^{4,3}, Armin Afchine⁴, Luca Di Liberto¹, Sergey Khaykin⁷, Lorenza Lucaferri⁵, Valentin Mitev⁶, Max Port³, Christian Rolf⁴, Marcel Snels¹, Nicole Spelten⁴, Ralf Weigel³, and Stephan Borrmann^{2,3}

¹Istituto di Scienze dell'Atmosfera e del Clima, Consiglio Nazionale delle Ricerche, Rome, Italy
²Particle Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany
³Institute for Atmospheric Physics, Johannes Gutenberg University, Mainz, Germany
⁴Institute of Energy and Climate Research – IEK7, Forschungszentrum Jülich, Jülich, Germany
⁵Dipartimento di Fisica, Università degli Studi di Roma "Tor Vergata", Rome, Italy
⁶Centre Suisse d'Electronique et de Microtechnique, Neuchâtel, Switzerland
⁷Laboratoire Atmosphères, Observations Spatiales (LATMOS), UVSQ, Sorbonne Université, CNRS, IPSL, Guyancourt, France

Correspondence: Francesco Cairo (f.cairo@isac.cnr.it)

Abstract. We analyze cirrus cloud measurements from two dual-instrument cloud spectrometers, two hygrometers and a backscattersonde in view to connect cirrus optical parameters usually accessible by remote sensing with microphysical size resolved and bulk properties accessible in situ. Specifically, we compare the particle backscattering coefficient and depolarization ratio to the particle size distribution, effective and mean radius, surface area density, particle aspherical fraction and ice water content. Data have been acquired by instruments on board the M55 Geophysica research aircraft during July and August 2017 during the Asian Monsoon campaign based in Kathmandu, Nepal, in the framework of the StratoClim (Stratospheric and upper tropospheric processes for better climate predictions) project. Cirrus have been observed over the Hymalaian region between 10 km and the tropopause, situated at 17-18 km. The observed particle number densities varied between 10 and 10^{-4} cm⁻³ in the dimensional range from 1.5 to 468.5 μ m in radius. Correspondingly, backscatter ratios from one tenth up to 50 have been

10 observed.

5

Optical scattering theory has been used to compare the backscattering coefficient computed from measured particle size distribution with those directly observed by the backscattersonde. The aspect ratio of the particles, modeled as spheroids for the T-matrix approach, was left as a free parameter to match the calculations to the optical measures. The computed backscattering coefficient can be set in good agreement with the observed one, but the match between simulated and determined depolarization

- 15 ratios is insufficient, however. Relationships between ice particle concentration, mean and effective radius, surface area density and ice water content with the measured backscattering coefficient are investigated for an estimate of the bulk microphysical parameters of cirrus clouds from remote sensing lidar data. The comparison between particle depolarization and aspherical fraction as measured by one of the cloud spectrometers equipped with a detector for polarization, represents a novelty since it was the first time the two instruments are operated simultaneously on aircraft. The analysis shows the difficulty of establishing
- an univocal link between depolarization values and the presence and amount of aspherical scatterers. This suggests the need

A Comparative Analysis of In-Situ Measurements of High Altitude Cirrus in the Tropics

Francesco Cairo¹, Martina Krämer^{2,3}, Armin Afchine², Guido Di Donfrancesco⁴, Luca Di Liberto¹, Sergey Khaykin⁶, Lorenza Lucaferri⁷, Valentin Mitev^{8,*}, Max Port^{3,**}, Christian Rolf², Marcel Snels¹, Nicole Spelten², Ralf Weigel³, and Stephan Borrmann^{3,5} ¹Istituto di Scienze dell'Atmosfera e del Clima, Consiglio Nazionale delle Ricerche, Rome, Italy ²Institute of Energy and Climate Research – IEK7, Forschungszentrum Jülich, Jülich, Germany ³Institute for Atmospheric Physics, Johannes Gutenberg University, Mainz, Germany ⁴UnitaTecnica Antartide, ENEA, Centro Ricerche Casaccia, Rome, Italy ⁵Particle Chemistry Department, Max Planck Institute for Chemistry, Mainz, Germany ⁶Laboratoire Atmosphères, Observations Spatiales (LATMOS), UVSQ, Sorbonne Université, CNRS, IPSL, Guyancourt, France ⁷Dipartimento di Fisica, Università degli Studi di Roma "Tor Vergata", Rome, Italy ⁸Centre Suisse d'Electronique et de Microtechnique, Neuchâtel, Switzerland ^{*}now retired ^{**}now at: Montessori Zentrum Hofheim, Hofheim, Germany

Correspondence: Francesco Cairo (f.cairo@isac.cnr.it)

Abstract. We analyze cirrus cloud measurements from two dual-instrument cloud spectrometers, two hygrometers and a backscattersonde with the goal of connecting cirrus optical parameters usually accessible by remote sensing with microphysical size resolved and bulk properties accessible in situ. Specifically, we compare the particle backscattering coefficient and depolarization ratio to the particle size distribution, effective and mean radius, surface area density, particle aspherical fraction

- and ice water content. Data have been acquired by instruments on board the M55 Geophysica research aircraft in July and August 2017 during the Asian Monsoon campaign based in Kathmandu, Nepal, in the framework of the StratoClim (Stratospheric and upper tropospheric processes for better climate predictions) project. Cirrus have been observed over the Himalayan region between 10 km and the tropopause, situated at 17-18 km. The observed particle number densities varied between 10 and 10^{-4} cm^{-3} in the dimensional range from 1.5 to 468.5 μ m in radius. Correspondingly, backscatter ratios from one tenth up to 50
- 10 have been observed.

Optical scattering theory has been used to compare the backscattering coefficient computed from measured particle size distribution with those directly observed by the backscattersonde. The aspect ratio of the particles, modeled as spheroids for the T-matrix approach, was left as a free parameter to match the calculations to the optical measures. The computed backscattering coefficient can be brought into good agreement with the observed one, but the match between simulated and

15 measured depolarization ratios is insufficient. Relationships between ice particle concentration, mean and effective radius, surface area density and ice water content with the measured backscattering coefficient are investigated for an estimate of the bulk microphysical parameters of cirrus clouds from remote sensing lidar data. The comparison between particle depolarization and aspherical fraction as measured by one of the cloud spectrometers equipped with a detector for polarization, represents a

of further investigation that could take into consideration not only the fraction of aspheric particles but also their predominant morphology.

Copyright statement. TEXT

1 Introduction

- 25 Cirrus are clouds composed of micron to millimeter sized ice crystals of various shapes that form in the upper troposphere between 4 and 20 km above sea level (Lynch et al., 2002). Cirrus higher elevations are usually found in the tropics, where their highest occurrence frequency is also recorded, and lower elevations are found in polar regions (Sassen et al., 2008, 2009). Tropical cirrus originate either from outflows from deep convective clouds (liquid-origin cirrus) or from vertical uplifting of air (in situ-origin cirrus) associated with Kelvin or gravity waves as well as with the synoptic-scale tropospheric tropical ascent
- 30 (Jensen et al., 1996; Pfister et al., 2001; Immler et al., 2008; Fujiwara et al., 2009; He et al., 2012; Krämer et al., 2016; Luebke et al., 2016; Wernli et al., 2016). Studies of these clouds are important for a better understanding of their impact on the climate as they play a crucial role in earth's radiation budget (Prabhakara et al., 1993; Campbell et al., 2016; Lolli, 2017; Krämer et al., 2020). Their impact is based on two effects (Stephens, 2002, 2005): i. a greenhouse potential that traps the outgoing long-wave radiation emitted by the earth and the atmosphere underneath; ii. their albedo that reflects the incoming solar radiation. The
- balance between the cirrus induced warming and cooling depends on their coverage, height, thickness, horizontal and vertical temperature distribution as well as ice crystal size and shape distributions within the clouds (Lynch, 1996; Boucher et al., 2013). Moreover, cirrus are an essential modulator of the water budget in the upper troposphere and in the stratosphere (Luo et al., 2003; MacKenzie et al., 2006; Corti et al., 2008; Fueglistaler et al., 2009).

The qualitative and quantitative assessment of the cirrus properties over large spatial and temporal scales require the use of satellite data. Methods exist to provide spaceborne retrievals of cirrus bulk and microphysical parameters. Passive visible or thermal, spectrally resolved measurements have been used to infer the cloud optical depth, ice particle effective radius (R_{eff}), Ice Water Path (Meyer and Platnick, 2010; Sourdeval et al., 2013; Guignard et al., 2012; Sourdeval et al., 2015) but may subsample clouds of small optical thickness. Active remote sensing by radars and lidars can be more sensitive to thin clouds and can still provide information on cirrus geometrical and optical properties with high spatial and temporal resolutions. Vertical

- profiles of cirrus extinction, Ice Water Content (IWC) and R_{eff} are retrieved using lidar and/or radar measurements (Austin et al., 2009; Delanoë and Hogan, 2010). Global mapping of cirrus properties is obtained from satellite borne instruments like the CALIOP (Cloud and Aerosol Lidar with Orthogonal Polarization) lidar aboard the CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation) polar orbiting satellite (Nazaryan et al., 2008). Such measurements have begun to include estimates of ice crystal number concentrations (N_{ice}). In fact, climate model parameterizations of ice cloud optical
- 50

properties are based on R_{eff} and IWC (Fu, 2007) but these two do not fully constrain the ice cloud Particle Size Distribution (PSDs) and its optical properties (Mitchell et al., 2011). Moreover, the knowledge of N_{ice} on a global scale would ameliorate

novelty since it was the first time the two instruments are operated simultaneously on aircraft. The analysis shows the difficulty

20 of establishing an univocal link between depolarization values and the presence and amount of aspherical scatterers. This suggests the need of further investigation that could take into consideration not only the fraction of aspheric particles but also their predominant morphology.

Copyright statement. TEXT

1 Introduction

- 25 Cirrus are high clouds existing between -35° and -85°C, composed of ice crystals, of micron to millimeter size (Lynch et al., 2002), that are fairly widely dispersed, usually resulting in relative transparency and whiteness and often producing halo phenomena not observed with other cloud forms (AMS, 2023). Cirrus higher elevations are usually found in the tropics, where their highest occurrence frequency is also recorded, and lower elevations are found in polar regions (Sassen et al., 2008, 2009). Tropical cirrus originate either from outflows from deep convective clouds (liquid-origin cirrus) or from vertical uplifting of
- air (in situ-origin cirrus) associated with Kelvin or gravity waves as well as with the synoptic-scale tropospheric tropical ascent (Jensen et al., 1996; Pfister et al., 2001; Immler et al., 2008; Fujiwara et al., 2009; He et al., 2012; Krämer et al., 2016; Luebke et al., 2016; Wernli et al., 2016). Studies of these clouds are important for a better understanding of their impact on the climate as they play a crucial role in earth's radiation budget (Prabhakara et al., 1993; Campbell et al., 2016; Lolli, 2017; Krämer et al., 2020). Their impact is based on two effects (Stephens, 2002, 2005): i. a greenhouse potential that traps the outgoing long-wave
- 35 radiation emitted by the earth and the atmosphere underneath; ii. their albedo that reflects the incoming solar radiation. The balance between the cirrus induced warming and cooling depends on their coverage, height, thickness, horizontal and vertical temperature distribution as well as ice crystal size and shape distributions within the clouds (Lynch, 1996; Boucher et al., 2013). Moreover, cirrus are an essential modulator of the water budget in the upper troposphere and in the stratosphere (Luo et al., 2003; MacKenzie et al., 2006; Corti et al., 2008; Fueglistaler et al., 2009).
- The qualitative and quantitative assessment of the cirrus properties over large spatial and temporal scales require the use of satellite data. Methods exist to provide spaceborne retrievals of cirrus bulk and microphysical parameters. Passive visible or thermal, spectrally resolved measurements have been used to infer the cloud optical depth, ice particle effective radius (R_{eff}), Ice Water Path (Meyer and Platnick, 2010; Sourdeval et al., 2013; Guignard et al., 2012; Sourdeval et al., 2015) but may subsample clouds of small optical thickness. Active remote sensing by radars and lidars can be more sensitive to thin clouds and
- can still provide information on cirrus geometrical and optical properties with high spatial and temporal resolutions. Vertical profiles of cirrus extinction, Ice Water Content (IWC) and R_{eff} are retrieved using lidar and/or radar measurements (Austin et al., 2009; Delanoë and Hogan, 2010). Global mapping of cirrus properties is obtained from satellite borne instruments like the CALIOP (Cloud and Aerosol Lidar with Orthogonal Polarization) lidar aboard the CALIPSO (Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation) polar orbiting satellite (Nazaryan et al., 2008). Such measurements have begun to

the understanding of ice nucleation and its parametrization in climate models. A refinement of the ice crystal nucleation rates would in turn improve the predictions of R_{eff} . Mitchell et al. (2018) use co-located observations from the Infrared Imaging Radiometer (IIR) and from CALIOP to retrieve N_{ice} , R_{eff} and IWC in semi-transparent cirrus clouds, while Sourdeval et al. (2018) employ combined lidar-radar measurement to provide satellite estimates of N_{ice} using a methodology that constrains moments of a parameterized Particle Size Distribution (PSD) through lidar extinction and radar reflectivity.

Cirrus clouds' microphysical properties have been characterized by in-situ measurements taken during several airborne observation campaigns (Thomas et al., 2002; Schiller et al., 2008; Krämer et al., 2009; Frey et al., 2011; Luebke et al., 2013; Frey et al., 2014; Krämer et al., 2016; Schumann et al., 2017). A review of these studies and of the challenges they present is
reported in Baumgardner et al. (2017) while Krämer et al. (2020) describe extensive statistics of meteorological parameters, IWC, N_{ice}, ice crystal mean radii (R_{mean}), relative humidities with respect to ice and water vapor mixing ratios from airborne in situ measurements performed during 150 flights in 24 campaigns, in midlatitudes and the tropics. Such observational activity, in addition to being essential for shedding light on the processes of formation, aging and dissipation of the clouds, is of great

55

and bulk parameters by comparing them with corresponding results of in situ measurements. In the present work we intend to compare optical measurements of cirrus clouds, i.e. particle backscatter coefficient β and total particle depolarization δ_{TA} with bulk and microphysical parameters observed by cloud spectrometers and hygrometers. The optical measurements are generally accessible to lidar probing, but in our case have been taken in situ by a backscattersonde,

help in the interpretation of satellite sensor data, allowing the calibration and validation of the retrievals of cirrus microphysical

henceforth they are directly comparable with the other data acquired in situ as all measurements originate from the same air

- 70 parcel. Measurements have been taken during the aircraft field campaign of the EU funded project StratoClim (Stratospheric and upper tropospheric processes for better climate predictions) carried out in South Asia in 2017. A full description of the campaign is provided by Stroh et al. (2022, same issue). The South Asian campaign of the high altitude research aircraft M55 Geophysica (Stefanutti et al., 1999) focused on detailed observations of atmospheric transport and physical-chemical processes which dominate the input of air and aerosols into the (sub-)tropical stratosphere. In the present work we make use of data from
- 75 seven flights, performed on 29 and 31 July and on 2, 4, 6, 8 and 10 August 2017. During these flights the airplane penetrated into cirrus clouds several times for nearly seven hours of observations in clouds over approximatively 35 flight hours. We verify the ability and limits of optical modeling in reproducing the results of remote sensing optical measurements from

the concomitant in-situ microphysical ones, thus assessing the compatibility of the two datasets.

- Closure studies between aerosol light scattering coefficients and aerosol PSD with optical modeling are common in atmospheric science, and they often use measurements from in situ nephelometers, and Mie theory applied to cloud spectrometers data (Wex et al., 2002; Teri et al., 2022). Similar approaches have been also attempted by comparing lidar or backscatter probe measurements with cloud spectrometers mounted on balloons or on airplanes, in the characterization of polar stratospheric clouds (Schreiner et al., 2002; Deshler et al., 2000; Scarchilli et al., 2005; Snels et al., 2021; Cairo et al., 2022). Comparatively fewer studies are present in comparing the size distributions of ice crystals in cirrus clouds, and their modeled and measured
 optical scattering properties, this is particularly true for backscattering measurements (Wagner and Delene, 2022). The reason probably lies in the greater difficulty to obtain significant and definitive confirmations from such optical closures. This is likely
 - 3

- include estimates of ice crystal number concentrations (N_{ice}). In fact, climate model parameterizations of ice cloud optical 50 properties are based on R_{eff} and IWC (Fu, 2007) but these two do not fully constrain the ice cloud Particle Size Distribution (PSDs) and its optical properties (Mitchell et al., 2011). Moreover, the knowledge of N_{ice} on a global scale would ameliorate the understanding of ice nucleation and its parametrization in climate models. A refinement of the ice crystal nucleation rates would in turn improve the predictions of R_{eff} . Mitchell et al. (2018) use co-located observations from the Infrared Imaging
- Radiometer (IIR) and from CALIOP to retrieve Nice, Reff and IWC in semi-transparent cirrus clouds, while Sourdeval et al. 55 (2018) employ combined lidar-radar measurement to provide satellite estimates of N_{ice} using a methodology that constrains moments of a parameterized Particle Size Distribution (PSD) through lidar extinction and radar reflectivity.

Cirrus clouds' microphysical properties have been characterized by in-situ measurements taken during several airborne observation campaigns (Thomas et al., 2002; Schiller et al., 2008; Krämer et al., 2009; Frey et al., 2011; Luebke et al., 2013;

- 60 Frey et al., 2014; Krämer et al., 2016; Schumann et al., 2017). A review of these studies and of the challenges they present is reported in Baumgardner et al. (2017) while Krämer et al. (2020) describe extensive statistics of meteorological parameters, IWC, N_{ice} , ice crystal mean radii (R_{mean}), relative humidities with respect to ice and water vapor mixing ratios from airborne in situ measurements performed during 150 flights in 24 campaigns, in midlatitudes and the tropics. Such observational activity, in addition to being essential for shedding light on the processes of formation, aging and dissipation of the clouds, is of great
- help in the interpretation of satellite sensor data, allowing the calibration and validation of the retrievals of cirrus microphysical 65 and bulk parameters by comparing them with corresponding results of in situ measurements.

In the present work we intend to compare optical measurements of cirrus clouds, i.e. particle backscatter coefficient β and total particle depolarization δ_{TA} with bulk and microphysical parameters observed by cloud spectrometers and hygrometers. The optical measurements are generally accessible to lidar probing, but in our case have been taken in situ by a backscattersonde,

- 70 henceforth they are directly comparable with the other data acquired in situ as all measurements originate from the same air parcel. Measurements have been taken during the aircraft field campaign of the EU funded project StratoClim (Stratospheric and upper tropospheric processes for better climate predictions) carried out in South Asia in 2017. A full description of the campaign is provided by Stroh et al. (2022, same issue). The South Asian campaign of the high altitude research aircraft M55 Geophysica (Stefanutti et al., 1999) focused on detailed observations of atmospheric transport and physical-chemical processes which dominate the input of air and aerosols into the (sub-)tropical stratosphere. 75

80

In the present work we make use of data from seven flights, performed on 29 and 31 July and on 2, 4, 6, 8 and 10 August 2017. During these flights the airplane penetrated into cirrus clouds several times for nearly six hours of observations in clouds over approximatively 35 flight hours. Most of the measurements during StratoClim were performed at temperatures < 205 K, corresponding to potential temperatures >355 K and altitudes >14 km, i.e., in the TTL. Krämer et al. (2020) reports a description of the clouds observed during the campaign. The first part of the campaign period suffered from very rare cloud passes at elevated altitudes, with comparatively low cloud particle number concentrations (below 1 cm^{-3}). In fact, during this period, the vast majority of clouds were encountered during ascent from or during approach to the Kathmandu airport. On 29 and 31 July, and on 02 August, most of the clouds were come across at pressure level of ~ 400 hPa (and higher) during ascent and descent, with cloud particle concentrations ranging from 100 to 1000 cm^{-3} . The second campaign period (flights on 04,

because of the uncertainties affecting both from the particle size distribution measurements and their optical modeling, arising from the large dimensional range of the ice crystals extending for more than three orders of magnitude, and the great variety of shapes that they can take simultaneously within the cloud (Bailey and Hallett, 2009). A further complication arises from the

- 90 presence of rough irregular surfaces, corners and edges in ice crystals (Schnaiter et al., 2011). Actually, modeling of the optical properties of cirrus clouds is a formidable problem with probably not always a valid solution, despite the numerous methods available for calculating scattering from aspherical particles (Mishchenko et al., 1999; Konoshonkin et al., 2017a). In fact many light-scattering computation methods have been employed to calculate the scattering properties of cirrus particles and we may quote the finite-difference time-domain (FDTD) method (Sun et al., 1999), the discrete dipole approximation (DDA) (Yurkin
- et al., 2007), the boundary element method (Groth et al., 2015), the pseudo-spectral time-domain method (Liu et al., 2012), the surface-integral equation method (Nakajima et al., 2009), and geometrical optics in various implementations: improved geometric optics method (IGOM) (Yang and Liou, 1996a), geometric optics integral equation (GOIE) (Ishimoto et al., 2013) and ray-tracing geometric optics method (GOM) (Macke et al., 1996). The T-matrix theory offers solution to the computation of electromagnetic scattering from axisymmetric particles and has practical advantages over other methods, largely due to its
- 100 analytical character and the exploitation of particle symmetries, which considerably simplifies the calculation, and has long been used to study the scattering properties of cirrus clouds. Mishchenko et al. (1997) used it to compute the backscattering from horizontally oriented ice platelets in cirrus, Baran et al. (2001) modeled absorption and extinction properties of the finite hexagonal ice columns and plates in random and preferred orientation, Liu et al. (2006) reported results on the scattering properties of small cirrus crystals modeled as mixtures of polydisperse, randomly oriented spheroids and cylinders with varying
- 105 aspect ratios, Bi and Yang (2014) employed it to compute the optical properties of randomly oriented ice crystals of various shapes including hexagonal columns, hollow columns, droxtals, bullet rosettes and aggregates. T- matrix theory has also been used to simulate the response of cloud spectrometers with forward scattering geometries, when exposed to clouds of aspherical particles (Borrmann and Luo, 2000).

Finally, the possibility of using microphysical observations taken along with optical observations usually accessible in remote

- sensing but here obtained with quasi in situ techniques (the air mass sampled by the backscattersonde is for all practical purposes the same sampled by hygrometers and cloud spectrometers) allows us establish relationships between β and δ_{TA} and cloud parameters retrieved from the measured PSDs, as N_{ice} , R_{mean} , R_{eff} , Surface Area Density (SAD), particle Aspherical Fraction (AF), and cloud IWC, this latter retrieved from a comparison of data from hygrometers that differentially measure the proportion of gaseous and total water. These empirical relationships are useful in interpreting lidar measurements of cirrus clouds in terms of their microphysical and bulk parameters.
- 115

Instruments and Data 2

We have compared and used data from the two dual instrument cloud spectrometers that flew aboard the aircraft Geophysica, the NIXE-CAPS and the CCP. The CCP and NIXE-CAPS instruments (see Section 2.2) are both developments by Droplet Measurement Technologies (DMT) and are nearly identical in their optical measurement technology, the sensors used for

- 85 06, 08, and 10 August 2017) provided extended fields of cirrus clouds of convective origin, with elevated particle densities and broad size distributions covering almost the entire detection size range of the different particle probes. The cloud particle measurements, mostly carried out over the southern slopes of the Himalayas captured high ice water content up to 2400 ppmv and ice particle aggregates exceeding 700 μ m in size. The observed ice particles were mainly of liquid origin, with only a small amount formed in situ. ERA5 reanalysis corroborates the presence of high IWC detrained from deep-convective. A
- 90 microphysical modeling study by Lamraoui et al. (2023) focuses on the flight of the 10 August, but its results can probably also be applied to the other cases of convectively generated cirrus measured during the second part of the campaign. The study predicts ice habits and reproduces the observed IWC, ice number concentration, and bimodal ice particle size distribution. The lower range of particle sizes is mostly represented by planar and columnar habits, while the upper range is dominated by aggregates with sizes between 600 and 800 μ m. The study suggests that most of measured ice particles are of liquid origin 95 with only a small amount formed in situ. These latter are associated with low values of IWC and number concentration, which
- makes them less influential in regulating the IWC which is, on the contrary, substantially influenced by planar ice particles of liquid origin. The difference in ice number concentration across habits can be up to 4 orders of magnitude, with aggregates occurring in much smaller numbers.

We verify the ability and limits of optical modeling in reproducing the results of remote sensing optical measurements

- from the concomitant in-situ microphysical ones, thus assessing the compatibility of the two datasets. Closure studies between aerosol light scattering coefficients and aerosol PSD with optical modeling are common in atmospheric science, and they often use measurements from in situ nephelometers, and Mie theory applied to particle spectrometers data (Wex et al., 2002; Teri et al., 2022). Similar approaches have been also attempted by comparing lidar or backscatter probe measurements with cloud spectrometers mounted on balloons or on airplanes, in the characterization of polar stratospheric clouds (Schreiner et al.,
- 2002; Deshler et al., 2000; Scarchilli et al., 2005; Snels et al., 2021; Cairo et al., 2023). Comparatively fewer studies are present in comparing the size distributions of ice crystals in cirrus clouds, and their modeled and measured optical scattering properties, this is particularly true for backscattering measurements (Wagner and Delene, 2022). The reason probably lies in the greater difficulty to obtain significant and definitive confirmations from such optical closures. This is likely because of the uncertainties affecting both from the particle size distribution measurements and their optical modeling, arising from
- the large dimensional range of the ice crystals extending for more than three orders of magnitude, and the great variety of shapes that they can take simultaneously within the cloud (Bailey and Hallett, 2009). A further complication arises from the presence of rough irregular surfaces, corners and edges in ice crystals (Schnaiter et al., 2011). Actually, modeling of the optical properties of cirrus clouds is a formidable problem with probably not always a valid solution, despite the numerous methods available for calculating scattering from aspherical particles (Mishchenko et al., 1999; Konoshonkin et al., 2017a). In fact many
- 115 light-scattering computation methods have been employed to calculate the scattering properties of cirrus particles and we may quote the finite-difference time-domain (FDTD) method (Sun et al., 1999), the discrete dipole approximation (DDA) (Yurkin et al., 2007), the boundary element method (Groth et al., 2015), the pseudo-spectral time-domain method (Liu et al., 2012), the surface-integral equation method (Nakajima et al., 2009), and geometrical optics in various implementations: improved geometric optics method (IGOM) (Yang and Liou, 1996a), geometric optics integral equation (GOIE) (Ishimoto et al., 2013)

120 pressure and temperature, and the flow measurement technology (Prandl's pitot tube system). The IWC has been derived from the total water hygrometer FISH and the water vapor hygrometer FLASH (see Section 2.3); the cloud optical parameters have been derived from the backscattersonde MAS (see Section 2.1) which measures in situ particle backscattering coefficient and depolarization ratio at 532 nm. Data from MAS are also compared on some cases of interest with the 532 nm elastic lidar MAL on board the same aircraft.

125 2.1 Particle backscatter measurements: MAS and MAL instruments

2.1.1 The MAS backscattersonde

Optical observations have been provided by the backscattersonde MAS, located in a bay beneath the pilot's cockpit, facing sideways on the right. It emits polarized laser light at 532 nmm and collects the light backscattered from the portion of atmosphere in close proximity (3-10 m) to the instrument, sensitive to optically detectable (i.e. whose diameter is greater than few tenths of μm) cloud particles and aerosols. Polarization resolved observations allow to detect the particle's asphericity, hence thermodynamic phase. The instrument is basically a polarization diversity Rayleigh lidar system that measures insitu the same atmospheric parameters which are accessible to lidars. The sampling volume is approximately $10^{-3} m^{-3}$, the resolution is 10 *s*, corresponding to a 1.5-1.9 *km* horizontal resolution along the aircraft trajectory, given the 154±39 $m s^{-1}$ aircraft speed at altitude (Weigel et al., 2021a, b).

The backscattered light is split according to parallel and perpendicular polarization with respect to the linearly polarized laser light, allowing the measurement of the Backscatter Ratio BR, the Volume Depolarization δ and the Total Aerosol Depolarization δ_{TA}. These optical parameters follows the usual definitions (Cairo et al., 1999) reported hereby for convenience; in the following the subscripts *mol* and *A* denote respectively the molecular and aerosols contribution to the optical coefficients, and *cross* and *par* denote the perpendicular and parallel polarization of the backscattering coefficient β (Collis and Russell, 1976).

$$BR = \frac{\beta_A^{cross} + \beta_{mol}^{par} + \beta_{mol}^{par}}{\gamma_{mol}^{par} + \beta_{mol}^{par}}$$
(1)

$$\delta_{T} = \frac{\beta_{mol}^{cross} + \beta_{Mol}^{cross}}{\beta_{mol}^{par} + \beta_{A}^{par} + \beta_{Mol}^{cross} + \beta_{A}^{cross}}$$

$$\delta_{A} = \frac{\beta_{A}^{cross}}{\beta_{A}^{par}}$$

$$(2)$$

$$(3)$$

$$\delta_{TA} = \frac{\beta_A^{cross}}{\beta_A^{cross} + \beta_A^{par}} \tag{4}$$

145

130

Note the different definitions of Particle Depolarization δ_A and Total Particle Depolarization δ_{TA} which will be used both in the following. Formulae to pass from one to the other definition can be found in Cairo et al. (1999).

The signal detected by the backscattersonde channels is directly proportional to $\beta_A^{par(cross)} + \beta_{mol}^{par(cross)}$. The backscatter ratio BR is derived by a calibration procedure that uses concomitant measurements of pressure and temperature to retrieve the air density, and defines a suitable constant K – taking into account the molecular scattering cross section as well as the

- and ray-tracing geometric optics method (GOM) (Macke et al., 1996). The T-matrix theory offers solution to the computation of electromagnetic scattering from axisymmetric particles and has practical advantages over other methods, largely due to its analytical character and the exploitation of particle symmetries, which considerably simplifies the calculation, and has long been used to study the scattering properties of cirrus clouds. Mishchenko et al. (1997) used it to compute the backscattering from horizontally oriented ice platelets in cirrus, Baran et al. (2001) modeled absorption and extinction properties of the
- finite hexagonal ice columns and plates in random and preferred orientation, Liu et al. (2006) reported results on the scattering properties of small cirrus crystals modeled as mixtures of polydisperse, randomly oriented spheroids and cylinders with varying aspect ratios, Bi and Yang (2014) employed it to compute the optical properties of randomly oriented ice crystals of various shapes including hexagonal columns, hollow columns, droxtals, bullet rosettes and aggregates. T- matrix theory has also been used to simulate the response of cloud spectrometers with forward scattering geometries, when exposed to clouds of aspherical properties (Bormann and Lue, 2000).
- 130 particles (Borrmann and Luo, 2000).

Finally, the possibility of using microphysical observations taken along with optical observations usually accessible in remote sensing but here obtained with quasi in situ techniques (the air mass sampled by the backscattersonde is for all practical purposes the same sampled by hygrometers and cloud spectrometers) allows us establish relationships between β and δ_{TA} and cloud parameters retrieved from the measured PSDs, as N_{ice} , R_{mean} , R_{eff} , Surface Area Density (SAD), particle Aspherical

135 Fraction (AF), and cloud IWC, this latter retrieved from a comparison of data from hygrometers that differentially measure the proportion of gaseous and total water. These empirical relationships are useful in interpreting lidar measurements of cirrus clouds in terms of their microphysical and bulk parameters.

2 Instruments and Data

We have compared and used data from the two dual instrument cloud spectrometers that flew aboard the aircraft Geophysica, the
NIXE-CAPS and the CCP. Both instruments (see Section 2.2) are both developments by Droplet Measurement Technologies (DMT) and are nearly identical in their technology, the sensors used for pressure and temperature, and the flow measurement technology (Prandl's pitot tube system). The IWC has been derived from the total water hygrometer FISH and the water vapor hygrometer FLASH (see Section 2.3); the cloud backscattering coefficient and depolarization ratio at 532 nm have been derived from the in situ backscattersonde MAS (see Section 2.1). Data from MAS are also compared on some cases of interest with the 532 nm elastic lidar MAL.

2.1 Particle backscatter measurements: MAS and MAL instruments

2.1.1 The MAS backscattersonde

Optical observations in-situ have been provided by the backscattersonde MAS which is basically a polarization diversity Rayleigh lidar that measures in-situ the same parameters accessible to lidars. It is located in a bay beneath the pilot's cockpit,

facing sideways on the right. It emits polarized laser light at 532 nmm and collects the light backscattered from the portion of

instrumental sensitivity - in order to achieve BR=1 from the signal detected in air masses where no particles are present. A 150 description of the instrument and data processing can be found in Cairo et al. (2011).

Figure 1 reports the 2D histogram of frequency distribution of BR with respect to geometrical altitude; the particle observations clusters in two regions, namely between 7.5 and 10 km, and above 13 km up to the tropopause region, the Cold Point Tropopause being at 17 km on average. In the following we will focus on observations taken in that upper region. We identify a cloud presence when BR>1.2.

155

Figure 2 exhibits δ_{TA} with respect to temperature, for clouds observed above 11 km. Total Particle Depolarization ranges from 20 to 50%. In the upper part of the troposphere, i.e. when temperatures fall below 200K, a negative trend with respect to temperature can be discerned; in that altitude range, an increase of δ_{TA} with altitude as has been often reported for cirrus (Sassen and Benson, 2001; Sunilkumar and Parameswaran, 2005; Cairo et al., 2021). However, at higher temperatures, namely at around 205K and 215K, observations of highly depolarizing clouds are also here reported.

160

165

2.1.2 The MAL lidars

The lidars MAL1 and MAL2 (Miniature Aerosol Lidars Mk1 and Mk2) are elastic backscatter-depolarization instruments. operating at 532 nm (Martucci et al., 2005). They operate with micropulse lasers, having pulse repetition rates of 4.5-5.5 kHz and are packed each in a single pressurized box. MAL1 is installed on M55 Geophysica for upward probing, while MAL2 is installed for downward probing. The measured values are the aerosol backscatter ratio and volume depolarization ratio. From the measured signals it is possible to obtain the aerosol backscattering coefficient and depolarisation ratio. The definition of the measured and obtained values follows eqs. (1-4). The high-repetition rate combined with low pulse energy of the laser gives

- the possibility to use a photon-counting detection/acquisition system with high dynamic range. This makes possible to start the lidar profile as close to the aircraft, as allowed by the geometrical overlap function, i.e. at 400 m from the platform delivering
- backscatter and depolarization profiles every minute, with a vertical resolution of 50m. The processing procedure is based on 170 comparison of the signal from the clouds with the molecular backscatter signal from aerosol-free parts of the lidar profile. In this it is similar to the one for MAS. The difference with MAS is that in MAL the aerosol free atmospheric volumes are identified in the lidar profile, respectively at some range above and below the aircraft, and not at the flight altitude. The lidars participated in a number of campaigns with M55 Geophysica (Cairo et al., 2004; Molleker et al., 2014; Mahnke et al., 2021).
- 175 This includes a comparison case with CALIPSO lidar CALIOP during RECONCILE campaign (Mitev et al., 2012).

2.2 **Cloud spectrometers**

The NIXE-CAPS instrument 2.2.1

The NIXE-CAPS (for details see Luebke et al. (2016); Costa et al. (2017)) is located below the aircraft wing and incorporates two instruments: the CAS-DPOL (Cloud and Aerosol Spectrometer with detector for polarization) and the NIXE-CIPgs (Cloud 180 Imaging Probe grey scale); it also includes an air speed sensor and a temperature probe. CAS-DPOL is a light-scattering probe

atmosphere in close proximity (3-10 m) to the instrument, sensitive to air molecules, cloud particles and aerosols. Polarization resolved observations allow to detect the particle's asphericity, hence thermodynamic phase. The sampling volume is approximately $10^{-3} m^{-3}$, the resolution is 10 s, corresponding to a 1.5-1.9 km horizontal resolution along the aircraft trajectory, given the $154\pm39 m s^{-1}$ aircraft speed at altitude (Weigel et al., 2021a, b).

The backscattered light is split into two channels according to polarization, allowing the measurement of the Backscatter Ratio BR and the Volume Depolarization δ , the Particle Depolarization δ_A and the Total Particle Depolarization δ_{TA} . These optical parameters follows the usual definitions (Cairo et al., 1999) reported hereby for convenience; the subscripts *mol* and *A* denote respectively the molecular and particle contribution to the measured parameters, and *cross* and *par* denote the perpendicular and parallel polarization of the backscattering coefficient β (Collis and Russell, 1976).

160	$BR = \frac{\beta_A^{cross} + \beta_{mol}^{cross} + \beta_A^{par} + \beta_{mol}^{par}}{\beta_{mol}^{cross} + \beta_{mol}^{par}}$	(1)
	$\delta = \frac{\beta_{mol}^{cross} + \beta_A^{cross}}{\beta_{mol}^{par} + \beta_A^{par}}$	(2)
	$\delta_T = \frac{\beta_{mol}^{cross} + \beta_A^{cross}}{\beta_{mol}^{par} + \beta_A^{par} + \beta_A^{cross} + \beta_A^{cross}}$	(3)
	$\delta_A = \frac{\beta_A^{cross}}{\beta_A^{par}}$	(4)
	$\delta_{TA} = \frac{\beta_A^{cross}}{\beta_A^{cross} + \beta_A^{par}}$	(5)

165

170

Note the different definitions of Particle Depolarization δ_A and Total Particle Depolarization δ_{TA} which will be both used in the following. To pass from one to the other definition, please refer to Cairo et al. (1999).

The signal detected in the parallel (cross) channel is directly proportional to $\beta_A^{par(cross)} + \beta_{mol}^{par(cross)}$. The BR is derived by a calibration procedure that uses colocated measurements of pressure and temperature to retrieve the air density, and defines a suitable constant K – taking into account the molecular scattering cross section as well as the instrumental sensitivity - in order to achieve BR=1 in air masses where no particles are present. A description of the instrument and data processing can be found in Cairo et al. (2011).

Figure 1 reports the 2D histogram of frequency distribution of BR with respect to geometrical altitude; the cloud observations clusters in two regions, namely between 7.5 and 10 km, and above 13 km up to the tropopause region, the Cold Point Tropopause being at 17 km on average. In the following we will focus on observations taken in that upper region. We identify

175 cloud presence when BR>1.2.

Figure 2 exhibits δ_{TA} with respect to temperature, for clouds observed above 11 km. In the upper part of the troposphere, i.e. when temperatures fall below 200K, a negative trend with respect to temperature can be discerned; in that altitude range, an increase of δ_{TA} with altitude as has been often reported for cirrus (Sassen and Benson, 2001; Sunilkumar and Parameswaran, 2005; Cairo et al., 2021). However, at higher temperatures, namely at around 205K and 215K, observations of highly depolar-

180 izing clouds are also here reported. To avoid contamination by aerosols or an excessive uncertainty on the depolarization for very low BR, we will restrict our analysis to observations with BR>1.2, for a total of 2132 data points. covering the particle size range of 0.3 to 25 μm in radius. Moreover, the CAS records the change of polarization in the backward-scattered light, thus giving information about the particle asphericity.

NIXE–CIPgs is an optical array probe that covers the particle size range between 7.5 and 468.5 μm . The instrument captures the image of a cloud particle by using a 64-element photodiode array (15 μ m resolution) to generate two-dimensional shadow images that can be analyzed for particle size and asphericity using various algorithms (Costa et al., 2017).

For aircraft speeds between 240 and 80 $m \, s^{-1}$, the instruments' sampling volumes limit the particle concentration measurements to concentrations above 0.02 to 0.05 cm^{-3} (NIXE–CAS–DPOL) and about 0.0001 to 0.001 cm^{-3} (NIXE–CIPgs).

The particle size distributions (PSDs) of CAS-DPOL and CIPgs are merged into a single PSD covering the range of 0.3 to 468.5 μm , where the size bins up to 20 μm are taken from the CAS-DPOL and those larger than 20 μm from the CIPgs. This

190

195

185

threshold is used since the CIPgs has a larger sampling volume than the CAS-DPOL, thus providing better particle sampling statistics. Particles larger than 1.5 μm in radius are classified as cloud, while the smaller particles are considered aerosols.

The PSDs are reported for each second. In the present work the PSDs have been averaged over 10s and synchronized to the backscattersonde data. In the following, in addition to the PSD, we have used the "combined" aspherical fraction AF_i which refers to fraction of aspherical particles detected in size channels both from NIXE-CAS and NIXE-CIPgs. The AFs are derived as described by Costa et al. (2017)

The lidar-based aerosol depolarization, being an optical measure, may be biased toward a particular size range of the PSD, depending on the respective particle number densities and absolute sizes, as many smaller ones can be as "active" as few bigger ones. In order to study its relationship with the microphysical measurements, we have used both the ordinary mean AF defined as:

200
$$AF = \frac{\sum_{i=1}^{max} Af_i n_i}{\sum_{i=1}^{max} n_i}$$
 (5)

and an "effective" mean AF_{eff} where in the mean the AF_i , detected in the single size bin *i* where n_i particles with radii between r_{i-1} and g_i have been counted, are weighted proportionally to the mean cross sectional area S_i :

$$S_i = \pi \left(\frac{r_{i-1} + r_i}{2}\right)^2 \tag{6}$$

relative to that bin.

205
$$AF_{eff} = \frac{\sum_{i=1}^{max} Af_i S_i n_i}{\sum_{i=1} S_i n_i}$$
(7)

In this way the AF_i are weighted not only with the numerical density of the particles in the bin *i*, but also in proportion to a proxy of their scattering cross section. In the subsequent analysis, however, we did not find notable differences when seeking a correlation between the observed depolarization and AF or AF_{eff} . Hence in the following we will report only the results relating AF.

2.1.2 The MAL lidars

The lidars MAL1 and MAL2 (Miniature Aerosol Lidars Mk1 and Mk2) are depolarization diversity Rayleigh lidar instruments operating at 532 nm (Martucci et al., 2005). They use micropulse lasers with pulse repetition rates of 4.5-5.5 kHz. MAL1 is

- installed on M55 Geophysica for upward probing, while MAL2 is for downward probing. The measured parameters are BR and δ . The definition of the measured and processed parameters follows eqs. (1-5). The high-repetition rate combined with low pulse energy of the laser allows the use of a photon-counting detection/acquisition system with high dynamic range. This makes possible to start the lidar profile as close to the aircraft, as allowed by the geometrical overlap function, i.e. at 400 m from the platform delivering backscatter and depolarization profiles every 2 minutes, with a vertical resolution of 50 m.
- The processing procedure is based on comparison of the signal from the clouds with the molecular backscatter signal from aerosol-free parts of the lidar profile. In this it is similar to the one for MAS, except that in MAL the particle free atmospheric volumes are identified in the lidar profile, at some range above and below the aircraft, and not at the flight altitude. The lidars participated in a number of campaigns with M55 Geophysica (Cairo et al., 2004; Molleker et al., 2014; Mahnke et al., 2021). This includes a comparison case with CALIPSO lidar CALIOP during RECONCILE campaign (Mitev et al., 2012).

195 2.2 Cloud spectrometers

210

2.2.1 The NIXE-CAPS instrument

The NIXE-CAPS (for details see Luebke et al. (2016); Costa et al. (2017)) is located below the aircraft wing and incorporates two instruments: the CAS–DPOL (Cloud and Aerosol Spectrometer with detector for polarization) and the NIXE–CIPgs (Cloud Imaging Probe grey scale); it also includes an air speed sensor and a temperature probe. CAS-DPOL is a light-scattering probe
covering the particle size range of 0.3 to 25 μm in radius. Moreover, it records the change of polarization in the backward-scattered light, thus giving information about the particle asphericity.

NIXE–CIPgs is an optical array probe that covers the particle size range between 7.5 and 468.5 μm . The instrument captures the image of a cloud particle by using a 64-element photodiode array (15 μ m resolution) to generate two-dimensional shadow images that can be analyzed for particle size and asphericity using various algorithms (Costa et al., 2017).

For aircraft speeds between 240 and 80 $m s^{-1}$, the instruments' sampling volumes limit the particle concentration measurements to concentrations above 0.02 to 0.05 cm^{-3} (NIXE–CAS–DPOL) and about 0.0001 to 0.001 cm^{-3} (NIXE–CIPgs).

The particle size distributions (PSDs) of CAS-DPOL and CIPgs are merged into a single PSD covering the range of 0.3 to 468.5 μm , where the size bins up to 20 μm are taken from the CAS-DPOL and those larger than 20 μm from the CIPgs. This threshold is used since the CIPgs has a larger sampling volume than the CAS-DPOL, thus providing better particle sampling statistics. Particles larger than 1.5 μm in radius are classified as cloud, while the smaller particles are considered aerosols.

The PSDs are reported for each second. In the present work the PSDs have been averaged over 10 s and synchronized to the backscattersonde data. In the following, in addition to the PSD, we have used the "combined" aspherical fraction AF_i which refers to fraction of aspherical particles detected in size channels both from NIXE-CAS and NIXE-CIPgs. The AFs are derived as described by Costa et al. (2017)

The CCP (Cloud Combination Probe) combines a CDP (Cloud Droplet Probe) with a CIPgs (Cloud Imaging Probe with grey scale), whose measurement technique as well as the characteristics of the measurement data analysis have already been described in detail (Frey et al., 2011; Molleker et al., 2014; Klingebiel et al., 2015; Grulich et al., 2021). The CCP-CDP is a light scattering probe comparable to the CAS-Dpol (or NIXE-CAS), but it covers the range of 2.5-46 μ m in particle diameter with a size resolution of 1 - 2 μ m (Mei et al., 2020), encompassing also the uppermost range of the aerosol's size spectrum. The sample area of the CCP-CIPgs was examined (Klingebiel et al., 2015), and this analysis yielded 0.27±0.025 mm^2 with an uncertainty of less than 10%. In contrast, the CCP-CIPgs is an optical array probe designed to detect cloud particles and hydro meteors with a resolution of about 15 μ m. CIPgs captures images of cloud elements using a 64-element photodiode array to obtain two-dimensional images with nominal detection diameters ranging from 15 to 960 μ m. CCP-CIPg and NIXE-CAS performances have been frequently validated by glass sphere calibrations. Before or after each flight, CCP-CIPgs and NIXE-CIP calibrations were performed using spinning disks carrying in-transparent spots sized to the particles to be detected. Particle concentration data measured with CCP are corrected for compression under measurement conditions using a thermodynamic approach developed by Weigel et al. (2016).

2.3 FISH and FLASH Hygrometers

The FISH (Fast In situ Stratospheric Hygrometer) instrument is a closed-path Lyman-α photofragment fluorescence hygrometer (Zöger et al., 1999; Meyer et al., 2015) used to measure H₂O_{tot} in the range of 1–1000 ppmv between 50 and 500 hPa with an accuracy and precision during StratoClim of 7% and 0.3 ppmv. FISH is calibrated versus a reference frost point mirror MBW 373 LX on ground before and after each flight to ensure high data quality. On board Geophysica, the inlet for the H₂O_{tot} hygrometer FISH is mounted on the side of the aircraft, heated and with a 90° bend to quickly evaporate ice crystals. In the present work the original 1s resolution data have been averaged over 10s.

FLASH (FLuorescent Airborne Stratospheric Hygrometer; for details see Khaykin et al. (2013, 2022)) also uses the Lyman- α photofragment fluorescence technique for the detection of water vapor, but its inlet is designed to sample only the gas phase. The detection range is 1–1000 ppmv with an accuracy and precision of <9% and 0.5 ppmv. The time resolution is 1 Hz but in the present work data have been averaged over 10s. The H_2O_{gas} hygrometer FLASH is mounted below the wing and equipped with its own inlet.

235

215

220

Clear-sky data from the two hygrometers have been inter-compared together and with those from a third hygrometer on board the M55-Geophysica, ChiWIS (Chicago Water Isotope) designed for airborne measurements of vapour phase water isotopologues in the dry Upper Troposphere - Lower Stratosphere with integrated cavity output absorption spectroscopy (Sarkozy et al., 2020). The comparison showed excellent agreement between these in situ instruments (Singer et al., 2022).

$$AF = \frac{\sum_{i=1}^{\max} AF_i n_i}{\sum_{i=1}^{\max} n_i}$$

2.2.2 The CCP instrument

The CCP (Cloud Combination Probe) combines a CDP (Cloud Droplet Probe) with a CIPgs (Cloud Imaging Probe with grey scale), whose measurement technique as well as the data analysis have been described in detail (Frey et al., 2011; Molleker et al., 2014; Klingebiel et al., 2015; Grulich et al., 2021). The CCP-CDP is a light scattering probe comparable to the CAS-DPOL (or NIXE-CAS), but it covers the range of 2.5-46 µm in particle diameter with a size resolution of 1 - 2 µm (Mei et al., 2020), encompassing also the uppermost range of the aerosol's size spectrum. The sampling area of the CCP-CDP was examined by Klingebiel et al. (2015) yelding 0.27±0.025 mm² with an uncertainty of less than 10%. In contrast, the CCP-CIPgs is an optical array probe designed to detect cloud particles and hydro meteors with a resolution of 15 µm. CIPgs captures images of cloud particles using a 64-element photodiode array to obtain two-dimensional images with nominal detection diameters ranging from 15 to 960 µm. CCP-CDP and NIXE-CAS performances have been frequently validated by glass sphere calibrations. Before or after each flight, CCP-CIPgs and NIXE-CIP calibrations were performed using spinning disks carrying opaque spots sized to the particle range to be detected. Particle concentration data measured with CCP are corrected for compression under measurement conditions using a thermodynamic approach developed by Weigel et al. (2016).

(6)

230 2.3 FISH and FLASH Hygrometers

The FISH (Fast In situ Stratospheric Hygrometer) instrument is a closed-path Lyman- α photofragment fluorescence hygrometer (Zöger et al., 1999; Meyer et al., 2015) used to measure H_2O_{tot} in the range of 1–1000 ppmv between 50 and 500 hPa with an accuracy and precision during StratoClim of 7% and 0.3 ppmv. FISH is calibrated versus a reference frost point mirror MBW 373 LX on ground before and after each flight to ensure high data quality. On board Geophysica, the inlet for the H_2O_{tot} hygrometer FISH is mounted on the side of the aircraft, heated and with a 90° bend to quickly evaporate ice crystals. In the present work the original 1 s resolution data have been averaged over 10 s.

FLASH (FLuorescent Airborne Stratospheric Hygrometer; for details see Khaykin et al. (2013, 2022)) also uses the Lyman- α photofragment fluorescence technique for the detection of water vapor, but its inlet is designed to sample only the gas phase. The detection range is 1–1000 ppmv with an accuracy and precision of <9% and 0.5 ppmv. The time resolution is 1 Hz here data have been averaged over 10 s. The H_2O_{gas} hygrometer FLASH is mounted below the wing and equipped with its own

240

inlet.

235

Clear-sky data from the two hygrometers have been inter-compared together and with those from a third hygrometer on board the M55-Geophysica, ChiWIS (Chicago Water Isotope) designed for airborne measurements of vapour phase water isotopologues in the dry Upper Troposphere - Lower Stratosphere with integrated cavity output absorption spectroscopy (Sarkozy

et al., 2020). The comparison showed excellent agreement between these in situ instruments (Singer et al., 2022).

240 3 Methods

245

250

We first compare the particle backscattering coefficient derived by computing it from the measured PSD, namely $\beta_{NIXE-CAPS}$ with the β measured by the backscattersonde MAS. We then present regressions between particle backscattering coefficient β and particle number density N_{ice} , mean radius R_{mean} , effective radius R_{eff} , SAD and IWC. We will assume that the dispersion of the measurements among the various instruments compared could be used as an estimate of the uncertainty to be attributed to such regressions. In addition, we investigate the relation between the particle aspherical fraction (AF) with the measured total particle depolarization δ_{TA} and with the other cloud microphysical and environmental parameters.

3.1 Optical Modelling

The backscattering coefficient and depolarization ratio have been computed with the GRASP (Generalized Retrieval of Aerosol and Surface Properties) Spheroid Package coupled with Mie scattering computations performed using a code available from the NASA's OceanColor Web site (*https://oceancolor.gsfc.nasa.gov/docs/ocssw/bhmie*₈*py*_s*ource.html*), one of the python's avatars of the Bohren and Huffman fortran code originally published in their book on light scattering (Bohren and Huffman, 2008).

GRASP is the first unified algorithm developed for characterizing atmospheric properties gathered from a variety of remote sensing observations (Dubovik et al., 2014), whose software packages are available on the Web project repository (*https*: //www.grasp - open.com/). The Spheroid Package allows fast, fairly accurate and flexible modeling of single scattering properties by randomly oriented spheroids with aspect ratio spanning from 0.3 to 3. The details of the scientific concept are described in the paper by Dubovik et al. (2006). The code uses kernel look-up tables including results of calculations using T-Matrix codes for particle size parameters where convergence was acquired, and geometric-optics-integral-equation code (Yang and Liou, 1996b, 2006) for greater size parameters where T-matrix codes did not converge. The two methods have been shown to produce comparable results over the size range in which both are applicable (Dubovik et al., 2006). The GRASP spheroid package thus provide backscattering coefficients for randomly oriented spheroids with Aspect Ratios (AR) from ~ 0.3 (flattened spheroids) to ~ 3.0 (elongated spheroids) and covering size parameters from ~ 0.01 to ~ 517 (when a wavelength of 532*nm* is used) for a wide range of the particle complex refractive index, the real part form 1.3 to

1.6, the imaginary part from 0.0005

265

to

0.5. Since in our case the size parameters of the particles detected by the cloud spectrometers extend for more than an order of magnitude beyond the GRASP limit, we were forced to extrapolate the GRASP results using an approximation that makes use of the Mie code, even for aspherical particles.

270

To justify this extrapolation, we recall some characteristics of the backscattering and of the depolarization from aspherical scatterers, depending on their size. According to T-Matrix computations on randomly oriented spheroids and cylinders, depolarization is negligible when their size parameters less than unity (given the wavelength used in our study, this corresponds to equivalent radius around 0.1 μ m) Then it grows to a maximum for size parameters of the order of ten (equivalent radius

3 Methods

250

We first compare the particle backscattering computed from the measured PSD, namely $\beta_{NIXE-CAPS}$ with the β measured by the backscattersonde MAS. We then present regressions between particle backscattering coefficient β and particle number density N_{ice} , mean radius R_{mean} , effective radius R_{eff} , SAD and IWC. We will assume that the dispersion of the data in the comparison could be used as an estimate of the uncertainty to be attributed to such regressions. In addition, we investigate the relation between the particle aspherical fraction (AF) with the measured total particle depolarization δ_{TA} and with other cloud microphysical and environmental parameters.

3.1 Optical Modelling

- The backscattering coefficient and depolarization ratio have been computed with the GRASP (Generalized Retrieval of Aerosol and Surface Properties) Spheroid Package coupled with Mie scattering computations performed using a code available from the NASA's OceanColor Web site (https://oceancolor.gsfc.nasa.gov/docs/ocssw/bhmie_8py_source.html), one of the python's avatars of the Bohren and Huffman fortran code originally published in their book on light scattering (Bohren and Huffman, 2008).
- GRASP is the first unified algorithm developed for characterizing atmospheric properties gathered from a variety of remote sensing observations (Dubovik et al., 2014), whose software packages are available on the Web project repository (https://www.grasp-open.com/). The Spheroid Package allows fast, fairly accurate and flexible modeling of single scattering properties by randomly oriented spheroids. The details of the scientific concept are described in the paper by Dubovik et al. (2006). The code uses kernel look-up tables including results of calculations using T-Matrix codes for particle size parameters
- where convergence was acquired, and geometric-optics-integral-equation code (Yang and Liou, 1996b, 2006) for greater size parameters where T-matrix codes did not converge. The two methods have been shown to produce comparable results over the size range in which both are applicable (Dubovik et al., 2006). The GRASP spheroid package thus provide backscattering coefficients for randomly oriented spheroids with Aspect Ratios (AR) from ~ 0.3 (flattened spheroids) to ~ 3.0 (elongated spheroids) and covering size parameters from ~ 0.01 to ~ 517 (when a wavelength of 532nm is used) for a wide range of the
- 270 particle complex refractive index, the real part from 1.3 to 1.6, the imaginary part from 0.0005 to 0.5. Since in our case the size parameters of the particles detected by the cloud spectrometers extend for more than an order of magnitude beyond the GRASP limit, we were forced to extrapolate the GRASP results using an approximation that makes use of the Mie code, even for aspherical particles.

To justify this extrapolation, we recall some characteristics of the backscattering and of the depolarization from aspherical

scatterers, depending on their size. According to T-Matrix computations on randomly oriented spheroids and cylinders, depolarization is negligible when their size parameters less than unity (given the wavelength used in our study, this corresponds to equivalent radius around 0.1 μ m). Then it grows to a maximum for size parameters of the order of ten (equivalent radius around 1 μ m), to decrease to an asymptotic value which stabilizes for size parameters approximately greater than one hundred (particle radius greater than 10 μ m). The maximum and the asymptotic values of the depolarization vary in dependence of the

specific AR. The variability of these values are unrelated, and there is no general relationship that links the peak and asymptotic

275

280

285

290

depolarization values to the AR of the spheroids.

For what concerns the backscattering, when the particle size parameter is below unity the T-matrix backscattering efficiency reproduces the Mie results for surface equivalent spheres, then reaches a minimum as the size parameter increases, to go up again and stabilize, at a few tens of size parameters, at values in a constant relationship with Mie's calculations. This constant is often less than unity; it can exceed the unit by a few decimal places in a few cases, for ARs around 0.5. (Mishchenko et al., 1996, 2002). The increase in backscattering for AR around 0.5, probably due to a sort of mirror reflection effect that predominates with respect to the backscattering of spheres of equivalent radius, does not exceed a factor of 1.5-2. Conversely the backscattering depression, which is a more general feature of aspherical scatterers, can be as large as a factor 10 or more. The dependence of the single particle's depolarization on its shape and size has been studied extensively by Liu and Mishchenko (2001).

The GRASP calculations extend well into the region of asymptotic values for the depolarization and for the spherical vs aspherical ratio of backscattering, so that we can extrapolate its results with confidence. Beyond the computational limits of GRASP we have in fact set the depolarization of the particles as constant, i.e. equal to its asymptotic value. Moreover we have used the Mie code for the calculation of the backscattering efficiencies, suitably rescaled by a constant factor so as to make the scaled Mie backscattering to overlap with the asymptotic values calculated by GRASP. Such constant factor was calculated from the ratio of Mie vs T-matrix efficiencies in the radius dimensional region from 5 to 14 μ m (60-180 size parameters). It turned out to be always less than unity.

We thus calculated, for 25 different Aspect Ratios from 0.3 to 3, the backscattering efficiencies Q_l^{AR} and the depolarizations δ_l^{AR} on a grid of 2000 radius points R_l equally spaced on a logarithmic scale and extending from 0.005 to 1000 μ m. We used the GRASP computations for radii below 14 μ m and we extrapolated these result with the Mie code for larger radii, as outlined above. We used the value of 1.31 as refractive index relative to ice. To get rid of the oscillating nature of the backscattering efficiency, for each radius R_l the values of Q_l^{AR} and δ_l^{AR} were actually obtained as averages over a narrow lognormal distribution centered at R_l , with variance 1.01. Such averages were computed over the lognormal with a finer subgrid of 500 points R_k , equally spaced on a logarithmic scale and extending over 1- σ from the lognormal center .

300

Figure A1 and A2 in the Appendix reports the results of such extrapolation. In those figures, particle backscattering coefficients and depolarization ratio have been calculated for a reference particle density of 1 cm⁻³, for different AR, and displayed for particle radius from 0.01 to 100 μ m.

These efficiencies were then used to calculate the backscattering coefficients associated with the PSD measurements. For each PSD size bin i, we computed the arithmetic average of the M_i lattice radius points falling within the size channel limit around 1 μ m), to decrease to an asymptotic value which stabilizes for size parameters approximately greater than one hundred (equivalent radius greater than 10 μ m). The maximum and the asymptotic values of the depolarization vary in dependence

of the specific AR. The variability of these values are unrelated, and there is no general relationship that links the peak and

280

asymptotic depolarization values to the AR of the spheroids.

For what concerns the backscattering, when the particle size parameter is below unity the T-matrix backscattering efficiency reproduces the Mie results for surface equivalent spheres, then reaches a minimum as the size parameter increases, to go up again and stabilize, at a few tens of size parameters, at values in a constant relationship with Mie's calculations. This constant

is often less than unity; it can exceed the unit by a few decimal places in a few cases, for ARs around 0.5. (Mishchenko et al., 1996, 2002). The increase in backscattering for AR around 0.5, probably due to a sort of mirror reflection effect that predominates with respect to the backscattering of spheres of equivalent radius, does not exceed a factor of 1.5-2. Conversely the backscattering depression, which is a more general feature of aspherical scatterers, can be as large as a factor 10 or more. The dependence of the single particle's depolarization on its shape and size has been studied extensively by Liu and Mishchenko (2001).

The GRASP calculations extend well into the region of asymptotic values for the depolarization and for the spherical vs aspherical ratio of backscattering, so that we can extrapolate its results with confidence. Beyond the computational limits of GRASP we have thus set the depolarization of the particles as constant, i.e. equal to its asymptotic value. Moreover we have used the Mie code for the calculation of the backscattering efficiencies, suitably rescaled by a constant factor so as to make the scaled Mie backscattering to overlap with the GRASP asymptotic values. Such constant factor was calculated from the ratio of Mie vs T-matrix efficiencies in the radius dimensional region from 5 to 14 μ m (60-180 size parameters). It turned out to be

always less than unity.

We thus calculated, for 25 different Aspect Ratios from 0.3 to 3, the backscattering efficiencies Q_l^{AR} and the depolarizations δ_l^{AR} on a grid of 2000 radius points R_l equally spaced on a logarithmic scale and extending from 0.005 to 1000 μ m. We 300 used the GRASP computations for radii below 14 μ m and we extrapolated these result with the Mie code for larger radii, as outlined above. We used the value of 1.31 as refractive index relative to ice. To get rid of the oscillating nature of the backscattering efficiency, for each radius R_l the values of Q_l^{AR} and δ_l^{AR} were actually obtained as averages over a narrow lognormal distribution centered at R_l , with variance 1.01. Such averages were computed over the lognormal with a finer subgrid of 500 points R_k , equally spaced on a logarithmic scale and extending over 1- σ from the lognormal center .

305

295

Figure A1 and A2 in the Supplementary Material reports the results of such extrapolation. In those figures, particle backscattering coefficients and depolarization ratio have been calculated for a reference particle density of 1 cm⁻³, for different AR, and displayed for particle radius from 0.01 to 100 μ m.

These efficiencies were then used to calculate the backscattering coefficients associated with the PSD measurements. For each PSD size bin i, we computed the arithmetic average of the M_i lattice radius points falling within the size channel limit

305 and the corresponding average efficiency and depolarization

$$r_{i} = \frac{\sum_{l=1}^{M_{i}} R_{l}}{M_{i}}$$

$$Q_{i}^{AR} = \frac{\sum_{l=1}^{M_{i}} Q_{l}^{AR}}{M_{i}}$$

$$\delta_{Ai}^{AR} = \frac{\sum_{l=1}^{M_{i}} \delta_{Al}^{AR}}{M_{i}}$$

$$(10)$$

then we used the concentration of particles n_i in the size bin *i* to derive the contribution of that bin to the backscattering coefficient and the depolarization, and summed over the bins. In the case of depolarization, the average was weighted with the backscattering coefficient of the same channel. This was repeated for all the 25 AR.

$$\beta_i^{AR} = n_i \pi r_i^2 Q_i^{AR} \tag{11}$$

$$\beta_{NC}^{AR} = \sum_{i=1}^{N} \beta_{i}^{AR}$$

$$\delta_{ANC}^{AR} = \frac{\sum_{i=1}^{max} \delta_{i}^{AR} \beta_{i}^{AR}}{\beta_{NC}^{AR}}$$
(12)
(13)

Uncertainties in the backscattering coefficient $\Delta(\beta_{NC}^{AR})$ and $\Delta(\delta_{ANC}^{AR})$ due to sizing and counting errors were derived using a weighted error propagation in quadrature method (Berendsen, 2011):

$$\Delta(\beta_{NC}^{AR}) = \sum_{i=1}^{max} \sqrt{\left(\pi r_i^2 Q_i^{AR} \Delta(n_i)\right)^2 + \left(2\pi r_i Q_i^{AR} \Delta(r_i)\right)^2} \tag{14}$$

$$\Delta(\delta_{ANC}^{AR}) = \frac{1}{\beta_{NC}^{AR}} \left(\sum_{i=1}^{max} \Delta_i^{AR} \sqrt{\left(\pi r_i^2 Q_i^{AR} \Delta(n_i)\right)^2 + \left(2\pi r_i Q_i^{AR} \Delta(r_i)\right)^2} + \delta_{NC}^{AR} \Delta(\beta_{NC}^{AR}) \right)$$

$$(15)$$

where $\Delta(n_i)$ is the uncertainty in concentration following Poisson statistics and $\Delta(r_i)$ is the uncertainty in particle radius for channel *i*, (Horvath et al., 1990, Baumgardner et al. 2014) here one half the channel width has been used as radius uncertainty. (Wagner and Delene, 2022).

$$\Delta(n_i) = \frac{\sqrt{n_i}}{n_i} \tag{16}$$
$$\Delta(r_i) = \frac{\Delta r_i}{2} \tag{17}$$

325

To perform the optical computations we use the complete dimensional range of NIXE-CAPS particle detectability, which extends lower down to 0.3 μ m in radius. This was done because the measurement of β made by the backscattersonde is sensitive to the backscattering from all the particles present in the sampled air mass. An estimation of the contribution of particles in the 0.3-1.5 range can be given by inspecting Figure A7 in the Appendix, where the occurrence histograms of the computation of β

310 and the corresponding average efficiency and depolarization

$$r_{i} = \frac{\sum_{l=1}^{M_{i}} R_{l}}{M_{i}}$$

$$Q_{i}^{AR} = \frac{\sum_{l=1}^{M_{i}} Q_{l}^{AR}}{M_{i}}$$

$$\delta_{Ai}^{AR} = \frac{\sum_{l=1}^{M_{i}} \delta_{Al}^{AR}}{M_{i}}$$

$$(9)$$

then we used the concentration of particles n_i in the size bin *i* to derive the contribution of that bin to the backscattering 315 coefficient and the depolarization, and summed over the bins. In the case of depolarization, the average was weighted with the backscattering coefficient of the same channel. This was repeated for all the 25 AR.

$$\beta_i^{AR} = n_i \pi r_i^2 Q_i^{AR}$$

$$\beta_{NC}^{AR} = \sum_{i=1}^{max} \beta_i^{AR}$$
(10)
(11)

$$\delta_{ANC}^{AR} = \frac{\sum_{i=1}^{max} \delta_{Ai}^{AR} \beta_i^{AR}}{\beta_{NC}^{AR}}$$
(12)

320 Uncertainties in the backscattering coefficient $\Delta(\beta_{NC}^{AR})$ and $\Delta(\delta_{ANC}^{AR})$ due to sizing and counting errors were derived using a weighted error propagation in quadrature method (Berendsen, 2011):

$$\Delta(\beta_{NC}^{AR}) = \sum_{i=1}^{max} \sqrt{\left(\pi r_i^2 Q_i^{AR} \Delta(n_i)\right)^2 + \left(2\pi r_i Q_i^{AR} \Delta(r_i)\right)^2} \tag{13}$$

$$\Delta(\delta_{ANC}^{AR}) = \frac{1}{\beta_{NC}^{AR}} \left(\sum_{i=1}^{max} \Delta_i^{AR} \sqrt{\left(\pi r_i^2 Q_i^{AR} \Delta(n_i)\right)^2 + \left(2\pi r_i Q_i^{AR} \Delta(r_i)\right)^2} + \delta_{NC}^{AR} \Delta(\beta_{NC}^{AR}) \right)$$
(14)

where $\Delta(n_i)$ is the uncertainty in concentration following Poisson statistics and $\Delta(r_i)$ is the uncertainty in particle radius for 325 channel *i*, (Horvath et al., 1990, Baumgardner et al. 2014) here one half the channel width has been used as radius uncertainty. (Wagner and Delene, 2022).

$$\Delta(n_i) = \frac{\sqrt{n_i}}{n_i} \tag{15}$$

$$\Delta(r_i) = \frac{\Delta r_i}{2} \tag{16}$$

330

For the optical computations we use the complete dimensional range of NIXE-CAPS particle detectability, which extends lower down to 0.3 μ m in radius. This because the β provided by the backscattersonde is sensitive to all the particles (i.e. clouds and aerosol) present in the air mass. An estimation of the contribution of particles in the 0.3-1.5 range is given in Fig. A3 in the Supplementary Material, where the occurrence histograms of the computation of β for NIXE-CAPS measuring from 0.3 for NIXE-CAPS measuring from 0.3 μ m (vertical axis) or 1.5 μ m (horizontal axis) is reported. Such contribution is always low and negligible, except in some cases at medium-to-low values of the β , where neglecting the 0.3-1.5 1.5 μ m part of the PSD can lead to a detectable underestimation of the β_{NC} , roughly of a factor 2.

330

335

3.2 Cirrus cloud microphysical parameters

For the comparison with the measured backscatter signals, we exclude the aerosol component of the particulate, by setting 1.5 μ m in radius as a lower limit for the cloud particle radius. We calculate the cloud particle concentration N_{ice} , mean mass radius R_{mean} (Krämer et al., 2009) calculated from IWC/ N_{ice} – the particles assumed as spheres, effective radius R_{eff} defined as in (18) (Schumann et al., 2011) where the definition on the second equality applies only to spherical particles and has not been used in the present work,

$$R_{eff} = \frac{3VD}{4SAD} = \frac{\int r^3 PSD(r) dr}{\int r^2 PSD(r) dr}$$
⁽¹⁸⁾

and SAD, VD stands for, respectively, Surface Area Density and Volume Density. To compute SAD and VD we have used the the m–D relation described in Krämer et al. (2016). To this end, the PSD were averaged over 10 s. To retrieve cirrus cloud IWC
the FLASH water vapor measurements were subtracted from the FISH total water data, once corrected for inlet enhancements (see Afchine et al. (2018)). This was done for 5 out of the 8 flights under assessment. For two flights (on 31 July and 4 August) when FISH was not operating, the IWC was computed from NIXE-CAPS particle volumes with an estimate of the ice density of 0.92 g cm ⁻³. An extensive assessment of the methodology to retrieve IWC from the two hygrometers and from the PSDs, as well as a comparison between the two approaches, can be found in Afchine et al. (2018) who demonstrated the equivalence
of the two methods.

4 Results

350

355

4.1 Cloud Spectrometers Comparison

We have first compared the PSD from the two cloud spectrometers in terms of N_{ice} , R_{mean} , SAD and VD, these latter evaluated in terms of spherical particle approximation. The particle backscattering from the PSD, namely $\beta_{NIXE-CAPS}$, and β_{CCP} were computed based on Mie theory and subsequently compared.

The NIXE-CAPS N_{ice} resulted to be lower, of roughly a factor 2, than the corresponding one measured by CCP in the low to medium particle concentration regime, while the two determinations resulted to be more on a 1:1 line in the high particle concentration regime. Conversely NIXE-CAPS R_{mean} resulted larger than the corresponding from what measured by CCP in the mid to high R_{mean} regime, of roughly a factor 2 (see Figures A3 and A4 in Appendix). With NIXE-CAPS measuring larger and, at low concentration, fewer particles, and wider PSD as well, not surprisingly we found a slight mismatch also in surfaces and volumes, with NIXE-CAPS measuring SAD and VD a factor 2 larger on the average than the corresponding CCP acquisitions. μ m (vertical axis) or 1.5 μ m (horizontal axis) is reported. Such contribution is always low and negligible, except in some cases at medium-to-low values of the β , where neglecting the 0.3-1.5 1.5 μ m part of the PSD can lead to an underestimation of the β_{NC} roughly of a factor 2.

335

340

3.2 Cirrus cloud microphysical parameters

In the estimation of the cloud microphysical parameters we exclude the aerosol component of the particulate by setting 1.5 μ m in radius as a lower limit for the cloud particle. We calculate the cloud particle concentration N_{ice} , mean mass radius R_{mean} (Krämer et al., 2009) calculated from IWC/ N_{ice} – the particles assumed as spheres, effective radius R_{eff} defined as in (17) (Schumann et al., 2011) where the definition on the second equality applies only to spherical particles and has not been used in the present work,

$$R_{eff} = \frac{3VD}{4SAD} = \frac{\int r^3 PSD(r) dr}{\int r^2 PSD(r) dr}$$
(17)

and SAD, VD stands for, respectively, Surface Area Density and Volume Density. To compute SAD and VD we have used the the m–D relation described in Krämer et al. (2016), i.e m=a $\cdot D^b$ where a = 0.001902, b = 1.802 for D > 240 μ m; a = 0.058000, b = 2.700 for D = 10–240 μ m; ice crystals are spheres for D < 10 μ m. The validity of the m-D relation is verified by Afchine et al. (2018) by comparison to thirteen others. The PSD were averaged over 10 s. To retrieve cirrus cloud IWC the FLASH water vapor measurements were subtracted from the FISH total water data, once corrected for inlet enhancements (see Afchine et al. (2018)). This was done for 5 out of the 8 flights under assessment. For two flights (on 31 July and 4 August) when FISH was not operating, the IWC was computed from NIXE-CAPS particle volumes with an estimate of the ice density of 0.92 g 350 cm⁻³. An extensive assessment of the methodology to retrieve IWC from the two hygrometers and from the PSDs, as well as a comparison between the two approaches, can be found in Afchine et al. (2018) who demonstrated the equivalence of the two methods.

4 Results

4.1 Cloud Spectrometers Comparison

355 We have first compared the PSD from the two cloud spectrometers in terms of N_{ice} , R_{mean} , SAD and VD, these latter evaluated in terms of spherical particle approximation. The particle backscattering from the PSD, namely $\beta_{NIXE-CAPS}$, and β_{CCP} were computed based on Mie theory and subsequently compared.

The NIXE-CAPS N_{ice} resulted to be lower, of roughly a factor 2, than the corresponding one measured by CCP in the low to medium particle concentration regime, while the two determinations resulted to be more on a 1:1 line in the high particle 360 concentration regime. Conversely NIXE-CAPS R_{mean} resulted larger than the corresponding from what measured by CCP in the mid to high R_{mean} regime, of roughly a factor 2 (see Figs. A4 and A5 in Supplementary Material). With NIXE-CAPS measuring larger and, at low concentration, fewer particles, and wider PSD as well, not surprisingly we found a slight mismatch

The comparison of the the particle backscattering $\beta_{NIXE-CAPS}$, and β_{CCP} computed from the PSD by use of Mie theory again produced $\beta_{NIXE-CAPS}$ larger than the corresponding β_{CCP} by roughly a factor 2 (see figure A5 in Appendix).

An in-depth analysis of the datasets from the two cloud spectrometers showed that in the common size range of 1.25 - 15 360 μ m particle radius, CAS and CDP agreed very well with each other. Conversely, an offset was found between NIXE-CAPS-CIPgs and CCP-CIPgs-data which was not a product of different evaluation methods or filter criteria of the image files but was very likely due to a hardware problem. A failure of the lasers' temperature stabilization was identified to lead to a loss of beam intensity at higher altitudes and therefore a less illuminated sample volume and diode array of the CCP-CIPgs. This 365 resulted in a slightly lower instrument particle detection sensitivity which could be only identified by a comparison of particle habits like size and number concentrations in the range of both CIPgs instruments measured at significantly elevated altitudes (Port, 2021). An implication of this finding on a similar comparison of the same two instruments from previous simultaneous measurements at lower altitudes (Mei et al., 2020) can be ruled out. Therefore for all further analyses of the optical and microphysical parameters, only the NIXE-CAPS data set was used.

370 4.2 **Optical modelling and measurements**

To compare the β_{NC}^{AR} backscattering calculated with optical modeling applied to the PSDs, we select point-by-point the AR that provides the best match with the β measured by the backscattersonde. In order to illustrate the results, in the upper panels of Figure 3 we report the time series of the β measured by the backscattersonde MAS during the flight on 10 Aug 2017 (red line), together with the best choice of the calculated β_{NC} (black line), with representative error bars in the first part of the time serie. The black dashed lines report the highest and lowest values of the β_{NC}^{AR} . The agreement between the measured backscattering β and the selected one among the optical computations β_{NC} can be made surprisingly good almost everywhere, except at its highest values - where the loss of linearity in the response of the backscattersonde could play a role - and at its lowest values - where it is possible that backscattering from particles below the minimum cloud spectrometer detectable size can become non-negligible. It is noticeable how in many parts of the time series the calculated value β_{NC} is able to reproduce even the finest structures of the measured β . From the figure we note that the selected β_{NC} is often in the lower range of variability of the β_{NC}^{AR} values. Moreover, the random error attributed to β_{NC} is an order of magnitude lower than the uncertainty range deriving from the lack of knowledge in the particulate AR. In fact, the particle AR is probably the main factor of uncertainty to be attributed to our optical calculations.

385

375

380

As a test for the arbitrariness of the choice of AR that ensures the best agreement with the measurements, we looked for possible correlations between the selected AR and i. the particle depolarization δ_A ; the fraction of aspheric particles AF. In the lower panels of Figure 3 the AR that provided the best match is reported (black line) together with the observed particle depolarization (red line) and particle AF, as measured by NIXE-CAPS (blue line). We note that the AR values that made the best match are arranged according to a certain temporal continuity, and we can identify well-defined time intervals in which their value remains almost constant. This is encouraging to think that the choice of AR that best matches the measures is not 390 exclusively an exercise of handpicking, but rather reflects characteristics of the average morphology of the measured particulate matter. It is of particular interest to note that regions with AR = 3 - that provides the lowest β_{NC}^{AR} - almost always coincide with

also in surfaces and volumes, with NIXE-CAPS measuring SAD and VD a factor 2 larger on the average than the corresponding CCP acquisitions.

The comparison of the the particle backscattering $\beta_{NIXE-CAPS}$, and β_{CCP} computed from the PSD by use of Mie theory 365 again produced $\beta_{NIXE-CAPS}$ larger than the corresponding β_{CCP} by roughly a factor 2 (see Fig. A6 in Supplementary Material).

An in-depth analysis of the datasets from the two cloud spectrometers showed that in the common size range of 1.25 - 15 μ m particle radius, CAS and CDP agreed very well with each other. Conversely, an offset was found between NIXE-CAPS-370 CIPgs and CCP-CIPgs data which was not a product of different evaluation methods or filter criteria of the image files but was very likely due to a hardware problem. A failure of the lasers' temperature stabilization was identified to lead to a loss of beam intensity at higher altitudes and therefore a less illuminated sample volume and diode array of the CCP-CIPgs. This resulted in a slightly lower instrument particle detection sensitivity which could be only identified by a comparison of particle habits like size and number concentrations in the range of both CIPgs instruments measured at significantly elevated altitudes 375 (Port, 2021). An implication of this finding on a similar comparison of the same two instruments from previous simultaneous measurements at lower altitudes (Mei et al., 2020) can be ruled out. Therefore for all further analyses of the optical and

4.2 **Optical modelling and measurements**

microphysical parameters, only the NIXE-CAPS data set was used.

To compare the β_{NC}^{AR} backscattering calculated with optical modeling applied to the PSDs, we select point-by-point the β_{NC}^{AR} 380 that provides the best match with the β measured by the backscattersonde and let β_{NC} be this best match. In order to illustrate the results, in the upper panels of Fig. 3 we report the time series of the β measured by the backscattersonde MAS during the flight on 10 Aug 2017 (red line), together with the best choice of the calculated β_{NC} (black line), with representative error bars in the first part of the time serie. The black dashed lines report the highest and lowest values of the β_{NC}^{AR} . The agreement between the measured backscattering β and the selected one among the optical computations β_{NC} can be made surprisingly 385 good almost everywhere, except at its highest values - where the loss of linearity in the response of the backscattersonde could play a role - and at its lowest values - where it is possible that backscattering from particles below the minimum cloud spectrometer detectable size can become non-negligible. It is noticeable how in many parts of the time series the calculated value β_{NC} is able to reproduce even the finest structures of the measured β . From the figure we note that the selected β_{NC} is often in the lower range of variability of the β_{NC}^{AR} values. Moreover, the random error attributed to β_{NC} is an order of 390 magnitude lower than the uncertainty range deriving from the lack of knowledge in the particulate AR. In fact, the particle AR is probably the main factor of uncertainty to be attributed to our optical calculations.

As a test for the arbitrariness of the choice of AR that ensures the best agreement with the measurements, we looked for possible correlations between the selected AR and i. the particle depolarization δ_A ; ii. the fraction of aspheric particles AF. In the lower panels of Fig. 3 the AR that provided the best match is reported (black line) together with the observed particle depolarization (red line) and particle AF, as measured by NIXE-CAPS (blue line). We note that the AR values that made the best match are arranged according to a certain temporal continuity, and we can identify well-defined time intervals in which

regions in which the NIXE-CAPS has observed the highest particle AF. This, in turn, is often correlated with high values of total depolarization.

395

We have explored a possible lack of linearity in the response of the backscattersonde at high backscattered signals as a possible cause for the discrepancy between the highest calculated and measured β . The linear response of the backscattersonde was checked in the laboratory before the campaign deployment. This was done by screening the receiving optics with a series of calibrated gray filters and controlling the response of the instrument to pulsed light of various intensity. This procedure was repeated at various levels of background light. For the present study a further test was carried out by comparing the backscattering coefficient observed during the campaign by the backscattersonde in some of the thickest clouds, on 8 August 400 2017 between 05:17 and 05:50 UTC (19000s and 21000s) and on 10 August 2017 between 10:00 and 10:33 UTC (36000s to and 38000s), with what observed by the two MAL lidars on board the Geophysica. These two lidars point respectively upward and downward providing profiles of Backscatter Ratio and depolarization. We have used signals from the closest lidar range, in its partial overlap region, and processed them as if they came from backscattersondes. The average of the two lidar backscattering coefficients observed by them at a distance of 500 meters upward and downward was then compared with what 405 observed by MAS. The data have been averaged over 60 s. The result is displayed in Figure 4. Despite the scattering of the data points, which might be attributed to a lack of vertical homogeneity of the cloud, it shows that there is a good correlation between the Backscattering coefficient measured in-situ and that measured at close ranges. The lidar backscatter signal is measured in photon counting mode. The linear dynamic range is identified to be below the count rate of approx. 1.5 MHz. The complete signal saturation is noticed at approx. 15 MHz count rate. For the cases reported here the signals are at the levels of 410 some tens of KHz, i.e. well inside the linearity range. Thus we may exclude any loss of linearity. So we are tempted to exclude a severe loss of linearity of the backscattersonde and a consequent significant underestimation of the largest backscattering coefficients observed by it. Further sustaining our conclusion, we note that the overestimation of the optical model appears also in regions where saturation of the backscattersonde signal should not be expected, as instance, around 34000s on the 10 August 2017 flight.

- 415 Figures 5 and 6 report the two-dimensional histograms of occurrence of the measured-calculated backscattering coefficients and particle depolarization. While for the β the agreement is good and the analysis on the whole campaign dataset comfirms what has already been shown in the previously discussed time series, i.e. a underestimation of the β_{NC} for values of β below $5 \ 10^{-5} \ \text{km}^{-1} \ \text{sr}^{-1}$, and a slight overestimation for high values in a relatively insignificant number of cases, the agreement for the depolarization is not good. From the inspection of Figure 6 we can see an attempt to achieve a 1-1 correlation between the
- 420 two datasets. However, for a considerable number of observations, the calculated depolarization remains around 10% while the measured one varies throughout its range of variability. Moreover the optical modelling completely fails to reproduce the measured depolarization. We have tested how to improve the agreement by linking the choice of AR to the matching with depolarization rather than with backscattering. Of course, the comparison between measured and modeled depolarization improves but in the same way leads to a larger disagreement in the backscatter, which in many cases is strongly overestimated by the optical model by up to 1-2 orders of magnitude. (see figure A6 in Appendix). Even where the agreement between the 425 calculated and the measured β is kept within an order of magnitude, it fails to reproduce effectively the experimental dataset,

their value remains almost constant. This is encouraging to think that the choice of AR that best matches the measures is not exclusively an exercise of handpicking, but rather reflects characteristics of the average morphology of the measured particles. It is of particular interest to note that regions with AR = 3 - that provides the lowest β_{NC}^{AR} - almost always coincide with regions

400

in which the NIXE-CAPS has observed the highest particle AF. This, in turn, is often correlated with high values of total depolarization. Moreover, looking at the entire dataset, we have noted a tendency to have ARs close to 1 associated with small β values and depolarization of less than 20%, while ARs greater than 1.5 and less than 0.5 favors medium-high β values and depolarization greater than 20% (see Fig. A7 in the Supplementary Material).

- We have explored a possible lack of linearity in the response of the backscattersonde at high backscattered signals as a 405 possible cause for the discrepancy between the highest calculated and measured β . The linear response of the backscattersonde was checked in the laboratory before the campaign deployment. This was done by screening the receiving optics with a series of calibrated gray filters and controlling the response of the instrument to pulsed light of various intensity. This procedure was repeated at various levels of background light. For the present study a further test was carried out by comparing the backscattering coefficient observed during the campaign by the backscattersonde in some of the thickest clouds, on 8 August 2017
- 410 between 05:17 and 05:50 UTC (19000s and 21000s, see Fig. A8 in Supplementary Material) and on 10 August 2017 between 10:00 and 10:33 UTC (36000s to and 38000s), with what observed by the two MAL lidars on board the Geophysica. These two lidars point respectively upward and downward providing profiles of Backscatter Ratio and depolarization. We have used signals from the closest lidar range, in its partial overlap region, and processed them as if they came from backscattersondes. The average of the two lidar backscattering coefficients observed by them at a distance of 500 meters upward and downward
- 415 was then compared with what observed by MAS. The data have been averaged over 60 s. The result is displayed in Fig. 4. Despite the scattering of the data points, which might be attributed to a lack of vertical homogeneity of the cloud, it shows that there is a good correlation between the backscattering coefficient measured in-situ and that measured at close range. The lidar backscatter signal is measured in photon counting mode. The linear dynamic range is identified to be below the count rate of approx. 1.5 MHz. The complete signal saturation is noticed at approx. 15 MHz count rate. For the cases reported here the
- 420 signals are at the levels of some tens of KHz, i.e. well inside the linearity range. Thus we may exclude any loss of linearity. So we are tempted to exclude a severe loss of linearity of the backscattersonde and a consequent significant underestimation of the largest backscattering coefficients observed by it. Further sustaining our conclusion, we note that the overestimation of the optical model appears also in regions where saturation of the backscattersonde signal should not be expected, for instance, around 34000s on the 10 August 2017 flight.
- Figures 5 and 6 report the two-dimensional histograms of occurrence of the measured-calculated backscattering coefficients and particle depolarization. While for the β the agreement is good and the analysis on the whole campaign dataset comfirms what has already been shown in the previously discussed time series, i.e. a underestimation of the β_{NC} for values of β below 5 10⁻⁵ km⁻¹ sr⁻¹, and a slight overestimation for high values in a relatively insignificant number of cases, the agreement for the depolarization is not good. From the inspection of Fig. 6 we see that there are few cases when the two depolarization
- 430 line up to a 1-1 correlation. However, for a considerable number of observations, the calculated depolarization remains around 10% while the measured one varies throughout its range of variability. Moreover the optical modelling performance is very

as reported in Figures 5 and 6. In some ways it is unfortunate not to be able to simulate backscattering and depolarization simultaneously with a single choice of AR. However, we are aware of the fact that the modeling of depolarization is an open and highly challenging problem, one that is not easy and perhaps impossible to solve given the wide variety of shapes that atmospheric ice crystals can take (Liou and Yang, 2016). For this reason we prefer to keep our comparison between optical measurements and calculations limited to the backscattering coefficient.

430

4.3 Backscattering versus Particle Size Distribution bulk parameters

Figures 7, 8, 9, 10 and 11 show N_{ice} , R_{mean} , R_{eff} , SAD and IWC as a function of β , respectively. In the graphs, the black lines represent regression curves, parametrized in the general form $X = A \cdot \beta^B$. The coefficients of the regression are reported in Table 1 together with the R-squared of the fit. The dataset has been fitted between the limits $2 \cdot 10^{-5}$ and $5 \cdot 10^{-3}$ km⁻¹ sr⁻¹.

- **435** in Table 1 together with the R-squared of the fit. The dataset has been fitted between the limits $2 \cdot 10^{-5}$ and $5 \cdot 10^{-3}$ km⁻¹ sr⁻¹, which have been chosen to maximize the goodness of the fit and try to avoid outliers at the extremes of the variability range. We can regard those regression curves as guidance for estimating the microphysics bulk parameters of the clouds and their variability range, from remote measurements of cloud optical parameters.
- The linearity between the β and N_{ice} (Fig. 7) is quite striking and indicates that β basically scales with the cloud particle number density N_{ice} , as seems to do SAD and IWC. Hence, this suggests that the various shapes of the PSD in our observations are hardly change the scattering properties of the clouds. This finding is further confirmed by the linearity seen between β and SAD or IWC (Figs.10, 11), and the lack of clear correlation between β and R_{mean} or R_{eff} (Figs. 8, 9).

This is most probably because the majority of measurements during StratoClim is at T < 200K, where there is only little water available thus in- situ origin ice particles cannot develop a variety in shapes – in fact most of them are so called "irregulars". Only the large liquid-origin ice particles can have distinct shapes, but they are rare.

4.4 Depolarization, Aspherical Fraction and PSD parameters

4.4.1 Depolarization vs Aspherical Fraction

In our study we have looked for and found no direct correlation between particle depolarization observed with the backscattersonde, and PSD Aspherical Fraction (AF). AF has peaks around 60% but on the average is maintained around values of
 20%, while the corresponding values of the depolarization spans its entire range of variability, giving an unclear relationship between the two quantities (see figure A8 in the Supplementary Material).

This is possibly because the maximum detectable AF of ice particles decreases with the size of the particles. The reason is that the depolarization signal becomes weaker with decreasing ice particle size and also the ice particles become lesser and lesser aspheric, and the backscattersonde and the cloud spectrometer have depolarization sensitivities that vary differently with the size of the scattering particle.



445

In order to display other parameters along with the former two, that may help to find and possibly disentangle a link between these two quantities, we choose to represent our dataset in a δ_T - BR space (in fact, for ease of scale, 1-1/BR is used instead of BR), and color code the points with AF, N, R_{mean} and temperature T as possible parameters of interest.

poor in reproducing the measured depolarization. We have tested how to improve the agreement by linking the choice of AR to the matching with depolarization rather than with backscattering. Of course, the comparison between measured and modeled depolarization improves but in the same way leads to a larger disagreement in the backscatter, which in many cases is strongly

435

overestimated by the optical model by up to 1-2 orders of magnitude. (see Fig. A9 in Supplementary Material). Even where the agreement between the calculated and the measured β is kept within an order of magnitude, it fails to reproduce effectively the experimental dataset, as reported in Figs. 5 and 6. In some ways it is unfortunate not to be able to simulate backscattering and depolarization simultaneously with a single choice of AR. However, we are aware of the fact that the modeling of depolarization is an open and highly challenging problem, one that is not easy and perhaps impossible to solve given the wide variety of shapes 440 that atmospheric ice crystals can take (Liou and Yang, 2016). For this reason we prefer to keep our comparison between optical measurements and calculations limited to the backscattering coefficient.

4.3 **Backscattering versus Particle Size Distribution bulk parameters**

Figures 7, 8, 9, 10 and 11 show N_{ice} , R_{mean} , R_{eff} , SAD and IWC as a function of β , respectively. In the graphs, the black lines represent regression curves, parametrized in the general form $X = A \cdot \beta^B$. The coefficients of the regression are reported in Table 1 together with the R-squared of the fit. The dataset has been fitted between the limits $2 \cdot 10^{-5}$ and $5 \cdot 10^{-3}$ km⁻¹ sr⁻¹, 445 which have been chosen to maximize the goodness of the fit and try to avoid outliers at the extremes of the variability range. We can regard those regression curves as guidance for estimating the microphysics bulk parameters of the clouds and their variability range, from remote measurements of cloud optical parameters.

The linearity between the β and N_{ice} (Fig. 7) is quite striking and indicates that β basically scales with the cloud particle number density N_{ice} , as well as SAD and IWC. Hence, this suggests that the various shapes of the PSD in our observations are 450 hardly changing the scattering properties of the clouds. This finding is further confirmed by the linearity seen between β and SAD or IWC (Figs.10, 11), and the lack of clear correlation between β and R_{mean} or R_{eff} (Figs. 8, 9).

With simple analytical calculations on various types of functional forms for the PSD (gamma, lognormal, etc.), and in the spherical ice approximation, it is easy to demonstrate that a dependence on the square of the modal radius – and hence of other 455 similar parameters linked to it, as the mean or the effective radius - as well as by the total number of particles, is indeed to be expected for β , i.e. $\beta \sim N_0 \cdot R_{mean}^2$, as the physical intuition would also suggest. In our case, such a dependency on R_{mean} , which varies by a factor of 2, is masked by the much wider variability of N_0 , which varies over five orders of magnitude.

4.4 **Depolarization, Aspherical Fraction and PSD parameters**

Depolarization vs Aspherical Fraction 4.4.1

460 In our study we have looked for and found no direct correlation between particle depolarization observed with the backscattersonde, and PSD Aspherical Fraction (AF). AF has peaks around 60% but on the average is maintained around values of 20%, while the corresponding values of the depolarization spans its entire range of variability, giving an unclear relationship between the two quantities (see Fig. A10 in the Supplementary Material). Figure 13 clearly shows again the positive correlation between We remind here that δ_T is not an intensive quantity as δ_{TA}, since it simultaneously depends on the average shape of the
cloud particles and on the backscattering of the whole particle distribution, hence on the particle number concentration, or
SAD, as instance. Hence, for a given and fixed particle shape and dimension, δ_T increases with the BR to a limiting value, which is δ_{TA}. It is the latter the true intensive quantity that depends solely on the particle's average morphology. In fact, Adachi et al. (2001) demonstrated that in a plot of Total Volume Depolarization δ_T towards 1-1/BR, the experimental points of clouds composed of particles sharing the same shape and size but with variable particle number density (i.e. variable BR), will arrange themselves along a straight line, starting at δ_{mol} for BR=1 (this is the case when no particles are present i.e. when the Total Volume Depolarization attains its molecular value δ_{mol} (Young, 1980)) and ending at a δ_{TA} for BR = ∞, this δ_{TA} depending on the particular shape and size common to all particles.

If we were to report data points from clouds composed of particles with different shapes and sizes, these points will be distributed within a triangle whose vertices are δ_{mol} for BR = 1, a δ_{TAmax} and δ_{TAmin} for $BR = \infty$. It should be noted that, if among the possible particle shapes the spherical one is included, then δ_{TAmin} would attain a 0 value.

In Figure 12 we report our dataset, with the data points from in-cloud measurements (i.e. BR>1.2, 1-1/BR > 0.17) color coded in terms of AF. The y axis intercepts at $BR = \infty$ gives us the range of variability of δ_{TA} , which for our cirrus ranges approximately from 20% to 50%.

- We remind again that in this (BR, δ_T) graph, ice clouds composed of particles with the same δ_{TA} and different BR (which we may take as a proxy for particle number concentration) would be distributed along a line starting at δ_{mol} for BR = 1 and and trending towards a particular δ_{TA} when extrapolated to $BR = \infty$. The value of such δ_{TA} depends on the average morphology of the cloud particles. Therefore we may imagine the data point distribution as composed of linear series of points starting at δ_{mol} and spanning the triangle: each line of data points represents the results of the measured variable particle number concentration at constant depolarization properties of the particles as encountered in the clouds.
- Inspecting Figure 12 we may argue that clouds with medium AF (cyan colour) are often linked to intermediate values of δ_{TA} over the whole BR variability range, while clouds with high AF (yellow and red colour), associated with high BR, show no clear correlation with δ_{TA} . Interestingly, clouds with low AF (blue colors) show up at both low BR (with low to intermediate depolarization) and high BR (with intermediate to high depolarization). The conditions at the highest BRs are in fact noteworthy: there, we have both the highest values of the AF for δ_{TAmin} and δ_{TAmax} , and the lowest values of the AF in a range spanning from δ_{TAmed} to δ_{TAmax} .

These relationships between BR, depolarization and AF are probably reflecting aspects of the morphology, size and numerical concentration of the cloud particulate, but are not straightforward to interpret unambiguously. To seek a better understanding, we present in similar plots the dependence of depolarization on N_{ice} and R_{mean} , and on temperature.

4.4.2 Depolarization vs Particle Number Concentration, mean Radius, Temperature

470

490 We report in Figure 13, using the same δ_T - BR framework, the data points color coded in terms of N_{ice} and in Figure 14 the same data points, color coded in terms of R_{mean} . Figure 13 clearly shows again the positive correlation between BR and N_{ice} , independent of the polarization δ_{TA} except at the highest values of BR, where it appears a positive correlation

BR and N_{ice} . This relationship is independent from the polarization δ_{TA} throughout the BR range, except at the highest BR

- values. There, we note how for the same extreme values of BR, low δ_{TA} are associated with low N_{ice} , and high δ_{TA} high N_{ice} . This is possibly because the maximum detectable AF of ice particles decreases with the size of the particles. The reason is that the depolarization signal becomes weaker with decreasing ice particle size and also the ice particles become lesser and lesser aspheric, and the backscattersonde and the cloud spectrometer have depolarization sensitivities that vary differently with the size of the scattering particle.
- In order to display other parameters along with the former two, that may help to find and possibly disentangle a link between these two quantities, we choose to represent our dataset in a δ_T - BR space (in fact, for ease of scale, 1-1/BR is used instead of BR), and color code the points with AF, N, R_{mean} and temperature T as possible parameters of interest.
- We remind here that δ_T is not an intensive quantity as δ_{TA} , since it simultaneously depends on the average shape of the cloud particles and on the backscattering of the whole particle distribution, hence on the particle number concentration, or SAD, as instance. Hence, for a given and fixed particle shape and dimension, δ_T increases with the BR to a limiting value, which is δ_{TA} . It is the latter the true intensive quantity that depends solely on the particle's average morphology. In fact, Adachi et al. (2001) demonstrated that in a plot of δ_T towards 1-1/BR, the experimental points of clouds composed of particles sharing the same shape and size but with variable particle number density (i.e. variable BR), will arrange themselves along a straight line, starting

at δ_{mol} for BR=1 (this is the case when no particles are present i.e. when δ_T attains its molecular value δ_{mol} (Young, 1980))

- and ending at a δ_{TA} for $BR = \infty$, this δ_{TA} depending on the particular shape and size common to all particles. So, in this (BR, δ_T) graph, ice clouds composed of particles with the same δ_{TA} and different BR (which we may take as a proxy for particle number concentration) would be distributed along a line starting at δ_{mol} for BR = 1 and trending towards a particular δ_{TA} when extrapolated to $BR = \infty$. The value of such δ_{TA} depends on the average morphology of the cloud particles. Therefore we may imagine the data point distribution as composed of linear series of points starting at δ_{mol} and spanning the triangle:
- each line of data points represents the results of the measured variable particle number concentration at constant depolarization properties of the particles as encountered in the clouds. Conversely, if we were to report data points from clouds composed of particles with different shapes and sizes, these points will be distributed within a triangle whose vertices are δ_{mol} for BR = 1, a δ_{TAmax} and δ_{TAmin} for $BR = \infty$. It should be noted that, if among the possible particle shapes the spherical one is included, then δ_{TAmin} would attain a 0 value.
- In Fig. 12 we report our dataset, with the data points from in-cloud measurements (i.e. BR>1.2, 1-1/BR > 0.17) color coded in terms of AF. The y axis intercepts at $BR = \infty$ gives us the range of variability of δ_{TA} , which for our cirrus ranges approximately from 20% to 50%. Inspecting the figure we may argue that clouds with medium AF (cyan colour) are often linked to intermediate values of δ_{TA} over the whole BR variability range, while clouds with high AF (yellow and red colour), associated with high BR, show no clear correlation with δ_{TA} . Interestingly, clouds with low AF (blue colors) show up at both
- low BR (with low to intermediate depolarization) and high BR (with intermediate to high depolarization). The conditions at the highest BRs are in fact noteworthy: there, we have both the highest values of the AF for δ_{TAmin} and δ_{TAmax} , and the lowest values of the AF in a range spanning from δ_{TAmed} to δ_{TAmax} .

between δ_{TA} and N_{ice} . Figure 14 shows how high δ_{TA} are often associated with high R_{mean} (red dots along the line $\delta_{mol}-\delta_{TAmax}$), and again peculiarly in the same region of highest values of BR, small R_{mean} coincides with medium to high δ_{TA}). **195** In Figure 15 we show the same dataset this time color coded in terms of temperature. The points at temperatures above 200K are showing both very high and, although in smaller numbers, very low depolarization with intermediate to high BR values. Points below 200K show the general tendency to an increase in depolarization at colder temperature, already commented in figure 2. Noteworthy, the observations in the region of very high BR and between δ_{TAmed} and δ_{TAmax} , where the coldest temperatures, large particle concentration, low AF and low mean radius have been met, are worth of a special mention. These all came from clouds encountered in a single flight, performed on 10 August 2017. On that flight, a convective overshooting updraft of probably liquid origin, with very large aggregates and significant number of freshly nucleated cloud particles was met, together with younger outflow of the overshoot with both growing small particles and sedimenting large ones (Krämer et al., 2020) see their Figure 10b, (Khaykin et al., 2022). This dynamic and therefore particularly variable situation makes the overall interpretation of the observed parameters exceptionally complicated.

505

As stated at the beginning of the paragraph, there seems to be no direct correlation between δ_{TA} and AF and although some clustering of the variables is possible, these clusters often overlaps. It is possible that additional information on particle shape in this analysis, which is missing in the present study (a measured aspect ratio would be an obvious candidate), would help disentangle the variables.

5 Discussion

- 510 As outlined in the previous paragraph, in our study the T-matrix has been used directly to model the scattering properties of particles up to a few hundred size parameters, and indirectly to estimate the depolarization, and the modification of Mie backscattering, for larger particles. We remind that while Mie theory strictly applies to spherical particles, it has no upper size limit and can be applied to the entire cloud particle size range.
- Despite the limitations outlined above, some conclusion can be deduced from our study. It is possible to make the measurements coincide with the optical modeling through a suitable and reasonable choice of the particle AR, and this reassures us about the compatibility of the two datasets. This choice generally favors high ARs (i.e high Mie backscattering depression) in regions of high backscattering, high depolarization and large presence of AF, while ARs near or less than unity are chosen when backscattering is medium to low and the presence of AF is not very pronounced. In general the choice of AR produces a high depression of the backscattering compared to Mie's predictions, and this depression increases with increasing presence of aspherical scatterers, reaching to exceed an order of magnitude when AF is large. In cases with particularly high backscattering, our optical model produces backscattering higher than those observed. If as seems appropriate non-linearity problems in the backscatter probe are excluded, it is quite possible that, to reduce the computed backscattering and reconcile it with

Moreover, it is clear from our simulations that the greatest ambiguity in the results of the comparison is linked to the choice of particles' morphology, rather than to uncertainties in the determination of the PSD or in the measurement of backscattering.

observations, we would need to use ARs beyond the limits considered in our optical model.

These relationships between BR, depolarization and AF are probably reflecting aspects of the morphology, size and numerical concentration of the cloud particulate, but are not straightforward to interpret unambiguously. To seek a better understanding, we present in similar plots the dependence of depolarization on N_{ice} and R_{mean} , and on temperature.

500

4.4.2 Depolarization vs Particle Number Concentration, mean Radius, Temperature

We report in Fig. 13, using the same δ_T - BR framework, the data points color coded in terms of N_{ice} and in Fig. 14 the same data points, color coded in terms of R_{mean} . Figure 13 clearly shows again the positive correlation between BR and N_{ice} , independent of the polarization δ_{TA} except at the highest values of BR, where it appears a positive correlation between δ_{TA} and N_{ice} . Figure 14 shows how high δ_{TA} are often associated with high R_{mean} (red dots along the line δ_{mol} δ_{TAmax}), and again peculiarly in the same region of highest values of BR, small R_{mean} coincides with medium to high δ_{TA} . In Fig. 15 we show the same dataset this time color coded in terms of temperature. The points at temperatures above 200K are showing both very high and, although in smaller numbers, very low depolarization with intermediate to high BR values. Points below 200K show the general tendency to an increase in depolarization at colder temperature, already commented in Fig. 2. Noteworthy, the observations in the region of very high BR and between δ_{TAmed} and δ_{TAmax} , where the coldest temperatures, large particle

- concentration, low AF and low mean radius have been met, are worth of a special mention. These all came from clouds encountered in a single flight, performed on 10 August 2017. On that flight, a convective overshooting updraft of probably liquid origin, with very large aggregates and significant number of freshly nucleated cloud particles was met, together with younger outflow of the overshoot with both growing small particles and sedimenting large ones (Krämer et al. (2020) see their
- 515 Fig. 10b, and Khaykin et al. (2022)). This dynamic and therefore particularly variable situation makes the overall interpretation of the observed parameters exceptionally complicated.

As stated at the beginning of the paragraph, there seems to be no direct correlation between δ_{TA} and AF and although some clustering of the variables is possible, these clusters often overlaps. It is possible that additional information on particle shape in this analysis, which is missing in the present study (a measured aspect ratio would be an obvious candidate), would help disentangle the variables.

5 Discussion

As outlined in the previous paragraph, in our study the T-matrix has been used directly to model the scattering properties of particles up to a few hundred size parameters, and indirectly to estimate the depolarization, and the modification of Mie backscattering, for larger particles. We note that while Mie theory strictly applies to spherical particles, it has no upper size limit and can be applied to the entire cloud particle size range.

525

520

Despite the limitations outlined above, some conclusion can be deduced from our study. It is possible to make the measurements coincide with the optical modeling through a suitable and reasonable choice of the particle AR, and this reassures us about the compatibility of the two datasets. This choice generally favors high ARs (i.e high Mie backscattering depression) in regions of high backscattering, high depolarization and large presence of AF, while ARs near or less than unity are chosen Another interesting result is the identification of the size range of the particulate matter that most contributes to the observed backscattering. For all the PSD under study we have examined the cumulative function (19)

$$\beta_{NC}^{Mie}(\mathbf{r}_i) = \frac{\sum_{i=1}^{r_i} \beta_i^{AR}}{\beta_{NC}^{Mie}} \tag{19}$$

530

535

540

of the respective β as a function of the particle radius, in order to determine the buildup of the final values of the particle backscattering coefficient in relationship to the particle radius. We have used Mie codes to give an upper limit to such computation. The histogram of the values of the cumulative distributions as a function of the particle radius is reported in Figure 16 . It can be seen, from the analysis of the histogram of cumulatives, that the values of the backscattering coefficient are formed mainly in the dimensional range below 300 μ m and particles with radii greater than 400 μ m do not significantly contribute to the β . This analysis gives us confidence in affirming that the size range of the detected particles is sufficient to fully characterize the backscattering coefficient.

The T-matrix computation of depolarization gives no satisfactory results, being the modelled depolarization an underestimation in comparison to the depolarization measured by MAS. It is possible that this is due to the specificity of our approach. Our computations are not able to produce δ_A higher than 40% in the large particle regime (i.e. radius grater than 10 μ m). Other theoretical depolarization ratios computed from a unified theory of light scattering by ice crystals for shapes including bullet rosettes, solid and hollow columns, Koch snowflakes, and plates, extend up to 60% and more (Liu and Mishchenko, 2001). As stated previously there seems to be no clear correlation between particle depolarization and AF. Particle depolarization increases with temperature, with N_{ice} at high backscattering. High depolarization is often, but not always, associated with high R_{mean} as there are cases when at high values of backscattering, small R_{mean} coincides with medium to high δ_{TA} . This study therefore suggest that, although some correlations can be discerned between the depolarization and the environmental condi-

545

tions in which the cloud is observed, or its microphysical and morphological average parameters, there are a lot of exceptions. These probably depend on the history of the formation and the instant of evolution of the cloud, leading to the coexistence of cloud particles with different morphology and sizes. Hence, the study does not allow us to formulate general rules that link depolarization to the microphysics of the cloud.

550

555

The regression of the backscattering coefficient with respect to the various bulk parameters of the cirrus clouds N_{ice} , R_{mean} , R_{eff} , SAD, IWC therefore is based on a solid foundation. Effectively, similar regressions were presented in Cairo et al. (2011) based on cirrus particle measurements from an FSSP-100. There, the upper detection limit of the instrument was 15.5 μ m in radius, while in the present study this limit has extended upward for more than an order of magnitude. So, although the present figures are qualitatively similar to those of Cairo et al. (2011), some difference is to be expected due to the larger range of particle radii that is now covered. The limitedness of the radius range was a caveat already advocated in the aforementioned work. In fact the larger detection threshold is impacting all the regressions, which results in a underestimation in Cairo et al. (2011) of N_{ice} , SAD and VD. In the present work, the detection of a wider particle dimensional range, including the radii which determine most of the measured backscatter coefficient, allows us to place more reliability on the regressions here presented.

The relative independence of β from R_{mean} and R_{eff} confirms N_{ice} as the main parameter governing the cirrus scattering properties at optical wavelengths. This finding would imply that the shapes of the PSD should not play a mayor role and should
when backscattering is medium to low and the presence of AF is not very pronounced. In general the choice of AR produces a high depression of the backscattering compared to Mie's predictions, and this depression increases with increasing presence of aspherical scatterers, reaching to exceed an order of magnitude when AF is large. In cases with particularly high backscattering, our optical model produces backscattering higher than those observed. If - as seems appropriate - non-linearity problems in the backscatter probe are excluded, it is quite possible that, to reduce the computed backscattering and reconcile it with observations, we would need to use ARs beyond the limits considered in our optical model.

Moreover, it is clear from our simulations that the greatest ambiguity in the results of the observed versus the calculated comparison is linked to the choice of particles' morphology, rather than to uncertainties in the determination of the PSD or in the measurement of backscattering.

Another interesting result is the identification of the size range of the particulate matter that most contributes to the observed backscattering. For all the PSD under study we have examined the cumulative function (18)

$$\beta_{NC}^{Mie}(r_j) = \frac{\sum_{i=1}^{r_j} \beta_i^{AR}}{\beta_{NC}^{Mie}} \tag{18}$$

of the respective β as a function of the particle radius, in order to determine the buildup of the final values of the particle backscattering coefficient in relationship to the particle radius. We have used Mie codes to give an upper limit to such computation. The histogram of the values of the cumulative distributions as a function of the particle radius is reported in Fig. 16.

- 545 It can be seen, from the analysis of the histogram of cumulatives, that the values of the backscattering coefficient are formed mainly in the dimensional range below 300 μ m and particles with radii greater than 400 μ m do not significantly contribute to the β . This analysis gives us confidence in affirming that the size range of the detected particles is sufficient to fully characterize the backscattering coefficient.
- The T-matrix computation of depolarization gives no satisfactory results, being the modelled depolarization an underestimation in comparison to the depolarization measured by MAS. It is possible that this is due to the specificity of our approach. Our computations are not able to produce δ_A higher than 40% in the large particle regime (i.e. radius grater than 10 µm). Other theoretical depolarization ratios computed from a unified theory of light scattering by ice crystals for shapes including bullet rosettes, solid and hollow columns, Koch snowflakes, and plates, extend up to 60% and more (Liu and Mishchenko, 2001). As stated previously there seems to be no clear correlation between particle depolarization and AF. Particle depolarization increases with temperature, with N_{ice} at high backscattering. High depolarization is often, but not always, associated with high R_{mean} as there are cases when at high values of backscattering, small R_{mean} coincides with medium to high δ_{TA}. This study therefore suggest that, although some correlations can be discerned between the depolarization and the environmental conditions in which the cloud is observed, or its microphysical and morphological average parameters, there are a lot of exceptions. These probably depend on the history of the formation and the instant of evolution of the cloud, leading to the coexistence of cloud particles with different morphology and sizes. Hence, the study does not allow us to formulate general rules that link

The regression of the backscattering coefficient with respect to the various bulk parameters of the cirrus clouds N_{ice} , R_{mean} , R_{eff} , SAD, IWC therefore is based on a solid foundation. Effectively, similar regressions were presented in Cairo et al. (2011)

depolarization to the microphysics of the cloud.

- all share a similar shape once normalized for the total number of particles, at least for low to medium β values. For high β values, the spreading of the corresponding PSD get larger, suggesting that such observations originated from both clouds of very large and few particles, and of small and more numerous particles. Konoshonkin et al. (2017b) computed backscattering Mueller matrix for the typical shapes of ice crystals of cirrus (hexagonal columns and plates, bullets and droxtals) in the case of their random orientations, for crystal size from 10 µm to 1000 µm with a physical-optics-approximation code and proposed to use the backscatter-to-IWC ratio (as well as extinction-to backscatter, (LR)) for inferring the crystal size in the
- clouds. Figure 17 reports such ratio versus R_{mean} which in fact shows such linear relationship for R_{mean} larger than 10 μ m, while for lower values a different linear behavior might be discerned. In the Figure, the black line is a regression of the form $\frac{beta}{IWC} = 1.52 \cdot 10^{-3} \cdot R_{mean}^{-2.34}$ which has been estimated for R_{mean} from 10 to 100 μ m. The R-squared value of the fit is 0.54, which does not allow us to conclude that such possible linear relationship is based on solid foundations. Because of this, such regression should be used with great caution to estimate R_{mean} when β and IWC are independently available.

Estimates of IWC can be obtained from lidar extinction σ measurements (Heymsfield et al., 2005). Such estimates assume a relationship of the form IWC=A σ^B between the two values, where A and B which in Heymsfield et al. (2005) were posed to A=119 and B=1.22. In a later study Heymsfield et al. (2014) these coefficients have been found to depend on the temperature, based on in situ measurement of IWC and particle PSD obtained during the field campaign of the Stratospheric–Climate Links with Emphasis on the Upper Troposphere and Lower Stratosphere (SCOUT) project, based out of Darwin, Australia, in November–December 2005 (de Reus et al., 2009).

575 I

The IWC estimates from the knowledge of the extinction σ could be compared with our IWC estimates based on β , if a suitable LR is chosen. Unfortunately, LR can vary from 10 to 40 sr in tropical cirrus clouds (Chen et al., 2002) thus making the comparison somewhat arbitrary. Using a LR = 30 sr and posing $\sigma = LR \cdot \beta$ we correlated the IWC measurements with their estimate obtained with the regression towards β presented in this work, and with one obtained from the formulation of Heymsfield et al. (2014). Although both are successful in predicting the IWC within the order of magnitude, the estimate obtained with regression using directly β yelds better results. This is probably due to having chosen the same LR value for all the clouds observed. It should also be noted that the σ -based estimation improves by using the formulation with coefficients that do not depend on the temperature, as in Heymsfield et al. (2005), see figure A9 in Appendix.

- To conclude, a comment on the representativeness of our study. The range of backscattering values observed during campaign activities is wide and covers the variability of possible lidar observations from satellite (Balmes et al., 2019). Regarding the type of cirrus clouds observed, during the campaign activities both in-situ and liquid-origin were sampled, but the second type is dominating since most of the observations came from penetration into the outflow regions of deep convective clouds. As there might be differences in the microphysical properties of the cirrus depending on the formation process (Krämer et al.,
- 590 2020) especially in the initial stage of their life cycle, this may induce an unquantified bias in our presented statistics.

based on cirrus particle measurements from an FSSP-100. There, the upper detection limit of the instrument was 15.5 μ m in

- radius, while in the present study this limit has extended upward for more than an order of magnitude. So, although the present 565 figures are qualitatively similar to those of Cairo et al. (2011), some difference is to be expected due to the larger range of particle radii that is now covered. The limitedness of the radius range was a caveat already advocated in the aforementioned work. In fact the larger detection threshold is impacting all the regressions, which results in a underestimation in Cairo et al. (2011) of N_{ice} , SAD and VD. In the present work, the detection of a wider particle dimensional range, including the radii which
- determine most of the measured backscatter coefficient, allows us to place more reliability on the regressions here presented. 570 The relative independence of β from R_{mean} and R_{eff} confirms N_{ice} as the main parameter governing the cirrus scattering properties at optical wavelengths. This finding would imply that the shapes of the PSD should not play a mayor role and should all share a similar shape once normalized for the total number of particles, at least for low to medium β values. For high β values, the spreading of the corresponding PSD get larger, suggesting that such observations originated from both clouds of
- very large and few particles, and of small and more numerous particles. Konoshonkin et al. (2017b) computed backscattering 575 Mueller matrix for the typical shapes of ice crystals of cirrus (hexagonal columns and plates, bullets and droxtals) in the case of their random orientations, for crystal size from 10 μ m to 1000 μ m with a physical-optics-approximation code and proposed to use the backscatter-to-IWC ratio (as well as extinction-to backscatter, (LR)) for inferring the crystal size in the clouds. Figure 17 reports such ratio versus R_{mean} which in fact shows such linear relationship for R_{mean} larger than 10 580
- μ m, while for lower values a different linear behavior might be discerned. In Fig. 17 the black line is a regression of the form $\frac{beta}{IWC} = 1.52 \cdot 10^{-3} \cdot R_{mean}^{-2.34}$ which has been estimated for R_{mean} from 10 to 100 μ m. The R-squared value of the fit is 0.54, which does not allow us to conclude that such possible relationship is based on solid foundations. Because of this, such

regression should be used with great caution to estimate R_{mean} when β and IWC are independently available.

- Several studies have provided an estimate of the dependence of the IWC on lidar extinction (Heymsfield et al., 2005; Avery et al., 2012; Heymsfield et al., 2014; Thornberry et al., 2017). They are based on in situ measurements of IWC and PSD, the 585 latter used to provide an estimate of the lidar extinction from optical modeling of the cloud particles. These IWC- σ relationships could be compared with our IWC estimates based on β , if a suitable extinction to backscatter ratio (a.k.a. Lidar Ratio) LR is chosen. Unfortunately, LR can vary from 10 to 40 sr in tropical cirrus clouds (Chen et al., 2002) thus making the comparison somewhat arbitrary. Using a LR = 30 sr as a most probable value Balmes et al. (2019) and posing $\sigma = LR^*\beta$ we can correlate our IWC measurements with σ Figure 18 is thus the analogue of Fig. 11, where this time IWC is reparametrized as a function 590
- of σ . The same figure shows the analytical relationship obtained in this work (solid black line), with those present in the literature, shown in Table 2. Although all parameterizations capture the IWC- σ trend and align with each other in the lower range of data variability, the result of our study is in agreement only with Avery et al. (2012) while it diverges from the other parameterizations, more severely for those that depend on the temperature. This especially in the upper range of data variability.
- 595 It should be noted that in this range, the data themselves have also a greater dispersion. We want to underline the limits of this comparison: in the case of the present study, they are caused by having chosen, rather arbitrarily, the same LR value for all the clouds observed, and in the case of the other parametrizations they are caused by having used an indirect determination of the

6 Conclusions

595

600

Measurements in cirrus clouds obtained during the StratoClim campaign by two dual-instrument cloud spectrometers, two hygrometers and a backscattersonde have been compared. The comparison of the microphysical data with the optical observations of the backscattersonde was performed by calculating the ice particle backscattering coefficient from the PSD by means of optical modelling. A proper adjustment of the modeled particle AR allows to match the optical computation with the backscattering observations. Relations have been obtained to link the ice particles backscatter coefficient to their concentration, their mean and effective radius, surface area density and ice water content. These results confirm and expand similar studies and allow to estimate within an order of magnitude, data of bulk cirrus microphysics from lidar remote sensing observations.

The comparison between particle depolarization from the backscattersonde and the Aspherical Fraction measured by one of the cloud spectrometers shows no univocal relationship between the two quantities and ask for further investigations, where additional information on particle morphology may be required.

Code availability. TEXT

Data availability. TEXT

Code and data availability. Water measurements (FLASH and FISH), meteorological data from UCSE and TDC, MAS backscatter and particle measurements from NIXE-CAPS from the StratoClim aircraft campaign are available on the HALO database at https://halodb.pa.op.dlr.de/mission/101 (DLR, 2022). Particle measurements from CCP are available upon request to the PIs. Lidar measurement from MAL are available upon request to the PI. Analysis and plotting scripts for this paper are available upon request to the corresponding author.

Sample availability. TEXT

Video supplement. TEXT

610 *Author contributions.* MS, LDL, FC prepared the MAS for campaign activity. FC performed and processed the MAS measurements. LL did the analysis scripts for data comparison. SK performed and processed the FLASH measurements. CR, AA, NS and MK performed and processed the FISH and NIXE-CAPS measurements. SB, MP performed and processed the CCP measurements. RW contributed to the

extinction, calculated from the PSDs. It would be very interesting to have simultaneous in situ observations of backscattering, extinction and IWC available in the future.

- To conclude, a comment on the representativeness of our study. The range of backscattering values observed during campaign activities is wide and covers the variability of possible lidar observations from satellite (Balmes et al., 2019). Regarding the type of cirrus clouds observed, during the campaign activities both in-situ and liquid-origin were sampled, but the second type is dominating since most of the observations came from penetration into the outflow regions of deep convective clouds. As there might be differences in the microphysical properties of the cirrus depending on the formation process (Lawson et al., 2019;
- 605 Krämer et al., 2020) especially in the initial stage of their life cycle, the results presented here are not necessarily representative of all cirrus that can be observed from space.

6 Conclusions

Measurements in cirrus clouds obtained during the StratoClim campaign by two dual-instrument cloud spectrometers, two hygrometers and a backscattersonde have been compared. The comparison of the microphysical data with the optical observations of the backscattersonde was performed by calculating the cirrus particle backscattering coefficient from the PSD by

- 610 vations of the backscattersonde was performed by calculating the cirrus particle backscattering coefficient from the PSD by means of optical modelling. A proper adjustment of the modeled particle AR allows to match the optical computation with the backscattering observations. Relations have been obtained to link the ice particles backscatter coefficient to their concentration, their mean and effective radius, surface area density and ice water content. These results confirm and expand similar studies and allow to estimate within an order of magnitude, data of bulk cirrus microphysics from lidar remote sensing observations.
- 615 The comparison between particle depolarization from the backscattersonde and the Aspherical Fraction measured by one of the cloud spectrometers shows no univocal relationship between the two quantities and ask for further investigations, where additional information on particle morphology may be required.

Code availability. TEXT

Data availability. TEXT

620 Code and data availability. Water measurements (FLASH and FISH), meteorological data from UCSE and TDC, MAS backscatter and particle measurements from NIXE-CAPS from the StratoClim aircraft campaign are available on the HALO database at https://halo-db.pa.op.dlr.de/mission/101 (DLR, 2022). Particle measurements from CCP are available upon request to the PIs. Lidar measurement from MAL are available upon request to the PI. Analysis and plotting scripts for this paper are available upon request to the corresponding author.

collection of the particle dataset. SB and FC conceived and planned the study. FC performed the analysis, prepared the figures, and wrote the manuscript. All authors commented on the manuscript.

615 Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer.

620

640

Acknowledgements. We gratefully thank the StratoClim coordination team and the Myasishchev Design Bureau for successfully conducting the field campaign. This research has been supported by the StratoClim project of the European Community's Seventh Framework Programme (FP7/2007–2013) under grant agreement no. 603557 and internal funds of the Max Planck institute for Chemistry. FC is grateful to the MPI for Chemistry for supporting his sabbatical stay in Mainz during which the study behind this publication was initiated. SK's work was partly supported by the Agence Nationale de la Recherche TTL-Xing ANR-17-CE01-0015 project.

Appendix A: SUPPLEMENTARY MATERIAL

- We include in this Appendix in figure A1 and A2 the particle backscattering coefficient and depolarization vs particle radius, parametrized with the particle AR. These optical parameters have been computed for a reference monodisperse PSD with 625 particle concentration of 1 cm⁻³. In figure A3 the 2D histogram of the particle concentration N_{ice} computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). A4 reports the 2D histogram of the particle mean radius R_{mean} computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). In both figure, for NIXE-CAPS only particles with with lower size limit at 1.5 μ m have been considered. A5 the comparison of the backscattering coefficients retrieved from the PSD measured by the two cloud spectrometers. In A6 results of the backscattering coefficient β measured by MAS on 10 August 2017 (red lines); Black solid line, β_{NC} corresponding to the best matching between measured δ and computed δ_{NC}^{AR} (this latter in not displayed); error bars are reported in the first part of the curve; black dashed lines, maximum and minimum values of the optical modeling β_{NC}^{AR} values. In the lower panel: Red line , particle depolarization measured by MAS; Blue line Particle Aspherical Faction measured by NIXE-CAPS; black line, AF values of the best matching between measured δ and computed δ_{NC}^{AR} .
- In figure A7 we report 2D histogram of the backscattering coefficient computed with PSD with lover size limit at 0.3 μ m (y-axis) and at 1.5 μ m (x-axis).

In figure A8 we report 2D histogram of Particle AF (y-axis) vs Particle depolarization (x-axis).

Finally, in figure A9 scatterplots of estimated vs measured IWC using different regressions are reported. In the left panel the regression from the present work is presented. This uses the measured β to estimate the IWC. In the middle and left panels, the regressions from Heymsfield et al. (2014) (middle panel) and from Heymsfield et al. (2005) (right panel) are reported.

625 Video supplement. TEXT

Author contributions. MS, LDL, FC, GDD prepared the MAS for campaign activity. FC performed and processed the MAS measurements. LL did the analysis scripts for data comparison. SK performed and processed the FLASH measurements. CR, AA, NS and MK performed and processed the FISH and NIXE-CAPS measurements. SB, MP performed and processed the CCP measurements. RW contributed to the collection of the particle dataset. SB and FC conceived and planned the study. FC performed the analysis, prepared the figures, and drafted the manuscript. All authors commented on the manuscript.

630

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer.

Acknowledgements. We gratefully thank the StratoClim coordination team and the Myasishchev Design Bureau for successfully conducting the field campaign. This research has been supported by the StratoClim project of the European Community's Seventh Framework Pro 635 gramme (FP7/2007–2013) under grant agreement no. 603557 and internal funds of the Max Planck institute for Chemistry. FC is grateful to the MPI for Chemistry for supporting his sabbatical stay in Mainz during which the study behind this publication was initiated. SK's work was partly supported by the Agence Nationale de la Recherche TTL-Xing ANR-17-CE01-0015 project.

Appendix A: SUPPLEMENTARY MATERIAL

We include in this Supplementary Material in Fig. A1 and A2 the particle backscattering coefficient and depolarization vs
particle radius, parametrized with the particle AR. These optical parameters have been computed for a reference monodisperse
PSD with particle concentration of 1 cm⁻³. In Fig. A3 the comparison of the backscattering coefficient computed with SD with
lower size limit at 0.3 (y-axis) and 1.5 (x-axis) µm. Mie codes have been used in teh comutation. In Fig. A4 and A5 the 2D
histogram of the particle concentration N_{ice} and particle mean radius R_{mean} computed from PSD from NIXE-CAPS (x-axis)
and CCP (y-axis). Figure A6 reports particle backscattering coefficient computation from NIXE-CAPS (x-axis) and CCP (y-axis), with Mie scattering codes. The Figure A7 reports the scatterplot of the Total Particle Depolarization vs the Aspect Ratio,
colour coded in terms of the backscatter coefficient. In A9 results of the backscattering coefficient β measured by MAS on

These two have been obtained by using the measured β to estimate the extinction σ , originally used in Heymsfield's works, and assuming a LR=30 sr. The colors code the temperature at the observation.

10 August 2017 (red lines); Black solid line, β_{NC} corresponding to the best matching between measured δ and computed δ_{NC}^{AR} (this latter in not displayed); error bars are reported in the first part of the curve; black dashed lines, maximum and minimum

650 values of the optical modeling β_{NC}^{AR} values. In the lower panel: Red line , particle depolarization measured by MAS; Blue line Particle Aspherical Faction measured by NIXE-CAPS; black line, AF values of the best matching between measured δ and computed δ_{NC}^{AR} .

In figure A10 we report 2D histogram of the backscattering coefficient computed with PSD with lover size limit at 0.3 μ m (y-axis) and at 1.5 μ m (x-axis).

655 In figure A10 we report 2D histogram of particle AF (y-axis) vs particle depolarization (x-axis).

References

650

Adachi, H., Shibata, T., Iwasaka, Y., and Fujiwara, M.: Calibration method for the lidar-observed stratospheric depolarization ratio in the

- 645 presence of liquid aerosol particles, Appl. Opt., 40, 6587–6595, https://doi.org/10.1364/AO.40.006587, http://www.osapublishing.org/ao/ abstract.cfm?URI=ao-40-36-6587, 2001.
 - Afchine, A., Rolf, C., Costa, A., Spelten, N., Riese, M., Buchholz, B., Ebert, V., Heller, R., Kaufmann, S., Minikin, A., Voigt, C., Zöger, M., Smith, J., Lawson, P., Lykov, A., Khaykin, S., and Krämer, M.: Ice particle sampling from aircraft influence of the probing position on the ice water content, Atmospheric Measurement Techniques, 11, 4015–4031, https://doi.org/10.5194/amt-11-4015-2018, https://amt. copernicus.org/articles/11/4015/2018/, 2018.
 - Austin, R. T., Heymsfield, A. J., and Stephens, G. L.: Retrieval of ice cloud microphysical parameters using the CloudSat millimeterwave radar and temperature, Journal of Geophysical Research: Atmospheres, 114, https://doi.org/https://doi.org/10.1029/2008JD010049, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008JD010049, 2009.
 - Bailey, M. P. and Hallett, J.: A Comprehensive Habit Diagram for Atmospheric Ice Crystals: Confirmation from the Laboratory, AIRS II,
- and Other Field Studies, Journal of the Atmospheric Sciences, 66, 2888 2899, https://doi.org/10.1175/2009JAS2883.1, https://journals. ametsoc.org/view/journals/atsc/66/9/2009jas2883.1.xml, 2009.
 - Balmes, K. A., Fu, Q., and Thorsen, T. J.: Differences in Ice Cloud Optical Depth From CALIPSO and Ground-Based Raman Lidar at the ARM SGP and TWP Sites, Journal of Geophysical Research: Atmospheres, 124, 1755–1778, https://doi.org/10.1029/2018JD028321, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD028321, 2019.
- 660 Baran, A. J., Francis, P. N., Havemann, S., and Yang, P.: A study of the absorption and extinction properties of hexagonal ice columns and plates in random and preferred orientation, using exact T-matrix theory and aircraft observations of cirrus, Journal of Quantitative Spectroscopy and Radiative Transfer, 70, 505–518, https://doi.org/https://doi.org/10.1016/S0022-4073(01)00025-5, https://www.sciencedirect. com/science/article/pii/S0022407301000255, light Scattering by Non-Spherical Particles, 2001.
- Baumgardner, D., Abel, S. J., Axisa, D., Cotton, R., Crosier, J., Field, P., Gurganus, C., Heymsfield, A., Korolev, A., Krämer, M., Lawson, P., McFarquhar, G., Ulanowski, Z., and Um, J.: Cloud Ice Properties: In Situ Measurement Challenges, Meteorological Monographs, 58, 9.1 9.23, https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0011.1, https://journals.ametsoc.org/view/journals/amsm/58/1/amsmonographs-d-16-0011.1.xml, 2017.
 - Berendsen, H. J. C.: A Student's Guide to Data and Error Analysis, Student's Guides, Cambridge University Press, https://doi.org/10.1017/CBO9780511921247, 2011.
- 670 Bi, L. and Yang, P.: Accurate simulation of the optical properties of atmospheric ice crystals with the invariant imbedding T-matrix method, jqsrt, 138, 17–35, https://doi.org/10.1016/j.jqsrt.2014.01.013, 2014.

Bohren, C. F. and Huffman, D. R.: Absorption and scattering of light by small particles, John Wiley & Sons, 2008.

- Borrmann, S. and Luo, B.: Application of the T-matrix method to the measurement of aspherical (ellipsoidal) particles with forward scattering optical particle counters, Journal of Aerosol Science, 31, 789–799, https://doi.org/10.1016/S0021-8502(99)00563-7, 2000.
- 675 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and aerosols, pp. 571–657, Cambridge University Press, Cambridge, UK, https://doi.org/10.1017/CBO9781107415324.016, 2013.
 - Cairo, F., Donfrancesco, G. D., Adriani, A., Pulvirenti, L., and Fierli, F.: Comparison of various linear depolarization parameters measured by lidar, Appl. Opt., 38, 4425–4432, https://doi.org/10.1364/AO.38.004425, http://ao.osa.org/abstract.cfm?URI=ao-38-21-4425, 1999.

References

- Adachi, H., Shibata, T., Iwasaka, Y., and Fujiwara, M.: Calibration method for the lidar-observed stratospheric depolarization ratio in the presence of liquid aerosol particles, Appl. Opt., 40, 6587–6595, https://doi.org/10.1364/AO.40.006587, http://www.osapublishing.org/ao/ abstract.cfm?URI=ao-40-36-6587, 2001.
- 660
 - Afchine, A., Rolf, C., Costa, A., Spelten, N., Riese, M., Buchholz, B., Ebert, V., Heller, R., Kaufmann, S., Minikin, A., Voigt, C., Zöger, M., Smith, J., Lawson, P., Lykov, A., Khavkin, S., and Krämer, M.: Ice particle sampling from aircraft – influence of the probing position on the ice water content, Atmospheric Measurement Techniques, 11, 4015-4031, https://doi.org/10.5194/amt-11-4015-2018, https://amt. copernicus.org/articles/11/4015/2018/, 2018.

665 AMS: Glossary of meteorology, 2023.

- Austin, R. T., Heymsfield, A. J., and Stephens, G. L.: Retrieval of ice cloud microphysical parameters using the CloudSat millimeterwave radar and temperature, Journal of Geophysical Research: Atmospheres, 114, https://doi.org/https://doi.org/10.1029/2008JD010049, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008JD010049, 2009.
- Avery, M., Winker, D., Heymsfield, A., Vaughan, M., Young, S., Hu, Y., and Trepte, C.: Cloud ice water content retrieved from the CALIOP 670 space-based lidar, Geophysical Research Letters, 39, 2012.
 - Bailey, M. P. and Hallett, J.: A Comprehensive Habit Diagram for Atmospheric Ice Crystals: Confirmation from the Laboratory, AIRS II, and Other Field Studies, Journal of the Atmospheric Sciences, 66, 2888 - 2899, https://doi.org/10.1175/2009JAS2883.1, https://journals. ametsoc.org/view/journals/atsc/66/9/2009jas2883.1.xml, 2009.
- Balmes, K. A., Fu, O., and Thorsen, T. J.: Differences in Ice Cloud Optical Depth From CALIPSO and Ground-675 Based Raman Lidar at the ARM SGP and TWP Sites, Journal of Geophysical Research: Atmospheres, 124, 1755–1778, https://doi.org/https://doi.org/10.1029/2018JD028321, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD028321, 2019.
- Baran, A. J., Francis, P. N., Havemann, S., and Yang, P.: A study of the absorption and extinction properties of hexagonal ice columns and plates in random and preferred orientation, using exact T-matrix theory and aircraft observations of cirrus, Journal of Quantitative Spectroscopy and Radiative Transfer, 70, 505-518, https://doi.org/https://doi.org/10.1016/S0022-4073(01)00025-5, https://www.sciencedirect. 680 com/science/article/pii/S0022407301000255, light Scattering by Non-Spherical Particles, 2001.
 - Baumgardner, D., Abel, S. J., Axisa, D., Cotton, R., Crosier, J., Field, P., Gurganus, C., Heymsfield, A., Korolev, A., Krämer, M., Lawson, P., McFarquhar, G., Ulanowski, Z., and Um, J.: Cloud Ice Properties: In Situ Measurement Challenges, Meteorological Monographs, 58, 9.1 - 9.23, https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0011.1, https://journals.ametsoc.org/view/journals/amsm/58/ 1/amsmonographs-d-16-0011.1.xml, 2017.
- 685 Berendsen, H. J. C.: A Student's Guide to Data and Error Analysis, Student's Guides, Cambridge University Press, https://doi.org/10.1017/CBO9780511921247, 2011.
 - Bi, L. and Yang, P.: Accurate simulation of the optical properties of atmospheric ice crystals with the invariant imbedding T-matrix method, jgsrt, 138, 17-35, https://doi.org/10.1016/j.jgsrt.2014.01.013, 2014.

Bohren, C. F. and Huffman, D. R.: Absorption and scattering of light by small particles, John Wiley & Sons, 2008.

690 Borrmann, S. and Luo, B.: Application of the T-matrix method to the measurement of aspherical (ellipsoidal) particles with forward scattering optical particle counters, Journal of Aerosol Science, 31, 789-799, https://doi.org/10.1016/S0021-8502(99)00563-7, 2000.

- Cairo, F., Adriani, A., Viterbini, M., Di Donfrancesco, G., Mitev, V., Matthey, R., Bastiano, M., Redaelli, G., Dragani, R., Ferretti, R., Rizi, V., Paolucci, T., Bernardini, L., Cacciani, M., Pace, G., and Fiocco, G.: Polar stratospheric clouds observed during the Airborne Polar Experiment–Geophysica Aircraft in Antarctica (APE-GAIA) campaign, Journal of Geophysical Research: Atmospheres, 109, https://doi.org/https://doi.org/10.1029/2003JD003930, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2003JD003930, 2004.
- Cairo, F., Di Donfrancesco, G., Snels, M., Fierli, F., Viterbini, M., Borrmann, S., and Frey, W.: A comparison of light backscattering and particle size distribution measurements in tropical cirrus clouds, Atmospheric Measurement Techniques, 4, 557–570, https://doi.org/10.5194/amt-4-557-2011, https://amt.copernicus.org/articles/4/557/2011, 2011.
 - Cairo, F., De Muro, M., Snels, M., Di Liberto, L., Bucci, S., Legras, B., Kottayil, A., Scoccione, A., and Ghisu, S.: Lidar observations of cirrus clouds in Palau (7°33' N, 134°48' E), Atmospheric Chemistry and Physics, 21, 7947–7961, https://doi.org/10.5194/acp-21-7947-2021, https://acp.copernicus.org/articles/21/7947/2021/, 2021.
- 690 Cairo, F., Deshler, T., Di Liberto, L., Scoccione, A., and Snels, M.: A study of optical scattering modelling for mixed phase Polar Stratospheric Clouds, EGUsphere, 2022, 1–21, https://doi.org/10.5194/egusphere-2022-972, https://egusphere.copernicus.org/preprints/
 - Campbell, J. R., Lolli, S., Lewis, J. R., Gu, Y., and Welton, E. J.: Daytime cirrus cloud top-of-atmosphere radiative forcing properties at a midlatitude site and their global consequence, J. Appl. Meteor. Climatol., 55, 1667–1679, 2016.
- 695 Chen, W.-N., Chiang, C.-W., and Nee, J.-B.: Lidar ratio and depolarization ratio for cirrus clouds, Appl. Opt., 41, 6470–6476, https://doi.org/10.1364/AO.41.006470, http://ao.osa.org/abstract.cfm?URI=ao-41-30-6470, 2002.
 - Collis, R. and Russell, P.: Lidar measurement of particles and gases by elastic backscattering and differential absorption, in: Laser monitoring of the atmosphere, pp. 71–151, Springer, 1976.
- Corti, T., Luo, B., De Reus, M., Brunner, D., Cairo, F., Mahoney, M., Martucci, G., Matthey, R., Mitev, V., Dos Santos, F., et al.: Unprecedented evidence for deep convection hydrating the tropical stratosphere, Geophysical Research Letters, 35, 2008.
- Costa, A., Meyer, J., Afchine, A., Luebke, A., Günther, G., Dorsey, J. R., Gallagher, M. W., Ehrlich, A., Wendisch, M., Baumgardner, D., Wex, H., and Krämer, M.: Classification of Arctic, midlatitude and tropical clouds in the mixed-phase temperature regime, Atmospheric Chemistry and Physics, 17, 12219–12238, https://doi.org/10.5194/acp-17-12219-2017, https://acp.copernicus.org/articles/17/12219/2017/, 2017.
- de Reus, M., Borrmann, S., Bansemer, A., Heymsfield, A. J., Weigel, R., Schiller, C., Mitev, V., Frey, W., Kunkel, D., Kürten, A., Curtius, J., Sitnikov, N. M., Ulanovsky, A., and Ravegnani, F.: Evidence for ice particles in the tropical stratosphere from in-situ measurements, Atmospheric Chemistry and Physics, 9, 6775–6792, https://doi.org/10.5194/acp-9-6775-2009, https://acp.copernicus.org/articles/9/6775/2009/, 2009.
- Delanoë, J. and Hogan, R. J.: Combined CloudSat-CALIPSO-MODIS retrievals of the properties of ice clouds, Journal of Geophysical Research: Atmospheres, 115, https://doi.org/https://doi.org/10.1029/2009JD012346, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.
 - Research: Atmospheres, 115, https://doi.org/https://doi.org/10.1029/2009JD012346, https://agupubs.onlinelibrary.wiley.com/doi/abs/10. 1029/2009JD012346, 2010.
 - Deshler, T., Nardi, B., Adriani, A., Cairo, F., Hansen, G., Fierli, F., Hauchecorne, A., and Pulvirenti, L.: Determining the index of refraction of polar stratospheric clouds above Andoya (69 N) by combining size-resolved concentration and optical scattering measurements, Journal of Geophysical Research: Atmospheres, 105, 3943–3953, 2000.
- 715 Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B. N., Mishchenko, M., Yang, P., Eck, T. F., Volten, H., Muñoz, O., Veihelmann, B., van der Zande, W. J., Leon, J.-F., Sorokin, M., and Slutsker, I.: Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust, J. Geophys. Res., 111, D11 208, https://doi.org/10.1029/2005JD006619, 2006.

- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S. K., Sherwood, S., Stevens, B., and Zhang, X. Y.: Clouds and aerosols, pp. 571–657, Cambridge University Press, Cambridge, UK, https://doi.org/10.1017/CBO9781107415324.016, 2013.
- 695 Cairo, F., Donfrancesco, G. D., Adriani, A., Pulvirenti, L., and Fierli, F.: Comparison of various linear depolarization parameters measured by lidar, Appl. Opt., 38, 4425–4432, https://doi.org/10.1364/AO.38.004425, http://ao.osa.org/abstract.cfm?URI=ao-38-21-4425, 1999.
 - Cairo, F., Adriani, A., Viterbini, M., Di Donfrancesco, G., Mitev, V., Matthey, R., Bastiano, M., Redaelli, G., Dragani, R., Ferretti, R., Rizi, V., Paolucci, T., Bernardini, L., Cacciani, M., Pace, G., and Fiocco, G.: Polar stratospheric clouds observed during the Airborne Polar Experiment–Geophysica Aircraft in Antarctica (APE-GAIA) campaign, Journal of Geophysical Research: Atmospheres, 109, https://doi.org/10.1029/2003JD003930, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2003JD003930, 2004.
 - Cairo, F., Di Donfrancesco, G., Snels, M., Fierli, F., Viterbini, M., Borrmann, S., and Frey, W.: A comparison of light backscattering and particle size distribution measurements in tropical cirrus clouds, Atmospheric Measurement Techniques, 4, 557–570, https://doi.org/10.5194/amt-4-557-2011, https://amt.copernicus.org/articles/4/557/2011/, 2011.
 - Cairo, F., De Muro, M., Snels, M., Di Liberto, L., Bucci, S., Legras, B., Kottayil, A., Scoccione, A., and Ghisu, S.: Lidar observations of
- cirrus clouds in Palau (7°33' N, 134°48' E), Atmospheric Chemistry and Physics, 21, 7947–7961, https://doi.org/10.5194/acp-21-7947-2021, https://acp.copernicus.org/articles/21/7947/2021/, 2021.
 - Cairo, F., Deshler, T., Di Liberto, L., Scoccione, A., and Snels, M.: A study of optical scattering modelling for mixed-phase polar stratospheric clouds, Atmospheric Measurement Techniques, 16, 419–431, https://doi.org/10.5194/amt-16-419-2023, https://amt.copernicus. org/articles/16/419/2023/, 2023.
- 710 Campbell, J. R., Lolli, S., Lewis, J. R., Gu, Y., and Welton, E. J.: Daytime cirrus cloud top-of-atmosphere radiative forcing properties at a midlatitude site and their global consequence, J. Appl. Meteor. Climatol., 55, 1667–1679, 2016.
 - Chen, W.-N., Chiang, C.-W., and Nee, J.-B.: Lidar ratio and depolarization ratio for cirrus clouds, Appl. Opt., 41, 6470–6476, https://doi.org/10.1364/AO.41.006470, http://ao.osa.org/abstract.cfm?URI=ao-41-30-6470, 2002.

Collis, R. and Russell, P.: Lidar measurement of particles and gases by elastic backscattering and differential absorption, in: Laser monitoring

715 of the atmosphere, pp. 71–151, Springer, 1976.

- Corti, T., Luo, B., De Reus, M., Brunner, D., Cairo, F., Mahoney, M., Martucci, G., Matthey, R., Mitev, V., Dos Santos, F., et al.: Unprecedented evidence for deep convection hydrating the tropical stratosphere, Geophysical Research Letters, 35, 2008.
- Costa, A., Meyer, J., Afchine, A., Luebke, A., Günther, G., Dorsey, J. R., Gallagher, M. W., Ehrlich, A., Wendisch, M., Baumgardner, D., Wex, H., and Krämer, M.: Classification of Arctic, midlatitude and tropical clouds in the mixed-phase temperature regime, Atmospheric Chem-
- 720 istry and Physics, 17, 12219–12238, https://doi.org/10.5194/acp-17-12219-2017, https://acp.copernicus.org/articles/17/12219/2017/, 2017.
 - Delanoë, J. and Hogan, R. J.: Combined CloudSat-CALIPSO-MODIS retrievals of the properties of ice clouds, Journal of Geophysical Research: Atmospheres, 115, https://doi.org/https://doi.org/10.1029/2009JD012346, https://agupubs.onlinelibrary.wiley.com/doi/abs/10. 1029/2009JD012346, 2010.
- 725 Deshler, T., Nardi, B., Adriani, A., Cairo, F., Hansen, G., Fierli, F., Hauchecorne, A., and Pulvirenti, L.: Determining the index of refraction of polar stratospheric clouds above Andoya (69 N) by combining size-resolved concentration and optical scattering measurements, Journal of Geophysical Research: Atmospheres, 105, 3943–3953, 2000.

Dubovik, O., Lapyonok, T., Litvinov, P., Herman, M., Fuertes, D., Ducos, F., Torres, B., Derimian, Y., Huang, X., Lopatin, A., Chaikovsky, A., Aspetsberger, M., and Federspiel, C.: GRASP: a versatile algorithm for characterizing the atmosphere, SPIE Newsroom, https://doi.org/10.1117/2.1201408.005558, http://spie.org/x109993.xml, 2014.

720

725

- Frey, W., Borrmann, S., Kunkel, D., Weigel, R., de Reus, M., Schlager, H., Roiger, A., Voigt, C., Hoor, P., Curtius, J., Krämer, M., Schiller, C., Volk, C. M., Homan, C. D., Fierli, F., Di Donfrancesco, G., Ulanovsky, A., Ravegnani, F., Sitnikov, N. M., Viciani, S., D'Amato, F., Shur, G. N., Belyaev, G. V., Law, K. S., and Cairo, F.: In situ measurements of tropical cloud properties in the West African Monsoon: upper tropospheric ice clouds, Mesoscale Convective System outflow, and subvisual cirrus, Atmospheric Chemistry and Physics, 11, 5569–5590, https://doi.org/10.5194/acp-11-5569-2011, https://acp.copernicus.org/articles/11/5569/2011/, 2011.
- Frey, W., Borrmann, S., Fierli, F., Weigel, R., Mitev, V., Matthey, R., Ravegnani, F., Sitnikov, N. M., Ulanovsky, A., and Cairo, F.: Tropical deep convective life cycle: Cb-anvil cloud microphysics from high-altitude aircraft observations, Atmospheric Chemistry and Physics, 14, 13 223–13 240, https://doi.org/10.5194/acp-14-13223-2014, https://acp.copernicus.org/articles/14/13223/2014/, 2014.
 - Fu, Q.: A New Parameterization of an Asymmetry Factor of Cirrus Clouds for Climate Models, Journal of the Atmospheric Sciences, 64, 4140 4150, https://doi.org/10.1175/2007JAS2289.1, https://journals.ametsoc.org/view/journals/atsc/64/11/2007jas2289.1.xml, 2007.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, Reviews of Geophysics, 47, https://doi.org/10.1029/2008RG000267, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008RG000267, 2009.
 - Fujiwara, M., Iwasaki, S., Shimizu, A., Inai, Y., Shiotani, M., Hasebe, F., Matsui, I., Sugimoto, N., Okamoto, H., Nishi, N., Hamada, A., Sakazaki, T., and Yoneyama, K.: Cirrus observations in the tropical tropopause layer over the western Pacific, Journal of Geo-
- 735 physical Research: Atmospheres, 114, https://doi.org/10.1029/2008JD011040, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2008JD011040, 2009.
 - Groth, S. P., Baran, A. J., Betcke, T., Havemann, S., and Śmigaj, W.: The boundary element method for light scattering by ice crystals and its implementation in BEM++, Journal of Quantitative Spectroscopy and Radiative Transfer, 167, 40–52, 2015.
- Grulich, L., Weigel, R., Hildebrandt, A., Wand, M., and Spichtinger, P.: Automatic shape detection of ice crystals, Journal of Computa tional Science, 54, 101 429, https://doi.org/https://doi.org/10.1016/j.jocs.2021.101429, https://www.sciencedirect.com/science/article/pii/
 \$1877750321001137, 2021.
 - Guignard, A., Stubenrauch, C. J., Baran, A. J., and Armante, R.: Bulk microphysical properties of semi-transparent cirrus from AIRS: a six year global climatology and statistical analysis in synergy with geometrical profiling data from CloudSat-CALIPSO, Atmospheric Chemistry and Physics, 12, 503–525, https://doi.org/10.5194/acp-12-503-2012, https://acp.copernicus.org/articles/12/503/2012/, 2012.
- 745 He, Q., LI, C., Ma, J., Wang, H., Shi, G., Liang, Z., Luan, Q., Geng, F., and Zhou, a.: The Properties and Formation of Cirrus Clouds over the Tibetan Plateau Based on Summertime Lidar Measurements, Journal of the Atmospheric Sciences, 70, https://doi.org/10.1175/JAS-D-12-0171.1, 2012.
 - Heymsfield, A., Winker, D., Avery, M., Vaughan, M., Diskin, G., Deng, M., Mitev, V., and Matthey, R.: Relationships between Ice Water Content and Volume Extinction Coefficient from In Situ Observations for Temperatures from 0° to -86°C: Implications for Spaceborne
- 750 Lidar Retrievals, Journal of Applied Meteorology and Climatology, 53, 479 505, https://doi.org/10.1175/JAMC-D-13-087.1, https: //journals.ametsoc.org/view/journals/apme/53/2/jamc-d-13-087.1.xml, 2014.
 - Heymsfield, A. J., Winker, D., and van Zadelhoff, G.-J.: Extinction-ice water content-effective radius algorithms for CALIPSO, Geophysical Research Letters, 32, https://doi.org/https://doi.org/10.1029/2005GL022742, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2005GL022742, 2005.

- Dubovik, O., Sinyuk, A., Lapyonok, T., Holben, B. N., Mishchenko, M., Yang, P., Eck, T. F., Volten, H., Muñoz, O., Veihelmann, B., van der Zande, W. J., Leon, J.-F., Sorokin, M., and Slutsker, I.: Application of spheroid models to account for aerosol particle nonsphericity in remote sensing of desert dust, J. Geophys. Res., 111, D11 208, https://doi.org/10.1029/2005JD006619, 2006.
- Dubovik, O., Lapyonok, T., Litvinov, P., Herman, M., Fuertes, D., Ducos, F., Torres, B., Derimian, Y., Huang, X., Lopatin, A., Chaikovsky, A., Aspetsberger, M., and Federspiel, C.: GRASP: a versatile algorithm for characterizing the atmosphere, SPIE Newsroom, https://doi.org/10.1117/2.1201408.005558, http://spie.org/x109993.xml, 2014.
- Frey, W., Borrmann, S., Kunkel, D., Weigel, R., de Reus, M., Schlager, H., Roiger, A., Voigt, C., Hoor, P., Curtius, J., Krämer, M., Schiller, C.,
- 735 Volk, C. M., Homan, C. D., Fierli, F., Di Donfrancesco, G., Ulanovsky, A., Ravegnani, F., Sitnikov, N. M., Viciani, S., D'Amato, F., Shur, G. N., Belyaev, G. V., Law, K. S., and Cairo, F.: In situ measurements of tropical cloud properties in the West African Monsoon: upper tropospheric ice clouds, Mesoscale Convective System outflow, and subvisual cirrus, Atmospheric Chemistry and Physics, 11, 5569–5590, https://doi.org/10.5194/acp-11-5569-2011, https://acp.copernicus.org/articles/11/5569/2011/, 2011.
- Frey, W., Borrmann, S., Fierli, F., Weigel, R., Mitev, V., Matthey, R., Ravegnani, F., Sitnikov, N. M., Ulanovsky, A., and Cairo, F.: Tropical
 deep convective life cycle: Cb-anvil cloud microphysics from high-altitude aircraft observations, Atmospheric Chemistry and Physics, 14, 13 223–13 240, https://doi.org/10.5194/acp-14-13223-2014, https://acp.copernicus.org/articles/14/13223/2014/, 2014.
 - Fu, Q.: A New Parameterization of an Asymmetry Factor of Cirrus Clouds for Climate Models, Journal of the Atmospheric Sciences, 64, 4140 – 4150, https://doi.org/10.1175/2007JAS2289.1, https://journals.ametsoc.org/view/journals/atsc/64/11/2007jas2289.1.xml, 2007.
- Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, Reviews of Geophysics, 47,
 https://doi.org/https://doi.org/10.1029/2008RG000267, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008RG000267, 2009.
- Fujiwara, M., Iwasaki, S., Shimizu, A., Inai, Y., Shiotani, M., Hasebe, F., Matsui, I., Sugimoto, N., Okamoto, H., Nishi, N., Hamada, A., Sakazaki, T., and Yoneyama, K.: Cirrus observations in the tropical tropopause layer over the western Pacific, Journal of Geophysical Research: Atmospheres, 114, https://doi.org/10.1029/2008JD011040, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2008JD011040, 2009.
- 750 Groth, S. P., Baran, A. J., Betcke, T., Havemann, S., and Śmigaj, W.: The boundary element method for light scattering by ice crystals and its implementation in BEM++, Journal of Quantitative Spectroscopy and Radiative Transfer, 167, 40–52, 2015.
 - Grulich, L., Weigel, R., Hildebrandt, A., Wand, M., and Spichtinger, P.: Automatic shape detection of ice crystals, Journal of Computational Science, 54, 101 429, https://doi.org/https://doi.org/10.1016/j.jocs.2021.101429, https://www.sciencedirect.com/science/article/pii/ S1877750321001137, 2021.
- 755 Guignard, A., Stubenrauch, C. J., Baran, A. J., and Armante, R.: Bulk microphysical properties of semi-transparent cirrus from AIRS: a six year global climatology and statistical analysis in synergy with geometrical profiling data from CloudSat-CALIPSO, Atmospheric Chemistry and Physics, 12, 503–525, https://doi.org/10.5194/acp-12-503-2012, https://acp.copernicus.org/articles/12/503/2012/, 2012.
 - He, Q., LI, C., Ma, J., Wang, H., Shi, G., Liang, Z., Luan, Q., Geng, F., and Zhou, a.: The Properties and Formation of Cirrus Clouds over the Tibetan Plateau Based on Summertime Lidar Measurements, Journal of the Atmospheric Sciences, 70, https://doi.org/10.1175/JAS-D-
- 760 12-0171.1, 2012.

730

Heymsfield, A., Winker, D., Avery, M., Vaughan, M., Diskin, G., Deng, M., Mitev, V., and Matthey, R.: Relationships between Ice Water Content and Volume Extinction Coefficient from In Situ Observations for Temperatures from 0° to -86°C: Implications for Spaceborne Lidar Retrievals, Journal of Applied Meteorology and Climatology, 53, 479 – 505, https://doi.org/10.1175/JAMC-D-13-087.1, https: //journals.ametsoc.org/view/journals/apme/53/2/jamc-d-13-087.1.xml, 2014.

- 755 Immler, F., Krüger, K., Fujiwara, M., Verver, G., Rex, M., and Schrems, O.: Correlation between equatorial Kelvin waves and the occurrence of extremely thin ice clouds at the tropical tropopause, Atmospheric Chemistry and Physics, 8, 4019–4026, https://doi.org/10.5194/acp-8-4019-2008, https://www.atmos-chem-phys.net/8/4019/2008/, 2008.
 - Ishimoto, H., Masuda, K., Mano, Y., Orikasa, N., and Uchiyama, A.: Optical modeling of irregularly shaped ice particles in convective cirrus, in: AIP Conference Proceedings, vol. 1531, pp. 184–187, American Institute of Physics, 2013.
- Jensen, E. J., Toon, O. B., Pfister, L., and Selkirk, H. B.: Dehydration of the upper troposphere and lower stratosphere by subvisible cirrus clouds near the tropical tropopause, Geophysical Research Letters, 23, 825–828, https://doi.org/https://doi.org/10.1029/96GL00722, https://doi.org/lo.1029/96GL00722, https://doi.org/lo.1029/96GL00722, 1996.
 - Khaykin, S., Engel, I., Voemel, H., Formanyuk, I., Kivi, R., Korshunov, L., Kraemer, M., Lykov, A., Meier, S., Naebert, T., Pitts, M., Santee, M., Spelten, N., Wienhold, F., Yushkov, V., and Peter, T.: Arctic stratospheric dehydration – Part 1: Unprecedented observation of vertical
- 765 redistribution of water, Atmospheric Chemistry and Physics Discussions, 13, 14 249–14 295, https://doi.org/10.5194/acpd-13-14249-2013, 2013.
- Khaykin, S. M., Moyer, E., Krämer, M., Clouser, B., Bucci, S., Legras, B., Lykov, A., Afchine, A., Cairo, F., Formanyuk, I., Mitev, V., Matthey, R., Rolf, C., Singer, C. E., Spelten, N., Volkov, V., Yushkov, V., and Stroh, F.: Persistence of moist plumes from overshooting convection in the Asian monsoon anticyclone, Atmospheric Chemistry and Physics, 22, 3169–3189, https://doi.org/10.5194/acp-22-3169-2022, https://acp.copernicus.org/articles/22/3169/2022/, 2022.
 - Klingebiel, M., de Lozar, A., Molleker, S., Weigel, R., Roth, A., Schmidt, L., Meyer, J., Ehrlich, A., Neuber, R., Wendisch, M., and Borrmann, S.: Arctic low-level boundary layer clouds: in situ measurements and simulations of mono- and bimodal supercooled droplet size distributions at the top layer of liquid phase clouds, Atmospheric Chemistry and Physics, 15, 617–631, https://doi.org/10.5194/acp-15-617-2015, https://acp.copernicus.org/articles/15/617/2015/, 2015.
- 775 Konoshonkin, A., Borovoi, A., Kustova, N., Okamoto, H., Ishimoto, H., Grynko, Y., and Förstner, J.: Light scattering by ice crystals of cirrus clouds: From exact numerical methods to physical-optics approximation, Journal of Quantitative Spectroscopy and Radia-tive Transfer, 195, 132–140, https://doi.org/https://doi.org/10.1016/j.jqsrt.2016.12.024, https://www.sciencedirect.com/science/article/pii/S0022407316305775, laser-light and Interactions with Particles 2016, 2017a.

- Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus, M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, Atmospheric Chemistry and Physics, 9, 3505–3522, https://doi.org/10.5194/acp-9-3505-2009, https://acp.copernicus.org/articles/9/3505/2009/, 2009.
- Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zöger, M., Smith, J., Herman, R. L., Buchholz, B., Ebert,
 V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A microphysics guide to cirrus clouds Part 1: Cirrus types, Atmospheric Chemistry and Physics, 16, 3463–3483, https://doi.org/10.5194/acp-16-3463-2016, https://acp.copernicus.org/articles/16/ 3463/2016/, 2016.
- Krämer. М., Rolf. C., Spelten. N., Afchine, Α., Fahey, D.. Jensen. E.. Khaykin, S.. Kuhn. Т.. Law-P., Lykov, А., L. L., Riese, М., Rollins, A., Stroh, F., Thornberry, Т., Wolf, V., son, Pan, 790 Woods. S.. Spichtinger, Р., Quaas, J., and Sourdeval, O.: А microphysics guide cirrus Part 2: to _ Climatologies of clouds and humidity from observations, Atmospheric Chemistry and Physics, 20, 12569-12608, https://doi.org/10.5194/acp-20-12569-2020, https://acp.copernicus.org/articles/20/12569/2020/, 2020.

Konoshonkin, A., Borovoi, A., Kustova, N., and Reichardt, J.: Power laws for backscattering by ice crystals of cirrus clouds, Opt. Express, 25, 22 341–22 346, https://doi.org/10.1364/OE.25.022341, http://opg.optica.org/oe/abstract.cfm?URI=oe-25-19-22341, 2017b.

- 765 Heymsfield, A. J., Winker, D., and van Zadelhoff, G.-J.: Extinction-ice water content-effective radius algorithms for CALIPSO, Geophysical Research Letters, 32, https://doi.org/https://doi.org/10.1029/2005GL022742, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2005GL022742, 2005.
 - Immler, F., Krüger, K., Fujiwara, M., Verver, G., Rex, M., and Schrems, O.: Correlation between equatorial Kelvin waves and the occurrence of extremely thin ice clouds at the tropical tropopause, Atmospheric Chemistry and Physics, 8, 4019–4026, https://doi.org/10.5194/acp-8-4019-2008, https://www.atmos-chem-phys.net/8/4019/2008/, 2008.
 - Ishimoto, H., Masuda, K., Mano, Y., Orikasa, N., and Uchiyama, A.: Optical modeling of irregularly shaped ice particles in convective cirrus, in: AIP Conference Proceedings, vol. 1531, pp. 184–187, American Institute of Physics, 2013.

770

- Jensen, E. J., Toon, O. B., Pfister, L., and Selkirk, H. B.: Dehydration of the upper troposphere and lower stratosphere by subvisible cirrus clouds near the tropical tropopause, Geophysical Research Letters, 23, 825–828, https://doi.org/https://doi.org/10.1029/96GL00722, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/96GL00722, 1996.
- Khaykin, S., Engel, I., Voemel, H., Formanyuk, I., Kivi, R., Korshunov, L., Kraemer, M., Lykov, A., Meier, S., Naebert, T., Pitts, M., Santee, M., Spelten, N., Wienhold, F., Yushkov, V., and Peter, T.: Arctic stratospheric dehydration – Part 1: Unprecedented observation of vertical redistribution of water, Atmospheric Chemistry and Physics Discussions, 13, 14 249–14 295, https://doi.org/10.5194/acpd-13-14249-2013, 2013.
- 780 Khaykin, S. M., Moyer, E., Krämer, M., Clouser, B., Bucci, S., Legras, B., Lykov, A., Afchine, A., Cairo, F., Formanyuk, I., Mitev, V., Matthey, R., Rolf, C., Singer, C. E., Spelten, N., Volkov, V., Yushkov, V., and Stroh, F.: Persistence of moist plumes from overshooting convection in the Asian monsoon anticyclone, Atmospheric Chemistry and Physics, 22, 3169–3189, https://doi.org/10.5194/acp-22-3169-2022, https://acp.copernicus.org/articles/22/3169/2022/, 2022.
 - Klingebiel, M., de Lozar, A., Molleker, S., Weigel, R., Roth, A., Schmidt, L., Meyer, J., Ehrlich, A., Neuber, R., Wendisch, M., and Borrmann,
- 785 S.: Arctic low-level boundary layer clouds: in situ measurements and simulations of mono- and bimodal supercooled droplet size distributions at the top layer of liquid phase clouds, Atmospheric Chemistry and Physics, 15, 617–631, https://doi.org/10.5194/acp-15-617-2015, https://acp.copernicus.org/articles/15/617/2015/, 2015.
 - Konoshonkin, A., Borovoi, A., Kustova, N., Okamoto, H., Ishimoto, H., Grynko, Y., and Förstner, J.: Light scattering by ice crystals of cirrus clouds: From exact numerical methods to physical-optics approximation, Journal of Quantitative Spectroscopy and Radia-
- 790 tive Transfer, 195, 132–140, https://doi.org/https://doi.org/10.1016/j.jqsrt.2016.12.024, https://www.sciencedirect.com/science/article/pii/ S0022407316305775, laser-light and Interactions with Particles 2016, 2017a.
 - Konoshonkin, A., Borovoi, A., Kustova, N., and Reichardt, J.: Power laws for backscattering by ice crystals of cirrus clouds, Opt. Express, 25, 22 341–22 346, https://doi.org/10.1364/OE.25.022341, http://opg.optica.org/oe/abstract.cfm?URI=oe-25-19-22341, 2017b.
- Krämer, M., Schiller, C., Afchine, A., Bauer, R., Gensch, I., Mangold, A., Schlicht, S., Spelten, N., Sitnikov, N., Borrmann, S., de Reus, 795 M., and Spichtinger, P.: Ice supersaturations and cirrus cloud crystal numbers, Atmospheric Chemistry and Physics, 9, 3505–3522,
 - https://doi.org/10.5194/acp-9-3505-2009, https://acp.copernicus.org/articles/9/3505/2009/, 2009.
- Krämer, M., Rolf, C., Luebke, A., Afchine, A., Spelten, N., Costa, A., Meyer, J., Zöger, M., Smith, J., Herman, R. L., Buchholz, B., Ebert, V., Baumgardner, D., Borrmann, S., Klingebiel, M., and Avallone, L.: A microphysics guide to cirrus clouds Part 1: Cirrus types, Atmospheric Chemistry and Physics, 16, 3463–3483, https://doi.org/10.5194/acp-16-3463-2016, https://acp.copernicus.org/articles/16/ 3463/2016/, 2016.
 - Krämer, М., Rolf, С., Spelten, N., Afchine, A., Fahey, D., Jensen, Е., Khaykin, S., Kuhn, Т., Law-Lykov, Pan, L. L., Riese, М., Rollins. Stroh, F.. Thornberry, Т., Wolf. V., son. Р... A., A.,

Liou, K.-N. and Yang, P.: Light scattering by ice crystals: Fundamentals and applications, https://doi.org/10.1017/CBO9781139030052, 2016.

- 795 Liu, C., Panetta, R. L., and Yang, P.: Application of the pseudo-spectral time domain method to compute particle single-scattering properties for size parameters up to 200, Journal of Quantitative Spectroscopy and Radiative Transfer, 113, 1728–1740, 2012.
 - Liu, L. and Mishchenko, M. I.: Constraints on PSC particle microphysics derived from lidar observations, Journal of Quantitative Spectroscopy and Radiative Transfer, 70, 817–831, 2001.
 - Liu, L., Mishchenko, M. I., Cairns, B., Carlson, B. E., and Travis, L. D.: Modeling single-scattering properties of small cirrus particles by
- 800 use of a size-shape distribution of ice spheroids and cylinders, Journal of Quantitative Spectroscopy and Radiative Transfer, 101, 488–497, https://doi.org/https://doi.org/10.1016/j.jqsrt.2006.02.040, https://www.sciencedirect.com/science/article/pii/S0022407306000501, light in Planetary Atmospheres and Other Particulate Media, 2006.
 - Lolli, S.: Daytime top-of-the-atmosphere cirrus cloud radiative forcing properties at Singapore, J. Appl. Meteor. Climatol., 56, 1249–1257, 2017.
- 805 Luebke, A., Avallone, L., Schiller, C., Meyer, J., Rolf, C., and Krämer, M.: Ice water content of Arctic, midlatitude, and tropical cirrus Part 2: Extension of the database and new statistical analysis, Atmospheric Chemistry and Physics, 13, https://doi.org/10.5194/acp-13-6447-2013, 2013.

810

- Luebke, A. E., Afchine, A., Costa, A., Grooß, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, Atmospheric Chemistry and Physics, 16, 5793–5809, https://doi.org/10.5194/acp-16-5793-2016, https://acp.copernicus.org/articles/16/5793/2016/, 2016.
- Luo, B. P., Peter, T., Fueglistaler, S., Wernli, H., Wirth, M., Kiemle, C., Flentje, H., Yushkov, V. A., Khattatov, V., Rudakov, V., Thomas, A., Borrmann, S., Toci, G., Mazzinghi, P., Beuermann, J., Schiller, C., Cairo, F., Di Donfrancesco, G., Adriani, A., Volk, C. M., Strom, J., Noone, K., Mitev, V., MacKenzie, R. A., Carslaw, K. S., Trautmann, T., Santacesaria, V., and Stefanutti, L.: Dehydration potential of ultrathin clouds at the tropical tropopause, Geophysical Research Letters, 30, https://doi.org/https://doi.org/10.1029/2002GL016737, https://doi.org/https://doi.org/10.1029/2002GL016737, 2003.
 - Lynch, D.: Cirrus clouds: Their role in climate and global change, Acta Astronautica, 38, 859-863. https://doi.org/https://doi.org/10.1016/S0094-5765(96)00098-7, https://www.sciencedirect.com/science/article/pii/S0094576596000987, 1996.
- Lynch, D., Sassen, K., Delgenio, A., Heymsfield, A., Minnis, P., Platt, M., Quante, M., Schumann, U., and Sundqvist, H.: Cirrus, Oxford University Press, https://doi.org/10.1093/oso/9780195130720.003.0025, 2002.

Macke, A., Mueller, J., and Raschke, E.: Single scattering properties of atmospheric ice crystals, Journal of Atmospheric Sciences, 53, 2813–2825, 1996.

- MacKenzie, A. R., Schiller, C., Peter, T., Adriani, A., Beuermann, J., Bujok, O., Cairo, F., Corti, T., DiDonfrancesco, G., Gensch, I., Kiemle, C., Krämer, M., Kröger, C., Merkulov, S., Oulanovsky, A., Ravegnani, F., Rohs, S., Rudakov, V., Salter, P., Santacesaria, V., Stefanutti,
- 825 L., and Yushkov, V.: Tropopause and hygropause variability over the equatorial Indian Ocean during February and March 1999, Journal of Geophysical Research: Atmospheres, 111, https://doi.org/https://doi.org/10.1029/2005JD006639, https://agupubs.onlinelibrary.wiley. com/doi/abs/10.1029/2005JD006639, 2006.
 - Mahnke, C., Weigel, R., Cairo, F., Vernier, J.-P., Afchine, A., Krämer, M., Mitev, V., Matthey, R., Viciani, S., D'Amato, F., Ploeger, F., Deshler, T., and Borrmann, S.: The Asian tropopause aerosol layer within the 2017 monsoon anticyclone: microphysical properties derived

Woods, S., Spichtinger, P., Quaas, J., and Sourdeval, O.: A microphysics guide to cirrus – Part 2: Climatologies of clouds and humidity from observations, Atmospheric Chemistry and Physics, 20, 12569–12608, https://doi.org/10.5194/acp-20-12569-2020, https://acp.copernicus.org/articles/20/12569/2020/, 2020.

- Lamraoui, F., Krämer, M., Afchine, A., Sokol, A. B., Khaykin, S., Pandey, A., and Kuang, Z.: Sensitivity of convectively driven tropical tropopause cirrus properties to ice habits in high-resolution simulations, Atmospheric Chemistry and Physics, 23, 2393–2419, https://doi.org/10.5194/acp-23-2393-2023, https://acp.copernicus.org/articles/23/2393/2023/, 2023.
- Lawson, R. P., Woods, S., Jensen, E., Erfani, E., Gurganus, C., Gallagher, M., Connolly, P., Whiteway, J., Baran, A. J., May, P., Heyms field, A., Schmitt, C. G., McFarquhar, G., Um, J., Protat, A., Bailey, M., Lance, S., Muehlbauer, A., Stith, J., Korolev, A., Toon, O. B., and Krämer, M.: A Review of Ice Particle Shapes in Cirrus formed In Situ and in Anvils, Journal of Geophysical Research: Atmospheres, 124, 10049–10090, https://doi.org/https://doi.org/10.1029/2018JD030122, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JD030122, 2019.

- Liu, C., Panetta, R. L., and Yang, P.: Application of the pseudo-spectral time domain method to compute particle single-scattering properties for size parameters up to 200, Journal of Quantitative Spectroscopy and Radiative Transfer, 113, 1728–1740, 2012.
- Liu, L. and Mishchenko, M. I.: Constraints on PSC particle microphysics derived from lidar observations, Journal of Quantitative Spectroscopy and Radiative Transfer, 70, 817–831, 2001.
- 820 Liu, L., Mishchenko, M. I., Cairns, B., Carlson, B. E., and Travis, L. D.: Modeling single-scattering properties of small cirrus particles by use of a size-shape distribution of ice spheroids and cylinders, Journal of Quantitative Spectroscopy and Radiative Transfer, 101, 488–497, https://doi.org/https://doi.org/10.1016/j.jqsrt.2006.02.040, https://www.sciencedirect.com/science/article/pii/S0022407306000501, light in Planetary Atmospheres and Other Particulate Media, 2006.

Lolli, S.: Daytime top-of-the-atmosphere cirrus cloud radiative forcing properties at Singapore, J. Appl. Meteor. Climatol., 56, 1249–1257,

825 2017.

805

- Luebke, A., Avallone, L., Schiller, C., Meyer, J., Rolf, C., and Krämer, M.: Ice water content of Arctic, midlatitude, and tropical cirrus Part 2: Extension of the database and new statistical analysis, Atmospheric Chemistry and Physics, 13, https://doi.org/10.5194/acp-13-6447-2013, 2013.
- Luebke, A. E., Afchine, A., Costa, A., Grooß, J.-U., Meyer, J., Rolf, C., Spelten, N., Avallone, L. M., Baumgardner, D., and Krämer, M.: 830 The origin of midlatitude ice clouds and the resulting influence on their microphysical properties, Atmospheric Chemistry and Physics,

16, 5793-5809, https://doi.org/10.5194/acp-16-5793-2016, https://acp.copernicus.org/articles/16/5793/2016/, 2016.

- Luo, B. P., Peter, T., Fueglistaler, S., Wernli, H., Wirth, M., Kiemle, C., Flentje, H., Yushkov, V. A., Khattatov, V., Rudakov, V., Thomas, A., Borrmann, S., Toci, G., Mazzinghi, P., Beuermann, J., Schiller, C., Cairo, F., Di Donfrancesco, G., Adriani, A., Volk, C. M., Strom, J., Noone, K., Mitev, V., MacKenzie, R. A., Carslaw, K. S., Trautmann, T., Santacesaria, V., and Stefanutti, L.: Dehydration potential
- of ultrathin clouds at the tropical tropopause, Geophysical Research Letters, 30, https://doi.org/https://doi.org/10.1029/2002GL016737, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2002GL016737, 2003.
 - Lynch, D.: Cirrus clouds: Their role global change, Acta 38, 859-863. in climate and Astronautica, https://doi.org/https://doi.org/10.1016/S0094-5765(96)00098-7, https://www.sciencedirect.com/science/article/pii/S0094576596000987, 1996.

Liou, K.-N. and Yang, P.: Light scattering by ice crystals: Fundamentals and applications, https://doi.org/10.1017/CBO9781139030052, 2016.

- 830 from aircraft-borne in situ measurements, Atmospheric Chemistry and Physics, 21, 15259–15282, https://doi.org/10.5194/acp-21-15259-2021, https://acp.copernicus.org/articles/21/15259/2021/, 2021.
 - Martucci, G., Mitev, V., Matthey, R., and Makarov, V.: Compact backscattter Lidar for cloud detection on a stratospheric aircraft, Proc SPIE, 5979, https://doi.org/10.1117/12.627644, 2005.
- Mei, F., Wang, J., Comstock, J. M., Weigel, R., Krämer, M., Mahnke, C., Shilling, J. E., Schneider, J., Schulz, C., Long, C. N.,
 Wendisch, M., Machado, L. A. T., Schmid, B., Krisna, T., Pekour, M., Hubbe, J., Giez, A., Weinzierl, B., Zoeger, M., Pöhlker, M. L.,
 Schlager, H., Cecchini, M. A., Andreae, M. O., Martin, S. T., de Sá, S. S., Fan, J., Tomlinson, J., Springston, S., Pöschl, U., Ar
 - taxo, P., Pöhlker, C., Klimach, T., Minikin, A., Afchine, A., and Borrmann, S.: Comparison of aircraft measurements during GoAmazon2014/5 and ACRIDICON-CHUVA, Atmospheric Measurement Techniques, 13, 661–684, https://doi.org/10.5194/amt-13-661-2020, https://amt.copernicus.org/articles/13/661/2020/, 2020.
- 840 Meyer, J., Rolf, C., Schiller, C., Rohs, S., Spelten, N., Afchine, A., Zoeger, M., Sitnikov, N., Thornberry, T., Rollins, A., Bozóki, Z., Tatrai, D., Ebert, V., Kühnreich, B., Mackrodt, P., Möhler, O., Saathoff, H., Rosenlof, K., and Krämer, M.: Two decades of water vapor measurements with the FISH fluorescence hygrometer: A review, Atmospheric Chemistry and Physics, 15, 7735–7782, https://doi.org/10.5194/acp-15-8521-2015, 2015.
- Meyer, K. and Platnick, S.: Utilizing the MODIS 1.38 μ m channel for cirrus cloud optical thickness retrievals: Algorithm and retrieval uncertainties, Journal of Geophysical Research, 115, https://doi.org/10.1029/2010JD014872, 2010.
 - Mishchenko, M. I., Travis, L. D., and Mackowski, D. W.: T-matrix computations of light scattering by nonspherical particles: A review, J. Quant. Spectrosc. Radiat. Transfer, 55, 535–575, https://doi.org/10.1016/0022-4073(96)00002-7, 1996.
 - Mishchenko, M. I., Wielaard, D. J., and Carlson, B. E.: T-matrix computations of zenith-enhanced lidar backscatter from horizontally oriented ice plates, Geophys. Res. Lett., 24, 771–774, https://doi.org/10.1029/97GL00545, 1997.
- 850 Mishchenko, M. I., Hovenier, J. W., and Travis, L. D.: Light scattering by nonspherical particles: theory, measurements, and applications, Elsevier, 1999.
 - Mishchenko, M. I., Travis, L. D., and Lacis, A. A.: Scattering, Absorption, and Emission of Light by Small Particles, Cambridge University Press, Cambridge, 2002.
 - Mitchell, D. L., Lawson, R. P., and Baker, B.: Understanding effective diameter and its application to terrestrial radiation in ice clouds,
- Atmospheric Chemistry and Physics, 11, 3417–3429, https://doi.org/10.5194/acp-11-3417-2011, https://acp.copernicus.org/articles/11/ 3417/2011/, 2011.
 - Mitchell, D. L., Garnier, A., Pelon, J., and Erfani, E.: CALIPSO (IIR–CALIOP) retrievals of cirrus cloud ice-particle concentrations, Atmospheric Chemistry and Physics, 18, 17 325–17 354, https://doi.org/10.5194/acp-18-17325-2018, https://acp.copernicus.org/articles/18/ 17325/2018/, 2018.
- 860 Mitev, V., Poole, L. R., Pitts, M. C., and Matthey, R.: Comparison Case Between CALIPSO Lidar and Mals On M55 Geophysica During Reconcile Campaign, in: Reviewed and Revised Papers Presented at the 26th International Laser Radar Conference, 26, pp. 729–732, ILRC, Porto Heli, Greece, 2012.
 - Molleker, S., Borrmann, S., Schlager, H., Luo, B., Frey, W., Klingebiel, M., Weigel, R., Ebert, M., Mitev, V., Matthey, R., Woiwode, W., Oelhaf, H., Dörnbrack, A., Stratmann, G., Grooß, J.-U., Günther, G., Vogel, B., Müller, R., Krämer, M., Meyer, J., and Cairo, F.:
- 865 Microphysical properties of synoptic-scale polar stratospheric clouds: in situ measurements of unexpectedly large HNO₃-containing particles in the Arctic vortex, Atmospheric Chemistry and Physics, 14, 10785–10801, https://doi.org/10.5194/acp-14-10785-2014, https://acp.copernicus.org/articles/14/10785/2014/, 2014.

- 840 Lynch, D., Sassen, K., Delgenio, A., Heymsfield, A., Minnis, P., Platt, M., Quante, M., Schumann, U., and Sundqvist, H.: Cirrus, Oxford University Press, https://doi.org/10.1093/oso/9780195130720.003.0025, 2002.
 - Macke, A., Mueller, J., and Raschke, E.: Single scattering properties of atmospheric ice crystals, Journal of Atmospheric Sciences, 53, 2813–2825, 1996.
 - MacKenzie, A. R., Schiller, C., Peter, T., Adriani, A., Beuermann, J., Bujok, O., Cairo, F., Corti, T., DiDonfrancesco, G., Gensch, I., Kiemle,
- 845 C., Krämer, M., Kröger, C., Merkulov, S., Oulanovsky, A., Ravegnani, F., Rohs, S., Rudakov, V., Salter, P., Santacesaria, V., Stefanutti, L., and Yushkov, V.: Tropopause and hygropause variability over the equatorial Indian Ocean during February and March 1999, Journal of Geophysical Research: Atmospheres, 111, https://doi.org/10.1029/2005JD006639, https://agupubs.onlinelibrary.wiley. com/doi/abs/10.1029/2005JD006639, 2006.
- Mahnke, C., Weigel, R., Cairo, F., Vernier, J.-P., Afchine, A., Krämer, M., Mitev, V., Matthey, R., Viciani, S., D'Amato, F., Ploeger, F.,
 Deshler, T., and Borrmann, S.: The Asian tropopause aerosol layer within the 2017 monsoon anticyclone: microphysical properties derived from aircraft-borne in situ measurements, Atmospheric Chemistry and Physics, 21, 15 259–15 282, https://doi.org/10.5194/acp-21-15259-2021, https://acp.copernicus.org/articles/21/15259/2021, 2021.
 - Martucci, G., Mitev, V., Matthey, R., and Makarov, V.: Compact backscattter Lidar for cloud detection on a stratospheric aircraft, Proc SPIE, 5979, https://doi.org/10.1117/12.627644, 2005.
- Mei, F., Wang, J., Comstock, J. M., Weigel, R., Krämer, M., Mahnke, C., Shilling, J. E., Schneider, J., Schulz, C., Long, C. N., Wendisch, M., Machado, L. A. T., Schmid, B., Krisna, T., Pekour, M., Hubbe, J., Giez, A., Weinzierl, B., Zoeger, M., Pöhlker, M. L., Schlager, H., Cecchini, M. A., Andreae, M. O., Martin, S. T., de Sá, S. S., Fan, J., Tomlinson, J., Springston, S., Pöschl, U., Artaxo, P., Pöhlker, C., Klimach, T., Minikin, A., Afchine, A., and Borrmann, S.: Comparison of aircraft measurements during GoAmazon2014/5 and ACRIDICON-CHUVA, Atmospheric Measurement Techniques, 13, 661–684, https://doi.org/10.5194/amt-13-661-2020, https://amt.copernicus.org/articles/13/661/2020/, 2020.
- Meyer, J., Rolf, C., Schiller, C., Rohs, S., Spelten, N., Afchine, A., Zoeger, M., Sitnikov, N., Thornberry, T., Rollins, A., Bozóki, Z., Tatrai, D., Ebert, V., Kühnreich, B., Mackrodt, P., Möhler, O., Saathoff, H., Rosenlof, K., and Krämer, M.: Two decades of water vapor measurements with the FISH fluorescence hygrometer: A review, Atmospheric Chemistry and Physics, 15, 7735–7782, https://doi.org/10.5194/acp-15-8521-2015, 2015.
- 865 Meyer, K. and Platnick, S.: Utilizing the MODIS 1.38 μm channel for cirrus cloud optical thickness retrievals: Algorithm and retrieval uncertainties, Journal of Geophysical Research, 115, https://doi.org/10.1029/2010JD014872, 2010.
 - Mishchenko, M. I., Travis, L. D., and Mackowski, D. W.: T-matrix computations of light scattering by nonspherical particles: A review, J. Quant. Spectrosc. Radiat. Transfer, 55, 535–575, https://doi.org/10.1016/0022-4073(96)00002-7, 1996.
- Mishchenko, M. I., Wielaard, D. J., and Carlson, B. E.: T-matrix computations of zenith-enhanced lidar backscatter from horizontally oriented
 ice plates, Geophys. Res. Lett., 24, 771–774, https://doi.org/10.1029/97GL00545, 1997.
 - Mishchenko, M. I., Hovenier, J. W., and Travis, L. D.: Light scattering by nonspherical particles: theory, measurements, and applications, Elsevier, 1999.
 - Mishchenko, M. I., Travis, L. D., and Lacis, A. A.: Scattering, Absorption, and Emission of Light by Small Particles, Cambridge University Press, Cambridge, 2002.
- 875 Mitchell, D. L., Lawson, R. P., and Baker, B.: Understanding effective diameter and its application to terrestrial radiation in ice clouds, Atmospheric Chemistry and Physics, 11, 3417–3429, https://doi.org/10.5194/acp-11-3417-2011, https://acp.copernicus.org/articles/11/ 3417/2011/, 2011.

- Nakajima, T. Y., Nakajima, T., Yoshimori, K., Mishra, S. K., and Tripathi, S. N.: Development of a light scattering solver applicable to particles of arbitrary shape on the basis of the surface-integral equations method of Müller type. I. Methodology, accuracy of calculation, and electromagnetic current on the particle surface. Applied optics, 48, 3526–3536, 2009.
- Nazaryan, H., Mccormick, M., and Menzel, W.: Global characterization of cirrus clouds using CALIPSO data, Journal of Geophysical Research, 113, https://doi.org/10.1029/2007JD009481, 2008.
 - Pfister, L., Selkirk, H. B., Jensen, E. J., Schoeberl, M. R., Toon, O. B., Browell, E. V., Grant, W. B., Gary, B., Mahoney, M. J., Bui, T. V., and Hintsa, E.: Aircraft observations of thin cirrus clouds near the tropical tropopause, Journal of Geophysical Research: Atmospheres, 106, 9765–9786. https://doi.org/10.1029/2000JD900648. https://agupubs.onlinelibrary.wilev.com/doi/abs/10.1029/2000JD900648. 2001.
- Port, M.: Flugzeuggetragene Messungen von Zirren und vereisten Wolken im Troposphären-Stratosphären-Übergangsbereich des asiatischen Sommermonsuns, Ph.D. thesis, Fachbereich Physik, Mathematik und Informatik der Johannes Gutenberg-Universität Mainz, 2021.
- Prabhakara, C., Kratz, D., Yoo, J.-M., Dalu, G., and Vernekar, A.: Optically thin cirrus clouds: Radiative impact on the warm pool, Journal of Quantitative Spectroscopy and Radiative Transfer, 49, 467–483, https://doi.org/https://doi.org/10.1016/0022-4073(93)90061-L, https://www.sciencedirect.com/science/article/pii/002240739390061L, 1993.
- Sarkozy, L. C., Clouser, B. W., Lamb, K. D., Stutz, E. J., Saathoff, H., Möhler, O., Ebert, V., and Moyer, E. J.: The Chicago Water Isotope Spectrometer (ChiWIS-lab): A tunable diode laser spectrometer for chamber-based measurements of water vapor isotopic evolution during cirrus formation, Review of Scientific Instruments, 91, 045 120, https://doi.org/10.1063/1.5139244, https://doi.org/10.1063/1.5139244, 2020.
- 885 Sassen, K. and Benson, S.: A midlatitude cirrus cloud climatology from the Facility for Atmospheric Remote Sensing. Part II: Microphysical properties derived from lidar depolarization, Journal of the Atmospheric Sciences, 58, 2103–2112, 2001.
 - Sassen, K., Wang, Z., and Liu, D.: Global distribution of cirrus clouds from CloudSat/Cloud-Aerosol Lidar and Pathfinder Satellite Observations (CALIPSO) measurements, Journal of Geophysical Research, 113, https://doi.org/10.1029/2008JD009972, 2008.
- Sassen, K., Wang, Z., and Liu, D.: Cirrus clouds and deep convection in the tropics: Insights from CALIPSO and CloudSat, Journal of Geophysical Research: Atmospheres, 114, https://doi.org/https://doi.org/10.1029/2009JD011916, https://agupubs.onlinelibrary.wiley.
 - Scarchilli, C., Adriani, A., Cairo, F., Di Donfrancesco, G., Buontempo, C., Snels, M., Moriconi, M. L., Deshler, T., Larsen, N., Luo, B., et al.: Determination of polar stratospheric cloud particle refractive indices by use of in situ optical measurements and T-matrix calculations, Applied optics, 44, 3302–3311, 2005.
- Schiller, C., Krämer, M., Afchine, A., Spelten, N., and Sitnikov, N.: Ice water content of Arctic, midlatitude, and tropical cirrus, Journal of Geophysical Research: Atmospheres, 113, https://doi.org/https://doi.org/10.1029/2008JD010342, https://agupubs.onlinelibrary.wiley. com/doi/abs/10.1029/2008JD010342, 2008.
 - Schnaiter, M., Kaye, P. H., Hirst, E., Ulanowski, Z., and Wagner, R.: Exploring the surface roughness of small ice crystals by measuring high resolution angular scattering patterns, Atti della Accademia Peloritana dei Pericolanti-Classe di Scienze Fisiche, Matematiche e Naturali,
- **900 8**9, 2011.

com/doi/abs/10.1029/2009JD011916, 2009.

870

875

880

Schreiner, J., Voigt, C., Weisser, C., Kohlmann, A., Mauersberger, K., Deshler, T., Kröger, C., Rosen, J., Kjome, N., Larsen, N., et al.: Chemical, microphysical, and optical properties of polar stratospheric clouds, Journal of Geophysical Research: Atmospheres, 107, SOL– 56, 2002. Mitchell, D. L., Garnier, A., Pelon, J., and Erfani, E.: CALIPSO (IIR–CALIOP) retrievals of cirrus cloud ice-particle concentrations, Atmospheric Chemistry and Physics, 18, 17 325–17 354, https://doi.org/10.5194/acp-18-17325-2018, https://acp.copernicus.org/articles/18/ 17325/2018/, 2018.

- Mitev, V., Poole, L. R., Pitts, M. C., and Matthey, R.: Comparison Case Between CALIPSO Lidar and Mals On M55 Geophysica During Reconcile Campaign, in: Reviewed and Revised Papers Presented at the 26th International Laser Radar Conference, 26, pp. 729–732, ILRC, Porto Heli, Greece, 2012.
- Molleker, S., Borrmann, S., Schlager, H., Luo, B., Frey, W., Klingebiel, M., Weigel, R., Ebert, M., Mitev, V., Matthey, R., Woiwode,
- 885 W., Oelhaf, H., Dörnbrack, A., Stratmann, G., Grooß, J.-U., Günther, G., Vogel, B., Müller, R., Krämer, M., Meyer, J., and Cairo, F.: Microphysical properties of synoptic-scale polar stratospheric clouds: in situ measurements of unexpectedly large HNO₃-containing particles in the Arctic vortex, Atmospheric Chemistry and Physics, 14, 10785–10801, https://doi.org/10.5194/acp-14-10785-2014, https://acp.copernicus.org/articles/14/10785/2014/, 2014.
- Nakajima, T. Y., Nakajima, T., Yoshimori, K., Mishra, S. K., and Tripathi, S. N.: Development of a light scattering solver applicable to
- particles of arbitrary shape on the basis of the surface-integral equations method of Müller type. I. Methodology, accuracy of calculation, and electromagnetic current on the particle surface, Applied optics, 48, 3526–3536, 2009.
 - Nazaryan, H., Mccormick, M., and Menzel, W.: Global characterization of cirrus clouds using CALIPSO data, Journal of Geophysical Research, 113, https://doi.org/10.1029/2007JD009481, 2008.
 - Pfister, L., Selkirk, H. B., Jensen, E. J., Schoeberl, M. R., Toon, O. B., Browell, E. V., Grant, W. B., Gary, B., Mahoney, M. J., Bui, T. V., and
- Hintsa, E.: Aircraft observations of thin cirrus clouds near the tropical tropopause, Journal of Geophysical Research: Atmospheres, 106, 9765–9786, https://doi.org/10.1029/2000JD900648, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2000JD900648, 2001.
 - Port, M.: Flugzeuggetragene Messungen von Zirren und vereisten Wolken im Troposphären-Stratosphären-Übergangsbereich des asiatischen Sommermonsuns, Ph.D. thesis, Fachbereich Physik, Mathematik und Informatik der Johannes Gutenberg-Universität Mainz, https://doi.org/10.25358/openscience-7538, https://openscience.ub.uni-mainz.de/handle/20.500.12030/7552, 2021.
- 900 Prabhakara, C., Kratz, D., Yoo, J.-M., Dalu, G., and Vernekar, A.: Optically thin cirrus clouds: Radiative impact on the warm pool, Journal of Quantitative Spectroscopy and Radiative Transfer, 49, 467–483, https://doi.org/https://doi.org/10.1016/0022-4073(93)90061-L, https://www.sciencedirect.com/science/article/pii/002240739390061L, 1993.
 - Sarkozy, L. C., Clouser, B. W., Lamb, K. D., Stutz, E. J., Saathoff, H., Möhler, O., Ebert, V., and Moyer, E. J.: The Chicago Water Isotope Spectrometer (ChiWIS-lab): A tunable diode laser spectrometer for chamber-based measurements of water vapor isotopic evolution during
- cirrus formation, Review of Scientific Instruments, 91, 045120, https://doi.org/10.1063/1.5139244, https://doi.org/10.1063/1.5139244, 2020.
 - Sassen, K. and Benson, S.: A midlatitude cirrus cloud climatology from the Facility for Atmospheric Remote Sensing. Part II: Microphysical properties derived from lidar depolarization, Journal of the Atmospheric Sciences, 58, 2103–2112, 2001.
- Sassen, K., Wang, Z., and Liu, D.: Global distribution of cirrus clouds from CloudSat/Cloud-Aerosol Lidar and Pathfinder Satellite Obser-910 vations (CALIPSO) measurements, Journal of Geophysical Research, 113, https://doi.org/10.1029/2008JD009972, 2008.
 - Sassen, K., Wang, Z., and Liu, D.: Cirrus clouds and deep convection in the tropics: Insights from CALIPSO and CloudSat, Journal of Geophysical Research: Atmospheres, 114, https://doi.org/https://doi.org/10.1029/2009JD011916, https://agupubs.onlinelibrary.wiley. com/doi/abs/10.1029/2009JD011916, 2009.

Schumann, U., Mayer, B., Gierens, K., Unterstrasser, S., Jessberger, P., Petzold, A., Voigt, C., and Gayet, J.-F.: Effective Radius of Ice
 Particles in Cirrus and Contrails, Journal of the Atmospheric Sciences, 68, 300 – 321, https://doi.org/10.1175/2010JAS3562.1, https://journals.ametsoc.org/view/journals/atsc/68/2/2010jas3562.1.xml, 2011.

- Schumann, U., Kiemle, C., Schlager, H., Weigel, R., Borrmann, S., D'Amato, F., Krämer, M., Matthey, R., Protat, A., Voigt, C., and Volk, C. M.: Long-lived contrails and convective cirrus above the tropical tropopause, Atmospheric Chemistry and Physics, 17, 2311–2346, https://doi.org/10.5194/acp-17-2311-2017, https://acp.copernicus.org/articles/17/2311/2017/, 2017.
- 910 Singer, C. E., Clouser, B., Khaykin, S., Krämer, M., Cairo, F., Peter, T., Lykov, A., Rolf, C., Spelten, N., Brunamonti, S., and Moyer, E. J.: Intercomparison of upper tropospheric and lower stratospheric water vapor measurements over the Asian Summer Monsoon during the StratoClim Campaign, Atmospheric Measurement Techniques Discussions, 2022, 1–23, https://doi.org/10.5194/amt-2022-13, https://amt.copernicus.org/preprints/amt-2022-13/, 2022.
 - Snels, M., Cairo, F., Di Liberto, L., Scoccione, A., Bracaglia, M., and Deshler, T.: Comparison of coincident optical particle counter and lidar
- measurements of polar stratospheric clouds above McMurdo (77.85 S, 166.67 E) from 1994 to 1999, Journal of Geophysical Research:
 Atmospheres, 126, e2020JD033 572, 2021.
 - Sourdeval, O., Labonnote, L. C., Brogniez, G., Jourdan, O., Pelon, J., and Garnier, A.: A variational approach for retrieving ice cloud properties from infrared measurements: application in the context of two IIR validation campaigns, Atmospheric Chemistry and Physics, 13, 8229–8244, https://doi.org/10.5194/acp-13-8229-2013, https://acp.copernicus.org/articles/13/8229/2013/, 2013.
- 920 Sourdeval, O., C.-Labonnote, L., Baran, A. J., and Brogniez, G.: A methodology for simultaneous retrieval of ice and liquid water cloud properties. Part I: Information content and case study, Quarterly Journal of the Royal Meteorological Society, 141, 870–882, https://doi.org/https://doi.org/10.1002/qj.2405, https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2405, 2015.
- Sourdeval, O., Gryspeerdt, E., Krämer, M., Goren, T., Delanoë, J., Afchine, A., Hemmer, F., and Quaas, J.: Ice crystal number concentration estimates from lidar–radar satellite remote sensing Part 1: Method and evaluation, Atmospheric Chemistry and Physics, 18, 14327–
 14 350, https://doi.org/10.5194/acp-18-14327-2018, https://acp.copernicus.org/articles/18/14327/2018, 2018.
- Stefanutti, L., Sokolov, L., Balestri, S., MacKenzie, A. R., and Khattatov, V.: The M-55 Geophysica as a Platform for the Airborne Polar Experiment, Journal of Atmospheric and Oceanic Technology, 16, 1303 – 1312, https://doi.org/10.1175/1520-0426(1999)016<1303:TMGAAP>2.0.CO;2, https://journals.ametsoc.org/view/journals/atot/16/10/1520-0426_1999_016_1303_tmgaap_ 2_0_co_2.xml, 1999.
- Stephens, G. L.: Cirrus, climate, and global change, Oxford University Press New York, 2002.
 Stephens, G. L.: Cloud feedbacks in the climate system: A critical review, Journal of climate, 18, 237–273, 2005.
 - Sun, W., Fu, Q., and Chen, Z.: Finite-difference time-domain solution of light scattering by dielectric particles with a perfectly matched layer absorbing boundary condition, Applied optics, 38, 3141–3151, 1999.
 - Sunilkumar, S. V. and Parameswaran, K.: Temperature dependence of tropical cirrus properties and radiative effects, Journal of Geo-
- 935 physical Research: Atmospheres, 110, https://doi.org/10.1029/2004JD005426, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2004JD005426, 2005.
 - Teri, M., Müller, T., Gasteiger, J., Valentini, S., Horvath, H., Vecchi, R., Bauer, P., Walser, A., and Weinzierl, B.: Impact of particle size, refractive index, and shape on the determination of the particle scattering coefficient–an optical closure study evaluating different nephelometer angular truncation and illumination corrections, Atmospheric Measurement Techniques, 15, 3161–3187, 2022.

Scarchilli, C., Adriani, A., Cairo, F., Di Donfrancesco, G., Buontempo, C., Snels, M., Moriconi, M. L., Deshler, T., Larsen, N., Luo, B., et al.:

- 915 Determination of polar stratospheric cloud particle refractive indices by use of in situ optical measurements and T-matrix calculations, Applied optics, 44, 3302–3311, 2005.
 - Schiller, C., Krämer, M., Afchine, A., Spelten, N., and Sitnikov, N.: Ice water content of Arctic, midlatitude, and tropical cirrus, Journal of Geophysical Research: Atmospheres, 113, https://doi.org/https://doi.org/10.1029/2008JD010342, https://agupubs.onlinelibrary.wiley. com/doi/abs/10.1029/2008JD010342, 2008.
- 920 Schnaiter, M., Kaye, P. H., Hirst, E., Ulanowski, Z., and Wagner, R.: Exploring the surface roughness of small ice crystals by measuring high resolution angular scattering patterns, Atti della Accademia Peloritana dei Pericolanti-Classe di Scienze Fisiche, Matematiche e Naturali, 89, 2011.
 - Schreiner, J., Voigt, C., Weisser, C., Kohlmann, A., Mauersberger, K., Deshler, T., Kröger, C., Rosen, J., Kjome, N., Larsen, N., et al.: Chemical, microphysical, and optical properties of polar stratospheric clouds, Journal of Geophysical Research: Atmospheres, 107, SOL-
- **925** 56, 2002.

- Schumann, U., Mayer, B., Gierens, K., Unterstrasser, S., Jessberger, P., Petzold, A., Voigt, C., and Gayet, J.-F.: Effective Radius of Ice Particles in Cirrus and Contrails, Journal of the Atmospheric Sciences, 68, 300 – 321, https://doi.org/10.1175/2010JAS3562.1, https: //journals.ametsoc.org/view/journals/atsc/68/2/2010jas3562.1.xml, 2011.
 - Schumann, U., Kiemle, C., Schlager, H., Weigel, R., Borrmann, S., D'Amato, F., Krämer, M., Matthey, R., Protat, A., Voigt, C., and Volk,
- 930 C. M.: Long-lived contrails and convective cirrus above the tropical tropopause, Atmospheric Chemistry and Physics, 17, 2311–2346, https://doi.org/10.5194/acp-17-2311-2017, https://acp.copernicus.org/articles/17/2311/2017/, 2017.
 - Singer, C. E., Clouser, B., Khaykin, S., Krämer, M., Cairo, F., Peter, T., Lykov, A., Rolf, C., Spelten, N., Brunamonti, S., and Moyer, E. J.: Intercomparison of upper tropospheric and lower stratospheric water vapor measurements over the Asian Summer Monsoon during the StratoClim Campaign, Atmospheric Measurement Techniques Discussions, 2022, 1–23, https://doi.org/10.5194/amt-2022-13, https: //amt.copernicus.org/preprints/amt-2022-13/, 2022.
 - Snels, M., Cairo, F., Di Liberto, L., Scoccione, A., Bracaglia, M., and Deshler, T.: Comparison of coincident optical particle counter and lidar measurements of polar stratospheric clouds above McMurdo (77.85 S, 166.67 E) from 1994 to 1999, Journal of Geophysical Research: Atmospheres, 126, e2020JD033 572, 2021.
 - Sourdeval, O., Labonnote, L. C., Brogniez, G., Jourdan, O., Pelon, J., and Garnier, A.: A variational approach for retrieving ice cloud
- 940 properties from infrared measurements: application in the context of two IIR validation campaigns, Atmospheric Chemistry and Physics, 13, 8229–8244, https://doi.org/10.5194/acp-13-8229-2013, https://acp.copernicus.org/articles/13/8229/2013/, 2013.
 - Sourdeval, O., C.-Labonnote, L., Baran, A. J., and Brogniez, G.: A methodology for simultaneous retrieval of ice and liquid water cloud properties. Part I: Information content and case study, Quarterly Journal of the Royal Meteorological Society, 141, 870–882, https://doi.org/10.1002/qj.2405, https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.2405, 2015.
- 945 Sourdeval, O., Gryspeerdt, E., Krämer, M., Goren, T., Delanoë, J., Afchine, A., Hemmer, F., and Quaas, J.: Ice crystal number concentration estimates from lidar–radar satellite remote sensing – Part 1: Method and evaluation, Atmospheric Chemistry and Physics, 18, 14327– 14350, https://doi.org/10.5194/acp-18-14327-2018, https://acp.copernicus.org/articles/18/14327/2018, 2018.
 - Stefanutti, L., Sokolov, L., Balestri, S., MacKenzie, A. R., and Khattatov, V.: The M-55 Geophysica as a Platform for the Airborne Polar Experiment, Journal of Atmospheric and Oceanic Technology, 16, 1303 1312, https://doi.org/10.1175/1520-0426(1999)016<1303;TMGAAP>2.0.CO;2, https://journals.ametsoc.org/view/journals/atot/16/10/1520-0426 1999 016 1303 tmgaap
- 950 0426(1999)016<1303:TMGAAP>2.0.CO;2, https://journals.ametsoc.org/view/journals/atot/16/10/1520-0426_1999_016_1303_tmgaap_ 2_0_co_2.xml, 1999.

- 940 Thomas, A., Borrmann, S., Kiemle, C., Cairo, F., Volk, C., Beuermann, J., Lepuchov, B., Santacesaria, V., Matthey, R., Rudakov, V., Yushkov, V., Mackenzie, A., and Stefanutti, L.: In Situ Measurements of Background Aerosol and Subvisible Cirrus in the Tropical Tropopause Region, Journal of Geophysical Research, v.107 (2002), 107, https://doi.org/10.1029/2001JD001385, 2002.
- Wagner, S. W. and Delene, D. J.: Technique for comparison of backscatter coefficients derived from in situ cloud probe measurements with concurrent airborne lidar, Atmospheric Measurement Techniques, 15, 6447–6466, https://doi.org/10.5194/amt-15-6447-2022, https:
 945 //amt.copernicus.org/articles/15/6447/2022/, 2022.
 - Weigel, R., Spichtinger, P., Mahnke, C., Klingebiel, M., Afchine, A., Petzold, A., Krämer, M., Costa, A., Molleker, S., Reutter, P., Szakáll, M., Port, M., Grulich, L., Jurkat, T., Minikin, A., and Borrmann, S.: Thermodynamic correction of particle concentrations measured by underwing probes on fast-flying aircraft, Atmospheric Measurement Techniques, 9, 5135–5162, https://doi.org/10.5194/amt-9-5135-2016, https://amt.copernicus.org/articles/9/5135/2016/, 2016.
- Weigel, R., Mahnke, C., Baumgartner, M., Dragoneas, A., Vogel, B., Ploeger, F., Viciani, S., D'Amato, F., Bucci, S., Legras, B., Luo, B., and Borrmann, S.: In situ observation of new particle formation (NPF) in the tropical tropopause layer of the 2017 Asian monsoon anticyclone – Part 1: Summary of StratoClim results, Atmospheric Chemistry and Physics, 21, 11689–11722, https://doi.org/10.5194/acp-21-11689-2021, https://acp.copernicus.org/articles/21/11689/2021/, 2021a.
 - Weigel, R., Mahnke, C., Baumgartner, M., Krämer, M., Spichtinger, P., Spelten, N., Afchine, A., Rolf, C., Viciani, S., D'Amato, F., Tost,
- 955 H., and Borrmann, S.: In situ observation of new particle formation (NPF) in the tropical tropopause layer of the 2017 Asian monsoon anticyclone – Part 2: NPF inside ice clouds, Atmospheric Chemistry and Physics, 21, 13455–13481, https://doi.org/10.5194/acp-21-13455-2021, https://acp.copernicus.org/articles/21/13455/2021/, 2021b.
- Wernli, H., Boettcher, M., Joos, H., Miltenberger, A. K., and Spichtinger, P.: A trajectory-based classification of ERA-Interim ice clouds in the region of the North Atlantic storm track, Geophysical Research Letters, 43, 6657–6664,
 https://doi.org/https://doi.org/10.1002/2016GL068922, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL068922, 2016.
- Wex, H., Neusüß, C., Wendisch, M., Stratmann, F., Koziar, C., Keil, A., Wiedensohler, A., and Ebert, M.: Particle scattering, backscattering, and absorption coefficients: An in situ closure and sensitivity study, Journal of Geophysical Research: Atmospheres, 107, LAC–4, 2002.
 - Yang, P. and Liou, K.: Geometric-optics-integral-equation method for light scattering by nonspherical ice crystals, Applied Optics, 35, 6568-6584, 1996a.
- Yang, P. and Liou, K. N.: Geometric-optics-integral-equation method for light scattering by nonspherical ice crystals, Appl. Opt., 35, 6568– 6584, https://doi.org/10.1364/AO.35.006568, http://opg.optica.org/ao/abstract.cfm?URI=ao-35-33-6568, 1996b.
 - Yang, P. and Liou, K.-N.: Light scattering and absorption by nonspherical ice crystals, pp. 31–71, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/3-540-37672-0_2, https://doi.org/10.1007/3-540-37672-0_2, 2006.
- Young, A. T.: Revised depolarization corrections for atmospheric extinction, Appl. Opt., 19, 3427–3428, 970 https://doi.org/10.1364/AO.19.003427, http://www.osapublishing.org/ao/abstract.cfm?URI=ao-19-20-3427, 1980.
 - Yurkin, M. A., Maltsev, V. P., and Hoekstra, A. G.: The discrete dipole approximation for simulation of light scattering by particles much larger than the wavelength, Journal of Quantitative Spectroscopy and Radiative Transfer, 106, 546–557, 2007.
 - Zöger, M., Afchine, A., Eicke, N., Gerhards, M.-T., Klein, E., McKenna, D. S., Mörschel, U., Schmidt, U., Tan, V., Tuitjer, F., Woyke, T., and Schiller, C.: Fast in situ stratospheric hygrometers: A new family of balloon-borne and airborne Lyman *α* photofragment fluorescence
- 975 hygrometers, Journal of Geophysical Research: Atmospheres, 104, 1807–1816, https://doi.org/https://doi.org/10.1029/1998JD100025, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/1998JD100025, 1999.

Stephens, G. L.: Cirrus, climate, and global change, Oxford University Press New York, 2002.

955

Stephens, G. L.: Cloud feedbacks in the climate system: A critical review, Journal of climate, 18, 237-273, 2005.

Sun, W., Fu, Q., and Chen, Z.: Finite-difference time-domain solution of light scattering by dielectric particles with a perfectly matched layer absorbing boundary condition, Applied optics, 38, 3141–3151, 1999.

- Sunilkumar, S. V. and Parameswaran, K.: Temperature dependence of tropical cirrus properties and radiative effects, Journal of Geophysical Research: Atmospheres, 110, https://doi.org/10.1029/2004JD005426, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/ 2004JD005426, 2005.
- Teri, M., Müller, T., Gasteiger, J., Valentini, S., Horvath, H., Vecchi, R., Bauer, P., Walser, A., and Weinzierl, B.: Impact of particle size,
 refractive index, and shape on the determination of the particle scattering coefficient–an optical closure study evaluating different nephelometer angular truncation and illumination corrections. Atmospheric Measurement Techniques, 15, 3161–3187, 2022.
 - Thomas, A., Borrmann, S., Kiemle, C., Cairo, F., Volk, C., Beuermann, J., Lepuchov, B., Santacesaria, V., Matthey, R., Rudakov, V., Yushkov, V., Mackenzie, A., and Stefanutti, L.: In Situ Measurements of Background Aerosol and Subvisible Cirrus in the Tropical Tropopause Region, Journal of Geophysical Research, v.107 (2002), 107, https://doi.org/10.1029/2001JD001385, 2002.
- 965 Thornberry, T. D., Rollins, A. W., Avery, M. A., Woods, S., Lawson, R. P., Bui, T. V., and Gao, R.-S.: Ice water content-extinction relationships and effective diameter for TTL cirrus derived from in situ measurements during ATTREX 2014, Journal of Geophysical Research: Atmospheres, 122, 4494–4507, https://doi.org/https://doi.org/10.1002/2016JD025948, https://agupubs.onlinelibrary.wiley.com/ doi/abs/10.1002/2016JD025948, 2017.
 - Wagner, S. W. and Delene, D. J.: Technique for comparison of backscatter coefficients derived from in situ cloud probe measurements
- 970 with concurrent airborne lidar, Atmospheric Measurement Techniques, 15, 6447–6466, https://doi.org/10.5194/amt-15-6447-2022, https: //amt.copernicus.org/articles/15/6447/2022/, 2022.
 - Weigel, R., Spichtinger, P., Mahnke, C., Klingebiel, M., Afchine, A., Petzold, A., Krämer, M., Costa, A., Molleker, S., Reutter, P., Szakáll, M., Port, M., Grulich, L., Jurkat, T., Minikin, A., and Borrmann, S.: Thermodynamic correction of particle concentrations measured by underwing probes on fast-flying aircraft, Atmospheric Measurement Techniques, 9, 5135–5162, https://doi.org/10.5194/amt-9-5135-2016, https://amt.copernicus.org/articles/9/5135/2016/, 2016.
- 975 https://amt.copernicus.org/articles/9/5135/2016/, 2016.
 - Weigel, R., Mahnke, C., Baumgartner, M., Dragoneas, A., Vogel, B., Ploeger, F., Viciani, S., D'Amato, F., Bucci, S., Legras, B., Luo, B., and Borrmann, S.: In situ observation of new particle formation (NPF) in the tropical tropopause layer of the 2017 Asian monsoon anticyclone – Part 1: Summary of StratoClim results, Atmospheric Chemistry and Physics, 21, 11689–11722, https://doi.org/10.5194/acp-21-11689-2021, https://acp.copernicus.org/articles/21/11689/2021/, 2021a.
- 980 Weigel, R., Mahnke, C., Baumgartner, M., Krämer, M., Spichtinger, P., Spelten, N., Afchine, A., Rolf, C., Viciani, S., D'Amato, F., Tost, H., and Borrmann, S.: In situ observation of new particle formation (NPF) in the tropical tropopause layer of the 2017 Asian monsoon anticyclone Part 2: NPF inside ice clouds, Atmospheric Chemistry and Physics, 21, 13455–13481, https://doi.org/10.5194/acp-21-13455-2021, https://acp.copernicus.org/articles/21/13455/2021/, 2021b.
 - Wernli, H., Boettcher, M., Joos, H., Miltenberger, A. K., and Spichtinger, P.: A trajectory-based classification of ERA-
- 985 Interim ice clouds in the region of the North Atlantic storm track, Geophysical Research Letters, 43, 6657–6664, https://doi.org/https://doi.org/10.1002/2016GL068922, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL068922, 2016.
 - Wex, H., Neusüß, C., Wendisch, M., Stratmann, F., Koziar, C., Keil, A., Wiedensohler, A., and Ebert, M.: Particle scattering, backscattering, and absorption coefficients: An in situ closure and sensitivity study, Journal of Geophysical Research: Atmospheres, 107, LAC-4, 2002.

Yang, P. and Liou, K.: Geometric-optics-integral-equation method for light scattering by nonspherical ice crystals, Applied Optics, 35,

- **990** 6568–6584, 1996a.
 - Yang, P. and Liou, K. N.: Geometric-optics-integral-equation method for light scattering by nonspherical ice crystals, Appl. Opt., 35, 6568– 6584, https://doi.org/10.1364/AO.35.006568, http://opg.optica.org/ao/abstract.cfm?URI=ao-35-33-6568, 1996b.
 - Yang, P. and Liou, K.-N.: Light scattering and absorption by nonspherical ice crystals, pp. 31–71, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/3-540-37672-0_2, https://doi.org/10.1007/3-540-37672-0_2, 2006.
- **995** Young, A. T.: Revised depolarization corrections for atmospheric extinction, Appl. Opt., 19, 3427–3428, https://doi.org/10.1364/AO.19.003427, http://www.osapublishing.org/ao/abstract.cfm?URI=ao-19-20-3427, 1980.
 - Yurkin, M. A., Maltsev, V. P., and Hoekstra, A. G.: The discrete dipole approximation for simulation of light scattering by particles much larger than the wavelength, Journal of Quantitative Spectroscopy and Radiative Transfer, 106, 546–557, 2007.
 - Zöger, M., Afchine, A., Eicke, N., Gerhards, M.-T., Klein, E., McKenna, D. S., Mörschel, U., Schmidt, U., Tan, V., Tuitjer, F., Woyke, T.,
- and Schiller, C.: Fast in situ stratospheric hygrometers: A new family of balloon-borne and airborne Lyman α photofragment fluorescence hygrometers, Journal of Geophysical Research: Atmospheres, 104, 1807–1816, https://doi.org/https://doi.org/10.1029/1998JD100025, https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/1998JD100025, 1999.



Figure 1. 2D histogram of Backscatter Ratio vs altitude. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 1. 2D histogram of Backscatter Ratio vs altitude. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin, representing 8477 data points.



Figure 2. 2D histogram of Total Particle Depolarization data vs temperature for altitudes above 11 km. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 2. 2D histogram of Total Particle Depolarization data vs temperature for altitudes above 11 km. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin, representing 2308 data point.



Figure 3. Upper panel: Red line, time series of particle backscattering coefficient β measured by MAS on 10 August 2017; Black solid line, best optical modelling match β_{NC} , error bars are displayed in the first part of the curve; black dashed lines, maximum and minimum values of the optical modelling β_{NC}^{AR} values. Lower panel: Red line, particle depolarization measured by MAS; Blue line Particle aspherical fraction measured by NIXE-CAPS; black dots, AF values of the best optical modeling match β_{NC} , the black line is a 5-points running average through them.



Figure 3. Upper panel: Red line, time series of particle backscattering coefficient β measured by MAS on 10 August 2017; Black solid line, best optical modelling match β_{NC} , error bars are displayed in the first part of the curve; black dashed lines, maximum and minimum values of the optical modelling β_{NC}^{AR} values. Lower panel: Red line, particle depolarization measured by MAS; Blue line Particle aspherical fraction measured by NIXE-CAPS; black dots, AF values of the best optical modeling match β_{NC} , the black line is a 5-points running average through them.



Figure 4. 2D histogram of particle backscatter coefficient observations from the backscattersonde MAS and the particle backscattering coefficients data from the two lidars MAL, pointing upward and downward, at 500m from the aircraft. The average of the two lidar backscattering coefficients has been used for the comparison. The data have been averaged over 60 s. In the graph are reported 500 data points acquired while crossing some of the thickest clouds observed during the campaign.


Figure 4. 2D histogram of particle backscatter coefficient observations from the backscattersonde MAS and the particle backscattering coefficients data from the two lidars MAL, pointing upward and downward, at 500m from the aircraft. The average of the two lidar backscattering coefficients has been used for the comparison. The data have been averaged over 60 s. In the graph are reported 500 data points acquired while crossing some of the thickest clouds observed during the campaign.



Figure 5. 2D histograms of occurrence of the measured and calculated backscattering coefficients. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 5. 2D histograms of occurrence of the measured and calculated backscattering coefficients. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 6. 2D histograms of occurrence of the measured and calculated particle depolarization. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 6. 2D histograms of occurrence of the measured and calculated particle depolarization. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 7. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle number concentration N. The black line represent the fit N=4.33·10² · $\beta^{1.32}$. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 7. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle number concentration N. The black line represent the fit N=3.96·10⁶ $\beta^{1.32}$. Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 8. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle mean radius R_{mean} . Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 8. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle mean radius R_{mean} . Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 9. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle effective radius R_{eff} . Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 9. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle effective radius R_{eff} . Data were acquired throughout the campaign by the MAS backscattersonde. The color codes the number of observations in the 2D bin.



Figure 10. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle Surface Area Density SAD. The black line represents the fit SAD= $3.24 \cdot 10^5 \cdot \beta^{1.11}$. The color codes the number of observations in the 2D bin.



Figure 10. 2D histogram of particle backscattering coefficient observations β from the backscattersonde MAS and particle Surface Area Density SAD. The black line represents the fit SAD=7.07.10⁸ $\cdot \beta^{1.11}$. The color codes the number of observations in the 2D bin.



Figure 11. 2D histogram of particle backscattering coefficient observations from the backscattersonde MAS and Ice Water Content IWC. The black line represents the fit IWC=1.17.10⁴ $\cdot \beta^{1.39}$. The color codes the number of observations in the 2D bin.



Figure 11. 2D histogram of particle backscattering coefficient observations from the backscattersonde MAS and Ice Water Content IWC. The black line represents the fit IWC= $1.78 \cdot 10^5 \cdot \beta^{1.39}$. The color codes the number of observations in the 2D bin.



Figure 12. Scatterplot of BR vs δ_T color coded in term of particle Aspherical Fraction.



Figure 12. Scatterplot of BR vs δ_T color coded in term of particle Aspherical Fraction.



Figure 13. Scatterplot of BR vs δ_T color coded in term of particle number concentration.



Figure 13. Scatterplot of BR vs δ_T color coded in term of particle number concentration.



Figure 14. Scatterplot of BR vs δ_T color coded in term of particle mean radius.



Figure 14. Scatterplot of BR vs δ_T color coded in term of particle mean radius.



Figure 15. Scatterplot of BR vs δ_T color coded in term of temperature.



Figure 16. 2D histogram of cumulative distributions of particle backscattering coefficient computation. The histogram shows the buildup of the particle backscattering coefficient with respect to the particle radius, for all the PSD under study. For the present graph, Mie scattering computation have been used.



Figure 15. Scatterplot of BR vs δ_T color coded in term of temperature.



Figure 16. 2D histogram of cumulative distributions of particle backscattering coefficient computation. The histogram shows the buildup of the particle backscattering coefficient with respect to the particle radius, for all the PSD under study. For the present graph, Mie scattering computation have been used.



Figure 17. 2D histogram of the ratio of the backscattering coefficient to IWC, versus particle mean radius. The black line report a regression computed for R_{mean} from 10 to 100 μ m. The fit is $\frac{beta}{IWC} = 1.52 \cdot 10^{-3} \cdot R_{mean}^{-2.34}$. The color codes the number of observations in the 2D bin.

Table 1. Linear relations linking the backscattering coefficient to particle number, surface (SAD) and Ice Water Content (IWC). Here β is expressed in km⁻¹ sr⁻¹ while N, SAD and IWC are expressed respectively in cm⁻³, μ m²cm⁻³, and mg m⁻³.

fit	R-squared
N=4.33.10 ² · $\beta^{1.32}$	0.71
$SAD = 3.24 \cdot 10^5 \cdot \beta^{1.11}$	0.73
IWC=1.17.10 ⁴ $\cdot \beta^{1.39}$	0.59



Figure 17. 2D histogram of the ratio of the backscattering coefficient to IWC, versus particle mean radius. The black line report a regression computed for R_{mean} from 10 to 100 μ m. The fit is $\frac{\beta}{IWC} = 1.53 \cdot R_{mean}^{-2.34}$, with an R-squared of 0.54. The color codes the number of observations in the 2D bin.



Figure 18. Scatterplot of measured IWC vs estimated extinction $\sigma = 30\beta$. The solid lines represent regressions from i. the present work, black; ii. Heymsfield et al. (2005), purple; iii. Avery et al., (2012), brown; Heymsfield et al. (2014), (a) yellow; (b) green; Thornberry et al. (2017), blue. Experimental points are color-coded in temperature of the observation.



Figure A1. Particle backscattering coefficient vs particle radius, for various choices of the Aspect Ratio. A particle concentration of 1 cm^{-3} has been considered. The black line refers to Mie computations (AR=1). For particle radius below 14 μ m, the computations were performed with the GRASP package. Beyond 14 μ m, for every given AR, the backscattering coefficients have been computed from Mie backscattering efficiencies, suitably rescaled with a constant factor. This scaling factor was chosen to make the scaled Mie efficiency to overlap with the GRASP T-matrix efficiency in the particle radius dimensional region from 5 to 14 μ m (60-180 size parameters).



Figure A2. Particle depolarization vs particle radius, for various choices of the Aspect Ratio. The black zero depolarization line refers to Mie computations (AR=1). For particle radius below 14 μ m, the computations were performed with the GRASP package. Beyond 14 μ m, for every given AR, the depolarization of the particles was set at its constant, asymptotic value, computed as its mean over the the particle radius dimensional region from 5 to 14 μ m (60-180 size parameters)

Table 1. Parametrizations linking the backscattering coefficient to particle number, surface (SAD) and Ice Water Content (IWC). Here β is expressed in m⁻¹ sr⁻¹ while N, SAD and IWC are expressed respectively in cm⁻³, μ m²cm⁻³, and g m⁻³.

fit	R-squared
N=3.96·10 ⁶ · $\beta^{1.32}$	0.71
SAD=SAD=7.07 $\cdot 10^8 \cdot \beta^{1.11}$	0.73
IWC= $1.78 \cdot 10^5 \cdot \beta^{1.39}$	0.59



Figure A3. 2D histogram of the particle concentration N_{ice} computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). For NIXE-CAPS only particles with with lower size limit at 1.5 μ m have been considered.



Figure A4. 2D histogram of the particle mean radius R_{mean} computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). For NIXE-CAPS only particles with with lower size limit at 1.5 μ m have been considered.

Table 2. IWC- σ parameterizations (adapted from Thornberry et al. (2017)). IWC is expressed in $g \cdot m^{-3}$, σ in m^{-1}

IWC - σ parametrization		
Reference	Functional Form	T range
Heymsfield et al. (2005)	IWC=119 $\cdot \sigma^{1.22}$	198-263 K
Avery et al. (2012)	IWC=238 $\cdot \sigma^{1.22}$	-
Heymsfield et al. (2014) (a)	IWC=a· σ^b a=0.00532 · $(T - 183)^{2.55}$, b=1.31· $e^{0.0047 \cdot (T - 273)}$	188-270 K
Heymsfield et al. (2014) (b)	$IWC = \sigma \cdot (0.91/3) \cdot 91744 \cdot e^{0.177 \cdot (T-273)}$ $IWC = \sigma \cdot (0.91/3) \cdot 83.3 \cdot e^{0.0184 \cdot (T-273)}$	202-217 K 188-202 K
Thornberry et al. (2017)	$IWC = \sigma \cdot (0.92/3) \cdot (40 + 0.53 \cdot (T - 192))$ $IWC = \sigma \cdot (0.92/3) \cdot (12 + 28 \cdot e^{0.65 \cdot (T - 192)})$	192-207 K 185-192 K
Present Work	IWC=1552 $\cdot \sigma^{1.39}$	-

Table 3. List of Instrument

Instrument	Parameter	Technique	Range	Sensitivity	Resolution
ССР	Particle Size Distribution	Laser Optical Particle Counter	Particle Diameter 3-47 μ m Concentration < 2000 cm ⁻³	Single Particle Detection	1 s
CIP	Particle Size Distribution	Laser Optical Particle Imager	Particle Diameter 25-1550 μ m Concentration < 500 cm ⁻³	Single Particle Detection	1 s
NIXE-CAPS	Particle Size Distribution Particle Asphericity	Laser Particle Spectrometer	Particle Diameter 0.61-937 μ m Concentration < 2000 cm ⁻³	Single Particle Detection	1 s
MAS	Backscattering Depolarization in-situ	Laser elastic scattering	backscatter coefficient β $5 \cdot 10^{-10} - 10^{-4}m^{-1}sr^{-1}$ volume depolarization δ 0-100 %	backscatter coefficient β $5 \cdot 10^{-10} m^{-1}$ volume depolarization δ 1 %	<u>10 s</u>
MAL	Backscattering Depolarization remote sensing	Laser elastic scattering	backscatter coefficient β 0.3 - 80 $\cdot 10^{-7}m^{-1}sr^{-1}$ volume depolarization δ 0-100 %	backscatter coefficient β $0.3 \cdot 10^{-7} m^{-1} sr^{-1}$ volume depolarization δ 2% nighttime 2 km above/below 19 km	120 s
FLASH	Total Water	Lyman- α	1-1000 ppmv	0.1 ppmv	1 s
FISH	Gas Phase Water	Lyman- α	1-1000 ppmv	0.1 ppmv	1 s



Figure A5. 2D histogram of the backscattering coefficient computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). For NIXE-CAPS only particles with with lower size limit at 1.5 μ m have been considered. Computation has been performed with Mie scattering codes.



Figure A6. Red line, time serie of particle backscattering coefficient β measured by MAS on 10 August 2017; Black solid line, β_{NC} corresponding to the best matching between measured δ and computed δ_{NC}^{AR} (not displayed), error bars are reported in the first part of the curve; black dashed lines, maximum and minimum values of the optical modeling β_{NC}^{AR} values. Lower panel: Red line, particle depolarization measured by MAS; Blue line Particle aspherical fraction measured by NIXE-CAPS; black line, AF values of the best matching between measured δ and computed δ_{NC}^{AR} .

Table 4. List of achronyms used in the text

Achronym	Definition
AF	Aspherical Factor
AR	Aspect Ratio
BR	Backscatter Ratio
β	backscattering coefficient
CALIOP	Cloud and Aerosol Lidar with Orthogonal Polarization
CALIPSO	Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation
CAS-DPOL	Cloud and Aerosol Spectrometer with detector for polarization
ССР	Cloud Combination Probe
CDP	Cloud Droplet Probe
ChiWIS	Chicago Water Isotope
CIPgs	Cloud Imaging Probe grey scale
δ	depolarization
DDA	Discrete Dipole approximation
DMT	Droplet Measurement Technologies
ERA5	ECMWF Reanalysis version 5
FISH	Fast In situ Stratospheric Hygrometer
FLASH	FLuorescent Airborne Stratospheric Hygrometer
GOIE	geometric optics integral equation
GOM	geometric optics method
GRASP	Generalized Retrieval of Aerosol and Surface Properties
IGOM	improved geometric optics method
FDTD	Finite Difference Time Domain
IIR	Infrared Imaging Radiometer
IWC	Ice Water Content
N_{ice}	ice particle concentration
MAL	Miniature Aerosol Lidar
MAS	Multiwavelenght Aerosol Scatterometer
NASA	National Aeronautics and Space Administration
NIXE-CAPS	New Ice eXpEriment - Cloud and Aerosol Particle Spectrometer
PSD	Particle Size Distribution
R_{eff}	effective radius
R_{mean}	mean radius
RECONCILE	Reconciliation of essential process parameters for enhanced predictability of Arctic stratospheric ozone loss and climate interactions
σ	extinction coefficient
SAD	Surface Area Density
VD	Volume Density



Figure A1. Particle backscattering coefficient vs particle radius, for various choices of the Aspect Ratio. A particle concentration of 1 cm^{-3} has been considered. The black line refers to Mie computations (AR=1). For particle radius below 14 μ m, the computations were performed with the GRASP package. Beyond 14 μ m, for every given AR, the backscattering coefficients have been computed from Mie backscattering efficiencies, suitably rescaled with a constant factor. This scaling factor was chosen to make the scaled Mie efficiency to overlap with the GRASP T-matrix efficiency in the particle radius dimensional region from 5 to 14 μ m (60-180 size parameters).



Figure A2. Particle depolarization vs particle radius, for various choices of the Aspect Ratio. The black zero depolarization line refers to Mie computations (AR=1). For particle radius below 14 μ m, the computations were performed with the GRASP package. Beyond 14 μ m, for every given AR, the depolarization of the particles was set at its constant, asymptotic value, computed as its mean over the the particle radius dimensional region from 5 to 14 μ m (60-180 size parameters)


Figure A3. 2D histogram of the backscattering coefficient computed with PSD with lower size limit at 0.3 μ m (y-axis) and at 1.5 μ m (x-axis). Spherical ice approximation has been used in the computation.



Figure A4. 2D histogram of the particle concentration N_{ice} computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). For NIXE-CAPS only particles with with lower size limit at 1.5 μ m have been considered.



Figure A5. 2D histogram of the particle mean radius R_{mean} computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). For NIXE-CAPS only particles with with lower size limit at 1.5 μ m have been considered.



Figure A6. 2D histogram of the backscattering coefficient computed from PSD from NIXE-CAPS (x-axis) and CCP (y-axis). For NIXE-CAPS only particles with with lower size limit at 1.5 μ m have been considered. Computation has been performed with Mie scattering codes.



Figure A7. Scatterplot of the Total Particle Depolarization vs the Aspect Ratio, colour coded in terms of the backscatter coefficient.



Figure A8. Red line, time serie of particle backscattering coefficient β measured by MAS on 10 August 2017; Black solid line, β_{NC} corresponding to the best matching between measured δ and computed δ_{NC}^{AR} (not displayed), error bars are reported in the first part of the curve; black dashed lines, maximum and minimum values of the optical modeling β_{NC}^{AR} values. Lower panel: Red line , particle depolarization measured by MAS; Blue line Particle aspherical fraction measured by NIXE-CAPS; black line, AF values of the best matching between measured δ and computed δ_{NC}^{AR} .



Figure A7. 2D histogram of the backscattering coefficient computed with PSD with lower size limit at 0.3 μ m (y-axis) and at 1.5 μ m (x-axis)



Figure A8. 2D histogram of Particle Aspherical Fraction (y-axis) vs Particle depolarization (x-axis)



Figure A9. Scatterplot of estimated vs measured IWC using regressions from the present work (left panel), from Heymsfield et al. (2014) (middle panel) and from Heymsfield et al. (2005) (right panel). The colors code the temperature at the observation.



Figure A9. Red line, time serie of particle backscattering coefficient β measured by MAS on 10 August 2017; Black solid line, β_{NC} corresponding to the best matching between measured δ and computed δ_{NC}^{AR} (not displayed), error bars are reported in the first part of the curve; black dashed lines, maximum and minimum values of the optical modeling β_{NC}^{AR} values. Lower panel: Red line , particle depolarization measured by MAS; Blue line Particle aspherical fraction measured by NIXE-CAPS; black line, AF values of the best matching between measured δ and computed δ_{NC}^{AR} .



Figure A10. 2D histogram of Particle Aspherical Fraction (y-axis) vs Particle depolarization (x-axis)