



How the extreme 2019-2020 Australian wildfires affected global circulation and adjustments

Fabian Senf¹, Bernd Heinold¹, Anne Kubin¹, Jason Müller¹, Roland Schrödner¹, and Ina Tegen¹

¹Leibniz Institute for Tropospheric Research, Leipzig, Germany.

Correspondence: Fabian Senf (senf@tropos.de)

Abstract. Wildfires are a significant source of absorbing aerosols in the atmosphere. Especially extreme fires, such as those during the 2019-2020 Australian wildfire season (Black Summer fires), can have considerable large-scale effects. In this context, the climate impact of extreme wildfires not only unfolds because of the emitted carbon dioxide, but also due to smoke aerosol released as high as the stratosphere. The overall aerosol effects depend on a variety of factors, such as the amount emitted, the injection height, and the composition of the burned material, and is therefore subject to considerable uncertainty. In the present study, we address the global impact caused by the exceptionally strong and high-reaching smoke emissions from the Australian wildfires using simulations with a global aerosol-climate model. We show that the absorption of solar radiation by the black carbon contained in the emitted smoke led to a shortwave radiative forcing of more than $+5 \text{ W m}^{-2}$ in the southern mid-latitudes of the lower stratosphere. Subsequent adjustment processes in the stratosphere slowed down the diabatically driven meridional circulation, thus redistributing the heating perturbation on a global scale. As a result of these stratospheric adjustments, a positive temperature perturbation developed in both hemispheres leading to additional longwave radiation emitted back to space. According to the model results, this adjustment occurred in the stratosphere within the first two months after the event. At the top of atmosphere (TOA), the net effective radiative forcing (ERF) in the southern hemisphere was initially dominated by the instantaneous positive radiative forcing of about $+0.5 \text{ W m}^{-2}$, for which the positive sign resulted mainly from the presence of clouds above the Southern Ocean. The longwave adjustments led to a compensation of the initially net positive TOA ERF, which is seen in the southern hemisphere, the tropics and the northern mid-latitudes. The changes in the lower stratosphere also affected the upper troposphere through a thermodynamic downward coupling mechanism in the model. Subsequently, increased temperatures were also obtained in the upper troposphere, causing a decrease in relative humidity, cirrus amount, and the ice water path. As a result, surface precipitation also decreased, which was accompanied by a weakening of the tropospheric circulation due to the given energetic constraints. In general, it appears that the radiative effects of smoke from single extreme wildfire events can lead to global impacts that affect the interplay of tropospheric and stratospheric cycles in complex ways. This emphasizes that future changes in extreme wildfires need to be included in projections of aerosol radiative forcing.



1 Introduction

25 Atmospheric aerosol particles from both anthropogenic and natural sources are important climate factors. The magnitudes of the different aerosol effects remain, however, highly uncertain (see Forster et al. (2021)). Aerosols influence the atmospheric radiation balance in different ways. The "direct radiative effect" refers to the change in the radiation budget of the Earth due to the scattering and absorption of incident solar radiation. An "indirect effect" of aerosol particles, on the other hand, results from the influence of aerosol particles as condensation nuclei for the formation of cloud droplets or for cloud ice and precipitation formation. While the overall average effect of aerosol particles on the radiation balance and surface temperatures is negative leading to reduction of surface temperatures (Forster et al., 2021; Thornhill et al., 2021), certain aerosol types, such as black carbon or soot particles, which originate from incomplete combustion processes and strongly absorb light in the spectral range of incident solar radiation, cause a positive radiative effect that counteracts the cooling caused by the non-absorbing aerosol types. This absorption of radiation results in heating rate changes within the aerosol-containing atmospheric layers. In turn, this heating causes rapid atmospheric adjustments to the instantaneous aerosol forcing, and has the potential to alter atmospheric dynamics and circulation. Such atmospheric adjustments are complex and include the effects of changes in heating rates on boundary layer stability or cloud cover. The various effects of the rapid adjustments or "semi-direct" absorbing aerosol effects have been summarized by Koch and Del Genio (2010). They can be studied with atmospheric circulation models on different scales that take into account the distribution of aerosol radiative properties. As an example, for a high resolution model experiment with grid resolutions of 312 and 625 m for Germany, Senf et al. (2021) found that the adjustment to heating by absorbing aerosol contributed to a positive radiative forcing of similar magnitude as the instantaneous radiative forcing. Apart from changes in radiation budget forcing, such adjustments can also affect atmospheric dynamics and circulation. As an example for changes with global scale impacts, the effects of absorbing aerosol over the Indian subcontinent on the monsoon circulations have been analyzed in several model studies (Sanap and Pandithurai, 2015).

45 Carbonaceous aerosol is produced in the industrial and agricultural regions mainly by burning fossil fuels or biomass (Bond et al., 2013). Sources comprise industrial and domestic emissions and traffic on the one hand, and vegetation fires including crop residue burning on the other hand. Climate change has been recognized as a new major driver for fire regimes worldwide (Jolly et al., 2015; Bowman et al., 2020). In addition to their immediate local disruptions of nature and lives and release of CO₂, wildfires produce carbonaceous particles that impact on the radiation balance (Sokolik et al., 2019). While anthropogenic aerosol from fossil fuel burning usually remains within the boundary layer, smoke from wildfires containing BC aerosol can be released into the atmosphere in several kilometers high, and thus is subject to long-distance transport in the free troposphere (Huang et al., 2015). This is especially the case for high-intensity wildfires leading to deep convection due to the fire heat and eventually pyrocumulonimbus (pyroCb) formation (Fromm et al., 2010, 2019), potentially injecting smoke aerosols as high as the lower stratosphere. Such deep pyroconvection from extreme wildfires can have similar atmospheric effects as aerosol from strong volcanic eruptions where the volcanic eruption plume reaches the stratosphere (Peterson et al., 2018).

55 Exceptionally strong vegetation fires occurred in Southeastern Australia in December and January 2019-2020 during the so-called Australia's Black Summer. During that fire season such pyroconvective smoke plumes were observed, injecting massive



smoke amounts into the lower stratosphere (Kablick et al., 2020; Khaykin et al., 2020). Mainly between 29 December 2019 and 4 January 2020, several intense pyroCb events transported between 0.3 and 2 Tg smoke particles (Peterson et al., 2021; Hirsch and Koren, 2021) up to 14–16 km height (Kablick et al., 2020; Boone et al., 2020). The resulting smoke layer extended across the southern mid and high latitudes, and could be detected in the stratosphere for as long as two years after the event (Ohneiser et al., 2022a). It had significant effects on the radiation budget. Its instantaneous positive radiative forcing in the southern hemisphere was estimated as high as $+0.5 \text{ W m}^{-2}$ by Heinold et al. (2022). The actual radiative effect by the smoke aerosol would however be moderated by longwave adjustments in the stratosphere (Yu et al., 2021; Liu et al., 2022).

Beside the above mentioned value, a variety of other results for radiative forcing (RF) by Australian smoke can be found in the current literature. Reported values range from around 0.8 W m^{-2} (cloud case in Sellitto et al., 2022) to -1.0 W m^{-2} (Hirsch and Koren, 2021). It has been argued by Sellitto et al. (2022) that the large spread can be attributed to uncertainties in the optical properties of the smoke, i.e., absorptivity and backscattered fraction. In our study, we aim to show in a complementary manner that the different assumptions used to estimate the radiative forcing, i.e., the consideration of cloud effects (Heinold et al., 2022), the contribution of adjustments to the forcing (Yu et al., 2021; Liu et al., 2022), and the spatial extent over which the averaged values were collected (applies e.g. to Hirsch and Koren, 2021; Liu et al., 2022), have significant influence on the estimated magnitude of the forcing value or even the sign of the radiative forcing. The dynamical mechanisms that led to the formation of a synoptic-scale anticyclonic smoke-induced vortex appear to be well understood (Khaykin et al., 2020; Kablick et al., 2020; Allen et al., 2020; Doglioni et al., 2022), but it is unclear how the global dispersion of smoke impacted on the global circulation of the stratosphere and whether the perturbations in the lower stratosphere could have affected the upper troposphere. This chain of effects could represent an important mechanism for the downward coupling between the middle and lower atmosphere and help to better understand the climate effects due to extreme fire events.

We address these issues in this study using different global modeling approaches, which are described in Section 2. Emission characteristics are prescribed in our global modeling setup such that smoke is introduced into the lower stratosphere during the Australian pyroCb events (Section 2.2.1). To describe global smoke dispersion and understand emerging effects and couplings, both nudged simulations that relax the flow pattern towards an observation-based reanalysis (Section 2.2.2) and free-running ensemble simulations are conducted (Section 2.2.3). With the help of the simulation data, the regional and global radiative forcing is estimated and analyzed in Section 3.1 and compared with existing findings from the literature (Section 3.1.2). Furthermore, adjustments of the thermodynamic structure and the global circulation of the stratosphere (Section 3.2) and adjustments in the troposphere with respect to the hydrological cycle (3.3) are analyzed. In Section 4 we discuss uncertainties, limitations and perspectives of our study and conclude in Section 5 with a summary and final remarks.

2 Data & Methods

2.1 Global Model ECHAM6.3-HAM2.3

ECHAM6.3-HAM2.3 is a global state-of-the-art aerosol-climate model (Tegen et al., 2019). In this study, we use the same model configuration already detailed in Heinold et al. (2022). Global simulations were initialized on 1st October 2019 and

integrated forward until 31st March 2020, i.e. six months ahead. The global model is set up and tuned for a horizontal resolution of approximately $1.875^\circ \times 1.875^\circ$ which results from the spectral truncation at T63. In the vertical, 47 levels are used from ground to 0.01 hPa. Model layers stretch in the vertical having around 70 m layer thickness close to ground and more than one kilometer in the lower stratosphere. This vertical resolution is quite coarse for resolving stratospheric processes, and a better vertical refinement of the middle atmosphere is envisioned for future studies.

The aerosol module HAM2.3 applies the aerosol microphysics model M7 (Vignati et al., 2004) for the prediction of several aerosol species, including sea salt, sulfate, mineral dust, organic carbon (OC) and black carbon (BC). The latter two are of special interest for the radiative impact of the investigated wildfire emissions. The aerosol microphysics scheme treats source processes (e.g. from fire emissions), transformations due to microphysics and atmospheric chemistry as well as loss processes, e.g. due to scavenging by clouds and precipitation.

Aerosol particles impact atmospheric radiation fluxes. The particle optical properties are interactively calculated during model run time, taking into account their respective size and chemical composition. The aerosol particles are treated as internal mixtures and optical aerosol properties such as extinction coefficient, single-scattering albedo, and asymmetry parameter are input from a pre-calculated lookup table.

Atmospheric radiation is computed within ECHAM6.3 using the PSrad/RRTMG (Rapid Radiative Transfer Model for GCMs; Iacono et al., 2008; Pincus and Stevens, 2013) radiation package. 30 spectral bands are considered with 16 in the shortwave part of the spectrum, mainly influenced by solar radiation, and 14 in the longwave part of the spectrum, mainly influenced by terrestrial radiation. In addition to aerosol particles, also optics of various gaseous atmospheric constituents and different hydrometeor types are also considered.

2.2 Setup of Global Simulation Experiments

2.2.1 Prescribed Wildfire Emissions

As in Heinold et al. (2022), the chosen global model setup applies prescribed wildfire emissions from the Global Fire Assimilation System (GFAS; Kaiser et al., 2012). GFAS assimilates observations of fire radiative power (FRP) based on MODIS instruments aboard the polar orbiting NASA satellites Terra and Aqua to retrieve estimates of wildfire and biomass burning emissions on a daily basis. The GFAS emission data are input into the model as external data and mapped onto sulfate, dimethyl sulfide (DMS), OC and BC.

Wildfire aerosols are emitted mainly in the boundary layer and directly above it in the default model configuration. However, as argued in Heinold et al. (2022), the exceptional pyroCb activity of the extreme Australian fire event required to prescribe that most of the fire aerosol had to be injected at tropopause levels. Therefore, this study uses the approach that during the two phases (29–31 December 2019 and 4 January 2020) with strong pyroCb activity (as reported by Kablick et al. (2020)), the fire aerosol is directly introduced in the model layer above the tropopause with emitted aerosol masses of 0.6 and 0.2 Tg, respectively, according to GFAS analysis. Around 6 to 7% of the total carbon load were attributed to highly absorbing black carbon. In the terminology of Heinold et al. (2022), the chosen emission setup is referred to as “TP+1”. Equivalent aerosol

effects were documented by Heinold et al. (2022) with other emission height specifications as long as most of the emitted
125 absorbing aerosol is injected above the tropopause.

In order to make more robust statements about the resulting global fire effects, simulations were not only performed with
the original GFAS emissions. As an extension, the fire emission strength during the pyroCb events was artificially increased
by the factors 2, 3 and 5. Corresponding increases concern both organic carbon and black carbon emissions. The emission
increases impact transport patterns and aerosol radiative forcings. The resulting additional simulation experiments were named
130 according to the respective scaling factors with FIRE2, FIRE3, and FIRE5. Further experiments include FIRE1, in which the
original GFAS emissions are used without scaling, and FIRE0 with zero Australian pyroCb emissions that is used as a reference
to analysis the overall perturbations by the Australian wildfires.

Assuming that the perturbations behave mainly linearly, the experiments with different scaling factors can be compared.
For this purpose, rescaling is applied, in which fire-induced perturbations are divided by the fire scaling factor, e.g. by two
135 for FIRE2 vs. FIRE0, and, if desirable, all rescaled experiments are subsequently averaged into a composite value to increase
statistical confidence.

2.2.2 Nudged Experiments

An important piece of evidence in our study originates from so-called *nudged experiments*. For nudging, data from the ECMWF
ERA5 reanalysis (Hersbach et al., 2020) were input as external data. Additional tendencies are introduced into the momentum
140 equations such that the horizontal winds simulated by ECHAM are relaxed towards ERA5 winds. Surface pressure is also
dragged towards ERA5 reanalyses. This strategy is applied to keep the model as close as possible to the observed meteorology.
It is expected that more realistic transport pathways result in the nudged experiments especially because prescribed fire emission
agree well with simulated fire weather. Temperature fields were chosen not to be directly impacted by nudging in accordance
to Zhang et al. (2014), thus keeping the energy budget free of artificial source and sink terms.

The nudged experiments in our study are equivalent to the "replay" simulations mentioned in Doglioni et al. (2022) for
145 the investigation of regional dynamical effects of absorbing smoke plumes. Similarly to Doglioni et al. (2022), freely running
simulations were conducted for each experiment type, which will be discussed and contrasted with respect to their dynamical
responses in the next subsection.

2.2.3 Ensemble Experiments

While the nudged simulations ensure a certain degree of realism of the resulting aerosol transport, they have their limitation
150 for studying the effects of the injected wildfire aerosol on the atmospheric dynamics. These can only be evaluated from freely
running *ensemble simulations*. However, this comes at the expense of probable discrepancies between the atmospheric state
and the aerosol emission. As an example, the model might simulate rainy weather in South East Australia at the dates of the
major wildfires and pyroCb events. Nonetheless, in order to obtain robust results of the dynamic response to the additional
155 heating in the lower stratosphere using a freely running model, an ensemble approach was chosen.



Technically, all ensemble members share the same initial conditions as they were restarted from the nudged simulation on 1 Oct 2019. Tiny differences between the members are introduced by a variation (of order 10^{-6}) of a parameter that governs the upward increase in stratospheric horizontal diffusion between adjacent layers, thus slightly perturbing mixing in the upper model layers. This perturbation does not alter the model climate, but introduces ensemble spread that fully develops within a few weeks after initialization.

For every fire scaling factor (0,1,2,3,5) an ensemble with 36 members was set up. Every member was run from October 2019 to March 2020 with the first two months being considered as spin-up time and, hence, excluded from the analyses.

All simulations, nudged and ensemble simulations share in common that prescribed sea surface temperatures and sea-ice concentrations were used at the lower boundary. These data were taken from the Atmospheric Model Intercomparison Project (AMIP) (Giorgetta et al., 2012). Long-lived greenhouse gases were defined following the Representative Concentration Pathway (RCP) 4.5 scenario.

2.3 Circulation Analysis

In the troposphere, the meridional overturning circulation is mainly fueled by strong meridional temperature gradients and is thus called thermally driven. In contrast, in the stratosphere the main driver that continuously induces a meridional transport of air is momentum deposition by breaking waves (Holton, 1997; Butchart, 2014). This is a key difference to the tropospheric circulation. As it turns out, the conventional Eulerian mean of motion does not sufficiently take into account this additional eddy transport (Butchart, 2014; Matsuno, 1980), also known as Stokes drift (Dunkerton, 1978; McIntyre, 1980).

As a result of this under-representation of eddy effects, the circulation, calculated with the Eulerian mean, deviates strongly from the Lagrangian circulation deduced from tracer and mass transport in the stratosphere. A very popular and well established solution to this problem is the so-called Transformed Eulerian Mean (TEM), derived by Andrews and McIntyre (1976, 1978) as reviewed by Butchart (2014).

The TEM circulation can be understood as the part of the meridional Eulerian mean circulation that is not balanced by the Stokes drift and it is, thus, also referred to as the residual circulation (Butchart, 2014). Defined in this way, the calculated residual circulation is in much closer agreement with the Lagrangian mean paths of tracers compared to the conventional Eulerian mean whenever eddy activity is important. This makes the TEM a very well suited tool for the investigation of the circulation as well as possible circulation adjustments in the stratosphere. Following Peixoto and Oort (1992) and Grotjahn (1993), the residual wind speeds \bar{v}^* and $\bar{\omega}^*$ are defined as

$$\bar{v}^* = \bar{v} - \partial_p (\overline{v'\theta'} \Gamma^{-1}), \quad (1)$$

$$\bar{\omega}^* = \bar{\omega} + \frac{\partial_\varphi (\cos \varphi \overline{v'\theta'} \Gamma^{-1})}{a \cos \varphi} \quad (2)$$

where \bar{v} and $\bar{\omega}$ are the zonally and monthly averaged meridional and vertical wind speeds (Eulerian sense), respectively. p denotes atmospheric pressure, θ is the potential temperature, stability is related to $\Gamma = \partial_p \bar{\theta}$, φ is latitude, ∂_p and ∂_φ are shortcuts for partial derivatives and a the radius of the Earth.



Using these residual velocities one can now define a residual meridional mass stream function to quantify and display the stratospheric meridional overturning circulation e.g (Peixoto and Oort, 1992):

$$190 \quad \bar{\psi}^* = 2\pi a \cos(\varphi) g^{-1} \int_p^{p_0} \bar{v}^* \partial p', \quad (3)$$

where g is the acceleration due to gravity. The following section takes a closer look at the impact of the Australian wildfires on the atmospheric state including the circulation variables $\bar{\omega}^*$, \bar{v}^* and $\bar{\psi}^*$.

3 Results

3.1 Radiative Perturbations

195 3.1.1 Indications of Global Adjustments to Smoke-Radiation Interactions

The Australian fire outbreak led to a significant input of black (BC) and organic carbon (OC) aerosol into the lower stratosphere. Estimates range from 0.3 to 2.1 ± 1.1 Tg (Khaykin et al., 2020; Peterson et al., 2021; Hirsch and Koren, 2021) which accommodates well the total emitted amount of 0.8 Tg used in our study. Once emitted from Southeast Australian fire hot spots, the smoke aerosol began to spread over the entire southern hemisphere reaching the southern tip of America after around ten
200 days (Ohneiser et al., 2020). The sudden increases in OC/BC burden lead to a significant absorption of shortwave radiation in the middle atmosphere (in our case 200 hPa to TOA) that mainly localized over the Southern Pacific Ocean during January 2020 (see Fig. 1a and d) with large-scale average values of shortwave radiation flux convergence larger than 5 W m^{-2} . Atmospheric mixing and transport lead to a more equal redistribution of stratospheric smoke aerosol in the subsequent months (Fig. 1b and c). Simulations with nudged meteorology indicate that significant amounts of smoke have been transported polewards with
205 the implications that the bright Antarctic surface may have been darkened by stratospheric smoke when viewed from TOA. However, the poleward transport pathway is not as clearly visible in the ensemble simulations (Fig. 1d-f). This could be explained by the fact that, firstly, the fire emissions in the variety of ensemble realizations were not necessarily carried out during suitable fire weather conditions, but imposed at externally determined times independent of large-scale conditions. Secondly, it is plausible that the coarse horizontal and vertical resolution of the climate model does not well represent the formation and
210 movement of the smoke-induced vortex, as detailed in Khaykin et al. (2020), such that the flow field that is super-imposed by nudging is able to more realistically represent the smoke transport. An extended discussion of this aspect can be found in Sect. 4.

The average longwave forcing of the middle atmosphere (see Fig. 1g-l) appears to be weaker in amplitude than the shortwave forcing. It is mainly the nudged runs that show a global negative longwave radiation response. The maximum amplitude
215 was reached in February 2020 (Fig. 1h). The ensemble simulations confirm a negative longwave response in the southern hemisphere that appears to be weaker than the corresponding shortwave forcing. Towards February 2020, spatial areas increase in size in which nudged as well as ensemble simulations agree in the negative sign of the stratospheric longwave response,



220

also in the northern hemisphere. We interpret this signal as a fingerprint of an interhemispheric change in the stratosphere that can only be realized by changes in the global circulation pattern and thus need to be interpreted as global adjustments to stratospheric smoke-radiation interactions.




Figure 1. Convergence of monthly average radiative flux perturbations (difference between FIRE1 and FIRE0) between TOA and 200 hPa, abbreviated as middle atmosphere (MATM) forcing. The upper two rows from (a) to (f) present shortwave MATM forcing, the lower two rows from (g) to (l) are longwave MATM forcing. Time is increasing from left to right as indicated in the sub-panel titles, from January to March 2020. Average results from nudged (1st and 3rd row) and ensemble (2nd and 4th row) simulations are compared. Hatching represents areas in which the ensemble mean perturbations are different from zero at 95% confidence level.

The situation becomes even more clear when larger smoke perturbations are considered (see Fig. 2). Increasingly larger radiative perturbations result from smoke emissions scaled with factors 1, 2, 3 and 5 (see Sect. 2.2.1). The time-average radiative flux convergence in the middle atmosphere is dominated by shortwave absorption in the southern hemisphere between



60 and 20°S (Fig. 2d). The stratosphere responds with a negative longwave flux convergence that appears to be rather constant
225 south of 45°S (Fig. 2e). However, we even find negative values of longwave forcing up to around 60°N. In the net, a positive
forcing still localizes at southern mid-latitudes, however, with a significantly smaller amplitude. This fact again indicates that
the highly localized shortwave forcing is compensated to a considerable degree by a non-local (w.r.t. the smoke location)
longwave forcing. In comparison to the middle atmosphere, forcings originating from the lower atmosphere are relatively
small (Fig. 2g-j). The only component that contains perturbation signals is the longwave forcing. Due to the high uncertainties,
230 it remains speculative whether a positive longwave forcing from the lower atmosphere compensates for some of the negative
stratospheric forcing in the southern mid-latitudes. At the Earth's surface, dimming of solar radiation due to extinction by
smoke is predominant in the southern mid-latitudes and has maximum amplitude between 40 and 50°S. In total, the shortwave
TOA flux is positive (Fig. 2a) because the absorbed flux in the middle atmosphere more than offsets the reduction of shortwave
fluxes at the surface. In relative terms, this reduction appears to be much smaller than reported elsewhere in literature (Hirsch
235 and Koren, 2021; Yu et al., 2021; Liu et al., 2022), however, we will show in the next section that the presence of clouds is
of major importance for the quantification of surface and TOA forcing of the Australian smoke event. The negative longwave
TOA flux compensates the shortwave TOA forcing such that in the net, the radiative forcing cannot be differentiated from zero
with confidence (Fig. 2b and c).

3.1.2 Effective Radiative Forcing of Smoke and the Role of Clouds

240 In the following, we quantify and discuss the effective radiative forcing of the Australian smoke event based on a composite of
all global simulation experiments where rescaled perturbations from different fire emission cases have been averaged. For this,
linearity of the atmospheric responses as assumed and perturbations divided by the respective fire scaling factor are averaged.
The composite average across all experimental setups allows for a much more robust assessment of underlying effects, although
non-linear responses may become more likely when fire emission strength is successively increased.

245 The effective radiative forcing and its components, instantaneous radiative forcing (IRF) and adjustments (Adj) are presented
in Fig. 3a for all-sky conditions. In the southern hemisphere, the shortwave ERF is around 0.5 W m^{-2} and dominated by
shortwave IRF. Adjustments become more and more important in the shortwave effects over time and contribute to around 30%
of the total ERF in March 2020. In contrast, longwave ERF is dominated by adjustments and maximizes between February and
March 2020. In comparison with the nudged simulations, the shortwave IRF appears to be smaller in the ensemble simulations
250 due to less aerosol transport towards the southern high latitudes (see also Fig. 1a vs. d). In the temporal evolution of shortwave
ERF, this mismatch seems to be compensated by stronger and more positive shortwave adjustments in the ensemble runs. In
the tropics, ERF and its components are weak and uncertain in January 2020. In later months, longwave adjustments become
more evident and contribute to a global negative longwave forcing. However, the net ERF in the tropics, for which substantial
negative values were obtained in the nudged simulations, remains uncertain in the ensemble runs. In the northern hemisphere,
255 the disagreement in ERF between nudged and ensemble simulation is largest which makes conclusions about the actually
realized ERF considerably more uncertain. While the nudged simulations show similar longwave adjustments there as in the
other regions, the ensemble simulations show a much weaker signal.




Figure 2. Ensemble and time zonal mean perturbations (JFM average) of radiative fluxes at TOA (1st row) and surface (4th row) and respective flux convergences between TOA and 200 hPa (2nd row), indicated as middle atmosphere (MATM) forcing, and between 200 hPa and surface (3rd row), indicated as lower atmosphere (LATM) forcing, as function of latitude. The colors represent deviations from the reference simulation (FIRE0) with different fire emission strength scaled by a factor of 1 (FIRE1: blue), 2 (FIRE2: green), 3 (FIRE3: orange) and 5 (FIRE5: red). The sum of shortwave (left column) and longwave (middle column) gives the net contribution (right column). Thick lines represent the ensemble mean value, whereas light shadings show the 95% confidence interval.

Literature values of the net TOA radiative forcing from Australian wildfires are mostly negative (see Tab. 1). At first glance, existing knowledge seems to be in conflict with our results. However, previous studies have mainly been limited to reporting
 260 either forcing for clear-sky situations, forcing that ignores atmospheric adjustments or forcing excluding the bright snow-covered surfaces in the polar regions. Previously, Heinold et al. (2022) had shown that the high total cloud cover over the Southern Ocean increases the planetary albedo so much that the dominance of scattering by smoke is lost. Thus, the original brightening of the dark ocean surface becomes a darkening of the bright, cloud-covered ocean and the negative shortwave forcing turns into a positive one (Bellouin et al., 2020). It is a known effect that increasing cloud cover also increases the
 265 instantaneous radiative forcing of absorbing aerosols that lie above the clouds (Chand et al., 2009). This effect is especially pronounced over the oceans. It becomes apparent from Fig. 3b in which the individual components of the clear-sky forcings have been subtracted from the respective all-sky forcings, in agreement with the definition of cloud-radiative effects (Stephens,




Figure 3. (a) All-sky forcings and (b) cloud effects averaged over specified geographical regions (outer interval on x-axis), over all months (inner interval on x-axis) and over all fire strength perturbations. The different components of the effective radiative forcing (ERF) are represented by differently colored bars. ERF is split into instantaneous radiative forcing (IRF) and adjustments (Adj) and also decomposed into shortwave and longwave components. Each bar represents the average of ensemble mean values for the four normalized fire perturbation experiments. Values for which the standard deviation among the four perturbation experiments is larger than the actual mean value are indicated by light colors. Data from the nudged experiments are shown as small gray horizontal lines for comparison and are also calculated as average over all normalized fire perturbations. Regions are split at $\varphi = 20^\circ\text{S}$ and $\varphi = 20^\circ\text{N}$ where φ represents latitude.

2005). The similarity between all-sky forcing and cloud effects is striking. It means that the clouds are the main driver of the positive forcings obtained in the southern hemisphere simply because of their presence, not regarding any cloud changes. In the
 270 tropics, the cloud effects significantly contribute to negative longwave adjustments. As an intermediate conclusion, it can be stated that just the presence of clouds has a significant impact on ERF estimates and turns shortwave ERF into positive values in contrast to previously reported clear-sky values.

3.2 Adjustments in the Stratosphere

Shortwave heating by the extreme Australian smoke event triggers local and non-local temperature increases in the stratosphere.
 275 As shown in Fig. 4, a net heating maximum between 0.03 and $0.04 \text{ K}(\text{day})^{-1}$ per fire scaling factor is reached after around one week after smoke injection considering the globally averaged lower stratosphere. The nudged simulations seem to reach higher heating maxima for smaller fire strengths. In the following adjustments, a chain of processes acts to reduce the net stratospheric heating rate down to zero again. The turning point is reached equally fast for each fire strength after around nine weeks for the ensemble simulations or after six weeks for the nudged simulations. After the early peak in net heating, the decay
 280 follows approximately an exponential curve with decay constants of 14 and 21 days for the nudged and ensemble simulations, respectively.

As part of the stratospheric adjustments, a positive temperature perturbation emerges (see Fig. 5). Initially the temperature perturbation develops in the southern mid-latitudes in both the nudged and ensemble simulations, and subsequently spreads



Table 1. Reported values of radiative forcing by Australian wildfire smoke are compared to our simulation results (first four rows). Our ERF estimates have been derived from the ensemble runs including the confidence interval with all values rescaled by the fire scaling factor. Literature values have been sorted in descending order.

Reference	Radiative Forcing / (W m^{-2})	ERF	Clear-sky	Comments
FIRE1	0.2 ± 0.2	*		
FIRE2	0.06 ± 0.11	*		rescaled, globally averaged, JFM season
FIRE3	0.03 ± 0.07	*		
FIRE5	-0.01 ± 0.04	*		
Sellitto et al. (2022)	0.8 ± 0.2			
Heinold et al. (2022)	0.44 ± 0.07			global modeling, JFM season
Yu et al. (2021)	-0.03	*	*	global modeling
Liu et al. (2022)	-0.17 ± 0.02	*	*	values averaged from 60°N to 60°S
Khaykin et al. (2020)	-0.31 ± 0.09		*	area-weighted global-equivalent; instantaneous
Sellitto et al. (2022)	-0.35 ± 0.09		*	area-weighted global-equivalent; instantaneous
Chang et al. (2021)	-0.5 ± 0.2		*	rough estimate based on MODIS; instantaneous
Fasullo et al. (2021)	-0.95 ± 0.15	*		likely no significant smoke emission into UTLS region
Hirsch and Koren (2021)	-1.0 ± 0.6		*	observation-based; oceanic areas; 20°S -to- 60°S belt




Figure 4. Net heating rate perturbation averaged globally for stratospheric altitudes between $380 < \theta < 500$ K as function of time. Colors indicate the different fire strength perturbations similar to Fig. 2. Thick solid lines are ensemble mean values and thin solid line represent averaged nudged data. The light gray horizontal line marks the zero line for orientation.

285 towards north and south in the following weeks. While the expansion of the positive temperature perturbation towards the south, i.e. in polar direction, can be mainly attributed to the transport of smoke aerosol and is therefore much more pronounced in the nudged runs, the expansion towards the tropics is related to the changes in the stratospheric circulation. The positive

temperature perturbation reaches northern mid-latitudes after 10 to 14 days. Progression further north can also be seen in the nudged runs. However, the considerable variability of the polar vortex in northern winter possibly hinders the attribution of any temperature effects in the ensemble simulations. A warmer northern hemisphere during 2020 was also identified in Rieger et al. (2021) based on observational data. Based on our analyses, we are wondering if the observed warming is not only a coincidence but may be explained in part by smoke-induced circulation changes. Another difference between the nudged and ensemble runs is that the temperature perturbation is also more confined in the vertical direction in the ensemble simulations. This could indicate that the vertical self-lofting is more pronounced in the nudged runs, consistent with Fig. 1a-f in which significantly larger shortwave forcings were found for the nudged runs.




Figure 5. Temporal evolution of the potential temperature perturbation for the FIRE1 case with respect to the FIRE0 case. Shown is the 1-K contour of the average potential temperature perturbation for the (a) nudged simulations and (b) ensemble simulation that is color-coded by time (see legend). Each second day is presented beginning with 1 Jan 2020. Please note that the pressure starts at 300 hPa at the vertical axis.

The stratospheric circulation changes become apparent from Fig. 6. In the southern tropics at around 10°S , upward motion lofts air towards the tropopause region. Between 100 and 200 hPa, the mean flow divides into a northward and into a southward directed branch of the global atmospheric circulation. The northward directed branch reaches high into the stratosphere and is known as Brewer-Dobson circulation (Butchart, 2014). Nudged and ensemble simulation agree with confidence that due to the effects of absorbing wildfire smoke, a positive anomaly of the residual streamfunction develops in southern mid-latitudes and in the tropics. This leads to a reduction of the strength of the poleward circulation in the southern hemisphere and to an increase of the circulation strength in the upper northern hemispheric branch. This increase propagates much further northward in the nudged simulations than in the ensemble simulations that are highly impacted by the variability of the winter vortex. Both, the nudged and the ensemble runs, also agree on a small weakening of the lower northward branch, but this finding is highly uncertain as well.




Figure 6. Average perturbations in residual circulation for (a) nudged simulations and (b) ensemble simulation as function of latitude and height. Contour lines indicate the state of the reference simulation FIRE0 with solid lines for positive values and dashed lines for negative values. Perturbations of the residual circulation in FIRE1 are presented in colored shading. Both, contours and shading have logarithmic spacing. Hatched regions have the same meaning as in Fig. 1.

305 Changes in the global circulation impact the energy budget of the stratosphere and especially how local diabatic cooling or heating by radiation is compensated by adiabatic heating or cooling from the global circulation. This aspect is visualized in Fig. 7. On average, lofting of air in the tropics leads to adiabatic cooling (dashed contours in Fig. 7) that counterbalances local diabatic heating by radiation and convection. Towards the poles, the average circulation acts to heat the stratosphere, thereby compensating the dominant longwave cooling. This cooling is much more pronounced in the winter hemisphere which is why
310 a stronger circulation is needed there for compensation. The circulation response (colored shading in Fig. 7) develops such that the excess of shortwave heating by smoke in the southern hemisphere is partly compensated due to adiabatic cooling. Moreover, the strengthening of the northward circulation causes additional adiabatic heating in the tropics and the northern mid-latitudes, thus redistributing the initial smoke-induced perturbation on a global scale. Such a temperature broadening mechanism by circulation changes has been recently analyzed in detail for volcanic impacts on tropical dynamics (Brown et al., 2022). It is
315 again highly uncertain how far this heating perturbation reaches northwards.

Due to adjustments, a new balance of the stratospheric state is established. Geostrophy is the strongest constraint for the zonally and monthly averaged dynamics. The positive temperature perturbation in the southern hemisphere causes the slope of the geopotential surfaces to increase toward the south pole, i.e. the perturbation of $\partial_{\varphi} \bar{\Phi} / a$, where φ denote latitude and $\bar{\Phi}$ the




Figure 7. Similar to Fig. 6, but for the average perturbations in adiabatic heating due to circulation changes.

zonally and monthly averaged geopotential, is getting more negative. The resulting pressure-gradient force is nearly exactly
 320 balanced by the meridional component of the Coriolis force $-f\bar{u}$. Since the Coriolis parameter f has negative values in the
 southern hemisphere, the perturbation of the mean zonal wind \bar{u} also becomes negative and reaches values of a few meters per
 second in 100 hPa (not shown). The ageostrophic wind components \bar{v}^* and \bar{w}^* describe the zonal mean residual circulation
 (see Sect. 2.3). As shown in Fig. 8a, \bar{v}^* is getting more positive in the southern hemisphere, thus the magnitude of the average
 poleward motion is reduced and the southern-hemispheric branch of the mean circulation is slowed down (see also again Fig. 6).
 325 The sign of $\Delta\bar{v}^*$ is less clear in the tropics and in the northern hemisphere, and especially in the former region the response
 depends on the affected circulation branch. The significant positive correlations in Fig. 8a-c make clear that the adjustment of
 the mean meridional wind is accompanied by an adjustment of the large-scale waves measured by the so-called Eliassen-Palm
 flux defined after Edmon et al. (1980). The effect is only present in the freely running ensemble simulation, whereas nudging
 destroys this important relation between the mean flow and the wave activity (not shown). The vertical component of the
 330 residual circulation is impacted differently by radiative heating depending on the region under consideration (Fig. 8d-f). In the
 southern hemisphere, the additional heating by shortwave smoke forcing induces increased upward motion (given by negative
 $\Delta\bar{w}^*$) that is strongest in January 2020 before longwave cooling starts to balance shortwave heating in the later months (Fig. 8d).
 Thus, the response of the southern hemisphere is thermally direct. The circulation change is responsible for an adiabatic cooling
 induced by $\approx |\Gamma| \Delta\bar{w}^*$ with $\Gamma = \partial_p \bar{\theta}$ that compensates the smoke perturbation. However, an indirect relationship between
 335 radiative heating and circulation adjustments is found in the tropics (Fig. 8e). Thus, circulation changes generate additional



adiabatic heating, which further amplifies the radiation-induced increase in temperature. Circulation changes remain uncertain in the northern hemisphere (Fig. 8f) and appear to be unrelated to radiative perturbations.




Figure 8. Relationship between components of the residual circulation and the forcing by large-scale waves (1st row) or the radiative heating (2nd row). Symbols mark monthly and regionally averaged anomalies with colors indicating fire strength from FIRE1 (blue) to FIRE5 (red) and symbol style for different months: January (circle), February (square) and March (triangle). Gray error bars indicate the 95% confidence interval. Regions are split by $\varphi = 20^\circ\text{S}$ and $\varphi = 20^\circ\text{N}$ where φ represents latitude.

3.3 Adjustments in the Troposphere

In the following, adjustments in the troposphere due to the Australian wildfire smoke are considered. Smoke-induced heating affects not only the lower stratosphere as described above, but also the upper troposphere. This is visualized by vertical profiles of radiative heating, potential temperature, relative humidity and cloud cover in Fig. 9. Again, the heating by smoke induces a positive temperature perturbation. This temperature anomaly not only travels upwards by self-lofting (as e.g. seen in Fig. 5), but also progresses downwards and subsequently enters the upper troposphere over the course of several months as it can be clearly seen in Fig. 9b. The warming of upper tropospheric layers leads to reduced relative humidity for which negative perturbations reach anomalous values of a few percent at 200 hPa (Fig. 9c). The reduction of supersaturated regions in the upper troposphere also reduces the coverage of the high clouds in the model, e.g. by about 1% at 200 hPa (Fig. 9d). Similar aerosol effects on cirrus clouds were described in Samset and Myhre (2015) who applied perturbations by black carbon at different atmospheric



layers. Furthermore, Samset and Myhre (2015) pointed out that instantaneous forcing and adjustments differ the most when absorbing aerosol perturbations are introduced higher up in the atmosphere.




Figure 9. Globally-averaged vertical profiles of anomalies of (a) net radiative heating, (b) potential temperature, (c) relative humidity and (d) cloud cover as function of pressure. Similar to Fig. 4, thick colored lines represent ensemble mean perturbations with respect to the FIRE0 case. Temporal evolution from January 2020 (leftmost sub-panels) to March 2020 (rightmost sub-panels) is plotted. Humidity is shown relative to saturation over liquid water. Please note that the x-scale is non-linear to improve visibility.

350 The simulated reduction in cirrus clouds impacts longwave as well as shortwave radiation. Due to the smaller amount of cirrus, a smaller amount of terrestrial longwave radiation is trapped in the troposphere. Thus, more longwave radiation is able to escape to space leading to a negative forcing of around -1 W m^{-2} for a cloud-cover reduction of 1% (shown for the tropics in Fig. 10a). The compensation between shortwave and longwave radiative perturbations happens in such a way that the net cloud radiative effects remain rather unchanged (Fig. 10b), in line with known neutral to slightly positive net effects of
 355 cirrus (e.g. Chen et al., 2000; Stephens, 2005; Senf et al., 2020). The reduction of ice-containing clouds is accompanied by a reduction of surface precipitation, see Fig. 10c which is dominated by the response of convective precipitation in the tropics. The link between cloud ice and precipitation is undermined by the known importance of ice and mixed-phase processes in



precipitation formation (Mülmenstädt et al., 2015). The precipitation behavior may be traced back to a dynamical response of the Earth system (see Fig. 10c). For $\Delta\bar{\omega}^*$ at mid-tropospheric levels of 500 hPa, a negative correlation with precipitation is obtained (-0.9). A larger $\Delta\bar{\omega}^*$ means that the upward motion in the tropics is reduced. This dynamical response causes the mean circulation to contribute less to the adiabatic cooling of the tropical troposphere, producing a net dynamical warming of $Q_{\text{dyn}} \approx |\Gamma|\Delta\bar{\omega}^*$. The greater this circulation-induced warming by Q_{dyn} , the more atmospheric cooling Q_{atm} is compensated and the less precipitation must contribute to the warming of the atmospheric column. This is known as energetic constraint on precipitation responses (Muller and O’Gorman, 2011), which highlights that atmospheric heating competes with dynamical heating. Similar to the results of Muller and O’Gorman (2011), a regression coefficient of around $-3300 \text{ W m}^{-2} \text{ Pa}^{-1} \text{ s}$ is obtained from Fig. 10c. In general, the analysis of the energetic perspective of precipitation changes has proven to be very useful for understanding atmospheric adjustments to various forcing agents (Richardson et al., 2018). In the context of stratospheric aerosol intervention strategies, Haywood et al. (2022) could show that continuous injection of absorbing aerosols in the tropical low stratosphere, does not only impact stratospheric temperatures and water vapor and ozone concentrations, but also affects the troposphere. Similar to our findings, absorbing aerosol in the stratosphere lead to a downward impact leading to a reduction in surface precipitation and considerable changes in climate variability modes.




Figure 10. Cirrus effects on TOA radiation and energetic constraints for precipitation changes in the tropical belt set here to $\varphi \in (20^\circ\text{S}, 20^\circ\text{N})$. Symbols and colors have the same meaning as in Fig. 8, which provides a legend to decipher the meaning of the symbols.

As seen above, not only cirrus cloud cover but the entire hydrological cycle is affected by the Australian fires, although the perturbations to evaporative and precipitation fluxes are quite uncertain (as indicated by the gray bars in Fig. 11a-c). Simulated total precipitation and evaporative fluxes are reduced by around 0.4% for February and March 2020 on average (Fig. 11a-c). For the different water phases, the clearest signal occurs for the ice water path which shows a systematic reduction of up to 1% in the FIRE5 run. Again, the plausible underlying cause is the temperature increase localized near the tropopause together with increased downwelling of longwave radiation. These cause a slight perturbation of ice formation conditions such that less areas are supersaturated with respect to ice.




Figure 11. Globally and monthly average perturbations of (a) surface evaporation fluxes, (b) convective precipitation, (c) large-scale precipitation, (d) total column water vapor Q_v , (e) liquid water path Q_c , and (f) ice water path Q_i as function of time. All perturbations are computed relative to the FIRE0 run as reference and presented in %. Colors from blue to red indicated perturbation resulting from fire perturbation runs FIRE1, FIRE2, FIRE3 and FIRE5. Bright colors indicate data from ensemble simulations, whereas light colors represent nudged data shown side-by-side for the respective runs.

4 Discussion

380 In the following, we will focus on the limitations of our study and discuss the consequences that follow. We deliberately chose
 different modeling approaches to compile our results. On the one hand, we were able to model the dispersion of Australian
 smoke in a realistic way using nudged runs. On the other hand, free-running ensemble simulations allowed us to simulate the
 interaction of different thermodynamical and dynamical adjustments and also to estimate their resulting uncertainties. How-
 ever, both approaches have their weaknesses and we aim to partially compensate them by using both approaches. One of the
 385 important weaknesses of the nudged runs is that they do not allow a free circulation response and thus reflect the dynamical
 adjustments in a distorted way. This is especially the case for the relation between meridional residual wind and Eliassen-Palm
 flux divergence, which usually reflects a first order approximation to one component of the momentum balance (Edmon et al.,
 1980). In the nudged setup, the relaxation towards the observation-based mean winds and synoptic-scale wave structures dom-
 inates here. The free-running ensemble simulations eliminate this inconsistency. In those simulations, however, now the exter-
 390 nally imposed fire emissions are not consistent with the meteorological situation. For example, precipitation may occur during
 the extreme fire days in some ensemble realizations and not all runs have convectively unstable environments. Furthermore,
 we cannot answer the question whether the respective synoptic-scale smoke-induced perturbation phenomena are realistically
 reproduced or whether the coarse model resolution prevents the formation and self-maintenance of the smoke-induced anti-
 cyclonic vortex. Doglioni et al. (2022) has shown that this phenomenon can also be described with coarser-resolution models
 395 which is promising. For our data, however, this aspect remains a future research need.



As demonstrated by Heinold et al. (2022), the injection height of the wildfire smoke aerosol is critical for its atmospheric lifetime as well as its spatial distribution and, hence, for its radiative impact. Common injection heights for biomass burning are far below the tropopause leading to quick wet removal (within a few days) of the injected aerosol with only limited horizontal transport and impacts. The strong upward lofting of smoke by exceptionally severe pyroconvection leads to injection of aerosol mass into the upper troposphere / lower stratosphere, where it has much longer life time. The location above most precipitating clouds hinders wet and dry removal and a wider and even global dispersion becomes possible. Self-lofting due to heating by absorbed solar radiation (Yu et al., 2019; Allen et al., 2020; Ohneiser et al., 2022b) further prolongs the presence of the smoke aerosol in the stratosphere and its impacts. Since we have no robust representation of pyroconvection and related transport processes in our model, the traditionally applied smoke plume injection was not able to reproduce the observed injection characteristics for the investigated wildfire event. For this reason, we had to make the compromise that the pyroCb emission heights had to be prescribed in the model for the four most severe Australian wildfire days following Heinold et al. (2022). However, in order to be able to correctly simulate the climate impact of severe wildfires in climate projections, parameterizations are needed that reliably produce the correct frequency and characteristics of pyroCb events.

It must be mentioned that considerable uncertainties exist with respect to the microphysical and optical properties of wildfire smoke. In the following, we reiterate the discussion detailed in Heinold et al. (2022) and again emphasize that even small variations in the content of highly-absorbing BC and in the morphology of smoke particles lead to significant changes in the global radiative forcing estimates (Sellitto et al., 2022). In our study, the ratio of BC to total carbon in Australian smoke is about 0.05-0.08, corresponding to a particle single scattering albedo (SSA) between 0.82-0.85 at 550 nm. In comparison to Bellouin et al. (2020) and Brown et al. (2021) these optical properties are similar to those used in other global aerosol-climate models. However, Brown et al. (2021) demonstrated using aircraft data that the absorptivity of aerosol particles from biomass burning is overestimated by the models due to an inadequate description of the mixing state. However, these are generic results averaged over all possible biomass burning events that may not be easily extrapolated to extreme wildfires, since both the burned biofuel and the time for chemical processing are unique in these cases. This may also be the reason why Haarig et al. (2018) were able to derive a rather low SSA of 0.80 from multispectral lidar observations for the intense Canadian fires of 2017. Similar low SSA values were also identified for the Australian extreme event (Ohneiser et al., 2022a). Furthermore, we also cannot exclude that secondary organic aerosol, which is not considered in our model, contributes significantly to the aerosol mass (especially in the later course) and leads to increased scattering by the aerosol mixture. Yet, the remarkable agreement of the various simulated smoke plume properties with observational data obtained by Heinold et al. (2022) further makes us confident that the absorption of solar radiation by the smoke aerosol is not massively overestimated in our model.

The Australian smoke impacted not only on stratospheric temperatures and circulation, but also on concentrations of stratospheric water vapor and ozone. During the period from August to December 2020, a reduction in the total ozone column of 10-20 Dobson units was estimated for the southern hemisphere (Yu et al., 2021). First, changes in the Brewer-Dobson circulation (Diallo et al., 2022) may be responsible. Second, heterogeneous ozone chemistry at the surface of the smoke particles was identified as the potential cause of the record-breaking ozone hole over Antarctica in September-November 2020 (Ohneiser et al., 2020). Yu et al. (2021) estimated the contribution of the latter mechanism to be about 50% with respect to the total ozone

reduction. Furthermore, a substantially moister lower stratosphere was attributed to the effects of the Australian fires by Diallo et al. (2022). However, since we do not sufficiently describe these aspects with our current model setup, and since they mainly occur in the second half of 2020, we only conducted simulation experiments up to and including March 2020. Therefore, we are only able to draw conclusions on the changes and underlying mechanisms during the first three months after the extreme
435 Australian wildfire event.

In general, global adjustments can produce very diverse responses to perturbations in forcing constituents. The occurrence and impacts of severe wildfires are, however, highly dependent on details of the dynamical and thermodynamical state of the atmosphere as well as on details of the properties of the biosphere and the Earth's surface. These interconnections might make fire impacts less tractable as part of the global adjustment concept which might complicate the robust assessment of severe fire
440 impacts in future climates.

5 Conclusions

Wildfires contribute to a significant amount to absorbing aerosol in the atmosphere (Bond et al., 2013; Sokolik et al., 2019). Absorption induces atmospheric adjustment processes that can have considerably diverse feedback on thermodynamic structure, cloud characteristics, and circulations (Koch and Del Genio, 2010; Bellouin et al., 2020). These adjustment processes can
445 cause both positive and negative climate forcings and are still subject to large uncertainties (Smith et al., 2018).

The Australian wildfires were a particularly extreme fire event during the turn of the year between 2019 and 2020, which had a terrible impact on nature and the human population (Borchers Arriagada et al., 2020; Ward et al., 2020; Wintle et al., 2020). Due to the meteorological situation and extreme heat, strong pyroconvection clouds formed, efficiently transporting wildfire aerosol from the source of the fire to the upper troposphere and lower stratosphere (Kablick et al., 2020; Peterson et al., 2021).
450 Once there, the fire aerosol had a significantly longer lifetime compared to smoke remaining in the lower troposphere, allowing it to disperse throughout the southern hemisphere (Ohneiser et al., 2022a). In this particular situation, localized causes had global and longer-term effects, which is why the Australian fires are compared to strong volcanic eruptions in the literature (Hirsch and Koren, 2021; Liu et al., 2022).

In contrast to aerosol of volcanic origin, the absorption of solar radiation by the black carbon contained in wildfire aerosol
455 lead to relatively rapid adjustment processes in the stratosphere. That these caused a short-term increase in stratospheric temperature for the Australian event and thus compensate for the positive radiative flux convergence in the stratosphere is sufficiently documented by existing literature (Rieger et al., 2021; Stocker et al., 2021; Yu et al., 2021; Liu et al., 2022). However, the mechanisms leading to the adjustment, the resulting interactions between the lower stratosphere and the upper troposphere, and the magnitude of the effective radiative forcing are not yet conclusively understood. Therefore, we would like to contribute
460 our perspective on these aspects.

To shed light on the global effect of fire aerosol from the Australian wildfires, simulations were performed in our study using the global climate chemistry model ECHAM-HAM. Two separate modeling approaches were chosen and compared to reduce the uncertainties that arise. In nudged runs, where the simulated winds were relaxed to ERA5 data, the transport of smoke



was well reproduced. Additional free-running ensemble runs allowed us to examine adjustments in the global circulation and
465 large-scale waves in more detail. Focusing on the smoke effects within the first three months after the Australian wildfire event,
the following important results could be derived:

- (i) The stratosphere responds to smoke-induced heating with a local and non-local warming leading to a temperature increase
in the order of a few Kelvin. Compensating longwave cooling can offset a significant portion of the heating from locally
absorbed solar radiation.
- 470 (ii) Adjustments in the stratosphere lead to changes in the global circulation. In the southern hemisphere, less adiabatic
heating by the residual circulation is required to compensate radiative cooling. Thus, the southern-hemispheric circulation
branch is weakened. In contrast, the northern-hemispheric circulation branch is slightly strengthened such that the energy
from the initially localized smoke-induced shortwave heating is redistributed to the tropics and the northern hemisphere.
- (iii) In the troposphere, the amount of cirrus is reduced by up to 1 % due to the warming effects of Australian smoke, which
475 is caused by a reduction in relative humidity in the upper troposphere. The resulting tropospheric adjustments impact
the hydrological cycle, with subsequently reduced amounts of ice water path, surface precipitation and evaporation. Due
to energetic constraints, tropospheric circulation is also affected, with a link between precipitation changes and tropical
upward motion leading to changes in adiabatic cooling in the tropics.
- (iv) Clouds, just by their presence, have a significant impact on aerosol effective radiative forcing (ERF) estimates at top of the
480 atmosphere (TOA) and turn shortwave ERF to positive values in contrast to previously reported clear-sky values. How-
ever, as net effect atmospheric adjustments nearly balance the initially positive instantaneous TOA forcing by Australian
smoke.

We showed that high reaching smoke plumes from pyroconvection can cause modifications in the stratospheric circulation.
The consequences of such large-scale circulation changes due to stratospheric smoke and potential impacts on the troposphere
485 are not yet fully understood and further research is needed to disentangle the complex interaction and coupling mechanisms.
As rapid climate change, affecting both the atmosphere and vegetation, increases the risk and intensity of wildfires (Jain et al.,
2022), the representation of deep pyroconvective events and associated transport of absorbing aerosols into the stratosphere
has to be improved in global climate models. Furthermore, the related consequences for the radiation budget, the changes in
the stratospheric circulation and the effects on the troposphere should be well understood in order to adequately consider these
490 possible interactions in climate projections.

Code availability. The ECHAM-HAMMOZ code is maintained and made available to the scientific community under <https://redmine.hammoz.ethz.ch>
(HAMMOZ community, 2022). The availability is regulated under the HAMMOZ Software Licence Agreement that can be downloaded from
https://redmine.hammoz.ethz.ch/attachments/download/291/License_ECHAM-HAMMOZ_June2012.pdf (last access: 15 January 2023). The
analysis source code has been made freely available to improve reproducibility of our results. The final plots for our paper were done with
495 Jupyter Notebooks which are published at <https://doi.org/10.5281/zenodo.7575809>.



Data availability. Analysis data have been collected and can be assessed under <https://doi.org/10.5281/zenodo.7568466>.

Author contributions. FS, BH and IT conceived the idea and led the study. FS and AK performed the model development and ran the simulations. FS and JM focused on the analysis and interpretation of model results. FS, IT, RS, AK and JM wrote the paper with contributions from all the co-authors. All the authors participated in the revision and editing of the paper.

500 *Competing interests.* The contact author has declared that none of the authors has any competing interests.

Acknowledgements. The ECHAM-HAMMOZ model is developed by a consortium composed of ETH Zürich, Max Planck Institute for Meteorology, Forschungszentrum Jülich, the University of Oxford, the Finnish Meteorological Institute and the Leibniz Institute for Tropospheric Research (TROPOS) and managed by the Center for Climate Systems Modelling (C2SM) at ETH Zürich. We acknowledge computational support for conducting the experiments and storing the data at the Deutsches Klimarechenzentrum (DKRZ), within the compute projects
505 bb1174, bb1262, bb1004 and bb1191. We further thank Kevin Ohneiser for his constructive feedback on an earlier manuscript.



References

- Allen, D. R., Fromm, M. D., III, G. P. K., and Nedoluha, G. E.: Smoke with Induced Rotation and Lofting (SWIRL) in the Stratosphere, *J. Atmos. Sci.*, 77, 4297 – 4316, <https://doi.org/10.1175/JAS-D-20-0131.1>, 2020.
- Andrews, D. G. and McIntyre, M. E.: Generalized Eliassen-Palm and Charney-Drazin theorems for waves on axisymmetric mean flows in compressible atmospheres, *J ATMOS SCI*, 35, 175–185, 1978.
- Andrews, G. D. and McIntyre, M. E.: Planetary waves in horizontal and vertical shear: The generalized Eliassen-Palm relation and the mean zonal acceleration, *J ATMOS SCI*, 33, 2031–2079, [https://doi.org/https://doi.org/10.1175/1520-0469\(1976\)033%3C2031:PWIHAV%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1976)033%3C2031:PWIHAV%3E2.0.CO;2), 1976.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding global aerosol radiative forcing of climate change, *Rev. Geophys.*, 58, e2019RG000 660, <https://doi.org/10.1029/2019RG000660>, 2020.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment, *J. Geophys. Res. Atmos.*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- Boone, C. D., Bernath, P. F., and Fromm, M. D.: Pyrocumulonimbus Stratospheric Plume Injections Measured by the ACE-FTS, *Geophys. Res. Lett.*, 47, e2020GL088 442, <https://doi.org/https://doi.org/10.1029/2020GL088442>, e2020GL088442 2020GL088442, 2020.
- Borchers Arriagada, N., Palmer, A. J., Bowman, D. M., Morgan, G. G., Jalaludin, B. B., and Johnston, F. H.: Unprecedented smoke-related health burden associated with the 2019–20 bushfires in eastern Australia, *Med. J. Aust.*, 213, 282–283, <https://doi.org/https://doi.org/10.5694/mja2.50545>, 2020.
- Bowman, D. M. J. S., Kolden, C. A., Abatzoglou, J. T., Johnston, F. H., van der Werf, G. R., and Flannigan, M.: Vegetation fires in the Anthropocene, *Nature Reviews Earth and Environment*, 1, 500–515, <https://doi.org/10.1038/s43017-020-0085-3>, 2020.
- Brown, F., Marshall, L., Haynes, P. H., Garcia, R. R., Birner, T., and Schmidt, A.: On the magnitude and sensitivity of the QBO response to a tropical volcanic eruption, *EGUsphere*, 2022, 1–37, <https://doi.org/10.5194/egusphere-2022-1096>, 2022.
- Brown, H., Liu, X., Pokhrel, R., Murphy, S., Lu, Z., Saleh, R., Mielonen, T., Kokkola, H., Bergman, T., Myhre, G., Skeie, R. B., Watson-Parris, D., Stier, P., Johnson, B., Bellouin, N., Schulz, M., Vakkari, V., Beukes, J. P., van Zyl, P. G., Liu, S., and Chand, D.: Biomass burning aerosols in most climate models are too absorbing, *Nat. Commun.*, 12, <https://doi.org/10.1038/s41467-020-20482-9>, 2021.
- Butchart, N.: The Brewer-Dobson circulation, *REV GEOPHYS*, 52, 157–184, 2014.
- Chand, D., Wood, R., Anderson, T. L., Satheesh, S. K., and Charlson, R. J.: Satellite-derived direct radiative effect of aerosols dependent on cloud cover, *Nat. Geosci.*, 2, 181–184, <https://doi.org/10.1038/ngeo437>, 2009.
- Chang, D. Y., Yoon, J., Lelieveld, J., Park, S. K., Yum, S. S., Kim, J., and Jeong, S.: Direct radiative forcing of biomass burning aerosols from the extensive Australian wildfires in 2019–2020, *Environ. Res. Lett.*, 16, 044 041, <https://doi.org/10.1088/1748-9326/abecfe>, 2021.
- Chen, T., Rossow, W. B., and Zhang, Y.: Radiative Effects of Cloud-Type Variations, *J. Climate*, 13, 264–286, 2000.



- Diallo, M. A., Ploeger, F., Hegglin, M., Ern, M., Grooß, J., Khaykin, S., and Riese, M.: Stratospheric water vapour and ozone response to the quasi-biennial oscillation disruptions in 2016 and 2020, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/ACP-22-14303-2022>, 2022.
- 545 Doglioni, G., Aquila, V., Das, S., Colarco, P. R., and Zardi, D.: Dynamical perturbation of the stratosphere by a pyrocumulonimbus injection of carbonaceous aerosols, *Atmos. Chem. Phys.*, 22, 11 049–11 064, <https://doi.org/10.5194/acp-22-11049-2022>, 2022.
- Dunkerton, T.: On the Mean Meridional Mass Motions of the Stratosphere and Mesosphere, *J ATMOS SCI*, 35, 2325 – 2333, [https://doi.org/10.1175/1520-0469\(1978\)035<2325:OTMMMM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2325:OTMMMM>2.0.CO;2), 1978.
- Edmon, H. J., Hoskins, B. J., and McIntyre, M. E.: Eliassen-Palm Cross Sections for the Troposphere, *J. Atmos. Sci.*, 37, 2600 – 2616, [https://doi.org/10.1175/1520-0469\(1980\)037<2600:EPCSFT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2600:EPCSFT>2.0.CO;2), 1980.
- 550 Fasullo, J. T., Rosenbloom, N., Buchholz, R. R., Danabasoglu, G., Lawrence, D. M., and Lamarque, J.-F.: Coupled Climate Responses to Recent Australian Wildfire and COVID-19 Emissions Anomalies Estimated in CESM2, *Geophys. Res. Lett.*, <https://doi.org/10.1029/2021GL093841>, 2021.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity, pp. 923–1054, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, 2021.
- 555 Fromm, M., Lindsey, D. T., Servranckx, R., Yue, G., Trickl, T., Sica, R., Doucet, P., and Godin-Beekmann, S.: The Untold Story of Pyrocumulonimbus, *Bull. Amer. Meteor. Soc.*, 91, 1193 – 1210, <https://doi.org/10.1175/2010BAMS3004.1>, 2010.
- Fromm, M., Peterson, D., and Di Girolamo, L.: The Primary Convective Pathway for Observed Wildfire Emissions in the Upper Troposphere and Lower Stratosphere: A Targeted Reinterpretation, *J. Geophys. Res. Atmos.*, 124, 13 254–13 272, <https://doi.org/https://doi.org/10.1029/2019JD031006>, 2019.
- 560 Giorgetta, M., Jungclaus, J., Reick, C., Legutke, S., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., Haak, H., Hollweg, H.-D., Kinne, S., Kornblueh, L., Matei, D., Mauritsen, T., Mikolajewicz, U., Müller, W., Notz, D., Raddatz, T., Rast, S., Roeckner, E., Salzmann, M., Schmidt, H., Schnur, R., Segschneider, J., Six, K., Stockhause, M., Wegner, J., Widmann, H., Wieners, K.-H., Claussen, M., Marotzke, J., and Stevens, B.: CMIP5 simulations of the Max Planck Institute for Meteorology (MPI-M) based on the MPI-ESM-LR model: The amip experiment, served by ESGF, Tech. rep., World Data Center for Climate (WDCC) at DKRZ, <https://doi.org/https://doi.org/10.1594/WDCC/CMIP5.MXELam>, 2012.
- Grotjahn, R.: *Global atmospheric circulations- Observations and theories*, 1993.
- Haarig, M., Ansmann, A., Baars, H., Jimenez, C., Veselovskii, I., Engelmann, R., and Althausen, D.: Depolarization and lidar ratios at 355, 532, and 1064 nm and microphysical properties of aged tropospheric and stratospheric Canadian wildfire smoke, *Atmos. Chem. Phys.*, 18, 11 847–11 861, <https://doi.org/10.5194/acp-18-11847-2018>, 2018.
- 570 Haywood, J. M., Jones, A., Johnson, B. T., and McFarlane Smith, W.: Assessing the consequences of including aerosol absorption in potential stratospheric aerosol injection climate intervention strategies, *Atmos. Chem. Phys.*, 22, 6135–6150, <https://doi.org/10.5194/acp-22-6135-2022>, 2022.
- Heinold, B., Baars, H., Barja, B., Christensen, M., Kubin, A., Ohneiser, K., Schepanski, K., Schutgens, N., Senf, F., Schrödner, R., Villanueva, D., and Tegen, I.: Important role of stratospheric injection height for the distribution and radiative forcing of smoke aerosol from the 2019–2020 Australian wildfires, *Atmos. Chem. Phys.*, 22, 9969–9985, <https://doi.org/10.5194/acp-22-9969-2022>, 2022.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E.,



- 580 Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quart. J. Roy. Meteor. Soc.*, 146, 1999–2049, <https://doi.org/https://doi.org/10.1002/qj.3803>, 2020.
- Hirsch, E. and Koren, I.: Record-breaking aerosol levels explained by smoke injection into the stratosphere, *Science*, 371, 1269–1274, <https://doi.org/10.1126/science.abe1415>, 2021.
- Holton, J.: Zonally Averaged Flow, in: The stratosphere and its role in the Climate System, edited by Brasseur, G. P., vol. Vol. 54 of *Series*
585 *1: Global Environmental Change*, Springer Science & Business Media, 1997.
- Huang, J., Guo, J., Wang, F., Liu, Z., Jeong, M.-J., Yu, H., and Zhang, Z.: CALIPSO inferred most probable heights of global dust and smoke layers, *J. Geophys. Res. Atmos.*, 120, 5085–5100, <https://doi.org/https://doi.org/10.1002/2014JD022898>, 2015.
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., and Collins, W. D.: Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models, *J. Geophys. Res. Atmos.*, 113, <https://doi.org/https://doi.org/10.1029/2008JD009944>, 2008.
- 590 Jain, P., Castellanos-Acuna, D., Coogan, S. C. P., Abatzoglou, J. T., and Flannigan, M. D.: Observed increases in extreme fire weather driven by atmospheric humidity and temperature, *Nature Climate Change*, 12, 63–70, <https://doi.org/10.1038/s41558-021-01224-1>, 2022.
- Jolly, W. M., Cochrane, M. A., Freeborn, P. H., Holden, Z. A., Brown, T. J., Williamson, G. J., and Bowman, D. M. J. S.: Climate-induced variations in global wildfire danger from 1979 to 2013, *Nat. Commun.*, 6, 7537, 2015.
- 595 Kablick, G., Allen, D. R., Fromm, M., and Nedoluha, G.: Australian PyroCb Smoke Generates Synoptic-Scale Stratospheric Anticyclones, *Geophys. Res. Lett.*, <https://doi.org/10.1029/2020GL088101>, 2020.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- 600 Khaykin, S., Legras, B., Bucci, S., Sellitto, P., Isaksen, L., Tencé, F., Bekki, S., Bourassa, A., Rieger, L., Zawada, D., Jumelet, J., and Godin-Beekmann, S.: The 2019/20 Australian wildfires generated a persistent smoke-charged vortex rising up to 35 km altitude, *Commun Earth Environ*, 1, 22, <https://doi.org/10.1038/s43247-020-00022-5>, 2020.
- Koch, D. and Del Genio, A. D.: Black carbon semi-direct effects on cloud cover: review and synthesis, *Atmos. Chem. Phys.*, 10, 7685–7696, <https://doi.org/10.5194/acp-10-7685-2010>, 2010.
- 605 Liu, C.-C., Portmann, R. W., Liu, S., Rosenlof, K. H., Peng, Y., and Yu, P.: Significant Effective Radiative Forcing of Stratospheric Wildfire Smoke, *Geophys. Res. Lett.*, 49, e2022GL100175, <https://doi.org/https://doi.org/10.1029/2022GL100175>, e2022GL100175 2022GL100175, 2022.
- Matsuno, T.: Lagrangian motion of air parcels in the stratosphere in the presence of planetary waves, *PURE APPL GEOPHYS*, 118, 189–216, <https://doi.org/https://doi.org/10.1007/BF01586451>, 1980.
- 610 McIntyre, M. E.: Towards a Lagrangian-mean description of stratospheric circulations and chemical transports, *PHILOS TR R SOC S-A*, 296, 129–148, 1980.
- Muller, C. J. and O’Gorman, P.: An energetic perspective on the regional response of precipitation to climate change, *Nature Clim Change*, 1, 266–271, 2011.
- Mülmenstädt, J., Sourdeval, O., Delanoë, J., and Quaas, J.: Frequency of occurrence of rain from liquid-, mixed-, and ice-phase clouds derived
615 from A-Train satellite retrievals, *Geophys. Res. Lett.*, 42, 6502–6509, <https://doi.org/https://doi.org/10.1002/2015GL064604>, 2015.
- Ohneiser, K., Ansmann, A., Baars, H., Seifert, P., Barja, B., Jimenez, C., Radenz, M., Teisseire, A., Floutsi, A. A., Haorig, M., Foth, A., Chudnovsky, A., Engelmann, R., Zamorano, F., Bühl, J., and Wandinger, U.: Smoke of extreme Australian bushfires observed in the



- stratosphere over Punta Arenas, Chile, in January 2020: optical thickness, lidar ratios, and depolarization ratios at 355 and 532 nm, *Atmos. Chem. Phys.*, <https://doi.org/10.5194/ACP-20-8003-2020>, 2020.
- 620 Ohneiser, K., Ansmann, A., Kaifler, B., Chudnovsky, A., Barja, B., Knopf, D. A., Kaifler, N., Baars, H., Seifert, P., Villanueva, D., Jimenez, C., Radenz, M., Engelmann, R., Veselovskii, I., and Zamorano, F.: Australian wildfire smoke in the stratosphere: the decay phase in 2020/2021 and impact on ozone depletion, *Atmos. Chem. Phys.*, 22, 7417–7442, <https://doi.org/10.5194/acp-22-7417-2022>, 2022a.
- Ohneiser, K., Ansmann, A., Witthuhn, J., Deneke, H., Chudnovsky, A., and Walter, G.: Self-lofting of wildfire smoke in the troposphere and stratosphere caused by radiative heating: simulations vs space lidar observations, *Atmos. Chem. Phys. Discuss.*, 2022, 1–41, <https://doi.org/10.5194/acp-2022-343>, 2022b.
- 625 Peixoto, J. P. and Oort, A. H.: *Physics of climate*, New York, NY (United States); American Institute of Physics, 1992.
- Peterson, D. A., Campbell, J. R., Hyer, E. J., Fromm, M. D., Kablick, G. P., Cossuth, J. H., and DeLand, M. T.: Wildfire-driven thunderstorms cause a volcano-like stratospheric injection of smoke, *npj Climate and Atmospheric Science*, 1, 30, 2018.
- Peterson, D. A., Fromm, M. D., McRae, R. H., Campbell, J. R., Hyer, E. J., Taha, G., Camacho, C. P., Kablick, G. P., Schmidt, C. C., and
630 DeLand, M. T.: Australia’s Black Summer pyrocumulonimbus super outbreak reveals potential for increasingly extreme stratospheric smoke events, *npj Clim Atmos Sci*, 4, 1–16, 2021.
- Pincus, R. and Stevens, B.: Paths to accuracy for radiation parameterizations in atmospheric models, *J. Adv. Model. Earth Syst.*, 5, 225–233, <https://doi.org/https://doi.org/10.1002/jame.20027>, 2013.
- Richardson, T. B., Forster, P. M., Andrews, T., Boucher, O., Faluvegi, G., Fläschner, D., Hodnebrog, Ø., Kasoar, M., Kirkevåg, A., Lamarque,
635 J.-F., Myhre, G., Olivie, D., Samset, B. H., Shawki, D., Shindell, D., Takemura, T., and Voulgarakis, A.: Drivers of Precipitation Change: An Energetic Understanding, *J. Climate*, 31, 9641 – 9657, <https://doi.org/10.1175/JCLI-D-17-0240.1>, 2018.
- Rieger, L. A., Randel, W. J., Bourassa, A. E., and Solomon, S.: Stratospheric Temperature and Ozone Anomalies Associated With the 2020 Australian New Year Fires, *Geophys. Res. Lett.*, 48, e2021GL095898, <https://doi.org/https://doi.org/10.1029/2021GL095898>, e2021GL095898 2021GL095898, 2021.
- 640 Samset, B. H. and Myhre, G.: Climate response to externally mixed black carbon as a function of altitude, *J. Geophys. Res. Atmos.*, 120, 2913–2927, <https://doi.org/https://doi.org/10.1002/2014JD022849>, 2015.
- Sanap, S. and Pandithurai, G.: The effect of absorbing aerosols on Indian monsoon circulation and rainfall: A review, *Atmos. Res.*, 164–165, 318–327, <https://doi.org/https://doi.org/10.1016/j.atmosres.2015.06.002>, 2015.
- Sellitto, P., Belhadji, R., Kloss, C., and Legras, B.: Radiative impacts of the Australian bushfires 2019–2020 – Part 1: Large-scale radiative
645 forcing, *Atmos. Chem. Phys.*, 22, 9299–9311, <https://doi.org/10.5194/acp-22-9299-2022>, 2022.
- Senf, F., Voigt, A., Clerbaux, N., Hünerbein, A., and Deneke, H.: Increasing Resolution and Resolving Convection Improve the Simulation of Cloud-Radiative Effects Over the North Atlantic, *J. Geophys. Res. Atmos.*, 125, e2020JD032667, <https://doi.org/10.1029/2020JD032667>, e2020JD032667 10.1029/2020JD032667, 2020.
- Senf, F., Quaas, J., and Tegen, I.: Absorbing aerosol decreases cloud cover in cloud-resolving simulations over Germany, *Quart. J. Roy. Meteor. Soc.*, 147, 4083–4100, <https://doi.org/https://doi.org/10.1002/qj.4169>, 2021.
- 650 Smith, C. J., Kramer, R. J., Myhre, G., Forster, P. M., Soden, B. J., Andrews, T., Boucher, O., Faluvegi, G., Fläschner, D., Hodnebrog, Ø., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Mülmenstädt, J., Olivie, D., Richardson, T., Samset, B. H., Shindell, D., Stier, P., Takemura, T., Voulgarakis, A., and Watson-Parris, D.: Understanding Rapid Adjustments to Diverse Forcing Agents, *Geophys. Res. Lett.*, 45, 12,023–12,031, <https://doi.org/10.1029/2018GL079826>, 2018.



- 655 Sokolik, I. N., Soja, A. J., DeMott, P. J., and Winker, D.: Progress and Challenges in Quantifying Wildfire Smoke Emissions, Their Properties, Transport, and Atmospheric Impacts, *J. Geophys. Res. Atmos.*, 124, 13 005–13 025, <https://doi.org/https://doi.org/10.1029/2018JD029878>, 2019.
- Stephens, G. L.: Cloud Feedbacks in the Climate System: A Critical Review, *J. Climate*, 18, 237–273, 2005.
- Stocker, M., Ladstädter, F., and Steiner, A.: Observing the climate impact of large wildfires on stratospheric temperature., *Scientific reports*, <https://doi.org/10.1038/S41598-021-02335-7>, 2021.
- 660 Tegen, I., Neubauer, D., Ferrachat, S., Siegenthaler-Le Drian, C., Bey, I., Schutgens, N., Stier, P., Watson-Parris, D., Stanelle, T., Schmidt, H., Rast, S., Kokkola, H., Schultz, M., Schroeder, S., Daskalakis, N., Barthel, S., Heinold, B., and Lohmann, U.: The global aerosol–climate model ECHAM6.3–HAM2.3 – Part 1: Aerosol evaluation, *Geosci. Model Dev.*, 12, 1643–1677, <https://doi.org/10.5194/gmd-12-1643-2019>, 2019.
- 665 Thornhill, G. D., Collins, W. J., Kramer, R. J., Olivieri, D., Skeie, R. B., O’Connor, F. M., Abraham, N. L., Checa-Garcia, R., Bauer, S. E., Deushi, M., Emmons, L. K., Forster, P. M., Horowitz, L. W., Johnson, B., Keeble, J., Lamarque, J.-F., Michou, M., Mills, M. J., Mulcahy, J. P., Myhre, G., Nabat, P., Naik, V., Oshima, N., Schulz, M., Smith, C. J., Takemura, T., Tilmes, S., Wu, T., Zeng, G., and Zhang, J.: Effective radiative forcing from emissions of reactive gases and aerosols – a multi-model comparison, *Atmos. Chem. Phys.*, 21, 853–874, <https://doi.org/10.5194/acp-21-853-2021>, 2021.
- 670 Vignati, E., Wilson, J., and Stier, P.: M7: An efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *J. Geophys. Res. Atmos.*, 109, <https://doi.org/https://doi.org/10.1029/2003JD004485>, 2004.
- Ward, M., Tulloch, A. I., Radford, J. Q., Williams, B. A., Reside, A. E., Macdonald, S. L., Mayfield, H. J., Maron, M., Possingham, H. P., Vine, S. J., et al.: Impact of 2019–2020 mega-fires on Australian fauna habitat, *Nat. Ecol. Evol.*, 4, 1321–1326, 2020.
- Wintle, B. A., Legge, S., and Woinarski, J. C.: After the Megafires: What Next for Australian Wildlife?, *Trends Ecol. Evol.*, 35, 753–757, <https://doi.org/https://doi.org/10.1016/j.tree.2020.06.009>, 2020.
- 675 Yu, P., Toon, O. B., Bardeen, C. G., Zhu, Y., Rosenlof, K. H., Portmann, R. W., Thornberry, T. D., Gao, R.-S., Davis, S. M., Wolf, E. T., Gouw, J. d., Peterson, D. A., Fromm, M. D., and Robock, A.: Black carbon lofts wildfire smoke high into the stratosphere to form a persistent plume., *Science*, <https://doi.org/10.1126/SCIENCE.AAX1748>, 2019.
- Yu, P., Davis, S. M., Toon, O. B., Portmann, R. W., Bardeen, C. G., Barnes, J. E., Telg, H., Maloney, C., and Rosenlof, K. H.: Persistent Stratospheric Warming Due to 2019–2020 Australian Wildfire Smoke, *Geophys. Res. Lett.*, 48, e2021GL092609, <https://doi.org/https://doi.org/10.1029/2021GL092609>, e2021GL092609 2021GL092609, 2021.
- 680 Zhang, K., Wan, H., Liu, X., Ghan, S. J., Kooperman, G. J., Ma, P.-L., Rasch, P. J., Neubauer, D., and Lohmann, U.: Technical Note: On the use of nudging for aerosol–climate model intercomparison studies, *Atmos. Chem. Phys.*, 14, 8631–8645, <https://doi.org/10.5194/acp-14-8631-2014>, 2014.