



Combined assimilation of NOAA surface and MIPAS satellite observations to constrain the global budget of carbonyl sulfide

Jin Ma¹, Linda M. J. Kooijmans², Norbert Glatthor³, Stephen A. Montzka⁴, Marc von Hobe⁵, Thomas Röckmann¹, and Maarten C. Krol^{1,2}

¹Institute for Marine and Atmospheric Research, Utrecht University, Utrecht, The Netherlands

²Meteorology and Air Quality, Wageningen University & Research, Wageningen, The Netherlands

³Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

⁴National Oceanic and Atmospheric Administration (NOAA), Boulder, CO, USA

⁵Forschungszentrum Jülich GmbH, Germany

Correspondence: J. Ma (j.ma@uu.nl)

Abstract.

Carbonyl sulfide (COS), a trace gas in our atmosphere that leads to the formation of aerosols in the stratosphere, is taken up by terrestrial ecosystems. Quantifying the biosphere uptake of COS could provide a useful quantity to estimate Gross Primary Productivity. Some COS sources and sinks still contain large uncertainties, and several top down estimates of the COS budget point to an underestimation of sources especially in the tropics. We extended the inverse model TM5-4DVAR to assimilate MIPAS satellite data, in addition to NOAA surface data as used in a previous study. To resolve possible discrepancies among the two observational datasets, a bias correction scheme was implemented. A set of inversions is presented that explores the influence of the different measurement instruments and the settings of the prior fluxes. To evaluate the performance of the inverse system, the HIAPER Pole-to-Pole Observations (HIPPO) aircraft observations and NOAA airborne profiles are used. All inversions reduce the COS biosphere uptake from a prior value of 1053 GgS a⁻¹ to much smaller values, depending on the inversion settings. These large adjustments of the biosphere uptake often turn parts of the Amazonia into a COS source. Only inversions that exclusively use MIPAS observations, or strongly reduce the prior errors on the biosphere flux maintain the Amazonia as a COS sink. Assimilating both NOAA surface data and MIPAS data requires a small bias correction for MIPAS data, mostly at higher latitudes, to correct for inconsistencies in the observational data and/or transport model errors. Analysis of the error reduction and posterior correlation between land and ocean fluxes indicates that co-assimilation of NOAA surface observations and MIPAS data better constrains the COS budget than assimilation of one individual dataset alone. Our inversions with bias corrections reduce the global biosphere uptake to respectively 570 and 687 GgS a⁻¹, depending on the prior biosphere error. Over the Amazonia, these inversions reduce the biosphere uptake from roughly 300 to 100 GgS a⁻¹, indicating a strongly overestimated prior uptake over the Amazonia. Although a recent study also reported reduced COS uptake over the Amazonia, we emphasize that a careful construction of prior fluxes and their associated errors remains important. For instance, an inversion that gives large freedom to adjust the anthropogenic and ocean fluxes of CS₂, an important COS precursor, also closes the budget satisfactorily with much smaller adjustments to the biosphere. Thus, a better characterization of biosphere and ocean fluxes by observations is urgently needed, especially over the data-poor tropics.



1 Introduction

25 Understanding sources and sinks of greenhouse gases is a scientific challenge (Friedlingstein et al., 2020, 2022a, b). One important climate-related process is the carbon uptake by terrestrial ecosystems, named gross primary productivity (GPP). However, accurate estimation of GPP is difficult, since the measured net flux of carbon at ecosystem level is determined by both GPP and ecosystem respiration. In the recent decade, GPP proxy methods have been developed with the aim to improve GPP estimates (Campbell et al., 2008; Wingate et al., 2010; Berry et al., 2013; Commane et al., 2015; Koren, 2021).

30 Carbonyl sulfide (COS), a low-abundant and long-lived trace gas with an average mixing ratio of about 500 pmol mol⁻¹ in the Earth's atmosphere (Montzka et al., 2007; Kremser et al., 2016), is biochemically coupled with CO₂ (Stimler et al., 2010; Campbell et al., 2008; Berry et al., 2013), and a major contributor to stratospheric sulfur aerosols (Crutzen, 1976; Turco et al., 1980; Brühl et al., 2012). The concentration of COS remains relatively constant and shows small inter-annual variability, implying that the sources and sinks of COS are roughly balanced (Montzka et al., 2007). A recent study based on vertical
35 column measurements in the Network for the Detection of Atmospheric Composition Change (NDACC) (Hannigan et al., 2022) reported a slightly increasing tropospheric trend in free tropospheric COS ranging from ~0.0 to 1.55±0.30% a⁻¹ in 2002–2016 that appears correlated with estimates of anthropogenic emissions (Hannigan et al., 2022). In the period 2016–2020, all NDACC stations show a free tropospheric decline in COS. Besides direct emissions of COS, carbon disulfide (CS₂) and dimethyl sulfide (DMS) are species that potentially account for substantial chemical production of COS in the troposphere
40 (Chin and Davis, 1993; Watts, 2000; Kettle et al., 2002; Ma et al., 2021; Jernigan et al., 2022).

COS is absorbed by terrestrial vegetation through photosynthesis sharing a similar pathway as CO₂, but it is not emitted from vegetation through respiration, which makes COS a promising diagnostic tracer to improve estimates of GPP globally (Wohlfahrt et al., 2012; Berry et al., 2013; Campbell et al., 2017; Whelan et al., 2018). Recently, COS observations have been used to estimate regional GPP over arctic North America and boreal regions (Hu et al., 2021), and over the Amazonia forest
45 (Stinecipher et al., 2022), and to detect city-level signals from the urban biosphere (Villalba et al., 2021). Besides, isotopologue measurement and modelling studies are being used to differentiate signals from various sources and sinks (Hattori et al., 2020; Davidson et al., 2021; Baartman et al., 2021; Nagori et al., 2022). In parallel, COS flux measurements combined with land surface models improve our understanding of the interaction of COS with plants and soils (Kooijmans et al., 2017, 2019; Spielmann et al., 2019; Kooijmans et al., 2021; Maignan et al., 2021; Wu et al., 2015; Ogée et al., 2016; Abadie et al., 2022).
50 Inverse modelling to better constrain the global COS budget is also readily progressing (Suntharalingam et al., 2008; Berry et al., 2013; Wang et al., 2016; Kuai et al., 2015; Ma et al., 2021; Remaud et al., 2022; Stinecipher et al., 2022).

The COS abundance and variability in the atmosphere is determined by multiple sources and sinks. Anthropogenic, oceanic and biomass burning emissions are the major sources of COS. The inventories of COS and CS₂ anthropogenic emissions are in the range of 223–586 GgS a⁻¹ (Campbell et al., 2015; Zumkehr et al., 2018). Specifically for China, anthropogenic
55 emission estimates were reported as 174 GgS a⁻¹ in 2015 (Yan et al., 2019), of which 31 GgS a⁻¹ from domestic coal combustion (Du et al., 2016). Global emissions from oceans of 600–800 GgS a⁻¹ would be required to balance the global COS budget (Berry et al., 2013; Glatthor et al., 2015; Kuai et al., 2015), but a mechanistic ocean box model and COS seawater



measurements suggest that direct COS oceanic emissions are unlikely to fully account for the large missing source (Lennartz et al., 2017, 2021, 2020). Biomass burning emissions of COS, estimated as 60 ± 37 GgS a^{-1} by Stinecipher et al. (2019) without accounting for biofuel, is unlikely to account for the missing source as well. More recently, a new chemical product of DMS oxidation known as hydroperoxymethyl thioformate (HPMTF) was discovered, and may play a role in accounting for the missing sources of COS in the troposphere (Wu et al., 2015; Veres et al., 2020; Fung et al., 2022; Jernigan et al., 2022).

On the uptake part of the budget, the largest sink of COS is uptake by terrestrial vegetation and soils, with estimates ranging from -500 to -1200 GgS a^{-1} (Campbell et al., 2008; Suntharalingam et al., 2008; Berry et al., 2013; Remaud et al., 2022). Kooijmans et al. (2021) recently updated the COS biosphere uptake based on the Simple Biosphere Model (SiB4) and the estimate is from -922 to -753 GgS a^{-1} by considering the linear dependency of the flux on the atmospheric COS mole fraction and COS soil production. This estimate is also in line with inverse modelling studies, which report fluxes that range from -1053 to -862 GgS a^{-1} (Ma et al., 2021). The uncertainty in COS biosphere uptake introduced by the soil flux was large (Whelan et al., 2013, 2016, 2018), and Abadie et al. (2022) reported a global net soil sink of only -30 GgS a^{-1} over the 2009–2016 period. Finally, chemical loss by OH-induced tropospheric oxidation and stratospheric photolysis contribute respectively about -100 GgS a^{-1} (Kettle et al., 2002; Montzka et al., 2007; Ma et al., 2021; Remaud et al., 2022) and -40 GgS a^{-1} (Barkley et al., 2008; Brühl et al., 2012; Krysztofiak et al., 2015; Ma et al., 2021) to the COS sink.

Satellites capable of observing COS include The Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) (Glatthor et al., 2017), the Tropospheric Emission Spectrometer (TES) (Kuai et al., 2014), The Atmospheric Chemistry Experiment (ACE-FTS) (Bernath et al., 2021) and the Infrared Atmospheric Sounding Interferometer (IASI) (Vincent and Dudhia, 2017; Camy-Peyret et al., 2017; Cartwright et al., 2021). MIPAS and TES ended their observational period in respectively April 2012 and January 2018, and IASI and ACE are the only operational satellites, the latter sensing numerous gases including COS and its isotopologues (Yousefi et al., 2019). In the past, TES data was assimilated in the GEOS-Chem model to constrain atmospheric COS in the tropics. The results pointed to a missing source over the tropical oceans (Kuai et al., 2015). Recently, MIPAS data was used in the same model to show that GPP over the Amazonia is likely in the lower range of previous estimates (Stinecipher et al., 2022). Satellite data have also served as validation data in COS modelling studies (Ma et al., 2021; Remaud et al., 2022; Cartwright et al., 2022; Wang et al., 2023).

In our previous inverse modelling study (Ma et al., 2021), we found that COS inverse modelling is an under-determined problem caused by the scarcity of ground-based observations, specifically in tropical areas. In that light, it would be of particular interest to investigate the application of satellite data to improve the global inversion of COS. In this study, we explore the use of MIPAS data in addition to the NOAA surface observation network for COS. Specifically, we will present inversions based on MIPAS only, NOAA only, and the combined data set. We attempt to explore a variety of data assimilation approaches to better separate the sources and sinks of COS. In addition, we explore the need for a bias correction scheme to account for the different measurement principles of the data streams, and for potential biases in the model. Previous studies implemented and explored bias correction in the TM5-4DVAR system (Basu et al., 2013; Houweling et al., 2014). In this paper, we will introduce the observations and inverse model in Sections 2 and 3, analyse inversion performance, validation and the COS global budget in Section 4. Discussion and conclusions follow in Sections 5 and 6.



2 Observations of COS

2.1 NOAA surface network

95 The Global Monitoring Laboratory (GML) of the National Oceanic and Atmospheric Administration (NOAA) conducts continuous surface measurements at many sites globally, and provides long-term ground-based measurement data that are suitable for inverse modelling studies. COS is measured over 14 observational sites since 2000 to present time (Montzka et al., 2007). The sites are depicted in Fig. 1.

Among the 14 measurement sites, 10 of them are located in the Northern Hemisphere (NH), and 4 in the Southern Hemisphere (SH). Three are purely oceanic sites (MLO, KUM, and SMO) suitable to sample oceanic emissions. SMO is located in the middle of the South Pacific between Hawaii and New Zealand. Another important station, MLO, is located on the north side of Mauna Loa Volcano, Hawaii, at an elevation of 3397 m a.s.l. The KUM station also samples air from the ocean at Hawaii, but at sea level. From the perspective of inverse modelling, locations in the SH or over oceans are important because they measure COS mole fractions from pristine areas or from the marine atmosphere. In contrast, sites over land are useful to trace COS observations influenced by the biosphere or soil exchange, and sources from anthropogenic emissions and biomass burning. The COS measurements are taken on a weekly or bi-weekly timescale, with an average measurement error of less than 8 pmol mol⁻¹ (Montzka et al., 2007). The measurements are calibrated to maintain data consistency amongst NOAA sites. It is worth noting that the coverage of the NOAA surface network for COS is sparse, which leads to challenges for inverse modelling. The total error used in inverse modelling includes the measurement error, the station dependent representation error, and the flux error (Ma et al., 2021).

2.2 MIPAS satellite

The Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) was a Fourier transform spectrometer that flew on the Environmental Satellite (Envisat) mission. MIPAS was a limb sounder, designed to detect a wide range of gases between the upper troposphere and lower thermosphere. The data sets retrieved by the European Space Agency from calibrated MIPAS spectra include profiles of temperature, H₂O, O₃, CH₄, N₂O, HNO₃, and NO₂ (Fischer et al., 2008). The MIPAS COS dataset used here (V5R_OCS_221/222) was retrieved with the level-2 data processor developed and operated by the Institute for Meteorology and Climate Research (IMK) in cooperation with the Instituto de Astrofísica de Andalucía (IAA). COS retrievals from MIPAS data were characterized and compared with other data sets, showing that COS retrievals from MIPAS are good quality measurements that reveal COS surface exchange processes (Glatthor et al., 2015, 2017). The data sets from MIPAS are available from July 1, 2002 to April 8, 2012.

The MIPAS COS data consist of volume mixing ratio profiles, averaging kernels and geophysical information (temperature, latitude, longitude, altitude, etc.). To compare modelled COS mixing ratios to MIPAS soundings the averaging kernel is applied as:



$$\mathbf{x}_{\text{conv}} = \mathbf{x}_{\text{prior}} + \mathbf{A}(\mathbf{x}_{\text{model}} - \mathbf{x}_{\text{prior}}), \quad (1)$$

125 where \mathbf{A} is the MIPAS averaging kernel (AK, with dimension 60×60) matrix that contains information about the vertical sensitivity, $\mathbf{x}_{\text{model}}$ is the profile sampled from a transport model, $\mathbf{x}_{\text{prior}}$ is the prior profile, and \mathbf{x}_{conv} the model profile convolved with the AK. In practice, the prior for MIPAS COS retrieval is a zero profile, so Eq. (1) simplifies to:

$$\mathbf{x}_{\text{conv}} = \mathbf{A}\mathbf{x}_{\text{model}}. \quad (2)$$

The convolved COS mixing ratios are compared to MIPAS retrievals and then included in the cost function for inverse
130 modelling. In this study with a focus on tropospheric COS, we only use the 8th level of the MIPAS COS retrievals. To make best use of the MIPAS data, we applied data quality control, and soundings that match one of the following criteria were removed and the purposes are indicated:

1. To avoid cloud contamination, soundings with a MIPAS visibility flag of zero are discarded.
- 135 2. To avoid low sensitivity indicated by the AK, soundings with a diagonal element of the 8th AK row smaller than 0.03 are removed.
3. To avoid artifacts due to orography, soundings with a difference of more than 6000 Pa between the MIPAS reported surface pressure and the TM5 surface pressure are removed.

Fig. 2 shows filtered MIPAS data based on the annual mean of 2009 and a time series from 2008 to 2010. Panel (a) demonstrates that the 8th row of the AK roughly peaks at around 200 hPa, indicating a retrieval sensitivity to the troposphere, at least
140 in the tropics. To motivate our selection for the 8th MIPAS level, panel (b) shows the latitudinal distribution of the MIPAS COS mole fraction at levels 5, 7, 8, 10, and 12. The MIPAS measurements for levels 5, 7, 8 are relatively similar, but COS mole fractions drop quickly for level 10 and above, due to enhanced sensitivity to stratospheric COS, as shown in the mean COS profile in panel (c). Panel (d) shows the total number of valid MIPAS data after applying the data quality control. It is evident that the number of valid MIPAS soundings drops when the level is closer to Earth's surface, due to the influence of clouds. In
145 summary, the 8th MIPAS level is chosen to best represent tropospheric COS, while still retaining a substantial number of valid measurements.

2.3 Validation observations

Since the COS inverse problem is data-constrained, independent data are crucial to further validate the inversion results, particularly with respect to the modelled vertical distribution. This is particularly true when assimilating surface observations
150 combined with MIPAS data, which has the highest sensitivity in the upper troposphere. For clarification, independent or validation data refers to observations that are not assimilated in the inverse model. In principle, evaluation with independent observations can help pinpointing potential model errors or poorly estimated flux adjustments.



155 HIAPER Pole-to-Pole Observations (HIPPO) data provides valuable and well-calibrated COS concentrations between January 2009 and September 2011, a period with valid MIPAS data. HIPPO data were collected during several campaigns (see Fig. 1) between the surface and the tropopause, and from the NH to the SH (Wofsy, 2011; Wofsy et al., 2017). HIPPO COS measurements were evaluated to be consistent with the NOAA surface network. We use data from HIPPO campaigns 1–3 to validate the inversion results. Next to HIPPO data, we use NOAA airborne profile data that are routinely collected. Most profile measurements are taken in North America, shown in Fig. 1 as red crosses.

3 Inverse model

160 Inverse modelling is a widely-used mathematical technique to estimate surface fluxes of trace gases, constrained by concentration observations at ground level or from space. The inverse model usually consists of a transport model and an observational system, coupled to an optimizer that minimises the difference between observations and model results by changing the emissions. The transport model describes physical and chemical processes: advection in three dimensions, parameterisation of deep convection and other sub-grid-scale mixing processes, and chemistry (Krol et al., 2005). The observational system in this study includes the NOAA surface network and MIPAS data, and these data provide constraints to estimate surface fluxes of COS and CS₂. Note that we do not optimise DMS surface fluxes in this study.

3.1 Prescribed surface fluxes of COS, CS₂ and DMS

170 In the model implementation, we used the prior surface fluxes of the three tracers from Ma et al. (2021). Here, four categories are predefined for COS surface fluxes: anthropogenic, biosphere, ocean and biomass burning. We briefly describe these fluxes below and prior flux information is listed in Table 1.

The reported anthropogenic COS emissions range between 223–586 GgS a⁻¹ (Zumkehr et al., 2018) in the period 2000–2012. Specifically for 2009, the focus year of this study, the emissions amount to 354.2 GgS a⁻¹. The anthropogenic sources are available as monthly estimates on a high spatial resolution resolution of 0.1° × 0.1°. Furthermore, the Zumkehr et al. (2018) inventory reports direct COS and CS₂ emissions, and we used the CS₂ to COS ratios of the various emission types to split emissions into direct COS emissions and indirect CS₂ emissions following Table 1 of Lee and Brimblecombe (2016). In addition, a small amount of DMS emissions over land (6 GgS a⁻¹) is also considered as an indirect COS source. DMS and CS₂ are treated as separate tracers that are quickly oxidized to COS (see below). The biosphere flux is the largest sink for COS and includes uptake by plants and soil at the surface. Here, we use the output generated by SiB4 model (Kooijmans et al., 2021) to drive the TM5 model. These fluxes are still evaluated assuming a constant atmospheric COS mole fraction of 500 pmol mol⁻¹. This leads to a prior flux estimate of –1053 GgS a⁻¹. Fluxes are provided as monthly fields on a 1° × 1° resolution. Ocean fluxes of COS, CS₂ and DMS are based on climatological fields (Kettle et al., 2002; Suntharalingam et al., 2008; Lana et al., 2011). The COS oceanic flux depends on sea surface temperature and thus can be either a source or sink, depending on the season. COS biomass burning emissions considered biomass burning and biofuel use, with an amount of 136 GgS a⁻¹ (Ma et al., 2021).



185 3.2 Chemistry

The chemical conversion of CS₂ and DMS is handled as described in (Ma et al., 2021). Briefly, CS₂ is converted to COS through reaction with OH assuming a yield of 0.83 (Stickel et al., 1993). DMS is converted to COS using a yield of 0.007 (Barnes et al., 1994) assuming an exponential decay to COS with a timescale of 1.2 days (Khan et al., 2016). Recent studies found a new DMS oxidation product HPMTF that potentially plays a role in COS production (Wu et al., 2015; Veres et al., 190 2020; Fung et al., 2022; Jernigan et al., 2022). Here, we still use the assumption of a 0.7% yield from DMS oxidation (Barnes et al., 1994). Note that the reaction rate of COS and OH quoted in Ma et al. (2021) contained a typo. Here we give the correct equation:



where T is temperature in K, and A is pre-exponential factor of Arrhenius equation ($1.13 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1} \text{ molecule}^{-1}$) for the reaction of COS with OH (Cheng and Lee, 1986). As in Ma et al. (2021) we used the OH climatology by Spivakovsky et al. (2000) multiplied with 0.92 in the troposphere.

3.3 TM5-4DVAR system

TM5-4DVAR is an offline global inverse model using 4DVAR data assimilation techniques (Krol et al., 2005, 2008; Meirink et al., 2008; Bergamaschi et al., 2010). The 3-hourly meteorological fields from the European Centre for Medium-Range 200 Weather Forecasts (ECMWF) ERA-Interim reanalysis is employed to drive the offline TM5 model (Dee et al., 2011). The model settings are generally the same as in Ma et al. (2021).

The forward model consists of a transport operator and an observation operator that acts on a state vector \mathbf{x} :

$$\mathbf{y} = \mathbf{H}(\mathbf{x}). \quad (3)$$

The TM5 model (\mathbf{H}) simulates the mole fraction of COS, after which the misfit between model and measurement (surface and/or satellite) is calculated, co-sampled in space and time. In this study the operator is linear, and Eq. (3) can be written as $\mathbf{H}\mathbf{x}$. The value of the cost function \mathbf{J} (Eq. (4)) is calculated with the forward model and the derivative of the cost function with respect to the state vector elements is calculated with the adjoint model \mathbf{H}^T . An optimisation algorithm is used to minimize the cost function up to a pre-defined gradient reduction. Eq. (4) is a classic form of a cost function in an inverse system (Tarantola, 2005; Brasseur and Jacob, 2017):

$$210 \quad \mathbf{J}(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \mathbf{x}_b)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + \frac{1}{2}(\mathbf{H}\mathbf{x} - \mathbf{y})^T \mathbf{R}^{-1}(\mathbf{H}\mathbf{x} - \mathbf{y}), \quad (4)$$

and the derivative of the cost function w.r.t the state vector \mathbf{x} reads:

$$\nabla_{\mathbf{x}} \mathbf{J}(\mathbf{x}) = \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + \mathbf{H}^T \mathbf{R}^{-1}(\mathbf{H}\mathbf{x} - \mathbf{y}), \quad (5)$$



where \mathbf{x} is the state vector to be optimized, \mathbf{x}_b is the prior state vector. The covariance matrix \mathbf{B} describes the uncertainty statistics associated with the state vector. The covariance matrix \mathbf{R} describes errors related to the misfit between observations and the model ($\mathbf{H}\mathbf{x} - \mathbf{y}$). While \mathbf{R} is (often) assumed to be diagonal, off-diagonal elements of \mathbf{B} are determined by user-prescribed spatial and temporal correlations of the fluxes (details of the used correlation lengths can be found in Ma et al. (2021)). The inverse of \mathbf{B} is not explicitly calculated owing to its huge size, but dealt with during pre-conditioning and eigenvalue decomposition (Meirink et al., 2008).

Our implementation of TM5-4DVAR runs on a spatial resolution of 6×4 degrees horizontally, and vertically on 25 pressure levels. Fluxes are optimized on a monthly time-scale. In this study, we focus on a 3-year inversion from January 1, 2008 to January 1, 2011. The initial condition is based on a previous 3-year inversion in which atmospheric COS was spun up using atmospheric surface observations. The inversion model optimises all COS and CS₂ fluxes, but keeps the DMS fluxes constant, as scenario S1 described in Ma et al. (2021). Since the inverse system remains linear, including the comparison to MIPAS satellite data, we use the CONGRAD algorithm (Lanczos, 1950) to optimize the fluxes until the gradient norm is reduced by a factor of 10^{-7} . We found that with this convergence criterion, the leading eigenvalue of the Hessian of the cost function is close to 1.0, which implies that both the fluxes, as well as the posterior errors can be reliably interpreted (Meirink et al., 2008). Using the posterior co-variance matrix, the error reduction (\mathbf{ER} , defined in Eq. (6)) and correlations of the optimized fluxes can be evaluated.

$$\mathbf{ER} = 1 - \frac{\mathbf{error}_{posterior}}{\mathbf{error}_{prior}} \quad (6)$$

We report correlations and error reductions of COS and CS₂ fluxes based on error aggregation over NH lands, NH oceans, SH lands and SH oceans during the period of 2009 in Sect. 4.5. The years 2008 and 2010 are considered as spin up and spin down period, respectively.

3.4 Bias correction

A bias may exist amongst different measurement streams, e.g. between the NOAA surface network and the MIPAS satellite data. This bias may be due to different instruments or calibration methods. In addition, since some of the sensitivity of the MIPAS data extends to the stratosphere (see Fig. 2(a)), some bias may be introduced by incorrect transport to, and chemistry in the stratosphere. A bias correction scheme may successfully correct for systematic satellite measurement biases (Monteil et al., 2013; Houweling et al., 2014). To correct for potential biases, we implemented a bias correction to MIPAS data with monthly and latitudinal variations. The prior values of the bias correction are assumed to be 1.0, i.e. no bias existing between different measurement systems. The cost function of the state vector of surface fluxes can be extended with a bias correction as:

$$\begin{aligned} J(\mathbf{x}, \boldsymbol{\beta}) = & \frac{1}{2}(\mathbf{x} - \mathbf{x}_b)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + \frac{1}{2}(\boldsymbol{\beta} - \boldsymbol{\beta}_b)^T \mathbf{B}_\beta^{-1}(\boldsymbol{\beta} - \boldsymbol{\beta}_b) + \\ & \frac{1}{2}(\mathbf{H}\mathbf{x} - \mathbf{y}_{\text{NOAA}})^T \mathbf{R}_{\text{NOAA}}^{-1}(\mathbf{H}\mathbf{x} - \mathbf{y}_{\text{NOAA}}) + \frac{\gamma}{2}(\mathbf{H}\mathbf{x} - \mathbf{y}_{\text{MIPAS}}\boldsymbol{\beta})^T \mathbf{R}_{\text{MIPAS}}^{-1}(\mathbf{H}\mathbf{x} - \mathbf{y}_{\text{MIPAS}}\boldsymbol{\beta}), \end{aligned} \quad (7)$$



$$\gamma = \frac{1}{\mathbf{e}\mathbf{i}}. \quad (8)$$

The derivative of the cost function with respect to the bias terms β then becomes:

$$\nabla_{\beta} \mathbf{J}(\mathbf{x}, \beta) = \mathbf{B}_{\beta}^{-1}(\beta - \beta_{\mathbf{b}}) - \gamma \mathbf{R}_{\text{MIPAS}}^{-1}(\mathbf{H}\mathbf{x} - \mathbf{y}_{\text{MIPAS}}(\beta)) \mathbf{y}_{\text{MIPAS}}. \quad (9)$$

245 In these equations $\mathbf{R}_{\text{MIPAS}}$ and \mathbf{R}_{NOAA} are the error covariance matrices associated with NOAA surface observations and MIPAS data, respectively. β is the bias correction parameter assigned to MIPAS measurements, and γ is a regularization factor, which scales the MIPAS data part of the cost function. Eq. (8) relates the MIPAS error inflation to γ . The assumption here is that NOAA surface observations and MIPAS profiles are all independent, so that the error covariance matrix for all data in Eq. (4) is diagonal and can be separated into two matrices. The regularisation factor is similar with the terminology defined
250 in Brasseur and Jacob (2017). In our case, the number of MIPAS measurements (127243 in 2009 after data quality control) overwhelms the number of NOAA surface observations (490 in 2009), and the error inflation is a factor that limits the weight of MIPAS data in the cost function. Note that the error inflation also can account for potential correlations between the various MIPAS soundings. We determined that an error inflation of 10 leads to a reasonable balance between MIPAS and NOAA terms in the cost function.

255 The derivative w.r.t β (Eq. (9)) is implemented in the data assimilation system, such that β is simultaneously optimized along with the fluxes. As discussed in Sect. 2, we selected the 8th level of MIPAS data near 200 hPa to be assimilated in the system. In the assimilation, a mass conservation mapping algorithm is used to map the 25 TM5 pressure levels to the 60 MIPAS levels.

The bias parameters β are multiplied to the MIPAS observations (Eq. (7)). The prior values of bias parameters are set to
260 1.0 ± 0.003 as shown in Table 1. The small error assigned to bias parameters is to prevent the MIPAS data from being adjusted too strongly. We furthermore assume that the MIPAS bias depends on month and latitude, and not on longitude. The monthly and latitudinal bias corrections are assumed to be correlated by decay functions that fall off in space and time. For the latitudinal variations, a Gaussian decay function is selected:

$$\text{corr}_{\mathbf{h}}(i_{\text{lat}}, j_{\text{lat}}) = \exp\left(\frac{-(i_{\text{lat}} \cdot \text{bin} - j_{\text{lat}} \cdot \text{bin})^2}{2\mathbf{L}^2}\right), \quad (10)$$

265 where i_{lat} and j_{lat} are the two correlated latitude indices, and bin is the bin size that determines how many bins taken into account. For MIPAS we take into account the latitude range from 80°S to 80°N and 16 latitude bins are applied, so bin equals to 10°. \mathbf{L} is the spatial correlation length, which is set to 5°. For the temporal correlation, an exponential decay is assumed:

$$\text{corr}_{\mathbf{t}}(i_{\text{month}}, j_{\text{month}}) = \exp\left(\frac{-|i_{\text{month}} - j_{\text{month}}|}{\tau}\right), \quad (11)$$

where i_{month} and j_{month} are the two correlated month indices, and τ is the temporal correlation length, taken as 9.5
270 months. Finally, the covariance matrix \mathbf{B}_{β} is written as a Kronecker product of Eqs. (10) and (11).



3.5 Evaluation metrics

To evaluate the performance of the inverse system, it is important to compare the model with assimilated and unassimilated data, e.g. either ground-based observations or satellite observations. We use the statistical metrics χ^2 to characterise the performance of the inversions, by comparing the surface network observations and inversion results.

275 When the observations are assumed independent, χ^2 is twice of the cost function (Tarantola, 2005):

$$\chi_N^2 = (\mathbf{x} - \mathbf{x}_b)^T \mathbf{B}^{-1} (\mathbf{x} - \mathbf{x}_b) + \sum_{k=1}^N \frac{(\mathbf{H}\mathbf{x} - \mathbf{y})^2}{\sigma^2} \quad (12)$$

In a well-balanced inverse problem, χ^2_N is equal to the number of observations N . Since the background part of the cost function is usually small, and we are interested in the performance of the inversion at different stations, we prefer to use the station-based χ^2 to characterize the performance of inversions. For comparison between model results and NOAA stations, the station-based χ^2 is defined as:

280

$$\chi^2 = \frac{1}{N} \sum_{i=1}^N \frac{(\mathbf{H}\mathbf{x} - \mathbf{y}_i)^2}{\sigma_i^2}, \quad (13)$$

where i denotes the i^{th} station, and N is the number of observations at that station. Note that a χ^2 of 1 denotes a statistically good fit to the available observations, i.e. the difference between model result and measurements is of the same size as the spread of the observations. For comparison between model and satellite data, χ^2 is also used to compare the modelled and MIPAS measured COS without taking the error inflation in order to avoid small values of the satellite χ^2 .

285

In this study we also analyse regional correlation and error reduction on various spatial and temporal scales, as outlined in Sect. 3.3. Aggregation to different scales is performed by accounting for the covariance from the full prior and posterior covariance matrices. Results will be presented in Sect. 4.4 and 4.5.

3.6 Overview of inversions

290 To explore various model settings and simulation scenarios, we have designed a set of inversions, shown in Tables 2 and 3. The inversion scenario labelled S1 represents similar setting as used in Ma et al. (2021), with 50% prior errors for the COS biosphere fluxes, and COS & CS₂ ocean fluxes. To investigate the impact of the prior error settings on the posterior biosphere fluxes, in scenario S0 the biosphere error was reduced to 10%, while the COS anthropogenic and biomass burning emission errors were increased from 10% to 50%. For scenario S1, we assimilate only NOAA surface observations (NOAA-only_S1), only MIPAS data (MIPAS-only_S1) and both observational datasets with (MIPAS+NOAA_bias=0.3%_S1) and without bias correction (MIPAS+NOAA_no-bias_S1). For scenario S0, we only consider assimilation of NOAA and MIPAS data with bias correction (MIPAS+NOAA_bias=0.3%_S0).

295



4 Results

4.1 Posterior fit to observations

300 To demonstrate the performance of each inversion, we first evaluate the posterior fit at the 14 NOAA surface stations. Fig. 3 compares the various inversions at 6 selected stations. At Alert (ALT, Fig. 3a), MIPAS-only_S1 severely overestimates COS observations, while the other inversions that assimilate NOAA data match the observations well. The same pattern is found at station LEF (Fig. 3b). However, the posterior fit is less good at oceanic stations MLO and SMO (Fig. 3c and 3d). Here, the best results are obtained for the NOAA-only_S1 inversion, which is logical because only NOAA surface data are assimilated. When
305 MIPAS data are also assimilated, the model fit at MLO and SMO deteriorates, which implies trade-offs between assimilation of surface and MIPAS data. At the Antarctica stations SPO and PSA (Fig. 3e and 3f) the model fit improves again. Overall, the MIPAS-only_S1 inversion overestimates COS mixing ratios at all surface stations. This is the result of the low sensitivity of MIPAS to COS that resides close to the surface. It also indicates that MIPAS information is biased high within the TM5-4DVAR system. As outlined above, this might imply a true bias of MIPAS compared to NOAA observations, or a bias in
310 the model transport and chemistry. There is also a high bias of MIPAS retrievals compared to ACE-FTS profiles in the upper troposphere in certain regions (Glatthor et al., 2017).

To evaluate the performance of the various inversions (except for MIPAS-Only) at each NOAA station, Fig. 4 shows χ^2 per station (Eq. 13). The stations are sorted from South to North. As expected, the best performance is found for NOAA-only_S1 (in blue). Values of χ^2 close to one are found for CGO, LEF and MHD. Reasonable results are obtained at the stations, with
315 χ^2 values less than 5. Although the agreement with observations improves considerably w.r.t. the prior (not shown), this could indicate that error settings for these station were too conservative, possibly related to unaccounted processes that influence the COS budget. Indeed we note considerable variability at remote ocean stations like MLO and SMO (see Fig. 3) that is not reproduced well by the model.

In Fig. 4, the worst performance is found for inversion MIPAS+NOAA_no-bias_S1 (excluding MIPAS-only inversion), with
320 a deteriorated model fit especially at SMO and MLO. When bias correction is applied in inversion MIPAS+NOAA_bias=0.3%_S1, the model fit improves at all NOAA stations. When the prior errors are reduced for the biosphere flux (and increased for the anthropogenic and biomass burning fluxes, inversion MIPAS+NOAA_bias=0.3%_S0), the posterior fit degrades at most stations, except for MLO. This indicates that adjustments of the biosphere fluxes play an important role in the goodness of the posterior fit. However, alternative choices of the prior error setting (e.g. scenario S0) still lead to satisfactory results.

325 Next we evaluate the posterior fit to MIPAS data. The posterior model fit with MIPAS data for inversion MIPAS+NOAA_bias=0.3%_S1 is shown in Fig. 5. The mismatch (panel c) shows that the model, after bias correction, still slightly underestimates MIPAS COS mixing ratios at high latitudes, and overestimates values over tropical regions. As expected from assimilated observations, the probability density function (panel d) indicates that the average modelled COS mixing ratio is relatively close to the MIPAS average. However, due to the large error on individual MIPAS soundings, the distribution of MIPAS is much
330 wider. Supplementary information Figs. S1 and S2 show the posterior model fit with MIPAS data for inversion MIPAS-only_S1 and MIPAS+NOAA_no-bias_S1, respectively, where the bias correction scheme is not applied. In the case MIPAS-



only_S1, the MIPAS-based χ^2 amounts to 16.4, and in the case MIPAS+NOAA_no-bias_S1 the χ^2 increased to 17.33. This implies that when both NOAA and MIPAS are assimilated, the fit is deteriorated to some extent. Interestingly, in the case MIPAS+NOAA_bias=0.3%_S1 the fit to MIPAS improved ($\chi^2=14.8$). In general, inversion MIPAS+NOAA_bias=0.3%_S1 reached an acceptable posterior model fit both at NOAA surface stations and to MIPAS data that are sensitive to the upper troposphere.

4.2 Optimized bias correction parameters

Fig. 6 shows the posterior bias parameters as a function of month and latitude for inversion MIPAS+NOAA_bias=0.3%_S1. The posterior bias parameters are in the range 0.94–1.00, indicating that bias correction slightly reduces MIPAS observations in order to reconcile the fit to surface observation. In the tropics, only a small bias correction is found, while higher latitudes show larger corrections. At these latitudes there is also a slight seasonal cycle in the bias parameters, with larger corrections in local winter. Since in local winter high latitude MIPAS observations are mostly sensitive to the stratosphere, this could indicate potential model errors in transporting COS to the stratosphere. The other inversion scenario (S0, not shown) shows similar posterior bias parameters within the range of 0.94–1.0 with similar seasonal and latitudinal variations.

To show the overall performance of all inversions, Table 4 breaks down the cost function in terms of background costs, NOAA costs, and MIPAS costs. The background part contains both the flux part and the costs associated with bias correction (see Eq. (7)). Inversion NOAA-only_S1 has the smallest posterior cost, since only deviations with surface observations are taken into account. For this inversion, the background costs are 15% of the model-data mismatch. Whenever MIPAS data is assimilated, the satellite cost function becomes the dominant part, even though an error inflation of 10 is considered. It is instructive to look at the increase in the background term when bias correction is considered (MIPAS+NOAA_bias=0.3%_S1 and MIPAS+NOAA_bias=0.3%_S0). As expected within an inverse modelling framework, this increase in background costs is more than compensated by reductions in both the MIPAS costs and surface costs. It also shows that additional adjustments to the fluxes are made to comply with both the surface and satellite data.

4.3 Validation with HIPPO and NOAA airborne profiles

In this section, independent HIPPO and NOAA airborne profiles observations are used to evaluate the inversions. The evaluation of HIPPO campaigns 1–3 is shown in Fig. 7. Notably, as can be seen in the first two rows of Fig. 7, the NOAA-only_S1 inversion underestimates COS in the troposphere, while inversion MIPAS-only_S1 generally overestimates COS in the lower troposphere. The other inversions that assimilate MIPAS data decrease the bias against HIPPO observations. One remarkable feature in the comparison with HIPPO campaign 1 in January 2009 is the model overestimate around the equator, that gets reduced in inversion MIPAS+NOAA_bias=0.3%_S0. This is caused by smaller biosphere flux adjustments that are the result of the tighter error settings (prior error reduced from 50% to 10%, see Table 2). The HIPPO 1 air samples were taken close to the Amazonia (Fig. 1). This indicates that the adjustments to the biosphere over the Amazonia in scenario S1 are likely too large. This will be addressed further in Sect. 4.4.



The vertical distributions of inversions and HIPPO observations are shown in supplementary Figs. S3 and S4. In HIPPO
365 campaign 1, the vertical profiles of the inversions are in good agreement with HIPPO vertical distribution, with differences less
than 20 pmol mol^{-1} below 10 km. For HIPPO campaigns 2 and 3, there is a negative bias between the inversions and HIPPO
observations. This points to underestimated oceanic contributions, as HIPPO campaigns 2 and 3 mainly covered the Pacific
Ocean.

The evaluation against NOAA airborne profiles is presented in Fig. 8. Note that the data were grouped to two regions:
370 Alaska and the rest of North America as shown in Fig. 1. All inversions (except for MIPAS-only_S1) fit the profiles relatively
well. Again, inversion MIPAS-only_S1 (in green) overestimates NOAA airborne observations significantly. Other inversions
are generally slightly lower than the NOAA airborne observations, with slightly larger deviations when MIPAS bias correction
is applied. The NOAA airborne profiles were mainly acquired over continents, where the inversions seem generally well-
constrained by the NOAA surface network. Assimilation of MIPAS data does not strongly influence the agreement with the
375 observed profiles. As shown in supplementary Fig. S5, we still find a low bias in our inversions, which however remains well
below 20 pmol mol^{-1} over North America.

4.4 The global and Amazonia COS budget

The global COS budgets of all inversions are graphically summarized in Fig. 9. Furthermore, the regional prior and posterior
COS and CS_2 fluxes, errors and error reductions are listed in Table 5.

380 The prior COS budget is not in balance with a net excess sink of -432 GgS a^{-1} , as described in Ma et al. (2021). The
net fluxes of all inversion experiments are close to zero, which demonstrates the capability of the inverse model to close the
gap in the global budget. All inversions reduce the biosphere uptake, albeit with different amounts depending on the inversion
settings. Sometimes the biosphere sink even turns into a source, e.g. over the Amazonia (supplementary Fig. S6 from inversion
NOAA-only_S1). Although the MIPAS-only_S1 inversion overestimates COS both at the surface (Fig. 3) and in the lower
385 troposphere (Fig. 8), the tropical biosphere remains a sink in this inversion (Fig. 10). When NOAA and MIPAS data are co-
assimilated, the tropical biosphere still turns locally into a source, but less strong than for inversion NOAA-only_S1 (Fig. 10).
From Table 5, it is clear that most of the error reduction is obtained for the biosphere flux. Ocean fluxes of both COS and CS_2
only see marginal error reductions of a few percent at most. The error reduction on anthropogenic emissions is also small for
the S1 inversions. However, as expected, error reductions become larger when their prior errors are increased from 10 % to 50
390 % in scenario S0. For instance, error reductions on the biomass burning emissions are increased from 0.4% in scenario S1 to
31.0% globally in scenario S0.

In scenario S1, by far the largest adjustments are made to the tropical biosphere to close the COS budget. It is questionable,
however, whether positive fluxes from the Amazonia are realistic. In inversion MIPAS+NOAA_bias=0.3%_S0, the biosphere
remains a sink, as shown in Fig. 10. By reducing the prior error on the biosphere from 50% to 10%, the global biosphere prior
395 error of 191.2 GgS a^{-1} in scenario S1 decreased to 38.2 GgS a^{-1} in scenario S0 (Table 5). To leverage the decrease of freedom,
the biomass burning and anthropogenic errors were increased to 50% in S0. Interestingly, despite the need for a tropical source,
biomass burning emission showed a decrease from 126.6 ± 27.5 to $95.8 \pm 18.9 \text{ GgS a}^{-1}$ in scenario S0. This might imply that



biomass burning is not responsible for the COS budget gap, in line with findings from Stinecipher et al. (2019). Instead, the budget is closed by increasing the anthropogenic CS₂ and COS emissions and CS₂ ocean emissions. For instance, the global
400 CS₂ ocean source increased from 83.0±12.0 to 171.2±11.6 GgS a⁻¹. Note that direct COS ocean emissions decreased in all scenarios (except for MIPAS-only) because COS ocean exchange occurs mostly at high latitudes. Note also that many of the adjustments, especially in scenario S0, are large compared to the error settings. For instance, the global biosphere COS flux is adjusted from -1053.0±38.2 to -686.5±22.9 GgS a⁻¹. A similar large adjustment is seen for the oceanic CS₂ emissions. This points to too tight prior error settings in scenario S0. However, the inversion system has to correct for the missing tropical
405 source, which might be a structural model error (i.e. missing processes, incorrect regional or temporal correlation settings of sources and sinks). Adjusting the tropical fluxes outside the predefined prior error ranges is the only way to close the budget. In scenario S1 the error settings seem more realistic. However, as discussed above, the biosphere often turns into a source in the S1 scenario. These results underscore the importance of a proper quantification of the prior fluxes and their associated errors in inverse systems.

410 To investigate how well the different fluxes can be separated in our inversions, Fig. 11 shows posterior correlations of total COS fluxes aggregated over the different regions. In defining the fluxes, we quantified positive spatial and temporal correlations in the prior fluxes. After inversion, however, the correlations often turn strongly negative when the inversions cannot properly separate the fluxes. It can be inferred from Fig. 11 that the posterior fluxes from NH land and ocean are less (negatively) correlated than the fluxes from SH land and ocean. This implies that the inverse model can better separate NH
415 land and ocean sources or sinks. This is because most NOAA surface observations are located in NH. The MIPAS-only_S1 and NOAA-only_S1 show high anti-correlations (-0.67 and -0.65) between SH land and ocean, which indicates poor separation of land and ocean fluxes. The case MIPAS+NOAA_no-bias_S1 and MIPAS+NOAA_bias=0.3%_S0 decrease the correlations between SH land and ocean to -0.55 and -0.38, respectively. This indicates that inclusion of MIPAS data in the inversion leads to a better separation of land and ocean fluxes. Also the prior error specification has a large impact on the posterior
420 correlations. Specifically, the MIPAS+NOAA_bias=0.3%_S0 inversion helps the separation of COS sources and sinks between land and ocean. The posterior CS₂ correlations generally remain similar to the prior (see supplementary Fig. S7), because error reductions for CS₂ fluxes are generally marginal, except in scenario S0 (MIPAS+NOAA_bias=0.3%_S0).

Table 6 summarizes the COS and CS₂ budgets, calculated for the Amazonia. The Amazonia COS budgets are aggregated by adding the regions South American Temperate and South American Tropical of the 22 Transcom regions (Fig. 2 in Feng et al.
425 (2011)). The prior biosphere flux of the Amazonia is -311.0 GgS a⁻¹, and all the inversions tend to decrease the COS uptake by the biosphere, implying an overestimated prior biosphere sink or an unknown source over the Amazonia. Notably, inversion NOAA-only_S1 obtained a posterior biosphere flux of -10.6±48.0 GgS a⁻¹. Interestingly, a fraction of the Northwestern Amazonia turned in a source (see Fig. 10) that may be questioned in light of the overestimated HIPPO 1 samples (Fig. 7). Inversion MIPAS-only_S1 obtained a larger posterior biosphere flux over the Amazonia of -145.7 GgS a⁻¹, which is however
430 still less than half of the prior sink. When MIPAS and NOAA observations are co-assimilated, inversion MIPAS+NOAA_no-bias_S1 reverses the biosphere sink into a source of +15.5±25.9 GgS a⁻¹. The biosphere again remains a sink when bias



correction is applied. Scenarios S1 and S0 with bias correction lead to biosphere flux of $-99.5 \pm 26.3 \text{ GgS a}^{-1}$ and $-108.2 \pm 15.8 \text{ GgS a}^{-1}$, respectively. Comparison to HIPPO 1 (Fig. 7) shows the smallest biases for scenario MIPAS+NOAA_bias=0.3%_S0.

4.5 Error reduction

435 In this section, maps of the error reduction are analysed. The error reduction on the grid-scale of the model, aggregated over all direct COS fluxes, is shown in Fig. 12. For reference, the regional and global error reductions are shown in parenthesis in Table 5. Inversion MIPAS-only_S1 mainly reduces the prior error over tropical regions, covering the Amazonia region, Africa and Central Asia. In contrast, inversion NOAA-only_S1 reduces the error mostly over the NH and mainly over North America, where the NOAA surface network is more dense. On the grid scale, the SH sees little error reduction, partly due to the lack of surface observational constraints, but also because the ocean surface fluxes are distributed over a larger area compared to the biosphere fluxes. Inversions MIPAS+NOAA_no-bias_S1 and MIPAS+NOAA_bias=0.3%_S1 reduce the errors again globally over the continents. This demonstrates the complementary effects of NOAA observations and MIPAS data on constraining the COS budget. Concerning error reductions for scenario S0, the error reduction is substantially smaller, because the prior error on the biosphere flux is lowered from 50% to 10%. This largely limits the degrees of freedom to adjust the biosphere. It should be remembered that errors in our inverse system are defined as percentage of the fluxes themselves. Biosphere fluxes are generally more localized and larger than ocean fluxes. As a result, the error reduction of the COS flux over the oceans is small compared to the error reduction of the land fluxes (see Fig. S8). Again, this points to the need to carefully construct the prior error covariance matrix \mathbf{B} (Eq. 4). For instance, if we enlarge the prior errors on the CS_2 ocean emissions to 150% based on the S0 scenario (with a biosphere error of 10%), we find that the CS_2 ocean emissions are enlarged to provide the required tropical COS source (see scenario of SCS2X in Fig. S9). In that solution, we find comparable agreement to independent data with a global biosphere flux of $-922.9 \pm 28.7 \text{ GgS a}^{-1}$ ($-178.4 \pm 17.7 \text{ GgS a}^{-1}$ over the Amazonia). Although the scenario is extreme (i.e. small freedom to adjust the tropical biosphere flux and large freedom to adjust the ocean CS_2 flux), this inversion shows the need for accurate prior fluxes, including realistic errors.

The error reduction in the CS_2 total flux (supplementary Fig. S10) is significantly smaller compared to COS (Fig. 12). Again, this is attributed to the small prior errors shown in supplementary Fig. S11. In scenario S0, in which the prior error on CS_2 anthropogenic emissions is increased to 50%, larger error reductions are seen over land, e.g. close to 30% over NH land areas. Interestingly, the CS_2 ocean fluxes roughly double in inversion S0, however, with only a small error reduction. This is clearly driven by the need for a tropical COS source, and the limited ability in scenario S0 to adjust the biosphere. It seems therefore logical to increase the prior errors on ocean CS_2 emissions in future inversions.

460 5 Discussion

A general feature of inversion results is that the tropical biosphere uptake is reduced, depending on the inversion settings. Recently, Stinecipher et al. (2022) reported that a smaller COS biosphere sink of $-81.1 \pm 28.1 \text{ GgS a}^{-1}$ over the Amazonia (i.e. a low GPP model) leads to the best agreement to MIPAS observations, which is consistent with our findings. However, many



of our inversions turn parts of the Amazonia into a net source, and we question the realism of this result. This can potentially
465 be the case that biosphere uptake is overestimated, or emissions from anthropogenic or biomass burning are underestimated.
Indeed, we show biases when we compare simulations using our optimized fluxes to HIPPO 1 observations (Fig. 7), which
disappear in a scenario in which we reduce the freedom to adjust the biosphere. This underscores the need to better characterize
the prior fluxes, including their error settings. Our scenario S0 currently optimizes fluxes well outside of the prior error settings.
Specifically, a good balance should be found in optimizing the tropical fluxes from land and ocean. Adjustments of the budget
470 terms outside of the prior errors could also point to structural model errors, such as unaccounted processes. For instance,
enhanced production of COS from DMS oxidation through the intermediate HPMTF as recently reported (e.g. Jernigan et al.
(2022)) could be such a process. The difficulty of our inverse system to properly fit observations over remote oceans (e.g. SMO
and MLO, Fig. 3) might be indicative of unresolved processes over the oceans. The enhanced CS₂ flux could also come from
DMS. Note that given the chemical mechanism of DMS oxidation, any COS from DMS enhancement would not scale with
475 the current prior flux, because in some regions with large DMS emissions, the atmospheric conditions almost exclude any COS
production, while in other regions (e.g. the tropical Pacific), the chemical conditions tend to favour COS production.

One of the limitations of the assimilation of MIPAS data in this work is that we only used one specific level of MIPAS
data to constrain the tropospheric COS mole fractions. We could exploit the MIPAS data more, e.g. by assimilating other
levels. However, at higher latitudes the averaging kernels of MIPAS level 8 already mixes information from the stratosphere
480 (Fig. 2), which makes the comparison vulnerable to model errors associated with transport and stratospheric removal. A recent
paper exploited TES data to analyse the seasonal cycle of COS over the Amazonia (Wang et al., 2023). Assimilation of
TES data, for instance in combination with MIPAS data, could be a possibility to further constrain the tropical COS budget.
Unfortunately, like MIPAS, the TES instrument is no longer operational. Thus, more COS observations to better characterise
COS concentrations and fluxes are still urgently needed.

485 6 Conclusions and Outlook

In this study we have implemented the combined data assimilation of surface observations and MIPAS satellite data within the
TM5-4DVAR system. The major conclusions are summarized as follows:

- 2008–2010 inversions show generally good agreement both to NOAA observations and MIPAS data, particularly with
the assistance of a bias correction scheme. The optimized bias parameters show that the bias correction reduced MIPAS
490 COS data at higher latitudes.
- In all inversions, the largest adjustments are made to the biosphere fluxes. As a result, the required tropical COS source
is obtained by strongly reducing the tropical biosphere, e.g. over the Amazonia. Although the large flux adjustments over
the Amazonia are not always consistent with independent HIPPO observations, a yearly Amazonia COS uptake of 100
GgS is much smaller than the SiB4-based prior flux of about -300 GgS a^{-1} and consistent with findings of a recent
495 study by Stinecipher et al. (2022).



- Analysis of the correlations between land and ocean fluxes indicates that assimilation of MIPAS data improves the separation between ocean sources and land sinks. Analysis of the error reduction shows that MIPAS data generally adds information compared to a NOAA-only inversion.
- A scenario in which the prior biosphere errors are reduced (S_0) leads to a more realistic solution of the inverse problem. However, in this scenario other fluxes are often optimized outside of the error settings. This could point to missing COS production terms in the model (e.g. from DMS), but also underscores the need to carefully construct the prior error covariance matrix \mathbf{B} (Eq. 4).

500

505

Overall, our study shows that, although MIPAS data provides useful additional information, closure of the COS budget remains to some extent unresolved. Additional information could come from assimilation of satellite data from TES, ACE-FTS, and IASI and data from NDACC (Hannigan et al., 2022). On top of that, additional characterization of the different COS sources (e.g. from oceans and DMS oxidation), and sinks (e.g. soils and biosphere) is recommended.

Code and data availability. MIPAS satellite COS measurements are available at <https://www.imk-asf.kit.edu/english/308.php>. NOAA surface measurement of COS is available at <https://www.esrl.noaa.gov/gmd/dv/data/>. HIPPO COS observations are available at https://www.eol.ucar.edu/field_projects/hippo. Model codes of TM5-4DVAR are available on the TM5-4DVAR website <https://sourceforge.net/projects/tm5/>.

510

Author contributions. JM and MC designed the study and implemented the algorithms. JM performed numerical experiments and data analysis. TR and MvH gave valuable discussions and inputs on this work. LMJK provided SiB4 biosphere flux for model input. NG provided MIPAS data and helpful discussions on usage of MIPAS data. SAM provided NOAA surface network. JM and MK wrote the manuscript with comments from all co-authors.

515

Competing interests. The authors declare that they have no conflict or competing of interests. At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

520

Acknowledgements. This work is carried out on the Dutch National Supercomputer Infrastructure with the support from Surfsara and Snelius. The authors acknowledge Michael Kiefer at KIT for the MIPAS data product, and Lei Hu at NOAA Global Monitoring Laboratory for NOAA airborne platform measurements. The authors acknowledge the NOAA team for providing the NOAA surface data and maintenance of the data. The research has been funded by the European Research Council (ERC) COS-OCS project with grant agreement No 742798 (<http://cos-ocs.eu>).



References

- Abadie, C., Maignan, F., Remaud, M., Ogée, J., Campbell, J. E., Whelan, M., Kitz, F., Spielmann, F., Wohlfahrt, G., Wehr, R., et al.: Global modelling of soil carbonyl sulfide exchanges, *Biogeosciences*, 19, 2427–2463, 2022.
- 525 Baartman, S. L., Krol, M. C., Röckmann, T., Hattori, S., Kamezaki, K., Yoshida, N., and Popa, M. E.: A GC-IRMS method for measuring sulfur isotope ratios of carbonyl sulfide from small air samples, *Open Research Europe*, 1, 105, 2021.
- Barkley, M. P., Palmer, P. I., Boone, C. D., Bernath, P. F., and Suntharalingam, P.: Global distributions of carbonyl sulfide in the upper troposphere and stratosphere, *Geophysical Research Letters*, 35, L14 810, <https://doi.org/10.1029/2008GL034270>, 2008.
- Barnes, I., Becker, K. H., and Patroescu, I.: The tropospheric oxidation of DMS: A new source of OCS, *Geophysical Research Letters*, 21, 2389–2392, <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/94GL02499>, 1994.
- 530 Basu, S., Guerlet, S., Butz, A., Houweling, S., Hasekamp, O., Aben, I., Krummel, P., Steele, P., Langenfelds, R., Torn, M., et al.: Global CO₂ fluxes estimated from GOSAT retrievals of total column CO₂, *Atmospheric Chemistry and Physics*, 13, 8695–8717, 2013.
- Bergamaschi, P., Krol, M., Meirink, J. F., Dentener, F., Segers, A., Van Aardenne, J., Monni, S., Vermeulen, A. T., Schmidt, M., Ramonet, M., Yver, C., Meinhardt, F., Nisbet, E. G., Fisher, R. E., O’Doherty, S., and Dlugokencky, E. J.: Inverse modeling of European CH₄ emissions 2001–2006, *Journal of Geophysical Research Atmospheres*, 115, <https://doi.org/10.1029/2010JD014180>, 2010.
- 535 Bernath, P., Crouse, J., Hughes, R., and Boone, C.: The Atmospheric Chemistry Experiment Fourier transform spectrometer (ACE-FTS) version 4.1 retrievals: Trends and seasonal distributions, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 259, 107 409, 2021.
- Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., Denning, A. S., Kawa, S. R., Montzka, S. A., Seibt, U., et al.: A coupled model of the global cycles of carbonyl sulfide and CO₂: A possible new window on the carbon cycle, *Journal of Geophysical Research: Biogeosciences*, 118, 842–852, 2013.
- 540 Brasseur, G. P. and Jacob, D. J.: *Inverse Modeling for Atmospheric Chemistry*, in: *Modeling of Atmospheric Chemistry*, Cambridge University Press, <https://doi.org/https://doi.org/10.1017/9781316544754.012>, 2017.
- Brühl, C., Lelieveld, J., Crutzen, P. J., and Tost, H.: The role of carbonyl sulphide as a source of stratospheric sulphate aerosol and its impact on climate, *Atmospheric Chemistry and Physics*, 12, 1239–1253, <https://doi.org/10.5194/acp-12-1239-2012>, 2012.
- Campbell, J., Whelan, M., Seibt, U., Smith, S. J., Berry, J., and Hilton, T. W.: Atmospheric carbonyl sulfide sources from anthropogenic activity: Implications for carbon cycle constraints, *Geophysical research letters*, 42, 3004–3010, 2015.
- 545 Campbell, J., Berry, J., Seibt, U., Smith, S. J., Montzka, S., Launois, T., Belviso, S., Bopp, L., and Laine, M.: Large historical growth in global terrestrial gross primary production, *Nature*, 544, 84–87, 2017.
- Campbell, J. E., Carmichael, G. R., Chai, T., Mena-Carrasco, M., Tang, Y., Blake, D., Blake, N., Vay, S. A., Collatz, G. J., Baker, I., et al.: Photosynthetic control of atmospheric carbonyl sulfide during the growing season, *Science*, 322, 1085–1088, 2008.
- 550 Camy-Peyret, C., Liuzzi, G., Masiello, G., Serio, C., Venafra, S., and Montzka, S.: Assessment of IASI capability for retrieving carbonyl sulphide (OCS), *Journal of Quantitative Spectroscopy and Radiative Transfer*, 201, 197–208, 2017.
- Cartwright, M. P., Harrison, J. J., and Moore, D. P.: Retrievals of Atmospheric Carbonyl Sulfide from IASI, in: *EGU General Assembly Conference Abstracts*, pp. EGU21–10 373, 2021.
- 555 Cartwright, M. P., Pope, R. J., Harrison, J. J., Chipperfield, M. P., Wilson, C., Feng, W., Moore, D. P., and Suntharalingam, P.: Modelling atmospheric carbonyl sulfide using gross primary productivity to constrain vegetative uptake, *Atmospheric Chemistry and Physics Discussions*, pp. 1–26, 2022.



- Cheng, B.-M. and Lee, Y.-P.: Rate constant of OH+ OCS reaction over the temperature range 255–483 K, *International journal of chemical kinetics*, 18, 1303–1314, 1986.
- Chin, M. and Davis, D.: Global sources and sinks of OCS and CS₂ and their distributions, *Global Biogeochemical Cycles*, 7, 321–337, 1993.
- 560 Commane, R., Meredith, L. K., Baker, I. T., Berry, J. A., Munger, J. W., Montzka, S. A., Templer, P. H., Juice, S. M., Zahniser, M. S., and Wofsy, S. C.: Seasonal fluxes of carbonyl sulfide in a midlatitude forest, *Proceedings of the National Academy of Sciences*, 112, 14 162–14 167, 2015.
- Crutzen, P. J.: The possible importance of CSO for the sulfate layer of the stratosphere, *Geophysical Research Letters*, 3, 73–76, <https://doi.org/10.1029/GL003i002p00073>, 1976.
- 565 Davidson, C., Amrani, A., and Angert, A.: Tropospheric carbonyl sulfide mass balance based on direct measurements of sulfur isotopes, *Proceedings of the National Academy of Sciences*, 118, 2021.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, d. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological society*, 137, 553–597, 2011.
- 570 Du, Q., Zhang, C., Mu, Y., Cheng, Y., Zhang, Y., Liu, C., Song, M., Tian, D., Liu, P., Liu, J., Xue, C., and Ye, C.: An important missing source of atmospheric carbonyl sulfide: Domestic coal combustion, *Geophysical Research Letters*, 43, 8720–8727, <https://doi.org/10.1002/2016GL070075>, 2016.
- Feng, L., Palmer, P. I., Yang, Y., Yantosca, R. M., Kawa, S. R., Paris, J.-D., Matsueda, H., and Machida, T.: Evaluating a 3-D transport model of atmospheric CO₂ using ground-based, aircraft, and space-borne data, *Atmospheric chemistry and physics*, 11, 2789–2803, 2011.
- 575 Fischer, H., Birk, M., Blom, C., Carli, B., Carlotti, M., Von Clarmann, T., Delbouille, L., Dudhia, A., Ehhalt, D., Endemann, M., et al.: MIPAS: an instrument for atmospheric and climate research, *Atmospheric Chemistry and Physics*, 8, 2151–2188, 2008.
- Friedlingstein, P., O’sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., et al.: Global carbon budget 2020, *Earth System Science Data*, 12, 3269–3340, 2020.
- Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, 580 J., et al.: Global carbon budget 2021, *Earth System Science Data*, 14, 1917–2005, 2022a.
- Friedlingstein, P., O’sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Le Quéré, C., Luijkx, I. T., Olsen, A., Peters, G. P., et al.: Global carbon budget 2022, *Earth System Science Data*, 14, 4811–4900, 2022b.
- Fung, K. M., Heald, C. L., Kroll, J. H., Wang, S., Jo, D. S., Gettelman, A., Lu, Z., Liu, X., Zaveri, R. A., Apel, E. C., et al.: Exploring dimethyl sulfide (DMS) oxidation and implications for global aerosol radiative forcing, *Atmospheric Chemistry and Physics*, 22, 1549–1573, 2022.
- 585 Glatthor, N., Höpfner, M., Baker, I. T., Berry, J., Campbell, J., Kawa, S. R., Krysztofiak, G., Leyser, A., Sinnhuber, B.-M., Stiller, G., et al.: Tropical sources and sinks of carbonyl sulfide observed from space, *Geophysical Research Letters*, 42, 10–082, 2015.
- Glatthor, N., Höpfner, M., Leyser, A., Stiller, G. P., von Clarmann, T., Grabowski, U., Kellmann, S., Linden, A., Sinnhuber, B.-M., Krysztofiak, G., et al.: Global carbonyl sulfide (OCS) measured by MIPAS/Envisat during 2002–2012, *Atmospheric Chemistry and Physics*, 17, 2631–2652, 2017.
- 590 Hannigan, J. W., Ortega, I., Shams, S. B., Blumenstock, T., Campbell, J. E., Conway, S., Flood, V., Garcia, O., Griffith, D., Grutter, M., et al.: Global atmospheric OCS trend analysis from 22 NDACC stations, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035 764, 2022.
- Hattori, S., Kamezaki, K., and Yoshida, N.: Constraining the atmospheric OCS budget from sulfur isotopes, *Proceedings of the National Academy of Sciences*, 117, 20 447–20 452, 2020.



- 595 Houweling, S., Krol, M., Bergamaschi, P., Frankenberg, C., Dlugokencky, E., Morino, I., Notholt, J., Sherlock, V., Wunch, D., Beck, V., et al.: A multi-year methane inversion using SCIAMACHY, accounting for systematic errors using TCCON measurements, *Atmospheric Chemistry and Physics*, 14, 3991–4012, 2014.
- Hu, L., Montzka, S. A., Kaushik, A., Andrews, A. E., Sweeney, C., Miller, J., Baker, I. T., Denning, S., Campbell, E., Shiga, Y. P., et al.: COS-derived GPP relationships with temperature and light help explain high-latitude atmospheric CO₂ seasonal cycle amplification, *Proceedings of the National Academy of Sciences*, 118, 2021.
- 600 Jernigan, C. M., Fite, C. H., Vereecken, L., Berkelhammer, M. B., Rollins, A. W., Rickly, P. S., Novelli, A., Taraborrelli, D., Holmes, C. D., and Bertram, T. H.: Efficient Production of Carbonyl Sulfide in the Low-NO_x Oxidation of Dimethyl Sulfide, *Geophysical Research Letters*, 49, e2021GL096838, 2022.
- Kettle, A., Kuhn, U., von Hobe, M. v., Kesselmeier, J., and Andreae, M.: Global budget of atmospheric carbonyl sulfide: Temporal and spatial variations of the dominant sources and sinks, *Journal of Geophysical Research: Atmospheres*, 107, ACH–25, 2002.
- 605 Khan, M. A., Gillespie, S. M., Razis, B., Xiao, P., Davies-Coleman, M. T., Percival, C. J., Derwent, R. G., Dyke, J. M., Ghosh, M. V., Lee, E. P., and Shallcross, D. E.: A modelling study of the atmospheric chemistry of DMS using the global model, STOCHEM-CRI, *Atmospheric Environment*, 127, 69–79, <https://doi.org/10.1016/j.atmosenv.2015.12.028>, 2016.
- Kooijmans, L. M., Maseyk, K., Seibt, U., Sun, W., Vesala, T., Mammarella, I., Kolari, P., Aalto, J., Franchin, A., Vecchi, R., Valli, G., and Chen, H.: Canopy uptake dominates nighttime carbonyl sulfide fluxes in a boreal forest, *Atmospheric Chemistry and Physics*, 17, 11453–11465, <https://doi.org/10.5194/acp-17-11453-2017>, 2017.
- 610 Kooijmans, L. M., Cho, A., Ma, J., Kaushik, A., Haynes, K. D., Baker, I., Lujckx, I. T., Groenink, M., Peters, W., Miller, J. B., et al.: Evaluation of carbonyl sulfide biosphere exchange in the Simple Biosphere Model (SiB4), *Biogeosciences*, 18, 6547–6565, 2021.
- Kooijmans, L. M. J., Sun, W., Aalto, J., Erkkilä, K.-M. M., Maseyk, K., Seibt, U., Vesala, T., Mammarella, I., and Chen, H.: Influences of light and humidity on carbonyl sulfide-based estimates of photosynthesis, *Proceedings of the National Academy of Sciences of the United States of America*, 116, 2470–2475, <https://doi.org/10.1073/pnas.1807600116>, 2019.
- 615 Koren, G.: Constraining photosynthesis with $\Delta 17\text{O}$ in CO₂, *Nature Reviews Earth & Environment*, 2, 744–744, 2021.
- Kremser, S., Thomason, L. W., von Hobe, M., Hermann, M., Deshler, T., Timmreck, C., Toohey, M., Stenke, A., Schwarz, J. P., Weigel, R., Fueglistaler, S., Prata, F. J., Vernier, J. P., Schlager, H., Barnes, J. E., Antuña-Marrero, J. C., Fairlie, D., Palm, M., Mahieu, E., Notholt, J., Rex, M., Bingen, C., Vanhellemont, F., Bourassa, A., Plane, J. M., Klocke, D., Carn, S. A., Clarisse, L., Trickl, T., Neely, R., James, A. D., Rieger, L., Wilson, J. C., and Meland, B.: Stratospheric aerosol—Observations, processes, and impact on climate, <https://doi.org/10.1002/2015RG000511>, 2016.
- 620 Krol, M., Houweling, S., Bregman, B., van den Broek, M., Segers, A., van Velthoven, P., Peters, W., Dentener, F., and Bergamaschi, P.: The two-way nested global chemistry-transport zoom model TM5: algorithm and applications, *Atmospheric Chemistry and Physics*, 5, 417–432, <https://doi.org/10.5194/acp-5-417-2005>, 2005.
- Krol, M. C., Meirink, J. F., Bergamaschi, P., Mak, J. E., Lowe, D., Jöckel, P., Houweling, S., and Röckmann, T.: What can ¹⁴C measurements tell us about OH?, *Atmospheric Chemistry and Physics*, 8, 5033–5044, <https://doi.org/10.5194/acp-8-5033-2008>, 2008.
- Krysztófiak, G., Té, Y. V., Catoire, V., Berthet, G., Toon, G. C., Jégou, F., Jeseck, P., and Robert, C.: Carbonyl sulphide (OCS) variability with latitude in the atmosphere, *Atmosphere - Ocean*, 53, 89–101, <https://doi.org/10.1080/07055900.2013.876609>, 2015.
- 630 Kuai, L., Worden, J., Kulawik, S., Montzka, S., and Liu, J.: Characterization of Aura TES carbonyl sulfide retrievals over ocean, *Atmospheric Measurement Techniques*, 7, 163–172, 2014.



- Kuai, L., Worden, J. R., Campbell, J. E., Kulawik, S. S., Li, K.-F., Lee, M., Weidner, R. J., Montzka, S. A., Moore, F. L., Berry, J. A., et al.: Estimate of carbonyl sulfide tropical oceanic surface fluxes using Aura Tropospheric Emission Spectrometer observations, *Journal of Geophysical Research: Atmospheres*, 120, 11–012, 2015.
- 635 Lana, A., Bell, T., Simó, R., Vallina, S., Ballabrera-Poy, J., Kettle, A., Dachs, J., Bopp, L., Saltzman, E., Stefels, J., et al.: An updated climatology of surface dimethylsulfide concentrations and emission fluxes in the global ocean, *Global Biogeochemical Cycles*, 25, 2011.
- Lanczos, C.: An iteration method for the solution of the eigenvalue problem of linear differential and integral operators, *Journal of Research of the National Bureau of Standards*, 45, 255, <https://doi.org/10.6028/jres.045.026>, 1950.
- Lee, C.-L. and Brimblecombe, P.: Anthropogenic contributions to global carbonyl sulfide, carbon disulfide and organosulfides fluxes, *Earth-Science Reviews*, 160, 1–18, 2016.
- 640 Lennartz, S. T., Marandino, C. A., Von Hobe, M., Cortes, P., Quack, B., Simo, R., Booge, D., Pozzer, A., Steinhoff, T., Arevalo-Martinez, D. L., et al.: Direct oceanic emissions unlikely to account for the missing source of atmospheric carbonyl sulfide, *Atmospheric chemistry and physics*, 17, 385–402, 2017.
- Lennartz, S. T., Marandino, C. A., Von Hobe, M., Andreae, M. O., Aranami, K., Atlas, E., Berkelhammer, M., Bingemer, H., Booge, D., 645 Cutter, G., et al.: Marine carbonyl sulfide (OCS) and carbon disulfide (CS₂): a compilation of measurements in seawater and the marine boundary layer, *Earth system science data*, 12, 591–609, 2020.
- Lennartz, S. T., Gauss, M., von Hobe, M., and Marandino, C. A.: Monthly resolved modelled oceanic emissions of carbonyl sulphide and carbon disulphide for the period 2000–2019, *Earth System Science Data*, 13, 2095–2110, 2021.
- Ma, J., Kooijmans, L. M., Cho, A., Montzka, S. A., Glatthor, N., Worden, J. R., Kuai, L., Atlas, E. L., and Krol, M. C.: Inverse modelling of 650 carbonyl sulfide: implementation, evaluation and implications for the global budget, *Atmospheric Chemistry and Physics*, 21, 3507–3529, 2021.
- Maignan, F., Abadie, C., Remaud, M., Kooijmans, L. M., Kohonen, K.-M., Commane, R., Wehr, R., Campbell, J. E., Belviso, S., Montzka, S. A., et al.: Carbonyl sulfide: comparing a mechanistic representation of the vegetation uptake in a land surface model and the leaf relative uptake approach, *Biogeosciences*, 18, 2917–2955, 2021.
- 655 Meirink, J. F., Bergamaschi, P., and Krol, M. C.: Four-dimensional variational data assimilation for inverse modelling of atmospheric methane emissions: method and comparison with synthesis inversion, *Atmospheric chemistry and physics*, 8, 6341–6353, 2008.
- Monteil, G., Houweling, S., Butz, A., Guerlet, S., Schepers, D., Hasekamp, O., Frankenberg, C., Scheepmaker, R., Aben, I., and Röckmann, T.: Comparison of CH₄ inversions based on 15 months of GOSAT and SCIAMACHY observations, *Journal of Geophysical Research Atmospheres*, 118, 11 807–11 823, <https://doi.org/10.1002/2013JD019760>, 2013.
- 660 Montzka, S., Calvert, P., Hall, B., Elkins, J., Conway, T., Tans, P., and Sweeney, C.: On the global distribution, seasonality, and budget of atmospheric carbonyl sulfide (COS) and some similarities to CO₂, *Journal of Geophysical Research: Atmospheres*, 112, 2007.
- Nagori, J., Nechita-Bândă, N., Oscar Danielache, S., Shinkai, M., Röckmann, T., and Krol, M.: Modelling the atmospheric 34 S-sulfur budget in a column model under volcanically quiescent conditions, *Atmospheric Chemistry and Physics Discussions*, pp. 1–30, 2022.
- Ogée, J., Sauze, J., Kesselmeier, J., Genty, B., Van Diest, H., Launois, T., and Wingate, L.: A new mechanistic framework to predict OCS 665 fluxes from soils, *Biogeosciences*, 13, 2221–2240, <https://doi.org/10.5194/bg-13-2221-2016>, 2016.
- Remaud, M., Chevallier, F., Maignan, F., Belviso, S., Berchet, A., Parouffe, A., Abadie, C., Bacour, C., Lennartz, S., and Peylin, P.: Plant gross primary production, plant respiration and carbonyl sulfide emissions over the globe inferred by atmospheric inverse modelling, *Atmospheric Chemistry and Physics*, 22, 2525–2552, 2022.



- Spielmann, F., Wohlfahrt, G., Hammerle, A., Kitz, F., Migliavacca, M., Alberti, G., Ibrom, A., El-Madany, T. S., Gerdel, K., Moreno, G.,
670 et al.: Gross primary productivity of four European ecosystems constrained by joint CO₂ and COS flux measurements, *Geophysical
research letters*, 46, 5284–5293, 2019.
- Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B., Horowitz, L. W., Fusco, A. C.,
Brenninkmeijer, C. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Three-dimensional climatological distribution of tropospheric
OH: Update and evaluation, <https://doi.org/10.1029/1999JD901006>, 2000.
- 675 Stichel, R. E., Chin, M., Daykin, E. P., Hynes, A. J., Wine, P. H., and Wallington, T. J.: MECHANISTIC STUDIES OF THE OH-
INITIATED OXIDATION OF CS₂ IN THE PRESENCE OF O-2, *Journal of Physical Chemistry*, 97, 13 653–13 661, <https://pubs.acs.org/sharingguidelines>, 1993.
- Stimler, K., Montzka, S. A., Berry, J. A., Rudich, Y., and Yakir, D.: Relationships between carbonyl sulfide (COS) and CO₂ during leaf gas
exchange, *New Phytologist*, 186, 869–878, 2010.
- 680 Stinecipher, J. R., Cameron-Smith, P. J., Blake, N. J., Kuai, L., Lejeune, B., Mahieu, E., Simpson, I. J., and Campbell, J. E.: Biomass Burning
Unlikely to Account for Missing Source of Carbonyl Sulfide, *Geophysical Research Letters*, <https://doi.org/10.1029/2019GL085567>,
2019.
- Stinecipher, J. R., Cameron-Smith, P., Kuai, L., Glatthor, N., Höpfner, M., Baker, I., Beer, C., Bowman, K., Lee, M., Miller, S. M., et al.:
Remotely Sensed Carbonyl Sulfide Constrains Model Estimates of Amazon Primary Productivity, *Geophysical Research Letters*, 49,
685 e2021GL096 802, 2022.
- Suntharalingam, P., Kettle, A. J., Montzka, S. M., and Jacob, D. J.: Global 3-D model analysis of the seasonal cycle of atmospheric carbonyl
sulfide: Implications for terrestrial vegetation uptake, *Geophysical Research Letters*, 35, L19 801, <https://doi.org/10.1029/2008GL034332>,
2008.
- Tarantola, A.: *Inverse problem theory and methods for model parameter estimation*, SIAM, 2005.
- 690 Turco, R., Whitten, R., Toon, O., Pollack, J., and Hamill, P.: OCS, stratospheric aerosols and climate, *Nature*, 283, 283–285, 1980.
- Veres, P. R., Neuman, J. A., Bertram, T. H., Assaf, E., Wolfe, G. M., Williamson, C. J., Weinzierl, B., Tilmes, S., Thompson, C. R., Thames,
A. B., et al.: Global airborne sampling reveals a previously unobserved dimethyl sulfide oxidation mechanism in the marine atmosphere,
Proceedings of the National Academy of Sciences, 117, 4505–4510, 2020.
- Villalba, G., Whelan, M., Montzka, S. A., Cameron-Smith, P. J., Fischer, M., Zumkehr, A., Hilton, T., Stinecipher, J., Baker, I., Bambha,
695 R. P., et al.: Exploring the Potential of Using Carbonyl Sulfide to Track the Urban Biosphere Signal, *Journal of Geophysical Research:
Atmospheres*, 126, e2020JD034 106, 2021.
- Vincent, R. A. and Dudhia, A.: Fast retrievals of tropospheric carbonyl sulfide with IASI, *Atmospheric Chemistry and Physics*, 17, 2981–
3000, 2017.
- Wang, X., Jiang, X., Li, K.-F., Liang, M.-C., Kuai, L., Tan, L., and Yung, Y. L.: Variations of Carbonyl sulfide during the dry/wet seasons
700 over the Amazon, *Geophysical Research Letters*, 50, e2022GL101 717, 2023.
- Wang, Y., Deutscher, N. M., Palm, M., Warneke, T., Notholt, J., Baker, I., Berry, J., Suntharalingam, P., Jones, N., Mahieu, E., et al.: Towards
understanding the variability in biospheric CO₂ fluxes: using FTIR spectrometry and a chemical transport model to investigate the sources
and sinks of carbonyl sulfide and its link to CO₂, *Atmospheric Chemistry and Physics*, 16, 2123–2138, 2016.
- Watts, S. F.: The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and hydrogen sulfide, *Atmospheric Environment*, 34,
705 761–779, [https://doi.org/10.1016/S1352-2310\(99\)00342-8](https://doi.org/10.1016/S1352-2310(99)00342-8), 2000.



- Whelan, M. E., Min, D. H., and Rhew, R. C.: Salt marsh vegetation as a carbonyl sulfide (COS) source to the atmosphere, *Atmospheric Environment*, 73, 131–137, <https://doi.org/10.1016/j.atmosenv.2013.02.048>, 2013.
- Whelan, M. E., Hilton, T. W., Berry, J. A., Berkelhammer, M., Desai, A. R., and Campbell, J. E.: Carbonyl sulfide exchange in soils for better estimates of ecosystem carbon uptake, *Atmospheric Chemistry and Physics*, 16, 3711–3726, 2016.
- 710 Whelan, M. E., Lennartz, S. T., Gimeno, T. E., Wehr, R., Wohlfahrt, G., Wang, Y., Kooijmans, L. M., Hilton, T. W., Belviso, S., Peylin, P., Commane, R., Sun, W., Chen, H., Kuai, L., Mammarella, I., Maseyk, K., Berkelhammer, M., Li, K. F., Yakir, D., Zumkehr, A., Katayama, Y., Oge, J., Spielmann, F. M., Kitz, F., Rastogi, B., Kesselmeier, J., Marshall, J., Erkkila, K. M., Wingate, L., Meredith, L. K., He, W., Bunk, R., Launois, T., Vesala, T., Schmidt, J. A., Fichot, C. G., Seibt, U., Saleska, S., Saltzman, E. S., Montzka, S. A., Berry, J. A., and Elliott Campbell, J.: Reviews and syntheses: Carbonyl sulfide as a multi-scale tracer for carbon and water cycles, *Biogeosciences*, 15, 715 3625–3657, <https://doi.org/10.5194/bg-15-3625-2018>, 2018.
- Wingate, L., Ogé, J., Burlett, R., Bosc, A., Devaux, M., Grace, J., Loustau, D., and Gessler, A.: Photosynthetic carbon isotope discrimination and its relationship to the carbon isotope signals of stem, soil and ecosystem respiration, *New Phytologist*, 188, 576–589, 2010.
- Wofsy, S., Daube, B., Jimenez, R., and Kort, E.: HIPPO Combined Discrete Flask and GC Sample GHG, Halocarbon, and Hydrocarbon Data. Version 1.0. UCAR/NCAR - Earth Observing Laboratory, https://doi.org/https://doi.org/10.3334/CDIAC/HIPPO_012, 2017.
- 720 Wofsy, S. C.: HIAPER Pole-to-Pole Observations (HIPPO): Fine-grained, global-scale measurements of climatically important atmospheric gases and aerosols, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369, 2073–2086, <https://doi.org/10.1098/rsta.2010.0313>, 2011.
- Wohlfahrt, G., Brilli, F., Hörtnagl, L., Xu, X., Bingemer, H., Hansel, A., and Loreto, F.: Carbonyl sulfide (COS) as a tracer for canopy photosynthesis, transpiration and stomatal conductance: potential and limitations, *Plant, Cell & Environment*, 35, 657–667, 2012.
- 725 Wu, R., Wang, S., and Wang, L.: New mechanism for the atmospheric oxidation of dimethyl sulfide. The importance of intramolecular hydrogen shift in a CH₃SCH₂O radical, *The Journal of Physical Chemistry A*, 119, 112–117, 2015.
- Yan, Y., Li, R., Peng, L., Yang, C., Liu, C., Cao, J., Yang, F., Li, Y., and Wu, J.: Emission inventory of carbonyl sulfide (COS) from primary anthropogenic sources in China, *Environmental Pollution*, 247, 745–751, 2019.
- Yousefi, M., Bernath, P. F., Boone, C. D., and Toon, G. C.: Global measurements of atmospheric carbonyl sulfide (OCS), OC34S and O13CS, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 238, 106554, 2019.
- 730 Zumkehr, A., Hilton, T. W., Whelan, M., Smith, S., Kuai, L., Worden, J., and Campbell, J. E.: Global gridded anthropogenic emissions inventory of carbonyl sulfide, *Atmospheric Environment*, 183, 11–19, <https://doi.org/10.1016/j.atmosenv.2018.03.063>, 2018.




Figure 1. Measurement data locations in this study. HIPPO campaigns 1–3 are shown as colored flight tracks. NOAA airborne data are shown as red crosses. NOAA stations are marked on the map with station names.




Figure 2. Filtered MIPAS satellite data annual mean in 2009: (a) The 60 MIPAS mean averaging kernel rows, colored by their representative pressure; (b) selected MIPAS COS mean mole fractions on MIPAS levels (5, 7, 8, 10, 12); (c) MIPAS mean COS vertical profile; (d) Number of valid measurements on selected MIPAS levels after data quality control. Note that the pressure scales on the y-axis in (a) and (c) are not linear.




Figure 3. NOAA surface measurements at selected stations, and modelled COS mole fractions of inversions. The NOAA surface observations are plotted as red dots with their total error, and modelled COS mole fractions are plotted as solid lines. The first and last 6 months are shaded in blue, as spin-up and spin-down periods, respectively.




Figure 4. Inversion metric χ^2 (Eq. 13) at each NOAA surface station for all inversions except MIPAS-only_S1. The red line denotes $\chi^2=1$.




Figure 5. Comparison of the MIPAS observations with the results of inversion MIPAS+NOAA_no-bias_S1. (a) modelled COS at MIPAS level 8; (b) bias corrected MIPAS observations in the upper troposphere (MIPAS level 8); (c) mismatch (a) minus (b); (d) probability density function of the modelled and MIPAS COS mole fractions.




Figure 6. Optimized bias correction parameters for inversion MIPAS+NOAA_bias=0.3%_S1.




Figure 7. Comparison of inversions with HIPPO observations as a function of latitude and altitude. The plotted results are modelled COS minus HIPPO observations. The rows denote the five inversions labeled on the left column.




Figure 8. Comparison of inversions with NOAA airborne profiles. The NOAA airborne data were separated to two regions: Alaska and North America. The airborne profiles are averaged vertically in 500 m intervals. The grey shading is the standard deviation of the NOAA airborne measurements within each vertical interval.




Figure 9. Global COS budgets for the different inversions. The error bars represent the prior or posterior errors which are aggregated to the global scale. Note that the prior errors on the fluxes are not plotted because scenarios S1 and S0 have different prior errors. Note further that the CS₂ and DMS budget terms are counted as indirect COS fluxes.




Figure 10. The prior (APRI) and optimized (APOS) COS biosphere flux of all inversions. The flux represents the annual mean over 2009.




Figure 11. Correlations of total COS fluxes between regions (NH lands and oceans, SH lands and oceans) for the prior and the inversions. The top left panel is the prior correlation between regions. The total COS flux is the sum of anthropogenic, oceanic, biomass burning and biosphere fluxes.




Figure 12. Error reduction (in %) on the TM5 model grid-scale for inversions as shown for the total COS flux. The total COS flux is the sum of anthropogenic, oceanic, biomass burning and biosphere fluxes.



Table 1. Description of inversion settings, prior fluxes and prior bias correction.

Prior flux	Model flux abbrev.	Budget (GgS a ⁻¹)	Hor. Res.(°)	Temp. Res.	Reference
Direct oceanic COS	COS ocean	+40	2.5×2	climatology	Kettle et al. (2002); Suntharalingam et al. (2008)
Indirect oceanic CS ₂ as COS	CS ₂ ocean	+81	2.5×2	climatology	Kettle et al. (2002); Suntharalingam et al. (2008)
Indirect oceanic DMS as COS	DMS ocean	+156	2.5×2	climatology	Kettle et al. (2002); Suntharalingam et al. (2008)
Direct anthropogenic COS	COS anthr	+155	0.1×0.1	monthly	Zumkehr et al. (2018)
Indirect anthropogenic CS ₂ as COS	CS ₂ anthr	+188	0.1×0.1	monthly	Zumkehr et al. (2018)
Indirect anthropogenic DMS as COS	DMS land	+6	1×1	monthly	Zumkehr et al. (2018)
Biomass burning	COS biobr	+136	0.25×0.25	monthly	Ma et al. (2021)
<i>Sources</i>	–	+762			
Destruction by OH	–	–101	6×4	monthly	Ma et al. (2021)
Destruction by photolysis	–	–40	6×4	monthly	Ma et al. (2021)
Uptake by plants and soil	COS biosp	–1053	0.5×0.5	monthly	Kooijmans et al. (2021)
<i>Sinks</i>	–	–1194			
Net total	–	–432			
Prior bias	–	1±0.003 (unitless)	16 lat bins	monthly	This study

Table 2. Prior errors applied in the inversion scenarios in this study. The percentage is the prior error assigned to the prior fluxes. DMS remains as the prior value and is not adjusted during optimisations. Note that the absolute grid-scale flux error depends also on the flux quantity, which means that 10% error to the biosphere is larger than the same error to anthropogenic COS.

Scenario	Biosphere	Ocean COS	Ocean CS ₂	Biomass burning	Anthropogenic COS and CS ₂	DMS
S1	50%	50%	50%	10%	10%	Prior
S0	10%	50%	50%	50%	50%	Prior

Table 3. Inversions carried out in this study. The parameters are presented as MIPAS or NOAA assimilation, bias correction and bias error. The inversion names are used to denote the experiment throughout the paper.

Inversion	MIPAS assimilation	NOAA assimilation	Bias correction	Bias error	Scenario
NOAA-only_S1	No	Yes	No	No	S1
MIPAS-only_S1	Yes	No	No	No	S1
MIPAS+NOAA_no-bias_S1	Yes	Yes	No	No	S1
MIPAS+NOAA_bias=0.3%_S1	Yes	Yes	Yes	0.3%	S1
MIPAS+NOAA_bias=0.3%_S0	Yes	Yes	Yes	0.3%	S0



Table 4. Breakdown of the cost functions of the five inversions. The costs are given for the NOAA network, MIPAS satellite data and the background term, respectively.

Inversion	NOAA network	Satellite	Background	Total cost function
NOAA-only_S1	2144.3	0.0	347.1	2491.4
MIPAS-only_S1	0.0	33678.0	314.2	33992.0
MIPAS+NOAA_no-bias_S1	2640.6	35655.0	765.0	39060.0
MIPAS+NOAA_bias=0.3%_S1	2254.3	30531.0	2515.7	35301.0
MIPAS+NOAA_bias=0.3%_S0	2488.1	30755.0	3136.6	36380.0



Table 6. The same as Table. 5 but for Amazonia based on the 22 Transcom regions (Fig. 2 in Feng et al. (2011)), the Amazonia fluxes are calculated as the sum of South American Temperate and South American Tropical.

Category	Region	S1 APRI	NOAA-only_S1	MIPAS-only_S1	MIPAS+NOAA_no-bias_S1	MIPAS+NOAA_bias=0.3%_S1	S0 APRI	MIPAS+NOAA_bias=0.3%_S0
Amazonia	COS anthr	2.2±0.2	2.2±0.1 (7.2%)	2.2±0.1 (25.5%)	2.2±0.1 (22.5%)	2.2±0.1 (21.4%)	2.2±0.8	2.2±0.8 (5.7%)
	COS ocean	0.7±0.3	0.8±0.3 (12.9%)	1.2±0.2 (26.7%)	0.7±0.2 (27.3%)	1.0±0.2 (26.2%)	0.7±0.3	1.1±0.3 (11.4%)
	COS biosp	-311.0±115.0	-10.6±48.0 (58.3%)	-145.7±28.4 (75.3%)	15.5±25.9 (77.5%)	-99.5±26.3 (77.1%)	-311.0±23.0	-108.2±15.8 (31.4%)
	COS biobr	8.0±0.6	7.8±0.5 (8.2%)	8.1±0.4 (29.6%)	8.2±0.4 (26.0%)	8.0±0.4 (24.7%)	8.0±3.0	9.7±2.7 (10.2%)
	COS total	-300.1±115.0	0.1±48.0 (58.3%)	-134.2±28.4 (75.3%)	26.5±25.9 (77.5%)	-88.4±26.3 (77.2%)	-300.1±23.2	-95.1±15.9 (31.4%)
	CS ₂ anthr	3.3±0.2	3.3±0.2 (0.0%)	3.4±0.2 (0.0%)	3.4±0.2 (0.0%)	3.3±0.2 (0.0%)	3.3±1.0	4.2±1.0 (0.2%)
	CS ₂ ocean	0.7±0.2	0.8±0.2 (0.1%)	0.9±0.2 (0.1%)	0.9±0.2 (0.3%)	1.1±0.2 (0.2%)	0.7±0.2	1.6±0.2 (0.5%)
	CS ₂ total	4.0±0.3	4.1±0.3 (0.1%)	4.3±0.3 (0.1%)	4.2±0.3 (0.2%)	4.4±0.3 (0.1%)	4.0±1.0	5.8±1.0 (0.3%)