

# Advancing the estimation of future climate impacts within the United States

Corinne Hartin<sup>1</sup>, Erin E. McDuffie<sup>1</sup>, Karen Noiva<sup>3</sup>, Marcus Sarofim<sup>1</sup>, Bryan Parthum<sup>2</sup>, Jeremy Martinich<sup>1</sup>, Sarah Barr<sup>1</sup>, Jim Neumann<sup>3</sup>, Jacqueline Willwerth<sup>3</sup>, [and](#) Allen Fawcett<sup>1</sup>

<sup>1</sup>Climate Change Division, Office of Atmospheric Protection, U.S. Environmental Protection Agency, Washington, DC, USA

<sup>2</sup>National Center for Environmental Economics, Office of Policy, U.S. Environmental Protection Agency, Washington, DC, USA

<sup>3</sup>Industrial Economics incorporated 2067 Massachusetts Ave, Cambridge, MA 02140

Correspondence to: Corinne Hartin ([hartin.corinne@epa.gov](mailto:hartin.corinne@epa.gov))


## Abstract

Evidence of the physical and economic impacts of climate change is a critical input to policy development and decision making. In addition to the magnitude of potential impacts, detailed estimates of where, when, and to whom those damages may occur, the types of impacts that will be most damaging, uncertainties in these damages, and the ability of adaptation to reduce potential risks are all interconnected and important considerations. This study utilizes the reduced-complexity model, the Framework for Evaluating Damages and Impacts (FrEDI), to rapidly ~~assess-project the~~ economic and physical impacts of climate change ~~across~~ [10,000 future scenarios](#) for [multiple impact sectors](#), regions, and populations ~~in~~ [within](#) the contiguous United States (U.S.). Results from FrEDI show that net national damages increase overtime, with mean climate-driven damages estimated to reach \$2.9 trillion USD (95% CI: \$510 billion to \$12 trillion) annually by 2090. ~~Detailed FrEDI results show that of the FrEDI-analysed sectors, the majority of annual long-term (e.g., 2090) climate-driven~~ damages are [associated with climate change impacts to human health](#), largest for the health category with the majority of damages in this category ~~from the valuation estimates of including~~ mortality attributable to climate-driven changes in [extreme temperature and air quality pollution](#) (O<sub>3</sub> and PM<sub>2.5</sub>) [exposure in 2090](#). Regional results ~~from FrEDI~~ also show that [annual long-term](#) climate-driven damages vary ~~by geographical region geographically, with the~~ Southeast [experiencing is projected to experience](#) the largest annual damages per capita (mean: \$9,300 per person annually, 95% CI: \$1,800-\$37,000 per person annually), whereas the smallest damages per capita are expected in the Southwest (mean: \$6,300 per person annually, 95% CI: \$840-\$27,000 per person annually). Climate change impacts may also broaden existing societal inequalities, [with, for example, with](#) Black or African Americans disproportionately affected by additional premature mortality from changes in air quality. Lastly, we extend FrEDI projections through 2300 to estimate the net present climate-driven damages within U.S. borders from marginal changes in greenhouse gas emissions. Combined, this [work-analysis provides the most detailed and comprehensive illustration to date of the distribution of climate change impacts within U.S. borders, i.e., significantly advances our](#)

35 understanding of the impacts from climate change to the U.S., in what U.S. regions, sectors and populations are being impacted are happening, what sectors are being impacted, and which population groups being impacted the most.

### Annual U.S. Climate-Driven Damages

Mean Damages by Year and Category (Trillions \$)



40 Graphical abstract figure.

### 1 Introduction

Evidence of the physical and economic impacts of climate change is a critical input to policy development and decision making. The information on the potential magnitude of climate change damages, where, when, and to whom those damages may occur across the United States (U.S.), the types of impacts that will be most damaging, and the potential for adaptation to ability of adaptation to reduce potential risks are all important and interconnected (Martinich et al., 2018). Information Understanding this rich set of information about potential damages in different U.S. sectors and regions can help federal decision makers identify significant climate risks, which is as an initial important first step toward prioritizing and managing such risks, especially through mitigation and adaptation actions (GAO, 2017). Specifically in the U.S., rResults of recent multi-sector impact analyses show complex patterns of projected climate-driven changes across the country, with annual damages in some impact sectors (for example, labor, extreme temperature -related mortality, and coastal property) estimated to range in the hundreds of billions of U.S. dollars annually by the end of the century (Martinich and Crimmins, 2019; Hsiang et al., 2017).

Climate economics research has also continued to leverage recent advancements to develop and improve our understanding of damage functions that represent climate-driven impacts in broader economic frameworks ; leveraging considerable advancements in the science and economics of estimating future climate change impacts (NAS, 2017). Advances For example, advances in our understanding of the historical relationships between climatic

variables and the economy have ~~also~~ enabled the development of methods to assess the economic effects from ~~future~~ climate change within the U.S. ~~under different scenario~~ (GAO 2017; Field et al., 2014). ~~For As one~~ example, the Climate Change Impacts and Risk Analysis (CIRA) project, ~~coordinated by the USEPA and involving researchers from government, academiaies, and consultants~~ the private sector, has used and continues to use detailed sectoral models to ~~quantifiesy~~ the physical ~~effects~~ and economic climate-driven ~~damages of climate change in the U.S. across individual impact sectors within the U.S., using detailed models of sectoral impacts~~ (e.g., human health, infrastructure, and water resources) (EPA, 2017a). ~~In addition~~ Another example is, the Climate Impact Lab ~~- is a~~ collaboration of more than 30 climate scientists, economists, and researchers from across the U.S. and which has focused its work on understanding the economic damages from climate change both within the U.S. (Hsiang et al., 2017) and across the globe, including impacts to human health (Carleton et al., 2022), agriculture (Rising and Devineni, 2020; Hultgren et al., 2022), coastal property (Depsky et al., 2022), and energy (Rode et al., 2021) ~~and labor (Burke et al., 2015).~~

~~Typically, these resource-intensive, bottom up impact studies typically use rely on a select number of the~~ large-scale global ~~scenarios-emission and warming scenarios~~ (e.g., the Representative Concentration Pathways) ~~and are not run under different future trajectories, limiting theirre ability to explore certain aspects of uncertainty associated with a wider range of alternative future trajectories and explore certain aspects of uncertainty.~~ Temperature binning provides an alternative framework for analysing sector-specific impacts by degree of warming, which complements traditional scenario-based approaches. This framing can improve communication of results, comparability between models and studies, and flexibility for new scenario analysis (Sarofim et al., 2021). ~~The As an alternative approach, the recently developed~~ Framework for Evaluating Damages and Impacts (FrEDI) (EPA, 2021b) utilizes this temperature binning approach to ~~draws upon information from these detailed sectoral impact studies to~~ rapidly assess U.S. economic and physical impacts of climate change ~~to the United States under any future warming scenario~~ within a common framework. FrEDI was developed using a transparent process, peer-reviewed methodologies, and is designed to be a flexible framework that is continually refined to FrEDI incorporates ~~advances in peer-reviewed economic damage functions, including the incorporation of new sectors and adaptation options. In this analysis, FrEDI -draws upon over 30 climate change impact models from existing peer-reviewed studies and climate change impact models~~ to develop relationships between mean surface temperature change and climate-driven impacts across 20 sectors within U.S. borders, through the end of ~~over~~ the 21<sup>st</sup> century, ~~providing an efficient and transparent damage estimation approach that operates independently of Integrated Assessment Models (IAMs).~~ FrEDI has the flexibility to use any custom warming scenario (which can be derived from a climate model (e.g., the Finite Amplitude Impulse Response model (FaIR)e.g., Figure 1)) and couple it with any accompanying socioeconomic projections (e.g., gross domestic product (GDP) and population). Due to this level of detail and flexibility, FrEDI provides an efficient and transparent damage estimation approach useful framework to explore a variety of future baseline trajectories or, emission reduction policies and, and thereby, can

provide policy-relevant information and complement the types of analyses and outputs provided by existing integrated assessment models.


95 In this study, we ~~utilize-use 10,000~~ paired probabilistic emissions and socioeconomic projections ~~and-with a~~ simple climate model to provide inputs for FrEDI, ~~which then quantify both the~~quantifies the annual physical and economic impacts of projected climate change through the end of the 21<sup>st</sup> century across the contiguous United States (CONUS). This framework allows us to investigate the potential range of projected long-term annual climate change impacts that are associated with uncertainty in climate model parameters, a wide range of future emissions and socioeconomic conditions, as well as structural uncertainty in select damage functions. We present  
100 annual damages overtime and discuss the differential impacts projected to occur across different sectors, regions, and populations within CONUS borders to illustrate the breadth of the potential climate change risks to the U.S. These impacts incorporate a broad range of sectoral detail as well as spatial detail across seven U.S. regions. Our  
Lastly, we extend our methodology ~~extends~~ out to the year 2300 to assess the net present damage in the U.S. resulting from an additional ton of CO<sub>2</sub>, CH<sub>4</sub> or N<sub>2</sub>O emissions. Summing-Aggregating net present damages across  
105 all sectors and regions within FrEDI provides a traceable estimate of the ~~physical and~~ economic damages within U.S. borders, from a marginal change in greenhouse gas emissions.

## 2 Methods

This study-analysis consists of three components, each representing recent scientific advances in their respective fields (Figure 1). ~~First, Probabilistic~~ projections of global greenhouse gas emissions (Figure 1, Input 1) are used as  
110 ~~inputs into-to~~ a simple climate model to derive trajectories of changes in global mean surface temperature (Figure 1, Output 1). These temperature trajectories are then passed to FrEDI (Figure 1, Input 2) alongside ~~probabilistic~~ projections of U.S. Population and GDP (Figure 1, Input 1) to model annual long-term climate damages across 20 impact sectors, seven U.S.-CONUS regions, multiple adaptation scenarios, and ~~the option to explore impacts to~~ socially vulnerable populations (Figure 1, Output 2).

115 ~~A unique feature of the probabilistic emissions and socioeconomic projections, and the reduced-complexity climate model, is the rich range of uncertainty and uncertain parameters underlying them.~~In this study Specifically, we use 10,000 randomly sampled scenarios of global greenhouse gas emissions (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O), U.S. population, and U.S. GDP from the Resources for the Future – Socioeconomic Projections (RFF-SPs) (Rennert et al., 2021) (Section 2.1). ~~Probabilistic e~~mission trajectories ~~of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O~~ are ~~used as inputs~~input to the Finite  
120 Amplitude Impulse Response (FaIR) model, a simple emissions-based climate model (v1.6.2) ~~calibrated based on historical data and IPCC AR6 assessed climate variables (Smith et al., 2018)~~that relates emissions to , to calculate the changes in global mean surface temperature (relative to 1850-1900 average), calibrated based on historical data and Intergovernmental Panel on Climate Change (IPCC) AR6 assessed climate variables (Smith et al., 2018). The resulting 10,000 global mean surface temperature projections, ~~-(~~along with corresponding population and GDP

125 projections from the RFF-SPs<sup>1</sup> are then passed to FrEDI (v3.0) to calculate the physical and economic climate-driven damages to the U.S. A unique feature of using probabilistic projections with a simple climate model in this approach is the rich range of uncertainty parameters that can be assessed. We describe each process in more detail below.



130 **Figure 1:** Flow diagram of the inputs and outputs needed to evaluate the economic damages within the U.S. Emission trajectories are passed as inputs into FaIR to calculate global mean surface temperature. Global mean surface temperature, population, and GDP are then passed as inputs to FrEDI to calculate sectoral climate impacts to the U.S. Not shown is the estimation of global mean sea level rise; these values are calculated within FrEDI using a semi-empirical approach from existing literature (Kopp et al., 2016) to calculate the impacts to the subset of FrEDI sectors that are impacted by sea level rise (i.e., high-tide flooding and traffic/transportation impacts from high tide flooding, and coastal properties) (EPA, 2021b).

## 2.1 Emissions and Socioeconomics

Socioeconomic and emissions projections from 2020 to 2300 were recently developed under the Resources for the Future Social Cost of Carbon Initiative (Rennert et al., 2021). These include multi-century probabilistic projections of country-level population, GDP, and global emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O. The projections represent a state-of-the-art set of probabilistic socioeconomic and emissions scenarios based on high-quality data, robust statistical techniques, and expert elicitation. In addition, these projections incorporate coupled uncertainty in the time-dependent relationship between GDP and emissions, while also explicitly accounting for potential future climate policy and its contribution to the economy-emissions relationship (Rennert et al., 2021).

## 2.2 The Climate Model

145 The Finite Amplitude Impulse Response model (FaIRv1.6.2)<sup>1</sup> calculates atmospheric concentrations of greenhouse gases, radiative forcing, and global mean surface temperature from emissions of greenhouse gases, aerosols, and other gases (Smith et al., 2018). Version 1.6.2 was calibrated to and extensively used within the Sixth Assessment Report (AR6) of the IPCC (Forster, P. et al., 2021), resulting in 2,237 calibrated sets of climate parameters (out of

<sup>1</sup> <https://github.com/OMS-NetZero/FAIR>

the full 1 million member ensemble). ~~We Here we use the Monte Carlo simulation capabilities of MimiGIVE.jl~~  
150 ~~(<https://github.com/rffscghg/MimiGIVE.jl>) to randomly sample the~~ accessed a random sample of 10,000  
~~trajectories from the~~ RFF-SP emission scenarios (consisting of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O) ~~paired with a and the random~~  
~~sample from the~~ calibrated set of uncertain parameters contained in FaIR.<sup>2</sup> ~~The Emissions of the~~ other gases and  
aerosols (e.g., HFCs, BC, OC, etc.) not included in the RFF-SP projections were set to the associated emissions in the  
SSP2-4.5 (Meinshausen et al., 2020) scenario, ~~for~~ which most closely matches the median ~~of the~~ RFF-SP emission  
155 trajectories (Rennert et al., 2022). From the 10,000 model simulations, the average change in global mean surface  
temperature relative to 1986-2005 (FrEDI baseline) is 1.9°C (95% confidence interval: 0.8°C to 3.5°C) by 2100 and  
increases to 3.1°C (95% CI: -0.2°C to 7.8°C) by 2300 (Figure A1-1).

### 2.3 Damages from Climate Change to the U.S.

The Framework for Evaluating Damages and Impacts (FrEDI) is a reduced complexity model that assesses and  
160 quantifies future impacts to the U.S. from a changing climate. As described in detail in the Technical  
Documentation (EPA, 2021b), FrEDI uses a temperature binning approach and data from previously published  
climate impact studies (Sarofim et al., 2021) to develop relationships between climate-driven changes in CONUS  
temperature or global mean sea level rise and the resulting physical and economic damages across 20 sectors  
(Table A2-14) in seven U.S. regions. ~~While FrEDI evaluates both negative and positive impacts of climate change~~  
165 ~~across sectors and regions, climate-driven damages outweigh the positive effects for all sectors at the national~~  
~~level.~~ FrEDI also provides insight into differences in impacts under various adaptation scenarios and contains a  
module that can be used to quantify impacts to socially vulnerable populations. The underlying studies in FrEDI  
consist of bottom-up detailed sectoral analyses from the CIRA project (EPA, 2017a) and other studies including  
those from the Climate Impact Lab (e.g., Hsiang et al., 2017) and the American Thoracic Society (e.g., Cromar et al.,  
170 2022). FrEDI was designed to fill the current need of monetizing a broad range of climate-driven impacts in the U.S.  
across various emission/socioeconomic trajectories, while doing so in a significantly shorter computational  
timeframe (e.g., seconds) relative to existing impact models. To achieve this objective, the detailed spatial  
resolution of the underlying studies is reduced within FrEDI ~~to the regional level,~~ for ease, flexibility, and speed.

FrEDI currently ~~consists includes of~~ 20 ~~impact~~ sectors for which damages are modelled as functions of a climate  
175 driver (CONUS temperature or sea-level rise), U.S. GDP, and regional population. The population projections from  
the RFF-SPs are ~~spatially aggregated~~ total U.S. population. For the analysis, we disaggregate national values to  
populations for each of the seven FrEDI regions based on the percentage of regional to total U.S. population in the

---

<sup>2</sup> ~~FaIR developers provide 2,237 calibrated parameter sets from the full 1 million member ensemble. We use the~~  
~~Monte Carlo simulation capabilities of MimiGIVE.jl (<https://github.com/rffscghg/MimiGIVE.jl>) to randomly sample~~  
~~the RFF-SPs and FaIR parameter sets.~~ See Rennert et al. (2022) for more detail on the RFF-SPs and FaIR parameter  
sets. Each of the 10,000 RFF-SPs are assumed equally likely.

years 2010-2090 using projected regional populations derived from ICLUS (EPA, 2017b). The proportions for each region are held constant after 2090. Figure A1-1 shows ~~the~~that the mean and 95<sup>th</sup> confidence intervals for U.S. population and time-averaged U.S. GDP per capita growth rates are 390 million (95% CI: 260-520 million) and 1.5% (95% CI: -0.4% to 4.0%), respectively in 2100.<sup>3</sup> By 2300, the average of all 10,000 trajectories for U.S. population and time-averaged U.S. GDP per capita growth rates are 370 million (95% CI: 43 million to 1.3 billion) and 0.9% (95%CI: -0.2% to 3.4%), respectively. The trends shown in Figure A1-1 reflect the aggregate of the 10,000 individual RFF-SP trajectories (each of which has a different, equally likely growth path).

For sectoral impacts driven by temperature change, damages in FrEDI are calculated as functions of CONUS degrees of warming over time, relative to a 1986-2005 average temperature baseline. In this analysis, CONUS mean temperature change is estimated for each FaIR-derived temperature projection (calculated from each RFF-SP emissions scenario), as CONUS temperature (°C) = 1.42 × Global Temperature (°C) (EPA, 2021b). This relationship between CONUS and global temperatures is relatively stable across GCMs and over time, allowing the use of these available datapoints to develop a generalized relationship between global and CONUS temperature anomalies. For sectoral impacts driven by sea level rise (i.e., coastal properties and high tide flooding and traffic transportation impacts from high tide flooding), global mean sea level is calculated within FrEDI from global mean surface temperature using a semi-empirical method that estimates global sea level change based on a statistical synthesis of a global database of regional sea-level reconstructions from Kopp et al. (2016). In FrEDI, for a given year, sea level-driven damages are calculated by interpolating between modelled damages at different sea level heights at that same point in time; this enables FrEDI to account for interactions between adaptation costs, increased coastal property values, and sea level rise over time (EPA, 2021b).

This analysis groups mean damages from each of 20 FrEDI sectors into six topical categories and uses the default FrEDI adaptation assumptions of ‘Reactive’, ‘Reasonably Anticipated Adaptation’, or ‘No Additional Adaptation’ (see Table A3-1) for each sector. As discussed further in Section A3, Reactive or Reasonably Anticipated Adaptation is where decision makers respond to climate change impacts by repairing damaged infrastructure, but do not take actions to prevent or mitigate future climate change impacts. No additional adaptation largely incorporates historical or current levels of adaptive mitigation that were in place during the time period of each underlying sectoral study

FrEDI also has the capability to investigate adaptation options in select sectors. Available adaptation options reflect the treatment of adaptation in the underlying sectoral studies. For most of these studies, because the implicit or explicit impact response functions are calibrated to historical or current data, historically practiced adaptation or

---

<sup>3</sup> All dollar values in this paper are presented in 2020 U.S. dollars. Any necessary transformations in the inputs (e.g., RFF-SPs are in 2011\$ and FrEDI takes 2015\$) are performed using the U.S. Bureau of Economic Analysis national data on annual implicit price deflators for U.S. GDP, the top row of BEA Table 1.1.9.

210 ~~hazard avoidance actions are “baked in”, while enhanced adaptation action or new (currently unknown) technologies are not considered. Exceptions include FrEDI’s coastal property and select other infrastructure sectors (e.g., roads, rail), where the underlying study adaptation options and scenarios from the underlying studies have been incorporated into FrEDI. Total damages in these sectors are sensitive to adaptation assumptions indicating that adaptation has the capacity to both exacerbate and ameliorate future climate-driven damages, with the latter being more common. These results are further explored below and in Section A3.~~

215 In addition to quantifying differential climate-driven damages across impact sectors, geographic regions, and adaptation options, FrEDI can also compare climate-driven damages ~~associated with select sectors~~ across different populations within the U.S. This ~~capability~~ ability is largely based on a recent EPA Report on Climate Change and Social Vulnerability in the United States (EPA, 2021a), ~~which considers differential climate change risks areas a function of exposure to where climate change impacts are projected to occur, and vulnerability, in terms of an individual’s capacity to prepare for, cope with, and recover from climate change impacts. This~~ FrEDI incorporates 220 ~~this approach by analysis uses~~ using data on where populations live (US Census, 2014) as an indicator of exposure, and for vulnerability, considers four categories for which there is evidence of differential vulnerability (Table A2-25), including low income, ethnicity and race<sup>4</sup>, educational attainment, and age.

225 ~~These differential impacts in each group are calculated in FrEDI at the Census tract level as a function of current population demographic patterns (i.e., percent of each group living in each census tract), projections of CONUS population (from ILCUS, U.S. Environmental Protection Agency, 2017), and projections of where climate driven impacts are projected to occur (i.e., using FrEDI temperature impact relationships) at the Census tract level. The relative percent of each socially vulnerable group in each Census tract are from the 2014-2018 U.S. Census American Community Survey dataset (U.S. Census) and are held constant overtime because robust and long term projections for local changes in demographics are not readily available.~~

230 ~~FrEDI was initially built to project damages through 2090 for temperature scenarios with a maximum value of 10°C of warming. For this work, we extend FrEDI through the year 2300 by linearly extrapolating the temperature-binned damage functions for each sector to account for the full range of the RFF-SPs temperature scenarios (some~~

---

<sup>4</sup> This ~~report~~ analysis uses the term BIPOC to refer to individuals identifying as Black or African American; American Indian or Alaska Native; Asian; Native Hawaiian or Other Pacific Islander; and/or Hispanic or Latino. It is acknowledged that there is no ‘one size fits all’ language when it comes to talking about race and ethnicity, and that no one term is going to be embraced by every member of a population or community. The use of BIPOC is intended to reinforce the fact that not all people of color have the same experience and cultural identity. This report therefore includes, where possible, results for individual racial and ethnic groups. ~~Note the SV report reported these values as minority.~~



235 ~~of which see degrees of warming above 10°C). See Section A1 for more information on the sector-specific changes implemented in this model extension.~~

## 2.4 Estimating Net Present Value of Future Damages per ton of GHG Emissions

240 ~~While FrEDI was initially built to project damages through 2090 for temperature scenarios with a maximum value of 10°C of warming. As described previously,~~ FrEDI was extended in this work to project climate damages out to 2300 to quantify the net present damages in the U.S. resulting from an additional tonne of CO<sub>2</sub>, CH<sub>4</sub> or N<sub>2</sub>O emissions. ~~As described further in Section A4, FrEDI is extended by linearly extrapolating its sector-specific, temperature-binned damage functions to account for the full range of temperature scenarios derived from the RFF-SP emission scenarios run through FaIR (some of which have degrees of warming above 10°C). For To quantify the net present damages, this analysis, all 10,000 RFF-SP-derived temperature and socioeconomic scenarios FrEDI is are run then run through FrEDI out to 2300 under two cases:~~ a baseline (emissions = RFF-SP emissions) and a perturbed case, where 1 GtC pulse of CO<sub>2</sub> (or CH<sub>4</sub> or N<sub>2</sub>O) is added to each of the RFF-SP emissions scenarios in the year 2020. The emissions are identical between the cases for all other years. The annual marginal climate-driven damages are calculated as the difference between the damages in the baseline and perturbed cases, summed across all sectors and all regions for each year. Lastly, these marginal annual damages are discounted to the year of emissions and then aggregated across the timeseries into a single net present damage estimate. The results are normalized by the pulse size and gas chemistry (e.g., C to CO<sub>2</sub>) and reported in 2020 U.S. dollars.

245 Future monetary impacts are generally discounted relative to present value. Circular A-4 (White House, 2003) recommends a constant value of 3% for the “social rate of time preference”, which is considered to be the appropriate discount rate to use for impacts on private consumption (which would include most environmental and health impacts). The discount rate of 3% was calibrated to the real rate of return for 10-year Treasury notes from 1973 through 2003. However, OMB Circular A-4 also noted that for intergenerational impacts (a category in which climate change clearly falls), discount rates lower than 3% might be appropriate. Moreover, recent real rates of return for Treasury notes have been lower than 3%, adding support for use of a discount rate smaller than 3% (CEA, 2017). A number of economists, as well as the National Academies of Sciences (NAS, 2017) have alternatively suggested the use of Ramsey discounting (Eq. 2,  $\rho$  is the rate of pure time preference,  $g$  is a time-varying measure of per capita consumption or income, and  $\eta$  is the elasticity of the marginal value of consumption with changes in  $g$ ) as an appropriate approach to discounting long-term problems such as climate change. The effect of Ramsey discounting is to value damages more highly in futures with less economic growth – e.g., future societies that have fewer resources available for adaptation, and vice versa. A recent study from Rennert et al. (2022) used a Ramsey

265 approach calibrated to a near-term target discount rate of 2%, with  $\rho = 0.2\%$  and  $\eta = 1.24$ .<sup>5</sup> Here we use this Ramsey discounting approach to calculate the net present value.

The net present value (NPV) for a constant discount rate ( $r$ ) is calculated such that

$$NPV(D(t)) = \sum_{t=2020}^{t=2300} \frac{D(t)}{(1+r)^t} \quad (1)$$

270 The net present value for a Ramsey discounting approach is calculated using a time-varying and state-specific discount rate<sup>6</sup> which is a function of per capita economic growth ( $g_t$ ):

$$r_t = \rho + \eta * g_t \quad (2)$$

and where this time varying rate is then used in the net present value calculation such that

$$NPV(D(t), g(t)) = \sum_{t=2020}^{t=2300} \frac{D(t)}{\prod_{x=2020}^{x=t} (1+r_x)^*} \quad (3)$$

275 ~~In this expression,  $g_t$  has also been adjusted to reflect climate damages, such that in any given year-time period  $g_t$  it is the per capita consumption as calculated by taking the exogenous RFF-SP GDP, subtracting the damages output by FrEDI, and dividing by total population, and recovering the GDP per capita growth rate net of climate damages.~~ Because most of the sectoral damages are proportional to GDP per capita (given that the default elasticity of VSL to GDP per capita is 1, all sectors with a mortality endpoint also qualify), ~~in extreme cases where damages become a substantial percentage of GDP~~ a correction can be made to account for this relationship (Nordhaus, 2017) . For this analysis, we use the equation

280

$$D(t, g(t)) = \frac{D_0(t)}{1 + \frac{D_0(t)}{GDP_0(t)}} \quad (4)$$

Where  $GDP_0(t)$  is the exogenous RFF-SP GDP, ~~and~~  $D_0(t)$  is the initial total damages output by FrEDI, ~~and~~  $D(t, g(t))$  are the resulting damages.

### 3 Results and Discussion

#### 285 3.1 Annual U.S. Climate-Driven Damages by the End of 21st Century

FrEDI was developed to quantify the physical and economic damages ~~in the U.S.~~ from climate change over the 21<sup>st</sup> century, within contiguous U.S. borders. Figure 2 shows the net annual ~~mean~~ economic climate-driven damages across 20 sectors in the U.S. in the years 2050, 2070, and 2090, as calculated by the mean from the 10,000 baseline

<sup>5</sup> For Ramsey discounting calibrated to near-term target discount rates of 1.5%, 2.5%, or 3%,  $\rho = 0.01\%$ , 0.5%, and 0.8% and  $\eta = 1.02$ , 1.42, and 1.57 respectively.


<sup>6</sup> Consistent with Rennert et al. [2022], we use a stochastic Ramsey discount factor to discount future climate-driven damages.

RFF-SP scenarios (i.e., emission, population, and GDP trajectories). [This analysis groups mean damages from each of 20 FrEDI sectors into six topical categories and uses the default FrEDI adaptation assumptions of ‘reactive’](#)

290

## Annual U.S. Climate-Driven Damages

Mean Damages by Year and Category (Trillions \$)



‘reasonably anticipated adaptation’<sup>7</sup> or ‘no additional adaptation’<sup>3, 1</sup> for each sector, and groups resulting mean damages from each of 20 FrEDI sectors into six topical categories. As discussed further in Section A3, rNo adaptation largely incorporates historical or current levels of adaptive mitigation that were in place during the time period of each underlying sectoral study. Proactive adaptation is where decision makers take adaptive action with the goal of preventing infrastructure repair costs associated with future climate change impacts. Section A3 in the Appendix describes the impacts of modeling different adaptation options in more detail. FrEDI does consider both the positive and negative effects of climate change across sectors, however, the negative effects of climate change outweigh the positive for all sectors at the national scale. Total annual damages throughout this analysis are shown in 2020 U.S. dollars, converted from FrEDI’s base units of \$2015 USD using Annual GDP Implicit Price Deflators [U.S. Bureau of Economic Analysis, 2023]. Figure 2 shows that net national damages increase overtime, with mean climate-driven damages estimated to reach \$2.9 trillion USD (95% CI: \$510 billion to \$12 trillion), or ~3% of U.S. GDP, annually by 2090 for a subset of total climate impacts. Given that the drop in GDP in 2009 during the Great Recession was 2.2%<sup>7</sup>, an annual decrease in GDP of over 3.0% per year by the end of the century (Figure 3) reflects substantial damage to the national economy (though it is relevant to recognize that much of the damages estimated in FrEDI are a result of mortality, which is not directly reflected in historical GDP estimates). Table 1 provides the 2090 annual mean damages and 95% confidence interval (CI) for each aggregate category. The individual sectors that contribute to each category are listed in Table A2-1.


295

300

305

<sup>7</sup> Data from <https://fred.stlouisfed.org/series/FYGDGP>, percentage decline in annual GDP from 2008 to 2009.

310 **Figure 2:** Annual mean U.S. climate-driven damages in 2050, 2070, and 2090. Damages are average values in billions of dollar (2020 USD) calculated from the 10,000 RFF-SPs. Sectors are grouped into six categories for visual purposes. The number of sectors included in each category is given in parenthesis in the legend. See Table A2-1 for the list of sectors in each category. [Note that this is only a subset of potential climate impacts to the U.S.](#)



315 **Figure 35:** [Share of U.S. GDP \(from the RFF-SPs\) of climate-driven damages over time for those impacts represented in FrEDI.](#) Mean (solid) and median (dashed) lines along with 5<sup>th</sup>-95<sup>th</sup> (dark shaded) and 1<sup>st</sup>-99<sup>th</sup> (light shaded) percentile bounds.

**Table 1: The 95% confidence interval (CI) and mean annual U.S. climate-driven damages in 2090 for the six categories shown in Figure 2. All values are in 2020 USD. Totals may not sum due to rounding.**

Category	Mean (billions)	95% CI (billions)
Health	\$2,600	\$350-\$11,000
Infrastructure	\$220	\$140-\$360
Labor	\$51	\$6.7-\$220
Electricity	\$22	\$9.3-\$35
Agriculture	\$6.1	\$0.42-\$19
Ecosystems + Recreation	\$4.0	\$1.6 - \$7.5
Total <a href="#">in FrEDI</a>	<a href="#">\$21,9700</a>	\$510 – \$12,000

320 Climate-driven damages ~~as estimated by~~ [from](#) FrEDI are largest for the health category. The majority of damages in this category are from the ~~estimated~~ valuation ~~estimates~~ of premature mortality attributable to climate-driven changes in ~~extreme~~ temperature and air quality (O<sub>3</sub> and PM<sub>2.5</sub>), but also include monetized health damages

attributable to Vvalley fever, southwest dust, wildfire smoke exposure and suppression costs, and crime incidents.


325 Another FrEDI category that includes the monetized value of directly estimated physical impacts (rather than a direct modelled relationship between temperature and monetized damages) is labor, which is the third largest category in 2090 and represents the damages resulting from lost hours of work when temperatures are too hot for workers to work outdoors or in unconditioned workplaces (e.g., warehouses). Table 2 provides the mean physical impacts from each of the sectors in the health and labor categories in 2090, along with the 95% CI. As shown in  
 330 Table 2, climate-driven changes in extreme temperature have the largest impact on premature mortality, resulting in nearly 50,000 additional deaths (95% CI: 19,00-91,000 deaths) annually by 2090, followed by climate-driven changes in air quality (5,100 deaths; 95% CI: 2,100-10,000 deaths) and exposure to wildfire smoke (1,100 deaths; 95% CI: 460-1,700 deaths).

335 **Table 2: The range of 2090 physical impact results across the 10,000 RFF-SP projections, including the 95% CI and mean. Totals may not add-sum due to rounding.**

Sector	Impact	95% CI	Mean
<u>Extreme</u> <u>Temperature</u> <u>Temperature</u> <u>Related Mortality</u>	Premature Mortality (deaths)	19,000 – 91,000	50,000
Air Quality		2,100 – 10,000	5,100
Wildfire		460 – 1,700	1,100
Southwest Dust		160 – 690	390
Valley Fever		130 – 480	300
Crime	Incidence (number of property and violent crimes)	-160 – 1,100	4,700
Labor	Work Hours lost (millions of hours)	170 – 830	430

To further illustrate the distribution of monetized damages contributions of across each individual sectors, Figure 4 shows the range of 2090 annual climate-driven damages in each of the 20 sectors in FrEDI, across all 10,000 RFF-SP emission, GDP, and population scenarios, in decreasing order of sectoral mean damages. Figure 4 shows that  
 340 national total damages in 2090 are primarily driven by the valuation of premature mortality attributable to climate-driven changes in extreme temperature (mean: \$2.3 trillion per year; 95% CI: \$0.31 – \$9.9 trillion per year). The next four sectors with the largest monetary climate-driven damages include premature mortality attributable to changes in air quality (mean: \$240 billion per year 95% CI: \$32-\$1000 billion per year), transportation impacts  
traffic delays associated with changes in high tide flooding (mean: \$140 billion per year; 95% CI: \$110-\$200 billion  
 345 per year), national labor hours lost (mean: \$51 billion per year; 95% CI: \$6.7-\$220 billion per year), and health


damages from wildfire smoke exposure and response costs from wildfire suppression (mean: \$51 billion per year, 95% CI: \$8.1-\$220 billion per year). Climate-driven damages to coastal properties associated with changes in tropical storm frequency and wind strength (mean: \$29 billion per year; 95% CI: \$12-\$49 billion per year), damages attributable to changes in rail (mean: \$19 billion per year; 95% CI: \$7.7-\$45 billion per year) and road systems (mean: \$17 billion per year; 95% CI: \$6.6-\$35 billion per year), health damages from changes in southwestern dust exposure (mean: \$18 billion per year ; 95% CI: \$2.5-\$77 billion per year), and the health burden of change in Valley fever incidence (mean: \$14 billion per year; 95% CI: \$2.0-\$58 billion per year), round out the top 10 sectors with the largest annual damages in 2090. Figure A2-12 provides the mean and 95% confidence interval total damages for each sector over the entire 2020-2100 timeseries. The large distribution of damages in each individual sector is driven by large range of RFF-SP emissions, population, and GDP projections and the dependence of the valuation approach for each sector on these parameters (as described in EPA, 2021b).



**Figure 43:** Annual U.S. damages in the year 2090 by sector, in order of decreasing mean damages, colored by six sector category groupings. Note the change in x-axis in each panel. Box and whiskers show the 2.5<sup>th</sup>, 25<sup>th</sup>, 50<sup>th</sup>, 75<sup>th</sup>, and 97.5<sup>th</sup> percentiles, and mean damages (diamonds) across all 10,000 projections. Damages are in billions of 2020USD.

Climate change also may broaden existing societal inequalities (EPA, 2021a). Both the effects of climate change and the distribution of populations within the U.S. vary geographically and therefore it is also possible to project how climate changes may impact overburdened and underserved populations. In addition to estimating national and regional sectoral impacts, FrEDI also contains a module to generate, and report results of disproportionate exposure and distributional physical effects across four categories of potentially socially vulnerable populations for a subset of six sectors with larger economic damages. For example, results from this module suggest Black or African Americans are disproportionately affected by additional premature mortality from changes in air quality, while Hispanic or Latino Americans are more affected by lost labor hours (Figure A3) under a changing climate.

Results from FrEDI also show that climate-driven damages across the national population vary by geographical region. Figure 54 shows a map of the damages per capita in each CONUS region in the year 2090, with pie charts showing the per capita damages in each region and the share of the four sectors with the largest damages (same figure for absolute regional damages in Figure A2-24). Based on the climate impacts included in FrEDI, Figure 54 shows that the Southeast will experience the largest annual damages per capita (mean: \$9,300 per person annually, 95% CI: \$1,800-\$37,000 per person annually), whereas the smallest damages per capita are expected in the Southwest region (mean: \$6,300 per person annually, 95% CI: \$840-\$27,000 per person annually). In each region, the largest monetary damages in 2090 are expected from premature mortality associated with changes in extreme temperature, ranging from \$4,500 per person in the Southwest to \$6,500 per person in the Southeast. Damages from ~~traffic delays due transportation impacts to from~~ high tide flooding and premature mortality attributable to climate-driven change in air quality are the second and third largest in the coastal Southeast and Northeast regions. In the Northwest and Southwest, the sectors with the second and third largest climate-driven monetized damages are air quality and wildfires. In the Southern Plains, high tide flooding ~~traffic delays transportation impacts~~ and labor hours lost are the second and third largest sectors, while rail and wildfires are the second and third largest in the Northern Plains, and labor and rail are the second and third largest in the Midwest. There are some regions and sectors projected to benefit from warming temperatures, including an expected reduction in air pollution attributable mortality in the Midwest under warmer conditions. Overall, however, the negative impacts of climate change outweigh the positives such that net losses are projected in each region.



390 **Figure 54:** Mean per capita annual climate-driven damages across the seven regions in 2090 for the subset of climate impacts included in FrEDI. Donut charts show the annual per capita damages (2020\$ per person annually) and the top four sectors with the largest damages in each region. All damages from remaining (non-top four) sectors are shown by the light gray wedges.

395 **3.2 Differential impacts**


Lastly, climate change ~~As illustrated in Error! Reference source not found. Figure 4, impacts resulting from climate change will not be distributed evenly across U.S. geographic regions or sectors, may also broaden existing societal inequalities (EPA, 2021a), and -~~ understanding the comparative risks to different populations ~~groups living in these areas is also critical for developing effective and equitable strategies for responding to climate change. As~~ described in Section 2,

400 ~~Lastly, climate change also may broaden existing societal inequalities (EPA, 2021a). Both the effects of climate change and the distribution of populations within the U.S. vary geographically and therefore it is also possible to project how climate changes may impact overburdened and underserved populations. In addition to estimating national and regional sectoral impacts, FrEDI also contains a module to generate, and report results of~~


405 ~~disproportionate exposure and distributional physical effects across four categories~~ groups of potentially socially vulnerable populations ~~for a subset of six sectors with larger economic damages.~~ For example, results from this module show that ~~Black or African Americans are more likely to be affected by additional premature mortality from climate-driven changes in air quality, while Hispanic or Latino Americans are more likely to experience lost labor hours affected than other racial and ethnic groups by lost labor hours (Figure 6A3) under a changing climate.~~



410 ~~As illustrated in Figure 4, impacts resulting from climate change will not be distributed evenly across U.S. geographic regions or sectors. Understanding the comparative risks to different population groups living in these areas is also critical for developing effective and equitable strategies for responding to climate change.~~



415 **Figure 6: Vulnerability to climate-driven changes in air quality attributable mortality and labor hours lost, by race and vulnerable groups in 2090. Top panels) Difference in risk in 2090 for four vulnerable populations. Bottom panels) Additional rates of impacts in 2090, by race and ethnicity.**




420 **Figure 4: Mean per capita annual climate driven damages across the seven regions in 2090. Donut charts show the annual per capita damages (2020\$ per person annually) and the top four sectors with the largest damages in each region. All damages from remaining (non-top four) sectors are shown by the light gray wedges.**

425 Confidence intervals presented throughout this analysis account for uncertainty associated with the range of future emission and socioeconomic projections across the 10,000 RFF-SP scenarios. These also incorporate climate parameter uncertainty as a Monte Carlo approach was used to sample the calibrated parameter set when running FaIR with the 10,000 RFF-SP emissions scenarios. In addition to these uncertainties and sensitivities to adaptation options, damage estimates within FrEDI are also sensitive to uncertainties in the underlying damage functions themselves. Similar to adaptation, FrEDI can incorporate this source of structural uncertainty when uncertainty estimates are available in the underlying study or when multiple damages functions are available for a single

430 sector. For example, FrEDI incorporates three studies of climate-driven temperature-related mortality, two of which include underlying uncertainty estimates. As shown in Table A3-2, there is a large range of damage estimates from temperature-related mortality across each study, however, these values all fall within the uncertainty range derived from the RFF-SP scenarios, presented in the main text.

435 Figure 5 shows annual total national damages as a percentage of GDP out to 2100. The mean total damage share in 2100 is 3.4% and a maximum up to 13.0%. Given that the drop in GDP in 2009 during the Great Recession was 2.2%<sup>8</sup>, an annual decrease in GDP of over 3.0% per year reflects substantial damage to the national economy (though it is relevant to recognize that much of the damages estimated in FrEDI are a result of mortality, which is not directly reflected in historical GDP estimates).

<sup>8</sup>Data from <https://fred.stlouisfed.org/series/FYGDP>, percentage decline in annual GDP from 2008 to 2009.



440 ~~Figure 5: Share of U.S. GDP of climate-driven damages over time. Mean (solid) and median (dashed) lines along with 5<sup>th</sup>-95<sup>th</sup> (dark shaded) and 1<sup>st</sup>-99<sup>th</sup> (light shaded) percentile bounds.~~

### 3.2 Comparison with SSPs

To place mean damages in context of alternative future storylines, Table 3 shows a comparison of annual national climate-driven damages in the U.S. in the year 2090 from a subset of four Shared Socioeconomic Pathways (SSPs), which represent projected socioeconomic global changes up to 2100 (O'Neill et al., 2017). Annual damages in Table 3 are calculated following the same approach as outlined in Figure 1, but using SSP trajectories of emissions, U.S. GDP, and U.S. population from the SSP Public Database (v2.0)<sup>9</sup>. These trajectories do not include uncertainty related to climate and so we present only one value for each trajectory. Table 3 shows that annual U.S. climate-driven damages in 2090 from all but the SSP5-8.5 scenario fall below mean U.S. annual damages as predicted by the RFF-SP scenarios (\$3.1 trillion). However, annual damages from all SSP scenarios fall within the 95% confidence interval (\$0.5-\$12.3 trillion-\$).

**Table 3: Comparison of FrEDI damages from SSP and RFF socioeconomic input scenarios in 2090 (billions \$2020 USD)**

Scenario	Annual U.S. Damages (billion \$2020USD)	Temperature Change in 2090 relative to FrEDI baseline (1986-2005 average)
SSP1-1.9	700	0.64
SSP2-4.5	1700	1.8
SSP3-7.0	1600	2.7
SSP5-8.5	7000	3.4
This study mean (95% CI)	2900 (510-12,000)	1.8 (0.80-3.2)

<sup>9</sup> <https://tntcat.iiasa.ac.at/SspDb/dsd?Action=htmlpage&page=80>

### 3.3 Net Present Damages per ton of GHG emissions

455 We extend FrEDI to project climate damages through to 2300 (Section A41 and Figure A5) to quantify the net present damages within the CONUS-US resulting from an additional tonne of CO<sub>2</sub>, CH<sub>4</sub>, or N<sub>2</sub>O emissions. As described in Section 2.4, the net present value is the discounted sum of a stream of future damages produced by an emissions pulse in 2020 over the entire 2020-2300 time period. We explore the sensitivity of the remaining estimates to discounting assumptions by using Ramsey discounting calibrated to near-term target rates of 1.5%, 2.0%, 2.5%. Figure 76 shows the average, median, and range of estimated values for each discounting approach.<sup>10</sup>




Figure 76. Net present value of future damages from one tonne of CO<sub>2</sub> for damages occurring only within the CONUS. Units are in dollars (2020 USD) per ton of CO<sub>2</sub> emitted. Whiskers represent the 2.5<sup>th</sup> and 97.5<sup>th</sup> percentiles, while boxes span the 25<sup>th</sup> to 75<sup>th</sup>. Mean values (stars and text) along with median values (vertical lines) are also shown.

465 ~~This methodology can also be extended to explore the net present value of future damages resulting from an additional tonne of CH<sub>4</sub> (500\$/ton of CH<sub>4</sub> under a 2% Ramsey discounting), N<sub>2</sub>O (9,700\$/ton of N<sub>2</sub>O under a 2% Ramsey discounting), or other greenhouse gas emissions.~~ These results show that even considering only the direct CONUS impacts as estimated by FrEDI, damages per tonne of CO<sub>2</sub> are almost 20% of a recently estimated global value (\$185 per tonne of CO<sub>2</sub> under a 2% Ramsey discounting, (Rennert et al. 2022)). This methodology can also be

470 extended to explore the net present value of future damages resulting from an additional tonne of CH<sub>4</sub> (500\$/ton of CH<sub>4</sub> under a 2% Ramsey discounting), N<sub>2</sub>O (9,700\$/ton of N<sub>2</sub>O under a 2% Ramsey discounting), or other greenhouse gas emissions.

<sup>10</sup> Figure A4-26 additionally compares these results to those using a constant discount rate of 3%, for a comparison with the historical approach in Circular A-4 (White House, 2003).

While the damages estimated within FrEDI are constrained to the 48 contiguous United States ~~that are included within the model~~, it is important to note that the appropriate climate damages to consider when evaluating policy-induced changes in a global pollutant such as greenhouse gases would be damages that account for impacts around the globe. For example, The National Academies of Sciences advised that “[i]t is important to consider what constitutes a domestic impact in the case of a global pollutant that could have international implications that affect the United States” (NAS, 2017). Impacts that occur outside of U.S. borders (and outside of FrEDI) will impact the welfare of U.S. residents and firms because of the interconnectedness of the global economy, international markets, trade, tourism, national security, political destabilization, additional spillover effects, and many other activities not yet captured in ~~FrEDI~~the model. Moreover, the act of international reciprocity has been highlighted as motivation for including damages occurring outside of U.S. borders in a social cost estimate of global pollutants (Carleton and Greenstone, 2022; Revesz et al., 2017; and references within). It has also been shown that accounting for global damages in domestic policymaking can be individually rational (Kotchen, 2018). Therefore, we emphasize the contribution of the damages estimated within FrEDI as providing a useful understanding of the channels through which climate change can affect U.S. citizens and residents and their relative magnitudes beyond what is currently possible in many global models yet remain a partial estimate of the total damages from greenhouse gas emissions.

#### 4 Conclusions

This study presents an evolving framework to quantify the damages of climate change to the U.S. economy, relying on more than a decade of research exploring individual sectoral impacts within the contiguous U.S. (EPA, 2021b). Impacts are dependent upon a change in global mean surface temperature, U.S. GDP and U.S. population, and assumptions about adaptation. Adaptation is relevant in many sectors when quantifying benefits (Section A3), however, there are some sectors within FrEDI that do not have explicit options to model adaptation. For example, the largest sector, premature mortality from ~~extreme~~temperature ~~changes~~, dominates the monetized damages across all regions. The mortality approach used in this paper is based on a well-regarded systematic review and meta-analysis of temperature-related mortality studies (Cromar et al., 2022). However, there is substantial uncertainty based both on difficulty of relating historical mortality to ~~extreme~~temperature ~~changess~~, but also the potential for future adaptive responses to reduce vulnerability to temperatures (Carleton et al., 2022; Lay et al., 2021).

While this work advances our understanding of climate-related impacts to the U.S., it is far from a comprehensive accounting of sectoral damages within the U.S. The FrEDI framework is dynamic, with new sectors being added to the framework on a continuous basis (including in the near term several types of health impacts including mental health, vibriosis, and health impacts of extreme storms), as well as broader coverage of direct and indirect impacts of inland flooding, ~~but reflecting the overall status of underlying literature.~~ However, the framework still omits

coverage of many nonmarket sectors such as biodiversity, ocean acidification, many other ecosystem service losses, climate-forced migration, conflict, etc. We anticipate that the inclusion of more sectors will increase the estimates of net present damages due to GHG emissions. This work also omits the impacts of tipping elements due to climate change, which may lead to abrupt and irreversible impacts (Armstrong McKay et al., 2022). This study does not explore tipping elements like permafrost thaw or Antarctic ice sheet instability. Future work may entail coupling BRICK to the framework to better explore the uncertainty within sea level rise (Wong et al., 2022, 2017). While CO<sub>2</sub> fertilization effects are included in the damage estimates for the agriculture sector, the work does not account for any other direct effects of GHGs, such as the health, agriculture, or ecosystem damages resulting from ozone produced by methane's reaction in the atmosphere. Lastly, this work does not account for interactions among sectors, interactions between non-U.S. and U.S. damages through global markets, and their feedback on the U.S. economy. While we focus on U.S. damages, we acknowledge that impacts resulting from GHG emissions, regardless of where they originate, are global in nature. The bulk of the economic damages from climate change will be outside of the U.S. and the U.S. may also experience indirect effects through trade, business, migration, etc. (NAS, 2017; Hsiang et al., 2017).

[Regardless of these limitations](#), this work significantly advances our understanding of the impacts from climate change to the U.S., in what U.S. regions impacts are happening, what sectors are being impacted, and which population groups being impacted the most. These results imply that there can be significant benefits to the U.S. from greenhouse gas mitigation, and significant benefits to the people of the U.S. [FrEDI can also quantify the benefits of mitigation policies by comparing two scenarios similar to the results presented in section 3.3](#). Due to FrEDI's flexible framework, it allows for the model to be continually updated as studies of impacts to new sectors, or updates to outdated sectoral studies become available. Since this work incorporates multiple disciplines, emission projections, climate modeling, impact modeling, and economic communities, it has the potential to be a useful tool in bridging the research gap between these communities and helping to address some of the omitted climate change risks currently within this field (Rising et al., 2022).

530

## Appendix

### Section A1: Detailed Inputs to FrEDI ~~Extrapolation method(s) past 2090~~

FrEDI was calibrated to estimate impacts for detailed 21<sup>st</sup>-century scenarios and trajectories, as described in Sarofim et al. (2021). Extending the FrEDI approach to 2300 requires two adjustments to adapt the sensitivity of the model to climate drivers and to socioeconomic conditions beyond the 21<sup>st</sup>-century. First, we consider how the sensitivity to climate drivers (temperatures and SLR inputs) might differ from 21<sup>st</sup>-century conditions. FrEDI damages were originally calibrated for temperatures from 0 to 6 degrees, relative to the 1986 to 2005 era, and SLR for 21<sup>st</sup>-century trajectories that result in 30 to 250 cm GMSL outcomes by 2100. The original framework only returns physical and economic damage estimates within those bounds. In the modified FrEDI framework used within this study, damage estimates for temperature or SLR below those bounds are set to zero, because FrEDI does not consider benefits from reductions in temperatures or sea level relative to the calibration baseline (although FrEDI does consider benefits of warming—for example in the current demonstration of the framework both the Air Quality and Marine Fisheries sectors return benefits of climate change at regional scales across the temperature trajectory examined). Damage estimates for temperature inputs above these bounds are calculated by extrapolating damages per degree using the change in damages between 5 and 6 degrees. SLR inputs above the bounds are extrapolated based on the damages per centimeter of SLR modelled by the two highest sea level scenarios in 2090.

Second, we consider how the sensitivity to socioeconomic drivers continues beyond 2090 through 2300 on a sector-specific basis (the supplement provides a full tabular summary of socioeconomic extensions for all FrEDI sectors). In general, health-based sectors (e.g., extreme temperature\_mortality) are driven by population which enters linearly; infrastructure sectors (e.g., rail) are driven by population for passenger use and GDP for freight use; and coastal infrastructure is adjusted by expected real property price appreciation, using GDP per capita and income elasticity of 0.45, consistent with the underlying Neumann et al., (2021). For many sectors, economic valuation is a separable process from the overall damage calculation in FrEDI—for example willingness to pay to reduce fatality risk (referred to as the value of statistical life or VSL) is adjusted based on the projection of GDP per capita and a default income elasticity of 1.0. Some sectors—which, in general, make up a small portion of overall damages—are not adjusted for socioeconomic projections but vary based only on sensitivity to projected temperature or sea level.

An additional consideration is FrEDI's application of custom scalars to account for sector-specific physical impact adjustments not directly tied to GDP or population, but that vary through the 21<sup>st</sup>-century and/or over space. For example, some morbidity outcomes are based on baseline incidence rates, while winter recreation activities depend on activity demand projections. For these scalars, we adopt region-specific 2090 values, which we consistently apply throughout the post-2090 period. This approach—i.e., using the 2090 conditions throughout the

565

post-2090 simulation—is also applied for sectors for which FrEDI is calibrated based solely on 2010 and 2090 socioeconomic boundary conditions (rather than as a direct function of varying GDP and population inputs).

570

Up to 6 degrees, FrEDI uses a piecewise linear function to estimate damages. This approach captures non-linearities from the underlying impact models. However, The assumption that for damages outside temperatures above of the calibration regime, FrEDI assumes increase linearly at the linear rate of change in damages equal to the rate of the increase over the change in damages from 5 degrees to 6 degrees is likely to be a conservative one. This assumptions is likely to be conservative: Hsiang et al. (2017) found that combined damages in the United States increased quadratically with temperature, and Weitzman (Weitzman, 2012) suggested that while a quadratic damage form might be reasonable for temperature changes up to 2.5 degrees C globally, for higher temperatures it would make sense for damages to increase more quickly, as standard damage functions are unlikely to capture the sheer magnitude of impacts resulting from the kind of dramatic changes the planet would undergo at temperature changes substantially higher than that.


575

**Table A1. Summary of Strategy for Extending FrEDI Sectoral Results from 2090 to 2300 Modeling Horizon. Impact column provides detail for subcategories of impacts estimated within the Framework. Wildfire sector subcategories include morbidity and mortality associated with air quality impacts and fire suppression response costs — these two classes of subcategories are listed separately because they employ different extension strategies.**

Extension Strategy	Sector	Impact	Notes
Impacts scale by Input Population and/or GDP	Air Quality	Ozone	- - - - No change to method, continue to scale with input socioeconomic trajectories - - - - Multiple Impact Years, use 2090 for all years past 2090
		PM <sub>2.5</sub>	
	Extreme Temperature Related Mortality (Cromar et al., 2022)	N/A	
	Labor	N/A	
	Valley Fever	Mortality	
		Morbidity	
		Lost Wages	
	Water Quality	N/A	
	Wildfire	Morbidity	
		Mortality	
	Winter Recreation	Alpine Skiing	
		Cross-Country Skiing	
		Snowmobiling	
	Southwest Dust	Acute Myocardial Infarction	
All Cardiovascular			
All Mortality			
All Respiratory			
Asthma-ER			
Develop New multipliers	Electricity Supply and Demand	N/A	Extend existing scalars linearly past 2090



	Electricity Transmission and Distribution	N/A	New multipliers scale results with GDP or GDP per capita	
	Roads	N/A		
	Coastal Properties	N/A		
	High-Tide Flooding and <u>Traffic Transportation Impacts from High Tide Flooding</u>	N/A		
	Rail	N/A		
No Time-Dependent Multipliers	Asphalt Roads	N/A	-	
	Inland Flooding	N/A	-	
	Urban Drainage	N/A	-	
	Wildfire	Response Costs	-	
	Wind Damage	N/A	-	
	Marine Fisheries	N/A	-	
	Agriculture	Cotton		-
		Maize		-
		Soybean		-
		Wheat		-
	Crime	Property		-
Violent			-	



580 **Figure A1-1.** Timeseries of global mean temperature (°C) relative to 1986-2005 baseline, U.S. population (millions), and average  
 585 U.S. GDP percapita growth rate (2020\$) for the 10,000 RFF-SP scenarios from 2020-2300. Temperature trajectories are derived  
 from FalR model runs of the 10,000 RFF-SP emission scenarios. Individual scenarios are shown by light gray lines. Medium and  
 dark gray shaded regions represent the 99<sup>th</sup> and 95<sup>th</sup> percent confidence intervals, respectively. The red line is the mean value  
 overtime.

**Section A2: Differential Impacts Across Populations**

FrEDI incorporates information from the recent EPA Report of Climate Change and Social Vulnerability in the United States: A Focus on Six Impacts (EPA 2021) (hereafter called the SV Report) to assess the differential climate-driven impacts in 2090 across different socially vulnerable groups. As described in the SV Report, this analysis

590 *Table A2-1. Four socially vulnerable groups considered in this analysis and the reference groups (adapted from U.S. Environmental Protection Agency (2021b))*

Categories	Group Name	Description	Reference Group
Income	Low income	Individuals living in households with income that is 200% of the poverty level or lower	Individuals living in households with income greater than 200% of the poverty level.
Age	65 and Older	Ages 65 and older	Under age 65
Race and ethnicity	BIPOC	Individuals identifying as one or more of the following: Black or African American, American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and/or Hispanic or Latino	Individuals identifying as White and/or non-Hispanic
Education	No High School Diploma	Individuals aged 25 and older with less than a high school diploma or equivalent	Individuals aged 25 or older with educational attainment of a high school diploma (or equivalent) or higher.

## Section A2: Detailed results to 2090

**Table A2-1. National Annual Damage Statistics (mean and 95% confidence interval) for the year 2090, in billions of 2020 USD, listed alphabetically by Sector**

Sector	Category	Default Adaptation or Variant	Impact Type	95% CI (\$billion/year)	Mean (\$billion/year)
Agriculture	Agriculture	With CO <sub>2</sub> fertilization	Revenue lost from changes in wheat, cotton, soybean, and maize crop yields	\$0.42-\$19	\$6.1
Coastal Property	Infrastructure	Reactive Adaptation	Damage to coastal property value	\$5.9-\$21	\$9.4
Electricity demand and supply	Electricity	No <a href="#">Additional</a> Adaptation	Increases in power sector costs (e.g., capital, fuel, variable operation and maintenance (O&M), and fixed O&M cost)	\$2.4-\$21	\$11
Electricity transmission and distribution	Electricity	Reactive Adaptation	Damages to transmission & distribution infrastructure	\$6.9-\$14	\$11
<a href="#">Extreme Related Mortality Temperature-related mortality</a>	Health	No Adaptation	Mortality from changes in hot and cold <a href="#">extreme</a> temperatures	\$310 -\$9,900	\$2300
Hightide Flooding and Traffic	Infrastructure	Reasonably Anticipated Adaptation	Costs of traffic delays from flooding and cost of related infrastructure improvements	\$110 -\$200	\$141
Inland Flooding (residential)	Infrastructure	No <a href="#">Additional</a> Adaptation	Damages from riverine flooding	\$0.1-\$1.6	\$0.74
Labor Allocation	Labor	No <a href="#">Additional</a> Adaptation	Damages from work hours lost	\$6.7-\$220	\$51
Marine Fisheries	Ecosystems + Recreation	No <a href="#">Additional</a> Adaptation	Changes in thermally available habitat for commercial fish species	-\$0.1-\$0	-\$0.06
Long-Term Air Quality Exposure	Health	2011 Precursor Emissions	Mortality from ozone and fine particulate matter exposure	\$32 -\$9,900	\$230
Property and Violent Crime	Health	No <a href="#">Additional</a> Adaptation	Change in the number of Property and Violent crimes	\$0.1-\$2.0	\$0.92

Rail Infrastructure	Infrastructure	Reactive Adaptation	Infrastructure costs associated with temperature-induced track buckling	\$7.7-\$45	\$19
Road Infrastructure	Infrastructure	Reactive Adaptation	Cost of road repair, user costs (vehicle damage), and road delays due to changes in road surface quality	\$6.6-\$35	\$17
Southwest Dust	Health	No <a href="#">Additional Adaptation</a>	Mortality from changes in fine and coarse dust particle exposure	\$2.5-\$77	\$18
Tropical Storm Wind Damage	Infrastructure	No <a href="#">Additional Adaptation</a>	Cost of changes in hurricane wind damage to coastal properties	\$12-\$49	\$28
Urban Drainage	Infrastructure	Proactive Adaptation	Costs of proactive urban drainage infrastructure adaptation	\$3.2-\$5.0	\$4.2
Water Quality	Ecosystems + Recreation	<a href="#">No Additional Adaptation</a>	Willingness to pay to avoid water quality changes	\$0.83-\$3.8	\$2.0
Wildfire Air Quality Health Effects and Suppression Costs	Health	No <a href="#">Additional Adaptation</a>	Mortality from wildfire emission exposure and response cost for fire suppression	\$8.1-\$210	\$51
Winter Recreation	Ecosystems + Recreation	Adaptation	Revenue lost from suppliers of alpine, cross-country skiing, and snowmobiling	\$0.83-\$3.7	\$2.0
Valley Fever	Health	No <a href="#">Additional Adaptation</a>	Mortality, morbidity, and lost wages	\$2.0-\$58	\$14

595

## Annual Climate-Driven Impacts






Figure A2-1. Timeseries of sectoral damages across all 10,000 projections through 2100. Ordered by decreasing mean damages

600 in the year 2090. The lower right panel shows total damages summed across all sectors. Dashed Lines show the mean (dashed) and median (solid) line shows the mean damages each year. Shaded areas show the 95% CI. Annual damages are in units of billions of 2020USD (trillions for total panel). Temporal trends are a function of the underlying temperature (or sea level rise) binned damage functions, as well as sector specific scalars (e.g., per capita income-dependent VSL). Note, The slight discontinuities in some of these sectors (e.g., agriculture) can occur either at the boundary between temperature bins (e.g., for agriculture and wind damage) or due to thresholds in the underlying studies. For example, the sharp increase in damages in the coastal property damage sector after 2080 are directly reflected in the underlying damage functions and correspond to a sharp increase in damages that occur after sea levels breach 100 cm.

605



**Figure A3: Disproportionate vulnerability to climate-driven changes in air quality attributable mortality and labor hours lost, by race and vulnerable groups in 2090. Top panels) Difference in risk in 2090 for four vulnerable populations. Bottom panels) Additional rates of impacts in 2090, by race.**



620 Figure A2-24. Map of mean **total** annual climate-driven damages for a subset of sectors across 10,000 projections in each of the seven U.S. regions in the year 2090 (undiscounted). Damages are in billions of 2020USD. Donut charts show the absolute **total** damages (in billions) in each region for those sectors included in FrEDI, (in billions) and the top four sectors with the largest annual climate-driven damages. The share of damages from all remaining sectors are shown by the light gray wedge.

625 FrEDI also has a module to incorporate information from the recent EPA Report of Climate Change and Social Vulnerability in the United States: A Focus on Six Impacts (EPA, 2021a) (EPA 2021) (hereafter called the SV Report) to assess the differential climate-driven impacts in 2090 across different socially vulnerable groups. As described in the SV Report, this analysis considers four categories for which there is evidence of differential vulnerability. These groups are listed in Table A2-2.

630 **Table A2-2. Four socially vulnerable groups considered in this analysis and the reference groups (adapted from U.S. Environmental Protection Agency (2021ba))**

Categories	Group Name	Description	Reference Group
<u>Income</u>	<u>Low income</u>	<u>Individuals living in households with income that is 200% of the poverty level or lower</u>	<u>Individuals living in households with income greater than 200% of the poverty level.</u>
<u>Age</u>	<u>65 and Older</u>	<u>Ages 65 and older</u>	<u>Under age 65</u>
<u>Race and ethnicity</u>	<u>BIPOC</u>	<u>Individuals identifying as one or more of the following: Black or African American,</u>	<u>Individuals identifying as White and/or non-Hispanic</u>



	<u>American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, and/or Hispanic or Latino</u>		
<u>Education</u>	<u>No High School Diploma</u>	<u>individuals aged 25 and older with less than a high school diploma or equivalent</u>	<u>Individuals aged 25 or older with educational attainment of a high school diploma (or equivalent) or higher.</u>

These differential impacts are calculated in FrEDI at the Census tract level as a function of current population demographic patterns (i.e., percent of each group living in each census tract), projections of CONUS population (from ILCUS, U.S. Environmental Protection Agency, 2017), and projections of where climate-driven impacts are projected to occur (i.e., using FrEDI temperature-impact relationships) at the Census tract level. The relative percent of each socially vulnerable group in each Census tract are from the 2014-2018 U.S. Census American Community Survey dataset (U.S. Census) and are held constant overtime because robust and long-term projections for local changes in demographics are not readily available.

635

640 **Section A3: [FrEDI Adaptation and Uncertainty Results](#)**

FrEDI also has the additional capability to investigate some of these adaptation options in select sectors by reflecting the treatment in the underlying sector studies. [FrEDI maintains adaptation assumptions from the underlying studies that form the basis of FrEDI’s temperature-driven sectoral damage functions. For most of these studies, because the implicit or explicit impact response function is calibrated to historical or current data, this means that historically practiced adaptation or hazard avoidance actions are “baked in” – but enhanced adaptation action, or new \(currently unknown\) technologies are not considered. The exceptions include coastal property and select other infrastructure sectors, where the underlying studies consider specific adaptation actions. These have been incorporated into FrEDI. For example, for the coastal flooding sector, FrEDI’s default adaptation assumption is a Reactive Adaptation scenario, as defined in Neumann et al. \(2021\), and includes the costs \(and reflects the hazard reduction benefits\) of elevation of properties, where and when the benefits exceed the costs of this measure and expanded beach nourishment at locations where it is currently practiced. No other measures are included. There is an option in FrEDI, however, for the user to select either a No Adaptation scenario for this sector, which excludes the option to elevate properties as well as measures that might hold back floodwaters, or a Proactive Adaptation scenario, where adaptation measures include elevation, beach nourishment, and armoring \(either with bulkheads in protected areas or more expensive seawalls in areas exposed to higher open ocean wave action\). It is difficult to comment on the realism of future action. There is some discussion in both Neumann et al. \(2021\) and Lorie et al. \(2020\), both of which make the point that even under current coastal hazards, cost-effective adaptation measures have not been adopted, probably because they involve short term capital investment to yield future, uncertain benefits. This is one reason why Proactive adaptation is not the default scenario in FrEDI. Table A3 lists the FrEDI sectors that currently include adaptation options and the climate-driven damages for each in the year 2090.](#)

~~The no adaptation scenario represents a “business as usual” scenario additional adaptation options, while ‘reactive’ or ‘reasonably anticipated’ adaptation options reflect options taken without advanced warning or foresight, and ‘proactive’ or ‘direct’ adaptation reflects damages where cost-effective adaptations are implemented with perfect foresight.~~

For econometrically based sectors (e.g., Labor), adaptation is included to the extent that adaptation is currently occurring (e.g., work-place safety procedures currently being utilized to protect against extreme temperatures; individual risk/damage avoidance behavior reflected in current practice). For infrastructure sectors (i.e., Rail, Roads, Electricity Transmission and Distribution Infrastructure, Coastal Properties, and [Transportation Impacts from High Tide Flooding](#)), a [no additional](#) adaptation approach to infrastructure management does not incorporate climate change risks into the maintenance and repair decision-making process beyond baseline expectations and practice. The infrastructure sectors include two adaptation scenarios, following Melvin et al. (2017): Reactive adaptation, where decision makers respond to climate change impacts by repairing damaged infrastructure, but do

not take actions to prevent or mitigate future climate change impacts (a variant on this scenario is the “Reasonably anticipated adaptation” option for the High-Tide Flooding and Traffic sector, which is defined similarly to the Reactive scenario); and Proactive adaptation, where decision makers take adaptive action with the goal of preventing infrastructure repair costs associated with future climate change impacts. This Proactive Adaptation scenario assumes well-timed infrastructure investments, which may be overly optimistic given that such investments have oftentimes been delayed and underfunded in the past, and because decisionmakers and the public are typically not fully aware of potential climate risks (these barriers to realizing full deployment of cost-effective adaptation are described in Chambwera et al., 2014).

Table A3-1 shows that climate damages are sensitive to assumptions in the adaptation scenarios with mean 2090 annual damages of up to 2 to nearly 500 times larger in proactive or direct adaptation scenarios relative to damages when considering no adaptation. This illustrates adaptation has the capacity to both exacerbate and ameliorate future climate-driven damages.

**Table A3-1: Annual mean (and 95% confidence interval) climate-driven damages in 2090 for sectors that include different adaptation options. Damages are in billions of dollars (2020 USD).**

Sector	Adaptation Option <sup>A</sup>	Mean (\$billions/year)	95% CI (\$billions/year)
Electricity Transmission and Distribution	No Adaptation	\$12	\$7.3-\$18
	Reactive Adaptation	\$11	\$7.0-\$14
	Proactive Adaptation	\$6.3	\$5.0-\$8.3
Rail	No Adaptation	\$21	\$7.2-\$55
	Reactive Adaptation	\$19	\$7.7-\$45
	Proactive Adaptation	\$1.5	\$0.28-\$3.9
Roads	No Adaptation	\$135	\$25-\$330
	Reactive Adaptation	\$17	\$6.6-\$35
	Proactive Adaptation	\$7.3	\$5.8-\$8.4
Coastal Properties	No Adaptation	\$17	\$10.1-\$39
	Reactive Adaptation	\$9.4	\$6.0-\$21
	Proactive Adaptation	\$7.5	\$7.0-\$8.3
High Tide Flooding and Traffic Transportation Impacts from High Tide Flooding	No Adaptation	\$890	\$680-\$1,200
	Reasonably Anticipated Adaptation	\$140	\$110-\$200
	Direct Adaptation	\$2.0	\$1.3-\$3.4

<sup>A</sup>Default adaptation assumption in FrEDI is the Reactive or Reasonably Anticipated Adaptation [variantoption](#)

In addition to adaptation scenarios, FrEDI also has the capability to explore the sensitivity of future climate damages to specific changes in additional sectors, including agricultural damages with and without CO<sub>2</sub> fertilization, a lower air quality precursor emissions scenario, and high and low confidence intervals associated with damages specially from extreme-temperature-related related mortality. [Table A3-2X provides a snapshot of the structural uncertainty within the temperature-related mortality estimates. These results are not discussed](#)

here, but further illustrate the breadth of information provided by FrEDI. Within FrEDI, there are currently three underlying temperature-related mortality studies. We evaluate with mean damages from the RFF-SPs as well as the 5<sup>th</sup> and 95<sup>th</sup> % confidence intervals from the mean RFF-SPs. These confidence intervals are taken from the uncertainty within the underlying study. We note that the underlying data in Hsiang et al., is calculated as the median and therefore we are taking the mean across the RFF-SPs and the median damages. The Mills et al., study evaluates two scenarios, one with adaption and one without adaptation.

**Table A34-2: Annual mean (5<sup>th</sup>-95<sup>th</sup> % confidence interval) climate-driven damages in 2090 for premature mortality from temperature across three separate studies. Damages are in billions of dollars (2020 USD). Cromar et al., is used for temperature-related mortality throughout this analysis.**

<u>2090 Temperature-Related Premature mortality – Billions 2020 USD</u>		
<u>Model/Underlying Study</u>	<u>95<sup>th</sup> CI</u>	<u>Mean</u>
<u>Cromar et al.,</u>	<u>\$300 - \$3,900</u>	<u>\$2,100</u>
<u>Hsiang et al.,</u>	<u>-\$280 - \$1,800</u>	<u>\$740</u>
<u>Mills et al., (w/ adaptation)</u>	<u>=</u>	<u>\$31.0</u>
<u>Mills et al., (w/o adaptation)</u>	<u>=</u>	<u>\$110</u>

705 **Section A4: FrEDI through 2300**

FrEDI was calibrated to estimate impacts for detailed 21<sup>st</sup> century scenarios and trajectories, as described in Sarofim et al. (2021). Extending the FrEDI approach to 2300 requires two adjustments to adapt the sensitivity of the model to climate drivers and to socioeconomic conditions beyond the 21<sup>st</sup> century. First, we consider how the sensitivity to climate drivers (temperatures and SLR inputs) might differ from 21<sup>st</sup> century conditions. FrEDI damages were originally calibrated for temperatures from 0 to 6-degrees, relative to the 1986 to 2005 era, and SLR for 21<sup>st</sup> century trajectories that result in 30 to 250 cm GMSL outcomes by 2100. The original framework only returns physical and economic damage estimates within those bounds. In the modified FrEDI damage estimates for temperature inputs above these bounds are calculated by extrapolating damages per degree using the change in damages between 5 and 6 degrees. SLR inputs above the bounds are extrapolated based on the damages per centimeter of SLR modelled by the two highest sea level scenarios in 2090.

4Up to 6 degrees, FrEDI uses a piecewise linear function to estimate damages. This approach captures non-linearities from the underlying impact models. However, for temperatures above the calibration regime, FrEDI assumes a linear rate of change in damages equal to the change in damages from 5 degrees to 6 degrees. This assumption is likely to be conservative: Hsiang et al. (2017) found that combined damages in the United States increased quadratically with temperature, and Weitzman (Weitzman, 2012) suggested that while a quadratic damage form might be reasonable for temperature changes up to 2.5 degrees C globally, for higher temperatures it would make sense for damages to increase more quickly, as standard damage functions are unlikely to capture the sheer magnitude of impacts resulting from the kind of dramatic changes the planet would undergo at temperature changes substantially higher than that.

Second, we consider how the sensitivity to socioeconomic drivers continues beyond 2090 through 2300 on a sector-specific basis (Table A4-1). Damage estimates in FrEDI reflect year-specific socioeconomic conditions. There are several ways these conditions are defined through 2090 and linked to the damage estimates for temperature-based damages. Treatment for 2090 through 2300 is explained after the description of the original definition for each category of adjustments.

1. **Impacts scale with population and/or GDP per capita.** For sectors with explicit links to population and GDP, temperature-based damage estimates are scaled based on the population and GDP trajectory for a defined run. This is most common for health sectors, where total cases scale linearly with population and valuation of cases scales with GDP per capita. For example, willingness to pay to reduce fatality risk (referred to as the value of statistical life or VSL) is adjusted based on the projection of GDP per capita and a default income elasticity of 1.0. 2090 through 2300: Defined population and GDP trajectories continue to scale damage estimates through 2300.

2. **Year-specific Adjustment Factors.** In sectors where population and/or GDP per capita enter the impact function in complex ways that cannot be extracted and replicated within the FrEDI framework, a series of

year-specific adjustment factors defined based on the underlying study are used to adjust damages over time and/or space. For example, changes in health outcomes over time driven by demographic composition (e.g., population by age group or geographic distribution within region, which affect baseline mortality rates or exposure) are incorporated in FrEDI as year-specific adjustment factors. These factors are derived from the underlying studies via two methods:

a. By comparing per capita damage rates from a constant population run to a run that incorporates population growth<sup>11</sup>, resulting in a time series of adjustment factors. 2090 through 2300: The time series of adjustment factors is either linearly extrapolated through 2300 or held constant at 2090 levels based on the observed trends 2010 through 2090 and the interpretation of the factor.

b. By comparing per capita damage rates for two constant population scenarios (i.e., 2010 and 2090) and interpolating for between years. 2090 through 2300: Per capita damage rate adjustments are held at 2090 levels through 2300.

3. **No time-dependent adjustments.** Some sectors – which, in general, make up a small portion of overall damages– are not adjusted for socioeconomic projections but vary based only on sensitivity to projected temperature (Table A4-1). 2090 through 2300: No additional adjustments necessary.

Some sectors utilize more than one method (e.g., southwest dust outcomes scale linearly with population, method 1 in list above, and per capita mortality rates are adjusted over time based on method 2a).

**Sea level rise-based damages** in FrEDI are derived from damages in the underlying studies that are year and sea level rise specific through 2100, thus no additional time-dependent adjustments are necessary for that timeframe. Damages in each year reflect real property prices and adaptation decisions made in previous periods. 2090 through 2300: Damages post-2100 are based on sea level rise-based damages from 2100 adjusted for real property price appreciation using GDP per capita and income elasticity of 0.45, consistent with the underlying Neumann et al., (2021).

**Table A4-1. Summary of Strategy for Extending FrEDI Sectoral Results from 2090 to 2300 Modeling Horizon. Impact column provides detail for subcategories of impacts estimated within the Framework. Wildfire sector subcategories include morbidity and mortality associated with air quality impacts and fire suppression response costs – these two classes of subcategories are listed separately because they employ different extension strategies.**

<u>Sector</u>	<u>Impact</u>	<u>Extension Strategy</u>
<u>Air Quality</u>	<u>Ozone</u>	-
	<u>PM<sub>2.5</sub></u>	-

<sup>11</sup> Another, less common method for calculating adjustment factors is to compare two runs with and without climate change, each with population growth, to baseline damages (e.g. no population growth and no climate change).

<a href="#">Temperature-Related Mortality (Cromar et al., 2022)</a>	<a href="#">N/A</a>	-
<a href="#">Labor</a>	<a href="#">N/A</a>	-
<a href="#">Valley Fever</a>	<a href="#">Mortality</a>	<a href="#">Impacts continue to scale with population and/or GDP per capita (<i>Adjustment 1 in list above</i>)</a>
	<a href="#">Morbidity</a>	
	<a href="#">Lost Wages</a>	
<a href="#">Water Quality</a>	<a href="#">N/A</a>	-
<a href="#">Wildfire</a>	<a href="#">Morbidity</a>	-
	<a href="#">Mortality</a>	-
<a href="#">Winter Recreation</a>	<a href="#">Alpine Skiing</a>	<a href="#">Impacts continue to scale with population and/or GDP per capita (<i>Adjustment 1</i>)</a> <a href="#">AND</a> <a href="#">Year-specific adjustment factors developed from two constant population scenarios: per capita damages rates from 2090 applied 2090-2300 (<i>Adjustment 2b</i>)</a>
	<a href="#">Cross-Country Skiing</a>	
	<a href="#">Snowmobiling</a>	
<a href="#">Southwest Dust</a>	<a href="#">Acute Myocardial Infarction</a>	<a href="#">Year-specific adjustment factors developed from two constant population scenarios: per capita damages rates from 2090 applied 2090-2300 (<i>Adjustment 2b</i>)</a>
	<a href="#">All Cardiovascular</a>	
	<a href="#">All Mortality</a>	
	<a href="#">All Respiratory</a>	
<a href="#">Asthma ER</a>		
<a href="#">Electricity Supply and Demand</a>	<a href="#">N/A</a>	<a href="#">Year-specific adjustment factors developed based on comparison of with and without population growth scenarios: extend existing scalars linearly past 2090 (<i>Adjustment 2a</i>)</a>
<a href="#">Electricity Transmission and Distribution</a>	<a href="#">N/A</a>	
<a href="#">Roads</a>	<a href="#">N/A</a>	
<a href="#">Rail</a>	<a href="#">N/A</a>	
<a href="#">Coastal Properties</a>	<a href="#">N/A</a>	<a href="#">Sea level rise-based sectors: post-2090 impacts scale with GDP or GDP per capita</a>
<a href="#">Transportation Impacts from High Tide Flooding</a>	<a href="#">N/A</a>	
<a href="#">Asphalt Roads</a>	<a href="#">N/A</a>	-
<a href="#">Inland Flooding</a>	<a href="#">N/A</a>	-
<a href="#">Urban Drainage</a>	<a href="#">N/A</a>	-
<a href="#">Wildfire</a>	<a href="#">Response Costs</a>	-
<a href="#">Wind Damage</a>	<a href="#">N/A</a>	-
<a href="#">Marine Fisheries</a>	<a href="#">N/A</a>	-
<a href="#">Agriculture</a>	<a href="#">Cotton</a>	-
	<a href="#">Maize</a>	-
	<a href="#">Soybean</a>	-
	<a href="#">Wheat</a>	-
<a href="#">Crime</a>	<a href="#">Property</a>	-
	<a href="#">Violent</a>	-

As described in the main text, FrEDI is run through 2300 (Figure A4-15) to calculate the net present damages associated with an additional pulse of 1 tonne of CO<sub>2</sub> in the year 2020. In addition to the Ramsey discounting

770 approach presented in the main text, Figure [A4-26](#) provides a comparison to the net present damages calculated using a constant discount rate of 3%, consistent with OMB Circular A-4 (White House, 2003).



## Annual Climate-Driven Impacts





Figure A45-1. Timeseries of sectoral damages across all 10,000 projections from 2100-2300. Ordered by decreasing mean

775

damages in the year 2300. The lower right panel shows total damages summed across all sectors. Dashed line (solid) shows the mean (median) damages each year. Shaded areas show the 95% CI. Annual damages are in units of billions of 2020USD (trillions for total panel).



780

Figure A45-26. Net present value of future damages from one tonne of CO<sub>2</sub> for damages occurring only within the CONUS. Units are in dollars (2020 USD) per ton of CO<sub>2</sub> emitted. 'CDR' refers to Constant Discount Rate. Whiskers represent the 2.5<sup>th</sup> and 97.5<sup>th</sup> percentiles, while boxes span the 25<sup>th</sup> to 75<sup>th</sup>. Mean values (stars and text) along with median values (vertical lines) are also shown.

## Code Availability

785 The Framework for Evaluating Damages and Impacts (FrEDI) is available on the U.S. EPA Enterprise GitHub  
[https://github.com/USEPA/FrEDI/releases/tag/FrEDI\\_2300](https://github.com/USEPA/FrEDI/releases/tag/FrEDI_2300). FaIR is available at <https://github.com/OMS-NetZero/FAIR>. The RFF SP projections are available at <https://zenodo.org/record/6016583> and the SSP projections are available at <https://tntcat.iiasa.ac.at/SspDb/>.

## Data Availability

All code and data associated with this study are available at [www.github.com/USEPA/FrEDI\\_NPD](http://www.github.com/USEPA/FrEDI_NPD).

## 790 Author Contributions

CH, EEM, MS drafted the manuscript text and figures with contributions from all co-authors. BP, EEM, KN, and CH conducted the computational analysis. KN and JW developed the FrEDI code. SB drafted ~~and~~ [Figure 1](#) and provided input on graphics and all authors contributed to the writing of this manuscript.

## Competing Interests

795 The authors declare that they have no conflict of interest

## Acknowledgements

The views presented in this manuscript are solely those of the authors and do not necessarily represent the views or policies of the U.S. Environmental Protection Agency. Support for Industrial Economics was provided under EPA contracts 47QFSA21D0002 and 140D0420A0002. The authors also wish to acknowledge research assistance and  
800 other analytic support from William Maddock, Hayley Kunkle, Anthony Gardella, and Charles Fant.

## References

- Armstrong McKay, D. I., Staal, A., Abrams, J. F., Winkelmann, R., Sakschewski, B., Loriani, S., Fetzer, I., Cornell, S. E., Rockström, J., and Lenton, T. M.: Exceeding 1.5°C global warming could trigger multiple climate tipping points, *Science*, 377, eabn7950, <https://doi.org/10.1126/science.abn7950>, 2022.
- 805 Carleton, T. and Greenstone, M.: A Guide to Updating the US Government’s Social Cost of Carbon, *Rev. Environ. Econ. Policy*, 16, 196–218, <https://doi.org/10.1086/720988>, 2022.
- Carleton, T., Jina, A., Delgado, M., Greenstone, M., Houser, T., Hsiang, S., Hultgren, A., Kopp, R. E., McCusker, K. E., Nath, I., Rising, J., Rode, A., Seo, H. K., Viaene, A., Yuan, J., and Zhang, A. T.: Valuing the Global Mortality Consequences of Climate Change Accounting for Adaptation Costs and Benefits\*, *Q. J. Econ.*, 137, 2037–2105,  
810 <https://doi.org/10.1093/qje/qjac020>, 2022.

CEA: Discounting for public policy" theory and recent evidence on the merits of updating the discount rate, 2017.

815 Cromar, K. R., Anenberg, S. C., Balmes, J. R., Fawcett, A. A., Ghazipura, M., Gohlke, J. M., Hashizume, M., Howard, P., Lavigne, E., Levy, K., Madrigano, J., Martinich, J. A., Mordecai, E. A., Rice, M. B., Saha, S., Scovronick, N. C., Sekercioglu, F., Svendsen, E. R., Zaitchik, B. F., and Ewart, G.: Global Health Impacts for Economic Models of Climate Change: A Systematic Review and Meta-Analysis, *Ann. Am. Thorac. Soc.*, 19, 1203–1212, <https://doi.org/10.1513/AnnalsATS.202110-1193OC>, 2022.

Depsky, N., Bolliger, I., Allen, D., Choi, J. H., Delgado, M., Greenstone, M., Hamidi, A., Houser, T., Kopp, R. E., and Hsiang, S.: DSCIM-Coastal v1.0: An Open-Source Modeling Platform for Global Impacts of Sea Level Rise, *EGUsphere*, 1–47, <https://doi.org/10.5194/egusphere-2022-198>, 2022.

820 EPA: Multi-Model Framework for Quantitative Sectoral Impacts Analysis: A Technical Report for the Fourth National Climate Assessment., , U.S. Environmental Protection Agency, EPA 430-R-17-001., 2017a.

EPA: Updates To The Demographic And Spatial Allocation Models To Produce Integrated Climate And Land Use Scenarios (Iclus), , EPA/600/R-16/366F, 2017b.

825 EPA: Climate Change and Social Vulnerability in the United States: A Focus on Six Impacts., , U.S. Environmental Protection Agency, EPA 430-R-21-003, 2021a.

EPA: Technical Documentation on the Framework for Evaluating Damages and Impacts (FrEDI), , U.S. Environmental Protection Agency, EPA 430-R-21-004, 2021b.

830 Forster, P., T. Storelvmo, K. Armour, W. Collins, J.-L. Dufresne, D. Frame, D.J. Lunt, T. Mauritsen, M.D. Palmer, M. Watanabe, M. Wild, and H. Zhang: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, 2021.

GAO: CLIMATE CHANGE: Information on Potential Economic Effects Could Help Guide Federal Efforts to Reduce Fiscal Exposure, 2017.

835 Hsiang, S., Kopp, R., Jina, A., Rising, J., Delgado, M., Mohan, S., Rasmussen, D. J., Muir-Wood, R., Wilson, P., Oppenheimer, M., Larsen, K., and Houser, T.: Estimating economic damage from climate change in the United States, *Science*, 356, 1362–1369, <https://doi.org/10.1126/science.aal4369>, 2017.

840 Hultgren, A., Carleton, T., Delgado, M., Gergel, D. R., Greenstone, M., Houser, T., Hsiang, S., Jina, A., Kopp, R. E., Malevich, S. B., McCusker, K. E., Mayer, T., Nath, I., Rising, J., Rode, A., and Yuan, J.: Estimating Global Impacts to Agriculture from Climate Change Accounting for Adaptation, <https://doi.org/10.2139/ssrn.4222020>, 16 September 2022.

Kopp, R. E., Kemp, A. C., Bittermann, K., Horton, B. P., Donnelly, J. P., Gehrels, W. R., Hay, C. C., Mitrovica, J. X., Morrow, E. D., and Rahmstorf, S.: Temperature-driven global sea-level variability in the Common Era, *Proc. Natl. Acad. Sci.*, 113, E1434–E1441, <https://doi.org/10.1073/pnas.1517056113>, 2016.

845 Kotchen, M. J.: Which Social Cost of Carbon? A Theoretical Perspective, *J. Assoc. Environ. Resour. Econ.*, 5, 673–694, <https://doi.org/10.1086/697241>, 2018.

Lay, C. R., Sarofim, M. C., Vodonos Zilberg, A., Mills, D. M., Jones, R. W., Schwartz, J., and Kinney, P. L.: City-level vulnerability to temperature-related mortality in the USA and future projections: a geographically clustered meta-regression, *Lancet Planet. Health*, 5, e338–e346, [https://doi.org/10.1016/S2542-5196\(21\)00058-9](https://doi.org/10.1016/S2542-5196(21)00058-9), 2021.

- 850 Martinich, J. and Crimmins, A.: Climate damages and adaptation potential across diverse sectors of the United States, *Nat. Clim. Change*, 9, 397–404, <https://doi.org/10.1038/s41558-019-0444-6>, 2019.
- Martinich, J., DeAngelo, B., Diaz, D., Ekwurzel, B., Franco, G., Frisch, C., McFarland, J., and O'Neill, B.: Reducing Risks Through Emissions Mitigation. In *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment*, 2018.
- 855 Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J.: The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500, *Geosci. Model Dev.*, 13, 3571–3605, <https://doi.org/10.5194/gmd-13-3571-2020>, 2020.
- 860 Melvin, A. M., Larsen, P., Boehlert, B., Neumann, J. E., Chinowsky, P., Espinet, X., Martinich, J., Baumann, M. S., Rennels, L., Bothner, A., Nicolsky, D. J., and Marchenko, S. S.: Climate change damages to Alaska public infrastructure and the economics of proactive adaptation, *Proc. Natl. Acad. Sci.*, 114, E122–E131, <https://doi.org/10.1073/pnas.1611056113>, 2017.
- NAS: Updating Climate Damages" Updating Estimations of the Social Cost of Carbon Dioxide, 2017.
- 865 Neumann, J. E., Chinowsky, P., Helman, J., Black, M., Fant, C., Strzepek, K., and Martinich, J.: Climate effects on US infrastructure: the economics of adaptation for rail, roads, and coastal development, *Clim. Change*, 167, 44, <https://doi.org/10.1007/s10584-021-03179-w>, 2021.
- Nordhaus, W. D.: Revisiting the social cost of carbon, *Proc. Natl. Acad. Sci.*, 114, 1518–1523, <https://doi.org/10.1073/pnas.1609244114>, 2017.
- 870 O'Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P., Birkmann, J., Kok, K., Levy, M., and Solecki, W.: The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century, *Glob. Environ. Change*, 42, 169–180, <https://doi.org/10.1016/j.gloenvcha.2015.01.004>, 2017.
- 875 Rennert, K., Prest, B. C., Pizer, W., Newell, R. G., Anthoff, D., Kingdon, C., Rennels, L., Cooke, R., Raftery, A. E., Sevcikova, Hana, and Errickson, F.: The Social Cost of Carbon: Advances in Long-Term Probabilistic Projections of Population, GDP, Emissions, and Discount Rates, *Resour. Future*, Working Paper 21-28, 2021.
- Rennert, K., Errickson, F., Prest, B. C., Rennels, L., Newell, R. G., Pizer, W., Kingdon, C., Wingenroth, J., Cooke, R., Parthum, B., Smith, D., Cromar, K., Diaz, D., Moore, F. C., Müller, U. K., Plevin, R. J., Raftery, A. E., Ševčíková, H., Sheets, H., Stock, J. H., Tan, T., Watson, M., Wong, T. E., and Anthoff, D.: Comprehensive Evidence Implies a Higher Social Cost of CO<sub>2</sub>, *Nature*, 1–3, <https://doi.org/10.1038/s41586-022-05224-9>, 2022.
- 880 Revesz, R., Greenstone, M., Hanemann, M., Livermore, M., Sterner, T., Grab, D., Howard, P., and Schwartz, J.: Best cost estimate of greenhouse gases, *Science*, 357, 655–655, <https://doi.org/10.1126/science.aao4322>, 2017.
- Rising, J. and Devineni, N.: Crop switching reduces agricultural losses from climate change in the United States by half under RCP 8.5, *Nat. Commun.*, 11, 4991, <https://doi.org/10.1038/s41467-020-18725-w>, 2020.
- 885 Rising, J., Tedesco, M., Piontek, F., and Stainforth, D. A.: The missing risks of climate change, *Nature*, 610, 643–651, <https://doi.org/10.1038/s41586-022-05243-6>, 2022.
- Rode, A., Carleton, T., Delgado, M., Greenstone, M., Houser, T., Hsiang, S., Hultgren, A., Jina, A., Kopp, R. E., McCusker, K. E., Nath, I., Rising, J., and Yuan, J.: Estimating a social cost of carbon for global energy consumption, *Nature*, 598, 308–314, <https://doi.org/10.1038/s41586-021-03883-8>, 2021.

- 890 Sarofim, M. C., Martinich, J., Neumann, J. E., Willwerth, J., Kerrich, Z., Kolian, M., Fant, C., and Hartin, C.: A temperature binning approach for multi-sector climate impact analysis, *Clim. Change*, 165, 22, <https://doi.org/10.1007/s10584-021-03048-6>, 2021.
- Smith, C. J., Forster, P. M., Allen, M., Leach, N., Millar, R. J., Passerello, G. A., and Regayre, L. A.: FAIR v1.3: a simple emissions-based impulse response and carbon cycle model, *Geosci. Model Dev.*, 11, 2273–2297, <https://doi.org/10.5194/gmd-11-2273-2018>, 2018.
- 895 US Census: American Community Survey, Available [Wwwdatacensus.gov](http://www.datacensus.gov), 2014.
- Weitzman, M.: GHG Targets as Insurance Against Catastrophic Climate Damages, *J. Public Econ. Theory*, 14, 221–244, 2012.
- White House: Circular A-4. Regulatory Analysis, , Washington: Office of Management and Budget, 2003.
- 900 Wong, T. E., Bakker, A. M. R., Ruckert, K., Applegate, P., Slangen, A. B. A., and Keller, K.: BRICK v0.2, a simple, accessible, and transparent model framework for climate and regional sea-level projections, *Geosci. Model Dev.*, 10, 2741–2760, <https://doi.org/10.5194/gmd-10-2741-2017>, 2017.
- Wong, T. E., Rennels, L., Errickson, F., Srikrishnan, V., Bakker, A., Keller, K., and Anthoff, D.: MimiBRICK.jl: A Julia package for the BRICK model for sea-level change in the Mimi integrated modeling framework, *J. Open Source Softw.*, 7, 4556, <https://doi.org/10.21105/joss.04556>, 2022.
- 905