

The climate in Poland (Central Europe) in the first half of the last millennium, revisited

Rajmund Przybylak^{1,2}; Piotr Oliński^{2,3}; Marcin Koprowski^{2,4}; Elżbieta Szychowska-Krąpiec⁵; Marek Krąpiec⁶; Aleksandra Pospieszyńska^{1,2}; Radosław Puchałka^{2,4}

5

¹Department of Meteorology and Climatology, Faculty of Earth Sciences and Spatial Management, Nicolaus Copernicus University, Toruń, Poland

²Centre for Climate Change Research, Nicolaus Copernicus University, Toruń, Poland

10 ³Department of Medieval History, Institute of History and Archival Sciences, Faculty of History, Nicolaus Copernicus University in Toruń, Poland

⁴Department of Ecology and Biogeography, Faculty of Biology and Environmental Protection, Nicolaus Copernicus University, Toruń, Poland

⁵Department of Environmental Analysis, Geological Mapping and Economic Geology, Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology, Kraków, Poland

15 ⁶Department of General Geology and Geotourism, Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology, Kraków, Poland

Correspondence: Rajmund Przybylak (rp11@umk.pl)

Abstract. The article presents **updated** ~~current state of~~ knowledge on climate change in Poland (Central Europe) in the first half of the last millennium (1001–1500). **This knowledge is required to delimit the existence, duration and scale of the Medieval Warm Period (MWP, also called the Medieval Climate Anomaly [MCA] here).** To this end, it employs all available quantitative climate reconstructions created **for Poland** in the last two decades and four new reconstructions using three dendrochronological series and an extensive database of historical source data on weather conditions. The growth of conifers in lowland and upland Poland depends on the temperature in the cold season, especially in February and March. All available reconstructions based on dendrochronology **data** represent this time of the year. Summer temperatures were reconstructed using biological proxies and documentary evidence. The latter, however, is limited to the 15th century only. Winter temperature was used as the proxy for annual temperature proxies instead of the more usual use of summer temperature. The **MWP** ~~Medieval Warm Period (MWP; also called the Medieval Climate Anomaly [MCA])~~ occurred in Poland probably from the late 12th century to the first halves of the 14th or 15th centuries. All the analysed quantitative reconstructions suggest that the MWP in Poland was comparable to or warmer than the **mean** ~~current~~ **temperature in the** ~~period (1951–2000).~~ The coldest conditions in the entire study period were noted in the first half of the 11th century (both winter and summer) and the second half of the 15th century (only winter). The greatest climate continentality occurred in the 15th century. Good agreement was found between the reconstructions of Poland's climate and many reconstructions available for Europe.

Keywords: Climate change; Medieval times; Documentary evidence; Dendrochronological data; Climate reconstructions; Extreme events

1. Introduction

Knowledge about weather and climate in the pre-instrumental period, particularly for the last millennium, is fundamental and necessary to identify past and potential future drivers of climate change (Brönnimann et al., 2019; Brönnimann, 2022; Sieber et al., 2022). The traditional scheme of climate changes in this time that was proposed by Lamb (1965, 1977, 1984), in particular the occurrence of the Medieval Warm Period (MWP, recently called also the Medieval Climate Anomaly, MCA), was recently questioned on a global scale (e.g., Hughes and Diaz, 1994; Brázdil et al., 2005; Diaz et al., 2011); nonetheless, it is still valid for central and western Europe and some other areas mainly in the Northern Hemisphere (Mann et al. 2009). More reliable climate reconstructions for the MCA and the Little Ice Age (LIA) are needed for many areas to obtain better knowledge about spatial temperature patterns. According to Mann et al. (2009), they are still poorly defined.

In the present paper, we focus on climate reconstruction for the area of modern-day Poland in the period 1001–1500, when the anthropogenic fingerprints were small (Hernández-Almeida et al., 2017). According to Niedźwiedz et al. (2015), this period encompasses the MCA (1001–1350) and the Transitional Period (TP, 1351–1500) in the Baltic Sea Basin, which includes Poland. Przybylak (2016), summarising climate knowledge for that time, found that the MCA in Poland began in the 11th century and probably ended in the 14th or early 15th century, and that the LIA started in the mid-16th century.

An increase in knowledge of the Polish climate from the so-called pre-instrumental period is observable in the last several decades. However, the majority of available works cover the entire period since 1500, but more often only short sub-periods thereof (e.g., Przybylak et al., 2005; Przybylak and Pospieszyńska, 2010 and references therein; Opała-Owczarek et al., 2021 and references therein). **There are significantly fewer studies dealing with the pre-1500 period compared to the post-1500 period.** Only a few papers also deal with a pre-1500 period (e.g., Maruszczak, 1988; Sadowski, 1991; Wójcik et al., 2000; Zielski et al., 2010; Przybylak, 2011, 2016; Koprowski et al., 2012; Hernández-Almeida et al., 2015, 2017; Balanzategui et al., 2017; Przybylak et al., 2020; Oliński, 2022).

Przybylak (2016) concluded that “the state of knowledge concerning changes in air temperature in Poland in the period 1001–1500 is very limited [...] existing reconstructions being very uncertain. In some periods there are even cases in which opposing trends for the course of air temperature are presented.” Significantly worse knowledge is available for other meteorological variables and related phenomena, like droughts (Przybylak et al. 2020) and floods (Ghazi et al., 2023). Such information about the climate of this period is also noted in many different areas of the world. Therefore in recent years, increasing attention has been directed to this period (Vogt et al., 2011; Goosse et al., 2012; Moreno et al., 2012; Pribyl, 2014; Camenisch, 2015; Camenisch et al., 2016).

In this paper we presented an updated knowledge on the Poland’s climate in the period 1001–1500 using multiproxy data (mainly documentary evidence and dendrochronological data). For this purpose, a few new quantitative climate reconstructions have been used. Based on documentary evidence, mean 10-year winter (DJF) and summer (JJA) air temperatures for the 15th century were estimated. On the other hand, the dendrochronological data allowed the mean late

winter–early spring temperature to be reconstructed encompassing the period since the 12th century. **The newly obtained knowledge allowed us to more precisely describe the climate in this period, including the occurrence and scale of warming during the MCA.**

2. Area, data and methods

2.1. Area

The analysis is presented for Poland in its contemporary boundaries (Fig. 1). Eight Natural-Forest Provinces (Zielony and Kliczkowska, 2012) with the location of dendrochronological sites, as well as all historical regions for which documentary evidence are available, are also shown on the map. In the case of historical regions whose boundaries were changeable in the past, some sources were taken into consideration that were written outside of the contemporary Polish territory.

2.2. Documentary evidence

The primary type of much of the valuable information about the climate of medieval Poland is narrative sources, e.g., chronicles, annals, memoirs and private correspondence. These sources usually describe weather conditions throughout the seasons, mainly in winter and summer. There is also a strong focus on severe droughts, floods and other weather, climate and water extremes (WCWs) like hails, thunderstorms, snowstorms, or sea, lake and river ice. In many cases, they describe the consequences for the economy and society and activities to mitigate their effects.

In our research, we performed a critical analysis of historical sources reporting on climatic phenomena, which allowed us to choose the most reliable “first-hand” sources for the present work.

Another type of source includes administrative and economic sources, i.e., official correspondence or business records containing direct references to WCWs. These also include requests for tax exemption or even financial support related to, for example, flood damage, such as the costs of repairing bridges, flood embankments. In addition to the descriptions of various events, numerical data on sizes or losses of harvests from some of these sources can be processed, and these can then be linked to the economic consequences of the WCWs expressed in quantitative data.

The search for historical sources from this period undertaken by us for the purpose of this work provided significantly more new information than Maruszczak (1991) and Sadowski (1991) used in their reconstructions. We obtained, among other things, many mentions referring to weather from Teutonic correspondence contained in the *Ordens-Brief Archiv* and were able to correct some information from medieval chronicles hitherto either incorrectly dated or uncritically treated as fully credible. It should be emphasised that the first of the aforementioned temperature reconstructions for Poland (Maruszczak, 1991) did not use any historical sources from Poland.

We analysed a total of 813 weather records, of which 145 from the period 1001–1360 (Table S1, Fig. 2) and 668 from the period 1361–1500 (Table S1, Fig. 2). **It should be emphasized that there was a significant increase in the number of**

100 available historical written references from the 1360s onwards, and therefore 1360 was chosen as the threshold year for delimiting the two subperiods. The sources used are listed in Table S2.

We considered it important to list the years from the oldest sub-period analysed in this work (AD 1001–1360) for which reliable historical sources describing weather conditions are available (including those documenting the occurrence of severe winters and hot summers or other seasons) and to make them available in Table S3. Based on these sources, the thermal and pluvial conditions for four seasons were classified jointly using the seven-degree index scale (Pfister et al., 1994). Examples of typical descriptions of weather in winter and summer based on which we classified thermal and precipitation conditions into seven categories (-3, ..., +3) are presented in Table S4. Of those first 360 years, the amount of weather information collected was, as is to be expected, greatest for the 14th century (75 items) and least for the 11th century (only 13 items) (Fig. 2). We collected the most weather notes for winter (39.3%), undetermined season (24.8%) and autumn 110 (14.5%), and the fewest for spring (9.0%).

For the period 1361–1500, significantly more weather records are available, especially for the last 50 years, when the number ranges around 100 per decade (excluding the decade 1481–90) (Table S1, Fig. 2). As in the previous period, the number of notes was greatest for winter (33.5%). The weather in summer was described with a slightly lower frequency (27.7%).

115 An indexation of thermal and precipitation conditions was made for the period 1361–1500, similarly as for the previous period. It was based on the information collected from the types of historical sources presented above, was supported by previous catalogues (Walawender, 1932; Girguś and Strupczewski, 1965 database of natural disasters: <http://pth.net.pl/projekty/bazy-danych/kleski-elementarne/do-1795>) and took into account earlier climate reconstructions for the 15th century (Polackówna, 1925; Sadowski, 1991; Przybylak, 2011). For both indexation periods, this is the first time 120 for Poland that pluvial conditions have been indexed and that air temperature has been indexed for the transitional seasons (e.g., spring and autumn).

The new indexation was used to update and significantly supplement the reconstruction of 10-year average winter and summer temperatures for the 15th century that were previously presented by Przybylak (2011). Standard methods used commonly in the historical climatology were utilised for this purpose. Pfister et al. (1994) proposed that, for the following index values: i) +3 and -3 should represent anomalies exceeding 2.0 standard deviations (SD) from the mean of the long-term period, ii) +2/-2 and +1/-1 should represent less extreme conditions, i.e. 1.41–2.00SD and 0.7–1.4SD, respectively, iii) 0 (>-0.7SD – <0.7SD) should represent the average climate of the long-term period or missing data. The above-mentioned criteria were slightly modified by us using results of calibration of documentary evidence with air temperature data from Warsaw for the period 1789–1850 done by Sadowski (1991). As a result, the seasons described in this paper by indices from 130 +3 to -3 fulfil the following criteria:

$$+3 \geq m + 1.5SD$$

$$m + 1.0SD \leq +2 < m + 1.5SD$$

$$m + 0.5SD \leq +1 < m + 1.0SD$$

$$m - 0.5SD < 0 < m + 0.5SD$$

$$m - 1.0SD < -1 \leq m - 0.5SD$$

$$m - 1.5SD < -2 \leq m - 1.0SD$$

$$-3 \leq m - 1.5SD$$

135

140

145

where m is the long-term mean (for calibration period 1789–1850) air temperature from the Warsaw series and SD is the standard deviation of that series. For more details, see Przybylak et al. (2005). The reader should be reminded of the possible biases related to climate reconstruction based on the documentary evidence such as, for example: i) the number of available sources, their quality and discontinuous structure, ii) the subjectivity of the indexation, iii) weaknesses of the reconstruction method used. A detailed overview of the strong points and the drawbacks of documentary data is presented by Brázdil et al. (2005). The method used for this reconstruction is described in a paper by Przybylak et al. (2005) and is therefore omitted here.

The amount of information available for the period 1361–1400 (Table S1, Fig. 2) does not currently permit a quantitative reconstruction. For summers, too, for the same reason, it was not possible to reconstruct the first and third decades of the 15th century.

150

A comparative indexation based on historical sources and dendrochronological data was also undertaken that indicated consistencies and disparities in the assessment of individual extreme years.

2.3. Dendrochronological data

2.3.1. Material and removal of age-related development trends

155

The regional tree-ring chronologies used in the study were developed by Zielski (1997), Krapiec (1998), Zielski and Krapiec (2004) and Szychowska-Krapiec (2010). The wood samples were taken from forest stands, historic architectural objects and archaeological excavations (Fig. 1). The cores from living trees were taken using incremental borers of 5 mm diameter at a height of ~1.5 m, dried and mounted on wooden holders (Cook and Kairiukstis, 1990). Samples were taken from historic buildings, in the form of 15-mm cores or discs. The measuring paths were cut along 2 or 3 core radii using a dissecting knife.

160

The width of annual rings was measured under microscope with an accuracy of 0.01 mm. Each dated series was checked for measurement errors and missing rings using the COFECHA software (Holmes, 1983). The averaged measurement series of growing trees and historical wood were combined into one regional chronology.

To remove age-related trends from raw grain width measurements, the procedure described in Melvin and Briffa (2008) was used, with processing in RCSigFree 45_v2b software (Cook et al., 2014). Variance was stabilised using an Rbar-

165 weighted method to adjust for changes in sample size (Osborn et al., 1997; Frank et al., 2007) followed by a robust bi-weight mean applied to develop a final indexed chronology (Cook, 1985; Cook and Kairiukstis, 1990).

The current Rbar and Expressed Population Signal (EPS) metrics were computed in 21-year windows with a 10-year overlap. These metrics were used to assess the strength and stability of the common signal between trees over time, and to determine the length of the final chronology used for the reconstruction. Rbar is a measure of the percentage of shared
170 variance between individual runs, with higher Rbar values indicating a stronger common signal (Briffa, 1995). EPS, by contrast, provides estimates of how well a finite number of samples represent a theoretically infinite population (Wigley et al., 1984). EPS values above 0.85 were taken as indicating a period when the ring-width data showed a strong common signal (Table 1).

The sensitivity of the tree-ring chronology of the examined trees to the climate was tested using long-term series of
175 average monthly air temperature values and monthly precipitation sums from Kraków and Toruń (Fig. 1) using the method proposed by Fritts (see Table 1). We confirmed (Table 1) the results presented earlier by Zielski (1997), who reveals statistically significant correlations between annual tree-ring widths (Kuyavia-Pomerania dendrochronology, 1891–1991) and the monthly mean air temperatures from the region, particularly from February and March, but also from January and April. Their values were equal to 0.47, 0.55, 0.26 and 0.18, respectively. This means that, in Poland, the low temperature
180 occurring at the end of winter and at the beginning of spring has a strong negative influence on the width of tree-rings. On the other hand, precipitation has a weaker influence than temperature, though only in June and July is this statistically significant. More information about climate signal in selected trees in Poland is provide in Table 7.1 in Zielski et al. (2010). After identifying the strongest and most time-stable relationship between ring width and the listed meteorological parameters, a transfer function based on linear regression was built using ring-width chronology as the climate predictor. To
185 assess the stability of the linear model, calibration/verification tests were performed at different times depending on the length of the climatic data and chronology (Table 1). The calculations were made using the R software (RCoreTeam, 2022), the dplR package (Bunn, 2008) and treeclim package (Zang and Biondi, 2015).

2.3.2. Moon rings, included sapwood

190 A good indicator of extreme cold thermal conditions during winter is moon rings (MR), also known as “included sapwood”, which are found in the wood of European oaks (*Quercus robur* and *Quercus petraea*). They are so called for their presentation as halos in the cross-section of the dark heartwood. The wood of an MR shows an absence or low number of tyloses in the vessels and a lower content of heartwood substances (Dujesiefken and Bauch, 1987; Dzbeński and Krutul, 1994). MR are caused by disturbances in starch management disrupting the heartwood formation process (Dujesiefken and Bauch, 1987) and is caused particularly by cold winters (Bolychevtsev, 1970; Dujesiefken and Liese, 1986; Krapiec, 1998).
195

Samples with MRs were identified from a collection of about 2,500 discs and cores from oaks from Holocene alluvial deposits in southern Poland, archaeological excavations and wood from buildings. MR zones were identified on a prepared cross-sectional area under a binocular magnifier. For each site, a correlation diagram of trunks with increments in

the MR zone marked was prepared. Since some of the samples come from the lower trunk, and the area of MR is smaller in this part (Krapiec, 1999; Dujesiefken and Liese, 1986), the dating of the last increment from the MR zone may differ by about 2–5 years from the actual date of the moon ring's formation.

2.3.3. Climate reconstruction pointer years

For climate reconstruction, we employed two approaches. The first involves the pointer years identified for all chronologies, both oak and coniferous trees (Table S5), and the second involves a regression model used for pine from the Kuyavia-Pomerania region and pine and oak from Lesser Poland.

Pointer years (Huber and Giertz-Siebenlist, 1969) were defined as those in which weather conditions cause the vast majority of trees to develop a narrower or wider ring than in the previous year. They appear as a result of various factors, both short-term (e.g., one-night, late-spring frosts) and long-term (e.g., droughts, severe winters, cold spells) (Schweingruber, 1992). Despite the weather conditions of individual pointer years sometimes being hard to explain, their usefulness in climate change research and dendrochronological dating is unquestionable. In this paper, the criterion for pointer years was that a 90% match be found for a threshold of ten or more trees.

2.3.4. Climate reconstruction regression model

Previous research proves that the correlation between Scots pine tree rings and climate in northern (Zielski, 1997; Zielski et al., 2010; Waszak et al., 2021) and southern (Szychowska-Krapiec, 2010) Poland is strongly dependent on temperature. Similar relationships were observed for fir from southern Poland (Szychowska-Krapiec, 2010). Hence, we assumed that the previous findings justify our temperature reconstruction. After the study of the regression model, the calibration period for three sites was chosen as the earlier one (Table 1). The highest correlation was observed for fir from Lesser Poland, with 0.57 for the calibration period and 0.49 for the whole period. Positive RE and CE values for verification intervals indicated that the chosen linear models have good predictive skill suitable for reconstruction. Furthermore, the RMSE test is very good, with values below 1.

~~Statistics of RE and CE tests yielded a reasonable model, and the RMSE test was very good, with values below 1.~~

3. Results

225 3.1. Documentary evidence

3.1.1. 1001–1360

For this period, there exists only one reconstruction of air temperature based on documentary evidence (Sadowski 1991). The author calculated the frequencies of severe winters and hot summers since the 13th century using weather descriptions available mainly in the chronicle *Annales seu cronici incliti regni Poloniae* of Jan Długosz (1415–80). He found 18 severe
230 winters and 18 hot summers in this time. An exceptionally high decadal value of hot summers (7) occurred in the 1330s. Only in one decade (the 1280s) did as many as three severe winters occurred.

The number of sources for the period 1001–1360 is small and has remained essentially unchanged for many years. The probability of finding new sources is negligible, as also shown by our query. As a result, the number of mentions of weather conditions we have collected is limited (Table S1, Fig. 2). They cannot be used to create a quantitative
235 reconstruction – nor even a full qualitative reconstruction – of changes in climatic conditions. Meanwhile, the scarcity of the available sources requires that their credibility should, all the more, be thoroughly reassessed. Below, we present only selected examples of extreme hydrometeorological events recorded in the sources. However, in the Table S3, we present all such events. The descriptions are decidedly dominated by information on floods and severe winters. Jan Długosz in his *Annales* was the first to mention the weather in Polish lands when describing the year 988. Then, after numerous floods,
240 there was an exceptionally hot summer.

For severe winters, the descriptions indicated rivers freezing, the Baltic Sea freezing, and the prevailing winter conditions in descriptions of military events. Several Western European sources tell, for example, of a severe winter in
Lusatia in 1069. These descriptions were associated with a military campaign being waged by the German emperor in that territory. Such weather conditions may also have prevailed in Poland at that time. Western sources also mention the severe
245 winter of 1076/77, during which the Elbe and Vistula rivers froze over. From sources established in Poland lands, we know of the severe winters of 1110/11, 1124/25, 1204/05, 1234/35, 1252/53, 1257/58, 1279/80, 1285/86, 1305/06 and 1322/23. However, not all of these severe winters are entirely certain. Some doubts as to the sources exist regarding the winters of 1234/35, 1252/53, 1257/58 and 1305/06. Meanwhile, it has been erroneously reported that there was a severe winter in Polish lands in 1224/25.

250 In the period 1001–1360 we have evidence for occurrence of 22 extremely severe and very severe (indices -3 and -2) winters, and only one extremely warm and very warm (3 and 2) summer.

~~Another frequent phenomenon reported in the oldest sources is floods, which were studied by Ghazi et al. (in review) and these are therefore not described here.~~

3.1.2. 1361–1500

255 Of the study period, our supplementary query significantly increased the number of historical sources describing weather conditions for only the second half of the 14th century onwards. Thus, the knowledge about the climate of Poland presented here is new, being the most complete and reliable record to date. Particularly valuable is the quantitative reconstruction of the winter and summer temperatures for the 15th century.

260 The frequency of winters (DJF) and summers (JJA) in Poland that were either extremely warm and wet or extremely cold and dry in the period from 1361 to 1500 is shown in Table 2. It is in line with expectations that sources reported the occurrence of extremely cold and very cold winters with a greater frequency (41) than extremely warm and very warm winters (10). For summers, on the other hand, extremely warm and very warm seasons were about twice as frequent as cold seasons. Large and very large amounts of precipitation were associated with extremely warm and very warm winters, and were significantly rarer during cold winters. On the other hand, the thermal character of summer (warm, cold) did not
265 differentiate the occurrence of heavy precipitation (Table 2). Time runs of seasonal frequencies of all categories of thermal and humidity extremes in Poland in 10-year periods are shown in Figs 3 and 4. In the study period, but in particularly in the 15th century, extremely cold and very cold winters dominated almost all decades (Fig. 3). Such winters were frequent in the 1430s and 1450s (7 cases each) and the 1490s (5). In the first two decades, springs were also very cold. Close to normal thermal conditions in summer were noted in the first 80 years (1361–1440). Later, a decade-to-decade warming of summers
270 accelerated, to reach a maximum in the 1470s (Fig. 3). In the two warmest summer decades (1461–70 and 1471–80) warming was also noted very often in autumn (Fig. 3).

The documentary evidence gives significantly less information about the wetness of the seasons. In line with expectations for Poland, such information dominates for summer and then for winter, but is only rarely available for the transitional seasons (Fig. 4). Very little information was gathered for the 14th century. In the 15th century, all categories of
275 wet winters dominated, with a maximum in the 1430s (4). This means that the 1430s in Poland were characterised by very cold and snowy winters. Except for last three decades of the 15th century, summers were also dominated by extremely wet and very wet conditions, in particular in the 1460s (5 cases). Radical dryness of the air occurred in the 1470s (Fig. 4).

According to the new reconstruction (Fig. 5), average winter temperatures were lowest in the 1450s. Then, the average winter temperature was $-6.7\text{ }^{\circ}\text{C}$ and was as much as $5.3\text{ }^{\circ}\text{C}$ lower than in the period 1951–2000 in Poland ($-1.4\text{ }^{\circ}\text{C}$,
280 Kożuchowski and Żmudzka, 2003) and $6.1\text{ }^{\circ}\text{C}$ lower than in 1991–2020 ($-0.6\text{ }^{\circ}\text{C}$, Tomczyk, 2022). **These two present mean temperatures for the area of Poland were calculated based on data taken from 45–50 and 40 weather stations, respectively.** The next coldest decades in terms of winter temperatures were the 1430s ($-5.5\text{ }^{\circ}\text{C}$) and the 1490s ($-4.8\text{ }^{\circ}\text{C}$). The warmest winters were in the decade 1481–90 ($-3.4\text{ }^{\circ}\text{C}$). In the contemporary climate of Poland, only the coldest winters (e.g., winter 1995/96 with an average value of $-5.1\text{ }^{\circ}\text{C}$, Tomczyk, 2022) reached values similar to the reconstructed 15th-century winter
285 temperatures.

The decadal average summer temperature in Poland in the 15th century ranged from 17.8 °C (1431–40) to 19.2 °C (1471–80), as compared to 17.0 °C for 1951–2000 (Kožuchowski and Źmudzka, 2003) and 17.6 °C in 1991–2020 (Tomczyk, 2022). In all decades of the 15th century, the decadal average summer temperature was significantly higher than at present, with positive anomalies (relative to 1951–2000) ranging from 0.8 °C to 2.2 °C (Fig. 5). In recent years, however, the summer temperature in Poland has been comparable to that in the warmest decade of the 15th century; for example, in the summer of 2019 it was 19.5 °C (Tomczyk, 2022). The presented results also clearly confirm the correctness of the assessment of the Polish climate as continental in this period, as calculated by Sadowski (1991, see his Fig. 6). He showed that the climate of the 15th century was the most continental, far more so than the contemporary current climate or that of the 13th and 14th centuries.

295

3.2. Dendrochronology

3.2.1. Moon rings (MR)

During the period 1001–1500, MRs, which represent cold winter conditions, were found for years 1141/42, 1314/15, 1328/29 in Nysa; 1186/87 in Czermno (Lublin Province); 1332/33 in Olsztyn; 1397/98 in Gieczno (Łódź); 1422/23 (24?) in Branice near Krakow; 1414/15(16?), 1426/27, 1440/41 (?), 1450/51, 1459/60, 1470/71, 1480/81, 1491/92 in Wrocław; 1370/71, 1443/44, 1458/59 in Kraków; 1480/81, 1490/91(92?), 1499/1500 in Kutno; 1408/9, 1453/54 in Domachowo (Greater Poland) and 1381/82 in Węgrów (Mazovia).

The time distribution exhibits a few MRs coinciding across a larger territory simultaneously. These events are limited to the 15th century, when they were found in southern Poland (Wrocław, Kraków) in the 1440s and 1450s, and in Wrocław and Kutno in the early 1480s and 1490s.

305

3.2.2. Pointer years

The range of oak chronologies for Greater Poland, Lower Silesia and Lesser Poland covered the entire analysed period of AD 1001–1500. Sequences of tree rings in specific regional chronologies were used to determine pointer years for Lesser Poland, Greater Poland and Lower Silesia (Table S5 and Figs 6 and 7). For the study period, only two pointer years were found to be common to all three regions – one negative (1401) and one positive (1186). In total, for the years 1001–1500, 34 pointer years were observed in Lower Silesia, 38 in Lesser Poland and 39 in Greater Poland (Figs 6 and 7).

For the pine chronology from Lesser Poland in the study period (1091–1500), 27 pointer years were found (Table S5, Figs 7 and 8), of which, 14 were negative pointer years.

315 This was fewer than in Lesser Poland's fir chronology (1109–1500), where 47 pointer years were found (Table S5, Figs 7 and 8). The distribution of pointer years is quite uniform between the centuries, except for the early 12th century and the exceptional 13th century, when as many as 21 pointer years were found.

In the case of pine from Kuyavia-Pomerania for the years 1168–1500, 25 pointer years were identified, of which 21 are negative (Table S5, Figs 7 and 8).

320

3.2.3. Regression model

Temperature reconstructions for the winter months based on the three constructed chronologies are presented in Fig. 9 and in a more generalised form in Fig. 10A–C. For the years AD 1091–1500, the average temperature of February–March (Fig. 9b) was reconstructed using the residual pine chronology, whereas for 1109–1500 the average temperature of December–March 325 was reconstructed (Fig. 9c) using the residual fir chronology as a predictor. In both reconstructions, the temperatures in the study period were lower than those recorded today (1951–2000), whether comparing the study period to the anomaly reconstruction based on the pine or that based on the fir chronology. However, in the reconstruction of average February–March temperatures using the chronology for 1168–1500 (Fig. 9a), warmer periods were observed in the 13th and 14th centuries, followed by a slight cooling.

330 4. Summary and discussion

To improve our knowledge concerning climate change in Poland from 1001–1500, we constructed three new late-winter–early-spring temperature reconstructions covering the period since the 12th century based on tree-ring widths (Figs 9 and 10A–C). In addition, winter and summer temperatures for the 15th century were reconstructed based on documentary evidence (Fig. 5). Unfortunately, the scarcity of historical sources (see Fig. 2) for the area of present-day Poland does not 335 allow such reconstructions for the earlier centuries. For the complete synthesis of Poland's climate in the study period and for comparison purposes, we used all available reconstructions in the literature of cold season temperatures based on tree-rings widths (mainly *P. sylvestris*) covering medieval times: Przybylak et al. (2001), Jan–Apr, 1170–1994, Szychowska-Krąpiec (2010), Dec–Mar 1091–2006 and Dec–Mar 1109–2004 [*A. alba*], Koprowski et al. (2012), Feb–Mar 1168–2000, and Balanzategui et al. (2017), Nov–Apr 1200–2010). See also the recently published inventory of all available 340 dendrochronologies for Poland (Opała-Owczarek et al., 2021 Table 5.2). Information about climate conditions occurring in Poland in the 11th century is available only from analysis of biological proxies taken from laminated (varved) sediments in Lake Żabińskie (NE Poland). Quantitative reconstructions of the cold season (Hernández-Almeida et al., 2015) and August temperatures (Hernández-Almeida et al., 2017) are available for the entire millennium (Fig. 10F,G).

345 The 11th century in Poland was the coldest in the entire study period both in light of chrysophyte-based reconstruction of a number of days below 4 °C (DB4 °C) representing the severity of winter and chironomid-based reconstruction of August temperatures (Fig. 10F,G). Particularly cold was the first half of the 11th century. This century was also colder in comparison to the contemporary period (1951–2000). For the 12th century, we have also data from old (Szychowska-Krapiec, 2010, Fig. 56) and new (see Figs 9 and 10B,C) reconstructions of air temperature from Lesser Poland
350 based on fir and pine chronologies. The 12th century was markedly warmer than the 11th century and also warmer than at present. In particular, the summer seasons in the first half of this century were very warm (August temperature about 18 °C, Fig. 10G), while in the second half they were close to the 1951–2000 norm (16.9 °C). As results from Fig. 10F,G, the 12th century was the warmest century in the entire study period in terms of summer temperature, while the cold-half year temperature was slightly colder (except the last 2–3 decades) than in the next three centuries. The reconstructions of winter
355 temperature for Lesser Poland based on dendrochronological data (Fig. 10B,C) is characterised by strong fluctuations oscillating from about 0 °C to -2 °C relative to present means. But on average, winter conditions were warmer than in the 13th, 14th and 15th centuries.

Proxies represented the entire cold season (see Fig. 10 D,F), revealing the existence of warm temperatures in the 13th century that were particularly high in its first half. The Scots pine reconstructions of late winter and early spring (Feb–
360 Mar) both for Lesser Poland and the Kuyavian-Pomeranian region (Figs 9 and 10A–B,D) generally also confirm this finding. The average temperature of this season was only slightly colder (by 0.2 °C) than at present. Summer temperatures in the 13th century oscillated below and above the long-term contemporary norm. Similarly to winter temperature, summers were also on average slightly colder than at present by the same value. However, this century was significantly colder (by 0.3–0.6 °C) than the 12th, 14th and 15th centuries.

365 Good agreement exists for the cold season temperature dendrochronological reconstructions for Poland in the 14th century. They indicate the existence of colder conditions in the 1330s and the last three decades of the century and warmer conditions in the middle of the century (see Fig. 10). On the other hand, values of the reconstructed number of DB4 °C are slightly lower than the norm for the period 1001–2000 (100.3 days), and significantly lower than the present norm (102.7). That is, winters were mild and thermally stable in this time (see Fig. 10F). Summer temperature was relatively high in
370 almost the entire century and very high in the last decade, when it was comparable to that in the first half of the 12th century (see Fig. 10G). On average, the temperature was only slightly lower than in the 12th century (by 0.1 °C), but it was higher than **in the latter half of the 20th century at present** (by 0.4 °C).

For the 15th century, we have many excerpts describing the weather. Based on these sources, winter and summer temperatures were reconstructed (see Fig. 5). They document the existence in that century of very cold winters and very
375 warm summers, which means that climate continentality was very high. Sadowski (1991) estimated that Poland's climate continentality greater in the 15th century than at any other time in the period 1201–1980 that he had analysed. The chironomid-based reconstruction of August temperatures (Fig. 10G) shows a similar time pattern as the summer temperature changes presented in Fig. 5. In both reconstructions, higher temperatures were observed in the second half of the century,

and particularly in the 1470s. In the case of winter temperatures, all dendrochronological reconstructions generally confirm
380 that they were usually lower than at present (Fig. 10), although large oscillations are seen. Some contradictions, however, are
noted regarding the occurrence of the two coldest decades in the reconstruction based on documentary evidence (the 1430s
and 1450s). The first cold decade was noted in Lesser Poland, but not in the northern and central Poland. In the latter areas,
very warm winters occurred in this time. Similar results are shown by the reconstruction of the number of DB4 °C. At this
time, they were fewest (only 96 days) of the entire study period (Fig. 10F) and even the entire millennium (Hernández-
385 Almeida et al., 2015). On the other hand, there is correspondence of the thermal character of winter between the
reconstruction based on documentary evidence and the number of DB4 °C for the second half of the 15th century. The
coldest winters (in the 1450s) were significantly better registered in trees growing in northern and central Poland, while in
southern Poland a warming was even noted (Fig. 10). The decrease in winter temperature in Poland in the 1430s and 1450s
was confirmed by the reconstruction of the ice winter index in the Western Baltic Sea for the 15th century (Kosłowski and
390 Schmelzer, 2007).

We observed that, for Scots pine from Lesser Poland and northern Poland, negative pointer years outnumbered
positive pointer years. This was probably caused by the influence of unfavourable climatic conditions prevailing in the study
period, and long and severe winters. For the period 1361–1500, 38 pointer years were found for which information about
weather conditions was available in the historical sources. For as many as 71% of pointer years, descriptions of extreme
395 weather conditions were found that favoured the annual tree growth being either low (e.g., severe and long winter, drought)
or high (e.g., warm winter, heavy rainfall).

The interpretation of climate based on pointer years in oaks is more difficult because they depend simultaneously on
temperature and precipitation. Krąpiec (1998) found that positive pointer years for oaks in Poland are associated mainly with
heavy rainfall in the growing season following a mild winter, while negative ones are linked to dry years with cold and
400 snowless winters, and to spring frosts damaging the cambium. For example, the negative pointer years of AD 1314 and 1317
in Lesser Poland were caused by severe cold weather that caused a famine in Bohemia and neighbouring areas (Brázdil and
Kotyza, 1997). The analysis of moon rings (MR) of oaks is easier to interpret, allowing long and severe winters to be
determined. In the period 1361–1500, 19 such winters were recorded, and, for almost half (47.4%), the presence of moon
rings was found.

405 As results from the above summary for Poland in medieval times, most of the proxies allow only for the
temperature reconstruction of the entire cold half-year (Nov–Apr) or some of its sub-periods. Information about the weather
in summer for the whole of the study period is limited to only one reconstruction of August temperature based on
chironomid assembles (Fig. 10G). For late medieval times, some information is also available from documentary evidence.
As a result, it is difficult to compare our results presented in the paper with those from the Northern Hemisphere and the
410 European territory since proxy series of data (tree rings in particular) for the mentioned areas are biased towards the warmer
seasons of the year (e.g., Table 1 in Ljungqvist, 2010). Fortunately, some comparisons are possible using existing

reconstructions for European regions based on documentary evidence. However, due to the scarcity of available historical sources, the comparison should be limited mainly to late medieval times.

415 Comparison of August temperature from NE Poland (Fig. 10G) against summer temperature (Jun–Aug)
reconstructed for Europe by Luterbacher et al. (2016) shows quite good correspondence. In both reconstructions, the coldest
temperatures in the period 1001–1500 occurred in the 11 century and the mid-15th century. Cold summers were also noted
from the mid-13th century to the mid-14th century. On the other hand, in both reconstructions, the markedly warmest
summers occurred in the 12th century. However, in Poland they ended earlier than in Europe (in the middle part of this
century) but started earlier, at the end of the 11th century. Smaller summer warmings in both areas were also noted at the turn
420 of the 15th century and in the second part of the 15th century.

In Europe, however, the scale of warming in these times was significantly smaller than in the 12th century, while in
Poland it was comparable across the periods, but the duration of the warming was shorter in both areas. The most
comparison of the results is most reliable for the 15th century, for which we reconstructed summer temperature (Fig. 5). The
second half of the century was clearly warmer than the first, not only in Poland but also in Czech Lands (Brázdil, 1994,
425 1996), Switzerland and in the entirety of Central Europe (Camenisch et al., 2016), as well as in Scandinavia (Gouirand et al.,
2008). The decade 1471–1480 was the warmest in the entire 15th century in Poland (Fig. 5) and also in the Low Countries
and Central Europe (Camenisch, 2015; Riemann et al., 2015; Camenisch et al., 2016), while the warmest summers occurred
in the 1480s in Czech Lands (Brázdil, 1994, 1996) and in the 1490s in Switzerland (Trachsel et al., 2010; Camenisch et al.,
2016). For the 15th century, there exists a correspondence between the run of the August temperature in Poland (Fig. 10G)
430 and reconstructions of summer temperature for some European areas based on documentary evidence and tree rings (Fig. 2;
see also reconstructions 2–5 in Camenisch et al., 2016). Summer temperature was colder in the first part of the century than
in the second.

Luterbacher et al. (2010) found that the correspondence of temperatures between Poland and Europe is lower for
summer than for winter. In the latter season, there exists a very high correlation (0.96, at interannual and multidecadal time
435 scales) between reconstructed Polish and (non-Polish) European mean temperatures (e.g., excluding the grid points
representing Poland) over the period 1500–2000. However, for Europe there are only some reconstructions based on
documentary evidence (e.g., Brázdil, 1994, 1996; Pfister et al., 1996, 1998; van Engelen et al., 2001; Shabalova and Van
Engelen, 2003; Glaser and Riemann, 2009; Camenisch, 2015; Camenisch et al., 2016; Retsö and Söderberg, 2019) and
modeling works (e.g., Goosse et al., 2006; Schimanke et al., 2012). The scarcity of historical sources for the study period,
440 particularly before the 15th century (e.g., Brázdil, 1996; Glaser and Riemann, 2009; Oliński, 2022; see also Fig. 2), means
that the reconstruction uncertainties can be significant. ~~For this reason, the history of the winter climate in Poland shows
better coherence with modeling reconstructions. But~~ Nevertheless, a correlation exists for some periods between Poland's
winter temperature and reconstructions for European areas based on documentary evidence.

For example, such agreement was observed in the 11th century, where also in western-central Europe, winters were
445 lower than average (Pfister et al., 1998). Similar results were also found for Europe north of the Alps (Alexandre, 1987 after

Brázdil 1996) and Czech Lands (Brázdil, 1996), while the opposite was found in the Low Countries (van Engelen et al., 2001). In the 12th century, there is still good agreement with the reconstruction of the DB4, but less agreement with the reconstructions of Feb–Mar temperatures (compare Fig. 10 in Pfister et al. (1998) and Fig. 9 here). On the other hand, all reconstructions reveal a significant winter warming at the end of the 12th century that has also been noted in Germany (Glaser and Riemann, 2009) and the Low Countries (van Engelen et al., 2001). In the 13th century, changes in the reconstructed temperature in Poland and west-central Europe differed much, particularly in the first half of the century. However, the warming of winters in this time occurred also in England, Ukraine and Russia, but not in Europe north of the Alps, Czech Lands (Fig. 4 in Brázdil, 1996), Germany (Glaser and Riemann, 2009) or the Baltic Sea Basin (Schimanke et al., 2012). On the other hand, a good coherence was usually stated for the 14th century, where in the two compared areas, colder conditions prevailed except in the middle of the century (compare Fig. 2 in Pfister et al. (1996)) with available reconstructions from Poland, including new ones presented in Fig. 9). Also agreement is seen for Europe north of Alps, but not for Czech Lands, Ukraine or Russia (Brázdil, 1996) and the Low Countries (van Engelen et al., 2001). The reconstructed winter (DJF) temperature in Poland for the 15th century based on documentary evidence (Fig. 5) matches well with the data presented for Czech Lands, Russia, England (Brázdil, 1994, 1996), Baltic Sea Basin (Schimanke et al., 2012), Sweden (Retsö and Söderberg, 2019), the Low Countries (Camenisch, 2015), Alps (Mangini et al., 2005), and the whole of Europe (Goosse et al., 2006), but only partly with data for Germany (Glaser and Riemann, 2009).

Ljungqvist (2010) found that the Northern Hemisphere proxy collection is biased toward the year's warmer seasons. The reason for this is that predominant among the proxies in the reconstruction process are tree rings taken mainly from mountain regions (see Pages2k Consortium 2013, 2017; Tardiff et al. 2019). Nevertheless, the reconstructed summer temperatures are frequently used to represent annual values (e.g., Cook et al., 2004; Moberg et al., 2005; Ljungqvist, 2010) due to the high correlation between seasons on decadal and longer time scales within the instrumental period. However, for smaller areas that are dominated by lowlands – like Poland – this is not the case. There is lack of correlation between mean summer and winter temperatures for Poland in 1901–2000. The correlation of summer temperature with the annual temperature is 0.4 and is two times smaller than the correlation of winter temperature (0.79). Therefore in Poland, winter temperature represents the annual temperature better than does summer temperature. It is thus justified to use winter temperatures as annual proxies and thus to use them to distinguish the occurrence or not of the MWP (MCA) in Poland.

Due to the coverage of the entire millennium and also of the whole cold season, the best proxy that can presently be used to delimit the MWP in Poland is DB4 (Fig. 10F). These data locate this period in the time period 1180–1440 (excluding the period 1260–1270, partly a consequence of a volcanic eruption in 1268), when the winters were generally shorter and warmer than today (1951–2000). The other proxies usually do not allow the beginning of the MWP to be distinguished (due to the reconstructed data series starting too late) but only its end. There exist, however, some essential differences between proxy results. The MWP is also suggested to have ended in the first half of the 15th century (e.g., in line with DB4 data) by Sadowski (1991). Some other authors concluded that the MWP in Poland ended at the beginning of the 14th century (Maruszczak, 1991; Kotarba, 2004; Balanzategui et al., 2017; Szychowska-Krapiec, 2010 only fir chronology) or even the

480 beginning of the 13th century (Koprowski et al., 2012). Based on new February–March reconstructions (Fig. 9), it is possible to distinguish the MWP (from the end of the 13th century to the mid-15th century) only in series representing the Kuyavia-Pomerania region. These are thus in correlation with DB4 data, except the first 100 years. All the mentioned quantitative reconstructions suggest that the MWP in Poland was comparable to or warmer than the current temperature (1951–2000). For Europe, various time frames are given for the MWP, but most suggest that this period ended in the 14th century (e.g., 485 Brázdil, 1996 for Czech Lands; Glaser and Riemann, 2009 for Germany; Millet et al., 2009 for northern French Alps; Niemann et al., 2012 for Swiss Alps; Niedźwiedz et al., 2015 for the Baltic Sea Basin). The simulation of mean winter European temperatures presented by Goose et al. (2006, their Fig. 3) locates the MWP as extending until the first half of the 15th century; however, when we take mean summer temperature, the MWP ended in the beginning of the 13th century. The lack of spatial coherence in timing of the MWP across the entire globe was also recently confirmed by Neukom et al. (2019) 490 based on proxy data published by Pages2k Consortium (2017). They locate the MWP from 751 to 1350, and according to their Fig. 3, within this time span the peak warming in Poland occurred between 800 and 1000. However, we need to add that the Pages2k databases contain not a single proxy data series taken from Poland.

The warm/cold periods observed in Poland during the study period are an effect of natural variability in the climatic system (see the course of the set of forcings presented in Fig. 1 in Goose et al. 2006). For example, the low temperatures in 495 the 11th century can be connected with negative anomalies of total solar irradiance (TSI), while in the 15th century they can also be associated with increased volcanic activity and the negative phase of the North Atlantic Oscillation (NAO) (Trouet et al., 2009). For example, the very cold decade of the 1450s observed both in Europe (Goose et al., 2006) and globally (e.g., Tardif et al., 2019) is most often explained by the Kuwae eruption of 1452 or 1453 (Gao et al., 2006). According to those authors “the Kuwae eruption was the largest stratospheric sulfate event of that period [since 1300, authors], probably 500 surpassing the total sulfate deposition of the Tambora eruption of 1815, which produced 59 kg SO₄/km² in Antarctica and 50 kg SO₄/km² in Greenland.” The eruption probably changed atmospheric circulation in the North Atlantic leading to the occurrence of the negative phase of the NAO (see Fig. 2 in Trouet et al. 2009), which is important in shaping the climate in Poland, as we show in the next paragraph.

Any extreme thermal condition in Poland in a season or entire year is most often caused by anomalous behaviour of 505 atmospheric circulation, and only infrequently by volcanic activity. According to investigations made by Przybylak et al. (2003) for the period 1500–1990, the main cause of cold winters in Poland in this time was the negative phase of the NAO (in which advection of cold air from the north and east dominates), while warm winters were observable when the NAO was in the positive phase (advection from the west dominates). So too, the occurrence of summers that are extremely warm or very warm in Poland is also connected with changes in atmospheric circulations. However, cold summers are caused by 510 advection from the west and north, and warm summers with southerly and easterly advection (Bartoszek, 2017, see his Table 3). It is reasonable to assume that the relationships observed in the contemporary summer climate of Poland also obtained in the study period.

5. Conclusions and final remarks

The main results of the present paper can be summarised as follows:

- 515 1) Since the 1990s, significantly more quantitative reconstructions of climate in Poland have been published using different kind of proxies covering the study period, including four new ones presented in the paper.
- 2) The areally averaged summer temperature for Poland in 1901–2000 correlates with the annual temperature at $r=0.4$, which is smaller than the correlation with winter temperature ($r=0.79$). Therefore in Poland, winter temperature better represents the annual temperature than does summer temperature. It is justified to use winter temperature as annual
520 proxies and thus to use them to distinguish the occurrence or not of the MWP (MCA) in Poland.
- 3) Analysis of all available reconstructions reveals the existence of the MWP in Poland from the late 12th century to the first halves of the 14th or 15th centuries. We found that the MWP in Poland was comparable to or warmer than the current temperature (1951–2000).
- 4) The coldest conditions in the entire study period were noted in the first half of the 11th century (both winter and
525 summer) and the second half of the 15th century (only winter).
- 5) The best and most reliable knowledge about climate in the study period exists for the 15th century, for which we present winter and summer reconstructions based on documentary evidence (Fig. 5). All reconstructed 10-year mean values of winters and summers were, correspondingly, colder and warmer than today. The coldest winters occurred in the 1450s, 1430s and 1490s, while the warmest summers occurred mainly in the 1470s. Thus our reconstruction confirms the
530 conclusion formulated earlier by Sadowski (1991) that the 15th century in Poland was characterised by the highest climate continentality in the entire millennium.
- 6) For the study period, information about wetness of seasons in Poland is available only based on documentary evidence, and mainly for the 15th century. In line with expectations, the most such information is available for summer and then for winter, while it is only very rarely available for the transitional seasons (Fig. 4). In the 15th century, all categories of wet winters dominated, with a maximum in 1430s (4). This means that the 1430s were characterised in Poland by very cold and snowy winters. Except for the last three decades of the 15th century, extremely wet and very wet conditions also dominated in summer, and especially in the 1460s (5 cases). Radical dryness of the air occurred in the 1470s, when five extremely dry or very dry summers were noted (Fig. 4).
- 7) Good agreement was found between the reconstructions of Poland's climate from 1001–1500 and many reconstructions
540 available for Europe, which is in line with findings presented by Luterbacher et al. (2010) for the period 1500–2000.

The further improvement of the knowledge of the climate in Poland in the first half of the last millennium presented here will be closely connected with findings based on new proxies from the natural archives – mainly tree-ring wood density and stable isotopes and other biological proxies. Based on the preliminary archival research and the survey of the library holdings, we have conducted in the last 20 years, we are certain that it is impossible to find a significant amount of further
545 information about weather in historical sources, particularly prior to the 15th century.

Author's contribution. Conceptualization: RPr, PO, MKo; Methodology: RPr, PO, MKo, MKr, ESK; Validation: RPr, PO, MKo, MKr, ESK; Formal analysis: RPr, PO, MKo, MKr; Investigation: RPr, PO, MKo, MKr, ESK, RPu, AP; Resources: RPr, PO, MKo, MKr, ESK, AP; Data Curation: RPr, PO, MKo, MKr, ESK, AP; Writing - Original Draft: RPr, PO, MKo, MKr, ESK; Writing - Review & Editing: RPr, PO, MKo, MKr, ESK, RPu, AP; Visualization: RPr, PO, MKo, MKr, RPu; Project administration: RPr; Funding acquisition: RPr, ESK and MKr.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Financial support. The research work of PO and RP was supported by grants funded by the National Science Centre, Poland (Grants Nos 2013/11/b/HS3/01458, 2020/37/B/ST10/00710). MK and RPu were supported by a grant funded by the National Science Centre, Poland (Grant No 2020/37/B/ST10/00710). ESK and MKr were supported by the grant of the Faculty of Geology, Geophysics and Environmental Protection of AGH University of Science and Technology in Cracow no. 16.16.140.315.

Data Availability. The tree-ring chronologies supporting the conclusion of this study are available from ESK, MK and MKr upon request.

References

- Alexandre, P.: Le climat en Europe au Moyen Age. Contribution à l'histoire des variations climatiques de 1000 à 1425, d'après les sources narratives de l'Europe occidentale, Ecole des hautes études en sciences sociales, Paris, 827 pp., 1987.
- Balanzategui, D., Knorr, A., Heussner, K.-U., Wazny, T., Beck, W., Słowiński, M., Helle, G., Buras, A., Wilmking, M., Van Der Maaten, E., Scharnweber, T., Dorado-Liñán, I., and Heinrich, I.: An 810-year history of cold season temperature variability for northern Poland, *Boreas*, 47, 443–453, <https://doi.org/10.1111/bor.12274>, 2017.
- Bartoszek K.: *Relacje między cyrkulacją atmosferyczną a sezonową temperaturą powietrza na obszarze Lubelszczyzny, Annales Universitatis Mariae Curie-Skłodowska*, LXXII, 1, 33–45, <https://journals.umcs.pl/b/article/view/4912>, 2017.
- Bolychevtsev, V. G.: Godnichnyje sloi u duba kak pokazatel' vekovykh ciklov kolebanij klimata (Annual ring of oak as evidence of secular climatic cycles), *Lesovedenie*, 1, 15–23, 1970.
- Brázdil, R.: Climatic fluctuations in the Czech lands during the Last Millennium, *GeoJournal*, 32, 199–205, 1994.

- Brázdil, R.: Reconstructions of past climate from historical sources in the Czech lands, in: Climatic variations and forcing mechanisms of the last 2000 years, vol. 41, edited by: Jones, P. D., Bradley, R. S., and Jouzel, J., Springer-Verlag, Berlin, 409–431, 1996.
- 580 Brázdil, R. and Kotyza, O.: Kolísání klimatu v Jeských zemích první polovině našeho tisíciletí, *Archeol. Rozhl.*, 49, 663–699, 1997.
- Brázdil, R., Pfister, C., Wanner, H., von Storch, H., and Luterbacher, J.: Historical climatology in Europe - the state of the art, *Clim. Change*, 70, 363–430, <https://doi.org/10.1007/s10584-005-5924-1>, 2005.
- Briffa, K. R.: Interpreting high-resolution proxy climate data - the example of dendroclimatology, in: *Analysis of Climate Variability*, edited by: von Storch, H. and Navarra, A., Springer, Berlin and Heidelberg, 77–94, https://doi.org/10.1007/978-3-662-03167-4_5, 1995.
- 585 Brönnimann, S.: From climate to weather reconstructions, *PLOS Clim.*, 1, e0000034, <https://doi.org/10.1371/journal.pclm.0000034>, 2022.
- Brönnimann, S., Allan, R., Ashcroft, L., Baer, S., Barriendos, M., Brázdil, R., Brugnara, Y., Brunet, M., Brunetti, M., Chimani, B., Cornes, R., Dominguez-Castro, F., Filipiak, J., Founda, D., Herrera, R., Gergis, J., Grab, S., Hannak, L., Huhtamaa, H., Jacobsen, K., Jones, P., Jourdain, S., Kiss, A., Lin, K., Lorrey, A., Lundstad, E., Luterbacher, J., Mauelshagen, F., Maugeri, M., Maughan, N., Moberg, A., Neukom, R., Nicholson, S., Noone, S., Nordli, Ø., Ólafsdóttir, K., Pearce, P., Pfister, L., Pribyl, K., Przybylak, R., Pudmenzky, C., Rasol, D., Reichenbach, D., Rezníčková, L., Rodrigo, F., Rohr, C., Skrynyk, O., Slonosky, V., Thorne, P., Valente, M., Vaquero, J., Westcott, N., Williamson, F., and Wyszyński, P.:
- 590 Unlocking pre-1850 instrumental meteorological records: A global inventory, *Bull. Am. Meteorol. Soc.*, 100, ES389–ES413, <https://doi.org/10.1175/BAMS-D-19-0040.1>, 2019.
- Bunn, A. G.: A dendrochronology program library in R (dplR), *Dendrochronologia*, 26, 115–124, <https://doi.org/10.1016/j.dendro.2008.01.002>, 2008.
- Camenisch, C.: Endless cold: a seasonal reconstruction of temperature and precipitation in the Burgundian Low Countries during the 15th century based on documentary evidence, *Clim. Past*, 11, 1049–1066, <https://doi.org/10.5194/cp-11-1049-2015>, 2015.
- 600 Camenisch, C., Keller, K., Salvisberg, M., Amann, B., Bauch, M., Blumer, S., Brázdil, R., Brönnimann, S., Buntgen, U., Campbell, B., Fernández-Donado, L., Fleitmann, D., Glaser, R., González-Rouco, F., Grosjean, M., Hoffmann, R., Huhtamaa, H., Joos, F., Kiss, A., Kotyza, O., Lehner, F., Luterbacher, J., Maughan, N., Neukom, R., Novy, T., Pribyl, K., Raible, C., Riemann, D., Schuh, M., Slavin, P., Werner, J., and Wetter, O.: The 1430s: a cold period of extraordinary internal climate variability during the early Spörer Minimum with social and economic impacts in north-western and central Europe, *Clim. Past*, 12, 2107–2126, 2016.
- Cook, E. and Kairiukstis, L.: *Methods of dendrochronology. Applications in the environmental sciences*, Springer, Dordrecht, 351 pp., 1990.
- 610 Cook, E., Esper, J., and D'Arrigo, R. D.: Extra-tropical Northern Hemisphere land temperature variability over the past 1000

- years, *Quat. Sci. Rev.*, 23, 2063–2074, <https://doi.org/10.1016/j.quascirev.2004.08.007>, 2004.
- Cook, E., Krusic, P., and Melvin, T. M.: Program RCSigFree. Lamont-Doherty Earth Observatory, Columbia University, Palisades, 2014.
- Cook, E. R.: *A Time Series Analysis Approach to Tree-ring Standardization*, The University of Arizona, Tucson, 170 pp., 1985.
- 615 Diaz, H., Trigo, R., Hughes, M., Mann, M., Xoplaki, E., and Barriopedro, D.: Spatial and temporal characteristics of climate in medieval times revisited, *Bull. Am. Meteorol. Soc.*, 92, 1487–1500, <https://doi.org/10.1175/BAMS-D-10-05003.1>, 2011.
- Dujesiefken, D. and Bauch, J.: Biologische Charakterisierung von Eichenholz mit Mondringen, *Holz als Roh-und Werkst.*, 45, 365–370, 1987.
- 620 Dujesiefken, D. and Liese, W.: Vorkommen und Entstehung der mondringe (*Quercus* spp.), *Forstwissenschaftliches Cent.*, 105, 137–155, 1986.
- Dzbeński, W. and Krutul, D.: Fizyko-chemiczne właściwości drewna dębowego z wewnętrznym bielem, in: *Materiały XVII Sympozjum “Ochrona drewna,”* 127–134, 1994.
- van Engelen, A. F. V., Buisman, J., and Ijnsen, F.: A millennium of weather, winds and water in the Low Countries, in: *History and Climate*, edited by: Jones, P. D. and Ogilvie, A., Kluwer Academic/Plenum Publishers, 101–124, 2001.
- 625 Frank, D., Esper, J., and Cook, E.: Adjustment for proxy number and coherence in a large-scale temperature reconstruction, *Geophys. Res. Lett.*, 34, L16709, <https://doi.org/10.1029/2007GL030571>, 2007.
- Fritts, H. C.: *Tree-ring and climate*, Academic Press, London, New York, San Francisco, 567 pp., 1976.
- Gao, C., Robock, A. Self, S., Witter, J. B. , Steffenson, J. P., Clausen, H. B., Siggaard-Andersen, M.-L., Johnsen, S.,
- 630 Mayewski, P. A., and Ammann, C.: The 1452 or 1453 A.D. Kuwae eruption signal derived from multiple ice core records: Greatest volcanic sulfate event of the past 700 years, *J. Geophys. Res.*, 111, D12107, [doi:10.1029/2005JD006710](https://doi.org/10.1029/2005JD006710), 2006
- Ghazi, B., Przybylak, R., Oliński, P., Bogdańska, K., and Pospieszńska, A.: The frequency, intensity, and origin of floods in Poland in the 11th–15th centuries based on documentary evidence, *J. Hydrol.*, 623, 2023 <https://doi.org/10.1016/j.jhydrol.2023.129778>.
- 635 Girguś, R. and Strupczewski, W.: Wyjątki ze źródeł historycznych o nadzwyczajnych zjawiskach hydro-logiczno-meteorologicznych na ziemiach polskich w wiekach od X do XVI, *Wydawnictwa Komunikacji i Łączności*, Warszawa, Warszawa, 1965.
- Glaser, R. and Riemann, D.: A thousand-year record of temperature variations for Germany and Central Europe based on documentary data, *J. Quat. Sci.*, 24, 437–449, <https://doi.org/https://doi.org/10.1002/jqs.1302>, 2009.
- 640 Goosse, H., Arzell, O., Luterbacher, J., Mann, M. E., Renssen, H., Riedwyl, N., Timmermann, A., Xoplaki, E., and Wanner, H.: The origin of the European “Medieval Warm Period,” *Clim. Past*, 2, 99–113, 2006.
- Goosse, H., Guiot, J., Mann, M. E., Dubinkina, S., and Sallaz-Damaz, Y.: The medieval climate anomaly in Europe: Comparison of the summer and annual mean signals in two reconstructions and in simulations with data assimilation, *Glob. Planet. Change*, 84–85, 35–47, <https://doi.org/10.1016/j.gloplacha.2011.07.002>, 2012.

- 645 Gouirand, I., Linderholm, H., Moberg, A., and Wohlfarth, B.: On the spatiotemporal characteristics of Fennoscandian tree-ring based summer temperature reconstructions, *Theor. Appl. Climatol.*, 91, 1–25, <https://doi.org/https://doi.org/10.1007/s00704-007-0311-7>, 2008.
- Hernández-Almeida, I., Grosjean, M., Przybylak, R., and Tylmann, W.: A chrysophyte-based quantitative reconstruction of winter severity from varved lake sediments in NE Poland during the past millennium and its relationship to natural climate variability, *Quat. Sci. Rev.*, 122, 74–88, <https://doi.org/10.1016/j.quascirev.2015.05.029>, 2015.
- 650 Hernández-Almeida, I., Grosjean, M., Gómez-Navarro, J. J., Larocque-Tobler, I., Bonk, A., Enters, D., Ustrzycka, A., Piotrowska, N., Przybylak, R., Wacnik, A., Witak, M., and Tylmann, W.: Resilience, rapid transitions and regime shifts: Fingerprinting the responses of Lake Żabińskie (NE Poland) to climate variability and human disturbance since AD 1000, *The Holocene*, 27, 258–270, <https://doi.org/10.1177/0959683616658529>, 2017.
- 655 Holmes, R. L.: Computer-assisted quality control in tree-ring dating and measurement, *Tree-Ring Bull.*, 43, 69–78, 1983.
- Huber, B. and Giertz-Siebenlist, V.: Unsere tausendjährige Eichen-Jahrringchronologie durchschnittlich 57 (10-150) – fach belegt, *Sitzungsberichte Österreichische Akad. der Wissenschaften, Math. Klasse*, 57, 37–42, 1969.
- Hughes, M. K. and Diaz, H. F.: Was there a “Medieval Warm Period”, and if so, where and when?, *Clim. Change*, 26, 109–142, 1994.
- 660 Koprowski, M., Przybylak, R., Zielski, A., and Pospieszynska, A.: Tree rings of Scots pine (*Pinus sylvestris* L.) as a source of information about past climate in northern Poland., *Int. J. Biometeorol.*, 56, 1–10, <https://doi.org/10.1007/s00484-010-0390-5>, 2012.
- Kosłowski, G. and Schmelzer, N.: Ice winter severity in the Western Baltic Sea in the period 1301-1500, *Berichte des Bundesamtes für Seeschifffahrt und Hydrogr.*, 42, 47–56, 2007.
- 665 Kotarba, A.: Geomorphological events in the High Tatra Mountains during the Little Ice Age, in: *Rola Mafej Epoki Lodowej w przekształcaniu środowiska przyrodniczego Tatry*, edited by: Kotarba, A., 9–55, 2004.
- Kożuchowski, K. and Żmudzka, E.: 100-year series of areally averaged temperatures and precipitation totals in Poland, *Acta Univ. Wratislav. Stud. Geogr.*, 75, 116–122, 2003.
- Krapiec, M.: Oak Dendrochronology of the Neoholocene in Poland, *Folia Quartheria*, 69, 5–133, 1998.
- 670 Krapiec, M.: Occurrence of Moon Rings in Oak from Poland During the Holocene, in: *Tree Ring Analysis, Biological, Methodological and Environmental Aspects*, edited by: Wimmer, R. and Vetter, R., CAB Int., 193–203, 1999.
- Lamb, H.: The early medieval warm epoch and its sequel, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 1, 13–37, 1965.
- Lamb, H.: *Climate: Present, Past and Future. vol. 2: Climatic History and the Future*, Methuen, Methuen, London, 835 pp., 1977.
- 675 Lamb, H.: Climate in the last thousand years: Natural climatic fluctuations and change, in: *The Climate of Europe: Past, Present and Future*, edited by: Flohn, H. and Fantechi, R., D. Reidel Publishing Company, Dordrecht, Boston, Lancaster, 25–64, 1984.
- Ljungqvist, F. C.: A new reconstruction of temperature variability in the extra-tropical Northern Hemisphere during the last

- two millennia, *Geogr. Ann. Ser. A, Phys. Geogr.*, 92, 339–351, 2010.
- 680 Luterbacher, J., Xoplaki, E., Kilttel, M., Zorita, E., Gonzalez-Rouco, J. F., Jones, P. D., Stossel, M., Rutishauser, T., Wanner, H., Wibig, J., and Przybylak, R.: Climate Change in Poland in the Past Centuries and Its Relationship to European Climate: Evidence From Reconstructions and Coupled Climate Models, in: *The Polish Climate in the European Context: An Historical Overview*, edited by: Przybylak, R., Majorowicz, J., Brázdil, R., and Kejna, M., Springer, Berlin Heidelberg New York, 3–39, 2010.
- 685 Luterbacher, J., Werner, J. P., Smerdon, J. E., Fernández-Donado, L., González-Rouco, F. J., Barriopedro, D., Ljungqvist, F. C., Büntgen, U., Zorita, E., Wagner, S., Esper, J., McCarroll, D., Toreti, A., Frank, D., Jungclauss, J. H., Barriendos, M., Bertolin, C., Bothe, O., Brázdil, R., Camuffo, D., Dobrovolný, P., Gagen, M., García-Bustamante, E., Ge, Q., Gómez-Navarro, J. J., Guiot, J., Hao, Z., Hegerl, G. C., Holmgren, K., Klimenko, V. V., Martín-Chivelet, J., Pfister, C., Roberts, N., Schindler, A., Schurer, A., Solomina, O., von Gunten, L., Wahl, E., Wanner, H., Wetter, O., Xoplaki, E., Yuan, N.,
- 690 Zanchettin, D., Zhang, H., and Zerefos, C.: European summer temperatures since Roman times, *Environ. Res. Lett.*, 11, 24001, <https://doi.org/10.1088/1748-9326/11/2/024001>, 2016.
- Mangini, A., Spötl, C., and Verdes, P.: Reconstruction of temperature in the Central Alps during the past 2000 yr from a d18O stalagmite record, *Earth Planet. Sci. Lett.*, 235, 741–751, 2005.
- Mann, M., Zhang, Z., Rutherford, S., Bradley, R. S., Hughes, M. K., Shindell, D., Ammann, C., Faluvegi, G., and Ni, F.:
- 695 Global signatures and dynamical origins of the Little Ice Age and medieval climate anomaly, *Science (80-.)*, 326, 1256–1260, 2009.
- Maruszczak, H.: Zmiany środowiska przyrodniczego kraju w czasach historycznych, in: *Przemiany środowiska geograficznego*, edited by: Starkel, L., Wszechnica PAN, Ossolineum, Wrocław, 109–135, 1988.
- Maruszczak, H.: Tendencje do zmian klimatu w ostatnim tysiącleciu, in: *Geografia Polski - środowisko przyrodnicze*, edited
- 700 by: Starkel, L., PWN, Warszawa, 182–190, 1991.
- Melvin, T. and Briffa, K.: A “signal-free” approach to dendroclimatic standardisation, *Dendrochronologia*, 26, 71–86, 2008.
- Millet, L., Arnaud, F., Heiri, O., Magny, M., Verneaux, V., and Desmet, M.: Late-Holocene summer temperature reconstruction from chironomid assemblages of Lake Anterne, northern French Alps, *The Holocene*, 19, 317–328, 2009.
- Moberg, A., Sonechkin, D. M., Holmgren, K., Datsenko, N. M., and Karlen, W.: Highly variable Northern Hemisphere
- 705 temperatures reconstructed from low- and high-resolution proxy data, *Nature*, 433, 613–617, 2005.
- Moreno, A., Perez, A., Frigola, J., Nieto-Moreno, V., Rodrigo-Gamiz, M., Martrat, B., Gonzalez-Samperiz, P., Morellon, M., Martin-Puertas, C., Corella, J. P., Belmonte, A., Sancho, C., Cacho, I., Herrera, G., Canals, M., Grimalt, J. O., Jimenez-Espejo, F. J., Martínez-Ruiz, F., Vegas-Vilarrubia, T., and Valero-Garcés, B. L.: The Medieval Climate Anomaly in the Iberian Peninsula reconstructed from marine and lake records, *Quat. Sci. Rev.*, 43, 16–32,
- 710 <https://doi.org/10.1016/j.quascirev.2012.04.007>, 2012.
- Niedźwiedz, T., Glaser, R., Hansson, D., Helama, S., Klimenko, V., Łupikasza, E., Małarzewski, Ł., Nordli, Ø., Przybylak, R., Riemann, D., and Solomina, O.: The historical time frame (past 1000 years), in: *Second assessment of climate change for*

- the Baltic Sea Basin, edited by: BACC_II_Author_Team, Springer, Berlin, Germany, 51–65, 2015.
- 715 Neukom, R., Steiger, N., Gómez-Navarro, J.J., Wang, J., Werner, J.P.: No evidence for globally coherent warm and cold periods over the preindustrial Common Era. *Nature* 571, 550–554. 2019, <https://www.nature.com/articles/s41586-019-1401-2>.
- Niemann, H., Stadnitskaia, A., Wirth, S. B., Gilli, A., Anselmetti, F. S., Damste, J. S. S., Schouten, S., Hopmans, E. C., and Lehmann, M. F.: Bacterial GDGTs in Holocene sediments and catchment soils of a high Alpine lake: application of the MBT/CBT-paleothermometer, *Clim. Past*, 8, 889–906, 2012.
- 720 Oliński, P.: Pogoda i klimat regionów południowobałtyckich od końca XIV wieku do początków XVI w. w źródłach narracyjnych, Wydawnictwo Naukowe Uniwersytetu Mikołaja Kopernika, Toruń, 299 pp., 2022.
- Opala-Owczarek, M., Niedźwiedź, T., Przybylak, R., and Tylmann, W.: Climate Change Before Instrumental Measurements, in: *Climate Change in Poland*, edited by: Falarz, M., Springer, 71–119, https://doi.org/10.1007/978-3-030-70328-8_5, 2021.
- 725 Osborn, T. J., Briffa, K. R., and Jones, P.: Adjusting variance for sample-size in tree-ring chronologies and other regional-mean time series, *Dendrochronologia*, 15, 88–99, 1997.
- PAGES 2k Consortium: Continental-scale temperature variability during the past two millennia, *Nat. Geosci.*, 6, 339–346, <https://doi.org/10.1038/ngeo1834>, 2013.
- PAGES 2k Consortium: A global multiproxy database for temperature reconstructions of the Common Era, *Sci. Data*, 4, 170088, <https://doi.org/10.1038/sdata.2017.88>, 2017.
- 730 Pfister, C., Kington, J., Kleinogel, G., Schule, H., and Siffert, E.: High resolution spatio-temporal reconstructions of past climate from direct meteorological observations and proxy-data, in: *Climatic Trends and Anomalies in Europe 1675-1715*, edited by: Frenzel, B., G. Fischer, Stuttgart, 329–375, 1994.
- Pfister, C., Schwarz-Zanetti, G., and Wegmann, M.: Winter severity in Europe: The fourteenth century, *Clim. Change*, 34, 91–108, 1996.
- 735 Pfister, C., Luterbacher, J., Schwarz-Zanetti, G., and Wegmann, M.: Winter air temperature variations in western Europe during the Early and High Middle Ages (AD 750-1300), *The Holocene*, 8, 535–552, 1998.
- Polackówna, M.: Wahania klimatyczne w Polsce w wiekach średnich (940-1500), *Pr. Geogr.*, 5, 65–126, 1925.
- Pribyl, K.: The study of the climate of medieval England: a review of historical climatology's past achievements and future potential, *Weather*, 69, 116–120, <https://doi.org/10.1002/wea.2317>, 2014.
- 740 Przybylak, R.: Changes in Poland's climate over the last millennium, *Czas. Geogr.*, 82, 23–48, 2011.
- Przybylak, R.: Poland's Climate in the Last Millennium, in: *Oxford Research Encyclopedia, Climate Science*, Oxford University Press USA, 36, <https://doi.org/10.1093/acrefore/9780190228620.013.2>, 2016.
- Przybylak, R. and Pospieszynska, A.: Air temperature in Wrocław (Breslau) in the period 1710-1721 based on measurements made by David von Grebner, *Acta Agrophysica, Rozpr. i Monogr.*, 184, 35–43, 2010.
- 745 Przybylak, R., Majorowicz, J., and Wójcik, G.: Changes of air temperature and atmospheric precipitation in Poland from the 16th to the 20th century, *Pr. i Stud. Geogr.*, 29, 79–92, 2001.

- Przybylak, R., Majorowicz, J., Wójcik, G., Zielski, A., Chorążyczewski, W., Marciniak, K., Nowosad, W., Oliński, P., and Syta, K.: Temperature changes in Poland from the 16th to the 20th centuries, *Int. J. Climatol.*, 25, 773–791, 2005.
- Przybylak, R., Oliński, P., Koprowski, M., Filipiak, J., Pospieszńska, A., Chorążyczewski, W., Puchałka, R., and Dąbrowski, H. P.: Droughts in the area of Poland in recent centuries in the light of multi-proxy data, *Clim. Past*, 16, 627–661, <https://doi.org/10.5194/cp-16-627-2020>, 2020.
- Przybylak, R., Wójcik, G., Marciniak, K.: Wpływ Oscylacji Północnoatlantyckiej i Arktycznej na warunki termiczne chłodej pory roku w Polsce w XVI-XX wiekach, [Eng. The influence of the North Atlantic oscillation and Arctic oscillation on thermal conditions in the cold season in Poland from the 16th to 20th centuries), *Przegląd Geofizyczny XLVIII*, 61–74, 2003 (in Polish).
- RCoreTeam: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>, 2022.
- Retsö, D. and Söderberg, J.: Winter severity in medieval Sweden: The documentary evidence, in: *The Dance of Death in Late Medieval and Renaissance Europe: Environmental Stress, Mortality and Social Response*, edited by: Kiss, A. and Pribyl, K., Routledge, Taylor & Francis Group, 25–45, 2019.
- Riemann, D., Glaser, R., Kahle, M., and Vogt, S.: The CRE tambora.org-new data and tools for collaborative research in climate and environmental history, *Geosci. Data J.*, 2, 63–77, <https://doi.org/10.1002/gdj3.30>, 2015.
- Sadowski, M.: Variability of extreme climatic events in Central Europe since the 13th century, *Zeitschrift für Meteorol.*, 41, 350–356, 1991.
- Schimanke, S., Meier, H. E. M., Kjellström, E., Strandberg, G., and Hordoier, R.: The climate of the Baltic Sea region during the last millennium simulated with a regional climate model, *Clim. Past*, 8, 1419–1433, 2012.
- Schweingruber, F. H.: Event years and pointer years, *Lundqua Rep.*, 34, 288–292, 1992.
- Shabalova, M. V and Van Engelen, A. F. V.: Evaluation of a reconstruction of winter and summer temperatures in the Low Countries, AD 764–1998, *Clim. Chang.*, 58, 219–242, 2003.
- Sieber, R., Slonosky, V. C., Ashcroft, L., and Pudmenzky, C.: Formalizing Trust in Historical Weather Data, *Weather. Clim. Soc.*, 14, 993–1007, <https://doi.org/10.1175/WCAS-D-21-0077.1>, 2022.
- Szychowska-Krapiec, E.: Long-term chronologies of pine (*Pinus sylvestris* L.) and fir (*Abies alba* Mill.) from the Małopolska region and their palaeoclimatic interpretation, *Folia Quat.*, 79, 1–124, 2010.
- Tardif, R., Hakim, G.J., Perkins, W.A., Horlick, K.A., Erb, M.P., Emile-Geay, J., Anderson, D.M., Steig, E.J., Noone, D.: Last millennium reanalysis with an expanded proxy database and seasonal proxy modeling. *Clim. Past* 15, 12, 2019. <https://cp.copernicus.org/articles/15/1251/2019/cp-15-1251-2019.pdf>.
- Tomczyk, A. M.: Temperatura powietrza, in: *Atlas klimatu Polski (1991-2020)*, edited by: Tomczyk, A. M. and Bednorz, E., Bogucki Wydawnictwo Naukowe, Poznań, 40–61, 2022.
- Trachsel, M., Grosjean, M., Larocque-Tobler, I., Schwikowski, M., Blass, A., and Sturm, M.: Quantitative summer temperature reconstruction derived from a combined biogenic Si and chironomid record from varved sediments of Lake

Silvaplana (southeastern Swiss Alps) back to AD 1177, *Quat. Sci. Rev.*, 29, 2719–2730, <https://doi.org/10.1016/j.quascirev.2010.06.026>, 2010.

Trouet, V., Esper, J., Graham, N.E., Baker, A., Scourse, J.D., and Frank, D.C.: Persistent Positive North Atlantic Oscillation Mode Dominated the Medieval Climate Anomaly, *Science* 324, 78–80,

785 <https://www.science.org/doi/abs/10.1126/science.1166349>, 2009.

Vogt, S., Glaser, R., Luterbacher, J., Riemann, D., Al Dyab, G., Schoenbein, J., and Garcia-Bustamante, E.: Assessing the Medieval Climate Anomaly in the Middle East: The potential of Arabic documentary sources, *PAGES news*, 19, 28–29, 2011.

Walawender, A.: *Kronika klęsk elementarnych w Polsce i w krajach sąsiednich w l. 1450-1586*, Warszawa, 1932.

790 Waszak, N., Robertson, I., Puchałka, R., Przybylak, R., Pospieszyńska, A., and Koprowski, M.: Investigating the Climate - Growth Response of Scots Pine (*Pinus sylvestris* L.) in Northern Poland, *Atmosphere (Basel)*, 12, 1690, <https://doi.org/10.3390/atmos12121690>, 2021.

Wigley, T., Briffa, K., and Jones, P.: On the average value of correlated time series, with applications in dendroclimatology and hydrometeorology., *J. Clim. Appl. Meteorol.*, 23, 201–213, 1984.

795 Wójcik, G., Majorowicz, J., Marciniak, K., Przybylak, R., Šafanda, J., and Zielski, A.: The last millennium climate change in northern Poland derived from well temperature profiles, tree – rings and instrumental data., *Pr. Geogr.*, 107, 137–148, 2000.

Zang, C. and Biondi, F.: treeclim: an R package for the numerical calibration of proxy-climate relationships, *Ecography (Cop.)*, 38, 431–436, <https://doi.org/10.1111/ecog.01335>, 2015.

800 Zielony, R. and Kliczkowska, A.: *Regionalizacja przyrodniczo-leśna Polski*, Centrum Informacyjne Lasów Państwowych, Warszawa, 2012.

Zielski, A.: *Uwarunkowania środowiskowe przyrostów radialnych sosny zwyczajnej (Pinus sylvestris L.) w Polsce północnej na podstawie wielowiekowej chronologii*, Wydawnictwo UMK, 127 pp., 1997.

Zielski, A. and Krąpiec, M.: *Dendrochronologia.*, PWN, Warszawa, 328 pp., 2004.


805 Zielski, A., Krąpiec, M., and Koprowski, M.: Dendrochronological Data, in: *The Polish Climate in the European Context: An Historical Overview.*, edited by: Przybylak, R., Majorowicz, J., Brázdil, R., and Kejna, M., Springer, Dordrecht, 191–217, <https://doi.org/10.1007/978-90-481-3167-9>, 2010.

Table 1. Statistics of chronology and climate reconstructions.


Chronology and its time span	Number of samples	eps	snr	Reconstructed parameter	Calibration Period	Calibration statistics	Verification period	Verification statistics			Model for whole period
								RE=	CE=	prediction RMSE=	
Scots pine. Kuyavia-Pomerania; 1168–2015	285		28.76	Feb–Mar temperature	1871–1943	r=0.477, p<0.05	1944–2015	0.159	0.133	0.153	0.45, p<0.05
Scots pine. Lesser Poland; 1091–2011	285	0.97	28.70	Feb–Mar temperature	1846–1960	r=0.44, p<0.05	1961–2000	0.134	0.116	0.384	0.39, p<0.05
Silver fir. Lesser Poland; 1109–2017	484	0.98	61.34	Dec–Mar temperature	1846–1960	r=0.57, p<0.05	1961–2000	0.036	0.028	0.312	0.49, p<0.05

825 **Table 2.** Frequency of occurrence of extremely warm and wet, as well as cold and dry, winters (DJF) and summers (JJA) in Poland from 1361 to 1500.

Period	Air temperature				Atmospheric precipitation				Extreme situations	
	DJF		JJA		DJF		JJA			
	2 & 3	-2 & -3	2 & 3	-2 & -3	2 & 3	-2 & -3	2 & 3	-2 & -3	Total	%
1361–1400	0	4	1	1	2	0	1	1	10	7.1
1401–50	5	18	4	4	7	3	8	4	53	37.6
1451–1500	5	19	16	5	7	3	10	13	78	55.3
1361–1500	10	41	21	10	16	6	19	18	141	
%	7.1	29.1	14.9	7.1	11.3	4.2	13.5	12.8		100




830 **Figure 1.** Study area with historical and natural-forest regionalisation and location of materials used for research. Natural-Forest Provinces (Zielony and Kliczkowska 2012): I – Baltic Coast province II – Masuria–Podlachia province, III – Greater Poland–Pomerania province, IV – Masovia–Podlachia province, V – Silesia province, VI – Lesser Poland province, VII – Sudetia province, VIII – Carpathia province. Dendrochronological sites: yellow dots – pine, red dots – oak, black dots – fir. Black squares – meteorological stations




835


Figure 2. Number of weather notes/excerpts for Poland available in historical sources in the periods: *above*, AD 1001–1360; *below*, AD 1361–1500



840 **Figure 3.** Decadal frequencies of occurrence of winters (DJF), springs (MAM), summers (JJA) and autumns (SON) that were extremely cold and very cold (indices -3 and -2) and extremely warm and very warm (indices 3 and 2) in Poland between 1361 and 1500



845 **Figure 4.** Decadal frequencies of occurrence of winters (DJF), springs (MAM), summers (JJA) and autumns (SON) that were extremely dry and very dry (indices -3 and -2) and extremely wet and very wet (indices 3 and 2) in Poland between 1361 and 1500



850 **Figure 5.** Anomalies of 10-year mean values of winter (DJF) and summer (JJA) air temperature in Poland in 15th century relative to means from 1951–2000 reference period. For reference period. Data were taken after Kożuchowski and Żmudzka (2003).



Figure 6. Pointer years in oak growing in Greater Poland, Lower Silesia and Lesser Poland, 1001–1500






Figure 7. Frequency of occurrence of pointer years in trees (Scots pine, silver fir and oak) in 50-year periods in Poland, 1001–1500



860 **Figure 8.** Pointer years in trees (Scots pine and silver fir) growing in Kuyavia-Pomerania and Lesser Poland, 1101–1500



865 **Figure 9.** Reconstruction of average air temperature (°C): for February–March in: (a) northern Poland and (b) southern Poland, based on the width of pine rings; and (c) for December–March in southern Poland based on fir-ring widths. Anomalies were calculated relative to the period 1951–2000. Standard deviations were calculated using data from 1951–2000




Figure 10. Reconstruction of air temperature in Poland in moving windows: A) Feb–Mar temperature anomaly in a Gaussian 20-year moving window for Kuyavia-Pomerania region. B) Feb–Mar temperature anomaly in a Gaussian 20-year moving window for the Lesser Poland region. C) Dec–Mar temperature anomaly in a Gaussian 20-year moving window for the Lesser Poland region. D) Feb–Mar temperature anomaly in a Gaussian 20-year moving window for Kuyavia-Pomerania (modified after Koprowski et al. 2012). E) Nov–Apr temperature anomaly in a Gaussian 20-year moving window for northern Poland (modified after Balanzategui et al. 2017). F) chrysophyte-based reconstruction of number of days below 4 °C (Hernández-Almeida et al. 2015). G) chironomid-based reconstruction of August temperature (Hernández-Almeida et al. 2017)