

Evaluation of WRF-Chem simulated meteorology and aerosols over northern India during the severe pollution episode of 2016

Prerita Agarwal¹, David S. Stevenson¹, Mathew R Heal²

¹School of GeoSciences, University of Edinburgh, Crew Building, Edinburgh, EH9 3FF, UK,

5 ²School of Chemistry, University of Edinburgh, Joseph Black Building, Edinburgh, EH9 3FJ, UK

Correspondence to: Prerita Agarwal (prerita.agarwal@ed.ac.uk), David Stevenson (david.s.stevenson@ed.ac.uk)

Abstract

10 We use a state-of-the-art regional chemistry transport model (WRF-Chem v4.2.1) to simulate particulate air pollution over northern India during September-November 2016. This period includes a severe air pollution episode marked by exceedingly high levels of hourly PM_{2.5} (particulate matter having an aerodynamic diameter ≤ 2.5 microns) during 30th October to 7th November, particularly over the wider Indo-Gangetic Plain (IGP). We provide a comprehensive evaluation of simulated seasonal
15 meteorology (nudged by ERA5 reanalysis products) and aerosol chemistry (PM_{2.5} and its black carbon (BC) component) using a range of ground-based, satellite and reanalysis products, with a focus on the November 2016 haze episode. We find the daily and diurnal features in simulated surface temperature show the best agreement followed by relative humidity, with the largest discrepancies being an overestimate of night-time wind-speeds (up to 1.5 m s⁻¹) confirmed by both ground and radiosonde
20 observations. Upper air meteorology comparisons with radiosonde observations show excellent model skill in reproducing the vertical temperature gradient ($r > 0.9$). We evaluate modelled PM_{2.5} at 20 observation sites across the IGP including eight in Delhi and compare simulated aerosol optical depth (AOD) with data from four AERONET sites. We also compare our model aerosol results with MERRA-2 reanalysis aerosol fields and MODIS satellite AOD. We find that the model captures many features
25 of the observed aerosol distributions but tends to overestimate PM_{2.5} during September (by a factor of 2) due to too much dust, and underestimate peak PM_{2.5} during the severe episode. Delhi experiences some of the highest daily mean PM_{2.5} concentrations within the study region, with dominant components nitrate (~25 %), dust (~25 %), secondary organic aerosols (~20 %), and ammonium (~10%). Modelled PM_{2.5} and BC spatially correlate well with MERRA-2 products across the whole
30 domain. High AOD at 550nm across the IGP is also well predicted by the model relative to MODIS satellite ($r \geq 0.8$) and ground-based AERONET observations ($r \geq 0.7$), except during September. Overall, the model realistically captures the seasonal and spatial variations of meteorology and ambient pollution over northern India. However, the observed underestimations in pollutant concentrations likely come from a combination of underestimated emissions, too much night-time dispersion, and some
35 missing or poorly represented aerosol chemistry processes. Nevertheless, we find the model is sufficiently accurate to be a useful tool for exploring the sources and processes that control PM_{2.5}-levels during severe pollution episodes.

Introduction

Atmospheric particle pollution in India is a persistent environmental issue and a leading health risk factor for its 1.4 billion population (Pandey et al., 2021). In 2019, ambient air pollution was estimated to cause almost a million premature deaths in India (Pandey et al., 2021). The State of Global Air 2022 (HEI, 2022) reports that over 90% of the Indian population resides in areas where the annual mean concentrations of PM_{2.5} (particulate matter having an aerodynamic diameter smaller than 2.5 microns) exceed even the minimal interim target of 35 $\mu\text{g m}^{-3}$ recommended by the World Health Organization Air Quality Guidelines (WHO 2021). The country is home to 18 of the 20 cities worldwide with the greatest rise in PM_{2.5} pollution in the last decade (HEI, 2022). This upward trend in degraded air quality is projected to continue across South Asia under current policies, including more frequent high pollution incidents over northern India (Kumar et al., 2018c; Paulot et al., 2022). These trends have huge consequences for the future life expectancy of the 400 million residents of this region which is currently reported to be reduced by more than 9 years under the current pollution burden (Greenstone and Fan 2020).

The Indo-Gangetic Plain (IGP) is situated south of the Himalayas and stretches from parts of Pakistan in the west, through north and east India and Nepal, to Bangladesh in the east. The IGP is a heavily populated region (home to over 40% of the total Indian population) with many rural, suburban and urban clusters (Fig. 1a). It is characterised by intensive multi-cropping systems, rapid industrialisation and a growing economy, which results in a heterogeneous mix of particle and gaseous pollutant emissions (Venkataraman et al., 2018; Kumar et al., 2020a). The region is a global centre for poor air quality (Singh et al., 2017), underpinned by India being one of the largest emitters of anthropogenic aerosols in the world (Lu et al., 2011). The anthropogenic sources include vehicles, industry, burning of crop-waste and garbage, residential cooking, and mining. The emissions contributions are dominantly composed of nitrate and sulfate precursors and carbonaceous aerosols, driven by a rapid increase in demand for energy (Lu et al., 2011). Black carbon (BC) is fine particulate matter's light-absorbing component (Lack and Cappa, 2010) and is released during incomplete combustion of carbon-containing fossil fuels like coal, oil and gas, and biofuels like wood, agricultural residues and forest fires. BC particles are short-term climate forcers with a net positive radiative forcing (Ramanathan et al., 2001; Bond et al., 2013; Wang et al., 2014). BC emissions from India are one of the highest globally, and significantly impact the Indian summer monsoon, regional climate, and human health (Ramanathan et al., 2001). Natural particle sources such as mineral dust also substantially influence the air quality over the IGP and broader northern India (Li et al., 2017). Additionally, air quality over the IGP region is greatly affected by the prevailing meteorology, topography and the long-range transport of pollutants (Kaskaoutis et al., 2014; Kumar et al., 2014; Schnell et al., 2018; Ojha et al., 2020).

In addition to the year-round poor air quality over the IGP region, recurring intense post-monsoon and winter haze episodes have been reported in numerous studies (Ram et al., 2016; Kumar et al., 2018a; Kanawade et al., 2020; Beig et al., 2019; Bharali et al., 2019; Thomas et al., 2019; Dhaka et al., 2020; Kumari et al., 2021; Gupta et al., 2022). Most of these severe episodes coincide with the biomass-burning period (mid-October to November), during which agricultural land is cleared in open fields by burning crop residues, primarily paddy (Singh et al., 2020). Although highly seasonal, the emissions from these multiple small to large fires emit large amounts of reactive gases and particles such as carbon monoxide (CO), nitrogen oxides (NO_x), volatile organic compounds (VOCs), carbonaceous particles and other components of PM_{2.5} (Singh et al., 2020; Kumar et al., 2021). One such severe haze event over northern India occurred between 30th October and 7th November 2016, leading to daily mean PM_{2.5} concentrations of 300-600 µg m⁻³, some 20-40 times greater than the 24-h WHO 2021 Air Quality Guideline of 15 µg m⁻³ (Mukherjee et al., 2018; Sawlani et al., 2019; Kanawade et al., 2020; Jethva et al., 2019). Jethva et al. (2019) reported that crop-residue fire counts over northwest India were particularly high in the 2016 post-monsoon period. Alongside crop biomass burning emissions, the unfavourable meteorology and accumulation of local urban emissions contributed to this week-long extremely high pollution episode (Kanawade et al., 2020; Sawlani et al., 2019).

Modelling studies characterising air pollution over India have utilised a variety of regional chemistry transport models (Nair et al., 2012; Kumar et al., 2012a, b; Moorthy et al., 2013; Pan et al., 2015; Srivastava et al., 2016; Schnell et al., 2018; Ghosh et al., 2023). These studies highlight various problems in simulating atmospheric composition over the Indian subcontinent, such as capturing the high aerosol loading, erroneous boundary-layer parametrizations, underestimations in emissions inventories, complex mountain topography and inaccurate moisture transport. This is especially true for simulations of surface BC concentrations, which utilise regional South Asian emissions inventories that are thought to underestimate the BC emissions (Kumar et al., 2015; Govardhan et al., 2019). Equally important is simulating the vertical distribution of BC particles and understanding their effect on atmospheric stability, for which only limited measurements have been made over India. These studies found high BC loadings vertically (up to 8 km altitude) over northwest and central India during different months (Babu et al., 2011; Bisht et al., 2016; Brooks et al., 2019). The role of these absorbing particles in modifying the vertical boundary layer structure during a haze episode over the northern Indian region has been poorly explored to date (Bharali et al., 2019).

This study aims to evaluate the WRF-Chem regional atmospheric chemistry transport model's ability to simulate the meteorology and aerosol chemistry across north India and the IGP in September-November 2016. Our choice to analyse the 2016 seasonality and the pollution episode differs from previous literature (Kumar et al., 2020b; Jena et al., 2020; Sengupta et al., 2022; Govardhan et al., 2023b) in several aspects, listed below.

110 (i) We use an updated WRF-Chem version (v4.2.1) and utilise the MOZART-MOSAIC chemical scheme (detailed in Section 2.1), which explicitly resolves the aerosols into four size bins and represents the chemistry of secondary organic and inorganic aerosols that make up the dominant components of PM_{2.5} in the post-monsoon season, as compared to the GOCART scheme employed in these earlier studies which lacks particle size information.

115 (ii) The 2016 pollution episode over the IGP was one of the worst for air quality (since 2004) and anomalous for the highest rice crop production (since 2002) in NW Indian states, resulting in high crop residue burning in that year (Voiland and Jethva, 2017; Jethva et al., 2019; Sembhi et al., 2020). As shown by multiple trend analyses, 2016 had the highest number of agricultural fires of the last decade during the post-monsoon season (Sarkar et al., 2018; Mukherjee et al., 2018; Thomas et al., 2019; Kulkarni et al., 2020; Sembhi et al., 2020; Liu et al., 2021; Jethva, 2022; Gupta et al., 2023). Moreover, although several modelling studies have analysed the air quality during intense post-monsoon pollution episodes in the years after 2016 (e.g., Dekker et al., 2019; Beig et al., 2019; Kulkarni et al., 2020; Roozitalab et al., 2021), studies for 2016 are fewer (Sembhi et al., 2020; Mukherjee et al., 2020). It is, therefore, necessary to understand the implications of this particularly extreme episode with a chemistry transport model whose performance at simulating prevailing seasonal meteorology over a sufficiently long period has been evaluated.

125 (iii) The use of hourly fifth-generation European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis (ERA5) data to drive the model meteorology and a comprehensive comparison of the simulated meteorology and biases across northern India is an additional novelty of this work, as is the use of a wide range of ground and satellite observations as well as reanalysis products in the evaluation.

130 We first evaluate WRF-Chem simulations of surface and vertical meteorology against multiple available observations from weather stations and radiosonde profiles and reanalysis datasets. We then evaluate the modelled chemistry and aerosol optical properties against ground-based measurements, reanalysis products, and satellite-retrieved data. The study focuses on characterising monsoon to post-monsoon changes in meteorology and the atmospheric chemical composition of modelled PM_{2.5} and BC in 2016.




Figure 1. a) Degree of urbanisation based on 2015 human population size and built-up area density data over India from GHS-SMOD (Schiavina et al., 2023) b) Locations of the observation sites used for comparison in this study; the legend indicates the different datasets (ASOS: automatic weather stations, RAOB: Radiosonde observations, CPCB: Indian Central Pollution Control Board PM_{2.5} ground monitoring stations, AERONET: Aerosol Robotic Network ground remote sensing observations). The inset figure is an enlarged map of Delhi capital and the geographical area falling under IGP region is highlighted in light blue colour.

2. Data and Methods

2.1 WRF-Chem model description and configuration

The Weather Research and Forecasting model (version 4.2.1) coupled with Chemistry (WRF-Chem) (Grell et al., 2005; Fast et al., 2006) is an atmospheric chemistry transport model widely applied to the South Asia region, including its development as an air quality early-warning system for Delhi (Jena et al., 2021; Kumar et al., 2020b). It has a terrain-following vertical coordinate system and is available with a range of physical parameterizations (Skamarock et al., 2008). The transport of trace gases and aerosol species in WRF-Chem uses identical vertical and horizontal coordinates, allowing for feedback between meteorology and chemistry via radiation and photolysis (Grell et al., 2005). This makes WRF-Chem well-suited for investigating and isolating the interactions between aerosols and meteorology.

The single domain for this study covers the northern part of South Asia (20 – 38° N and 66 – 90° E) at 12 km horizontal resolution (Fig. 1b), with 33 vertical levels from the surface to the model top which is fixed at 50 hPa. The lowest ten levels are below 1 km. The configurations of WRF-Chem dynamical and chemical parametrizations used in this study are adopted from the literature available for South Asia and are summarized in Supplementary Table S1. ERA5 data at a horizontal resolution of 0.25° x

0.25° is used for initializing the meteorology, boundary conditions and nudging in the model (Hersbach et al., 2020). Temperature, winds and water vapour are nudged towards ERA5 values above the planetary boundary layer (PBL) every 6 hours, using grid-nudging with a nudging coefficient of $6 \times 10^{-4} \text{ s}^{-1}$ (Stauffer and Seaman, 1994). Terrestrial and land-use data are static and obtained from the MODIS IGBP 21-category land-cover classification (Friedl et al., 2002).

Time-varying boundary conditions for chemistry are taken from simulations of the global 6-hourly Model for Ozone and Related Chemical Tracers (MOZART-4)/Goddard Earth Observing System Model version 5 (GEOS-5) (National Center for Atmospheric Research 2016). The simulation of gas-phase chemistry in WRF-Chem is provided by the updated MOZART-4 scheme (Emmons et al., 2010, 2020) which includes treatment of biogenic hydrocarbons and aromatics (Hodzic and Jimenez, 2011; Knote et al., 2014). Aerosol chemistry is simulated using the Model for Simulating Aerosol Interactions and Chemistry (MOSAIC) 4-bin scheme (Zaveri et al., 2008). MOSAIC includes detailed solid, liquid and mixed-phase equilibria and thermodynamic gas-particle partitioning to compute aerosol composition, and a simple parameterisation of SOA aqueous chemistry using glyoxal (Knote et al., 2014), but does not explicitly include detailed aqueous-phase chemistry, such as that described in Acharja et al. (2023). The aerosol processes in the mechanism include aerosol transport, dry and wet removal, water uptake, nucleation, coagulation, and condensation processes. The MOSAIC scheme uses a sectional approach to divide dry aerosol diameter into four discrete bins: 0.039–0.156 μm , 0.156–0.625 μm , 0.625–2.5 μm and 2.5–10 μm (the coarse PM bin) (Zaveri et al., 2008). The aerosol distribution scheme includes both in-cloud and impaction scavenging and assumes aerosols to be internally mixed within the same bin and externally mixed between the bins (Riemer et al., 2019). MOSAIC simulates sulfate (SO_4^{2-}), nitrate (NO_3^-), ammonium (NH_4^+), calcium (Ca^{2+}), carbonate (CO_3^{2-}), black carbon (BC), primary organic mass (OM), liquid water (H_2O), sea salt (NaCl) and other inorganic species such as minerals and trace metals (Zaveri et al., 2008). The Fast Tropospheric Ultraviolet–Visible (FTUV) photolysis scheme (Tie et al., 2003) provides photolysis rates and accounts for the aerosol feedback on photolysis (Hodzic and Knote, 2014).

Monthly anthropogenic emissions at $0.1^\circ \times 0.1^\circ$ horizontal resolution are obtained from the 2010 Emission Database for Global Atmospheric Research for Hemispheric Transport of Air Pollution version 2.2 (EDGAR-HTAPv2.2, https://edgar.jrc.ec.europa.eu/dataset_htap_v2). The emission sectors included in EDGAR-HTAPv2.2 are industrial, residential, transportation, agriculture, shipping, energy, and aviation. For emissions from India, EDGAR-HTAPv2.2 incorporates the regional emissions inventory from the Model Inter-Comparison Study for Asia Phase III (MICS-Asia III) to derive emissions maps at a common grid resolution of $0.1^\circ \times 0.1^\circ$ (Janssens-Maenhout et al., 2015). Within MICS-Asia III, a mosaic of regional anthropogenic emission inventories was developed by combining the nationally reported estimates by Argonne National Laboratory (ANL-India) and REAS2 (Regional Emission inventory in Asia) (Lu et al., 2011; Li et al., 2017). The total emissions for SO_2 , NO_x , NH_3 ,

PM₁₀, PM_{2.5}, BC, OC and non-methane volatile organic compounds (NMVOCs) are speciated in the model following the MOSAIC-MOZART chemistry mechanism. The anthropogenic emissions exhibit a diurnal variation, with a simple transition between day-time and night-time values at 05:30 and 17:30 local time for each emitted pollutant, as specified by the preprocessor tool anthro-emiss utility in the model (<https://www2.acom.ucar.edu/wrf-chem/wrf-chem-tools-community>). The use of the 2010 EDGAR-HTAPv2.2 inventory to model air quality during 2016 adds some uncertainties to the model results as the emissions over India evolved from 2010 to 2016. Emissions of OC, CO, NO_x, SO₂, and NMVOC from anthropogenic sectors such as industrial and energy sectors increased because of rapidly increasing demand, whilst primary particulate emissions of BC, OC and PM_{2.5} from residential and informal industry sectors reduced due to cleaner fuel policies (such as the Ujjawala scheme; <http://www.pmujjwalayojana.in/>) (McDuffie et al., 2020). The estimates derived from the global CEDS inventory reported by McDuffie et al. (2020) show a combined increase from road transport, energy, industry, and agricultural sectors in annual NH₃, SO₂, NO_x, and NMVOC emissions over India between 2010 to 2016. The reported increases in emissions of these pollutants across India are ~17%, ~11%, ~12 % and ~10 %, respectively. These changes in emissions may mean our model simulations underestimate the BC, primary OC, and secondary aerosol contributions to total PM. However, it is challenging to fully isolate the impact of these changes in an atmospheric chemistry model because the model output also depends substantially on other factors, such as the meteorology, which drives online emissions. Compared with other global inventories of coarser resolution (e.g., ECLIPSE), the use of EDGAR-HTAPv2.2 has been found to simulate air quality over India with a greater local heterogeneity and to show slightly smaller overall biases when compared against reanalysis and satellite products. (e.g., Upadhyay et al., 2020). Hence, although the EDGAR-HTAPv2.2 emissions are from 2010, we believe that they have been widely evaluated and are amongst the best available for our simulations.

In India, the post-harvest agricultural residue is largely cleared by burning it in open fields, and this is a dominant contributor to Indian PM_{2.5}, BC, OC, SO₂, NO_x and NMVOC emissions (Venkataraman et al., 2018). As EDGAR emissions do not include any biomass burning emissions (from agricultural fires, wildfires or prescribed fires), these are derived from the Fire Inventory from NCAR, version 1.5 (FINNv1.5) (Wiedinmyer et al., 2011). The emissions are based on satellite-measured locations of active fires and emission factors relevant to the underlying land cover (Akagi et al., 2011). The FINNv1.5 fire emissions inputs are distributed at 1 km spatial and hourly temporal resolution for 2016 (<https://www.acom.ucar.edu/Data/fire/>).

Biogenic emissions are calculated online (updated every 30 minutes) using the Model of Emissions of Gases and Aerosol from Nature (MEGAN v2.0) (Guenther et al., 2006). MEGAN uses satellite-driven land cover and modelled meteorological information (e.g., temperature and photosynthetically available radiation, PAR) to estimate VOCs, NO_x and CO from vegetation at spatial resolution. Dust emissions are generated online by incorporating the Goddard Global Ozone Chemistry Aerosol Radiation and

Transport (GOCART) scheme from terrain data and modelled meteorology (Chin et al., 2002). The
225 GOCART scheme, described in detail elsewhere (Ginoux et al., 2001; Zhao et al., 2010, 2013), utilises
the information about 10 m wind speed, threshold wind velocity (minimum value to reach for the dust
emission to occur) and potential dust source region factors to calculate the dust emission flux. The total
dust emission fluxes are calculated by multiplying with an empirical dimensional constant which is
taken from Ginoux et al. (2001). The GOCART scheme then distributes the emitted dust particles into
230 4 size bins (described earlier).

For our evaluation of WRF-Chem performance, hourly simulations are conducted for 01 September to
30 November 2016, allowing six days of spin-up (from 25 – 30 August). September falls within the
south-west (SW) monsoon season (its withdrawal typically begins in mid-September), whilst October
and November are in the post-monsoon season (Annual report 2016, India Meteorological Department).
235 This permits a comparative assessment of meteorology and air quality between the two seasons.
Although 2015-2016 was widely recorded as subject to a pronounced El Niño event, its effects over
India lasted only until the summer of 2016 (India Meteorological Department, Govt. of India Ministry
of Earth Sciences, 2017) and therefore should not significantly impact the study period. In terms of
general climatology, the 2016 SW monsoon rainfall was recorded to be normal over the country, aside
240 from a deficit in rainfall over parts of northwest India.

2.2 Meteorological data

WRF-Chem simulated meteorology is compared with observational networks measuring daily surface
weather (Iowa Environmental Mesonet-Automated Surface Observing System; IEM-ASOS Network)
and atmospheric soundings (radiosonde observations (RAOB), University of Wyoming). Figure 1b
245 shows the locations of the observation sites from these networks. The data links and access details are
given in Table S2.

The IEM-ASOS network is an archive of global automated airport weather observations from weather
stations operated by national agencies and airport authorities. Hourly 2-m air temperature (T2), relative
humidity (RH), wind speed (WS) and wind direction (WD) data for 49 observation sites (Table S4)
250 within the study domain are used. Processing and general quality control of the data is undertaken by
the IEM-Network so the downloaded data was only checked for missing values before comparison with
model output.

Radiosonde measurements for vertical meteorology profile comparison are available for eight sites
within the model domain. Pilot balloon soundings are undertaken by the India Meteorological
255 Department, and rigorous quality checks are performed before making them freely available (Durre et
al., 2006). The radiosonde measurements are available each day at 00:00 UTC (05.30 and 17.30 Indian
Standard Time (IST), respectively). No station has complete soundings for the entire study period, so

model-measurement comparisons include only times when observations are available. The sounding observations are vertically interpolated to the model's pressure levels from 1000 hPa to 100 hPa. The average vertical temperature, virtual potential temperature (VPT), WS and RH profiles are compared for individual sites and temporal variability (as standard deviation) is reported for the entire period across all the pressure levels.

The spatial features of modelled meteorology are compared against the global MERRA-2 reanalysis (Gelaro et al., 2017) dataset available at a latitude-longitude grid resolution of $0.5^\circ \times 0.625^\circ$ and 72-eta hybrid levels at 6-h frequency. MERRA-2 reanalysis data is provided by NASA's Global Modelling and Assimilation Office (GMAO). The meteorological variables are re-gridded to WRF-Chem spatial resolution (12 km), and comparison was undertaken for T2, 10 m WS, water vapour mixing ratio (QV) and planetary boundary layer height (PBLH) variables.

2.3 Ground-based PM_{2.5}

We evaluate the performance of WRF-Chem in simulating aerosols by comparing modelled PM_{2.5} mass concentrations and aerosol optical depth (AOD) at 550 nm with observations and reanalysis products. The measurements of surface PM_{2.5} used for model comparison are undertaken by the Central Pollution Control Board of India (CPCB), accessed via the OpenAQ platform (Table S1, Fig. 1b). In addition to general quality control procedures applied by CPCB, the hourly PM_{2.5} mass concentration data for 20 stations in the study domain was filtered for missing, zero and negative values. Days with <40 % of hourly measurements were also removed before comparing with the modelled PM_{2.5} mass concentrations. Since Delhi has many more individual sites than other states in the domain, the data is grouped into two categories: all sites within the Delhi region ($n = 8$), and the remaining sites (referred to as 'Others', $n = 12$), the majority of which are located within the IGP region (Fig. 1b, Table S4).

2.4 Reanalysis PM_{2.5} and Black Carbon concentrations

The spatial distributions of modelled surface PM_{2.5} and BC concentrations are compared with MERRA-2 global reanalysis products, which is based on the GOCART scheme employed in GEOS-5 atmospheric model (Randles et al., 2017). The GOCART model in MERRA-2 employs the online coupling of radiatively-active aerosols with meteorology in the GEOS-5 model. GOCART in MERRA-2 simulates OC, BC, sea salt, dust, and sulfate aerosols which are used to derive the total PM_{2.5} mass concentrations but it lacks information on size distribution and composition of aerosols. Additionally, the aerosols in the GOCART scheme are externally mixed and exclude the treatment of nitrate, and secondary organic aerosols (Randles et al., 2017) due to which the MERRA-2 PM_{2.5} is underestimated during high pollution events (Buchard et al., 2017). Whilst the MERRA-2 AOD is directly constrained by the assimilation of observations, the aerosol diagnostics (such as PM_{2.5}) also partly depend upon systematic biases and assumptions of aerosol speciation and optical properties in the GEOS-

5/GOCART model used in MERRA-2, as noted by (Buchard et al., 2017). This is likely to influence the total PM_{2.5} comparisons between WRF-Chem and MERRA-2, as WRF-Chem simulates a wider range of aerosol species. However, the utilisation of MERRA-2 reanalysis data still serves as a useful reference for assessment of spatial and seasonal trends of aerosols between the two modelled datasets.

AOD in MERRA-2 is assimilated using multiple satellite and ground-based observation data, including bias-corrected AOD from the Moderate Resolution Imaging Spectroradiometer (MODIS), Advanced Very High-Resolution Radiometer (AVHRR) instruments, Multi-angle Imaging Spectroradiometer (MISR) and Aerosol Robotic Network (AERONET). The aerosol assimilation uses satellite radiance and albedo from observing sensors and bias-corrected AOD, described in detail in Randles et al. (2017).

Based on past studies and recommendations, the PM_{2.5} concentration from MERRA-2 produced aerosol fields is calculated via the following summation of aerosol components in the size bin $\leq 2.5 \mu\text{m}$ diameter.

$$[\text{PM}_{2.5}] = [\text{BC}] + 1.6 \times [\text{OC}] + 1.375 \times [\text{SO}_4^{2-}] + [\text{Dust}] + [\text{Sea Salt}]$$

The multiplication factor of 1.375 on the sulfate ion concentration is based on the assumption in MERRA-2 that sulfate is primarily present as neutralised ammonium sulfate (Buchard et al., 2016; Provencal et al., 2017; Song et al., 2018). OC in MERRA-2 is scaled up to organic matter concentration using values ranging from 1.2 - 2.6, and this study uses the factor 1.6, which is commonly used for urban carbonaceous particles (Chow et al., 2015; Buchard et al., 2016; Provencal et al., 2017; Song et al., 2018).

2.5 Satellite and ground-based AOD data

WRF-Chem AOD at 550 nm is compared with satellite observations from the MODIS sensor on board the Terra and Aqua polar orbiting satellites. The AOD products from MODIS have a 10 km horizontal resolution at equatorial local overpass times of 10.30 (Terra) and 13.30 (Aqua). AOD retrievals from MODIS are based on combined Dark Target (DT: retrieval algorithm over dark land and ocean surfaces) and Dark Blue algorithms (DB: bright land surface) and re-gridded to the WRF-Chem resolution of 12 km. AOD in WRF-Chem is simulated between wavelengths 300 - 1000 nm and interpolated to 550 nm using the Ångström power law (Ångström, 1964; Kumar et al., 2014). In addition, ground-based AERONET version 2 level 2.0 (quality-assured and cloud-screened) AOD is available at four locations (Fig. 1b) within the study domain and is used for comparison with modelled results. AERONET is a global network (Holben et al., 1998) that has been extensively used for validating satellite observations over South Asia (Sayer et al., 2014; Mhawish et al., 2017).




Figure 2. Daily-mean time series (left) and mean diurnal cycle (right) of observed (black) and modelled (red) meteorological variables from 01 September – 30 Nov 2016 averaged across the 49 ASOS measurement sites shown in Figure 1b. From top to bottom: daily mean 2-m temperature, relative humidity, wind speed, and wind direction. The shaded regions indicate the standard deviation in the spatial variability in the model and measured variables. The vertical dashed lines delineate the period of severe high pollution between 30 October and 7 November.

2.5 Statistical Metrics

325 Statistical metrics used here for the evaluation of model performance include mean bias (MB), normalized mean bias (NMB), mean absolute error (MAE), root mean square error (RMSE) and Pearson's correlation coefficient (r). Definitions of these metrics are provided in Supplementary Material Table S3.

Table 1. Summary of statistical comparison of modelled and observed meteorology variables derived from hourly data between September to November 2016 and averaged across the 49 ASOS measurement sites shown in Figure 1b. The statistical metrics used for comparison are mean bias (MB), normalized mean bias (NMB), mean absolute error (MAE), root mean square error (RMSE) and Pearson's correlation coefficient (r).

330

Month/Variable	MB	NMB	MAE	RMSE	r
($N = 49$)					
<u>Temperature ($^{\circ}\text{C}$)</u>					
September	-0.28	-0.01	1.5	2.2	0.86
October	-0.75	-0.03	1.8	2.6	0.90
November	-0.84	-0.04	2.2	3.0	0.87
<u>RH (%)</u>					
September	-1.90	-0.03	7.8	10.0	0.75
October	-4.10	-0.07	10.1	13.2	0.79
November	-8.20	-0.15	12.8	17.7	0.65
<u>Wind Speed (m s^{-1})</u>					
September	0.40	0.20	0.8	1.07	0.62
October	0.54	0.36	1.0	1.26	0.30
November	0.81	0.61	1.1	1.37	0.40

Table 2. Summary of statistical comparison of WRF-Chem and MERRA-2 derived meteorology variables from September to November 2016. The statistical metrics used for comparison are mean bias (MB), normalized mean bias (NMB), mean absolute error (MAE), root mean square error (RMSE) and Pearson’s correlation coefficient (r).

Month	MB	NMB	MAE	RMSE	r
Temperature (°C)					
September	-0.57	-0.03	1.4	2.12	0.99
October	-1.5	-0.10	1.9	2.67	0.99
November	-2.4	-0.22	2.7	3.30	0.99
Wind Speed (m s⁻¹)					
September	-0.17	-0.09	0.57	0.73	0.85
October	-0.23	-0.12	0.64	0.85	0.76
November	-0.17	-0.08	0.79	1.09	0.73
QV2 (g kg⁻¹)					
September	0.56	0.05	0.94	1.38	0.98
October	0.19	0.02	0.79	1.11	0.98
November	-0.07	-0.01	0.65	0.99	0.97
PBLH (m)					
September	-324	-0.28	355	430	0.69
October	-477	-0.37	481	550	0.67
November	-344	-0.36	356	446	0.70
PM_{2.5} (µg m⁻³)					
September	54	1.9	55.1	72	0.87
October	20	0.49	21.7	30	0.87
November	-8.4	-0.12	13.8	23	0.95
BC (µg m⁻³)					
September	0.52	0.65	0.57	0.93	0.91
October	0.24	0.19	0.44	0.79	0.91
November	-0.78	-0.28	0.89	1.42	0.92

3. Meteorology evaluation results

3.1 Near-surface meteorological fields

Figure 2 shows modelled and measured time series of daily means and mean diurnal cycles T2, RH, WS and WD derived from hourly data and averaged across all the observational sites. The statistical comparison metrics for the three months are provided in Table 1. As the exact measurement heights at individual sites are not known, the comparisons are made assuming the standard above-ground heights of 2 m for temperature and RH and 10 m for wind speed and direction. Daily average T2 variability correlates well between the model and observations for all the months ($r > 0.85$), with maxima and minima captured well (Fig. 2). Model MB for T2 is slightly lower but by less than $-0.8\text{ }^{\circ}\text{C}$ for all months. The T2 diurnal profile is also well represented by the model, with differences slightly larger (up to $2\text{ }^{\circ}\text{C}$) during night-time.

The general day-to-day variability in modelled surface RH also compares reasonably well with the observations (r range across the months: $0.65 - 0.79$) with slight underestimations that gradually increase from -1.9% in September to -8.2% in November mainly due to underestimations seen in the night-time RH peaks. The observed diurnal RH cycle is also well simulated by the model, although as for T2 with larger differences during the night when RH is greatest.

The differences in simulated 10 m wind patterns are relatively higher than those for T2 and RH, with modelled WS showing a relatively poor correlation of $r \leq 0.4$ and overestimations of about $0.5 - 0.8\text{ m s}^{-1}$ ($36 - 61\%$) in October and November. However, better correlation ($r = 0.62$) and lower biases (MB = 0.4 m s^{-1} and NMB = 0.2) are observed for September. The diurnal variation of WS during daytime is captured quite well by the model, while the bias is higher at night (up to 1.5 m s^{-1}); this is the reason for the observed large biases in modelled daily variabilities in WS. Since local WD is highly variable across sites in different regions, it is hard for a model to capture the daily variabilities. Differences between modelled and observed WD are smallest during the daytime when the general wind direction is south-westerly, and largest at night.

Table 2 provides the statistical evaluation results from the comparison of WRF-Chem and MERRA-2 global reanalysis data for mean T2, 10-m wind speed, water vapour mixing ratio (QV) and planetary boundary layer height (PBLH). The spatial maps of these variables are presented in Figures S1 and S2. Except for PBLH, the meteorological variables generally show good spatiotemporal agreement between the model and MERRA-2, with the best agreement for T2 and QV, as reflected in the high spatial correlations ($r \geq 0.97$). However, regional heterogeneities exist between the two datasets which are generally more evident temporally across all the variables. The largest spatial differences are seen for WS and QV which show overall underestimations by WRF-Chem for WS (in contrast to observed overestimations as compared to the measured data) and overestimations for QV (in contrast to observed

underestimations as compared to the measured data) in the wider IGP region. There is a stronger west-east gradient in PBLH in MERRA-2 compared to WRF-Chem which possibly influences the PM_{2.5} concentrations in MERRA-2.

370 A seasonal dry bias in the WRF model over the Indian region due to possible errors in moisture fluxes has been reported previously (Kumar et al., 2012b; Conibear et al., 2018), and night-time underestimations in modelled RH similar in magnitude to this study were noted by Gunwani and Mohan (2017). A comparison of modelled results, including ERA5 (used to drive WRF-Chem here) and independent MERRA-2 global reanalysis datasets with hourly ground observations (Fig. S3) shows the highest positive bias in RH in ERA5 during all the months, while WRF-Chem and MERRA-2 tend to
375 underestimate RH across all the months. This may affect the model's ability to capture the diurnal evolution of secondary aerosols by hygroscopic growth, particularly at night.

The observed positive bias in simulated 10 m WS (also seen in Fig. S1 meteorology comparison with ERA5 and MERRA-2) is well known, and the observed magnitude of the bias is largely consistent with previous studies (Zhang et al., 2016; Mues et al., 2018; Gunwani and Mohan, 2017; Wang et al., 2021).
380 First, this could be in part due to inaccurate land-surface parameterizations (such as roughness length or surface drag and urban canopy) yielding smaller friction velocities and stronger winds in the model. Second, it could also be due to unknown differences in heights of measured and modelled WS . However, the afternoon simulated WS are close to the observations, which suggests there are underlying weaknesses in nocturnal stable boundary layer decoupling in the model. The associated turbulent fluxes
385 and thermodynamic exchanges occurring in the atmospheric boundary layer are important for model simulated PBL and pollutant dispersal (Shen et al., 2023; Nelli et al., 2020). However, during the extreme pollution episode (30 October to 7 November) both model and observations agree on a reduction in WS (although with varying magnitudes) and a shift in WD. These changes highlight the role of stagnant meteorology in greatly enhancing the near-surface pollution lasting over a week.

390






Figure 3. Top to bottom: Comparisons of vertical profiles of temperature ($^{\circ}\text{C}$), virtual potential temperature (VPT, $^{\circ}\text{C}$) and wind speed (m s^{-1}) between the model (red) and radiosonde observations (black) for 8 sites at 00 UTC (5.30 IST) averaged for September – November 2016. The horizontal lines show the standard deviation in the day-to-day temporal variability during the comparison period.




Figure 4. Top to bottom: Comparisons of vertical profiles of temperature (°C), virtual potential temperature (VPT, °C) and wind speed (m s⁻¹) between the model (red) and radiosonde observations (black) for 8 sites at 12 UTC (17.30 IST) averaged for September – November 2016. The horizontal lines show the standard deviation in the day-to-day temporal variability during the comparison period.

395 3.2 Vertical Profiles

Figures 3 and 4 show averaged modelled and observed sounding profiles over individual RAOB sites (Fig. 1) for temperature (T), virtual potential temperature (VPT) and wind speed (WS) at 05.30 IST (00 UTC) and 17.30 IST (12 UTC), respectively. The corresponding summaries of statistical metrics are presented in Tables 3 and 4. The upper air meteorology and thermodynamic structure are crucial parameters of the atmosphere as they impact the transport and convective distribution of pollutants. Of all the meteorological quantities examined here, vertical profiles of T and VPT are represented best by the model, with correlations of $r \geq 0.95$ across all the sites and $r = 1.0$ for most of the sites at both times. At 05.30 IST, modelled T profiles show a warm bias of up to 1.5 °C at 6 sites and a cold bias of up to 2 °C at Delhi and Gwalior sites up to about 980 hPa (Fig. 3a), which gradually decreases with altitude. The model also captures the observed marked inversion near the surface in morning T and VPT profiles reasonably well at most sites. Agreement at 17.30 IST is even better (Fig. 4a): biases in modelled T profiles are less than 0.5 °C below 980 hPa at all sites except Ranchi and negligible aloft. Overall, across all sites, the average MB, NMB, and RMSE values are generally lower for VPT compared to T at both times (Tables 3 and 4).

The simulated WS vertical profiles have larger variations across most of the sites at both times as compared to T and VPT profiles (Figures 3c and 4c). Consistent with the 10 m WS comparisons, the model tends to overestimate WS vertically by up to 4 m s⁻¹ at 05.30 IST and up to 3 m s⁻¹ at 17.30 IST in the lower layers but better captures it aloft (above ~900 hPa) with only slight differences across all the sites (Fig. S4). Despite the considerable positive bias within the bottom layers, the model reproduces the observed higher WS at higher altitudes reasonably well, resulting in good correlations of $r \geq 0.77$ at 05.30 IST and $r \geq 0.95$ at 17.30 IST. As an exception, the modelled WS profiles are very well represented over the Patna site (in the east) during both times. The results here differ from those of Mohan and Bhati (2011), who noted increased deviation in simulated WS at higher altitudes over Delhi during the summer months.

The simulated RH profiles were also evaluated (Fig. S4) and show underestimations by up to 20 % in the lower layers of the model across most sites at both times, which decreases in magnitude at higher altitudes except at Gorakhpur. These biases vertically are generally more negative at 05.30 IST compared to 17.30 IST, indicating a dry bias in the early morning hours in the model, consistent with the ground observation comparisons.

VPT profiles are particularly useful in understanding the stability and turbulence of the atmosphere, which helps in the dilution of the pollutants within the mixed boundary layer. By accounting for moisture and temperature, a VPT profile indicates buoyancy and stability in the atmosphere and can be used to derive planetary boundary layer heights (Liu et al., 2019; Voogelezang and Holtslag, 1996). Figure 3 shows that, at all sites, observed and simulated temperature inversion layers close to the surface

430 **Table 3.** Summary of statistical comparison of modelled and observed 0.5.30 IST profiles derived from radiosonde data for the individual RAOB stations shown in Figure 1b averaged from September to November 2016. The statistical metrics used for comparison are mean bias (MB), normalized mean bias (NMB), mean absolute error (MAE), root mean square error (RMSE) and Pearson's correlation coefficient (*r*).

435

Station Name	MB	NMB	MAE	RMSE	<i>r</i>
Temperature (°C)					
Calcutta	-0.22	0.03	0.58	0.83	1.00
Delhi	0.14	-0.02	0.67	1.01	1.00
Gorakhpur	0.14	-0.02	1.14	1.68	1.00
Gwalior	-0.08	0.01	0.67	1.12	1.00
Jodhpur	-0.25	0.03	0.92	1.92	1.00
Lucknow	-0.81	0.10	1.54	7.70	0.96
Patna	-0.15	0.02	0.75	1.06	1.00
Ranchi	-1.08	0.14	1.70	7.56	0.96
VPT (°C)					
Calcutta	-0.31	-0.01	0.70	0.98	1.00
Delhi	0.09	0.00	0.74	1.09	1.00
Gorakhpur	-0.02	0.00	1.37	1.99	0.99
Gwalior	-0.16	0.00	0.78	1.29	1.00
Jodhpur	-0.37	-0.01	1.13	2.65	0.99
Lucknow	0.14	0.00	1.24	3.87	0.98
Patna	-0.27	-0.01	0.90	1.25	1.00
Ranchi	0.68	0.01	1.70	5.99	0.95
WS (m s⁻¹)					
Calcutta	-0.35	-0.04	1.34	2.02	0.96
Delhi	-0.29	-0.02	1.57	2.08	0.99
Gorakhpur	-0.78	-0.07	1.80	2.42	0.98
Gwalior	-0.33	-0.03	1.60	2.37	0.97
Jodhpur	-0.74	-0.06	1.82	2.41	0.98
Lucknow	0.15	0.01	2.37	4.92	0.89
Patna	-0.59	-0.06	1.53	2.13	0.98
Ranchi	0.58	0.07	2.65	5.33	0.77
RH (%)					
Calcutta	-1.09	-0.02	7.56	11.9	0.93
Delhi	-2.20	-0.08	6.23	10.0	0.92
Gorakhpur	-10.5	-0.21	15.4	19.4	0.88
Gwalior	-1.87	-0.06	7.21	10.9	0.93
Jodhpur	1.54	0.07	8.59	11.7	0.89
Lucknow	-2.99	-0.08	10.4	15.1	0.87
Patna	-0.84	-0.02	9.37	14.4	0.92
Ranchi	-3.12	-0.07	10.6	16.5	0.89

at 5.30 IST, demonstrating the typical formation of an urban nocturnal stable boundary layer. In contrast, at 17.30 IST (Fig. 4), both the observed and modelled VPTs exhibit a typical well-mixed late afternoon profile due to surface heating, with higher values of VPT near the surface (33–36 °C surface) that remains nearly constant up to about 850 hPa across most sites. The negligible biases and error statistics in T and VPT profiles (Tables 3 - 4) across all sites provide high confidence in model skill in simulating the thermodynamic structure of the atmosphere. This is an improvement on Mues et al. (2018), who reported larger biases in T profiles (up to 3 °C and 7 °C at 05.30 IST and 17.30 IST, respectively) at the Delhi site in winter and summer 2013. As noted in Section 3.1, and elsewhere (Mohan and Bhati, 2011; Gunwani and Mohan, 2017), errors in simulated WS are highly sensitive to local roughness length and model topography and are thus subject to greater noise. Given these limitations, we find the model performance statistics comparable to previous studies (Mohan and Bhati, 2011; Kumar et al., 2012b) and close to the benchmarks provided by Emery and Tai (2001).

4 Chemistry evaluation results

4.1 Ground-based PM_{2.5}

Figure 5 compares the modelled and measured daily averaged time series (left) and diurnal variability (right) of surface PM_{2.5} concentrations from hourly samples from September to November 2016. The observations are spatially averaged across 8 sites in Delhi and 12 sites across the rest of the domain (referred to as ‘Others’). The statistical summary is presented in Table 5. The model adequately captures the day-to-day variation of PM_{2.5} for October-November, when it is biased low, while it fails to reproduce the daily variability during September when it is strongly biased high. On average, during September, the model overestimates surface PM_{2.5} concentrations by more than a factor of two (NMB range: 1.69 to 1.91) across all the sites and underestimates in November by 26 % over Delhi and by 14 % over Others. Overall, the model and observed daily surface PM_{2.5} correlate reasonably well during October (Delhi: $r = 0.65$, Others: $r = 0.53$) and November (Delhi: $r = 0.76$, Others: $r = 0.66$). Correlation for these months is better across Delhi sites but shows relatively larger mean biases (+17.7 to – 73.2 $\mu\text{g m}^{-3}$) and NMBs (+0.13 to – 0.26) compared to Others. Additionally, the model tends to predict PM_{2.5} concentrations with a fairly broad range of monthly RMSE values (56.3 – 138 $\mu\text{g m}^{-3}$).

The spatially averaged diurnal cycle for modelled surface PM_{2.5} shows a pronounced diurnal trend matching observations for Delhi sites, while the diurnal cycle is less pronounced at Others sites. Generally, diurnal trends are in good agreement across all sites, although on average, the model tends to underpredict the afternoon dips and night-time peaks compared to the observations, indicating missing anthropogenic activities from the simplified diurnal emissions patterns derived from monthly estimates used in the model. The lack of a representation of a realistic diurnal activity cycle in the anthropogenic emissions highlights meteorology could be driving the modelled PM_{2.5} variation.

Although, this might partly be affected by the imperfectly represented diurnal variability of WS in the model (Section 3.1).

Table 4. Same as Table 3, but for 17.30 IST profiles.

475

Station Name	MB	NMB	MAE	RMSE	<i>r</i>
Temperature (°C)					
Calcutta	-0.16	0.02	0.57	0.82	1.00
Delhi	0.04	-0.01	0.63	0.85	1.00
Gorakhpur	-0.02	0.00	1.24	1.89	1.00
Gwalior	-0.17	0.02	0.68	1.15	1.00
Jodhpur	-0.11	0.01	0.74	1.02	1.00
Lucknow	0.01	0.00	0.95	1.73	1.00
Patna	0.06	-0.01	0.75	1.20	1.00
Ranchi	-1.22	0.18	2.15	8.45	0.96
VPT (°C)					
Calcutta	-0.24	0.00	0.68	1.00	1.00
Delhi	-0.02	0.00	0.74	0.97	1.00
Gorakhpur	-0.31	-0.01	1.50	2.19	0.99
Gwalior	-0.26	0.00	0.81	1.39	1.00
Jodhpur	-0.20	0.00	0.91	1.25	1.00
Lucknow	-0.12	0.00	1.16	2.13	0.99
Patna	-0.01	0.00	0.85	1.34	1.00
Ranchi	0.65	0.01	1.89	5.59	0.96
WS (m s⁻¹)					
Calcutta	-0.30	-0.04	1.30	1.78	0.97
Delhi	-0.24	-0.02	1.48	1.91	0.99
Gorakhpur	-0.71	-0.06	1.76	2.32	0.98
Gwalior	-0.27	-0.02	1.50	1.97	0.98
Jodhpur	-0.64	-0.06	1.76	2.27	0.98
Lucknow	-0.40	-0.03	1.81	2.42	0.98
Patna	-0.68	-0.06	1.49	2.01	0.98
Ranchi	-0.15	-0.02	1.88	2.74	0.95
RH (%)					
Calcutta	-2.13	-0.04	8.06	12.7	0.93
Delhi	-1.47	-0.06	6.73	10.6	0.90
Gorakhpur	-9.62	-0.20	14.9	18.8	0.86
Gwalior	-0.58	-0.02	7.80	12.0	0.90
Jodhpur	2.29	0.12	8.54	11.4	0.86
Lucknow	-2.12	-0.06	10.9	14.9	0.85
Patna	-0.96	-0.02	9.44	14.0	0.91
Ranchi	1.73	0.04	10.3	15.3	0.89




Figure 5. Time series of daily means (left) and mean diurnal cycles (right) of observed and modelled $PM_{2.5}$, averaged across 8 sites in Delhi and 12 sites over the rest of the domain (labelled ‘Others’) from September–November 2016. The shaded area in both panels shows standard deviation of the spatial variability of the model and measured $PM_{2.5}$. The locations of the ground measurement sites are shown in Figure 1b. The vertical dashed lines delineate the period of severe high pollution between 30 October and 7 November.




Figure 6. Time series of daily means (left) and mean diurnal cycles (right) of modelled individual $PM_{2.5}$ components averaged across 8 stations in Delhi and 12 stations over the rest of the domain (labelled ‘Others’) from September – November 2016. The individual species contribution abbreviations are: SOA (secondary organic aerosol), POA (primary organic aerosol), SO_4^{2-} (sulfate), NH_4^+ (ammonium), NO_3^- (nitrate), BC (black carbon). The vertical dashed lines delineate the period of severe high pollution between 30 October and 7 November.

During the 30 October –7 November pollution episode, both observations and model show the highest daily mean surface PM_{2.5} (Observed: 300 – 750 $\mu\text{g m}^{-3}$, Modelled: 150 – 420 $\mu\text{g m}^{-3}$) across Delhi, while relatively lower concentrations are seen across Others sites during this period (observed and modelled: < 200 $\mu\text{g m}^{-3}$) (Fig. 5). The observed daily mean PM_{2.5} concentrations exceed the 24-h average 2021 WHO air quality guideline of 15 $\mu\text{g m}^{-3}$ (WHO 2021) by nearly 50 times and the predicted concentrations exceed by nearly 28 times. The maximum negative differences (up to 350 $\mu\text{g m}^{-3}$) between the daily mean modelled and observed PM_{2.5} also occur during this episode. During this period, the observed hourly PM_{2.5} concentrations exceed 1000 $\mu\text{g m}^{-3}$ (mostly at night) at the Delhi US embassy site (in central Delhi) and exceed 800 $\mu\text{g m}^{-3}$ at all the sites across Delhi and two downwind stations in the lower IGP (Lucknow and Kanpur). The corresponding modelled hourly concentrations at these locations and times underestimate PM_{2.5} by a factor of 2-3 (380 – 520 $\mu\text{g m}^{-3}$), in part attributable to overestimated surface WS. One study characterising this 2016 high pollution episode over Delhi reported exceptionally high night-time mean PM_{2.5} concentrations of 2924 $\mu\text{g m}^{-3}$ on 30 October (Diwali festival night), 1520 $\mu\text{g m}^{-3}$ on 5 November, and daytime mean values of nearly 1500 $\mu\text{g m}^{-3}$ on 6 November (Sawhani et al., 2019). The modelled and observed daily average PM_{2.5} across downwind Others sites peaks (> 250 $\mu\text{g m}^{-3}$) only towards the end of the high pollution episode, suggesting a regional distribution of PM_{2.5} over time. The observed and simulated near-surface meteorology during this time over northern India shows stagnant conditions conducive for the build-up of pollutants: smaller WS (1-1.5 m s^{-1}), lower PBLH (< 500 m) and a nearly 2 - 3 °C drop in near-surface temperature leading to atmospheric inversion (Fig. 2). These stagnant conditions combined with regional and local anthropogenic emissions facilitate pollution accumulation within the shallow continental boundary layer over wider northern India. After the extreme pollution days (9 November onwards), the model captures the magnitude of daily PM_{2.5} variation well everywhere except for an observed peak across Delhi on 17 November.

4.2 Modelled PM_{2.5} Composition

The daily time series and average diurnal variability of modelled mean surface PM_{2.5} composition over observation sites in Delhi and Others are shown in Figure 6. Due to the lack of observed PM_{2.5} speciation data for this period, only modelled results are presented here. These are qualitatively compared with literature for other years as the aerosol loading over the Indian region exhibits stronger intra-annual variabilities than interannual variabilities (Conibear et al., 2018; Mhawish et al., 2021). The largest variations in daily PM_{2.5} components across all months are observed for secondary organic aerosol (SOA) and secondary inorganic aerosol (SIA) (sulfate, nitrate and ammonium) over all the sites. The concentration of fine dust particles dominates most evidently at the beginning of September and reduces to almost half in October and November but remains a non-negligible contributor to total PM_{2.5} on average (15 - 25 %) across all sites. The fine dust component is mainly responsible for the overestimations seen in modelled PM_{2.5} in September compared to the measurement. Another notable

Table 5. Statistical summary of comparisons of –modelled and observed PM_{2.5} concentrations derived from hourly data between September to November 2016 for Delhi (top) and Other stations (bottom). The statistical metrics are mean bias (MB), normalized mean bias (NMB), mean absolute error (MAE), root mean square error (RMSE) and Pearson’s correlation coefficient (*r*). *n* denotes number of available measurement stations in the group.

Month	MB	NMB	MAE	RMSE	<i>r</i>	Obs_mean	Mod_mean
PM_{2.5} (µg m⁻³)							
Delhi sites (<i>n</i> = 8)							
September	111	1.91	111	124	0.17	58.7	170
October	17.7	0.13	58.1	74.2	0.65	141	159
November	-73.2	-0.26	95.2	138	0.76	279	206
Others sites (<i>n</i> = 12)							
September	69.9	1.69	70.26	89.5	0.44	41.3	111
October	10.9	0.11	40.71	56.3	0.53	102	113
November	-23.8	-0.14	54.94	73	0.66	172	148

Table 6. Statistical summary of comparisons of concentrations (µg m⁻³) of PM_{2.5} and black carbon from the WRF-Chem model and MERRA-2 from September to November 2016. The statistical metrics are mean bias (MB), normalized mean bias (NMB), mean absolute error (MAE), root mean square error (RMSE) and Pearson’s correlation coefficient (*r*).

Month	MB	NMB	MAE	RMSE	<i>r</i>
PM_{2.5} (µg m⁻³)					
September	54	1.9	55.1	72	0.87
October	20	0.49	21.7	30	0.87
November	-8.4	-0.12	13.8	23	0.95
BC (µg m⁻³)					
September	0.52	0.65	0.57	0.93	0.91
October	0.24	0.19	0.44	0.79	0.91
November	-0.78	-0.28	0.89	1.42	0.92




Figure 7. Spatial distributions of monthly mean concentrations ($\mu\text{g m}^{-3}$) of a) $\text{PM}_{2.5}$ and b) black carbon from the WRF-Chem model and MERRA-2 for September to November 2016. The monthly mean $\text{PM}_{2.5}$ at the measurement sites are shown in circles in a).

change is in the nitrate component which dramatically peaks during the high pollution period, together with SOA, ammonium, and primary aerosols (OC, BC). The modelled peaks in $\text{PM}_{2.5}$ and its components largely follow the observed $\text{PM}_{2.5}$ trend (Oct - Nov period) which highlights the model's skill in representing the diversity of aerosols during dramatic shifts in surface particle pollution and is more clearly seen across Delhi sites than Others. Among SIA, the $\text{PM}_{2.5}$ composition in November is dominated by nitrate aerosols (10 - 30 %) which are comparable to reported measurements. For example, a high nitrate fraction (20 - 27 %) in post-monsoon months has been reported in various measurement studies over India (Ram and Sarin, 2011; Schnell et al., 2018; Patel et al., 2021; Talukdar et al., 2021). The average modelled BC contribution over Delhi during September ($3 \mu\text{g m}^{-3}$), October ($8.2 \mu\text{g m}^{-3}$) and November ($13.2 \mu\text{g m}^{-3}$) are comparable to the measured EC (assumed to be equivalent to modelled BC) concentrations ($\sim 3 \mu\text{g m}^{-3}$, $\sim 6 \mu\text{g m}^{-3}$ and $\sim 12 \mu\text{g m}^{-3}$, respectively) reported by Sharma et al. (2018). The dominance of secondary particle contribution to modelled $\text{PM}_{2.5}$ during post-monsoon months is fully consistent with other studies (Gani et al., 2019; Talukdar et al., 2021) although the relative abundance is lower. The diurnal variation of $\text{PM}_{2.5}$ components over Delhi show more pronounced dips in primary and secondary inorganics, suggesting influence of local emissions while the fine dust component remains relatively stable, suggesting both local and natural non-local emissions influence.




Figure 8. Spatial variation of monthly mean AOD at 550 nm derived from the model and MODIS sampled at local overpass times of 10.30 (Terra) and 13.30 (Aqua) for September to November 2016. Absolute differences of model minus satellite AOD are shown in the bottom row.




Figure 9. Scatterplots of monthly averaged model versus MODIS-derived AOD at 550 nm for the months (from left to right) September, October, and November 2016. The 2:1, 1:1 and 1:2 lines (red dashed lines), the best-fit line (black line) and Pearson's correlation coefficient r are also shown for each month.

535 4.3 Comparison of PM_{2.5} and Black Carbon distribution with reanalysis products

Figure 7 compares the monthly averaged spatial distribution of WRF-Chem modelled and MERRA-2 reanalysis derived surface PM_{2.5} and BC concentrations. The corresponding domain-averaged performance statistics are summarised in Table 6. The overall spatial agreement between the model and MERRA-2 is excellent for both PM_{2.5} and BC ($r > 0.87$, Fig. S6). However, on a regional scale, 540 the modelled PM_{2.5} is biased high over parts of arid western India and eastern Pakistan in September, resulting in a domain-wide NMB of 1.9. The model shows a stronger west-east gradient in PM_{2.5} than MERRA-2 with the highest modelled concentrations of $> 250 \mu\text{g m}^{-3}$ in the western and north-western regions. Agreement between the model and MERRA-2 improves for October-November.

The high simulated PM_{2.5} loading over some parts of north-western India during September is most 545 likely due to erroneous dust uplift by overestimated winds from the Thar Desert in the west (Fig. S6), the major seasonal natural dust source region (Bali et al., 2021; Kumar et al., 2018a). This overestimation could further be enhanced by the underestimation of dust deposition in the model arising from a dry bias over the land region in the domain (Ratnam and Kumar, 2005; Conibear et al., 2018). The notable change in modelled PM_{2.5} over the dust source region along the western borders from 550 September to November shows a strong seasonality in dust emissions in the model. Compared to WRF-Chem, MERRA-2 shows a slightly better comparison with monthly mean surface PM_{2.5} (Fig. 7a) for individual monitoring sites with smaller differences between model-measured mean than WRF-Chem (especially for September).

In contrast, the highest BC concentrations occur along the IGP for all the months and increases from 555 September to November (Fig. 7b). During October and November, the northwest and eastern parts of the IGP exhibit the highest PM_{2.5} and BC concentrations in both datasets. Compared to MERRA-2, modelled BC shows more distinguishable spatial features including localised hotspots coinciding with densely populated major metropolitan and industrial cities with clusters of coal-fired power plants (Singh et al., 2018). For instance, conspicuous localised regions appear over dense urban centres like 560 Ahmedabad, Delhi, Kolkata, the steel industrial city of Jamshedpur, Raipur with heavy mining, Singrauli with ore-processing industries in the upper central domain, and Jharia coal belts in the east having clusters of coal-fired power plants. Overall, the spatial variabilities of BC and PM_{2.5} are quite similar in both WRF-Chem and MERRA-2 with WRF-Chem estimating slightly lower PM_{2.5} and BC in November over the majority of the IGP except over Delhi.

565




Figure 10. Time series (left panel) and scatter plots (right panel) of modelled and AERONET daily averaged AOD at 550 nm over the 4 AERONET stations shown in Figure 1b for the period September to November 2016.

4.4 Evaluation of aerosol optical depth with satellite and AERONET observations

Figure 8 compares WRF-Chem simulated and MODIS (Aqua) retrieved monthly averaged distribution of AOD at 550 nm. The unitless quantity AOD is a measure of particle extinction within the atmospheric column from the surface to the top atmosphere and provides a useful spatial estimate of particle loading using satellite instruments. The spatial distributions of modelled and MODIS AOD agree well for all months ($r \geq 0.72$ Table 7) although regional biases similar to the MERRA-2 comparisons occur over northern and western parts of the domain. As with MERRA-2 $PM_{2.5}$ comparisons, during September the model captures well the high AOD (up to 1.2) over north-western India and along the borders with north-east Pakistan but predicts higher AOD over the western arid region (Fig. 8), indicated by the overall NMB of 0.69. The statistical evaluation metrics for all the months (Table 7) show there is a good overall agreement between modelled and satellite AOD which gradually improves from September to November. In both model and satellite data, AOD values are generally low (<0.5) outside of the broader IGP region in all the months. Although the satellite AOD shows higher spatial variability, a good spatial correlation exists between the two datasets in October-November ($r = 0.80$ and 0.86 , respectively) (Fig. 9). The domain averaged modelled AOD (0.39 and 0.34, respectively) during these months are comparable to satellite retrieved AOD (0.32 and 0.34, respectively). Despite the overall underestimations during the biomass burning period of (mid-October to mid-November), the model

captures high AOD values over some small, localised parts in Punjab and Haryana in northern India and north-eastern Pakistan, although with slightly lower magnitudes. The higher AOD along the entire IGP region is more apparent from the satellite observations in November, which show AOD values reaching ~ 2.0 (underestimated in the model by about 10 %) over parts of Punjab in the north and Uttar Pradesh and Bihar in the east (AOD >1.8). Interestingly, the regional hotspots along the IGP region, over eastern Uttar Pradesh and eastern Bihar as observed in modelled PM_{2.5} maps during October- November are evident in MODIS AOD distribution but less discernible in modelled AOD maps (Fig. 8). It is important to note that the MODIS satellite overpass times of 10.30 and 13.30 local time limits comparisons to the afternoon each day. Therefore, it is the modelled meteorological conditions typical to daytime (deep PBL height, increased WS) that affect the modelled AOD column. In a similar model set-up over northern India, Roozitalab et al. (2021) and Kulkarni et al. (2020) found comparable estimates of modelled AOD distribution during the 2017 post-monsoon high pollution event.

To further evaluate model skill in predicting the optical properties of aerosols, the modelled daily averaged AOD at 550 nm is compared in Figure 10 against the four AERONET sites (Fig. 1b) in the study domain. There are missing data at all the sites with Kanpur in the east and Jaipur in the west (both dense urban locations) having the most data coverage. The daily variabilities of AOD comparison with point observations show similar trends as previously noted for comparison with satellite AOD and ground-based and MERRA-2 PM_{2.5} comparisons. The model evaluation against AERONET AOD largely agrees with the PM_{2.5} evaluations including higher disparities seen for September with a positive MB (0.02 to 0.43) across all the sites. However, the high daily averaged AERONET AOD (>1.0) at all sites during the high pollution event at the start of November is captured reasonably well by the model except in Lahore, a large city in eastern Pakistan, where the model underestimates AOD the most. Of the four sites, crop residue burning occurs in Lahore (Kulkarni et al., 2020), which is also situated close to other biomass-burning regions of northwestern India. This AERONET site shows the highest observed (~3.0) and modelled (~2.0) AOD values during the high pollution episode.

Additionally, to check for consistency between satellite and ground measurements, the time series of satellite, AERONET and modelled AOD at 550 nm at the four observation locations are shown in Figure 11. To compare the three datasets, the data points corresponding to the local overpass time of MODIS are selected from the hourly AERONET and WRF-Chem datasets. The satellite AOD generally matches more closely with AERONET at lower values and misses the magnitude of high AOD during high pollution days. Earlier studies have attributed the inaccuracies in MODIS AOD retrievals due to dense haze hanging over north India and the IGP region during severe pollution days (Mhawish et al., 2022). The modelled AOD captures the hourly AOD trend quite well but also underestimates AOD in absolute magnitude during high pollution days across the sites. Overall, the modelled AOD agrees well with satellite and ground observations during October and November despite some underestimations in absolute magnitudes.

620 **4.5 Discussion**

The discrepancies in model-observation particulate matter comparisons for September have also been noted in other studies for India and suggest inaccuracies in modelling moisture transport during the monsoon season, which affects particle deposition and washout (Conibear et al., 2018; Mogno et al., 2021). Furthermore, in 2016, almost all the ground stations were in urban locations of the IGP region, which prevents the evaluation of the model at more spatially representative rural locations. In addition, nearly all the measurement sites are in or near dense urban areas with heavy influences from traffic and local anthropogenic activities (for example, trash burning and residential cooking), which may not be fully reflected in the monthly anthropogenic emission inputs. The sudden jumps in particulate matter during an extreme pollution event are especially difficult to capture within the model (despite adequate meteorological fields) without updated emissions estimates and knowledge of dynamic local activity data (for instance, diurnal activity profiles specific to Indian regions). For example, residential emissions are a major contributor to poor air quality in rural and suburban areas in northern India, with an estimated 16 % to 80 % contribution towards SOA components of PM_{2.5} (Rooney et al., 2019).




Figure 11. Time series of MODIS-retrieved (green), modelled (red) and AERONET (blue) AOD at 550 nm sampled at 13.30 IST over the 4 AERONET stations shown in Fig. 1b: from top to bottom, Jaipur, Kanpur Gandhi College and Lahore.

635

Table 7. Statistical summary of comparisons modelled and observed AOD at 550 nm derived from MODIS and at the 4 AERONET stations at an hourly temporal resolution between September to November 2016. The statistical metrics are mean bias (MB), normalized mean bias (NMB), mean absolute error (MAE), root mean square error (RMSE) and Pearson’s correlation coefficient (*r*).

640

Month	MB	NMB	MAE	RMSE	<i>r</i>
AOD (MODIS)					
September	0.25	0.69	0.28	0.34	0.72
October	0.06	0.20	0.11	0.15	0.80
November	0.00	-0.01	0.09	0.13	0.86
AERONET					
Jaipur					
September	0.43	0.96	0.43	0.45	0.38
October	0.10	0.31	0.14	0.17	0.04
November	-0.03	-0.06	0.17	0.26	0.83
Kanpur					
September	0.30	0.66	0.32	0.37	0.60
October	-0.01	-0.02	0.19	0.25	0.64
November	-0.15	-0.21	0.20	0.26	0.72
Gandhi College					
September	0.02	0.04	0.22	0.27	-0.08
October	-0.03	-0.05	0.13	0.17	0.69
Lahore					
September	0.39	0.49	0.49	0.57	0.15
October	-0.14	-0.13	0.38	0.50	0.26
November	-0.37	-0.37	0.44	0.65	0.75

645

650

655

660

665

Diurnally, the nocturnal biases in wind speeds in the model allow for an increased dispersal of the pollutants near the surface leading to underestimated PM_{2.5} concentrations during the night. Furthermore, the inaccuracies in simulating individual fractions of total PM_{2.5} also add to the observed model biases; for example, high contributions from dust in the MOSAIC scheme could lead to overestimations (Georgiou et al., 2018). Overestimated modelled dust aerosols were also observed by Kalenderski et al. (2013) and Zhao et al. (2010), who tuned the online dust emission flux calculation based on region specific AERONET measurements for a dust event and found modelled AOD estimates to improve. The lack of aqueous-phase chemistry in our model framework further adds some biases in reproducing accurate amounts of secondary aerosol components of PM (such as SO₄²⁻, NH₄⁺, and NO₃⁻) and their subsequent scavenging by aqueous chemistry in the cloud or water droplets (Tuccella et al., 2012; Balzarini et al., 2015). Furthermore, underestimations in modelled PM_{2.5} concentrations across Delhi could also be due to the lack of input emissions of hydrogen chloride (HCl) gas, typically from seasonal local rubbish and crop residue burning, which adds substantial chloride aerosols to total PM_{2.5} by its partitioning between gas and aerosol phases (Cash et al., 2021; Lalchandani et al., 2022; Pawar

et al., 2023). The temporal mismatch due to changes in emissions between the anthropogenic emissions inventory –(2010) and the simulated year (2016) also contributes to the likely underestimation in the modelled PM_{2.5} and its components for our study period. Notably, a few studies

670 also report positive seasonal biases in simulating surface and column concentrations of trace gases like NO_x (Kumar et al., 2012a) and concentrations of SO₂ (Conibear et al., 2018) over urban areas in India . These biases further contribute to uncertainties in simulating reactive trace gas and secondary PM_{2.5} subspecies.

Significant post-harvest crop residue burning takes place in north-western states of India from late
675 October to mid-November (Jethva et al., 2019), which impacts the air quality locally as well as in downwind regions of central and eastern IGP (Bhardwaj et al., 2016; Kanawade et al., 2020; Kulkarni et al., 2020; Singh et al., 2021; Mhawish et al., 2022; Govardhan et al., 2023a). Other uncertainties in simulating PM_{2.5} concentrations arise from errors in scaling biomass burning emissions estimates, which largely depend on the limited number of daily satellite-based retrievals and are sometimes
680 compromised by dense smoke from fires being misrepresented as cloud cover in the detection algorithm (Cusworth et al., 2018). In their study, Singh et al. (2021) report the annual mean contribution of biomass burning to PM_{2.5} over India to be 8%, but with a strong seasonal dependence (up to 39 % in October-November in Delhi). As previously discussed in the literature, MODIS fire detection is susceptible to missing small fires like agricultural burning (Cusworth et al., 2018; Roozitalab et al.,
685 2021). In addition to the surface measurements, comparisons with MERRA-2 products highlight a good agreement between the WRF-Chem simulations and the reanalysis approach of employing satellite data assimilations. Navinya et al. (2020) and others, however, find MERRA-2 to underestimate simulated PM_{2.5} over India in comparison to the measurements. As noted in Section 2.5, several factors need to

be considered when comparing the WRF-Chem aerosol concentrations with MERRA-2 reanalysis products. These include (but are not limited to) a coarser resolution of MERRA-2, limited observations data available for assimilation, input anthropogenic emissions lacking seasonal variability, the GOCART chemistry scheme missing nitrate and SOA treatment and the quality of aerosol transport dynamics from the GEOS-5/GOCART model wherein the reanalysed aerosol mass concentrations are not directly constrained by the observations as is the case with the AOD (Buchard et al., 2017). These
690 points, therefore, suggest the aerosol assimilation capabilities used in combination with the chemistry representing complex aerosol processes is potentially a more accurate way to predict air quality across polluted regions such as northern India.

Overall, the evaluation of the WRF-Chem simulated chemistry demonstrates adequate performance during October and November for PM_{2.5} and is found to be suitable for investigating the atmospheric
700 dynamics during extreme pollution events. The modelled results presented here, and in other studies of pollution episodes and aerosol climatology over India, clearly show that the October-November period has higher aerosol loading over most of the domain compared to other times of the year. A mix of

factors like emission patterns, meteorology shifts and topography intensify the existing high pollution levels in some parts of India (Kulkarni et al., 2020; Kumar et al., 2018a; Mhawish et al., 2022; Kanawade et al., 2020). Sawlani et al. (2019) and Kanawade et al. (2020) attribute the 2016 haze episode to a mix of coinciding factors: local emissions from fireworks, enhanced fire counts from agricultural crop residue burning in northwestern states, stagnant conditions resulting from low temperatures, shallow PBL, weaker northwesterly winds, and high ambient RH. The crop residue burning in 2016 (over Punjab, Haryana and Uttar Pradesh in northwest India and Pakistan) detected by combined VIIRS and MODIS sensors reveal higher total burning events by up to 30% and 41% compared to 2017 and 2018, respectively (Chhabra et al., 2019). Similar high pollution events have been reported during post-monsoon months in later years (Dekker et al., 2019; Kulkarni et al., 2020; Takigawa et al., 2020; Roozitalab et al., 2021; Beig et al., 2021; Mhawish et al., 2022). Additionally, a few studies also report a layer of biomass-burning smoke aerosols at 2-3 km altitude above the IGP region using CALIPSO (Cloud–Aerosol Lidar and Infrared Pathfinder Satellite Observations) retrievals (Shaik et al., 2019; Kumar et al., 2018b).

5. Conclusions

We comprehensively compare the WRF-Chem v4.2.1 modelled meteorology and aerosol chemistry with a wide range of observational data that includes ground-based, satellite and reanalysis products over northern India. The simulations are performed at a spatial resolution of 12 km for the 2016 monsoon (September) to post-monsoon (October–November) transition, with a focus on the severe haze pollution episode from 30 October to 7 November.

The meteorological fields show strong seasonal and spatial variability over the IGP region with a marked decrease in temperature, WS, and PBLH from monsoon to post-monsoon, most notably for PBLH. Overall, we find that the model accurately represents meteorology during the afternoon hours. The surface daily and diurnal trend in temperature is best reproduced by the model, followed by relative humidity, with negligible biases across all sites. In contrast, daily mean model wind speed is widely biased high (by $0.5 - 0.8 \text{ m s}^{-1}$) largely due to strong night-time overestimations (up to 1.5 m s^{-1}), while the afternoon WS is reasonably reproduced by the model. This suggests a potential model failure in surface layer decoupling at night.

Comparison of upper air meteorology with radiosonde profiles shows negligible biases and excellent correlations for temperature and virtual potential temperature ($r > 0.9$) across all sites. The model overestimates wind speed in the lowest layers, consistent with surface observation comparisons whilst matching well with observed WS aloft. In comparison to MERRA-2 reanalysis products, modelled PBLH generally has negative mean bias of $> 25 \%$ in all the months but agrees well spatially.

Modelled and observed PM_{2.5} concentrations show good agreement (except during September) with overall slightly better correlations for eight sites averaged across Delhi ($r > 0.6$) compared to twelve sites across the remaining domain ($r > 0.5$). In September, model concentrations show large biases due to overestimation in dust generation over the western arid region and possible long-range transport across the measurement sites.

740

The model simulates the high pollution episode with notable peaks in daily mean PM_{2.5} concentrations but underestimates the exceptionally high observed daily PM_{2.5} (300 – 750 $\mu\text{g m}^{-3}$) by a factor of 2-3. Despite the accurate representation of the vertical temperature gradient, the model underestimates high surface PM_{2.5} concentrations due to stronger simulated WS favouring the dispersion of the surface pollutants, together with uncertainties in the emissions inventories. Both the model and surface measurements show that Delhi experiences the highest PM_{2.5} concentrations during the severe pollution episode followed by regional dispersal of pollutants downwind. During the episode, daily simulated anthropogenic PM_{2.5} composition comprised high fractions of nitrate (5 - 25 %) and secondary organic aerosols (10 - 20 %), consistent with previous measurement and modelling studies. The contribution of BC and primary organic matter to the total simulated PM_{2.5} mass also increases in November.

745

750

Comparison with MERRA-2 reanalysis data shows the spatiotemporal distribution of surface PM_{2.5} to have systematic high biases in September along the dry western region of the domain and low bias in October-November in the IGP region. However, the model captures quite well the high PM_{2.5} and BC concentrations over the IGP, including Delhi and upwind biomass burning regions during November. Variability in modelled AOD compared with satellite retrievals from MODIS is captured very well with $r \geq 0.8$ in October-November. The model likewise compares well with ground-based AERONET measurements of daily AOD ($r \geq 0.7$) across all sites except during September.

755

Our evaluations consistently reveal the best performance of the model in simulating PM_{2.5} and BC concentrations is for November followed by October, with model underestimations largely stemming from the extreme episodic nature of the pollution event. The lack of measurement data for individual PM_{2.5} components and the limited spatial coverage of measurement sites restricts the extent of the evaluation of this period. Overall, however, the model is found adequate for investigation of the vertical distribution of particle components and their interactions with meteorology through sensitivity simulations, including investigation of different emissions datasets, which forms part of our future work. Our results also suggest that improved diurnal characterisation of boundary layer processes could considerably enhance the model performance over this region.

760

765

Code and data availability

770 All the data sets used for comparison and source codes for model simulations are openly available.
WRF-Chem source code can be obtained from https://www2.mmm.ucar.edu/wrf/users/download/get_source.html. The ERA5 input data were
downloaded from <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels?tab=overview>. The chemical boundary conditions from MOZART are available at
775 <https://www2.acom.ucar.edu/gcm/mozart>. All the emissions inputs and pre-processor tools were
obtained from <https://www2.acom.ucar.edu/wrf-chem/wrf-chem-tools-community>. The links for
openly available ground, satellite and reanalysis datasets used for evaluation are provided in Table
S2.

780 Author contributions

PA, DSS and MRH conceptualised the study. PA compiled the measurement datasets performed formal
model simulations, and data analyses, curated the data and wrote the text with discussions and
supervision by MRH and DSS. DSS and MRH edited and commented on the text.

Competing interests. The authors declare that they have no conflict of interest.

785 Acknowledgments

This work was carried out on the Cirrus UK National Tier-2 HPC Service at EPCC
(<http://www.cirrus.ac.uk>) funded by the University of Edinburgh and EPSRC (EP/P020267/1). We
acknowledge the WRF-Chem community and the Atmospheric Chemistry Observations and Modeling
Lab (ACOM) of NCAR for providing the preprocessor tools {mozbc, fire_emis, anthro_emis,
790 bio_emiss} used in this study (<https://www2.acom.ucar.edu/wrf-chem/wrf-chem-tools-community> last
accessed 29 May 2023). We also acknowledge the use of data provided by GHSL - Global Human
Settlement Layer (<https://ghsl.jrc.ec.europa.eu/index.php> last accessed 29 May 2023). The geographical
maps were downloaded from https://github.com/IDFCInstitute/IndiaMap_Data (last accessed 29 May
2023). PA is thankful to Alaa Mhawish for providing the MODIS data. The use of open software
795 packages and python libraries are also gratefully acknowledged.

Financial support

PA acknowledges UoE scholarships (Principal's Career Development Scholarships and Edinburgh
Global Research Scholarship). DSS acknowledges support from Natural Environment Research
Council (grant no. NE/S009019/1).

800

References

- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, <https://doi.org/10.5194/acp-11-4039-2011>, 2011.
- 805 Ångström, A.: The parameters of atmospheric turbidity, *Tellus*, 16, 64–75, <https://doi.org/10.3402/tellusa.v16i1.8885>, 1964.
- Babu, S. S., Moorthy, K. K., Manchanda, R. K., Sinha, P. R., Satheesh, S. K., Vajja, D. P., Srinivasan, S., and Kumar, V. H. A.: Free tropospheric black carbon aerosol measurements using high altitude balloon: Do BC layers build “their own homes” up in the atmosphere?: FREE TROPOSPHERIC
810 BLACK CARBON AEROSOL, *Geophys. Res. Lett.*, 38, n/a–n/a, <https://doi.org/10.1029/2011GL046654>, 2011.
- Bali, K., Dey, S., and Ganguly, D.: Diurnal patterns in ambient PM_{2.5} exposure over India using MERRA-2 reanalysis data, *Atmospheric Environment*, 248, 118180, <https://doi.org/10.1016/j.atmosenv.2020.118180>, 2021.
- 815 Balzarini, A., Pirovano, G., Honzak, L., Žabkar, R., Curci, G., Forkel, R., Hirtl, M., San José, R., Tuccella, P., and Grell, G. A.: WRF-Chem model sensitivity to chemical mechanisms choice in reconstructing aerosol optical properties, *Atmospheric Environment*, 115, 604–619, <https://doi.org/10.1016/j.atmosenv.2014.12.033>, 2015.
- Beig, G., Srinivas, R., Parkhi, N. S., Carmichael, G. R., Singh, S., Sahu, S. K., Rathod, A., and Maji, S.: Anatomy of the winter 2017 air quality emergency in Delhi, *Science of The Total Environment*, 681, 305–311, <https://doi.org/10.1016/j.scitotenv.2019.04.347>, 2019.
- 820 Beig, G., Sahu, S. K., Rathod, A., Tikle, S., Singh, V., and Sandeepan, B. S.: Role of meteorological regime in mitigating biomass induced extreme air pollution events, *Urban Climate*, 35, 100756, <https://doi.org/10.1016/j.uclim.2020.100756>, 2021.
- 825 Bharali, C., Nair, V. S., Chutia, L., and Babu, S. S.: Modeling of the Effects of Wintertime Aerosols on Boundary Layer Properties Over the Indo Gangetic Plain, *Journal of Geophysical Research: Atmospheres*, 124, 4141–4157, <https://doi.org/10.1029/2018JD029758>, 2019.
- Bhardwaj, P., Naja, M., Kumar, R., and Chandola, H. C.: Seasonal, interannual, and long-term variabilities in biomass burning activity over South Asia, *Environ Sci Pollut Res*, 23, 4397–4410, <https://doi.org/10.1007/s11356-015-5629-6>, 2016.
- 830 Bisht, D. S., Tiwari, S., Dumka, U. C., Srivastava, A. K., Safai, P. D., Ghude, S. D., Chate, D. M., Rao, P. S. P., Ali, K., Prabhakaran, T., Panickar, A. S., Soni, V. K., Attri, S. D., Tunved, P., Chakrabarty, R. K., and Hopke, P. K.: Tethered balloon-born and ground-based measurements of black carbon and particulate profiles within the lower troposphere during the foggy period in Delhi, India, *Science of The Total Environment*, 573, 894–905, <https://doi.org/10.1016/j.scitotenv.2016.08.185>, 2016.
- 835 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment: BLACK CARBON IN THE CLIMATE SYSTEM, *J. Geophys. Res. Atmos.*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- 840

- 845 Brooks, J., Liu, D., Allan, J. D., Williams, P. I., Haywood, J., Highwood, E. J., Kompalli, S. K., Babu, S. S., Satheesh, S. K., Turner, A. G., and Coe, H.: Black Carbon physical and optical properties across northern Indiaduring pre-monsoon and monsoon seasons, *Aerosols/Field Measurements/Troposphere/Chemistry* (chemical composition and reactions), <https://doi.org/10.5194/acp-2019-505>, 2019.
- 850 Buchard, V., da Silva, A. M., Randles, C. A., Colarco, P., Ferrare, R., Hair, J., Hostetler, C., Tackett, J., and Winker, D.: Evaluation of the surface PM_{2.5} in Version 1 of the NASA MERRA Aerosol Reanalysis over the United States, *Atmospheric Environment*, 125, 100–111, <https://doi.org/10.1016/j.atmosenv.2015.11.004>, 2016.
- 855 Buchard, V., Randles, C. A., Da Silva, A. M., Darmenov, A., Colarco, P. R., Govindaraju, R., Ferrare, R., Hair, J., Beyersdorf, A. J., Ziemba, L. D., and Yu, H.: The MERRA-2 Aerosol Reanalysis, 1980 Onward. Part II: Evaluation and Case Studies, *J. Climate*, 30, 6851–6872, <https://doi.org/10.1175/JCLI-D-16-0613.1>, 2017.
- 860 Cash, J. M., Langford, B., Di Marco, C., Mullinger, N. J., Allan, J., Reyes-Villegas, E., Joshi, R., Heal, M. R., Acton, W. J. F., Hewitt, C. N., Misztal, P. K., Drysdale, W., Mandal, T. K., Shivani, Gadi, R., Gurjar, B. R., and Nemitz, E.: Seasonal analysis of submicron aerosol in Old Delhi using high-resolution aerosol mass spectrometry: chemical characterisation, source apportionment and new marker identification, *Atmospheric Chemistry and Physics*, 21, 10133–10158, <https://doi.org/10.5194/acp-21-10133-2021>, 2021.
- 865 Chhabra, A., Sehgal, V. K., Dhakar, R., Jain, N., and Verma, R.: MONITORING OF ACTIVE FIRE EVENTS DUE TO PADDY RESIDUE BURNING IN INDO-GANGETIC PLAINS USING THERMAL REMOTE SENSING, *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.*, XLII-3/W6, 649–657, <https://doi.org/10.5194/isprs-archives-XLII-3-W6-649-2019>, 2019.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric Aerosol Optical Thickness from the GOCART Model and Comparisons with Satellite and Sun Photometer Measurements, *JOURNAL OF THE ATMOSPHERIC SCIENCES*, 59, 23, 2002.
- 870 Chow, J. C., Lowenthal, D. H., Chen, L.-W. A., Wang, X., and Watson, J. G.: Mass reconstruction methods for PM_{2.5}: a review, *Air Qual Atmos Health*, 8, 243–263, <https://doi.org/10.1007/s11869-015-0338-3>, 2015.
- 875 Conibear, L., Butt, E. W., Knote, C., Arnold, S. R., and Spracklen, D. V.: Residential energy use emissions dominate health impacts from exposure to ambient particulate matter in India, *Nat Commun*, 9, 617, <https://doi.org/10.1038/s41467-018-02986-7>, 2018.
- Cusworth, D. H., Mickley, L. J., Sulprizio, M. P., Liu, T., Marlier, M. E., DeFries, R. S., Guttikunda, S. K., and Gupta, P.: Quantifying the influence of agricultural fires in northwest India on urban air pollution in Delhi, India, *Environ. Res. Lett.*, 13, 044018, <https://doi.org/10.1088/1748-9326/aab303>, 2018.
- 880 Dekker, I. N., Houweling, S., Pandey, S., Krol, M., Röckmann, T., Borsdorff, T., Landgraf, J., and Aben, I.: What caused the extreme CO concentrations during the 2017 high-pollution episode in India?, *Atmos. Chem. Phys.*, 19, 3433–3445, <https://doi.org/10.5194/acp-19-3433-2019>, 2019.
- 885 Dhaka, S. K., Chetna, Kumar, V., Panwar, V., Dimri, A. P., Singh, N., Patra, P. K., Matsumi, Y., Takigawa, M., Nakayama, T., Yamaji, K., Kajino, M., Misra, P., and Hayashida, S.: PM_{2.5} diminution and haze events over Delhi during the COVID-19 lockdown period: an interplay between the baseline pollution and meteorology, *Sci Rep*, 10, 13442, <https://doi.org/10.1038/s41598-020-70179-8>, 2020.

- Durre, I., Vose, R. S., and Wuertz, D. B.: Overview of the Integrated Global Radiosonde Archive, *Journal of Climate*, 19, 53–68, <https://doi.org/10.1175/JCLI3594.1>, 2006.
- 890 Emery, C. and Tai, E.: Enhanced Meteorological Modeling and Performance Evaluation for Two Texas Ozone Episodes, 2001.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., Laepple, T., Orlando, J., Tie, X., Tyndall, G., Wiedinmyer, C., Baughcum, S. L., and Kloster, S.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43–67, <https://doi.org/10.5194/gmd-3-43-2010>, 2010.
- 895 Emmons, L. K., Schwantes, R. H., Orlando, J. J., Tyndall, G., Kinnison, D., Lamarque, J.-F., Marsh, D., Mills, M. J., Tilmes, S., Bardeen, C., Buchholz, R. R., Conley, A., Gettelman, A., Garcia, R., Simpson, I., Blake, D. R., Meinardi, S., and Pétron, G.: The Chemistry Mechanism in the Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001882, <https://doi.org/10.1029/2019MS001882>, 2020.
- 900 Fast, J. D., Gustafson Jr., W. I., Easter, R. C., Zaveri, R. A., Barnard, J. C., Chapman, E. G., Grell, G. A., and Peckham, S. E.: Evolution of ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006721>, 2006.
- 905 Friedl, M. A., McIver, D. K., Hodges, J. C. F., Zhang, X. Y., Muchoney, D., Strahler, A. H., Woodcock, C. E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F., and Schaaf, C.: Global land cover mapping from MODIS: algorithms and early results, *Remote Sensing of Environment*, 83, 287–302, [https://doi.org/10.1016/S0034-4257\(02\)00078-0](https://doi.org/10.1016/S0034-4257(02)00078-0), 2002.
- 910 Gani, S., Bhandari, S., Seraj, S., Wang, D. S., Patel, K., Soni, P., Arub, Z., Habib, G., Hildebrandt Ruiz, L., and Apte, J. S.: Submicron aerosol composition in the world's most polluted megacity: the Delhi Aerosol Supersite study, *Atmospheric Chemistry and Physics*, 19, 6843–6859, <https://doi.org/10.5194/acp-19-6843-2019>, 2019.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *J. Climate*, 30, 5419–5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- 915 Georgiou, G. K., Christoudias, T., Proestos, Y., Kushta, J., Hadjinicolaou, P., and Lelieveld, J.: Air quality modelling in the summer over the eastern Mediterranean using WRF-Chem: chemistry and aerosol mechanism intercomparison, *Atmospheric Chemistry and Physics*, 18, 1555–1571, <https://doi.org/10.5194/acp-18-1555-2018>, 2018.
- 920 Ghosh, S., Dey, S., Das, S., Riemer, N., Giuliani, G., Ganguly, D., Venkataraman, C., Giorgi, F., Tripathi, S. N., Ramachandran, S., Rajesh, T. A., Gadhavi, H., and Srivastava, A. K.: Towards an improved representation of carbonaceous aerosols over the Indian monsoon region in a regional climate model: RegCM, *Geosci. Model Dev.*, 16, 1–15, <https://doi.org/10.5194/gmd-16-1-2023>, 2023.
- 925 Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S.-J.: Sources and distributions of dust aerosols simulated with the GOCART model, *Journal of Geophysical Research: Atmospheres*, 106, 20255–20273, <https://doi.org/10.1029/2000JD000053>, 2001.

- 930 Govardhan, G., Satheesh, S. K., Moorthy, K. K., and Nanjundiah, R.: Simulations of black carbon over the Indian region: improvements and implications of diurnality in emissions, *Atmos. Chem. Phys.*, 19, 8229–8241, <https://doi.org/10.5194/acp-19-8229-2019>, 2019.
- Govardhan, G., Ghude, S. D., Kumar, R., Sharma, S., Gunwani, P., Jena, C., Yadav, P., Ingle, S., Debnath, S., Pawar, P., Acharja, P., Jat, R., Kalita, G., Ambulkar, R., Kulkarni, S., Kaginalkar, A., Soni, V. K., Nanjundiah, R. S., and Rajeevan, M.: Decision Support System version 1.0 (DSS v1.0) for air quality management in Delhi, India, *Atmospheric sciences*, <https://doi.org/10.5194/gmd-2022-300>, 2023a.
- 935 Govardhan, G., Ambulkar, R., Kulkarni, S., Vishnoi, A., Yadav, P., Choudhury, B. A., Khare, M., and Ghude, S. D.: Stubble-burning activities in north-western India in 2021: Contribution to air pollution in Delhi, *Heliyon*, 9, e16939, <https://doi.org/10.1016/j.heliyon.2023.e16939>, 2023b.
- Greenstone, M. and Fan, C.: Air Quality Life Index, Annual Update, 2020.
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmospheric Environment*, 39, 6957–6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.
- 945 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 6, 3181–3210, <https://doi.org/10.5194/acp-6-3181-2006>, 2006.
- Gunwani, P. and Mohan, M.: Sensitivity of WRF model estimates to various PBL parameterizations in different climatic zones over India, *Atmospheric Research*, 194, 43–65, <https://doi.org/10.1016/j.atmosres.2017.04.026>, 2017.
- 950 Gupta, P., Christopher, S. A., Patadia, F., and Rastogi, N.: The unusual stubble burning season of 2020 in northern India: a satellite perspective, *International Journal of Remote Sensing*, 44, 6882–6896, <https://doi.org/10.1080/01431161.2023.2277160>, 2023.
- 955 Gupta, T., Rajeev, P., and Rajput, R.: Emerging Major Role of Organic Aerosols in Explaining the Occurrence, Frequency, and Magnitude of Haze and Fog Episodes during Wintertime in the Indo Gangetic Plain, *ACS Omega*, 7, 1575–1584, <https://doi.org/10.1021/acsomega.1c05467>, 2022.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 965 Hodzic, A. and Jimenez, J. L.: Modeling anthropogenically controlled secondary organic aerosols in a megacity: a simplified framework for global and climate models, *Geosci. Model Dev.*, 4, 901–917, <https://doi.org/10.5194/gmd-4-901-2011>, 2011.
- Hodzic, A. and Knote, C.: WRF-Chem 3.6.1: MOZART gas-phase chemistry with MOSAIC aerosols, 2014.
- 970 Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenue, F., Jankowiak, I., and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive for Aerosol Characterization, *Remote Sensing of Environment*, 66, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.

India Meteorological Department, Govt. of India Ministry of Earth Sciences: Annual Report 2016, 2017.

- 975 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Dentener, F., Muntean, M., Pouliot, G., Keating, T., Zhang, Q., Kurokawa, J., Wankmüller, R., Denier van der Gon, H., Kuenen, J. J. P., Klimont, Z., Frost, G., Darras, S., Koffi, B., and Li, M.: HTAP_v2.2: a mosaic of regional and global emission grid maps for 2008 and 2010 to study hemispheric transport of air pollution, *Atmos. Chem. Phys.*, 15, 11411–11432, <https://doi.org/10.5194/acp-15-11411-2015>, 2015.
- 980 Jena, C., Ghude, S. D., Kulkarni, R., Debnath, S., Kumar, R., Soni, V. K., Acharja, P., Kulkarni, S. H., Khare, M., Kaginalkar, A. J., Chate, D. M., Ali, K., Nanjundiah, R. S., and Rajeevan, M. N.: Evaluating the sensitivity of fine particulate matter (PM_{2.5}) simulations to chemical mechanism in Delhi, *Aerosols/Atmospheric Modelling/Troposphere/Chemistry (chemical composition and reactions)*, <https://doi.org/10.5194/acp-2020-673>, 2020.
- 985 Jena, C., Ghude, S. D., Kumar, R., Debnath, S., Govardhan, G., Soni, V. K., Kulkarni, S. H., Beig, G., Nanjundiah, R. S., and Rajeevan, M.: Performance of high resolution (400 m) PM_{2.5} forecast over Delhi, *Sci Rep*, 11, 4104, <https://doi.org/10.1038/s41598-021-83467-8>, 2021.
- Jethva, H.: Assessing predictability of post-monsoon crop residue fires in Northwestern India, *Frontiers in Earth Science*, 10, 2022.
- 990 Jethva, H., Torres, O., Field, R. D., Lyapustin, A., Gautam, R., and Kayetha, V.: Connecting Crop Productivity, Residue Fires, and Air Quality over Northern India, *Sci Rep*, 9, 16594, <https://doi.org/10.1038/s41598-019-52799-x>, 2019.
- Kalenderski, S., Stenchikov, G., and Zhao, C.: Modeling a typical winter-time dust event over the Arabian Peninsula and the Red Sea, *Atmospheric Chemistry and Physics*, 13, 1999–2014, <https://doi.org/10.5194/acp-13-1999-2013>, 2013.
- 995 Kanawade, V. P., Srivastava, A. K., Ram, K., Asmi, E., Vakkari, V., Soni, V. K., Varaprasad, V., and Sarangi, C.: What caused severe air pollution episode of November 2016 in New Delhi?, *Atmospheric Environment*, 222, 117125, <https://doi.org/10.1016/j.atmosenv.2019.117125>, 2020.
- 1000 Kaskaoutis, D. G., Kumar, S., Sharma, D., Singh, R. P., Kharol, S. K., Sharma, M., Singh, A. K., Singh, S., Singh, A., and Singh, D.: Effects of crop residue burning on aerosol properties, plume characteristics, and long-range transport over northern India: Effects of crop residue burning, *J. Geophys. Res. Atmos.*, 119, 5424–5444, <https://doi.org/10.1002/2013JD021357>, 2014.
- 1005 Knote, C., Hodzic, A., Jimenez, J. L., Volkamer, R., Orlando, J. J., Baidar, S., Brioude, J., Fast, J., Gentner, D. R., Goldstein, A. H., Hayes, P. L., Knighton, W. B., Oetjen, H., Setyan, A., Stark, H., Thalman, R., Tyndall, G., Washenfelder, R., Waxman, E., and Zhang, Q.: Simulation of semi-explicit mechanisms of SOA formation from glyoxal in aerosol in a 3-D model, *Atmos. Chem. Phys.*, 14, 6213–6239, <https://doi.org/10.5194/acp-14-6213-2014>, 2014.
- 1010 Kulkarni, S. H., Ghude, S. D., Jena, C., Karumuri, R. K., Sinha, B., Sinha, V., Kumar, R., Soni, V. K., and Khare, M.: How Much Does Large-Scale Crop Residue Burning Affect the Air Quality in Delhi?, *Environ. Sci. Technol.*, 54, 4790–4799, <https://doi.org/10.1021/acs.est.0c00329>, 2020.
- Kumar, A., Sinha, V., Shabin, M., Hakkim, H., Bonsang, B., and Gros, V.: Non-methane hydrocarbon (NMHC) fingerprints of major urban and agricultural emission sources for use in source apportionment studies, *Atmos. Chem. Phys.*, 20, 12133–12152, <https://doi.org/10.5194/acp-20-12133-2020>, 2020a.
- 1015 Kumar, A., Hakkim, H., Sinha, B., and Sinha, V.: Gridded 1 km × 1 km emission inventory for paddy stubble burning emissions over north-west India constrained by measured emission factors of 77 VOCs

and district-wise crop yield data, *Science of The Total Environment*, 789, 148064, <https://doi.org/10.1016/j.scitotenv.2021.148064>, 2021.

1020 Kumar, M., Parmar, K. S., Kumar, D. B., Mhawish, A., Broday, D. M., Mall, R. K., and Banerjee, T.: Long-term aerosol climatology over Indo-Gangetic Plain: Trend, prediction and potential source fields, *Atmospheric Environment*, 180, 37–50, <https://doi.org/10.1016/j.atmosenv.2018.02.027>, 2018a.

Kumar, M., Parmar, K. S., Kumar, D. B., Mhawish, A., Broday, D. M., Mall, R. K., and Banerjee, T.: Long-term aerosol climatology over Indo-Gangetic Plain: Trend, prediction and potential source fields, *Atmospheric Environment*, 180, 37–50, <https://doi.org/10.1016/j.atmosenv.2018.02.027>, 2018b.

1025 Kumar, R., Naja, M., Pfister, G. G., Barth, M. C., Wiedinmyer, C., and Brasseur, G. P.: Simulations over South Asia using the Weather Research and Forecasting model with Chemistry (WRF-Chem): chemistry evaluation and initial results, *Geoscientific Model Development*, 5, 619–648, <https://doi.org/10.5194/gmd-5-619-2012>, 2012a.

1030 Kumar, R., Naja, M., Pfister, G. G., Barth, M. C., and Brasseur, G. P.: Simulations over South Asia using the Weather Research and Forecasting model with Chemistry (WRF-Chem): set-up and meteorological evaluation, *Geoscientific Model Development*, 5, 321–343, <https://doi.org/10.5194/gmd-5-321-2012>, 2012b.

1035 Kumar, R., Barth, M. C., Pfister, G. G., Naja, M., and Brasseur, G. P.: WRF-Chem simulations of a typical pre-monsoon dust storm in northern India: influences on aerosol optical properties and radiation budget, *Atmospheric Chemistry and Physics*, 14, 2431–2446, <https://doi.org/10.5194/acp-14-2431-2014>, 2014.

Kumar, R., Barth, M. C., Pfister, G. G., Nair, V. S., Ghude, S. D., and Ojha, N.: What controls the seasonal cycle of black carbon aerosols in India?, *J. Geophys. Res. Atmos.*, 120, 7788–7812, <https://doi.org/10.1002/2015JD023298>, 2015.

1040 Kumar, R., Barth, M. C., Pfister, G. G., Monache, L. D., Lamarque, J. F., Archer-Nicholls, S., Tilmes, S., Ghude, S. D., Wiedinmyer, C., Naja, M., and Walters, S.: How Will Air Quality Change in South Asia by 2050?, *Journal of Geophysical Research: Atmospheres*, 123, 1840–1864, <https://doi.org/10.1002/2017JD027357>, 2018c.

1045 Kumar, R., Ghude, S. D., Biswas, M., Jena, C., Alessandrini, S., Debnath, S., Kulkarni, S., Sperati, S., Soni, V. K., Nanjundiah, R. S., and Rajeevan, M.: Enhancing Accuracy of Air Quality and Temperature Forecasts During Paddy Crop Residue Burning Season in Delhi Via Chemical Data Assimilation, *J. Geophys. Res. Atmos.*, 125, <https://doi.org/10.1029/2020JD033019>, 2020b.

1050 Kumari, S., Verma, N., Lakhani, A., and Kumari, K. M.: Severe haze events in the Indo-Gangetic Plain during post-monsoon: Synergetic effect of synoptic meteorology and crop residue burning emission, *Science of The Total Environment*, 768, 145479, <https://doi.org/10.1016/j.scitotenv.2021.145479>, 2021.

Lack, D. A. and Cappa, C. D.: Impact of brown and clear carbon on light absorption enhancement, single scatter albedo and absorption wavelength dependence of black carbon, *Atmospheric Chemistry and Physics*, 10, 4207–4220, <https://doi.org/10.5194/acp-10-4207-2010>, 2010.

1055 Lalchandani, V., Srivastava, D., Dave, J., Mishra, S., Tripathi, N., Shukla, A. K., Sahu, R., Thamban, N. M., Gaddamidi, S., Dixit, K., Ganguly, D., Tiwari, S., Srivastava, A. K., Sahu, L., Rastogi, N., Gargava, P., and Tripathi, S. N.: Effect of Biomass Burning on PM_{2.5} Composition and Secondary Aerosol Formation During Post-Monsoon and Winter Haze Episodes in Delhi, *JGR Atmospheres*, 127, <https://doi.org/10.1029/2021JD035232>, 2022.

- 1060 Li, M., Zhang, Q., Kurokawa, J., Woo, J.-H., He, K., Lu, Z., Ohara, T., Song, Y., Streets, D. G., Carmichael, G. R., Cheng, Y., Hong, C., Huo, H., Jiang, X., Kang, S., Liu, F., Su, H., and Zheng, B.: MIX: a mosaic Asian anthropogenic emission inventory under the international collaboration framework of the MICS-Asia and HTAP, *Atmos. Chem. Phys.*, 17, 935–963, <https://doi.org/10.5194/acp-17-935-2017>, 2017.
- 1065 Liu, N., Zhou, S., Liu, C., and Guo, J.: Synoptic circulation pattern and boundary layer structure associated with PM_{2.5} during wintertime haze pollution episodes in Shanghai, *Atmospheric Research*, 228, 186–195, <https://doi.org/10.1016/j.atmosres.2019.06.001>, 2019.
- Liu, T., Mickley, L. J., Gautam, R., Singh, M. K., DeFries, R. S., and Marlier, M. E.: Detection of delay in post-monsoon agricultural burning across Punjab, India: potential drivers and consequences for air quality, *Environ. Res. Lett.*, 16, 014014, <https://doi.org/10.1088/1748-9326/abcc28>, 2021.
- 1070 Lu, Z., Zhang, Q., and Streets, D. G.: Sulfur dioxide and primary carbonaceous aerosol emissions in China and India, 1996–2010, *Atmos. Chem. Phys.*, 11, 9839–9864, <https://doi.org/10.5194/acp-11-9839-2011>, 2011.
- 1075 McDuffie, E. E., Smith, S. J., O'Rourke, P., Tibrewal, K., Venkataraman, C., Marais, E. A., Zheng, B., Crippa, M., Brauer, M., and Martin, R. V.: A global anthropogenic emission inventory of atmospheric pollutants from sector- and fuel-specific sources (1970–2017): an application of the Community Emissions Data System (CEDS), *Earth System Science Data*, 12, 3413–3442, <https://doi.org/10.5194/essd-12-3413-2020>, 2020.
- 1080 Mhawish, A., Banerjee, T., Broday, D. M., Misra, A., and Tripathi, S. N.: Evaluation of MODIS Collection 6 aerosol retrieval algorithms over Indo-Gangetic Plain: Implications of aerosols types and mass loading, *Remote Sensing of Environment*, 201, 297–313, <https://doi.org/10.1016/j.rse.2017.09.016>, 2017.
- 1085 Mhawish, A., Sorek-Hamer, M., Chatfield, R., Banerjee, T., Bilal, M., Kumar, M., Sarangi, C., Franklin, M., Chau, K., Garay, M., and Kalashnikova, O.: Aerosol characteristics from earth observation systems: A comprehensive investigation over South Asia (2000–2019), *Remote Sensing of Environment*, 259, 112410, <https://doi.org/10.1016/j.rse.2021.112410>, 2021.
- Mhawish, A., Sarangi, C., Babu, P., Kumar, M., Bilal, M., and Qiu, Z.: Observational evidence of elevated smoke layers during crop residue burning season over Delhi: Potential implications on associated heterogeneous PM_{2.5} enhancements, *Remote Sensing of Environment*, 280, 113167, <https://doi.org/10.1016/j.rse.2022.113167>, 2022.
- 1090 Mogno, C., Palmer, P. I., Knote, C., Yao, F., and Wallington, T. J.: Seasonal distribution and drivers of surface fine particulate matter and organic aerosol over the Indo-Gangetic Plain, *Aerosols/Atmospheric Modelling/Troposphere/Chemistry (chemical composition and reactions)*, <https://doi.org/10.5194/acp-2021-69>, 2021.
- 1095 Mohan, M. and Bhati, S.: Analysis of WRF Model Performance over Subtropical Region of Delhi, India, *Advances in Meteorology*, 2011, 1–13, <https://doi.org/10.1155/2011/621235>, 2011.
- Moorthy, K. K., Beegum, S. N., Srivastava, N., Satheesh, S. K., Chin, M., Blond, N., Babu, S. S., and Singh, S.: Performance evaluation of chemistry transport models over India, *Atmospheric Environment*, 71, 210–225, <https://doi.org/10.1016/j.atmosenv.2013.01.056>, 2013.
- 1100 Mues, A., Lauer, A., Lupascu, A., Rupakheti, M., Kuik, F., and Lawrence, M. G.: WRF and WRF-Chem v3.5.1 simulations of meteorology and black carbon concentrations in the Kathmandu Valley, *Geosci. Model Dev.*, 11, 2067–2091, <https://doi.org/10.5194/gmd-11-2067-2018>, 2018.

- Mukherjee, T., Asutosh, A., Pandey, S. K., Yang, L., Gogoi, P. P., Panwar, A., and Vinoj, V.: Increasing Potential for Air Pollution over Megacity New Delhi: A Study Based on 2016 Diwali Episode, *Aerosol Air Qual. Res.*, 18, 2510–2518, <https://doi.org/10.4209/aaqr.2017.11.0440>, 2018.
- 1105 Mukherjee, T., Vinoj, V., Midya, S. K., Puppala, S. P., and Adhikary, B.: Numerical simulations of different sectoral contributions to post monsoon pollution over Delhi, *Heliyon*, 6, e03548, <https://doi.org/10.1016/j.heliyon.2020.e03548>, 2020.
- Nair, V. S., Solmon, F., Giorgi, F., Mariotti, L., Babu, S. S., and Moorthy, K. K.: Simulation of South Asian aerosols for regional climate studies, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016711>, 2012.
- 1110 Navinya, C. D., Vinoj, V., and Pandey, S. K.: Evaluation of PM_{2.5} Surface Concentrations Simulated by NASA's MERRA Version 2 Aerosol Reanalysis over India and its Relation to the Air Quality Index, *Aerosol Air Qual. Res.*, 20, 1329–1339, <https://doi.org/10.4209/aaqr.2019.12.0615>, 2020.
- Nelli, N. R., Temimi, M., Fonseca, R. M., Weston, M. J., Thota, M. S., Valappil, V. K., Branch, O., Wulfmeyer, V., Wehbe, Y., Al Hosary, T., Shalaby, A., Al Shamsi, N., and Al Naqbi, H.: Impact of Roughness Length on WRF Simulated Land-Atmosphere Interactions Over a Hyper-Arid Region, *Earth and Space Science*, 7, e2020EA001165, <https://doi.org/10.1029/2020EA001165>, 2020.
- Ojha, N., Sharma, A., Kumar, M., Girach, I., Ansari, T. U., Sharma, S. K., Singh, N., Pozzer, A., and Gunthe, S. S.: On the widespread enhancement in fine particulate matter across the Indo-Gangetic Plain towards winter, *Sci Rep*, 10, 5862, <https://doi.org/10.1038/s41598-020-62710-8>, 2020.
- 1120 Pan, X., Chin, M., Gautam, R., Bian, H., Kim, D., Colarco, P. R., Diehl, T. L., Takemura, T., Pozzoli, L., Tsigaridis, K., Bauer, S., and Bellouin, N.: A multi-model evaluation of aerosols over South Asia: common problems and possible causes, *Atmos. Chem. Phys.*, 15, 5903–5928, <https://doi.org/10.5194/acp-15-5903-2015>, 2015.
- 1125 Pandey, A., Brauer, M., Cropper, M. L., Balakrishnan, K., Mathur, P., Dey, S., Turkoglu, B., Kumar, G. A., Khare, M., Beig, G., Gupta, T., Krishnankutty, R. P., Causey, K., Cohen, A. J., Bhargava, S., Aggarwal, A. N., Agrawal, A., Awasthi, S., Bennitt, F., Bhagwat, S., Bhanumati, P., Burkart, K., Chakma, J. K., Chiles, T. C., Chowdhury, S., Christopher, D. J., Dey, S., Fisher, S., Fraumeni, B., Fuller, R., Ghoshal, A. G., Golechha, M. J., Gupta, P. C., Gupta, R., Gupta, R., Gupta, S., Guttikunda, S., Hanrahan, D., Harikrishnan, S., Jeemon, P., Joshi, T. K., Kant, R., Kant, S., Kaur, T., Koul, P. A., Kumar, P., Kumar, R., Larson, S. L., Lodha, R., Madhipatla, K. K., Mahesh, P. A., Malhotra, R., Managi, S., Martin, K., Mathai, M., Mathew, J. L., Mehrotra, R., Mohan, B. V. M., Mohan, V., Mukhopadhyay, S., Mutreja, P., Naik, N., Nair, S., Pandian, J. D., Pant, P., Perianayagam, A., Prabhakaran, D., Prabhakaran, P., Rath, G. K., Ravi, S., Roy, A., Sabde, Y. D., Salvi, S., Sambandam, S., Sharma, B., Sharma, M., Sharma, S., Sharma, R. S., Shrivastava, A., Singh, S., Singh, V., Smith, R., Stanaway, J. D., Taghian, G., Tandon, N., Thakur, J. S., Thomas, N. J., Toteja, G. S., Varghese, C. M., Venkataraman, C., Venugopal, K. N., Walker, K. D., Watson, A. Y., Wozniak, S., Xavier, D., Yadama, G. N., Yadav, G., Shukla, D. K., Bekedam, H. J., et al.: Health and economic impact of air pollution in the states of India: the Global Burden of Disease Study 2019, *The Lancet Planetary Health*, 5, e25–e38, [https://doi.org/10.1016/S2542-5196\(20\)30298-9](https://doi.org/10.1016/S2542-5196(20)30298-9), 2021.
- 1130 1135 1140 Patel, K., Bhandari, S., Gani, S., Campmier, M. J., Kumar, P., Habib, G., Apte, J., and Hildebrandt Ruiz, L.: Sources and Dynamics of Submicron Aerosol during the Autumn Onset of the Air Pollution Season in Delhi, India, *ACS Earth Space Chem.*, 5, 118–128, <https://doi.org/10.1021/acsearthspacechem.0c00340>, 2021.
- 1145 Paulot, F., Naik, V., and W. Horowitz, L.: Reduction in Near-Surface Wind Speeds With Increasing CO₂ May Worsen Winter Air Quality in the Indo-Gangetic Plain, *Geophysical Research Letters*, 49, e2022GL099039, <https://doi.org/10.1029/2022GL099039>, 2022.

- 1150 Pawar, P. V., Ghude, S. D., Govardhan, G., Acharja, P., Kulkarni, R., Kumar, R., Sinha, B., Sinha, V., Jena, C., Gunwani, P., Adhya, T. K., Nemitz, E., and Sutton, M. A.: Chloride (HCl  Cl⁻) dominates inorganic aerosol formation from ammonia in the Indo-Gangetic Plain during winter: modeling and comparison with observations, *Atmospheric Chemistry and Physics*, 23, 41–59, <https://doi.org/10.5194/acp-23-41-2023>, 2023.
- 1155 Provencal, S., Buchard, V., Silva, A. M. da, Leduc, R., Barrette, N., Elhacham, E., and Wang, S.-H.: Evaluation of PM_{2.5} Surface Concentrations Simulated by Version 1 of NASA’s MERRA Aerosol Reanalysis over Israel and Taiwan, *Aerosol Air Qual. Res.*, 17, 253–261, <https://doi.org/10.4209/aaqr.2016.04.0145>, 2017.
- Ram, K. and Sarin, M. M.: Day–night variability of EC, OC, WSOC and inorganic ions in urban environment of Indo-Gangetic Plain: Implications to secondary aerosol formation, *Atmospheric Environment*, 45, 460–468, <https://doi.org/10.1016/j.atmosenv.2010.09.055>, 2011.
- 1160 Ram, K., Singh, S., Sarin, M. M., Srivastava, A. K., and Tripathi, S. N.: Variability in aerosol optical properties over an urban site, Kanpur, in the Indo-Gangetic Plain: A case study of haze and dust events, *Atmospheric Research*, 174–175, 52–61, <https://doi.org/10.1016/j.atmosres.2016.01.014>, 2016.
- Ramanathan, V., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119–2124, <https://doi.org/10.1126/science.1064034>, 2001.
- 1165 Randles, C. A., Da Silva, A. M., Buchard, V., Colarco, P. R., Darmenov, A., Govindaraju, R., Smirnov, A., Holben, B., Ferrare, R., Hair, J., Shinozuka, Y., and Flynn, C. J.: The MERRA-2 Aerosol Reanalysis, 1980 - onward, Part I: System Description and Data Assimilation Evaluation, *J Clim*, 30, 6823–6850, <https://doi.org/10.1175/JCLI-D-16-0609.1>, 2017.
- 1170 Ratnam, J. V. and Kumar, K. K.: Sensitivity of the Simulated Monsoons of 1987 and 1988 to Convective Parameterization Schemes in MM5, *Journal of Climate*, 18, 2724–2743, <https://doi.org/10.1175/JCLI3390.1>, 2005.
- Riemer, N., Ault, A. P., West, M., Craig, R. L., and Curtis, J. H.: Aerosol Mixing State: Measurements, Modeling, and Impacts, *Rev. Geophys.*, 57, 187–249, <https://doi.org/10.1029/2018RG000615>, 2019.
- 1175 Rooney, B., Zhao, R., Wang, Y., Bates, K. H., Pillarisetti, A., Sharma, S., Kundu, S., Bond, T. C., Lam, N. L., Ozaltun, B., Xu, L., Goel, V., Fleming, L. T., Weltman, R., Meinardi, S., Blake, D. R., Nizkorodov, S. A., Edwards, R. D., Yadav, A., Arora, N. K., Smith, K. R., and Seinfeld, J. H.: Impacts of household sources on air pollution at village and regional scales in India, *Atmos. Chem. Phys.*, 19, 7719–7742, <https://doi.org/10.5194/acp-19-7719-2019>, 2019.
- 1180 Roozitalab, B., Carmichael, G. R., and Guttikunda, S. K.: Improving regional air quality predictions in the Indo-Gangetic Plain – case study of an intensive pollution episode in November 2017, *Atmos. Chem. Phys.*, 21, 2837–2860, <https://doi.org/10.5194/acp-21-2837-2021>, 2021.
- Sarkar, S., Singh, R. P., and Chauhan, A.: Crop Residue Burning in Northern India: Increasing Threat to Greater India, *J. Geophys. Res. Atmos.*, 123, 6920–6934, <https://doi.org/10.1029/2018JD028428>, 2018.
- 1185 Sawlani, R., Agnihotri, R., Sharma, C., Patra, P. K., Dimri, A. P., Ram, K., and Verma, R. L.: The severe Delhi SMOG of 2016: A case of delayed crop residue burning, coincident firecracker emissions, and atypical meteorology, *Atmospheric Pollution Research*, 10, 868–879, <https://doi.org/10.1016/j.apr.2018.12.015>, 2019.
- 1190 Sayer, A. M., Munchak, L. A., Hsu, N. C., Levy, R. C., Bettenhausen, C., and Jeong, M.-J.: MODIS Collection 6 aerosol products: Comparison between Aqua’s e-Deep Blue, Dark Target, and “merged”

- data sets, and usage recommendations, *Journal of Geophysical Research: Atmospheres*, 119, 13,965–13,989, <https://doi.org/10.1002/2014JD022453>, 2014.
- 1195 Schnell, J. L., Naik, V., Horowitz, L. W., Paulot, F., Mao, J., Ginoux, P., Zhao, M., and Ram, K.: Exploring the relationship between surface PM_{2.5} and meteorology in Northern India, *Atmospheric Chemistry and Physics*, 18, 10157–10175, <https://doi.org/10.5194/acp-18-10157-2018>, 2018.
- Sembhi, H., Wooster, M., Zhang, T., Sharma, S., Singh, N., Agarwal, S., Boesch, H., Gupta, S., Misra, A., Tripathi, S. N., Mor, S., and Khaiwal, R.: Post-monsoon air quality degradation across Northern India: assessing the impact of policy-related shifts in timing and amount of crop residue burnt, *Environ. Res. Lett.*, 15, 104067, <https://doi.org/10.1088/1748-9326/aba714>, 2020.
- 1200 Sengupta, A., Govardhan, G., Debnath, S., Yadav, P., Kulkarni, S. H., Parde, A. N., Lonkar, P., Dhangar, N., Gunwani, P., Wagh, S., Nivdange, S., Jena, C., Kumar, R., and Ghude, S. D.: Probing into the wintertime meteorology and particulate matter (PM_{2.5} and PM₁₀) forecast over Delhi, *Atmospheric Pollution Research*, 13, 101426, <https://doi.org/10.1016/j.apr.2022.101426>, 2022.
- 1205 Shaik, D. S., Kant, Y., Mitra, D., Singh, A., Chandola, H. C., Sateesh, M., Babu, S. S., and Chauhan, P.: Impact of biomass burning on regional aerosol optical properties: A case study over northern India, *Journal of Environmental Management*, 244, 328–343, <https://doi.org/10.1016/j.jenvman.2019.04.025>, 2019.
- 1210 Sharma, S. K., Mandal, T. K., Sharma, A., Jain, S., and Saraswati: Carbonaceous Species of PM_{2.5} in Megacity Delhi, India During 2012–2016, *Bull Environ Contam Toxicol*, 100, 695–701, <https://doi.org/10.1007/s00128-018-2313-9>, 2018.
- Shen, C., Liu, Y., Shen, A., Cui, Y., Chen, X., Fan, Q., Chan, P., Tian, C., Xie, Z., Wang, C., Lan, J., Li, X., Wu, J., and Yang, Y.: Spatializing the roughness length of heterogeneous urban surfaces to improve the WRF simulation-Part 2: Impacts on the thermodynamic environment, *Atmospheric Environment*, 294, 119464, <https://doi.org/10.1016/j.atmosenv.2022.119464>, 2023.
- 1215 Singh, N., Murari, V., Kumar, M., Barman, S. C., and Banerjee, T.: Fine particulates over South Asia: Review and meta-analysis of PM_{2.5} source apportionment through receptor model, *Environmental Pollution*, 223, 121–136, <https://doi.org/10.1016/j.envpol.2016.12.071>, 2017.
- 1220 Singh, N., Agarwal, S., Sharma, S., Chatani, S., and Ramanathan, V.: Air Pollution Over India: Causal Factors for the High Pollution with Implications for Mitigation, *ACS Earth Space Chem.*, 5, 3297–3312, <https://doi.org/10.1021/acsearthspacechem.1c00170>, 2021.
- Singh, R., Kumar, S., and Singh, A.: Elevated Black Carbon Concentrations and Atmospheric Pollution around Singrauli Coal-Fired Thermal Power Plants (India) Using Ground and Satellite Data, *IJERPH*, 15, 2472, <https://doi.org/10.3390/ijerph15112472>, 2018.
- 1225 Singh, T., Biswal, A., Mor, S., Ravindra, K., Singh, V., and Mor, S.: A high-resolution emission inventory of air pollutants from primary crop residue burning over Northern India based on VIIRS thermal anomalies, *Environmental Pollution*, 266, 115132, <https://doi.org/10.1016/j.envpol.2020.115132>, 2020.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 3, 125, n.d.
- 1230 Song, Z., Fu, D., Zhang, X., Wu, Y., Xia, X., He, J., Han, X., Zhang, R., and Che, H.: Diurnal and seasonal variability of PM_{2.5} and AOD in North China plain: Comparison of MERRA-2 products and ground measurements, *Atmospheric Environment*, 191, 70–78, <https://doi.org/10.1016/j.atmosenv.2018.08.012>, 2018.

- 1235 Srivastava, N., Satheesh, S. K., Blond, N., and Krishna Moorthy, K.: Simulation of Aerosol Fields over South Asia Using CHIMERE - Part-II: Performance Evaluation, *Current Science*, 111, 83, <https://doi.org/10.18520/cs/v111/i1/83-92>, 2016.
- Stauffer, D. R. and Seaman, N. L.: Multiscale Four-Dimensional Data Assimilation, *J. Appl. Meteor.*, 33, 416–434, [https://doi.org/10.1175/1520-0450\(1994\)033<0416:MFDDA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1994)033<0416:MFDDA>2.0.CO;2), 1994.
- 1240 Takigawa, M., Patra, P. K., Matsumi, Y., Dhaka, S. K., Nakayama, T., Yamaji, K., Kajino, M., and Hayashida, S.: Can Delhi's Pollution be Affected by Crop Fires in the Punjab Region?, *SOLA*, 16, 86–91, <https://doi.org/10.2151/sola.2020-015>, 2020.
- 1245 Talukdar, S., Tripathi, S. N., Lalchandani, V., Rupakheti, M., Bhowmik, H. S., Shukla, A. K., Murari, V., Sahu, R., Jain, V., Tripathi, N., Dave, J., Rastogi, N., and Sahu, L.: Air Pollution in New Delhi during Late Winter: An Overview of a Group of Campaign Studies Focusing on Composition and Sources, *Atmosphere*, 12, 1432, <https://doi.org/10.3390/atmos12111432>, 2021.
- Thomas, A., Sarangi, C., and Kanawade, V. P.: Recent Increase in Winter Hazy Days over Central India and the Arabian Sea, *Sci Rep*, 9, 17406, <https://doi.org/10.1038/s41598-019-53630-3>, 2019.
- 1250 Tie, X., Madronich, S., Walters, S., Zhang, R., Rasch, P., and Collins, W.: Effect of clouds on photolysis and oxidants in the troposphere, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2003JD003659>, 2003.
- Tuccella, P., Curci, G., Visconti, G., Bessagnet, B., Menut, L., and Park, R. J.: Modeling of gas and aerosol with WRF/Chem over Europe: Evaluation and sensitivity study, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016302>, 2012.
- 1255 Upadhyay, A., Dey, S., and Goyal, P.: A comparative assessment of regional representativeness of EDGAR and ECLIPSE emission inventories for air quality studies in India, *Atmospheric Environment*, 223, 117182, <https://doi.org/10.1016/j.atmosenv.2019.117182>, 2020.
- 1260 Venkataraman, C., Brauer, M., Tibrewal, K., Sadavarte, P., Ma, Q., Cohen, A., Chaliyakunnel, S., Frostad, J., Klimont, Z., Martin, R. V., Millet, D. B., Philip, S., Walker, K., and Wang, S.: Source influence on emission pathways and ambient PM_{2.5} pollution over India (2015–2050), *Atmos. Chem. Phys.*, 18, 8017–8039, <https://doi.org/10.5194/acp-18-8017-2018>, 2018.
- Vogelezang, D. H. P. and Holtslag, A. A. M.: Evaluation and model impacts of alternative boundary-layer height formulations, *Boundary-Layer Meteorol.*, 81, 245–269, <https://doi.org/10.1007/BF02430331>, 1996.
- 1265 Voiland and Jethva: *Earth Matters - Using Satellites to Size Up the Severity of Crop Fires in Northern India*, 2017.
- Wang, K., Zhang, Y., and Yahya, K.: Decadal application of WRF/Chem over the continental U.S.: Simulation design, sensitivity simulations, and climatological model evaluation, *Atmospheric Environment*, 253, 118331, <https://doi.org/10.1016/j.atmosenv.2021.118331>, 2021.
- 1270 Wang, R., Tao, S., Shen, H., Huang, Y., Chen, H., Balkanski, Y., Boucher, O., Ciais, P., Shen, G., Li, W., Zhang, Y., Chen, Y., Lin, N., Su, S., Li, B., Liu, J., and Liu, W.: Trend in Global Black Carbon Emissions from 1960 to 2007, *Environ. Sci. Technol.*, 48, 6780–6787, <https://doi.org/10.1021/es5021422>, 2014.
- Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions

- 1275 from open burning, *Geoscientific Model Development*, 4, 625–641, <https://doi.org/10.5194/gmd-4-625-2011>, 2011.
- Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for Simulating Aerosol Interactions and Chemistry (MOSAIC), *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008782>, 2008.
- 1280 Zhang, Y., Zhang, X., Wang, L., Zhang, Q., Duan, F., and He, K.: Application of WRF/Chem over East Asia: Part I. Model evaluation and intercomparison with MM5/CMAQ, *Atmospheric Environment*, 124, 285–300, <https://doi.org/10.1016/j.atmosenv.2015.07.022>, 2016.
- Zhao, C., Liu, X., Leung, L. R., Johnson, B., McFarlane, S. A., Gustafson, W. I. J., Fast, J. D., and Easter, R.: The spatial distribution of mineral dust and its shortwave radiative forcing over North Africa: modeling sensitivities to dust emissions and aerosol size treatments, *Atmospheric Chemistry and Physics*, 10, 8821–8838, <https://doi.org/10.5194/acp-10-8821-2010>, 2010.
- 1285 Zhao, C., Chen, S., Leung, L. R., Qian, Y., Kok, J. F., Zaveri, R. A., and Huang, J.: Uncertainty in modeling dust mass balance and radiative forcing from size parameterization, *Atmospheric Chemistry and Physics*, 13, 10733–10753, <https://doi.org/10.5194/acp-13-10733-2013>, 2013.

1290