

Investigating the differences in calculating global mean surface CO₂ abundance: the impact of analysis methodologies and site selection

Zhendong Wu^{1,2,*}, Alex Vermeulen^{2,*}, Yousuke Sawa³, Ute Karstens^{1,2}, Wouter Peters^{4,5}, Remco de Kok⁶, Xin Lan^{7,8}, Yasuyuki Nagai³, Akinori Ogi³, Oksana Tarasova⁹

¹ICOS Carbon Portal at Lund University, Lund, Sweden

²ICOS ERIC, Carbon Portal, Lund, Sweden

³Japan Meteorological Agency (JMA), Tokyo, Japan

⁴Wageningen University, Wageningen, The Netherlands

⁵University of Groningen, Groningen, The Netherlands

⁶ICOS Carbon Portal at Wageningen University, Wageningen, The Netherlands

⁷NOAA Global Monitoring Laboratory (GML), Boulder, USA

⁸Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, USA

⁹WMO, Geneva, Switzerland

Correspondence to: Alex Vermeulen (alex.vermeulen@icos-ri.eu), and Zhendong Wu (zhendong.wu@nateko.lu.se)

Abstract. The World Meteorological Organization (WMO) Global Atmosphere Watch (GAW) coordinates high-quality atmospheric greenhouse gas observations globally and provides these observations through the WMO World Data Centre for Greenhouse Gases (WDCGG) supported by Japan Meteorological Agency. The WDCGG and the National Oceanic and Atmospheric Administration (NOAA) analyse these measurements using different methodologies and site selection to calculate global annual mean surface CO₂ and its growth rate as a headline climate indicator. This study introduces a third hybrid method named GFIT, which serves as an independent validation and open-source alternative to the methods described by NOAA and WDCGG. We apply GFIT to incorporate observations from most WMO GAW stations and 3D modelled CO₂ fields from CarbonTracker Europe (CTE). We find that different observational networks (i.e., NOAA, GAW, and CTE networks) and analysis methods result in differences in the calculated global surface CO₂ mole fractions equivalent to the current atmospheric growth rate over a three-month period. However, the CO₂ growth rate derived from these networks and CTE model output shows good agreement. Over the long-term period (40 years), both networks with and without continental sites exhibit the same trend in the growth rate (0.030 ± 0.002 ppm per year each year). However, a clear difference emerges in the short-term (one-month) change in the growth rate. The network that includes continental sites improves the early detection of changes in biogenic emissions.

40 1 Introduction

41 Global mean surface temperature averaged over 2011–2020 has increased by about 1.09°C relative to the average temperature
42 of 1850–1900 (Gulev et al., 2021). The increasing amount of atmospheric carbon dioxide (CO₂), together with increases in
43 other greenhouse gases, is the main driver of the warming (Eyring et al., 2021). After being relatively stable between 180 ppm
44 (ice age) and 280 ppm (interglacial) for the last 800,000 years (Lüthi et al., 2008), the annual average CO₂ level of the
45 atmosphere has increased since the industrial revolution from roughly 277 ppm in 1750 to 415.7±0.2 ppm in 2021 (WMO,
46 2022), due to emissions of CO₂ related to human activities like burning of fossil fuels and land use changes (Friedlingstein et
47 al., 2022). Mean global atmospheric CO₂ annual growth rate (G_{ATM}) is an important constraint on the global carbon cycle.
48 Based on the most recent Global Carbon Budget (GCB) analysis (Friedlingstein et al., 2022), the total emission of CO₂ due to
49 human activities was $10.2 \pm 0.8 \text{ GtC yr}^{-1}$ in 2020, of which $3.0 \pm 0.4 \text{ GtC yr}^{-1}$ was captured by the ocean sink and $2.9 \pm 1 \text{ GtC}$
50 yr^{-1} by the terrestrial sink, leaving a net increase of $5.0 \pm 0.2 \text{ GtC yr}^{-1}$ of CO₂ in the atmosphere, corresponding to an
51 atmospheric CO₂ mole fraction increase of $2.4 \pm 0.1 \text{ ppm yr}^{-1}$ (the conversion factor comes from Ballantyne et al. (2012)).

52 As the atmosphere mixes the contributions of all sources and sinks, an observational global average CO₂ mole fraction can be
53 constructed if there are enough observations to represent the spatial and temporal variation across the globe. Since most land
54 masses are concentrated in the Northern Hemisphere, and the highest anthropogenic emissions (e.g. during winter) occur in
55 the relatively narrow latitudinal band between 30 °N and 60 °N, relatively large spatial and temporal gradients in CO₂ mole
56 fraction exist in and around that region. Due to convective and advective mixing, the average mixing time of air within the
57 same latitudinal bands varies from several weeks to a month. However, mixing between latitudinal bands is slower, especially
58 the exchange between the northern and southern hemispheres, which has an approximate interhemispheric transport time of
59 1.4 ± 0.2 years (Patra et al., 2011). The interplay of the latitudinal and interhemispheric differences in fossil fuel emissions
60 and seasonal exchange with land biota (Denning et al., 1995) creates a latitudinal and interhemispheric gradient that requires
61 a sufficiently dense network to capture a representative global annual mean.


62 However, measurement stations that are close to sources or sinks may not be representative of a large atmospheric volume and
63 the average signal at their latitude. Therefore, inclusion of these observations might introduce biases on the global mean CO₂
64 and its growth rate. These biases can be avoided by filtering of data and a careful selection of spatially representative stations,
65 as done by NOAA in their use of 43 stations (Fig. 1) that are considered to be representative for the Marine Boundary Layer
66 (MBL reference network, <https://www.esrl.noaa.gov/gmd/ccgg/mbl/mbl.html>). An additional data processing step developed
67 by NOAA to further avoid biases due to unrepresentative local signals is filtering and smoothing, by using a combination of a
68 low pass filter and decomposition into a fitted long-term trend and seasonal cycle (Thoning et al., 1989), hereafter referred to
69 as the NOAA analysis. These fits can also be used to fill gaps for missing data, though care must be taken to avoid extrapolation
70 errors before and beyond the time covered by the data record of the station. The WMO Global Atmosphere Watch (GAW)
71 World Data Centre for Greenhouse Gases (WDCGG) publishes global averages mole fraction for CO₂ and the other major
72 greenhouse gases in the annual WMO GAW Greenhouse Gas Bulletin (latest version: WMO, 2022). They use curve fitting
73 and filter methods that are very similar to those developed by NOAA, but WDCGG includes continental locations that are
74 potentially more influenced by local sources and sinks (Tsutsumi et al., 2009).

75 The NOAA MBL observations are all part of the NOAA cooperative global air sampling network and analysed in the same
76 laboratory. All NOAA flask-air observations are traceable to the current scale WMO-CO₂-X2019 (Hall et al., 2021) that is
77 maintained by NOAA Global Monitoring Laboratory (GML). In contrast, the WDCGG data originate from multiple
78 independent laboratories (including NOAA GML), that together form a network of hundreds of stations coordinated by WMO
79 GAW [<http://gawsis.meteoswiss.ch>]. Having a multitude of independent laboratories carries an additional risk of biases due to
80 differences in sampling, measurement, and analysis methods, for example calibration scales, although much care is taken to
81 avoid these by coordination in the network and use of a common calibration scale from WMO Central Calibration Laboratories

82 (CCL) guided by a set of strict measurement compatibility goals (WMO, 2022). The different selection of stations results in a
83 larger seasonal cycle amplitude in WDCGG results compared to those of NOAA and a small but quite consistent bias in global
84 surface annual mean CO₂ mole fraction (Tsutsumi et al., 2009). The NOAA estimate of global surface annual mean CO₂ mole
85 fraction is expected to be lower (e.g. ~0.35 ppm lower than the WDCGG estimate, Tsutsumi et al., 2009) compared to a full
86 global surface average because areas with large sources are not represented. However, the two aforementioned approaches
87 neither represent the parts of the atmosphere with low CO₂ mole fraction levels (i.e., the full troposphere, up to ~8-15 km
88 altitude, and the stratosphere), nor do they cover the regions of the world with substantial observational gaps.

89 In this paper, we propose a data integration method to estimate the global mean surface CO₂ and its growth rate, named GFIT.
90 This method serves as an independent validation of the methods as described by NOAA and WDCGG through a completely
91 independent and open-source implementation. The global mean surface CO₂ refers to the mean CO₂ mole fraction within the
92 planetary boundary layer, which extends from the Earth's surface up to a few hundred or thousand meters in height. We apply
93 the GFIT methodology to incorporate CO₂ data from the GAW network (139 stations, Fig. 1) and the modelled CO₂ distribution
94 from a well-established 3D global transport model (TM5: Transport Model 5, Peters et al., 2004, Krol et al., 2005). We
95 investigate the influence of small differences between the three methodologies and whether these are significant or not for
96 calculating the global mean surface CO₂ and its growth rate, how consistent the GFIT and WDCGG approaches are with each
97 other, and how they compare with NOAA analysis and estimates derived from a CO₂ simulation with the 3D transport model
98 TM5. These 3D CO₂ results for 2001-2020 using TM5 are performed in the CarbonTracker Europe framework (CTE, Peters
99 et al., 2004, van der Laan-Luijkx et al., 2017), where the CO₂ uptake and emission fluxes are optimized by the inversion system
100 to minimize the mismatch between the *in situ* observations and the modelled CO₂ mole fraction. CTE generally has a good
101 representation of the CO₂ field, with mean biases with respect to independent aircraft measurements of generally less than 0.5
102 ppm (Friedlingstein et al., 2022). Furthermore, the inferred CO₂ fluxes from CTE fit well within the ensemble of those of other
103 inversions used for the evaluation of Global Carbon Budget (e.g. Friedlingstein et al., 2022).

104 2 Methods and data



105
106 **Figure 1. Three observation networks are employed to assess the impact of continental site inclusion when calculating**
107 **global CO₂ mole fraction and its growth rates. The NOAA network (43 sites, yellow stars) comprises MBL sites only.**
108 **The selected GAW global network (139 sites, red dots) includes both MBL sites and continental sites, for example from**
109 **the Advanced Global Atmospheric Gases Experiment (AGAGE) and European ICOS contribution network. The CTE**
110 **network serves as the global network for the CTE model evaluations (230 sites, blue dots), comprises MBL sites and a**
111 **more extensive inclusion of continental sites.**

112 **2.1 The WMO GAW observations and WDCGG analysis method**

113 The WMO GAW network measurements are archived and distributed by WDCGG, hosted by the Japan Meteorological
114 Agency. The GAW observations used in this study originate from 139 selected stations of the GAW network, and all
115 observations are on the WMO standard scale WMO-CO₂-X2019. The details on the station selection are described in Tsutsumi
116 et al., (2009), which mainly excludes stations located in the northern hemisphere that show large standard deviations from the
117 latitudinal fitted curve. The remaining 139 stations show a more reasonable latitudinal scatter range (Fig. 1).

118 The WDCGG global analysis method (hereafter WDCGG method), as described in Tsutsumi et al., (2009), includes the
119 mentioned station selection, a data fitting and filter (involves data interpolation and extrapolation), and calculation of the zonal
120 and global mean mole fractions, trends, and growth rates. The procedure is also summarized in Text S1. The output from the
121 global analysis by the WDCGG method is used to compare against an alternative method (GFIT) that we designed to follow
122 as closely as possible the fit and filter method (Conway et al., 1994) deployed by NOAA and is described in the section 2.3.

123 **2.2 CTE model output and station observations**

124 CarbonTracker Europe (CTE) is a global model of atmospheric CO₂ and designed to keep track of CO₂ uptake and release at
125 the Earth's surface over time (van der Laan-Luijkx et al., 2017). CTE incorporates an off-line atmospheric transport module
126 (TM5, Peters et al., 2004, Krol et al., 2005) driven by ECMWF ERA5 data, and there are four prescribed fluxes (i.e. from
127 ocean, biosphere, fire and fossil fuel), which are transported in the model, together with the transported initial CO₂ field. CTE
128 also includes a data assimilation system that applies an ensemble Kalman filter to optimize the biogenic and ocean fluxes for
129 a combination of plant-functional types and climate zones to improve the fit of the simulated concentrations with observations.
130 The optimized fluxes from the data assimilation have been used in Global Carbon Project (GCP) 2021, and the comparison of
131 CTE CO₂ product to the other data assimilation systems used in GCP shows good agreement (within 0.8 ppm at all latitude
132 bands) (Friedlingstein et al., 2022).

133 The CTE model data used here consists of simulated monthly CO₂ mole fraction at 1x1 degree horizontal resolution and 25
134 levels in the vertical, the data period ranges from 2001 to 2020 which has no influence of model spin-up (Krol et al., 2018).
135 From the CTE output a set of simulated synthetic atmospheric CO₂ mole fractions with monthly resolution can be extracted
136 within grid cells where stations are situated. This study analyses monthly observation data (1980-2020) and synthetic time
137 series (2001-2020) by using the GFIT method (section 2.3) and attempts to estimate global mean CO₂ mole fraction and its
138 growth rate. The observed CO₂ mole fractions are taken from 230 out of 290 global-wide distributed stations (Fig. 1, the station
139 selection is summarized in Text S2), the data come from the GLOBALVIEW-plus V8 ObsPack data product (Schuldt et al.,
140 2022), and include surface-based, shipboard-based and tower-based measurements.

141 **2.3 The GFIT method**

142 The temporal pattern of CO₂ measurement records at locations around the globe can be explained as the combination of roughly
143 three components: a long-term trend, a non-sinusoidal yearly cycle (or seasonality), and short-term variations. This study
144 synchronizes monthly CO₂ records with the fitting and filter method developed at the NOAA Global Monitoring Laboratory
145 (Thoning et al., 1989, Conway et al., 1994), without extrapolation. The station selection and CO₂ averaging method are kept
146 the same as in the WDCGG method (Text S1). This method will be referred to as the GFIT method and will be compared to
147 the WDCGG method without extrapolation. The only difference from WDCGG method without extrapolation is the fitting
148 and filter method. All code for the method described here was developed in Python and is available as a Jupyter notebook
149 under a GPL license [<https://doi.org/10.18160/Q788-9081>]. The GFIT method can be summarized and illustrated by the
150 following three steps.

151 2.3.1. Fitting and filter

152 CO₂ records from each station can be abstracted as a combination of long-term trend and seasonality, which can be fitted by a
153 function consisting of polynomial and harmonics. We applied a linear regression analysis based on 3 polynomial coefficients
154 and 4 harmonics (Eq. 1) to fit CO₂ data using general linear least-squares fit (LFIT, Press et al., 1988).

$$155 \quad f(x) = a_0 + a_1 t + a_2 t^2 + \dots + a_k t^k + \sum_{n=1}^{n_h} (A_n \cos 2\pi n t + B_n \sin 2\pi n t) \quad (1)$$

156 where a_k , A_n and B_n are fitted parameters, t is the time from the beginning of the observation and it is in months and expressed
157 as a decimal of its year. k denotes polynomial number, $k = 2$. n_h denotes harmonic number, $n_h = 4$. Fig. 2 illustrates the
158 function fit to CO₂ data to obtain the annual oscillation (red line in Fig. 2a), is a combination of a polynomial fit to the trend
159 (blue line in Fig. 2a) and harmonic fit to the seasonality (green line in Fig. 2b).


160 The residuals are the difference between raw data and the function fit (black dots in Fig. 2c). The filtering method is based on
161 Thoning et al. (1989) which transforms CO₂ data from time domain to frequency domain using a Fast Fourier Transform
162 (FFT), then applies a low pass filter to the frequency data to remove high-frequency variations, and then transforms the filtered
163 data back to the time domain using an inverse FFT. The short-term (a cut-off value of 80 days, red line in Fig. 2c) and long-
164 term (a cut-off value of 667 days, blue line in Fig. 2c) filters used here are the same as in NOAA method, and applied to obtain
165 the short term and interannual variations that are not determined by the fit function. The original code is also available as
166 Python code from the NOAA website [<https://gml.noaa.gov/aftp/user/thoning/ccgcrv/>].

167 2.3.2. Calculate smoothed CO₂ and long-term trend

168 The result of filtering residuals is added to the fitted curve to obtain smoothed CO₂ and its long-term trend. The smoothed CO₂
169 comprises fitted trend, fitted seasonality and smoothed residuals (red line in Fig. 2d), the latter removes only short-term
170 variations or noise. The long-term trend comprises fitted trend and residual trend, which removes seasonal cycle and noise
171 (blue line in Fig. 2d).

172 2.3.3. Calculate CO₂ growth rate, G_{ATM}

173 The CO₂ growth rate (G_{ATM}) is determined by taking the first derivative of the long-term trend. However, the growth is made
174 up of discrete points, e.g. the black dots in Fig. 3a shows the trend points. In this case, a cubic spline interpolation is applied
175 to the trend points, in which the spline curve passes through each trend points, as the blue line in Fig. 3a. G_{ATM} is obtained by
176 taking the derivative of the spline at each trend point (Fig. 3b).



177

178

179

180

181


182

183

184

185

Figure 2. Example of analysed CO₂ data from station Pallas (PAL, Finland), illustrating GFIT curve fitting and filter method. Panel (a) shows monthly averaged CO₂ (dots), curve fitting with 2-degree polynomial and 4-degree harmonics (red line), and long-term trend estimated by a 2-degree polynomial (blue line). Panel (b) shows seasonality estimated by 4-degree harmonics. Panel (c) shows the residuals of raw data from the function fit (black dots), the red line is obtained by the short-term filter and the blue line is obtained by the long-term filter. The cyan dots show the residuals of raw data from the sum of fitted curve and smoothed residuals. Panel (d) shows final processed CO₂, which comprises fitted trend, fitted seasonality and smoothed residuals (red line). The blue line shows the final trend which comprises fitted trend and residuals trend.



186

187 **Figure 3.** Example of CO₂ growth rate, the raw data is the same as used in Fig. 2 from station Pallas (PAL, Finland).
 188 Panel (a) shows the trend points (black dots) and its cubic spline interpolation (blue line). Panel (b) shows the G_{ATM} at
 189 each trend point.

190 **3 Results**

191 Global averaged surface CO₂ and its G_{ATM} are calculated using the WDCGG method and our GFIT method based on the data
 192 from the GAW and CTE networks (Fig. 1). The different observation networks and their analysis methods are listed in Table
 193 1. We calculated the global means and its G_{ATM} by area-weighted averaging the zonal means over each latitudinal band (30°),
 194 following the same CO₂ averaging method as described in Tsutsumi et al. (2009). A bootstrap method is used to estimate the
 195 uncertainties of global CO₂ mean and its G_{ATM} , which is an almost identical uncertainty analysis as presented by Conway et
 196 al. (1994) who constructed 100 bootstrap networks for the NOAA analysis. We construct 200 bootstrap networks, consistent
 197 with the WDCGG analysis in Tsutsumi et al., (2009). For each bootstrap network, we randomly draw the same number of sites
 198 as the actual network (e.g. 139 sites for GAW network) with replacement from the actual network, which means some sites
 199 are missing whereas others will be represented twice or more often. We calculate global mean CO₂ mole fraction and its G_{ATM}
 200 for each network, and then calculate the statistics (i.e. mean and 68% confidence interval, CI) on the 200 networks. All
 201 uncertainties in this paper are reported as \pm 68% CI.


202 **Table 1. Description of the three observation networks and their analysis methods.**

Terminology	Description
NOAA network	NOAA network comprises MBL sites only (43 sites).
GAW network	The selected GAW global network (139 sites) includes both MBL sites and some continental sites.

CTE network	The CTE network serves as the global network for the CTE model evaluations (230 sites), comprises MBL sites and a more extensive inclusion of continental sites.
GAW (GFIT)	GAW network observations analysed using the GFIT method
GAW (WDCGG)	GAW network observations analysed using the WDCGG method without extrapolation
GAW (WDCGG+)	GAW network observations analysed using the WDCGG method with extrapolation
CTE_obs (GFIT)	CTE network observations analysed using the GFIT method. The observations come from the ObsPack data product (Schuldt et al., 2022)
CTE_output (GFIT)	CTE model output at the 230 sites (sampled at the same location, altitude, and time) analysed using the GFIT method
CTE_global (GFIT)	CTE model output for full global grids (averaged over the first three levels, 0 to 0.35 km Alt.) analysed using the GFIT method
MLO (GFIT)	Mauna Loa (MLO) observations analysed using the GFIT method
SPO (GFIT)	South Pole (SPO) observations analysed using the GFIT method

203

204 **3.1 Globally averaged surface CO₂ mole fraction and its G_{ATM}**



205

206

207

Figure 4. Comparison of globally and locally averaged CO₂ mole fraction (a) and its G_{ATM} (b) from 1980 to 2020. Panel (a) shows the global monthly CO₂ mole fraction from 139 GAW sites (estimated from observations only), 43 NOAA

208 **MBL sites and those from 230 sites used in CTE (either from observations or model output). The two local CO₂ mole**
209 **fractions are from Mauna Loa (MLO, cyan line) and South Pole (SPO, magenta line) stations, analysed using the GFIT**
210 **method. The red and blue lines show the CO₂ derived from GAW (GFIT) and GAW (WDCGG), respectively. The**
211 **green and orange lines show the CO₂ derived from CTE_obs (GFIT) and CTE_output (GFIT), respectively. The right**
212 **y-axis shows their difference from NOAA CO₂ mole fraction, and the dashed lines show the mean of the difference over**
213 **the available period. Panel (b) compares the corresponding global and local CO₂ growth rate, the legend refers to panel**
214 **(a). The shadow area shows the uncertainty as 68% confidence interval obtained by the bootstrap analysis.**

215 Fig. 4 presents a monthly comparison of globally and locally averaged CO₂ mole fractions and their G_{ATM} from 1980 to 2020.
216 The statistical metrics assessing the agreement of these monthly comparisons are available in Fig. 5 (for 2001-2020) and Fig.
217 S1 (for 1980-2020). The statistical metrics for the annual comparisons can be found in Fig. S2 (for 2001-2020) and Fig. S3
218 (for 1980-2020). They exhibit a similar pattern to the monthly comparisons (i.e. Fig.5 and Fig. S1).


219 Globally averaged monthly surface CO₂ mole fractions, derived from the GAW network (GAW (GFIT) or GAW (WDCGG)),
220 are significantly ($p < 0.05$) higher by 0.329-0.335 ppm during 1980-2020 (Fig. S1a) and 0.370-0.390 ppm during 2001-2020
221 (Fig. 5a) when compared to the NOAA analysis (Fig. 4a). This finding aligns with Tsutsumi et al., (2009), who reported a
222 0.350 ppm higher global average in the GAW network during 1983-2006. The higher estimate from the GAW network can be
223 attributed to the inclusion of more diverse sites, encompassing not only NOAA's MBL sites but also additional continental
224 sites (Fig. 1).

225 Both global CO₂ and its G_{ATM} derived from the GAW (GFIT) and GAW (WDCGG) are nearly overlapping (the red and blue
226 lines) in Fig. 4a and 4b. The statistical metrics (Fig. 5 and S1) indicate a high agreement ($ME < 0.020$ ppm, $RMSE < 0.145$ ppm,
227 $r > 0.999$ for CO₂ mole fraction; $ME < 0.005$ ppm yr⁻¹, $RMSE < 0.108$ ppm yr⁻¹, $r > 0.982$ for G_{ATM}) between these two methods,
228 which confirms that the GFIT method agrees well with WDCGG method without extrapolation. The WDCGG method with
229 extrapolation (i.e. GAW (WDCGG+)), which involves extrapolating the long-term trend of each station to match the period
230 of the most long-running station and adding it to the average seasonal variation to synchronize data period of all stations
231 (Tsutsumi et al., 2009), produces 0.096 ppm significantly ($p < 0.05$) higher values than the global monthly surface CO₂ mole
232 fraction derived from the GAW (WDCGG) during the common period 1984-2020 (Table S1). However, the extrapolation has
233 a minimal effect ($RMSE = 0.076$ ppm yr⁻¹, $ME = -0.011$ ppm yr⁻¹, Table S1) on the CO₂ growth rate.


234 Globally averaged monthly surface CO₂ derived from CTE_obs (GFIT) and CTE_output (GFIT) are 0.422 ppm (1980-2020,
235 Fig. S1) and 0.668 ppm (2001-2020, Fig. 5) significantly ($p < 0.05$) higher compared to the NOAA analysis, respectively (Fig.
236 4a). Comparing the global mean of CTE_obs (GFIT) with CTE_output (GFIT) during the common period of 2001-2020, we
237 observe a low bias (0.069 ppm in CTE_output, Fig 5a), which suggests that the CTE model results can reasonably reproduce
238 the global mean CO₂ levels. The global annual CO₂ mole fraction from CTE_obs (GFIT), CTE_output (GFIT) and CTE_global
239 (GFIT) is 0.367, 0.299 and 0.186 ppm significantly ($p < 0.05$) higher than the result of the GAW (GFIT), respectively (Fig 5a).
240 The higher global mean from CTE_obs (GFIT) and CTE_output (GFIT) can be attributed to the presence of more sites in the
241 Northern Hemisphere within the CTE network compared to the GAW network. The lower bias observed between GAW (GFIT)
242 and CTE_global (GFIT) suggests that the GAW network provides a good representation of the low-level atmosphere (i.e. 0 to
243 0.35 km altitude) at global scale, or the CTE model performs well in the low-level atmosphere.

244 A common approach to estimate global surface CO₂ mole fraction is by using one or two representative sites, such as Mauna
245 Loa (MLO) and South Pole (SPO). The globally averaged monthly surface CO₂ mole fractions, derived from the GAW, CTE,
246 and NOAA networks, are significantly ($p < 0.05$) lower by 0.46-0.88 ppm during 1980-2020 (Fig. S1a) and 0.45-1.19 during
247 2001-2020 (Fig. 5a) than the local CO₂ estimates solely based on MLO measurements. Conversely, these global monthly CO₂
248 mole fractions are significantly ($p < 0.05$) higher by 1.91-2.24 ppm during 1980-2020 (Fig. S1a) and 2.21-2.94 during 2001-
249 2020 (Fig. 5a) when compared to local measurements at SPO site. Furthermore, the global seasonal cycle leads the local cycle

250 at MLO by approximately one month (estimated by averaging the time difference between the peaks of their seasonal cycles).
 251 In contrast, the local cycle at SPO is not evident and is opposite to the global seasonal cycle (Fig. 4a).



252
 253 **Figure 5. Pair-wise statistical metrics assess the agreement of monthly global and local CO₂ mole fraction (ppm) and**
 254 **its G_{ATM} (ppm yr⁻¹) across various networks and methodologies (see Table 1 and Fig. 4) for the period 2001-2020. Panel**
 255 **(a) presents the Mean Error (ME) quantifying the difference for each pair, focusing on CO₂ mole fraction, while panel**
 256 **(b) does the same for G_{ATM}. The significance levels of paired t-test for ME are indicated as follows: * p<0.1, ** p<0.05,**
 257 ***** p<0.01. Panel (c) and (d) present the Root Mean Squared Error (RMSE) for CO₂ mole fraction and G_{ATM},**
 258 **respectively. Panel (e) and (f) present the Pearson Correlation Coefficient (r) for CO₂ mole fraction and G_{ATM},**
 259 **respectively.**



260
 261 **Figure 6. Trend analysis of the global CO₂ growth rate from 1980 to 2020. Panel (a) shows the trends of CO₂ growth**
 262 **rate for the GAW network (red line), the CTE network (green line) and the NOAA network (black line) during the**
 263 **whole period 1980-2020, the CO₂ growth rate is derived from GAW (GFIT), CTE_obs (GFIT) and NOAA analysis (Fig.**
 264 **4b). Panel (b) shows the trend of CO₂ growth rate for each month during 1980-2020, calculated as the derivative of the**
 265 **growth rate. The grey bands mark the period of three strong El Niño events, i.e. 1987-1988, 1997-1998 and 2014-2016.**

266 Despite differences in the global averaged surface CO₂ mole fractions derived from different networks and analysis methods,
 267 the G_{ATM} derived from GAW network, CTE network and its model output, and NOAA network exhibits strong agreement
 268 during 1980-2020 ($ME < 0.031 \text{ ppm yr}^{-1}$, $RMSE < 0.217 \text{ ppm yr}^{-1}$, $r > 0.948$, Fig. 4b and S1). The differences in the G_{ATM} remain
 269 below $0.023 \text{ ppm yr}^{-1}$ during 2001-2020, with low or no significance level (Fig. 5b), especially when comparing the annual
 270 G_{ATM} (Fig. S2b). Furthermore, over the long-term period of 40 years, the estimated local growth rate at MLO ($ME < 0.046 \text{ ppm}$
 271 yr^{-1} higher, $RMSE < 0.272 \text{ ppm yr}^{-1}$, $r > 0.915$) and SPO ($ME < 0.049 \text{ ppm yr}^{-1}$ lower, $RMSE < 0.305 \text{ ppm yr}^{-1}$, $r > 0.888$) behaves
 272 similarly to the G_{ATM} derived from GAW, CTE and NOAA network (Fig. 4b and S1). However, noticeable monthly differences
 273 between the local and global growth rates, deviating up to approximately 0.8 ppm yr^{-1} , and time shifts are observed (Fig. 4b).
 274 The trend analysis reveals that with development of continental sites, the slope of the trend of annual global CO₂ mole fraction
 275 changes from NOAA network ($1.832 \pm 0.029 \text{ ppm yr}^{-1}$) to CTE network ($1.859 \pm 0.029 \text{ ppm yr}^{-1}$) during 1980-2020 (Fig. S4).
 276 However, the G_{ATM} increased steadily at a rate of $0.030 \pm 0.002 \text{ ppm per year}$ each year from 1980 to 2020 (Fig. 6a), based on
 277 the observations from the three networks (i.e. GAW, CTE and NOAA). This implies that over long-term period (here 40 years),
 278 the networks with and without continental sites exhibits the same trend of the G_{ATM} and has little effect on the transient change
 279 in the rate of CO₂ increase in the atmosphere. Hence, the role of CO₂ advective transport and mixing in estimating the long-
 280 term change of the G_{ATM} appears negligible. However, a notable difference emerges in the short-term (here one month) change
 281 of the G_{ATM} between the networks with and without continental sites (Fig. 6b). El Niño events are known to diminish net global


282 C uptake (due to factors such as droughts, floods and fires) while increasing global CO₂ growth rate (Sarmiento et al., 2010).
 283 During three strong El Niño events, which are marked as grey bands in Fig. 6b, the G_{ATM} derived from the GAW and CTE
 284 network (red and green lines) begins to increase approximately 1-2 months (Table S2) earlier before the El Niño events (marked
 285 as blue circles in Fig. 6b) and reaches its peak approximately 1-2 months (Table S2) earlier during the El Niño events (marked
 286 as orange circles in Fig. 6b), compared to the G_{ATM} derived from the NOAA network (black line). This suggests that continental
 287 sites can aid in the early detection of G_{ATM} changes resulting from changes in biogenic emission or uptake. The CTE network
 288 (green line) even detects the change one month earlier than the GAW network (red line) e.g. for the El Niño 1997-1998 event
 289 (Fig. 6b, Table S2). This earlier detection is attributed to the inclusion of even more continental sites in the CTE network (Fig.
 290 1), although the more continental sites also induce the greater variability.

291 Table 2 presents the global annual CO₂ mole fraction and its G_{ATM} derived from GAW (GFIT), along with the uncertainty
 292 estimates using the bootstrap method. The global average surface CO₂ mole fraction has increased from 339.17±0.38 ppm in
 293 1980 to 413.06±0.16 ppm in 2020. Notably, the uncertainty is greater before 1990, primarily due to the limited number of
 294 measurement stations worldwide during that period. The average G_{ATM} for the two decades before 2000 is approximately
 295 1.54±0.08 ppm yr⁻¹. However, in the following two decades, it has experienced increases, reaching 1.91±0.05 ppm yr⁻¹ during
 296 2000-2009 and further rising to 2.41±0.06 ppm yr⁻¹ during 2010-2019 (Table 2, Fig. S5).

297 **Table 2. Annual global averaged CO₂ mole fraction (Mean, ppm) and its G_{ATM} (ppm yr⁻¹) derived from GAW**
 298 **observations using GFIT method. U(Mean) and U(G_{ATM}) respectively indicate the uncertainty of Mean and its G_{ATM} as**
 299 **68% confidence interval. The annual value is averaged over the monthly values of the year.**

Year	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989
<i>Mean</i>	339.17	340.16	341.03	342.59	344.46	345.69	347.08	348.99	351.45	353.15
<i>U(Mean)</i>	0.38	0.24	0.19	0.24	0.26	0.22	0.14	0.15	0.12	0.15
<i>G_{ATM}</i>	1.65	1.07	0.88	2.02	1.32	1.38	1.55	2.38	2.08	1.23
<i>U(G_{ATM})</i>	0.12	0.10	0.15	0.13	0.08	0.11	0.14	0.08	0.09	0.06
Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
<i>Mean</i>	354.22	355.64	356.37	357.09	358.51	360.52	362.27	363.40	366.14	368.10
<i>U(Mean)</i>	0.10	0.11	0.10	0.10	0.11	0.12	0.12	0.10	0.10	0.10
<i>G_{ATM}</i>	1.41	1.03	0.65	1.22	1.72	2.06	1.16	1.82	2.89	1.34
<i>U(G_{ATM})</i>	0.08	0.06	0.05	0.05	0.05	0.08	0.07	0.05	0.05	0.05
Year	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
<i>Mean</i>	369.30	370.77	372.92	375.45	377.22	379.28	381.38	383.20	385.26	386.78
<i>U(Mean)</i>	0.12	0.11	0.10	0.10	0.10	0.10	0.09	0.10	0.10	0.11
<i>G_{ATM}</i>	1.58	1.58	2.33	2.17	1.66	2.42	1.75	2.20	1.71	1.68
<i>U(G_{ATM})</i>	0.05	0.06	0.06	0.04	0.04	0.03	0.05	0.04	0.05	0.04
Year	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
<i>Mean</i>	389.01	390.97	393.14	396.00	397.79	400.12	403.47	405.70	407.93	410.57
<i>U(Mean)</i>	0.12	0.12	0.14	0.11	0.10	0.10	0.11	0.09	0.10	0.13
<i>G_{ATM}</i>	2.32	1.73	2.74	2.30	1.91	2.98	2.95	2.04	2.50	2.61
<i>U(G_{ATM})</i>	0.05	0.06	0.09	0.05	0.04	0.05	0.06	0.06	0.07	0.05
Year	2020									
<i>Mean</i>	413.06									
<i>U(Mean)</i>	0.16									
<i>G_{ATM}</i>	2.60									

300 **3.2 Vertical profile of global CO₂ mole fraction**




301
 302 **Figure 7. Global vertical profile of CO₂ mole fraction derived from CTE model output. Panel (a) presents the vertical**
 303 **profile in 2020. Panel (b) presents the difference of the vertical profile between 2001 and 2020. Panel (c) presents the**
 304 **annual mean vertical profile from 2001 to 2020, the dots mark CTE vertical level heights and lines are the linear**
 305 **interpolation between the heights.**

306 The CTE model simulates CO₂ mole fraction on global 3D grids, enabling us to visualize the modelled vertical CO₂ profile.
 307 In the lower atmosphere, highest CO₂ mole fraction is found in the Northern mid-latitude region (dark red between 30 °N and
 308 40 °N, Fig. 7a). This area experiences more anthropogenic emissions, which are subsequently transported towards both
 309 northern and southern latitudes. The latitudinal and interhemispheric gradient of atmospheric CO₂, as shown in Fig. 7a, is
 310 influenced not only by differences in the latitudinal and interhemispheric fossil fuel emissions and seasonal exchanges with
 311 terrestrial biota (Denning et al., 1995), but also by atmospheric transport (Patra et al., 2011). As altitude increases, the gradient
 312 between the Northern and Southern hemisphere becomes small and levels out at higher altitudes (e.g. >50 km). When

313 comparing the vertical profile change between 2001 and 2020 (Fig. 7b and 7c), we observe that the CO₂ mole fraction increases
 314 slowly in the higher atmosphere (>25 km altitude) compared to the lower atmosphere (<25 km altitude). Fig. 7c shows that the
 315 vertical gradient (difference between 50 km and 0.05 km) changes from approximately 5 ppm in 2001 to around 13 ppm in
 316 2020. The high vertical gradient in 2020 reflects the accumulation of CO₂ in the lower atmosphere, resulting from continuous
 317 CO₂ emissions from the surface during 2001-2020 and slow vertical transport. The low vertical gradient in 2001 is partly due
 318 to lower surface emissions.

319 Pressure-weighted average CO₂ mole fraction in the lower atmosphere (0 to 0.35 km altitude) and the entire atmosphere are
 320 calculated from CTE output. The annual absolute change in CO₂ mole fraction, computed as the difference between annual
 321 means, is more pronounced in the lower atmosphere (orange bars in Fig. S6a) than in the entire atmosphere (blue bars in Fig.
 322 S6a). The reason is that the entire atmosphere has a larger air volume than the lower atmosphere, and changes in the surface
 323 CO₂ sinks and sources are diluted due to atmospheric horizontal and vertical transport. The CO₂ annual absolute change derived
 324 from GAW (GFIT), GAW (WDCGG) and NOAA (represented by red, purple and brown bars in Fig S6a) shows small positive
 325 or negative differences from the CTE_output (GFIT) and CTE_global (GFIT) across different years. However, over the long
 326 term (e.g. on a decadal scale, 2001-2010 and 2011-2020), the CTE model-derived changes in lower and entire atmospheric
 327 CO₂ shows good agreement (<0.09 ppm yr⁻¹) with the surface observation-based estimate, especially for lower atmospheric
 328 CO₂ (<0.07 ppm yr⁻¹). In Fig. S6b, the interannual variability (IAV) of CO₂ mole fraction derived from CTE model follows a
 329 similar temporal pattern as the observation-based IAV derived from the GAW and NOAA network, especially the IAV of the
 330 low-level atmosphere (orange bars) exhibits strong agreement with the observation-based IAV (r>0.971, RMSE<0.178 ppm).

331 3.3 Relationship between the surface CO₂ mole fraction and atmospheric CO₂ mass



332
 333 **Figure 8. Relationship between the monthly surface CO₂ mole fraction and atmospheric CO₂ mass. The atmospheric**
 334 **CO₂ mass calculated from the 3D CTE output. In panel (a), the monthly surface CO₂ derived from the CTE_output**
 335 **(GFIT), GAW (GFIT) and NOAA analysis, presented as blue, red and green dots, respectively. Panel (b) compares the**
 336 **corresponding interannual variability (IAV) of the atmospheric CO₂ mass and the surface CO₂. The IAV is calculated**
 337 **as the anomaly departure from a quadratic trend.**

338 The atmospheric CO₂ mass, calculated from the CTE output as a function of air mass and CO₂ concentration (Text S3), has
 339 increased from 789.46 PgC in 2001 to 877.88 PgC in 2020 (Fig. S7a). The spatial distribution of the atmospheric CO₂ mass is
 340 presented in Fig. S7b and Fig. S7c. Monthly global surface CO₂ mole fraction derived from CTE_output (GFIT) and GAW
 341 (GFIT), represented as red and blue dots in Fig. 8a, exhibit a similar linear relationship with the monthly atmospheric CO₂
 342 total mass, both showing the same slope of 2.08 ± 0.01 PgC ppm⁻¹. Similarly, NOAA CO₂ (green dots, Fig. 8a) also demonstrates

343 a comparable linear relationship with a slope of 2.09 ± 0.01 PgC ppm⁻¹. Notably, the slopes or conversion factors in Fig. 8a are
344 slightly lower than the factor 2.12 PgC ppm⁻¹ used in Ballantyne et al. (2012) for the period 1980-2010. This minor difference
345 in the conversion factor is expected, considering the different model and data used.
346 We further compare the interannual variability (IAV), calculated as the anomaly departure from a quadratic trend, of the
347 atmospheric CO₂ mass and the surface CO₂ (Fig. 8b). The coefficient of the linear relationship closely approaches ~ 1.0 ,
348 indicating the temporal changes in atmospheric CO₂ mass align with the temporal changes in surface CO₂ mole fraction. The
349 CO₂ IAV based on the NOAA network exhibits a slightly closer relationship ($r=0.938$) with the CTE atmospheric CO₂ mass
350 estimates than the GAW ($r=0.861$) and CTE ($r=0.812$) networks. This finding is consistent with the long atmospheric residence
351 time and well-mixed nature of CO₂ in the NOAA network. Overall, the relationship found in Fig. 8 implies that the current
352 surface CO₂ network can effectively serve as an indicator of the CO₂ mass changes throughout the entire atmosphere through
353 a linear relationship.

354 **4 Discussion**

355 Over the past few decades, observational networks have been extended beyond the NOAA MBL network to include more
356 continental sites, such as in the GAW and CTE networks (Fig. 1). These expansions aim to better monitor global CO₂
357 concentrations and quantify CO₂ sources and sinks. While the continental observations encompass contributions from both
358 substantial sources of anthropogenic emissions and sources/sinks from terrestrial vegetation and soil, these continental
359 observations consistently yield a higher global surface CO₂ mole fraction in the overall global CO₂ analysis, indicating that
360 they are influenced by a bigger net source. We find that the global mean derived from the GAW network is consistently 0.329
361 (GFIT method) or 0.335 (WDCGG method) ppm higher than that derived from the NOAA network during 1980-2020.
362 Similarly, Tsutsumi et al. (2009) reported a roughly 0.350 ppm higher mole fraction in the GAW network for years 1983-2006.
363 Notably, the CTE network leads to an even higher global mean (0.422 ppm during 1980-2020), which is likely due to more
364 observational sites locate in the Northern Hemisphere, where the highest anthropogenic emissions occur. This also explains
365 the large fluctuation of CO₂ concentrations observed during the winters and summers during 2001-2020 (Fig. 4a). In the future,
366 with the addition of new observation sites, particularly in the Northern Hemisphere, to the existing observational network (e.g.
367 GAW network), we expect that this would lead to higher global surface CO₂ levels and a greater amplitude in the global CO₂
368 seasonal cycle in the global CO₂ analysis.

369 Although Friedlingstein et al. (2022) reported a 5.4% drop (~ 0.52 PgC) in fossil fuel CO₂ emissions in 2020 (due to restrictions
370 on e.g. transport, industry, power etc during the COVID-19 pandemic), the increase in annual CO₂ from 2019 to 2020
371 (2.60 ± 0.16 ppm yr⁻¹) remains at a similar level as from 2018 to 2019 (2.61 ± 0.05 ppm yr⁻¹). In principle, an equivalent drop of
372 roughly 0.25 ppm yr⁻¹ (according to the conversion factor 2.08 PgC ppm⁻¹ in Fig. 8a) or roughly 0.13 ppm yr⁻¹ (according to
373 the annual absolute change, red bars in Fig. S6a) in the growth rate should be visible for period 2019-2020 due to the declined
374 CO₂ emissions. However, such a short-term human activity induced change in the CO₂ growth rate may be hidden by the
375 natural variability. The bootstrap analysis is used in this study (also in Conway et al., (1994) and Tsutsumi et al., (2009)) to
376 estimate the uncertainty of the CO₂ temporal mean and its growth rate and to assess how sensitive the global value is to the
377 distribution of sampling sites. The relatively large uncertainty (± 0.16 ppm yr⁻¹) at the end of 2020 compared to previous years
378 (Table 2) is likely due to an end-effect associated with the curve fitting and filter procedure. The end-effect is a tendency for
379 the growth rate to converge toward the mean value at the end of the record (Conway et al., 1994). Therefore, Conway et al.
380 (1994) suggested that the growth rate curves for the last 6 months should be viewed with caution. Reducing the end-effect
381 requires further study, such as using machine learning or bias-correction methods to extrapolate the smoothed trend for a short
382 period (e.g. one year) before and after. This extrapolated portion is used exclusively for calculating local mole fraction and
383 growth rate, while it is not included in the global or zonal average, as it could introduce additional uncertainty.

384 Extrapolation beyond the measurement period extends knowledge gained from a limited period of measurements. During a
385 limited measurement period, we can define the average seasonality, long-term trend, and short-term variation at a measurement
386 site. The long-term trend of an individual site can be extrapolated by various methods, such as referring to the latitude reference
387 time series (Masarie and Tans, 1995) or calculating the mean long-term trend over sites within a certain latitudinal zone (e.g.
388 30°) (Tsutsumi et al., 2009). This extrapolated trend is then combined with the average seasonality to produce estimates beyond
389 the measurement period. However, the extrapolation process relies on the assumption that the relationship of an individual site
390 to the latitude reference remains invariant in time, while in reality the relationship between nearby sites is continuously
391 changing (Masarie and Tans, 1995). In addition, the short-term variation is often ignored or estimated from nearby sites,
392 introducing extra uncertainty into the extrapolation process. In this study, we find that the WDCGG method with extrapolation
393 (GAW (WDCGG+)) results in a global surface CO₂ mole fraction approximately 0.096 ppm higher than the WDCGG method
394 without extrapolation (GAW (WDCGG)) using the same GAW observations, although the extrapolation has a minor effect on
395 the growth rate (Table S1). Therefore, we chose not to use extrapolation beyond the measurement period in our analysis. As
396 the number of long-term measurements increases, the need for such extrapolation becomes less necessary.

397 Our analysis shows that basing the CO₂ growth rate on GAW surface observations does not introduce a large bias (with an
398 average agreement within 0.016 ppm yr⁻¹) compared to a full atmospheric analysis (Fig. 4b and 5). This full atmosphere CO₂
399 was provided by the CTE model, in which the global annual mean CO₂ is significantly overestimated compared to GAW
400 observations (e.g. 0.299 ppm higher in CTE_output (GFIT), or 0.186 ppm higher in the CTE_global (GFIT) during 2001-
401 2020). The overestimate derived from the CTE_output (GFIT) is mainly due to more sites in the Northern Hemisphere in the
402 CTE network than in the GAW network. The lower overestimate derived from the CTE_global (GFIT) implies that the biases
403 in CTE outputs are not uniform spatially and tend to balance out. We estimate the CTE bias by comparing the observations
404 and CTE outputs at the same sites, which results in a 0.069 ppm low bias derived from the CTE outputs in calculating the
405 global surface CO₂ mole fraction.

406 The local growth rate at MLO and SPO generally behaves similarly to the global growth rate derived from the GAW, CTE,
407 and NOAA networks (Fig. 4b and S1). However, the local CO₂ mole fraction and its seasonal cycle noticeably differ from
408 global estimates derived from different observational networks. In this regard, the utilization of individual sites for the
409 evaluation of the global average mole fraction and its growth rate is not precise and can only be used for illustration rather
410 than as a substitute for the proper global average calculation. The local observation sites, often situated away from significant
411 local sources and sinks, such as MLO, provide long-term and high-quality data, serving as reference data for global CO₂ mole
412 fraction. However, a single observation site cannot capture the CO₂ spatial variability, transport, and mixing. To overcome
413 these limitations, global CO₂ trends and variations are best assessed by integrating data from multiple sources and locations.

414 Different observational networks (i.e. NOAA network, GAW network and CTE network) are analysed in this study, revealing
415 differences in calculated global surface CO₂ mole fractions equivalent to the current atmospheric growth rate over a three-
416 month period. This suggests that the station selection, especially if and how many continental observations are used, has some
417 influence on the derived global surface CO₂ levels, but it is not particularly strong. Nowadays, an increasing number of
418 continental observations are established to monitor biogenic sources and sinks, providing further insight into the climate
419 change and the associated ecosystem processes (Ciais et al., 2005, Ramonet et al., 2020). Such continental observations carry
420 more variability in measurements than the marine observations, which requires caution when including them in the mix of
421 stations used to determine global surface CO₂ mole fraction. Our study demonstrates that continental sites can help early detect
422 changes in CO₂ growth rate caused by biogenic emission change, such as those resulting from El Niño events. Furthermore,
423 the current observational networks (with and without continental sites) and CTE model show a good agreement on the global
424 CO₂ growth rate, with low or no significant differences within 0.023 ppm yr⁻¹ during 2001-2020 and 0.031 ppm yr⁻¹ during
425 1980-2020. This implies that the current observation networks (as shown in Fig. 1, represent for various ecosystems, sinks,

426 sources, and latitudes) have a similar good capacity to capture changes in the global surface CO₂, although there is the spatial
427 and temporal variability in the CO₂ growth rate (e.g. Conway et al. 1994).

428 We also notice that the uncertainty in global CO₂ growth rate is approximately 0.07 ppm yr⁻¹, as derived from GAW (GFIT)
429 and averaged over 1980-2020 (Table 2). To reduce the uncertainty to 0.02 ppm yr⁻¹ (equivalent to 1% of the global CO₂
430 growth rate), in principle it would theoretically requires adding more stations to the current observation network. We conducted
431 an experiment that demonstrates how the uncertainty of the global CO₂ growth rate exponentially increases as the number of
432 land observation sites decreased (Fig. S8). According to our experiment, to achieve the goal of reducing the uncertainty to 0.02
433 ppm yr⁻¹, 332 land observation sites are required (Fig. S8). However, the required number of sites also depends on their
434 measurement accuracy, consistency, and geographical distribution (i.e. CO₂ footprint coverage of observation network, and
435 the importance of the network design was addressed by Storm et al. (2022)).

436 **5 Conclusions**

437 The WMO GAW CO₂ network documents the gradual global accumulation of CO₂ in the atmosphere due to human activities.
438 It has been used to assess the large-scale and long-term environmental consequence of fossil CO₂ emission and land use
439 changes. The high-quality observations conducted by the WMO GAW network include not only background stations (most of
440 NOAA MBL stations) but also continental stations. This comprehensive network enables proper global average calculation.
441 Furthermore, the WMO has initiated a new program, Global Greenhouse Gas Watch (GGGW), with the aim of establishing a
442 reference network. This network will be built on the high-quality observations already performed under the WMO GAW
443 program that follows consistent good practices and standards. Although the current monitoring networks have limitations in
444 terms of geographical coverage, data consistency, and long-term measurements, they are well-equipped and have the capacity
445 to effectively represent global surface CO₂ mole fraction and its growth rate and trends in atmospheric CO₂ mass changes. The
446 three different analysis methods yield very similar global CO₂ increases from 2001 to 2020, which gives confidence in using
447 any one of them in climate change studies. Continuous monitoring of atmospheric CO₂, based on the current GAW network
448 together with reliable global data integration methods, provides essential information. This includes understanding trends in
449 atmospheric CO₂ concentration, assessing the impacts of past policies, identifying high-emission areas, informing climate
450 models, forecasting future scenarios, and raising public awareness. Policymakers can rely on this information to support their
451 efforts in mitigating the global warming.

452 **6 Data and Code Availability**

453 All data and code necessary to calculate the global mean surface CO₂ mole fraction and Atmospheric CO₂ mass is freely
454 available from ICOS Carbon Portal [<https://doi.org/10.18160/Q788-9081>]. The file list of results and code can be found in
455 Text S4.

456 **Acknowledgments**

457 We acknowledge Ingrid Lujikx for providing the TM5 data, WMO GAW Principal Investigators of the WMO GAW station
458 network for providing the observational data, Ed Dlugokencky for providing NOAA data and comments. Thanks to the support
459 from ICOS, GAW, NOAA and CTE group.

460 **Author Contributions**

461 A.V. and Z.W. designed this study in discussion with Y.S., O.T and U.K..

462 Z.W. performed analysis and led the writing.
463 Y.S., Y.N. and A.O. provided the GAW data, and commented on the manuscript.
464 W.P. and R.K. provided CTE model results and relevant ObsPack data and commented on the manuscript.
465 X.L. provided NOAA data and commented on the manuscript.
466 All authors contributed to the writing of the paper and interpretation of the results.

467 **Competing Interests**

468 The authors declare no competing interests.

469

470 **Financial support**

471 This research is a part of ICOS core work, there is no grant reference no.

472 **References**

- 473 Ballantyne, A. á., Alden, C. á., Miller, J. á., Tans, P. á. & White, J. 2012. Increase in observed net carbon dioxide uptake by
474 land and oceans during the past 50 years. *Nature*, 488, 70-72. <https://doi.org/10.1038/nature11299>
- 475 Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C. & Carrara,
476 A. 2005. Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature*, 437, 529-
477 533. <https://doi.org/10.1038/nature03972>
- 478 Conway, T. J., Tans, P. P., Waterman, L. S., Thoning, K. W., Kitzis, D. R., Masarie, K. A. & Zhang, N. 1994. Evidence for
479 interannual variability of the carbon cycle from the National Oceanic and Atmospheric Administration/Climate
480 Monitoring and Diagnostics Laboratory global air sampling network. *Journal of Geophysical Research: Atmospheres*,
481 99, 22831-22855. <https://doi.org/10.1029/94JD01951>
- 482 Denning, A. S., Fung, I. Y. & Randall, D. 1995. Latitudinal gradient of atmospheric CO₂ due to seasonal exchange with land
483 biota. *Nature*, 376, 240-243. <https://doi.org/10.1038/376240a0>
- 484 Eyring, V., Gillett, K., Achuta Rao, R., Barimalala, M., Barreiro Parrillo, N., Bellouin, C., Cassou, P., Durack, Y., Kosaka, S.
485 & Mcgregor, S. 2021. Human Influence on the Climate System. *Climate Change 2021: The Physical Science Basis.*
486 *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.*
487 <https://www.ipcc.ch/report/ar6/wg1/>
- 488 Friedlingstein, P., Jones, M. W., O'sullivan, M., Andrew, R. M., Bakker, D. C., Hauck, J., Le Quéré, C., Peters, G. P., Peters,
489 W. & Pongratz, J. 2022. Global carbon budget 2021. *Earth System Science Data*, 14, 1917-2005.
490 <https://doi.org/10.5194/essd-14-1917-2022>
- 491 Gulev, S., Thorne, P., Ahn, J., Dentener, F., Domingues, C. M., Gong, S. G. D., Kaufman, D., Nnamchi, H., Rivera, J. &
492 Sathyendranath, S. 2021. Changing state of the climate system. *Climate Change 2021: The Physical Science Basis.*
493 *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.*
494 <https://www.ipcc.ch/report/ar6/wg1/>
- 495 Hall, B. D., Crotwell, A. M., Kitzis, D. R., Mefford, T., Miller, B. R., Schibig, M. F. & Tans, P. P. 2021. Revision of the World
496 Meteorological Organization Global Atmosphere Watch (WMO/GAW) CO₂ calibration scale. *Atmospheric*
497 *Measurement Techniques*, 14, 3015-3032. <https://doi.org/10.5194/amt-14-3015-2021>
- 498 Krol, M., De Bruine, M., Killaars, L., Ouwersloot, H., Pozzer, A., Yin, Y., Chevallier, F., Bousquet, P., Patra, P. & Belikov,
499 D. 2018. Age of air as a diagnostic for transport timescales in global models. *Geoscientific Model Development*, 11,
500 3109-3130. <https://doi.org/10.5194/gmd-11-3109-2018>
- 501 Krol, M., Houweling, S., Bregman, B., Van Den Broek, M., Segers, A., Van Velthoven, P., Peters, W., Dentener, F. &
502 Bergamaschi, P. 2005. The two-way nested global chemistry-transport zoom model TM5: algorithm and applications.
503 *Atmospheric Chemistry and Physics*, 5, 417-432. <https://doi.org/10.5194/acp-5-417-2005>
- 504 Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H. &
505 Kawamura, K. 2008. High-resolution carbon dioxide concentration record 650,000–800,000 years before present.
506 *nature*, 453, 379-382. <http://doi.org/10.1038/nature06949>
- 507 Masarie, K. A. & Tans, P. P. 1995. Extension and integration of atmospheric carbon dioxide data into a globally consistent
508 measurement record. *Journal of Geophysical Research: Atmospheres*, 100, 11593-11610.
509 <https://doi.org/10.1029/95JD00859>
- 510 Patra, P. K., Houweling, S., Krol, M., Bousquet, P., Belikov, D., Bergmann, D., Bian, H., Cameron-Smith, P., Chipperfield,
511 M. P. & Corbin, K. 2011. TransCom model simulations of CH₄ and related species: linking transport, surface flux
512 and chemical loss with CH₄ variability in the troposphere and lower stratosphere. *Atmospheric Chemistry and*
513 *Physics*, 11, 12813-12837. <https://doi.org/10.5194/acp-11-12813-2011>

514 Peters, W., Krol, M., Dlugokencky, E., Dentener, F., Bergamaschi, P., Dutton, G., Velthoven, P. V., Miller, J., Bruhwiler, L.
515 & Tans, P. 2004. Toward regional-scale modeling using the two-way nested global model TM5: Characterization of
516 transport using SF6. *Journal of Geophysical Research: Atmospheres*, 109. <https://doi.org/10.1029/2004JD005020>

517 Press, W. H., Teukolsky, S. A., Vetterling, W. T. & Flannery, B. P. 1988. *Numerical recipes in C 1st edition: The art of*
518 *scientific computing*, Cambridge university press.
519 <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=e05e217a58481314e070b6c8899791faa91a3e27>

520 Ramonet, M., Ciais, P., Apadula, F., Bartyzel, J., Bastos, A., Bergamaschi, P., Blanc, P., Brunner, D., Caracciolo Di
521 Torchiarolo, L. & Calzolari, F. 2020. The fingerprint of the summer 2018 drought in Europe on ground-based
522 atmospheric CO2 measurements. *Philosophical Transactions of the Royal Society B*, 375, 20190513.
523 <https://doi.org/10.1098/rstb.2019.0513>

524 Sarmiento, J. L., Gloor, M., Gruber, N., Beaulieu, C., Jacobson, A. R., Mikaloff Fletcher, S. E., Pacala, S. & Rodgers, K. 2010.
525 Trends and regional distributions of land and ocean carbon sinks. *Biogeosciences*, 7, 2351-2367.
526 <https://doi.org/10.5194/bg-7-2351-2010>

527 Schuldt, K., Mund, J. & Luijkx, I. 2022. Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1957–
528 2021; obspack_co2_1_GLOBALVIEWplus_v8. 0_2022-08-27, NOAA Global Monitoring Laboratory [data set].
529 *NOAA Global Monitoring Laboratory [data set]*, 10, 20220808. <http://doi.org/10.25925/20220808>

530 Storm, I., Karstens, U., D'onofrio, C., Vermeulen, A. & Peters, W. 2022. A view of the European carbon flux landscape through
531 the lens of the ICOS atmospheric observation network. *Atmospheric Chemistry and Physics Discussions*, 1-25.
532 <https://doi.org/10.5194/acp-23-4993-2023>

533 Tans, P. P., Thoning, K. W., Elliott, W. P. & Conway, T. J. 1990. Error estimates of background atmospheric CO2 patterns
534 from weekly flask samples. *Journal of Geophysical Research: Atmospheres*, 95, 14063-14070.
535 <https://doi.org/10.1029/JD095iD09p14063>

536 Thoning, K. W., Tans, P. P. & Komhyr, W. D. 1989. Atmospheric carbon dioxide at Mauna Loa Observatory: 2. Analysis of
537 the NOAA GMCC data, 1974–1985. *Journal of Geophysical Research: Atmospheres*, 94, 8549-8565.
538 <https://doi.org/10.1029/JD094iD06p08549>

539 Tsutsumi, Y., Mori, K., Hirahara, T., Ikegami, M. & Conway, T. J. 2009. Technical Report of Global Analysis Method for
540 Major Greenhouse Gases by the World Data Center for Greenhouse Gases (WMO/TD-No. 1473). *GAW Report No.*
541 *184. Geneva, WMO*, 1-23. https://library.wmo.int/index.php?lvl=notice_display&id=12631

542 Van Der Laan-Luijkx, I. T., Van Der Velde, I. R., Van Der Veen, E., Tsuruta, A., Stanislawska, K., Babenhauserheide, A.,
543 Zhang, H. F., Liu, Y., He, W. & Chen, H. 2017. The CarbonTracker Data Assimilation Shell (CTDAS) v1. 0:
544 implementation and global carbon balance 2001–2015. *Geoscientific Model Development*, 10, 2785-2800.
545 <https://doi.org/10.5194/gmd-10-2785-2017>

546 WMO, G. 2022. The state of greenhouse gases in the atmosphere based on global observations through 2021. *WMO*
547 *Greenhouse Gas Bulletin*. https://library.wmo.int/doc_num.php?explnum_id=11352

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567