

Discriminating viscous creep features (rock glaciers) in mountain permafrost from debris-covered glaciers – a commented test at the Gruben and Yerba Loca sites, Swiss Alps and Chilean Andes

Deleted: D

5 Wilfried Haebler¹, Lukas U. Arenson², Julie Wee³, Christian Hauck³, and Nico Mölg⁴

¹Geography Department, University of Zurich, 8057 Zurich, Switzerland

²BGC Engineering, Suite 500 – 980 Howe Street, Vancouver, BC V6Z 0C8, Canada

³Department of Geosciences, University of Fribourg, 1700 Fribourg, Switzerland

⁴enveo, Fürstenweg 176, 6020 Innsbruck, Austria

10

Correspondence to: wilfried.haebler@geo.uzh.ch

Abstract. Viscous flow features in perennially frozen talus/debris called rock glaciers are being systematically inventoried as part of global climate-related monitoring of mountain permafrost. In order to avoid duplication and confusion, guidelines were developed by the International Permafrost Association for discriminating between the permafrost-related landform “rock glacier” and the glacier-related landform “debris-covered glacier”. In two regions covered by detailed field measurements, the corresponding data- and physics-based concepts are tested and shown to be adequate. Key physical aspects, which cause the striking morphological and dynamic difference between the two phenomena/landforms concern:

15

- tight mechanical coupling of the surface material to the frozen rock-ice mixture in the case of rock glaciers as contrasting with essential non-coupling of debris to glaciers they cover;
- talus-type advancing fronts of rock glaciers exposing fresh debris material from inside the moving frozen bodies as opposed to massive surface ice exposed by **increasingly rare** advancing fronts of debris-covered glaciers; and
- increasing creep rates and continued advance of rock glaciers as convex landforms with structured surfaces versus predominant slowing down and disintegration of debris-covered glaciers as **often** concave landforms with primarily chaotic surface structure.

20

25

Where debris-covered surface ice is, or has recently been, in contact with thermally-controlled subsurface ice in permafrost, complex conditions and interactions can develop morphologies beyond simple “either-or”-type landform classification. In such cases, remains of buried surface ice mostly tend to be smaller than the lower size limit of “glaciers” as applied in glacier inventories, and to be far thinner than the permafrost in which they are embedded.

30 1 Introduction

Application of modern technologies enables climate-related science and long-term monitoring of mountain permafrost on Earth to evolve in a rapid, fascinating and important way. Absolute age determination using radiocarbon, luminescence or exposure dating (e.g., Haeberli et al., 2003; Fuchs et al., 2013; Krainer et al., 2015; Nesje et al., 2021; Amschwand et al., 2021) document the multi-millennial, typically Holocene time scale related to the often-spectacular landforms usually called “rock glaciers”, which reflect cumulative deformation through slow viscous creep of perennially frozen talus/debris rich in ice (Wahrhaftig and Cox, 1959; Haeberli, 1985; Barsch, 1996; Haeberli et al., 2006; Berthling, 2011; Kääh, 2013). Sophisticated borehole observations and laboratory experiments together with results from high-precision geodetic-photogrammetric and interferometric measurements (e.g., Arenson et al., 2002; Roer et al., 2008; Springman et al., 2012; Cicoira et al., 2021; Kääh et al., 2021; Kaufmann et al., 2021; Cusicanqui et al., 2021; Noetzli et al., 2021; Thibert and Bodin, 2022; Fleischer et al., 2023) provide detailed and crucial quantitative information about the corresponding flow mechanisms and flow fields with remarkable indications of recent warming-related flow acceleration.

Increasing creep rates of rock glacier permafrost can be attributed to permafrost warming down to tens of meters below surface as systematically measured in boreholes (Harris et al., 2009; Etzelmüller et al., 2020; Hoelzle et al., 2022). Warming-induced reduction in strength of subsurface frozen materials parallels increasing amounts of unfrozen water as reflected by repeat geophysics (Mollaret et al., 2019; Buckel et al., 2023), a phenomenon and long-term climate impact which also affects the stability of perennially frozen rock walls (Davies et al., 2001; Gruber and Haeberli, 2007; Krautblatter et al., 2013; Mamot et al., 2021; Shugar et al., 2021). In striking contrast to enhanced creep rates in rock glacier permafrost, flow dynamics in lower parts of debris-covered glaciers are decreasing as a consequence of stronger thinning in their upper boundary zone with related reduction of driving stresses (Clayton, 1964; Anderson et al., 2021) and of continued warming-induced ice loss.

In its responsibility for the Global Terrestrial Network for Permafrost (GTN-P) as part of the Global Climate Observing System (GCOS), the International Permafrost Association (IPA) undertakes focused efforts via an Action Group (Rock Glacier Inventories and Kinematics; RGIK/IPA, 2022a, 2022b) to internationally coordinate the development and compilation of rock glacier inventories and the long-term monitoring of rock glacier kinematics (<https://www.unifr.ch/geo/geomorphology/en/research/ipa-action-group-rock-glacier/>). This action group consisting of numerous experts from around the world develops, through consensus, technical guidelines including recommendations on how to discriminate the landform “rock glacier” as expression of viscous creep in perennially frozen subsurface materials (Haeberli et al., 2006; Cicoira et al., 2021; Arenson et al., 2021) from debris-covered glaciers with their surface ice monitored as part of the Global Terrestrial Network for Glaciers (GTN-G). An objective way of differentiating corresponding landforms and kinematics is essential in creating clarity when utilizing such landforms to assess where and how climate change impacts our planet, specifically the cryosphere, or when used in a regulatory/legal context, for example in view of hydrological significance; or generally, to avoid confusion and duplication. The present note attempts at testing the proposed technical

Deleted: 3

65 recommendations/guidelines and at commenting on them using two practical examples: i) the Gruben site in the Swiss Alps
(Figure 1) and ii) the Yerba Loca Valley in the Chilean Andes (Figure 5). At the Gruben site, detailed and comprehensive field
investigations of permafrost, glaciers and complex contacts between the two have been carried out for more than half a century
in connection with hazardous lakes and flood-protection work (see for details Haeberli et al., 2001; Gärtner-Roer et al., 2022;
Wee et al., 2022). While less information exists for the Yerba Loca Valley, it is a valuable site for which some data is available,
and it helps demonstrating that similar conditions exist in different mountain ranges.

70

2. Terminology, characteristics and guidelines:

The RGIK documents provide rich, detailed and comprehensive explanations based on consensus of experts from around the world. As a short summary of this important source, the following terms and guidelines are adopted here and completed with brief notes on principles of surface and subsurface ice with their geophysical characteristics.

75 *Terms:*

Landforms: Rock glaciers are debris landforms generated by the former or current creep of frozen ground (permafrost), detectable in the landscape with the following morphologies: front, lateral margins and optionally ridge-and-furrow surface topography. In coherence with global glacier inventory standards, the minimum rock glacier size to be included into a global compilation should be 0.01 km². Rock glaciers should not be confused with debris-covered glaciers, which are glaciers partially or completely covered by supraglacial debris. The discussion in our invited perspective focusses (a) on patterns of ridges and furrows as expressions of cohesive flow, and (b) on frontal characteristics rather than lateral margins.

80

Surface and subsurface ice:

The term "glacier" is explicitly defined as "on the land surface" (Cogley et al., 2011), i.e., as surface ice. Therefore, debris-covered glaciers as contained in glacier inventories are, by definition, surface ice. Characteristic forms of surface ice, here also in the sense of debris-covered surface ice, are differentiated between glaciers, glacierets, perennial ice patches and dead ice bodies, because the term "glacier" is not appropriate for most of the commonly small landforms in question for reasons of size (area, elevation range) and dynamics. The definition of the term "permafrost" explicitly relates to thermal conditions of "ground (soil or rock and included ice or organic material)" (International Permafrost Association [IPA], 2023), i.e., of subsurface materials. Ice contained in permafrost is, therefore, by definition subsurface ice or ground ice, independent of its spatial extent.

85

90 The term permafrost is thereby independent of material characteristics, i.e., it is not a "type of ground" but a definition of its thermal state.

90

Deleted: Terms

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic, English (UK)

95 Confusion sometimes arises from the use of the term “glacier” in the misleading but historically established and today generally accepted term “rock glacier” as applied to a landform created by subsurface ice under thermal conditions of permafrost. Such confusion can be avoided by accompanying the term “rock glacier” with process- and material-related expressions like “viscous creep features in mountain permafrost”, as done in the title of the present contribution.

Formatted: Font: Not Italic

Geophysical characteristics:

100 Geophysical characteristics of perennially frozen subsurface materials and of massive (debris-covered) surface ice show marked differences which can be summarized as follows:

Formatted: Font: Not Italic

105 Thermal conditions of rock-glacier permafrost are measured in boreholes (Noetzli et al., 2021), by using miniature data loggers at shallow depth (PERMOS, 2023), or they can be approximated through applying numerical models based on climate data (e.g., Haq and Baral, 2019; Baral and Haq, 2020; Li et al., 2023) optimally in combination and, if possible, supported by geodetic measurements of flow characteristics to define activity levels (cf. Bertone et al. 2023). Results from extended time series document ongoing subsurface warming trends (Etzelmüller et al., 2020). Frozen conditions mostly reach down to depths of tens of meters to more than 100 meters. Vertical temperature gradients and heat flow values at depth are strongly reduced due to historical and ongoing surface warming. Debris-covered surface ice can be temperate, polythermal or cold. Perennial ice patches from avalanche cones or glacierets are most common in connection with rock-glacier permafrost. Such small/thin and rather static surface ice bodies can be assumed to be predominantly cold, because their ice cannot warm up above 0°C during the warm season but cool down far below 0°C during the cold part of the year.

Deleted: calculated

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

115 In view of geophysical soundings (e.g., Haeberli and Vonder Mühl, 1996; Hausmann et al., 2007; Hauck and Kneisel, 2008; Merz et al., 2015; Pavoni et al. 2021; Halla et al., 2021; de Pasquale et al., 2022), ice-sediment mixtures in rock-glacier permafrost tend to produce

Formatted: Font: Not Italic

Deleted: ; Hilbich et al., 2022

Formatted: Font: Not Italic

- strong scatter causing reduced transparency for electromagnetic waves, while homogenous/massive surface ice – especially if cold – is highly transparent for radio-echo soundings;
- heterogenous patterns of seismic P-wave velocities with characteristic values varying mostly between about 2,500 and 4,500 m/s, while homogenous/massive surface ice exhibits more uniform values close to 3600 m/s;
- heterogenous patterns of electrical D.C. resistivities with characteristic values ranging from about 10 kΩm to near 1 MΩm, Herring et al., 2023), in contrast to massive surface ice with characteristic values of 1 to 10 MΩm for small ice patches and glacierets primarily consisting of superimposed ice, and of >100 MΩm for glacier ice from warm/wet firn metamorphosis with efficient ion evacuation by percolating meltwater.

Formatted: List Paragraph, Bulleted + Level: 1 + Aligned at: 0.63 cm + Indent at: 1.27 cm

Formatted: Left, Bulleted + Level: 1 + Aligned at: 0.63 cm + Indent at: 1.27 cm

Formatted: Line spacing: 1.5 lines

Formatted: Font: Not Italic

Deleted:

Formatted: Font: Not Italic

155 complete thawing of the ice-rich rock glacier permafrost would most likely need centuries if not millennia (cf. early
calculations reported by Haeberli, 1985).

160 In strong contrast, the continuously slowing-down and decaying debris-covered cold tongue of *Gruben glacier*
(Figure 1) with its progressive loss of coherent flow is a concave landform with a chaotic rather than regular surface pattern,
with diffuse and hardly recognizable frontal and lateral margins but with, in places, local ice cliffs, visible crevasses exposing
ice and thermokarst/surface lakes. Detailed long-term measurements of horizontal and vertical surface displacements (Gärtner-
165 Roer et al., 2022) for the past decades document characteristic subsidence rates in the range of tens of centimeters per year due
to ice melting. From earlier measured radio-echo soundings (Haeberli and Fisch, 1984) and borehole temperatures (Haeberli,
1976), the remaining ice underneath its debris cover can be estimated to be about -1 to -2°C cold and its present-day thickness
to decrease from a maximum of about 20 m at the upper end of continuous debris cover to zero at the diffuse margins close to
Lake 1 (Figure 2). Complete vanishing of the presently existing remains of the glacier tongue is most likely a matter of a few
170 decades. Subglacial permafrost may presently penetrate into the thick sediment bed (Haeberli and Fisch, 1984) underneath the
rapidly thinning and vanishing cold glacier remains.

The differences are obvious, and the morphological criteria ([steep front and lateral margins, surface pattern with
ridges and furrows for rock glaciers, etc.](#)) proposed by the IPA action group are adequate: The subsurface ice contained in the
striking viscous creep feature of *Gruben rock glacier* is a permafrost (periglacial) phenomenon and as such part of [global
175 permafrost monitoring](#). The increasingly indistinctive feature of the down-wasting debris-covered *Gruben glacier* tongue is a
(glacial) phenomenon of surface ice and as such part of [global glacier monitoring](#). Comparison between the measured
kinematics of the two features at this site documents strikingly different characteristics and dynamics of the two domains under
conditions of global warming and highlights the importance of separating these two different features when developing
inventories. The situation is clear and the differentiated treatment of the two phenomena within the framework of
180 internationally coordinated global climate observation perfectly appropriate. Comments are, however, needed concerning two
special aspects:

- complex contact zones between debris-covered surface ice and thermally controlled subsurface ice; and
- the sometimes still maintained idea that covering of surface ice by debris alone can produce striking viscous flow
features called “rock glaciers”.

180 [4. Complex zones with contacts between surface and subsurface ice](#)

Where mean annual air temperatures are, or have until recently been, below zero centigrade, various types of surface ice -
especially cold perennial snow fields or ice patches, glacierets and mostly small, cold to polythermal glaciers - can be in contact
with thermally controlled, creeping ice-rich mountain permafrost, contributing debris and in cases remains of buried ice to
deeply frozen materials. This had already been recognized by Wahrhaftig and Cox (1959), Fisch et al. (1978), Haeberli (1985)
185 or Barsch (1996) and in the meantime became the subject of intense and detailed quantitative investigations by various authors

Deleted: 3

(cf., for instance, Kneisel and Käab, 2006; Bosson and Lambiel, 2016; Monnier and Kinnard, 2017; Bolch et al., 2018; Falatkova et al., 2019; Kunz et al., 2021; Vivero et al., 2021; Wee and Delaloye, 2022; Wee et al., 2022). At the Gruben site, a former contact zone exists between the polythermal Gruben glacier during its maximum LIA/Holocene advances and the permafrost of the Gruben rock glacier (Figure 1; cf., Käab et al., 1997; Gärtner-Roer et al., 2022). Contacts and interactions between glaciers and rock glaciers can give rise to a diverse range of landforms, exhibiting a wide spectrum of characteristics. These landforms encompass structured surface morphology, such as glacitectonics or back-creep, which can either align with or oppose the original stress exerted by the advancing glacier. They can also encompass diffuse and often chaotic topography with thermokarst features in degrading or aggrading permafrost. Furthermore, the local surface coarseness can be smoothed due to the deposition of fine-grained sediments (Wee and Delaloye, 2022; Kneisel and Käab, 2007).

Concerning the “transfer” of surface ice to creeping (rock glacier) permafrost, there is no simple or straightforward general solution. The Gruben and Yerba Loca examples, however, provide some indications. As mentioned in the caption of Figure Sup.-1, the isolated bodies with resistivities in the low MΩm range, still existing today on top of near 0°C permafrost in the former marginal zone of the LIA glacier are most probably dead ice from the small northern tributary underneath the Senggchuppa slope but could also be remains of a buried and frozen avalanche cone at the origin of the photogrammetrically defined flowlines. The earlier visible surface ice at Yerba Loca cannot be called “glaciers” for reasons of size but are/have been perennial ice patches, mostly from avalanche cones. In both cases, Gruben as well as Yerba Loca, the buried ice bodies are more or less passively riding on top of thick perennially frozen sediments. The persistence of MΩm-buried ice in lower parts of rock glaciers are rare exceptions.

Such complex and highly variable landforms can mostly not be attributed in a straightforward “either-or” scheme to the terms “rock glacier” or “debris-covered glacier” but constitute what could better be called “complex contact zones of (viscous creep in ice-rich) permafrost with remains of buried surface ice”. Various forms of surface ice must thereby be treated in a differentiated way – not every piece of surface ice is a “glacier”. A number of potentially interconnected effects and processes can take place as related to loading/unloading, reorientation of stress and flow fields, intermittent burial of surface ice in frozen debris with, in cases, thermokarst development due to its warming-induced vanishing, penetration of permafrost into previously unfrozen parts of now exposed glacier beds, etc. Understanding the full spatiotemporal complexity of relations and interactions and heat fluxes between surface and ground ice in such areas by far exceeds possibilities of intuitive/simplistic landform interpretation from visual or remotely-sensed surface inspection alone. Combinations of sophisticated quantitative measurements are needed, especially geophysical soundings, geotechnical drillings with undisturbed core extraction and geodetic measurements, in order to define subsurface physical conditions (thermal, hydrological and mechanical; stress conditions), material properties and characteristics (especially ice content, unfrozen water, cohesion and internal friction) and related physical processes (creep, shearing, cumulative long-term deformation, infiltration/advection, latent heat exchange) with their changes in time.

Neither the term “rock glacier” nor the term “debris-covered glacier” would be appropriate for such complex contact zones with their characteristically diffuse landforms. A practicable approach for internationally coordinated inventory work to

Deleted: 20

Deleted: 20

Formatted: Font: Not Italic

Formatted: Indent: First line: 1.27 cm

Formatted: Font: Not Italic

Formatted: Font: Not Italic

225 deal with such overlapping/combined contact zones still needs development. An important if not decisive aspect may thereby be that remains of surface ice buried in permafrost are – as the Gruben and Yerba examples document – mostly smaller than the lower size limit applied to the term “glacier” in glacier inventories and much thinner than related permafrost depths (cf. ~~the Kintole conditions documented in Haerberli, 1985).~~

Deleted: also

5. Debris-covered glaciers remain debris-covered glaciers

Deleted: 4

230 Few authors continue postulating that the burial of massive surface ice alone, i.e., in the absence ~~or independently~~ of long-term freezing at depth ~~with its fundamental impact on subsurface material properties~~, can produce rock glaciers, or even that rock glaciers are nothing else than debris-covered LIA glaciers (cf. ~~the literature overview provided by Janke and Bolch, 2022, and the community comments related to our contribution by Harrison and Whalley/Azizi, with response from our side~~). ~~A~~ never discussed implication of these beliefs ~~from the side of intuitive landform interpretation~~ is the rather astonishing, tacit assumption that debris-covered glaciers have two options of possible developments:

Deleted: (Anderson et al., 2018)

Deleted: (Whalley, 2020, 2023)

Deleted: Such concepts primarily stem from the side of intuitive landform interpretation as it had been more common before the development of readily available measurement techniques that allowed data- and physics-based approaches for quantitative research on mountain glaciers and permafrost (cf. the literature overview provided by Janke and Bolch, 2022). They were never supported by results from adequate, quantitative field measurements about subsurface thermal or ice conditions with their fundamental effects on geophysical/geotechnical properties, and essentially remained entirely speculative beliefs.

235 (a) they can remain *debris-covered glaciers* with predominantly chaotic surface structure, slowing down their flow, and decaying/disintegrating under conditions of global warming as increasingly concave landforms with diffuse margins; or

Deleted:

(b) they can become *rock glaciers* by maintaining or even accelerating their flow and advance under conditions of atmospheric temperature rise, adopting a coherent flow pattern of their surface debris and becoming convex landforms with strikingly ~~“organized”~~ surface structures and ~~over-steepened talus-type margins/fronts~~ continuously exposing fresh debris (instead of massive ice) from their inner parts.

Deleted: lavastream-like

240 Option (a) is documented in a rich, quantitative scientific literature (e.g., Iwata et al., 1980; Anderson and Anderson, 2018; Mölg et al., 2020; Shokory and Lane, 2023) and needs no further discussion. Version (b), on the other hand, ~~is in full~~ contradiction with the wealth of quantitative information available from decade-long sophisticated measurements all over the world. ~~It is nevertheless useful to understand the reason why version (b) does not seem to occur in nature, and what the physical~~ causes are for the striking differences in appearance and dynamic evolution of rock glaciers and debris-covered glaciers. Could perhaps permafrost and cold ice conditions constitute a possibility for debris-covered glaciers to “learn” how to turn into something so strikingly different from what they are initially?

Deleted: has never been safely documented in nature by any adequate subsurface prospection and

Deleted: even

Deleted: (cf., the community comments in The Cryosphere related to Whalley, 2023)

Deleted: N

Deleted: , it is

Deleted: this is the case,

Deleted: ,

250 Debris covers on both, rock glaciers and glaciers, are usually predominantly coarse-grained. Frost-susceptible, fine material tends to be washed out and/or to accumulate at the bottom of the debris cover/permafrost active layer. In addition, the source zones of the debris, which are typically steeper rock faces, favour the generation of coarse material. Excess ice formation inducing cohesion and reduced internal friction can therefore not take place throughout coarse-grained surface layers. As a consequence, material properties of the debris cover themselves can hardly make a decisive difference. ~~Material properties exert strong influences where frost-susceptible fines freeze, build up excess ice, and remain frozen. The key for understanding the striking difference between the involved processes and structures, therefore, relates to the formation of the~~ ice-debris

Deleted: The key for understanding the striking difference between the involved processes and structures relates to the formation of the

combination and the corresponding mechanical and thermal coupling between the surface layer of debris and the moving mass underneath (Figure 3). While freezing processes induce the formation of ice inside the rock material and, hence, induce a tightly interconnected mixture of ice and rock particles, largely pure surface ice in debris-covered glaciers or ice patches forms first, and rock components covering it are added independently in a second step as a fundamentally different process and without any tight connection to the ice.

4. The key role of ice-debris coupling ...

On *rock glaciers*, the rock components in the lower part of the “cover”, which includes the thermally defined active layer, reach down beyond the permafrost table and are firmly frozen within the creeping mass. Unless the permafrost is in an advanced state of degradation, mean annual temperatures just below the permafrost table are substantially colder than zero centigrade, adding to the strength of this mixed rock-ice layer. The coherent movement pattern of the perennially frozen mass at depth with its excess ice in fine material, strong cohesion and reduced internal friction is thereby directly transmitted to the rock components of the active layer. Components of the upper active layer, which may intermittently reach positive temperatures during summertime, are firmly interlocked with the deeper, frozen-in components and, hence, directly coupled with the creep movement of the frozen body underneath. This effect is especially strong in zones of compressing flow, which generally occurs in lower and flatter reaches of rock glaciers, causing buckle folding and the creation of characteristic arcuate transverse ridges and furrows (Frehner et al., 2015). Where movement speeds increase along the flow direction, especially in steep root zones of rock glacier flow, extending flow with tensile stresses weakens the interlocking effect between rock components at the surface, thereby often making surface structures somewhat diffuse or producing longitudinal rather than transverse ridge-and-furrow structures. The decisive fact related to the striking viscous appearance of rock glaciers is the large-scale stress coupling which is transferred from the cohesive creep of the thick perennially frozen body at depth to the partially frozen-in and interlocked surface layer.

Components of the debris cover resting on the surface of *massive ice* have hardly any contact with the rare major rock components inside the ice underneath. They are, therefore, not directly coupled with the movement of the ice underneath but typically remain freely mobile at the ice-debris interface (Figure 3). Relative displacements of individual components at smallest spatial scales are easily possible and tend to follow spatially variable, local melt patterns of the underlying ice (cf., Alean et al., 2020), which are further impacted by water flow patterns that develop at the surface of the ice, and thermokarst (cryokarst) processes leading to the collapse of the overlying ice (and debris) layers (Thompson et al., 2016; Mölg et al., 2020). Rather than coherent flow patterns, the debris cover on glaciers or on other buried massive ice tends to develop chaotic surface structures following small-scale features of ground ice melt (cf. Kneib et al., 2023). The mostly chaotic surface structure of debris-covered glaciers is due to the non-transmission of large-scale stress-coupling from the massive ice body at depth to its loose and unconstrained debris cover. This also applies to cold or polythermal ice conditions such as they exist at the Gruben site (cf. also Miles et al., 2018 for Khumbu glacier), because the thickness of the debris layer on glaciers (typically decimeters)

Deleted: 5

315 is mostly much smaller than that of the thermally controlled active layer (typically meters) in the permafrost of rock glaciers.
With other words: debris-covered glaciers have no choice – they remain debris-covered glaciers.

7... as visible at advancing fronts ...

A key element for understanding and recognizing the differences between creeping frozen debris and glaciers covered by debris is the morphology of over-steepened, advancing fronts (Figure 4).

320 Advancing fronts of creeping perennally frozen debris are presently widespread under conditions of atmospheric temperature rise and related permafrost warming/softening with resulting creep acceleration. Ongoing movements create their characteristic talus-like morphology with freshly exposed debris from inside the creeping mass in their over-steepened upper parts (cf., the detailed analyses by Wahrhaftig and Cox, 1959; Kääh and Reichmuth, 2005; Kummert et al., 2021). Rock glacier fronts show over-steepening in the upper section of the front because of the fines and the suction that is generated within the unsaturated sediments of the active layer. The slope of the lower part of the rock glacier front, as clearly visible on Gruben rock glacier (Figure 4), is typically at the angle of repose of the debris as these sediments accumulate in response to them being released from the upper part of the slope. Direct exposures of frozen materials are only observed in connection with extraordinary erosional processes such as detachment slides (Haeberli and Vonder Mühl, 1996; Arenson and Jakob, 2015) or extreme fluvial erosion (Elconin and LaChapelle, 1997).

330 Advancing debris-covered glaciers have become exceptional under conditions of atmospheric temperature rise and predominating glacier shrinkage. In such increasingly rare cases, massive ice of the flowing glacier is usually visible at near-vertical fronts where debris cannot accumulate (Figure 4). Massive ice is also often visible at terminal margins of shrinking debris-covered glaciers.

The advantages of interpreting over-steepened fronts of creeping frozen talus/debris can be illustrated with an example from the Andes of Chile (Yerba Loca Valley; cf. Marangunic et al., 2022). Numerous typical advancing fronts (yellow arrows in Figure 5) document active creep behavior of perennally frozen talus as confirmed by animations using recent aerial lidar scans (cf., supplement). Near points 4 and 5 in Figure 5, at an elevation of about 4150 m, a mean annual air temperature of -4.4°C was measured in 2014 and -2.5°C at 0.4 m depth (in the active layer). These temperatures are significantly colder than at Gruben. Occurrences of buried massive ice with thermokarst ponds are documented by visual observations and core drilling (Ugalde et al., 2018). The landforms 1, 2 and 3 can clearly be defined to be “rock glaciers” with steep fronts/lateral margins and recognizable ridge-and-furrow surface morphology, while the large debris mass marked with the numbers 4 and 5 neither shows clear characteristics of the landform “rock glacier” nor of the landform “debris-covered glacier”. Here, thermokarst ponds develop in remains of massive surface ice from perennial ice patches/avalanche cones and small glacierets on top of creeping and hydraulically impermeable, perennally frozen materials in deep-reaching permafrost. Three primary conclusions can be drawn from this example:

Deleted: 6

Moved down [3]: Rock glacier fronts show over-steepening in the upper section of the front because of the fines and the suction that is generated within the unsaturated sediments of the active layer. The slope of the lower part of the rock glacier front, as clearly visible on Gruben rock glacier (Figure 4), is typically at the angle of repose of the debris as these sediments accumulate in response to them being released from the upper part of the slope.

Formatted: Indent: First line: 0 cm

Moved (insertion) [3]

Deleted: ¶

Deleted: adequately

Formatted: Font: Not Italic

- Complex or doubtful landforms with combinations of surface ice and frozen ground represent conditions in nature that vary beyond limits of “either-or”-type landform categorization.
- Definition and measurement of material properties, ground thermal regimes and physical processes can provide more differentiated insights than applying simple landform schemes.
- Creep phenomena in mountain permafrost are more widespread than the occurrence of specific rock glacier landforms.

8... and its effects on ice loss as a response to long-term warming trends

Ice melting and ground thawing processes are again fundamentally different in thermally controlled permafrost as compared to debris-covered massive ice (Figure 3; cf. Haeberli and Vonder Mühll, 1996). As a response to surface warming caused by changes in time of atmospheric conditions, by creeping out of shadows towards warmer areas in the sun and/or at lower altitudes, or by a combination of both, active layer thickness on rock glacier permafrost increases and thereby adjusts to (near-) equilibrium conditions (cf. long-term measurement series reported by PERMOS, 2023). Thawing into the *perennially frozen ice-rock mixture* underneath sets free the necessary debris for active-layer thickening. Thaw settlement primarily results from melting of excess-ice within the frozen debris, i.e., the ice occupying space within the frozen subsurface matter that exceeds the naturally available pore space, while melting of ice within its pore space alone would not lead to settlement. This process coupling is among the principal reasons for the remarkably low subsidence rates typically observed during recent decades on rock glacier permafrost (Fey and Krainer, 2018; Vivero et al., 2021; Gärtner-Roer et al., 2022). Latent heat effects (energy required to melt ice), together with the increasing thermal protection of permafrost as active layer thickness increases, make the response to climate change through degradation of ice-rich permafrost extremely slow. Further, a thickening of the coarse active layer has a substantial impact on the heat transfer between the atmosphere and the permafrost. First, the thermal resistance of the active layer increases as the thickness of the typically unsaturated debris layer increases. Air has a much lower thermal conductivity than ice or frozen ground (e.g., Andersland and Ladanyi, 2003; Arenson et al., 2021), which is why the thermal conductivity of the active layer tends to decrease as a result of permafrost degradation, contrasting the cover of a debris-covered glacier that cannot change its thermal resistance over time. Secondly, and potentially more importantly, a thickening of the dry and coarse active layer allows increased air flow and with that additional cooling through air convection (Wicky and Hauck, 2020). The Rayleigh number, which describes the potential and the strength of natural convection in porous media (Kane et al., 2001; Niell and Bejan, 2017), is directly dependent on the thickness of the active layer. As illustrated in Figure 3, natural convection can increase, or start to form over time in thickening active layers of degrading rock glacier permafrost, but remains unchanged for a debris covered glacier.

The debris cover on *glaciers or buried massive ice* tends to be much thinner than the active layer on permafrost (Figure 4, cf. McCarthy et al., 2022), providing much smaller insulation capacity and potential for cooling effects by air convection. Most importantly, however, it can generally not significantly increase in thickness as a consequence of ice melting

Deleted: 7

Deleted: e.g.

Deleted: as

Deleted: ,

Deleted: may

Deleted: in

Deleted:

Deleted: n

Deleted: .

Commented [MOU1]: Lukas: Please expand here a bit on thermal resistance and Rayleigh number. Thanks.

Commented [LAZR1]: See added text.

Deleted: a

Deleted: shown

Deleted: the

Deleted: a

Deleted:

Deleted: over time

405 alone, because no significant mass of debris is provided from the vanishing of the underlying massive ice. Assuming 0.1% average debris content inside the ice as an upper-end value based on Bozhinskiy et al. (1986), Nakawo et al. (1986), Kirkbride and Deline (2013), Miles et al. (2021) or Anderson et al. (2021), the melting of a 100m column of ice would result in a thickening of surface debris by only 10cm. Continued warming therefore unavoidably leads to marked thermal imbalance and irreversible ice melting with enhanced surface subsidence until complete vanishing of the covered massive ice. The melting of ice thereby has a heterogenous spatial pattern as it directly responds to the highly variable thickness of the debris cover with its spatially heterogenous sources from rock falls, debris flows or various local displacements, furthered through ice melting
410 caused by thermal and mechanical erosion through water flow at the ice surface (Iwata et al., 1980; Mölg et al., 2020). This is in strong contrast with the thermally controlled active layer of rock glacier permafrost, which responds to climatic conditions in a much more homogeneous way. As a logical consequence, for which more measured evidence is still needed, bodies of massive ice buried in rock glacier permafrost are expected to disappear more quickly in response to warming surface conditions than surrounding frozen debris which can continue to exist over extended time periods (centuries, millennia). This may be the
415 primary reason why such massive buried ice seems to primarily occur, if at all, in upper reaches of presently existing rock glaciers (Haeberli and Vonder Mühll 1996; cf. Bosson and Lambiel, 2016) and hardly ever appears at advancing fronts.

Deleted: Moelg

The here-described material-process coupling causes the striking difference in the response of rock glaciers and debris-covered glaciers to ongoing atmospheric warming trends. It constitutes another reason why debris-covered glaciers have no choice – they can be in contact with creeping frozen materials but cannot, by themselves, become rock glaciers. They
420 remain debris-covered glaciers even under permafrost conditions with cold or polythermal ice, and can quite easily be discriminated from the remarkable viscous creep features usually called rock glaciers. The technical guidelines developed by RGIK/IPA (2022a, 2022b) rightly state that confusion between the two phenomena must be avoided. This is of particular importance when assessing and projecting how different components of the cryosphere respond to climate change and in the context of their hydrological role within a watershed.

Deleted: and safely

425 9. Summary and recommendation

Deleted: 8

A combination of striking morphological, thermal and dynamic characteristics makes the difference between rock glaciers as viscous creep features in mountain permafrost and debris-covered glaciers (and smaller forms of surface ice) under conditions of ongoing global warming: convex versus concave shape, sharp versus diffuse edges, structured versus chaotic surfaces, continued coherent flow and advance versus slowing-down, disintegration and down-wasting. The test at Gruben and Yerba
430 Loca illustrates the applicability of such criteria in concrete climate-related inventory and monitoring work and confirms limits and complexities of landform interpretation needing further exploration.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font colour: Text 1, English (CAN), Kern at 8 pt

The rich available quantitative knowledge basis from borehole and geophysical data in combination with advanced material-/process-related understanding enables safe and straightforward discrimination between rock glaciers as viscous creep

phenomena in ice-rich mountain permafrost and debris-covered glaciers. The corresponding strategies recommended by experts of the International Permafrost Association are informed by and developed based on the process understanding and rich quantitative knowledge basis from numerous sophisticated field investigations using advanced technologies. They are clear and easy to follow, and may be especially helpful in cases when inventories are being compiled without comprehensive site investigations including geophysical soundings or boreholes. The treatment within internationally coordinated global climate observation of thermally controlled subsurface ice in rock glaciers as part of the Terrestrial Network for Permafrost (GTN-P) and of surface ice in debris-covered glaciers as part of the Terrestrial Network for Glaciers (GTN-G) is fully appropriate and such a differentiation shall be followed into the future.

Moved (insertion) [1]

Deleted: clear and easy to follow. s

Formatted: Font: Not Italic

Moved up [1]: recommended by experts of the International Permafrost Association are clear and easy to follow.

Deleted:

Complex contact zones of surface ice (mostly thin perennial snow and ice patches, glacierets or small glaciers) with creep phenomena in ice-rich permafrost, however, in cases constitute diffuse landforms beyond “either-or”- type landform classification. Investigating the thereby involved relations and interactions is a challenging research field beyond attribution of simplistic “origins” to landforms. Exploring contacts and combinations of surface and subsurface ice with their strikingly different response characteristics concerning atmospheric warming is now indeed a growing field of advanced research. It involves quantitative treatment of the involved material properties and processes. This by far exceeds the possibilities of speculative interpretations based alone on visual surface inspection. A recent example illustrating the potential of multimethod field measurements to be used in such complex cases is the comprehensive investigation at the Chauvet site in the French Alps (Cusicanqui et al. 2023).

Deleted: materials and processes with their

Deleted: simplistic landform

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

The key physical phenomenon relates to the fundamentally different distribution of ice and debris: subsurface frozen ice-rock mixtures in rock glaciers versus massive ice with debris on top in debris-covered surface ice. The related mechanical coupling between the moving body at depth and the surface layer of debris is tight in the case of perennially frozen rock glaciers but virtually inexistent in the case of surface ice with a debris cover. The strikingly different morphologies of advancing fronts of rock glaciers and of debris-covered glaciers result from this difference and are in most cases easily recognizable. The difference in ice-debris distribution also explains the extreme contrast in reactions to global warming: Slow thaw subsidence, increasing creep rates and continued advance in the case of warmed-up and softened frozen debris/talus in rock glaciers versus down-wasting, slowing down and disintegration/collapse/vanishing of debris-covered glaciers.

Deleted: ¶

Formatted: Font: Italic, Font colour: Text 1

Formatted: Indent: First line: 0 cm

Deleted: surface

Deleted: glaciers

Data availability: Results from ongoing field measurements carried out by JW at the Gruben site are planned to soon be published in detail.

Team list and author contribution: WH and LA developed the original concept and prepared a first draft. JW and CH provided information about the most recent field campaign at the Gruben site. All co-authors were actively involved in the discussion and in the preparation of the submitted version.

Deleted: , including NM,

480

Competing interests. At least one of the (co-)authors is a member of the editorial board of The Cryosphere.

Acknowledgements

485

We appreciate and support the ongoing efforts undertaken by the International Permafrost Association (IPA) to compile systematic and worldwide inventories of viscous creep features in mountain permafrost – the so-called “rock glaciers” with their warming-related kinematics. The intention of the present contribution is to publicly and critically reflect and strengthen the data- and physics-based concepts thereby developed and their application within the framework of the Global Terrestrial Network for Permafrost (GTN-P/GCOS) as part of integrated global cryosphere observation. We especially thank the numerous colleagues who continue contributing to this important work by documenting solid facts using advanced technologies and process understanding. [An anonymous reviewer and Adriano Ribolini provided constructive feedback to our contribution, and Tobias Bolch edited it.](#)

References

495

Alean, J., Schwendener, L. and Zemp, M.: Migrating boulders on the surface of Alpine valley glaciers. *Geografiska Annaler Series A, Physical Geography*, 103 (21), 1-16. <https://doi.org/10.1080/04353676.2020.1850064>, 2020.

Amschwand, D., Ivy-Ochs, S., Frehner, M., Steinemann, O., Christl, M. and Vockenhuber, C.: Deciphering the evolution of the Bleis Marscha rock glacier (Val d’Err, eastern Switzerland) with cosmogenic nuclide exposure dating, aerial image correlation, and finite-element modelling. *The Cryosphere*, 15, 2057–2081. <https://doi.org/10.5194/tc-15-2057-2021>, 2021.

[Andersland, O.B. and Ladanyi, B.: Frozen Ground Engineering \(2nd ed.\). Wiley, 2003.](#)

Anderson, L.S. and Anderson, R.S.: Debris thickness patterns on debris-covered glaciers. *Geomorphology*, 311, 1-12. <https://doi.org/10.1016/j.geomorph.2018.03.014>, 2018.

505 Anderson, L.S., Armstrong, W.H., Anderson, R.S., Scherler, D. and Peter, E.: The causes of debris-covered glacier thinning: Evidence for the importance of ice dynamics from Kennicott Glacier, Alaska. *Frontiers in Earth Science*, 9:680995. <https://doi.org/10.3389/feart.2021.680995>, 2021.

[Arenson, L.U. and Jakob, M.: Periglacial Geohazard Risks and Ground Temperature Increases. In: Engineering Geology for](#)

Deleted: ¶

Deleted: doi:10

Deleted: ¶

Deleted: doi.org/10

Deleted: ,

Deleted: . 2003.

Deleted: .

Deleted: doi:10

Formatted: mixed-citation

Deleted: doi.org/10

Deleted: Anderson, R.S., Anderson, L.S., Armstrong, W.H., Rossi, M.W. and Crump, S.E.: Glaciation of alpine valleys: The glacier – debris-covered glacier – rock glacier continuum. *Geomorphology* 311, 127-142. doi.org/10.1016/j.geomorph.2018.03.015, 2018. ¶

Society and Territory - Volume 1, pp. 233-237, https://doi.org/10.1007/978-3-319-09300-0_44, 2015.

525 Arenson, L.U., Hoelzle M., and Springman, S.: Borehole deformation measurements and internal structure of some rock glaciers in Switzerland. *Permafrost and Periglacial Processes*, 13 (2), 117-135, 2002.

Arenson, L.U., Colgan, W. and Marshall, H.P.: Physical, thermal and mechanical properties of snow, ice, and permafrost. In: Haeberli, W., Whiteman, C. (Eds.), *Snow and Ice-Related Hazards, Risks, and Disasters*. Elsevier, 35–71, 2021.

530 Baral, P. and Haq, M.A.: Spatial prediction of permafrost occurrence in Sikkim Himalayas using logistic regression, random forests, support vector machines and neural networks. *Geomorphology*, 371, 107331. <https://doi.org/10.1016/j.geomorph.2020.107331>, 2020.

Barsch, D.: *Rockglaciers: Indicators for the Present and Former Geoecology in High Mountain Environments*. Springer, Berlin, 331pp, 1996.

Berthling, I. Beyond confusion: Rock glaciers as cryo-conditioned landforms. *Geomorphology*, 131, 98–106. 535 <https://doi.org/10.1016/j.geomorph.2011.05.002>, 2011.

Bertone, A., Jones, N., Mair, V., Scotti, R., Strozzi, T. and Brardinoni, F.: A climate-driven, altitudinal transition in rock glacier dynamics detected through integration of geomorphological mapping and InSAR-based kinematics. *The Cryosphere Discussion*. <https://doi.org/10.5194/tc-2023-143>, 2023.

540 Bolch, T., Rohrbach, N., Kutusov, S., Robson, B.A. and Osmonov, A.: Occurrence, evolution and ice content of ice-debris complexes in the Ak-Shiirak, Central Tien Shan, revealed by geophysical and remotely-sensed investigations. *Earth Surface Processes and Landforms*, 44 (1), 129-143. <https://doi.org/10.1002/esp.4487>, 2018.

Bosson, J.B. and Lambiel, C.: Internal structure and current evolution of very small debris-covered glacier systems located in Alpine permafrost environments. *Frontiers in Earth Science*, 4 (39). <https://doi.org/10.3389/feart.2016.00039>, 2016.

545 Bozhinskiy, A.N., Krass, M.S. and Popovnin, V.V.: Role of debris cover in the thermal physics of glaciers. *Journal of Glaciology*, 32 (111), 255-266. <https://doi.org/10.3189/S0022143000015598>, 1986.

Buckel, J., Mudler, J., Gardeweg, R., Hauck, C., Hilbich, C., Frauenfelder, R., Kneisel, C., Buchelt, S., Blöthe, J.H., Hördt, A. and Bückner, M.: Identifying mountain permafrost degradation by repeating historical ERT-measurements. *The Cryosphere*, 17, 2919-2940, <https://doi.org/10.5194/tc-17-2919-2023>, 2023.

Cicoira, A., Marcer, M., Gärtner-Roer, I., Bodin, X., Arenson, L.U. and Vieli, A.: A general theory of rock glacier creep based

Deleted: (

Deleted:)

Deleted: doi.org/10

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Deleted: doi.org/10

Formatted: Font colour: Auto

Deleted: doi.org/10

Deleted: doi: 10

Deleted: doi.org/10

Deleted: Discussions.

Deleted: doi.org/10

Deleted: 022

Deleted: 7

on in situ and remote sensing observations. *Permafrost and Periglacial Processes*, 32 (1), 139-153. <https://doi.org/10.1002/ppp.2090>, 2021.

Clayton, L.: Karst topography on stagnant glaciers. *Journal of Glaciology*, 5, 107-112. <https://doi.org/10.3189/s0022143000028628>, 1964.

565 [Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser, G., Möller, M., Nicholson, L., and Zemp, M.: Glossary of Glacier Mass Balance and Related Terms, IHP-VII Technical Documents in Hydrology No. 86, IACS Contribution No. 2. P. UNESCO-IHP, Paris, 2011.](#)

[Cusicanqui, D., Bodin, X., Duvillard, P.-A., Schoeneich, P., Revil, A., Assier, A., Berthet, J., Peyron, M., Roudnitska, S. and Rabatel, A.: Glacier, permafrost and thermokarst interactions in Alpine terrain: Insights from seven decades of reconstructed dynamics of the Chauvet glacial and periglacial system \(Southern French Alps\). *Earth Surface Processes and Landforms*, 48 \(13\), 2595-2612. <https://doi.org/10.1002/esp.5650>, 2023.](#)

570 [Cusicanqui, D., Rabatel, A., Vincent, C., Bodin, X., Thibert, E. and Francou, B.: Interpretation of volume and flux changes of the Laurichard rock glacier between 1952 and 2019, French Alps. *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006161. <https://doi.org/10.1029/2021JF006161>, 2021.](#)

575 [Davies, M.C.R., Hamza, O. and Harris C.: The effect of rise in mean annual temperature on the stability of rock slopes containing ice-filled discontinuities. *Permafrost and Periglacial Processes*, 12, 69-77, 2021.](#)

[de Pasquale, G., Valois, R., Schaffer, N. and MacDonell, S. : Contrasting geophysical signatures of a relict and an intact Andean rock glacier. *The Cryosphere*, 16 \(5\), 1579-1596. <https://doi.org/10.5194/tc-16-1579-2022>, 2022.](#)

[Halla, C., Blöthe, J.H., Tapia Baldis, C., Trombotto Liaudat, D., Hilbich, C., Hauck, C. and Schrott, L.: Ice content and interannual water storage changes of an active rock glacier in the dry Andes of Argentina. *The Cryosphere*, 15 \(2\), 1187-1213. <https://doi.org/10.5194/tc-15-1187-2021>, 2021.](#)

580 [Haq, M.A. and Baral, P. : Study of permafrost distribution in Sikkim Himalayas using Sentinel-2 satellite images and logistic regression modelling. *Geomorphology*, 333, 123-136. <https://doi.org/10.1016/j.geomorph.2019.02.024>, 2019.](#)

[Hauck, C., & Kneisel, C. \(2008\). *Applied Geophysics in Periglacial Environments*. Cambridge University Press. <https://doi.org/10.1017/cbo9780511535628>.](#)

585 [Elconin, R.F. and LaChapelle, E.R.: Flow and internal structure of a rock glacier. *Journal of Glaciology*, 43 \(144\): 238-244, 1997.](#)

Deleted: doi.org/10

Deleted: doi:10

Formatted: No underline

Formatted: English (CAN)

Deleted: doi.org/10

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: (Default) +Headings (Times New Roman), Not Italic

Formatted: Font: (Default) +Headings (Times New Roman), Not Italic

Formatted: Font: (Default) +Headings (Times New Roman), Not Italic

Formatted: Font: (Default) +Headings (Times New Roman), Not Italic

Formatted: Font: (Default) +Headings (Times New Roman), Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font colour: Blue, English (CAN)

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: (Default) +Headings (Times New Roman)

- Etzelmüller, B., Guglielmin, M., Hauck, C., Hilbich, C., Hoelzle, M., Isaksen, K., Noetzi, J., Oliva, M. and Ramos, M.: Twenty years of European mountain permafrost dynamics – the PACE legacy. *Environmental Research Letters*, 15, 104070. <https://doi.org/10.1088/1748-9326/abae9d>, 2020.
- 595 Falatkova, K., Šobr, M., Neureiter, A., Schöner, W., Janský, B., Häusler, H., Engel, Z. and Beneš, V.: Development of proglacial lakes and evaluation of related outburst susceptibility at the Adygin ice-debris complex, northern Tien Shan. *Earth Surface Dynamics*, 7, 301–320. <https://doi.org/10.5194/esurf-7-301-2019>, 2019.
- Fey, C. and Krainer, K.: Analyses of UAV and GNSS based flow velocity variations of the rock glacier Lazaun (Ötztal Alps, South Tyrol, Italy). *Geomorphology*, 365, 107261. <https://doi.org/10.1016/j.geomorph.2020.107261>, 2018.
- 600 Fisch, W. sen., Fisch, W. jun. and Haeblerli, W.: Electrical D.C. resistivity soundings with long profiles on rock glaciers and moraines in the Alps of Switzerland. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 13 (1/2), 239-260, 1978.
- Fleischer, F., Haas, F., Altmann, M., Rom, J., Knoflach, B. and Becht, M.: Combination of historical and modern data to decipher the geomorphologic evolution of the Innere Ölgruben rock glacier, Kaunertal, Austria, over almost a century (1922-2021). *Permafrost and Periglacial Processes*, 34, 3-21. <https://doi.org/10.1002/ppp.2178>, 2023.
- 605 Frehner, M., Ling, A.H.M. and Gärtner-Roer, I.: Furrow-and-ridge morphology on rockglaciers explained by gravity-driven buckle folding: a case study from the Murtèl rockglacier (Switzerland). *Permafrost and Periglacial Processes*, 26 (1), 57–66. <https://doi.org/10.1002/ppp.1831>, 2015.
- Fuchs, M.C., Böhlert, R., Krbetschek, M., Preusser, F. and Egli, M.: Exploring the potential of luminescence methods for dating Alpine rock glaciers. *Quaternary Geochronology*, 18, 17-33. <https://doi.org/10.1016/j.quageo.2013.07.001>, 2013.
- 610 Gärtner-Roer, I., Brunner, N., Delaloye, R., Haeblerli, W., Käab, A. and Thee, P.: Glacier-permafrost relations in a high-mountain environment: 5 decades of kinematic monitoring at the Gruben site, Swiss Alps. *The Cryosphere*, 16, 2083-2101. <https://doi.org/10.5194/tc-16-2083-2022>, 2022.
- Gruber, S., and Haeblerli, W.: Permafrost in steep bedrock slopes and its temperature-related destabilization following climate change. *Journal of Geophysical Research*, 112, F02S18. <https://doi.org/10.1029/2006JF000547>, 2007.
- 615 Haeblerli, W.: Eistemperaturen in den Alpen. *Zeitschrift für Gletscherkunde und Glazialgeologie*, XI/2, 203-220, 1976.
- Haeblerli, W.: Creep of mountain permafrost: internal structure and flow of Alpine rock glaciers. *Mitteilung VAW/ETHZ*, 77, 142pp, 1985.

Deleted: doi.org/10

Deleted: doi.org/10

Deleted: doi.org/10

Deleted: doi:10

Deleted: doi.org/10

Deleted: doi.org/10

Deleted: doi.org/10

Deleted: doi:10

Formatted: German (Switzerland)

Field Code Changed

- Haerberli, W. and Fisch, W.: Electrical resistivity soundings of glacier beds: a test study on Grubengletscher, Wallis, Swiss Alps. *Journal of Glaciology*, 30 (106), 373-376, 1984.
- Haerberli, W. and Vonder Mühl, D.: On the characteristics and possible origins of ice in rock glacier permafrost. *Zeitschrift für Geomorphologie, N.F.*, 104, 43-57, 1996.
- 630 Haerberli, W., Kääh, A., Vonder Mühl, D. and Teyseire, Ph.: Prevention of outburst floods from periglacial lakes at Grubengletscher, Valais, Swiss Alps. *Journal of Glaciology*, 47 (156), 111-122, 2001.
- Haerberli, W., Brandová, D., Burga, C., Egli, M., Frauenfelder, R., Kääh, A. and Maisch, M.: Methods for absolute and relative age dating of rock-glacier surfaces in alpine permafrost. *ICOP 2003 Permafrost: Proceedings of the Eighth International Conference on Permafrost*, 21-25 July 2003, Zurich, Switzerland, A.A. Balkema Publishers, Vol. 1, 343-348, 2003.
- 635 Haerberli, W., Hallet, B., Arenson, L., Elconin, R., Humlum, O., Kääh, A., Kaufmann, V., Ladanyi, B., Matsuoka, N., Springman, S. and Vonder Mühl, D.: Permafrost creep and rock glacier dynamics. *Permafrost and Periglacial Processes*, 17 (3), 189-214. <https://doi.org/10.1002/ppp>, 2006.
- 640 Halla, C., Blöthe, J.H., Tapia Baldis, C., Trombotto Liaudat, D., Hilbich, C., Hauck, C. and Schrott, L.: Ice content and interannual water storage changes of an active rock glacier in the dry Andes of Argentina. *The Cryosphere*, 15 (2), 1187-1213. <https://doi.org/10.5194/tc-15-1187-2021>.
- Harris, C., Arenson, L.U., Christiansen, H.H., Etzelmüller, B., Frauenfelder, R., Gruber, S., Haerberli, W., Hauck, C., Hoelzle, M., Humlum, O., Isaksen, K., Kääh, A., Kern-Lütschg, M.A., Lehning, M., Matsuoka, N., Murton, J.B., Nötzli, J., Phillips, M., Ross, N., Seppälä, M., Springman, S.M. and Vonder Mühl, D.: Permafrost and climate in Europe: Monitoring and modelling thermal, geomorphological and geotechnical responses. *Earth-Science Reviews*, 92, 117-171, 2009.
- 645 Hauck, C., and Kneisel, C.: *Applied Geophysics in Periglacial Environments*. Cambridge University Press. <https://doi.org/10.1017/cbo9780511535628>.
- Hausmann, H., Krainer, K., Brückl, E., and Mostler, W. : Internal structure and ice content of Reichenkar rock glacier (Stubai Alps, Austria) assessed by geophysical investigations. *Permafrost and Periglacial Processes*, 18(4), 351-367, 2007.
- 650 Herring, T., Lewkowicz, A.G., Hauck, C., Hilbich, C., Mollaret, C., Oldenborger, G.A., Uhlemann, S., Farzaman, M., Camels, F. and Scandroglio, R.: Best practices for using electrical resistivity tomography to investigate permafrost. *Permafrost and Periglacial Processes*, 34 (4), 494-512. <https://doi.org/10.1002/ppp.2207>.

Deleted: doi: 10

Formatted: Indent: Left: 0 cm, Hanging: 1 cm, Space After: 12 pt, No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: No underline

Formatted: No underline

Formatted: Font colour: Auto, English (UK)

Deleted: ¶

Moved (insertion) [2]

Deleted: doi.org/10

Field Code Changed

Formatted: Font: Not Italic

Deleted: ¶

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: No underline

Formatted: English (CAN)

Hilbich, C., Hauck, C., Mollaret, C., Wainstein, P.A. and Arenson, L.U.: Towards accurate quantification of ice content in permafrost of the Central Andes – Part 1: Geophysics-based estimates from three different regions. *The Cryosphere*, 16 (5), 1845-1872. <https://doi.org/10.5194/tc-16-1845-2022>, 2022.

660 Hoelzle, M., Hauck, C., Mathys, T., Noetzi, J., Pellet, C. and Scherler, M.: Long-term energy balance measurements at three different mountain permafrost sites in the Swiss Alps. *Earth System Science Data*, 14, 1531-1547. <https://doi.org/10.5194/essd-14-1531-2022>.

International Permafrost Association [IPA] (2023). What is Permafrost? <https://www.permafrost.org/what-is-permafrost>, accessed, August 2023.

665 Jwata, S., Watanabe, O. and Fushimi, H.: Surface morphology in the ablation area of the Khumbu glacier. *Seppyo, Journal of the Japanese Society of Snow and Ice, Special Issue 41*, 9-17. https://doi.org/10.5331/seppyo.41.Special_9, 1980.

Janke, J.R. and Bolch, T. Rock Glaciers, in: Shroder, J.F. (Ed.), *Treatise on Geomorphology (Second Edition)*. Academic Press, Oxford, 75–118. <https://doi.org/10.1016/B978-0-12-818234-5.00187-5>, 2022.

670 Kääh, A. Rock glaciers and protalus forms. In: Elias, S.A. (Ed.), *The Encyclopedia of Quaternary Science 3*. Elsevier, Amsterdam, 535–541, 2013.

Kääh, A. and Reichmuth, T.: Advance mechanisms of rock glaciers. *Permafrost and Periglacial Processes*, 16 (2), 187-193, 2005.

Kääh, A., Haeberli, W. and Gudmundsson, G.H.: Analysing the creep of mountain permafrost using high precision aerial photogrammetry: 25 years of monitoring Gruben rock glacier, Swiss Alps. *Permafrost and Periglacial Processes*, 8 (4), 409 – 426, 1997.

675 Kääh, A., Strozzi, T., Bolch, T., Caduff, R., Trefall, H., Stoffel, M. and Kokarev, A.: Inventory and changes of rock glacier creep speeds in Ile Alatau and Kungöy Ala-Too, northern Tien Shan, since the 1950s. *The Cryosphere*, 15, 927-949. <https://doi.org/10.5194/tc-15-927-2021>, 2021.

Kane, D.L., Hinkel, K.M., Goering, D.J., Hinzman, L.D., and Outcalt, S.I.: Non-conductive heat transfer associated with frozen soils. *Global and Planetary Change*, 29, 275–292. [https://doi.org/10.1016/S0921-8181\(01\)00095-99](https://doi.org/10.1016/S0921-8181(01)00095-99), 2001.

680 Kaufmann, V., Kellerer-Pirklbauer, A. and Seier, G.: Conventional and UAV-Based aerial surveys for long-term monitoring (1954–2020) of a highly active rock glacier in Austria. *Frontiers in Remote Sensing*, 2, 732744. <https://doi.org/10.3389/frsen.2021.732744>, 2021.

Formatted: English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: English (UK)

Moved up [2]: ¶

Deleted: Hilbich, C., Hauck, C., Mollaret, C., Wainstein, P.A., and Arenson, L.U. Towards accurate quantification of ice content in permafrost of the Central Andes – Part 1: Geophysics-based estimates from three different regions. *The Cryosphere*, 16(5), 1845-1872. doi.org/10.5194/tc-16-1845-2022, 2022. ¶
 Hoelzle, M., Hauck, C., Mathys, T., Noetzi, J., Pellet, C. and Scherler, M.: Long-term energy balance measurements at three different mountain permafrost sites in the Swiss Alps. *Earth System Science Data* 14, 1531-1547. doi.org/10.5194/essd-14-1531-2022, 2022. ¶

Deleted: doi.org/10

Deleted: doi:10

Deleted: doi.org/10

Deleted: .

Deleted: .

Deleted: doi: 10

Deleted: r

Deleted: doi: 10

705 Kirkbride, M. and Deline, P.: The formation of supraglacial debris covers by primary dispersal from transverse englacial debris bands. *Earth Surface Processes and Landforms*, 38, 15. <https://doi.org/10.1002/esp.3416>, 2013.

[Kneib, M., Fyffe, C.L., Miles, E.S., Lindemann, S., Shaw, T.E., Buri, P., McCarthy, M., Ouvry, B., Vieli, A., Sato, Y., Kraaijenbrink, P.D.A., Thao, C., Molnar, P. and Pellicciotti, F.: Controls on ice cliff distribution and characteristics on debris-covered glaciers. *Geophysical Research Letters*, 50, e2022GL102444. <https://doi.org/10.1029/2022GL102444>, 2023.](#)

710 Kneisel, C. and Käab, A.: Mountain permafrost dynamics within a recently exposed glacier forefield inferred by a combined geomorphological, geophysical and photogrammetrical approach. *Earth Surface Processes and Landforms*, 32 (12), 1797 – 1810. <https://doi.org/10.1002/esp.1488>, 2007.

Krainer, K., Bressan, D., Dietre, B. Haas, J.N., Hajdas, I., Lang, K., Mair, V., Nickus, U., Reidl, D., Thies, H. and Tonidandel, D.: A 10,300-year-old permafrost core from the active rock glacier Lazaun, southern Ötztal Alps (South Tyrol, northern Italy). *Quaternary Research*, 83 (2), 324–335. <https://doi.org/10.1016/j.yqres.2014.12.005>, 2014.

715 Krautblatter, M., Funk, D. and Günzel, F. K.: Why permafrost rocks become unstable: A rock-ice-mechanical model in time and space. *Earth Surface Processes and Landforms*, 38 (8), 876–887. <https://doi.org/10.1002/esp.3374>, 2013.

Kummert, M., Bodin, X., Braillard, L. and Delaloye, R.: Pluri-decadal evolution of rock glaciers surface velocity and its impacts on sediment export rates towards high-alpine torrents. *Earth Surface Processes and Landforms*, 46, 3217-3227. <https://doi.org/10.1002/esp.5231>, 2021.

720 Kunz, J., Ullmann, T. and Kneisel, C.: Internal structure and recent dynamics of a moraine complex in an alpine glacier forefield revealed by geophysical surveying and Sentinel-1 InSAR time series. *Geomorphology*, 398. <https://doi.org/10.1016/j.geomorph.2021.108052>, 2021.

[Li, M., Yang, Y., Peng, Z., Liu, G.: Assessment of rock glaciers and their water storage in Guokalariju, Tibetan Plateau. *The Cryosphere Discussion*. <https://doi.org/10.5194/tc-2022-178>, 2023.](#)

725 Mamot, P., Weber, S., Eppinger, S. and Krautblatter, M.: A temperature-dependent mechanical model to assess the stability of degrading permafrost rock slopes. *Earth Surface Dynamics*, 9, 1125-1151. <https://doi.org/10.5194/esurf-9-1125-2021>, 2021.

Marangunic, C., Ugalde, F., Apsey, A., Armendáris, I., Bustamante, M. and Peralta, C.: Glaciares en la Cuenca alta del río Mapocho: Variaciones y características principales. *Ecosistemas de Montaña de la Cuenca Alta del Río Mapocho*,

730

Deleted: doi.org/10

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Deleted: doi: 10

Deleted: doi.org/10

Deleted: doi.org/10

Deleted: doi:10

Deleted: doi.org/10

Formatted: Font: No underline, Font colour: Auto

Deleted: doi:10

740 McCarthy, M., Miles, M., Kneib, M., Buri, P., Fugger, S., and Pellicciotti, F.: Supraglacial debris thickness and supply rate in High-Mountain Asia. *Communications Earth & Environment*, 3, 269, 2022.

Deleted: ,

Merz, K., Maurer, H., Buchli, T., Horstmeyer, H., Green, A. G., and Springman, S. M. : Evaluation of ground-based and helicopter ground-penetrating radar data acquired across an Alpine rock glacier. *Permafrost and Periglacial Processes*, 26 (1), 13-27. <https://doi.org/10.1002/ppp.1836>, 2015.

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: mixed-citation

Formatted: Font: Not Italic

745 Miles, K., Hubbard, B., Quincey, D.J., Miles, E.S., Sherpa, T.C., Rowan, A.V. and Doyle, S.H.: Polythermal structure of a Himalayan debris-covered glacier revealed by borehole thermometry. *Scientific Reports*, 8 (16825), 1-9. <https://doi.org/10.1038/s41598-018-34327-5>, 2018.

Deleted: doi.org/10

Field Code Changed

Miles, K., Hubbard, B., Miles, E.S., Quincey, D.J., Rowan, A.V., Kirkbride, M. and Hornsey, J.: Continuous borehole optical televiewing reveals variable englacial debris concentrations at Khumbu Glacier, Nepal. *Communications Earth & Environment*, 2. <https://doi.org/10.1038/s43247-020-00070-x>, 2021.

Deleted: doi.org/10

750 Mölg, N., Ferguson, J., Bolch, T. and Vieli, A.: On the influence of debris cover on glacier morphology: How high-relief structures evolve from smooth surfaces. *Geomorphology*, 357, 107092. <https://doi.org/10.1016/j.geomorph.2020.107092>, 2020.

Deleted: doi/10

755 Mollaret, C., Hilbich, C., Pellet, C., Flores-Orozco, A., Delaloye, R. and Hauck, C.: Mountain permafrost degradation documented through a network of permanent electrical resistivity tomography sites. *The Cryosphere*, 13, 2557–2578. <https://doi.org/10.5194/tc-13-2557-2019>, 2019.

Deleted: doi.org/10

Monnier, S. and Kinnard, C. Pluri-decadal (1955-2014) evolution of glacier-rock glacier transitional landforms in the Central Andes of Chile (30-33°S). *Earth Surface Dynamics*, 5, 493-509. <https://doi.org/10.5194/esurf-5-493-2017>, 2017.

Deleted: doi.org/10

Nakawo, M., Iwata, S., Watanabe, O. and Yoshida, M.: Processes which distribute supraglacial debris on the Khumbu Glacier, Nepal Himalaya. *Annals of Glaciology*, 8, 129-131. <https://doi.org/10.3189/S0260305500001294>, 1986.

Deleted: doi.org/10

760 Nesje, A., Matthews, J.A., Linge, H., Bredal, M., Wilson, P. and Winkler, S.: New evidence for active talus-foot rock glaciers at Øyberget, southern Norway, and their development during the Holocene. *The Holocene*, 31 (11-12), 1786–1796. <https://doi.org/10.1177/09596836211033226>, 2021.

Deleted: doi.org/10

Nield, D.A., and Bejan, A. *Convection in Porous Media*. Cham: Springer. 2017.

Deleted: ¶

775 Noetzli, J., Arenson, L.U., Bast, A., Beutel, J., Delaloye, R., Farinotti, D., Gruber, S., Gubler, H., Haeberli, W., Hasler, A.,
Hauck, C., Hiller, M., Hoelzle, M., Lambiel, C., Pellet, C., Springman, S.M., Vonder Muehll, D. and Phillips, M.: Best
practice for measuring permafrost temperature in boreholes based on the experience in the Swiss Alps. *Frontiers in
Earth Science*, 9, 607875. <https://doi.org/10.3389/feart.2021.60787>, 2021.

Pavoni, M., Sirch, F., and Boaga, J.: *Electrical and electromagnetic geophysical prospecting for the monitoring of rock glaciers
in the Dolomites, Northeast Italy. Sensors*, 21 (4), 1294. <https://doi.org/10.3390/s21041294>, 2021.

780 PERMOS: Swiss Permafrost Bulletin 2022. Noetzli, J. and Pellet, C. (eds.), No. 4, 23 pp. [https://doi.org/10.13093/permos-
bull-2023](https://doi.org/10.13093/permos-
bull-2023), 2023.

Phillips, M., Haberkorn, A., Kenner, R. and Nötzli, J.: Current changes in mountain permafrost based on observations in the
Swiss Alps. *Swiss-Bulletin für angewandte Geologie*, 25 (1-2), 53-63, 2020.

Rock Glacier Inventories and Kinematics [RGIK]: Towards standard guidelines for inventorying rock glaciers: Baseline
concepts (version 4.2.2). IPA Action Group Rock glacier inventories and kinematics, 13 p.
785 https://bigweb.unifr.ch/Science/Geosciences/Geomorphology/Pub/Website/IPA/CurrentVersion/Current_Baseline_Concepts_Inventorizing_Rock_Glaciers.pdf, 2022a.

Rock Glacier Inventories and Kinematics [RGIK]: Towards standard guidelines for inventorying rock glaciers: Practical
concepts (version 2.0). IPA Action Group Rock glacier inventories and kinematics, 10 p.
790 https://bigweb.unifr.ch/Science/Geosciences/Geomorphology/Pub/Website/IPA/CurrentVersion/Current_Practical_Concepts_Inventorizing_Rock_Glaciers.pdf, 2022b

Roer, I., Haeberli, W., Avian, M., Kaufmann, V., Delaloye, R., Lambiel, C. and Kääh, A.: Observations and considerations on
destabilizing active rock glaciers in the European Alps. In: Kane, D.L. and Hinkel, K.M. (eds): Ninth International
Conference on Permafrost, Institute of Northern Engineering, University of Alaska Fairbanks, Vol 2, 1505-1510, 2008.

Shokory, J.A.N. and Lane, S.: Patterns and drivers of glacier debris-cover development in the Afghanistan Hindu Kush
795 Himalaya. *Journal of Glaciology*. <https://doi.org/10.1017/jog.2023.14>, 2023.

Shugar, D.H. and 52 co-authors: A massive rock and ice avalanche caused the 2021 disaster at Chamoli, Indian Himalaya.
Science, 10.1126/science.abh4455, 2021.

Springman, S., Arenson, L., Yamamoto, Y., Maurer, H., Kos, A., Buchli, T. and Derungs, G: Multidisciplinary investigations
on three rock glaciers in the Swiss Alps: Legacies and future perspectives. *Geografiska Annaler, Series A*, 215-243.

Deleted: doi:10

Deleted: ¶

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Not Highlight

Deleted: doi:10

Formatted: Font colour: Text 1

Deleted: doi. Org/10

Deleted:

805 <https://doi.org/10.1111/j.1468-0459.2012.00464.x>, 2012.

Deleted: doi.org/10

Thibert, E. and Bodin, X.: Changes in surface velocities over four decades on the Laurichard rock glacier (French Alps). *Permafrost and Periglacial Processes*, 33 (3), 323-335, <https://doi.org/10.1002/ppp.2159>, 2022.

Deleted: .

Deleted: doi: 10

Thompson, S., Benn, D. I., Mertes, J., and Luckman, A.: Stagnation and mass loss on a Himalayan debris-covered glacier: Processes, patterns and rates. *Journal of Glaciology*, 62, 467-485. <https://doi.org/10.1017/jog.2016.37>, 2016.

Deleted: .

Formatted: German (Switzerland)

Deleted: doi:10

810 Ugalde, F., Marangunic, C. and Becerra, J.: Estructura interna y evolución del glaciar rocoso Littoria en el estero de Yerba Loca, Chile central. In: XV Congreso Geológico Chileno "Geociencias Hacia La Comunidad" Concepción, Chile, 2018.

Vivero, S., Bodin, X., Farias-Barahona, D., MacDonell, S., Schaffer, N., Robson, B.A. and Lambiel, C.: Combination of aerial, satellite, and UAV photogrammetry for quantifying rock glacier kinematics in the Dry Andes of Chile (30°S) since the 1950s. *Frontiers in Remote Sensing*, 2, 784015. <https://doi.org/10.3389/frsen.2021.784015>, 2021.

Deleted: doi: 10

815 Wahrhaftig, C. and Cox, A.: Rock glaciers in the Alaska Range. *Geological Society of America Bulletin* 70, 383-436, 1959.

Wee, J. and Delaloye, R.: Post-glacial dynamics of an alpine Little Ice Age glacetonezoned frozen landform (Aget, western Swiss Alps). *Permafrost and Periglacial Processes*, 33 (4), 370-385. <https://doi.org/10.1002/ppp.2158>, 2022.

Deleted: doi:10


Wee, J., Hauck, C. and Lambiel, C.: Assessing the properties of ground ice and its influence on surface dynamics at Gruben, Swiss Alps. Abstract 11.8, Swiss Geoscience Meeting 2022, 2022.

820 Wicky, J. and Hauck, C.: Air convection in the active layer of rock glaciers. *Frontiers in Earth Science*, 8. <https://doi.org/10.3389/feart.2020.00335>, 2020.


Deleted: Whalley, W.B. Gruben glacier and rock glacier, Wallis, Switzerland: glacier ice exposures and their interpretation. *Geografiska Annaler A* 102, 141-161. doi.org/10.1080/04353676.2020.1765578, 2020. Whalley, W. B.: Comment on "Ice content and interannual water storage changes of an active rock glacier in the dry Andes of Argentina" by Halla et al. (2021). *The Cryosphere* 17, 699-700. doi.org/10.5194/tc-17-699-2023, 2023.

Deleted: doi.org/10

825




845 Figure 1: Continued viscous creep of perennially frozen debris, rich in ground ice at Gruben rock glacier (left), decaying debris-covered, cold tongue of Gruben glacier (right), and former contact zone (center) between the polythermal Little Ice Age (LIA) glacier and the creeping permafrost of the Gruben rock glacier. World Imagery available from Global Mapper V23.1; recent, but exact date unknown.



850
 855
 860

Figure 2: Schematic longitudinal profile through the Gruben rock glacier (top) and the cold debris-covered tongue of Gruben glacier (bottom) based on extensive geophysical prospection (electrical resistivity, seismic refraction, radio-echo-sounding as compiled by Haerberli et al., 2001; Gärtner-Roer et al., 2022, and on recent soundings by Wee et al., 2022). The upper part of the rock glacier presentation reflects the complex contact zone between the polythermal LIA Gruben glacier and the rock glacier permafrost (Kääb et al., 1997); flow direction in this zone is not parallel to the profile but somewhat towards the viewer, leading to the orographic right LIA moraine of Gruben glacier rather than to the advancing rock glacier front. Quantitative information about the debris-covered part of Gruben glacier is from a combination of earlier glacier-bed resistivity and radio echo soundings (Haerberli and Fisch, 1984) with measurements of long-term changes in surface elevation as reported by Gärtner-Roer et al. (2022). Vertical exaggeration is by a factor of 2. Legend: (1) maximum Little Ice Age extent of Gruben glacier; (2) talus, debris; (3) permafrost conditions; (4) buried massive ice of uncertain origin (Gruben rock glacier) and glacier ice (debris-covered part of Gruben glacier); and (5) bedrock after unpublished seismic refraction (Gruben rock glacier) and glacier-bed resistivity soundings (Gruben glacier).

Deleted ;
 Deleted ;



865

Figure 3: Types of ice-debris coupling and adjustment to warming effects in the cases of perennially frozen debris with excess ice (left), and debris-covered massive ice (right). As the ice melts, the thermal insulation in the form of the active layer increases for the permafrost case, but remains unchanged in the case of a debris-covered glacier, where the debris thickness largely remains constant. Modified from Haerberli and Vonder Mühl (1996).

870



Figure 4: Advancing fronts of creeping perennially frozen debris at Gruben rock glacier (top; fall 2009, © Google Earth) and of the debris-covered Belvedere glacier in the Italian Alps during its intermittent advance around the turn of the century (bottom; summer 2005).

|875

Deleted: a

Deleted: glacier at

880




Figure 5: Creep phenomena in the upper Yerba Loca Valley, Andes of Chile. Yellow arrows point to over-steepened advancing fronts of creeping perennially frozen talus/debris, numbers in circles point to landforms as discussed in the text. © DigitalGlobe available in GlobalMapper, modified.

890

Page 5: [1] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [2] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [3] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [4] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [5] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [6] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [7] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [8] Formatted **Lukas Arenson** **28/10/2023 18:15:00**

Centred, Space Before: 3 pt, After: 3 pt, Line spacing: single, Widow/Orphan control, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Page 5: [9] Deleted **Lukas Arenson** **28/10/2023 18:17:00**