

Supplements for : High resolution LGM climate over Europe and the Alpine region using the regional climate model WRF

Emmanuele Russo^{1,2,3}, Jonathan Buzan^{1,2}, Sebastian Lienert^{1,2}, Guillaume Jouvét^{4,5}, Patricio Velasquez Alvarez^{1,2,3}, Basil Davis⁵, Patrick Ludwig⁶, Fortunat Joos^{1,2}, and Christoph C. Raible^{1,2}

¹Climate and Environmental Physics, University of Bern, Sidlerstrasse 5, 3012, Bern, Switzerland.

²Oeschger Centre for Climate Change Research, University of Bern, Hochschulstrasse 4, 3012, Bern, Switzerland.

³now at: Institute for Atmospheric and Climate Science (IAC), ETH Zurich, Universitätsstrasse 16, 8092 Zurich, Switzerland.

⁴Department of Geography University of Zurich, Zurich, Switzerland

⁵Institute of Earth Surface Dynamics, Geopolis, University of Lausanne, Switzerland

⁶Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

Correspondence: emmanuele.russo@env.ethz.ch




Figure S1. Bias in summer precipitation calculated for the coarser resolution domain d01, between the reference simulation and the other members of the ensemble. The dots in each of the figures represent the points for which the values differ significantly, at a significance level of 0.05, according to the results of a KS significance test.