

Opinion: Aerosol Remote Sensing Over The Next Twenty Years

Lorraine A. Remer¹, Robert C. Levy², J. Vanderlei Martins³

¹Goddard Earth Sciences Technology and Research 2 (GESTAR 2), University of Maryland Baltimore County, Baltimore MD, 21228, USA

5 ²NASA Goddard Space Flight Center, Greenbelt MD, 20771, USA

³Department of Physics, University of Maryland Baltimore County, Baltimore MD, 21250, USA

Correspondence to: Lorraine A. Remer (rem@umbc.edu)

10 **Abstract.** More than two decades ago, aerosol remote sensing underwent a revolution with the launch of the Terra and Aqua satellites. Advancement continued via additional launches carrying new passive and active sensors. Capable of retrieving parameters characterizing aerosol loading, rudimentary particle properties and in some cases aerosol layer height, the satellite view of Earth's aerosol system came into focus. The modeling communities have made similar advances. Now the efforts have
15 continued long enough that we can see developing trends in both remote sensing and modeling communities, allowing us to speculate about the future and how the community will approach aerosol remote sensing twenty years from now. We anticipate technology that will replace today's standard multi-wavelength radiometers with hyperspectral and/or polarimetry all viewing in multiple angles. These will be supported by advanced active sensors with the ability to measure profiles of aerosol extinction in addition to backscatter. The result will be greater insight into aerosol particle properties.
20 Algorithms will move from being primarily physically-based to include an increasing degree of Machine Learning methods, but physically-based techniques will not go extinct. However, the practice of applying algorithms to a single sensor will no longer exist. Retrieval algorithms will encompass multiple sensors and all available ground measurements into a unifying framework, and these inverted products will be ingested directly into assimilation systems, becoming "cyborgs": half observations,
25 half model. In twenty years we will see a true democratization in space with nations large and small, private organizations and commercial entities of all sizes launching space sensors. With this increasing amount of data and aerosol products available, there will be a lot of bad data. User communities will organize to set standards and the large national space agencies will lead the effort to maintain quality by deploying and maintaining validation ground networks and focused field experiments. Through it all,
30 interest will remain high in the global aerosol system and how that system affects climate, clouds, precipitation and dynamics, air quality, the environment and public health, transport of pathogens and fertilization of ecosystems, and how these processes are adapting to a changing climate.

1 NASA initiates a revolution with the Terra and Aqua missions

35 As NASA's Terra and Aqua satellites launched at the end of 1999 and in mid-2002, respectively, they ushered in a new era of aerosol remote sensing. These satellites carrying the MODerate resolution

Deleted: Twenty years ago

Deleted: ,

Deleted: followed by

Deleted: information about

Deleted: Today we

Deleted: trends

Deleted: in the aerosol remote sensing and modeling communities that allow...

Deleted: concept

Deleted: s

Imaging Spectroradiometer (MODIS) (Salomonson et al., 1989) and the Multiangle Imaging SpectroRadiometer (MISR, on Terra only)(Diner et al., 1998) offered an unprecedented view of the global aerosol system (Kaufman et al., 2002). The products derived from MODIS and MISR observations defined an aerosol climatology (Remer et al., 2008, Voss and Evan, 2020), identified regional trends (Yu et al., 2020, Zhang and Reid, 2010; [Hammer et al., 2020](#)), advanced understanding of aerosol direct radiative forcing (Christopher and Zhang, 2002, Zhang and Christopher, 2003, Yu et al., 2006, Remer and Kaufman, 2006, Bellouin et al. 2008, [Li et al., 2016](#)), [aerosol indirect radiative forcing \(Peng et al., 2016\)](#)), discovered associations between aerosols, clouds and precipitation that shifted our perception of the water cycle (Kaufman et al., 2005, Koren et al., 2005, 2008, Yuan et al., 2011a,b, [Niu and Li, 2012](#)), proved the importance of aerosol intercontinental transport and nutrient deposition (Gao et al., 2001, Yu et al., 2013, 2019), and gave us important new tools to determine aerosol plume heights (Kahn et al., 2007, 2008, Val Martin et al., 2018) and air quality monitoring and mitigation (Engel-Cox, 2004, Van Donkelaar et al., 2006, 2016, Gupta and Christopher, 2009ab).

[During this period, aerosol remote sensing progressed as a discipline.](#) The MODIS and MISR algorithms had to run in an operational environment, [which meant they had to](#) work for all types of aerosols, [and](#) in all types of environments. The MODIS and MISR algorithms are rooted in the physics of radiative transfer and the mathematics of inversions, but they also incorporate empirical assumptions that tie results to the real world (Kaufman et al., 1997, Tanré et al. 1997, Martonchik et al., 1998, 2002, Kahn et al., 1998). Assumptions are necessary because of the fundamental lack of information content in the MODIS and MISR observations (Tanré et al. 1996, Kahn et al., 1998). MODIS provides a multi-spectral look at each Earth scene at a single viewing geometry. The original MODIS aerosol algorithms, Dark Target (Remer et al., 2005) and Deep Blue (Hsu et al. 2006), make use of no more than six wavelengths for their retrieval, although they use additional wavelengths for cloud masking and other peripheral needs. The MODIS Dark Target over ocean retrieval makes use of the most spectral bands, six. These six bands contain only 2-3 pieces of independent aerosol information (Tanré et al., 1996). Over land, fewer bands are used and the information content available to characterize aerosol is even less (Levy et al., 2007). MISR employs fewer wavelengths, only four from 0.486 μm to 0.867 μm , but adds significant additional information by viewing each scene from 9 different angles. MISR's angular information has shown to be immensely valuable in constraining aerosol type and in determining aerosol plume height (Kahn et al., 2007, 2008, Val Martin et al., 2018, Kahn and Gaitley, 2015).

[However, in addition to the advances leading to the global satellite retrievals in an operational setting, these products also had to be validated and interpreted. The development of MODIS, MISR, and then the active sensors led to significant advances in ground-based remote sensing \(e.g AERONET; \[Holben\]\(#\)](#)

Deleted: ¶

Deleted: At the same time deriving these products from basic satellite observations pushed the field of aerosol remote sensing forward. At the same time, it wasn't just about the basic products.

Deleted: . The algorithms had to

Deleted: and the products required validation

Deleted: huge

Deleted: XXXX

et al., 1998 for validation and algorithm improvements). At the same time, they also led to advances in the modeling and assimilation communities. By providing accurate global products across all types of environments, now models have (a) something to compare with, and (b) something to assimilate.

Deleted: ir

Deleted: and

Deleted:

MODIS and MISR created a revolution in aerosol remote sensing when they were launched in 1999, but the reasons for the revolution involved more than new satellite technology married to state-of-the-science algorithms. A combination of factors came together at just the right time. These factors include:

- A policy for open data access,
- An investment in infrastructure for processing algorithms and distributing data,
- A commitment to long-term science teams for maintenance, improvement and validation of data products including on-orbit sensor calibration,
- The vision and continuous support of the AERONET program for product validation (Holben et al., 1998),
- The overall growth of the international aerosol community and expanded interest in the global aerosol system as part of Earth system science.

We find the first bullet point of particular interest for the celebration of the twentieth anniversary of the journal *Atmospheric Chemistry and Physics*, which itself propelled a revolution in open access and information sharing at roughly the same time.

Thus, the lessons learned from the previous 20 years in aerosol remote sensing is that technical innovation leads to more observable information that leads to more complete aerosol characterization at higher accuracy. But that chain of advancement will only seed new scientific endeavors if the aerosol products are accessible, validated, accompanied by documentation that a user can actually understand and part of a larger data effort to characterize not only the global aerosol system but the Earth system as a whole. The infrastructure created for MODIS and MISR (LAADS DAAC, 2023, ASDC, 2023) continued to absorb and produce data products derived from other sensors and missions over the past two decades: the Aura mission, Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) data from Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO), the Polarization and Anisotropy of Reflectances for Atmospheric Sciences coupled with Observations from a Lidar (PARASOL) mission. Other agencies adopted NASA's open access policies for their missions and products: Visible Infrared Imaging Radiometer Suite (VIIRS), Himawari, Sentinels, Geostationary Ocean Color Instrument (GOCI) etc. Today, a user needing local or global aerosol characterization may feel overwhelmed with the number of aerosol products available to use, both those derived directly from satellite observations and those inferred from models that use satellite information in assimilation systems (Zhang et al., 2008, Benedetti et al., 2009, Gelaro et al., 2017). The perspective from 2023 is

Deleted: 2015

130 that of an apparently data rich era that could hardly have been imagined 23 years ago when Terra
launched.

But what will aerosol remote sensing and data products look like over the next 20 years?

135

140 **2 New technology leads to MORE, MORE, MORE information**

The future of aerosol remote sensing will be propelled by advances in satellite payload technology. Instead of imaging multi-wavelength radiometers, like MODIS, we will expect the additional angular information pioneered by MISR plus the advantages of polarization. Imaging multi-wavelength multi-angle polarimeters were introduced to the community with POLarization and Directionality of the Earth's Reflectances (POLDER)/PARASOL (Deuzé et al., 2000, 2001; [Chen et al., 2020](#)). Future imaging polarimeters will improve upon the POLDER/PARASOL technology with better polarization accuracy, hyper spectral and hyper angle capabilities, decreases in pixel size, increase in the number of wavelengths with polarization and still allow for imaging and enough view angles to optimize aerosol characterization. The Multi-viewing, Multi-channel, Multi-polarisation imaging mission (3MI) is the European follow-on to POLDER/PARASOL that will fly on EUMETSAT's 2nd generation polar-orbiting Meteorological Observing satellite series (MetOp SG) (Fougnie et al. 2018). The first of this series [is planned for launch in 2024](#), [The HyperAngle Rainbow Polarimeter 2 \(HARP2\) and the Spectro-polarimeter for Planetary Exploration One \(SPEXone\) will both fly on NASA's Plankton, Aerosol, Clouds, ocean Ecosystem \(PACE\) mission to be launched in 2024](#) (Werdell et al. 2019), and the Multi Angle Imager for Aerosols (MAIA; [Diner et al., 2018](#)) [is scheduled for launch soon after](#). [China has the Directional Polarized Camera \(DPC\) already in orbit aboard the Daqi-1 \(DQ-1\) satellite, with similar sensors repeated to launch until 2033. Finally, India is planning an aerosol mission, Nanosatellite for Earth Monitoring and Observation-Aerosol Monitoring \(NEMO-AM\), that will carry a multiangle polarimeter the Multi Angle Dual Polarisation Imaging sensor \(MADPI\).](#)

Similar instruments, [with extended wavelength ranges](#), are [planned for NASA's Atmospheric Observing System \(AOS, 2023\) launching later this decade](#). Beyond AOS, we expect more hyperspectral capability, broader wavelength ranges including the thermal infrared, finer spatial resolution, broader swaths and more angles. Twenty years from now these imaging multi-angle and hyperspectral polarimeters will be the norm for passive aerosol characterizing sensors. [Simple single-view multi-wavelength radiometers will still be flying through the 2030s, as they offer products across multiple disciplines: Visible InfraRed Suite \(VIIRS\), Ocean and Land Colour Instrument \(OLCI\), MultiSpectral Instrument \(MSI\), Medium Resolution Spectral Imager \(MERSI\), but the cutting edge](#)

165

Moved down [1]: Table 1 presents a non-comprehensive list of planned space missions from the U.S., Europe and China that have identified an aerosol objective. The list is not comprehensive by any measure, as many missions not specifically designed to characterize aerosol find a means to do so anyway and many additional countries are finding resources to launch space observatories as space becomes more democratized, as we will discuss below

Deleted: 2 Already in the queue!

What satellite missions, already in the queue at national space agencies, will be producing aerosol products? Table 1 presents a non-comprehensive list of planned space missions from the U.S., Europe and China that have identified an aerosol objective. The list is not comprehensive by any measure, as many missions not specifically designed to characterize aerosol find a means to do so anyway and many additional countries are finding resources to launch space observatories as space becomes more democratized, as we will discuss below.

Deleted: Table 1. Partial list of planned missions from major space agencies having an aerosol objective or with reference showing aerosol application without a specific objective! ... [1]
Country

Deleted: China launched the Atmospheric Environment Monitoring Satellite (AEMS) loaded with its first space-borne atmospheric lidar, Aerosol and Carbon Detection Lidar (ACDL) on April 16, 2022 [12, 13] (see Figures S1 and S2). ACDL carries two lidar instruments on a single platform in a 705-km solar synchronous orbit. One is the aerosol-cloud high-spectral-resolution lidar (ACHSRL), and the other (... [2]

Deleted: with the benefits of the added information content only apparent recently (Chen et al., 2020).

Deleted: will be

Deleted: ed

Deleted: the 2023-

Deleted: time frame

Deleted: Other instruments such as the

Deleted:

Deleted: scheduled

Deleted:) will be launched in the early mid 2020s (

Deleted: currently

Deleted: a mission that will be

Deleted: so that Chinese measurements of this type are planned

Deleted: continue

Deleted: with

Deleted: part of the planning for

Deleted: with launch of its first observatory before the end of the current decade with observations expected throughout the 2030s.

Deleted: and simple

Deleted: be antiques

230 aerosol characterization will have moved to the polarimeters by the end of the 2030s. The exception for radiometers will be the geostationary fleet of multispectral radiometers that we discuss below.

Deleted: .

235 In the thermal infrared, single-view instruments will still have a role (DeSouza-Machado, 2010, Klüser et al. 2011). Hyperspectral thermal infrared spectrometers such as the Atmospheric Infrared Sounder (AIRS) have been used to detect and quantify dust AOD and layer height beginning with the Aqua mission (DeSouza-Machado et al., 2006) and advancing to other hyperspectral sounders in this spectral range such as the Infrared Atmospheric Sounding Interferometer (IASI) (Clarisse et al., 2019). The community has recently directed greater focus to dust's contribution to Earth energy balance through its perturbation of the radiation field at infrared wavelengths (Zheng et al., 2022, Song et al., 2022). New technology for hyperspectral thermal infrared spectrometers will improve signal, reduce spatial footprints and are our best hope to achieve mineral speciation of airborne dust from space.

Deleted: even

240 Along with advances in passive remote sensing imagers, we expect new capabilities in active lidars. The straightforward backscatter lidar in space (CALIOP) has demonstrated the unique aerosol information that only a lidar can provide. By 2043 standard aerosol missions will include lidars and these lidars will surpass CALIOP with additional wavelengths, High Spectral Resolution Lidar (HSRL) and/or Raman capability, as well as multiple beams to expand coverage. Ground-based and airborne lidars with these enhanced capabilities are demonstrating that like polarimetry, information content from enhanced lidar design allows for aerosol characterization in ways previously unattainable (Müller et al., 2014). Post-

Deleted: will

Deleted: be

245 CALIOP lidars flying now or in the near future include; the Chinese Aerosol and Carbon Detection Lidar (ACDL; Ke et al., 2022) and the European Atmospheric backscatter LIDar (ATLID; Groß et al., 2015). both have HSRL capability. These are small steps forward, and we understand the expense of launching high capability lidars in space, as the HSRL planned for NASA's AOS mission has recently been descopeed because of cost. Yet, we remain optimistic. By 2043, these modest steps into lidar capability expansion could grow to achieve the ideal for aerosol characterization (Müller et al., 2014). This means that there could be at least one space payload offering 3 wavelengths of backscattering, two wavelengths of extinction profiles using HSRL technology and one wavelength providing depolarization. Aerosol scientists have been advocating for this array of lidar payload capability for more than a decade, and despite the next set of planned aerosol missions falling short, we foresee the dream become reality in 20 years.

Deleted: and

Deleted: Lidars on the books for the future include

Deleted: a 2-wavelength lidar with one HSRL channel for NASA's AOS mission (AOS, 2023), and the

Deleted: a one-wavelength in the ultraviolet lidar with

Deleted: capability for ESA's EARTH CARE mission (Groß et al., 2015)...

Deleted: will

Deleted: $3\beta + 2\alpha + 1\delta$

Deleted: will

Deleted: (β)

Deleted: (α)

Deleted: (δ)

Deleted: will see

265 The lessons we learned from the past twenty years in terms of the advantages of formation flying will guide mission planning far into the future. Like the A-Train of the 2000s and 2010s, missions will be constructed to purposely match passive polarimeters with active lidars in ways that promote synthesized data processing, algorithms, science and applications. The workhorse sun-synchronous polar orbiting mission that has been the mainstay for aerosol characterization will be supplemented with inclined orbits that have been more typical of precipitation-centered missions but have much to offer for aerosols. Inclined orbits allow for a multitude of geometries and time-of-day sampling of the aerosol system over the course of weeks as well as multiple coincidences a day with other satellites looking simultaneously at the same target at the ground. This provides a peek into aerosol situations never encountered by MODIS, MISR or their ilk. Part of this expanding constellation of aerosol observations

290 from satellite will be the geostationary imagers (GEO). Already capability from GEO platforms has evolved from the two or three broad uncalibrated wavelengths that were ubiquitous at the time of Terra launch into multi-wavelength moderate resolution sensors that mimic most of MODIS's capabilities. By 2043, these modern-day Advanced Baseline Imagers (ABI) and Advanced Himawari Imager (AHI), FengYu (FY) and Meteosat Third Generation (MTG) will be upgraded with spectrometers, as per the NASA/NOAA joint Geostationary Extended Observations (GEO-XO) program (Frost et al., 2020). At that time we expect studies in progress to adapt polarimeters to GEO orbit. Other GEO missions include the Geostationary Extended Observations Imager (GXI), Flexible Combined Imager (FCI), Geostationary High speed Imager (GHI) and the Advanced Geostationary Radiation Imager (AGRI), Lidar at that altitude may still be pending. With the equator crowded with observatories in GEO orbit, overlap between sensors provides both a means to cross-calibrate and also multi-angle looks at each scene to increase information content for aerosol retrieval (Bian et al., 2021).

305 Table 1 presents a partial list of planned space missions from the national agencies of U.S., Europe and China that have identified an aerosol objective. The list is not comprehensive by any measure, as many missions not specifically designed to characterize aerosol find a means to do so anyway. For example, the Earth surface Mineral dust source Investigation (EMIT) mission will advance characterization of dust aerosol by identifying mineral content of dust on the ground, before the particles are airborne. EMIT will help with aerosol characterization but is not included here because it is not designed to retrieve airborne aerosol parameters. Table 1 is also missing a comprehensive list of missions because many additional countries are finding resources to launch space observatories as space becomes more democratized, as we will discuss below. Already India, Japan and Korea are planning for future aerosol-capable missions into the next decades.

315 Table 1. Partial list of planned missions from major space agencies in the U.S., China, Europe and one each from India and South Korea having an aerosol objective or aerosol-capable without a specific objective. Compiled from WMO OSCAR (2011-2023).

Country	Mission	Satellite (duration)	Sensor: type
U.S.	Joint Polar Satellite System (JPSS) Sun synchronous	NOAA-21, JPSS-3, -4 (2022-2030), (2027-2036), (2032-2039)	VIIRS: multispectral radiometer
U.S.	Plankton, Aerosol, Clouds, ocean Ecosystems (PACE) Sun synchronous	PACE (2024 – 2028)	OCI: hyperspectral radiometer HARP2 and SPEXone: multiangle imaging polarimeters
U.S.	Multi Angle Imager for Aerosols (MAIA) Sun synchronous	MAIA (2024-2027)	MAIA: multiangle imaging polarimeter
U.S.	Atmosphere Observing System (AOS)	AOS-polar (2029-2039)	Multiangle imaging polarimeter;

Deleted: geosynchronous

Deleted: and a scanning backscatter lidar

Moved (insertion) [1]

Deleted: non-comprehensive

Deleted: further

Deleted: will not

Deleted: and

Deleted: ¶

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

	<u>Sun synchronous</u>		<u>Backscatter lidar</u>
<u>U.S.</u>	<u>Atmosphere Observing System (AOS)</u>	<u>AOS-inclined</u> <u>(2030-2040)</u>	<u>Backscatter lidar</u>
	<u>Inclined</u>		
<u>U.S.</u>	<u>Geostationary Extended Observations (GeoXO)</u>	<u>GeoXO-East, GeoXO-West</u> <u>(2032-2047),</u> <u>2035-2050)</u>	<u>GXI: multiangle radiometer</u>
	<u>Geostationary</u>		
<u>U.S.</u>	<u>Tropospheric Emissions: Monitoring of Pollution (TEMPO)</u>	<u>2023-2026</u>	<u>Hyperspectral in GEO orbit</u>
<u>China</u>	<u>Gao-Fen (GF)-5</u> <u>Sun synchronous</u>	<u>GF-01A, 02</u> <u>(2022-2030),</u> <u>(2021-2028),</u> <u>respectively</u>	<u>DPC: multiangle imaging polarimeter</u>
<u>China</u>	<u>Daqi (DQ)</u> <u>Sun synchronous</u>	<u>DQ-1, DQ-2</u> <u>(2022-2030),</u> <u>(2025-2033)</u>	<u>DPC: multiangle imaging polarimeter; ACDL: high spectral resolution lidar</u>
<u>China</u>	<u>FengYun (FY)-3</u> <u>Sun synchronous</u>	<u>FY-3D, 3F, 3H</u> <u>(2012-2024),</u> <u>(2023-2032),</u> <u>(2024-2029), respectively</u>	<u>MERSI: multispectral radiometer</u>
<u>China</u>	<u>FengYun (FY)-4</u> <u>Geostationary</u>	<u>FY-4B</u> <u>(2021 – 2028)</u>	<u>GHI: multispectral radiometer</u>
<u>China</u>	<u>FengYun (FY)-4</u> <u>Geostationary</u>	<u>FY-4C, 4D, 4E</u> <u>(2025-2031),</u> <u>(2026-2033),</u> <u>(2027-2034), respectively</u>	<u>AGRI: multispectral radiometer</u>
<u>Europe</u>	<u>Earthcare</u> <u>Sun synchronous</u>	<u>Earthcare</u> <u>(2024-2027)</u>	<u>ATLID: backscatter lidar</u> <u>MSI: multispectral radiometer</u>
<u>Europe</u>	<u>Sentinel-2</u> <u>Sun synchronous</u>	<u>Sentinel-2C, 2D</u> <u>(2024-2031),</u> <u>(2025-2032)</u>	<u>MSI: multispectral radiometer</u>
<u>Europe</u>	<u>Sentinel-3</u> <u>Sun synchronous</u>	<u>Sentinel-3C, 3D</u> <u>(2024-2034),</u> <u>(2028-2038)</u>	<u>OLCI: multispectral radiometer</u>
<u>Europe</u>	<u>EPS Second Generation</u> <u>Sun synchronous</u>	<u>MetOp-SG-A1, A2, A3</u> <u>(2025-2032),</u>	<u>3MI: multiangle imaging polarimeter</u>

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

		<u>(2032-2039)</u> <u>(2039-2046)</u>	<u>METimage: multispectral radiometer</u>
<u>Europe</u>	<u>MeteoSat Third Generation (MTG) Geostationary</u>	<u>MTG-11, 12, 13, 14</u> <u>(2022-2030),</u> <u>2026-2034),</u> <u>(2032-2040),</u> <u>(2036-2044)</u>	<u>FCI: multispectral radiometer</u>
<u>India</u>	<u>Nanosatellite for Earth Monitoring and Observation-Aerosol Monitoring (NEMO-AM)</u> <u>Sun synchronous</u>	<u>NEMO-AM (2025-2029)</u>	<u>MADPI:</u> <u>Multiangle imaging polarimeter</u>
<u>South Korea</u>	<u>Geostationary Korea MultiPurpose Satellite (GEO-KOMPSAT)</u>	<u>GEO-KOMPSAT-2A</u> <u>(2018-2029)</u> <u>GEO-KOMPSAT-2B</u> <u>(2020-2031)</u>	<u>Advanced Meteorological Instrument (AMI): multispectral radiometer</u> <u>Geostationary Environmental Monitoring Spectrometer (GEMS): Hyperspectral in GEO orbit</u> <u>Geostationary Ocean Color Imager II (GOCI-II): multispectral radiometer</u>

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt, Not Bold

Formatted: Font: (Default) +Body (Times New Roman), 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt, Not Bold

Formatted: Font: 11 pt

325

The tag line for technology in 2043 will be MORE, MORE, MORE! Each sensor will be collecting orders of magnitude more data than does MODIS today. Polarized measurements require three times the data per wavelength than a simple radiometer to account for the three angles of polarization. Multiply that increase by the number of view angles beyond the single MODIS-like view and we are likely now producing 15X to 60X the amount of data being produced today. The implementation of hyperspectral sensors or improvements in pixel size while maintaining broad swaths to view the entire globe further increases data volume. Advancing lidars with additional wavelengths, HSRL, multiple beams and quicker pulses also magnifies data volume quickly. As part of technology evolution will be the need for spacecraft and especially spacecraft electronics, data systems and transmission to keep up with data volumes. Optical transmission will replace radio frequencies and infrastructure in space will serve as way stations to stage downlinking to overworked ground stations. For example, today the Ka-band is the radio frequency of choice for satellite missions. NASA is promoting “new and improved” Ka-band technology, available for licensing, that achieves a data rate of 130 megabits per second (130 Mbps) (NASA Technology Solution Communications, 2023). In parallel, NASA recently announced achieving 200 gigabytes per second (Gbps) with a space-to-ground optical link (NASA Press Release, 2023). That is three orders of magnitude more data coming down with an optical system. On Earth,

330

Deleted: information

335

Deleted: Laser

340

345 computing power will be fully cloud-based and ever expanding to hold the proliferation of collected
data.

3 Evolution of Algorithms

350 Today, the majority of aerosol retrieval algorithms interpret the signal measured by the satellite sensor
in terms of radiative transfer through the atmosphere. The process is governed by this equation,

$$\rho_{TOA} = \rho_a + \frac{T\rho_s}{(1 - s\rho_s)}$$

355 Where ρ_{TOA} is the solar radiance reflected by the Earth system measured at the top of the atmosphere by
the satellite sensor, ρ_a is the contribution by the atmosphere alone, ρ_s is the reflectance of Earth's
surface that transmits through the atmosphere with transmittance, T , and s is the spherical albedo with
the $(1 - s\rho_s)$ term representing multiple scattering between the atmosphere and Earth's surface. Each
360 term is for a specific wavelength, sun, and sensor geometries, but that notation has been omitted for
simplicity. The terms depend on surface properties (ρ_s and s) and the amount and optical properties of
the gases and aerosols in the atmosphere (ρ_a and T). By making physically-based assumptions on
some surface, gas and aerosol properties and calculating ρ_{TOA} , the calculated ρ_{TOA} can be compared
with the measured ρ_{TOA} and assumptions adjusted until the calculated and measured ρ_{TOA} are the same
to within an acceptable error, consistent with measurement uncertainty.

365 The simplest physically-based algorithms use one wavelength, one geometry at a time. These simple
retrievals hold all assumptions constant except for the optical measure of aerosol loading, the aerosol
optical depth (AOD). Thus, the retrievals use one piece of information to retrieve one parameter, the
AOD. By minimizing the difference between measured and calculated ρ_{TOA} simultaneously for more
than one wavelength, or more than one angle, or more than one polarization state, more information is
370 introduced and additional assumptions can become free parameters leading to retrievals of aerosol
parameters beyond AOD. For example, the MODIS dark target retrieval is based on the knowledge that
independent, multi-spectral observations provide more information. However, not all wavelengths are
completely independent, so that seven MODIS wavelengths from 0.47 μm to 2.1 μm yield, at most,
three pieces of information (Tanré et al., 1996). Likewise, additional geometries and/or polarization
375 states can be introduced, freeing even more parameters for retrieval. The simplest algorithms have
employed radiative transfer models run in the forward direction with assumed aerosol, gases, and

Deleted: .

Deleted: Additional information content introduced by adding wavelengths is dependent on whether the additional wavelengths are independent. ...S

Deleted: S

Deleted: because of the dependencies between wavelengths

Deleted: , if those additional measurements are independent

Deleted: .

surface parameters resulting in a Look Up Table (LUT) stored for future use. The retrieval then can move quickly through the LUT, interpolating between entries, and return the retrieved parameters in an operational setting.

390 Today, as information content increases and computer power grows to meet demand, LUTs are being replaced by other methods to solve the radiative transfer equation. Cost-function minimization methods and other multivariate tools are being used to simultaneously retrieve multiple aerosol, gases, and surface parameters. These more flexible algorithms provide an alternative framework for retrieval, but they are still physically-based in radiative transfer. As we march towards 2043, the LUT methods will be permanently jettisoned, because although radiative transfer becomes more sophisticated with better representation of aerosol optical properties including irregularly shaped particles with complex composition, the dimensionality of the problem would need an infinitely large LUT. Thus, the trend towards optimal estimation, despite the need for an initial guess and the risk of finding local minima. With parallel improvements (and greater sampling) of ground-based and airborne observations including in situ sampling more properties can be constrained. The in situ measurements are especially important to characterize irregularly shaped particles. Algorithms will be designed to ingest data from multiple collocated sensors, especially combinations of passive and active sensors (Liu et al., 2017), and combinations of space-borne and ground-based sensor observations (Lopatin et al., 2013; Li et al., 2019). We expect joint retrievals of aerosol and trace gases, aerosol and land properties, aerosol and ocean color properties, and aerosol/gas/surface retrievals. Eventually there will be sufficient information by 2043 for retrieved properties to include vertical profiles of aerosol extinction and particle number concentration (Schlosser et al. 2022), retrievals over clouds (Torres et al., 2012; Jethva et al., 2013) and over all land surface types including snow and ice (Shi et al., 2019; Mei et al., 2020; Zhang et al., 2023). Characterization will include particle spectral absorption (Xu et al., 2021) and make progress towards constraining particle composition (Kacenenbogen, et al., 2022).

415 By 2043, although radiative transfer and scattering processes will be better understood, completely physically-based algorithms will be rare. Development of Machine Learning (ML) retrieval algorithms will accelerate, such that there will be algorithms that produce a full array of aerosol characteristics by pumping satellite-measured quantities through ML-derived models. These models have no physical basis and do not rely on the radiative transfer equation (Eq. 1). Instead, the models are created by training on a formulation data set that matches satellite-measured radiances with measured aerosol characteristics. For example, today we can train a ML model using satellite measurements collocated with AERONET observations (Lary et al., 2009, Di Noia and Hasekamp, 2018, Kang et al., 2022). In this way we can obtain aerosol characteristics such as AOD that match AERONET with great accuracy

Deleted: produce more parameters and require fewer assumptions

Deleted: than the old LUT algorithms

Deleted: the jettison of

Deleted: will become complete

Deleted: even

Deleted: as

Deleted: Fortunately, with

Deleted: and other satellites, some or

Deleted: Thus, aAlgorithms

Deleted: reach a plateau of usefulness and then become incorporated within physically-based algorithms. In 2043

and precision (Lary et al., 2009), but we lose all physical understanding of how that AOD is retrieved. The challenge for exclusive ML methods like these cited will be the identification of accurate measurements of aerosol characteristics, other than AOD, in sufficient quantity to train the ML models. However, ML algorithms will not recognize a physical situation that falls outside of its training data set, and they still need to accommodate measurement uncertainty.

Deleted:

There are other uses of ML in aerosol remote sensing besides directly obtaining aerosol parameters from satellite observations. Within a physically-based algorithm there are many assumptions that can be aided by ML methods. For example, running a radiative transfer model is computationally expensive. A ML model derived from radiative transfer runs will be more accurate than current interpolations of LUTs and save time in an operational setting (Gao et al., 2021, Ukkonen 2022). In other work, ML methods are used to classify specific aerosol types, such as dust by training on other satellite data sets (CALIPSO) (Lee et al., 2021). ML is used to bias-correct aerosol products derived using standard methods (Lipponen et al. 2022), and it can help with interpolation, constraining non-retrievable aerosol optical properties, cloud (Wei et al., 2020) or snow masking, and surface reflectance determination (Su et al., 2020). In 2043 aerosol retrieval algorithms will continue to be physically-based, but when examined carefully will contain many elements of indispensable ML within.

Deleted: almost all

4 The rise of the cyborg: Blends of measurements and models

The biggest noticeable change over the next twenty years will be the gradual extinction of “satellite aerosol products”. The entire purpose of what we know now as a satellite data product will be to feed major assimilation systems that produce a complete representation of the global aerosol system within the context of the complex Earth system (Zhang et al., 2008, Benedetti et al., 2009, Gelaro et al., 2017). We already see the widespread acceptance of assimilation products, as noted by the number of citations earned by key papers describing each system. For example, using Clarivate’s Web of Science data base (Clarivate, referenced January 2023)) and searching on “satellite AND aerosol” we find the first returned reference to be Holben et al. (1998), the seminal AERONET paper with 5265 citations accumulated over 24 years. Next up is Gelaro et al. (2017), the fundamental reference for the Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2) assimilation products, with 3270 citations accumulated over only nine years. We understand that not all citations to MERRA2 will involve applications to MERRA2 aerosol products and that having a single fundamental reference concentrates users’ citations to a single paper, but the rate of acceptance of MERRA2 and other systems’ assimilation products over the past five years is astounding. See Figure 1.

Deleted: 2015

Deleted: 2015



470 Assimilation systems are analogous to the science fiction concept of the “cyborg”, defined by the
Oxford Dictionary of Science Fiction as “A hybrid being: half human, half machine (a contraction of
‘cybernetic organism’)” (Prucher, 2006). In our context, the assimilation systems are a hybrid of half
observations, half models. Satellites measure electromagnetic radiation that is processed through aerosol
retrieval algorithms to determine aerosol parameters. Assimilation systems intercept parameters from
475 the retrieval algorithms during processing to ingest that data into a global model that adjusts model
representation of aerosol parameters to agree with the observations. Some assimilation systems draw
from the aerosol retrieval algorithms close to the satellite observations, making use of radiance or
reflectance directly (Randles et al., 2017). Some systems ingest the final retrieval products such as AOD
(Zhang et al., 2008). In all cases the global model is drawn closer to actual measurements at regular
480 time steps and is expected to represent the actual aerosol system better than a model that is initialized
and run forward from aerosol source inputs. The advantage of assimilation aerosol products is that users
can access aerosol parameters for any time period, at any location, at any altitude. The parameters will
have the benefit of ALL available data from ALL satellites simultaneously. User’s will not need to
patch together products from different algorithms. The model does that for you and presents the results
in an Earth system governed by physical restraints.

The disadvantages of assimilation are that the resulting aerosol field can be variable, changing abruptly
with model time step or with the available of different aerosol input fields. For example, there will be
much more data available during the day than at night, which may create day/night discontinuities.
490 Furthermore, by drawing the model nearer to the observed variables there can be arbitrary introduction
or subtraction of aerosol mass. This can create biases in mass budget parameters, such as mass
deposition. Furthermore, model parameterizations such as chemical transformation and interaction with
clouds remain uncertain, and likely will remain uncertain in 2043. Nevertheless, assimilation data sets
will become the norm for a majority of aerosol data product consumers. The assimilation systems will
495 run in the cloud and will be called on to produce the cyborg data at user’s will.

We note that satellite-retrieved aerosol products can interact with models without being assimilated.
They can serve as validation and identify and characterize aerosol point sources such as smoke and
volcanic aerosols.

Moved down [2]: The assimilation systems will run in the cloud and will be called on to produce the cyborg data at user’s will.

Moved (insertion) [2]



Deleted:

Deleted: 2015

Deleted: and

Deleted: and

Figure 1. Number of citations received by Gelaro et al. (2017), the fundamental reference for the NASA MERRA2 assimilation system products (green) and the combination of citations received by Levy et al. (2013), Hsu et al. (2013) and Lyapustin et al. (2018), representing the Dark Target, Deep Blue and MAIAC aerosol algorithm products for MODIS (blue). Source: Clarivate Web of Science data base. The point is to note the rapid acceleration of MERRA2 citations as an indicator of the speed of the research community's acceptance of assimilation products, while traditional retrieval products such as MODIS aerosol products maintain a consistent presence in the research community. We anticipate that over the next twenty years, assimilation product use will continue to grow while use of standard products will plateau and decrease.

505 5 Satellites, satellites everywhere

A gold rush to space has begun and will intensify over the next 20 years as space becomes crowded with a variety of sensors in various Earth orbits. During this period we will realize a true democracy in space as non-traditional players such as the space agencies from the developing world and from a wide

range of commercial sector companies join traditional national agencies in blanketing the skies with satellites and sensors. While only a fraction of these satellite sensors will be Earth-viewing and aerosol-capable, the result will yield a tremendous volume of data useful for aerosol retrieval.

Satellites will come in all sizes. Traditional agencies will still launch large, expensive, robust flagship missions with multiple platforms and multiple sensors on each platform. The flagship missions will continue demonstrating lower risk, higher accuracy, greater reliability and longevity. The flagship missions will be supplemented by and integrated with constellations of smaller satellites and nano-sized cubesats. The smaller satellites will be orders of magnitude less expensive, and thus riskier and with more limited capability than the larger observatories. However, the multitude of the smaller units renders the loss of any particular unit as insignificant to the greater objectives. With so many satellites flying, observing Earth and returning measurements for aerosol retrieval, each satellite will be seeking a particular niche. All orbit types will be covered including polar-orbiting, inclined orbits and geosynchronous, and near real time observations will be achieved.

The challenge of so many data sources is the maintenance of data quality. There will be a lot of data, and there will be a lot of bad data. Eventually the scientific and user communities will organize to set standards and protocols to regulate data quality. Such standards will rely on the cohesion of the data communities for enforcement. An example of a community establishing standards that are enforced through peer review are the protocols published by the International Ocean Colour Coordinating Group (IOCCG) for satellite ocean colour sensor validation. Not all data will be free and open access. Commercialization of data products will aid in enforcement of standards, as different entities will be competing to sell their products based on many characteristics including quality. Evaluation of that quality may move into the hands of independent companies or coordinating groups. Commercial entities that launch satellites intend to produce specialized data products for specific customers. Sometimes these customers will be national agencies looking for input to their assimilation systems. Other times commercial entities will be producing their own assimilation system cyborg products and selling those results to their customers.

6 Calibration and Validation systems will need to keep up

Calibration will be a continuing concern with the proliferation of small and nano satellite data flooding the market. As stated above, there will be a lot of bad data, much of it due to poor calibration of each individual sensor, as evidenced by today's fleet of mostly poorly calibrated small satellite sensors. This will change. Calibration takes many forms and includes radiometric, polarimetric, geometric, spectral

Deleted: Validation

550 components and each component requires its own intense scrutiny that not all satellite-launching entities
are ready to provide. To mitigate some of the chaos, a commercial pre- and post-launch calibration
industry will spring up. These companies will offer pre-launch calibration facilities and take
responsibility for a sensor's calibration, even verifying that calibration using vicarious methods once the
sensor is on orbit. Eventually, either governmental or space industry bodies will move to a certification
process similar to the Underwriters Laboratories (UL) or Conformance Europeenne (CE) marks common
today to denote adherence to a specified set of standards for electronic systems. There will also be a
need for in-space calibration options, such as was proposed for the Climate Absolute Radiance and
555 Refractivity Observatory (CLARREO) mission. CLARREO would maintain highly accurate, traceable
reflectance measurements from the Earth and Moon through its lifetime, which could be transferred to
other instruments in orbit.

560 Validation of aerosol products is another issue. With the democratization of space, the leading national
space agencies will no longer hold a monopoly on space access. Ironically, instead these government
agencies that pioneered aerosol characterization from space will take the lead on *suborbital*
measurements to support the validation side of producing high quality aerosol products. Government-
sponsored field experiments and ground networks will provide the basis for the entire international
community, including the commercial sector, to evaluate their products and improve their algorithms

565 Today's basis for validation of aerosol data sets is the ground-based AERONET network (Holben et al.,
1998, Giles et al. 2019). The AERONET concept will continue for decades into the future, but the
technology itself will evolve. Instruments will move from finite wavelength bands to hyperspectral
spectrometers and will continue AERONET's current trend into the shortwave infrared. Originally,
570 AERONET instruments had invested in polarimetry, which turned out to be a capability ahead of its
time. The community has caught up and AERONET will expand the polarimetric capabilities of their
networks and incorporate polarimetry into their retrievals. Data capacity, storage and transmission will
expand, as will the network itself. AERONET's subsidiary, the Marine Aerosol Network (MAN) will
continue to collect data on ocean-going cruises and ships of opportunity. The MAN instruments will
575 also upgrade their technology and data capabilities. ~~The AERONET retrieval that produces reliable~~
aerosol characterization in addition to AOD, including particle size distribution, single scattering
albedo, complex refractive indices, and non-sphericity, will become a more important asset over time as
aerosol satellite remote sensing advances in information content, requiring validation of a wide range of
aerosol parameters.

Formatted: Font: (Default) +Body (Times New Roman), 12 pt

Formatted: Font: (Default) +Body (Times New Roman), 12 pt

Deleted: without losing accuracy

The AERONET concept of a widespread, high quality, aerosol observing ground-network will proliferate, as we are already witnessing with the Interagency Monitoring of Protected Visual Environments (IMPROVE: Malm and Hand, 2007), the European Aerosol Research Lidar Network (EARLINET: Pappalardo et al., 2014), the Aerosol, Cloud and Trace gases Research Initiative (ACTRIS, 2023), the Surface Particulate Matter Network (SPARTAN: Snider et al., 2015) and other networks. The different networks offer different types of aerosol characterization necessary to keep up with the expansion of retrieved aerosol parameters from space. In-situ measurements of aerosol aerodynamic properties will continue to be essential for the assessment of the aerosol health impact and to link aerodynamic to optical measurements, which form the basis of satellite products. Ground LIDAR networks (Welton et al., 2001; Pappalardo et al., 2014) assisted by improved airport ceilometers will expand high temporal characterization of the aerosol vertical profile. As satellite retrieval algorithms upgrade their radiative transfer, refine their assumptions, incorporate ML and become cyborgs, so will the ground network retrieval algorithms. More and more often validation data sets will include the resulting aerosol characterization obtained by simultaneous retrieval of several collocated instruments. Ideally this will include passive measurements with collocated lidar and ground-level or balloon-borne in situ measurements.

Field experiments featuring aircraft-borne measurements will be indispensable for validating the satellite observations at the radiance/polarization level (Level 1) and the retrieved aerosol particle properties in regions and situations of high scientific interest or societal relevance. Aircraft fly in three dimensions, sampling situations between network stations and providing vertical profiles of parameters where lidar is absent. Field experiments will also be essential for confirming ground-based validation sites, as now these sites will be producing a suite of retrieved parameters that require their own validation. We will find that field experiment measurements will be the only truly independent data set for validation. With the abundance of suborbital data, quality again becomes an issue, and national agencies will be the only community players with the resources to maintain high quality, long-term validation data sets.

At the opposite end of the spectrum will be the proliferation of low-cost citizen science networks, including use of smart phone imagery. These commercial networks will expand across populated areas of developed countries and will contribute to characterizing the global aerosol system simply by the abundance of available observations. However, there will not be the same oversight to maintain quality of these data sets and users will need to calibrate the data from citizen scientist measurements with government-maintained and quality-assured sources.

Deleted: also

Deleted: to

Deleted: be everywhere

7 Unanswered science questions will drive the new systems

620 We have described a vibrant global measurement system focused on aerosol remote sensing as we move forward into the next twenty years. This picture assumes a continuing need for aerosol characterization, which will be driven by pressing science questions and societal priorities. What will be the aerosol science questions of the coming decades?

625 Quantifying changes to the aerosol system will be essential as we continue to struggle to catch up to climate change. By 2043 we will have 70 years of Total Ozone Measuring System (TOMS), 60 years of AVHRR and 40 years of MODIS + VIIRS aerosol time series. While most of these time series offer insight into global and regional changes to aerosol loading, inferring changes to aerosol types and particle properties will be especially important.

630 We will still be working towards understanding aerosol-cloud-precipitation processes, how these have changed in time and how do these processes interact with atmospheric dynamics.

635 The role of aerosols in transporting nutrients around the globe and depositing them into terrestrial and oceanic ecosystems will continue to be a pressing issue in the coming decades. Likewise, the role of aerosols in transporting pathogens will still be under investigations in the coming years.

640 Maintaining healthy air quality for all the world's populations will continue to call for understanding of global aerosol distributions and characterization in a way that will require satellite remote sensing.

Underlying these issues will be the goal of using the characterization of the global aerosol system to make our planet healthier and environmentally just.

8 So what will 2043 look like?

645 Despite today's assessment that we are "data rich", by 2043 we will look back at 2023 as a primitive data desert. ~~Aerosol~~ products that are actually cyborgs produced by assimilation systems will represent the global aerosol, producing near real time representation of aerosol fields at fine temporal and spatial scales. ~~Assimilation~~ aerosol product users will forget that clouds (real clouds not data clouds) exist, as all aerosol fields will be interpolated through traditionally cloudy scenes by the ~~models~~. Likewise, ~~assimilation aerosol data fields~~ will ~~have~~ no gaps due to difficult surfaces such as sun glint, snow or ice.

650 However, the challenge will be the quality checking and synthesis of so much data. There will be

Deleted: Satellite-derived

Deleted: a

Deleted: assimilation

Deleted: there

Deleted: be

commercial aerosol data ambassadors who, for a fee, will create custom data packages from the overwhelming supply of public and private data product sources.

660 Today's retrieval algorithms will appear quaint to aerosol remote sensing scientists in 2043. The days of a single inversion applied to a single sensor will be long gone. Even physically-based algorithms will be designed to invert the observations from multiple sensors and platforms simultaneously. This includes joint inversions between satellite and ground-based sensors to produce full characterization of atmospheric constituents (aerosols and gases), as well as complete characterization of the surface beneath whether that surface be a cloud, the ocean or a snow field.

665 Few people will care about aerosol optical depth. The assimilation systems will offer aerosol characterizations in forms most useful for individual users. Air quality managers will automatically obtain particulate mass concentration. Aerosol-cloud-precipitation scientists will obtain vertically-resolved number concentration or heating rates due to aerosol absorption. If anybody is still interested
670 in aerosol radiative forcing, those numbers in Wm^{-2} will come automatically out of some assimilation system. There will be attempts to produce deposition mass flux, but the error bars will still be large. Likewise, attempts to reliably retrieve chemical composition directly from remote sensing likely will fail, although aerosol type will be constrained very well. The only interest in aerosol optical depth will be users working towards continuity with the old sensors for long-term trend analysis.

675 In 2043 validation networks must be in place and major research efforts will switch from algorithm development to designing and implementing a strategy that reserves sufficient independent data for validation. If the assimilation is producing a cyborg vertically-resolved distribution of particle number concentration near or in clouds, then there needs to be a validation strategy to confirm and put error bars
680 on that product. Investment will be needed to develop the instruments and platforms needed to implement a validation strategy. True validation of some difficult parameters such as vertically-resolved particle number validation may be measured only infrequently during dedicated field campaigns. In some ways validation in 2043 may look like a step backward from what aerosol scientists have come to expect today. Instead of 1 million collocations that we achieve now when we validate AOD with
685 AERONET, for some retrieved aerosol parameters, we will be satisfied with 10 to 50 validation points hard-earned in the field.

Where will the funding come from? From the same tax base that has propelled major space agency missions in the past, but now directed more to assimilation systems, validation, quality assurance and

Formatted: Font: (Default) +Body (Times New Roman), 12 pt

Formatted: Font: (Default) +Body (Times New Roman), 12 pt

Formatted: Space Before: Auto, After: 6 pt, Line spacing: single

690 buying data from commercial entities who can sell it cheaper than the agencies can produce themselves. In addition, there will be private concerns, investors and customers, who see value in data products tailored to particular concerns: the insurance industry, the agricultural industry, local air quality jurisdictions.

Formatted: Font: (Default) +Body (Times New Roman), Font color: Blue

695 In summary, 2043 will be an exciting time for aerosol remote sensing. There will be plenty of data for answering science questions and plenty of work to do to assure the integrity of that data. We are sorry to say, though, that we cannot promise flying hover boards as seen in the *Back To The Future* movies (Zemeckis and Gale, 1989, Schildhouse, 2014), even in 2043.

700

References

ACTRIS (Aerosol, Cloud and Trace gases Research Initiative) <https://www.actris.eu>, referenced 29 May 2023

AOS (Atmospheric Observation System) <https://aos.gsfc.nasa.gov/mission.htm>, referenced 05 February 2023.

ASDC (Atmospheric Science Data Center) <https://asdc.larc.nasa.gov/about>, referenced 05 February 2023

710 **Bellouin, N.**, Jones, A., Haywood, J., and Christopher, S. A., Updated estimate of aerosol direct radiative forcing from satellite observations and comparison against the Hadley Centre climate model, *J. Geophys. Res.*, 113, D10205, doi:10.1029/2007JD009385, 2008.

Formatted: Highlight

715 **Benedetti, A.**, Morcrette, J.J., Boucher, O., Dethof, A., Engelen, R.J., Fisher, M., Flentje, H., Huneeus, N., Jones, L., Kaiser, J.W., and Kinne, S.: Aerosol analysis and forecast in the European centre for medium-range weather forecasts integrated forecast system: 2. Data assimilation. *J. Geophys. Res. Atmos.*, 114, D13205, doi:10.1029/2008JD011115, 2009.

Formatted: Highlight

720 **Bian, Q.**, Kreidenweis, S., Chiu, J.C., Miller, S.D., Xu, X., Wang, J., Kahn, R.A., Limbacher, J.A., Remer, L.A. and Levy, R.C.: Constraining Aerosol Phase Function Using Dual-View Geostationary Satellites. *J. Geophys. Res. Atmos.*, 126, doi:10.1029/2021JD035209, 2021.

Formatted: Highlight

Chen, C., Dubovik, O., Fuertes, D., Litvinov, P., Lapyonok, T., Lopatin, A., Ducos, F., Derimian, Y., Herman, M., Tanré, D., Remer, L. A., Lyapustin, A., Sayer, A. M., Levy, R. C., Hsu, N. C., Descloitres, J., Li, L., Torres, B., Karol, Y., Herrera, M., Herreras, M., Aspetsberger, M., Wanzenboeck, M., Bindreiter, L., Marth, D., Hangler, A., and Federspiel, C.: Validation of GRASP algorithm product from

Formatted: Highlight

725 POLDER/PARASOL data and assessment of multi-angular polarimetry potential for aerosol monitoring. *Earth Syst. Sci. Data*, 12, 3573–3620, doi.org/10.5194/essd-12-3573-2020, 2020.

Clarivate. <https://clarivate.com/webofsciencelgroup/solutions/web-of-science/>, 2023

Formatted: Highlight

Deuzé, J.L., Goloub, P., Herman, M., Marchand, A., Perry, G., Susana, S. and Tanré, D.: Estimate of the aerosol properties over the ocean with POLDER. *J. Geophys. Res. Atmos.*, 105, 15329-15346, doi.org/10.1029/2000JD900148, 2000.

Formatted: Highlight

730

Deuzé, J.L., Bréon, F.M., Devaux, C., Goloub, P.H., Herman, M., Lafrance, B., Maignan, F., Marchand, A., Nadal, F., Perry, G. and Tanré, D.: Remote sensing of aerosols over land surfaces from POLDER-ADEOS-1 polarized measurements. *J. Geophys. Res. Atmos.*, 106, 4913-4926, doi.org/10.1029/2000JD900364, 2001.

Formatted: Highlight

735 **Christopher, S.A. and Zhang, J.**: Shortwave aerosol radiative forcing from MODIS and CERES observations over the oceans. *Geophys. Res. Lett.*, 29, 6-1, doi.org/10.1029/2002GL014803, 2002.

Formatted: Highlight

Clarisse, L., Clerbaux, C., Franco, B., Hadji-Lazaro, J., Whitburn, S., Kopp, A. K., et al.: A decadal data set of global atmospheric dust retrieved from IASI satellite measurements. *J. Geophys. Res. Atmos.*, 124, 1618– 1647. doi.org/10.1029/2018JD029701, 2019.

Formatted: Highlight

740 **DeSouza-Machado, S. G., Strow, L. L., Hannon, S. E., and Motteler, H. E.**: Infrared dust spectral signatures from AIRS, *Geophys. Res. Lett.*, 33, L03801, doi:10.1029/2005GL024364, 2006.

Formatted: Highlight

DeSouza-Machado, S.G., Strow, L.L., Imbiriba, B., McCann, K., Hoff, R.M., Hannon, S.E., Martins, J.V., Tanré, D., Deuzé, J.L., Ducos, F. and Torres, O.: Infrared retrievals of dust using AIRS: Comparisons of optical depths and heights derived for a North African dust storm to other collocated EOS A-Train and surface observations. *J. Geophys. Res. Atmos.*, 115, doi.org/10.1029/2009JD012842, 2010.

Formatted: Highlight

Diner, D.J., Boland, S.W., Brauer, M., Bruegge, C., Burke, K.A., Chipman, R., Di Girolamo, L., Garay, M.J., Hasheminassab, S., Hyer, E. and Jerrett, M.: Advances in multiangle satellite remote sensing of speciated airborne particulate matter and association with adverse health effects: from MISR to MAIA. *J. Appl. Rem. Sens.*, 12, 042603-042603, doi.org/10.1117/1.JRS.12.042603, 2018.

Formatted: Highlight

750

Diner, D.J., Beckert, J.C., Reilly, T.H., Bruegge, C.J., Conel, J.E., Kahn, R.A., Martonchik, J.V., Ackerman, T.P., Davies, R., Gerstl, S.A. and Gordon, H.R.: 1998. Multi-angle Imaging SpectroRadiometer (MISR) instrument description and experiment overview. *IEEE Tran. Geosci. Rem. Sens.*, 36, 1072-1087, 10.1109/36.700992, 1998.

Formatted: Highlight

755

Di Noia, A. and Hasekamp, O.P.: Neural networks and support vector machines and their application to aerosol and cloud remote sensing: A review. *Springer Series in Light Scattering: Volume 1: Multiple Light Scattering, Radiative Transfer and Remote Sensing*, 279-329, doi.org/10.1007/978-3-319-70796-9_4, 2018.

Formatted: Highlight

760

765 | **Engel-Cox, J.A.**, Holloman, C.H., Coutant, B.W., and Hoff, R.M.: Qualitative and quantitative evaluation of MODIS satellite sensor data for regional and urban scale air quality. *Atmos. Environ.*, 38, 2495-2509, doi.org/10.1016/j.atmosenv.2004.01.039, 2004.

Formatted: Highlight

| **Fougnie, B.**, Marbach, T., Lacan, A., Lang, R., Schlüssel, P., Poli, G., Munro, R. and Couto, A.B.: The multi-viewing multi-channel multi-polarisation imager—Overview of the 3MI polarimetric mission for aerosol and cloud characterization. *J. Quant. Spec. Rad. Trans.*, 219, 23-32, doi.org/10.1016/j.jqsrt.2018.07.008, 2018.

Formatted: Highlight

770 | **Frost, G.J.** and co-authors: A Value Assessment of an Atmospheric Composition Capability on the NOAA Next-Generation Geostationary and Extended Orbits (GEO-XO) Missions. NOAA technical report OAR CPO, 8. <https://doi.org/10.25923/1s4s-t405>, 2020.

Formatted: Highlight

| **Gao, M.**, Franz, B.A., Knobelspiesse, K., Zhai, P.W., Martins, V., Burton, S., Cairns, B., Ferrare, R., Gales, J., Hasekamp, O. and Hu, Y.: Efficient multi-angle polarimetric inversion of aerosols and ocean color powered by a deep neural network forward model. *Atmos. Meas. Tech.* 14, 4083-4110, 2021.

Formatted: Highlight

775 | **Gao, Y.**, Kaufman, Y. J., Tanré, D., Kolber, D., Falkowski, P. G. Seasonal distributions of aeolian iron fluxes to the global ocean. *Geophys. Res. Lett.*, 28, 29–32. <https://doi.org/10.1029/2000GL011926>, 2001.

Formatted: Highlight

780 | **Gelaro, R.** and Coauthors. The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2). *J. Climate*, 30, 5419–5454, <https://doi-org.proxy-bc.researchport.umd.edu/10.1175/JCLI-D-16-0758.1>, 2017.

Formatted: Highlight

| **Giles, D.M.**, Sinyuk, A., Sorokin, M.G., Schafer, J.S., Smirnov, A., Slutsker, I., Eck, T.F., Holben, B.N., Lewis, J.R., Campbell, J.R. and Welton, E.J., Advancements in the Aerosol Robotic Network (AERONET) Version 3 database—automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements. *Atmos. Meas. Tech.*, 12(1), 169-209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.

Formatted: Highlight

790 | **Groß, S.**, Freudenthaler, V., Wirth, M. and Weinzierl, B., Towards an aerosol classification scheme for future EarthCARE lidar observations and implications for research needs. *Atmospheric Science Letters*, 16(1), 77-82, doi.org/10.1002/asl2.524, 2015.

Formatted: Highlight

| **Gupta, P.**, Christopher, S. A. Particulate matter air quality assessment using integrated surface, satellite, and meteorological products: Multiple regression approach, *J. Geophys. Res.*, 114, D14205, doi:10.1029/2008JD011496, 2009a.

Formatted: Highlight

795 | **Gupta, P.**, Christopher, S. A. Particulate matter air quality assessment using integrated surface, satellite, and meteorological products: 2. A neural network approach, *J. Geophys. Res.*, 114, D20205, doi:10.1029/2008JD011497, 2009b.

Formatted: Highlight

800 | **Hammer, M.S.**, van Donkelaar, A. Li, C., Lyapustin, A., Sayer, A.M., Hsu, N.C., Levy, R.C., Garay, M.J., Kalashnikova, O.V., Kahn, R.A., Brauer, M., Apte, J.S., Henze, D.K., Zhang, L., Zhang, Q., Ford, B., Pierce, J.R., Martin, R.V. Global estimates and long-term trends of

Formatted: Highlight

fine particulate matter concentrations (1998–2018), Environ. Sci. Technol., 54, 7879-7890, 10.1021/acs.est.0c01764, 2020.

805 Jethva, H. T., Torres, O., Remer, L., and Bhartia, P.K., A color ratio method for simultaneous retrieval of aerosol and cloud optical thickness of above-cloud absorbing aerosols from passive sensors: Application to MODIS measurements. *IEEE Transactions on Geoscience and Remote Sensing* 51, 3870 [10.1109/TGRS.2012.2230008, 2013.

810 Holben, B. N., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A., AERONET – A federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, 66, 1–16, doi:10.1016/S0034-4257(98)00031-5, 1998.

815 Hsu, N.C., Tsay, S.C., King, M.D. and Herman, J.R., Deep blue retrievals of Asian aerosol properties during ACE-Asia. *IEEE Transactions on Geoscience and Remote Sensing*, 44(11), 3180-3195, 10.1109/TGRS.2006.879540, 2006.

820 Hsu, N.C., Jeong, M.J., Bettenhausen, C., Sayer, A.M., Hansell, R., Seftor, C.S., Huang, J. and Tsay, S.C., Enhanced Deep Blue aerosol retrieval algorithm: The second generation. *J. Geophys. Res. Atmos.*, 118(16), 9296-9315, doi.org/10.1002/jgrd.50712, 2013.

825 Kacenenbogen, M. S. F., Tan, Q., Burton, S. P., Hasekamp, O. P., Froyd, K. D., Shinozuka, Y., Beyersdorf, A. J., Ziemba, L., Thornhill, K. L., Dibb, J. E., Shingler, T., Sorooshian, A., Espinosa, R. W., Martins, V., Jimenez, J. L., Campuzano-Jost, P., Schwarz, J. P., Johnson, M. S., Redemann, J., and Schuster, G. L.: Identifying chemical aerosol signatures using optical suborbital observations: how much can optical properties tell us about aerosol composition?, *Atmos. Chem. Phys.*, 22, 3713–3742, <https://doi.org/10.5194/acp-22-3713-2022>, 2022.

830 Kahn, R., Banerjee, P., McDonald, D. and Diner, D.J., Sensitivity of multiangle imaging to aerosol optical depth and to pure-particle size distribution and composition over ocean. *J. Geophys. Res. Atmos.*, 103(D24), 32195-32213, doi.org/10.1029/98JD01752, 1998.

835 Kahn, R. A., Li, W.-H., Moroney, C., Diner, D. J., Martonchik, J. V., and Fishbein, E. Aerosol source plume physical characteristics from space-based multiangle imaging, *J. Geophys. Res.*, 112, D11205, doi:10.1029/2006JD007647, 2007.

Kahn, R. A., Chen, Y., Nelson, D. L., Leung, F.-Y., Li, Q., Diner, D. J., and Logan, J. A., Wildfire smoke injection heights: Two perspectives from space, *Geophys. Res. Lett.*, 35, L04809, doi:10.1029/2007GL032165, 2008.

Formatted: Font: Not Bold, Highlight

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Highlight

Formatted: Font color: Custom Color(RGB(46,46,46))

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

| **Kahn, R.A. and Gaitley, B.J.**, An analysis of global aerosol type as retrieved by MISR. *J. Geophys. Res. Atmos.*, 120(9), 4248-4281, doi.org/10.1002/2015JD023322, 2015.

Formatted: Highlight

| **Kang, Y.**, Kim, M., Kang, E., Cho, D. and Im, J., Improved retrievals of aerosol optical depth and fine mode fraction from GOCI geostationary satellite data using machine learning over East Asia. *ISPRS Journal of Photogrammetry and Remote Sensing*, 183, 253-268, 845 https://doi.org/10.1016/j.isprsjprs.2021.11.016, 2022.

Formatted: Highlight

| **Kaufman, Y. J.**, Tanré, D., Remer, L. A., Vermote, E. F., Chu, A., and Holben, B. N., Operational remote sensing of tropospheric aerosol over land from EOS moderate resolution imaging spectroradiometer, *J. Geophys. Res.*, 102 (D14), 17051–17067, doi:10.1029/96JD03988, 1997.

Formatted: Highlight

850 | **Kaufman, Y.**, Tanré, D. and Boucher, O. A satellite view of aerosols in the climate system. *Nature* 419, 215–223 doi.org/10.1038/nature01091, 2002.

Formatted: Highlight

| **Kaufman, Y. J.**, Koren, I., Remer, L. A., Rosenfeld, D., and Rudich, Y.: The effect of smoke, dust, and pollution aerosol on shallow cloud development over the Atlantic Ocean, *P. Natl. Acad. Sci. USA*, 102, 11207–11212, https://doi.org/10.1073/PNAS.0505191102, 2005.

Formatted: Highlight

| **Ke J., Sun Y., Dong C., Zhang X., Wang Z., Lyu L., Zhu W., Ansmann A., Su L., Bu L., et al.** Development of China's first space-borne aerosol-cloud high-spectral-resolution lidar: retrieval algorithm and airborne demonstration *PhotonIX*, 3 (1) (2022), p. 17

Formatted: Highlight

| **Klüser, L.**, Martynenko, D. and Holzer-Popp, T., Thermal infrared remote sensing of mineral dust over land and ocean: a spectral SVD based retrieval approach for IASI. *Atmos. Meas. Tech.*, 4(1), 461-509, 865 doi.org/10.5194/amt-4-757-2011, 2011.

Deleted: ¶

| **Koren, I.**, Kaufman, Y. J., Rosenfeld, D., Remer, L. A., and Rudich, Y. , Aerosol invigoration and restructuring of Atlantic convective clouds, *Geophys. Res. Lett.*, 32, L14828, doi:10.1029/2005GL023187, 2005.

Formatted: Highlight

870 | **Koren, I.**, Martins, J. V., Remer, L. A., and Afargan, H.: Smoke Invigoration Versus Inhibition of Clouds over the Amazon, *Science*, 321, 946–949, https://doi.org/10.1126/science.1159185, 2008.

Formatted: Highlight

875 | LAADS DAAC (Level-1 and Atmosphere Archive & Distribution System Distributed Active Archive Center) https://ladsweb.modaps.eosdis.nasa.gov/about/, referenced 05 February 2023.

| **Lary, D.J.**, Remer, L.A., MacNeill, D., Roscoe, B. and Paradise, S., Machine learning and bias correction of MODIS aerosol optical depth. *IEEE Geoscience and Remote Sensing Letters*, 6(4), 694-698, 10.1109/LGRS.2009.2023605, 2009.

Formatted: Highlight

Lee, J., Shi, Y.R., Cai, C., Ciren, P., Wang, J., Gangopadhyay, A. and Zhang, Z., Machine learning based algorithms for global dust aerosol detection from satellite images: inter-comparisons and evaluation. *Remote Sensing*, 13(3), 456, doi.org/10.3390/rs13030456, 2021.

Formatted: Highlight

885 Levy, R.C., Remer, L.A., Mattoo, S., Vermote, E.F. and Kaufman, Y.J., Second-Generation Operational Algorithm: Retrieval of Aerosol Properties over Land from Inversion of Moderate Resolution Imaging Spectroradiometer Spectral Reflectance. *J. Geophys. Res. Atmos.*, 112, D13211, <https://doi.org/10.1029/2006JD007811>, 2007.

Formatted: Highlight

890 Levy, R.C., Mattoo, S., Munchak, L.A., Remer, L.A., Sayer, A.M., Patadia, F. and Hsu, N.C., The Collection 6 MODIS aerosol products over land and ocean. *Atmos. Meas. Tech.*, 6(11), 2989-3034, doi.org/10.5194/amt-6-2989-2013 2013.

Formatted: Highlight

895 Li, L., Dubovik, O., Derimian, Y., Schuster, G. L., Lapyonok, T., Litvinov, P., Ducos, F., Fuertes, D., Chen, C., Li, Z., Lopatin, A., Torres, B., and Che, H.: Retrieval of aerosol components directly from satellite and ground-based measurements, *Atmos. Chem. Phys.*, 19, 13409-13443, <https://doi.org/10.5194/acp-19-13409-2019>, 2019.

Formatted: Highlight

900 Li, Z., Lau, W. K.-M., Ramanathan, V., Wu, G., Ding, Y., Manoj, M.G., Liu, J., Qian, Y., Li, J., Zhou, T., Fan, J., Rosenfeld, D., Ming, Y., Wang, Y., Huang, ., Wang, B., Xu, X., Lee, S.-S., Cribb, M.C., Zhang, F., Yang, X., Zhao, C., Takemura, T., Wang, K., Xia, X., Yin, Y., Zhang, H., Guo, J., Zhai, P.M., Sugimoto, N., Babu, S.S. and Brasseur, G.P., Aerosol and monsoon interactions in Asia, *Rev. Geophys.*, doi:10.1002/2015 RG000500, 2016.

Formatted: Highlight

905 Lipponen, A., Reinvald, J., Väisänen, A., Taskinen, H., Lähivaara, T., Sogacheva, L., Kolmonen, P., Lehtinen, K., Arola, A. and Kolehmainen, V., Deep-learning-based post-process correction of the aerosol parameters in the high-resolution Sentinel-3 Level-2 Synergy product. *Atmos. Meas. Tech.*, 15(4), 895-914, doi.org/10.5194/amt-15-895-2022, 2022.

Formatted: Highlight

910 Liu, X., Stamnes, S., Burton, S., Ferrare, R., Hostetler, C., Chemyakin, E., Mueller, D., Cairns, B.: A Combined Polarimeter and Lidar Optimal Estimation Algorithm to Improve Aerosol Microphysical Property Retrievals. *NASA Technical Reports Server, Document ID 20200009791*, <https://ntrs.nasa.gov/citations/20200009791>, 2017.

Formatted: Highlight

915 Lopatin, A., Dubovik, O., Chaikovsky, A., Goloub, P., Lapyonok, T., Tarré, D., and Litvinov, P.: Enhancement of aerosol characterization using synergy of lidar and sun-photometer

Formatted: Highlight

coincident observations: the GARRLIC algorithm, *Atmos. Meas. Tech.*, 6, 2065–2088, <https://doi.org/10.5194/amt-6-2065-2013>, 2013.

Deleted: ¶

920 **Lyapustin, A.**, Wang, Y., Korkin, S., and Huang, D.: MODIS Collection 6 MAIAC algorithm, *Atmos. Meas. Tech.*, 11, 5741–5765, <https://doi.org/10.5194/amt-11-5741-2018>, 2018.

Formatted: Highlight

925 **Malm, W.C. and Hand, J.L.**, An examination of the physical and optical properties of aerosols collected in the IMPROVE program. *Atmospheric Environment*, 41(16), 3407-3427, doi.org/10.1016/j.atmosenv.2006.12.012, 2007.

Deleted: ¶

Formatted: Highlight

930 **Martonchik, J.V.**, Diner, D.J., Kahn, R.A., Ackerman, T.P., Verstraete, M.M., Pinty, B. and Gordon, H.R., Techniques for the retrieval of aerosol properties over land and ocean using multiangle imaging. *IEEE Transactions on Geoscience and Remote Sensing*, 36(4), 1212-1227, [10.1109/36.701027](https://doi.org/10.1109/36.701027), 1998.

Formatted: Highlight

935 **Martonchik, J.V.**, Diner, D.J., Crean, K.A. and Bull, M.A., Regional aerosol retrieval results from MISR. *IEEE Transactions on geoscience and remote sensing*, 40(7), 1520-1531, [10.1109/TGRS.2002.801142](https://doi.org/10.1109/TGRS.2002.801142), 2002.

Formatted: Highlight

Mei, L., Vandenbussche, S., Rozanov, V., Proestakis, E., Amiridis, V., Callewaert, S., Vountas, M. and Burrows, J.P.: On the retrieval of aerosol optical depth over cryosphere using passive remote sensing. *Remote Sens. Environ.*, 241, 111731, <https://doi.org/10.1016/j.rse.2020.111731>, 2020.

Formatted: Highlight

940 **Müller, D.**, Hostetler, C.A., Ferrare, R.A., Burton, S.P., Chemyakin, E., Kolgotin, A., Hair, J.W., Cook, A.L., Harper, D.B., Rogers, R.R. and Hare, R.W., Airborne multiwavelength high spectral resolution lidar (HSRL-2) observations during TCAP 2012: vertical profiles of optical and microphysical properties of a smoke/urban haze plume over the northeastern coast of the US. *Atmos. Meas. Tech.*, 7(10), pp.3487-3496, doi.org/10.5194/amt-7-3487-2014, 2014.

Formatted: Highlight

945 **Niu, F.** and Li, Z., Systematic variations of cloud top temperature and precipitation rate with aerosols over the global tropics. *Atmos. Chem. Phys.*, 12, 8491-8498, 2012.

Formatted: Highlight

950 [NASA Press Release, https://www.nasa.gov/feature/ames/tbird-milestone](https://www.nasa.gov/feature/ames/tbird-milestone), 2023

955 [NASA Technology Solution Communications, https://ntts-prod.s3.amazonaws.com/t2p/prod/t2media/tops/pdf/GSC-TOPS-49.pdf](https://ntts-prod.s3.amazonaws.com/t2p/prod/t2media/tops/pdf/GSC-TOPS-49.pdf), 2023

960 **Pappalardo, G.**, Amodeo, A., Apituley, A., Comeron, A., Freudenthaler, V., Linné, H., Ansmann, A., Bösenberg, J., D'Amico, G., Mattis, I. and Mona, L., EARLINET: towards an advanced sustainable European aerosol lidar network. *Atmos. Meas Tech.*, 7(8), pp.2389-2409, doi.org/10.5194/amt-7-2389-2014, 2014.

Formatted: Highlight

965 **Peng, J.**, Li, Z., Zhang, H., Liu, J. and Cribb, M., Systematic changes in cloud radiative forcing with aerosol loading for deep clouds in the tropics, *J. Atmos. Sci.*, 73, 231-249, doi:10.1175/JAS-D-15-0080.1, 2016.

Formatted: Highlight

970 **Prucher, Jeff**, 2006. The Oxford Dictionary of Science Fiction. Oxford University Press. ISBN-13: 9780195305678. On line: DOI:10.1093/acref/9780195305678.001.0001, accessed 29 May 2023.

Formatted: Highlight

Randles, C.A., Da Silva, A.M., Buchard, V., Colarco, P.R., Darmenov, A., Govindaraju, R., Smirnov, A., Holben, B., Ferrare, R., Hair, J. and Shinozuka, Y., The MERRA-2 aerosol reanalysis, 1980 onward. Part I: System description and data assimilation evaluation. *Journal of Climate*, 30(17), 6823-6850, doi.org/10.1175/JCLI-D-16-0609.1, 2017.

Formatted: Highlight

975 **Remer, L. A.**, Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R. R., Ichoku, C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N., The MODIS aerosol algorithm, products and validation, *J. Atmos. Sci.*, 62, 947-973, <https://doi.org/10.1175/JAS3385.1>, 2005.

Formatted: Highlight

980 **Remer, L. A., Kaufman, Y. J.** Aerosol direct radiative effect at the top of the atmosphere over cloud free ocean derived from four years of MODIS data. *Atmos. Chem. Phys.*, 6, 237-253, doi.org/10.5194/acp-6-237-2006, 2006.

Formatted: Highlight

985 **Remer, L. A.**, Kleidman, R.G., Levy, R.C., Kaufman, Y.J., Tanré, D., Mattoo, S., Martins, J.V., Ichoku, C., Koren, I., Yu, H., and Holben, B.N., Global aerosol climatology from the MODIS satellite sensors, *J. Geophys. Res.*, 113, D14S07, doi:10.1029/2007JD009661, 2008.

Formatted: Highlight

990 **Salomonson, V.V.**, Barnes, W.L., Maymon, P.W., Montgomery, H.E. and Ostrow, H., MODIS: Advanced facility instrument for studies of the Earth as a system. *IEEE Transactions on Geoscience and Remote Sensing*, 27(2), 145-153, 10.1109/36.20292, 1989.

Formatted: Highlight

Schildhause, Chloe (November 25, 2014). "Back to the Future II's Art Director Tells Us How They Developed The Film's Somewhat Misguided Predictions". *Uproxx*. Retrieved May 2023.

Formatted: Highlight

995 **Schlosser, J. S.**, Stamnes, S., Burton, S. P., Cairns, B., Crosbie, E., Van Diedenhoven, B., Diskin, G., Dimitrovic, S., Ferrare, R., Hair, J. W., Hostetler, C. A., Hu, Y., Liu, X., Moore, R. H., Shingler, T., Shook, M. A., Thornhill, K. L., Winstead, E., Ziemba, L., and Sorooshian, A., Polarimeter + Lidar – Derived

Formatted: Highlight

Aerosol Particle Number Concentration, *Frontiers in Remote Sensing*, 3, 885332, <https://doi.org/10.3389/frsen.2022.885332>, 2022.

1000 Shi, Z., Xing, T., Guang, J., Xue, Y., Che, Y., Aerosol Optical Depth over the Arctic Snow-Covered Regions Derived from Dual-Viewing Satellite Observations, *Remote Sens.* 11, 891, <https://doi.org/10.3390/rs11080891>, 2019.

Formatted: Highlight

Formatted: Font: Not Italic

Formatted: Font: Not Italic

1005 Snider, G., Weagle, C.L., Martin, R.V., Van Donkelaar, A., Conrad, K., Cunningham, D., Gordon, C., Zwicker, M., Akoshile, C., Artaxo, P. and Anh, N.X., SPARTAN: a global network to evaluate and enhance satellite-based estimates of ground-level particulate matter for global health applications. *Atmos. Meas. Tech.*, 8(1), 505-521, doi.org/10.5194/amt-8-505-2015, 2015.

Formatted: Highlight

1010 Song, Q., Zhang, Z., Yu, H., Kok, J. F., Di Biagio, C., Albani, S., Zheng, J., and Ding, J., Size-resolved dust direct radiative effect efficiency derived from satellite observations, *Atmos. Chem. Phys.*, 22, 13115–13135, <https://doi.org/10.5194/acp-22-13115-2022>, 2022.

Formatted: Highlight

1015 Su, T., Laszlo, I., Li, Z., Wei, J. and Kalluri, S., Refining aerosol optical depth retrievals over land by constructing the relationship of spectral surface reflectances through deep learning: application to Himawari-8, *Remote Sens. Environ.*, 251, 112093, [doi:10.1016/j.rse.2020.112093](https://doi.org/10.1016/j.rse.2020.112093), 2020.

Formatted: Highlight

1020 Tanré, D., M. Herman, Y. J. Kaufman, Information on the aerosol size distribution contained in the solar reflected spectral radiances, *J. Geophys. Res.*, 101, 19043– 19060, <https://doi.org/10.1029/96JD00333>, 1996.

Formatted: Highlight

Tanré, D., Kaufman, Y. J., Herman, M., and Mattoo, S., Remote sensing of aerosol properties over oceans using the MODIS/EOS spectral radiances, *J. Geophys. Res.*, 102 (D14), 16971– 16988, [doi:10.1029/96JD03437](https://doi.org/10.1029/96JD03437), 1997.

Formatted: Highlight

1025 Torres, O., Jethva, H.T. and Bhartia, P.K., Retrieval of Aerosol Optical Depth above Clouds from OMI Observations: Sensitivity Analysis and Case Studies, *J. Atmos. Sci.*, 69 (3), 1037-1053, [10.1175/JAS-D-11-0130.1](https://doi.org/10.1175/JAS-D-11-0130.1), 2012.

Formatted: Font: Not Bold, Highlight

Formatted: Highlight

Formatted: Font: Not Bold, Highlight

Formatted: Font: Not Bold

Formatted: Font: Not Italic

Formatted: Font: Not Bold

Formatted: Font: Bold

1030 **Ukkonen**, P. Exploring pathways to more accurate machine learning emulation of atmospheric radiative transfer. *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002875. doi.org/10.1029/2021MS002875, 2022

Formatted: Highlight

1035 **Val Martin** M, Kahn RA, Tosca MG. A Global Analysis of Wildfire Smoke Injection Heights Derived from Space-Based Multi-Angle Imaging. *Remote Sensing*, 10(10), 1609. doi-org.proxy-bc.researchport.umd.edu/10.3390/rs10101609, 2018.

Formatted: Highlight

1040 **Van Donkelaar**, A., Martin, R. V., Park, R. J. Estimating ground-level PM_{2.5} using aerosol optical depth determined from satellite remote sensing. *J. Geophys. Res.*, 111, D21201, doi:10.1029/2005JD006996, 2006.

Formatted: Highlight

1045 **Van Donkelaar**, A., Martin, R.V., Brauer, M., Hsu, N.C., Kahn, R.A., Levy, R.C., Lyapustin, A., Sayer, A.M., Winker, D.M. Global estimates of fine particulate matter using a combined geophysical-statistical method with information from satellites, models, and monitors. *Environ. Sci. Tech.*, 50, 3762-3772, doi.org/10.1021/acs.est.5b05833, 2016.

Formatted: Highlight

1050 **Voss, K.K. and Evan, A.T.**, A new satellite-based global climatology of dust aerosol optical depth. *J. Appl. Meteorol. Clim.*, 59(1), 83-102, doi.org/10.1175/JAMC-D-19-0194.1, 2020.

Formatted: Highlight

1055 **Wei, J., Huang, W., Li, Z., Sun, L., Zhu, X., Yuan, Q., Liu, L. and Cribb, M.C.**, [Cloud detection for Landsat imagery by combining the random forest and superpixels extracted via energy-driven sampling segmentation approaches](https://doi.org/10.1016/j.rse.2020.112005), *Remote Sens. Environ.*, 248, 112005, doi:10.1016/j.rse.2020.112005, 2020.

Formatted: Highlight

1060 **Welton**, Ellsworth J., Campbell, James R., James D. Spinhirne, and V. Stanley Scott III. [Global monitoring of clouds and aerosols using a network of micropulse lidar systems](https://doi.org/10.1117/12.417040). *Proc. SPIE 4153. Lidar Remote Sensing for Industry and Environment Monitoring*, https://doi.org/10.1117/12.417040, 2001.

Formatted: Highlight

1065 **Werdell, P.J.**, Behrenfeld, M.J., Bontempi, P.S., Boss, E., Cairns, B., Davis, G.T., Franz, B.A., Gliese, U.B., Gorman, E.T., Hasekamp, O., Knobelspiesse, K.D., Mannino, A., Martins, J.V., McClain, C.R., Meister, G., and Remer, L.A., The Plankton, Aerosol, Cloud, ocean Ecosystem mission: status, science, advances. *Bulletin of the American Meteorological Society*, 100(9), 1775-1794, doi.org/10.1175/BAMS-D-18-0056.1, 2019.

Formatted: Highlight

1070 **WMO OSCAR**, [World Meteorological Organization Observing Systems Capability Analysis and Review Tool](https://space.oscar.wmo.int) https://space.oscar.wmo.int, 2011-2023.

Formatted: Font color: Text 1, Highlight

Formatted: Font color: Text 1

Formatted: Hyperlink, Font: (Default) Times New Roman, Font color: Auto, Pattern: Clear

Field Code Changed

1070 Xu, F., Gao, L., Redemann, J., Flynn, C.J., Espinosa, W.R., da Silva, A.M., Stamnes, S., Burton, S.P., Liu, X., Ferrare, R., Cairns, B., Dubovik, O.: A combined Lidar-Polarimeter inversion approach for aerosol remote sensing over ocean, *Frontiers in Remote Sensing*, 2, <https://doi.org/10.3389/frsen.2021.620871>, 2021.

Formatted: Highlight

1075 Yu, H., Kaufman, Y. J., Chin, M., Feingold, G., Remer, L. A., Anderson, T. L., Balkanski, Y., Bellouin, N., Boucher, O., Christopher, S., DeCola, P., Kahn, R., Koch, D., Loeb, N., Reddy, M. S., Schulz, M., Takemura, T., Zhou, M. A review of measurement-based assessment of the aerosol direct radiative effect and forcing., *Atmos. Chem. Phys.*, 6, 613-666, doi.org/10.5194/acp-6-613-2006, 2006.

Formatted: Font color: Text 1

Formatted: Highlight

1080 Yu, H., Remer, L. A., Kahn, R. A., Chin, M., & Zhang, Y., Satellite perspective of aerosol intercontinental transport: From qualitative tracking to quantitative characterization, *Atmospheric Research*, 124, 73– 100. <https://doi.org/10.1016/j.atmosres.2012.12.013>, 2013.

Formatted: Highlight

1085 Yu, H., Tan, Q., Chin, M., Remer, L. A., Kahn, R. A., Bian, H., et al., Estimates of African dust deposition along the trans-Atlantic transit using the decadelong record of aerosol measurements from CALIOP, MODIS, MISR, and IASI. *J. Geophys. Res. Atmos.*, 124, 7975– 7996, <https://doi.org/10.1029/2019JD030574>, 2019.

Formatted: Highlight

1090 Yu, H., Yang, Y., Wang, H., Tan, Q., Chin, M., Levy, R. C., Remer, L. A., Smith, S. J., Yuan, T., and Shi, Y.: Interannual variability and trends of combustion aerosol and dust in major continental outflows revealed by MODIS retrievals and CAM5 simulations during 2003–2017, *Atmos. Chem. Phys.*, 20, 139–161, <https://doi.org/10.5194/acp-20-139-2020>, 2020.

Formatted: Highlight

1095 Yuan, T., Remer, L. A., Pickering, K. E., and Yu, H.: Observational evidence of aerosol enhancement of lightning activity and convective invigoration, *Geophys. Res. Lett.*, 38, L04701, <https://doi.org/10.1029/2010GL046052>, 2011a.

Formatted: Highlight

1100 Yuan, T., Remer, L. A., and Yu, H.: Microphysical, macrophysical and radiative signatures of volcanic aerosols in trade wind cumulus observed by the A-Train, *Atmos. Chem. Phys.*, 11, 7119–7132, <https://doi.org/10.5194/acp-11-7119-2011>, 2011b.

Formatted: Highlight

Zemeckis, Robert and Gale, Bob, 1989. *Back To the Future Part II*. Amblin Entertainment and Universal Pictures. Produced by Bob Gale and Neil Canton. 108 minutes.

Formatted: Highlight

Zhang, J., Christopher, S.A. Longwave radiative forcing of Saharan dust aerosols estimated from MODIS, MISR, and CERES observations on Terra. *Geophys. Res. Lett.*, 30, 2188, doi:10.1029/2003GL018479, 2003.

Formatted: Highlight

1105 | **Zhang, J.**, Reid, J.S., Westphal, D.L., Baker, N.L., Hyer, E.J. A system for operational aerosol optical depth data assimilation over global oceans. *J. Geophys. Res. Atmos.*, 113, D10208, doi:10.1029/2007JD009065, 2008.

Formatted: Highlight

| **Zhang, J. and Reid, J. S.:** A decadal regional and global trend analysis of the aerosol optical depth using a data-assimilation grade over-water MODIS and Level 2 MISR aerosol products, *Atmos. Chem. Phys.*, 10, 10949–10963, doi.org/10.5194/acp-10-10949-2010, 2010.

Formatted: Highlight

1110 | **Zhang, Z., Fu, G., and Hasekamp, O.:** Aerosol retrieval over snow using RemoTAP, *Atmos. Meas. Tech. Discuss.* [preprint], <https://doi.org/10.5194/amt-2023-127>, in review, 2023.

Formatted: Highlight

| **Zheng, J., Zhang, Z., Garnier, A., Yu, H., Song, Q., Wang, C., Dubuisson, P., Di Biagio, C.,** The thermal infrared optical depth of mineral dust retrieved from integrated CALIOP and IIR observations, *Remote Sensing of Environment*, 270, (112841), 10.1016/j.rse.2021.112841, 2022.

Formatted: Highlight

1115

Page 4: [1] Deleted Lorraine Remer 8/30/23 5:02:00 PM

Page 4: [2] Deleted Lorraine Remer 8/30/23 5:01:00 PM

