

1 Mulch application as the overarching factor explaining increase in soil organic carbon  
2 stocks under conservation agriculture in two 8-year-old experiments in Zimbabwe  
3 ~~Conservation agriculture increases soil organic carbon stocks but not soil CO<sub>2</sub> efflux in two~~  
4 ~~8-year-old experiments in Zimbabwe~~

5 Armwell Shumba<sup>a,b,c\*</sup>, Regis Chikowo<sup>a,d</sup>, Christian Thierfelder<sup>e</sup>, Marc Corbeels<sup>f,g</sup>, Johan Six<sup>h</sup>,  
6 Rémi Cardinael<sup>a,b,f</sup>

7 <sup>a</sup>Department of Plant Production Sciences and Technologies, University of Zimbabwe, Harare,  
8 Zimbabwe

9 <sup>b</sup>CIRAD, UPR AIDA, Harare, Zimbabwe

10 <sup>c</sup>Fertilizer, Farm Feeds and Remedies Institute, Department of Research and Specialist  
11 Services, Ministry of Lands, Agriculture, Fisheries, Water and Rural Development, Harare,  
12 Zimbabwe

13 <sup>d</sup>Plant, Soil and Microbial Sciences Department, Michigan State University, East Lansing, MI  
14 48824, USA

15 <sup>e</sup>International Maize and Wheat Improvement Center (CIMMYT), P.O. Box MP 163, Mount  
16 Pleasant, Harare, Zimbabwe

17 <sup>f</sup>AIDA, Univ Montpellier, CIRAD, Montpellier, France

18 <sup>g</sup>IITA, International Institute of Tropical Agriculture, PO Box 30772, Nairobi, 00100, Kenya

19 <sup>h</sup>Department of Environmental Systems Science, ETH Zurich, 8092 Zürich, Switzerland

20 \* Corresponding author. Email: [armwellshumba123@gmail.com](mailto:armwellshumba123@gmail.com)

21  
22  
23  
24

25 **Abstract**

26 Conservation agriculture (CA), combining reduced or no tillage, permanent soil cover and  
27 improved rotations, is often promoted as a climate-smart practice. However, our understanding  
28 about the impact of CA and its respective three principles on top and sub-soil organic carbon  
29 (SOC) stocks ~~and on soil CO<sub>2</sub> efflux~~ in low input cropping systems of sub-Saharan Africa is  
30 rather limited. The study was conducted at two long-term experimental sites established in  
31 2013 in Zimbabwe. The soil types were abruptic Lixisols at Domboshava Training Centre  
32 (DTC) and xanthic Ferralsol at the University of Zimbabwe farm (UZF). Six treatments,  
33 replicated four times were investigated: conventional tillage (CT), conventional tillage with  
34 rotation (CTR), NT, no-tillage with mulch (NTM), no-tillage with rotation (NTR), no-tillage  
35 with mulch and rotation (NTMR). Maize (*Zea mays* L.) was the main crop and treatments with  
36 rotation included cowpea (*Vigna unguiculata* L. Walp.). SOC concentration and bulk density  
37 were determined for samples taken from the 0-5, 5-10, 10-15, 15-20, 20-30, 30-40, 40-50, 50-  
38 75 and 75-100 cm depths. Cumulative organic inputs to the soil were also estimated in all  
39 treatments. Gas samples were regularly collected using the static chamber method during the  
40 2019/20 and 2020/21 cropping seasons and during the 2020/21 dry season. SOC stocks at  
41 equivalent soil mass were significantly ( $p < 0.05$ ) higher under NTM, NTR and NTMR  
42 compared to NT and CT in top 5 and 10 cm layers at UZF, while SOC stocks were only  
43 significantly higher under NTM and NTMR compared to NT and CT in top 5 cm at DTC. NT  
44 alone had a slightly negative impact on top SOC stocks. Cumulative SOC stocks were not  
45 significantly different between treatments when considering the whole 100 cm soil profile.  
46 ~~Regardless of larger organic carbon inputs in mulch treatments, there were no significant~~  
47 ~~differences in CO<sub>2</sub> efflux between treatments, but it was higher in maize rows than in inter-~~  
48 ~~rows as a result of autotrophic respiration from maize roots.~~ Our results ~~showed~~ show the  
49 overarching role of crop residue mulching in CA cropping systems in enhancing SOC ~~storage~~

50 stocks but that this effect ~~is~~was limited to the topsoil. The highest cumulative organic carbon  
51 inputs to the soil were observed in NTM treatments at the two sites, and this could probably  
52 explain the positive effect on SOC stocks. Our results also showed that the combination of at  
53 least two CA principles including mulch is required to increase SOC stocks in these low  
54 nitrogen input cropping systems.

55 **Key words:** organic inputs, climate change mitigation, climate-smart agriculture, deep soil  
56 organic carbon, ~~mulch~~, sustainable intensification

## 58 1. Introduction

59 Soil organic carbon (SOC) is an important determinant of soil fertility, productivity and  
60 sustainability, and is a useful indicator of soil quality in tropical agricultural systems where  
61 nutrient poor and highly weathered soils are managed with little external inputs (Lal, 1997;  
62 Feller and Beare, 1997; Chivenge et al., 2007). Therefore, rebuilding depleted SOC stocks in  
63 such soils holds potential to contribute to climate change mitigation (Bossio et al., 2020;  
64 Minasny et al., 2017; Swanepoel et al., 2016) through sustainable management of agricultural  
65 soils (Paustian et al., 2016; Dignac et al., 2017).

66 Conservation agriculture (CA), based on minimum soil disturbance, crop residue retention and  
67 crop rotation, has been known to improve surface SOC, with beneficial effects on soil  
68 functioning such as improved water infiltration (Thierfelder and Wall, 2012, 2009) and better  
69 aggregate stability (Six et al., 1999; Thierfelder and Wall, 2012). The potential of CA to  
70 increase SOC stocks and thereby mitigate climate change has, however, been much debated  
71 (Corbeels et al., 2020a). ~~The~~but the general understanding is that, this potential is relatively  
72 low (Du et al., 2017; Powlson et al., 2014, 2016; Cheesman et al., 2016; Corbeels et al., 2020a).

73 ~~which is well demonstrated in sub-Saharan Africa (SSA) (Cheesman et al., 2016; Corbeels et~~  
74 ~~al., 2019; Powlson et al., 2016).~~ In fact, soil C storage has often been over-estimated for CA  
75 due to shallow soil sampling. Compared to conventional tillage systems, no-tillage redistributes  
76 SOC in the soil profile, with higher concentrations in the topsoil but potentially lower  
77 concentrations below, which can result in no differences in whole profile SOC stocks between  
78 no-tillage and conventional tillage (Angers and Eriksen-Hamel, 2008). However, this lack of  
79 significant differences in many studies assessing whole profile SOC stocks suffer from not  
80 enough statistical power to accurately assess the potential significant SOC changes  
81 (Kravchenko and Robertson, 2011).

82 CA can potentially build SOC in deeper soil layers from e.g. the use of cover crops with more  
83 extended root systems (Kell, 2011; Thorup-Kristensen et al., 2020; Yang et al., 2023).  
84 However, proper soil sampling strategies, to account for both topsoil and subsoil (> 30 cm  
85 depth) SOC stocks must, therefore, be prioritized. Soil sampling has often been limited to the  
86 top soil plough layer (0-30 cm) in the past two decades (Dube et al., 2012; Powlson et al., 2016;  
87 Patra et al., 2019; Yost and Hartemink, 2020), where SOC concentrations, root densities  
88 (Chikowo et al., 2003) and microbial activities (Mtambanengwe et al., 2004) are generally  
89 largest (Rumpel et al., 2012) and which is the minimum default soil depth recommended by  
90 the Intergovernmental Panel for Climate Change (IPCC, 2019). In a meta-analysis of SOC  
91 stocks in the top 1 m of soils, Balesdent et al. (2018) found that soils below 0.3 m contain on  
92 average 47 % of total SOC stock in the 1 m soil depth and accounts for 19 % of the SOC that  
93 has been recently incorporated. Therefore, focusing on topsoil only, could underestimate the  
94 potential of agricultural management practices to store SOC (Cardinael et al., 2015). In turn,  
95 this can give wrong conclusions on the climate change mitigation potential of agricultural  
96 management practices.

97 There ~~has~~have been many studies on the effects of CA on crop productivity and soil health  
98 benefits (Corbeels, Naudin, et al., 2020; Kimaro et al., 2016; Mhlanga et al., 2022a; Swanepoel  
99 et al., 2018; Thierfelder et al., 2015, 2017; Thierfelder & Mhlanga, 2022), and other studies  
100 have fuelled the debate on CA practicality and adoption in SSA (Giller et al., 2009, 2015;  
101 Kassam et al., 2019). However, the effects of CA on SOC dynamics ~~and soil CO<sub>2</sub> efflux~~  
102 ~~have~~has not been widely investigated in SSA. Thierfelder et al., (2017) have alluded to the fact  
103 that, data on climate change mitigation potential of CA in southern Africa is scanty hence the  
104 need for more research to better quantify the mitigation effects of CA as a climate-smart  
105 technology. It has also been observed that depending on the socio-economic and biophysical  
106 conditions, farmers may find it easier to adopt certain CA principles and/or their different  
107 combinations (Mbanyele et al., 2021; Baudron et al., 2012), although this also opened up new  
108 debates (Thierfelder et al., 2018). Therefore, in this study, the focus was on the individual  
109 versus combined effects of CA principles (no-tillage, crop residue retention, crop rotation) on  
110 SOC stocks ~~and soil CO<sub>2</sub> effluxes~~.

111 As changes in SOC stocks take time to be detected, long-term experiments are crucial, but are  
112 rare, especially in Africa (Bationo et al., 2013; Cardinael et al., 2022; Powlson et al., 2016;  
113 Thierfelder and Mhlanga, 2022). This study was conducted on two long-term experiments  
114 established in 2013 in Zimbabwe. We hypothesized that the full combination of CA  
115 components would be associated with ~~more rapid~~higher increases of SOC stocks ~~and soil CO<sub>2</sub>~~  
116 ~~efflux~~ than adoption of only one component, and that this increase would mainly be due to  
117 increased C inputs to the soil and minimum soil disturbance. However, C inputs due to crop  
118 rotation might be indirect and we therefore hypothesise that the productivity of the crops is  
119 enhanced due to reduction on biotic pressure (pests and diseases), and therefore C inputs to the  
120 soil might be increased too. Secondly, cereals in a cereals-legumes rotations tend to benefit  
121 from added soil nitrogen through biological nitrogen fixation by the preceding legume crop

122 hence more productivity. The third hypothesis is that crop diversification enhances soil  
123 biological processes via different root systems, and enhanced microfauna diversity and/or  
124 abundance (e.g. mycorrhizae) that could improve aggregate stability and therefore physical  
125 protection of soil carbon. Lastly, high quality residues (from the legume crop) have been shown  
126 to be preferentially stabilized in the soil due to a higher carbon use efficiency of soil microbes  
127 (Cotrufo et al., 2013; Kopittke et al., 2018).


## 129 2. Materials and methods

### 130 2.1 Study sites

131 The study was conducted at two long-term experimental sites established in November 2013  
132 by CIMMYT. The site at the University of Zimbabwe Farm (UZF) is located about 12 km north  
133 of Harare city centre (31° 00' 48" E; 17° 42' 24" S), while the site at the Domboshava Training  
134 Centre (DTC) is located about 30 km north-east of Harare (31° 07' 33" E; 17° 35' 17" S). UZF  
135 soils are dolerite-derived xanthic *Ferralsols* (FAO classification) and are medium-textured  
136 sandy clay loams (34 % clay) in the top 20 cm with a subsoil (20-40 cm) of slightly higher clay  
137 content (38 %). DTC soils are granite-derived abruptic *Lixisols* (FAO classification) and are  
138 light-textured sandy loams (15 % clay) in the 0-20 cm layer, overlying abruptly a heavier-  
139 textured subsoil (20-40 cm) of 30 % clay (Figure 1).

140 The two study sites have a sub-tropical climate with cool, dry winters and hot, wet summers  
141 with mean annual minimum and maximum temperatures of 12°C and 25°C, respectively  
142 (Mapanda et al., 2010). The rainy season starts in November and tails off in March with a mean  
143 annual rainfall of 826 and 814 mm at UZF and DTC, respectively (Mhlanga, et al., 2022b).  
144 Cumulative seasonal rainfall in 2019/20 (474 mm) was almost half of rainfall received in the

145 [2020/21 season \(932 mm\) at DTC \(Shumba et al., 2023b\). At UZF, cumulative seasonal rainfall](#)  
 146 [was 551 mm and 637 mm in the 2019/20 and 2020/21 cropping seasons. On average, minimum](#)  
 147 [and maximum temperatures were 16.9 and 28.1 °C in 2019/20 and 15.5 and 27.2 °C in 2020/21](#)  
 148 [at DTC and 15.5 and 28.6 °C in 2019/20 and 15.3 and 27.5 °C in 2020/21 cropping season at](#)  
 149 [UZF.](#)



150 **Figure 1**—Soil texture to 1 m soil depth at the DTC (top) and UZF (bottom) sites in Zimbabwe.  
 151  
 152 Error bars represent standard errors (N = 4).

153

154 ~~The two study sites have a sub-tropical climate with cool, dry winters and hot, wet summers~~  
155 ~~with mean annual minimum and maximum temperatures of 12°C and 25°C, respectively~~  
156 ~~(Mapanda et al., 2010). The rainy season starts in November and tails off in March with a mean~~  
157 ~~annual rainfall of 826 and 814 mm at UZF and DTC, respectively (Mhlanga, et al., 2022b).~~  
158 ~~Cumulative seasonal rainfall in 2019/20 (474 mm) was almost half of rainfall received in the~~  
159 ~~2020/21 season (932 mm) at DTC (Shumba et al., 2023b)(Figure S1). At UZF, cumulative~~  
160 ~~seasonal rainfall was 551 mm and 637 mm in the 2019/20 and 2020/21 cropping seasons. On~~  
161 ~~average, minimum and maximum temperatures were 16.9 and 28.1 °C in 2019/20 and 15.5 and~~  
162 ~~27.2 °C in 2020/21 at DTC and 15.5 and 28.6 °C in 2019/20 and 15.3 and 27.5 °C in 2020/21~~  
163 ~~cropping season at UZF.~~

164

## 165 **2.2 Experimental treatments and crop management**

166 Two identical experiments were set up at the study sites and treatments were maintained every  
167 season since November 2013. The experiments were set up in a randomised complete block  
168 design (RCBD) with eight treatments replicated in four blocks. However, in this study we  
169 investigated only six of these treatments. All crop residues were removed soon after harvesting  
170 in all treatments, stored and then applied prior to planting in treatments with mulch. The six  
171 treatments in our study were:

- 172 i. Conventional tillage (CT) – land preparation was done through digging with a hand hoe  
173 and maize (*Zea mays* L.) was sown as a sole crop in rip lines that were created  
174 afterwards using an animal-drawn Magoye ripper (a traditional plough with the



175 mouldboard replaced with a ripper tine) at DTC, and in planting basins (approximately  
176 10 cm diameter and 10 cm depth) created using a hand hoe at UZF.

177 ii. Conventional tillage with rotation (CTR) – land preparation was done as in the CT  
178 treatment and maize was rotated with cowpea (*Vigna unguiculata* L.).

179 iii. No-tillage (NT) – sole maize was sown in rip lines created using an animal-drawn  
180 Magoye ripper (no further soil disturbance was done) at DTC, and in planting basins  
181 (approximately 15 cm diameter and 15 cm depth) created using a hand hoe at UZF.

182 iv. No-tillage with mulch (NTM) – maize was sown as in the NT treatment and maize  
183 residues from the previous season were applied on the soil surface between maize  
184 rows at planting at a rate of 2.5 t DM ha<sup>-1</sup>.

185 v. No-tillage with rotation (NTR) – maize was sown in rip lines and rotated with cowpea.

186 vi. No-tillage with mulch and rotation (NTMR) – maize was sown in rip lines and rotated  
187 with cowpea and maize residues were applied on the soil surface between maize rows  
188 at planting at a rate of 2.5 t DM ha<sup>-1</sup>.

189 Crop residues were removed every year after harvest and weighed in again to maintain the  
190 exact 2.5 t ha<sup>-1</sup> residue weight year after year. There was a total of 24 plots at each site which  
191 were 6 m wide and 12 m long (72 m<sup>2</sup>). Treatments with rotation (CTR, NTR, NTMR) were  
192 split into 6 m wide and 6 m long (36 m<sup>2</sup>) subplots where maize and cowpea were grown  
193 interchangeably every season (maize was sown on one side of the plot while cowpea on the  
194 other side).

195 The inter-row spacing was 90 cm and 45 cm for maize and cowpea, respectively, and the in-  
196 row spacing was 25 cm for both crops which translated to 44,444 and 88,888 plants ha<sup>-1</sup>,  
197 respectively. Three seeds were planted per planting station and thinned to one after emergence.

198 Basal dressing of nitrogen (N), phosphorus (P) and potassium (K) was applied within 5 cm of

199 the seeds in the form of compound fertilizer for both maize and cowpea at 11.6 kg N ha<sup>-1</sup>, 10.6  
200 kg P ha<sup>-1</sup> and 9.6 kg K ha<sup>-1</sup>, respectively. Nitrogen top dressing to maize only, was applied at  
201 4 and 8 weeks after emergence (WAE) in two equal splits of 23.1 kg N ha<sup>-1</sup> each, as ammonium  
202 nitrate when soil moisture was adequate. However, in the 2019/20 cropping season at both sites  
203 and in the 2020/21 cropping season at UZF, the first N top dressing was delayed by an average  
204 of 4.5 and 2.0 weeks, respectively, due to mid-season dry spells. Ammonium nitrate was side  
205 dressed within 5 cm of the maize stems. Glyphosate [N-(phosphono-methyl) glycine], a pre-  
206 emergent non-selective herbicide was applied at 1.025 L active ingredient ha<sup>-1</sup> soon after  
207 sowing to control weeds. This was followed by manual hoe weeding whenever weeds reached  
208 a maximum of 10 cm height or 10 cm in diameter for stoloniferous weeds to achieve a weed  
209 clean field. More details about the experiment can be found in Shumba et al., (2023b) and  
210 Mhlanga et al., (2022a).

211

### 212 **2.3 Soil sampling for bulk density determination and soil organic carbon analysis**

213 Soil sampling was done in May 2021 at both sites. For each treatment and replicate, two  
214 sampling points in the maize rows and two sampling points in the middle of the inter-rows  
215 were randomly selected. The two samples from the rows were pooled into one sample per  
216 depth, similarly to the two samples taken in the inter-rows. The following nine depth  
217 increments were considered for both SOC and bulk density (BD) measurements: 0-5, 5-10, 10-  
218 15, 15-20, 20-30, 30-40, 40-50, 50-75 and 75-100 cm. Undisturbed soil samples using a metal  
219 cylinder (5 cm diameter and 5 cm height) were taken from the following depth ranges 0-5, 5-  
220 10, 10-15, 15-20 and 20-30 cm for both SOC and BD measurements. A motorized, hand-held  
221 soil corer, with an inside diameter of 10 cm, was used to take samples for the 30-40, 40-50, 50-  
222 75 and 75-100 cm depths for SOC analysis from the same positions where undisturbed samples

223 were taken. As no significant differences in BD were found below 20 cm between the different  
224 treatments at the two sites (see results section) and to avoid too much destruction of the  
225 experimental plots, two soil pits were opened at the edges of the experimental plots (also  
226 cropped with maize since 2013) at each site to take BD samples for the 30-40, 40-50, 50-75  
227 and 75-100 cm depths. As a result, BD below 30 cm depth was assumed the same across the  
228 treatments.

229 Soil samples were crumbled and fresh weight was determined using a field scale. Soil moisture  
230 was determined on a sub-sample by drying it in an oven at 105°C for 48 hours. All samples  
231 were then air-dried and sieved through a 2 mm sieve to determine the mass proportion of coarse  
232 soil (> 2 mm) as a fraction of the whole sample. Bulk density (BD) was determined by dividing  
233 the dry mass of soil by the volume of the cylinder. Subsamples from the  $\leq 2$  mm soil fraction  
234 were grinded to < 200  $\mu\text{m}$  for SOC analysis. SOC ~~concentration~~ was analysed ~~with a CHN~~  
235 ~~elemental analyser in the ISO9001:2015-certified IRD LAMA's laboratory in Dakar by dry~~  
236 ~~combustion on 100-mg aliquots of soil (ground to < 200  $\mu\text{m}$ ) using a CHN elemental analyser~~  
237 ~~(Thermo Finnigan Flash EA1112, Milan, Italy).~~

238

#### 239 **2.4 Soil organic carbon stocks calculation**

240 The mass proportion of the coarse fraction (> 2 mm) was removed to calculate SOC stocks.

241 The equivalent soil mass (ESM) approach was used to determine SOC stocks to avoid  
242 systematic bias in SOC calculation when using the fixed depth method (Ellert and Bettany,  
243 1995; Wendt and Hauser, 2013; von Haden et al., 2020). We defined reference soil mass  
244 profiles for each site, based on the lowest cumulative soil mass obtained for each replicate. For  
245 these references, cumulative soil mass layers were 0-650, 650-1340, 1340-2060, 2060-4160,  
246 4160-5590, 5590-7040, 7040-10550, 10550-13770 Mg soil ha<sup>-1</sup> at DTC and 0-460, 460-870,

247 870-1330, 1330-1840, 1840-2760, 2760-4030, 4030-5300, 5300-8190, 8190-11050 Mg soil  
248 ha<sup>-1</sup> at UZF, which roughly corresponded to soil depth layers of 0-5, 5-10, 10-15, 15-20, 20-  
249 30, 30-40, 40-50, 50-75, 75-100 cm, respectively. SOC stocks were calculated on the basis of  
250 the same soil mass as the reference profile but different soil depth layers which varied by < 1.5  
251 and < 0.6 cm at DTC and UZF, respectively. As mulch was only applied between maize rows,  
252 and fertilizer was only applied on maize rows, it was estimated that the row and interrow space  
253 represented 22 and 78 % respectively, hence SOC calculations were weighted accordingly  
254 (Shumba et al., 2023b). Change in cumulative SOC stock between treatments for a given soil  
255 depth was determined using the CT treatment as the reference treatment:

$$256 \Delta SOC \text{ stock} = SOC \text{ stock}_{treatment(i)} - SOC \text{ stock}_{CT(i)}, \text{ (Equation 1)}$$

257 where SOC stock<sub>treatment</sub> is the cumulative SOC stock per treatment (CTR, NT, NTM, NRT,  
258 NTMR) at a given soil layer and (i) representing 0-5, 0-10, 0-15, 0-20, 0-30, 0-40, 0-50, 50-  
259 75, 75-100 cm.

260 SOC accumulation or loss rates (kg C ha<sup>-1</sup> yr<sup>-1</sup>) were calculated by dividing the change in  
261 stocks by the number of years between the establishment of the experiment and the time of soil  
262 sampling (8 years):

$$263 SOC \text{ accumulation/loss rate} = \frac{\Delta SOC \text{ stocks}}{8} \times 1000, \text{ (Equation 2)}$$

264

## 265 **2.5 Estimation of organic carbon inputs to the soil**

266 Maize and cowpea yield and aboveground biomass were measured since the inception of the  
267 experiment, except for cowpea during the 2013/14 season. This data gap was filled by using  
268 the average cowpea yield and aboveground biomass values across seasons (from 2013/14 to

269 2020/21). We assumed that 5 % of the maize aboveground vegetative biomass remained in the  
270 field because maize stalk slashing at harvesting did not remove the whole stem. A root:shoot  
271 ratio of 0.16 and 0.06 for maize and cowpea, respectively (Amos and Walters, 2006; Kahn and  
272 Schroeder, 1999; Kimiti, 2011) was used to estimate the contribution of roots to organic C  
273 inputs to the soils. Organic C input contribution from weeds was assumed insignificant since  
274 there was effective control of weeds through the use of pre-emergence herbicide (glyphosate)  
275 and timely manual weeding throughout the cropping season. We also assumed that the relative  
276 amounts of organic C transferred through rhizodeposition was the same for maize and cowpea  
277 (i.e. 0.45 x root C biomass (Balesdent et al., 2011) and that the organic C content of all plant  
278 parts was 430 g kg<sup>-1</sup> (Ma et al., 2018). Cumulative organic C inputs to the soil were then  
279 estimated for each treatment (Cardinael et al., 2022).

280

## 281 **2.6 Gas sampling, analyses and flux determinations**

282 ~~The static chamber methodology was used for CO<sub>2</sub> gas sampling. The static chambers had PVC~~  
283 ~~base rings (height = 0.1 m and inside radius = 0.1 m) and PVC cylindrical lids (height = 0.2 m~~  
284 ~~and inside diameter = 0.1 m). Base rings were semi-permanently driven 0.07 m into the soil to~~  
285 ~~avoid possible gas leakages and contamination by lateral diffusion (Abalos et al., 2013; Clough~~  
286 ~~et al., 2020). The lids had an airtight and self-sealing rubber septum on top through which gas~~  
287 ~~was sampled. During gas sampling, the lids were inserted about 0.02 m into the base rings and~~  
288 ~~the contact area between the base rings and the lids was always smeared with petroleum jelly to~~  
289 ~~avoid possible leakages of trapped gas. The static chambers were painted white to minimize~~  
290 ~~temperature changes in the chamber headspace from the sun's radiative heat.~~

291 ~~Surface area coverage for each chamber was 0.0314 m<sup>2</sup> and headspace volume of 0.006 m<sup>3</sup>. Gas~~  
292 ~~sampling was done simultaneously in the row and interrow spaces, each replicate having a~~

293 chamber in the row and in the middle of the inter-row (Shumba et al., 2023b). It should be noted  
294 that, CO<sub>2</sub> measured in this study consisted of effluxes coming both from autotrophic and  
295 heterotrophic respiration.

296 A 20 mL syringe was used to collect gas samples at time 0 (immediately after securing the  
297 chamber) and after 48 minutes of gas trapping. The gas samples were pressurised into pre-  
298 evacuated 12 mL Exetainer glass vials (Labco Ltd., Lampeter SA48, United Kingdom). Linearity  
299 tests were carried out at both sites by collecting gas samples at times 0, 15, 30, 48 and 60 minutes  
300 of gas trapping. Results showed that CO<sub>2</sub> emissions increased linearly with time, suggesting that  
301 two gas samplings at 0 and 48 minutes were relevant for this study since no saturation was  
302 observed (data not shown). Gas sampling was done between 10 am and 12 pm on every sampling  
303 day.

304 CO<sub>2</sub> efflux measurements were carried out during the cropping season (November to April) in  
305 2019/20 and 2020/21, but in 2021, CO<sub>2</sub> efflux measurements were extended into the dry season  
306 (May to September). Gas sampling was done at least every two weeks during the cropping  
307 season, with additional sampling following fertilizer applications and rainfall events (Shumba et  
308 al., 2023b).

309 CO<sub>2</sub> was quantified at ETH Zurich by gas chromatography using the thermal conductivity  
310 detector and CO<sub>2</sub> fluxes were calculated as the differences in concentration between the 0 and 48  
311 minutes sampling times:

$$312 F = \frac{(GC_f - GC_o) \times V}{T \times A}, \text{ (Equation 4)}$$

313 where  $F$  is the gas flux (mg CO<sub>2</sub> m<sup>-2</sup> hr<sup>-1</sup>),  $GC_f$  and  $GC_o$  are the gas concentration (ppm) at end  
314 (time 48 minutes) and start (time 0 minutes) of chamber closure,  $V$  is the chamber volume (mL),  
315

~~$T$  is the duration of the chamber closure (hours) and  $A$  is the surface area covered by the static chamber ( $m^2$ ).~~

## ~~2.7 Cumulative soil CO<sub>2</sub>-C emissions~~

~~Cumulative CO<sub>2</sub>-C emissions were determined using linear interpolation between sampling points by multiplying the mean flux of two successive sampling dates by the length of the period between sampling and adding that amount to the previous cumulative total (Dorich et al., 2020). Cumulative efflux per treatment was computed as the weighted contribution from row and inter-row effluxes (Shumba et al., 2023b).~~

## ~~2.8.6 Data analysis~~

~~The full dataset is available in the CIRAD repository (Shumba et al., 2023a). Statistical analyses were performed using R software, version 4.0.0 (R Core Team 2020). Prior to analysis, CO<sub>2</sub> data were checked for normality by both visual inspection (Quantile-Quantile plots and density distributions) and with the Shapiro-Wilk test. Linear mixed-effect models were fitted to daily CO<sub>2</sub> emissions using the *lmer* function from the *lme4* package (Bates, 2010), using as fixed effects the site (DTC, UZF), the season (2019/20, 2020/21), the treatment (CT, CTR, NT, NTM, NTR, NTMR) and the chamber position (row vs inter-row). The chamber number nested in the replicate was considered as random factor. The final models were chosen based on the lowest Akaike information criterion (AIC) and on the lowest Bayesian information criterion (BIC). An analysis of variance (ANOVA) was then done on the fitted models. Separation of means was done using the post hoc Tukey test at 5 % significance level using the *emmeans* function from the *emmeans* package (Bolker et al., 2009).~~



339 ~~For soil data, n~~Normality was tested by the Kolmogorov-Smirnov test. After confirming that data  
340 were normally distributed, analyses of variance (ANOVA) was carried out to establish any  
341 significant treatment effects on BD, SOC concentration, and SOC stock. Separation of means  
342 was done using the post hoc Tukey test at 5 % significance level using the *emmeans* function  
343 from the *emmeans* package (Bolker et al., 2009).~~Subsequent mean separation was done using~~  
344 ~~Tukey's test.~~

### 345 3. Results





#### 346 3.1 Soil bulk density

347 The different cropping systems (CT, CTR, NT, NTM, NTR, NTMR) had no significant ( $p >$   
348 0.05) effect on BD across all soil depths except in the 5-10 cm depth in the inter-row at DTC  
349 (Figure 2) where BD was on average 5 % lower in the conventional tillage treatments (CT, CTR)  
350 than in the no-tillage treatments (NT, NTR). However, soil depth and location (row or inter-row),  
351 and the soil depth x location interaction had significant ( $p < 0.001$ ) effects on BD. In the tillage  
352 layer (0-15 cm), BD was at least 2 % higher in the inter-rows than in the rows at both sites. In  
353 the deeper soil layer (15 – 30 cm), there were no significant ( $p > 0.05$ ) differences. BD for depths  
354 below 30 cm were the same across treatments since it was determined from pits outside the  
355 experiment. It ranged between 1.47 – 1.51 and 1.47 – 1.49 g cm<sup>-3</sup> (Table S1) in the subsoil (30 –  
356 100 cm layers) at DTC and UZF, respectively.





357  
358  
359  
360  
361  
362






364 **Figure 2.** Top soil bulk density (0-30 cm) at the Domboshava Training Centre (DTC) and  
365 University of Zimbabwe Farm (UZF) experimental sites in Zimbabwe. CT: conventional tillage,  
366 CTR: conventional tillage with rotation, NT: no-tillage, NTM: no-tillage with mulch, NTR: no-  
367 tillage with rotation, NTMR: no-tillage with mulch and rotation. Error bars represent standard  
368 errors (N = 4).

### 369 **3.2 SOC concentrations**

370 SOC concentration decreased significantly ( $p < 0.001$ ) with soil depth (Figure 3, Table S2) and  
371 was highest in the surface soil (0-5 cm) for all treatments (Table S2). There were significant  
372 treatment effects in the 0-5 cm ( $p = 0.001$ ) and 5-10 cm ( $p = 0.005$ ) soil layers at DTC and in  
373 the 0-5 cm layer ( $p < 0.001$ ) only, at UZF. NTM had significantly ( $p < 0.05$ ) higher SOC  
374 concentration compared to conventional tillage treatments (CT, CTR) and NT in the 0-5 cm  
375 soil layer at both sites (Figure 3); the increase in SOC concentration ranged between 31 to 46  
376 % and 14 to 22 % at DTC and UZF, respectively. However, SOC concentration in NTM was  
377 equal ( $p > 0.05$ ) to NTR and NTMR treatments at both sites.

378 In the 5-10 cm soil layer of DTC, SOC concentrations in NTM and NTR were at least 19 %  
379 higher ( $p = 0.005$ ) than in NT and CT (Table S2). There were no significant ( $p > 0.05$ ) treatment  
380 effects on SOC concentration below 10 cm soil depth at DTC and below 5 cm depth at UZF.





382  
 383 **Figure 3.** Soil depth distribution of organic carbon concentration for the different experimental  
 384 treatments at the Domboshava Training Centre (DTC) and University of Zimbabwe Farm  
 385 (UZF) experimental sites in Zimbabwe. Error bars represent standard errors (N = 4). CT:  
 386 conventional tillage, CTR: conventional tillage with rotation, NT: no-tillage, NTM: no-tillage  
 387 with mulch, NTR: no-tillage with rotation, NTMR: no-tillage with mulch and rotation.

Mis en forme : Police :Non Gras

388

389 **3.3 SOC stocks**

390 There were significant ( $p < 0.05$ ) treatment effects on SOC stocks per soil layer in the [top 5 at](#)  
 391 [UZF and 10 cm 0-5 and 5-10 cm soil layers at DTC and at DTC the 0-5 cm soil layer at UZF](#)  
 392 (Table S3). Compared to CT, CTR and NT, NTM had at least 1.1 and 1.3 times more SOC  
 393 stocks in the top 5 and 10 cm layers at UZF and DTC, respectively. In terms of cumulative

394 SOC stocks, significant ( $p < 0.05$ ) treatment effects were limited to the top 30 cm soil layer at  
395 DTC and the 20 cm layer at UZF, where no tillage with mulching (NTM) increased SOC stocks  
396 (Table 1). There were no significant ( $p > 0.05$ ) tillage effects on SOC stocks (CT vs NT) for  
397 both sites. The rotation component had no significant ( $p > 0.05$ ) effects on SOC stocks when  
398 comparing CTR and NTR at DTC. However, the maize-cowpea rotation under NT (NTR) had  
399 at least 16 % higher SOC stocks in the top 30 cm compared to NT. In contrast, NTR had at  
400 least 7 % more SOC stocks than CTR in the top 10 cm soil layer at UZF, though there were no  
401 significant ( $p > 0.05$ ) differences in SOC stocks between NTR and NT. Compared to NT and  
402 CT, the mulching component significantly ( $p < 0.05$ ) increased SOC stocks by at least 8 % at  
403 UZF and 13 % at DTC in the top 20 and 30 cm soil layers, respectively. SOC stocks in the full  
404 CA treatment (NTMR) were not significantly ( $p > 0.05$ ) different with the other combinations  
405 of CA principles (NTM, NTR) and CTR at DTC. At UZF, the full CA treatment had similar  
406 SOC stocks as all the other NT treatments (NT, NTM, NTR).

407 SOC stocks for the whole soil profile for this study (0-100 cm) were at least 8.1, 3.5 and 2.1  
408 times higher at DTC and 11.6, 4.4 and 2.4 times higher at UZF than the SOC stocks in the 0-5  
409 cm (surface soil), 0-15 cm (tillage depth) and 0-30 cm (IPCC standard depth) layers. SOC  
410 stocks for the subsoil (30-100 cm) ranged from 18.4 to 51.4 Mg C ha<sup>-1</sup> at DTC and 41.9 to  
411 124.9 Mg C ha<sup>-1</sup> at UZF. Therefore, subsoil represented more than half (at least 53 % at DTC  
412 and 58 % at UZF) of SOC stocks for the whole 100 cm soil profile.

413 ~~Table 1: Cumulative SOC stocks at the Domboshava Training Centre (DTC) and University~~  
414 ~~of Zimbabwe Farm (UZF) after 8 years of different tillage, residue and crop management~~  
415 ~~systems. Means in the same row followed by different superscript letters are significantly~~  
416 ~~different and associated errors are standard errors (N = 4). CT: conventional tillage, CTR:~~



419 **Table 1:** Cumulative SOC stocks at the Domboshava Training Centre (DTC) and University  
 420 of Zimbabwe Farm (UZF) after 8 years of different tillage, residue and crop management  
 421 systems. Means in the same row followed by different superscript letters are significantly  
 422 different and associated errors are standard errors (N = 4). CT: conventional tillage, CTR:  
 423 conventional tillage with rotation, NT: no-tillage, NTM: no-tillage with mulch, NTR: no-tillage  
 424 with rotation, NTMR: no-tillage with mulch and rotation.

#### 425 3.4 SOC accumulation and loss rates

426 SOC accumulation rates at UZF differed significantly ( $p < 0.05$ ) with soil depth where top soil  
 427 layers (0-5, 0-10, 0-15, 0-20 and 0-30 cm) had SOC accumulation rates that were at least 6.9  
 428 times less than when considering the 0-100 cm soil profile ~~at UZF~~ (Table 2). In contrast, there  
 429 were no significant ( $p > 0.05$ ) differences, at DTC in SOC accumulation rates with depth ~~at~~  
 430 ~~DTC~~. On average, SOC accumulation rates ranged between 0.13 and 0.08 Mg C ha<sup>-1</sup> yr<sup>-1</sup> in the  
 431 top soil (0-5 cm) to 0.33 and 1.16 Mg C ha<sup>-1</sup> yr<sup>-1</sup> for the whole 1 m soil profile at DTC and  
 432 UZF, respectively. The depth and treatment interaction had no significant ( $p > 0.05$ ) effects at  
 433 both sites.

434 On the other hand, the different treatments in this study had significant ( $p < 0.05$ ) effects in  
 435 SOC accumulation /loss rates in the top 20 cm soil layer at both sites (Table 2). At DTC, NT  
 436 had significant ( $p < 0.05$ ) net loss of SOC in the 0-20 cm layer, ranging between -0.09 and -  
 437 0.02 Mg C ha<sup>-1</sup> yr<sup>-1</sup>, whereas ~~the NT treatments with different combinations of CA principles~~  
 438 ~~under NT~~ (NTM, NTR, NTMR) has had SOC accumulation rates ranging from 0.17 to 0.38  
 439 Mg C ha<sup>-1</sup> yr<sup>-1</sup>. However, maize stover mulching (NTM) had significantly ( $p < 0.05$ ) higher  
 440 SOC accumulation rates than CTR (2.9 – 4.2 times) and NT (5.2 – 13.5 times) in the top 15 cm  
 441 and 20 cm layers, respectively. The different combinations of mulching and rotation under NT  
 442 (NTM, NTR and NTMR) had no significant ( $p > 0.05$ ) differences in SOC accumulation rates.



443 Similarly, rotation treatments (CTR, NTR, NTMR) showed no significant ( $p > 0.05$ )  
 444 differences in SOC accumulation rates. Thus, the full CA treatment had similar SOC  
 445 accumulations rates to treatments with at least 2 combinations of CA principles (NTM and  
 446 NTR) and to CTR.

447 In contrast, at UZF, CTR had significant ( $p < 0.05$ ) net loss of SOC in the top 20 cm (Table 2).  
 448 The no-tillage treatments (NT, NTM, NTR, NTMR) showed significantly ( $p < 0.05$ ) higher  
 449 SOC accumulation rates ( $0.05 - 0.25 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ ) than CTR which ranged between  $-0.07$  to  
 450  $-0.03 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$  in top 10 cm soil layer. NTM had the highest SOC accumulations rates  
 451 ( $0.28$  to  $0.32 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ ) when considering the 0-15 and 0-20 cm soil layers. SOC  
 452 accumulation rates in NTM were at least 3.8, 3.5 and 2.4 times higher than CTR, NT and NTR  
 453 in the top 20 cm. The full CA treatment (NTMR) had significantly ( $p < 0.05$ ) higher SOC  
 454 accumulation rates compared to CTR (2.5 – 5.3 times) in the top 10 cm and lower SOC  
 455 accumulation rate to NTM in the 0-10 cm (2.3 times) and 0-20 cm (10.6 times) soil layer.  
 456 However, there were no significant ( $p > 0.05$ ) differences in SOC accumulation rates between  
 457 treatments beyond 20 cm soil layer at both sites.

458 ~~Table 2: SOC change rates ( $\pm$  standard error,  $N = 4$ ) of the different treatments compared to~~  
 459 ~~CT (conventional tillage) at Domboshava Training Centre (DTC) and University of Zimbabwe~~  
 460 ~~farm (UZF). Means in the same row followed by different superscripts are significantly~~  
 461 ~~different. CTR: conventional tillage with rotation, NT: no tillage, NTM: no tillage with mulch,~~  
 462 ~~NTR: no tillage with rotation, NTMR: no tillage with mulch and rotation, LSD = least~~  
 463 ~~significance difference, ns = not significant, Sig = significance, \*\* =  $p < 0.05$ , \*\*\* =  $p < 0.001$ .~~

Site	Approximate soil depth (cm)	SOC accumulation or loss rate ( $\text{Mg C ha}^{-1} \text{ yr}^{-1}$ )					LSD
		CTR	NT	NTM	NTR	NTMR	
DTC	0-5	$0.06 \pm 0.05^{\text{bc}}$	$-0.02 \pm 0.02^{\text{c}}$	$0.25 \pm 0.05^{\text{a}}$	$0.17 \pm 0.02^{\text{ab}}$	$0.19 \pm 0.04^{\text{ab}}$	0.13
	0-10	$0.10 \pm 0.09^{\text{bc}}$	$-0.04 \pm 0.04^{\text{c}}$	$0.31 \pm 0.09^{\text{a}}$	$0.24 \pm 0.01^{\text{ab}}$	$0.25 \pm 0.08^{\text{ab}}$	0.16
	0-15	$0.12 \pm 0.13^{\text{bc}}$	$-0.07 \pm 0.05^{\text{c}}$	$0.35 \pm 0.13^{\text{a}}$	$0.30 \pm 0.01^{\text{ab}}$	$0.27 \pm 0.12^{\text{ab}}$	0.23

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Tableau mis en forme

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice

Mis en forme : Non Expositant/ Indice




480 rates of about 1.3 to 1.6 Mg C ha<sup>-1</sup> season<sup>-1</sup>yr<sup>-1</sup> for NTMR and 2.0 Mg C ha<sup>-1</sup> season<sup>-1</sup>yr<sup>-1</sup> for  
481 NTM. The other treatments had mean seasonal-annual OC input rates  $\leq$  1.0 Mg C ha<sup>-1</sup> season<sup>-1</sup>yr<sup>-1</sup>  
482 <sup>1</sup>.

483

### 484 **3.6 Soil CO<sub>2</sub>-C efflux and cumulative emissions**

485 ~~Daily soil CO<sub>2</sub> fluxes were significantly ( $p < 0.05$ ) higher in the maize rows than the inter-rows~~  
486 ~~at both sites (Figure 5 and S2). However, there were no significant ( $p > 0.05$ ) differences in~~  
487 ~~daily CO<sub>2</sub> fluxes between treatments. Fluxes of CO<sub>2</sub> spiked at maximum maize vegetative stage~~  
488 ~~(from approximately 25 to 100 days after germination) in the rainy season and tailed off to  $<$~~   
489 ~~50 mg CO<sub>2</sub>-C m<sup>-2</sup> hr<sup>-1</sup> after harvesting and into the dry season (May to September 2021).~~


490 ~~There were no significant ( $p > 0.05$ ) differences in cumulative CO<sub>2</sub>-C emissions for both~~  
491 ~~seasons and sites (Figure 6). Cumulative CO<sub>2</sub>-C emissions ranged from 5.0 to 6.2 Mg CO<sub>2</sub>-C~~  
492 ~~ha<sup>-1</sup> yr<sup>-1</sup> and 5.9 to 7.5 Mg CO<sub>2</sub>-C ha<sup>-1</sup> yr<sup>-1</sup> at DTC and UZF, respectively, in the 2019/20~~  
493 ~~cropping season. In the 2020/21 season, cumulative CO<sub>2</sub>-C emissions ranged between 4.3 to~~  
494 ~~6.3 Mg CO<sub>2</sub>-C ha<sup>-1</sup> yr<sup>-1</sup> and 5.8 to 7.5 Mg CO<sub>2</sub>-C ha<sup>-1</sup> yr<sup>-1</sup> at DTC and UZF, respectively.~~




495

496 **Figure 4**— Estimated cumulative organic carbon (OC) inputs to the soil from the 2013/14 to  
 497 the 2020/21 cropping season for the different treatments at the Domboshava Training Centre  
 498 (DTC) and the University of Zimbabwe Farm (UZF) experimental sites. Error bars represent  
 499 standard errors (n = 4) for the cumulative OC. CT: conventional tillage, CTR: conventional  
 500 tillage with rotation, NT: no-tillage, NTM: no-tillage with mulch, NTR: no-tillage with  
 501 rotation, NTMR: no-tillage with mulch and rotation, ABG: aboveground biomass.

Mis en forme : Police :Non Gras



502  
 503 **Figure 5:** Daily CO<sub>2</sub>-C fluxes during the 2019/2020 (orange) and 2020/21 (black) seasons for  
 504 the different treatments at Domboshava Training Centre (DTC) and University of Zimbabwe  
 505 Farm (UZF) experimental sites. Solid and dotted lines represent fluxes measured in the maize  
 506 intra-row and inter-row spaces respectively; CT: conventional tillage, CTR: conventional tillage  
 507 with rotation, NT: no tillage, NTM: no tillage with mulch, NTR: no tillage with rotation, NTMR:  
 508 no tillage with mulch and rotation, S = sowing and H = harvesting.



509  
 510 **Figure 6:** Cumulative total CO<sub>2</sub>-C emissions for the different treatments in the 2019/20  
 511 cropping season, the 2020/21 cropping season, and the whole year (2020/21) at Domboshava  
 512 Training Centre (DTC) and University of Zimbabwe Farm (UZF). Bars represent standard  
 513 errors (N = 4). CT: conventional tillage, CTR: conventional tillage with rotation, NT: no  
 514 tillage, NTM: no tillage with mulch, NTR: no tillage with rotation, NTMR: no tillage with  
 515 mulch and rotation.

516

517 **4. Discussion**

518 **4.1 Role of soil texture in SOC accumulation**

519 Soil texture is widely recognized to influence SOC stocks (Sun et al., 2020) through physical  
520 and chemical protection of SOC against microbially mediated decomposition (Chivenge et al.,  
521 2007; Mtambanengwe et al., 2004). In our study, the main difference between the two study  
522 sites is soil texture in the top soil (0-30 cm), where DTC had light textured (sandy loams) soils  
523 and UZF had medium to heavy textured (sandy-clay-loams) soils (Figure 1). These soil textural  
524 differences explain why there were no differences in SOC stocks, changes and accumulation  
525 rates between NTM, NTR and NTMR at DTC regardless of higher cumulative OC inputs in  
526 NTM and NTMR (Figure 4). The direct SOC inputs in the top soil at DTC, where SOC was  
527 more concentrated (Table S2, Figure 3) was subject to mineralization because of low clay  
528 content and thus low protection by soil micro-aggregates (Chivenge et al., 2007;  
529 Mtambanengwe et al., 2004; Sun et al., 2020), such that the differences in OC inputs had little  
530 effect. Light textured soils have large pores which cannot protect SOC against microbial  
531 decomposition (Mtambanengwe et al., 2004; Christensen, 1987; Sun et al., 2020; Kravchenko  
532 and Guber, 2017). Additionally, the low clay content meant less surface area for SOC  
533 adsorption (Han et al., 2016; Churchman et al., 2020) which is another mechanism for SOC  
534 protection from mineralization. In contrast, there were differences between NTM and NTR at  
535 UZF in the top soil layers and intermediate between NTM and NTMR. Cumulative OC inputs  
536 in NTMR (12.4 Mg C ha<sup>-1</sup>) were about 75 % of cumulative OC inputs in NTM (16.2 Mg C ha<sup>-</sup>  
537 <sup>1</sup>) (Figure 4) after 8 seasons. The added C, especially from maize stover mulch, most likely  
538 was protected by clay particles as well as formation of organo-mineral complexes (Malepfane  
539 et al., 2022; Chivenge et al., 2007; Jephitha et al., 2023) which protects SOC from mineralization

(Dunjana et al., 2012; Shumba et al., 2020; Button et al., 2022; Rumpel et al., 2012; Sanaullah et al., 2016).

#### 4.1.2 SOC distribution across soil depth

Cumulative SOC stocks for the whole soil profile (0-100 cm) for this study were at least 8.0, 4.0 and 2.0 times higher than the 0-5 (surface soil), 0-15 (tillage depth for the study) and 0-30 (IPCC standard depth for SOC studies), for the two sites, respectively. This means that, ~~>50~~ >50 %over half of SOC stocks for this study were in the sub-soil (30-100 cm) which reflects on the importance of sub-soil SOC stocks. Significant SOC stocks in the sub-soil have also been reported by other authors (Yost and Hartemink, 2020; Balesdent et al., 2018; Cardinael et al., 2015; Harrison et al., 2011; Lal, 2018). Significant ~~treatment effects of mulch and/or rotation under NT effects~~ were restricted to the top 30 cm in our study as well as other studies in SSA (Dube et al., 2012; Powlson et al., 2016) and the world at large (Balesdent et al., 2018; Yost and Hartemink, 2020), which is most likely why the default soil depth for IPCC for SOC studies is 0-30 cm (IPCC, 2019). However, this underestimates whole soil profile C storage (Harrison et al., 2011; Singh et al., 2018; Lorenz and Lal, 2005). ~~Therefore, it is crucial and hence the need to consider depth differentiated assessments of whole soil profiles-profile sampling when monitoring SOC changes-storage in agricultural ecosystems if long term to determine SOC their C sequestration storage is to be effective in potential in~~ the pursuit of climate change mitigation (Malepfane et al., 2022). ~~The differentiated soil depth assessments of SOC also show soil depth sections that are sensitive to disturbance (tillage) and OC inputs through above-, below-ground biomass and organic soil fertility amendments like manure and compost.~~ SOC mineralization is relatively low in the sub-soil due to lack of oxygen and physical protection of SOC (aggregate protected C) (Rumpel et al., 2012; Sanaullah et al., 2016; Shumba et al., 2020; Button et al., 2022). Therefore, in the pursuit to improve subsoil (> 30 cm) SOC stocks through root mortality



564 and exudates, crop varieties with higher root-length densities (Chikowo et al., 2003) in the  
565 subsoil are recommended. Therefore, crop varieties with deep rooting systems are encouraged  
566 to be developed in the pursuit of increasing subsoil OC inputs through root mortality and  
567 exudates.

568 In this study, there is a conspicuous rotation effect at UZF in the subsoil (60 – 100 cm) on SOC  
569 concentration which is however, not significant. However, there is a block effect at UZF on  
570 SOC concentration in the subsoil as shown in Figure S2 (see Figure S1 for DTC where no  
571 block effect was observed) where there seems to be “outliers” in blocks 3 and 4. We decided  
572 not to exclude the “outliers” since we thought it was a block effect rather than a treatment  
573 effect. As a result, there are no significant differences in SOC concentration between treatments  
574 in deep soils. If we exclude the “outliers” in deep soils the graph for SOC concentration is as  
575 shown in Figure S3 where the rotation effects tend to diminish. Nonetheless, all raw data of  
576 this paper can be freely accessed on the CIRAD database repository and linked to this paper  
577 (Shumba et al., 2023a).

#### 578 **4.2.3 Cumulative SOC stocks and accumulation rates**

579 Overall, the insignificant differences in cumulative SOC stocks and SOC accumulation / loss  
580 rates between the different cropping systems in this and other studies (Laura et al., 2022; Leal  
581 et al., 2020) when considering whole soil profile (0-100 cm) is alluded to the dilution effect  
582 due to low OC inputs beyond 30 cm. Organic C inputs in the subsoil (> 30 cm) through crop  
583 root mortality and root exudates are highly reduced due to low root biomass (Button et al.,  
584 2022; Chikowo et al., 2003). Several authors have also reported similar cumulative SOC stocks  
585 for soil profile depth > 30 cm between different tillage and residue management practices  
586 (Angers et al., 1997; Doran et al., 1998; Lal, 2015; Powlson et al., 2014). This can be alluded  
587 to an accumulation of uncertainty when cumulating SOC stocks in several soil layers with their

588 respective error of measurement. This weakens the power of detecting statistically significant  
589 differences even where such differences exist (Kravchenko and Robertson, 2011). Kravchenko  
590 and Robertson, (2011) bemoaned the lack of enough replication when sampling deep soil  
591 horizons to reduce variability and the importance of post hoc power analysis to reduce Type II  
592 error. This study was limited to four replicates which might not have enough statistical power  
593 to detect significant differences between treatments when considering the whole soil profile.

Mis en forme : Police :Non Gras

#### 594 4.3.1 Mulching

595 The overarching role of mulching in cumulative SOC stocks and accumulation / loss rates at  
596 both sites (Tables 1 and 2), albeit, in the top soil (< 30 cm) has been shown in this study.  
597 Cumulative SOC stocks (Table 1) and SOC accumulation / loss rates (Table 2) did not differ  
598 with residue management under NT systems (NTM, NTMR) in the top soil at DTC regardless  
599 of high external OC inputs through maize residue application in mulch treatments (Figure 4,  
600 Table S4). This was attributed to low clay content (< 15 % clay) in the top 20 cm hence low  
601 physical SOC protection (Chivenge et al., 2007; Mtambanengwe et al., 2004; Sun et al., 2020).  
602 such that the differences in OC inputs had little effect. Alternatively, SOC can be protected  
603 from mineralization through adsorption to clay particles (Han et al., 2016; Churchman et al.,  
604 2020). However, there was low surface area for SOC adsorption due to low clay content in the  
605 top soil at DTC. Conversely, maize residue mulching effects were significant at UZF though  
606 NTMR was indifferent when compared to the rest of the NT treatments. Cumulative OC inputs  
607 in NTMR (12.4 Mg C ha<sup>-1</sup>) were about 77 % of cumulative OC inputs in NTM (16.2 Mg C ha<sup>-</sup>  
608 <sup>1</sup>) but at least 57 % higher OC inputs than NT and NTR after 8 seasons (Figure 4). This was  
609 alluded to SOC adsorption and physical protection due to higher clay content at UZF.  
610 Nonetheless, several studies have shown that aboveground biomass is less effective in  
611 sustaining SOC stocks compared to belowground biomass (Hirte et al., 2018, 2021; Jones et

612 al., 2009; Villarino et al., 2021) and we attribute that to the insignificant cumulative SOC stocks  
613 and accumulation rates between the NT treatments other than NTM regardless of higher  
614 aboveground biomass in NTR and NMTR than NT (Figure 4).

#### 615 4.3.2 Tillage

616 Significant changes in SOC stocks and accumulation rates were restricted to the top 20 cm at  
617 both sites below which there were no differences between treatments (Table 2). The  
618 consistently low SOC stocks under NT, CT and CTR and SOC accumulation rates (CTR and  
619 NT) at both sites was attributed to low OC inputs (Table S4, Figure 4). NT and CT had  
620 generally low yields in terms of grain and vegetative aboveground biomass (Mhlanga et al.,  
621 2021; Shumba et al., 2023b) since the establishment of the experiment in the 2013/14 season  
622 and hence low OC inputs through stubble, root mortality and root exudates. Our results  
623 dovetails with studies done elsewhere (Du et al., 2017; Koga and Tsuji, 2009) and meta-  
624 analyses and reviews (Corbeels et al., 2020a; Lal, 2018, 2015) where the authors found that  
625 NT alone does not significantly improve SOC. However, higher SOC stocks were observed  
626 when NT was combined with at least two CA principles (mulching and rotation) at DTC in the  
627 top 20 cm (Table 1). It has been reported that NT cropping systems ~~does not necessarily add~~  
628 ~~SOC but their contribution to enhance~~ SOC accumulation ~~is largely accomplished by~~ through  
629 increasing C inputs in the top layers and reducing erosion through minimum soil disturbance  
630 (Six et al., 2000; Lal, 2015, 2018; Bai et al., 2019; Cai et al., 2022). Minimum soil disturbance  
631 through NT also physically protects SOC in microaggregates from exposure to oxidative losses  
632 (Shumba et al., 2020; Six et al., 2002; Dolan et al., 2006; Liang et al., 2020). ~~Thus~~ However,  
633 NT without mulch is a nonentity compared to other combinations of CA principles for long-  
634 term sustainability in cropping systems (Nyamangara et al., 2013; Kodzwa et al., 2020;  
635 Mhlanga et al., 2021; Li et al., 2020; Bohoussou et al., 2022) and NT is only effective in

636 increasing SOC stocks when it is associated with other CA principles, especially mulch. On  
637 the other hand, our study suggest that NT can achieve the same results of SOC storage as NTR  
638 and NTMR since they had similar SOC stocks at UZF. This can be explained by the low  
639 aboveground OC inputs in rotation treatments during the season when cowpeas were grown.

640 ~~Therefore, the differences in the response to SOC changes and accumulation rates to the~~  
641 ~~treatments in this study, where NT and CTR had the lowest SOC accumulation rates at DTC~~  
642 ~~and UZF respectively, suggests that SOC storage is, as expected, site specific.~~

#### 643 4.3.3 Maize-cowpea rotation

644 Legume rotations have been found to improve SOC accumulation rates and subsequent soil  
645 structural improvement (aggregation) induced by the addition of organic residues with  
646 favourable C/N ratio (Virk et al., 2022; Laub et al., 2023; Jephita et al., 2023). However, in our  
647 study, cowpea rotation benefits on SOC accumulation rates were not ~~observed in comparison~~  
648 ~~to monocropping under NTM~~ significant at DTC. Maize-cowpea rotation had no significant  
649 effects on maize yield (Shumba et al., 2023b; Mhlanga et al., 2021) which corresponded to low  
650 belowground biomass as well. Instead, maize stover mulching improved maize yields at DTC.  
651 Nevertheless, benefits from cowpea rotation under NT cropping systems (NTR, NTMR)  
652 compared to CT cropping systems (CTR) were significant, albeit only in the top 10 cm, at UZF;  
653 CTR had a net loss of SOC ( $-0.07 \pm 0.04$  to  $0.03 \pm 0.03$  Mg C ha<sup>-1</sup> yr<sup>-1</sup>). Significantly higher  
654 maize yields in rotation treatments were observed at UZF (Shumba et al., 2023b; Mhlanga et  
655 al., 2021) and were attributed to more soil mineral N due to biological nitrogen fixation from  
656 the preceding cowpeas. Higher aboveground biomass is positively related to below ground  
657 biomass resulting in significant belowground OC inputs, of higher quality in the rotation  
658 treatments in the season when maize is grown. However, In addition to low OC inputs (Figure  
659 4), the net SOC loss in CTR at UZF was due to seasonal exposure to oxidative losses (SOC

660 mineralization) through disruption of soil macroaggregates by tillage as alluded by Bai et al.,  
661 (2019); Cambardella and Elliott, (1993) and Lal, (2018). We underscore that maize-cowpea  
662 rotation under NT improved SOC accumulation in the top soil due to reduced soil disturbance  
663 and alternate OC inputs of high (cowpeas) and low quality (maize). High quality OC inputs  
664 have a positive priming effect (Chen et al., 2014) which have been shown to be preferentially  
665 stabilized in the soil due to a higher carbon use efficiency of soil microbes (Cotrufo et al., 2013;  
666 Kopittke et al., 2018). This explains significant improvement in SOC stocks under the  
667 combination of NT and alternate high- and low-quality OC inputs (maize-cowpea rotation) to  
668 the soil in medium to heavy textured soils at UZF and vice versa at DTC.

669 ~~Overall, the insignificant differences in cumulative SOC stocks and SOC accumulation / loss~~  
670 ~~rates between the different cropping systems in this and other studies (Laura et al., 2022; Leal~~  
671 ~~et al., 2020) when considering whole soil profile (0-100 cm) is alluded to the dilution effect~~  
672 ~~due to low OC inputs beyond 30 cm. Organic C inputs in the subsoil (> 30 cm) through crop~~  
673 ~~root mortality and root exudates are highly reduced due to low root biomass (Button et al.,~~  
674 ~~2022; Chikowo et al., 2003). Several authors have also reported similar cumulative SOC stocks~~  
675 ~~for soil profile depth > 30 cm between different tillage and residue management practices~~  
676 ~~(Angers et al., 1997; Doran et al., 1998; Lal, 2015; Powlson et al., 2014). This can be alluded~~  
677 ~~to an accumulation of uncertainty when cumulating SOC stocks in several soil layers with their~~  
678 ~~respective error of measurement., which complicates the task~~This weakens the power of  
679 ~~detecting statistically significant differences even where such differences exist (Kravchenko~~  
680 ~~and Robertson, 2011). Kravchenko and Robertson, (2011) bemoaned the lack of enough~~  
681 ~~replication when sampling deep soil horizons to reduce variability and the importance of post~~  
682 ~~hoc power analysis to reduce Type II error.~~ This study was limited to four replicates which  
683 might not have enough statistical power to detect statistically significant differences between  
684 treatments.

685

#### 686 **4.3 Role of soil texture in SOC accumulation**

687 Soil texture is widely recognized to influence SOC stocks (Sun et al., 2020) through physical  
688 and chemical protection of SOC against microbially mediated decomposition (Chivenge et al.,  
689 2007; Mtambanengwe et al., 2004). In our study, the main difference between the two study  
690 sites is soil texture in the top soil (0–30 cm), where DTC had light textured (sandy loams) soils  
691 and UZF had medium to heavy textured (sandy clay loams) soils (Figure 1). These soil textural  
692 differences explain why there were no differences in SOC stocks, changes and accumulation  
693 rates between NTM, NTR and NTMR at DTC regardless of higher cumulative OC inputs in  
694 NTM and NTMR (Figure 4). The direct SOC inputs in the top soil, where SOC was more  
695 concentrated (Table S2, Figure 3) was subject to mineralization because of low clay content  
696 and thus low protection by soil micro-aggregates (Chivenge et al., 2007; Mtambanengwe et al.,  
697 2004; Sun et al., 2020), such that the differences in OC inputs had little effect. Light textured  
698 soils have large pores which cannot protect SOC against microbial decomposition  
699 (Mtambanengwe et al., 2004; Christensen, 1987; Sun et al., 2020; Kravchenko and Guber,  
700 2017). In contrast, there were differences between NTM and NTR at UZF in the top soil layers  
701 and intermediate between NTM and NTMR. Cumulative OC inputs in NTMR (12.4 Mg C ha<sup>-1</sup>)  
702 were about 75 % of cumulative OC inputs in NTM (16.2 Mg C ha<sup>-1</sup>) (Figure 4) after 8  
703 seasons. The added C, especially from maize stover mulch, most likely was protected by clay  
704 particles as well as formation of organo-mineral complexes (Malepfane et al., 2022; Chivenge  
705 et al., 2007; Jephita et al., 2023) which protects SOC from mineralization (Dunjana et al., 2012;  
706 Shumba et al., 2020; Button et al., 2022; Rumpel et al., 2012; Sanauallah et al., 2016).

707

#### 708 **4.4 Soil CO<sub>2</sub> fluxes and cumulative CO<sub>2</sub> emissions**

709 ~~Despite the higher OC inputs in NT cropping systems with mulch, CO<sub>2</sub> fluxes and cumulative~~  
710 ~~emissions were similar between treatments. This is attributed to the fact that the CO<sub>2</sub> fluxes in~~  
711 ~~this study were the sum of autotrophic and heterotrophic respiration (Heinemeyer et al., 2007);~~  
712 ~~hence possible treatment effects on heterotrophic respiration were most likely masked. Maize~~  
713 ~~root respiration (autotrophic respiration) has been shown to contribute an average of about 45~~  
714 ~~% (Hao and Jiang, 2014) to total soil respiration (heterotrophic and autotrophic respiration). In~~  
715 ~~contrast to other studies (Carbonell-Bojollo et al., 2019; Chatskikh and Olesen, 2007;~~  
716 ~~McDonald et al., 2019; O'Dell et al., 2020), no higher fluxes and emissions were observed~~  
717 ~~following top soil disturbance in the CT treatments (Figures 5 and 6). This was attributed to~~  
718 ~~low SOC stocks in CT treatments in the top 15 cm which was the plough depth in this study.~~

719

## 720 **5. Conclusions**

721 Our study has shown the overarching importance of mulching and of combining at least two  
722 CA principles to improve top SOC stocks. ~~Mulching under no tillage system (NTM) improves~~  
723 ~~SOC stocks in the top soil though the same can be achieved by the full CA (NTMR), or no~~  
724 ~~tillage plus rotation (NTR) cropping system on a sandy soil. The absence of No tillage (NT)~~  
725 ~~alone (NT) could not increase SOC stocks, and even lead led to a slight decrease compared to~~  
726 CT, due to lower crop productivity in NT and therefore reduced OC inputs to the soil.  
727 Nevertheless, whole profile (0-100 cm) SOC stocks ~~waswere~~ the same between all the  
728 treatment. Our study also showed that sampling the entire soil profile is necessary for a more  
729 accurate view of SOC accumulation potential among different cropping systems.

## 730 **6. Data availability**

731 All data are freely available on the CIRAD data repository  
732 <https://doi.org/10.18167/DVN1/VPOCHN> (Shumba et al., 2023a).

Mis en forme : Police :Gras

733

734 **6.7. Author contributions**

735 CT designed, established and maintained the experiments since 2013;

736 AS, RCa, RCh were involved in ~~various gas and~~ soil sampling campaigns; ~~JS was involved in~~  
737 ~~laboratory analysis of gas samples.~~

738 AS, RCa performed the statistical analyses, graphics and drafting the manuscript;

739 AS, RCa, RCh, MC, JS, CT reviewed and edited the manuscript.

740

741 **7.8. Competing interests**

742 One of the co-authors is a member of the editorial board of the SOIL journal. The other authors  
743 have no competing interests to declare.

744

745 **8.9. Acknowledgements**

746 This study was funded by the DSCATT project “Agricultural Intensification and Dynamics of  
747 Soil Carbon Sequestration in Tropical and Temperate Farming Systems” (N<sup>o</sup> AF 1802-001,  
748 N<sup>o</sup> FT C002181), supported by the Agropolis Foundation (“Programme d’Investissement  
749 d’Avenir” Labex Agro, ANR-10-LABX- 0001-01) and by the TOTAL Foundation within a  
750 patronage agreement. Authors are grateful to the International Maize and Wheat Improvement  
751 Center (CIMMYT) for the setup and running of the experiment. We also acknowledge the  
752 donors of the MAIZE CGIAR Research Program ([www.maize.org](http://www.maize.org)) and the Ukama Ustawi  
753 Regional CGIAR Initiative who supported the trials up to 2018 and staff time until 2023. We  
754 thank Britta Jahn-Humphrey for carrying out the gas analyses at ETH Zürich. We also thank  
755 Admire Muwati for his help in gas sampling. Special thanks go to the technical personnel at

Mis en forme : Police : (Par défaut) Times New Roman,  
12 pt

Mis en forme : Paragraphe de liste



756 each experimental locations namely Tarirai Muoni, Sign Phiri, Herbert Chipara and Connie  
757 Madembo who continuously assisted in trial establishment and management.

758

759 **9.10. References**

760 Amos, B. and Walters, D. T.: Maize Root Biomass and Net Rhizodeposited Carbon, *Soil Sci.*  
761 *Soc. Am. J.*, 70, 1489–1503, <https://doi.org/10.2136/sssaj2005.0216>, 2006.

762 Angers, D. A. and Eriksen-Hamel, N. S.: Full-inversion tillage and organic carbon distribution  
763 in soil profiles: A meta-analysis, *Soil Sci. Soc. Am. J.*, 72, 1370–1374,  
764 <https://doi.org/10.2136/sssaj2007.0342>, 2008.

765 Angers, D. A., Bolinder, M. A., Carter, M. R., Gregorich, E. G., Drury, C. F., Liang, B. C.,  
766 Voroney, R. P., Simard, R. R., Donald, R. G., Bevaert, R. P., and Martel, J.: Impact of tillage  
767 practices on organic carbon and nitrogen storage in cool, humid soils of eastern Canada, *Soil*  
768 *Tillage Res.*, 41, 191–201, <https://doi.org/10.4141/S96-111>, 1997.

769 Bai, X., Huang, Y., Ren, W., Coyne, M., Jacinthe, P.-A., Bo, T., Hui, D., Yang, J., and  
770 Matocha, C.: Responses of soil carbon sequestration to climate-smart agriculture practices: A  
771 meta-analysis, 25, 2591–2606, <https://doi.org/10.1111/gcb.14658>, 2019.

772 Balesdent, J., Derrien, D., Fontaine, S., Kirman, S., Klumpp, K., Loiseau, P., Marol, C.,  
773 Nguyen, C., Pean, M., Personeni, E., and Robin, C.: Contribution de la rhizodéposition aux  
774 matières organiques du sol, quelques implications pour la modélisation de la dynamique du  
775 carbone, *Etude Gest. des sols*, 18 (3), 201–216, 2011.

776 Balesdent, J., Basile-Doelsch, I., Chadoeuf, J., Cornu, S., Derrien, D., Fekiacova, Z., and Hatté,  
777 C.: Atmosphere–soil carbon transfer as a function of soil depth, *Nature*, 559, 599–602,  
778 <https://doi.org/10.1038/s41586-018-0328-3>, 2018.

779 Bationo, A., Kihara, J., Vanlauwe, B., Waswa, B., Adolwa, I., and Saidou, K.: Overview of  
780 Long Term Experiments in Africa, in: *Lessons Learned from Long-Term Soil Fertility*  
781 *Management Experiments in Africa.*, Springer Science+Business Media Dordrecht, 1–26,  
782 <https://doi.org/10.1007/978-94-007-2938-4>, 2013.

783 Baudron, F., Andersson, J. A., Corbeels, M., and Giller, K. E.: Failing to Yield? Ploughs,  
784 Conservation Agriculture and the Problem of Agricultural Intensification: An Example from  
785 the Zambezi Valley, Zimbabwe, *J. Dev. Stud.*, 48, 393–412,  
786 <https://doi.org/10.1080/00220388.2011.587509>, 2012.

787 Bohoussou, Y. N. D., Kou, Y. H., Yu, W. B., Lin, B. jian, Virk, A. L., Zhao, X., Dang, Y. P.,  
788 and Zhang, H. L.: Impacts of the components of conservation agriculture on soil organic carbon  
789 and total nitrogen storage: A global meta-analysis, *Sci. Total Environ.*, 842, 156822,  
790 <https://doi.org/10.1016/J.SCITOTENV.2022.156822>, 2022.

791 Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H.,  
792 and White, J. S. S.: Generalized linear mixed models: a practical guide for ecology and  
793 evolution, *Trends Ecol. Evol.*, 24, 127–135, <https://doi.org/10.1016/j.tree.2008.10.008>, 2009.

- 794 Bossio, D. A., Cook-Patton, S. C., Ellis, P. W., Fargione, J., Sanderman, J., Smith, P., Wood,  
795 S., Zomer, R. J., von Unger, M., Emmer, I. M., and Griscom, B. W.: The role of soil carbon in  
796 natural climate solutions, *Nat. Sustain.*, 3, 391–398, [https://doi.org/10.1038/s41893-020-0491-](https://doi.org/10.1038/s41893-020-0491-z)  
797 [z](https://doi.org/10.1038/s41893-020-0491-z), 2020.
- 798 Button, E. S., Pett-Ridge, J., Murphy, D. V., Kuzyakov, Y., Chadwick, D. R., and Jones, D. L.:  
799 Deep-C storage: Biological, chemical and physical strategies to enhance carbon stocks in  
800 agricultural subsoils, *Soil Biol. Biochem.*, 170, 108697,  
801 <https://doi.org/10.1016/j.soilbio.2022.108697>, 2022.
- 802 Cai, A., Han, T., Ren, T., Sanderman, J., Rui, Y., Wang, B., Smith, P., Xu, M., and Yu'e Li:  
803 Declines in soil carbon storage under no tillage can be alleviated in the long run, 425, 1–3,  
804 <https://doi.org/10.1016/j.geoderma.2022.116028>, 2022.
- 805 Cambardella, C. A. and Elliott, E. T.: Carbon and Nitrogen Distribution in Aggregates from  
806 Cultivated and Native Grassland Soils, *Soil Sci. Soc. Am. J.*, 57, 1071–1076,  
807 <https://doi.org/10.2136/SSSAJ1993.03615995005700040032X>, 1993.
- 808 Cardinael, R., Chevallier, T., Barthès, B. G., Saby, N. P. A., Parent, T., Dupraz, C., Bernoux,  
809 M., and Chenu, C.: Impact of alley cropping agroforestry on stocks, forms and spatial  
810 distribution of soil organic carbon - A case study in a Mediterranean context, *Geoderma*, 259–  
811 260, 288–299, <https://doi.org/10.1016/j.geoderma.2015.06.015>, 2015.
- 812 Cardinael, R., Guibert, H., Kouassi Brédoumy, S. T., Gigou, J., N’Goran, K. E., and Corbeels,  
813 M.: Sustaining maize yields and soil carbon following land clearing in the forest–savannah  
814 transition zone of West Africa: Results from a 20-year experiment, *F. Crop. Res.*, 275,  
815 <https://doi.org/10.1016/j.fcr.2021.108335>, 2022.
- 816 Cheesman, S., Thierfelder, C., Eash, N. S., Kassie, G. T., and Frossard, E.: Soil carbon stocks  
817 in conservation agriculture systems of Southern Africa, *Soil Tillage Res.*, 156, 99–109,  
818 <https://doi.org/10.1016/j.still.2015.09.018>, 2016.
- 819 Chen, R., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Lin, X., Blagodatskaya,  
820 E., and Kuzyakov, Y.: Soil C and N availability determine the priming effect: Microbial N  
821 mining and stoichiometric decomposition theories, *Glob. Chang. Biol.*, 20, 2356–2367,  
822 <https://doi.org/10.1111/gcb.12475>, 2014.
- 823 Chikowo, R., Mapfumo, P., Nyamugafata, P., Nyamadzawo, G., and Giller, K. E.: Nitrate-N  
824 dynamics following improved fallows and maize root development in a Zimbabwean sandy  
825 clay loam, *Agrofor. Syst.*, 59, 187–195,  
826 <https://doi.org/10.1023/B:AGFO.0000005219.07409.a0>, 2003.
- 827 Chivenge, P. P., Murwira, H. K., Giller, K. E., Mapfumo, P., and Six, J.: Long-term impact of  
828 reduced tillage and residue management on soil carbon stabilization: Implications for  
829 conservation agriculture on contrasting soils, *Soil Tillage Res.*, 94, 328–337,  
830 <https://doi.org/10.1016/j.still.2006.08.006>, 2007.
- 831 Christensen, B. T.: Decomposability of organic matter in particle size fractions from field soils  
832 with straw incorporation, *Soil Biol. Biochem.*, 19, 429–435, [https://doi.org/10.1016/0038-](https://doi.org/10.1016/0038-0717(87)90034-4)  
833 [0717\(87\)90034-4](https://doi.org/10.1016/0038-0717(87)90034-4), 1987.
- 834 Churchman, G. J., Singh, M., Schapel, A., Sarkar, B., and Bolan, N.: Clay Minerals As the Key  
835 To the Sequestration of Carbon in Soils, *Clays Clay Miner.*, 68, 135–143,

836 <https://doi.org/10.1007/s42860-020-00071-z>, 2020.

837 Corbeels, M., Cardinael, R., Powlson, D., Chikowo, R., and Gerard, B.: Carbon sequestration  
838 potential through conservation agriculture in Africa has been largely overestimated: Comment  
839 on: “Meta-analysis on carbon sequestration through conservation agriculture in Africa,” *Soil*  
840 *Tillage Res.*, 196, 104300, <https://doi.org/10.1016/j.still.2019.104300>, 2020a.

841 Corbeels, M., Naudin, K., Whitbread, A. M., Kühne, R., and Letourmy, P.: Limits of  
842 conservation agriculture to overcome low crop yields in sub-Saharan Africa, *Nat. Food*, 1, 447–  
843 454, <https://doi.org/10.1038/s43016-020-0114-x>, 2020b.

844 Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Deneff, K., and Paul, E.: The Microbial  
845 Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with  
846 soil organic matter stabilization: Do labile plant inputs form stable soil organic matter?, *Glob.*  
847 *Chang. Biol.*, 19, 988–995, <https://doi.org/10.1111/gcb.12113>, 2013.

848 Dignac, M. F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T.,  
849 Freschet, G. T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron, P. A.,  
850 Nunan, N., Roumet, C., and Basile-Doelsch, I.: Increasing soil carbon storage: mechanisms,  
851 effects of agricultural practices and proxies. A review, *Agron. Sustain. Dev.*, 37,  
852 <https://doi.org/10.1007/s13593-017-0421-2>, 2017.

853 Dolan, M. S., Clapp, C. E., Allmaras, R. R., Baker, J. M., and Molina, J. A. E.: Soil organic  
854 carbon and nitrogen in a Minnesota soil as related to tillage, residue and nitrogen management,  
855 *Soil Tillage Res.*, 89, 221–231, <https://doi.org/10.1016/j.still.2005.07.015>, 2006.

856 Doran, J. W., Elliott, E. T., and Paustian, K.: Soil microbial activity, nitrogen cycling, and long-  
857 term changes in organic carbon pools as related to fallow tillage management, *Soil Tillage*  
858 *Res.*, 49, 3–18, [https://doi.org/10.1016/S0167-1987\(98\)00150-0](https://doi.org/10.1016/S0167-1987(98)00150-0), 1998.

859 Du, Z., Angers, D. A., Ren, T., Zhang, Q., and Li, G.: The effect of no-till on organic C storage  
860 in Chinese soils should not be overemphasized: A meta-analysis, *Agric. Ecosyst. Environ.*,  
861 236, 1–11, <https://doi.org/10.1016/j.agee.2016.11.007>, 2017.

862 Dube, E., Chiduza, C., and Muchaonyerwa, P.: Conservation agriculture effects on soil organic  
863 matter on a Haplic Cambisol after four years of maize-oat and maize-grazing vetch rotations  
864 in South Africa, *Soil Tillage Res.*, 123, 21–28, <https://doi.org/10.1016/j.still.2012.02.008>,  
865 2012.

866 Dunjana, N., Nyamugafata, P., Shumba, A., Nyamangara, J., and Zingore, S.: Effects of cattle  
867 manure on selected soil physical properties of smallholder farms on two soils of Murewa,  
868 Zimbabwe, *Soil Use Manag.*, 28, <https://doi.org/10.1111/j.1475-2743.2012.00394.x>, 2012.

869 Ellert, B. H. and Bettany, J. R.: Calculation of organic matter and nutrients stored in soils under  
870 contrasting management regimes, *Can. J. Soil Sci.*, 75, 529–538,  
871 <https://doi.org/10.4141/cjss95-075>, 1995.

872 Feller, C. and Beare, M. H.: Physical control of soil organic matter dynamics in the tropics,  
873 *Geoderma*, 79, 69–116, [https://doi.org/10.1016/S0016-7061\(97\)00039-6](https://doi.org/10.1016/S0016-7061(97)00039-6), 1997.

874 Giller, K. E., Witter, E., Corbeels, M., and Tittonell, P.: Conservation agriculture and  
875 smallholder farming in Africa: The heretics’ view, *F. Crop. Res.*, 114, 23–34,  
876 <https://doi.org/10.1016/j.fcr.2009.06.017>, 2009.

877 Giller, K. E., Andersson, J. A., Corbeels, M., Kirkegaard, J., Mortensen, D., Erenstein, O., and  
878 Vanlauwe, B.: Beyond conservation agriculture, *Front. Plant Sci.*, 6, 1–14,  
879 <https://doi.org/10.3389/fpls.2015.00870>, 2015.

880 von Haden, A. C., Yang, W. H., and DeLucia, E. H.: Soils' dirty little secret: Depth-based  
881 comparisons can be inadequate for quantifying changes in soil organic carbon and other  
882 mineral soil properties, *Glob. Chang. Biol.*, 26, 3759–3770, <https://doi.org/10.1111/gcb.15124>,  
883 2020.

884 Han, L., Sun, K., Jin, J., and Xing, B.: Some concepts of soil organic carbon characteristics and  
885 mineral interaction from a review of literature, *Soil Biol. Biochem.*, 94, 107–121,  
886 <https://doi.org/10.1016/j.soilbio.2015.11.023>, 2016.

887 Harrison, R. B., Footen, P. W., Harrison, R. B., Footen, P. W., and Strahm, B. D.: Deep Soil  
888 Horizons: Contribution and Importance to Soil Carbon Pools and in Assessing Whole-  
889 Ecosystem Response to Management and Global Change, *For. Sci.*, 57, 67–76, 2011.

890 Hirte, J., Leifeld, J., Abiven, S., Oberholzer, H. R., and Mayer, J.: Below ground carbon inputs  
891 to soil via root biomass and rhizodeposition of field-grown maize and wheat at harvest are  
892 independent of net primary productivity, *Agric. Ecosyst. Environ.*, 265, 556–566,  
893 <https://doi.org/10.1016/j.agee.2018.07.010>, 2018.

894 Hirte, J., Walder, F., Hess, J., Büchi, L., Colombi, T., van der Heijden, M. G., and Mayer, J.:  
895 Enhanced root carbon allocation through organic farming is restricted to topsoils, *Sci. Total  
896 Environ.*, 755, 143551, <https://doi.org/10.1016/j.scitotenv.2020.143551>, 2021.

897 IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas  
898 Inventories., 1–19 pp., 2019.

899 Jephitha, G., Jefline, K., Willis, G., and Justice, N.: Carbon stock, aggregate stability and  
900 hydraulic properties of soils under tillage, crop rotation and mineral fertiliser application in  
901 sub-humid Zimbabwe, *Heliyon*, 9, <https://doi.org/10.1016/j.heliyon.2023.e15846>, 2023.

902 Jones, D. L., Nguyen, C., and Finlay, R. D.: Carbon flow in the rhizosphere : carbon trading at  
903 the soil – root interface, *Plant Soil*, 321, 5–33, <https://doi.org/10.1007/s11104-009-9925-0>,  
904 2009.

905 Kahn, B. A. and Schroeder, J. L.: Root Characteristics and Seed Yields of Cowpeas Grown  
906 with and without Added Nitrogen Fertilizer, *Hortscience A Publ. Am. Soc. Horticultural Sci.*,  
907 34, 1238–1239, <https://doi.org/10.21273/hortsci.34.7.1238>, 1999.

908 Kassam, A., Friedrich, T., and Derpsch, R.: Global spread of Conservation Agriculture, *Int. J.  
909 Environ. Stud.*, 76, 29–51, <https://doi.org/10.1080/00207233.2018.1494927>, 2019.

910 Kell, D. B.: Breeding crop plants with deep roots: their role in sustainable carbon , nutrient and  
911 water sequestration, *Ann. Bot.*, 108, 407–418, <https://doi.org/10.1093/aob/mcr175>, 2011.

912 Kimaro, A. A., Mpanda, M., Rioux, J., Aynekulu, E., Shaba, S., Thiong'o, M., Mutuo, P.,  
913 Abwanda, S., Shepherd, K., Neufeldt, H., and Rosenstock, T. S.: Is conservation agriculture  
914 'climate-smart' for maize farmers in the highlands of Tanzania?, *Nutr. Cycl. Agroecosystems*,  
915 105, 217–228, <https://doi.org/10.1007/s10705-015-9711-8>, 2016.

916 Kimiti, J. M.: Influence of integrated soil nutrient management on cowpea root growth in the  
917 semi-arid Eastern Kenya, *African J. Agric. Res.*, 6, 3084–3091,

918 <https://doi.org/10.5897/AJAR10.1023>, 2011.

919 Kodzwa, J. J., Gotosa, J., and Nyamangara, J.: Mulching is the most important of the three  
920 conservation agriculture principles in increasing crop yield in the short term, under sub humid  
921 tropical conditions in Zimbabwe, *Soil Tillage Res.*, 197, 104515,  
922 <https://doi.org/10.1016/j.still.2019.104515>, 2020.

923 Koga, N. and Tsuji, H.: Effects of reduced tillage, crop residue management and manure  
924 application practices on crop yields and soil carbon sequestration on an Andisol in northern  
925 Japan, *Soil Sci. Plant Nutr.*, 55, 546–557, <https://doi.org/10.1111/j.1747-0765.2009.00385.x>,  
926 2009.

927 Kopittke, P. M., Hernandez-Soriano, M. C., Dalal, R. C., Finn, D., Menzies, N. W., Hoeschen,  
928 C., and Mueller, C. W.: Nitrogen-rich microbial products provide new organo-mineral  
929 associations for the stabilization of soil organic matter, *Glob. Chang. Biol.*, 24, 1762–1770,  
930 <https://doi.org/10.1111/gcb.14009>, 2018.

931 Kravchenko, A. N. and Guber, A. K.: Soil pores and their contributions to soil carbon  
932 processes, *Geoderma*, 287, 31–39, <https://doi.org/10.1016/j.geoderma.2016.06.027>, 2017.

933 Kravchenko, A. N. and Robertson, G. P.: Whole-Profile Soil Carbon Stocks: The Danger of  
934 Assuming Too Much from Analyses of Too Little, *Soil Sci. Soc. Am. J.*, 75, 235–240,  
935 <https://doi.org/10.2136/sssaj2010.0076>, 2011.

936 Lal, R.: Residue management, conservation tillage and soil restoration for mitigating  
937 greenhouse effect by CO<sub>2</sub>-enrichment, *Soil Tillage Res.*, 43, 81–107, 1997.

938 Lal, R.: Sequestering carbon and increasing productivity by conservation agriculture, *J. Soil*  
939 *Water Conserv.*, 70, 55A-62A, <https://doi.org/10.2489/jswc.70.3.55A>, 2015.

940 Lal, R.: Digging deeper: A holistic perspective of factors affecting soil organic carbon  
941 sequestration in agroecosystems, *Glob. Chang. Biol.*, 24, 3285–3301,  
942 <https://doi.org/10.1111/gcb.14054>, 2018.

943 Laub, M., Corbeels, M., Couëdel, A., Ndungu, S. M., Mucheru-muna, M. W., Mugendi, D.,  
944 Necpalova, M., Waswa, W., Van de Broek, M., Vanlauwe, B., and Six, J.: Managing soil  
945 organic carbon in tropical agroecosystems: evidence from four long-term experiments in  
946 Kenya, *SOIL*, 9, 301–323, <https://doi.org/10.5194/soil-9-301-2023>, 2023.

947 Laura, van der P. K., Andy, R., Meagan, S., Francisco, C. J., Wallenstein, M. D., and Cotrufo,  
948 M. F.: Addressing the soil carbon dilemma: Legumes in intensified rotations regenerate soil  
949 carbon while maintaining yields in semi-arid dryland wheat farms., *Agric. Ecosyst. Environ.*,  
950 330, <https://doi.org/10.1016/j.agee.2022.107906>, 2022.

951 Leal, O. A., Amado, T. J. C., Fiorin, J. E., Keller, C., Reimche, G. B., Rice, C. W., Nicoloso,  
952 R. S., Bortolotto, R. P., and Schwalbert, R.: Linking cover crop residue quality and tillage  
953 system to  $\text{CO}_2\text{-e-C}$  emission, soil c and n stocks and crop yield based on a long-term  
954 experiment, *Agronomy*, 10, <https://doi.org/10.3390/agronomy10121848>, 2020.

955 Li, Y., Li, Z., Chang, S. X., Cui, S., Jagadamma, S., Zhang, Q., and Cai, Y.: Residue retention  
956 promotes soil carbon accumulation in minimum tillage systems: Implications for conservation  
957 agriculture, *Sci. Total Environ.*, <https://doi.org/10.1016/j.scitotenv.2020.140147>, 2020.

958 Liang, B. C., VandenBygaart, A. J., MacDonald, J. D., Cerkowniak, D., McConkey, B. G.,

959 Desjardins, R. L., and Angers, D. A.: Revisiting no-till's impact on soil organic carbon storage  
960 in Canada, *Soil Tillage Res.*, 198, 104529, <https://doi.org/10.1016/j.still.2019.104529>, 2020.

961 Lorenz, K. and Lal, R.: The Depth Distribution of Soil Organic Carbon in Relation to Land  
962 Use and Management and the Potential of Carbon Sequestration in Subsoil Horizons, *Adv.*  
963 *Agron.*, 88, 35–66, [https://doi.org/10.1016/S0065-2113\(05\)88002-2](https://doi.org/10.1016/S0065-2113(05)88002-2), 2005.

964 Ma, S., He, F., Tian, D., Zou, D., Yan, Z., Yang, Y., Zhou, T., Huang, K., Shen, H., and Fang,  
965 J.: Variations and determinants of carbon content in plants: A global synthesis, *Biogeosciences*,  
966 15, 693–702, <https://doi.org/10.5194/bg-15-693-2018>, 2018.

967 Malepfane, N. M., Muchaonyerwa, P., Hughes, J. C., and Zengeni, R.: Land use and site effects  
968 on the distribution of carbon in some humic soil profiles of KwaZulu-Natal, South Africa,  
969 *Heliyon*, 8, <https://doi.org/10.1016/j.heliyon.2021.e08709>, 2022.

970 Mapanda, F., Mupini, J., Wuta, M., Nyamangara, J., and Rees, R. M.: A cross-ecosystem  
971 assessment of the effects of land cover and land use on soil emission of selected greenhouse  
972 gases and related soil properties in Zimbabwe, *Eur. J. Soil Sci.*, 61, 721–733,  
973 <https://doi.org/10.1111/j.1365-2389.2010.01266.x>, 2010.

974 Mbanyele, V., Mtambanengwe, F., Nezomba, H., Groot, J. C. J., and Mapfumo, P.:  
975 Comparative short-term performance of soil water management options for increased  
976 productivity of maize-cowpea intercropping in semi-arid Zimbabwe, *J. Agric. Food Res.*, 5,  
977 100189, <https://doi.org/10.1016/j.jafr.2021.100189>, 2021.

978 Mhlanga, B., Ercoli, L., Pellegrino, E., Onofri, A., and Thierfelder, C.: The crucial role of  
979 mulch to enhance the stability and resilience of cropping systems in southern Africa, *Agron.*  
980 *Sustain. Dev.*, 41, <https://doi.org/10.1007/s13593-021-00687-y>, 2021.

981 Mhlanga, B., Pellegrino, E., Thierfelder, C., and Ercoli, L.: Conservation agriculture practices  
982 drive maize yield by regulating soil nutrient availability, arbuscular mycorrhizas, and plant  
983 nutrient uptake, *F. Crop. Res.*, 277, 108403, <https://doi.org/10.1016/j.fcr.2021.108403>, 2022a.

984 Mhlanga, B., Ercoli, L., Thierfelder, C., and Pellegrino, E.: Conservation agriculture practices  
985 lead to diverse weed communities and higher maize grain yield in Southern Africa, *F. Crop.*  
986 *Res.*, 289, 108724, <https://doi.org/10.1016/j.fcr.2022.108724>, 2022b.

987 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A.,  
988 Chaplot, V., Chen, Z. S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong,  
989 S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke,  
990 S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I.,  
991 Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C. C., Vågen, T. G., van Wesemael, B.,  
992 and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86,  
993 <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.

994 Mtambanengwe, F., Mapfumo, P., and Kirchmann, H.: Decomposition of organic matter in soil  
995 as influenced by texture and pore size distribution, in: *Managing Nutrient Cycles to Sustain*  
996 *Soil Fertility in Sub-Saharan Africa*, edited by: Bationo, A., Academy Science Publishers and  
997 TSBF CIAT, Nairobi, 261–276, 2004.

998 Nyamangara, J., Masvaya, E. N., Tirivavi, R., and Nyengerai, K.: Effect of hand-hoe based  
999 conservation agriculture on soil fertility and maize yield in selected smallholder areas in  
1000 Zimbabwe, *Soil Tillage Res.*, 126, 19–25, <https://doi.org/10.1016/j.still.2012.07.018>, 2013.

- 1001 Patra, S., Julich, S., Feger, K. H., Jat, M. L., Sharma, P. C., and Schwärzel, K.: Effect of  
 1002 conservation agriculture on stratification of soil organic matter under cereal-based cropping  
 1003 systems, *Arch. Agron. Soil Sci.*, 65, 2013–2028,  
 1004 <https://doi.org/10.1080/03650340.2019.1588462>, 2019.
- 1005 Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., and Smith, P.: Climate-smart  
 1006 soils, *Nature*, 532, 49–57, <https://doi.org/10.1038/nature17174>, 2016.
- 1007 Powlson, D. S., Stirling, C. M., Jat, M. L., Gerard, B. G., Palm, C. A., Sanchez, P. A., and  
 1008 Cassman, K. G.: Limited potential of no-till agriculture for climate change mitigation, *Nat.*  
 1009 *Clim. Chang.*, 4, 678–683, <https://doi.org/10.1038/nclimate2292>, 2014.
- 1010 Powlson, D. S., Stirling, C. M., Thierfelder, C., White, R. P., and Jat, M. L.: Does conservation  
 1011 agriculture deliver climate change mitigation through soil carbon sequestration in tropical agro-  
 1012 ecosystems?, *"Agriculture, Ecosyst. Environ.*, 220, 164–174,  
 1013 <https://doi.org/10.1016/j.agee.2016.01.005>, 2016.
- 1014 [R Core Team, 2020. R: A Language and Environment for Statistical Computing. R Foundation  
 1015 for Statistical Computing, Vienna, Austria. https://www.R-project.org/.](https://www.R-project.org/)
- 1016 Rumpel, C., Chabbi, A., and Marschner, B.: Carbon Storage and Sequestration in Subsoil  
 1017 Horizons: Knowledge, Gaps and Potentials, in: *Recarbonization of the Biosphere: Ecosystems  
 1018 and the Global Carbon Cycle*, edited by: Lal, R., Lorenz, K., Hüttl, R. F., Schneider, B. U., and  
 1019 Von Braun, J., Springer Science+Business Media B.V, 445–464, <https://doi.org/10.1007/978-94-007-4159-1>, 2012.
- 1021 Sanaullah, M., Chabbi, A., Maron, P. A., Baumann, K., Tardy, V., Blagodatskaya, E.,  
 1022 Kuzyakov, Y., and Rumpel, C.: How do microbial communities in top- and subsoil respond to  
 1023 root litter addition under field conditions?, *Soil Biol. Biochem.*, 103, 28–38,  
 1024 <https://doi.org/10.1016/j.soilbio.2016.07.017>, 2016.
- 1025 Shumba, A., Dunjana, N., Nyamasoka, B., Nyamugafata, P., Madyiwa, S., and Nyamangara,  
 1026 J.: Maize (*Zea mays*) yield and its relationship to soil properties under integrated fertility,  
 1027 mulch and tillage management in urban agriculture, *South African J. Plant Soil*, 1–10,  
 1028 <https://doi.org/10.1080/02571862.2019.1678686>, 2020.
- 1029 Shumba, A., Chikowo, R., Thierfelder, C., Corbeels, M., Six, J., and Cardinael, R.: [Data for  
 1030 "Mulch application as the overarching factor explaining increase in soil organic carbon stocks  
 1031 under conservation agriculture in two 8-year-old experiments in Zimbabwe".  
 1032 https://doi.org/10.18167/DVNI/VPOCHN, CIRAD Dataverse, V2 Conservation agriculture  
 1033 increases soil organic carbon stocks but not soil CO<sub>2</sub> efflux in two 8-year-old experiments in  
 1034 Zimbabwe, \*EGU sphere\*, 2023, 1–40, <https://doi.org/10.18167/DVNI/VPOCHN>, 2023a.](https://doi.org/10.18167/DVNI/VPOCHN)
- 1035 Shumba, A., Chikowo, R., Corbeels, M., Six, J., Thierfelder, C., and Cardinael, R.: Long-term  
 1036 tillage, residue management and crop rotation impacts on N<sub>2</sub>O and CH<sub>4</sub> emissions on two  
 1037 contrasting soils in sub-humid Zimbabwe, *Agric. Ecosyst. Environ.*, 341,  
 1038 <https://doi.org/10.1016/j.agee.2022.108207>, 2023b.
- 1039 Singh, G., Schoonover, J. E., Williard, K. W. J., Kaur, G., and Crim, J.: Carbon and Nitrogen  
 1040 Pools in Deep Soil Horizons at Different Landscape Positions, *Soil Sci. Soc. Am. J.*, 82, 1512–  
 1041 1525, <https://doi.org/10.2136/sssaj2018.03.0092>, 2018.
- 1042 Six, J., Paustian, K., and Elliott, E.: Aggregate and soil organic matter dynamics under

Mis en forme : Police :Italique

Mis en forme : Indice

Mis en forme : Indice

- 1043 conventional and no-tillage systems, *Soil Sci. Soc. Am. J.*, 63, 1350–1358, 1999.
- 1044 Six, J., Elliott, E. T., and Paustian, K.: Soil macroaggregate turnover and microaggregate  
1045 formation: A mechanism for C sequestration under no-tillage agriculture, *Soil Biol. Biochem.*,  
1046 32, 2099–2103, [https://doi.org/10.1016/S0038-0717\(00\)00179-6](https://doi.org/10.1016/S0038-0717(00)00179-6), 2000.
- 1047 Six, J., Conant, R. T., Paul, E. A., and Paustian, K.: Stabilization mechanisms of soil organic  
1048 matter : Implications for C-saturation of soils, 155–176, 2002.
- 1049 Sun, W., Canadell, J. G., Yu, L. L., Yu, L. L., Zhang, W., Smith, P., Fischer, T., and Huang,  
1050 Y.: Climate drives global soil carbon sequestration and crop yield changes under conservation  
1051 agriculture, *Glob. Chang. Biol.*, 26, 3325–3335, <https://doi.org/10.1111/gcb.15001>, 2020.
- 1052 Swanepoel, C. M., van der Laan, M., Weepener, H. L., du Preez, C. C., and Annandale, J. G.:  
1053 Review and meta-analysis of organic matter in cultivated soils in southern Africa, *Nutr. Cycl.*  
1054 *Agroecosystems*, 104, 107–123, <https://doi.org/10.1007/s10705-016-9763-4>, 2016.
- 1055 Swanepoel, C. M., Rötter, R. P., van der Laan, M., Annandale, J. G., Beukes, D. J., du Preez,  
1056 C. C., Swanepoel, L. H., van der Merwe, A., and Hoffmann, M. P.: The benefits of conservation  
1057 agriculture on soil organic carbon and yield in southern Africa are site-specific, *Soil Tillage*  
1058 *Res.*, 183, 72–82, <https://doi.org/10.1016/j.still.2018.05.016>, 2018.
- 1059 Thierfelder, C. and Mhlanga, B.: Short-term yield gains or long-term sustainability? – a  
1060 synthesis of Conservation Agriculture long-term experiments in Southern Africa, *Agric.*  
1061 *Ecosyst. Environ.*, 326, 107812, <https://doi.org/10.1016/j.agee.2021.107812>, 2022.
- 1062 Thierfelder, C. and Wall, P. C.: Effects of conservation agriculture techniques on infiltration  
1063 and soil water content in Zambia and Zimbabwe, *Soil Tillage Res.*, 105, 217–227,  
1064 <https://doi.org/10.1016/j.still.2009.07.007>, 2009.
- 1065 Thierfelder, C. and Wall, P. C.: Effects of conservation agriculture on soil quality and  
1066 productivity in contrasting agro-ecological environments of Zimbabwe, *Soil Use Manag.*, 28,  
1067 209–220, <https://doi.org/10.1111/j.1475-2743.2012.00406.x>, 2012.
- 1068 Thierfelder, C., Matemba-Mutasa, R., and Rusinamhodzi, L.: Yield response of maize (*Zea*  
1069 *mays* L.) to conservation agriculture cropping system in Southern Africa, *Soil Tillage Res.*,  
1070 146, 230–242, <https://doi.org/10.1016/j.still.2014.10.015>, 2015.
- 1071 Thierfelder, C., Chivenge, P., Mupangwa, W., Rosenstock, T. S., Lamanna, C., and Eyre, J. X.:  
1072 How climate-smart is conservation agriculture (CA)? – its potential to deliver on adaptation,  
1073 mitigation and productivity on smallholder farms in southern Africa, *Food Secur.*, 9, 537–560,  
1074 <https://doi.org/10.1007/s12571-017-0665-3>, 2017.
- 1075 Thierfelder, C., Baudron, F., Setimela, P., Nyagumbo, I., Mupangwa, W., Mhlanga, B., Lee,  
1076 N., and Gérard, B.: Complementary practices supporting conservation agriculture in southern  
1077 Africa. A review, *Agron. Sustain. Dev.*, 38, <https://doi.org/10.1007/s13593-018-0492-8>, 2018.
- 1078 Thorup-Kristensen, K., Halberg, N., Nicolaisen, M., Olesen, J. E., Crews, T. E., Hinsinger, P.,  
1079 Kirkegaard, J., Pierret, A., and Dresbøll, D. B.: Digging Deeper for Agricultural Resources,  
1080 the Value of Deep Rooting, *Trends Plant Sci.*, 25, 406–417,  
1081 <https://doi.org/10.1016/j.tplants.2019.12.007>, 2020.
- 1082 Villarino, S. H., Pinto, P., Jackson, R. B., and Piñeiro, G.: Plant rhizodeposition : A key factor  
1083 for soil organic matter formation in stable fractions, *Sci. Adv.*, 7, 1–14,

Mis en forme : Police :Italique



1084 <https://doi.org/10.1126/sciadv.abd3176>, 2021.

1085 Virk, A. L., Lin, B. J., Kan, Z. R., Qi, J. Y., Dang, Y. P., Lal, R., Zhao, X., and Zhang, H. L.:  
1086 Simultaneous effects of legume cultivation on carbon and nitrogen accumulation in soil, *Adv.*  
1087 *Agron.*, 171, 75–110, <https://doi.org/10.1016/bs.agron.2021.08.002>, 2022.

1088 Wendt, J. W. and Hauser, S.: An equivalent soil mass procedure for monitoring soil organic  
1089 carbon in multiple soil layers, *Eur. J. Soil Sci.*, 64, 58–65, <https://doi.org/10.1111/ejss.12002>,  
1090 2013.

1091 Yang, L., Luo, Y., Lu, B., Zhou, G., Chang, D., Gao, S., Zhang, J., Che, Z., and Cao, W.: Long-  
1092 term maize and pea intercropping improved subsoil carbon storage while reduced greenhouse  
1093 gas emissions, *Agric. Ecosyst. Environ.*, 349, 108444,  
1094 <https://doi.org/10.1016/J.AGEE.2023.108444>, 2023.

1095 Yost, J. L. and Hartemink, A. E.: How deep is the soil studied – an analysis of four soil science  
1096 journals, *Plant Soil*, 452, 5–18, <https://doi.org/10.1007/s11104-020-04550-z>, 2020.

1097