

1 **Mulch application as the overarching factor explaining increase in soil organic carbon**
2 **stocks under conservation agriculture in two 8-year-old experiments in Zimbabwe**

3 Armwell Shumba^{a,b,c*}, Regis Chikowo^{a,d}, Christian Thierfelder^e, Marc Corbeels^{f,g}, Johan Six^h,
4 Rémi Cardinael^{a,b,f}

5 ^aDepartment of Plant Production Sciences and Technologies, University of Zimbabwe, Harare,
6 Zimbabwe

7 ^bCIRAD, UPR AIDA, Harare, Zimbabwe

8 ^cFertilizer, Farm Feeds and Remedies Institute, Department of Research and Specialist
9 Services, Ministry of Lands, Agriculture, Fisheries, Water and Rural Development, Harare,
10 Zimbabwe

11 ^dPlant, Soil and Microbial Sciences Department, Michigan State University, East Lansing, MI
12 48824, USA

13 ^eInternational Maize and Wheat Improvement Center (CIMMYT), P.O. Box MP 163, Mount
14 Pleasant, Harare, Zimbabwe

15 ^fAIDA, Univ Montpellier, CIRAD, Montpellier, France

16 ^gIITA, International Institute of Tropical Agriculture, PO Box 30772, Nairobi, 00100, Kenya

17 ^hDepartment of Environmental Systems Science, ETH Zurich, 8092 Zürich, Switzerland

18 * Corresponding author. Email: armwellshumba123@gmail.com

19

20

21

22

23

24

25

26 **Abstract**

27 Conservation agriculture (CA), combining reduced or no tillage, permanent soil cover and
28 improved rotations, is often promoted as a climate-smart practice. However, our understanding
29 about the impact of CA and its respective three principles on top and sub-soil organic carbon
30 (SOC) stocks in low input cropping systems of sub-Saharan Africa is rather limited. The study
31 was conducted at two long-term experimental sites established in 2013 in Zimbabwe. The soil
32 types were abruptic Lixisols at Domboshava Training Centre (DTC) and xanthic Ferralsol at
33 the University of Zimbabwe farm (UZF). Six treatments, replicated four times were
34 investigated: conventional tillage (CT), conventional tillage with rotation (CTR), NT, no-
35 tillage with mulch (NTM), no-tillage with rotation (NTR), no-tillage with mulch and rotation
36 (NTMR). Maize (*Zea mays* L.) was the main crop and treatments with rotation included cowpea
37 (*Vigna unguiculata* L. Walp.). SOC concentration and bulk density were determined for
38 samples taken from the 0-5, 5-10, 10-15, 15-20, 20-30, 30-40, 40-50, 50-75 and 75-100 cm
39 depths. Cumulative organic inputs to the soil were also estimated in all treatments. SOC stocks
40 at equivalent soil mass were significantly ($p < 0.05$) higher under NTM, NTR and NTMR
41 compared to NT and CT in top 5 and 10 cm layers at UZF, while SOC stocks were only
42 significantly higher under NTM and NTMR compared to NT and CT in top 5 cm at DTC. NT
43 alone had a slightly negative impact on top SOC stocks. Cumulative SOC stocks were not
44 significantly different between treatments when considering the whole 100 cm soil profile. Our
45 results showed the overarching role of crop residue mulching in CA cropping systems in
46 enhancing SOC stocks but that this effect was limited to the topsoil. The highest cumulative
47 organic carbon inputs to the soil were observed in NTM treatments at the two sites, and this
48 could probably explain the positive effect on SOC stocks. Our results also showed that the
49 combination of at least two CA principles including mulch is required to increase SOC stocks
50 in these low nitrogen input cropping systems.

51 **Key words:** organic inputs, climate change mitigation, climate-smart agriculture, deep soil
52 organic carbon, sustainable intensification

53

54 **1. Introduction**

55 Soil organic carbon (SOC) is an important determinant of soil fertility, productivity and
56 sustainability, and is a useful indicator of soil quality in tropical agricultural systems where
57 nutrient poor and highly weathered soils are managed with little external inputs (Lal, 1997;
58 Feller and Beare, 1997; Chivenge et al., 2007). Therefore, rebuilding depleted SOC stocks in
59 such soils holds potential to contribute to climate change mitigation (Bossio et al., 2020;
60 Minasny et al., 2017; Swanepoel et al., 2016) through sustainable management of agricultural
61 soils (Paustian et al., 2016; Dignac et al., 2017).

62 Conservation agriculture (CA), based on minimum soil disturbance, crop residue retention and
63 crop rotation, has been known to improve surface SOC, with beneficial effects on soil
64 functioning such as improved water infiltration (Thierfelder and Wall, 2012, 2009) and better
65 aggregate stability (Six et al., 1999; Thierfelder and Wall, 2012). The potential of CA to
66 increase SOC stocks and thereby mitigate climate change has, however, been much debated
67 (Corbeels et al., 2020a) but the general understanding is that, this potential is relatively low
68 (Du et al., 2017; Powlson et al., 2014, 2016; Cheesman et al., 2016; Corbeels et al., 2020a). In
69 fact, soil C storage has often been over-estimated for CA due to shallow soil sampling.
70 Compared to conventional tillage systems, no-tillage redistributes SOC in the soil profile, with
71 higher concentrations in the topsoil but potentially lower concentrations below, which can
72 result in no differences in whole profile SOC stocks between no-tillage and conventional tillage
73 (Angers and Eriksen-Hamel, 2008). However, this lack of significant differences in many

74 studies assessing whole profile SOC stocks suffer from not enough statistical power to
75 accurately assess the potential significant SOC changes (Kravchenko and Robertson, 2011).

76 CA can potentially build SOC in deeper soil layers from e.g. the use of cover crops with more
77 extended root systems (Kell, 2011; Thorup-Kristensen et al., 2020; Yang et al., 2023).
78 However, proper soil sampling strategies, to account for both topsoil and subsoil (> 30 cm
79 depth) SOC stocks must, therefore, be prioritized. Soil sampling has often been limited to the
80 top soil plough layer (0-30 cm) in the past two decades (Dube et al., 2012; Powlson et al., 2016;
81 Patra et al., 2019; Yost and Hartemink, 2020), where SOC concentrations, root densities
82 (Chikowo et al., 2003) and microbial activities (Mtambanengwe et al., 2004) are generally
83 largest (Rumpel et al., 2012) and which is the minimum default soil depth recommended by
84 the Intergovernmental Panel for Climate Change (IPCC, 2019). In a meta-analysis of SOC
85 stocks in the top 1 m of soils, Balesdent et al. (2018) found that soils below 0.3 m contain on
86 average 47 % of total SOC stock in the 1 m soil depth and accounts for 19 % of the SOC that
87 has been recently incorporated. Therefore, focusing on topsoil only, could underestimate the
88 potential of agricultural management practices to store SOC (Cardinael et al., 2015). In turn,
89 this can give wrong conclusions on the climate change mitigation potential of agricultural
90 management practices.

91 There have been many studies on the effects of CA on crop productivity and soil health benefits
92 (Corbeels, Naudin, et al., 2020; Kimaro et al., 2016; Mhlanga et al., 2022a; Swanepoel et al.,
93 2018; Thierfelder et al., 2015, 2017; Thierfelder & Mhlanga, 2022), and other studies have
94 fuelled the debate on CA practicality and adoption in SSA (Giller et al., 2009, 2015; Kassam
95 et al., 2019). However, the effects of CA on SOC dynamics has not been widely investigated
96 in SSA. Thierfelder et al., (2017) have alluded to the fact that, data on climate change mitigation
97 potential of CA in southern Africa is scanty hence the need for more research to better quantify

98 the mitigation effects of CA as a climate-smart technology. It has also been observed that
99 depending on the socio-economic and biophysical conditions, farmers may find it easier to
100 adopt certain CA principles and/or their different combinations (Mbanyele et al., 2021;
101 Baudron et al., 2012), although this also opened up new debates (Thierfelder et al., 2018).
102 Therefore, in this study, the focus was on the individual versus combined effects of CA
103 principles (no-tillage, crop residue retention, crop rotation) on SOC stocks.

104 As changes in SOC stocks take time to be detected, long-term experiments are crucial, but are
105 rare, especially in Africa (Bationo et al., 2013; Cardinael et al., 2022; Powlson et al., 2016;
106 Thierfelder and Mhlanga, 2022). This study was conducted on two long-term experiments
107 established in 2013 in Zimbabwe. We hypothesized that the full combination of CA
108 components would be associated with higher increases in SOC stocks than adoption of only
109 one component. This increase in SOC stocks could mainly be due to increased C inputs to the
110 soil, especially under minimum soil disturbance. However, C inputs due to crop rotation could
111 be indirect through increased crop productivity due to reduction on biotic pressure (pests and
112 diseases), and therefore C inputs to the soil might be increased too. Cereals, in cereal-legume
113 rotations may benefit from added soil nitrogen through biological nitrogen fixation from the
114 preceding legume crop enhancing their productivity. Crop diversification, on the other hand,
115 can enhance soil biological processes by increasing the diversity and/or abundance of
116 microfauna like mycorrhizae. This, in turn, improves aggregate stability and offers physical
117 protection for SOC. Lastly, high quality residues (from the legume crop) have been shown to
118 be preferentially stabilized in the soil due to a higher carbon use efficiency of soil microbes
119 (Cotrufo et al., 2013; Kopittke et al., 2018).

120


121 **2. Materials and methods**

122 2.1 Study sites

123 The study was conducted at two long-term experimental sites established in November 2013
124 by CIMMYT. The site at the University of Zimbabwe Farm (UZF) is located about 12 km north
125 of Harare city centre (31° 00' 48" E; 17° 42' 24" S), while the site at the Domboshava Training
126 Centre (DTC) is located about 30 km north-east of Harare (31° 07' 33" E; 17° 35' 17" S). UZF
127 soils are dolerite-derived xanthic *Ferralsols* (FAO classification) and are medium-textured
128 sandy clay loams (34 % clay) in the top 20 cm with a subsoil (20-40 cm) of slightly higher clay
129 content (38 %). DTC soils are granite-derived abruptic *Lixisols* (FAO classification) and are
130 light-textured sandy loams (15 % clay) in the 0-20 cm layer, overlying abruptly a heavier-
131 textured subsoil (20-40 cm) of 30 % clay (Figure 1).

132 The two study sites have a sub-tropical climate with cool, dry winters and hot, wet summers
133 with mean annual minimum and maximum temperatures of 12°C and 25°C, respectively
134 (Mapanda et al., 2010). The rainy season starts in November and tails off in March with a mean
135 annual rainfall of 826 and 814 mm at UZF and DTC, respectively (Mhlanga, et al., 2022b).
136 Cumulative seasonal rainfall in 2019/20 (474 mm) was almost half of rainfall received in the
137 2020/21 season (932 mm) at DTC (Shumba et al., 2023b). At UZF, cumulative seasonal rainfall
138 was 551 mm and 637 mm in the 2019/20 and 2020/21 cropping seasons. On average, minimum
139 and maximum temperatures were 16.9 and 28.1 °C in 2019/20 and 15.5 and 27.2 °C in 2020/21
140 at DTC and 15.5 and 28.6 °C in 2019/20 and 15.3 and 27.5 °C in 2020/21 cropping season at
141 UZF.

142



143 **Figure 1.** Soil texture to 1 m soil depth at the DTC (top) and UZF (bottom) sites in Zimbabwe.

145 Error bars represent standard errors (N = 4).

146

147 **2.2 Experimental treatments and crop management**

148 Two identical experiments were set up at the study sites and treatments were maintained every
 149 season since November 2013. The experiments were set up in a randomised complete block

150 design (RCBD) with eight treatments replicated in four blocks. However, in this study we
151 investigated only six of these treatments. All crop residues were removed soon after harvesting
152 in all treatments, stored and then applied prior to planting in treatments with mulch. The six
153 treatments in our study were:

- 154 i. Conventional tillage (CT) – land preparation was done through digging with a hand hoe
155 and maize (*Zea mays* L.) was sown as a sole crop in rip lines that were created
156 afterwards using an animal-drawn Magoye ripper (a traditional plough with the
157 mouldboard replaced with a ripper tine) at DTC, and in planting basins (approximately
158 10 cm diameter and 10 cm depth) created using a hand hoe at UZF.
- 159 ii. Conventional tillage with rotation (CTR) – land preparation was done as in the CT
160 treatment and maize was rotated with cowpea (*Vigna unguiculata* L.).
- 161 iii. No-tillage (NT) – sole maize was sown in rip lines created using an animal-drawn
162 Magoye ripper (no further soil disturbance was done) at DTC, and in planting basins
163 (approximately 15 cm diameter and 15 cm depth) created using a hand hoe at UZF.
- 164 iv. No-tillage with mulch (NTM) – maize was sown as in the NT treatment and maize
165 residues from the previous season were applied on the soil surface between maize
166 rows at planting at a rate of 2.5 t DM ha⁻¹.
- 167 v. No-tillage with rotation (NTR) – maize was sown in rip lines and rotated with cowpea.
- 168 vi. No-tillage with mulch and rotation (NTMR) – maize was sown in rip lines and rotated
169 with cowpea and maize residues were applied on the soil surface between maize rows
170 at planting at a rate of 2.5 t DM ha⁻¹.

171 Crop residues were removed every year after harvest and weighed in again to maintain the
172 exact 2.5 t ha⁻¹ residue weight year after year. There was a total of 24 plots at each site which
173 were 6 m wide and 12 m long (72 m²). Treatments with rotation (CTR, NTR, NTMR) were

174 split into 6 m wide and 6 m long (36 m²) subplots where maize and cowpea were grown
175 interchangeably every season (maize was sown on one side of the plot while cowpea on the
176 other side).

177 The inter-row spacing was 90 cm and 45 cm for maize and cowpea, respectively, and the in-
178 row spacing was 25 cm for both crops which translated to 44,444 and 88,888 plants ha⁻¹,
179 respectively. Three seeds were planted per planting station and thinned to one after emergence.
180 Basal dressing of nitrogen (N), phosphorus (P) and potassium (K) was applied within 5 cm of
181 the seeds in the form of compound fertilizer for both maize and cowpea at 11.6 kg N ha⁻¹, 10.6
182 kg P ha⁻¹ and 9.6 kg K ha⁻¹, respectively. Nitrogen top dressing to maize only, was applied at
183 4 and 8 weeks after emergence (WAE) in two equal splits of 23.1 kg N ha⁻¹ each, as ammonium
184 nitrate when soil moisture was adequate. However, in the 2019/20 cropping season at both sites
185 and in the 2020/21 cropping season at UZF, the first N top dressing was delayed by an average
186 of 4.5 and 2.0 weeks, respectively, due to mid-season dry spells. Ammonium nitrate was side
187 dressed within 5 cm of the maize stems. Glyphosate [N-(phosphono-methyl) glycine], a pre-
188 emergent non-selective herbicide was applied at 1.025 L active ingredient ha⁻¹ soon after
189 sowing to control weeds. This was followed by manual hoe weeding whenever weeds reached
190 a maximum of 10 cm height or 10 cm in diameter for stoloniferous weeds to achieve a weed
191 clean field. More details about the experiment can be found in Shumba et al., (2023b) and
192 Mhlanga et al., (2022a).

193

194 **2.3 Soil sampling for bulk density determination and soil organic carbon analysis**

195 Soil sampling was done in May 2021 at both sites. For each treatment and replicate, two
196 sampling points in the maize rows and two sampling points in the middle of the inter-rows

197 were randomly selected. The two samples from the rows were pooled into one sample per
198 depth, similarly to the two samples taken in the inter-rows. The following nine depth
199 increments were considered for both SOC and bulk density (BD) measurements: 0-5, 5-10, 10-
200 15, 15-20, 20-30, 30-40, 40-50, 50-75 and 75-100 cm. Undisturbed soil samples using a metal
201 cylinder (5 cm diameter and 5 cm height) were taken from the following depth ranges 0-5, 5-
202 10, 10-15, 15-20 and 20-30 cm for both SOC and BD measurements. A motorized, hand-held
203 soil corer, with an inside diameter of 10 cm, was used to take samples for the 30-40, 40-50, 50-
204 75 and 75-100 cm depths for SOC analysis from the same positions where undisturbed samples
205 were taken. As no significant differences in BD were found below 20 cm between the different
206 treatments at the two sites (see results section) and to avoid too much destruction of the
207 experimental plots, two soil pits were opened at the edges of the experimental plots (also
208 cropped with maize since 2013) at each site to take BD samples for the 30-40, 40-50, 50-75
209 and 75-100 cm depths. As a result, BD below 30 cm depth was assumed the same across the
210 treatments.

211 Soil samples were crumbled and fresh weight was determined using a field scale. Soil moisture
212 was determined on a sub-sample by drying it in an oven at 105°C for 48 hours. All samples
213 were then air-dried and sieved through a 2 mm sieve to determine the mass proportion of coarse
214 soil (> 2 mm) as a fraction of the whole sample. Bulk density (BD) was determined by dividing
215 the dry mass of soil by the volume of the cylinder. Subsamples from the ≤ 2 mm soil fraction
216 were grinded to < 200 μm for SOC analysis. SOC was analysed in the ISO9001:2015-certified
217 IRD LAMA's laboratory in Dakar by dry combustion on 100-mg aliquots of soil (ground to <
218 200 μm) using a CHN elemental analyser (Thermo Finnigan Flash EA1112, Milan, Italy).

219

220 **2.4 Soil organic carbon stocks calculation**

221 The mass proportion of the coarse fraction (> 2 mm) was removed to calculate SOC stocks.
222 The equivalent soil mass (ESM) approach was used to determine SOC stocks to avoid
223 systematic bias in SOC calculation when using the fixed depth method (Ellert and Bettany,
224 1995; Wendt and Hauser, 2013; von Haden et al., 2020). We defined reference soil mass
225 profiles for each site, based on the lowest cumulative soil mass obtained for each replicate. For
226 these references, cumulative soil mass layers were 0-650, 650-1340, 1340-2060, 2060-4160,
227 4160-5590, 5590-7040, 7040-10550, 10550-13770 Mg soil ha⁻¹ at DTC and 0-460, 460-870,
228 870-1330, 1330-1840, 1840-2760, 2760-4030, 4030-5300, 5300-8190, 8190-11050 Mg soil
229 ha⁻¹ at UZF, which roughly corresponded to soil depth layers of 0-5, 5-10, 10-15, 15-20, 20-
230 30, 30-40, 40-50, 50-75, 75-100 cm, respectively. SOC stocks were calculated on the basis of
231 the same soil mass as the reference profile but different soil depth layers which varied by < 1.5
232 and < 0.6 cm at DTC and UZF, respectively. As mulch was only applied between maize rows,
233 and fertilizer was only applied on maize rows, it was estimated that the row and interrow space
234 represented 22 and 78 % respectively, hence SOC calculations were weighted accordingly
235 (Shumba et al., 2023b). Change in cumulative SOC stock between treatments for a given soil
236 depth was determined using the CT treatment as the reference treatment:

$$237 \Delta SOC \text{ stock} = SOC \text{ stock}_{treatment(i)} - SOC \text{ stock}_{CT(i)}, \text{ (Equation 1)}$$

238 where $SOC \text{ stock}_{treatment}$ is the cumulative SOC stock per treatment (CTR, NT, NTM, NRT,
239 NTMR) at a given soil layer and (*i*) representing 0-5, 0-10, 0-15, 0-20, 0-30, 0-40, 0-50, 50-
240 75, 75-100 cm.

241 SOC accumulation or loss rates (kg C ha⁻¹ yr⁻¹) were calculated by dividing the change in
242 stocks by the number of years between the establishment of the experiment and the time of soil
243 sampling (8 years):

244 $SOC\ accumulation/loss\ rate = \frac{\Delta SOC\ stocks}{8} \times 1000$, (Equation 2)

245

246 **2.5 Estimation of organic carbon inputs to the soil**

247 Maize and cowpea yield and aboveground biomass were measured since the inception of the
248 experiment, except for cowpea during the 2013/14 season. This data gap was filled by using
249 the average cowpea yield and aboveground biomass values across seasons (from 2013/14 to
250 2020/21). We assumed that 5 % of the maize aboveground vegetative biomass remained in the
251 field because maize stalk slashing at harvesting did not remove the whole stem. A root:shoot
252 ratio of 0.16 and 0.06 for maize and cowpea, respectively (Amos and Walters, 2006; Kahn and
253 Schroeder, 1999; Kimiti, 2011) was used to estimate the contribution of roots to organic C
254 inputs to the soils. Organic C input contribution from weeds was assumed insignificant since
255 there was effective control of weeds through the use of pre-emergence herbicide (glyphosate)
256 and timely manual weeding throughout the cropping season. We also assumed that the relative
257 amounts of organic C transferred through rhizodeposition was the same for maize and cowpea
258 (i.e. 0.45 x root C biomass (Balesdent et al., 2011) and that the organic C content of all plant
259 parts was 430 g kg⁻¹ (Ma et al., 2018). Cumulative organic C inputs to the soil were then
260 estimated for each treatment (Cardinael et al., 2022).

261

262 **2.6 Data analysis**

263 The full dataset is available in the CIRAD repository (Shumba et al., 2023a). Statistical analyses
264 were performed using R software, version 4.0.0 (R Core Team 2020). Normality was tested by
265 the Kolmogorov-Smirnov test. After confirming that data were normally distributed, analyses of


266 variance (ANOVA) was carried out to establish any significant treatment effects on BD, SOC
267 concentration, and SOC stock. Separation of means was done using the post hoc Tukey test at 5
268 % significance level using the *emmeans* function from the *emmeans* package (Bolker et al.,
269 2009).

270

271 **3. Results**

272 **3.1 Soil bulk density**

273 The different cropping systems (CT, CTR, NT, NTM, NTR, NTMR) had no significant ($p >$
274 0.05) effect on BD across all soil depths except in the 5-10 cm depth in the inter-row at DTC
275 (Figure 2) where BD was on average 5 % lower in the conventional tillage treatments (CT, CTR)
276 than in the no-tillage treatments (NT, NTR). However, soil depth and location (row or inter-row),
277 and the soil depth x location interaction had significant ($p < 0.001$) effects on BD. In the tillage
278 layer (0-15 cm), BD was at least 2 % higher in the inter-rows than in the rows at both sites. In
279 the deeper soil layer (15 – 30 cm), there were no significant ($p > 0.05$) differences. BD for depths
280 below 30 cm were the same across treatments since it was determined from pits outside the
281 experiment. It ranged between 1.47 – 1.51 and 1.47 – 1.49 g cm⁻³ (Table S1) in the subsoil (30 –
282 100 cm layers) at DTC and UZF, respectively.



283



284 **Figure 2.** Top soil bulk density (0-30 cm) at the Domboshava Training Centre (DTC) and

285 University of Zimbabwe Farm (UZF) experimental sites in Zimbabwe. CT: conventional tillage,
286 CTR: conventional tillage with rotation, NT: no-tillage, NTM: no-tillage with mulch, NTR: no-
287 tillage with rotation, NTMR: no-tillage with mulch and rotation. Error bars represent standard
288 errors (N = 4).

289 **3.2 SOC concentration**

290 SOC concentration decreased significantly ($p < 0.001$) with soil depth (Figure 3, Table S2) and
291 was highest in the surface soil (0-5 cm) for all treatments (Table S2). There were significant
292 treatment effects in the 0-5 cm ($p = 0.001$) and 5-10 cm ($p = 0.005$) soil layers at DTC and in
293 the 0-5 cm layer ($p < 0.001$) only, at UZF. NTM had significantly ($p < 0.05$) higher SOC
294 concentration compared to conventional tillage treatments (CT, CTR) and NT in the 0-5 cm
295 soil layer at both sites (Figure 3); the increase in SOC concentration ranged between 31 to 46
296 % and 14 to 22 % at DTC and UZF, respectively. However, SOC concentration in NTM was
297 equal ($p > 0.05$) to NTR and NTMR treatments at both sites.

298 In the 5-10 cm soil layer of DTC, SOC concentrations in NTM and NTR were at least 19 %
299 higher ($p = 0.005$) than in NT and CT (Table S2). There were no significant ($p > 0.05$) treatment
300 effects on SOC concentration below 10 cm soil depth at DTC and below 5 cm depth at UZF.



301

302 **Figure 3.** Soil depth distribution of organic carbon concentration for the different experimental
 303 treatments at the Domboshava Training Centre (DTC) and University of Zimbabwe Farm
 304 (UZF) experimental sites in Zimbabwe. Error bars represent standard errors (N = 4). CT:
 305 conventional tillage, CTR: conventional tillage with rotation, NT: no-tillage, NTM: no-tillage
 306 with mulch, NTR: no-tillage with rotation, NTMR: no-tillage with mulch and rotation.

307

308 **3.3 SOC stock**

309 There were significant ($p < 0.05$) treatment effects on SOC stocks per soil layer in the top 5 at
 310 UZF and 10 cm at DTC (Table S3). Compared to CT, CTR and NT, NTM had at least 1.1 and
 311 1.3 times more SOC stocks in the top 5 and 10 cm layers at UZF and DTC, respectively. In
 312 terms of cumulative SOC stocks, significant ($p < 0.05$) treatment effects were limited to the top
 313 30 cm at DTC and the 20 cm at UZF, where no tillage with mulching (NTM) increased SOC
 314 stocks (Table 1). There were no significant ($p > 0.05$) tillage effects on SOC stocks (CT vs NT)
 315 for both sites. The rotation component had no significant ($p > 0.05$) effects on SOC stocks
 316 when comparing CTR and NTR at DTC. However, the maize-cowpea rotation under NT (NTR)
 317 had at least 16 % higher SOC stocks in the top 30 cm compared to NT. In contrast, NTR had
 318 at least 7 % more SOC stocks than CTR in the top 10 cm at UZF. Compared to NT and CT,
 319 the mulching component significantly ($p < 0.05$) increased SOC stocks by at least 8 % at UZF
 320 and 13 % at DTC in the top 20 and 30 cm soil layers, respectively. SOC stocks in the full CA
 321 treatment (NTMR) were not significantly ($p > 0.05$) different with the other combinations of
 322 CA principles (NTM, NTR) and CTR at DTC. At UZF, the full CA treatment had similar SOC
 323 stocks as all the other NT treatments (NT, NTM, NTR).

324 SOC stocks for the whole soil profile for this study (0-100 cm) were at least 8.1, 3.5 and 2.1
 325 times higher at DTC and 11.6, 4.4 and 2.4 times higher at UZF than the SOC stocks in the 0-5
 326 cm (surface soil), 0-15 cm (tillage depth) and 0-30 cm (IPCC standard depth) layers. SOC
 327 stocks for the subsoil (30-100 cm) ranged from 18.4 to 51.4 Mg C ha⁻¹ at DTC and 41.9 to
 328 124.9 Mg C ha⁻¹ at UZF. Therefore, subsoil represented more than half (at least 53 % at DTC
 329 and 58 % at UZF) of SOC stocks for the whole 100 cm soil profile.

330

Significance	$p < 0.001$	$p < 0.001$	$p < 0.05$	$p < 0.05$	$p < 0.05$	ns	ns	ns	ns	$p < 0.001$	$p < 0.05$	$p < 0.05$	$p < 0.05$	ns	ns	ns	ns	ns

Site	Cumulative ESM (Mg ha ⁻¹)	Approximate soil depth (cm)	Cumulative SOC stocks (Mg C ha ⁻¹)								LSD
			CT	CTR	NT	NTM	NTR	NTMR			
DTC	650	0-5	3.9 ± 0.8c	4.4 ± 0.4bc	3.8 ± 0.7c	5.9 ± 1.2a	5.3 ± 1.1ab	5.4 ± 0.7ab	0.9		
	1340	0-10	7.4 ± 1.3c	8.2 ± 0.6bc	7.1 ± 1.1c	9.9 ± 1.8a	9.4 ± 1.5ab	9.4 ± 1.1ab	1.2		
	2060	0-15	10.7 ± 1.6c	11.6 ± 0.8bc	10.1 ± 1.3c	13.5 ± 2.0a	13.1 ± 1.7ab	12.9 ± 1.2ab	1.7		
	2760	0-20	13.6 ± 1.7b	14.6 ± 1.0ab	12.9 ± 1.4b	16.7 ± 2.1a	16.2 ± 1.9a	15.8 ± 1.4a	2.1		
	4160	0-30	19.4 ± 1.9ab	20.5 ± 1.2ab	18.4 ± 1.6b	22.3 ± 2.2a	22.0 ± 1.9a	22.0 ± 1.5a	2.7		
	5590	0-40	24.9 ± 2.0a	26.6 ± 1.3a	23.7 ± 1.7a	27.5 ± 2.3a	27.9 ± 2.0a	26.9 ± 1.6a	3.1		
	7040	0-50	29.6 ± 1.9a	31.2 ± 1.3a	28.0 ± 1.8a	32.0 ± 2.4a	32.7 ± 2.1a	31.7 ± 1.7a	3.4		
	10550	0-75	38.5 ± 2.0a	42.6 ± 1.3a	37.3 ± 2.0a	39.5 ± 2.4a	42.7 ± 2.1a	42.6 ± 1.9a	5.2		
	13770	0-100	46.5 ± 2.0a	51.4 ± 1.3a	44.8 ± 2.0a	47.5 ± 2.4a	51.1 ± 2.2a	50.7 ± 2.0a	6.3		
	UZF	460	0-5	8.2 ± 0.9cd	7.9 ± 0.5d	8.6 ± 0.6bc	9.6 ± 1.0a	8.8 ± 0.9bc	9.2 ± 0.9ab	0.7	
870		0-10	15.4 ± 1.5bc	14.8 ± 1.0c	15.9 ± 1.3b	17.3 ± 1.7a	15.9 ± 1.6b	16.3 ± 1.4ab	1.1		
1330		0-15	22.9 ± 1.9b	22.1 ± 1.6b	23.4 ± 1.8b	25.1 ± 2.1a	23.2 ± 1.9b	23.6 ± 1.7ab	1.7		
1840		0-20	30.8 ± 2.2b	29.9 ± 2.1b	31.3 ± 2.0ab	33.3 ± 2.4a	30.9 ± 2.2b	31.0 ± 2.1b	2		
2760		0-30	42.3 ± 2.4a	42.8 ± 2.2a	44.1 ± 2.1a	46.4 ± 2.8a	41.9 ± 2.7a	43.3 ± 2.7a	3.3		
4030		0-40	55.2 ± 2.6a	58.1 ± 2.6a	57.2 ± 2.2a	61.0 ± 3.3a	56.7 ± 3.0a	57.5 ± 3.2a	4.8		
5300		0-50	66.3 ± 2.7a	70.4 ± 3.0a	67.5 ± 2.3a	73.1 ± 3.9a	68.8 ± 3.1a	69.7 ± 3.3a	6.6		
8190		0-75	89.3 ± 3.1a	95.9 ± 3.3a	90.0 ± 2.7a	89.9 ± 4.6a	93.7 ± 3.9a	98.4 ± 4.3a	17		
11050		0-100	107.8 ± 3.5a	119.1 ± 3.7a	109.8 ± 3.3a	110.9 ± 5.2a	116.1 ± 4.9a	124.9 ± 5.6a	19		

331 **Table 1.** Cumulative SOC stocks at the Domboshava Training Centre (DTC) and University
332 of Zimbabwe Farm (UZF) after 8 years of different tillage, residue and crop management
333 systems. Means in the same row followed by different superscript letters are significantly
334 different and associated errors are standard errors (N = 4). CT: conventional tillage, CTR:

335 conventional tillage with rotation, NT: no-tillage, NTM: no-tillage with mulch, NTR: no-tillage
336 with rotation, NTMR: no-tillage with mulch and rotation.

337

338 **3.4 SOC accumulation and loss rates**

339 SOC accumulation rates at UZF differed significantly ($p < 0.05$) with soil depth where top soil
340 layers (0-5, 0-10, 0-15, 0-20 and 0-30 cm) had SOC accumulation rates that were at least 6.9
341 times less than when considering the 0-100 cm soil profile (Table 2). In contrast, there were no
342 significant ($p > 0.05$) differences, at DTC in SOC accumulation rates with depth. On average,
343 SOC accumulation rates ranged between 0.13 and 0.08 Mg C ha⁻¹ yr⁻¹ in the top soil (0-5 cm)
344 to 0.33 and 1.16 Mg C ha⁻¹ yr⁻¹ for the whole 1 m soil profile at DTC and UZF, respectively.
345 The depth and treatment interaction had no significant ($p > 0.05$) effects at both sites.

346 On the other hand, the different treatments in this study had significant ($p < 0.05$) effects in
347 SOC accumulation / loss rates in the top 20 cm soil layer at both sites (Table 2). At DTC, NT
348 had significant ($p < 0.05$) net loss of SOC in the 0-20 cm layer, ranging between -0.09 and -
349 0.02 Mg C ha⁻¹ yr⁻¹, whereas NT treatments (NTM, NTR, NTMR) had SOC accumulation rates
350 ranging from 0.17 to 0.38 Mg C ha⁻¹ yr⁻¹. However, maize stover mulching (NTM) had
351 significantly ($p < 0.05$) higher SOC accumulation rates than CTR (2.9 – 4.2 times) and NT (5.2
352 – 13.5 times) in the top 15 cm and 20 cm layers, respectively. The different combinations of
353 mulching and rotation under NT (NTM, NTR and NTMR) had no significant ($p > 0.05$)
354 differences in SOC accumulation rates. Similarly, rotation treatments (CTR, NTR, NTMR)
355 showed no significant ($p > 0.05$) differences in SOC accumulation rates. Thus, the full CA
356 treatment had similar SOC accumulations rates to treatments with at least 2 combinations of
357 CA principles (NTM and NTR) and to CTR.

358 In contrast, at UZF, CTR had significant ($p < 0.05$) net loss of SOC in the top 20 cm (Table 2).
 359 The no-tillage treatments (NT, NTM, NTR, NTMR) showed significantly ($p < 0.05$) higher
 360 SOC accumulation rates ($0.05 - 0.25 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$) than CTR which ranged between -0.07 to
 361 $-0.03 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ in top 10 cm soil layer. NTM had the highest SOC accumulations rates
 362 (0.28 to $0.32 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$) when considering the 0-15 and 0-20 cm soil layers. SOC
 363 accumulation rates in NTM were at least 3.8, 3.5 and 2.4 times higher than CTR, NT and NTR
 364 in the top 20 cm. The full CA treatment (NTMR) had significantly ($p < 0.05$) higher SOC
 365 accumulation rates compared to CTR ($2.5 - 5.3$ times) in the top 10 cm and lower SOC
 366 accumulation rate to NTM in the 0-10 cm (2.3 times) and 0-20 cm (10.6 times) soil layer.
 367 However, there were no significant ($p > 0.05$) differences in SOC accumulation rates between
 368 treatments beyond 20 cm soil layer at both sites.

Site	Approximate soil depth (cm)	SOC accumulation or loss rate ($\text{Mg C ha}^{-1} \text{ yr}^{-1}$)					LSD	Sig
		CTR	NT	NTM	NTR	NTMR		
DTC	0-5	$0.06 \pm 0.05bc$	$-0.02 \pm 0.02c$	$0.25 \pm 0.05a$	$0.17 \pm 0.02ab$	$0.19 \pm 0.04ab$	0.13	***
	0-10	$0.10 \pm 0.09bc$	$-0.04 \pm 0.04c$	$0.31 \pm 0.09a$	$0.24 \pm 0.01ab$	$0.25 \pm 0.08ab$	0.16	***
	0-15	$0.12 \pm 0.13bc$	$-0.07 \pm 0.05c$	$0.35 \pm 0.13a$	$0.30 \pm 0.01ab$	$0.27 \pm 0.12ab$	0.23	**
	0-20	$0.12 \pm 0.17ab$	$-0.09 \pm 0.05b$	$0.38 \pm 0.17a$	$0.32 \pm 0.01a$	$0.27 \pm 0.16a$	0.29	**
	0-30	$0.13 \pm 0.25a$	$-0.13 \pm 0.07a$	$0.36 \pm 0.25a$	$0.32 \pm 0.08a$	$0.33 \pm 0.20a$	0.35	ns
	0-40	$0.22 \pm 0.25a$	$-0.15 \pm 0.07a$	$0.33 \pm 0.25a$	$0.38 \pm 0.07a$	$0.25 \pm 0.23a$	0.41	ns
	0-50	$0.20 \pm 0.27a$	$-0.20 \pm 0.14a$	$0.30 \pm 0.27a$	$0.40 \pm 0.09a$	$0.26 \pm 0.22a$	0.46	ns
	0-75	$0.51 \pm 0.28a$	$-0.15 \pm 0.28a$	$0.13 \pm 0.28a$	$0.53 \pm 0.13a$	$0.51 \pm 0.20a$	0.73	ns
	0-100	$0.62 \pm 0.32a$	$-0.20 \pm 0.37a$	$0.13 \pm 0.32a$	$0.58 \pm 0.29a$	$0.53 \pm 0.20a$	0.86	ns
UZF	0-5	$-0.03 \pm 0.03c$	$0.05 \pm 0.04b$	$0.17 \pm 0.05a$	$0.07 \pm 0.04b$	$0.13 \pm 0.06ab$	0.08	***
	0-10	$-0.07 \pm 0.04c$	$0.07 \pm 0.08b$	$0.25 \pm 0.09a$	$0.07 \pm 0.07b$	$0.11 \pm 0.08b$	0.13	**
	0-15	$-0.10 \pm 0.03b$	$0.06 \pm 0.11b$	$0.28 \pm 0.13a$	$0.04 \pm 0.07b$	$0.09 \pm 0.11ab$	0.22	**
	0-20	$-0.11 \pm 0.07b$	$0.06 \pm 0.14b$	$0.32 \pm 0.17a$	$0.02 \pm 0.11b$	$0.03 \pm 0.12b$	0.25	**
	0-30	$0.06 \pm 0.15a$	$0.22 \pm 0.25a$	$0.51 \pm 0.25a$	$-0.05 \pm 0.18a$	$0.12 \pm 0.16a$	0.44	ns
	0-40	$0.37 \pm 0.11a$	$0.25 \pm 0.27a$	$0.72 \pm 0.25a$	$0.19 \pm 0.28a$	$0.29 \pm 0.14a$	0.65	ns
	0-50	$0.51 \pm 0.20a$	$0.15 \pm 0.34a$	$0.85 \pm 0.27a$	$0.31 \pm 0.41a$	$0.43 \pm 0.10a$	0.88	ns
	0-75	$0.83 \pm 0.56a$	$0.08 \pm 0.55a$	$0.08 \pm 0.28a$	$0.55 \pm 0.76a$	$1.14 \pm 0.44a$	1.17	ns
	0-100	$1.41 \pm 0.86a$	$0.25 \pm 0.75a$	$0.98 \pm 0.32a$	$1.03 \pm 1.26a$	$2.14 \pm 0.99a$	2.31	ns


369 **Table 2.** SOC change rates (\pm standard error, $N = 4$) of the different treatments compared to
 370 CT (conventional tillage) at Domboshava Training Centre (DTC) and University of Zimbabwe

371 farm (UZF). Means in the same row followed by different superscripts are significantly
372 different. CTR: conventional tillage with rotation, NT: no-tillage, NTM: no-tillage with mulch,
373 NTR: no-tillage with rotation, NTMR: no-tillage with mulch and rotation, LSD = least
374 significance difference, ns = not significant, Sig = significance, ** = $p < 0.05$, *** = $p < 0.001$.

375

376 **3.5 Organic carbon inputs via crops residues, root mortality and rhizodeposition to the** 377 **soil**

378 There were significant ($p < 0.001$) differences in cumulative OC inputs between treatments
379 (Figure 4). Cumulative OC inputs were at least 1.5 times higher in mulch treatments (NTM,
380 NTMR) than in treatments without mulch. However, the mulch plus rotation treatment
381 (NTMR) had significantly ($p < 0.001$) lower cumulative OC inputs than continuous mulching
382 (NTM). Cumulative OC input after 8 seasons was as high as 16.0 and 10.5 Mg C ha⁻¹ at DTC
383 and 16.2 and 12.4 Mg C ha⁻¹ at UZF in NTM and NTMR, respectively (Figure 4), resulting in
384 mean annual OC input rates of about 1.3 to 1.6 Mg C ha⁻¹ yr⁻¹ for NTMR and 2.0 Mg C ha⁻¹ yr⁻¹
385 ¹ for NTM. The other treatments had mean annual OC input rates ≤ 1.0 Mg C ha⁻¹ yr⁻¹.



386

387 **Figure 4.** Estimated cumulative organic carbon (OC) inputs to the soil from the 2013/14 to the
 388 2020/21 cropping season for the different treatments at the Domboshava Training Centre
 389 (DTC) and the University of Zimbabwe Farm (UZF) experimental sites. Error bars represent
 390 standard errors ($n = 4$) for the cumulative OC. CT: conventional tillage, CTR: conventional
 391 tillage with rotation, NT: no-tillage, NTM: no-tillage with mulch, NTR: no-tillage with

392 rotation, NTMR: no-tillage with mulch and rotation, ABG: aboveground biomass
393 (corresponding to an estimated 5 % of the maize aboveground vegetative biomass remained in
394 the field because maize stalk slashing at harvesting did not remove the whole stem).

395

396 **4. Discussion**

397 **4.1 SOC distribution across soil depth**

398 Cumulative SOC stocks for the whole soil profile (0-100 cm) for this study were at least 8.0,
399 4.0 and 2.0 times higher than the 0-5 (surface soil), 0-15 (tillage depth for the study) and 0-30
400 (IPCC standard depth for SOC studies), for the two sites, respectively. This means that, over
401 half of SOC stocks for this study were in the sub-soil (30-100 cm) which reflects on the
402 importance of sub-soil SOC stocks. Significant SOC stocks in the sub-soil have also been
403 reported by other authors (Yost and Hartemink, 2020; Balesdent et al., 2018; Cardinael et al.,
404 2015; Harrison et al., 2011; Lal, 2018). Significant effects of mulch and/or rotation under NT
405 were restricted to the top 30 cm in our study as well as other studies in SSA (Dube et al., 2012;
406 Powlson et al., 2016) and the world at large (Balesdent et al., 2018; Yost and Hartemink, 2020),
407 which is most likely why the default soil depth for IPCC for SOC studies is 0-30 cm (IPCC,
408 2019). However, this underestimates whole soil profile C storage (Harrison et al., 2011; Singh
409 et al., 2018; Lorenz and Lal, 2005). Therefore, it is crucial to consider whole soil profile
410 sampling when monitoring SOC storage in agricultural ecosystems to determine their C
411 sequestration potential in the pursuit of climate change mitigation (Malepfane et al., 2022).
412 SOC mineralization is relatively low in the sub-soil due to lack of oxygen and physical
413 protection of SOC (aggregate protected C) (Rumpel et al., 2012; Sanaullah et al., 2016; Shumba
414 et al., 2020; Button et al., 2022). Therefore, in the pursuit to improve subsoil (> 30 cm) SOC

415 stocks through root mortality and exudates, crop varieties with higher root-length densities
416 (Chikowo et al., 2003) in the subsoil are recommended.

417

418 **4.2 Cumulative SOC stocks and accumulation rates**

419 Overall, the insignificant differences in cumulative SOC stocks and SOC accumulation / loss
420 rates between the different cropping systems in this and other studies (Laura et al., 2022; Leal
421 et al., 2020) when considering whole soil profile (0-100 cm) is alluded to the dilution effect
422 due to low OC inputs beyond 30 cm. Organic C inputs in the subsoil (> 30 cm) through crop
423 root mortality and root exudates are highly reduced due to low root biomass (Button et al.,
424 2022; Chikowo et al., 2003). Several authors have also reported similar cumulative SOC stocks
425 for soil profile depth > 30 cm between different tillage and residue management practices
426 (Angers et al., 1997; Doran et al., 1998; Lal, 2015; Powlson et al., 2014). This can be alluded
427 to an accumulation of uncertainty when cumulating SOC stocks in several soil layers with their
428 respective error of measurement. This weakens the power of detecting statistically significant
429 differences even where such differences exist (Kravchenko and Robertson, 2011). Kravchenko
430 and Robertson, (2011) bemoaned the lack of enough replication when sampling deep soil
431 horizons to reduce variability and the importance of post hoc power analysis to reduce Type II
432 error. This study was limited to four replicates which might not have enough statistical power
433 to detect significant differences between treatments when considering the whole soil profile.

434 **4.2.1 Mulching**

435 The overarching role of mulching in cumulative SOC stocks and accumulation / loss rates at
436 both sites (Tables 1 and 2), albeit, in the top soil (< 30 cm) has been shown in this study.
437 Cumulative SOC stocks (Table 1) and SOC accumulation / loss rates (Table 2) did not differ

438 with residue management under NT systems (NTM, NTMR) in the top soil at DTC regardless
439 of high external OC inputs through maize residue application in mulch treatments (Figure 4,
440 Table S4). This was attributed to low clay content (< 15 % clay) in the top 20 cm hence low
441 physical SOC protection (Chivenge et al., 2007; Mtambanengwe et al., 2004; Sun et al., 2020),
442 such that the differences in OC inputs had little effect. Alternatively, SOC can be protected
443 from mineralization through adsorption to clay particles (Han et al., 2016; Churchman et al.,
444 2020). However, there was low surface area for SOC adsorption due to low clay content in the
445 top soil at DTC. Conversely, maize residue mulching effects were significant at UZF though
446 NTMR was indifferent when compared to the rest of the NT treatments. Cumulative OC inputs
447 in NTMR (12.4 Mg C ha⁻¹) were about 77 % of cumulative OC inputs in NTM (16.2 Mg C ha⁻¹)
448 but at least 57 % higher OC inputs than NT and NTR after 8 seasons (Figure 4). This was
449 alluded to SOC adsorption and physical protection due to higher clay content at UZF.
450 Nonetheless, several studies have shown that aboveground biomass is less effective in
451 sustaining SOC stocks compared to belowground biomass (Hirte et al., 2018, 2021; Jones et
452 al., 2009; Villarino et al., 2021) and we attribute that to the insignificant cumulative SOC stocks
453 and accumulation rates between the NT treatments other than NTM regardless of higher
454 aboveground biomass in NTR and NMTR than NT (Figure 4).

455 **4.2.2 Tillage**

456 Significant changes in SOC stocks and accumulation rates were restricted to the top 20 cm at
457 both sites below which there were no differences between treatments (Table 2). The
458 consistently low SOC stocks under NT, CT and CTR and SOC accumulation rates (CTR and
459 NT) at both sites was attributed to low OC inputs (Table S4, Figure 4). NT and CT had
460 generally low yields in terms of grain and vegetative aboveground biomass (Mhlanga et al.,
461 2021; Shumba et al., 2023b) since the establishment of the experiment in the 2013/14 season

462 and hence low OC inputs through stubble, root mortality and root exudates. Our results
463 dovetails with studies done elsewhere (Du et al., 2017; Koga and Tsuji, 2009) and meta-
464 analyses and reviews (Corbeels et al., 2020a; Lal, 2018, 2015) where the authors found that
465 NT alone does not significantly improve SOC. However, higher SOC stocks were observed
466 when NT was combined with at least two CA principles (mulching and rotation) at DTC in the
467 top 20 cm (Table 1). It has been reported that NT cropping systems enhance SOC accumulation
468 through increasing C inputs in the top layers and reducing erosion through minimum soil
469 disturbance (Six et al., 2000; Lal, 2015, 2018; Bai et al., 2019; Cai et al., 2022). Minimum soil
470 disturbance through NT also physically protects SOC in microaggregates from exposure to
471 oxidative losses (Shumba et al., 2020; Six et al., 2002; Dolan et al., 2006; Liang et al., 2020).
472 However, NT without mulch is a nonentity compared to other combinations of CA principles
473 for long-term sustainability in cropping systems (Nyamangara et al., 2013; Kodzwa et al., 2020;
474 Mhlanga et al., 2021; Li et al., 2020; Bohoussou et al., 2022) and NT is only effective in
475 increasing SOC stocks when it is associated with other CA principles, especially mulch. On
476 the other hand, our study suggest that NT can achieve the same results of SOC storage as NTR
477 and NTMR since they had similar SOC stocks at UZF. This can be explained by the low
478 aboveground OC inputs in rotation treatments during the season when cowpeas were grown.

479 **4.2.3 Maize-cowpea rotation**

480 Legume rotations have been found to improve SOC accumulation rates and subsequent soil
481 structural improvement (aggregation) induced by the addition of organic residues with
482 favourable C/N ratio (Virk et al., 2022; Laub et al., 2023; Jephita et al., 2023). However, in our
483 study, cowpea rotation benefits on SOC accumulation rates were not significant at DTC.
484 Maize-cowpea rotation had no significant effects on maize yield (Shumba et al., 2023b;
485 Mhlanga et al., 2021) which corresponded to low belowground biomass as well. Instead, maize

486 stover mulching improved maize yields at DTC. Nevertheless, benefits from cowpea rotation
487 under NT cropping systems (NTR, NTMR) compared to CT cropping systems (CTR) were
488 significant, albeit only in the top 10 cm, at UZF; CTR had a net loss of SOC (-0.07 ± 0.04 to
489 $0.03 \pm 0.03 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$). Significantly higher maize yields in rotation treatments were
490 observed at UZF (Shumba et al., 2023b; Mhlanga et al., 2021) and were attributed to more soil
491 mineral N due to biological nitrogen fixation from the preceding cowpeas. Higher aboveground
492 biomass is positively related to below ground biomass resulting in significant belowground OC
493 inputs, of higher quality in the rotation treatments in the season when maize is grown. However,
494 the net SOC loss in CTR at UZF was due to seasonal exposure to oxidative losses (SOC
495 mineralization) through disruption of soil macroaggregates by tillage as alluded by Bai et al.,
496 (2019); Cambardella and Elliott, (1993) and Lal, (2018). We underscore that maize-cowpea
497 rotation under NT improved SOC accumulation in the top soil due to reduced soil disturbance
498 and alternate OC inputs of high (cowpeas) and low quality (maize). High quality OC inputs
499 have a positive priming effect (Chen et al., 2014) which have been shown to be preferentially
500 stabilized in the soil due to a higher carbon use efficiency of soil microbes (Cotrufo et al., 2013;
501 Kopittke et al., 2018). This explains significant improvement in SOC stocks under the
502 combination of NT and alternate high- and low-quality OC inputs (maize-cowpea rotation) to
503 the soil in medium to heavy textured soils at UZF and vice versa at DTC.

504

505 **4.3. Role of soil texture in SOC accumulation**

506 Soil texture is widely recognized to influence SOC stocks (Sun et al., 2020) through physical
507 and chemical protection of SOC against microbially mediated decomposition (Chivenge et al.,
508 2007; Mtambanengwe et al., 2004). In our study, the main difference between the two study
509 sites is soil texture in the top soil (0-30 cm), where DTC had light textured (sandy loams) soils

510 and UZF had medium to heavy textured (sandy-clay-loams) soils (Figure 1). These soil textural
511 differences explain why there were no differences in SOC stocks, changes and accumulation
512 rates between NTM, NTR and NTMR at DTC regardless of higher cumulative OC inputs in
513 NTM and NTMR (Figure 4). The direct SOC inputs in the top soil at DTC, where SOC was
514 more concentrated (Table S2, Figure 3) was subject to mineralization because of low clay
515 content and thus low protection by soil micro-aggregates (Chivenge et al., 2007;
516 Mtambanengwe et al., 2004; Sun et al., 2020), such that the differences in OC inputs had little
517 effect. Light textured soils have large pores which cannot protect SOC against microbial
518 decomposition (Mtambanengwe et al., 2004; Christensen, 1987; Sun et al., 2020; Kravchenko
519 and Guber, 2017). Additionally, the low clay content meant less surface area for SOC
520 adsorption (Han et al., 2016; Churchman et al., 2020) which is another mechanism for SOC
521 protection from mineralization. In contrast, there were differences between NTM and NTR at
522 UZF in the top soil layers and intermediate between NTM and NTMR. Cumulative OC inputs
523 in NTMR (12.4 Mg C ha⁻¹) were about 75 % of cumulative OC inputs in NTM (16.2 Mg C ha⁻¹)
524 (Figure 4) after 8 seasons. The added C, especially from maize stover mulch, most likely
525 was protected by clay particles as well as formation of organo-mineral complexes (Malepfane
526 et al., 2022; Chivenge et al., 2007; Jephitha et al., 2023) which protects SOC from mineralization
527 (Dunjana et al., 2012; Shumba et al., 2020; Button et al., 2022; Rumpel et al., 2012; Sanaullah
528 et al., 2016).

529

530 **5. Conclusions**

531 Our study has shown the overarching importance of mulching and of combining at least two
532 CA principles to improve top SOC stocks. No tillage (NT) alone could not increase SOC stocks,
533 and even led to a slight decrease compared to CT, due to lower crop productivity in NT and
534 therefore reduced OC inputs to the soil. Nevertheless, whole profile (0-100 cm) SOC stocks

535 were the same between all the treatment. Our study also showed that sampling the entire soil
536 profile is necessary for a more accurate view of SOC accumulation potential among different
537 cropping systems.

538

539 **6. Data availability**

540 All data are freely available on the CIRAD data repository
541 <https://doi.org/10.18167/DVN1/VPOCHN> (Shumba et al., 2023a).

542

543 **7. Author contributions**

544 CT designed, established and maintained the experiments since 2013; RCa, RCh were involved
545 in soil sampling campaigns; AS, RCa performed the statistical analyses, graphics and drafting
546 the manuscript; AS, RCa, RCh, MC, JS, CT reviewed and edited the manuscript.

547

548 **8. Competing interests**

549 One of the co-authors is a member of the editorial board of the SOIL journal. The other authors
550 have no competing interests to declare.

551

552 **9. Acknowledgements**

553 This study was funded by the DSCATT project “Agricultural Intensification and Dynamics of
554 Soil Carbon Sequestration in Tropical and Temperate Farming Systems” (N° AF 1802-001, N°
555 FT C002181), supported by the Agropolis Foundation (“Programme d’Investissement
556 d’Avenir” Labex Agro, ANR-10-LABX- 0001-01) and by the TOTAL Foundation within a
557 patronage agreement. Authors are grateful to the International Maize and Wheat Improvement

558 Center (CIMMYT) for the setup and running of the experiment. We also acknowledge the
559 donors of the MAIZE CGIAR Research Program (www.maize.org) and the Ukama Ustawi
560 Regional CGIAR Initiative who supported the trials up to 2018 and staff time until 2023. We
561 thank Britta Jahn-Humphrey for carrying out the gas analyses at ETH Zürich. We also thank
562 Admire Muwati for his help in gas sampling. Special thanks go to the technical personnel at
563 each experimental locations namely Tarirai Muoni, Sign Phiri, Herbert Chipara and Connie
564 Madembo who continuously assisted in trial establishment and management.

565

566 **10. References**

- 567 Amos, B. and Walters, D. T.: Maize Root Biomass and Net Rhizodeposited Carbon, *Soil Sci.*
568 *Soc. Am. J.*, 70, 1489–1503, <https://doi.org/10.2136/sssaj2005.0216>, 2006.
- 569 Angers, D. A. and Eriksen-Hamel, N. S.: Full-inversion tillage and organic carbon distribution
570 in soil profiles: A meta-analysis, *Soil Sci. Soc. Am. J.*, 72, 1370–1374,
571 <https://doi.org/10.2136/sssaj2007.0342>, 2008.
- 572 Angers, D. A., Bolinder, M. A., Carter, M. R., Gregorich, E. G., Drury, C. F., Liang, B. C.,
573 Voroney, R. P., Simard, R. R., Donald, R. G., Bevaert, R. P., and Martel, J.: Impact of tillage
574 practices on organic carbon and nitrogen storage in cool, humid soils of eastern Canada, *Soil*
575 *Tillage Res.*, 41, 191–201, <https://doi.org/10.4141/S96-111>, 1997.
- 576 Bai, X., Huang, Y., Ren, W., Coyne, M., Jacinthe, P.-A., Bo, T., Hui, D., Yang, J., and
577 Matocha, C.: Responses of soil carbon sequestration to climate-smart agriculture practices: A
578 meta-analysis, 25, 2591–2606, <https://doi.org/10.1111/gcb.14658>, 2019.
- 579 Balesdent, J., Derrien, D., Fontaine, S., Kirman, S., Klumpp, K., Loiseau, P., Marol, C.,
580 Nguyen, C., Pean, M., Personeni, E., and Robin, C.: Contribution de la rhizodéposition aux
581 matières organiques du sol, quelques implications pour la modélisation de la dynamique du
582 carbone, *Etude Gest. des sols*, 18 (3), 201–216, 2011.
- 583 Balesdent, J., Basile-Doelsch, I., Chadoeuf, J., Cornu, S., Derrien, D., Fekiacova, Z., and Hatté,
584 C.: Atmosphere–soil carbon transfer as a function of soil depth, *Nature*, 559, 599–602,
585 <https://doi.org/10.1038/s41586-018-0328-3>, 2018.
- 586 Bationo, A., Kihara, J., Vanlauwe, B., Waswa, B., Adolwa, I., and Saidou, K.: Overview of
587 Long Term Experiments in Africa, in: *Lessons Learned from Long-Term Soil Fertility*
588 *Management Experiments in Africa.*, Springer Science+Business Media Dordrecht, 1–26,
589 <https://doi.org/10.1007/978-94-007-2938-4>, 2013.
- 590 Baudron, F., Andersson, J. A., Corbeels, M., and Giller, K. E.: Failing to Yield? Ploughs,
591 Conservation Agriculture and the Problem of Agricultural Intensification: An Example from
592 the Zambezi Valley, Zimbabwe, *J. Dev. Stud.*, 48, 393–412,

593 <https://doi.org/10.1080/00220388.2011.587509>, 2012.

594 Bohoussou, Y. N. D., Kou, Y. H., Yu, W. B., Lin, B. jian, Virk, A. L., Zhao, X., Dang, Y. P.,
595 and Zhang, H. L.: Impacts of the components of conservation agriculture on soil organic carbon
596 and total nitrogen storage: A global meta-analysis, *Sci. Total Environ.*, 842, 156822,
597 <https://doi.org/10.1016/J.SCITOTENV.2022.156822>, 2022.

598 Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H.,
599 and White, J. S. S.: Generalized linear mixed models: a practical guide for ecology and
600 evolution, *Trends Ecol. Evol.*, 24, 127–135, <https://doi.org/10.1016/j.tree.2008.10.008>, 2009.

601 Bossio, D. A., Cook-Patton, S. C., Ellis, P. W., Fargione, J., Sanderman, J., Smith, P., Wood,
602 S., Zomer, R. J., von Unger, M., Emmer, I. M., and Griscom, B. W.: The role of soil carbon in
603 natural climate solutions, *Nat. Sustain.*, 3, 391–398, [https://doi.org/10.1038/s41893-020-0491-](https://doi.org/10.1038/s41893-020-0491-z)
604 [z](https://doi.org/10.1038/s41893-020-0491-z), 2020.

605 Button, E. S., Pett-Ridge, J., Murphy, D. V., Kuzyakov, Y., Chadwick, D. R., and Jones, D. L.:
606 Deep-C storage: Biological, chemical and physical strategies to enhance carbon stocks in
607 agricultural subsoils, *Soil Biol. Biochem.*, 170, 108697,
608 <https://doi.org/10.1016/j.soilbio.2022.108697>, 2022.

609 Cai, A., Han, T., Ren, T., Sanderman, J., Rui, Y., Wang, B., Smith, P., Xu, M., and Yu'e Li:
610 Declines in soil carbon storage under no tillage can be alleviated in the long run, 425, 1–3,
611 <https://doi.org/10.1016/j.geoderma.2022.116028>, 2022.

612 Cambardella, C. A. and Elliott, E. T.: Carbon and Nitrogen Distribution in Aggregates from
613 Cultivated and Native Grassland Soils, *Soil Sci. Soc. Am. J.*, 57, 1071–1076,
614 <https://doi.org/10.2136/SSSAJ1993.03615995005700040032X>, 1993.

615 Cardinael, R., Chevallier, T., Barthès, B. G., Saby, N. P. A., Parent, T., Dupraz, C., Bernoux,
616 M., and Chenu, C.: Impact of alley cropping agroforestry on stocks, forms and spatial
617 distribution of soil organic carbon - A case study in a Mediterranean context, *Geoderma*, 259–
618 260, 288–299, <https://doi.org/10.1016/j.geoderma.2015.06.015>, 2015.

619 Cardinael, R., Guibert, H., Kouassi Brédoumy, S. T., Gigou, J., N’Goran, K. E., and Corbeels,
620 M.: Sustaining maize yields and soil carbon following land clearing in the forest–savannah
621 transition zone of West Africa: Results from a 20-year experiment, *F. Crop. Res.*, 275,
622 <https://doi.org/10.1016/j.fcr.2021.108335>, 2022.

623 Cheesman, S., Thierfelder, C., Eash, N. S., Kassie, G. T., and Frossard, E.: Soil carbon stocks
624 in conservation agriculture systems of Southern Africa, *Soil Tillage Res.*, 156, 99–109,
625 <https://doi.org/10.1016/j.still.2015.09.018>, 2016.

626 Chen, R., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Lin, X., Blagodatskaya,
627 E., and Kuzyakov, Y.: Soil C and N availability determine the priming effect: Microbial N
628 mining and stoichiometric decomposition theories, *Glob. Chang. Biol.*, 20, 2356–2367,
629 <https://doi.org/10.1111/gcb.12475>, 2014.

630 Chikowo, R., Mapfumo, P., Nyamugafata, P., Nyamadzawo, G., and Giller, K. E.: Nitrate-N
631 dynamics following improved fallows and maize root development in a Zimbabwean sandy
632 clay loam, *Agrofor. Syst.*, 59, 187–195,
633 <https://doi.org/10.1023/B:AGFO.0000005219.07409.a0>, 2003.

- 634 Chivenge, P. P., Murwira, H. K., Giller, K. E., Mapfumo, P., and Six, J.: Long-term impact of
635 reduced tillage and residue management on soil carbon stabilization: Implications for
636 conservation agriculture on contrasting soils, *Soil Tillage Res.*, 94, 328–337,
637 <https://doi.org/10.1016/j.still.2006.08.006>, 2007.
- 638 Christensen, B. T.: Decomposability of organic matter in particle size fractions from field soils
639 with straw incorporation, *Soil Biol. Biochem.*, 19, 429–435, [https://doi.org/10.1016/0038-](https://doi.org/10.1016/0038-0717(87)90034-4)
640 [0717\(87\)90034-4](https://doi.org/10.1016/0038-0717(87)90034-4), 1987.
- 641 Churchman, G. J., Singh, M., Schapel, A., Sarkar, B., and Bolan, N.: Clay Minerals As the Key
642 To the Sequestration of Carbon in Soils, *Clays Clay Miner.*, 68, 135–143,
643 <https://doi.org/10.1007/s42860-020-00071-z>, 2020.
- 644 Corbeels, M., Cardinael, R., Powlson, D., Chikowo, R., and Gerard, B.: Carbon sequestration
645 potential through conservation agriculture in Africa has been largely overestimated: Comment
646 on: “Meta-analysis on carbon sequestration through conservation agriculture in Africa,” *Soil*
647 *Tillage Res.*, 196, 104300, <https://doi.org/10.1016/j.still.2019.104300>, 2020a.
- 648 Corbeels, M., Naudin, K., Whitbread, A. M., Kühne, R., and Letourmy, P.: Limits of
649 conservation agriculture to overcome low crop yields in sub-Saharan Africa, *Nat. Food*, 1, 447–
650 454, <https://doi.org/10.1038/s43016-020-0114-x>, 2020b.
- 651 Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K., and Paul, E.: The Microbial
652 Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with
653 soil organic matter stabilization: Do labile plant inputs form stable soil organic matter?, *Glob.*
654 *Chang. Biol.*, 19, 988–995, <https://doi.org/10.1111/gcb.12113>, 2013.
- 655 Dignac, M. F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T.,
656 Freschet, G. T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron, P. A.,
657 Nunan, N., Roumet, C., and Basile-Doelsch, I.: Increasing soil carbon storage: mechanisms,
658 effects of agricultural practices and proxies. A review, *Agron. Sustain. Dev.*, 37,
659 <https://doi.org/10.1007/s13593-017-0421-2>, 2017.
- 660 Dolan, M. S., Clapp, C. E., Allmaras, R. R., Baker, J. M., and Molina, J. A. E.: Soil organic
661 carbon and nitrogen in a Minnesota soil as related to tillage, residue and nitrogen management,
662 *Soil Tillage Res.*, 89, 221–231, <https://doi.org/10.1016/j.still.2005.07.015>, 2006.
- 663 Doran, J. W., Elliott, E. T., and Paustian, K.: Soil microbial activity, nitrogen cycling, and long-
664 term changes in organic carbon pools as related to fallow tillage management, *Soil Tillage*
665 *Res.*, 49, 3–18, [https://doi.org/10.1016/S0167-1987\(98\)00150-0](https://doi.org/10.1016/S0167-1987(98)00150-0), 1998.
- 666 Du, Z., Angers, D. A., Ren, T., Zhang, Q., and Li, G.: The effect of no-till on organic C storage
667 in Chinese soils should not be overemphasized: A meta-analysis, *Agric. Ecosyst. Environ.*,
668 236, 1–11, <https://doi.org/10.1016/j.agee.2016.11.007>, 2017.
- 669 Dube, E., Chiduzza, C., and Muchaonyerwa, P.: Conservation agriculture effects on soil organic
670 matter on a Haplic Cambisol after four years of maize-oat and maize-grazing vetch rotations
671 in South Africa, *Soil Tillage Res.*, 123, 21–28, <https://doi.org/10.1016/j.still.2012.02.008>,
672 2012.
- 673 Dunjana, N., Nyamugafata, P., Shumba, A., Nyamangara, J., and Zingore, S.: Effects of cattle
674 manure on selected soil physical properties of smallholder farms on two soils of Murewa,
675 Zimbabwe, *Soil Use Manag.*, 28, <https://doi.org/10.1111/j.1475-2743.2012.00394.x>, 2012.

676 Ellert, B. H. and Bettany, J. R.: Calculation of organic matter and nutrients stored in soils under
677 contrasting management regimes, *Can. J. Soil Sci.*, 75, 529–538,
678 <https://doi.org/10.4141/cjss95-075>, 1995.

679 Feller, C. and Beare, M. H.: Physical control of soil organic matter dynamics in the tropics,
680 *Geoderma*, 79, 69–116, [https://doi.org/10.1016/S0016-7061\(97\)00039-6](https://doi.org/10.1016/S0016-7061(97)00039-6), 1997.

681 Giller, K. E., Witter, E., Corbeels, M., and Tittonell, P.: Conservation agriculture and
682 smallholder farming in Africa: The heretics' view, *F. Crop. Res.*, 114, 23–34,
683 <https://doi.org/10.1016/j.fcr.2009.06.017>, 2009.

684 Giller, K. E., Andersson, J. A., Corbeels, M., Kirkegaard, J., Mortensen, D., Erenstein, O., and
685 Vanlauwe, B.: Beyond conservation agriculture, *Front. Plant Sci.*, 6, 1–14,
686 <https://doi.org/10.3389/fpls.2015.00870>, 2015.

687 von Haden, A. C., Yang, W. H., and DeLucia, E. H.: Soils' dirty little secret: Depth-based
688 comparisons can be inadequate for quantifying changes in soil organic carbon and other
689 mineral soil properties, *Glob. Chang. Biol.*, 26, 3759–3770, <https://doi.org/10.1111/gcb.15124>,
690 2020.

691 Han, L., Sun, K., Jin, J., and Xing, B.: Some concepts of soil organic carbon characteristics and
692 mineral interaction from a review of literature, *Soil Biol. Biochem.*, 94, 107–121,
693 <https://doi.org/10.1016/j.soilbio.2015.11.023>, 2016.

694 Harrison, R. B., Footen, P. W., Harrison, R. B., Footen, P. W., and Strahm, B. D.: Deep Soil
695 Horizons: Contribution and Importance to Soil Carbon Pools and in Assessing Whole-
696 Ecosystem Response to Management and Global Change, *For. Sci.*, 57, 67–76, 2011.

697 Hirte, J., Leifeld, J., Abiven, S., Oberholzer, H. R., and Mayer, J.: Below ground carbon inputs
698 to soil via root biomass and rhizodeposition of field-grown maize and wheat at harvest are
699 independent of net primary productivity, *Agric. Ecosyst. Environ.*, 265, 556–566,
700 <https://doi.org/10.1016/j.agee.2018.07.010>, 2018.

701 Hirte, J., Walder, F., Hess, J., Büchi, L., Colombi, T., van der Heijden, M. G., and Mayer, J.:
702 Enhanced root carbon allocation through organic farming is restricted to topsoils, *Sci. Total*
703 *Environ.*, 755, 143551, <https://doi.org/10.1016/j.scitotenv.2020.143551>, 2021.

704 IPCC: 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas
705 Inventories., 1–19 pp., 2019.

706 Jephitha, G., Jefline, K., Willis, G., and Justice, N.: Carbon stock, aggregate stability and
707 hydraulic properties of soils under tillage, crop rotation and mineral fertiliser application in
708 sub-humid Zimbabwe, *Heliyon*, 9, <https://doi.org/10.1016/j.heliyon.2023.e15846>, 2023.

709 Jones, D. L., Nguyen, C., and Finlay, R. D.: Carbon flow in the rhizosphere : carbon trading at
710 the soil – root interface, *Plant Soil*, 321, 5–33, <https://doi.org/10.1007/s11104-009-9925-0>,
711 2009.

712 Kahn, B. A. and Schroeder, J. L.: Root Characteristics and Seed Yields of Cowpeas Grown
713 with and without Added Nitrogen Fertilizer, *Hortscience A Publ. Am. Soc. Horticultural Sci.*,
714 34, 1238–1239, <https://doi.org/10.21273/hortsci.34.7.1238>, 1999.

715 Kassam, A., Friedrich, T., and Derpsch, R.: Global spread of Conservation Agriculture, *Int. J.*
716 *Environ. Stud.*, 76, 29–51, <https://doi.org/10.1080/00207233.2018.1494927>, 2019.

- 717 Kell, D. B.: Breeding crop plants with deep roots: their role in sustainable carbon , nutrient and
718 water sequestration, *Ann. Bot.*, 108, 407–418, <https://doi.org/10.1093/aob/mcr175>, 2011.
- 719 Kimaro, A. A., Mpanda, M., Rioux, J., Aynekulu, E., Shaba, S., Thiong’o, M., Mutuo, P.,
720 Abwanda, S., Shepherd, K., Neufeldt, H., and Rosenstock, T. S.: Is conservation agriculture
721 ‘climate-smart’ for maize farmers in the highlands of Tanzania?, *Nutr. Cycl. Agroecosystems*,
722 105, 217–228, <https://doi.org/10.1007/s10705-015-9711-8>, 2016.
- 723 Kimiti, J. M.: Influence of integrated soil nutrient management on cowpea root growth in the
724 semi-arid Eastern Kenya, *African J. Agric. Res.*, 6, 3084–3091,
725 <https://doi.org/10.5897/AJAR10.1023>, 2011.
- 726 Kodzwa, J. J., Gotosa, J., and Nyamangara, J.: Mulching is the most important of the three
727 conservation agriculture principles in increasing crop yield in the short term, under sub humid
728 tropical conditions in Zimbabwe, *Soil Tillage Res.*, 197, 104515,
729 <https://doi.org/10.1016/j.still.2019.104515>, 2020.
- 730 Koga, N. and Tsuji, H.: Effects of reduced tillage, crop residue management and manure
731 application practices on crop yields and soil carbon sequestration on an Andisol in northern
732 Japan, *Soil Sci. Plant Nutr.*, 55, 546–557, <https://doi.org/10.1111/j.1747-0765.2009.00385.x>,
733 2009.
- 734 Kopittke, P. M., Hernandez-Soriano, M. C., Dalal, R. C., Finn, D., Menzies, N. W., Hoeschen,
735 C., and Mueller, C. W.: Nitrogen-rich microbial products provide new organo-mineral
736 associations for the stabilization of soil organic matter, *Glob. Chang. Biol.*, 24, 1762–1770,
737 <https://doi.org/10.1111/gcb.14009>, 2018.
- 738 Kravchenko, A. N. and Guber, A. K.: Soil pores and their contributions to soil carbon
739 processes, *Geoderma*, 287, 31–39, <https://doi.org/10.1016/j.geoderma.2016.06.027>, 2017.
- 740 Kravchenko, A. N. and Robertson, G. P.: Whole-Profile Soil Carbon Stocks: The Danger of
741 Assuming Too Much from Analyses of Too Little, *Soil Sci. Soc. Am. J.*, 75, 235–240,
742 <https://doi.org/10.2136/sssaj2010.0076>, 2011.
- 743 Lal, R.: Residue management, conservation tillage and soil restoration for mitigating
744 greenhouse effect by CO₂-enrichment, *Soil Tillage Res.*, 43, 81–107, 1997.
- 745 Lal, R.: Sequestering carbon and increasing productivity by conservation agriculture, *J. Soil
746 Water Conserv.*, 70, 55A-62A, <https://doi.org/10.2489/jswc.70.3.55A>, 2015.
- 747 Lal, R.: Digging deeper: A holistic perspective of factors affecting soil organic carbon
748 sequestration in agroecosystems, *Glob. Chang. Biol.*, 24, 3285–3301,
749 <https://doi.org/10.1111/gcb.14054>, 2018.
- 750 Laub, M., Corbeels, M., Couëdel, A., Ndungu, S. M., Mucheru-muna, M. W., Mugendi, D.,
751 Nepalova, M., Waswa, W., Van de Broek, M., Vanlauwe, B., and Six, J.: Managing soil
752 organic carbon in tropical agroecosystems: evidence from four long-term experiments in
753 Kenya, *SOIL*, 9, 301–323, <https://doi.org/10.5194/soil-9-301-2023>, 2023.
- 754 Laura, van der P. K., Andy, R., Meagan, S., Francisco, C. J., Wallenstein, M. D., and Cotrufo,
755 M. F.: Addressing the soil carbon dilemma: Legumes in intensified rotations regenerate soil
756 carbon while maintaining yields in semi-arid dryland wheat farms., *Agric. Ecosyst. Environ.*,
757 330, <https://doi.org/10.1016/j.agee.2022.107906>, 2022.

758 Leal, O. A., Amado, T. J. C., Fiorin, J. E., Keller, C., Reimche, G. B., Rice, C. W., Nicoloso,
759 R. S., Bortolotto, R. P., and Schwalbert, R.: Linking cover crop residue quality and tillage
760 system to CO₂-C emission, soil C and N stocks and crop yield based on a long-term experiment,
761 *Agronomy*, 10, <https://doi.org/10.3390/agronomy10121848>, 2020.

762 Li, Y., Li, Z., Chang, S. X., Cui, S., Jagadamma, S., Zhang, Q., and Cai, Y.: Residue retention
763 promotes soil carbon accumulation in minimum tillage systems: Implications for conservation
764 agriculture, *Sci. Total Environ.*, <https://doi.org/10.1016/j.scitotenv.2020.140147>, 2020.

765 Liang, B. C., VandenBygaart, A. J., MacDonald, J. D., Cerkowniak, D., McConkey, B. G.,
766 Desjardins, R. L., and Angers, D. A.: Revisiting no-till's impact on soil organic carbon storage
767 in Canada, *Soil Tillage Res.*, 198, 104529, <https://doi.org/10.1016/j.still.2019.104529>, 2020.

768 Lorenz, K. and Lal, R.: The Depth Distribution of Soil Organic Carbon in Relation to Land
769 Use and Management and the Potential of Carbon Sequestration in Subsoil Horizons, *Adv.*
770 *Agron.*, 88, 35–66, [https://doi.org/10.1016/S0065-2113\(05\)88002-2](https://doi.org/10.1016/S0065-2113(05)88002-2), 2005.

771 Ma, S., He, F., Tian, D., Zou, D., Yan, Z., Yang, Y., Zhou, T., Huang, K., Shen, H., and Fang,
772 J.: Variations and determinants of carbon content in plants: A global synthesis, *Biogeosciences*,
773 15, 693–702, <https://doi.org/10.5194/bg-15-693-2018>, 2018.

774 Malepfane, N. M., Muchaonyerwa, P., Hughes, J. C., and Zengeni, R.: Land use and site effects
775 on the distribution of carbon in some humic soil profiles of KwaZulu-Natal, South Africa,
776 *Heliyon*, 8, <https://doi.org/10.1016/j.heliyon.2021.e08709>, 2022.

777 Mapanda, F., Mupini, J., Wuta, M., Nyamangara, J., and Rees, R. M.: A cross-ecosystem
778 assessment of the effects of land cover and land use on soil emission of selected greenhouse
779 gases and related soil properties in Zimbabwe, *Eur. J. Soil Sci.*, 61, 721–733,
780 <https://doi.org/10.1111/j.1365-2389.2010.01266.x>, 2010.

781 Mbanyele, V., Mtambanengwe, F., Nezomba, H., Groot, J. C. J., and Mapfumo, P.:
782 Comparative short-term performance of soil water management options for increased
783 productivity of maize-cowpea intercropping in semi-arid Zimbabwe, *J. Agric. Food Res.*, 5,
784 100189, <https://doi.org/10.1016/j.jafr.2021.100189>, 2021.

785 Mhlanga, B., Ercoli, L., Pellegrino, E., Onofri, A., and Thierfelder, C.: The crucial role of
786 mulch to enhance the stability and resilience of cropping systems in southern Africa, *Agron.*
787 *Sustain. Dev.*, 41, <https://doi.org/10.1007/s13593-021-00687-y>, 2021.

788 Mhlanga, B., Pellegrino, E., Thierfelder, C., and Ercoli, L.: Conservation agriculture practices
789 drive maize yield by regulating soil nutrient availability, arbuscular mycorrhizas, and plant
790 nutrient uptake, *F. Crop. Res.*, 277, 108403, <https://doi.org/10.1016/j.fcr.2021.108403>, 2022a.

791 Mhlanga, B., Ercoli, L., Thierfelder, C., and Pellegrino, E.: Conservation agriculture practices
792 lead to diverse weed communities and higher maize grain yield in Southern Africa, *F. Crop.*
793 *Res.*, 289, 108724, <https://doi.org/10.1016/j.fcr.2022.108724>, 2022b.

794 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A.,
795 Chaplot, V., Chen, Z. S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong,
796 S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke,
797 S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I.,
798 Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C. C., Vågen, T. G., van Wesemael, B.,
799 and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86,

800 <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.

801 Mtambanengwe, F., Mapfumo, P., and Kirchmann, H.: Decomposition of organic matter in soil
802 as influenced by texture and pore size distribution, in: *Managing Nutrient Cycles to Sustain*
803 *Soil Fertility in Sub-Saharan Africa*, edited by: Bationo, A., Academy Science Publishers and
804 TSBF CIAT, Nairobi, 261–276, 2004.

805 Nyamangara, J., Masvaya, E. N., Tirivavi, R., and Nyengerai, K.: Effect of hand-hoe based
806 conservation agriculture on soil fertility and maize yield in selected smallholder areas in
807 Zimbabwe, *Soil Tillage Res.*, 126, 19–25, <https://doi.org/10.1016/j.still.2012.07.018>, 2013.

808 Patra, S., Julich, S., Feger, K. H., Jat, M. L., Sharma, P. C., and Schwärzel, K.: Effect of
809 conservation agriculture on stratification of soil organic matter under cereal-based cropping
810 systems, *Arch. Agron. Soil Sci.*, 65, 2013–2028,
811 <https://doi.org/10.1080/03650340.2019.1588462>, 2019.

812 Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., and Smith, P.: Climate-smart
813 soils, *Nature*, 532, 49–57, <https://doi.org/10.1038/nature17174>, 2016.

814 Powlson, D. S., Stirling, C. M., Jat, M. L., Gerard, B. G., Palm, C. A., Sanchez, P. A., and
815 Cassman, K. G.: Limited potential of no-till agriculture for climate change mitigation, *Nat.*
816 *Clim. Chang.*, 4, 678–683, <https://doi.org/10.1038/nclimate2292>, 2014.

817 Powlson, D. S., Stirling, C. M., Thierfelder, C., White, R. P., and Jat, M. L.: Does conservation
818 agriculture deliver climate change mitigation through soil carbon sequestration in tropical agro-
819 ecosystems ?, *"Agriculture, Ecosyst. Environ.*, 220, 164–174,
820 <https://doi.org/10.1016/j.agee.2016.01.005>, 2016.

821 Rumpel, C., Chabbi, A., and Marschner, B.: Carbon Storage and Sequestration in Subsoil
822 Horizons: Knowledge, Gaps and Potentials, in: *Recarbonization of the Biosphere: Ecosystems*
823 *and the Global Carbon Cycle*, edited by: Lal, R., Lorenz, K., Hüttl, R. F., Schneider, B. U., and
824 Von Braun, J., Springer Science+Business Media B.V, 445–464, [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-94-007-978-94-007-4159-1)
825 [94-007-4159-1](https://doi.org/10.1007/978-94-007-4159-1), 2012.

826 Sanaullah, M., Chabbi, A., Maron, P. A., Baumann, K., Tardy, V., Blagodatskaya, E.,
827 Kuzyakov, Y., and Rumpel, C.: How do microbial communities in top- and subsoil respond to
828 root litter addition under field conditions?, *Soil Biol. Biochem.*, 103, 28–38,
829 <https://doi.org/10.1016/j.soilbio.2016.07.017>, 2016.

830 Shumba, A., Dunjana, N., Nyamasoka, B., Nyamugafata, P., Madyiwa, S., and Nyamangara,
831 J.: Maize (*Zea mays*) yield and its relationship to soil properties under integrated fertility,
832 mulch and tillage management in urban agriculture, *South African J. Plant Soil*, 1–10,
833 <https://doi.org/10.1080/02571862.2019.1678686>, 2020.

834 Shumba, A., Chikowo, R., Thierfelder, C., Corbeels, M., Six, J., and Cardinael, R.:
835 Conservation agriculture increases soil organic carbon stocks but not soil CO₂ efflux in two
836 8-year-old experiments in Zimbabwe, *EGUsphere*, 2023, 1–40,
837 <https://doi.org/10.18167/DVN1/VPOCHN>, 2023a.

838 Shumba, A., Chikowo, R., Corbeels, M., Six, J., Thierfelder, C., and Cardinael, R.: Long-term
839 tillage, residue management and crop rotation impacts on N₂O and CH₄ emissions on two
840 contrasting soils in sub-humid Zimbabwe, *Agric. Ecosyst. Environ.*, 341,
841 <https://doi.org/10.1016/j.agee.2022.108207>, 2023b.

842 Singh, G., Schoonover, J. E., Williard, K. W. J., Kaur, G., and Crim, J.: Carbon and Nitrogen
843 Pools in Deep Soil Horizons at Different Landscape Positions, *Soil Sci. Soc. Am. J.*, 82, 1512–
844 1525, <https://doi.org/10.2136/sssaj2018.03.0092>, 2018.

845 Six, J., Paustian, K., and Elliott, E.: Aggregate and soil organic matter dynamics under
846 conventional and no-tillage systems, *Soil Sci. Soc. Am. J.*, 63, 1350–1358, 1999.

847 Six, J., Elliott, E. T., and Paustian, K.: Soil macroaggregate turnover and microaggregate
848 formation: A mechanism for C sequestration under no-tillage agriculture, *Soil Biol. Biochem.*,
849 32, 2099–2103, [https://doi.org/10.1016/S0038-0717\(00\)00179-6](https://doi.org/10.1016/S0038-0717(00)00179-6), 2000.

850 Six, J., Conant, R. T., Paul, E. A., and Paustian, K.: Stabilization mechanisms of soil organic
851 matter : Implications for C-saturation of soils, 155–176, 2002.

852 Sun, W., Canadell, J. G., Yu, L. L., Yu, L. L., Zhang, W., Smith, P., Fischer, T., and Huang,
853 Y.: Climate drives global soil carbon sequestration and crop yield changes under conservation
854 agriculture, *Glob. Chang. Biol.*, 26, 3325–3335, <https://doi.org/10.1111/gcb.15001>, 2020.

855 Swanepoel, C. M., van der Laan, M., Weepener, H. L., du Preez, C. C., and Annandale, J. G.:
856 Review and meta-analysis of organic matter in cultivated soils in southern Africa, *Nutr. Cycl.*
857 *Agroecosystems*, 104, 107–123, <https://doi.org/10.1007/s10705-016-9763-4>, 2016.

858 Swanepoel, C. M., Rötter, R. P., van der Laan, M., Annandale, J. G., Beukes, D. J., du Preez,
859 C. C., Swanepoel, L. H., van der Merwe, A., and Hoffmann, M. P.: The benefits of conservation
860 agriculture on soil organic carbon and yield in southern Africa are site-specific, *Soil Tillage*
861 *Res.*, 183, 72–82, <https://doi.org/10.1016/j.still.2018.05.016>, 2018.

862 Thierfelder, C. and Mhlanga, B.: Short-term yield gains or long-term sustainability? – a
863 synthesis of Conservation Agriculture long-term experiments in Southern Africa, *Agric.*
864 *Ecosyst. Environ.*, 326, 107812, <https://doi.org/10.1016/j.agee.2021.107812>, 2022.

865 Thierfelder, C. and Wall, P. C.: Effects of conservation agriculture techniques on infiltration
866 and soil water content in Zambia and Zimbabwe, *Soil Tillage Res.*, 105, 217–227,
867 <https://doi.org/10.1016/j.still.2009.07.007>, 2009.

868 Thierfelder, C. and Wall, P. C.: Effects of conservation agriculture on soil quality and
869 productivity in contrasting agro-ecological environments of Zimbabwe, *Soil Use Manag.*, 28,
870 209–220, <https://doi.org/10.1111/j.1475-2743.2012.00406.x>, 2012.

871 Thierfelder, C., Matemba-Mutasa, R., and Rusinamhodzi, L.: Yield response of maize (*Zea*
872 *mays* L.) to conservation agriculture cropping system in Southern Africa, *Soil Tillage Res.*,
873 146, 230–242, <https://doi.org/10.1016/j.still.2014.10.015>, 2015.

874 Thierfelder, C., Chivenge, P., Mupangwa, W., Rosenstock, T. S., Lamanna, C., and Eyre, J. X.:
875 How climate-smart is conservation agriculture (CA)? – its potential to deliver on adaptation,
876 mitigation and productivity on smallholder farms in southern Africa, *Food Secur.*, 9, 537–560,
877 <https://doi.org/10.1007/s12571-017-0665-3>, 2017.

878 Thierfelder, C., Baudron, F., Setimela, P., Nyagumbo, I., Mupangwa, W., Mhlanga, B., Lee,
879 N., and Gérard, B.: Complementary practices supporting conservation agriculture in southern
880 Africa. A review, *Agron. Sustain. Dev.*, 38, <https://doi.org/10.1007/s13593-018-0492-8>, 2018.

881 Thorup-Kristensen, K., Halberg, N., Nicolaisen, M., Olesen, J. E., Crews, T. E., Hinsinger, P.,
882 Kirkegaard, J., Pierret, A., and Dresbøll, D. B.: Digging Deeper for Agricultural Resources,

883 the Value of Deep Rooting, *Trends Plant Sci.*, 25, 406–417,
884 <https://doi.org/10.1016/j.tplants.2019.12.007>, 2020.

885 Villarino, S. H., Pinto, P., Jackson, R. B., and Piñeiro, G.: Plant rhizodeposition : A key factor
886 for soil organic matter formation in stable fractions, *Sci. Adv.*, 7, 1–14,
887 <https://doi.org/10.1126/sciadv.abd3176>, 2021.

888 Virk, A. L., Lin, B. J., Kan, Z. R., Qi, J. Y., Dang, Y. P., Lal, R., Zhao, X., and Zhang, H. L.:
889 Simultaneous effects of legume cultivation on carbon and nitrogen accumulation in soil, *Adv.*
890 *Agron.*, 171, 75–110, <https://doi.org/10.1016/bs.agron.2021.08.002>, 2022.

891 Wendt, J. W. and Hauser, S.: An equivalent soil mass procedure for monitoring soil organic
892 carbon in multiple soil layers, *Eur. J. Soil Sci.*, 64, 58–65, <https://doi.org/10.1111/ejss.12002>,
893 2013.

894 Yang, L., Luo, Y., Lu, B., Zhou, G., Chang, D., Gao, S., Zhang, J., Che, Z., and Cao, W.: Long-
895 term maize and pea intercropping improved subsoil carbon storage while reduced greenhouse
896 gas emissions, *Agric. Ecosyst. Environ.*, 349, 108444,
897 <https://doi.org/10.1016/J.AGEE.2023.108444>, 2023.

898 Yost, J. L. and Hartemink, A. E.: How deep is the soil studied – an analysis of four soil science
899 journals, *Plant Soil*, 452, 5–18, <https://doi.org/10.1007/s11104-020-04550-z>, 2020.

900