

Estimation of anthropogenic and volcanic SO₂ emissions from satellite data in the presence of snow/ice on the ground

Vitali E. Fioletov¹, Chris A. McLinden¹, Debora Griffin¹, Nikolay Krotkov², Can Li^{2,3}, Joanna Joiner², Nicolas Theys⁴, and Simon Carn⁵

5 ¹Air Quality Research Division, Environment and Climate Change Canada, Toronto, ON, Canada

²Atmospheric Chemistry and Dynamics Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA

³Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD, USA

⁴Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

10 ⁵Department of Geological and Mining Engineering and Sciences, Michigan Technological University, Houghton, MI 49931, USA

Correspondence to: Vitali.Fioletov@outlook.com or Vitali.Fioletov@ec.gc.ca

Abstract. Early versions of satellite nadir-viewing UV SO₂ data products did not explicitly account for the effects of snow/ice on retrievals. Snow covered terrain, with its high reflectance in the UV, typically enhances satellite sensitivity to boundary layer pollution. However, a significant fraction of high-quality cloud-free measurements over snow is currently excluded from
15 analyses. This leads to increased uncertainties of satellite emissions estimates and potential seasonal biases due to the lack of data in winter months for some high-latitude sources. In this study, we investigated how OMI and TROPOMI satellite SO₂ measurements over snow-covered surfaces can be used to improve the annual emissions reported in our SO₂ emissions catalogue (version 2, Fioletov et al., 2023). Only 100 out of 759 sources listed in the catalogue have 10% or more of the observations over snow. However, for 40 high-latitude sources, more than 30% of measurements suitable for emission
20 calculations were made over snow-covered surfaces. For example, in the case of Norilsk, the world's largest SO₂ point source, annual emissions estimates in the SO₂ catalogue were based only on 3-4 summer months, while addition of data for snow conditions extends that period to 7 months. Emissions in the SO₂ catalogue were based on satellite measurements of SO₂ slant column densities (SCDs) that were converted to vertical column densities (VCDs) using site-specific clear-sky air mass factors (AMFs), calculated for snow-free conditions. The same approach was applied to measurements with snow on the ground
25 whereby a new set of constant, site-specific, clear sky with snow AMFs was created, and these were applied to the measured SCDs. Annual emissions were then estimated for each source considering (i) only clear-sky snow-free days, (ii) only clear-sky with snow days and (iii) a merged dataset (snow and snow-free conditions). For individual sources, the difference between emissions estimated for snow and snow-free conditions is within $\pm 20\%$ for three quarters of smelters and oil and gas sources, and with practically no systematic bias. This is excellent consistency given that there is typically a factor of 3-5 difference
30 between AMFs for snow and snow-free conditions. For coal-fired power plants, however, emissions estimated for snow conditions are on average 25% higher than for snow-free conditions; this difference is likely real and due to larger production (consumption of coal) and emissions in wintertime.

1 Introduction

Sulfur dioxide (SO₂) is a major air pollutant that plays an important role in atmospheric chemistry, contributes to aerosol formation, adversely affects the environment and human health, and impacts climate (Fischer et al., 2019; Hansell and Oppenheimer, 2010; Longo et al., 2010; Robock, 2000). SO₂ also leads to acid deposition that affects terrestrial ecosystems (Dentener et al., 2006; Hutchinson and Whitby, 1977; Vet et al., 2014; Fedkin et al., 2018). Most of atmospheric SO₂ is a result of human activity and the major sources of anthropogenic emissions are coal-burning power plants, oil refineries and smelters (Klimont et al., 2013; Smith et al., 2011), while volcanoes are the primary natural source of SO₂ (Carn et al., 2017; Oppenheimer et al., 2011). A proper account for SO₂ content in the atmosphere is necessary for climate, weather, and air quality models (e.g., Liu et al., 2018; Stenchikov et al., 2021; Ukhov et al., 2020). Such models require information about SO₂ emissions from anthropogenic and natural (mostly volcanic) sources.

The history of satellite SO₂ measurements in the Earth's atmosphere goes back to 1982, when volcanic SO₂ from the El Chichón eruption in 1982 was first retrieved from measurements of the Total Ozone Mapping Spectrometer (TOMS) and the Solar Backscattered Ultraviolet (SBUV) instruments on the Nimbus 7 satellite (Krueger, 1983; McPeters et al., 1984). Over the last two decades, several satellite UV-visible instruments have been used for the monitoring of atmospheric SO₂. They include the Global Ozone Monitoring Experiment (GOME) in 1995-2011 (Eisinger and Burrows, 1998; Khokhar et al., 2008); the SCanning Imaging Absorption spectroMeter for Atmospheric CHartographY (SCIAMACHY) in 2002-2012 (Bovensmann et al., 1999; Lee et al., 2009); the Global Ozone Monitoring Experiment-2 (GOME-2) instrument, 2006-present, (Bramstedt et al., 2004; Callies et al., 2000; Nowlan et al., 2011; Rix et al., 2012), and the Ozone Mapping and Profiler Suite (OMPS), 2011- present (Zhang et al., 2017). However, the spatial resolution of these instruments was not very high, above about 40 km, that made it difficult to monitor small emission sources.

Satellite SO₂ measurements with high spatial resolution that is sufficient to detect hundreds of SO₂ emissions sources started with the launch of the Dutch-Finnish Ozone Monitoring Instrument (OMI) (Levelt et al., 2018, 2006) on NASA's Earth Observing System (EOS) -Aura spacecraft (Schoeberl et al., 2006) in 2004. OMI has the spatial resolution as fine as 13 × 24 km² at nadir and is able to provide daily, nearly global maps of SO₂ VCDs (Krotkov et al., 2016; Li et al., 2020). The TROPOspheric Monitoring Instrument (TROPOMI) (Theys et al., 2017) on the ESA Copernicus Sentinel-5 Precursor (S-5P) spacecraft (Veefkind et al., 2012) launched in 2017 has even higher spatial resolution, 5.5 × 3.5 km² and also is able to provide daily global coverage.

Satellite measurements of SO₂ are available in the form of vertical column density (VCD) that represent the total number of molecules in a vertical column per a unit of area. VCDs and SO₂ emissions estimated from them are used for multiple applications such as evaluation of long-term changes and trends on a regional and global scale in SO₂ VCDs (Krotkov et al., 2016) and surface concentrations (Kharol et al., 2017) and for assessment of the efficiency of industrial clean technology solutions in reducing of SO₂ emissions (Ialongo et al., 2018; McLinden et al., 2020). Such measurements are also an important source of information about volcanic SO₂ (Carn et al., 2013; Inness et al., 2022; Krueger et al., 1995; McCormick et al., 2013)

and volcanic SO₂ is used to estimate volcanic carbon dioxide (CO₂) fluxes (Fischer et al., 2019). Satellite SO₂ VCDs are also used to estimate emissions from anthropogenic SO₂ sources (Fioletov et al., 2013, 2015; McLinden et al., 2012, 2020; Qu et al., 2019) and to update the available emissions inventories (Liu et al., 2018; Ukhov et al., 2020).

70 A global catalogue of large SO₂ sources (hereafter referred to as the “SO₂ catalogue”) and their emissions estimated from satellite SO₂ VCD observations was developed (Fioletov et al., 2016; McLinden et al., 2016). The SO₂ catalogue has been updated to version 2 recently (Fioletov et al., 2023). This version 2 covers the 2005-2022 period and contains information on a total of 759 point sources emitting from about 10 kt yr⁻¹ to more than 4000 kt yr⁻¹ of SO₂. The SO₂ catalogue also includes information about the country and source type, so data can be grouped and summed by them. The SO₂ catalogue contains annual emission estimates for 106 volcanoes, 477 power plants, 74 smelters, and 102 sources related to the oil and gas industry.
75 Satellite data from version 2 of OMI and OMPS as well as from a recent version of the SO₂ TROPOMI dataset were used to estimate emissions for the version 2 SO₂ catalogue.

Early versions of satellite SO₂ data products were based on retrieval algorithms that estimated SO₂ slant column density (SCD) first and then converted it to VCD using a single conversion factor that corresponds to typical snow-free surface conditions (Khokhar et al., 2005; Lee et al., 2008; Thomas et al., 2005). For example, in the original OMI algorithm, all SCD
80 values were then converted to planetary boundary layer (PBL) VCDs by applying a constant AMF = 0.36 that was appropriate for anthropogenic pollution in the eastern US in summer (Krotkov et al., 2008, 2006; Li et al., 2013). Subsequently, PBL AMFs were calculated using a climatological SO₂ profile and information about surface reflectivity, solar zenith angle, and other characteristics (Lee et al., 2009; Li et al., 2020c; Theys et al., 2017). For example, Lee et al. (2009) used shapes of SO₂ profiles determined from a global 3-D model of tropospheric chemistry (GEOS-Chem) along with a surface reflectivity
85 climatology, and information about cloud cover and aerosols. McLinden et al. (2014) calculated site-specific AMFs based on information about the local climatological surface reflectivity, aerosols, and boundary layer height assuming that all SO₂ is in the boundary layer. Based on that approach, estimated emissions reported in the Versions 1 and 2 SO₂ catalogues were calculated using a single site-specific AMF that corresponds to measurements under snow-free conditions only (Fioletov et al., 2016, 2023). Thus, potentially high-quality measurements collected under snow conditions were excluded from the
90 emission estimates increasing the uncertainties of the estimates and introducing potential biases due to the lack of data in winter months for some high-latitude sources.

More recently, NASA version 2 (V2) of the principal component analysis (PCA) algorithm for OMI/OMPS data processing used updated a priori SO₂ vertical profiles that were calculated using the Goddard Earth Observing System Version 5 (GEOS-5) global model and the corresponding OMI pixel-specific AMFs (Li et al., 2020). Thus, V2 SO₂ data included
95 VCDs calculated over snow and ice using OMI-measured Lambertian Equivalent scene Reflectivity (LER) and are in a better agreement with summertime values (Li et al., 2020). In this study, we consider two data sets of satellite SO₂ measurements taken under snow on the ground conditions: (1) the NASA OMI V2 SO₂ VCD dataset and (2) the dataset based on OMI and TROPOMI SCDs converted to VCDs using two site-specific time-independent AMF values calculated for each source: one for conditions with snow and another for conditions without snow on the ground. Both data sets are used to calculate SO₂ point

100 source emissions in order to improve the existing global catalogue of large SO₂ sources and their emissions (Fioletov et al., 2023). The study is focused on OMI data analysis because of its 18-year long record of observations with spatial resolution suitable for emissions estimates for hundreds of sources. TROPOMI data have a superior spatial resolution, but a much shorter record and were mostly used for some illustrations. We did not use OMPS data in this study because the number of measurements is not large enough to reliably estimate emissions under snow conditions for most of the sources.

105 This article is organized as follows: the data sets and AMF values are described in **Section 2**. **Section 3** discusses the differences between emissions estimated for conditions with and without snow on the ground and provides an overview of the estimated emissions using all data (with and without snow on the ground). **Section 4** summarizes the findings of the study.

2 Data sets and emissions estimates

The OMI and TROPOMI Level 2 SCD data sets, data selection criteria, and the emission calculation algorithm used in this study are identical to those in Fioletov et al. (2023) and we just briefly review them here. However, we included measurements taken in presence of snow on the ground that were previously excluded from the emissions calculations. The estimated annual emission rates are given in metric kilotonnes of SO₂ per year (kt y⁻¹) and the VCD values are given in Dobson Units (DU, 1 DU = 2.69•10¹⁶ molec•cm⁻²).

2.1 OMI data

115 OMI was launched on NASA's Earth Observing System (EOS - Chemistry) Aura satellite on 15 July 2004 (Schoeberl et al., 2006). Aura is on a sun-synchronous polar orbit that crosses the equator at about 13:45 local mean time. The OMI spatial resolution is 13 km × 24 km at nadir and up to 30 km × 100 km at the swath edges (de Graaf et al., 2016). The OMI wide swath (~2600 km) is divided into 60 cross-track positions; the first and last 10 cross-track positions were excluded from our analysis to limit the across-track pixel width from 24 km to about 40 km. About half of OMI pixels have been affected by a field-of-view blockage and stray light (the so-called "row anomaly") since 2007 (Levelt et al., 2018), and they were also excluded from our analysis.

In this study, we used the V2 OMI PCA SO₂ (OMSO2 V2) data (Li et al., 2020) aimed at anthropogenic SO₂ pollution for the period 2005-2022. Only measurements with solar zenith angles (SZA) less than 70° are used. In the V2, a priori SO₂ vertical profiles are based on average GEOS-5 global model simulations (72 vertical layers, 0.5° latitude by 0.667° longitude resolution) for the period 2004–2014. Monthly climatological profiles are calculated for each grid cell and then used as a priori profiles in the SO₂ VCD retrievals. There are two potential issues in this approach. First, the period may not reflect the present emission levels as SO₂ emission sources, and their emission levels are changing over time. Second, such climatological profiles represent "average" conditions within the model grid cell, that could be different from plume profiles near pollution sources (i.e., the signals near the emissions sources are diluted by the relatively coarse spatial resolution as compared with the resolution of OMI pixels). While such profiles near the source assume high concentrations in the boundary layer, if we move

away from the source, the model simulated SO₂ profile is more heavily weighted towards the free troposphere due to less emissions in the boundary layer and chemical loss and deposition in the boundary layer. As a result, the Air Mass Factors (AMF = SCD/VCD) depend on the distance from the source: they are lower over the source and getting larger further away from it. Both of these issues affect the emissions estimates based on OMSO₂ V2 VCD data.

135 Another data set of VCDs used for emission estimates in this study is also based on OMSO₂ V2 data but uses a different approach to calculate VCDs. OMSO₂ V2 also provides SO₂ SCDs produced by spectral fitting using SO₂ absorption cross sections. In the SO₂ catalogue, these SCDs were converted to VCD using a constant over time site-specific air mass factor (AMF_{source} = SCD/VCD) calculated at Environment and Climate Change Canada (ECCC) as discussed in Section 2.4 (McLinden et al., 2014). The same approach is used here; however, two AMFs are calculated for each source: for snow and
140 snow-free conditions. We will refer to these AMFs as to “ECCC AMFs”.

A +10% empirical correction is applied to OMI VCDs for the version 2 SO₂ catalogue (Fioletov et al., 2023). This is done to match the values from the original SO₂ catalogue, that were validated against available emission inventories. This is a relatively small correction compared to uncertainties in AMF and lifetime (18% and 35%, respectively (Fioletov et al., 2023)).

2.2 TROPOMI data

145 While this study is largely focused on OMI data, we also use TROPOMI measurements for some illustrations and comparisons. TROPOMI was launched on the ESA Copernicus Sentinel 5-Precursor satellite on October 13, 2017 (Veefkind et al., 2012). The instrument contains a UV spectrometer that is used to retrieve SO₂. The TROPOMI detector has 450 cross-track positions. We excluded the first and last 20 of them since their measurements have a relatively high noise level (Fioletov et al., 2020). The TROPOMI spatial resolution for the centre of the swath was approximately 7 km × 3.5 km until August 6, 2019 when it
150 was improved to 5.5 km × 3.5 km.

Here, we use TROPOMI data processed by the Covariance-Based Retrieval Algorithm (COBRA) (Theys et al., 2021) for the period from April 1, 2018, to December 31, 2022. Similar to the OMI analysis, only data with SZA < 70° are used for the study. Note that TROPOMI and OMI SO₂ SCD retrieval algorithms use different SO₂ cross-section temperature (203K and 293K respectively). As a result, reported TROPOMI SCDs are 22% lower than they would be retrieved using cross-section for
155 293K (see Theys et al., 2017, their Figure 6). For this reason, we increase TROPOMI SCDs used this study by 22% (see also Fioletov et al., 2020; Theys et al., 2017). For TROPOMI, the same conversion of SCD to VCD with the same site-specific ECCC AMFs as for OMI is used.

2.3 Emissions estimates

The emission calculation algorithm was identical to that used to calculate emissions for the SO₂ catalogue (Fioletov et al.,
160 2015; 2016; 2023). It is based on the fitting algorithm to estimate the average total SO₂ mass at and downwind of the source in order to derive emissions, assuming a constant lifetime. The pixel-based SO₂ VCDs were fit to a two-dimensional exponentially-modified Gaussian plume dispersion function (Fioletov et al., 2016) that depends on the wind speed and

direction at the pixel location on the time of satellite overpass. The parameter of the fit represents the total SO₂ mass, emitted from the source. As we are interested in annual emissions, the fitting was done using all pixels near the source collected over one year. The exact fitting area depends on the wind speed and emission strength as described by Fioletov et al. (2016). The wind speed and direction are obtained from most recent ERA5 reanalysis wind data (C3S, 2017). A recent study shows that ERA5 data are reliable and have practically no bias on offshore and flat onshore locations, although wind uncertainties are greater at mountainous and coastal sites (Gualtieri, 2022). Available from ERA5, U- and V- (west-east and south-north, respectively) wind component data are grouped into 1 km-thick layers and the mean wind speed and direction are calculated for each level as done for the original catalogue. The winds for the layer that corresponds to the height of a source are used by the algorithm.

As in the SO₂ catalogue, the fitting function depends on two prescribed non-linear parameters: the plume width and SO₂ effective lifetime (or decay time, τ). The plume width is related to the size of satellite pixels and the spatial extent of the source plume, and it should not depend on the presence of snow. The same values as in the SO₂ catalogue were used for the plume width (20 km for OMI and 15 km for TROPOMI). We also used the same value of $\tau=6$ hours for the lifetime for both snow and snow-free conditions for all sources. The lifetime uncertainty is one of the major contributors to the overall uncertainty of the emission estimates. For snow-free conditions (i.e., for the SO₂ catalogue), it was estimated that for 80% of all sources τ is between 3 and 9.5 h with the mean of about 6 h (Fioletov et al., 2016). If τ is included as one of the fitting parameters, owing to its non-linear nature, the emission uncertainties may become too large, particularly for small sources. A single prescribed τ value reduced random uncertainties of emission estimates, although it potentially introduces a multiplicative bias in estimated values for some sources (Fioletov et al., 2016). For snow conditions, we tested the same value of $\tau=6$ hours and found that, on average, the estimated emissions have only a small (3-4%) bias compared to these for snow-free conditions (see also section 3.3). A single value for snow and snow-free conditions also substantially simplifies the algorithms because otherwise it is not clear how to treat days with partial snow cover around the source.

The chemical SO₂ lifetime is much longer than $\tau=6$ hours. It is 13-19 h in summer and about 50 h in winter according to in situ measurements and model calculations (Lee et al., 2011). This suggests that the used decay time $\tau = 6$ h is more related to the plume dispersion that brings SO₂ VCDs below the instrument sensitivity level, rather than to the SO₂ chemical destruction. If the dispersion is not very different in winter and summer, then the decay times could be also similar in the two seasons. Or, perhaps, the lifetime in the plume is substantially different from that under background conditions. It is difficult to obtain τ directly from the satellite SO₂ VCD data fitting, because the number of sources with snow conditions and the number of measurements for snow conditions is rather limited. Norilsk is probably the only place where such estimates can be reliably done where all three parameters of the fitting (see Fioletov et al., 2015) are estimated from the data. The calculations for Norilsk show τ values of $8.4 \text{ h} \pm 0.5 \text{ h}$ and $7.5 \text{ h} \pm 0.7 \text{ h}$ (2- σ intervals) for snow and snow-free conditions respectively, i.e., different by about 10%. As τ is a parameter of the fitting function, the overall impact of this difference in the τ value to the estimated emission is even less; the emissions for snow and snow-free calculated with these τ values are different by only

about 3.5%, and they are different by about 6% from the emission values calculated for the $\tau = 6$ h value using the standard SO₂ catalogue one-parameter algorithm. Estimates of τ using TROPOMI data show similar results: $\tau = 7.4 \pm 1.3$ h for snow and 6.2 ± 0.6 h for snow-free conditions in 2018-2022. This shows that an approximation of 6 h for the SO₂ lifetime is reasonable and ensures consistency throughout the emission estimates.

As part of the OMI level 1B to level 2 processing, before the SO₂ retrieval, cloud radiance fraction (CRF), cloud optical centroid pressure, and effective scene pressure are derived using the OMCLDRR algorithm (Joiner and Vasilkov, 2006; Vasilkov et al., 2008). This information is essential to correctly interpret the SO₂ SCDs and to calculate the AMF. OMI itself cannot discriminate between snow and clouds as both act as strong reflectors of UV/visible light. Hence, the OMI cloud algorithm uses external snow/ice flag from Near-real-time Ice and Snow Extent (NISE) dataset. Misdiagnosing the presence of snow, and the reflectivity of snow if it is present, can lead to errors in the cloud fraction and dependent products (see e.g., Vasilkov et al., 2010).

Emissions estimates were done for pixels with and without snow on the ground under “clear sky” conditions. For snow-free pixels, clear sky conditions were defined as those with CRF of less than 0.3. For snow-covered pixels, the clear sky conditions were determined from comparisons of terrain and effective scene pressures (ESP). In OMI version 2 data, the ESP from the OMCLDRR algorithm is compared with the terrain pressure. If the difference is within 50 hPa, the pixels was considered as cloud-free and CRF was set to zero (Li et al., 2020). If it is greater than 100 hPa, then the pixel was likely cloudy, and CRF was set to one. For pixels having scene and terrain pressure differences between 50 and 100 hPa, “unambiguous cloud detection is not possible” (Li et al., 2020), but CRF was set to 0. We tested both the 50 and 100 hPa cut-off limits for determination of the clear sky conditions for several sources. It appears that there is no systematic difference in estimated emissions between the two limits, however the number of clear sky pixels is 20%-40% larger for the 100 hPa limit than for the 50 hPa limit making the emission uncertainties smaller. For this reason, we used the 100 hPa limit in this study.

Similarly, TROPOMI COBRA data contain information about the terrain and cloud top pressure taken from the TROPOMI cloud product (Loyola et al., 2018). In case of TROPOMI, we also applied the condition that the difference between the terrain and cloud top pressure should be less than 100 hPa. It should be noted that OMI scene pressure may be different from the TROPOMI cloud top pressure data product, however the comparison of these two data products is outside the scope of this paper.

2.4. Site-specific AMFs

As in the previous studies (Fioletov et al., 2016; McLinden et al., 2016), pre-calculated site-specific and time-independent AMF values were used to calculate the SO₂ VCDs and emissions for each source location. Here, two different AMF values are calculated for each emission source: one for snow-free conditions (as in the previous publications) and another for the snow on the ground conditions. The AMF calculation method is based on a general approach from McLinden et al. (2014) modified as described in Fioletov et al. (2022). The AMFs were calculated for a subsample of OMI observations within 100 km of the emission source coordinates. Every 100th observation from every 3rd year were used that yields several thousand

230 observations and is sufficient to represent conditions (cloud fraction, viewing and solar geometry, and seasonal sampling) for a given source. As before, AMFs were calculated using the MODIS MCD43C3 (v6.1) reflectance data (Schaaf et al., 2002). MODIS LER (43C3) albedo at 477 nm is used, with albedo calculated separately for snow and snow-free conditions. For snow-free conditions, an annually varying monthly mean was used considering only snow free MODIS pixels and the MODIS 477 nm albedo is mapped to SO₂ wavelengths using OMI 0.5 degree climatology at ratio of 342 nm to 477 nm albedos (see
235 McLinden et al., 2014, sec 3.2.2). For snow conditions, the 477 nm albedo is used as is.

For snow conditions, a single snow albedo map was created by considering all years and months using pixels with a snow flag of 1. The final, site-specific AMFs were the average over these individual AMFs, separating them into snow and snow-free categories. Note that the SO₂ profile is estimated based on the elevation of the source and a climatological boundary-layer height (as a function of latitude, longitude, month, and UTC hour) from von Engel and Teixeira, (2013), assuming a
240 constant mixing ratio in the boundary layer and zero elsewhere.

The snow cover information was obtained from the Interactive Multisensor Snow and Ice Mapping System measurements (IMS) (Helfrich et al., 2007) which has been shown to be one of the best for identifying snow on the ground (Cooper et al. 2018). While the OMI datafiles contain information about snow/ice obtained from the NISE database, this product is known to frequently misidentify snow as no-snow when the layer is thin (McLinden et al., 2014; Cooper et al.,
245 2018). Thus, the IMS snow data were used for both OMI (24-km data product) and TROPOMI (4 km data product). The use of a single, consistent product is preferred to avoid possible sampling differences when data from other instruments are used for emission calculations for the merged SO₂ catalogue.

Although the NASA VCD retrievals do not explicitly calculate AMFs, we obtained an effective AMF by taking the ratio of SCD to VCD in order to better understand the differences between emissions estimates from the two data sets. “NASA
250 AMFs” were calculated for each source as a median value of the SCD to VCD ratios for all pixels within a 25 km radius around the source. As mentioned in Section 2.1, the NASA AMFs depend on the distance from the source: they are lower over the source and getting larger further away from it. We used a small area around the source with a 25 km radius to have a better representation for AMF in the plume. Further reduction of the radius does not change the AMF value but increases the estimate uncertainty. For the ECCC AMFs, the profiles are determined by the PBL height and therefore is nearly constant within 200-
255 300 km around an emission source used for satellite data fitting.

It should be also mentioned that snow-covered pixels were included in the previous OMI and TROPOMI NO₂ data products and their analysis using an approach that is similar to presented in this study (O'Byrne et al., 2010; van der A et al., 2020; van Geffen et al., 2022).

3. Results

260 3.1 Source statistics

At present, there are 759 sources of emissions listed in the SO₂ catalogue (Fioletov et al., 2023), but many of them are located at low latitudes where snow conditions are very rare. **Figure 1** shows a map of the sources where OMI measurements are available, in the presence of snow on the ground. The size of the dots on the map is proportional to the fraction of measurements under snow conditions among all measurements suitable for emission estimates. Most of the industrial sources are in Russia, 265 Canada, Kazakhstan, and a region in north-west China. There are also several volcanic sources in Alaska and Kamchatka. The largest fraction, about 60%, is at four sites in northern Russia including Norilsk, the world's largest anthropogenic SO₂ point source. The fraction is less than 10% for most of the US and European sources. For a total of 100 sources the fraction is more than 10%, and we focus this study on the analysis of data from these sources. Note that the utilized snow cover information from IMS does not cover the southern hemisphere. This is not a problem since sources in the southern hemisphere do not 270 normally have snow on the ground. The only exceptions are the Erebus volcano in Antarctica and the South Sandwich Island volcanoes (Michael and Montagu), but they are always covered by snow. There are also several volcanic sources in the northern and southern latitudes located on small islands that we did not consider in this study: while the volcanoes themselves may have snow cover, the islands are small and surrounded by open ocean.

As snow reflects more light than snow-free surfaces, the SCD values over snow are higher, and VCD values are 275 overestimated if AMF for a snow-free albedo is used to convert SCD to VCD. This overestimation is illustrated in **Figure 2** that shows the 2005-2022 mean SO₂ SCDs and VCDs for snow and snow free conditions for Norilsk. SCDs for snow conditions are about 7 times greater than those for snow-free conditions; however, VCDs are very similar.

The ratio of AMF for snow conditions to AMF for snow-free conditions is calculated to estimate potential biases if measurements over snow are used to calculate VCD using a snow-free AMF. If the ratio is close to 1, then there is no need to 280 introduce an additional AMF for snow conditions. **Figure 3** shows the map of such ratios calculated for ECCC AMFs. The ratios are higher over the plains in the US and Kazakhstan as well as over Canadian and Russian tundra and lower over boreal forests in Alaska, Canada, and Russia. The results for NASA AMFs are similar (not shown), although there are also some differences. The absolute values of the ratios are typically higher for ECCC AMFs (the range is from 1 to 8) than for NASA AMFs (the range is from 1 to 6). Note that NASA AMFs the NISE snow/ice flag is known to miss some snow. There are also 285 two sources in northern Russia (marked by a red ellipse), where the difference between NASA and ECCC ratios is particularly large. These sources came online after 2012, and their emissions are likely not included in the GEOS-5 model simulations from which the a priori vertical profile are derived for the NASA OMSO2 algorithm. Therefore, modelled profiles used in the standard OMSO2 V2 retrievals over that region would not have any surface enhancements and this leads to errors in AMF.

To investigate these ratios and their distribution further, we plotted scatter diagrams of the NASA and ECCC AMFs 290 and a scatter diagram of NASA and ECCC ratios for all sources (**Figure 4**). On average, ECCC ratios are smaller than NASA ratios: the average values for all sources are 2.2 and 3.8, respectively. For sources located above 2000 m Above Sea Level

(ASL), the AMFs for conditions with and without snow are not very different, and there are no large differences between NASA and ECCC AMFs. The snow to snow-free AMF ratios for such sources are not very large; all are between 1.3 and 2.2 for both NASA and ECCC data sets. NASA and ECCC AMFs for snow conditions are not very different from the snow-free AMFs with majority of the values between 1 and 2. The largest difference between NASA and ECCC AMFs for no-snow conditions can be seen for sources located below 300 m ASL (the light green diamonds in **Figure 4**). While ECCC AMFs for these sources are between 0.2 and 0.4, the NASA AMFs range from 0.2 to 1.3. This difference probably occurs because no-snow, clear sky NASA AMFs are more sensitive to the choice of a priori profiles for low altitude sources. The two largest NASA AMFs belong to the sources in northern Russia mentioned above where a priori profiles are unreliable.

Thus, there are some difference between mean ECCC and NASA AMFs that are largely caused by different assumptions regarding a priori SO₂ profile shapes. The ECCC AMF assumptions are more suitable for emission estimates since they better represent the SO₂ profile within a plume, while the NASA AMF assumptions are more suitable for average conditions. Since this study is focused on emissions, the presented results are based on ECCC AMFs, except for some case study illustrations, where results for both AMFs are shown to further demonstrate the differences in estimated emissions.

3.2 Case studies

Figure 5 shows four examples of SO₂ annual emissions time series estimated for snow and snow-free conditions using NASA and ECCC VCDs derived from OMI and TROPOMI data. The green lines represent emission estimates for snow-free conditions, i.e., the same data as in the Version 2 SO₂ catalogue. The cyan lines show emissions estimated using data with snow on the ground, i.e., data that were not used in the SO₂ catalogue. There are also two estimates of emissions based on all data. The red line is the average of the estimates for conditions with and without snow weighted using an inverse-variance weighting method. This approach can be considered as a quick correction of the existing SO₂ catalogue when emissions are estimated using only pixels with snow and then those emission estimates are merged with the existing “snow free” catalogue values. The blue line represents the emission estimates based on all data (with and without snow on the ground).

Kliuchevskoi is a volcanic source in Kamchatka, where observations over snow account for 35% of all measurements suitable for emission estimates. Volcanic degassing emissions are variable, nevertheless both snow and snow-free data show a similar variability except, perhaps in 2018, when emissions under snow conditions were about 5 times larger than under no-snow (**Figure 5 a-c**). This could be explained by high emission levels in the winter-spring time. TROPOMI data do not show the same high emission values in 2018; however, no TROPOMI data were available from January to mid-April. This example also shows that the weighted average of emission estimates under snow and no-snow conditions may not reflect the total annual emissions correctly. The no-snow estimates show low emissions with small spread, while estimates for snow conditions were, probably, highly variable with large standard deviations and therefore had a low weight. Emission estimates based on all data provide a value close to the mean of emissions under snow and no-snow conditions. As Kliuchevskoi is about 5000 m high, the ECCC and NASA AMF values as well as emission estimates are very similar for these two data versions.

325 Karabash is a copper smelter in the Ural region of Russia, where observations over snow accounts for 31% of all measurements. It is expected that smelter emissions would not change very much with the season. **Figure 5e** confirms this, as the ECCC-AMF-based estimates with and without snow are almost identical in most years. The standard deviation of the difference between these estimates is only 14 kt y⁻¹ (about 12%). Estimates based on NASA data demonstrate larger (even twice larger in 2011) emissions when data for snow conditions are used.

330 East Lambeishor is an oil treatment plant in northern Russia that processes fluid mixtures of oil, gas, and water from oil wells, removes hydrogen sulfide, and prepares the oil for further use. It is one of the two sources mentioned in Section 2.1 where the difference between NASA and ECCC ratios is particularly large. The plant started its operation in about 2014, and therefore its emissions were not properly reflected in the inventories used to derive model SO₂ profiles for the NASA retrievals. NASA AMFs for snow-free conditions are almost the same as for snow conditions (1.3 and 1.8, respectively), and the ratio is only 1.4, while it is 5 for ECCC AMFs. Snow and snow-free conditions each account for half of the total number of measurements. **Figure 5h** shows that ECCC estimates for snow and snow-free conditions are similar, while NASA estimates for snow conditions in 2015-2020 are 2.5 or more times larger than that for snow-free conditions (**Figure 5g**). The ECCC AMF-based estimates show similar emissions for both conditions for most years. The emission uncertainties for snow-free conditions are much larger than for snow conditions. In 2019, the emissions for snow-free conditions were about zero, although the uncertainty of this estimate was large, about 50 kt y⁻¹. The 2019 weighted average (the red line in **Figure 5h**) is determined by the emission value for snow conditions and nearly identical to the estimated annual emissions in the three previous years. TROPOMI data also show that emissions for snow-free conditions in 2019 are about 80 kt y⁻¹, i.e., very close to OMI estimated for snow conditions. This example demonstrates that additional measurements under snow conditions can substantially improve the satellite annual emission inventory.

345 The proper way to validate the estimated emissions for snow and snow-free conditions is to compare them with reliable inventories of reported point-source emissions. Such inventories are available for the USA and Europe, but as **Figure 1** shows, the number of measurements for snow conditions there is too small for validation. There is, however, information about annual emissions for Canadian and some Russian sources. Three examples are given below.

350 The Angarsk source (**Figure 5 j-l**) represents a cluster of two coal-fired power plants (Thermal Power Stations 9 and 10) located 8 km apart in the Irkutsk region of Russia. These plants provide steam, heat, and electricity, and it is expected that their emissions are higher during the cold season when the plant outputs are used to heat buildings in local residential and industrial areas. As **Figure 5 j, k** shows, emissions for snow conditions are 2-5 times greater than those for snow-free conditions for both ECCC and NASA AMF-based data sets. Uncertainties and inter annual variations in emissions estimates are large for this source. It is interesting to note that the total emissions are rather stable. This example illustrates that there are sources with very large seasonal variations of emissions. Note that there are also other industrial sources in the region, e.g., the Angarsk oil refinery, that also contribute to total emissions from the Angarsk source. TROPOMI data show (**Figure 5 l**), in general, total emissions that are similar to those from OMI, although the spread between snow and no-snow estimates is not as large as for OMI. The average 2005-2022 estimated Angarsk SO₂ emissions are 122, 31, and 66 kt y⁻¹ for snow, snow-free,

and all conditions respectively (for ECCC AMFs). The reported annual total emissions from the two power plants were available for 2018-2020 from the Reports on the State and protection of the environment in the Irkutsk Region (in Russian, available online from <https://www.ecoindustry.ru/gosdoklad/view/570.html> accessed on November 23, 2022). They were from 75 to 97 kt y⁻¹. Thus, emissions based on OMI snow-free data (on average, 33 kt y⁻¹ in 2018-2020) are largely underestimated, and additional data for the snow conditions, that account for 35% of all measurements, produce more realistic annual emission estimates (on average, 72 kt y⁻¹ in 2018-2020).

Norilsk smelters are the world's largest industrial point source, and column SO₂ values over Norilsk are high (**Figure 2**). High Norilsk SO₂ VCDs are clearly seen by satellite instruments working in the IR- and UV-spectral intervals (Bauduin et al., 2014; Fioletov et al., 2013; Li et al., 2020c; Walter et al., 2012). Norilsk is located in Central Siberia at 69.36° N (above the Arctic Circle). The polar night there lasts 45 days, and observations with SZA < 70° are available only from March to September. The mean annual temperature is around -10°C with only 84 days a year with the temperature above 0°C. The snow cover on average lasts for 247 days (Shiklomanov and Laruelle, 2017), and only three summer months are typically snow-free. Measurements over snow account for 60% of all suitable measurements. Thus, the annual emission estimates in the SO₂ catalogue are actually based on just 3-4 months of data and the addition of snow-covered days to the emission estimates doubles that length.

Norilsk emissions are published annually in the Reports on the state and protection of the environment in the Krasnoyarsk Region (in Russian, available online from <http://krasecology.ru> (e.g., <http://krasecology.ru/Data/Docs/Сводный%20Доклад%20-%202021.pdf> accessed on November 23, 2022)). Reported and estimated emissions for Norilsk are shown in **Figure 6 a**. For the ECCC AMF-based data, the average difference between estimated and reported emissions is -3%, 0%, and -6.4% for all conditions, snow-free and snow conditions respectively, while the standard deviations are 9.3%, 10.3%, and 15.0% respectively. Estimates based on all data are typically closer to reported emissions compared with estimates based on either data with or without snow (**Figure 6a**). The biases between reported and estimated emissions are within the 2-σ uncertainties and are not unexpected since several factors could produce a multiplicative bias in the estimated emissions (Fioletov et al., 2016).

Flin Flon copper and zinc smelter was one of the largest SO₂ emission sources in North America, although its emissions were nearly 10 times less than the Norilsk emissions. The Canadian annual point-source SO₂ emission data are available from the National Pollutant Release Inventory (NPRI, <https://www.canada.ca/en/services/environment/pollution-waste-management/national-pollutant-release-inventory.html> accessed on December 05, 2022). Smelter operations ceased on June 11, 2010. Flin Flon is located at 54.77° N and typically has snow cover from November to April and a snow-free surface in May-September; therefore, about 37% of all OMI data over Flin Flon suitable for emission estimates are from snow conditions. Our estimated emissions for Flin Flon are typically lower than the reported ones. In the case of all data and ECCC AMFs, the difference is about 15% (**Figure 6b**). The difference is even larger (about 20%) if only snow-free data are used. Nevertheless, all estimated emissions show high values in 2005-2009, a steep decline in 2010, and nearly zero values thereafter.

3.3 Overall statistics

Annual emissions were estimated for conditions with and without snow, and these estimates were compared with the results summarized in **Table 1**. First, the mean annual emissions for the two types of conditions (snow and snow-free) were calculated for every source, and then these mean values were averaged among all sources within that source category (power plant, smelter together with oil and gas, volcano). Only sources with mean annual emissions exceeding 20 kt y⁻¹ over the 2005-2022 period for both snow and snow-free conditions were used to avoid the impact of small sources with high emission uncertainties. Finally, the percent difference between the mean values for conditions with and without snow were calculated for each group. The calculations were done using ECCC AMF-based data.

As **Table 1** shows, there is a 3%-4% average difference (estimates for the snow conditions are larger) between snow and snow-free conditions for smelters and oil and gas-related sources as well as for volcanos. This difference is relatively small taking into account that the estimates are based on two independent subsets of data and is within its statistical uncertainty. As for individual sources, for three quarters of smelters and oil and gas-related sources, the difference is within $\pm 20\%$. For power plants, however, the average difference, at about 25%, is much larger than for other types of sources, i.e., power plants have additional 20% difference between snow and snow-free conditions compared to, e.g., smelters. This is consistent with the fact that most of them serve as a source of heat during the cold season and/or responding to a higher energy demand during the cold season and therefore produce more emissions during that time than during the warm season. The largest difference between the snow and snow-free-based estimates occurs at Angarsk discussed in Section 3.2.

The overall impact of additional measurements for snow conditions on estimated total annual emissions for four main regions is shown in **Figure 7**. The largest differences between the current SO₂ catalogue (i.e., for snow-free conditions) and the emission estimates that include observations over snow can be seen for power plants in the region of Russia plus Kazakhstan. As discussed, it is likely due to larger power generation during the cold season. The difference is particularly large for the thermoelectrical power station Angarsk (see section 3.2). Overall the annual power plants emissions in that region are 15% higher (13% without Angarsk) if measurements over snow are included. In contrast, smelters and oil and gas-related sources in that region show only a 2% difference in estimated emissions when data for snow conditions are added, although the fraction of measurements over snow is nearly the same for both cases (about 30%). The standard deviation of the difference is also nearly the same in both cases (about 6%). The total emissions from five Ukrainian power plants are also higher, although the mean difference is only 6% due to warmer winter conditions as illustrated by the total number of days with snow on the ground (10%). All sources in China with 10% or more observations over snow are in the north-western part of the country. Although most of them are power plants, the addition of data with snow on the ground practically does not change the average emission estimates (the difference is only 1.5%). For Canada, the mean difference is about 6%, and it is largely caused by the difference for two major sources: Flin Flon and Thompson smelters, that were shut down in 2010 and 2018, respectively. Aside from some biases, the estimates based on snow-free data and all data are consistent: The standard deviation of the difference between annual emissions for them is between 6% and 12% for all regions.

4. Summary and discussion

425 In this study we investigated how satellite SO₂ measurements over snow-covered surfaces can be used to improve the “top
down” emissions reported in the snow-free SO₂ catalogue. OMI data for 2005-2022 were used in the study and the same
emission estimation algorithm as in the SO₂ catalogue (Fioletov et al., 2016, 2023) was applied. Only 100 out of 759 sources,
listed in the catalogue, have 10% or more of the observations over snow. However, for 40 sources, more than 30% of
430 measurements suitable for emission calculations were taken over snow-covered surfaces. The addition of data for snow
conditions is particularly important at high-latitude sites. For example, in the case of Norilsk smelters, the world’s largest
SO₂ point source, annual emission estimates in the original SO₂ catalogue were based only on 3-4 summer months; additional
data for snow conditions extends that period to 7 months.

The SO₂ catalogue was based on satellite measurements of SO₂ SCDs that were converted to VCDs using single site-
specific constant ECCC AMFs, calculated for snow-free conditions. The same approach was applied to measurements with
435 snow on the ground; a new set of constant site-specific AMFs was created and applied to SCDs to obtain VCDs for snow
conditions. Then, the emissions were estimated using these VCDs for snow conditions as well as using a merged dataset of
VCDs for snow and snow-free conditions. It was found that emissions estimated for snow conditions using ECCC AMFs on
average are consistent with estimated for snow-free conditions for volcanic sources, smelters, and oil and gas-related sources
(there is a small, 3-4% bias between them). For power plants, emissions estimated for snow conditions are on average 25%
440 higher than for snow-free conditions. This is likely because many power plants are also used as a heat source in the wintertime.

Similar emissions estimates based on NASA OMI V2 VCD data show larger biases snow and snow-free conditions.
Biases are about 20% for volcanic sources, 38% for smelter and oil and gas sources, and more than 60% for power plants.
Analysis of the AMF factors (calculated as a median ratio of SCD to VCD) suggests that the difference between such NASA
AMFs for snow and snow-free conditions is typically 1.5-2 times less than that for ECCC AMFs. The reason for that is
445 relatively high NASA AMFs for some sources at low elevation under snow-free conditions that can be explained by non-
optimal a priori profiles in the retrievals.

The approach to establish a priori SO₂ profiles in the NASA dataset is based on a climatology from CTM model
output may not be optimal for emissions estimates, particularly over the areas around isolated point sources. It is important
that the AMFs represent the vertical profile in the plumes correctly in order for the emission estimation algorithms to be
450 accurate. For the NASA dataset, cases are combined when SO₂ is in the plume and then it is outside the plume, i.e., without
SO₂ in the boundary layer. Therefore, the vertical profile represents the “average” conditions that may not describe the vertical
distribution in the plume correctly. This effect is particularly important in the areas where SO₂ emission sources are not
properly accounted for in the CTM model or in the areas where the spatial resolution of a global model does not sufficiently
resolve the gradient in SO₂ in the boundary layer (e.g., around major point sources). Examples of oil and gas sources in northern
455 Russia suggest that AMFs created from incorrect a priori SO₂ profiles can lead to unrealistic emission estimates (e.g., can be
overestimated by a factor of four). On the other hand, if the SO₂ is in the free troposphere or stratosphere, e.g., for emissions

from volcanic eruptions, an incorrect a priori profile would underestimate volcanic VCDs. The better approach is to make assumptions about the vertical profile based on the source of SO₂, e.g., from local pollutions or from remote volcanic eruptions, to calculate the final retrieved SO₂ VCD value.

460 In summary, it is worth adding data in snow-covered conditions for the annual emission estimates. If we use just two
ECCC AMFs (one for snow, the other for snow-free conditions), these AMFs always have some uncertainties due to
uncertainties in the input parameters. Moreover, the snow albedo changes depending on snow depth and age. Thus, there could
be some systematic differences between SO₂ values (and emissions) retrieved over snow and snow-free conditions.
Nevertheless, as shown for individual sources, the difference between emissions estimated for snow and snow-free conditions
465 is within $\pm 20\%$ for three quarters of smelters and oil and gas sources. This is a reasonable agreement given that there is typically
a 3-5 times difference between AMFs for conditions with and without snow (**Figure 3 c**). With just a few exceptions, such as
Norilsk, the snow-covered observation conditions represent less than 40% of all observations for typical sites with snow cover
suitable for emission estimates. Therefore, adding snow conditions introduces the bias that is typically less than $20\% \times 0.4$
=8% that is not very large compared to the overall uncertainties of OMI-based annual emissions estimates that are about 50%
470 (Fioletov et al., 2016). We also saw that the difference between winter and summer emissions could be as large as 100% for
some power plants. Thus, adding observations with snow-covered conditions could be important to account for seasonal
changes in emissions. We are planning to include data for snow conditions for emission estimates in the next version of the
SO₂ catalogue. The emission estimates could be further improved in the future if more accurate AMF estimates for different
albedo conditions and better estimates of winter and summer decay times become available.

475 5. Data availability

The OMI version 2 SO₂ data are publicly available from the Goddard Earth Sciences (GES) Data and Information Services
Center (DISC) (https://disc.gsfc.nasa.gov/datasets/OMSO2_003/summary accessed on June 1, 2023 (Li et al., 2020a). The
TROPOMI COBRA SO₂ data set is presently available from available from BIRA-IASB through distributions.aeronomie.be
(accessed on April 19, 2023) and will be available from the S5P-PAL Data Porta (<https://data-portal.s5p-pal.com/> accessed on
480 June 1, 2023). The version 2 SO₂ point source catalogue is available from the NASA Global Sulfur Dioxide Monitoring Home
Page (<https://so2.gsfc.nasa.gov/measures.html> accessed on April 19, 2023). The direct link to the data set is
https://so2.gsfc.nasa.gov/kml/Catalogue_SO2_2022.xls. DOI identifier
https://disc.gsfc.nasa.gov/datasets/MSAQSO2L4_2/summary (DOI: 10.5067/MEASURES/SO2/DATA406). The IMS snow
data are available from the National Snow and Ice Data Center, University of Colorado
485 (<ftp://sidads.colorado.edu/pub/DATASETS/NOAA/G02156/> accessed on September 27, 2023).

Author Contributions. VF prepared the paper and figures with contributions from all the co-authors. VF and CM developed
the emission estimation algorithm. CM calculated the AMF factors. VF, CM and DG processed satellite data and estimated

the emissions. CL, NK, and JJ developed the OMI PCA V2 algorithms and provided data. NT developed the TROPOMI COBRA algorithms and provided data. SC contributed to cataloging of volcanic sources and interpreted the estimated volcanic emissions.

Competing interests. The authors declare that they have no conflict of interest. At least one of the (co-)authors is a member of the editorial board of Atmospheric Measurement Techniques.

Acknowledgments

Can Li, Nickolay Krotkov, and Joanna Joiner acknowledge support from the NASA Earth Science Division Aura Science Team and US participating investigator programs. OMI PCA SO₂ data used in this study have been publicly released as part of the Aura OMI Sulfur Dioxide Data Product - OMSO2. We thank EU/ESA/KNMI/DLR for providing the TROPOMI/S5P Level 1 products. Nicolas Theys acknowledges support from ESA (S5P-PAL and S5P MPC projects) and BELSO Prodex-Trace S5P.

References

- 500 Bauduin, S., Clarisse, L., Clerbaux, C., Hurtmans, D. and Coheur, P.-F.: IASI observations of sulfur dioxide (SO₂) in the boundary layer of Norilsk, *J. Geophys. Res. Atmos.*, 119, 4253–4263, doi:10.1002/2013JD021405, 2014.
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V. and Goede, A. P. H.: SCIAMACHY: Mission objectives and measurement modes, *J. Atmos. Sci.*, 56(2), 127–150, 1999.
- Bramstedt, K., Richter, A., Van Roozendaal, M. and De Smedt, I.: Comparisons of SCIAMACHY sulfur dioxide observations, 505 *Eur. Sp. Agency, (Special Publ. ESA SP, 2004(562), 85–88, 2004.*
- Callies, J., Corpaccioli, E., Eisinger, M., Hahne, A. and Lefebvre, A.: GOME-2-Metop’s second-generation sensor for operational ozone monitoring., 2000.
- Carn, S. A., Krotkov, N. A., Yang, K. and Krueger, A. J.: Measuring global volcanic degassing with the Ozone Monitoring Instrument (OMI), *Geol. Soc. London, Spec. Publ.*, 380(1), 229–257, doi:10.1144/SP380.12, 2013.
- 510 Carn, S. A., Fioletov, V. E., McLinden, C. A., Li, C. and Krotkov, N. A.: A decade of global volcanic SO₂ emissions measured from space, *Sci. Rep.*, 7, 44095, doi:10.1038/srep44095, 2017.
- Cooper, M. J., Martin, R. V., Lyapustin, A. I., and McLinden, C. A.: Assessing snow extent data sets over North America to inform and improve trace gas retrievals from solar backscatter, *Atmos. Meas. Tech.*, 11, 2983–2994, <https://doi.org/10.5194/amt-11-2983-2018>, 2018.
- 515 Dentener, F., Drevet, J., Lamarque, J. F., Bey, I., Eickhout, B., Fiore, A. M., Hauglustaine, D., Horowitz, L. W., Krol, M., Kulshrestha, U. C., Lawrence, M., Galy-Lacaux, C., Rast, S., Shindell, D., Stevenson, D., Van Noije, T., Atherton, C., Bell, N., Bergman, D., Butler, T., Cofala, J., Collins, B., Doherty, R., Ellingsen, K., Galloway, J., Gauss, M., Montanaro, V., Müller, J. F., Pitari, G., Rodriguez, J., Sanderson, M., Solomon, F., Strahan, S., Schultz, M., Sudo, K., Szopa, S. and Wild, O.: Nitrogen and sulfur deposition on regional and global scales: A multimodel evaluation, *Global*
- 520 *Biogeochem. Cycles*, 20(4), doi:10.1029/2005GB002672, 2006.
- Eisinger, M. and Burrows, J. P.: Tropospheric sulfur dioxide observed by the ERS-2 GOME instrument, *Geophys. Res. Lett.*, 25(22), 4177–4180, 1998.
- Fedkin, N., Li, C., Dickerson, R.R., Canty, T., Krotkov, N.A., Linking improvements in sulfur dioxide emissions to decreasing sulfate wet deposition by combining satellite and surface observations with trajectory analysis, *Atmospheric*
- 525 *Environment*, doi: <https://doi.org/10.1016/j.atmosenv.2018.11.039>, 2018)
- Fioletov, V., McLinden, C., Griffin, D., Theys, N., Loyola, D., Hedelt, P., Krotkov, N. and Li, C.: Anthropogenic and volcanic point source SO₂ emissions derived from TROPOMI onboard Sentinel 5 Precursor: first results, *Atmos. Chem. Phys. Discuss.*, (2018), 1–30, doi:10.5194/acp-2019-1095, 2020.
- Fioletov, V. E., McLinden, C. A., Krotkov, N., Yang, K., Loyola, D. G., Valks, P., Theys, N., Van Roozendaal, M., Nowlan, 530 C. R., Chance, K., Liu, X., Lee, C. and Martin, R. V.: Application of OMI, SCIAMACHY, and GOME-2 satellite SO₂ retrievals for detection of large emission sources, *J. Geophys. Res. Atmos.*, 118(19), 11,399-11,418,

doi:10.1002/jgrd.50826, 2013.

- Fioletov, V. E., McLinden, C. A., Krotkov, N. and Li, C.: Lifetimes and emissions of SO₂ from point sources estimated from OMI, *Geophys. Res. Lett.*, 42(6), 1969–1976, doi:10.1002/2015GL063148, 2015.
- 535 Fioletov, V. E., McLinden, C. A., Krotkov, N. A., Li, C., Joiner, J., Theys, N., Carn, S. and Moran, M. D.: A global catalogue of large SO₂ sources and emissions derived from the Ozone Monitoring Instrument, *Atmos. Chem. Phys.*, 16(18), 11497–11519, doi:doi:10.5194/acp-16-11497-2016, 2016.
- Fioletov, V. E., Mclinden, C. A., Griffin, D., Abboud, I., Krotkov, N., Leonard, P. J. T., Li, C., Joiner, J., Theys, N. and Carn, S.: Version 2 of the global catalogue of large anthropogenic and volcanic SO₂ sources and emissions derived from
540 satellite measurements, *Earth Syst. Sci. Data*, 15, 75–93 [online] Available from: <https://doi.org/10.5194/essd-2022-281>, 2023.
- Fischer, T. P., Arellano, S., Carn, S., Aiuppa, A., Galle, B., Allard, P., Lopez, T., Shinohara, H., Kelly, P., Werner, C., Cardellini, C. and Chiodini, G.: The emissions of CO₂ and other volatiles from the world's subaerial volcanoes, *Sci. Rep.*, 9(1), 1–11, doi:10.1038/s41598-019-54682-1, 2019.
- 545 de Graaf, M., Sihler, H., Tilstra, L. G. and Stammes, P.: How big is an OMI pixel?, *Atmos. Meas. Tech.*, 9(8), 3607–3618, doi:10.5194/amt-9-3607-2016, 2016.
- Gualtieri, G., Analysing the uncertainties of reanalysis data used for wind resource assessment: A critical review, *Renewable and Sustainable Energy Reviews*, 167, 112741, ISSN 1364-0321, <https://doi.org/10.1016/j.rser.2022.112741>, 2022.
- Hansell, A. and Oppenheimer, C.: Health Hazards from Volcanic Gases: A Systematic Literature Review, *Arch. Environ. Heal. An Int. J.*, 59(12), 628–639, doi:10.1080/00039890409602947, 2010.
- 550 Helfrich, S. R., McNamara, D., Ramsay, B. H., Baldwin, T. and Kasheta, T.: Enhancements to, and forthcoming developments in the Interactive Multisensor Snow and Ice Mapping System (IMS), *Hydrol. Process*, 21, 1576–1586, doi:doi:10.1002/hyp.6720, 2007.
- Hutchinson, T. C. and Whitby, L. M.: The effects of acid rainfall and heavy metal particulates on a boreal Forest ecosystem near the sudbury smelting region of Canada, *Water. Air. Soil Pollut.*, 7(4), 421–438, doi:10.1007/BF00285542, 1977.
- 555 Ialongo, I., Fioletov, V., Mclinden, C., Jåfs, M., Krotkov, N., Li, C. and Tamminen, J.: Application of satellite-based sulfur dioxide observations to support the cleantech sector: Detecting emission reduction from copper smelters, *Environ. Technol. Innov.*, 12, 172–179, doi:10.1016/j.eti.2018.08.006, 2018.
- Inness, A., Ades, M., Balis, D., Efremenko, D., Flemming, J., Hedelt, P., Koukouli, M.-E., Loyola, D. and Ribas, R.: Evaluating
560 the assimilation of S5P/TROPOMI near real-time SO₂ columns and layer height data into the CAMS integrated forecasting system (CY47R1), based on a case study of the 2019 Raikoke eruption, *Geosci. Model Dev.*, 15(3), 971–994, doi:10.5194/GMD-15-971-2022, 2022.
- Joiner, J. and Vasilkov, A. P.: First results from the OMI rotational- Raman scattering cloud pressure algorithm, *IEEE Trans. Geophys. Remote Sens.*, 44, 1272–1282, 2006.
- 565 Kharol, S. K., McLinden, C. A., Sioris, C. E., Shephard, M. W., Fioletov, V., van Donkelaar, A., Philip, S. and Martin, R. V.:

- OMI satellite observations of decadal changes in ground-level sulfur dioxide over North America, *Atmos. Chem. Phys.*, 17(9), 5921–5929, doi:10.5194/acp-17-5921-2017, 2017.
- Khokhar, M. F., Frankenberg, C., Van Roozendaal, M., Beirle, S., Köhl, S., Richter, a., Platt, U. and Wagner, T.: Satellite observations of atmospheric SO₂ from volcanic eruptions during the time-period of 1996–2002, *Adv. Sp. Res.*, 36(5), 879–887, doi:10.1016/j.asr.2005.04.114, 2005.
- Khokhar, M. F., Platt, U. and Wagner, T.: Temporal trends of anthropogenic SO₂ emitted by non-ferrous metal smelters in Peru and Russia estimated from Satellite observations, *Atmos. Chem. Phys. Discuss.*, 8(5), 17393–17422, 2008.
- Klimont, Z., Smith, S. J. and Cofala, J.: The last decade of global anthropogenic sulfur dioxide: 2000–2011 emissions, *Environ. Res. Lett.*, 8(1), 014003, doi:10.1088/1748-9326/8/1/014003, 2013.
- 575 Krotkov, N. a., McClure, B., Dickerson, R. R., Carn, S. a., Li, C., Bhartia, P. K., Yang, K., Krueger, A. J., Li, Z., Levelt, P. F., Chen, H., Wang, P. and Lu, D.: Validation of SO₂ retrievals from the Ozone Monitoring Instrument over NE China, *J. Geophys. Res.*, 113(D16), D16S40, doi:10.1029/2007JD008818, 2008.
- Krotkov, N. A., Carn, S. A., Krueger, A. J., Bhartia, P. K. and Yang, K.: Band Residual Difference Algorithm for Retrieval of SO₂ From the Aura Ozone Monitoring Instrument (OMI), *IEEE Trans. Geosci. Remote Sens.*, 44(5), 1259–1266, 580 2006.
- Krotkov, N. A., McLinden, C. A., Li, C., Lamsal, L. N., Celarier, E. A., Marchenko, S. V., Swartz, W. H., Bucsela, E. J., Joiner, J., Duncan, B. N., Boersma, K. F., Veefkind, J. P., Levelt, P. F., Fioletov, V. E., Dickerson, R. R., He, H., Lu, Z. and Streets, D. G.: Aura OMI observations of regional SO₂ and NO₂ pollution changes from 2005 to 2015, *Atmos. Chem. Phys.*, 16(7), 4605–4629, doi:10.5194/acp-16-4605-2016, 2016.
- 585 Krueger, A. J.: Sighting of El Chichón sulfur dioxide clouds with the Nimbus 7 Total Ozone Mapping Spectrometer, *Science.*, 220, 1377–1378, 1983.
- Krueger, A. J., Walter, L. S., Bhartia, P. K., Schnetzler, C. C., Krotkov, N. A., Sprod, I. and Bluth, G. J. S.: Volcanic sulfur dioxide measurements from the Total Ozone Mapping Spectrometer instruments, *J. Geophys. Res.*, 100(D7), 14057–14076, doi:10.1029/95JD01222, 1995.
- 590 Lee, C., Richter, A., Lee, H., Kim, Y. J., Burrows, J. P., Lee, Y. G. and Choi, B. C.: Impact of transport of sulfur dioxide from the Asian continent on the air quality over Korea during May 2005, *Atmos. Environ.*, 42(7), 1461–1475, doi:10.1016/J.ATMOSENV.2007.11.006, 2008.
- Lee, C., Martin, R. V., van Donkelaar, A., O’Byrne, G., Krotkov, N., Richter, A., Huey, L. G. and Holloway, J. S.: Retrieval of vertical columns of sulfur dioxide from SCIAMACHY and OMI: Air mass factor algorithm development, 595 validation, and error analysis, *J. Geophys. Res.*, 114(D22), D22303, doi:10.1029/2009JD012123, 2009.
- Lee, C., Martin, R. V., Van Donkelaar, A., Lee, H., Dickerson, R. R., Hains, J. C., Krotkov, N., Richter, A., Vinnikov, K. and Schwab, J. J.: SO₂ emissions and lifetimes: Estimates from inverse modeling using in situ and global, space-based (SCIAMACHY and OMI) observations, *J. Geophys. Res.*, 116(D6), D06304, doi:10.1029/2010JD014758, 2011.
- Levelt, P., Joiner, J., Tamminen, J., Veefkind, P., Bhartia, P. K., Stein Zweers, D., Duncan, B. N., Streets, D. G., Eskes, H.,

- 600 van der A, R., McLinden, C., Fioletov, V., Carn, S., de Laat, J., DeLand, M., Marchenko, S., McPeters, R., Ziemke, J., Fu, D., Liu, X., Pickering, K., Apituley, A., Gonz ales Abad, G., Arola, A., Boersma, F., Chan Miller, C., Chance, K., de Graaf, M., Hakkarainen, J., Hassinen, S., Ialongo, I., Kleipool, Q., Krotkov, N., Li, C., Lamsal, L., Newman, P., Nowlan, C., Suileiman, R., Tilstra, L. G., Torres, O., Wang, H. and Wargan, K.: The Ozone Monitoring Instrument: Overview of twelve years in space, *Atmos. Chem. Phys. Discuss.*, 1–61, doi:10.5194/acp-2017-487, 2017.
- 605 Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Malkki, A., Stammes, P., Lundell, J. O. V. and Saari, H.: The Ozone Monitoring Instrument, *IEEE Trans. Geosci. Remote Sens.*, 44(5), 1093–1101, doi:10.1109/TGRS.2006.872333, 2006.
- Li, C., Joiner, J., Krotkov, N. A. and Bhartia, P. K.: A fast and sensitive new satellite SO₂ retrieval algorithm based on principal component analysis: Application to the ozone monitoring instrument, *Geophys. Res. Lett.*, 40(23), 6314–6318, doi:10.1002/2013GL058134, 2013.
- 610 Li, C., Krotkov, N. A., Leonard, P. J. T. and Joiner, J.: OMI/Aura Sulphur Dioxide (SO₂) Total Column 1-orbit L2 Swath 13x24 km V003, Greenbelt, MD, USA, Goddard Earth Sci. Data Inf. Serv. Cent. (GES DISC), doi:10.5067/Aura/OMI/DATA2022, 2020a.
- Li, C., Krotkov, N. A., Leonard, P. J. T. and Joiner, J.: OMPS/NPP PCA SO₂ Total Column 1-Orbit L2 Swath 50x50km V2, 615 Greenbelt, MD, USA, Goddard Earth Sci. Data Inf. Serv. Cent. (GES DISC), doi:10.5067/MEASURES/SO2/DATA205, 2020b.
- Li, C., Krotkov, N. A., Leonard, P. J. T., Carn, S., Joiner, J., Spurr, R. J. D. and Vasilkov, A.: Version 2 Ozone Monitoring Instrument SO₂ product (OMSO₂ V2): New anthropogenic SO₂ vertical column density dataset, *Atmos. Meas. Tech.*, 13(11), 6175–6191, doi:10.5194/amt-13-6175-2020, 2020c.
- 620 Liu, F., Choi, S., Li, C., Fioletov, V. E., McLinden, C. A. and Joiner, J.: A new global anthropogenic SO₂ emission inventory for the last decade : a mosaic of satellite-derived and bottom-up emissions, *Atmos. Chem. Phys.*, 18, 16571–16586, 2018.
- Longo, B. M., Yang, W., Green, J. B., Crosby, F. L. and Crosby, V. L.: Acute health effects associated with exposure to volcanic air pollution (vog) from increased activity at Kilauea Volcano in 2008., *J. Toxicol. Environ. Health. A*, 625 73(20), 1370–81, doi:10.1080/15287394.2010.497440, 2010.
- Loyola, D. G., Gimeno Garc a, S., Lutz, R., Argyrouli, A., Romahn, F., Spurr, R. J. D., Pedernana, M., Doicu, A., Molina Garc a, V., and Sch ussler, O.: The operational cloud retrieval algorithms from TROPOMI on board Sentinel-5 Precursor, *Atmos. Meas. Tech.*, 11, 409–427, <https://doi.org/10.5194/amt-11-409-2018>, 2018.
- McCormick, B. T., Edmonds, M., Mather, T. A., Champion, R., Hayer, C. S. L. L., Thomas, H. E. and Carn, S. A.: Volcano monitoring applications of the Ozone Monitoring Instrument, *Geol. Soc. London, Spec. Publ.*, 380(1), 259–291, doi:10.1144/SP380.11, 2013.
- 630 McLinden, C. A., Fioletov, V., Boersma, K. F., Krotkov, N., Sioris, C. E., Veefkind, J. P. and Yang, K.: Air quality over the Canadian oil sands: A first assessment using satellite observations, *Geophys. Res. Lett.*, 39(4), 1–8,

doi:10.1029/2011GL050273, 2012.

- 635 McLinden, C. A., Fioletov, V., Boersma, K. F., Kharol, S. K., Krotkov, N., Lamsal, L., Makar, P. A., Martin, R. V., Veefkind, J. P. and Yang, K.: Improved satellite retrievals of NO₂ and SO₂ over the Canadian oil sands and comparisons with surface measurements, *Atmos. Chem. Phys.*, 14(7), 3637–3656, doi:10.5194/acp-14-3637-2014, 2014.
- McLinden, C. A., Fioletov, V., Shephard, M. W., Krotkov, N., Li, C., Martin, R. V., Moran, M. D. and Joiner, J.: Space-based detection of missing sulfur dioxide sources of global air pollution, *Nat. Geosci.*, 9(7), 496–500, doi:10.1038/ngeo2724, 2016.
- 640 McLinden, C. A., Adams, C. L. F., Fioletov, V., Griffin, D., Makar, P. A., Zhao, X., Kovachik, A., Dickson, N. M., Brown, C., Krotkov, N., Li, C., Theys, N., Hedelt, P. and Loyola, D. G.: Inconsistencies in sulphur dioxide emissions from the Canadian oil sands and potential implications, *Environ. Res. Lett.*, 16, 014012, doi:10.1088/1748-9326/abcbbb, 2020.
- 645 McPeters, R. D., Heath, D. F. and Schlesinger, B. M.: Satellite observation of SO₂ from El Chichon: identification and measurement., *Geophys. Res. Lett.*, 11(12), 1203–1206, 1984.
- Nowlan, C. R., Liu, X., Chance, K., Cai, Z., Kurosu, T. P., Lee, C. and Martin, R. V.: Retrievals of sulfur dioxide from the Global Ozone Monitoring Experiment 2 (GOME-2) using an optimal estimation approach: Algorithm and initial validation, *J. Geophys. Res.*, 116(D18), D18301, doi:10.1029/2011JD015808, 2011.
- 650 O'Byrne, G., Martin, R. V., van Donkelaar, A., Joiner, J., and Celarier, E. A. (2010), Surface reflectivity from the Ozone Monitoring Instrument using the Moderate Resolution Imaging Spectroradiometer to eliminate clouds: Effects of snow on ultraviolet and visible trace gas retrievals, *J. Geophys. Res.*, 115, D17305, doi:[10.1029/2009JD013079](https://doi.org/10.1029/2009JD013079).
- Oppenheimer, C., Scaillet, B. and Martin, R. S.: Sulfur Degassing From Volcanoes: Source Conditions, Surveillance, Plume Chemistry and Earth System Impacts, *Rev. Mineral. Geochemistry*, 73(1), 363–421, doi:10.2138/RMG.2011.73.13, 2011.
- 655 Qu, Z., Henze, D. K., Li, C., Theys, N., Wang, Y., Wang, J., Wang, W., Han, J., Shim, C., Dickerson, R. R. and Ren, X.: SO₂ Emission Estimates Using OMI SO₂ Retrievals for 2005–2017, *J. Geophys. Res. Atmos.*, 124(14), 8336–8359, doi:10.1029/2019JD030243, 2019.
- Rix, M., Valks, P., Hao, N., Loyola, D., Schlager, H., Huntrieser, H., Flemming, J., Koehler, U., Schumann, U. and Inness, A.: Volcanic SO₂, BrO and plume height estimations using GOME-2 satellite measurements during the eruption of Eyjafjallajökull in May 2010, *J. Geophys. Res. Atmos.*, 117, doi:10.1029/2011JD016718, 2012.
- 660 Robock, A.: Volcanic eruptions and climate, *Rev. Geophys.*, 38(2), 191–219, doi:10.1029/1998RG000054, 2000.
- Schaaf, C. B., Gao, F., Strahler, A. H., Lucht, W., Li, X., Tsang, T., Strugnell, N. C., Zhang, X., Jin, Y., Muller, J.-P., Lewis, P., Barnsley, M., Hobson, P., Disney, M., Roberts, G., Dunderdale, M., Doll, C., d'Entremont, R. P., Hu, B., Liang, S., Privette, J. L., and Roy, D.: First operational BRDF, albedo nadir reflectance products from MODIS, *Remote Sens. Environ.*, 83, 135–148, 2002.
- 665 Schoeberl, M. R., Douglass, A. R., Hilsenrath, E., Bhartia, P. K., Beer, R., Waters, J. W., Gunson, M. R., Froidevaux, L., Gille,

- J. C., Barnett, J. J., Levelt, P. F. and DeCola, P.: Overview of the EOS Aura mission, *IEEE Trans. Geosci. Remote Sens.*, 44(5), 1066–1074, doi:10.1109/TGRS.2005.861950, 2006.
- 670 Shiklomanov, N. I. and Laruelle, M.: A truly Arctic city: an introduction to the special issue on the city of Norilsk, Russia, *Polar Geogr.*, 40(4), 251–256, doi:10.1080/1088937X.2017.1387823, 2017.
- Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A. and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, *Atmos. Chem. Phys.*, 11(3), 1101–1116, doi:10.5194/acp-11-1101-2011, 2011.
- Stenchikov, G., Ukhov, A., Osipov, S., Ahmadov, R., Grell, G., Cady- Pereira, K., et al. How does a Pinatubo-size volcanic
675 cloud reach the middle stratosphere? *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033829. <https://doi.org/10.1029/2020JD033829>, 2021.
- Theys, N., De Smedt, I., Yu, H., Danckaert, T., van Gent, J., Hörmann, C., Wagner, T., Hedelt, P., Bauer, H., Romahn, F., Pedergnana, M., Loyola, D. and Van Roozendael, M.: Sulfur dioxide retrievals from TROPOMI onboard Sentinel-5 Precursor: algorithm theoretical basis, *Atmos. Meas. Tech.*, 10(1), 119–153, doi:10.5194/amt-10-119-2017, 2017.
- 680 Theys, N., Fioletov, V., Li, C., De Smedt, I., Lerot, C., McLinden, C., Krotkov, N., Griffin, D., Clarisse, L., Hedelt, P., Loyola, D., Wagner, T., Kumar, V., Innes, A., Ribas, R., Hendrick, F., Vlietinck, J., Brenot, H. and Van Roozendael, M.: A Sulfur Dioxide Covariance-Based Retrieval Algorithm (COBRA): application to TROPOMI reveals new emission sources, *Atmos. Chem. Phys.*, 1–42, doi:10.5194/acp-2021-294, 2021.
- Thomas, W., Erbetseder, T., Ruppert, T., Roozendael, M. Van, Verdebout, J., Balis, D., Meleti, C. and Zerefos, C.: On the
685 retrieval of volcanic sulfur dioxide emissions from GOME backscatter measurements, *J. Atmos. Chem.*, 50(3), 295–320, doi:10.1007/s10874-005-5544-1, 2005.
- Ukhov, A., Mostamandi, S., Krotkov, N., Flemming, J., da Silva, A., Li, C., Fioletov, V., McLinden, C., Anisimov, A., Alshehri, Y. M. and Stenchikov, G.: Study of SO₂ Pollution in the Middle East Using MERRA-2, CAMS Data Assimilation Products, and High-Resolution WRF-Chem Simulations, *J. Geophys. Res. Atmos.*, 125(6),
690 e2019JD031993, doi:10.1029/2019JD031993, 2020.
- van der A, R.J., de Laat, A.T.J., Ding, J. *et al.* Connecting the dots: NO_x emissions along a West Siberian natural gas pipeline. *npj Clim Atmos Sci* 3, 16 , <https://doi.org/10.1038/s41612-020-0119-z>, 2020.
- van Geffen, J. H. G. M., Eskes, H. J., Boersma, K. F., and Veefkind, J. P. TROPOMI ATBD of the total and tropospheric NO₂ data products, S5P-KNMI-L2-0005-RP, issue 2.4.0 (available online at
695 <https://sentinel.esa.int/documents/247904/2476257/sentinel-5p-tropomi-atbd-no2-data-products>), 2022.
- Vasilkov, A., Joiner, J., Spurr, R., Bhartia, P. K., Levelt, P., and Stephens, G.: Evaluation of the OMI cloud pressures derived from rotational Raman scattering by comparisons with other satellite data and radiative transfer simulations, *J. Geophys. Res.*, 113, D15S19, doi:[10.1029/2007JD008689](https://doi.org/10.1029/2007JD008689), 2008.
- Vasilkov, A. P., Joiner, J., Haffner, D., Bhartia, P. K., and Spurr, R. J. D.: What do satellite backscatter ultraviolet and visible
700 spectrometers see over snow and ice? A study of clouds and ozone using the A-train, *Atmos. Meas. Tech.*, 3, 619–629, <https://doi.org/10.5194/amt-3-619-2010>, 2010

- 705 Veefkind, J. P. P., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H. J. J., de Haan, J. F. F., Kleipool, Q., van Weele, M., Hasekamp, O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H. and Levelt, P. F. F.: TROPOMI on the ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer applications, *Remote Sens. Environ.*, 120(2012), 70–83, doi:10.1016/j.rse.2011.09.027, 2012.
- 710 Vet, R., Artz, R. S., Carou, S., Shaw, M., Ro, C. U., Aas, W., Baker, A., Bowersox, V. C., Dentener, F., Galy-Lacaux, C., Hou, A., Pienaar, J. J., Gillett, R., Forti, M. C., Gromov, S., Hara, H., Khodzher, T., Mahowald, N. M., Nickovic, S., Rao, P. S. P. and Reid, N. W.: A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus, *Atmos. Environ.*, 93, 3–100, doi:10.1016/j.atmosenv.2013.10.060, 2014.
- Walter, D., Heue, K.-P., Rauthe-Schöch, A., Brenninkmeijer, C. A. M., Lamsal, L. N., Krotkov, N. A. and Platt, U.: Flux calculation using CARIBIC DOAS aircraft measurements: SO₂ emission of Norilsk, *J. Geophys. Res.*, 117(D11), D11305, doi:10.1029/2011JD017335, 2012.
- 715 Zhang, Y., Li, C., Krotkov, N. A., Joiner, J., Fioletov, V. and McLinden, C.: Continuation of long-term global SO₂ pollution monitoring from OMI to OMPS, *Atmos. Meas. Tech.*, 10(4), 1495–1509, doi:10.5194/amt-10-1495-2017, 2017.

720 **Table 1.** The average and standard deviation of the difference (%) between emissions estimates based on OMI measurements for snow and snow-free conditions calculated using ECCC AMFs. The difference between the mean emissions for the period 2005-2021 was calculated first for each source and then the average of these differences was calculated. The total number of sources is also shown. Only sources with mean emissions exceeding 20 kt y⁻¹ for both snow and snow-free conditions were included.

Source Type	Number of sources	Average difference (%)	Standard deviation of the difference (%)
Power plants	27	25.8	33.3
Smelters and Oil and Gas	34	4.2	20.6
Volcanos	7	3.2	27.3

725




Figure 1. Emission sources detected with OMI observations over snow or ice. The map shows the sites and the fraction of OMI satellite pixels that are suitable for emission estimates (with $SZA < 70$ and $CF < 0.3$) for these sites as a percentage of all satellite pixels suitable for such estimates. The symbol size is proportional to that fraction.





Figure 2. Mean OMI SO₂ slant (a, b) and vertical (c, d) column density over Norilsk for snow-free (a, c) and snow (b, d) conditions for the period 2005-2022. ECCC AMFs were used for the VCD calculations. Data are smoothed by the oversampling technique with a 25 km averaging radius. Slant column values for snow conditions are about 7 times greater than those for snow-free conditions.



740 **Figure 3.** The ratio between AMFs for snow condition to AMFs for snow-free conditions for ECCC AMFs. Two recently built sources that are not included in the inventories and therefore do not have reliable NASA AMFs are marked by the red circle.






Figure 4. Scatter plots of NASA and ECCC AMFs for conditions (a) without and (b) with snow on the ground. (c) Scatter plot of the AMF enhancement due to snow (the ratio between AMF for conditions with snow to that without snow) for NASA and ECCC AMFs. Different colors represent different elevations of the emission sources (in m) as shown in the legend. Two recently built sources that are not included in the inventories and therefore do not have reliable NASA AMFs are marked by the red circles and ellipses.



750

Figure 5. Time series of estimated annual emissions from OMI NASA dataset (the left column), OMI ECCC AMF-based dataset (the middle column) and TROPOMI ECCC AMF-based dataset (the right column). The lines represent estimates based on all data (blue), on data for snow-free conditions (green), on data for snow conditions (cyan), and the weighted average of the two latter datasets (red) with 2- σ error bars. Note that TROPOMI records started in April 2018 and 2018 estimates over snow are not comparable with OMI estimates for that year.

755



760

Figure 6. Reported and OMI-based estimated SO₂ emissions for Norilsk (a) and Flin Flon (b). Reported emissions (solid dark red line) and emissions estimated by applying ECCC AMFs and using all data (solid green line), only pixels without (dashed orange line) and with (dashed blue line) snow on the ground. The error bars represent 2- σ uncertainties.

765




Figure 7. Total emissions for different regions and source types estimated using (blue line) all data (i.e., with and without snow on the ground) and (orange line) only snow-free data. The mean difference between the two estimates, standard deviation of the difference (sigma), and the fraction of satellite measurements taken over snow-covered surface are also shown. Only emission sources where at least 10% of total measurements were taken over snow are included in this analysis. Estimates based on OMI data are shown.

775