

Aerosol optical depth climatology from the high-resolution MAIAC product over Europe: differences between major European cities and their surrounding environments

5 Ludovico Di Antonio¹, Claudia Di Biagio², Gilles Foret¹, Paola Formenti², Guillaume Siour¹, Jean-François Doussin¹, and Matthias Beekmann²

¹Univ Paris Est Creteil and Université Paris Cité, CNRS, LISA, F-94010 Créteil, France

²Université Paris Cité and Univ Paris Est Creteil, CNRS, LISA, F-75013 Paris, France

10 *Correspondence to:* Ludovico Di Antonio (ludovico.diantonio@lisa.ipsl.fr), Matthias Beekmann (matthias.beekmann@lisa.ipsl.fr)

Abstract. The aerosol optical depth (AOD) is a derived measurement useful to investigate the aerosol load and its distribution at different spatio-temporal scales. In this work we use long-term (2000–2021) MAIAC (Multi-Angle Implementation of Atmospheric Correction) retrievals with 1 km resolution to investigate the climatological AOD variability and trends at
15 different scales in Europe: a continental (30–60°N; 20°W–40°E), a regional (100x100 km²) and an urban local scale (3x3 km²). The AOD climatology at the continental scale shows the highest values during summer (JJA) and the lowest during winter (DJF) seasons. Regional and urban local scales are investigated for twenty-one cities in Europe including capitals and large urban agglomerations. Analyses show AOD average (550 nm) values between 0.06 and 0.16 at the urban local scale, while
20 also displaying a strong north-south gradient. This gradient corresponds to a similar one in the European background, with higher AOD being located over the Po-Valley, the Mediterranean basin, and Eastern Europe. Average enhancements of the local with respect to regional AOD of 57%, 55%, 39% and 32% are found for large metropolitan centers such as Barcelona, Lisbon, Paris and Athens respectively, suggesting a non-negligible enhancement to the aerosol burden through local emissions. Negative average deviations are observed for other cities, such as Amsterdam (–17%) and Brussels (–6%) indicating higher
25 regional background signal and suggesting a heterogeneous aerosol spatial distribution that conceals the urban local signal. Finally, negative statistically significant AOD trends for the entire European continent are observed. A stronger decrease rate at the regional scale with respect to the local scale ~~one~~ occurs for most of the cities under investigation.

1. Introduction

Climate change and air quality preservation represent two of the greatest challenges of our times, especially in densely
30 populated areas. Aerosol particles have been shown to play a key role in climate change and to affect air quality over many regions of the world (Robotto et al., 2022; Viana et al., 2014; Fiore et al., 2012). Aerosols affect the radiative budget both directly, by scattering and absorption of solar and thermal radiation (the aerosol radiation-interactions, ARI) or indirectly, by influencing the cloud formation and properties (aerosol-cloud interactions effect, ACI) (Bellouin et al., 2020). Constraining the aerosol contribution to climate and its change is still a challenge (Bender, 2020) as further demonstrated by the Climate
35 Change 2021 IPCC report indicating ~~still huge large remaining~~ -spread in ARI and ACI estimations (Masson-Delmotte et al., 2021). Atmospheric aerosols are also a concern for air quality and human health (Yang et al., 2018; Li et al., 2017, 2016; Dockery, 2009). Millions of people in Europe and around the world, especially over dense urban agglomerations, industrial areas and rural environment, ~~experience a significant particulate matter exposure~~ ~~are everyday exposed to significant aerosol levels~~ (Sicard et al., 2021). Under favorable weather conditions such as high radiation levels, high temperature, low

40 precipitations and low winds during summer, or temperature inversions and low planetary boundary layer height during winter, primary and secondary aerosol local formation have been shown to build up to create the so called “aerosol pollution episodes” (Foret et al., 2022; Groot Zwaftink et al., 2022; Diémoz et al., 2019). These episodes correspond to daily average PM levels above the European threshold of $50 \mu\text{g m}^{-3}$ and last ~~foring~~ several consecutively days. ~~If such episodes occur frequently,~~ they lead to significant air quality and visibility degradation (Majewski et al., 2021; Singh et al., 2020) and increase the potential health risk (Luo et al., 2021; Grigorieva and Lukyanets, 2021). However, the aerosol anthropogenic precursors, abundant in urban agglomerations, can also spread around emission hot spots and affect larger areas, including rural and forested environments, leading to situations of mixed anthropogenic–biogenic scenarios (Xu et al., 2021). This would lead aerosols to have different chemical, physical and radiative properties and therefore to potentially different impact both on human health (Tuet et al., 2017; Liu et al., 2009) , and the environment (Nascimento et al., 2021; Shrivastava et al., 2019; Martin et al., 50 2016). In this regard, how the local and regional scale anthropogenic and biogenic precursors, their mixing and their processing affect aerosol loading and properties, in particular around major city agglomerations, is still unknown and is a matter of scientific investigation (Cantrell and Michoud, 2022; Liu et al., 2021; Ma et al., 2021).

The Aerosol Optical Depth (AOD) is a key parameter to investigate aerosol load, ~~properties and~~ and distribution over local to large scale areas (Bai et al., 2022; Faisal et al., 2022; Raptis et al., 2020; Sun et al., 2019; Just et al., 2015; Smirnov et al., 55 2002). The AOD is defined as the integral of the aerosol extinction coefficient (units of length^{-1}) over the whole atmospheric column and it depends on the aerosol mass concentration, size distribution, shape and complex refractive index. Measurements of AOD are used to improve the air quality forecasts since they can be assimilated in regional or global models (Lee et al., 2022; Ha et al., 2020; Kondragunta et al., 2008) and they can be also linked to visibility measurements (Aman et al., 2022; Zhang et al., 2016; Boers et al., 2015; Kessner et al., 2013; Bäumer et al., 2008). Moreover, the AOD spectral variability can also be used to discern among different aerosol types and help source apportionment (Tuccella et al., 2020; Bahadur et al., 60 2012). However, since AOD observations are vertically integrated, the correlation with surface aerosol measurements may not be straightforward (He et al., 2021; Grgurić et al., 2014; Segura et al., 2017; Guo et al., 2009; Schaap et al., 2009; Schäfer et al., 2008). In fact, AOD is sensitive to dust and biomass burning plumes transported at high altitude, which may not affect surface measurements (Eck et al., 2023; Gkikas et al., 2022; Song et al., 2009). Different studies reported AOD trends on a global scale (Gupta et al., 2023, 2022; Zhao et al., 2017; He et al., 2016; Mehta et al., 2016; Mao et al., 2014) supporting a decreasing AOD trend over Europe (Gupta et al., 2023, 2022; Filonchik et al., 2020b; Alpert et al., 2012). The overall decreasing trend at the European regional scale has been attributed to mitigation policies applied in recent years for the aerosol and the aerosol precursor emissions (Gupta et al., 2022; Hammer et al., 2020; Provençal et al., 2017; Zhao et al., 2017).

The AOD is routinely retrieved across the globe by both ground–based sun photometers measurements, such those of the 70 widespread AERONET network (Aerosol Robotic Network) (Giles et al., 2019), and by satellite sensors, among them the Moderate Resolution Imaging Spectroradiometer (MODIS). Three complementary algorithms, developed at NASA, exist for the MODIS ~~aerosol~~-AOD retrieval: the Deep Blue (DB) (Hsu et al., 2004), the Dark Target (DT)(Remer et al., 2020, 2005) and the more recent Multi–Angle Implementation of Atmospheric Correction (MAIAC) algorithm (Lyapustin et al., 2018). The DB and DT algorithms, extensively used in literature (e.g., Shi et al., 2021; Spencer et al., 2019; Sayer et al., 2018; Lee 75 et al., 2017; Hsu et al., 2017), provide aerosol retrievals at the spatial resolution of 3km and 10 km. The MAIAC algorithm provides atmospheric retrievals of AOD at 470 and 550 nm at the more advanced spatial resolution of 1 km. As a matter of fact, an accurate estimation of surface reflectance, discerning among atmospheric and surface contributions, is necessary to provide the best quality AOD retrievals (Bilal et al., 2019). In this regards, the MAIAC algorithm benefits of the multi–angle satellite observations, retaining in memory up to 16 days of consecutive satellite overpasses, to better constrain the surface 80 reflectance, improving the AOD retrievals in particular over complex scenes as urban areas (Chen et al., 2021; Gupta et al.,

2016; Wang et al., 2010). The MAIAC aerosol algorithm uses eight different background aerosol models over land (Look Up Tables, LUT) and it has developed a more stable algorithm that reduces the AOD bias over bright surfaces (in absence of smoke and dust), typical for the DT and DB algorithms (Lyapustin et al., 2018). Furthermore, MAIAC can retrieve AOD over partial cloudy conditions and distinguish between smoke and dust scenes (Lyapustin et al., 2012). The AOD from the MAIAC algorithm has been validated over different areas of the world and shown to perform better with respect to the DT and DB algorithms when compared to AERONET observations (Su et al., 2023; Falah et al., 2021; Qin et al., 2021; Martins et al., 2019; Zhang et al., 2019; Tao et al., 2019; Mhawish et al., 2019; Martins et al., 2017; Just et al., 2015). The estimated expected error (EE) for MAIAC AOD is evaluated at $\pm(0.05 + 0.1AOD)$, but it is shown to vary as a function of surface reflectivity, aerosol loading and size, as well as aerosol type (Falah et al., 2021). Because of its 1 km resolution and good performances, the MAIAC AOD product has increasing use in air quality studies (Pedde et al., 2022; Gladson et al., 2022; Yang et al., 2022; van Donkelaar et al., 2021; Jung et al., 2021; Hough et al., 2021).

In this paper, we benefit from the high-spatial resolution MAIAC long-term data (from 2000 to 2021) to investigate AOD over Europe. This work is part of the ACROSS (Atmospheric ChemistRy Of the Suburban foreSt, <https://across.cnrs.fr/>) project, whose objective is to deepen the current physical-chemical knowledge of the interaction between anthropogenic emissions in the Paris area and its surrounding environment, through an intensive field campaign which took place in the summer 2022 (Cantrell and Michoud, 2022). Within the ACROSS context, this study ~~wants to achieve~~aims to achieve three different objectives:

- Investigate the urban local vs regional scale aerosol optical depth variability starting from a broader context over the European domain (20°W–40°E,30–60°N) up to the urban local scale (3x3 km²) around major urban agglomerations in Europe;
- explore the long-term trends at the urban local (3x3 km²), regional (100x100 km²) and continental scales (20°W–40°E,30–60°N);
- contextualize the results for the Paris agglomeration with respect to other European cities.

The manuscript is organized as follows. The MAIAC product and its use are described in Section 2. Previous validation studies of the MAIAC product in Europe have been performed in Italy (Stafoggia et al., 2017), the Moscow metropolitan area (Zhdanova et al., 2020) and Germany (Falah et al., 2021), but no analysis have considered the entire European continent. Therefore, a validation analysis for Europe is also provided in Section 2. The discussion of the AOD climatology and trends over Europe and local/regional analysis will be presented and discussed in Section 3, before giving Conclusions in section 4.

2. Methods

2.1 MAIAC dataset extraction and analysis

The daily MCD19A2 product (Lyapustin and Wang, 2018) providing the AOD at 470 nm and 550 nm has been used over the period February 2000–August 2021. All the observations are delivered in the HDF4 format and stored at 1 km resolution in sinusoidal grid mapping. The product, distributed on a daily basis, contains the collection of each MODIS Aqua and Terra satellites overpasses, whose number varies according to the latitude. The uncertainty attributed to the MAIAC AOD retrievals has been defined through the expected error $EE=\pm(0.1AOD_{AERONET} + 0.05)$, indicating the percentage of AOD_{MAIAC} retrievals falling in the envelope (expressed in %). The EE has been established following (Falah et al., 2021) and (Lyapustin et al., 2018), considering both absolute and relative errors and by attributing an absolute error of 0.05 and a relative error of 0.1. As discussed in the next Section, the validation against AERONET will be considered to reevaluate the EE over Europe

120 and subsequently update the MAIAC uncertainty. In this product, in order to merge the satellite data to perform the climatological averages at the European scale (20°W–40°E,30–60°N), the daily average of each tile has been taken, followed by horizontal and vertical concatenation over the different MODIS tiles of interest. Only data classified as best–quality AOD (quality check flag “0000”) have been used in the following analysis. Although this choice reduces the number of available data, it guarantees the quality of the retrieval which is an important aspect to perform high resolution studies over urban areas.

Starting from the merged MAIAC data, the following treatment is applied:

- 125 ▪ Sinusoidal to WGS 84 grid coordinate system conversion.
- AOD daily averages are calculated for each grid point using available cloud-free observations taking into account available observations in the day from Terra and Aqua (i.e. 2 to 5 observations per day are available for the different grid points with Terra and Aqua overpasses times between 9 AM and 2 PM local time).
- 130 ▪ Local and regional scale AOD extractions have been performed to investigate the effect of the aerosol formation and city emissions over the surrounding areas. To this aim a list of 21 cities has been established, including European capital cities and big agglomerates with more that 1 million inhabitants. Those cities are listed in Table 1 and their location is plotted in Figure 1. The MAIAC AOD data have been extracted around the city locations using two different concentric kernels (centered on the nearest pixel to the longitude and latitude values of each city in Table 1): 3x3 km² (9 km² area) for the local scale and 100x100 km² (10000 km²) for the regional scale. The regional domain was chosen large enough to minimize effects of city’s pollution was chosen large enough in order to avoid effects due to the city and its plume, i.e. the local scale product occupies only ~0.09% of its regional background. Days for which a minimum of 40% spatial data coverage is available are considered both for the local and the regional scale, the others are discarded for the analysis. The local–to–regional AOD ratio (LTRR) has been calculated for each available kernel extraction (after averaging over all the available AOD data satisfying the spatial threshold in the kernels), to quantify the local scale enhancement to the regional AOD by using the following formula:
- 135

140
$$LTRR = \frac{AOD_{local}}{AOD_{regional}} - 1 \quad (1)$$

Positive deviations of the LTRR highlights the positive contribution of the urban local scale to the regional background signal, considering that AOD_{local} intrinsically represents the sum of the local production and the possible regional advected AOD fractions. Conversely, negative deviations can be linked to the presence of a non–homogenous spatial aerosol distribution at the regional scale, as well as to a possible local sink of pollution. The former may result in a stronger regional background signal related to different aerosol sources surrounding the city which may conceal the urban local signal and reduce the pollution gradients.

145

Trend assessment on AOD has been conducted over annual averages of daily AOD data using the Original Mann–Kendall test (Hussain and Mahmud, 2019). Annual AOD averages are performed if at least 50 AOD daily data are available in the year, and trend evaluations are performed if at least 5 years data are available in the dataset. The output of the Mann–Kendall test provides the significance of the test (p–value) and the Theil–Sen slope (Theil, 1992; Sen, 1968). All the tests have been calculated assuming a significance level (α) of 0.05 and the trend is considered significant if p–value< α . The relative change has been calculated following (Colette et al., 2016):

150

$$RC(\% \text{ year}^{-1}) = \frac{s}{y_0} \quad (2)$$

where s is the Theil–Sen slope and y_0 is the first available year for the trend evaluation. ~~The uncertainty attributed to the MAIAC AOD retrievals has been defined through the expected error (EE) considering both absolute and relative errors by attributing an absolute error of 0.05 and a relative error of 0.1 following (Falah et al., 2021; Lyapustin et al., 2018). As discussed in the next Section, the validation against AERONET will be considered to reevaluate the EE over Europe and subsequently update the MAIAC uncertainty.~~

2.2 Validation against AERONET observations and revised MAIAC estimated error (EE) for Europe

The MAIAC AOD validation has been performed by comparing the 550 nm AOD with all the available acquisitions (207 sites) in the AERONET Version 3 ground-based sun photometers network over continental Europe (Giles et al., 2019). Version 3 Level 2 AERONET data have been used (last access: 16 May 2023). AERONET provides AOD measurements at four different wavelengths: 440nm, 675nm, 860nm, 1020nm. The AOD at 550 nm has been extrapolated by assuming a power law relationship with the Angstrom exponent α (Ångström, 1929; Schuster et al., 2006) calculated between 440 nm and 675 nm:

$$AOD_{550} = AOD_{675nm} \left(\frac{455040}{675} \right)^{-\alpha} \quad (3)$$

$$\alpha = - \frac{\log \left(\frac{AOD_{440nm}}{AOD_{675nm}} \right)}{\log \left(\frac{440}{675} \right)} \quad (4)$$

Since the AERONET measurements are taken at different elevation angles depending on the sun elevation over the horizon, the measurements may be considered representative of a larger area around the point of acquisition (Chen et al., 2020; Schutgens, 2020; Kinne et al., 2013). ~~In order to improve the meaningfulness against the AERONET observations, the MAIAC AOD have been additionally extracted by taking an arbitrary average area of $0.06^\circ \times 0.06^\circ$ over the AERONET site, corresponding to $\sim 7 \times 7$ km², chosen between the 1×1 km² and 9×9 km² boxes for which (Falah et al., 2021) show that MAIAC – AERONET comparisons give similar results. In order to improve the meaningfulness against the AERONET observations, the MAIAC AOD have been additionally extracted by taking an average area of $0.06^\circ \times 0.06^\circ$ over the AERONET site, corresponding to $\sim 7 \times 7$ km². Indeed (Falah et al., 2021) show that MAIAC – AERONET comparisons given similar results for boxes between 1×1 km² and 9×9 km².~~ Furthermore, AERONET AOD data between $\pm 1H$ of the satellite passage have been considered to compare with MAIAC. The uncertainty on the AERONET AOD is $\Delta AOD_{AERONET} = 0.02$ linked to calibration uncertainty (Sinyuk et al., 2020). As the differences between MAIAC and AERONET observations are attributed entirely to MAIAC uncertainty, the derived MAIAC expected error is conservative. Different statistical indicators have been calculated to evaluate the comparison between MAIAC and AERONET AOD data. Those the Mean Bias Error (MBE), the Normalized Mean Bias (NMB), the Root Mean Square Error (RMSE), and the fraction of data within a factor of two (FAC2), as defined below (N is the number of data points):

$$MBE = \frac{1}{N} (AOD_{MAIAC} - AOD_{AERONET}) \quad (5)$$

$$NMB = \frac{\sum (AOD_{MAIAC} - AOD_{AERONET})}{\sum AOD_{MAIAC}} \quad (6)$$

$$RMSE = \sqrt{\frac{\sum (AOD_{MAIAC} - AOD_{AERONET})^2}{N}} \quad (7)$$

185

$$FAC2 (\%) = \text{fraction of data satisfying } 0.5 \leq \frac{AOD_{MAIAC}}{AOD_{AERONET}} \leq 2 \quad (8)$$

The correlation between AOD_{MAIAC} and $AOD_{AERONET}$ has been evaluated through the Pearson correlation coefficient R . The slope and the intercept of the regression line have been calculated taking into account the uncertainty in both coordinates using the York regression (York et al., 2004). The comparison between MAIAC and AERONET AOD at 550 nm for all available European AERONET measurements from 2000 to 2021 is shown in Figure 2. The overall validation performed considering the entire dataset (panel a) shows a slight underestimation of the AOD from MAIAC with respect to AERONET, with a MBE (−0.02) and a RMSE (0.06) values similar to those retrieved in previous validation studies (Chen et al., 2020; Lyapustin et al., 2018; Martins et al., 2017). The probability density function (PDF) of the MAIAC–AERONET absolute differences (panel b in Figure 2), shows a mean value and a sigma of −0.02 and 0.06, respectively. 77% of the AOD retrievals fall in an $EE = \pm(0.05AOD + 0.05)$, with a relative error lower than the validation $EE = \pm(0.1AOD + 0.05)$ from (Falah et al., 2021) accounting for observations in Northern Africa, California and Germany, but comparable to the EE envelope (~74% of points falling within the EE) obtained in (Qin et al., 2021) over the Köppen climate classification of normally humid and warm climate (Cf) region, including part of the European domain.

Since dependency of EE on aerosol type and size has been evidenced by (Falah et al., 2021) a further detailed validation depending on the Angstrom Exponent (AE) between 440 nm and 870 nm has been performed and presented in Figure 2 (panels c, d, e). The AE, combined with AOD, is an indicator of the particle type and size. AE values lower than 1 can be associated to coarse-mode aerosols (sea-salt and dust), whereas AE values higher than 2 to fine-mode aerosols (urban pollution and smoke) (Schuster et al., 2006). In this regards, three different classes depending on AE have been identified, respectively referred to coarse, mixed and fine aerosol particles: $AE < 0.5$, $0.5 \leq AE < 1.5$, $AE \geq 1.5$. The MAIAC validation shows an R value of 0.84 for the overall validation (panel a) comparable with the 0.85, 0.81 and 0.87 for the coarse, mixed and fine classes respectively. The validation for the mixed- and fine-dominated classes show a satisfactory accuracy of the product, with an MBE of −0.02 for both and 79% and 81% of the points, respectively, respectively within the envelope of $EE = \pm(0.1AOD + 0.05)$ from (Falah et al., 2021). However, for the coarse-mode, the MAIAC validation shows an MBE of −0.08 and 46% of points, within the envelope of $EE = \pm(0.1AOD + 0.05)$ from (Falah et al., 2021), significantly lower with respect to the other two classes. In case of $AOD < 0.25$ (84% of points in the coarse-mode validation), attributable to a marine-dominated aerosols scene (Toledano et al., 2007), 51% of points are within the EE, whereas for $AOD > 0.25$ (16% of points in the coarse-mode validation), attributable to dust-dominated aerosols (Rogozovsky et al., 2023; Bodenheimer et al., 2021; Toledano et al., 2007), the % of points within EE is significantly lower than 1 sigma (i.e. 68% of points falling in the EE envelope) However, for the coarse-mode the MBE (−0.08) and EE (46%) are significantly lower with respect to the other two classes. In case of $AOD < 0.25$ (84% of points in the coarse-mode validation), attributable to a marine-dominated aerosols scene, the EE is 51%, whereas for $AOD \geq 0.25$ (16% of points in the coarse-mode validation), attributable to dust-dominated aerosols, the EE is significantly lower than 1 sigma. As a matter of fact, a higher EE is needed to contain the 68% of the MAIAC–AERONET differences for the coarse-mode validation. (Rogozovsky et al., 2023; Qin et al., 2021) show that the MAIAC algorithm is sensible to the aerosol size. (Rogozovsky et al., 2023) observed that the underestimation of MAIAC compared to AERONET is related to the presence of dust (characterized by high depolarization ratio and small AE). This result suggests that, further improvements are needed in case of coarse-mode dominated classification. As a matter of fact, a higher EE is needed to contain the 68% of the MAIAC–AERONET differences for the coarse-mode validation. (Qin et al., 2021) show as MAIAC regional background models seem to be affected by local aerosol properties as they are retrieved by AERONET climatology, suggesting that, further improvements are needed in case of coarse-mode dominated classification.

In summary, results of the validation against AERONET suggest that the EE for MAIAC for observations over Europe between
225 2000 and 2021 can be estimated at $EE = \pm(0.05AOD + 0.05)$, lower than the EE estimated by (Falah et al., 2021). The total
MAIAC AOD uncertainty has been therefore revised to take into account this new estimation.

3. Results and discussions

3.1 European scale analysis

Before looking at the fine scale variability of the major European cities (section 3.2), we address here the question of their
230 AOD European background levels and their seasonal variation, as seen by multi-year MAIAC observations. We place our
findings in the context of previous analysis mainly based on spatially less refined MODIS observations (Gupta et al., 2023;
Filonchyk et al., 2020a; Wei et al., 2019), introducing an analysis based on two decades of data, extending and validating
studies performed on a shorter time periods.

The aerosol optical depth variability at European scale is shown in Figure 3, reporting seasonal averages, and in Figure S1 in
235 the Supplementary Information reporting the monthly averages of the AOD at 550 nm. The summer (JJA) season shows the
highest AOD values ranging between 0.12–0.22 in the 30°N–60°N band, whereas DJF shows the lowest AOD values ranging
between 0.06–0.09. Figure S1 depicts maps of monthly AOD averages and shows maximum values between April and July,
and minimum between November and January. A North–South latitudinal gradient is present for all the seasons, as shown in
Figure 3 and Figure S2, with maximum gradient during the summer (JJA) and minimum during the winter (DJF) season.
240 According to Figure S2, seasonal AOD averages range between 0.06–0.11 and 0.09–0.22 and in the 55°N–60°N and
30°N–35°N bands, respectively.

These findings with the MAIAC dataset are broadly in line with a previous analysis of MODIS and MISR data (Gupta et al.,
2023, 2022; Filonchyk et al., 2020a; Mehta et al., 2016). A North–South AOD gradient over Europe has been also found in
other MODIS studies (Merdji et al., 2023; Floutsi et al., 2016; Israelevich et al., 2012; Barnaba and Gobbi, 2004). Averages
245 over Western Europe and for the 2007–2016 period, (Zhao et al., 2018) find a broad spring summer AOD maximum extending
from April to July around 0.2 for MODIS Aqua and Terra and around 0.15 for MISR, and a winter December and January
minimum between 0.08 and 0.10. (Ma and Yu, 2015) attribute simulated spring maximum over Southern France and Corsica
over the western Mediterranean basin, especially to sulfate and dust, while other primary aerosol species (sea salt, primary
carbonaceous aerosol) show lower contributions and a flat seasonal variation. For secondary aerosol as sulfates, these larger
250 AOD values during the late spring, summer season are attributed to stronger photochemical activity due to increased oxidant
capacity (enhanced OH and ozone levels), whereas the contrary is expected for the late autumn, early winter minima. However,
these authors did not take into account secondary organic aerosol, it can be expected to be maximum during summer caused
by the higher biogenic volatile organic compounds (BVOCs), and increased photochemical activity during this season (Gao et
al., 2022). Possible fire events can also affect summer AOD peaks over Europe since they are more frequent during this period
255 (European Commission et al., 2022; Zhao et al., 2018).

Different aerosol hotspots, as previously identified in the literature (Coelho et al., 2022; Backes et al., 2016; Gkikas et al.,
2016; Bovchaliuk et al., 2013; Vecchi et al., 2009) are also visible in Figure 3, especially the Mediterranean Sea, the Po Valley
and Eastern Europe. The Mediterranean basin (6°W, 36E, 30°N, 46°N) is affected both by anthropogenic, biogenic and dust
emissions (Dayan et al., 2020; Chazette et al., 2019; Michoud et al., 2017; Pace et al., 2006). Its AOD seasonal cycle shown
260 in Figure 3 and ranging between 0.07–0.19 (average values obtained from the ocean part of the basin), follows the Saharan
dust transport cycle for the southern part, whereas the northern part is mostly dominated by human activities. In fact, the high

AOD MAM values (between 0.2 and 0.3) shown over the South–South East part of the Mediterranean basin are caused by the on–set of the Saharan dust transport due to a low–pressure system (the Sharav cyclone), which pushes the dust plumes to the eastward basin (Floutsi et al., 2016; Moulin et al., 1998). During summer, the Balearic cyclogenesis is causing the spreading of the dust plumes northwards from the Saharan source region, explaining the high JJA values (AOD>0.2) over large areas of the Southern part of the basin (Formenti et al., 2018; Floutsi et al., 2016; Moulin et al., 1998). The AOD average over February 2000–August 2021 period over the Mediterranean basin (6°W, 36E,30°N, 46°N) resulted in an AOD of 0.13 at 550 nm, comparable to the result obtained in (Chiapello et al., 2021) at 865 nm based on POLDER–3 observations.

In the Po Valley (7°E, 12°30'E, 43°36'N, 46°12'N), the seasonal cycle is ranging between 0.09 (for DJF) to 0.15 (for JJA) with maxima in June, July (AOD>0.16). Particulate matter (PM) composition measurements at the ground show to be dominated by traffic, biomass burning emissions, as well as ammonium nitrate and sulfate formation (Scotto et al., 2021) and the largest ground PM values can occur in DJF and SON seasons due to recurrent low temperatures and possible intense residential biomass burning (Pietrogrande et al., 2015) and ammonium nitrate precursor emissions (Scotto et al., 2021; Vecchi et al., 2018). The MAM and JJA levels at the ground can be caused by agricultural local sources (e.g. burning of pruning and fertilizers) (Scotto et al., 2021; Bucci et al., 2018; Clarisse et al., 2009). In summertime high levels of secondary organic aerosols in presence of stagnation conditions have been also observed in (Sandrini et al., 2016). Since AOD values are columnar values, the difference observed in the seasonal cycle between the ground (mainly DJF–MAM peaks) and AOD (JJA) can be attributed to different reasons: 1) the planetary boundary layer height (PBL), lower in the winter and higher in the summer, which conversely to ground PM is not affecting the AOD measurements and 2) possible dust events and biomass burning fires that can contribute to the stronger AOD levels during the spring and summer seasons. In addition, high AOD levels are also favored by insufficient pollution dispersion and removal, the valley being surrounded by mountains (the Alps and the Apennines), especially under stable weather conditions, promoting pollutants accumulation and air masses stagnation (Putaud et al., 2014). As a matter of fact, this reasoning is general and not restricted to Po valley.

For what concerns Eastern Europe (13°E, 30°E, 42°N, 55°N), Figure 3 shows a strong seasonal variability of AOD for regions like Poland and Serbia, with maximum AOD of up to 0.2 during the JJA season and minimum over the DJF season with AOD generally below 0.15. Seasonal cycle with a maximum over summer and spring has been also observed in (Chubarova et al., 2016) for the 2001–2014, studying the Moscow AERONET site. Furthermore, (Bovchaliuk et al., 2013), found AOD values ranging between 0.05 and 0.2 at 870 nm for the 2003–2011 period, with peaks over the spring, and which the authors explain by agricultural fires correlated with an observed increase in the fine fraction mode particles.

Finally, also the Western Europe (11°W, 18°E, 35°N, 60°N) shows an AOD seasonal variability over land in the [0.06–0.12] range with the maximum on the JJA season. A summer AOD maximum attributable to dust, smoke and sea–spray aerosols has been also found in (Zhao et al., 2018) for this area. These values are lower compared to other regions within Europe. This result can be justified by its proximity to the Atlantic Ocean, which contribute to expose these areas to more humid and less polluted air masses as well as to a greater pollutants dispersion capability.

3.2 City scale analysis

The 1 km resolution MAIAC AOD data are used to explore AOD levels over cities and evaluating and quantifying the extra different AOD levels -of cities with respect to their surrounding areas. Figure 4 shows the distribution and the heterogeneity of the aerosol optical depth over the different sites. European cities are ordered by increasing 50th percentile values of the local scale AOD from left to right. Table 1 gives the coordinates of city centers. An example of AOD time series for some of the cities is shown in Figure S3. In line with Section 3.1, a North–South gradient can be found among the cities as well,

highlighting that European cities located at more northerly latitudes have AOD levels at 550 nm in general lower compared to cities at more southerly latitudes: Oslo, Dublin and London, show AOD median values (the 25th and 75th percentiles are also reported in brackets) of 0.06 [0.03–0.10], 0.06 [0.04–0.10] and 0.07 [0.04–0.12] respectively, whereas more southerly located Bologna, Milan and Athens show 0.13 [0.08–0.19], 0.14 [0.07–0.23], 0.16 [0.11–0.23] respectively. Sites located in the middle range of the Figure 4, like Lisbon, Berlin and Amsterdam, show AOD median values 0.09 [0.07–0.12], 0.10 [0.05–0.16], 0.09 [0.06–0.16]. The AOD values in the northern cities are only rarely exceeding a threshold of 0.3, which we arbitrarily relate to pollution (anthropogenic, dust, fires...) events (0.6%, 2.5% and 4.0% of the total observations for Oslo, Dublin and London respectively, see Table 2). For more southerly cities like Milan and Athens, this fraction is 14.7% and 10.6% respectively. Looking more closely to the timing of these occurrences, 18% and 10% of these “high pollution” cases occurred before and after 2010 respectively for Milan, and 14% and 8% for Athens.

Figure 4 shows that the city center local scale AOD levels are most of the cities considered in our study time (58.8%) larger than the regional AOD. As well, an increase in the frequency of AOD>0.3 can be also observed for the city AOD (See Table 2). The local-to-regional ratio (LTRR) calculated using Eq. (1) for the 2000–2021 period and for the different cities is summarized in Table 2. Again, positive LTRR values are characteristic for an urban scale contribution to the aerosol burden on top of the regional one, highlighting the importance of local anthropogenic emissions and atmospheric conditions favorable to pollutants accumulation. For instance, a LTRR value of 1 would correspond to a 50% contribution of local urban aerosol to total AOD, while a value of 0.5 to a contribution of a third. It should be noted that the local contribution to surface PM is necessarily stronger than that to AOD, as the importance of the regional background is more important for the vertical column.

On the contrary, negative LTRR values indicate a lower local city AOD than the regional one, suggesting a possible inhomogeneity in AOD within the rather large (100 x 100 km²) regional domain since the observed negative LTRR values were in general very small (in the order of some %). This could be true especially for coastal sites, or partly mountainous sites, where topography plays an important role. Furthermore, this inhomogeneity may be related to i) the spatial extent of the city, which may impact the AOD levels of the regional scale, ii) the different location of emission sources, such as the location of industrial areas, which, combined with favorable meteorological conditions can lead to inhomogeneous spatial patterns in the regional domain.

An alternative explanation to negative LTRR values would be local aerosol loss at urban scale. However, systematic urban loss processes are not easy to identify. Sedimentation and dry deposition processes are not expected to be particularly enhanced over urban areas, nor is precipitation, compared to its regional surrounding. On the other hand, the urban heat island with increased temperatures could lead to evaporation of particles. For instance, (Pirhadi et al., 2020) finds that due to its semi-volatile character, about 50% of ambient PM_{2.5} aerosol mass evaporated when heated up in a thermo-denuder from ambient temperature (~13°C in winter, 23°C in summer and up to 50°C). The urban heat island effect depends on the size and additional heat production within an urban area. It is restricted to light wind meteorological conditions and it is more pronounced during night, while MAIAC observations are made during daytime. For these reasons, we consider that evaporation of semi-volatile aerosol under higher urban temperatures could only play a limited effect in our dataset. In the frame of the present analysis it is in general difficult to distinguish between these two loss and inhomogeneity effects.

On the contrary, negative LTRR values indicate a lower local city AOD than the regional one, suggesting a local aerosol loss at urban scale. However, systematic urban loss processes are not easy to identify. Sedimentation and dry deposition processes are not expected to be particularly enhanced over urban areas, nor is precipitation, compared to its regional surrounding. On the other hand, the urban heat island with increased temperatures could lead to evaporation of particles. For instance, (Pirhadi et al., 2020) finds that due to its semi-volatile character, about 50% of ambient PM_{2.5} aerosol evaporated when heated up in a

~~thermo-denuder from ambient temperature (-13°C in winter, 23°C in summer and up to 50°C). The urban heat island effect depends on the size and additional heat production within an urban area. It is restricted to light wind meteorological conditions and it is more pronounced during night, while MAIAC observations are made during daytime. For these reasons, we consider that evaporation of semi-volatile aerosol under higher urban temperatures could only play a limited effect in our dataset.~~

345 ~~Since the observed negative LTRR values were in general very small (in the order of some %) an alternative explanation to negative LTRR values is a possible inhomogeneity in AOD within the rather large ($100 \times 100 \text{ km}^2$) regional domain. This could be true especially for coastal sites, or partly mountainous sites. In the frame of the present analysis it is in general difficult to distinguish between these two loss and inhomogeneity effects.~~

Maximum mean values of LTRR are shown for Barcelona (0.57 ± 0.02), Lisbon (0.55 ± 0.03), Paris (0.39 ± 0.02) and Athens
350 (0.32 ± 0.01). On the contrary, significantly negative LTRR values are shown for Brussels (-0.06 ± 0.01), Amsterdam (-0.17 ± 0.01), Berlin (-0.03 ± 0.01). The uncertainty has been calculated here as the standard error of the mean: σ/\sqrt{N} , where σ is the standard deviation of the LTRR distribution, and N is the number of points available over the 2000–2021 period. The most negative LTRR is found for Amsterdam. For this coastal city, larger AODs are observed over the sea than over the continent (see Figure 3 especially for the spring and autumn seasons) which could be caused by enhanced sea-salt, but possibly also by
355 slight differences in the retrieval algorithm for sea and land surfaces. Thus, the regional background cannot be considered as homogeneous for this case.

In this study we focus on the Paris area which shows a LTRR of 0.39 ± 0.02 . The aim of this interest is supporting the preceding climatological studies performed for the ACROSS field campaign (Cantrell and Michoud, 2022). Paris represents a strongly centralized agglomeration with about 11 million inhabitants. It is located in Western Europe, in a rural area without strong
360 orography, and at some 200 km from the Atlantic Ocean. This leads to generally favorable pollutant dispersion conditions (Vautard et al., 2003). The median local AOD value at 550 nm is 0.10 [0.07–0.15] for Paris which falls slightly over the median of the cities distribution in Figure 4. Results from MEGAPOLI (Beekmann et al., 2010) campaign have shown that large fraction of fine PM at the ground is transported from the European continent and southern France towards Paris, while local emissions represent a smaller fraction (Beekmann et al., 2015; Bressi et al., 2014). Later studies with multi-year data
365 sets (mid 2011–mid 2013, (Petit et al., 2015)) or pointing to specific pollution episodes (December 2016, (Foret et al., 2022)) make evident the local emission contribution to fine aerosol pollution peaks. (Skylakou et al., 2014) shows by source apportionment that primary organic aerosol (POA) and elemental carbon (BC) are controlling the PM_{2.5} fraction of the Paris local emissions, whereas regional advection is controlling the secondary PM_{2.5} fraction. Organic aerosols have been shown to play a key role in the Paris air quality assessment (Zhang et al., 2019; Petit et al., 2015; Bressi et al., 2014). Sulfate and
370 secondary organic aerosols are mainly attributed to long-range transport (Foret et al., 2022; Skylakou et al., 2014). For the period 2000–2021, the % of days with AOD>0.3 is found to be 4% at the Paris local scale.

Barcelona shows a local/regional distribution rather similar to Paris (Figure 4) although its LTRR is larger (0.57 ± 0.02 , actually the largest in our study) and the geomorphology of the two sites is significantly different. ~~Barcelona shows the largest LTRR (0.57 ± 0.02) among the cities studied.~~ It is located on the northeast part of the Iberian Peninsula, bordering the Mediterranean
375 Sea and the foot hills of the Pyrenees mountains. Re-circulation caused by mountain winds and sea breeze (Jaén et al., 2021; Pérez et al., 2004) could enhance the local urban AOD. In addition, regional background over the mountainous area next to Barcelona is relatively low with respect to that over the Mediterranean sea (Figure 3), which could contribute to the large LTRR.

The Bologna and Milan surrounding is the well-known Po-Valley, as previously discussed, where recirculation and stagnation
380 events of aerosol and precursors may occur and cause enhanced pollution levels (Putaud et al., 2014). (Vecchi et al., 2018)

showed by source apportionment analysis that, during the winter season, major PM contribution to light extinction for the Milan urban area is nitrate (42%), followed by sulfate, primary aerosol due to traffic and biomass burning related organic aerosol. In another study, secondary inorganic aerosols have been also shown to contribute with 35% on PM over the Milan urban area on the annual average (Amato et al., 2016). As a consequence, the large regional PM background leads to comparatively small additional local contributions and small LTRR values for these both cities.

For what concern Athens with a LTRR of 0.32 ± 0.01 , aerosols of anthropogenic–origin have been shown to dominate. Indeed (Taghvaei et al., 2019) showed by source apportionment analysis that traffic emissions, SOA and biomass burning correspond to major sources to PM_{2.5} samples, contributing respectively 44%, 16% and 9%, with higher PM values during summer than winter. During the latter season high PM_{2.5} episodes are linked to dust and biomass burning episodes (Raptis et al., 2020). Furthermore, the organic aerosol concentrations in Athens have been shown to be dominated by regional SOA during summertime (Tsiflikiotou et al., 2019), highlighting also the importance of long–range transport in this area (Manousakas et al., 2020). As a conclusion of this discussion, PM_{2.5} sources over the Athens region are a mixture of regional and local origin, which is reflected its LTRR value.

For the seasonal variation of LTRR values, Figure 5 shows the scatter plot between the local and regional AOD as a function of the season for all cities. The fitting line considering all seasons shows a slope of 1.08 and a linear correlation of 0.893, highlighting the overall average positive contribution on air quality degradation of the local on the regional scale over the different seasons. Figure 5 shows that the fraction of points where $AOD_{local} > AOD_{regional}$ is 84 %, 65 %, 75 %, 97 % for DJF, MAM, JJA, SON respectively. This result suggests that the local contribution is higher during winter and lower during summer. In order to explain this difference, it should be considered that during summer time, favorable weather condition, stronger photochemistry activity and enhanced BVOC emissions can lead to increased secondary aerosol formation, and increase the secondary to primary aerosol ratio. As secondary aerosol formation is a regional phenomenon (Beekmann et al., 2015; Skyllakou et al., 2014; Karl et al., 2009), the regional contribution to AOD is increased. Furthermore, possible dust and fire events can also contribute to the increase of the regional signal during summer and spring over Europe. However, during wintertime secondary aerosol formation is less pronounced, in addition primary aerosol emissions are increased due increased heating demand.

3.3 Trend Analysis

The analysis of the high resolution MAIAC product can contribute to further investigate the aerosol optical depth tendency over the European region. Statistically significant ($pvalue < 0.05$) absolute and relative AOD trends over the European continent are reported in Figure 6. Negative AOD trends have been found over the domain of interest, in the $[-3; -0.6]$ %year⁻¹ range, representing the 5th and 95th percentile respectively of the Figure 6b. Furthermore, more negative trends are mostly found over the regional hotspots outlined in section 3.1 (Po valley, Mediterranean basin, parts of eastern Europe). Decreasing relative and absolute trends of -1.34 ± 0.29 %year⁻¹ and -0.0021 ± 0.0005 units year⁻¹ for the Mediterranean Basin has been found for the 2001–2021 period. A decreasing absolute trend of -0.003 units year⁻¹ for the 2002–2014 period has been also found with the MAIAC data in agreement to the -0.003 units year⁻¹ observed in (Floutsi et al., 2016). A trend of -1.66 ± 0.58 %year⁻¹ at 550nm has been estimated for the Po Valley, lower than what has been observed at the Ispra AERONET site in the period 2004–2010 (Putaud et al., 2014) where they estimated a decreasing trend of -4.0 ± 1.8 and -2.5 ± 1.3 %year⁻¹ for the 440nm and 675nm respectively. Negative AOD tendency has been also registered for the Benelux and the Peloponnese area of -2.46 ± 0.96 %year⁻¹ and -1.49 ± 0.45 %year⁻¹ respectively. A statistically absolute significant trend of -0.003 ± 0.002 units year⁻¹ has been observed for the Eastern Europe area, in line to what observed in (Filonchik et al., 2020a) for the 2002–2018 period,

420 where values in the range of $[-0.0025; -0.0028]$ units year⁻¹ are observed for Czech Republic, Bulgaria, Slovakia and Hungary
with MODIS TERRA data. However stronger trends for the 2002–2019 period, in the range of $[-0.0031; -0.0076]$ units year⁻¹,
425 are observed in (Filonchik et al., 2020b) for several cities in the Eastern Europe, by using MODIS AQUA data. (Chubarova
et al., 2016) attributes the significant negative trends observed in Moscow for the 2001–2014 period to the strong decrease in
SO_x, non-methane volatile organic compounds (NMVOC) and NO_x emissions. As a matter of fact, (Tsyro et al., 2022)
430 predicted, through a six-models ensemble approach, a decreasing of surface PM_{2.5} and PM₁₀ over Europe for the 1990–2010
period, attributing a large impact to sulfate, ammonium and nitrate precursor emission reductions. Nevertheless, this decrease
appeared to be more impacting over Central and Eastern Europe. For instance, trends stronger than -2.5 %year⁻¹ are observed
over Germany, Czech Republic, Hungary, Slovakia and Ukraine both for PM_{2.5} and PM₁₀.

Taking advantage of the high spatial resolution of the MAIAC product, the analysis has been extended to a local scale to
430 estimate the AOD trends over the cities listed in the Table 2 and to compare them to the trends for the surrounding regional
background. Results are shown and summarized in Table 3. For most of the sites, a significant negative trend can be identified
consistently both within the city center (3x3km²) and in the surrounding area (100x100km²). For instance, Athens, Prague,
Vienna, Milan, Zagreb and Bologna show AOD trends of in the range of $[-0.9; -1.7]$ %year⁻¹ and $[-1.3; -2.0]$ %year⁻¹ for the
435 regional and local scale respectively. This result is in line with the aforementioned observations at the European scale and with
other studies focusing on European megacities (Papachristopoulou et al., 2022; Gupta et al., 2022; Zhao et al., 2017).
(Papachristopoulou et al., 2022) observed a decrease in AOD up to -0.03 units decade⁻¹ over the 2003–2020 period and for
European megacities like Paris, Barcelona, Madrid and London. This result is comparable to the range of $[-0.01; -0.03]$ units
decade⁻¹ observed for the European cities analyzed in this study. Conversely, cities where AOD levels are relatively low
(positioned in the leftmost part of the Figure 4) do generally not show statistically significant results. Among all the cities,
440 Prague shows the strongest relative trend at both regional and local scales of -2.0 %year⁻¹ and -1.7 %year⁻¹ respectively. The
absolute value obtained in this study for the local scale is comparable to -0.0022 obtained in (Filonchik et al., 2020a) for the
2000–2018 period. Moreover, the -1.0 %year⁻¹ estimated trend over Athens is in line with the -1.1 %year⁻¹ obtained at 440
nm at the AERONET urban site in (Raptis et al., 2020) for the 2000–2018 period. In the case of Paris, a trend of -1.5 %year⁻¹
445 is obtained for the regional scale, while the city center trend is not significant. Interestingly, the regional relative trend is for
most cities (9 of 11 in Table 3) stronger (i.e. more negative) than the local one. One possible reason of this outcome could be
a stronger decrease of secondary aerosols due to stringent pollution control of precursors (SO₂, NO_x, VOC) than that of
primary aerosol, as found for several French EMEP/MERA network sites (Font et al., 2023). Indeed, the primary to secondary
aerosol ratio is expected to be larger for urban than for regional background sites.

4. Conclusions

450 This study presents a quantitative estimation of the aerosol optical depth variability in Europe using long-term measurements
(2000–2021) from the MAIAC algorithm applied to MODIS satellite observations. The MAIAC validation, performed at the
European scale against ground-based sun photometer data, demonstrates a slight underestimation of MAIAC AOD, showing
a MBE of -0.02 and a RMSE of 0.06 respectively. An expected error $EE = \pm(0.05AOD + 0.05)$ has been found for the
European continent, lower with respect to what suggested by (Lyapustin et al., 2018). Moreover, according to the AERONET
455 AE splitting analysis, the validation, which provides satisfactory results for mixed- and fine- aerosol mode, is less performant
in presence of coarse-dominated aerosols. This suggests that further improvements of the MAIAC algorithm are needed for
scenes dominated by dust or other coarse-sized particles.

Regarding the AOD seasonal climatology over the European continent, the AOD exhibits maximum and minimum values during the summer (JJA) and winter (DJF) seasons, respectively, showing a strong North–South latitudinal gradient. Values of AOD in the range of 0.12–0.22 (JJA) and 0.06–0.09 (DJF) are observed in the 30°N–60°N band.

Concerning the link between regional and local scales air quality across the main European cities that was the main objective of this work both the regional background and city level AOD show a general north–south gradient with increasing AOD and several hotspots over the Po valley and the Mediterranean Sea. The local–to–regional analysis shows that most of the cities contribute to enhance the AOD loading with respect to their regional background. On the contrary, for some cities a slightly negative LTRR could be either explained by specific losses or by an inhomogeneity of the regional background. On a relative scale the city contribution to regional AOD is maximum during the winter season, because the primary vs. secondary aerosol ratio is expected to be the largest. Concerning the Paris area, most of pollution has been considered as transported from the European continent in previous studies (Beekmann et al., 2015). In fact, Paris represents an important isolated agglomeration with respect to the surrounding area. Indeed, the long–term analysis conducted in this work indicates an average local–to–regional excess ratio of 39%, suggesting a non–negligible impact of the city emissions in addition to the regional aerosol burden in Paris. Further investigation is needed to understand the nature of this AOD difference. As a matter of fact, the interaction between the regional background and the local emissions cannot be exploited through AOD measurements directly, although we know that changes in the chemical and optical properties lead to changes in the aerosol extinction profile. Further investigation on the interaction between biogenic and anthropogenic local and regional air masses and the impact on aerosol properties will be provided in the ACROSS project (Cantrell and Michoud, 2022).

Different studies have already shown a negative decreasing AOD trends over the European continent using in particular MODIS satellite data. However, most of the time with a broader spatial resolution with respect to the product used in this study. The MAIAC high spatial resolution product has been exploited to investigate the AOD trends both at European and local city scale. The result showed a general AOD decrease over the all European continent, consistent with the recent literature and in connection with the mitigation policies over the European countries. In particular, spatial homogeneous trends have been found over known European hotspots (e.g. $-1.34 \text{ \%year}^{-1}$, $-1.66\% \text{ year}^{-1}$, for the Mediterranean Sea, the Po Valley respectively). In addition, taking advantage of the high spatial resolution, the analysis has been extended also at the city level, showing a statistically significant yearly decrease during the last two decades in AOD at 550 nm in the range of $[-0.5; -1.7] \text{ \%year}^{-1}$ and at city level and $[-0.9; -2.0] \text{ \%year}^{-1}$ in the surroundings. This result highlights the faster decrease in regional AOD levels with respect to those at the urban local scale. Nevertheless, over the Paris area, we observed a statistically significant negative trend only at the regional scale. A potential explanation could be linked to the more stringent control of aerosol precursors emissions (SO_2 , NO_x , VOC) with respect to direct aerosol emissions (Font et al., 2023).

Author contributions

LDA, CDB, and MB designed the study and discussed the results. LDA performed the data analysis with contributions by CDB and MB. GF, PF, GS, and JFD contributed to the discussion of the results. LDA, CDB, and MB wrote the paper with contribution from all co–authors.

Data access

The MAIAC MCD19A2 data used in this study are accessible at: <https://lpdaac.usgs.gov/products/mcd19a2v006/> (last access: June 2023). The AERONET data are available at <https://aeronet.gsfc.nasa.gov/> (last access: June 2023). Population data

495 reported in Table 1 can be accessed at the following link:
<https://ec.europa.eu/eurostat/cache/RCI/#?vis=city.statistics&lang=en> (last access: June 2023). The data and the codes that support the findings of this study are available upon request from the corresponding authors.

Competing interests:

The authors declare that they have no conflict of interest.

500 Acknowledgements

This work has been supported by the ACROSS and the RI-URBANS projects. The ACROSS project has received funding from the French National Research Agency (ANR) under the investments programme integrated into France 2030, with the reference ANR-17-MPGA-0002, and it was supported by the French National program LEFE (Les Enveloppes Fluides et l'Environnement). The RI-URBANS project has received funding from the European Union's Horizon 2020 research and
505 innovation program under grant agreement No 101036245. We thank all the AERONET PIs and their staff for establishing and maintaining all the sites used in this investigation. Useful discussions with M. Mallet, Y. Derimian, and J. C. Raut are gratefully acknowledged. We thank C. Cantrell and V. Michoud, PIs of the ACROSS project. [Helpful comments by the reviewers are gratefully acknowledged.](#)

References

510 Alpert, P., Shvainshtein, O., and Kishcha, P.: AOD Trends over Megacities Based on Space Monitoring Using MODIS and MISR, 2012, <https://doi.org/10.4236/ajcc.2012.13010>, 2012.

Aman, N., Manomaiphiboon, K., Suwattiga, P., Assareh, N., Limpaseni, W., Suwanathada, P., Soonsin, V., and Wang, Y.: Visibility, aerosol optical depth, and low-visibility events in Bangkok during the dry season and associated local weather and synoptic patterns, *Environ. Monit. Assess.*, 194, 322,
515 <https://doi.org/10.1007/s10661-022-09880-2>, 2022.

Amato, F., Alastuey, A., Karanasiou, A., Lucarelli, F., Nava, S., Calzolari, G., Severi, M., Becagli, S., Gianelle, V. L., Colombi, C., Alves, C., Custódio, D., Nunes, T., Cerqueira, M., Pio, C., Eleftheriadis, K., Diapouli, E., Reche, C., Minguillón, M. C., Manousakas, M.-I., Maggos, T., Vratolis, S., Harrison, R. M., and Querol, X.: AIRUSE-LIFE+: a
520 *Phys.*, 16, 3289–3309, <https://doi.org/10.5194/acp-16-3289-2016>, 2016.

Ångström, A.: On the Atmospheric Transmission of Sun Radiation and on Dust in the Air, *Geogr. Ann.*, 11, 156–166, <https://doi.org/10.1080/20014422.1929.11880498>, 1929.

Backes, A. M., Aulinger, A., Bieser, J., Matthias, V., and Quante, M.: Ammonia emissions in Europe, part II: How ammonia emission abatement strategies affect secondary aerosols, *Atmos. Environ.*, 126, 153–161,
525 <https://doi.org/10.1016/j.atmosenv.2015.11.039>, 2016.

Bahadur, R., Praveen, P. S., Xu, Y., and Ramanathan, V.: Solar absorption by elemental and brown carbon determined from spectral observations, *Proc. Natl. Acad. Sci.*, 109, 17366–17371, <https://doi.org/10.1073/pnas.1205910109>, 2012.

Bai, R., Xue, Y., Jiang, X., Jin, C., and Sun, Y.: Retrieval of High-Resolution Aerosol Optical Depth for Urban Air
530 Pollution Monitoring, *Atmosphere*, 13, 756, <https://doi.org/10.3390/atmos13050756>, 2022.

- Barnaba, F. and Gobbi, G. P.: Aerosol seasonal variability over the Mediterranean region and relative impact of maritime, continental and Saharan dust particles over the basin from MODIS data in the year 2001, *Atmospheric Chem. Phys.*, 4, 2367–2391, <https://doi.org/10.5194/acp-4-2367-2004>, 2004.
- 535 Bäumer, D., Vogel, B., Versick, S., Rinke, R., Möhler, O., and Schnaiter, M.: Relationship of visibility, aerosol optical thickness and aerosol size distribution in an ageing air mass over South-West Germany, *Atmos. Environ.*, 42, 989–998, <https://doi.org/10.1016/j.atmosenv.2007.10.017>, 2008.
- 540 Beekmann, M., Chiappini, L., Favez, O., Aymoz, G., Bessagnet, B., Rouil, L., and Rossignol, S.: The megapoli paris campaign for urban aerosol characterisation - a comprehensive data set for air quality model evaluation, in: 13. International Conference on Harmonisation within Atmospheric Dispersion Modelling for Regulatory Purposes (HARMO 13), Paris, France, 519–523, 2010.
- 545 Beekmann, M., Prévôt, A. S. H., Drewnick, F., Sciare, J., Pandis, S. N., Denier van der Gon, H. a. C., Crippa, M., Freutel, F., Poulain, L., Gherzi, V., Rodriguez, E., Beirle, S., Zotter, P., von der Weiden-Reinmüller, S.-L., Bressi, M., Fountoukis, C., Petetin, H., Szidat, S., Schneider, J., Rosso, A., El Haddad, I., Megaritis, A., Zhang, Q. J., Michoud, V., Slowik, J. G., Moukhtar, S., Kolmonen, P., Stohl, A., Eckhardt, S., Borbon, A., Gros, V., Marchand, N., Jaffrezo, J. L., Schwarzenboeck, A., Colomb, A., Wiedensohler, A., Borrmann, S., Lawrence, M., Baklanov, A., and Baltensperger, U.: In situ, satellite measurement and model evidence on the dominant regional contribution to fine particulate matter levels in the Paris megacity, *Atmospheric Chem. Phys.*, 15, 9577–9591, <https://doi.org/10.5194/acp-15-9577-2015>, 2015.
- 550 Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, *Rev. Geophys.*, 58, e2019RG000660, <https://doi.org/10.1029/2019RG000660>, 2020.
- 555 Bender, F. A.-M.: Aerosol Forcing: Still Uncertain, Still Relevant, *AGU Adv.*, 1, e2019AV000128, <https://doi.org/10.1029/2019AV000128>, 2020.
- Bilal, M., Nazeer, M., Nichol, J. E., Bleiweiss, M. P., Qiu, Z., Jäkel, E., Campbell, J. R., Atique, L., Huang, X., and Lolli, S.: A Simplified and Robust Surface Reflectance Estimation Method (SREM) for Use over Diverse Land Surfaces Using Multi-Sensor Data, *Remote Sens.*, 11, 1344, <https://doi.org/10.3390/rs11111344>, 2019.
- 560 Bodenheimer, S., Nirel, R., Lensky, I. M., and Dayan, U.: The synoptic skill of aerosol optical depth and angstrom exponent levels over the Mediterranean Basin, *Int. J. Climatol.*, 41, 1801–1820, <https://doi.org/10.1002/joc.6931>, 2021.
- 565 Boers, R., Weele, M. van, Meijgaard, E. van, Savenije, M., Siebesma, A. P., Bosveld, F., and Stammes, P.: Observations and projections of visibility and aerosol optical thickness (1956–2100) in the Netherlands: impacts of time-varying aerosol composition and hygroscopicity, *Environ. Res. Lett.*, 10, 015003, <https://doi.org/10.1088/1748-9326/10/1/015003>, 2015.
- Bovchaliuk, A., Milinevsky, G., Danylevsky, V., Goloub, P., Dubovik, O., Holdak, A., Ducos, F., and Sosenkin, M.: Variability of aerosol properties over Eastern Europe observed from ground and satellites in the period from 2003 to 2011, *Atmospheric Chem. Phys.*, 13, 6587–6602, <https://doi.org/10.5194/acp-13-6587-2013>, 2013.
- 570 Bressi, M., Sciare, J., Gherzi, V., Mihalopoulos, N., Petit, J.-E., Nicolas, J. B., Moukhtar, S., Rosso, A., Féron, A., Bonnaire, N., Poulakis, E., and Theodosi, C.: Sources and geographical origins of fine aerosols in Paris (France), *Atmospheric Chem. Phys.*, 14, 8813–8839, <https://doi.org/10.5194/acp-14-8813-2014>, 2014.
- Bucci, S., Cristofanelli, P., Decesari, S., Marinoni, A., Sandrini, S., Größ, J., Wiedensohler, A., Di Marco, C. F., Nemitz, E., Cairo, F., Di Liberto, L., and Fierli, F.: Vertical distribution of aerosol optical properties in the Po

- 575 Valley during the 2012 summer campaigns, *Atmospheric Chem. Phys.*, **18**, 5371–5389, <https://doi.org/10.5194/acp-18-5371-2018>, 2018.
- Cantrell, C. and Michoud, V.: ACROSS: A Field Experiment to Study Atmospheric Oxidation Chemistry and Physics of Mixed Anthropogenic-Biogenic Air Masses in the Greater Paris Area, *Bull. Am. Meteorol. Soc.*, **1**, <https://doi.org/10.1175/BAMS-D-21-0115.1>, 2022.
- 580 Chazette, P., Totems, J., and Shang, X.: Transport of aerosols over the French Riviera – link between ground-based lidar and spaceborne observations, *Atmospheric Chem. Phys.*, **19**, 3885–3904, <https://doi.org/10.5194/acp-19-3885-2019>, 2019.
- Chen, C., Dubovik, O., Fuertes, D., Litvinov, P., Lapyonok, T., Lopatin, A., Ducos, F., Derimian, Y., Herman, M., Tanré, D., Remer, L. A., Lyapustin, A., Sayer, A. M., Levy, R. C., Hsu, N. C., Descloitres, J., Li, L., Torres, B., Karol, Y., Herrera, M., Herreras, M., Aspetsberger, M., Wanzenboeck, M., Bindreiter, L., Marth, D., Hangler, A., and Federspiel, C.: Validation of GRASP algorithm product from POLDER/PARASOL data and assessment of multi-angular polarimetry potential for aerosol monitoring, *Earth Syst. Sci. Data*, **12**, 3573–3620, <https://doi.org/10.5194/essd-12-3573-2020>, 2020.
- 585
- Chen, L., Wang, R., Wei, G., Han, J., and Zha, Y.: A surface reflectance correction model to improve the retrieval of MISR aerosol optical depth supported by MODIS data, *Adv. Space Res.*, **67**, 858–867, <https://doi.org/10.1016/j.asr.2020.10.033>, 2021.
- 590
- Chiapello, I., Formenti, P., Mbemba Kabuiku, L., Ducos, F., Tanré, D., and Dulac, F.: Aerosol optical properties derived from POLDER-3/PARASOL (2005–2013) over the Western Mediterranean Sea – Part 2: Spatial distribution and temporal variability, *Atmospheric Chem. Phys.*, **21**, 12715–12737, <https://doi.org/10.5194/acp-21-12715-2021>, 2021.
- 595
- Chubarova, N. Y., Poliukhov, A. A., and Gorlova, I. D.: Long-term variability of aerosol optical thickness in Eastern Europe over 2001–2014 according to the measurements at the Moscow MSU MO AERONET site with additional cloud and NO₂ correction, *Atmospheric Meas. Tech.*, **9**, 313–334, <https://doi.org/10.5194/amt-9-313-2016>, 2016.
- 600
- Clarisse, L., Clerbaux, C., Dentener, F., Hurtmans, D., and Coheur, P.-F.: Global ammonia distribution derived from infrared satellite observations, *Nat. Geosci.*, **2**, 479–483, <https://doi.org/10.1038/ngeo551>, 2009.
- Coelho, S., Ferreira, J., Rodrigues, V., and Lopes, M.: Source apportionment of air pollution in European urban areas: Lessons from the ClairCity project, *J. Environ. Manage.*, **320**, 115899, <https://doi.org/10.1016/j.jenvman.2022.115899>, 2022.
- 605
- Colette, A., Aas, W., Banin, L., Braban, C. F., Ferm, M., Gonzalez Ortiz, A., Ilyin, I., Mar, K., Pandolfi, M., Putaud, J.-P., Shatalov, V., Solberg, S., Spindler, G., Tarasova, O., Vana, M., Adani, M., Almodovar, P., Berton, E., Bessagnet, B., Bohlin-Nizzetto, P., Boruvkova, J., Breivik, K., Briganti, G., Cappelletti, A., Cuvelier, K., Derwent, R., D’Isidoro, M., Fagerli, H., Funk, C., Garcia Vivanco, M., Haeuber, R., Hueglin, C., Jenkins, S., Kerr, J., de Leeuw, F., Lynch, J., Manders, A., Mircea, M., Pay, M. T., Pritula, D., Querol, X., Raffort, V., Reiss, I., Roustan, Y., Sauvage, S., Scavo, K., Simpson, D., Smith, R. I., Tang, Y. S., Theobald, M., Torseth, K., Tsyro, S., van Pul, A., Vidic, S., Wallasch, M., and Wind, P.: Air pollution trends in the EMEP region between 1990 and 2012, 2016.
- 610
- Dayan, C., Fredj, E., Misztal, P. K., Gabay, M., Guenther, A. B., and Tas, E.: Emission of biogenic volatile organic compounds from warm and oligotrophic seawater in the Eastern Mediterranean, *Atmospheric Chem. Phys.*, **20**, 12741–12759, <https://doi.org/10.5194/acp-20-12741-2020>, 2020.
- 615
- Diémoz, H., Barnaba, F., Magri, T., Pession, G., Dionisi, D., Pittavino, S., Tombolato, I. K. F., Campanelli, M., Della Ceca, L. S., Hervo, M., Di Liberto, L., Ferrero, L., and Gobbi, G. P.: Transport of Po Valley aerosol pollution to the northwestern Alps – Part 1: Phenomenology, *Atmospheric Chem. Phys.*, **19**, 3065–3095, <https://doi.org/10.5194/acp-19-3065-2019>, 2019.

- 620 Dockery, D. W.: Health Effects of Particulate Air Pollution, *Ann. Epidemiol.*, 19, 257–263, <https://doi.org/10.1016/j.annepidem.2009.01.018>, 2009.
- van Donkelaar, A., Hammer, M. S., Bindle, L., Brauer, M., Brook, J. R., Garay, M. J., Hsu, N. C., Kalashnikova, O. V., Kahn, R. A., Lee, C., Levy, R. C., Lyapustin, A., Sayer, A. M., and Martin, R. V.: Monthly Global Estimates of Fine Particulate Matter and Their Uncertainty, *Environ. Sci. Technol.*, 55, 15287–15300, <https://doi.org/10.1021/acs.est.1c05309>, 2021.
- 625 Eck, T. F., Holben, B. N., Reid, J. S., Sinyuk, A., Giles, D. M., Arola, A., Slutsker, I., Schafer, J. S., Sorokin, M. G., Smirnov, A., LaRosa, A. D., Kraft, J., Reid, E. A., O'Neill, N. T., Welton, E. J., and Menendez, A. R.: The extreme forest fires in California/Oregon in 2020: Aerosol optical and physical properties and comparisons of aged versus fresh smoke, *Atmos. Environ.*, 305, 119798, <https://doi.org/10.1016/j.atmosenv.2023.119798>, 2023.
- 630 European Commission, Joint Research Centre, San-Miguel-Ayanz, J., Durrant, T., Boca, R., Maianti, P., Libertá, G., Artés-Vivancos, T., Oom, D., Branco, A., Rigo, D., Ferrari, D., Pfeiffer, H., Grecchi, R., and Nuijten, D.: Advance report on wildfires in Europe, Middle East and North Africa 2021, Publications Office of the European Union, <https://doi.org/10.2760/039729>, 2022.
- 635 Faisal, A.-A., Rahman, M. M., and Haque, S.: Retrieving spatial variation of aerosol level over urban mixed land surfaces using Landsat imageries: Degree of air pollution in Dhaka Metropolitan Area, *Phys. Chem. Earth Parts ABC*, 126, 103074, <https://doi.org/10.1016/j.pce.2021.103074>, 2022.
- Falah, S., Mhawish, A., Sorek-Hamer, M., Lyapustin, A. I., Kloog, I., Banerjee, T., Kizel, F., and Broday, D. M.: Impact of environmental attributes on the uncertainty in MAIAC/MODIS AOD retrievals: A comparative analysis, *Atmos. Environ.*, 262, 118659, <https://doi.org/10.1016/j.atmosenv.2021.118659>, 2021.
- 640 Filonchyk, M., Hurynovich, V., Yan, H., Zhou, L., and Gusev, A.: Climatology of aerosol optical depth over Eastern Europe based on 19 years (2000–2018) MODIS TERRA data, *Int. J. Climatol.*, 40, 3531–3549, <https://doi.org/10.1002/joc.6412>, 2020a.
- Filonchyk, M., Hurynovich, V., and Yan, H.: Trends in aerosol optical properties over Eastern Europe based on MODIS-Aqua, *Geosci. Front.*, 11, 2169–2181, <https://doi.org/10.1016/j.gsf.2020.03.014>, 2020b.
- 645 Fiore, A. M., Naik, V., Spracklen, D. V., Steiner, A., Unger, N., Prather, M., Bergmann, D., Cameron-Smith, P. J., Cionni, I., Collins, W. J., Dalsøren, S., Eyring, V., Folberth, G. A., Ginoux, P., Horowitz, L. W., Josse, B., Lamarque, J.-F., MacKenzie, I. A., Nagashima, T., O'Connor, F. M., Righi, M., Rumbold, S. T., Shindell, D. T., Skeie, R. B., Sudo, K., Szopa, S., Takemura, T., and Zeng, G.: Global air quality and climate, *Chem. Soc. Rev.*, 41, 6663–6683, <https://doi.org/10.1039/C2CS35095E>, 2012.
- 650 Floutsi, A. A., Korrás-Carraca, M. B., Matsoukas, C., Hatzianastassiou, N., and Biskos, G.: Climatology and trends of aerosol optical depth over the Mediterranean basin during the last 12 years (2002–2014) based on Collection 006 MODIS-Aqua data, *Sci. Total Environ.*, 551–552, 292–303, <https://doi.org/10.1016/j.scitotenv.2016.01.192>, 2016.
- 655 Font, A., Bourin, A., Gouillou, C., Debevec, C., Bonnaire, N., Sauvage, S., Brito, J. F. de, and Riffault, V.: Aerosol composition at EMEP remote sites in France : mass balance and de-weathered trends of PM_{2.5} and its main components, Copernicus Meetings, <https://doi.org/10.5194/egusphere-egu23-940>, 2023.
- 660 Foret, G., Michoud, V., Kotthaus, S., Petit, J.-E., Baudic, A., Siour, G., Kim, Y., Doussin, J.-F., Dupont, J.-C., Formenti, P., Gaimoz, C., Gherzi, V., Gratien, A., Gros, V., Jaffrezo, J.-L., Haefelin, M., Kreitz, M., Ravetta, F., Sartelet, K., Simon, L., Té, Y., Uzu, G., Zhang, S., Favez, O., and Beekmann, M.: The December 2016 extreme weather and particulate matter pollution episode in the Paris region (France), *Atmos. Environ.*, 291, 119386, <https://doi.org/10.1016/j.atmosenv.2022.119386>, 2022.

- Formenti, P., Mbemba Kabuiku, L., Chiapello, I., Ducos, F., Dulac, F., and Tanré, D.: Aerosol optical properties derived from POLDER-3/PARASOL (2005–2013) over the western Mediterranean Sea – Part 1: Quality assessment with AERONET and in situ airborne observations, *Atmospheric Meas. Tech.*, 11, 6761–6784, <https://doi.org/10.5194/amt-11-6761-2018>, 2018.
- 665 Gao, Y., Ma, M., Yan, F., Su, H., Wang, S., Liao, H., Zhao, B., Wang, X., Sun, Y., Hopkins, J. R., Chen, Q., Fu, P., Lewis, A. C., Qiu, Q., Yao, X., and Gao, H.: Impacts of biogenic emissions from urban landscapes on summer ozone and secondary organic aerosol formation in megacities, *Sci. Total Environ.*, 814, 152654, <https://doi.org/10.1016/j.scitotenv.2021.152654>, 2022.
- 670 Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korokin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database – automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements, *Atmospheric Meas. Tech.*, 12, 169–209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- 675 Gkikas, A., Hatzianastassiou, N., Mihalopoulos, N., and Torres, O.: Characterization of aerosol episodes in the greater Mediterranean Sea area from satellite observations (2000–2007), *Atmos. Environ.*, 128, 286–304, <https://doi.org/10.1016/j.atmosenv.2015.11.056>, 2016.
- 680 Gkikas, A., Proestakis, E., Amiridis, V., Kazadzis, S., Di Tomaso, E., Marinou, E., Hatzianastassiou, N., Kok, J. F., and García-Pando, C. P.: Quantification of the dust optical depth across spatiotemporal scales with the MIDAS global dataset (2003–2017), *Atmospheric Chem. Phys.*, 22, 3553–3578, <https://doi.org/10.5194/acp-22-3553-2022>, 2022.
- Gladson, L., Garcia, N., Bi, J., Liu, Y., Lee, H. J., and Cromar, K.: Evaluating the Utility of High-Resolution Spatiotemporal Air Pollution Data in Estimating Local PM_{2.5} Exposures in California from 2015–2018, *Atmosphere*, 13, 85, <https://doi.org/10.3390/atmos13010085>, 2022.
- 685 Grgurić, S., Križan, J., Gašparac, G., Antonić, O., Špirić, Z., Mamouri, R. E., Christodoulou, A., Nisantzi, A., Agapiou, A., Themistocleous, K., Fedra, K., Panayiotou, C., and Hadjimitsis, D.: Relationship between MODIS based Aerosol Optical Depth and PM₁₀ over Croatia, *Cent. Eur. J. Geosci.*, 6, 2–16, <https://doi.org/10.2478/s13533-012-0135-6>, 2014.
- Grigorieva, E. and Lukyanets, A.: Combined Effect of Hot Weather and Outdoor Air Pollution on Respiratory Health: Literature Review, *Atmosphere*, 12, 790, <https://doi.org/10.3390/atmos12060790>, 2021.
- 690 Groot Zwaaftink, C. D., Aas, W., Eckhardt, S., Evangeliou, N., Hamer, P., Johnsrud, M., Kylling, A., Platt, S. M., Stebel, K., Uggerud, H., and Yttri, K. E.: What caused a record high PM₁₀ episode in northern Europe in October 2020?, *Atmospheric Chem. Phys.*, 22, 3789–3810, <https://doi.org/10.5194/acp-22-3789-2022>, 2022.
- 695 Guo, J.-P., Zhang, X.-Y., Che, H.-Z., Gong, S.-L., An, X., Cao, C.-X., Guang, J., Zhang, H., Wang, Y.-Q., Zhang, X.-C., Xue, M., and Li, X.-W.: Correlation between PM concentrations and aerosol optical depth in eastern China, *Atmos. Environ.*, 43, 5876–5886, <https://doi.org/10.1016/j.atmosenv.2009.08.026>, 2009.
- Gupta, G., Venkat Ratnam, M., Madhavan, B. L., and Narayanamurthy, C. S.: Long-term trends in Aerosol Optical Depth obtained across the globe using multi-satellite measurements, *Atmos. Environ.*, 273, 118953, <https://doi.org/10.1016/j.atmosenv.2022.118953>, 2022.
- 700 Gupta, G., Venkat Ratnam, M., Madhavan, B. L., and Jayaraman, A.: Global trends in the aerosol optical, physical, and morphological properties obtained using multi-sensor measurements, *Atmos. Environ.*, 295, 119569, <https://doi.org/10.1016/j.atmosenv.2022.119569>, 2023.

- Gupta, P., Levy, R. C., Mattoo, S., Remer, L. A., and Munchak, L. A.: A surface reflectance scheme for retrieving aerosol optical depth over urban surfaces in MODIS Dark Target retrieval algorithm, *Atmospheric Meas. Tech.*, 9, 3293–3308, <https://doi.org/10.5194/amt-9-3293-2016>, 2016.
- 705 Ha, S., Liu, Z., Sun, W., Lee, Y., and Chang, L.: Improving air quality forecasting with the assimilation of GOCI aerosol optical depth (AOD) retrievals during the KORUS-AQ period, *Atmospheric Chem. Phys.*, 20, 6015–6036, <https://doi.org/10.5194/acp-20-6015-2020>, 2020.
- Hammer, M. S., van Donkelaar, A., Li, C., Lyapustin, A., Sayer, A. M., Hsu, N. C., Levy, R. C., Garay, M. J., Kalashnikova, O. V., Kahn, R. A., Brauer, M., Apte, J. S., Henze, D. K., Zhang, L., Zhang, Q., Ford, B., Pierce, J. R., and Martin, R. V.: Global Estimates and Long-Term Trends of Fine Particulate Matter Concentrations (1998–2018), *Environ. Sci. Technol.*, 54, 7879–7890, <https://doi.org/10.1021/acs.est.0c01764>, 2020.
- 710 He, Q., Zhang, M., and Huang, B.: Spatio-temporal variation and impact factors analysis of satellite-based aerosol optical depth over China from 2002 to 2015, *Atmos. Environ.*, 129, 79–90, <https://doi.org/10.1016/j.atmosenv.2016.01.002>, 2016.
- 715 He, Q., Wang, M., and Yim, S. H. L.: The spatiotemporal relationship between PM_{2.5} and aerosol optical depth in China: influencing factors and implications for satellite PM_{2.5} estimations using MAIAC aerosol optical depth, *Atmospheric Chem. Phys.*, 21, 18375–18391, <https://doi.org/10.5194/acp-21-18375-2021>, 2021.
- Hough, I., Sarafian, R., Shtein, A., Zhou, B., Lepeule, J., and Kloog, I.: Gaussian Markov random fields improve ensemble predictions of daily 1 km PM_{2.5} and PM₁₀ across France, *Atmos. Environ.*, 264, 118693, <https://doi.org/10.1016/j.atmosenv.2021.118693>, 2021.
- 720 Hsu, N. C., Tsay, S.-C., King, M. D., and Herman, J. R.: Aerosol properties over bright-reflecting source regions, *IEEE Trans. Geosci. Remote Sens.*, 42, 557–569, <https://doi.org/10.1109/TGRS.2004.824067>, 2004.
- Hsu, N. C., Lee, J., Sayer, A. M., Carletta, N., Chen, S.-H., Tucker, C. J., Holben, B. N., and Tsay, S.-C.: Retrieving near-global aerosol loading over land and ocean from AVHRR, *J. Geophys. Res. Atmospheres*, 122, 9968–9989, <https://doi.org/10.1002/2017JD026932>, 2017.
- 725 Hussain, M. M. and Mahmud, I.: pyMannKendall: a python package for non parametric Mann Kendall family of trend tests., *J. Open Source Softw.*, 4, 1556, <https://doi.org/10.21105/joss.01556>, 2019.
- Israelevich, P., Ganor, E., Alpert, P., Kishcha, P., and Stupp, A.: Predominant transport paths of Saharan dust over the Mediterranean Sea to Europe, *J. Geophys. Res. Atmospheres*, 117, <https://doi.org/10.1029/2011JD016482>, 2012.
- 730 Jaén, C., Udina, M., and Bech, J.: Analysis of two heat wave driven ozone episodes in Barcelona and surrounding region: Meteorological and photochemical modeling, *Atmos. Environ.*, 246, 118037, <https://doi.org/10.1016/j.atmosenv.2020.118037>, 2021.
- Jung, C.-R., Chen, W.-T., and Nakayama, S. F.: A National-Scale 1-km Resolution PM_{2.5} Estimation Model over Japan Using MAIAC AOD and a Two-Stage Random Forest Model, *Remote Sens.*, 13, 3657, <https://doi.org/10.3390/rs13183657>, 2021.
- 735 Just, A. C., Wright, R. O., Schwartz, J., Coull, B. A., Baccarelli, A. A., Tellez-Rojo, M. M., Moody, E., Wang, Y., Lyapustin, A., and Kloog, I.: Using high-resolution satellite aerosol optical depth to estimate daily PM_{2.5} geographical distribution in Mexico City, *Environ. Sci. Technol.*, 49, 8576–8584, <https://doi.org/10.1021/acs.est.5b00859>, 2015.
- 740 Karl, M., Tsigaridis, K., Vignati, E., and Dentener, F.: Formation of secondary organic aerosol from isoprene oxidation over Europe, *Atmospheric Chem. Phys.*, 9, 7003–7030, <https://doi.org/10.5194/acp-9-7003-2009>, 2009.

- 745 Kessner, A. L., Wang, J., Levy, R. C., and Colarco, P. R.: Remote sensing of surface visibility from space: A look at the United States East Coast, *Atmos. Environ.*, 81, 136–147, <https://doi.org/10.1016/j.atmosenv.2013.08.050>, 2013.
- Kinne, S., O'Donnel, D., Stier, P., Kloster, S., Zhang, K., Schmidt, H., Rast, S., Giorgetta, M., Eck, T. F., and Stevens, B.: MAC-v1: A new global aerosol climatology for climate studies, *J. Adv. Model. Earth Syst.*, 5, 704–740, <https://doi.org/10.1002/jame.20035>, 2013.
- 750 Kondragunta, S., Lee, P., McQueen, J., Kittaka, C., Prados, A. I., Ciren, P., Laszlo, I., Pierce, R. B., Hoff, R., and Szykman, J. J.: Air Quality Forecast Verification Using Satellite Data, *J. Appl. Meteorol. Climatol.*, 47, 425–442, 2008.
- Lee, J., Hsu, N. C., Sayer, A. M., Bettenhausen, C., and Yang, P.: AERONET-Based Nonspherical Dust Optical Models and Effects on the VIIRS Deep Blue/SOAR Over Water Aerosol Product, *J. Geophys. Res. Atmospheres*, 755 122, 10,384–10,401, <https://doi.org/10.1002/2017JD027258>, 2017.
- Lee, S., Park, S., Lee, M.-I., Kim, G., Im, J., and Song, C.-K.: Air Quality Forecasts Improved by Combining Data Assimilation and Machine Learning With Satellite AOD, *Geophys. Res. Lett.*, 49, e2021GL096066, <https://doi.org/10.1029/2021GL096066>, 2022.
- 760 Li, M.-H., Fan, L.-C., Mao, B., Yang, J.-W., Choi, A. M. K., Cao, W.-J., and Xu, J.-F.: Short-term Exposure to Ambient Fine Particulate Matter Increases Hospitalizations and Mortality in COPD: A Systematic Review and Meta-analysis, *Chest*, 149, 447–458, <https://doi.org/10.1378/chest.15-0513>, 2016.
- Li, Z., Wen, Q., and Zhang, R.: Sources, health effects and control strategies of indoor fine particulate matter (PM_{2.5}): A review, *Sci. Total Environ.*, 586, 610–622, <https://doi.org/10.1016/j.scitotenv.2017.02.029>, 2017.
- 765 Liu, J., Mauzerall, D. L., and Horowitz, L. W.: Evaluating inter-continental transport of fine aerosols:(2) Global health impact, *Atmos. Environ.*, 43, 4339–4347, <https://doi.org/10.1016/j.atmosenv.2009.05.032>, 2009.
- Liu, Y., Liu, Y., Wang, M., Dong, X., Zheng, Y., Shrivastava, M., Qian, Y., Bai, H., Li, X., and Yang, X.-Q.: Anthropogenic–biogenic interaction amplifies warming from emission reduction over the southeastern US, *Environ. Res. Lett.*, 16, 124046, <https://doi.org/10.1088/1748-9326/ac3285>, 2021.
- 770 Luo, M., Ji, Y., Ren, Y., Gao, F., Zhang, H., Zhang, L., Yu, Y., and Li, H.: Characteristics and Health Risk Assessment of PM_{2.5}-Bound PAHs during Heavy Air Pollution Episodes in Winter in Urban Area of Beijing, China, *Atmosphere*, 12, 323, <https://doi.org/10.3390/atmos12030323>, 2021.
- Lyapustin, A. and Wang, Y.: MCD19A2 MODIS/Terra+Aqua Land Aerosol Optical Depth Daily L2G Global 1km SIN Grid V006, <https://doi.org/10.5067/MODIS/MCD19A2.006>, 2018.
- 775 Lyapustin, A., Wang, Y., Laszlo, I., and Korkin, S.: Improved cloud and snow screening in MAIAC aerosol retrievals using spectral and spatial analysis, *Atmospheric Meas. Tech.*, 5, 843–850, <https://doi.org/10.5194/amt-5-843-2012>, 2012.
- Lyapustin, A., Wang, Y., Korkin, S., and Huang, D.: MODIS Collection 6 MAIAC algorithm, *Atmospheric Meas. Tech.*, 11, 5741–5765, <https://doi.org/10.5194/amt-11-5741-2018>, 2018.
- 780 Ma, X. and Yu, F.: Seasonal and spatial variations of global aerosol optical depth: multi-year modelling with GEOS-Chem-APM and comparisons with multiple-platform observations, *Tellus B Chem. Phys. Meteorol.*, 67, 25115, <https://doi.org/10.3402/tellusb.v67.25115>, 2015.
- Ma, Y., Xin, J., Zhang, W., Gong, C., Wen, T., Wu, X., Wang, Y., Wang, L., Wu, F., and Ding, X.: Uncertainties of Simulated Aerosol Direct Radiative Effect Induced by Aerosol Chemical Components: A Measurement-Based

- 785 Perspective From Urban-Forest Transition Region in East China, *J. Geophys. Res. Atmospheres*, 126, e2020JD033688, <https://doi.org/10.1029/2020JD033688>, 2021.
- Majewski, G., Szeląg, B., Mach, T., Rogula-Kozłowska, W., Anioł, E., Białowicz, J., Dmochowska, A., and Białowicz, J. S.: Predicting the Number of Days With Visibility in a Specific Range in Warsaw (Poland) Based on Meteorological and Air Quality Data, *Front. Environ. Sci.*, 9, 2021.
- 790 Manousakas, M. I., Florou, K., and Pandis, S. N.: Source Apportionment of Fine Organic and Inorganic Atmospheric Aerosol in an Urban Background Area in Greece, *Atmosphere*, 11, 330, <https://doi.org/10.3390/atmos11040330>, 2020.
- Mao, K. B., Ma, Y., Xia, L., Chen, W. Y., Shen, X. Y., He, T. J., and Xu, T. R.: Global aerosol change in the last decade: An analysis based on MODIS data, *Atmos. Environ.*, 94, 680–686, <https://doi.org/10.1016/j.atmosenv.2014.04.053>, 2014.
- 795 Martin, S. T., Artaxo, P., Machado, L. a. T., Manzi, A. O., Souza, R. a. F., Schumacher, C., Wang, J., Andreae, M. O., Barbosa, H. M. J., Fan, J., Fisch, G., Goldstein, A. H., Guenther, A., Jimenez, J. L., Pöschl, U., Silva Dias, M. A., Smith, J. N., and Wendisch, M.: Introduction: Observations and Modeling of the Green Ocean Amazon (GoAmazon2014/5), *Atmospheric Chem. Phys.*, 16, 4785–4797, <https://doi.org/10.5194/acp-16-4785-2016>, 2016.
- 800 Martins, V. S., Lyapustin, A., de Carvalho, L. a. S., Barbosa, C. C. F., and Novo, E. M. L. M.: Validation of high-resolution MAIAC aerosol product over South America, *J. Geophys. Res. Atmospheres*, 122, 7537–7559, <https://doi.org/10.1002/2016JD026301>, 2017.
- 805 Martins, V. S., Lyapustin, A., Wang, Y., Giles, D. M., Smirnov, A., Slutsker, I., and Korokin, S.: Global validation of columnar water vapor derived from EOS MODIS-MAIAC algorithm against the ground-based AERONET observations, *Atmospheric Res.*, 225, 181–192, <https://doi.org/10.1016/j.atmosres.2019.04.005>, 2019.
- 810 Masson-Delmotte, Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, Ö., Yu, R., and Zhou, B. (Eds.): *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896>, 2021.
- Mehta, M., Singh, R., Singh, A., Singh, N., and Anshumali: Recent global aerosol optical depth variations and trends — A comparative study using MODIS and MISR level 3 datasets, *Remote Sens. Environ.*, 181, 137–150, <https://doi.org/10.1016/j.rse.2016.04.004>, 2016.
- 815 Merdji, A. B., Lu, C., Xu, X., and Mhawish, A.: Long-term three-dimensional distribution and transport of Saharan dust: Observation from CALIPSO, MODIS, and reanalysis data, *Atmospheric Res.*, 286, 106658, <https://doi.org/10.1016/j.atmosres.2023.106658>, 2023.
- Met Office: *Cartopy: a cartographic python library with a Matplotlib interface*, Exeter, Devon, 2010.
- 820 Mhawish, A., Banerjee, T., Sorek-Hamer, M., Lyapustin, A., Broday, D. M., and Chatfield, R.: Comparison and evaluation of MODIS Multi-angle Implementation of Atmospheric Correction (MAIAC) aerosol product over South Asia, *Remote Sens. Environ.*, 224, 12–28, <https://doi.org/10.1016/j.rse.2019.01.033>, 2019.
- 825 Michoud, V., Sciare, J., Sauvage, S., Dusanter, S., Léonardis, T., Gros, V., Kalogridis, C., Zannoni, N., Féron, A., Petit, J.-E., Crenn, V., Baisnée, D., Sarda-Estève, R., Bonnaire, N., Marchand, N., DeWitt, H. L., Pey, J., Colomb, A., Gheusi, F., Szidat, S., Stavroulas, I., Borbon, A., and Locoge, N.: Organic carbon at a remote site of the western Mediterranean Basin: sources and chemistry during the ChArMEx SOP2 field experiment, *Atmospheric Chem. Phys.*, 17, 8837–8865, <https://doi.org/10.5194/acp-17-8837-2017>, 2017.

- Moulin, C., Lambert, C. E., Dayan, U., Masson, V., Ramonet, M., Bousquet, P., Legrand, M., Balkanski, Y. J., Guelle, W., Marticorena, B., Bergametti, G., and Dulac, F.: Satellite climatology of African dust transport in the Mediterranean atmosphere, *J. Geophys. Res. Atmospheres*, 103, 13137–13144, <https://doi.org/10.1029/98JD00171>, 1998.
- 830 Nascimento, J. P., Bela, M. M., Meller, B. B., Banducci, A. L., Rizzo, L. V., Vara-Vela, A. L., Barbosa, H. M. J., Gomes, H., Rafee, S. A. A., Franco, M. A., Carbone, S., Cirino, G. G., Souza, R. A. F., McKeen, S. A., and Artaxo, P.: Aerosols from anthropogenic and biogenic sources and their interactions – modeling aerosol formation, optical properties, and impacts over the central Amazon basin, *Atmospheric Chem. Phys.*, 21, 6755–6779, <https://doi.org/10.5194/acp-21-6755-2021>, 2021.
- 835 Pace, G., di Sarra, A., Meloni, D., Piacentino, S., and Chamard, P.: Aerosol optical properties at Lampedusa (Central Mediterranean). 1. Influence of transport and identification of different aerosol types, *Atmospheric Chem. Phys.*, 6, 697–713, <https://doi.org/10.5194/acp-6-697-2006>, 2006.
- Papachristopoulou, K., Raptis, I.-P., Gkikas, A., Fountoulakis, I., Masoom, A., and Kazadzis, S.: Aerosol optical depth regime over megacities of the world, *Atmospheric Chem. Phys.*, 22, 15703–15727, <https://doi.org/10.5194/acp-22-15703-2022>, 2022.
- 840 Pedde, M., Kloog, I., Szpiro, A., Dorman, M., Larson, T. V., and Adar, S. D.: Estimating long-term PM_{10-2.5} concentrations in six US cities using satellite-based aerosol optical depth data, *Atmos. Environ.*, 272, 118945, <https://doi.org/10.1016/j.atmosenv.2022.118945>, 2022.
- Pérez, C., Sicard, M., Jorba, O., Comerón, A., and Baldasano, J. M.: Summertime re-circulations of air pollutants over the north-eastern Iberian coast observed from systematic EARLINET lidar measurements in Barcelona, *Atmos. Environ.*, 38, 3983–4000, <https://doi.org/10.1016/j.atmosenv.2004.04.010>, 2004.
- 845 Petit, J.-E., Favez, O., Sciare, J., Crenn, V., Sarda-Estève, R., Bonnaire, N., Močnik, G., Dupont, J.-C., Haeffelin, M., and Leoz-Garziandia, E.: Two years of near real-time chemical composition of submicron aerosols in the region of Paris using an Aerosol Chemical Speciation Monitor (ACSM) and a multi-wavelength Aethalometer, *Atmospheric Chem. Phys.*, 15, 2985–3005, <https://doi.org/10.5194/acp-15-2985-2015>, 2015.
- 850 Pietrogrande, M. C., Bacco, D., Ferrari, S., Kaipainen, J., Ricciardelli, I., Riekkola, M.-L., Trentini, A., and Visentin, M.: Characterization of atmospheric aerosols in the Po valley during the supersito campaigns — Part 3: Contribution of wood combustion to wintertime atmospheric aerosols in Emilia Romagna region (Northern Italy), *Atmos. Environ.*, 122, 291–305, <https://doi.org/10.1016/j.atmosenv.2015.09.059>, 2015.
- 855 Pirhadi, M., Mousavi, A., Taghvaei, S., Shafer, M. M., and Sioutas, C.: Semi-volatile components of PM_{2.5} in an urban environment: volatility profiles and associated oxidative potential, *Atmospheric Environ. Oxf. Engl.* 1994, 223, 117197, <https://doi.org/10.1016/j.atmosenv.2019.117197>, 2020.
- Provençal, S., Kishcha, P., da Silva, A. M., Elhacham, E., and Alpert, P.: AOD distributions and trends of major aerosol species over a selection of the world's most populated cities based on the 1st version of NASA's MERRA Aerosol Reanalysis, *Urban Clim.*, 20, 168–191, <https://doi.org/10.1016/j.uclim.2017.04.001>, 2017.
- 860 Putaud, J. P., Cavalli, F., Martins dos Santos, S., and Dell'Acqua, A.: Long-term trends in aerosol optical characteristics in the Po Valley, Italy, *Atmospheric Chem. Phys.*, 14, 9129–9136, <https://doi.org/10.5194/acp-14-9129-2014>, 2014.
- Qin, W., Fang, H., Wang, L., Wei, J., Zhang, M., Su, X., Bilal, M., and Liang, X.: MODIS high-resolution MAIAC aerosol product: Global validation and analysis, *Atmos. Environ.*, 264, 118684, <https://doi.org/10.1016/j.atmosenv.2021.118684>, 2021.

- Raptis, I.-P., Kazadzis, S., Amiridis, V., Gkikas, A., Gerasopoulos, E., and Mihalopoulos, N.: A Decade of Aerosol Optical Properties Measurements over Athens, Greece, *Atmosphere*, **11**, 154, <https://doi.org/10.3390/atmos11020154>, 2020.
- 870 Remer, L. A., Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R.-R., Ichoku, C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS Aerosol Algorithm, Products, and Validation, *J. Atmospheric Sci.*, **62**, 947–973, <https://doi.org/10.1175/JAS3385.1>, 2005.
- Remer, L. A., Levy, R. C., Mattoo, S., Tanré, D., Gupta, P., Shi, Y., Sawyer, V., Munchak, L. A., Zhou, Y., Kim, M., Ichoku, C., Patadia, F., Li, R.-R., Gassó, S., Kleidman, R. G., and Holben, B. N.: The Dark Target Algorithm for
875 Observing the Global Aerosol System: Past, Present, and Future, *Remote Sens.*, **12**, 2900, <https://doi.org/10.3390/rs12182900>, 2020.
- Robotto, A., Barbero, S., Bracco, P., Cremonini, R., Ravina, M., and Brizio, E.: Improving Air Quality Standards in Europe: Comparative Analysis of Regional Differences, with a Focus on Northern Italy, *Atmosphere*, **13**, 642, <https://doi.org/10.3390/atmos13050642>, 2022.
- 880 Rogozovsky, I., Ohneiser, K., Lyapustin, A., Ansmann, A., and Chudnovsky, A.: The impact of different aerosol layering conditions on the high-resolution MODIS/MAIAC AOD retrieval bias: The uncertainty analysis, *Atmos. Environ.*, **309**, 119930, <https://doi.org/10.1016/j.atmosenv.2023.119930>, 2023.
- Sandrini, S., van Pinxteren, D., Giulianelli, L., Herrmann, H., Poulain, L., Facchini, M. C., Gilardoni, S., Rinaldi, M., Paglione, M., Turpin, B. J., Pollini, F., Bucci, S., Zanca, N., and Decesari, S.: Size-resolved aerosol composition at
885 an urban and a rural site in the Po Valley in summertime: implications for secondary aerosol formation, *Atmospheric Chem. Phys.*, **16**, 10879–10897, <https://doi.org/10.5194/acp-16-10879-2016>, 2016.
- Sayer, A. M., Hsu, N. C., Lee, J., Bettenhausen, C., Kim, W. V., and Smirnov, A.: Satellite Ocean Aerosol Retrieval (SOAR) Algorithm Extension to S-NPP VIIRS as Part of the “Deep Blue” Aerosol Project, *J. Geophys. Res. Atmospheres*, **123**, 380–400, <https://doi.org/10.1002/2017JD027412>, 2018.
- 890 Schaap, M., Apituley, A., Timmermans, R. M. A., Koelemeijer, R. B. A., and de Leeuw, G.: Exploring the relation between aerosol optical depth and PM_{2.5} at Cabauw, the Netherlands, *Atmospheric Chem. Phys.*, **9**, 909–925, <https://doi.org/10.5194/acp-9-909-2009>, 2009.
- Schäfer, K., Harbusch, A., Emeis, S., Koepke, P., and Wiegner, M.: Correlation of aerosol mass near the ground with aerosol optical depth during two seasons in Munich, *Atmos. Environ.*, **42**, 4036–4046,
895 <https://doi.org/10.1016/j.atmosenv.2008.01.060>, 2008.
- Schuster, G. L., Dubovik, O., and Holben, B. N.: Angstrom exponent and bimodal aerosol size distributions, *J. Geophys. Res. Atmospheres*, **111**, <https://doi.org/10.1029/2005JD006328>, 2006.
- Schutgens, N. A. J.: Site representativity of AERONET and GAW remotely sensed aerosol optical thickness and absorbing aerosol optical thickness observations, *Atmospheric Chem. Phys.*, **20**, 7473–7488,
900 <https://doi.org/10.5194/acp-20-7473-2020>, 2020.
- Scotto, F., Bacco, D., Lasagni, S., Trentini, A., Poluzzi, V., and Vecchi, R.: A multi-year source apportionment of PM_{2.5} at multiple sites in the southern Po Valley (Italy), *Atmospheric Pollut. Res.*, **12**, 101192, <https://doi.org/10.1016/j.apr.2021.101192>, 2021.
- Segura, S., Estellés, V., Utrillas, M. P., and Martínez-Lozano, J. A.: Long term analysis of the columnar and
905 surface aerosol relationship at an urban European coastal site, *Atmos. Environ.*, **167**, 309–322, <https://doi.org/10.1016/j.atmosenv.2017.08.012>, 2017.

- Shi, Y., Levy, R., Yang, L., Remer, L., Mattoo, S., and Dubovik, O.: A Dark Target research aerosol algorithm for MODIS observations over eastern China: increasing coverage while maintaining accuracy at high aerosol loading, *Atmospheric Meas. Tech.*, 14, <https://doi.org/10.5194/amt-14-3449-2021>, 2021.
- 910 Shrivastava, M., Andreae, M. O., Artaxo, P., Barbosa, H. M. J., Berg, L. K., Brito, J., Ching, J., Easter, R. C., Fan, J., Fast, J. D., Feng, Z., Fuentes, J. D., Glasius, M., Goldstein, A. H., Alves, E. G., Gomes, H., Gu, D., Guenther, A., Jathar, S. H., Kim, S., Liu, Y., Lou, S., Martin, S. T., McNeill, V. F., Medeiros, A., de Sá, S. S., Shilling, J. E., Springston, S. R., Souza, R. a. F., Thornton, J. A., Isaacman-VanWertz, G., Yee, L. D., Ynoue, R., Zaveri, R. A., Zelenyuk, A., and Zhao, C.: Urban pollution greatly enhances formation of natural aerosols over the Amazon rainforest, *Nat. Commun.*, 10, 1046, <https://doi.org/10.1038/s41467-019-08909-4>, 2019.
- 915 Sicard, P., Agathokleous, E., De Marco, A., Paoletti, E., and Calatayud, V.: Urban population exposure to air pollution in Europe over the last decades, *Environ. Sci. Eur.*, 33, 28, <https://doi.org/10.1186/s12302-020-00450-2>, 2021.
- Singh, A., Avis, W. R., and Pope, F. D.: Visibility as a proxy for air quality in East Africa, *Environ. Res. Lett.*, 15, 084002, <https://doi.org/10.1088/1748-9326/ab8b12>, 2020.
- 920 Sinyuk, A., Holben, B. N., Eck, T. F., Giles, D. M., Slutsker, I., Korkin, S., Schafer, J. S., Smirnov, A., Sorokin, M., and Lyapustin, A.: The AERONET Version 3 aerosol retrieval algorithm, associated uncertainties and comparisons to Version 2, *Atmospheric Meas. Tech.*, 13, 3375–3411, <https://doi.org/10.5194/amt-13-3375-2020>, 2020.
- Skyllakou, K., Murphy, B. N., Megaritis, A. G., Fountoukis, C., and Pandis, S. N.: Contributions of local and regional sources to fine PM in the megacity of Paris, *Atmospheric Chem. Phys.*, 14, 2343–2352, <https://doi.org/10.5194/acp-14-2343-2014>, 2014.
- 925 Smirnov, A., Holben, B. N., Eck, T. F., Slutsker, I., Chatenet, B., and Pinker, R. T.: Diurnal variability of aerosol optical depth observed at AERONET (Aerosol Robotic Network) sites, *Geophys. Res. Lett.*, 29, 30-1-30–4, <https://doi.org/10.1029/2002GL016305>, 2002.
- 930 Song, C. K., Ho, C. H., Park, R. J., Choi, Y. S., Kim, J., Gong, D. Y., and Lee, Y. B.: Spatial and seasonal variations of surface PM10 concentration and MODIS aerosol optical depth over China, *Asia-Pac. J. Atmospheric Sci.*, 45, 33–43, 2009.
- Spencer, R. S., Levy, R. C., Remer, L. A., Mattoo, S., Arnold, G. T., Hlavka, D. L., Meyer, K. G., Marshak, A., Wilcox, E. M., and Platnick, S. E.: Exploring Aerosols Near Clouds With High-Spatial-Resolution Aircraft Remote Sensing During SEAC4RS, *J. Geophys. Res. Atmospheres*, 124, 2148–2173, <https://doi.org/10.1029/2018JD028989>, 2019.
- 935 Stafoggia, M., Schwartz, J., Badaloni, C., Bellander, T., Alessandrini, E., Cattani, G., de' Donato, F., Gaeta, A., Leone, G., Lyapustin, A., Sorek-Hamer, M., de Hoogh, K., Di, Q., Forastiere, F., and Kloog, I.: Estimation of daily PM10 concentrations in Italy (2006–2012) using finely resolved satellite data, land use variables and meteorology, *Environ. Int.*, 99, 234–244, <https://doi.org/10.1016/j.envint.2016.11.024>, 2017.
- 940 Su, X., Cao, M., Wang, L., Gui, X., Zhang, M., Huang, Y., and Zhao, Y.: Validation, inter-comparison, and usage recommendation of six latest VIIRS and MODIS aerosol products over the ocean and land on the global and regional scales, *Sci. Total Environ.*, 884, 163794, <https://doi.org/10.1016/j.scitotenv.2023.163794>, 2023.
- Sun, E., Xu, X., Che, H., Tang, Z., Gui, K., An, L., Lu, C., and Shi, G.: Variation in MERRA-2 aerosol optical depth and absorption aerosol optical depth over China from 1980 to 2017, *J. Atmospheric Sol.-Terr. Phys.*, 186, 8–19, <https://doi.org/10.1016/j.jastp.2019.01.019>, 2019.
- 945 Taghvaei, S., Sowlat, M. H., Diapouli, E., Manousakas, M. I., Vasilatou, V., Eleftheriadis, K., and Sioutas, C.: Source apportionment of the oxidative potential of fine ambient particulate matter (PM2.5) in Athens, Greece, *Sci. Total Environ.*, 653, 1407–1416, <https://doi.org/10.1016/j.scitotenv.2018.11.016>, 2019.

- Tao, M., Wang, J., Li, R., Wang, L., Wang, L., Wang, Z., Tao, J., Che, H., and Chen, L.: Performance of MODIS high-resolution MAIAC aerosol algorithm in China: Characterization and limitation, *Atmos. Environ.*, 213, 159–169, <https://doi.org/10.1016/j.atmosenv.2019.06.004>, 2019.
- Toledano, C., Cachorro, V. E., Berjon, A., de Frutos, A. M., Sorribas, M., de la Morena, B. A., and Goloub, P.: Aerosol optical depth and Ångström exponent climatology at El Arenosillo AERONET site (Huelva, Spain), *Q. J. R. Meteorol. Soc.*, 133, 795–807, <https://doi.org/10.1002/qj.54>, 2007.
- 955 Tsiflikiotou, M. A., Kostenidou, E., Papanastasiou, D. K., Patoulas, D., Zampas, P., Paraskevopoulou, D., Diapouli, E., Kaltsonoudis, C., Florou, K., Bougiatioti, A., Stavroulas, I., Theodosi, C., Kouvarakis, G., Vasilatou, V., Siakavaras, D., Biskos, G., Pilinis, C., Eleftheriadis, K., Gerasopoulos, E., Mihalopoulos, N., and Pandis, S. N.: Summertime particulate matter and its composition in Greece, *Atmos. Environ.*, 213, 597–607, <https://doi.org/10.1016/j.atmosenv.2019.06.013>, 2019.
- 960 Tsyro, S., Aas, W., Colette, A., Andersson, C., Bessagnet, B., Ciarelli, G., Couvidat, F., Cuvelier, K., Manders, A., Mar, K., Mircea, M., Otero, N., Pay, M.-T., Raffort, V., Roustan, Y., Theobald, M. R., Vivanco, M. G., Fagerli, H., Wind, P., Briganti, G., Cappelletti, A., D’Isidoro, M., and Adani, M.: Eurodelta multi-model simulated and observed particulate matter trends in Europe in the period of 1990–2010, *Atmospheric Chem. Phys.*, 22, 7207–7257, <https://doi.org/10.5194/acp-22-7207-2022>, 2022.
- 965 Tuccella, P., Curci, G., Pitari, G., Lee, S., and Jo, D. S.: Direct Radiative Effect of Absorbing Aerosols: Sensitivity to Mixing State, Brown Carbon, and Soil Dust Refractive Index and Shape, *J. Geophys. Res. Atmospheres*, 125, e2019JD030967, <https://doi.org/10.1029/2019JD030967>, 2020.
- Tuet, W. Y., Chen, Y., Fok, S., Champion, J. A., and Ng, N. L.: Inflammatory responses to secondary organic aerosols (SOA) generated from biogenic and anthropogenic precursors, *Atmospheric Chem. Phys.*, 17, 11423–11440, <https://doi.org/10.5194/acp-17-11423-2017>, 2017.
- 970 Vautard, R., Menut, L., Beekmann, M., Chazette, P., Flamant, P. H., Gombert, D., Guédalia, D., Kley, D., Lefebvre, M.-P., Martin, D., Mégie, G., Perros, P., and Toupance, G.: A synthesis of the Air Pollution Over the Paris Region (ESQUIF) field campaign, *J. Geophys. Res. Atmospheres*, 108, <https://doi.org/10.1029/2003JD003380>, 2003.
- Vecchi, R., Bernardoni, V., Fermo, P., Lucarelli, F., Mazzei, F., Nava, S., Prati, P., Piazzalunga, A., and Valli, G.: 4-hours resolution data to study PM₁₀ in a “hot spot” area in Europe, *Environ. Monit. Assess.*, 154, 283–300, <https://doi.org/10.1007/s10661-008-0396-1>, 2009.
- Vecchi, R., Bernardoni, V., Valentini, S., Piazzalunga, A., Fermo, P., and Valli, G.: Assessment of light extinction at a European polluted urban area during wintertime: Impact of PM₁ composition and sources, *Environ. Pollut.*, 233, 679–689, <https://doi.org/10.1016/j.envpol.2017.10.059>, 2018.
- 980 Viana, M., Pey, J., Querol, X., Alastuey, A., de Leeuw, F., and Lükewille, A.: Natural sources of atmospheric aerosols influencing air quality across Europe, *Sci. Total Environ.*, 472, 825–833, <https://doi.org/10.1016/j.scitotenv.2013.11.140>, 2014.
- Wang, Y., Lyapustin, A. I., Privette, J. L., Cook, R. B., SanthanaVannan, S. K., Vermote, E. F., and Schaaf, C. L.: Assessment of biases in MODIS surface reflectance due to Lambertian approximation, *Remote Sens. Environ.*, 985, 114, 2791–2801, <https://doi.org/10.1016/j.rse.2010.06.013>, 2010.
- Wei, J., Li, Z., Peng, Y., and Sun, L.: MODIS Collection 6.1 aerosol optical depth products over land and ocean: validation and comparison, *Atmos. Environ.*, 201, 428–440, <https://doi.org/10.1016/j.atmosenv.2018.12.004>, 2019.
- Xu, L., Du, L., Tsona, N. T., and Ge, M.: Anthropogenic Effects on Biogenic Secondary Organic Aerosol Formation, *Adv. Atmospheric Sci.*, 38, 1053–1084, <https://doi.org/10.1007/s00376-020-0284-3>, 2021.
- 990

- 995 Yang, B.-Y., Qian, Z. (Min), Li, S., Chen, G., Bloom, M. S., Elliott, M., Syberg, K. W., Heinrich, J., Markevych, I., Wang, S.-Q., Chen, D., Ma, H., Chen, D.-H., Liu, Y., Komppula, M., Leskinen, A., Liu, K.-K., Zeng, X.-W., Hu, L.-W., Guo, Y., and Dong, G.-H.: Ambient air pollution in relation to diabetes and glucose-homoeostasis markers in China: a cross-sectional study with findings from the 33 Communities Chinese Health Study, *Lancet Planet. Health*, 2, e64–e73, [https://doi.org/10.1016/S2542-5196\(18\)30001-9](https://doi.org/10.1016/S2542-5196(18)30001-9), 2018.
- Yang, X., Xiao, D., Fan, L., Li, F., Wang, W., Bai, H., and Tang, J.: Spatiotemporal estimates of daily PM_{2.5} concentrations based on 1-km resolution MAIAC AOD in the Beijing–Tianjin–Hebei, China, *Environ. Chall.*, 8, 100548, <https://doi.org/10.1016/j.envc.2022.100548>, 2022.
- 1000 York, D., Evensen, N. M., Martínez, M. L., and De Basabe Delgado, J.: Unified equations for the slope, intercept, and standard errors of the best straight line, *Am. J. Phys.*, 72, 367–375, <https://doi.org/10.1119/1.1632486>, 2004.
- 1005 Zhang, Y., Favez, O., Petit, J.-E., Canonaco, F., Truong, F., Bonnaire, N., Crenn, V., Amodeo, T., Prévôt, A. S. H., Sciare, J., Gros, V., and Albinet, A.: Six-year source apportionment of submicron organic aerosols from near-continuous highly time-resolved measurements at SIRTa (Paris area, France), *Atmospheric Chem. Phys.*, 19, 14755–14776, <https://doi.org/10.5194/acp-19-14755-2019>, 2019.
- Zhang, Z. Y., Wong, M. S., and Lee, K. H.: Evaluation of the representativeness of ground-based visibility for analysing the spatial and temporal variability of aerosol optical thickness in China, *Atmos. Environ.*, 147, 31–45, <https://doi.org/10.1016/j.atmosenv.2016.09.060>, 2016.
- 1010 Zhao, B., Jiang, J. H., Gu, Y., Diner, D., Worden, J., Liou, K.-N., Su, H., Xing, J., Garay, M., and Huang, L.: Decadal-scale trends in regional aerosol particle properties and their linkage to emission changes, *Environ. Res. Lett.*, 12, 054021, <https://doi.org/10.1088/1748-9326/aa6cb2>, 2017.
- 1015 Zhao, B., Jiang, J. H., Diner, D. J., Su, H., Gu, Y., Liou, K.-N., Jiang, Z., Huang, L., Takano, Y., Fan, X., and Omar, A. H.: Intra-annual variations of regional aerosol optical depth, vertical distribution, and particle types from multiple satellite and ground-based observational datasets, *Atmospheric Chem. Phys.*, 18, 11247–11260, <https://doi.org/10.5194/acp-18-11247-2018>, 2018.
- Zhdanova, E. Y., Chubarova, N. Y., and Lyapustin, A. I.: Assessment of urban aerosol pollution over the Moscow megacity by the MAIAC aerosol product, *Atmospheric Meas. Tech.*, 13, 877–891, <https://doi.org/10.5194/amt-13-877-2020>, 2020.


1020

1025


Figures



1035 **Figure 1: Localization of European cities used for the local-to-regional analysis. Map created with Cartopy (Met Office, 2010).**



1045 **Figure 2: Scatterplot of the MAIAC against AERONET observations considering a) all the available data points in**
In Europe, and points selected based on the Angstrom Exponent (AE) assuming c) $AE < 0.5$ d) $0.5 \leq AE < 1.5$ e) $AE \geq 1.5$. Panel
 b) shows the PDF of the difference between MAIAC and AERONET values in reference to data points in a). Acronyms
 indicate: total number of points (NTOT), correlation coefficient (R), Root Mean Square Error (RMSE), Mean Bias
 Error (MBE), Normalized Mean Bias (NMB), fraction of y data between 0.5 and 2 times x (FAC2) expressed in %,
 1050 fraction of retrievals within the expected error ($EE = 0.05 + 0.05AOD$) and the equation of the regression line. Vertical
 and horizontal bars represent the x and y errors. Red solid lines represent the straight lines passing through the origin
with 2, 1 and 0.5 slope coefficients respectively. The green dotted line represents the regression line.



1055

Figure 3: Climatological seasonal mean of the Aerosol Optical Depth at 550 nm from MAIAC algorithm over the period 2000–2021. Seasons are ordered as follows: (a) March–April–May, (b) June–July–August, (c) September–October–November, and (d) December–January–February. The right side of each figure shows the latitudinal average of AOD.

1060




Figure 4: Climatology of the Aerosol Optical Depth at 550nm from MAIAC algorithm at different cities in Europe for (red) the local scale, and (orange) the regional scale, as defined in Sect. 2.1. The location of the cities is shown in Figure 1. The figure aims to enhance the contribution to the AOD enhancing due to the local source of pollution. The yellow triangles represent ~~the mean of the boxplot~~ the AOD mean, whereas the median has been reported as the line crossing the boxplot. Black dots represent the outliers.

1065






Figure 5: Scatter plot Local Vs Regional at different seasons (December–January–February (DJF), March–April–May (MAM), June–July–August (JJA), September–October–November (SON)). During the DJF and MAM seasons, the largest differences between local and regional are found. Vertical and horizontal bars represent the x and y errors. Solid and dashed lines represent the 1:1 and regression lines respectively.

1070



1075 **Figure 6: Theil–Sen (a) absolute and (b) relative change of Aerosol Optical Depth at 550 nm over the European domain for the 2000–2021 period. Only the significant (p value<0.05) pixels are reported.**

Tables

1080

LONGITUDE	LATITUDE	CITY
4.88	52.37	Amsterdam
23.72	37.98	Athens
2.15	41.39	Barcelona
20.43	44.80	Belgrade
13.40	52.52	Berlin
11.32	44.49	Bologna
4.38	50.83	Brussels
12.57	55.68	Copenhagen
-6.26	53.349	Dublin
-9.13	38.72	Lisbon
-0.12	51.50	London
-3.70	40.41	Madrid
5.4	43.3	Marseille
9.18	45.46	Milan
10.75	59.91	Oslo
2.33	48.86	Paris
14.43	50.07	Prague
12.49	41.90	Rome
18.06	59.33	Stockholm
16.36	48.21	Vienne
15.97	45.81	Zagreb

Table 1: List of European cities used for the city scale analysis. City names in bold are the cities (not metropolitan regions) with more than 1 million of inhabitants according to the Eurostat database (<https://ec.europa.eu/eurostat/cache/RCI/#?vis=city.statistics&lang=en>).

1085

1090

1095

	AOD>0.3 (%)	LOWER/UPPER BOUND [25 th /75 th]	AOD>0.3 (%)	LOWER/UPPER BOUND [25 th /75 th]	MEAN ± STD	LOWER/UPPER BOUND [25 th /75 th]	
N DAYS	AOD LOCAL		AOD REGIONAL		LTRR		CITY
886	5.3	0.06/0.16	4.1	0.07/0.16	-0.17 ± 0.01	-0.31/-0.04	Amsterdam
4081	10.6	0.11/0.23	3.2	0.07/0.18	0.32 ± 0.01	-0.04/0.52	Athens
3121	3.5	0.06/0.16	1.6	0.04/0.13	0.57 ± 0.02	0.02/0.93	Barcelona
2424	5.2	0.07/0.17	5.9	0.07/0.17	0.07 ± 0.01	-0.15/0.24	Belgrade
1631	5.3	0.05/0.16	4.1	0.06/0.15	-0.03 ± 0.01	-0.17/0.11	Berlin
3222	7.1	0.08/0.19	4.8	0.07/0.17	0.14 ± 0.01	-0.07/0.28	Bologna
1389	4.2	0.05/0.14	4.1	0.05/0.13	-0.06 ± 0.01	-0.21/0.07	Brussels
1037	3.1	0.05/0.12	2.4	0.05/0.14	-0.01 ± 0.02	-0.25/0.18	Copenhagen
910	2.5	0.04/0.10	2.6	0.04/0.10	-0.01 ± 0.02	-0.32/0.16	Dublin
445	1.3	0.07/0.12	1.5	0.05/0.11	0.55 ± 0.03	0.15/0.88	Lisbon
1080	4.0	0.04/0.12	3.1	0.04/0.10	0.13 ± 0.02	-0.08/0.29	London
3049	1.5	0.06/0.13	1.6	0.06/0.14	0.14 ± 0.01	-0.04/0.29	Madrid
4394	2.4	0.06/0.15	1.2	0.05/0.12	0.26 ± 0.01	-0.13/0.48	Marseille
3220	14.7	0.07/0.23	12.5	0.07/0.22	-0.01 ± 0.01	-0.18/0.15	Milan
1042	0.6	0.03/0.10	0.6	0.03/0.07	0.07 ± 0.02	-0.11/0.19	Oslo
1293	4.4	0.07/0.15	2.5	0.05/0.12	0.39 ± 0.02	0.01/0.64	Paris
1502	4.5	0.06/0.16	3.6	0.06/0.15	-0.03 ± 0.01	-0.21/0.14	Prague
3670	1.9	0.06/0.14	1.2	0.05/0.12	0.10 ± 0.01	-0.13/0.27	Rome
1119	1.8	0.05/0.12	1.8	0.04/0.10	0.04 ± 0.02	-0.10/0.15	Stockholm
2015	4.9	0.06/0.16	3.5	0.06/0.15	0.03 ± 0.01	-0.19/0.2	Vienna
2531	5.2	0.07/0.17	3.8	0.06/0.16	0.08 ± 0.01	-0.10/0.24	Zagreb

1100

Table 2: Aerosol Optical Depth statistics at 550 nm from MAIAC algorithm at different sites both for local and regional scale: number of days AOD>0.3, 25th and 75th distribution percentiles, AOD local-to-regional ratio mean ± its standard deviation, and AOD local-to-regional ratio 25th and 75th distribution percentiles are reported.

1105

1110

1115

TREND				
AOD LOCAL		AOD REGIONAL		CITY
ABSOLUTE (units year ⁻¹)	RELATIVE (% year ⁻¹)	ABSOLUTE (units year ⁻¹)	RELATIVE (% year ⁻¹)	
–	–	–	–	Amsterdam
–0.0017	–1.0	–0.0020	–1.3	Athens
–0.0010	–0.9	–0.0017	–1.5	Barcelona
–0.0016	–1.6	–0.0014	–1.2	Belgrade
–0.0015	–1.4	–	–	Berlin
–0.0021	–1.4	–0.0025	–1.8	Bologna
–0.0020	–1.5	–0.021	–1.5	Brussels
–	–	–	–	Copenhagen
–	–	–	–	Dublin
–	–	–	–	Lisbon
–	–	–	–	London
–	–	–	–	Madrid
–0.0005	–0.5	–0.0014	–0.9	Marseille
–0.0034	–1.4	–0.0033	–1.7	Milan
–	–	–	–	Oslo
–	–	–0.0015	–1.5	Paris
–0.0030	–1.7	–0.0030	–2.0	Prague
–0.0012	–1.1	–0.0014	–1.3	Rome
–	–	–	–	Stockholm
–0.0011	–0.9	–0.0025	–1.9	Vienna
–0.0020	–1.6	–0.0022	–1.8	Zagreb

Table 3: Optical Depth trends at local and regional scale for the different analyzed cities. Only significant trends are shown (pvalue<0.05).