

# Effect of boundary layer low-level jet on fog fast spatial propagation

Shuqi Yan<sup>1</sup>, Hongbin Wang<sup>1,\*</sup>, Xiaohui Liu<sup>2</sup>, Fan Zu<sup>1</sup>, Duanyang Liu<sup>1,\*</sup>

<sup>1</sup>Key Laboratory of Transportation Meteorology of China Meteorological Administration, Nanjing Joint Institute for Atmospheric Sciences, Nanjing, 210041, China

<sup>2</sup>Merchant Marine College, Shanghai Maritime University, Shanghai, 201306, China

*Correspondence to:* Hongbin Wang (kaihren@163.com); Duanyang Liu (liuduanyang2001@126.com)

**Abstract.** The spatiotemporal variation of fog reflects the complex interactions among fog, boundary layer thermodynamics and synoptic systems. Previous studies revealed that fog can present fast spatial propagation feature and attribute it to boundary layer low-level jet (BLLJ), but the effect of BLLJ on fog propagation is not quantitatively understood. Here we analyze a large-scale fog event in Jiangsu, China from 20 to 21 January 2020. Satellite retrievals show that fog propagates from southeast coastal area to northwest inland with the speed of 9.6 m/s, which is three times larger than the ground wind speeds. The ground meteorologies are insufficient to explain the fog fast propagation, which is further investigated by WRF simulations. The fog fast propagation could be attributed to the BLLJ occurring between 50 and 500 m, because the wind speeds (10 m/s) and directions (southeast) of BLLJ core are consistent with fog propagation. Through sensitive experiments and process analysis, three possible mechanisms of BLLJ are revealed: 1) The abundant oceanic moisture is transported inland, increasing the humidity of boundary layer and promoting condensation; 2) The oceanic warm air is transported inland, enhancing the inversion layer and favouring moisture accumulation; 3) The moisture advection probably promotes low stratus formation, and later it subsides to be ground fog by turbulent mixing of fog droplets. The fog propagation speed would decrease notably by 6.4m/s (66%) in the model if the BLLJ-related moisture and warm advections are turned off.

## 1. Introduction

Fog is a kind of low-visibility weather phenomenon that occurs at near surface, causing adverse impacts on traffic transportation. The formation, development and dissipation of fog are the comprehensive results of the interactions among radiation, moisture, microphysics, turbulence, aerosols and other factors (Gultepe et al., 2007; Koraćin et al., 2014; Nakanishi, 2000). The relations of fog with meteorological factors are highly variable under different conditions. Therefore, the mechanism of fog evolution needs to be intensively studied.

Under favourable conditions, the fog intensity or its spatial extent can develop extraordinarily fast with time. Field observations conducted at single site reveal that visibility in fog can deteriorate drastically, from about 1km to less than 200m within 30min (Li et al., 2019). It is referred to as fog burst reinforcement, which is firstly raised by Korb et al. (1970) and systematically reviewed by Liu et al. (2012) and Li et al. (2019). Fog burst reinforcement is accompanied by the drastic formation of fog droplets, sudden increase of fog liquid water and broadening of droplet spectrum (Liu et al., 2017; Liu et al., 2021). Additionally, fog can develop rather fast in spatial extent, i.e., the fast spatial propagation of fog (Zhu et al., 2022). It is reflected by the successive visibility dropping in space along a certain direction. The influencing factors of fast

35 spatial propagation could be more complex than that of the burst reinforcement at single site, which have received fewer  
36 quantitative studies recently.

37 Synoptic systems and planetary boundary layer (PBL) thermodynamic structures are key to understanding the cause of fog  
38 burst reinforcement and fast propagation. Weak cold air invasion and radiative cooling is an important factor for fog burst  
39 reinforcement and fast propagation (Liu et al., 2011; Wang et al., 2020). Dhangar et al. (2021) demonstrated that the radia-  
40 tive cooling at surface and fog top can increase supersaturation and promote fog vertical development. Shen et al. (2022)  
41 found that the different cooling rates at two nearby stations lead to a remarkable difference in fog formation time, fog du-  
42 ration and vertical extent. Sufficient water supply is also an important factor. Wobrock et al. (1992) revealed that the role  
43 of moisture advection outweighs radiative cooling in large-scale fog events. Pu et al. (2008) found that two layers of mois-  
44 ture advection enhance fog development and maintenance. Under stable synoptic systems, the PBL thermodynamic can  
45 also favour fog burst reinforcement and fast fog propagation. The formation of dense fog is usually accompanied by strong  
46 inversion layer, of which the intensity could reach 16K/100m (Pu et al., 2008; Liu et al., 2012). Liu et al. (2016) found that  
47 upper-level warm advection and low-level cold advection significantly enhance inversion intensity and promote fog de-  
48 velopment. The vapor advection resulting from southerly winds further increases fog intensity. Appropriate turbulence also  
49 facilitates fog formation and enhancement (Ye et al., 2015; Zhou and Ferrier, 2008). Turbulent results in the exchange of  
50 heat and moisture within PBL, e.g., the downward entrainment of vapor and cold air can promote condensation and droplet  
51 formation (Liu et al., 2016; Zhang et al., 2005). Other studies highlight the role of hygroscopic aerosols and aerosol indi-  
52 rect effects in strong fog events (Boutle et al., 2017; Quan et al., 2021; Shao et al., 2023; Wang et al., 2023; Yan et al.,  
53 2021).

54 Previous studies find that the large-scale fog events are accompanied by boundary layer low-level jet (BLLJ), and try to  
55 attribute the spatial propagation of fog to BLLJ. The causes of BLLJ include such as synoptic systems, terrain effect and  
56 inertial oscillation (Kraus et al., 1985). Tian et al. (2019) demonstrated that the warm-and-wet southerly BLLJ favours wa-  
57 ter vapor transportation and inversion layer construction, and later the fog is triggered by a weak cold front invasion. Wu et  
58 al. (2020) found that strong northerly BLLJ associated with cold air can destroy inversion layer and lead to early dissipa-  
59 tion of fog, while weak BLLJ can promote fog maintenance. Li et al. (2012) revealed that the strengthened turbulence gen-  
60 erated by BLLJ wind shear promotes vertical mixing and facilitates fog development. However, the relations between  
61 BLLJ and fog propagation and the key synoptic factors have not been quantitatively addressed. Also, the current horizontal  
62 and vertical observations are not sufficient to reveal the mechanism of fog propagation. It requires further investigation by  
63 numerical models.

64 In this work, we study a large-scale fog event with fast propagation feature occurring in Jiangsu Province, China from 20  
65 to 21 January 2020. By combination of observations and numerical simulations, we aim to quantitatively reveal the BLLJ  
66 effect on fast fog propagation to and identify the key impact factors and mechanisms. This work is expected to better un-  
67 derstand the complex interactions among synoptic systems, PBL thermodynamics and fog spatial propagation, as well as  
68 provide prediction indicators for operational fog forecast. The study is organized as follows: Section 2 describes the data,  
69 methods and numerical models of this study. Sections 3.1 to 3.4 analyze the fog propagation feature and PBL characteris-

70 tics. Section 3.5 quantitatively study the BLLJ effect on fast fog propagation and identifies key influencing factors. Section  
71 4 concludes the findings of this study.

## 72 **2. Data, methods and model configuration**

### 73 **2.1 Data and study area**

74 This study focuses on the Jiangsu area, China (Figure 1), where a large-scale fog event occurred from 20 to 21 January  
75 2020. We collected the data from 70 ground automatic weather stations (AWS) in Jiangsu Province, China. The data is  
76 recorded by every 10 minutes, including visibility, temperature, relative humidity (RH), wind direction and wind speed.  
77 This data is used to analyze the temporal variation of meteorology, as well as evaluate the model performance on tempera-  
78 ture, RH and wind. Additionally, the Sheyang (SY; 120.25°E, 33.76°N; 3m) station is a sounding station that used for  
79 model evaluation in the vertical direction. The sounding observations include temperature, RH, wind direction and wind  
80 speed which are sampled each second. It is conducted twice a day (00UTC and 12UTC).

81 The geostationary satellite Himawari 8 (<https://www.eorc.jaxa.jp/ptree/index.html>) is used to retrieve nighttime fog area  
82 and evaluate the model performance of fog simulation. The high spatiotemporal resolution (2km in space and 1h in time) is  
83 suitable for detecting the fast evolution of fog area. This satellite observation includes 16 bands, and the bands at 3.9 and  
84 11.2  $\mu\text{m}$  are used.

85 The ERA5 reanalysis data (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels>) is used to  
86 analyze synoptic conditions and provide initial & boundary fields for model simulation. The grid resolution is 0.125°  
87 (about 12.5km) and the time interval is 6h. All the time in this study is local time (UTC+8).

### 88 **2.2 Methods**

#### 89 2.2.1 Satellite fog retrieval

90 Since the ground AWS stations are not sufficiently fine in spatial resolution, the high spatiotemporal resolution product of  
91 Himawari 8 is suitable to study the propagation of fog. Nighttime fog has notable different optical properties at the bands  
92 of 3.9 $\mu\text{m}$  and 11.2 $\mu\text{m}$ , so it can be indicated by the dual-band brightness temperature difference ( $T_{bb3.9}$  minus  $T_{bb11.2}$ )  
93 lower than a threshold (Cermak et al., 2008). In this study, the threshold is determined to be -2 K following the dynamic  
94 threshold algorithm proposed by Di Vittorio et al. (2002). Daytime fog after 08:00 is not retrieved because we mainly fo-  
95 cus on the formation and development stage of fog before 08:00.

#### 96 2.2.2 Fog propagation speed calculation

97 We calculate the propagation speed according to satellite retrieved fog area. At 22:00 on 20 January 2020, a tiny fog area  
98 appeared at Nantong and Yanchen coastal region with an area smaller than 50km<sup>2</sup> (figure not shown). The center of this  
99 fog area is set as point A (120.6°E, 32.9°N). We draw a line starting from A with an arbitrary direction, and find its inter-  
100 section with the fog boundary area at 07:00 next day (point B). Then the propagation speed in this direction can be calcu-

101 lated by the distance from A to B divided by 9 hours (22:00~07:00). By looping from 0 to 360 with the interval of 1°,  
102 propagation speeds in all directions are calculated, and the maximum speed is defined as the fog propagation speed.

103 The fog propagation speed is verified by AWS data. We select three representative stations along the fog propagation di-  
104 rection, Dafeng (DF; 120.48°E, 33.20°N, 14m), Baoying (BY; 119.30°E, 33.23°N, 15m), Sihong (SH; 118.22°E, 33.48°N,  
105 13m) (Figure 1). According to their distances and the time differences when visibility drops to 200m, the propagation  
106 speed between two adjacent stations is calculated.

### 107 2.2.3 Process analysis on fog

108 The simulated fog is indicated by fog liquid water content (LWC). Process analysis is used to quantify the contribution of  
109 each physical process to LWC variation (Schwenkel et al., 2019; Yan et al., 2020). The variation of LWC is related to the  
110 following terms:

$$111 \quad \frac{\partial \text{LWC}}{\partial t} = - \underbrace{\left( u \frac{\partial}{\partial x} + v \frac{\partial}{\partial y} + w \frac{\partial}{\partial z} \right)}_{\text{Adv}c} \text{LWC} + \left( \frac{\partial \text{LWC}}{\partial t} \right)_{\text{Vmix}} + \left( \frac{\partial \text{LWC}}{\partial t} \right)_{\text{Cond}} + \left( \frac{\partial \text{LWC}}{\partial t} \right)_{\text{Sedi}} + \left( \frac{\partial \text{LWC}}{\partial t} \right)_{\text{other}}$$

112 where Advc includes horizontal and vertical advection, Vmix is associated with the fog droplet vertical exchange by tur-  
113 bulent mixing, Cond is the vapor condensation (negative means droplets evaporation), Sedi is fog droplets sedimentation.  
114 Other microphysical processes include autoconversion, accretion and cold phase processes. They are much smaller than the  
115 previous four processes, so they can be safely ignored.

## 116 2.3 Model configuration and experiments

117 The Weather Research and Forecasting model (WRF) is implemented to study the fast spatial propagation of fog events.  
118 Two domains are set up (Figure 1). The parent domain covers East China, with the grid size of 181×181 and grid interval  
119 of 9 km. The nested domain covers Jiangsu Province and its coastal area, with the grid size of 199×199 and grid interval of  
120 3 km. To simulate the turbulent process more reasonably, the vertical levels are refined to 42 levels, with 25 levels under  
121 1500m and 9 levels under 100m (Yang et al., 2019; Yan et al., 2020). The first model level is about 4m. The model is driv-  
122 en by the initial and boundary field from ERA5 Reanalysis. The simulation starts at 08:00 on 19 January and ends at 08:00  
123 on 21 January 2020, with the first 24h as spin-up period. All the time in this study is local time (UTC+8).

124 Fog is hard to be simulated or predicted well (Zhou et al., 2010, 2012), which is sensitive to the choice of parameterization  
125 schemes (Steenefeld et al., 2014; van der Velde et al., 2010). Through massive tests, the QNSE boundary layer scheme  
126 (Sukoriansky et al., 2005) and Pleim-Xiu land surface scheme (Pleim et al., 2009) yield the best simulation performance.  
127 Other parameterization schemes are listed in Table 1. The simulated fog is indicated by the liquid water content (LWC)  
128 greater than 0.015g/kg under the height of 500m, which corresponds to horizontal visibility less than 1km (Kunkel, 1983).

129 Apart from the base experiment, three sensitive experiments are performed to elucidate the mechanism of fast fog propa-  
130 gation (Table 1). The experiment "Tadv0" turns off the temperature advection within PBL during the fog period. The ex-  
131 periment "QvAdv0" and "QcAdv0" are the same as "Tadv0" except that turning off water vapor advection and cloud water

132 advection, respectively. The experiment "NoAdv" turns off all the advectons above. Therefore, the differences of the base  
133 experiment with Tadv0, QvAdv0, and Qcadv0 represent the effect of temperature advection, moister advection, and cloud  
134 water advection, respectively. The reasons and results of the sensitive experiments will be discussed in Section 3.5.

## 135 **3. Results**

### 136 **3.1 Fog overview and synoptic background**

137 The studied fog event occurs at the night of 20 January and dissipates in the daytime of 21 January 2020 (Figure 2). Figure  
138 3 shows the synoptic situations at 08:00 and 20:00 on 20 January. At 500hpa, a frontal zone is located north of 38°N. The  
139 Jiangsu area is dominated by prevailing westerly flows with no obvious troughs. At 850hpa, a ridge moves eastward and  
140 controls Jiangsu area. The descending motions associated with the ridge and the nocturnal radiative cooling at ground fa-  
141 vour the establishment of inversions. At ground level, a weak cold high pressure moves eastward with the central pressure  
142 of 1030hpa. The Jiangsu area is dominated by uniform pressure field with small wind speeds, which strengthens atmos-  
143 pheric stratification stability and promotes the accumulation of aerosols and moisture. The moisture condition in Jiangsu is  
144 additionally favoured by the water vapor transportation from ocean by easterly winds at 20:00. Under this conductive situ-  
145 ation, the fog event occurred from nighttime of 20 to daytime of 21 January over Jiangsu Province (Figure 2).

### 146 **3.2 Fog and ground meteorology variation**

147 Hourly Himawari 8 satellite image clearly shows the spatial propagation of fog (Figure 2). The fog initials at 22:00 on 20  
148 January in Nantong and Yanchen coastal region with an area smaller than 50km<sup>2</sup>. Later, this small fog area expands to a  
149 large-scale fog. Specifically, the southeast side of fog area varies relatively slowly, but the northwest side expands re-  
150 markably, indicating a large propagation speed. At 07:00 on 21 January, the front of fog expands to Anhui Province. After  
151 07:00, the fog begins to dissipate, and it fully disappears at 11:00 (figure not shown). Figure 4 quantitatively describes the  
152 propagation direction and speed of fog. From the east to south directions (the fourth quadrant), fog propagation speed is  
153 less than 3m/s. In the west-northwest and west directions, fog propagation speed is larger than 6m/s, and the maximum  
154 propagation speed is 9.6m/s occurring at 160° direction (in Cartesian coordinate system). The fast propagation of fog is  
155 also reported previously in Jiangsu area (Gao et al., 2023; Zhu et al., 2022), where the fog propagates from coastal area to  
156 west boundary of Jiangsu within about 10h.

157 Visibilities at three representative stations, Dafeng (DF), Baoying (BY) and Sihong (SH) are used to verify the fog propa-  
158 gation speed calculated by satellite (Table 2; Figure 5). At DF, fog forms (visibility less than 1km) early at 19:45 on 20  
159 January. The visibility drops sharply at 23:15 and reaches the minimum at about 00:15. At BY and SH, fog forms in turn,  
160 and their visibilities also have burst decreasing feature at 03:40 and 07:00, respectively. We calculate the fog propagation  
161 speed by the distances among stations and the time differences when visibility drops to 200m. The propagation speed is 7.6  
162 m/s between DF and BY and 8.3 m/s between BY and SH. These values correspond to the speed calculated by satellite  
163 observation.

164 Figure 5 shows the variation of other meteorological fields. We focus on the characteristics from fog formation to the burst  
165 visibility dropping (indicated by yellow dashed lines). At DF, the northerly wind decreases to lower than 1.5m/s at fog  
166 formation, which causes the weak cold advection and temperature decreasing. The temperature keeps decreasing and fa-  
167 vours the burst reduction of visibility at 23:15. The vapor content (indicated by dew point) increases sharply before 17:00  
168 and decreases slightly since then, so the RH increasing after fog formation is caused by temperature drop. At BY and SH,  
169 the wind directions are dominantly southeast and the speeds are generally less than 2m/s before fog formation. The tem-  
170 perature keeps decreasing and vapor content keeps increasing, leading to the further reduction of visibility. Later, the  
171 southeasterly winds obviously enhance by about 1m/s, which may contribute to the burst visibility dropping due to the in-  
172 tensified vapor advection from ocean.

173 The preliminary cause of fog formation and intensification are summarized. As located near the ocean, the moisture at DF  
174 reaches the maximum prior to fog formation, so the fog formation and intensification are largely caused by radiative cool-  
175 ing and weak cold advection. At BY and SH, the temperature cooling rate is weaker than DF, which is partly due to the  
176 weak warm advection by southeasterly winds. The vapor advection by southeasterly winds favours fog development, and  
177 the burst decrease in visibility coincides with the increase in wind speed. Therefore, deduced from BY and SH, the vapor  
178 transportation associated with southeasterly winds could be an important reason for northwesterly propagation of fog.  
179 However, it is obvious that the ground wind speed is rather small compared with fog propagation speed. Statistics on AWS  
180 stations show that although wind direction (east, southeast and south winds at 70% stations) is generally in accordance  
181 with fog propagation direction, wind speed is lower than 3m/s at 97% stations from 22:00 to 07:00, which is about  
182 one-third of the fog propagation speed. Therefore, the ground meteorological field is insufficient to explain the fast propa-  
183 gation of fog. The fog PBL characteristics and the key influencing factors need to be investigated by numerical simula-  
184 tions.

### 185 **3.3 Model evaluation**

186 Figure 6a evaluates the model performance on temperature, relative humidity (RH) and wind field at surface. The simulat-  
187 ed temperature and RH agree well with observations, with the root mean square error (RMSE) of 1.0K and 11%, respec-  
188 tively. The simulation reasonably captures the wind direction transition from north to east, and the RMSE is less than 1m/s.

189 Figure 6b evaluates the model performance on temperature, RH and wind field in the vertical direction at SY sounding  
190 station. The temperature profile is simulated well by the model, with the mean bias of less than 1K. The RH bias is rela-  
191 tively small below about 200m, while it is a bit larger above 200m at 08:00 on 21 January. The simulated wind speed and  
192 direction are basically consistent with observation. The large winds (greater than 6m/s) at about 200m are well reproduced  
193 by the model, indicating that the model reasonably simulates boundary layer low-level jet. Studies on boundary layer  
194 low-level jet are presented in next sections.

195 Figure 2 compares the satellite observed and simulated fog area. The simulation is only evaluated before 07:00, because  
196 the dissipation of fog after 08:00 is not the focus in this study. The model reasonably captures the spatiotemporal evolution  
197 of fog, with a slight overestimation of 5~10% in fog area.

198 Overall, the simulation reasonably captures the temporal variation of meteorology and reproduces the spatial propagation  
199 of fog. It establishes the basis for discerning the mechanism of fog propagation.

### 200 **3.4 Characteristics of fog and PBL structure**

201 The thermodynamic variation of PBL is crucial for understanding the propagation of fog. Figure 7a shows the temporal  
202 variation of horizontal winds in vertical directions. The simulated wind speed is consistently smaller than 4m/s under about  
203 30m, while it remarkably increases with height. At 18:00 on 20 January, a large wind speed zone (>6m/s) forms at the  
204 height between 50 and 500m in the east of 120°E. Since then, the large wind zone moves westward quickly accompanied  
205 by wind speed increasing. During the fog period, the average wind speed exceeds 6m/s at the height between 50 to 500m  
206 (Figure 7b), which is commonly larger than the wind speed in most fog events. Here, we refer to this large wind speed  
207 zone as boundary layer low-level jet (BLLJ). The existence of BLLJ is supported by ERA5 reanalysis on 1000hpa and  
208 975hpa levels (Figure 7b).

209 The formation of BLLJ is likely caused by the easterly movement of a high pressure at 1000hpa over East China. The cen-  
210 tral pressure gets enhanced, which strengthens the pressure gradient over Jiangsu area and favours wind speed increasing  
211 (figure not shown). The jet core (maximum wind speed) occurs at about 1000hpa (200m), with the time-averaged speed of  
212 10m/s (Figure 7b). At that level, the dominant wind direction is southeast and the wind speed over fog area is 8~16m/s  
213 (Figure 7c), which can fit the propagation direction and speed of fog. Also, the expansion speed of vertical fog zone is  
214 comparable to the movement speed of jet core (Figure 7a). Therefore, we hypothesize that the southeasterly BLLJ could  
215 account for the fast propagation of fog.

216 Previous studies reveal that southerly BLLJ can transport abundant water vapor to China inland and thus promote fog for-  
217 mation (Liu et al., 2016; Tian et al., 2019). Figure 8 shows the temporal variation of water vapor mixing ratio ( $Q_v$ ) profiles.  
218 Since the vapor content over the ocean is higher, it is transported to inland areas by southeasterly BLLJ. The BLLJ can  
219 further increase the  $Q_v$  in PBL by wind speed horizontal convergency and vertical shear. The larger wind speed in BLLJ  
220 zone and lower wind speed outside BLLJ zone cause wind speed convergence, which favours the increase in PBL moisture.  
221 Additionally, the turbulence generated by vertical shear of wind speed can promote vapor turbulent mixing, leading to the  
222 higher  $Q_v$  above surface being entrained downward and increasing the ground  $Q_v$  (Gao et al., 2007). The  $Q_v$  under 300m  
223 is generally higher than 3g/kg under the effect of BLLJ. Wu et al. (2020) also found that BLLJ continuously transports  
224 water vapor to fog layer, resulting in surface  $Q_v$  higher than 3g/kg. It is notable that the expansion of vertical fog area co-  
225 incides with the movement of the zone of  $Q_v > 4g/kg$ . Therefore, moister advection by BLLJ could be an important reason  
226 for fast fog propagation.

227 BLLJ is reported to result in warm advection and deepen inversion layer previously (Tian et al., 2019), and inversion layer  
228 is an important reason for fog burst reinforcement in most fog cases (e.g., Li et al., 2019; Liu et al., 2012; Jiao et al., 2016).  
229 Figure 9 shows the temporal variation of temperature profile and inversion layer. The inversion layer here refers to the  
230 height above ground where temperature monotonically decreases with height. Since 20:00 on 20 January, the ground tem-  
231 perature keeps decreasing due to radiative cooling. Within the fog area, the temperature drop is more significant, which is  
232 due to the longwave radiative cooling by fog droplets (Bott, 1991; Jia et al., 2018). Approximately above the fog top, there

233 is an obvious warm air mass transported from ocean to inland areas. The BLLJ-induced warm advection increases vertical  
234 temperature gradient and strengthens atmospheric stability. Accordingly, the inversion height over non-fog areas basically  
235 keeps increasing. The approximate inversion layer height is about 100~300m, which is consistent with previous studies  
236 (Dorman et al., 2021; Li et al., 2019). The maximum inversion intensity of 15K/100m, which is also reported in a dense  
237 fog event (16K/100m) by Pu et al. (2008). It favours the accumulation of vapor and condensation nuclei, which is also a  
238 possible reason for fog formation in the downstream area.

239 Additionally seen from Figure 9, the west boundary of vertical fog region below about 100m has a negative slope, i.e., fog  
240 forms at upper level ahead of forming at ground. The upper-level fog with no ground contact is referred to as low stratus.  
241 The height at which fog/low stratus firstly forms is shown in Figure 10. An initial fog area forms at ground level before  
242 00:00 on 21 January. Since then, low stratus forms at upper level (about 10~66m) over the downstream area, while the  
243 ground fog in downstream area forms about 0~20min later than low stratus. The formation of low stratus may also be  
244 caused by the BLLJ-induced moisture advection. In addition, the cloud water advection (Section 2.2.3) to downstream area  
245 by BLLJ could also be a potential reason. We hypothesize that the formation of ground fog is partly favoured by the stratus  
246 lowering, which has been reported by previous studies (e.g., Haeffelin et al., 2010; Liu et al., 2012); the base height of  
247 stratus can be smaller than 100m before fog formation (Dupont et al., 2012; Fathalli et al., 2022), which is basically close  
248 to our results (10~66m in Figure 10). While in this event, the stratus lowering phenomenon remains to be verified by addi-  
249 tional high-spatiotemporal resolution vertical observations.

250 According to above results, three potential factors for fog propagation are raised: BLLJ-related temperature advection,  
251 moisture advection and cloud water advection. These advections possibly promote low stratus formation within 100m  
252 above surface, and subsequently the low stratus could subside to be ground fog by the turbulent mixing or sedimentation of  
253 cloud droplets. Currently, their contributions to fog propagation have not been quantitatively revealed. Therefore, it will be  
254 addressed in the next section.

### 255 **3.5 Quantitative reasons for fast fog propagation**

256 Four sensitive experiments, Tadv0, QvAdv0, QcAdv0 and NoAdv0 (Section 2.3) are conducted to quantify the respective  
257 contributions of temperature advection, moisture advection, cloud water advection and all these advections to fog propaga-  
258 tion (Figure 11). Under the condition with no advections (Figure 11a-d), there is a 80% decrease in fog area and a 6.4m/s  
259 (66%) decrease in propagation speed, which highlights the role of BLLJ-related advections. When turning off temperature  
260 advection (Tadv0) (Figure 11e-h), the original fog area in the base experiment shrinks 50% in size and breaks into separate  
261 fog patches, and the propagation speed decreases by about 5.2m/s (54%). When turning off moisture advection (QvAdv0)  
262 (Figure 11i-l), the fog area shrinks by 62% in size and the propagation speed decreases by about 4.6m/s (48%). When  
263 turning off cloud water advection (QcAdv0) (Figure 11m-p), the fog area nearly keeps unchanged during 00:00~04:00 and  
264 decreases moderately in size (about 25%) at 06:00. The propagation speed decreases moderately by 2.4m/s (25%). Deduced  
265 from the changes in fog area and propagation speed under various experiments, we can infer that the BLLJ-related warm  
266 and moisture advection, especially moisture advection, could be the major cause of fast spatial propagation, while cloud  
267 water advection has a minor contribution.



268 We further perform process analysis on LWC (Section 2.2.3) to illustrate the mechanism of fog propagation (Figure 12).  
269 The horizontal and vertical values of Advc and Sedi are at least one order of magnitude smaller than that of Cond and Sedi,  
270 indicating that cloud water transportation to downstream areas and droplet sedimentation to ground are not the causes of  
271 fog propagation. At 00:00 on ground level, Cond is positive over the newly formed fog area (blue and cyan colors sur-  
272 rounding the fog area), indicating that fog firstly forms at ground by radiative cooling before 00:00. After 02:00, Cond is  
273 almost negative over the entire fog area, indicating that fog does not firstly form at the ground level (otherwise Cond  
274 would have positive values). The formation of ground fog may be contributed by the LWC turbulent entrainment from up-  
275 per level, because Vmix shows significant positive values after 02:00. In the vertical direction, Vmix and Cond are still  
276 two dominant physical processes (Figure 12b), and their signs show opposite patterns. At lower level (0~30m), Cond is  
277 negative and Vmix is positive, which is the same as their ground characteristics. At upper level (30~200m), Cond is posi-  
278 tive and Vmix is negative instead, indicating that cloud water is produced by vapor condensation at upper level and then  
279 being entrained to ground. The significant positive Cond supports that BLLJ-related moisture advection promotes vapor  
280 condensation and low stratus formation above surface, and the significant positive Vmix may indicate that the low stratus  
281 favours ground fog formation by turbulent exchange of LWC.

## 282 4. Discussions

283 Previous studies have elucidated the qualitative reasons for fog propagation. In this study, we describe the feature of fast  
284 fog propagation and identify its key impact factors more quantitatively. Figure 13 summarizes the mechanism of fog prop-  
285 agation. During the nighttime, a southerly BLLJ controls the study region, and the jet core intensity is about 10m/s which  
286 occurs at about 200m. The ground fog propagates northwestward with the speed of 9.6m/s. The BLLJ favours the fast fog  
287 propagation by three possible mechanisms: 1) BLLJ transports sufficient vapor from ocean to inland area. The turbulence  
288 strengthened by wind speed shear further moistens the PBL and promotes vapor condensation. This could be the dominant  
289 mechanism. 2) BLLJ transports warmer air from ocean to inland area and deepens the inversion layer. The strengthened  
290 inversion favours the accumulation of vapor and condensation nuclei. 3) The strong moisture advection could promote the  
291 low stratus formation in the downstream area, and later it subsides to be ground fog by turbulent exchange of cloud drop-  
292 lets. The stratus lowering phenomenon needs to be verified by additional observations.

293 The results could facilitate the understanding of cloud formation and development. Clouds, such as convective clouds, can  
294 develop and expand extraordinarily fast under strong synoptic forcing or unstable conditions. Fog can be viewed as a kind  
295 of near-surface stratus cloud, which usually forms under stable conditions with weak synoptic forcings. However, as re-  
296 vealed in this study, it can also develop and propagate fast under the effect of BLLJ. The quantitative relations between  
297 BLLJ and fog fast propagation may have implications on the cloud formation and development mechanism under stable  
298 synoptic conditions.

## 299 5. Conclusions

300 Previous studies have found that the spatial propagation of fog could be rather fast under favourable conditions, and the  
301 boundary layer low-level jet (BLLJ) could be a potential reason. In this study, we analyze the fast spatial propagation fea-  
302 ture of a large-scale fog event in Jiangsu Province, China by high spatiotemporal resolution ground and satellite observa-  
303 tions. The key impact factors and mechanisms of the BLLJ effect on fast spatial propagation are quantitatively revealed by  
304 WRF model simulations. Results show that:

305 The fog initials at 22:00 on 20 January 2020 over Jiangsu coastal area, and it reaches the west boundary of Jiangsu at 07:00  
306 next day. Satellite retrievals show that the southeast side of fog area varies slightly but the northwest side expands fast,  
307 with the maximum propagation speed of 9.6m/s. During the fog period, the ground wind direction is consistent with fog  
308 propagation, which favours the vapor transportation from ocean and promotes fog formation. However, the wind speed  
309 ( $<3\text{m/s}$ ) is at least one-third less than the fog propagation speed. Therefore, the ground meteorologies are insufficient to  
310 explain the fast propagation of fog. The influencing factors and mechanisms need to be investigated by exploring the PBL  
311 characteristics through numerical simulations.

312 The WRF model well simulates the temporal variation of meteorologies and reproduces the spatiotemporal evolution of  
313 fog area. A BLLJ ( $>6\text{m/s}$ ) exists at the height between 50 and 500m. The jet core occurs at 1000hpa (200m) with the  
314 southeasterly winds of 10m/s, which can fit the propagation direction and speed of fog. Therefore, the southeasterly BLLJ  
315 is hypothesized to be the cause of fast propagation. BLLJ creates favourable PBL conditions by transporting moisture and  
316 warm air from ocean. The moisture advection and the vapor turbulent mixing generated by wind speed shear increase the  
317 humidity within PBL, and the propagation of fog area coincides with the movement of high humidity zone (vapor mixing  
318 ratio  $>4\text{g/kg}$ ). The warm advection from ocean deepens inversion layer and additionally favours the accumulation of mois-  
319 ture and condensation nuclei. Additionally, it is found that low stratus could form above surface and subsides to be ground  
320 fog within 0~20min. The moisture advection is also responsible for the formation of low stratus.

321 Sensitive experiments quantitatively reveal the contributions of moisture advection and temperature advection to fog  
322 propagation. When moisture (temperature) advection is turned off, the fog area decreases by 62% (50%) and the propaga-  
323 tion speed decrease by about 4.6m/s (5.2m/s). Process analysis on fog liquid water content (LWC) further illustrates the  
324 mechanism of fog propagation. Condensation (Cond) and LWC turbulent exchange ( $V_{\text{mix}}$ ) are two important physical  
325 processes. At upper level (30~200m), Cond is positive and  $V_{\text{mix}}$  is negative. It indicates that BLLJ-related moisture ad-  
326 vection significantly promotes condensation and probably favours low stratus formation. At ground and lower level  
327 (0~30m), Cond is basically negative and  $V_{\text{mix}}$  is positive. It indicates that cloud droplets at upper level are entrained  
328 downward by turbulent mixing, leading to the subsequent formation of ground fog. The stratus lowering phenomenon  
329 needs to be verified by additional observations.

330 In this study, by combination of observations and simulations, we have revealed the effect of southeasterly BLLJ on fog  
331 propagation, and quantified the contributions of BLLJ-related moisture advection and temperature advection to fog propa-  
332 gation. Three possible mechanisms are concluded: 1) Moisture advection from ocean promotes vapor condensation in  
333 downstream area, which could be the dominant cause; 2) Warm advection from ocean deepens inversion layer and addi-

334 tionally promote vapor accumulation within PBL. 3) The moisture advection probably promotes low stratus formation first,  
335 and later it subsides to be ground fog by turbulent mixing of cloud droplets. The coexistence of fast fog propagation and  
336 BLLJ is not a common phenomenon, so finding more cases requires additional work. It should be addressed in future stud-  
337 ies in order to deeply understand the relationships between fog propagation and BLLJ under different regions and synoptic  
338 conditions. Their quantitative relationships could facilitate the understanding of cloud formation and development under  
339 stable synoptic conditions, since fog can be viewed as near-surface stratus cloud that can potentially propagate fast under  
340 stable conditions.

341

342 *Code and data availability.* Some of the data repositories have been listed in Section 2. The other data, model outputs and  
343 codes can be accessed by contacting Duanyang Liu via liuduanyang2001@126.com.

344 *Author contributions.* SY performed the model simulation, data analysis and manuscript writing. HW and DL proposed the  
345 idea, supervised this work and revised the manuscript. XL helped the revision of the manuscript. FZ provided and analyzed  
346 the observation data.

347 *Competing interests.* The authors declare that they have no conflict of interest.

348 *Acknowledgements.* This work is supported by the Special Project of Innovative Development of CMA (CXFZ2023J022),  
349 Open Research Foundation of Jiangsu Marine Meteorology (HYQX2022), Beijige Foundation (BJG202307), Research  
350 Foundation of Jiangsu Meteorology Bureau (KM202307), Basic Research Fund of CAMS (2022Y025).

351

## 352 **References**

- 353 Andreas, E.L., Claffy, K.J., and Makshtas, A.P.: Low-level atmospheric jets and inversions over the western Weddell Sea,  
354 *Boundary Layer Meteorol*, 97(3), 459–486, <https://doi.org/10.1023/A:1002793831076>, 2000.
- 355 Bott, A.: On the influence of the physico-chemical properties of aerosols on the life cycle of radiation fogs, *J. Aerosol Sci*,  
356 21(1-2), 1–31, <https://doi.org/10.1007/BF00119960>, 1991.
- 357 Boutle, I., Price, J., Kudzotsa, I., Kokkola, H., and Romakkaniemi, S.: Aerosol-fog interaction and the transition to well-mixed  
358 radiation fog, *Atmos. Chem. Phys.*, 18(11), 1–19, <https://doi.org/10.5194/acp-18-7827-2018>, 2017.
- 359 Cermak, J. and Bendix, J.: A novel approach to fog/low stratus detection using Meteosat 8 data, *Atmos. Res.*, 87(3-4), 279–292,  
360 <https://doi.org/10.1016/j.atmosres.2007.11.009>, 2008.
- 361 Dhangar, N. G., Lal, D. M., Ghude, S. D., Kulkarni, R., and Rajeevan, M.: On the conditions for onset and development of fog  
362 over new delhi: an observational study from the wifex, *Pure Appl. Geophys*, 673, 1-20,  
363 <https://doi.org/10.1007/s00024-021-02800-4>, 2021.
- 364 Dorman, C.E., Hoch, S.W., Gultepe, I., Wang, Q., Yamaguchi, R., Fernando, H., Krishnamurthy, R.: Large-Scale Synoptic Sys-  
365 tems and Fog During the C-FOG Field Experiment. *Boundary-Layer Meteorol*, 181, 171–202,  
366 <https://doi.org/10.1007/s10546-021-00641-1>, 2021.
- 367 Dupont, J., Haeffelin, M., Protat, A., Bouniol, D., Boyouk, N., and Morille, Y.: Stratus–Fog Formation and Dissipation: A 6-Day  
368 Case Study, *Boundary-Layer Meteorol*, 143, 207–225, <https://doi.org/10.1007/s10546-012-9699-4>, 2012.
- 369 Di Vittorio, A. V. and Emery, W. J.: An automated, dynamic threshold cloud-masking algorithm for daytime AVHRR images over  
370 land, *IEEE Trans. Geosci*, 40, 1682–1694, <https://doi.org/10.1109/TGRS.2002.802455>, 2002.

371 Fathalli, M., Lac, C., Burneta, F., and Vié, B.: Fog due to stratus lowering: Experimental and modelling case study, *Q. J. R. Me-*  
372 *teorol*, 148(746), 2299–2324. <https://doi.org/10.1002/qj.4304>, 2022.

373 Gao, S., Lin, H., Shen, B., and Fu, G.: A heavy sea fog event over the Yellow Sea in March 2005: Analysis and numerical mod-  
374 eling, *Adv Atmos Sci*, 24, 65-81, <https://doi.org/10.1007/s00376-007-0065-2>, 2007.

375 Gao, Y., Liu, D., Yan, S., Zhou, W., Wang, H., Zu, F., Mei, Q., Yi, C., and Sheng, Y.: Influence of sea-land breeze on formation  
376 and dissipation of severe dense fog and its explosive enhancement in the Yellow Sea Coastal Area, *Sci. China Earth Sci.*,  
377 2023. (accepted; under translation into English)

378 Gultepe, I., Tardif, R., Michaelides, S. C., Cermak, J., Bott, A., Bendix, J., Muller, M. D., et al.: Fog research: a review of past  
379 achievements and future perspectives, *Pure Appl. Geophys*, 164, 1121–1159, <https://doi.org/10.1007/s00024-007-0211-x>,  
380 2007.

381 Haeffelin, M., Bergot, T., Elias, T., Tardif, R., Carrer, D., Chazette, P., and Zhang, X.: PARISFOG: Shedding new light on fog  
382 physical processes, *Bull Am Meteorol Soc*, 91(6), 767-783, <https://doi.org/10.1175/2009bams2671.1>, 2010.

383 Jia, X., Quan, J., Zheng, Z., Liu, X., Liu, Q., He, H., and Liu, Y.: Impacts of anthropogenic aerosols on fog in North China Plain,  
384 *J. Geophys. Res.-Atmos.*, 124, 252–265, <https://doi.org/10.1029/2018jd029437>, 2018.

385 Jiao, S., Zhu, C., Zhu, Y., Yuan, C., Zu, F., and Sun, K.: A discussion on the reason for a rare persistent heavy fog event in Jiang-  
386 su Province, *Acta Meteorol. Sin*, 74, 200–212, <https://doi.org/10.11676/qxxb2016.015>, 2016.

387 Koraćin, D., Dorman, C. E., Lewis, J. M., Hudson, J. G., Wilcox, E. M., and Torregrosa, A.: Marine fog: A review, *Atmos. Res.*,  
388 143, 142-175, <https://doi.org/10.1016/j.atmosres.2013.12.012>, 2014.

389 Korb, G. and Zdunkowski, W.: Distribution of radiative energy in ground fog, *Tellus*, 22(3), 298-320,  
390 <https://doi.org/10.3402/tellusa.v22i3.10223>, 1970.

391 Kraus, H., Malcher, J., and Schaller, E.: A nocturnal low level jet during PUKK, *Boundary Layer Meteorol*, 31, 187-195,  
392 <https://doi.org/10.1007/BF00121177>, 1985.

393 Kunkel, B. A.: Parameterization of Droplet Terminal Velocity and Extinction Coefficient in Fog Models, *J. Appl. Meteorol*, 23(1),  
394 34–41, [https://doi.org/10.1175/1520-0450\(1984\)023<0034:PODTVA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1984)023<0034:PODTVA>2.0.CO;2), 1983.

395 Li, P. and Fu, G.: The Formation Mechanism of a Spring Sea Fog Event over the Yellow Sea Associated with a Low-Level Jet,  
396 *Weather and Forecasting*, 27(6), 1538-1553, <https://doi.org/10.1175/WAF-D-11-00152.1>, 2012.

397 Li, Z. H., Liu, D. Y., Yang, J.: The microphysical processes and macroscopic conditions of the radiation fog droplet spectrum  
398 broadening, *Chinese J. Atmospheric Sci.*, 35, 41–54, <https://doi.org/10.3878/j.issn.1006-9895.2011.01.04>, 2011. (in Chinese)

399 Li, Z., Liu, D., Yan, W., Wang, H., Zhu, C., Zhu, Y., and Zu, F.: Dense fog burst reinforcement over Eastern China: A review,  
400 *Atmos. Res.*, 230(D19), 104639, <https://doi.org/10.1016/j.atmosres.2019.104639>, 2019.

401 Liu, D., Yang, J., Niu, S., and Li, Z.: On the Evolution and Structure of a Radiation Fog Event in Nanjing, *Adv Atmos Sci*, 28(1),  
402 223-237, <https://doi.org/10.1007/s00376-010-0017-0>, 2011.

403 Liu, D. Y., Niu, S. J., Yang, J., Zhao L., Lv, J., and Lu, C.: Summary of a 4-year fog field study in Northern Nanjing, part 1: fog  
404 boundary layer, *Pure Appl. Geophys*, 169, 809–819, <https://doi.org/10.1007/s00024-011-0343-x>, 2012.

405 Liu, D., Yan, W., Yang, J., Pu, M., Niu, S., Li, Z.: A Study of the Physical Processes of an Advection Fog Boundary Layer,  
406 *Boundary Layer Meteorol*, 158(1), 125-138, <https://doi.org/10.1007/s10546-015-0076-y>, 2016.

407 Liu, D., Li, Z., Yan, W., and Li, Y.: Advances in fog microphysics research in China, *Asia Pac J Atmos Sci*, 53(1), 131–148,  
408 <https://doi.org/10.1007/s13143-016-0028-6>, 2017.

409 Liu, Q., Wang, Z. Y., Wu, B. G., Liu, J. L., and Gultepe, I.: Microphysics of fog bursting in polluted urban air, *Atmospheric En-*  
410 *viron.*, 10, 118357, <https://doi.org/10.1016/j.atmosenv.2021.118357>, 2021.

411 Nakanishi, M.: Large-eddy simulation of radiation fog. *Boundary Layer Meteorol*, 94, 461-493,  
412 <https://doi.org/10.1023/A:1002490423389>, 2000.

413 Pleim, J. E., Gilliam, R.: An indirect data assimilation scheme for deep soil temperature in the Pleim-Xiu land surface model, *J.*  
414 *Appl. Meteorol*, 48, 1362-1376, <https://doi.org/10.1175/2009JAMC2053.1>, 2009.

415 Pu, M. J., Zhang, G. Z., Yan, W. L., and Li, Z. H.: Features of a rare advection-radiation fog event, *Sci. China Earth Sci.*, 51(7),  
416 1044–1052, <https://doi.org/10.1007/s11430-008-0071-y>, 2008.

417 Quan, J., Liu, Y., Jia, X., Liu, L., Dou, Y., Xin, J., and Seinfeld, J. H.: Anthropogenic aerosols prolong fog lifetime in China, *En-*  
418 *viron. Res. Lett.*, 16(4), 044048, <https://doi.org/10.1088/1748-9326/abef32>, 2021.

419 Schwenkel, J. and Maronga, B.: Large-eddy simulation of radiation fog with comprehensive two-moment bulk microphysics:  
420 impact of different aerosol activation and condensation parameterizations, *Atmos. Chem. Phys.*, 19(10), 1-23,  
421 <https://doi.org/10.5194/acp-19-7165-2019>, 2018.

422 Shao, N., Lu, C., Jia, X., Wang, Y., Li, Y., Yin, Y., Zhu, B., Zhao, T., Liu, D., Niu, S., Fan, S., Yan, S., and Lv, J.: Radiation fog  
423 properties in two consecutive events under polluted and clean conditions in the Yangtze River Delta, China: a simulation  
424 study, *Atmos. Chem. Phys.*, 23, 9873–9890, <https://doi.org/10.5194/acp-23-9873-2023>, 2023.

425 Shen, P., Liu, D., Gultepe, I., Lin, H., Cai, N., and Cao, S.: Boundary layer features of one winter fog in the Yangtze River Delta,  
426 China, *Pure Appl. Geophys.*, 179(9), 3463-3480, <https://doi.org/10.1007/s00024-022-03119-4>, 2022.

427 Steeneveld, G. J., Ronda, R. J., and Holtslag, A. A. M.: The Challenge of Forecasting the Onset and Development of Radiation  
428 Fog Using Mesoscale Atmospheric Models, *Boundary Layer Meteorol.*, 154(2), 265–289,  
429 <https://doi.org/10.1007/s10546-014-9973-8>, 2014.

430 Sukoriansky, S., Galperin, B., Perov, V.: Application of a new spectral model of stratified turbulence to the atmospheric boundary  
431 layer over sea ice, *Boundary Layer Meteorol.*, 117, 231–257, <https://doi.org/10.1007/s10546-004-6848-4>, 2005.

432 Tian, M., Wu, B., Huang, H., Zhang, H., Zhang, W., and Wang, Z.: Impact of water vapor transfer on a Circum-Bohai-Sea heavy  
433 fog Observation and numerical simulation, *Atmos. Res.*, 229, 1-22, <https://doi.org/10.1016/j.atmosres.2019.06.008>, 2019.

434 van der Velde, I. R., Steeneveld, G. J., Wichers Schreur, B. G. J., and Holtslag, A. A. M.: Modeling and Forecasting the Onset  
435 and Duration of Severe Radiation Fog under Frost Conditions, *Mon Weather Rev.*, 138(11), 4237–4253,  
436 <https://doi.org/10.1175/2010mwr3427.1>, 2010.

437 Wang, H., Zhang, Z., Liu, D., Zhu, Y., Zhang, X., and Yuan, C.: Study on a Large-Scale Persistent Strong Dense Fog Event in  
438 Central and Eastern China, *Adv. Meteorol.*, 4, 1–15, <https://doi.org/10.1155/2020/8872334>, 2020.

439 Wang, Y., Lu, C., Niu, S., Lv, J., Jia, X., Xu, X., et al.: Diverse dispersion effects and parameterization of relative dispersion in  
440 urban fog in eastern China, *J. Geophys. Res.-Atmos.*, 128, e2022JD037514, <https://doi.org/10.1029/2022JD037514>, 2023.

441 Wei, W., Zhang, H. S., Ye, X. X.: Comparison of low-level jets along the north coast of China in summer, *J. Geophys.*  
442 *Res.-Atmos.*, 119(16), 9692–9706, <https://doi.org/10.1002/2014JD021476>, 2014.

443 Wobrock, W., Schell, D., Maser, R., Kessel, M., Jaeschke, W., Fuzzi, S., and Bendix, J.: Meteorological characteristics of the Po  
444 Valley fog, *Tellus B*, 44(5), 469-488, <https://doi.org/10.3402/tellusb.v44i5.15562>, 1992.

445 Wu, B., Li, Z., Ju, T., and Zhang, H.: Characteristics of Low-level jets during 2015–2016 and the effect on fog in Tianjin, *Atmos.*  
446 *Res.*, 245, 105102, <https://doi.org/10.1016/j.atmosres.2020.105102>, 2020.

447 Yan, S., Zhu, B., Zhu, T., Shi, C., Liu, D., Kang, H., Lu, W., and Lu, C.: The effect of aerosols on fog lifetime: observational ev-  
448 idence and model simulations, *Geophys. Res. Lett.*, 48(2), e2020GL61803, <https://doi.org/10.1029/2020GL091156>, 2021.

449 Yang, Y., Hu, X.-M., Gao, S., and Wang, Y.: Sensitivity of WRF simulations with the YSU PBL scheme to the lowest model level  
450 height for a sea fog event over the Yellow Sea, *Atmos. Res.*, 215, 253–267, <https://doi.org/10.1016/j.atmosres.2018.09.004>,  
451 2019.

452 Ye, X., Wu, B., and Zhang, H.: The turbulent structure and transport in fog layers observed over the Tianjin area, *Atmos. Res.*,  
453 153, 217-234, <https://doi.org/10.1016/j.atmosres.2014.08.003>, 2015.

454 Zhang, G., Bian, L., Wang, J., Yang, Y., Yao, W., and Xu, X.: The boundary layer characteristics in the heavy fog formation pro-  
455 cess over Beijing and its adjacent areas, *Sci. China Earth Sci.*, 48, 88-101, <https://doi.org/10.1360/05yd0029>, 2005.

456 Zhou, B. and Ferrier, B.: Asymptotic Analysis of Equilibrium in Radiation Fog, *J. Appl. Meteorol. and Climatology*, 47,  
457 1704-1722, <https://doi.org/10.1175/2007JAMC1685.1>, 2008.

458 Zhou, B. and Du, J.: Fog prediction from a multimodel mesoscale ensemble prediction system, *Weather and Forecasting*, 25(1),  
459 303-322, <https://doi.org/10.1175/2009WAF2222289.1>, 2010.

460 Zhou, B., Du, J., Gultepe, I., and Dimego, G.: Forecast of low visibility and fog from NCEP: Current status and efforts, *Pure*  
461 *Appl. Geophys.*, 169, 895-909, <https://doi.org/10.1007/s00024-011-0327-x>, 2012.

462 Zhu, Y., Zhu, C., Zu, F., Wang, H., Liu, Q., Qi, M., and Wang, Y.: A persistent fog event involving heavy pollutants in Yancheng  
463 area of Jiangsu Province, *Adv. Meteorol.*, 2018, 2512138, <https://doi.org/10.1155/2018/2512138>, 2018.

464 Zhu, Y., Li, Z., Zu, F., Wang, H., Liu, Q., Qi, M., and Wang, Y.: The propagation of fog and its related pollutants in the Central  
465 and Eastern China in winter, *Atmos. Res.*, 265, 105914, <https://doi.org/10.1016/j.atmosres.2021.105914>, 2022.

467 Table 1. Model parameterization schemes and sensitive experiments

Physical scheme	Option
Boundary layer	QNSE
Microphysics	Lin double moment
Longwave radiation	RRTM
Shortwave radiation	Goddard
Land surface	Pleim-Xiu
Cumulus	Grell-3D
Grid nudging	Off
Observation nudging	Off
Experiment	Description
Base	The base condition
Tadv0	Turning off temperature advection
QvAdv0	Turning off water vapor advection
QcAdv0	Turning off cloud water advection
NoAdv	Turning off all advections above

468


469 Table 2. The times when visibility reaches 1000m, 500m and 200m at three representative stations. (DF:Dafeng,  
470 BY:Baoying, SH:Sihong).

Station	Location	Formation (Vis=1000m)		Vis=500m		Vis=200m	
		Time	Wind	Time	Wind	Time	Wind
DF	120.48°E,33.20°N	19:45	1.3m/s, E	22:55	1.2m/s, E	23:45	1.3m/s, E
BY	119.30°E,33.23°N	01:25	1.2m/s, ESE	03:15	1.4m/s, ESE	03:45	1.3m/s, SE
SH	118.22°E,33.48°N	04:50	1.6m/s, ESE	06:10	1.3m/s, ESE	07:15	2.4m/s, ESE
Distance (km)		Time difference (h)		Time difference (h)		Time difference (h)	
DF-BY	110	4.7		4.3		4.0	
BY-SH	105	3.4		2.9		3.5	

471

472


473



475

476 Figure 1. The parent and nest model domain. The shaded color is terrain height. The red points are automatic weather stations  
 477 in Jiangsu, China. The three larger circle points are Sihong (SH), Baoying (BY), and Dafeng (DF) stations, and the square point  
 478 is Sheyang (SY) sounding station. The black labels are some province or city names. (JS:Jiangsu Province; AH:Anhui Province;  
 479 YC:Yanchen; NT:Nantong).

480




481

482 Figure 2. The spatial evolution of fog. The black dots are simulated fog areas. The shaded colors are satellite observed bright-  
 483 ness temperature difference ( $3.9\mu\text{m}$  minus  $11.2\mu\text{m}$ ), where the blue colors (smaller than -2 K) indicate the fog areas.

484

485




486

487 Figure 3. The synoptic background of 500hpa (first row), 850hpa (second row) and surface (third row) at 08:00 and 20:00 on  
 488 20 January 2020.

489






491


492 Figure 4. The colored curves are the fog boundaries (satellite retrievals) from 23:00 on 20 January to 07:00 next day every 2  
 493 hours. The gray straight line indicates the fog propagation direction, and the vertical features of meteorologies at this line will be  
 494 analyzed in Figures 7, 8, and 9. The lower-left polar plot is the fog propagation speed at 16 directions (22.5° interval), and the  
 495 narrow blue bar highlights the maximum propagation speed (9.6m/s) occurring at 160° direction (in Cartesian coordinate system)  
 496 (from southeast to northwest).

497



498


499 Figure 5. The temporal variation of ground visibility (Vis; black line), wind speed (Ws; green line), wind direction (vectors),  
 500 temperature (Tem; red line), dew point (Td; violet line) and relative humidity (RH; blue line) at Dafeng, Baoying, and Sihong  
 501 stations. The horizontal dashed lines are visibilities of 1000m and 200m. The vertical dashed lines mark the times of fog forma-  
 502 tion and visibility burst dropping.  
 503



505

506 Figure 6. (a) The model performance on 2m Temperature (Tem), 2m Relative humidity (RH) and 10m wind speed and direction.  
 507 The red color is simulation and black color is observation. The time is from 14:00 on 20 January 2020 to 11:00 next day. (b) The  
 508 model performance on temperature (red), RH (blue) and wind (barbs) profiles at Sheyang sounding station. For temperature and  
 509 RH, the observations are scatters and simulations are solid lines. For wind barbs, the left column is observations and the right  
 510 column is simulations. The scatters and barbs are interpolated onto 0~600m every 100m.


511



512


513 Figure 7. (a) The height-longitude variation of horizontal wind direction (vectors) and wind speed (shaded colors) at the cross-  
 514 ing line in Figure 4. The lower-right black polygons are the fog area. The times are from 18:00 on 20 January to 04:00 next day.  
 515 (b) The averaged wind speed profile at the crossing line during 23:00~07:00. The two red points are the wind speed calculated  
 516 from ERA5 reanalysis. (c) The averaged wind direction (vectors) and wind speed (shaded colors) at 1000hpa during  
 517 23:00~07:00.

518



519


520 Figure 8. The height-longitude distribution of water vapor mixing ratio (g/kg) at the crossing line in Figure 4. The deep black  
 521 polygons are the fog area. The light black lines are the region of water vapor mixing ratio larger than 4g/kg. The times are from  
 522 20:00 on 20 January to 06:00 next day.  
 523



524

525 Figure 9. Same as the previous figure, but for the temperature. The bold black polygons are the fog area. The thin black lines  
 526 are the top of inversion layer.

527




528

529 Figure 10. (a) The height (shaded color) at which fog/low stratus firstly forms. The black contours are the ground fog areas at  
 530 00:00 on 21 January 2020. The colorbar represents the model level and the corresponding height above surface. For example, the  
 531 cyan colors indicate that fog firstly forms at the surface level with the corresponding height of about 4m. The red colors indicate  
 532 that low stratus firstly forms at the 5<sup>th</sup> to 7<sup>th</sup> model level with the corresponding height of about 36~66m. (b) The time differences  
 533 between ground fog formation and low stratus formation. For example, the cyan colors indicate that fog firstly forms at ground.  
 534 The blue colors indicate that the ground fog forms 0~10min later than the low stratus formation.

535


536



538


539 Figure 11. The temporal variation of ground fog area under different experiments from 00:00 to 06:00 on 21 January. The black  
 540 color is the base experiment. The Tadv0 (red), QvAdv0 (green) and QcAdv0 (blue) are the experiments turning off temperature  
 541 advection, moisture advection and cloud water advection, respectively. The NoAdv (pink) is the experiment turning off all of the  
 542 above advections.

543



545

546 Figure 12. (a) The spatial distribution of the four process tendencies contributing to LWC variation at ground level. (b) The  
 547 vertical profiles of the process tendencies averaged in fog area. The times are from 00:00 to 06:00 on 21 January.  
 548 (Cond:condensation or evaporation; Sedi:sedimentation; Vmix:turbulent exchange; Advc:horizontal and vertical advection).  
 549



551

552 Figure 13. The concept diagram of fog propagation. The ground wind speed (short orange arrows) is generally less than 3m/s.  
 553 A southeasterly BLLJ exists at the height from 50 to 500m, and the jet core intensity is 10m/s at 200m (the long orange arrow).  
 554 The updraft arrows represent the warm and wet air from ocean. The two cloud shapes are fog areas at two adjacent times, and the  
 555 white arrow indicates the fog propagation speed (9.6m/s). The fog propagation is probably caused by three approaches: 1) Moisture  
 556 advection from ocean promotes vapor condensation in the downstream area, which could be the dominant cause (the blue  
 557 fancy arrow); 2) Warm advection from ocean deepens inversion layer and additionally promotes vapor accumulation within PBL  
 558 (the red fancy arrow); 3) The moisture advection probably result in the low stratus formation, and later it subsides to ground by  
 559 turbulent mixing of cloud droplets (the blue water drops and circular arrows). Note that warm and moisture advections occur at  
 560 nearly all heights below 500m, not merely at the height indicated by arrows.