

# Extreme melting at Greenland's largest floating ice tongue

Ole Zeising<sup>1,2</sup>, Niklas Neckel<sup>1</sup>, Nils Dörr<sup>3,4</sup>, Veit Helm<sup>1</sup>, Daniel Steinhage<sup>1</sup>, Ralph Timmermann<sup>1</sup>, and Angelika Humbert<sup>1,2</sup>

<sup>1</sup>Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

<sup>2</sup>Department of Geosciences, University of Bremen, Bremen, Germany

<sup>3</sup>Institute of Geosciences, Kiel University, Kiel, Germany

<sup>4</sup>Institute of Photogrammetry and Remote Sensing, Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

**Correspondence:** Ole Zeising (ole.zeising@awi.de)

**Abstract.** The 79° North Glacier (Nioghalvfjærdsbrae, 79NG) is one of three remaining glaciers with a floating tongue in Greenland. Although the glacier was considered exceptionally stable in the past, earlier studies indicate that the ice tongue has thinned in recent decades. By conducting high-resolution ground-based and airborne radar measurements in conjunction with satellite remote sensing observations, we find significant changes in the geometry of 79NG. In the vicinity of the grounding line, a 500 m high subglacial channel has grown since ~2010 and caused surface lowering of up to  $7.6 \text{ m a}^{-1}$ . Our results show extreme basal melt rates exceeding  $150 \text{ m a}^{-1}$  over a period of 17 d within a distance of 5 km from the grounding line, where the ice has thinned by ~~42%~~ 32% since 1998. We found a heterogeneous distribution of melt rates likely due to variability in water column thickness and channelization of the ice base. Time series of melt rates show a decrease in basal melting since 2018, indicating an inflow of colder water into the cavity below 79NG. We discuss the processes that have led to the changes in geometry and conclude that the inflow of warm ocean currents has led to the extensive thinning of 79NG's floating ice tongue near the grounding line in the last two decades. In contrast, we hypothesize that the growth of the channel results from increased subglacial discharge due to a considerably enlarged area of summer surface melt due to the warming of the atmosphere.

## 1 Introduction

The mass loss of the Greenland Ice Sheet over the last decades as a result of a warming atmosphere and ocean has accelerated (Shepherd et al., 2020) and contributed to recent sea-level rise by  $1.4 \text{ mm a}^{-1}$  (Khan et al., 2022a). Half of the mass loss is caused by ice-sheet discharge through marine-terminating glaciers (Shepherd et al., 2020), mainly due to the retreat of glacier fronts (King et al., 2020) as the floating ice tongues restrain the outflow of the grounded ice (Fürst et al., 2016). The largest of the three remaining floating tongues in Greenland is the one of Nioghalvfjærdsbrae (79NG). Together with its neighboring Zachariæ Isstrøm (ZI), it is the main outlet glacier of the Northeast Greenland Ice Stream (NEGIS; Fig. 1a), the largest ice stream of the Greenland Ice Sheet (Fahnestock et al., 2001). After the collapse of ZI's floating tongue in 2002, the glacier itself (Khan et al., 2014; Mouginot et al., 2015), as well as NEGIS, have shown an extensive speed-up (Khan et al., 2022b). In contrast, only minor acceleration rates have been observed at 79NG (Mouginot et al., 2015; Vijay et al., 2019).

Ice sheet simulations indicate that 79NG remains stable within this century and will experience only a minor grounding line retreat as bedrock rises inland (Choi et al., 2017). Its stability is attributed to pinning points at the calving front (Thomsen et al., 1997), lateral resistance from shear margins (Mayer et al., 2000; Rathmann et al., 2017; Mayer et al., 2018) and confinement of the glacier leading to lateral compression. However, thinning has occurred during the last two decades (Helm et al., 2014; Kjeldsen et al., 2015; Mouginot et al., 2015; Mayer et al., 2018) and cracks have formed at the calving front that might be a precursor of disintegration (~~(?)~~(Humbert et al., 2023)).

Observations and modeling show that the inflow of warm Atlantic Intermediate Water (AIW, temperatures exceeding 1 °C) into the cavity below 79NG (Straneo et al., 2012; Wilson and Straneo, 2015; Lindeman et al., 2020; Schaffer et al., 2020) and its variability are connected to the ocean currents in Fram Strait (Münchow et al., 2020; von Albedyll et al., 2021). ~~Bentley et al. (2023) gives evidence~~ Based on temperature and salinity measurements in an epishelf lake, Bentley et al. (2023) show that the AIW reaches the grounding line area of 79NG a few months after ~~the inflow into the~~ entering the cavity.

The observed oceanic heat transport into the sub-ice cavity (Schaffer et al., 2020) has been suggested to maintain intense basal melting (~~Mayer et al., 2018; Lindeman et al., 2020; Schaffer et al., 2020~~)(Mayer et al., 2018; Lindeman et al., 2020; Schaffer et al., 2020). Similar processes might have led to the disintegration and retreat of the floating ice tongues of Jakobshavn Isbræ (Motyka et al., 2011). In the future, basal melt rates are expected to increase most pronouncedly in the northeastern part of Greenland towards the end of the 21st century (Slater et al., 2020).

However, the supply of fresh water from ~~glacier surface melting was~~ glacial surface melting has been found to alter circulation in fjords and basal melting of glaciers (~~Straneo et al., 2016~~)by increasing buoyancy-driven circulation and decreasing shelf-forced circulation (Straneo et al., 2016). Subglacial water discharge from beneath the grounded ice is often linked to the location of basal channels in the floating ice shelves caused by locally enhanced melting (Le Brocq et al., 2013). Such channels can be up to a few kilometers in width and up to a few hundred meters in height (Rignot and Steffen, 2008).

The spatial distribution of basal melt rates can be investigated using repeated measurements with the phase-sensitive Radio Echo Sounder (pRES). The same device can be operated in an autonomous mode (henceforth ApRES) to perform measurements over a longer period of time with a defined interval. Previous studies used pRES and ApRES measurements to investigate the spatial distribution and temporal variability of basal melting inside basal channels: At the Ross Ice Shelf, Antarctica, Marsh et al. (2016) found enhanced melting inside a channel near the grounding line which reduced in the downstream direction. Humbert et al. (2022) revealed for a channel at Filchner Ice Shelf, Antarctica that melt rates inside the channel decrease in the direction of ice flow and fall below those outside the channel, causing the channel height to decrease. While they found no pronounced seasonality of melting inside the channel, Washam et al. (2019) detected a significant increase in melting inside a channel at Petermann Glacier, Greenland, during the surface melt period in summer. They linked the seasonality to the increased subglacial discharge that enhanced the inflow of warmer ocean currents into the cavity (Shroyer et al., 2017; Washam et al., 2019). Whether basal channels stabilize or weaken shelf ice is not fully understood yet (Alley et al., 2016). Numerical models indicate that the existence of channels can decrease the mean basal melt rate (Millgate et al., 2013), at the same time, the channels can structurally weaken the ice shelf (Vaughan et al., 2012). ~~In the future, submarine melt rates are expected to increase most pronouncedly in the northeastern part of Greenland towards the end of the 21st century (Slater et al., 2020)~~.

Observations of basal melt rates and their influence on the ice thickness are ~~thus~~ considered key to understanding the dynamics of the system. Basal melt rates of 79NG have been estimated based on indirect satellite remote sensing retrievals (Wilson et al., 2017), which are accompanied by considerable uncertainties ~~and limited to freely floating parts. Especially in the~~. Particularly within a few kilometers from the grounding line and above basal channels, the ice is in hydrostatic imbalance, limiting the analysis of melt rates based on changes in surface elevation (Chartrand and Howat, 2023). Thus, other methods must be used to monitor changes in ice thickness and to understand the underlying processes, especially in the area of the grounding line of 79NG where higher basal melt rates are expected due to thick (reduced melting temperature) ice getting into contact with warm ocean waters, ~~other methods must be used to monitor changes in ice thickness and to understand the underlying processes.~~

In this study, we investigate the recent changes in ice thickness of the 79NG from in-situ and airborne as well as satellite remote sensing observations. We analyze a spatial distribution of thinning and basal melt rates focusing on the vicinity of the grounding line of 79NG. Finally, we discuss the processes that explain the observations and how these have changed in the past decades.

## 2 Data

In order to obtain a time series of surface elevations of the 79NG grounding line area we generated 96 Digital Elevation Models (DEMs) from bistatic TanDEM-X SAR interferometry, which span the period December 2010 to April 2021. Additionally, we acquired airborne and ground-based radar measurements at the 79NG under the framework of the *Greenland Ice Sheet – Ocean Interaction* (GROCE, <https://www.groce.de>) project. The airborne radar measurements were performed in April 2018 and July 2021 with ~~the~~ AWI's ultra-wideband (UWB; Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, AWI) radar in order to determine the basal geometry of the 79NG. We obtained a spatial distribution of Lagrangian thinning rates from a repeat survey of ~~ground-based phase-sensitive radar (pRES)~~ pRES measurements in July 2017 and 2018. In July 2017, we marked the measurement location on the surface ~~by~~ with 4 m long bamboo stakes drilled into the ice in order to be able to repeat the measurement in 2018 at exactly the same location in the Lagrangian frame. The majority of the measurement locations were distributed within 8 km distance from the grounding line (Fig. 1b). Here, we operated ~~autonomous pRES~~ ApRES stations at three locations (ApRES1-3) until ~~June 2022~~ September 2023 that move with the ice to derive year-round time series of basal melt rates in a Lagrangian reference frame. In Summer 2018, we relocated ApRES2 to its starting position from 2016 in order to repeat the measurements on the same flowline. These stations are labeled as ApRES2a (2016–2018) and ApRES2b (2018–2019).

### 3 Methods

#### 3.1 Time series of surface elevations from TanDEM-X SAR interferometry

DEMs were generated from bistatic TanDEM-X SAR interferometry closely following the methods described by Neckel et al. (2013). Interferograms were formed from co-registered Single-look Slant range Complex (CoSSC) data employing a 4×4 multi-looking step. Prior to phase unwrapping we subtracted a simulated phase from the global TanDEM-X DEM at ~~30 m~~ 30 m resolution (Wessel et al., 2016). The latter was done to reduce unwrapping errors and the simulated phase was added back afterwards. The final DEMs were geocoded and spatially adjusted to the global TanDEM-X DEM ~~over stable bedrock following the methods described by Nuth and Kääb (2011),~~ by calculating the standard deviation and the normalized median absolute deviation (NMAD) for all DEMs over stable ground (Nuth and Kääb, 2011; Wessel et al., 2016). The NMAD is considered to be more robust to outliers than the standard deviation (Höhle and Höhle, 2009). For the entire stack of DEMs we obtain a standard deviation of 0.96 m and an NMAD of 0.55 m which is in agreement with the TanDEM-X mission requirements (Wessel et al., 2018). Surface elevation changes between 2010 and 2021 were estimated by fitting a linear trend to every pixel of the co-registered stack of 96 DEMs (e.g. Berthier et al., 2016).

#### 3.2 Ultra-wideband (UWB) airborne radar

The UWB is a multichannel coherent airborne radar that consists of an eight-element antenna array with a total transmit power of 6 kW (Hale et al., 2016). The antennas operate in the frequency band of 150 – 520 MHz, with a pulse repetition frequency of 10 kHz and a sampling frequency of 1.6 GHz. The characteristics of the transmitted waveform and the recording settings can be manually adjusted. We used alternating sequences of different transmission/recording settings (waveforms) to increase the dynamic range: short pulses (1  $\mu$ s) and low receiver gain (11 – 13 dB) to image the glacier surface, and longer pulses (3 – 10  $\mu$ s) with higher receiver gain (48 dB) to image internal features and the ice base. The waveforms were defined with regard to the glacier thickness. Additionally, we used two different frequency bands in the survey: 180 – 210 MHz and 150 – 520 MHz. The theoretical range resolution in ice after ~~pulse-compression-pulse compression~~ pulse compression for the two bandwidths is about 2.8 m and 0.23 m, respectively. Recorded traces were ~~presumed-pre-stacked~~ pre-stacked in the hardware by a factor between 2 and 16, depending on the pulse length. In order to reduce range side lobes, the transmitted and the received signals were tapered using a Tukey window and the received signal spectrum was filtered with a Hanning window (The MathWorks Inc., 2022). We recorded the position of the aircraft with four NovAtel GPS receivers, which were mounted on the wings and the fuselage.

Post-flight processing included pulse compression in the range direction, synthetic aperture radar (SAR) focusing in the along-track direction, and array processing ~~in the cross-track direction to~~ to increase signal-to-noise ratio and to suppress off-nadir echoes. The SAR focus is set to achieve a ground resolution of 10 m in the along-track direction.

To transform two-way travel time to depth, we used a propagation velocity for the electromagnetic wave of  $168.914 \text{ m } \mu\text{s}^{-1}$  ~~that~~ which refers to a relative permittivity of  $\epsilon_r = 3.15$  for pure ice. No firn correction was applied since the predominant part of the glacier is located in the ablation zone. We concatenated the echograms of the alternating waveforms to obtain the final echograms covering the glacier from the surface to the base with a high dynamic range. Finally, the surface return of the radar

120 echo was aligned with a high-resolution elevation model created with a vertical accuracy of 0.1 m, which was determined from simultaneously acquired laser scanner data.

### 3.3 Phase-sensitive Radio Echo Sounder (pRES)

#### 3.3.1 Technical Background

The phase-sensitive radio echo sounder (pRES) pRES is a ground-penetrating frequency-modulated continuous-wave (FMCW) radar, transmitting chirps with a frequency bandwidth of 200 MHz and a center frequency of 300 MHz (Brennan et al., 2014; Nicholls et al., 2015). While the repeat pRES survey was performed with 100 chirps and two skeleton slot antennas, the ApRES stations consist ~~consisted~~ of two bow-tie antennas and recorded 20 chirps with a measuring interval between one and six hours. For processing the raw data, we calculated pairwise correlation coefficients of all chirps, rejected chirps with low correlation coefficients, and stacked the remaining ones. We followed Brennan et al. (2014) for processing to get amplitude- and phase-depth profiles. We assumed a relative permittivity of  $\epsilon_r = 3.15$  in ice for the time-to-depth conversion.

#### 3.3.2 Thinning rates from single-repeated pRES measurements

The estimation of the Lagrangian thinning rate is based on the change in ice thickness along the flow of the same ice particles. The ice base is assumed to be responsible for strong peaks in the radar signal due to the high contrast in relative permittivity between ice and seawater. In the case of a flat ice base, the nadir reflection has the shortest two-way travel time of all basal reflections in a radius defined by the antenna beamwidth. However, steep basal gradients such as those of basal channels can cause off-nadir reflections which might appear before the nadir basal return. The steeper the basal gradient between the nadir ice base and the location of the off-nadir reflection increases on average, the earlier the off-nadir reflection occurs. Thus, if an off-nadir reflection appears before the nadir basal return depends on the location of the measurement relative to the surrounding basal slopes and their gradients.

In order to identify nadir and off-nadir returns, we used the ice base—ice surface—ice base multiple which we assume to be the first multiple reflections from the ice base, which were characterized by twice the two-way travel time since they originated from the reflections at the ice base, the ice surface, and again at the ice base. Here we assume that the multiple is strongest for the nadir reflections. The reflection since, in the case of a flat ice surface, most of the reflected energy from a far-off-nadir reflection will be mostly is reflected in the opposite direction in the case of a flat ice surface.

Therefore, multiples from off-nadir reflections will be weaker compared to nadir reflections. Additionally, we used the ice thickness distribution derived from UWB echograms nearby near the location of the pRES observations that can reveal the ice thickness and, furthermore, give a hint for the origin of the recorded off-nadir reflection. At pRES location g4\_p02, UWB echograms from 2018 show a growing subglacial channel in the immediate vicinity of both the first and repeated pRES measurement (Figs. 1c, 2). Based on these UWB echograms, we link the origin of the off-nadir reflection to the basal channel.

In Appendix A, we give further examples of amplitude profiles from repeated pRES measurements where the first basal return was identified to be an off-nadir return. The distinction between nadir and off-nadir returns is important as the precise local change in ice thickness can only be revealed from nadir returns. Following Stewart et al. (2019), we applied a cross-correlation of the amplitude and the phase of their basal segments ranging from  $-9$  to  $+1$  m around the identified return. The uncertainty in this case is below  $0.01$  m.

Even if no distinction between nadir and off-nadir reflection can be made, the ice thickness change can be estimated with the following approach: Using the first basal return in both measurements would always result in an underestimation of the change in ice thickness at least at one of both locations where the (off-nadir) first basal reflections occurred (Appendix A). This means that a thinning rate exists somewhere in the scattering area that is as large as this amount or even more. Note, this should not be understood as the minimum rate in the scattered area (marked in Fig. 1 for off-nadir thinning rates), as there can also be lower thinning rates at the same time. Thus, at stations where we could not distinguish reliably between a nadir and an off-nadir reflection, we used the first strong increase in amplitude for the ice thickness calculation and interpret this as an off-nadir return. The range to the off-nadir basal reflector differs from the ice thickness above the reflector, which can be derived by  $H^\alpha = R \cos \alpha$ , where  $R$  is the range of the basal reflector and  $H^\alpha$  the off-nadir ice thickness viewed at an angle  $\alpha$ . The resulting thinning rate  $\Delta \dot{H}^\alpha$  (positive values correspond to thinning) is

$$\Delta \dot{H}^\alpha = -\frac{(R_2 - R_1) \cos \alpha}{\Delta t}, \quad (1)$$

where  $R_1$  is the range to the off-nadir basal reflector of the first and  $R_2$  of the second measurement after the time period  $\Delta t$ . Since the off-nadir angle  $\alpha$  is often unknown, we assume that it ranges from  $0^\circ$  to a maximum of  $30^\circ$  (Brennan et al., 2014) and calculated the average of both angles. The spread in ice thickness difference from both  $\alpha$  together with the inaccuracy of the signal propagation speed in the ice of  $\sim 1\%$  (Fujita et al., 2000) is represented in the uncertainty of  $\Delta \dot{H}^\alpha$ . At those stations at which we identified nadir and off-nadir reflections, we determined both thinning rates. We rejected those measurements where the depth of the first basal return was unclear. ~~Since the~~ The estimation of vertical strain was not possible with single-repeated pRES measurements due to the low correlation of the amplitude profiles. We attribute this to water-saturated layers on the surface whose reflections also overlap those of deeper layers. Thus, we could not calculate the basal melt rate.

### 175 3.3.3 Basal melt rates from ApRES time series

The calculation of basal melt rates follows previously described methods (Corr et al., 2002; Jenkins et al., 2006; Stewart et al., 2019). Several quantities cause changes to the range  $R$  to the basal reflector within the time period  $\Delta t$ : ablation  $\Delta R_s$ , strain  $\Delta R_\varepsilon$  and basal melting  $\Delta R_b$ :

$$\frac{\Delta R}{\Delta t} = \frac{\Delta R_s}{\Delta t} + \frac{\Delta R_\varepsilon}{\Delta t} + \frac{\Delta R_b}{\Delta t} \quad (2)$$

180 With the ApRES time series (Fig. 3a, Appendix B), all of these quantities can be estimated in order to obtain the basal melt rate. Since the estimation is based on the detection of the vertical displacement of layers, we divided the first echogram in  $6$  m long segments with  $5$  m overlap starting at a depth of  $20$  m. For each segment, we derived displacements from complex cross-

correlation of the phase of all pairwise time-consecutive measurements (Stewart, 2018). ~~Afterward~~Afterwards, we calculated the daily mean values of the displacements.

185 In the first step, we used the time-mean vertical displacement of internal reflectors to calculate the vertical strain profile ~~:~~ (Fig. 3b). Here only those segments between 20m below the surface and 20m above the basal return at the last measurement were considered. In addition, we only considered measurements between October and May to avoid the influence of ablation on the calculation of the strain. The vertical strain is the depth derivative of the vertical displacement  $u_z$

$$\varepsilon_{zz} = \frac{\partial u_z}{\partial z}, \quad (3)$$

190 which we derived from a linear fit that best matches the vertical displacements. Although one of the ApRES stations was located within the hinge zone in which bending might affect the strain distribution (Jenkins et al., 2006), none of the displacement distributions indicated a deviation from a linear function over depth.

For a nadir basal reflection, the estimation of the range shift due to ice deformation  $\Delta R_\varepsilon^n$  is only affected by the vertical strain

$$195 \Delta R_\varepsilon^n = \int_{\underline{00m}}^R \varepsilon_{zz} dz, \quad (4)$$

from the surface at  $z = 0m$  to the ice base at  $z = R$ . Calculating the displacement of off-nadir reflectors due to deformation is more complex because the two horizontal strain components  $\varepsilon_{xx}$  and  $\varepsilon_{yy}$  must also be considered (see Appendix B1). Since we can only determine the vertical strain component  $\varepsilon_{zz}$  with ApRES measurements, we have to make assumptions to estimate  $\Delta R_\varepsilon$  viewed at an angle  $\alpha$ . In Appendix B1, we show that for small off-nadir angles of  $\alpha \leq 30^\circ$ , the absolute value of  $\Delta R_\varepsilon$  is always smaller than or equal to the absolute value of the nadir displacement  $\Delta R_\varepsilon^n$ :

$$0 \leq |\Delta R_\varepsilon| \leq |\Delta R_\varepsilon^n|, \quad (5)$$

At all ApRES sites, we found  $\varepsilon_{zz} > 0$  (see Figure 3b and Appendix B2), so that Equation 5 can be simplified further to

$$0 < \Delta R_\varepsilon \leq \Delta R_\varepsilon^n. \quad (6)$$

Thus, assuming that the reflection occurred from a nadir reflector, we cannot underestimate the deformation. The largest  $\Delta R_\varepsilon^n$  was found to be 2.7m for  $\Delta t = 1a$  at ApRES2b, ~~which results in an underestimated melt rate of  $\leq 2.7 \text{ m a}^{-1}$  due to deformation.~~ In case the change in ice thickness is based on an off-nadir basal reflection, the correction with the nadir range shift due to ice deformation underestimates the melt rate by  $\leq 2.7 \text{ m a}^{-1}$ .

Next, we use the displacement time series of the segment centered at a range of 50m ( $u_z^{50}$ ) to correct for ablation ~~:~~ (Fig. 3c). Since the ice above is affected by ice deformation, we subtract this contribution from the displacement

$$210 \Delta R_s^n = u_z^{50} - \int_{\underline{00m}}^{5050m} \varepsilon_{zz} dz. \quad (7)$$

Here,  $\Delta R_s^n$  is the vertical (nadir) ~~ablation~~displacement caused by ablation (negative for ablation). In order to determine the contribution of the ablation to the range difference to an off-nadir reflector,  $\Delta R_s^n$ , we need to correct for the angle  $\alpha$ :

$$\Delta R_s = \frac{\Delta R_s^n}{\cos \alpha}. \quad (8)$$

Since  $\alpha$  is still unknown, we use the extremes  $0^\circ$  and  $30^\circ$  and average both values. The difference from the mean is used as  
215 uncertainty, which corresponds up to  $2 \text{ m a}^{-1}$  in summer and near zero in winter. We removed outliers defined by ablation rates  $> 0.1 \text{ m d}^{-1}$ .


To finally derive the basal melt rate  $a_b$  in the vertical direction, we subtract  $\Delta R_\varepsilon$  and  $\Delta R_s$  from the displacement of a basal reflector

$$a_b = -\frac{\Delta R_b}{\Delta t} = -\frac{\Delta R - \Delta R_s - \Delta R_\varepsilon}{\Delta t}. \quad (9)$$


220 Similar to ~~(Vaňková et al., 2021)~~ Vaňková et al. (2021), we analyze  $\Delta R$  for all segments within a range of 50 m below the first basal return to obtain the nadir and off-nadir basal melt rates ~~-(Fig. 3d)~~. This range was chosen since at the ApRES sites, all strong basal reflections occurred within 50 m. To represent the variability within a time series, we calculated the median melt rate next to the 25%, 75%, and 95% quantile for each time step. ~~Afterward~~ Afterwards, a 7-day moving average filter was used to smooth the time series ~~-(Fig. 3e)~~.

225 The largest uncertainty of the melt rate estimate arises from the unknown off-nadir angle  $\alpha$ , which affects the ablation and strain correction. The sum of both uncertainties is up to  $8 \text{ m a}^{-1}$  in summer and significantly less in winter. In addition, the estimate of the melt rate quantifies the rate at which the ice base has approached the ApRES through melting. This can differ from the melt rate in the normal or vertical direction at the basal reflector. ~~A~~ In the case of a flat ice base, the measured nadir melt rate is equal to the melt rate in the normal and vertical direction. For an inclined ice base, the measured nadir melt rate is equal to the melt rate in the vertical direction, which is different from that in the normal direction. A measured off-nadir melt rate can differ from both the melt rate in the normal and in the vertical direction. A further uncertainty arises from the inaccuracy of the signal propagation speed in the ice ~~, which corresponds to  $\sim 1\%$  resulting in an inaccuracy~~ of the melt rate of  $\sim 1\%$  (Fujita et al., 2000).  
230






**Figure 1.** (a) Map of northern Greenland with drainage basins (black lines) of 79NG and Zachariæ Isstrøm (ZI) (Krieger et al., 2020) and surface velocities (Joughin et al., 2018) showing the Northeast Greenland Ice Stream (NEGIS) and Petermann Glacier (PG). (b) Sentinel-2 mosaic of 79NG with thinning rates derived from *in-situ* pRES measurements in 2017 and 2018 (box in (a)). (c) Enlargement of the 79NG hinge zone (box in (b)) with surface elevation change rates ( $dh/dt$ ) derived from TanDEM-X satellite data between 2010 and 2021. Dots and squares scattering areas showing nadir and off-nadir thinning rates with paths of Lagrangian pRES measurement location between July 2017 and July 2018. White stars mark the starting and black stars the ending position of ApRES stations. The upper flexure limit (grounding line) and lower flexure limit are based on interferometry and mark the area where the ice is being bent by the tides. Copernicus Sentinel data from 2018, retrieved from Copernicus SciHub on 16 August 2021.



**Figure 2.** Growing basal channel from pRES and UWB echograms. (a,b) UWB echograms from the across-flow profiles from 2018. The center of both is the location of a Lagrangian pRES measurement in 2017 ((a), vertical blue line) and 2018 ((b), vertical red line). Possible origins of nadir and off-nadir reflections, discovered in the pRES echograms (c), are represented by dashed lines. The suggested locations at which the reflections occurred are marked by circles. (c) pRES echograms from 2017 and 2018 with the identified nadir and off-nadir reflections. Location is shown in Fig. 1c.



**Figure 3.** Analysis of ApRES1 time series. (a) Time-echogram of a Lagrangian measurement at ApRES1 recorded between August 2016 and June 2022. In 2016 and 2017, several ApRES malfunctions caused data gaps. The black outline marks the first 50 m below the basal return. (b) Mean vertical displacement of englacial segments (dots). The gray shaded area marks the range between the 25% and 75% quantile. Segments between 20 m and 20 m above the first basal return at the end of the measurement period (red dots) were used to calculate the change in ice thickness due to vertical strain by fitting a linear function (black line). (c) Time series of ablation rate (negative for ablation). The grey shaded area marks the uncertainty due to the off-nadir correction. (d) Time series of the determined melt rate (color) within the first 50 m below the basal return, corresponding to the area marked by black lines in (a). (e) Time series of basal melt rate. The dashed line shows the 95% quantile, the solid line the median, and the shaded area marks the range between the 25% and 75% quantile.


## 4 Results

### 235 4.1 Growing subglacial channel causes local surface lowering


The DEM time series reveals that the surface elevation of the 79NG has decreased along the grounding line by a rate of  $-2.0 \pm 1.4 \text{ m a}^{-1}$  (mean value  $\pm$  standard deviation), corresponding to a surface lowering of  $-20 \pm 14 \text{ m}$  between December 2010 and April 2021 (Fig. 4a). The maximum surface-lowering of  $-56.9 \pm 0.1 \text{ m}$  (or  $-5.5 \pm 0.1 \text{ m a}^{-1}$ ) of the same time period is evident in a graben-like structure in the center of the grounding line (Fig. 4a). This remarkable area of enhanced lowering is located downstream of a supraglacial lake and extends from 4 km upstream to 4 km downstream of the grounding line. Its width decreases in flow direction from a maximum of 1 km roughly 2 km upstream from the grounding line to 500 m within 5 km in the ice flow direction (Fig. 1c). While a hill was present in the ice surface in the central part near to and upstream of the grounding line until 2015, this turned into a depression (Fig. 4b,c) due to enhanced surface lowering rates compared to those rates outside the graben-like structure (Fig. 4e,f). The average elevation change upstream of the grounding line was  $-2.1 \pm 0.1 \text{ m a}^{-1}$  between December 2010 and April 2015 and has increased to  $-6.6 \pm 0.1 \text{ m a}^{-1}$  until the end of the time series in April 2021 (Fig. 4b,e). Outside this area, the surface lowered at a significantly smaller rate of  $-1.1 \pm 0.1 \text{ m a}^{-1}$ . ~~Further downstream~~ Five kilometers downstream of the grounding line behind the lower flexure limit where the ice is freely floating (Fig. 1d), this sink already existed in 2010 and the lowering rate was smaller (Fig. 4d,g). At both locations downstream from the grounding line, the (Eulerian) surface elevation change rate suddenly changed in late 2019 and became less strong (Fig. 4f) and even turned into thickening (Fig. 4g), similar to what we have observed until 2013.

In order to investigate what causes this drop in surface elevation, we recorded flight profiles with the UWB airborne radar. Near the grounding line, these airborne radargrams reveal the existence of several subglacial channels (Figs Fig. 5 and Appendix Figs. C1 and C2). The by far largest channel with a height of 500 m and a width of 1 km is found in the central flow line near the grounding line and extends 5 km upstream from the grounding line. Above this channel, only 190 m of ice is left, which is 30% of the surrounding ice thickness. The location of this channel is in good agreement with the lowering of the surface observed from TanDEM-X satellite data (Fig. 4). However, upstream from the grounding line, the tip of the basal channel is located up to 400 m in the northwestern direction from the center of the surface depression. Between 2018 and 2021, the channel has grown, especially in the upstream area, where the channel height has increased by almost 200 m (Fig. 5a,b). In contrast, no significant change in ice thickness or channel height occurred downstream of the grounding line (Fig. 5c). ~~Seismic measurements~~ Airborne radar data from 1998 conducted by ~~Mayer et al. (2000)~~ Niels Reeh and Erik Lintz Christensen with the DTU (Technical University of Denmark) Space 60 MHz ice sounder show no channel near the grounding line and a small 120 m high channel, located 5 km downstream the grounding line (Fig. 5d).

The ice has also thinned considerably outside the central channel. Along an across ice flow profile 600 m downstream from the grounding line, the average ice thickness in 2021 was 38 m less than in 2018 (Fig. 5c). Compared to 1998, the ice thickness 5 km downstream from the grounding line has ~~even~~ decreased by more than ~~230 m or 42%, which corresponds to a rate of  $10 \text{ m a}^{-1}$ . The maximum thinning of 80% took place above the subglacial channel~~ 162 m or 32%. Above the large subglacial channel, the glacier thinned by 67%.



**Figure 4.** Surface elevation above sea level (EGM2008) from TanDEM-X satellite data between 2010 and 2021 (a) in ice flow direction and (b–d) across ice flow direction. The distance in (a) is relative to the grounding line and in (b–d) relative to the profile in (a). The location of all profiles is shown in Fig-Figure 1 and here marked by dashed lines. (e–f) Time series of surface elevation since 2010 at the three crossings above (black) and outside (gray) the graben-like structure. The numbers represent the gradient of the linear regression.



**Figure 5.** Sketch-ice thickness evolution of the 79NG ice thickness between 1998 and 2021 (a) in ice flow direction and (b–d) across ice flow. The sketch-bed topography in (a) and (d) is based on active seismic measurements by [\(Mayer et al., 2000\)](#) [Mayer et al. \(2000\)](#) from [1998 and UWB-1998](#). The ice geometry in (a) – (d) is based on airborne radar measurements from 1998 (Reeh), 2018 (UWB) and [2021](#), [/or 2021 \(UWB\)](#). The [IceBridge BedMachine Greenland, Version 5 \(Morlighem et al., 2017, 2022\)](#) ice geometry is shown for the two across ice flow sections (c) and (d) of the floating part. The distance in (a) is relative to the grounding line and in (b–d) relative to the profile in (a). The locations of all profiles are shown in [Fig-Figure 1](#) and [are](#) here marked by dashed lines. [Fig-Figure 4](#) shows the surface elevation change above these profiles. Appendix C shows the [UWB data](#) this sketch-figure is based on.


## 4.2 Extreme subglacial melting at the floating ice tongue

An analysis of the change in ice thickness at a given location (Eulerian perspective), as in the previous section, reveals changes in the geometry of the glacier. However, because the ice is flowing, considering the Lagrangian perspective [in addition to the Eulerian](#) is necessary for a full understanding of the process that causes these changes. The repeat UWB profile D–D' from July 2021 is the Lagrangian repeat of the profile C–C' from April [2018–2018 \(Fig. 1c\)](#). On average, the ice thickness at profile D–D' in 2021 was reduced by 193m, corresponding to a mean annual rate of  $59 \text{ m a}^{-1}$ . Since the surface ablation is typical  $< 2 \text{ m a}^{-1}$  (Zeising et al., 2020) and the dynamic thinning due to strain is small ([as shown by all ApRES measurements \(Fig. 3, Appendix B2\)](#)), most of this thinning is attributed to basal melting.

In order to investigate the spatial distribution of Lagrangian thinning, we analyzed pRES measurements performed in July 2017 and 2018 at the same surface point. Figure 1b shows the spatial distribution of the thinning rates of all repeated pRES measurements (colored dots), while these are separated in Figure 1c into nadir (colored dots with a line showing the flow path) and off-nadir thinning rates (colored area). The marker shape of the off-nadir thinning rates [in Figure 1c](#) corresponds to the scattering area from where the off-nadir basal reflections could have occurred. The thinning rates are between  $1.7 \pm 0.1$  and  $134 \pm 21 \text{ m a}^{-1}$  for locations spread over the entire ice tongue of 79NG (Fig. 6). The highest (off-nadir) thinning rates of  $126 \pm 20$  and  $134 \pm 21 \text{ m a}^{-1}$  were found at the most downstream bulge of the grounding line, next to the central subglacial channel where the ice draft is large. However, moderate thinning rates of  $< 21 \text{ m a}^{-1}$  were observed at a similar distance to the grounding line and for a similar draft (Fig. 6). Further downstream, but still within the hinge zone, we observed predominantly high thinning rates ( $> 50 \text{ m a}^{-1}$ ) spread across the entire width of the ice tongue. In general, thinning rates are observed to be below  $30 \text{ m a}^{-1}$  several kilometers downstream from the grounding line, declining towards the calving front to between  $1.7 \pm 0.1$  and  $3.2 \pm 0.1 \text{ m a}^{-1}$  (Figs. 1 and 6).

Variability on small spatial scales is accessible using a combination of nadir and off-nadir returns. At the pRES measurement location g4\_p02 (Fig. 2), where we link the origin of the off-nadir reflection to a small subglacial channel, we derived two estimates of thinning rates: One is based on the repeated nadir reflection outside the channel ( $6.4 \pm 0.1 \text{ m a}^{-1}$ ) while the other is based on both first basal (off-nadir) reflections in 2017 and 2018 within the channel ( $73 \pm 10 \text{ m a}^{-1}$ ). This comparison indicates a growth of the subglacial channel by more than  $66 \text{ m a}^{-1}$ .

The ApRES time series show a strong spatial and temporal variability of basal melt rates without a clear seasonal cycle. All three ApRES recorded high melt rates between October 2017 and July 2018 of  $> 50 \text{ m a}^{-1}$  on average, which reduced to  $\sim 30 \text{ m a}^{-1}$  until April 2019 and stayed low until the end of the record in [January 2020 \(ApRES2b\)](#), July 2022 ([ApRES1](#)), and [September 2023 \(ApRES4\), respectively](#) (Fig. 7). This change is particularly pronounced at ApRES1, which is located on the southeastern side of the glacier. Between April 2018 and April 2019, when ApRES1 was about 8 – 9km downstream from the grounding line, the melt rate dropped from  $137 \pm 2 \text{ m a}^{-1}$  (95% quantile) to just  $30 \pm 1 \text{ m a}^{-1}$  (Fig. 7a). After two periods with higher melt [in rates in the](#) summer and autumn [of 2020](#), the basal melt rate [reduces-reduced](#) to zero for almost all of the remaining 18 months. In early 2017, melt rates  $> 120 \text{ m a}^{-1}$  (95% quantile) were recorded 5.5km downstream from the grounding line at the north-western side by ApRES2a at the first basal return, whereas at the same time, the median




**Figure 6.** Distribution of the thinning rate as a function of distance from the grounding line. Color-coded draft of the floating tongue derived from pRES measurements and separated for nadir and off-nadir thinning rates. Most uncertainties are too small to visualize.

melt rate was below  $50 \text{ m a}^{-1}$  (Fig. 7b). After the relocation of ApRES2a (now named ApRES2b) to its starting point in the summer of 2018, ApRES2b recorded a 50 m lower ice thickness and a 50% lower melt rate (95% quantile) than ApRES2a two years before. Furthermore, the spatial variability (difference between the median and 95% quantile) of ApRES2b was greatly reduced.

The highest melt rates of  $150 - 168 \pm 5 \text{ m a}^{-1}$  lasting 17 d were recorded at ApRES3 at the beginning of the time series in July and August 2017. At that time, the ApRES3 was located 3 km from the grounding line next to the large central basal channel. After these high melt rates dropped to roughly  $50 \text{ m a}^{-1}$  after the summer in 2017, the basal melt rate showed, in general, a steady decrease to  $\sim 20 \text{ m a}^{-1}$  until the end of the time series in [July 2022](#). [September 2023](#).





**Figure 7.** Basal melt rate time series of all ApRES measurements: (a) ApRES1, (b) ApRES2a and ApRES2b, (c) ApRES3. The dashed line shows the 95% quantile, the solid line the median, and the shaded area marks the range between the 25% and 75% quantile. The dotted line in (b) represents the mean melt rate of ApRES2b shifted in time for a comparison with ApRES2a from an Eulerian perspective.

An analysis of the change in 79NG's geometry between 1998 (~~Mayer et al., 2000~~) (~~Reeh's airborne radar~~) and 2021 (this study) reveals a thinning by ~~42%~~ 32% in a narrow region 5 km from the grounding line and an ice base that became channelized, especially in the vicinity of the grounding line (Fig. ~~5d~~ 5). Compared to the 30% thinning observed by Mouginit et al. (2015) for the period 1999 to 2014, the thinning has continued without accelerating. The onset of steep basal slopes has been shifted several kilometers in the upstream direction, especially within the large central channel (Fig. 5a). We associate this shift with enhanced basal melt rates that are above those required for a steady-state ice thickness ~~and~~ thus causing steep basal gradients. A remarkably similar change in geometry was found for a melt channel at Petermann Glacier between 2002 and 2010 (Münchow et al., 2014).

For the initialization of ice sheet models, ice geometries such as the one from ~~BedMachine~~ IceBridge ~~BedMachine Greenland~~ (Morlighem et al., 2017) are often used, which are based on a compilation of airborne radar measurements. At 79NG, ice thickness measurements since 1993 have been taken into account in ~~BedMachine-V5~~ IceBridge ~~BedMachine Greenland, Version 5~~ (Morlighem et al., 2022), also the 2018 UWB data have been included here. The comparison of the 2021 UWB ice thicknesses with ~~BedMachine-V5~~, Version 5 at the two across-ice flow sections C–C' and D–D' shows differences of  $-91 \pm 108$  m and  $-188 \pm 56$  m, respectively (Fig. 5c,d). This illustrates that the ice thickness is difficult to represent of those glaciers which change significantly in a few years due to the warming of the ocean and atmosphere. The impact of a more accurate, current ice thickness distribution on the simulated evolution of floating ice tongues needs to be explored in regional studies such as the one from Choi et al. (2017) for 79NG, which is beyond the scope of this study.

At Petermann Glacier, Washam et al. (2019) observed strong seasonal variations of basal melt rates beneath the floating tongue with summer melt rates more than four times larger than in winter. ~~They linked the seasonality to warmer ocean currents reaching the ice base of Petermann Glacier (Shroyer et al., 2017; Washam et al., 2019).~~ In contrast to this, we see no evidence of seasonality in the melt rate time series for 79NG, despite the increase in melting at ApRES1 in July 2020. The absence of a summer increase of basal melt rates is consistent with in-situ measurements of ocean temperatures and velocities between September 2016 and September 2017 (Schaffer et al., 2020), showing persistent inflow of warm AIW into the cavity and an overlying outflow of cold-modified AIW throughout the year without a clear seasonal signal. ~~Also, the calving front position of 79NG shows no pronounced seasonality (Loebel et al., 2023).~~

Combining the findings of this study with the observed inflow (Schaffer et al., 2020) and modeled (Reinert et al., 2023) currents below the 79NG shapes a full picture of the ice–ocean interaction at 79NG. Warm AIW flows over the sill into the cavity as a dense and saline bottom plume. As the keel of thick ice near the grounding line is exposed to this warm water, large amounts of heat are supplied to the ice base, ~~which results in the observed high thinning and melt rates (see Appendix D).~~ The meltwater rises along the basal slope as a positively buoyant plume that may drive turbulent mixing with the warm AIW and thus intensify basal melting (~~Jenkins and Doake, 1991; Jenkins, 2011; Schaffer et al., 2020~~) (Jenkins and Doake, 1991; Jenkins, 2011; Schaffer et al., 2020).

345 To melt ice at a rate of  $140 \text{ m a}^{-1}$  requires a heat flux between  $1360$  and  $1600 \text{ W m}^{-2}$  (see Appendix D) depending on the range of the glaciers temperatures which we assume to be between  $\sim 0 \text{ K}$  (temperate ice) and  $30 \text{ K}$  below the pressure melting point. This heat flux must be provided by the water in the cavity below 79NG. We assume a salinity of  $34.5 \text{ psu}$  and an ice draft of  $320 \text{ m}$ , estimated for the location of ApRES2a, where the highest melt rates of  $140 \text{ m a}^{-1}$  were determined during winter. Measurements of the inflow temperatures exceeded  $1.2^\circ \text{ C}$  at the calving front (Schaffer et al., 2020), corresponding to  $2.9 \text{ K}$  above the pressure melting point at the position of the observation. In order to produce a sufficiently high turbulent heat flux into the boundary layer for this given temperature, an ambient velocity of  $0.22 \text{ m s}^{-1}$  is required for temperate ice and 350  $0.26 \text{ m s}^{-1}$  for ice of  $30 \text{ K}$  below the pressure melting point (see Appendix D). Previously simulated velocities of a buoyant plume rising along the ice base of 79NG indicate velocities of up to  $0.22 \text{ m s}^{-1}$  (Reinert et al., 2023). From these numbers, we conclude that the ocean currents underneath 79NG are able to supply a heat flux that is high enough to explain even the highest observed annual mean melt rates if they get in contact with the ice base.

The spread of thinning rates near the grounding line from near zero to  $> 100 \text{ m a}^{-1}$  may be related to the water column 355 thickness distribution. A water column thickness of  $50$  to  $140 \text{ m}$  (Mayer et al., 2000) was found where we observe the highest basal melt rates and where the grounding line reaches farthest downstream. We do not have any information on water column thickness elsewhere. However, the southeastern part of the grounding line is situated on a mountainous landform. We hypothesize that only a shallow water column exists here, which prevents the flow of warm ocean currents toward the grounding line, resulting in the observed low thinning rates.

360 Further downstream, the plume loses heat to the melting of ice and buoyancy by entrainment of ambient water. Thus, it cools down and eventually detaches from the ice base, leading to a strong decrease in basal melting for the thinner, more gently sloped areas of the floating ice tongue (Reinert et al., 2023). This concept is consistent with the low melt rates and glacially modified AIW observed at the calving front, where the outflowing water is  $0.9^\circ \text{ C}$   $0.9 \text{ K}$  cooler than the inflowing AIW (Schaffer et al., 2020).

365 While this picture accounts for the first-order, quasi-twodimensional distribution of melt rates as well as the observed hydrography, it does not explain the existence and growth of basal channels. In previous studies, the existence and location of basal channels have been linked to subglacial water discharge that rises along the basal slope inside a pre-existing basal channel and intensifies basal melting (Le Brocq et al., 2013; Marsh et al., 2016; Washam et al., 2019). We hypothesize that the same applies to the large basal channel at 79NG, where subglacial discharge might have caused the channel's growth in the upstream 370 direction due to extreme basal melting. ~~Although there are no measurements of basal~~

~~Unfortunately, we have no observations of nadir melt rates within a channel since meltwater in summer accumulated in the surface depression above, preventing the deployment of an ApRES. We analyzed the hydrostatic imbalance of the ice above the channel to assess the possibility of determining melt rates based on Lagrangian surface elevation changes. Therefore, we calculated the mean vertical ice density from the ice thickness and the surface elevation, recorded during the flight campaign 375 in 2021 using the UWB airborne radar and laser scanner. The result showed significantly lower densities of the ice above the channels, suggesting that the ice is not in hydrostatic equilibrium, which confirms the findings of Chartrand and Howat (2023). Since this prevents the analysis of melt rates using satellite remote sensing data, we can only draw conclusions from the~~


channel, we can draw conclusions about them by considering the basal geometry and its changes in time. Thus temporal changes using UWB airborne radar. High-resolution measurements of the basal topography at Petermann and Thwaites glaciers using underwater vehicles have revealed steep-sided terraces and heterogeneous melting (Dutrieux et al., 2014; Schmidt et al., 2023). Since the UWB airborne radar does not allow us to resolve the base in a similar resolution, we consider only the average basal slope and thus interpret the average melt pattern.

At 79NG, the high melt rates occur primarily near the origin of the channel, where the greatest basal slope exists. With decreasing basal slopes inside the channel, the melt rate also decreases. This results in an upstream shift in the melt pattern compared to the outside of the channel: (i) Upstream the grounding line and downstream where a low water column exists, higher melt rates occur inside the channel than outside where the ice is grounded or where a low water column exists. (ii) In the vicinity of the grounding line, lower melt rates occur in the channel than outside where the ice is in contact with warm ocean currents, lower melt rates occur in the channel than outside. This pattern is consistent with observations from a basal channel at the Filchner Ice Shelf, Antarctica (Humbert et al., 2022). In addition, it explains the small-scale variability in melt rates we observe at some pRES measurement locations (e.g. at g4\_p02 in Figs. 1c, 2).

The steepening of an ice base is indicating indicates that the melt rates and the ice transport are not in equilibrium, which seems to be the case at 79NG in the past at the same time when the inflow of warm AIW was present. However, we found indications for reduced heat transport into the cavity of 79NG since 2018. In that year, we observed a strong decrease in the melt rate at all ApRES sites that remained low since. Additionally, the repeat of the ApRES2 measurements after two years shows that high melt rates between October 2016 and July 2018 have reduced the ice thickness at the starting location of the ApRES measurement by 50 m (Eulerian perspective). Due to the lower melt rates from July 2018 onward, the ice thinned less than before (Lagrangian perspective). As a result, the ice thickness at the location where the measurement of ApRES2b stopped in December 2019 was even thicker than two years before.

Besides a warm AIW inflow into the cavity, the melt rates can also be enhanced by an increase in subglacial discharge. Subglacial discharge has a seasonal component, including supraglacial lake drainage. The drainage of supraglacial lakes is taking place on a short time scale, even within only one day (Neckel et al., 2020). The lag between lake drainage and discharge across the grounding line is not well known, but it is reasonable to assume that the subglacial hydrological system is buffering water. Drainage of supraglacial lakes is not restricted to the summer period, as the study of Schröder et al. (2020) also detected events in winter, which affects the timing of subglacial discharge. It is to be expected that this subglacial discharge mainly affects the melt rates in the basal channels (Le Brocq et al., 2013). This appears to have been the case for 79NG over the past decade as the central channel has grown and evolved in the upstream direction. In order to quantify if an increase in subglacial discharge has occurred, we roughly estimated the upstream extent of the average surface melt area from a simple analysis of the median summer skin temperatures (July and August) from the Copernicus Arctic Regional Reanalysis (Schyberg et al., 2020) along the ice flow line upstream of the large central channel (Figs Fig. 8, E1). This revealed a substantial increase of the median temperature by up to 1.9°C since the year 2000. The area in which the temperature was positive above 0°C on 50% of the days ranged 32 km upstream until the period 2000–2004, but over 70 km since 2005–2009. Thus, intense warming of the atmosphere in the early 2000s increased the area of summer surface melt and most likely increased subglacial discharge.

This is consistent with the finding at Petermann Glacier, where the subglacial water discharge has doubled since 2001 (Ciraci et al., 2023).



**Figure 8.** Development of skin temperature from Copernicus Arctic Regional Reanalysis (Schyberg et al., 2020) along the central flow line since 1991. The temperature shown is the median skin temperature between 01 July and 31 August (at 15:00 UTC) for the given years. Maps of the median skin temperature are shown in Appendix E.

## 415 6 Conclusions

By combining geophysical in-situ and remote sensing methods, we revealed changes in the ice geometry of the 79NG: the ice near the grounding line has become channelized and significantly thinner in the last two decades. Large, 500 m high subglacial channels originate several kilometers upstream of the grounding line. Here, higher melt rates occur inside the channel than outside, while downstream of the ground-grounding line we found evidence of higher melt rates outside the channels. These high melt rates of  $> 100 \text{ m a}^{-1}$  are caused by thick ice that is in contact with the warm bottom-plume and result in steep basal slopes because they exceed such melt rates, which are necessary water masses at the bottom of the cavity. Since these melt rates are above those required for a steady-state ice thickness, this leads to ice thinning in the Eulerian perspective and thus a steeper base slope. However, we also found low melt rates and small basal gradients under thick ice, particularly off the center of the glacier, which we attribute to a shallow water column thickness that prevents the flow of warm ocean currents toward the grounding line. As the ice thins in the downstream direction, the basal slope and the melt rates drop sharply, resulting in low values at the calving front. The temporal variation since September 2016 shows a non-seasonal variability and significantly decreasing Lagrangian melt rates with increasing distance to the grounding line. Since 2018, these time series show a decrease in melt rates, suggesting a recent inflow of colder water into the cavity beneath the glacier. We conclude that warmer ocean inflow and increased subglacial discharge have caused the changes in ice geometry in the vicinity of the grounding line by forcing high basal melt rates. The consequences of the thinning on However, based on our findings of thinning and upstream

progression of subglacial channels, we cannot assess their impact on future stability. It would require numerical models, as well as longer observational time series to evaluate the stability of ~~the~~-79NG and the Northeast Greenland Ice Stream ~~should be investigated~~-which should be addressed in further studies.

435 *Data availability.* ApRES time-series of basal melt rates, (<https://doi.pangaea.de/10.1594/PANGAEA.928903>; Zeising et al., 2023b) thinning rates derived from single repeated pRES measurements (<https://doi.pangaea.de/10.1594/PANGAEA.928541>; Zeising et al., 2023c) and ice thickness data from the 2021 UWB survey (<https://doi.pangaea.de/10.1594/PANGAEA.963752>; Zeising et al., 2023a) are submitted to the World Data Center PANGAEA. Stake surface ablation/accumulation measurements from 2017 to 2018 (<https://doi.org/10.1594/PANGAEA.922131>; Zeising et al., 2020) are available at the World Data Center PANGAEA.

## Appendix A: Occurrence and identification of nadir and off-nadir reflection

440 We have identified different cases of how nearby basal channels affect the origin of the first recorded basal reflection in repeated pRES echograms (Tab. A1 and Fig. A1). All have in common that the derived range differences of two measurements ( $\Delta R$  derived) underestimate the nadir ice thickness ( $\Delta H$  nadir) or the off-nadir ice thickness ( $\Delta H$  off-nadir).


**Table A1.** Possibilities of how basal channels affect the recording of nadir and off-nadir reflections. Notation:  $t_1$ : time of first measurement,  $t_2$ : time of repeated measurement,  $H_1$ : ice thickness at  $t_1$ ,  $H_2$ : ice thickness at  $t_2$ ,  $\Delta H$  nadir: difference in ice thickness nadir,  $\Delta H$  off-nadir: difference in ice thickness at off-nadir location,  $\Delta H$  derived: difference in depth at nadir projection.

Case	Figure	$t_1$	$t_2$	$\Delta H$
A	Fig. A1a	<b>nadir</b>	<b>off-nadir</b>	$\Delta H$ <b>nadir</b> < $\Delta R$ <b>derived</b> < $\Delta H$ <b>off-nadir</b> Basal channel did not exist or was too small to be detected at $t_1$ . At $t_2$ , the growth of the channel is significantly larger than the ice thickness reduction nadir of the measurement device.
B	Fig. A1b	<b>off-nadir</b>	<b>off-nadir</b>	$\Delta R$ <b>derived</b> < $\Delta H$ <b>off-nadir</b> Basal channel exists at $t_1$ . At $t_2$ , the ice thickness reduction nadir of the measurement device is not significantly larger than the growth of the channel.
C	Fig. A1c	<b>off-nadir</b>	<b>nadir</b>	$\Delta H$ <b>off-nadir</b> < $\Delta R$ <b>derived</b> < $\Delta H$ <b>nadir</b> Basal channel exists at $t_1$ . At $t_2$ , the ice thickness reduction nadir of the measurement device is significantly larger than the growth of the channel.
D	Fig. A1d	<b>off-nadir</b>	<b>off-nadir</b>	$\Delta R$ <b>derived</b> < $\Delta H$ <b>off-nadir</b> Two basal channel exist at $t_1$ . At $t_2$ , the ice thickness reduction nadir of the measurement device is not significantly larger than the growth of at least one of both channels. This type can not be distinguished from Case B without known geometry.

Several pRES echograms indicate the occurrence of numerous strong basal reflections (Fig. A2). For steep basal gradients, the off-nadir reflection may occur prior to the nadir reflection. We interpret the first basal reflection as an off-nadir reflection, as long as no further information reveals the true nadir reflection. ~~Herewith, The first off-nadir basal reflection in the first and the repeated measurement can have occurred at two different locations (locations “A” and “B”). If this is the case, we can conclude that the melt rate has been higher at location B than at location A as otherwise, the first basal reflection would have occurred at location A in both measurements. However, when we compare the range to A and to B, we know the true change in ice thickness at B has been higher. Thus, we underestimate the thinning and the melt rate. If the second basal return occurred~~ ~~at an off-nadir angle, the estimated melt rate is below the vertical melt rate at that location. Still, the resulting basal melting in the vicinity of the measurement is always underestimated, although the nadir melt rate might be lower can be even lower, but we cannot determine this melt rate.~~


445

450



**Figure A1.** Sketch of off-nadir reflections and their influence on basal melt rates. The solid lines refer to the time of the first measurement (blue),  $t_1$ , and the dotted lines refer to the time of the repeat measurement (red),  $t_2$ . The yellow triangles mark the measurement positions. The red and blue straight lines mark the closest distance from the measurement to the ice base. The segments of a circle (up to  $30^\circ$  to nadir) correspond to the possible positions of the reflector with the shortest distance. The lengths of the bars on the right reflect the thinning of the ice between  $t_1$  and  $t_2$  for the position of the measurement ( $\Delta H$  nadir), for the position of the closest reflector at  $t_2$  ( $\Delta H$  off-nadir), and for the range difference of the blue and red lines ( $\Delta R$  derived). Note that at least one of  $\Delta H$  nadir or  $\Delta H$  off-nadir is always larger than  $\Delta R$  derived.





**Figure A2.** pRES measurements with the identified nadir and off-nadir reflections. Echograms from the first measurement are shown in blue and from the repeated measurement in red. Vertical dashed lines mark the nadir basal return and thus represent the ice thickness  $H$ .

## Appendix B: ApRES time series analysis

### B1 Estimation of ice deformation

455 Ice deformation affects the range  $R$  from the ApRES to an englacial reflector located at  $x_0$ ,  $y_0$ , and  $z_0$  relative to the measurement location. ApRES measurements allow us to determine the depth profile of the vertical displacement of englacial reflectors relative to the surface and thus to compute the vertical strain  $\varepsilon_{zz}$ . The range displacement of an off-nadir reflector viewed at an angle  $\alpha$  due to ice deformation  $\Delta R_\varepsilon$  is also affected by the two horizontal normal  $\varepsilon_{xx}$  and  $\varepsilon_{yy}$  as well as the shear components  $\varepsilon_{xz}$ ,  $\varepsilon_{yz}$ ,  $\varepsilon_{zx}$  and  $\varepsilon_{zy}$ . Accordingly, the location of a reflector shifts to  $x_0 + \int_0^{x_0} (\varepsilon_{xx} + \varepsilon_{xz}) dx$ ,  $y_0 + \int_0^{y_0} (\varepsilon_{yy} + \varepsilon_{yz}) dy$  and  
 460  $z_0 + \int_0^{z_0} (\varepsilon_{zz} + \varepsilon_{zx} + \varepsilon_{zy}) dz$  at the time of a second measurement so that  $\Delta R_\varepsilon$  can be calculated as follows:

$$\Delta R_\varepsilon = \sqrt{\left(x_0 + \int_0^{x_0} (\varepsilon_{xx} + \varepsilon_{xz}) dx\right)^2 + \left(y_0 + \int_0^{y_0} (\varepsilon_{yy} + \varepsilon_{yz}) dy\right)^2 + \left(z_0 + \int_0^{z_0} (\varepsilon_{zz} + \varepsilon_{zx} + \varepsilon_{zy}) dz\right)^2} - \sqrt{x_0^2 + y_0^2 + z_0^2} \quad (\text{B1})$$

For a nadir reflection ( $\alpha = 0$ ) where  $x_0 = 0$  and  $y_0 = 0$ , ~~the displacement due to deformation depends only on the vertical strain~~  
we assume that shear terms are negligible

$$\Delta R_\varepsilon^n = \int_0^{z_0} \varepsilon_{zz} dz, \quad (\text{B2})$$

465 where the range  $R$  and thus  $z_0$  equals the ice thickness  $H$ .

The estimation of  $\Delta R_\varepsilon$  in the case of an off-nadir reflection requires the quantification of the normal and shear components as well as of  $\alpha$  and  $\beta$ , which are unknown. In the following, we consider the shear terms to be small, as investigation of a melt channel on Filchner Ice Shelf (Humbert et al., 2022) has shown that the elastic shear strain is an order of magnitude lower than the strain in normal direction. With channels appearing during our measurement period, the instantaneous elastic component  
 470 is the one to be considered here. From the continuity equation (e.g. Cuffey and Paterson, 2010), we find that


$$\varepsilon_{zz} = -(\varepsilon_{xx} + \varepsilon_{yy}) \quad (\text{B3})$$

which is the case for incompressible ice. Additionally, we know that  $\alpha \leq 30^\circ$  for the ApRES system so that the sum of the horizontal distances between the ApRES and the reflector is smaller or equals the vertical distance:  $x_0 + y_0 \leq z_0$ . Thus, we can do the following quantification

$$475 \quad 0 \leq |\Delta R_\varepsilon| \leq |\Delta R_\varepsilon^n|, \quad (\text{B4})$$


where  $\Delta R_\varepsilon$  and  $\Delta R_\varepsilon^n$  have always the same sign. This shows that strain thinning or thickening cannot be overestimated by assuming a reflection occurred from a nadir scatterer.

## **B2 ApRES echograms**



**Figure B1.** Analysis of [ApRES1 echograms](#) [ApRES2a time series](#). (a) Time-echogram of a Lagrangian measurement at [ApRES1](#) [ApRES2a](#) recorded between August 2016 and [June 2022](#). In 2016 and 2017, several ApRES malfunctions caused data gaps. (b) Selected echograms smoothed with a 5 m moving average filter. [July 2018](#). The blue shaded range corresponds to [black outline marks](#) the displacement of the first 50 m below the basal return within the measurement period. (c) Time series of ablation rate (negative for ablation). The grey shaded area marks the uncertainty due to the off-nadir correction. (d) Average time series of the determined melt rate of (color) within the first 50 m below the basal return, corresponding depth (to the area marked by black lines in (a)). (e) Time series of basal melt rate. The gray dashed line shows the 95% quantile, the solid line the median, and the shaded area marks the range between the 25% and 75% quantile.

Same as Figure ?? but for ApRES2a.



**Figure B2.** Same as Figure [??-B1](#) but for ApRES2b between July 2018 and January 2020.






Figure B3. Same as Figure [??-B1](#) but for ApRES3 [between July 2017 and September 2023](#).

## Appendix C: Subglacial channel observed by airborne radar



**Figure C1.** Map of [UWB seismic locations from 1998 \(Mayer et al., 2000\)](#) and [airborne radar data from 1998 \(Reeh's\), 2018 \(UWB\) and 2021](#) and [seismic locations from 1998 \(Mayer et al., 2000\)\(UWB\)](#).



**Figure C2.** UWB airborne echograms across basal channel (a–h) from 2018 and (g–q) from 2021. For location see Fig. C1.



## 480 Appendix D: Oceanic heat flux

In order to estimate the oceanic heat flux  $q_w$  required to sustain the basal melt rates  $a_b$  [ $\text{m s}^{-1}$ ] derived in this study, we separate the heat flux  $q_w$  into two components: the heat flux  $q_m$  to melt the ice and the heat flux into the glacier interior  $q_i$  that is required for heating the ice by  $\Delta T$  to the pressure melting point:


$$q_w = \underbrace{\rho_i a_b L}_{q_m} + \underbrace{\rho_i c_i(T) a_b \Delta T}_{q_i}. \quad (\text{D1})$$

485 The heat fluxes depend on the density of the ice,  $\rho_i = 917 \text{ kg m}^{-3}$ , the latent heat of fusion,  $L = 334000 \text{ J kg}^{-1}$ , and the specific heat capacity for ice,  $c_i(T) = 146.3 + 7.253 \cdot T[\text{K}] \text{ J kg}^{-1} \text{ K}^{-1}$  with the temperature  $T$  in Kelvin (Ritz, 1987). ~~For an assumed range of glacier interior temperatures between  $\sim 0 \text{ K}$  (temperate ice) and  $30 \text{ K}$  below the pressure melting point, a basal melt rate of  $140 \text{ m a}^{-1}$  (as it was observed at all three ApRES sites) requires a heat flux between  $1360$  and  $1600 \text{ W m}^{-2}$ . This heat flux must be provided by the water in the cavity below 79NG.~~

490 To obtain an estimate of the oceanic heat flux [that the ocean can provide](#), we follow the approach implemented in the Finite Element Sea ice-Ocean Model (FESOM; Timmermann et al., 2012). Here, a three-equation system is used that determines the temperature and salinity of a thin boundary layer along the ice-shelf base from its heat and freshwater exchange with the ice and the ambient ocean (Hellmer and Olbers, 1989; Holland and Jenkins, 1999). Besides the ocean temperature, the heat flux into this boundary layer is determined by the flow velocity in the ambient ocean, as the latter determines the friction and thus  
495 defines the turbulent fluxes of heat and salt (Jenkins and Doake, 1991).

~~As an example case for the 79NG sub-ice cavity, we assume a salinity of  $34.5 \text{ psu}$  and an ice draft of  $320 \text{ m}$ , estimated for the location of ApRES2a, where the highest melt rates of  $140 \text{ m a}^{-1}$  were determined during winter. Measurements of the inflow temperatures exceed  $1.2^\circ \text{ C}$  at the calving front (Schaffer et al., 2020), corresponding to  $2.9 \text{ K}$  above the pressure melting point at the position of the observation. In order to produce a sufficiently high turbulent heat flux into the boundary layer for this  
500 given temperature, an ambient velocity of  $0.22 \text{ m s}^{-1}$  is required for temperate ice and  $0.27 \text{ m s}^{-1}$  for ice of  $30 \text{ K}$  below the pressure melting point. Previously simulated velocities of a buoyant plume rising along the ice base of 79NG indicate velocities of up to  $0.22 \text{ m s}^{-1}$  (Reinert et al., 2023). From these numbers, we conclude that the ocean currents underneath 79NG are able to supply a heat flux that is high enough to explain even the maximum observed melt rates.~~

## Appendix E: Surface skin temperature



**Figure E1.** Skin temperature of 79NG between (a) 1991 – 1994, (b) 1995 – 1999, (c) 2000 – 2004, (d) 2005 – 2009, (e) 2010 – 2014, (f) 2015 – 2019 and (g) 2020 – 2022 from Copernicus Arctic Regional Reanalysis (Schyberg et al., 2020). The shown temperatures is the median skin temperature between 01 July and 31 August (at 15:00 UTC) of the given years.

505 *Author contributions.* OZ, DS, NN, and AH conducted the field expeditions, and AH and VH the airborne campaigns. AH has designed the study and planned field expeditions and airborne campaigns. OZ processed the (A)pRES data, and estimated and analyzed the resulting thinning and basal melt rates. ND processed the UWB 2018 data and discovered the central channel. VH processed the laser scanner data and the UWB 2021 data. NN processed all satellite data and determined the elevation changes and grounding line location. RT provided oceanographic expertise. OZ wrote the manuscript with contributions from all coauthors.

510 *Competing interests.* The contact author has declared that neither they nor their co-authors have any competing interests.

*Acknowledgements.* This study is part of the GROCE1 and GROCE2 projects funded by the German Federal Ministry of Research and Education under Grant No. 03F0778A and 03F0855A. NN has received funding from the European Union’s Horizon 2020 research and innovation program under grant agreement No 689443 via project iCUPE (Integrative and Comprehensive Understanding on Polar Environments). [Airborne DTU Space 60 MHz ice sounder raw data was processed into radargramms and into ice surface as well as ice base data by Steen Savstrup Kristensen](#). The airborne data were acquired as part of AWI’s RESURV79 (2018) and 79NG-EC (2021) campaigns with AWI’s polar aircrafts (Wesche et al., 2016). We want to thank Tobias Binder and Martin Gehrman, the aircraft crews, Will Wilson, Dean Emberly, Stewart Clark, Marc-André Verner, Luke Cirtwill, Ryan Schrader, the team of Villum Research Station, the Arctic Command and the team Station Nord in 2018 and 2021 for their support during the expeditions. TanDEM-X and TerraSAR-X data were made available through German Aerospace Center proposals GLAC7208 and HYD2059. The authors would like to thank Emerson E&P Software, Emerson Automation Solutions, for providing licenses in the scope of the Emerson Academic Program. We like to thank Steven Franke for his support in handling UWB data. We want to thank Jens Köhler and Graham Niven for their support in the field. We acknowledge the effort of Shfaqat Abbas Khan for servicing our ApRES during the lockdown in 2020 and retrieving our SD cards.

## References

- Alley, K. E., Scambos, T. A., Siegfried, M. R., and Fricker, H. A.: Impacts of warm water on Antarctic ice shelf stability through basal channel formation, *Nature Geoscience*, 9, 290–293, <https://doi.org/10.1038/ngeo2675>, 2016.
- Bentley, M. J., Smith, J. A., Jamieson, S. S. R., Lindeman, M. R., Rea, B. R., Humbert, A., Lane, T. P., Darvill, C. M., Lloyd, J. M., Straneo, F., Helm, V., and Roberts, D. H.: Direct measurement of warm Atlantic Intermediate Water close to the grounding line of Nioghalvfjærdsfjorden (79° N) Glacier, northeast Greenland, *The Cryosphere*, 17, 1821–1837, <https://doi.org/10.5194/tc-17-1821-2023>, 2023.
- Berthier, E., Cabot, V., Vincent, C., and Six, D.: Decadal Region-Wide and Glacier-Wide Mass Balances Derived from Multi-Temporal ASTER Satellite Digital Elevation Models. Validation over the Mont-Blanc Area, *Frontiers in Earth Science*, 4, 63, <https://doi.org/10.3389/feart.2016.00063>, 2016.
- Brennan, P. V., Lok, L. B., Nicholls, K., and Corr, H.: Phase-sensitive FMCW radar system for high-precision Antarctic ice shelf profile monitoring, *IET Radar, Sonar & Navigation*, 8, 776–786, <https://doi.org/10.1049/iet-rsn.2013.0053>, 2014.
- Burchard, H., Bolding, K., Jenkins, A., Losch, M., Reinert, M., and Umlauf, L.: The Vertical Structure and Entrainment of Subglacial Melt Water Plumes, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002925, <https://doi.org/10.1029/2021MS002925>, 2022.
- Chartrand, A. M. and Howat, I. M.: A comparison of contemporaneous airborne altimetry and ice-thickness measurements of Antarctic ice shelves, *Journal of Glaciology*, p. 1–14, <https://doi.org/10.1017/jog.2023.49>, 2023.
- Choi, Y., Morlighem, M., Rignot, E., Mouginot, J., and Wood, M.: Modeling the response of Nioghalvfjærdsfjorden and Zachariae Isstrøm Glaciers, Greenland, to ocean forcing over the next century, *Geophysical Research Letters*, 44, 11 071–11 079, <https://doi.org/10.1002/2017GL075174>, 2017.
- Ciraci, E., Rignot, E., Scheuchl, B., Tolpekin, V., Wollersheim, M., An, L., Milillo, P., Bueso-Bello, J.-L., Rizzoli, P., and Dini, L.: Melt rates in the kilometer-size grounding zone of Petermann Glacier, Greenland, before and during a retreat, *Proceedings of the National Academy of Sciences*, 120, e2220924 120, <https://doi.org/10.1073/pnas.2220924120>, 2023.
- Corr, H. F., Jenkins, A., Nicholls, K. W., and Doake, C.: Precise measurement of changes in ice-shelf thickness by phase-sensitive radar to determine basal melt rates, *Geophysical Research Letters*, 29, 73–1–73–4, <https://doi.org/10.1029/2001GL014618>, 2002.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers - 4th ed.*, Elsevier, 2010.
- Dutrieux, P., Stewart, C., Jenkins, A., Nicholls, K. W., Corr, H. F., Rignot, E., and Steffen, K.: Basal terraces on melting ice shelves, *Geophysical Research Letters*, 41, 5506–5513, <https://doi.org/10.1002/2014GL060618>, 2014.
- Fahnestock, M. A., Joughin, I., Scambos, T. A., Kwok, R., Krabill, W. B., and Gogineni, S.: Ice-stream-related patterns of ice flow in the interior of northeast Greenland, *Journal of Geophysical Research: Atmospheres*, 106, 34 035–34 045, <https://doi.org/10.1029/2001JD900194>, 2001.
- Fujita, S., Matsuoka, T., Ishida, T., Matsuoka, K., and Mae, S.: A summary of the complex dielectric permittivity of ice in the megahertz range and its applications for radar sounding of polar ice sheets, in: *Physics of ice core records*, pp. 185–212, Hokkaido University Press, <http://hdl.handle.net/2115/32469>, (last access: 28 January 2023), 2000.
- Fürst, J. J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M., and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nature Climate Change*, 6, 479, <https://doi.org/10.1038/nclimate2912>, 2016.

- Hale, R., Miller, H., Gogineni, S., Yan, J.-B., Rodriguez-Morales, F., Leuschen, C., Paden, J., Li, J., Binder, T., Steinhage, D., et al.: Multi-channel ultra-wideband radar sounder and imager, in: 2016 IEEE International Geoscience and Remote Sensing Symposium (IGARSS), pp. 2112–2115, IEEE, 2016.
- Hellmer, H. H. and Olbers, D. J.: A two-dimensional model for the thermohaline circulation under an ice shelf, *Antarctic Science*, 1, 325–336, <https://doi.org/10.1017/S0954102089000490>, 1989.
- Helm, V., Humbert, A., and Miller, H.: Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *The Cryosphere*, 8, 1539–1559, <https://doi.org/10.5194/tc-8-1539-2014>, 2014.
- Höhle, J. and Höhle, M.: Accuracy assessment of digital elevation models by means of robust statistical methods, *ISPRS Journal of Photogrammetry and Remote Sensing*, 64, 398–406, <https://doi.org/10.1016/j.isprsjprs.2009.02.003>, 2009.
- Holland, D. M. and Jenkins, A.: Modeling thermodynamic ice–ocean interactions at the base of an ice shelf, *Journal of Physical Oceanography*, 29, 1787–1800, [https://doi.org/10.1175/1520-0485\(1999\)029<1787:MTIOIA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<1787:MTIOIA>2.0.CO;2), 1999.
- Humbert, A., Christmann, J., Corr, H. F. J., Helm, V., Höyns, L.-S., Hofstede, C., Müller, R., Neckel, N., Nicholls, K. W., Schultz, T., Steinhage, D., Wolovick, M., and Zeising, O.: On the evolution of an ice shelf melt channel at the base of Filchner Ice Shelf, from observations and viscoelastic modeling, *The Cryosphere*, 16, 4107–4139, <https://doi.org/10.5194/tc-16-4107-2022>, 2022.
- Humbert, A., Helm, V., Neckel, N., Zeising, O., Rückamp, M., Khan, S. A., Loebel, E., Brauchle, J., Stebner, K., Gross, D., Sonderhaus, R., and Müller, R.: Precursor of disintegration of Greenland’s largest floating ice tongue, *The Cryosphere*, 17, 2851–2870, <https://doi.org/10.5194/tc-17-2851-2023>, 2023.
- Jenkins, A.: Convection-driven melting near the grounding lines of ice shelves and tidewater glaciers, *Journal of Physical Oceanography*, 41, 2279–2294, <https://doi.org/10.1175/JPO-D-11-03.1>, 2011.
- Jenkins, A. and Doake, C.: Ice-ocean interaction on Ronne Ice Shelf, Antarctica, *Journal of Geophysical Research: Oceans*, 96, 791–813, <https://doi.org/10.1029/90JC01952>, 1991.
- Jenkins, A., Corr, H. F., Nicholls, K. W., Stewart, C. L., and Doake, C. S.: Interactions between ice and ocean observed with phase-sensitive radar near an Antarctic ice-shelf grounding line, *Journal of Glaciology*, 52, 325–346, <https://doi.org/10.3189/172756506781828502>, 2006.
- Joughin, I., Smith, B. E., and Howat, I. M.: A complete map of Greenland ice velocity derived from satellite data collected over 20 years, *Journal of Glaciology*, 64, 1–11, <https://doi.org/10.1017/jog.2017.73>, 2018.
- Khan, S. A., Kjær, K. H., Bevis, M., Bamber, J. L., Wahr, J., Kjeldsen, K. K., Bjørk, A. A., Korsgaard, N. J., Stearns, L. A., Van Den Broeke, M. R., et al.: Sustained mass loss of the northeast Greenland ice sheet triggered by regional warming, *Nature Climate Change*, 4, 292–299, <https://doi.org/10.1038/nclimate2161>, 2014.
- Khan, S. A., Bamber, J. L., Rignot, E., Helm, V., Aschwanden, A., Holland, D. M., van den Broeke, M., King, M., Noël, B., Truffer, M., Humbert, A., Colgan, W., Vijay, S., and Kuipers Munneke, P.: Greenland Mass Trends From Airborne and Satellite Altimetry During 2011–2020, *Journal of Geophysical Research: Earth Surface*, 127, e2021JF006505, <https://doi.org/10.1029/2021JF006505>, 2022a.
- Khan, S. A., Choi, Y., Morlighem, M., Rignot, E., Helm, V., Humbert, A., Mouginot, J., Millan, R., Kjær, K. H., and Bjørk, A. A.: Extensive inland thinning and speed-up of Northeast Greenland Ice Stream, *Nature*, 611, 727–732, <https://doi.org/10.1038/s41586-022-05301-z>, 2022b.
- King, M. D., Howat, I. M., Candela, S. G., Noh, M. J., Jeong, S., Noël, B. P. Y., van den Broeke, M. R., Wouters, B., and Negrete, A.: Dynamic ice loss from the Greenland Ice Sheet driven by sustained glacier retreat, *Communications Earth & Environment*, 1, 1, <https://doi.org/10.1038/s43247-020-0001-2>, 2020.

- 595 Kjeldsen, K. K., Korsgaard, N. J., Bjørk, A. A., Khan, S. A., Box, J. E., Funder, S., Larsen, N. K., Bamber, J. L., Colgan, W., Van Den Broeke, M., et al.: Spatial and temporal distribution of mass loss from the Greenland Ice Sheet since AD 1900, *Nature*, 528, 396–400, <https://doi.org/10.1038/nature16183>, 2015.
- Krieger, L., Floricioiu, D., and Neckel, N.: Drainage basin delineation for outlet glaciers of Northeast Greenland based on Sentinel-1 ice velocities and TanDEM-X elevations, *Remote Sensing of Environment*, 237, 111 483, <https://doi.org/10.1016/j.rse.2019.111483>, 2020.
- 600 Le Brocq, A. M., Ross, N., Griggs, J. A., Bingham, R. G., Corr, H. F., Ferraccioli, F., Jenkins, A., Jordan, T. A., Payne, A. J., Rippin, D. M., et al.: Evidence from ice shelves for channelized meltwater flow beneath the Antarctic Ice Sheet, *Nature Geoscience*, 6, 945–948, <https://doi.org/10.1038/ngeo1977>, 2013.
- Lindeman, M. R., Straneo, F., Wilson, N. J., Toole, J. M., Krishfield, R. A., Beird, N. L., Kanzow, T., and Schaffer, J.: Ocean Circulation and Variability Beneath Nioghalvfjærdsbræ (79 North Glacier) Ice Tongue, *Journal of Geophysical Research: Oceans*, 125, e2020JC016 091, <https://doi.org/10.1029/2020JC016091>, 2020.
- 605 Loebel, E., Scheinert, M., Horwath, M., Humbert, A., Sohn, J., Heidler, K., Liebezeit, C., and Zhu, X. X.: Calving front monitoring at sub-seasonal resolution: a deep learning application to Greenland glaciers, *The Cryosphere Discussions*, 2023, 1–21, <https://doi.org/10.5194/tc-2023-52>, 2023.
- Marsh, O. J., Fricker, H. A., Siegfried, M. R., Christianson, K., Nicholls, K. W., Corr, H. F., and Catania, G.: High basal melting forming a channel at the grounding line of Ross Ice Shelf, Antarctica, *Geophysical Research Letters*, 43, 250–255, <https://doi.org/10.1002/2015GL066612>, 2016.
- 610 Mayer, C., Reeh, N., Jung-Rothenhäusler, F., Huybrechts, P., and Oerter, H.: The subglacial cavity and implied dynamics under Nioghalvfjærdsfjorden Glacier, NE-Greenland, *Geophysical Research Letters*, 27, 2289–2292, <https://doi.org/10.1029/2000GL011514>, 2000.
- Mayer, C., Schaffer, J., Hattermann, T., Floricioiu, D., Krieger, L., Dodd, P. A., Kanzow, T., Licciulli, C., and Schannwell, C.: Large ice loss variability at Nioghalvfjærdsfjorden Glacier, Northeast-Greenland, *Nature Communications*, 9, 2768, <https://doi.org/10.1038/s41467-018-05180-x>, 2018.
- 615 Millgate, T., Holland, P. R., Jenkins, A., and Johnson, H. L.: The effect of basal channels on oceanic ice-shelf melting, *Journal of Geophysical Research: Oceans*, 118, 6951–6964, <https://doi.org/10.1002/2013JC009402>, 2013.
- Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K. B.: BedMachine v3: Complete bed topography and ocean bathymetry mapping of Greenland from multibeam echo sounding combined with mass conservation, *Geophysical Research Letters*, 44, 11 051–11 061, <https://doi.org/10.1002/2017GL074954>, 2017.
- 620 Morlighem, M., Williams, C., Rignot, E., An, L., Arndt, J. E., Bamber, J., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B., O’Cofaigh, C., Palmer, S. J., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglensen, K.: IceBridge BedMachine Greenland, Version 5, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/GMEVBWFLWA7X>, 2022.
- 625 Motyka, R. J., Truffer, M., Fahnestock, M., Mortensen, J., Rysgaard, S., and Howat, I.: Submarine melting of the 1985 Jakobshavn Isbræ floating tongue and the triggering of the current retreat, *Journal of Geophysical Research: Earth Surface*, 116, F01007, <https://doi.org/10.1029/2009JF001632>, 2011.
- 630

- Mouginot, J., Rignot, E., Scheuchl, B., Fenty, I., Khazendar, A., Morlighem, M., Buzzi, A., and Paden, J.: Fast retreat of Zachariæ Isstrøm, northeast Greenland, *Science*, 350, 1357–1361, <https://doi.org/10.1126/science.aac7111>, 2015.
- 635 Münchow, A., Padman, L., and Fricker, H. A.: Interannual changes of the floating ice shelf of Petermann Gletscher, North Greenland, from 2000 to 2012, *Journal of Glaciology*, 60, 489–499, <https://doi.org/10.3189/2014JoG13J135>, 2014.
- Münchow, A., Schaffer, J., and Kanzow, T.: Ocean Circulation Connecting Fram Strait to Glaciers off Northeast Greenland: Mean Flows, Topographic Rossby Waves, and Their Forcing, *Journal of Physical Oceanography*, 50, 509 – 530, <https://doi.org/doi.org/10.1175/JPO-D-19-0085.1>, 2020.
- 640 Neckel, N., Braun, A., Kropáček, J., and Hochschild, V.: Recent mass balance of the Purogangri Ice Cap, central Tibetan Plateau, by means of differential X-band SAR interferometry, *The Cryosphere*, 7, 1623–1633, <https://doi.org/10.5194/tc-7-1623-2013>, 2013.
- Neckel, N., Zeising, O., Steinhage, D., Helm, V., and Humbert, A.: Seasonal observations at 79° N Glacier (Greenland) from remote sensing and in-situ measurements, *Frontiers in Earth Science*, 8, 142, <https://doi.org/10.3389/feart.2020.00142>, 2020.
- Nicholls, K. W., Corr, H. F., Stewart, C. L., Lok, L. B., Brennan, P. V., and Vaughan, D. G.: A ground-based radar for measuring vertical strain rates and time-varying basal melt rates in ice sheets and shelves, *Journal of Glaciology*, 61, 1079–1087, <https://doi.org/10.3189/2015JoG15J073>, 2015.
- 645 Nuth, C. and Kääb, A.: Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change, *The Cryosphere*, 5, 271–290, <https://doi.org/10.5194/tc-5-271-2011>, 2011.
- Rathmann, N., Hvidberg, C., Solgaard, A., Grinsted, A., Gudmundsson, G. H., Langen, P. L., Nielsen, K., and Kusk, A.: Highly temporally resolved response to seasonal surface melt of the Zachariae and 79N outlet glaciers in northeast Greenland, *Geophysical Research Letters*, 44, 9805–9814, <https://doi.org/10.1002/2017GL074368>, 2017.
- Reinert, M., Lorenz, M., Klingbeil, K., Büchmann, B., and Burchard, H.: High-Resolution Simulations of the Plume Dynamics in an Idealized 79°N Glacier Cavity Using Adaptive Vertical Coordinates, *Journal of Advances in Modeling Earth Systems*, 15, e2023MS003721, <https://doi.org/10.1029/2023MS003721>, 2023.
- 655 Rignot, E. and Steffen, K.: Channelized bottom melting and stability of floating ice shelves, *Geophysical Research Letters*, 35, L02 503, <https://doi.org/10.1029/2007GL031765>, 2008.
- Ritz, C.: Time dependent boundary conditions for calculation of temperature fields in ice sheets, in: *The Physical Basis of Ice Sheet Modelling*, edited by Waddington, E. D. and Walder, J. S., IAHS Publication No. 170, pp. 207–216, IAHS Press, Wallingford, UK, 1987.
- Schaffer, J., Kanzow, T., von Appen, W.-J., von Albedyll, L., Arndt, J. E., and Roberts, D. H.: Bathymetry constrains ocean heat supply to Greenland’s largest glacier tongue, *Nature Geoscience*, 13, 227–231, <https://doi.org/10.1038/s41561-019-0529-x>, 2020.
- 660 Schmidt, B. E., Washam, P., Davis, P. E. D., Nicholls, K. W., Holland, D. M., Lawrence, J. D., Riverman, K. L., Smith, J. A., Spears, A., Dichek, D. J. G., Mullen, A. D., Clyne, E., Yeager, B., Anker, P., Meister, M. R., Hurwitz, B. C., Quartini, E. S., Bryson, F. E., Basinski-Ferris, A., Thomas, C., Wake, J., Vaughan, D. G., Anandkrishnan, S., Rignot, E., Paden, J., and Makinson, K.: Heterogeneous melting near the Thwaites Glacier grounding line, *Nature*, 614, 471–478, <https://doi.org/10.1038/s41586-022-05691-0>, 2023.
- 665 Schröder, L., Neckel, N., Zindler, R., and Humbert, A.: Perennial Supraglacial Lakes in Northeast Greenland Observed by Polarimetric SAR, *Remote Sensing*, 12, 2798, <https://doi.org/10.3390/rs12172798>, 2020.
- Schyberg, H., Yang, X., Køltzow, M. A. O., Amstrup, B., Bakketun, A., Bazile, E., Bojarova, J., Box, J. E., Dahlgren, P., Hagelin, S., Homleid, M., Horányi, A., Høyer, J., Johansson, A., Killie, M. A., Körnich, H., Le Moigne, P., Lindskog, M., Manninen, T., Nielsen Englyst, P., Nielsen, K. P., Olsson, E., Palmason, B., Peralta Aros, C., Randriamampianina, R., Samuelsson, P., Stappers, R., Støylen, E., Thorsteinsson,

- 670 S., Valkonen, T., and Wang, Z. Q.: Arctic regional reanalysis on single levels from 1991 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.713858f6>, 2020.
- Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., A. G., Agosta, C., Ahlstrøm, A., Babonis, G., Barletta, V. R., Björk, A. A., Blazquez, A., Bonin, J., Colgan, W., Csatho, B., Cullather, R., Engdahl, M. E., Felikson, D., Fettweis, X., Forsberg, R., Hogg, A. E., Gallee, H., Gardner, A.,  
675 Gilbert, L., Gourmelen, N., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P. L., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mottram, R., Mouginit, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noël, B., Otosaka, I., Pattle, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Sandberg Sørensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., van de Berg, W. J., van der Wal, W., van Wessem, M., Vishwakarma,  
680 B. D., Wiese, D., Wilton, D., Wagner, T., Wouters, B., and Wuite, J.: Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, <https://doi.org/10.1038/s41586-019-1855-2>, 2020.
- Shroyer, E. L., Padman, L., Samelson, R., Münchow, A., and Stearns, L. A.: Seasonal control of Petermann Gletscher ice-shelf melt by the ocean's response to sea-ice cover in Nares Strait, *Journal of Glaciology*, 63, 324–330, <https://doi.org/10.1017/jog.2016.140>, 2017.
- Slater, D. A., Felikson, D., Straneo, F., Goelzer, H., Little, C. M., Morlighem, M., Fettweis, X., and Nowicki, S.: Twenty-first century ocean  
685 forcing of the Greenland ice sheet for modelling of sea level contribution, *The Cryosphere*, 14, 985–1008, <https://doi.org/10.5194/tc-14-985-2020>, 2020.
- Stewart, C. L.: Ice-ocean interactions beneath the north-western Ross Ice Shelf, Antarctica, Ph.D. thesis, University of Cambridge, <https://doi.org/10.17863/CAM.21483>, 2018.
- Stewart, C. L., Christoffersen, P., Nicholls, K. W., Williams, M. J., and Dowdeswell, J. A.: Basal melting of Ross Ice Shelf from solar heat  
690 absorption in an ice-front polynya, *Nature Geoscience*, 12, 435–440, <https://doi.org/10.1038/s41561-019-0356-0>, 2019.
- Straneo, F., Sutherland, D. A., Holland, D., Gladish, C., Hamilton, G. S., Johnson, H. L., Rignot, E., Xu, Y., and Koppes, M.: Characteristics of ocean waters reaching Greenland's glaciers, *Annals of Glaciology*, 53, 202–210, <https://doi.org/10.3189/2012AoG60A059>, 2012.
- Straneo, F., Hamilton, G. S., Stearns, L. A., and Sutherland, D. A.: Connecting the Greenland Ice Sheet and the Ocean: A Case Study of Helheim Glacier and Sermilik Fjord, *Oceanography*, 29, 34–45, <https://doi.org/10.5670/oceanog.2016.97>, 2016.
- 695 The MathWorks Inc.: MATLAB version: 9.13.0 (R2022b), The MathWorks Inc., <https://www.mathworks.com>, 2022.
- Thomsen, H. H., Reeh, N., Olesen, O. B., Bøggilde, C. E., Starzer, W., Weidick, A., and Higgins, A.: The Nioghalvfjærdsfjorden glacier project, North-East Greenland: a study of ice sheet response to climatic change, *Review of Greenland activities*, pp. 95–103, <https://doi.org/10.34194/ggub.v176.5073>, 1997.
- Timmermann, R., Wang, Q., and Hellmer, H.: Ice-shelf basal melting in a global finite-element sea-ice/ice-shelf/ocean model, *Annals of  
700 Glaciology*, 53, 303–314, <https://doi.org/10.3189/2012AoG60A156>, 2012.
- Vaughan, D. G., Corr, H. F., Bindschadler, R. A., Dutrieux, P., Gudmundsson, G. H., Jenkins, A., Newman, T., Vornberger, P., and Wingham, D. J.: Subglacial melt channels and fracture in the floating part of Pine Island Glacier, Antarctica, *Journal of Geophysical Research: Earth Surface*, 117, F03 012, <https://doi.org/10.1029/2012JF002360>, 2012.
- Vaňková, I., Cook, S., Winberry, J. P., Nicholls, K. W., and Galton-Fenzi, B. K.: Deriving Melt Rates at a Complex Ice Shelf Base Using In  
705 Situ Radar: Application to Totten Ice Shelf, *Geophysical Research Letters*, 48, e2021GL092 692, <https://doi.org/10.1029/2021GL092692>, 2021.



- Vijay, S., Khan, S. A., Kusk, A., Solgaard, A. M., Moon, T., and Bjørk, A. A.: Resolving Seasonal Ice Velocity of 45 Greenlandic Glaciers With Very High Temporal Details, *Geophysical Research Letters*, 46, 1485–1495, <https://doi.org/10.1029/2018GL081503>, 2019.
- 710 von Albedyll, L., Schaffer, J., and Kanzow, T.: Ocean Variability at Greenland’s Largest Glacier Tongue Linked to Continental Shelf Circulation, *Journal of Geophysical Research: Oceans*, 126, e2020JC017080, <https://doi.org/10.1029/2020JC017080>, 2021.
- Washam, P., Nicholls, K. W., Münchow, A., and Padman, L.: Summer surface melt thins Petermann Gletscher Ice Shelf by enhancing channelized basal melt, *Journal of Glaciology*, 65, 662–674, <https://doi.org/10.1017/jog.2019.43>, 2019.
- Wesche, C., Steinhage, D., and Nixdorf, U.: Polar aircraft Polar5 and Polar6 operated by the Alfred Wegener Institute, *Journal of large-scale research facilities*, 2, 1–7, <https://doi.org/10.17815/jlsrf-2-153>, 2016.
- 715 Wessel, B., Bertram, A., Gruber, A., Bemm, S., and Dech, S.: A new high-resolution elevation model of Greenland derived from TanDEM-X, in: XXIII ISPRS Congress, vol. III-7, pp. 9–16, <https://doi.org/10.5194/isprs-annals-III-7-9-2016>, 2016.
- Wessel, B., Huber, M., Wohlfart, C., Marschalk, U., Kosmann, D., and Roth, A.: Accuracy assessment of the global TanDEM-X Digital Elevation Model with GPS data, *ISPRS Journal of Photogrammetry and Remote Sensing*, 139, 171–182, <https://doi.org/10.1016/j.isprsjprs.2018.02.017>, 2018.
- 720 Wilson, N. and Straneo, F.: Water exchange between the continental shelf and the cavity beneath Nioghalvfjærdsbræ (79°N Glacier), *Geophysical Research Letters*, 42, 7648–7654, <https://doi.org/10.1002/2015GL064944>, 2015.
- Wilson, N., Straneo, F., and Heimbach, P.: Satellite-derived submarine melt rates and mass balance (2011–2015) for Greenland’s largest remaining ice tongues, *The Cryosphere*, 11, 2773–2782, <https://doi.org/10.5194/tc-11-2773-2017>, 2017.
- Zeising, O., Neckel, N., Steinhage, D., Scheinert, M., and Humbert, A.: Stake surface ablation/accumulation measurements at 79°N Glacier  
725 between 2017 and 2018, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.922131>, 2020.
- Zeising, O., Helm, V., Gupta, A., Steinhage, D., and Humbert, A.: Ice surface elevation and ice thickness at 79°N Glacier, Greenland from airborne measurements in July 2021, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.963752>, 2023a.
- Zeising, O., Steinhage, D., Neckel, N., and Humbert, A.: Time series of Lagrangian basal melt rates at 79°N Glacier since 2016, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.928903>, 2023b.
- 730 Zeising, O., Steinhage, D., Neckel, N., and Humbert, A.: Lagrangian basal melt rates at 79°N Glacier between 2017 and 2018, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.928541>, 2023c.