

INCHEM-Py v1.2: A community box model for indoor air chemistry

David R. Shaw¹, Toby J. Carter¹, Helen L. Davies¹, Ellen Harding-Smith^{1,2}, Elliott C. Crocker³, Georgia Beel^{1,4}, Zixu Wang¹, and Nicola Carslaw¹

¹Department of Environment and Geography, University of York, York, YO10 5NG, United Kingdom

²Wolfson Atmospheric Chemistry Laboratories, University of York, York, YO10 5DD, United Kingdom

³Department of Chemistry, University of York, York, YO10 5DD, United Kingdom

⁴UK Centre for Ecology and Hydrology, Edinburgh, EH26 0QB, United Kingdom

Correspondence: David R. Shaw (david.shaw@york.ac.uk)

Abstract. The Indoor CHEMical model in Python, INCHEM-Py, is an open-source and accessible box model for the simulation of the indoor atmosphere, and is a refactor and significant development of the INdoor Detailed Chemical Model (INDCM). INCHEM-Py creates and solves a system of coupled ordinary differential equations that include gas-phase chemistry, surface deposition, indoor/outdoor air change, indoor photolysis processes and gas-to-particle partitioning for three common terpenes. It is optimised for ease of installation and simple modification for inexperienced users, while also providing unfettered access to customise the physical and chemical processes for more advanced users. A detailed user manual is included with the model and updated with each version release. In this paper, INCHEM-Py v1.2 is introduced, the modelled processes are described in detail, with benchmarking between simulated data and published experimental results presented, alongside discussion of the parameters and assumptions used. It is shown that INCHEM-Py achieves excellent agreement with measurements from ~~two experimental campaigns~~ [an experimental campaign](#) which investigate the effects of ~~people and~~ different surfaces on the concentrations of different indoor air pollutants. In addition, INCHEM-Py shows closer agreement to experimental data than INDCM. This is due to the increased functionality of INCHEM-Py to model additional processes, such as deposition-induced surface emissions. [A comparative analysis with AtChem2 verifies the solution of the gas-phase chemistry.](#) Published community use-cases of INCHEM-Py are also presented to show the variety of applications for which this model is valuable to further our understanding of indoor air chemistry.

1 Introduction

In recent years, the quality of the air we breathe has gained increased attention. The World Health Organisation (WHO) recently stated that exposure to indoor and outdoor air pollution was one of the greatest risks to human health, and that improving air quality was necessary to reduce the global incidence and impact of diseases such as lung cancer, stroke and asthma (World Health Organisation, 2001). Much of the focus in developed countries to date has been on outdoor air quality, particularly in relation to attaining regulatory guidelines. However, exposure to air pollution indoors is arguably more important, given we are estimated to spend around 90% of our time indoors in high income countries like the United Kingdom (Klepeis et al., 2001).

Moreover, the recent COVID-19 pandemic has highlighted the importance of good indoor air quality, particularly around virus transmission.

25 Indoor air is a complex mixture of gas and particles, primarily as a result of numerous emission sources. These include activities such as cooking, cleaning, air freshener or scented candle use, household improvement activities (such as painting), emissions from building and furnishing materials (Uhde and Salthammer, 2007; Weschler, 2009), and from building occupants through breathing and interactions with compounds ~~in from~~ skin (Nazaroff and Weschler, 2004; Wisthaler and Weschler, 2010). Typical indoor pollutants include Volatile Organic Compounds (VOCs), nitrogen oxides (NO_x) and particulate matter (PM). In
30 addition, VOCs produced from indoor sources, or via ingress into buildings through windows, doors and cracks in the building fabric, can evolve over time through reactions with oxidant species, such as ozone (O₃) (Liu et al., 2021), hydroxyl radicals (OH) (Carslaw et al., 2017) and nitrate radicals (NO₃) (Arata et al., 2018). This secondary chemistry can result in the formation of a wide range of secondary products, such as formaldehyde and secondary organic aerosol (SOA), which can have a high potential for causing health issues, when compared to the parent VOC. For example, formaldehyde is a known carcinogen,
35 and SOA are ~~ultra-fine particles, which are~~ associated with a range of cardiovascular and respiratory conditions (Schraufnagel, 2020; WHO Air quality and health Guidelines Review Committee, 2010). Given that long-term monitoring of such processes has been limited to date, indoor air chemistry models provide a useful tool to aid further understanding, particularly of the evolution of secondary chemistry over time.

In recent years, there have been significant advances in our understanding of indoor air quality. This is due, in part, to
40 some larger-scale, intensive indoor measurement campaigns in test-houses, involving a suite of advanced instrumentation more typically used in outdoor field campaigns (see the overview in Farmer et al. (2019)). The results from these campaigns have demonstrated that indoor air quality is a complex, multidimensional problem, where indoor and outdoor sources, transformation processes, and building design, management and use are the driving physical and chemical factors for air quality. Human behaviour then adds further complexity. Ideally, one would perform numerous measurements in numerous buildings and for
45 long periods of time to understand these processes. However, such a task would be expensive, time consuming and logistically challenging. For instance, how would representative buildings be selected? An alternate approach is to simulate these processes in a model, coupled with evaluation using experimental data. It is then possible to use the evaluated model to provide forecasts and to explore hypothetical scenarios not possible through measurements, thus providing a deeper insight into the processes of interest. ~~To achieve this goal requires a thorough understanding of the fundamental processes that underpin indoor air quality~~
50 ~~in buildings.~~

There are numerous processes that need to be considered in a model of indoor air quality, these include direct and secondary emissions from surfaces (building materials, furnishings and people), physical building parameters (such as ventilation rates, temperature, humidity and light), gas- and particle-phase chemistry, surface deposition, and the effect of occupants. These factors will combine to determine the occupant exposure to air pollution under any set of building conditions. Note that
55 microbial emissions also play a role in occupant exposure to indoor air pollution, but they have not been considered in this work.

The INdoor CHEMical Model in Python (INCHEM-Py) is a zero dimensional community box model that includes many of these processes to enable investigation of the evolution of indoor air chemistry over time. It is a refactor and improvement of the INdoor Detailed Chemical Model (INDCM) developed by Carslaw (2007) ~~and is the most detailed chemical model that~~
60 ~~currently exists for exploring the chemistry of indoor atmospheres. It allows the user to understand the change in concentrations~~
~~of indoor species over time and the key reaction pathways that describe the formation, transformation and loss of indoor species.~~
~~Processes included within INCHEM-Py have been developed and implemented as our understanding of the indoor environment~~
~~has improved, when experimental data have supported new developments, and according to the the research questions being~~
65 ~~asked. The model is continuously being improved, this paper gives a snapshot of the current version of the model while also~~
~~defining the foundation upon which future model developments will build upon. INCHEM-Py~~ v1.1 has been described in basic
detail, covering accessibility and broad function in Shaw and Carslaw (2021). This paper describes the model in more detail,
including the developments included in the latest version (v1.2), and gives some examples of the ways in which it can be used
by the scientific community.

2 Model description

70 INCHEM-Py creates and solves a series of first order Ordinary Differential Equations (ODEs), with each equation representing
the rate of change of a species concentration (molecule cm^{-3}) with time. The general equation for these ODEs is shown for
the concentration (C) of species i in Eq. (1).

$$\frac{dC_i}{dt} = \sum R_{ij} + (\lambda_r C_{i,out} - \lambda_r C_i) - \sum \nu_d \left(\frac{A}{V} \right) C_i \pm k_t C_i \quad (1)$$

The chemical mechanisms are represented with the first term on the right hand side of Eq. (1) as a sum of all reaction rates
75 involving species i , with j representing other species. Each reaction between species takes place at a rate, R (s^{-1}), dependent
on the concentrations of the reactant species and the rate constant. The second term of each ODE accounts for indoor/outdoor
exchange, which depends on the air change rate, λ_r (s^{-1}), and the species concentrations both indoors (C_i (molecule cm^{-3}))
and outdoors ($C_{i,out}$ (molecule cm^{-3})). The third accounts for surface deposition, dependent on the species deposition velocity,
 ν_d (cm s^{-1}), and surface to volume ratio (A/V) of multiple surfaces (A/V (cm^{-1})), while the final term is for timed emissions,
80 where k_t (s^{-1}) is the emission/loss rate of species i at time t .

Rate coefficients and reactions are imported in the most part from the Master Chemical Mechanism (MCM) (Jenkin et al.,
1997, 2003, 2015, 2018, 2020; Saunders et al., 2003; Bloss et al., 2005), an almost explicit chemical mechanism following the
atmospheric degradation of 135 VOCs. Additional schemes and mechanisms for species that are not included in the MCM, but
which do occur indoors, are parsed from an additional file. Schemes have been developed for some terpenoids that are present
85 in many scented items found and used indoors (Carslaw, 2007, 2013; Carslaw et al., 2012, 2017; Terry et al., 2014). Chlorine
schemes have also been designed and implemented, since it is a key pollutant from bleach cleaning products (Wong et al.,
2017). The full list of additional schemes is available in the user manual. The model also ~~includes the function for~~ ~~allows~~ users
to add their own chemical mechanisms to supplement those already included, ~~thereby allowing or to use a mechanism other~~




Figure 1. Flow chart showing the order of operations within INCHEM-Py. The modules and parameters within the dashed box are solved within the function that is passed to the integrator and are therefore called/calculated at every integrator internal time step.

than the MCM (although any new mechanisms need to be in the same format as that adopted in INCHEM-Py). Such flexibility will facilitate reaction mechanism development, and community-driven model development.

All species within the model are assumed to egress with a specified air change rate (λ_r) as a function of their indoor concentration (C_i). Some species also ingress at the same air change rate, depending on their constant or diurnal outdoor concentrations ($C_{i,out}$) as discussed in Sec. 2.3. Species are also assumed to irreversibly deposit to surfaces at a rate dependent on individual deposition velocities (ν_d , mostly taken from Carslaw et al. (2012)) and the surface area to volume ratio (A/V) of the simulated space, as in Carslaw (2007). An optional mechanism for O₃ and H₂O₂ surface deposition and subsequent emission has been developed with a detailed discussion in Sect. 2.8. ~~Some emissions from surfaces are included in the new surface mechanisms, but additional emissions.~~ Additional emissions, both constant and intermittent, can be user defined. ~~The maximum Gas-to-particle partitioning reactions are included for limonene, α -pinene and β -pinene (discussed in Sect. 2.4).~~ The total size of the chemical mechanism solved by INCHEM-Py (v1.2), before any additional user mechanisms are added, is 6507 chemical species and particles undergoing 19581 reactions. Figure 1 shows how input parameters are utilised within the modules of INCHEM-Py, and how they produce the system of equations and parameters required to predict the evolution of indoor atmospheric constituents with time.

2.1 ODE formation and solution

All gas-phase reactions solved in INCHEM-Py are of the general form

$$C_i + \dots + C_j \xrightarrow{k} C_x + \dots + C_y \quad (2)$$

where C_i remains the concentration of an individual species and k is the rate constant of the reaction, which may be constant, or depend on other variables such as photolysis. In each case, the concentration of species on the right hand side of the equation are increased and the species on the left hand side (lhs) of the equation are decreased at the same rate (R) of

$$R_{ij} = \sum \left(\prod C_{lhs} \right) k \quad (3)$$

110 where k will be negative for the reactants and positive for the products. Therefore, by calculating $R_{\pm R}$ for each species in each reaction, the total change in an individual species concentration only through gas-phase reactions is given by the sum of all of the individual reaction rates, as follows:

$$\frac{dC_{i,Gas}}{dt} = \sum R_{ij} \quad (4)$$

~~Particles are discussed in Sect. 2.4, but the partitioning equations follow the same form as Eq. (2) and are, therefore, implemented into the ODEs in Stoichiometries are dealt with in the same way, with an ODE for each particle concentration.~~
 115 MCM export by repeating the species within the reaction, i.e. the photolysis reaction $H_2O_2 = 2OH$ is represented as $H_2O_2 = OH + OH$.
This method is used throughout INCHEM-Py. Each species with the potential to partition to the particle-phase has a corresponding term that tracks the change in its particle-phase concentration over time. This change in concentration for each partitioning species is determined by the equilibrium between the absorption to and desorption from the particle bulk based on its individual
 120 properties (Carslaw, 2007). Therefore, the particle-phase concentration of each species is represented by its own ODE in the system and individual species contributions to total particle concentration can be output (particles are discussed in detail in Sect. 2.4).

The ODEs for each species are solved simultaneously using integrate.ode from the Scipy Python library and the wrapped LSODA function from the Fortran solver package ODEPACK (Hindmarsh, 1983). LSODA was chosen as the solution method
 125 as it automatically switches between the implicit Adams-Moulton formulae for non-stiff problems and Backward Differentiation Formulae (BDF) for stiff problems. Due to the size of the system, and its highly-coupled structure, the problem is stiff in most cases (note that if using only a small selection of the MCM mechanism it may be non-stiff in some instances). LSODA makes use of the Jacobian of the system to ease concentration predictions and is calculated in full as an array of partial differential equations. ~~Representing Eq. (1) as $f(t, C_i)$ the (i,j)th entry in the Jacobian is~~

130 $J_{ij} = \frac{\partial f_i}{\partial C_{S_j}}$

by INCHEM-Py. The time step between outputs is set by the user in the settings file. The integrator will predict and output the solution to the ODE system at each time step. Default integrator parameters are given in the user manual and can be user adjusted. Verification of the solution of the gas-phase photochemistry is shown in Sec. (3.1).




Figure 2. Relationship between declination angle (Dec), latitude (Lat) and solar zenith angle (θ). C is the centre of the Earth, the line CS is a line connecting the centre of the Earth to the centre of the Sun and Z represents the zenith of the location being simulated.

2.2 Photolysis

135 Time is given in local solar time in INCHEM-Py and determined using the date and latitude of the simulation. A date input is used to calculate the solar declination angle (Dec) as

$$\text{Dec} = -23.45 \times \cos\left(\frac{360}{365.25} \times (d + 10)\right) \quad (5)$$

where d is the number of days since the 1st of January that year, -23.45 ($^{\circ}$) is the solar declination angle at the winter solstice, 360 ($^{\circ}$) is a complete angle, 365.25 is the number of days in the year and 10 is the approximate number of days between the winter solstice and January 1st. Figure 2 shows the relationship between the declination and the solar zenith angle, θ , which is used to calculate the outdoor photolysis rates. In Figure Fig. 2, CS represents a line connecting the centre of the Earth (C) with the centre of the Sun, Z represents the zenith of the location of the simulation, Lat is the latitude of the location of the simulation, and LHA is the local hour angle which is 0 at solar noon. Using these known values and the spherical law of cosines the solar zenith angle is calculated as

$$\cos(\theta) = \sin(\text{Lat}) \sin(\text{Dec}) + \cos(\text{Lat}) \cos(\text{Dec}) \cos(\text{LHA}) \quad (6)$$

The solar zenith angle is used to calculate the solar photolysis portion of the overall photolysis rate (J), given by Eq. (7), for the 43 photolysis reaction rates currently included within the model. A description of the scattering method employed for this calculation is taken from Hayman (1997) and utilised in Saunders et al. (2003), where parameters l , m and n are optimised as per the discussion in Jenkin et al. (1997). These values are then attenuated according to the glass type chosen by the user, through the use of a transmission factor (ψ).




Figure 3. Indoor OH concentrations as a function of (a) latitude and (b) time of year. In both cases, only one setting is varied from default, with the latitude variation being on [21/06/2020-21-06-2020](#) and the seasonal variation being at 45° N. [Note that \$1 \times 10^6\$ molecule \$\text{cm}^{-3}\$ is approximately 0.04 ppt.](#)

Indoor photolysis rate values (ϕ) are available for seven light types, and these are ~~are~~ summed with the attenuated outdoor values to give the total ~~indoor~~ photolysis rate as shown in Eq. (7), ~~where parameters l , m and n are optimised as per the discussion in Jenkin et al. (1997).~~

$$J = ((l \cos(\theta))^m \exp(-n \sec(\theta)) \psi) + \phi \quad (7)$$

- 155 Values for ψ and ϕ are calculated and discussed in [Wang et al. \(2022\)](#) [Wang et al. \(2022\)](#). In brief, for the transmission factor, ψ , three typical glass types were selected from the analysis in Blocquet et al. (2018), covering a range of transmission values in both the UV (300 - 400 nm) and visible (400 - 800 nm) spectral ranges. The wavelength ranges were split into 10 nm intervals and the percentage of light transmitted through each glass type and for each wavelength interval was defined. The absorption cross-section and quantum yields for each photolysing species ~~are~~ were used to calculate the weighted transmission
- 160 [factor, \$\psi\$, for each wavelength range as described in detail in Wang et al. \(2022\). The weighted transmission factor accounts for the proportion of the wavelength range transmitted by a particular glass type that is absorbed by each individual molecule.](#) For the indoor photolysis rate, ϕ , the same wavelength ranges, intervals, quantum yields and cross-sections were used, with spherically integrated photon fluxes from Kowal et al. (2017). Seven indoor light types are included in INCHEM-Py, covering a range of transmission spectra. The lights can be set to be on or off by the user for any period, over single or multiple days.
- 165 Details of the specific lights simulated are found in Kowal et al. (2017); Wang et al. (2022).

The impact of the different photolysis parameters is shown in Fig. 3 and Fig. 4, where latitude and date settings are varied while all others remain at their default values ~~(given in appendix ??)~~. Figure 3(a) shows the effect of latitude variation in the northern hemisphere in summer on indoor OH concentrations. In moving away from the equator, the length of the days increases, but the intensity of sunlight decreases. Therefore, this interplay results in peak indoor OH concentrations at mid




Figure 4. Indoor OH concentrations as a function of (a) indoor light source type and (b) glass type. All other values remained at their default settings.

170 latitudes. At the northern extremes, the Sun does not set, therefore, OH is produced through indoor photolysis reactions for the full 24-hour period but at a reduced rate due to the lower intensity of solar radiation. Figure 3(b) shows the seasonal change of indoor OH in the northern hemisphere. ~~As the outdoor concentrations of species are identical in each simulation the OH concentration difference is solely from photolysis. Reactant concentrations which form OH through photolysis are therefore very similar, resulting in comparable peak concentrations between the simulations~~ Although the outdoor concentration of OH varies with latitude and season, its lifetime is too short for it to ingress indoors. OH production indoors is driven by the reaction of HO₂ with NO in this instance, so the OH profile indoors is driven by the NO concentration. Therefore the tails of OH concentration in Fig. 3(b) follow the same minimum profile, only deviating with photolysis input. This minimum profile can be seen in Fig. 4(b) as the "No sunlight" trace.

180 Figure 4(a) shows the minimal impact of indoor lighting on the concentration of OH indoors, which contrasts with the large effects seen when the glass type is varied in Fig. 4(b). The different glass types impact the production of OH indoors, mainly through the photolysis of HONO and the reaction of HO₂ and NO. Glass C lets through the most sunlight and, therefore, allows the most photolysis reactions to occur. At the other end of the scale is the low-emissivity glass with a reflective film that blocks out the majority of wavelengths and is very close to not having any sunlight enter the room at all. Without sunlight there is very little OH produced by the indoor lighting, which is set by default to come on at 7am and go off at 7pm.

185 2.3 Outdoor air exchange

In INCHEM-Py only indoor species are predicted, and in some cases their concentrations will depend heavily on influx from, and efflux to, outdoors (Kruza et al., 2021). All indoor species are set to decay at a rate of $\lambda_r C_i$ and to increase at a rate of $\lambda_r C_{i,out}$ as in Eq. (1), however only species for which measured, representative outdoor concentrations are available, are as-

signed an outdoor concentration in the model. All other species (mostly short-lived intermediates) have outdoor concentrations set to zero.

Concentrations of OH, HO₂ and CH₃O₂ outdoors are photolysis driven and typically show strong diurnal variation outdoors. As in Carslaw (2007), the outdoor concentrations are set to peak at solar noon in the model, at 5×10^6 , 1×10^8 and 2.5×10^7 molecules cm⁻³, respectively, in line with measurements taken by Platt et al. (2002) and Emmerson et al. (2005). HONO shows the opposite trend, as it is photolysed during the day and is set to a minimum of ≈ 20 ppt at noon, peaking overnight at ≈ 300 ppt (Alicke, 2003).

NO, NO₂ and O₃ have four possible diurnal profiles which can be chosen by the user, depending on their specific requirements. Profiles are provided for urban London (UK), suburban London (UK), and urban Bergen (Norway) locations, based on measured data from 2018 in the EEA (2018) air quality database. Details of these sites and their exact identifiers, including latitude and longitude, are given in the INCHEM-Py user manual. The data from each site were provided as hourly averages and in local time (UTC) by the meteorological station. Using the station longitude, each data point was shifted to solar time and quarter three (Q3) data extracted. Q3 (July, August, September) was chosen because not all data sets had annual data for all species. The daily measurements were then overlaid and an average data point for each hour was used to fit a trigonometric Fourier function for each location and species. The same process was used for outdoor concentrations of PM_{2.5}, which can also be included within the model when the particle module is enabled. The fourth location is Milan (Italy) which is included from Terry et al. (2014) as a particularly polluted two-week period in August 2003. The measurements taken underwent the same procedure of averaging and fitting as the other three locations. The functions that have been input into the model are shown in the user manual. For INCHEM-Py v1.2 the constant outdoor concentrations have been updated, as shown in Table A1. These values have been sourced from published literature and measurement databases as referenced.

The outdoor concentrations used by INCHEM-Py can be adjusted to fit the requirements of the user. The default profiles of NO, NO₂ and O₃ from four European cities in the summer are provided as indicative locations with sufficient data to create the required diurnal profiles from the EEA (2018) air quality database. Further discussion on how the original profiles were created, how they may be adjusted, and how to create additional profiles is provided in the user manual.

The rate of air change (ACR) is a user defined variable that is input in units of acs⁻¹ (air changes per second), and will depend on how airtight the space being simulated is and any intentional ventilation. Weschler (2000) gives typical values of between 0.2 and 2 ach⁻¹ (air changes per hour) for ~~loosely-constructed and tightly~~ tightly constructed and loosely constructed residential properties respectively. The default value for INCHEM-Py is 1.38×10^{-4} acs⁻¹ (0.5 ach⁻¹). How the air change rate ~~effects~~ affects indoor OH concentrations (with all other values set to default) is shown in Fig. 5(a).

~~In general, increased ACR results in increased OH concentrations. The lifetime of OH is too short to survive ingress through a building and any OH indoors is made indoors through chemical reaction.~~ Through the analysis of the reaction rates, as detailed in Sect. 2.7, OH is mainly produced through HO₂ + NO → OH + NO₂, and the photolysis reaction HONO → OH + NO, where HO₂ is the hydroperoxy radical that is primarily produced via the oxidation of VOCs. Most indoor NO comes from the photolysis reaction NO₂ → NO + O in the absence of indoor sources, and NO₂ is mainly produced by the reaction of NO + O₃ → NO₂. In general, increased ACR results in increased OH concentrations, but the lifetime of OH is too short to




Figure 5. Indoor (a) OH and (b) NO concentrations as a function of air exchange rate. All other settings are the default values with the city outdoor concentration profile set to urban Bergen.

225 survive ingress through a building and most OH indoors is made indoors through the above chemical reactions. The influx of outdoor OH is less than 0.009 % of the total production rate of OH at 0.5 ach⁻¹. NO, NO₂ and O₃ are also sourced from outdoors, so as the ACR increases, their concentrations increase, as do the rates of the reactions that produce higher OH as described above. Increasing ACR will also increase the loss rate of OH to outdoors, however, due to its reactivity this is less than 0.0025 % of the total loss rate of OH at 0.5 ach⁻¹.

230 In the 0.2 ach⁻¹ case, there is a small peak in OH concentration at 20:00 which is not seen for the other ACRs. Analysing the reaction rates revealed that as ACR is reduced, indoor O₃ concentrations are reduced (as less is available to ingress from outdoors), which decreases the loss of NO through the reaction with O₃. This reduced NO consumption results in increased NO concentrations later in the day, as shown in Fig. 5(b), compared to the higher ACR cases. Increased NO concentrations increase the rate of the NO + HO₂ → OH + NO₂ reaction relative to the other simulations, therefore, causing the peak in OH concentrations later in the day in the low ACR case.

235 2.4 Particles

~~Gas-to-particle partitioning is included within INCHEM-Py for~~ includes gas-to-particle partitioning for the oxidation schemes of limonene, α -pinene and β -pinene. Carslaw et al. (2012) provide, producing 610 species that partition between the gas- and aerosol-phases. Carslaw et al. (2012) provides a full description of the method used to calculate the particle partitioning parameters, which is based on absorptive partitioning from Pankow (1994), whereby the phase of the species is determined by thermodynamic equilibrium. ~~It is described briefly here as implemented in the model~~ We assume that the particles are all in the PM_{2.5} range and focus only on chemical composition, with no representation of how these particles evolve dynamically over time.

The method relies on a balance between the rate of VOC absorption to (k_{on}) and the rate of desorption from (k_{off}) particles. The partitioning coefficient, K_p constant, K_p ($m^3 \mu g^{-1}$) is the gas:particle ratio for a species.

$$K_p = \frac{k_{on}}{k_{off}}$$

This can be calculated using k_{on} and k_{off} for each species that partitions to the particle phase as

$$K_p = \frac{7.501RT}{W_{om}10^9P_l} \quad (8)$$

where R is the ideal gas constant ($8.314 \text{ J K}^{-1} \text{ mol}^{-1}$), T is the temperature (K), W_{om} is the mean molecular weight of the particle (g mol^{-1}) and P_l is the liquid vapour pressure of the species (Torr) (Pankow, 1994; Carslaw et al., 2012).

k_{on} is set at a temperature independent value of $6.2 \times 10^{-3} \text{ m}^3 \mu \text{g}^{-1} \text{ s}^{-1}$, which is based on Jenkin (2004) and Johnson et al. (2006). Rearranging Eq. (9) and substituting Eq. (8) allows the determination of k_{off} for each species. W_{om} is set to 120 g mol^{-1} initially, but is calculated for each subsequent integration step based on the composition of the formed particles. Assuming equilibrium conditions exist (Leungsakul et al., 2005), Eq. (9) allows the determination of k_{off} for each species.

$$K_p = \frac{k_{on}}{k_{off}} \quad (9)$$

The rates k_{on} and k_{off} are used to track each species partitioning to and from the particle phase and are solved in the ODE alongside the gas-phase reactions. Each species is tracked individually in the particle phase, and summed to produce a total number ($\text{tsp, molecule cm}^{-3}$) and concentration ($\text{tsp, } \mu \text{g m}^{-3}$) of suspended particles.

This method assumes particles already exist within the model for the initial absorption reactions to take place. A small number of seed particles are present within the model, set within the initial conditions file as "seed". These seed particles do not change their number density but allow for partitioning to take place. Particles can also enter from outdoors with the outdoor particles assumed to be 30 % organic and 70 % inorganic. The organic fraction from outdoors is then also used as a seed on which new indoor particles can form. Each species is tracked individually in the particle phase, and summed to produce a total number ($\text{tsp, molecule cm}^{-3}$) and concentration ($\text{tsp, } \mu \text{g m}^{-3}$) of suspended particles. Particles are set to irreversibly deposit onto surfaces at a rate of 0.004 cm s^{-1} (Sarwar and Corsi, 2007).

2.5 Temperature

Most chemical reactions in INCHEM-Py have a temperature dependence. Indoor temperatures in most scenarios will have minimal variation but in some cases might vary throughout a day. The model allows for three methods of setting the indoor temperature: a constant value, a linear interpolation between given temperatures, or a zero-degree B-Spline interpolation between given temperatures. For both interpolation methods, a minimum of two temperatures at two distinct times must be given. INCHEM-Py compares the given times to the length of the simulation and duplicates points before and after the simulated period, if required. This ensures that there is an interpolated or given temperature value at all time points. Full details of the methods used are given in the INCHEM-Py user manual. Three different temperature profiles using the three different methods




Figure 6. Results of the three different temperature input methods on (a) temperature and (b) OH concentration. Only the "spline" variable was adjusted from default values, the constant temperature method and value is the default option.

are shown in Fig. 6, alongside their impact on the OH concentrations. Both interpolated methods used the same times and temperature inputs.

275 Temperature is a variable that changes the rate coefficients of many of the reactions in INCHEM-Py: some will increase with temperature, while others will decrease. Although the three temperature profiles are in good agreement around the middle of the day, there is a large difference earlier on given how the three methods work. This difference in methods leads to different concentrations for some of the precursors for OH, allowing them to have a differential impact on OH chemistry later, depending on their lifetimes. The combination of the varying reaction rates and resultant concentrations produce the final profile in Fig.
 280 6(b).

2.6 Timed emissions

Term four in Eq. (1) accounts for user-defined timed emissions that can be input to simulate a release of a chemical species over a specific period of time. Only species that are solved by the model can be input, these include chemical and particle species. Outdoor species cannot be input this way, as outdoor concentrations are not solved, but instead are input as time dependent
 285 functions or constants which can be adjusted in the outdoor concentrations file. An example of this function is shown in Fig. 7, where an emission of limonene and β -pinene simulates the use of a cleaning product. Both species were input at a rate of 5×10^8 molecule $\text{cm}^{-3} \text{s}^{-1}$ (~~2 ppt~~0.02 ppb s^{-1}) for 10 minutes at 1 pm, while all other settings remained at their default values. Figure 7(a) shows that, the terpene concentrations increase at the same rate and peak at very similar values. The small differences in the subsequent terpene decays, is due to their different reactivities. For example, the reaction rate coefficients at
 290 293K for limonene with O_3 ($k_{\text{O}_3} = 2.0 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) and OH ($k_{\text{OH}} = 1.7 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) are higher than for β -pinene ($k_{\text{O}_3} = 1.8 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, $k_{\text{OH}} = 8.1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) (Jenkin et al., 2018). This explains the lower peak concentration, and the faster decay, for limonene, compared to β -pinene.




Figure 7. (a) Limonene and β -pinene timed emissions as a simulated cleaning event with (b) formaldehyde secondary emission compared to background default values.

Figure 7(b) shows the secondary production of formaldehyde following the timed emission of limonene and β -pinene, compared with a background default value with no emissions. Using this method, we can simulate events where multiple chemical species interact, to inform what proportions of an experimentally measured chemical are produced by primary emission or via secondary chemistry.

2.7 Reaction rate outputs

When analysing chemical transformations it is useful to know the rate at which a species is being produced or lost, and the relative importance of the individual reactions that are contributing to that rate. INCHEM-Py has an option to output the rates of all reactions at all time points. Using the rate coefficients, and the calculated species concentrations, the reaction rates of individual reactions can be calculated at any time point in the simulation using Eq. (3). This function was developed for, and used by, Lakey et al. (2021) to identify important reactions and develop a reduced chemical mechanism for use in an indoor 3D fluid model.

In the case of the cleaning event shown in Fig. 7, the reaction rates from INCHEM-Py at each time step can be used to track the pathways linking the primary pollutants (limonene and β -pinene) to the secondary production of formaldehyde. This is visualised in Fig. 8, where a snapshot of the reaction pathways at the peak of the pollutant concentrations is shown. This method was used to analyse the contribution of surface cleaner formulations to indoor pollutants in Carslaw and Shaw (2022).

2.8 Surface Deposition

Surface deposition of gas-phase species is an important aspect of indoor air chemistry, and is key to include in indoor air models. Chemical species can be emitted from surfaces, either as primary emissions, or as secondary pollutants formed from




Figure 8. A flow chart showing the degradation pathways of limonene (orange) and β -pinene (blue) to final secondary products of interest (green). The fastest 95% of reactions occurring at the time point where their concentrations are highest in Fig. 7 are shown. The thickness of the line is proportional to the absolute reaction rate, where the thicker the line, the faster the reaction. Nodes show species created, and all species are labelled in MCM format.

gas-phase transformations which occur at surface level. O_3 can deposit onto a range of surfaces and induce oxidation, releasing secondary pollutants as surface emissions (Gall et al., 2013; Cros et al., 2012). The rate of deposition to surfaces is surface-specific, and is determined by mass transportation to the surface of a pollutant, and the uptake potential of the pollutant onto that specific surface (Reiss et al., 1994). The deposition rate of an oxidant onto a surface is also influenced by the air change rate, the bulk indoor concentration of the oxidant, and the surface-to-volume ratios in the indoor environment (Coleman et al., 2008).

INCHEM-Py v1.1 simulates the irreversible deposition of 3371 indoor gas-phase species. However, it does not incorporate the secondary pollutants emitted from the surface. It also does not consider different surface materials, and only calculates surface deposition based on estimated deposition velocities and a total surface to volume ratio (Carslaw, 2007).

320 [A-These are retained in v1.2 alongside a new surface deposition mechanism onto multiple surfaces that has been developed for INCHEM-Py v1.2, based on the work of Kruza et al. \(2017\) and considering the rates of deposition and secondary pollutant emissions following individual \$O_3\$ and \$H_2O_2\$ deposition \(Carter et al., 2023\). Given that the nature of such interactions is likely to be complex and currently not fully understood we consider the process occurs in 2 steps: deposition to the surface and loss of the oxidant, followed by emission of new species from that surface.](#) The surface removal rate of O_3 and H_2O_2 in the model is determined by Eq. (10) and Eq. (11), respectively:

$$F_{O_3} = \nu_{d_{O_3}} \frac{A}{V} \quad (10)$$

$$F_{H_2O_2} = \nu_{d_{H_2O_2}} \frac{A}{V} \quad (11)$$

where F_{O_3} and $F_{H_2O_2}$ represent the deposition rates (s^{-1}) of O_3 and H_2O_2 onto a surface, respectively. ν_d represents the surface deposition velocity of an oxidant in $cm\ s^{-1}$, A is the surface area of an indoor surface or material in cm^2 and V is the total volume of the indoor environment in cm^3 .

The emission rate calculations of secondary pollutants emitted as a result of oxidant surface deposition has been adapted from studies conducted by Morrison and Nazaroff (2002) and Kruza et al. (2017). The emission rates can be determined by solving Eq. (12) and Eq. (13):

$$E_i = \frac{A\nu_{dO_3}Y_iC_{O_3}}{V} \quad (12)$$

$$E_i = \frac{A\nu_{dH_2O_2}Y_iC_{H_2O_2}}{V} \quad (13)$$

where E_i represents the emission of secondary pollutants from a surface ($molecule\ cm^{-3}\ s^{-1}$). Y is the production yield of gas-phase species following deposition (dimensionless), and C_{O_3} and $C_{H_2O_2}$ represent the bulk concentrations of indoor O_3 and H_2O_2 , respectively ($molecule\ cm^{-3}$).

Surfaces included in the O_3 and H_2O_2 surface deposition mechanisms for INCHEM-Py v1.2 are soft fabric, painted, human skin, wood, metal, concrete, paper, plastic and glass. However, there are no data for H_2O_2 onto plastic, glass and skin surfaces. The mechanisms were constructed using surface specific deposition velocities of O_3 and H_2O_2 and respective production yields from a range of experimental literature (Sabersky et al., 1973; Lin and Hsu, 2015; Klenø et al., 2001; Grøntoft, 2002; Abbass et al., 2017; Gall et al., 2013; Tamás et al., 2006; Cros et al., 2012; Coleman et al., 2008; Ye et al., 2020; Lamble et al., 2011; Rim et al., 2016; Poppendieck et al., 2007; Wang and Morrison, 2010, 2006; Nicolas et al., 2007; Morrison and Nazaroff, 2000; Fadeyi et al., 2013; Yao et al., 2020; Di et al., 2017; Rai et al., 2014; Fischer et al., 2013; Wisthaler and Weschler, 2010; Schripp et al., 2012; Mueller et al., 1973; Cox and Penkett, 1972; Grøntoft and Raychaudhuri, 2004; Simmons and Colbeck, 1990; Cano-Ruiz et al., 1993; Poppendieck et al., 2021). These mechanisms ensure secondary species, primarily aldehydes, are emitted from specific surfaces as a result of oxidant deposition. A deposition velocity taken from Carslaw (2007) has been added for plastic, glass and skin surfaces to account for gas-phase deposition of H_2O_2 . Deposition of H_2O_2 onto these three surfaces do not produce any aldehyde emissions in our model.

A detailed description of the oxidant surface deposition mechanisms is found in a corresponding study conducted by Carter et al. (2023). Using these methods, further deposition mechanisms can be developed for other species and surfaces in the future, as relevant experimental data ~~become~~becomes available.

Since the Carter et al. (2023) study, the deposition module for INCHEM-Py has been updated to include O_3 deposition onto linoleum surfaces, using data from Kruza et al. (2017). The deposition velocity of O_3 onto plastic surfaces has also been updated as a result of an ongoing review of the literature. The deposition velocities of O_3 onto linoleum and plastic surfaces are now 0.0070 and $0.1225\ cm\ s^{-1}$ respectively (Coleman et al., 2008; Kruza et al., 2017; Klenø et al., 2001; Poppendieck et al., 2007; Nicolas et al., 2007; Wang and Morrison, 2006).

Table 1. Rates of breath and skin emissions for adults and children

Emitted species	Adult rate (molecule $\text{cm}^{-3} \text{s}^{-1}$)	Child rate (molecules $\text{cm}^{-3} \text{s}^{-1}$)
Acetone	2.534×10^7	4.781×10^6
Ethanol	1.988×10^7	3.009×10^6
Methanol	8.512×10^6	3.108×10^6
Isopropanol	3.862×10^6	6.593×10^5
Isoprene	5.412×10^6	5.953×10^5

2.9 Direct emissions

Breath emission values from humans are optionally included in the model, according to occupancy status (Kruza and Carslaw, 2019; Weschler et al., 2007). The number of adults and children can be specified in the settings file and a calculated emission in molecule $\text{cm}^{-3} \text{s}^{-1}$ is used, as shown in Table 1. These emissions are constant for the duration of the simulation.

365 3 Model output and evaluation

In this section, the INCHEM-Py model is used to simulate previously published results from both experimental and modelling efforts. Settings and outputs from all of the model runs are linked in the data availability section of this paper.

3.1 ~~Skin-oil emissions~~ Gas-phase verification

~~In Kruza and Carslaw (2019), the experimental results showing 4-oxopentanal (4-OPA) production from human skin from Weschler et al. (2007) and Wisthaler and Weschler (2010) were simulated using the precursor to INCHEM-Py, the INDCM. The experiment involved two adults (without showering or having used personal care products) being placed in a 28.5 m³ chamber with a high level of O₃. Concentrations of 4-OPA produced from surface reactions on human skin and clothing were then measured. The original work used units of ppb so outputs from 2). To validate the solution of this core component of the model INCHEM-Py were~~
375 ~~duly converted for this comparison.~~

~~Two simulations were run in was set up to compare with AtChem2 (Sommariva et al., 2020). AtChem2 is an outdoor model that also utilises the MCM. INCHEM-Py was set up by disabling the features that are not replicatable easily in AtChem2, including; the additional chemistry developed specifically for INCHEM-Py. The first was optimised to achieve a constant background O₃ concentration of 32 ppb to match that measured by Wisthaler and Weschler (2010). As the experimental chamber was located inside a laboratory, the diurnal outdoor concentrations were disabled in the model, and an emission of O₃ was created. The experimental chamber, a former aircraft cabin, had painted plastic walls and a carpeted floor giving surface area to volume ratios of 0.004 cm⁻¹ for soft furnishings and 0.0158 cm⁻¹ for painted walls. An emission of 9×10^8 molecule $\text{cm}^{-3} \text{s}^{-1}$; particle processes; air change; indoor lighting and attenuation of outdoor light; and surface deposition and emission. A~~
380

simulation was prepared in each model of a two day period representing 21-06-2020 at 45 °N, with no additional emissions. The percentage deviation ($\sigma_{i,t}$) for i at time t between the two models was calculated by Eq. (14), taken from O'Meara et al. (2020), where it was similarly used to validate gas-phase chemistry between AtChem2 and PyCHAM.

$$\sigma_{i,t} = \left(\frac{s_{i,t} - b_{i,t}}{v(b_i)} \right) 100 \quad (14)$$

In Eq. (0.037 ppb s⁻¹) of O₃ was needed to maintain a 31.53 ppb O₃ concentration with an air change rate of 1 h⁻¹. To match the experimental conditions, the simulated chamber was set to have constant indoor lighting from incandescent bulbs and no sunlight, the temperature was set to 296.15 K, and the relative humidity was 23 %.

With the background simulation optimised, a second simulation was run using the background simulation outputs as initial concentrations. From 10 am to 2 pm, two humans with a combined surface to volume ratio of 0.0014 cm⁻¹ (assuming a skin area of 2 m² per person) and associated breath emissions (according to Tab. 1), were simulated to be present in the room. The decrease in O₃ and production of 4-OPA from the skin surface chemistry from experiment and 14) the two models is shown in Fig. ???. The INCHEM-Py model output shows good agreement with the experimental results with the O₃ decreasing at the same rate but not for quite as long. When comparing the results from INCHEM-Py and the INDCM, the O₃ concentrations share the same shape, but due to the improved surface mechanism, a balance between O₃ deposition and 4-OPA yield results in much closer alignment to the experimental data. The 4-OPA measurements increase more slowly than the INCHEM-Py concentration, which is probably a result of the lack of spatial representation in result is represented by s and the model. Comparison of experimental (Wisthaler, grey lines) and simulated results (INCHEM-Py, black lines, and Kruza, yellow lines) of 4-OPA production from skin and clothing due to O₃ surface chemistry. AtChem2 result represented by b . $v(b_i)$ is the maximum value of component i and scales the difference between models to the individual component being compared.

With no diurnal variation in Figure 9(a) shows the calculated differences for key species. INCHEM-Py simulation, the ratio of O₃ to 4-OPA will eventually reach equilibrium, determined by the surface to volume ratio, the surface deposition velocity, and the 4-OPA yield from O₃ surface interactions as performs well when compared to AtChem2, with the largest differences being during rapid increases or decreases in species concentration. The deviation is due to the calculation of photolysis parameters in Python based INCHEM-Py vs the calculations in Fortran based AtChem. Values for Dec , $\cos(Lat)\cos(Dec)$ and $\sin(Lat)\sin(Dec)$ are identical to the thousandth between the two simulations propagating a small difference in the calculation of $\cos(\theta)$ in Eq. (12). Therefore, using 6). The larger deviation is due to the horizon placed within each model, a negative value for $\cos(\theta)$ indicates that the sun is below the horizon. AtChem2 limits the minimum value for $\cos(\theta)$ to 0.01, whereas INCHEM-Py and repeated experimental measurements of different surfaces, the values for these constants can be validated using the ratio of O₃ to product at equilibrium. Given experimental uncertainties, the values currently present within the model give a good estimation of 4-OPA production with minimal changes to the has a limit of 0. This means that the sun begins to rise between 8 and 10 minutes earlier in INCHEM-Py settings file., allowing photolysis reactions to begin to occur earlier and go on later. As shown in Fig. 9(b), the value for solar noon shows no deviation.




Figure 9. (a) [Percentage deviation in gas-phase chemistry solutions between INCHEM-Py and AtChem2. Species are named as in the MCM.](#) (b) [The percentage deviation between the solar zenith angles as calculated by INCHEM-Py and AtChem2. Percentage deviation calculated by Eq. \(14\)](#)

3.2 H₂O₂ emission and fate

Zhou et al. (2020) studied the impact of non-bleach cleaning events in a combined chamber and modelling study. In this study, the INDCM model was used to explore secondary pollutant formation, and the impact of different lighting levels on their concentrations. [Again, the The INDCM had less complete representations of surface interactions, indoor photolysis and indoor-outdoor exchange.](#) The original work was completed in ppb and so outputs from INCHEM-Py have been converted for the comparison.

The Indoor Environmental Quality chamber at the Building Energy and Environmental Systems Laboratory at Syracuse University had a mock residential room built within it. This 29.1 m³ room had a wooden frame, painted walls and a vinyl floor with no furniture present. Within this chamber a non-bleach H₂O₂ cleaning spray was used on a 0.75 m² area (12 squirts, 15 mL total) and then wiped dry over a period of 1-2 minutes. The surface to volume ratio of the experiment simulated here is 2 m⁻¹ and the floor is assumed to be 1/6 of the total surface area.

The air change rate was constant at 0.51±0.004 h⁻¹ and was sourced from outdoors, the temperature was controlled at 25.7±0.9 °C and relative humidity averaged at 25.8±9.5 % during the experiments. The indoor lighting was from uncovered fluorescent tubes and a solar illuminator was used to provide outdoor lighting through the window of the experimental room. The solar illuminator removed the diurnal variation of the photolysis but not of the outdoor concentrations. In the experiment, the solar flux was measured using a spectrometer, and photolysis coefficients of key species (H₂O₂, NO₂, HONO, NO₃, O₃ and




Figure 10. Comparison of experimental results of a H₂O₂ cleaning event from Zhou et al. (2020) and simulated results using INCHEM-Py.

formaldehyde) were calculated and input into the INDCM. In the INCHEM-Py simulations, to account for the constant solar photolysis, the local hour angle (LHA `in-inchem-main.py`, described in section 2.2) was set to 0 to give constant noon photolysis values. The indoor lighting was set to uncovered fluorescent tubes. A timed emission rate of 5.5×10^{10} molecule $\text{cm}^{-3} \text{s}^{-1}$ (2.25 ppb s^{-1}) for 3 minutes was used to simulate the cleaning spray, based on the measured values. The surface to volume ratios were set to represent the experiment and all other values were default, including the outdoor species concentrations.

Figure 10 shows the H₂O₂ mixing ratio of the experiment from Zhou et al. (2020) and two INCHEM-Py model runs - one simulating the experimental conditions of a plastic floor and painted walls (plastic/paint), and a second where the floor is changed to concrete and the walls to wood (concrete/wood). The plastic/paint scenario shows good agreement between the simulated and measured H₂O₂ mixing ratios. At peak H₂O₂ concentration, the main loss of H₂O₂ was to the painted walls (1.54×10^{10} molecule $\text{cm}^{-3} \text{s}^{-1}$), followed by loss to the plastic floor and loss to outdoors (1.43×10^9 and 1.16×10^9 molecule $\text{cm}^{-3} \text{s}^{-1}$, respectively). Compared to the plastic/paint scenario, the predicted H₂O₂ mixing ratio peaks at a lower value, and drops more quickly after the cleaning, in the concrete/wood scenario. This is due to a larger loss rate to the wooden walls (1.64×10^{10} molecule $\text{cm}^{-3} \text{s}^{-1}$) and concrete floor (5.96×10^9 molecule $\text{cm}^{-3} \text{s}^{-1}$) at peak H₂O₂ concentration.

Following the cleaning event, formaldehyde is produced as a secondary product. Using INCHEM-Py, the maximum increase in formaldehyde as a result of secondary chemistry in the plastic/paint scenario was predicted to be 0.72 ppb. In the experiment (Zhou et al. (2020)), the secondary emission peak was masked by the background formaldehyde concentration of over 50 ppb, therefore, this shows the importance of modelling for delineating between primary and secondary emissions. In the concrete/wood scenario, the increase in formaldehyde as a result of secondary chemistry was much higher, at 3.05 ppb. This




Figure 11. Comparison of run time of the default simulation on two different computers.

is due to higher H_2O_2 deposition velocities, and higher resulting surface emissions from concrete/wood surfaces, compared to the plastic/paint combination (Carter et al., 2023).

4 Practical details

URL to model: <https://github.com/DrDaveShaw/INCHEM-Py>

455 License: [GNU General Public License, v3](#)

INCHEM-Py requires Python 3 and some additional packages to run. The additional packages and their known working versions are given in the user manual. The integration process is set by default to be limited to 4 threads, as this is the only parallelised function within the model. The number of threads can be adjusted by changing the threadpool limits in inchem_main, as described in the user manual. The default model took 18 minutes and 41 seconds on a Windows desktop with an AMD Ryzen 5 5600X 3.7 GHz processor, and 14 minutes 54 seconds on an Apple laptop with a M1 Max 3.2 GHz processor. Figure 11 shows the simulation run times for these two processors. The total run time of a simulation is printed to the console at the end of the model run, and a file showing the time taken for each integration output is saved to the output folder and is included in the data set for all model runs in this manuscript. This function allows users to determine where the simulation may be struggling and where improvements to speed up run time might be made. In the default case, the slowest integration steps are at sunrise and sunset due to very small values for photolysis parameters when the Sun begins to rise or in the final stages of setting. Memory usage peaks at less than 2 GB for these simulations.

460

465

5 Conclusions

INCHEM-Py (the INdoor CHEMical model in Python) has been presented as a tool for the analysis and interrogation of the atmospheric chemistry of the indoor environment. We have presented the main modules developed for v1.2 within this paper, and have given a further detailed description in the user manual submitted alongside. INCHEM-Py has been developed as an open and accessible piece of software, has no hidden or proprietary code, and requires minimal previous coding or modelling experience to install or run. It utilises core Python libraries, including Numpy (Harris et al., 2020) and Scipy (Virtanen et al., 2020), keeping installation of additional libraries to a minimum, and capitalising on the maintenance of the Python ecosystem.

INCHEM-Py has been validated against experimental measurements and has shown improved accuracy in comparison with the INDCM, from which it was refactored. INCHEM-Py has been developed with a focus on predicting secondary chemistry that is not feasible, or in some cases possible, to measure. Outputs of species concentrations with time are given alongside key model parameters such as surface deposition rates, seasonal photolysis rates and diurnal outdoor concentrations. Interrogating these outputs allows for a detailed understanding of the atmospheric chemical processes that occur indoors, including temporally resolved reaction rates which can be used to identify the important pathways that should be included in models with reduced chemical mechanisms (Lakey et al., 2021). For example, there are generally two types of models used to solve indoor air chemistry; box models such as INCHEM-Py that have no spatial dimensions but very complex chemical mechanisms (> 6000 species, > 19,000 reactions); and computational fluid dynamics (CFD) models that solve the evolution of very few species (< 20 species, < 15 reactions), but can track them spatially (Lakey et al., 2021). Neither model is able to fully represent the complexities of the indoor environment. The main assumption of box models is that the atmosphere is well mixed, and that all species are available to react with each other at all times. However, CFD models simply cannot capture the complex chemistry of the indoor environment due to computational constraints. Each model type attempts to fill a different knowledge gap. It is up to the user to appropriately define the parameters of the models to gain the most effective insight into the processes occurring indoors (Shiraiwa et al., 2019).

The utility of INCHEM-Py is further demonstrated by the publications that have used the model since the Shaw and Carslaw (2021) release of v1.1. Published articles include ~~Lakey et al. (2021); Wang et al. (2022); Carter et al. (2023); Beel et al. (2023)~~ Lakey et al. (2021); Wang et al. (2022); Carter et al. (2023); Beel et al. (2023); L. Davies et al. (2023), with several additional articles in preparation that discuss emissions from cooking and cleaning, UV induced emissions from plastics, and future indoor air pollution scenarios. Each case exhibits the versatility of INCHEM-Py and the ability of users to simulate custom experimental scenarios to expand our understanding of indoor air chemistry.

Future developments of INCHEM-Py will likely include an expansion of the gas-to-particle partitioning module to include more source species and prediction of particle size distributions. Chemical mechanisms and analysis tools will continue to be added as they are developed, including a Python package for the analysis of INCHEM-Py outputs. In the long term a multi-box approach may be taken to simulate adjoining spaces, or a hybrid approach may be used to add spatial dimensions.

Appendix A: Outdoor constant concentrations and source of measurements on which they are based

Species	v1.1 value (molecule/cm ³)	v1.2 value (molecules/cm ³)	Ref. (v1.2)
Formaldehyde	9.13×10^{10}	6.017×10^{10}	Uchiyama et al. (2015)
Acetaldehyde	7.15×10^{10}	3.896×10^{10}	Uchiyama et al. (2015)
Propanal	2.02×10^{10}	9.332×10^{09}	Uchiyama et al. (2015)
3-Methylbutanal	-	1.049×10^{09}	Uchiyama et al. (2015)
Acrolein	4.94×10^{10}	2.685×10^{09}	Uchiyama et al. (2015)
Methacrolein	-	2.792×10^{09}	Baudic et al. (2016)
Crotonaldehyde	-	1.718×10^{09}	Uchiyama et al. (2015)
Pentanal	-	2.447×10^{09}	Uchiyama et al. (2015)
Hexanal	9.25×10^{09}	2.706×10^{09}	Uchiyama et al. (2015)
Heptanal	3.75×10^{09}	1.846×10^{09}	Uchiyama et al. (2015)
Octanal	7.25×10^{09}	2.349×10^{09}	Uchiyama et al. (2015)
Nonanal	2.5×10^{10}	1.482×10^{10}	Uchiyama et al. (2015)
Decanal	2.75×10^{09}	4.047×10^{09}	Uchiyama et al. (2015)
2-Nonenal	-	1.288×10^{09}	Uchiyama et al. (2015)
Acetone	1.3×10^{10}	4.977×10^{10}	Uchiyama et al. (2015)
2-Butanone (MEK)	2.41×10^{09}	5.429×10^{09}	Uchiyama et al. (2015)
3-Buten-2-one (MVK)	1.78×10^{10}	2.792×10^{09}	Baudic et al. (2016)
Cyclohexanone	9×10^{08}	1.706×10^{10}	Lü et al. (2006)
Benzaldehyde	6.13×10^{10}	1.419×10^{09}	Uchiyama et al. (2015)
o-Tolualdehyde	-	1.253×10^{09}	Uchiyama et al. (2015)
m-Tolualdehyde	-	2.005×10^{09}	Uchiyama et al. (2015)
p-Tolualdehyde	-	2.005×10^{09}	Uchiyama et al. (2015)
2,5-Dimethylbenzaldehyde	-	7.854×10^{09}	Uchiyama et al. (2015)
Benzene	5.9×10^{09}	9.637×10^{09}	Uchiyama et al. (2015)
Toluene	2×10^{10}	4.085×10^{10}	Uchiyama et al. (2015)
p-Xylene	6.5×10^{09}	6.098×10^{09}	Uchiyama et al. (2015)
m-Xylene	6.5×10^{09}	6.098×10^{09}	Uchiyama et al. (2015)
o-Xylene	1.3×10^{10}	4.254×10^{09}	Uchiyama et al. (2015)
Ethylbenzene	3.4×10^{09}	8.792×10^{09}	Uchiyama et al. (2015)
Propylbenzene	-	4.008×10^{09}	Mentese and Bas (2020)

continued on next page

Species	v1.1 value (molecule/cm ³)	v1.2 value (molecules/cm ³)	Ref. (v1.2)
2-Ethyltoluene	-	2.505×10^{08}	Bari and Kindzierski (2018)
3-Ethyltoluene	-	6.013×10^{08}	Bari and Kindzierski (2018)
4-Ethyltoluene	-	3.006×10^{08}	Bari and Kindzierski (2018)
1,3,5-Trimethylbenzene	2.2×10^{10}	1.754×10^{09}	Uchiyama et al. (2015)
1,2,4-Trimethylbenzene	2.23×10^{09}	5.511×10^{09}	Uchiyama et al. (2015)
1,2,3-Trimethylbenzene	8.0×10^{09}	1.253×10^{09}	Uchiyama et al. (2015)
p-Dichlorobenzene	-	1.229×10^{10}	Uchiyama et al. (2015)
Styrene	5.67×10^{09}	2.313×10^{09}	Mentese and Bas (2020)
Cumene	1.56×10^{08}	3.006×10^{09}	Mentese and Bas (2020)
Phenol	5×10^{10}	1.747×10^{10}	Sturaro et al. (2010)
Ethane	2.08×10^{10}	9.133×10^{10}	Baudic et al. (2016)
Propane	1.25×10^{10}	3.797×10^{10}	Baudic et al. (2016)
Butane	3.33×10^{10}	3.471×10^{10}	Baudic et al. (2016)
Isobutane	-	2.321×10^{10}	Baudic et al. (2016)
2,2-Dimethylbutane	-	2.027×10^{09}	Bari et al. (2016)
2,3-Dimethylbutane	-	2.586×10^{09}	Bari et al. (2016)
Pentane	9.13×10^{09}	8.681×10^{09}	Baudic et al. (2016)
2-Methylpentane	-	3.844×10^{09}	Bari et al. (2016)
3-Methylpentane	-	2.446×10^{09}	Bari et al. (2016)
Isopentane	-	1.469×10^{10}	Bari and Kindzierski (2018)
Hexane	9.15×10^{09}	1.118×10^{10}	Uchiyama et al. (2015)
2-Methylhexane	-	2.464×10^{09}	Bari et al. (2016)
3-Methylhexane	-	3.125×10^{09}	Bari et al. (2016)
Heptane	2.58×10^{09}	6.010×10^{08}	Uchiyama et al. (2015)
Octane	7.5×10^{09}	5.272×10^{08}	Uchiyama et al. (2015)
Nonane	1×10^{10}	3.052×10^{09}	Uchiyama et al. (2015)
Decane	1.96×10^{09}	9.946×10^{09}	Uchiyama et al. (2015)
Undecane	1.95×10^{09}	1.445×10^{10}	Uchiyama et al. (2015)
Dodecane	5.22×10^{08}	1.061×10^{09}	Mentese and Bas (2020)
Cyclohexane	1.19×10^{09}	6.440×10^{08}	Bari and Kindzierski (2018)
Ethene	1.25×10^{10}	3.327×10^{10}	Baudic et al. (2016)

continued on next page

Species	v1.1 value (molecule/cm ³)	v1.2 value (molecules/cm ³)	Ref. (v1.2)
Propene	4.3×10^{09}	9.159×10^{09}	Baudic et al. (2016)
1-Butene	-	3.971×10^{09}	Bari et al. (2016)
cis-2-Butene	3.5×10^{09}	4.293×10^{08}	Bari and Kindzierski (2018)
trans-2-Butene	4×10^{09}	5.367×10^{08}	Bari and Kindzierski (2018)
2-Methyl-1-butene	-	5.152×10^{08}	Bari and Kindzierski (2018)
2-Methyl-2-butene	7×10^{09}	4.293×10^{08}	Bari and Kindzierski (2018)
Isoprene	1.0×10^{10}	2.299×10^{09}	Baudic et al. (2016)
1,3-Butadiene	2.5×10^{09}	5.567×10^{08}	Bari and Kindzierski (2018)
trans-2-Pentene	-	4.293×10^{08}	Bari and Kindzierski (2018)
cis-2-Pentene	-	2.576×10^{08}	Bari and Kindzierski (2018)
Ethyne	-	1.573×10^{10}	Baudic et al. (2016)
Methanol	1.3×10^{11}	1.107×10^{11}	Baudic et al. (2016)
Ethanol	1.2×10^{12}	1.613×10^{11}	Gallego et al. (2016)
Isopropanol	2.0×10^{10}	9.239×10^{10}	Gallego et al. (2016)
1-Propanol	1.2×10^{09}	1.243×10^{10}	Gallego et al. (2016)
1-Butanol	1.3×10^{10}	2.519×10^{10}	Gallego et al. (2016)
1-Pentanol	-	5.658×10^{07}	Hellén et al. (2018)
1-Hexanol	-	3.014×10^{07}	Hellén et al. (2018)
2-Butoxyethanol	5.19×10^{09}	2.507×10^{10}	Gallego et al. (2016)
Linalool	-	1.292×10^{07}	Hellén et al. (2018)
Chloroform	2.93×10^{08}	7.567×10^{08}	Uchiyama et al. (2015)
Methylchloroform	8.33×10^{10}	7.674×10^{09}	Brickus et al. (1998)
Dichloromethane	1×10^{09}	2.340×10^{09}	Bari and Kindzierski (2018)
Trichloroethylene	3×10^{07}	9.075×10^{09}	Gallego et al. (2016)
Tetrachloroethylene	2×10^{08}	5.084×10^{08}	Bari and Kindzierski (2018)
1,2-Dichloroethane	-	4.260×10^{08}	Bari and Kindzierski (2018)
Chloromethane	-	1.396×10^{10}	Bari and Kindzierski (2018)
Hydrogen Chloride	-	3.716×10^{10}	Uchiyama et al. (2015)
Ethyl Acetate	-	2.392×10^{09}	Uchiyama et al. (2015)
Butyl Acetate	-	1.296×10^{09}	Uchiyama et al. (2015)
α -Pinene	1.45×10^{09}	3.094×10^{09}	Uchiyama et al. (2015)

continued on next page

Species	v1.1 value (molecule/cm ³)	v1.2 value (molecules/cm ³)	Ref. (v1.2)
β -Pinene	2.5×10^{07}	1.238×10^{09}	Gallego et al. (2016)
Limonene	9×10^{08}	2.431×10^{09}	Uchiyama et al. (2015)
Δ 3-Carene	9×10^{08}	2.718×10^{09}	Hakola et al. (2009)
Camphene	5×10^{08}	3.978×10^{08}	Hakola et al. (2009)
Formic Acid	-	1.832×10^{11}	Uchiyama et al. (2015)
Acetic Acid	-	3.861×10^{11}	Uchiyama et al. (2015)
Propanoic Acid	-	1.873×10^{09}	Hellén et al. (2018)
Butanoic Acid	-	1.381×10^{09}	Hellén et al. (2018)
Pentanoic Acid	-	7.534×10^{08}	Hellén et al. (2018)
Heptanoic Acid	-	9.932×10^{07}	Hellén et al. (2018)
Hydrogen Peroxide	5×10^{10}	3.13×10^{10}	He et al. (2010)
β -Caryophyllene	2.5×10^{07}	9.348×10^{07}	Hellén et al. (2018)
Methane (CH ₄)	4.63×10^{13}	4.652×10^{13}	Dlugokencky (2022)
Carbon Monoxide (CO)	2.5×10^{12}	$6.6424.797 \times 10^{12}$	Naghizadeh et al. (2019) Median value, U
Sulfur Dioxide (SO ₂)	-	1.715×10^{10}	EEA (2018)
Nitrogen Oxide (NO)	2.59×10^{10}	Average	EEA (2018)
Nitrogen Dioxide (NO ₂)	9.52×10^{10}	Average	EEA (2018)
Ozone (O ₃)	7.68×10^{11}	Average	EEA (2018)
Nitric Acid (HNO ₃)	5×10^{10}	9.557×10^{09}	Vichi et al. (2016)
Nitrous Acid (HONO)	1.6×10^{09}	1.588×10^{10}	Vichi et al. (2016)
Hydroxyl Radical (OH)	1×10^{06}	1.09×10^{06}	Li et al. (2018)
Peroxyacetyl Nitrates (PAN)	1.51×10^{10}	5.449×10^{10}	Liu et al. (2018)
Total Suspended Particles (TSP)	1.4×10^{11}	Average	EEA (2018)

Table A1. Outdoor constant concentrations used in INCHEM-Py including the references used to obtain the v1.2 values. Values given as "Average" are averages of the diurnal profiles used in the model. The cited values are taken from multiple papers covering multiple locations, times of year and lengths of study that may or may not be applicable to individual user requirements. A summary of these details is given in Tab. A1.

Appendix B: **Default settings**

505

```

1: filename = 'mcm_v331.fac'
2: -
3: particles = True
4: -

```

```

510 5: INCHEM_additional = True
6: -
7: eustom = False
8: -
9: spline = 293.
515 10: temperatures = [[25200, 288.15], [50400, 294.15]]
11: -
12: rel_humidity = 50.
13: M = 2.51e+19
14: -
520 15: const_diet = {
16: —— 'O2': 0.2095 * M,
17: —— 'N2': 0.7809 * M,
18: —— 'H2': 550e-9 * M,
19: —— 'saero': 1.3e-2}
525 20: -
21: ACRate = {0 : 0.5/3600,
22: —— 3600 * 24 : 1/3600,
23: —— 3600 * 48 : 2/3600}
24: diurnal = True
530 25: city = "Bergen_urban"
26: -
27: date = "21-06-2020"
28: lat = 45
29: light_type = "Incand"
535 30: light_on_times = [[7, 19], [31, 43], [55, 67], [79, 91]]
31: glass = "glass_C"
32: -
33: AV = 0.02
34: -
540 35: surfaces_AV = {
36: —— 'AVSOFT' : 0.0035,
37: —— 'AVPAINT' : 0.0114,
38: —— 'AVWOOD' : 0.0061,
39: —— 'AVMETAL' : 0.0025,

```

```
545 40: ————'AVCONCRETE': 0.0001,
41: ————'AVPAPER': 0.0006,
42: ————'AVLINO': 0.0000,
43: ————'AVPLASTIC': 0.0048,
44: ————'AVGLASS': 0.0009,
550 45: ————'AVHUMAN': 0.0017}
46: —
47: H2O2_dep = True
48: O3_dep = True
49: —
555 50: adults = 0
51: children = 0
52: —
53: initials_from_run = False
54: —
560 55: initial_conditions_gas = 'initial_concentrations.txt'
56: —
57: timed_emissions = False
58: —
59: timed_inputs = {"LIMONENE": [[46800, 47400, 5e8], [107600, 108000, 5e8]],
565 60: —————"BPINENE": [[46800, 47400, 5e8]]}
61: —
62: dt = 120
63: t0 = 0
64: seconds_to_integrate = 86400
570 65: —
66: custom_name = "Bergen_urban"
67: —
68: reactions_output = True
69: —
575 70: output_graph = True
71: output_species = ['LIMONENE', 'APINENE']
```

Table A1. Locations, times of year and lengths of study for literature used to update outdoor concentrations in INCHEM-Py v1.2

<u>Reference</u>	<u>Location</u>	<u>Time of year</u>	<u>Length of study</u>
<u>Uchiyama et al. (2015)</u>	<u>Japan</u>	<u>Winter (January - March) and Summer (July to September)</u>	<u>3 years</u>
<u>Baudic et al. (2016)</u>	<u>France</u>	<u>All Year</u>	<u>11 months</u>
<u>Lü et al. (2006)</u>	<u>China</u>	<u>Winter (January - March)</u>	<u>2.5 months</u>
<u>Mentese and Bas (2020)</u>	<u>Turkey</u>	<u>All Year</u>	<u>1 year</u>
<u>Bari and Kindzierski (2018)</u>	<u>Canada (Calgary)</u>	<u>All Year</u>	<u>5 years</u>
<u>Sturaro et al. (2010)</u>	<u>Italy</u>	<u>February - November</u>	<u>10 months</u>
<u>Bari et al. (2016)</u>	<u>Canada (Fort McKay)</u>	<u>All Year</u>	<u>12 years</u>
<u>Gallego et al. (2016)</u>	<u>Spain</u>	<u>July - November</u>	<u>5 months</u>
<u>Hellén et al. (2018)</u>	<u>Finland</u>	<u>All Year</u>	<u>2 years</u>
<u>Hakola et al. (2009)</u>	<u>Finland</u>	<u>All Year</u>	<u>8 years</u>
<u>He et al. (2010)</u>	<u>China</u>	<u>July - September</u>	<u>2 months</u>
<u>Dlugokencky (2022)</u>	<u>Global</u>	<u>All Year</u>	<u>1 year</u>
<u>Vichi et al. (2016)</u>	<u>Switzerland and Czech Republic</u>	<u>Spring, Summer and Winter</u>	<u>4 years</u>
<u>Li et al. (2018)</u>	<u>Global</u>	<u>All Year</u>	<u>8 years</u>
<u>Liu et al. (2018)</u>	<u>China</u>	<u>November - July</u>	<u>9 months</u>

Code and data availability. INCHEM-Py, including the user manual, is available on github at <https://github.com/DrDaveShaw/INCHEM-Py> with an archived submission version available at Shaw et al. (2023). The full data used to produce all the figures and data within this paper is available at <https://doi.org/10.15124/b68c1c34-8974-46d8-8728-05c6cd6e9e8b>.

Author contributions. DRS conceptualisation, data curation, formal analysis, investigation, methodology, software, validation, visualisation, writing - original draft preparation, writing - review and editing. TJC data curation, investigation, software, methodology, writing - original draft preparation, writing - review and editing. HLD, EHS and GB validation, writing - review and editing. ECC validation. ZW methodology. NC conceptualisation, formal analysis, investigation, methodology, project administration, funding acquisition, resources, software, supervision, writing - review and editing.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The development of this model has been funded by grants from the Alfred P. Sloan Foundation COMMODIAC (2018-10083) and MOCCIE2 (2019-12306) for DRS and MOCCIE3 for TJC (2020-13912). Conclusions reached or positions taken by researchers or other grantees represent the views of the grantees themselves and not those of the Alfred P. Sloan Foundation or its trustees, officers, or staff. The authors would like to thank Tara Kahan and Shan Zhou for providing the raw data from Zhou et al. (2020) for comparative analysis. We would also like to thank Roberto Sommariva for his feedback during model development [and for running AtChem2 for the comparative study](#). For the purpose of open access a Creative Commons Attribution (CC BY) licence is applied to any Author Accepted Manuscript version arising from this submission.

References

- 595 Abbass, O. A., Sailor, D. J., and Gall, E. T.: Effect of fiber material on ozone removal and carbonyl production from carpets, *Atmospheric Environment*, 148, 42–48, <https://doi.org/10.1016/j.atmosenv.2016.10.034>, 2017.
- Alicke, B.: OH formation by HONO photolysis during the BERLIOZ experiment, *Journal of Geophysical Research*, 108, 8247, <https://doi.org/10.1029/2001JD000579>, 2003.
- Arata, C., Zarzana, K. J., Misztal, P. K., Liu, Y., Brown, S. S., Nazaroff, W. W., and Goldstein, A. H.: Measurement of NO
600 ₃ and N₂O₅ in a Residential Kitchen, *Environmental Science & Technology Letters*, 5, 595–599, <https://doi.org/10.1021/acs.estlett.8b00415>, 2018.
- Bari, M. A. and Kindzierski, W. B.: Ambient volatile organic compounds (VOCs) in Calgary, Alberta: Sources and screening health risk assessment, *Science of the Total Environment*, 631–632, 627–640, <https://doi.org/10.1016/j.scitotenv.2018.03.023>, 2018.
- Bari, M. A., Kindzierski, W. B., and Spink, D.: Twelve-year trends in ambient concentrations of volatile organic compounds in a community
605 of the Alberta Oil Sands Region, Canada, *Environment International*, 91, 40–50, <https://doi.org/10.1016/j.envint.2016.02.015>, 2016.
- Baudic, A., Gros, V., Sauvage, S., Locoge, N., Sanchez, O., Sarda-Estève, R., Kalogridis, C., Petit, J. E., Bonnaire, N., Baisnée, D., Favez, O., Albinet, A., Sciare, J., and Bonsang, B.: Seasonal variability and source apportionment of volatile organic compounds (VOCs) in the Paris megacity (France), *Atmospheric Chemistry and Physics*, 16, 11 961–11 989, <https://doi.org/10.5194/acp-16-11961-2016>, 2016.
- Beel, G., Langford, B., Carslaw, N., Shaw, D., and Cowan, N.: Temperature driven variations in VOC emissions from plastic
610 products and their fate indoors: A chamber experiment and modelling study, *Science of The Total Environment*, 881, 163 497, <https://doi.org/10.1016/j.scitotenv.2023.163497>, 2023.
- Blocquet, M., Guo, F., Mendez, M., Ward, M., Coudert, S., Batut, S., Hecquet, C., Blond, N., Fittschen, C., and Schoemaeker, C.: Impact of the spectral and spatial properties of natural light on indoor gas-phase chemistry: Experimental and modeling study, *Indoor Air*, 28, 426–440, <https://doi.org/10.1111/ina.12450>, 2018.
- 615 Bloss, C., Wagner, V., Jenkin, M. E., Volkamer, R., Bloss, W. J., Lee, J. D., Heard, D. E., Wirtz, K., Martin-Reviejo, M., Rea, G., Wenger, J. C., and Pilling, M. J.: Development of a detailed chemical mechanism (MCMv3.1) for the atmospheric oxidation of aromatic hydrocarbons, *Atmospheric Chemistry and Physics*, 5, 641–664, <https://doi.org/10.5194/acp-5-641-2005>, 2005.
- Brickus, L. S., Cardoso, J. N., and De Aquino Neto, F. R.: Distributions of indoor and outdoor air pollutants in Rio de Janeiro, Brazil: Implications to indoor air quality in bayside offices, *Environmental Science and Technology*, 32, 3485–3490, <https://doi.org/10.1021/es980336x>,
620 1998.
- Cano-Ruiz, J., Kong, D., Balas, R., and Nazaroff, W. W.: Removal of reactive gases at indoor surfaces: Combining mass transport and surface kinetics, *Atmospheric Environment. Part A. General Topics*, 27, 2039–2050, 1993.
- Carslaw, N.: A new detailed chemical model for indoor air pollution, *Atmospheric Environment*, 41, 1164–1179, <https://doi.org/10.1016/j.atmosenv.2006.09.038>, 2007.
- 625 Carslaw, N.: A mechanistic study of limonene oxidation products and pathways following cleaning activities, *Atmospheric Environment*, 80, 507–513, <https://doi.org/10.1016/j.atmosenv.2013.08.034>, 2013.
- Carslaw, N. and Shaw, D.: Modification of cleaning product formulations could improve indoor air quality, *Indoor Air*, 32, <https://doi.org/10.1111/ina.13021>, 2022.
- Carslaw, N., Mota, T., Jenkin, M. E., Barley, M. H., and McFiggans, G.: A Significant role for nitrate and peroxide groups on indoor secondary
630 organic aerosol, *Environmental Science and Technology*, 46, 9290–9298, <https://doi.org/10.1021/es301350x>, 2012.

- Carslaw, N., Fletcher, L., Heard, D., Ingham, T., and Walker, H.: Significant OH production under surface cleaning and air cleaning conditions: Impact on indoor air quality, *Indoor Air*, 27, 1091–1100, <https://doi.org/10.1111/ina.12394>, 2017.
- Carter, T. J., Poppendieck, D. G., Shaw, D., and Carslaw, N.: A Modelling Study of Indoor Air Chemistry: The Surface Interactions of Ozone and Hydrogen Peroxide, *Atmospheric Environment*, 297, 119 598, <https://doi.org/10.1016/j.atmosenv.2023.119598>, 2023.
- 635 Coleman, B. K., Destailhats, H., Hodgson, A. T., and Nazaroff, W. W.: Ozone consumption and volatile byproduct formation from surface reactions with aircraft cabin materials and clothing fabrics, *Atmospheric Environment*, 42, 642–654, <https://doi.org/10.1016/j.atmosenv.2007.10.001>, 2008.
- Cox, R. A. and Penkett, S. A.: Effect of relative humidity on the disappearance of ozone and sulphur dioxide in contained systems, *Atmospheric Environment* (1967), 6, 365–368, [https://doi.org/10.1016/0004-6981\(72\)90203-X](https://doi.org/10.1016/0004-6981(72)90203-X), 1972.
- 640 Cros, C. J., Morrison, G. C., Siegel, J. A., and Corsi, R. L.: Long-term performance of passive materials for removal of ozone from indoor air, *Indoor Air*, 22, 43–53, <https://doi.org/10.1111/j.1600-0668.2011.00734.x>, 2012.
- Di, Y., Mo, J., Zhang, Y., and Deng, J.: Ozone deposition velocities on cotton clothing surface determined by the field and laboratory emission cell, *Indoor and Built Environment*, 26, 631–641, <https://doi.org/10.1177/1420326X16628315>, 2017.
- Dlugokencky, E.: NOAA/GML CH4 Trends, https://gml.noaa.gov/webdata/ccgg/trends/ch4/ch4_mm_gl.txt, 2022.
- 645 EEA: European Air Quality Portal, <https://eeadmz1-cws-wp-air02.azurewebsites.net/>, 2018.
- Emmerson, K. M., Carslaw, N., and Pilling, M. J.: Urban Atmospheric Chemistry During the PUMA Campaign 2: Radical Budgets for OH, HO₂ and RO₂, *Journal of Atmospheric Chemistry*, 52, 165–183, <https://doi.org/10.1007/s10874-005-1323-2>, 2005.
- Fadeyi, M. O., Weschler, C. J., Tham, K. W., Wu, W. Y., and Sultan, Z. M.: Impact of human presence on secondary organic aerosols derived from ozone-initiated chemistry in a simulated office environment, *Environmental Science and Technology*, 47, 3933–3941, <https://doi.org/10.1021/es3050828>, 2013.
- 650 Farmer, D. K., Vance, M. E., Abbatt, J. P. D., Abeleira, A., Alves, M. R., Arata, C., Boedicker, E., Bourne, S., Cardoso-Saldaña, F., Corsi, R., DeCarlo, P. F., Goldstein, A. H., Grassian, V. H., Hildebrandt Ruiz, L., Jimenez, J. L., Kahan, T. F., Katz, E. F., Mattila, J. M., Nazaroff, W. W., Novoselac, A., O'Brien, R. E., Or, V. W., Patel, S., Sankhyan, S., Stevens, P. S., Tian, Y., Wade, M., Wang, C., Zhou, S., and Zhou, Y.: Overview of HOMEChem: House Observations of Microbial and Environmental Chemistry, *Environmental Science: Processes & Impacts*, 21, 1280–1300, <https://doi.org/10.1039/C9EM00228F>, 2019.
- Fischer, A., Ljungström, E., and Langer, S.: Ozone removal by occupants in a classroom, *Atmospheric Environment*, 81, 11–17, <https://doi.org/10.1016/j.atmosenv.2013.08.054>, 2013.
- Gall, E., Darling, E., Siegel, J. A., Morrison, G. C., and Corsi, R. L.: Evaluation of three common green building materials for ozone removal, and primary and secondary emissions of aldehydes, *Atmospheric Environment*, 77, 910–918, <https://doi.org/10.1016/j.atmosenv.2013.06.014>, 2013.
- 660 Gallego, E., Roca, F. J., Perales, J. F., Guardino, X., Gadea, E., and Garrote, P.: Impact of formaldehyde and VOCs from waste treatment plants upon the ambient air nearby an urban area (Spain), *Science of the Total Environment*, 568, 369–380, <https://doi.org/10.1016/j.scitotenv.2016.06.007>, 2016.
- Grøntoft, T.: Dry deposition of ozone on building materials. Chamber measurements and modelling of the time-dependent deposition, *Atmospheric Environment*, 36, 5661–5670, [https://doi.org/10.1016/S1352-2310\(02\)00701-X](https://doi.org/10.1016/S1352-2310(02)00701-X), 2002.
- 665 Grøntoft, T. and Raychaudhuri, M. R.: Compilation of tables of surface deposition velocities for O₃, NO₂ and SO₂ to a range of indoor surfaces, *Atmospheric Environment*, 38, 533–544, <https://doi.org/10.1016/j.atmosenv.2003.10.010>, 2004.

- Hakola, H., Hellén, H., Tarvainen, V., Bäck, J., Patokoski, J., and Rinne, J.: Annual variations of atmospheric VOC concentrations in a boreal forest, *Boreal Environment Research*, 14, 722–730, 2009.
- 670 Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk, M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., Gérard-Marchant, P., Sheppard, K., Reddy, T., Weckesser, W., Abbasi, H., Gohlke, C., and Oliphant, T. E.: Array programming with NumPy, *Nature*, 585, 357–362, <https://doi.org/10.1038/s41586-020-2649-2>, 2020.
- Hayman, G.: Effects of pollution control on UV Exposure, Tech. rep., Oxfordshire, 1997.
- 675 He, S. Z., Chen, Z. M., Zhang, X., Zhao, Y., Huang, D. M., Zhao, J. N., Zhu, T., Hu, M., and Zeng, L. M.: Measurement of atmospheric hydrogen peroxide and organic peroxides in Beijing before and during the 2008 Olympic Games: Chemical and physical factors influencing their concentrations, *Journal of Geophysical Research*, 115, D17 307, <https://doi.org/10.1029/2009JD013544>, 2010.
- Hellén, H., Praplan, A. P., Tykkä, T., Ylivinkka, I., Vakkari, V., Bäck, J., Petäjä, T., Kulmala, M., and Hakola, H.: Long-term measurements of volatile organic compounds highlight the importance of sesquiterpenes for the atmospheric chemistry of a boreal forest, *Atmospheric*
- 680 *Chemistry and Physics*, 18, 13 839–13 863, <https://doi.org/10.5194/acp-18-13839-2018>, 2018.
- Hindmarsh, A. C.: ODEPACK, A Systematized Collection of ODE Solvers, *IMACS Transactions on Scientific Computation*, 1, 55–64, <https://www3.nd.edu/~powers/ame.60636/hindmarsh1983.pdf>, 1983.
- Jenkin, M. E.: Modelling the formation and composition of secondary organic aerosol from α - and β -pinene ozonolysis using MCM v3, *Atmospheric Chemistry and Physics*, 4, 1741–1757, <https://doi.org/10.5194/acp-4-1741-2004>, 2004.
- 685 Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: A protocol for mechanism development, *Atmospheric Environment*, 31, 81–104, [https://doi.org/10.1016/S1352-2310\(96\)00105-7](https://doi.org/10.1016/S1352-2310(96)00105-7), 1997.
- Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, *Atmospheric Chemistry and Physics*, 3, 181–193, <https://doi.org/10.5194/acp-3-181-2003>, 2003.
- 690 Jenkin, M. E., Young, J. C., and Rickard, A. R.: The MCM v3.3.1 degradation scheme for isoprene, *Atmospheric Chemistry and Physics*, 15, 11 433–11 459, <https://doi.org/10.5194/acp-15-11433-2015>, 2015.
- Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., and Wallington, T. J.: Estimation of rate coefficients and branching ratios for gas-phase reactions of OH with aromatic organic compounds for use in automated mechanism construction, *Atmospheric Chemistry and Physics*, 18, 9329–9349, <https://doi.org/10.5194/acp-18-9329-2018>, 2018.
- 695 Jenkin, M. E., Valorso, R., Aumont, B., Newland, M. J., and Rickard, A. R.: Estimation of rate coefficients for the reactions of O₂ with unsaturated organic compounds for use in automated mechanism construction, *Atmospheric Chemistry and Physics*, 20, 12 921–12 937, <https://doi.org/10.5194/acp-20-12921-2020>, 2020.
- Johnson, D., Utembe, S. R., Jenkin, M. E., Derwent, R. G., Hayman, G. D., Alfarra, M. R., Coe, H., and McFiggans, G.: Simulating regional scale secondary organic aerosol formation during the TORCH 2003 campaign in the southern UK, *Atmospheric Chemistry and Physics*,
- 700 6, 403–418, <https://doi.org/10.5194/acp-6-403-2006>, 2006.
- Klenø, J. G., Clausen, P. A., Weschler, C. J., and Wolkoff, P.: Determination of ozone removal rates by selected building products using the FLEC emission cell, *Environmental Science and Technology*, 35, 2548–2553, <https://doi.org/10.1021/es000284n>, 2001.
- Klepeis, N. E., Nelson, W. C., Ott, W. R., Robinson, J. P., Tsang, A. M., Switzer, P., Behar, J. V., Hern, S. C., and Engelmann, W. H.: The National Human Activity Pattern Survey (NHAPS): a resource for assessing exposure to environmental pollutants, *Journal of Exposure*
- 705 *Science & Environmental Epidemiology*, 11, 231–252, <https://doi.org/10.1038/sj.jea.7500165>, 2001.

- Kowal, S. F., Allen, S. R., and Kahan, T. F.: Wavelength-Resolved Photon Fluxes of Indoor Light Sources: Implications for HO_x Production, *Environmental Science and Technology*, 51, 10423–10430, <https://doi.org/10.1021/acs.est.7b02015>, 2017.
- Kruza, M. and Carslaw, N.: How do breath and skin emissions impact indoor air chemistry?, *Indoor Air*, 29, 369–379, <https://doi.org/10.1111/ina.12539>, 2019.
- 710 Kruza, M., Lewis, A. C., Morrison, G. C., and Carslaw, N.: Impact of surface ozone interactions on indoor air chemistry: A modeling study, *Indoor Air*, 27, 1001–1011, <https://doi.org/10.1111/ina.12381>, 2017.
- Kruza, M., Shaw, D., Shaw, J., and Carslaw, N.: Towards improved models for indoor air chemistry: A Monte Carlo simulation study, *Atmospheric Environment*, 262, 118625, <https://doi.org/10.1016/j.atmosenv.2021.118625>, 2021.
- Lakey, P. S. J., Won, Y., Shaw, D., Østerstrøm, F. F., Mattila, J., Reidy, E., Bottorff, B., Rosales, C., Wang, C., Ampollini, L., Zhou, S.,
715 Novoselac, A., Kahan, T. F., DeCarlo, P. F., Abbatt, J. P. D., Stevens, P. S., Farmer, D. K., Carslaw, N., Rim, D., and Shiraiwa, M.: Spatial and temporal scales of variability for indoor air constituents, *Communications Chemistry*, 4, 1–7, <https://doi.org/10.1038/s42004-021-00548-5>, 2021.
- Lamble, S. P., Corsi, R. L., and Morrison, G. C.: Ozone deposition velocities, reaction probabilities and product yields for green building materials, *Atmospheric Environment*, 45, 6965–6972, <https://doi.org/10.1016/j.atmosenv.2011.09.025>, 2011.
- 720 Leungsakul, S., Jaoui, M., and Kamens, R. M.: Kinetic Mechanism for Predicting Secondary Organic Aerosol Formation from the Reaction of d-Limonene with Ozone, *Environmental Science & Technology*, 39, 9583–9594, <https://doi.org/10.1021/es0492687>, publisher: American Chemical Society, 2005.
- Li, M., Karu, E., Brenninkmeijer, C., Fischer, H., Lelieveld, J., and Williams, J.: Tropospheric OH and stratospheric OH and Cl concentrations determined from CH₄, CH₃Cl, and SF₆ measurements, *npj Climate and Atmospheric Science*, 1, 1–7, [https://doi.org/10.1038/s41612-018-](https://doi.org/10.1038/s41612-018-0041-9)
725 0041-9, 2018.
- Lin, C. C. and Hsu, S. C.: Deposition velocities and impact of physical properties on ozone removal for building materials, *Atmospheric Environment*, 101, 194–199, <https://doi.org/10.1016/j.atmosenv.2014.11.029>, 2015.
- Liu, L., Wang, X., Chen, J., Xue, L., Wang, W., Wen, L., Li, D., and Chen, T.: Understanding unusually high levels of peroxyacetyl nitrate (PAN) in winter in Urban Jinan, China, *Journal of Environmental Sciences (China)*, 71, 249–260,
730 <https://doi.org/10.1016/j.jes.2018.05.015>, 2018.
- Liu, Y., Misztal, P. K., Arata, C., Weschler, C. J., Nazaroff, W. W., and Goldstein, A. H.: Observing ozone chemistry in an occupied residence, *Proceedings of the National Academy of Sciences*, 118, <https://doi.org/10.1073/pnas.2018140118>, 2021.
- Lü, H., Wen, S., Feng, Y., Wang, X., Bi, X., Sheng, G., and Fu, J.: Indoor and outdoor carbonyl compounds and BTEX in the hospitals of Guangzhou, China, *Science of the Total Environment*, 368, 574–584, <https://doi.org/10.1016/j.scitotenv.2006.03.044>, 2006.
- 735 L. Davies, H., O’Leary, C., Dillon, T., R. Shaw, D., Shaw, M., Mehra, A., Phillips, G., and Carslaw, N.: A measurement and modelling investigation of the indoor air chemistry following cooking activities, *Environmental Science: Processes & Impacts*, 25, 1532–1548, <https://doi.org/10.1039/D3EM00167A>, publisher: Royal Society of Chemistry, 2023.
- Mentese, S. and Bas, B.: A year-round motoring of ambient volatile organic compounds across Dardanelles strait, *Journal of Chemical Metrology*, 14, 177–189, <https://doi.org/10.25135/jcm.51.20.09.1816>, 2020.
- 740 Morrison, G. C. and Nazaroff, W. W.: The Rate of Ozone Uptake on Carpets: Experimental Studies, *Environmental Science & Technology*, 34, 4963–4968, <https://doi.org/10.1021/es001361h>, 2000.
- Morrison, G. C. and Nazaroff, W. W.: Ozone interactions with carpet: Secondary emissions of aldehydes, *Environmental Science and Technology*, 36, 2185–2192, <https://doi.org/10.1021/es0113089>, 2002.

- Mueller, F. X., Loeb, L., and Mapes, W. H.: Decomposition Rates of Ozone in Living Areas, *Environmental Science and Technology*, 7, 745 342–346, <https://doi.org/10.1021/es60076a003>, 1973.
- Naghizadeh, A., Sharifzadeh, G., Tabatabaei, F., Afzali, A., Yari, A. R., Geravandi, S., and Mohammadi, M. J.: Assessment of carbon monoxide concentration in indoor/outdoor air of Sarayan city, Khorasan Province of Iran, *Environmental Geochemistry and Health*, 41, 1875–1880, <https://doi.org/10.1007/s10653-018-0226-5>, 2019.
- Nazaroff, W. W. and Weschler, C. J.: Cleaning products and air fresheners: Exposure to primary and secondary air pollutants, *Atmospheric Environment*, 38, 2841–2865, <https://doi.org/10.1016/j.atmosenv.2004.02.040>, 2004.
- Nicolas, M., Ramalho, O., and Maupetit, F.: Reactions between ozone and building products: Impact on primary and secondary emissions, *Atmospheric Environment*, 41, 3129–3138, <https://doi.org/10.1016/j.atmosenv.2006.06.062>, 2007.
- O’Meara, S., Xu, S., Topping, D., Capes, G., Lowe, D., Alfara, M., and McFiggans, G.: PyCHAM: CHemistry with Aerosol Microphysics in Python, *Journal of Open Source Software*, 5, 1918, <https://doi.org/10.21105/joss.01918>, 2020.
- 755 Pankow, J. F.: An absorption model of the gas/aerosol partitioning involved in the formation of secondary organic aerosol, *Atmospheric Environment*, 28, 189–193, [https://doi.org/10.1016/1352-2310\(94\)90094-9](https://doi.org/10.1016/1352-2310(94)90094-9), 1994.
- Platt, U., Alicke, B., Dubois, R., Geyer, A., Hofzumahaus, A., Holland, F., Martinez, M., Mihelcic, D., Klüpfel, T., Lohrmann, B., Pätz, W., Perner, D., Rohrer, F., Schäfer, J., and Stutz, J.: Free Radicals and Fast Photochemistry during BERLIOZ, in: *Tropospheric Chemistry*, pp. 359–394, Springer Netherlands, Dordrecht, https://doi.org/10.1007/978-94-010-0399-5_15, 2002.
- 760 Poppendieck, D., Hubbard, H., Ward, M., Weschler, C., and Corsi, R. L.: Ozone reactions with indoor materials during building disinfection, *Atmospheric Environment*, 41, 3166–3176, <https://doi.org/10.1016/j.atmosenv.2006.06.060>, 2007.
- Poppendieck, D., Hubbard, H., and Corsi, R. L.: Hydrogen Peroxide Vapor as an Indoor Disinfectant: Removal to Indoor Materials and Associated Emissions of Organic Compounds, *Environmental Science and Technology Letters*, 8, 320–325, <https://doi.org/10.1021/acs.estlett.0c00948>, 2021.
- 765 Rai, A. C., Guo, B., Lin, C. H., Zhang, J., Pei, J., and Chen, Q.: Ozone reaction with clothing and its initiated VOC emissions in an environmental chamber, *Indoor Air*, 24, 49–58, <https://doi.org/10.1111/ina.12058>, 2014.
- Reiss, R., Ryan, P. B., and Koutrakis, P.: Modeling Ozone Deposition onto Indoor Residential Surfaces, *Environmental Science and Technology*, 28, 504–513, <https://doi.org/10.1021/es00052a025>, 1994.
- Rim, D., Gall, E. T., Maddalena, R. L., and Nazaroff, W. W.: Ozone reaction with interior building materials: Influence of diurnal ozone variation, temperature and humidity, *Atmospheric Environment*, 125, 15–23, <https://doi.org/10.1016/j.atmosenv.2015.10.093>, 2016.
- 770 Sabersky, R. H., Sinema, D. A., and Shair, F. H.: Concentrations, Decay Rates, and Removal of Ozone and Their Relation to Establishing Clean Indoor Air, *Environmental Science and Technology*, 7, 347–353, <https://doi.org/10.1021/es60076a001>, 1973.
- Sarwar, G. and Corsi, R.: The effects of ozone/limonene reactions on indoor secondary organic aerosols, *Atmospheric Environment*, 41, 959–973, <https://doi.org/10.1016/j.atmosenv.2006.09.032>, 2007.
- 775 Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmospheric Chemistry and Physics*, 3, 161–180, <https://doi.org/10.5194/acp-3-161-2003>, 2003.
- Schraufnagel, D. E.: The health effects of ultrafine particles, *Experimental & Molecular Medicine*, 52, 311–317, <https://doi.org/10.1038/s12276-020-0403-3>, 2020.
- 780 Schripp, T., Langer, S., and Salthammer, T.: Interaction of ozone with wooden building products, treated wood samples and exotic wood species, *Atmospheric Environment*, 54, 365–372, <https://doi.org/10.1016/j.atmosenv.2012.02.064>, 2012.

- Shaw, D. and Carslaw, N.: INCHEM-Py: An open source Python box model for indoor air chemistry, *Journal of Open Source Software*, 6, 3224, <https://doi.org/10.21105/joss.03224>, 2021.
- 785 Shaw, D. R., Carter, T. J., Davies, H. L., Harding-Smith, E., Crocker, E. C., Beel, G., Wang, Z., and Carslaw, N.: INCHEM-Py v1.2: A community box model for indoor air chemistry, <https://doi.org/10.5281/ZENODO.8046598>, 2023.
- Shiraiwa, M., Carslaw, N., Tobias, D. J., Waring, M. S., Rim, D., Morrison, G., Lakey, P. S. J., Kruza, M., Von Domaros, M., Cummings, B. E., and Won, Y.: Modelling consortium for chemistry of indoor environments (MOCCIE): Integrating chemical processes from molecular to room scales, *Environmental Science: Processes and Impacts*, 21, 1240–1254, <https://doi.org/10.1039/c9em00123a>, publisher: Royal Society of Chemistry, 2019.
- 790 Simmons, A. and Colbeck, I.: Resistance of various building materials to ozone deposition, *Environmental Technology (United Kingdom)*, 11, 973–978, <https://doi.org/10.1080/09593339009384949>, 1990.
- Sommariva, R., Cox, S., Martin, C., Borońska, K., Young, J., Jimack, P. K., Pilling, M. J., Matthaios, V. N., Nelson, B. S., Newland, M. J., Panagi, M., Bloss, W. J., Monks, P. S., and Rickard, A. R.: AtChem (version 1), an open-source box model for the Master Chemical Mechanism, *Geoscientific Model Development*, 13, 169–183, <https://doi.org/10.5194/gmd-13-169-2020>, publisher: Copernicus GmbH, 795 2020.
- Sturaro, A., Rella, R., Parvoli, G., and Ferrara, D.: Long-term phenol, cresols and BTEX monitoring in urban air, *Environmental Monitoring and Assessment*, 164, 93–100, <https://doi.org/10.1007/s10661-009-0877-x>, 2010.
- Tamás, G., Weschler, C. J., Bakó-Biró, Z., Wyon, D. P., and Strøm-Tejsen, P.: Factors affecting ozone removal rates in a simulated aircraft cabin environment, *Atmospheric Environment*, 40, 6122–6133, <https://doi.org/10.1016/j.atmosenv.2006.05.034>, 2006.
- 800 Terry, A. C., Carslaw, N., Ashmore, M., Dimitroulopoulou, S., and Carslaw, D. C.: Occupant exposure to indoor air pollutants in modern European offices: An integrated modelling approach, *Atmospheric Environment*, 82, 9–16, <https://doi.org/10.1016/j.atmosenv.2013.09.042>, 2014.
- Uchiyama, S., Tomizawa, T., Tokoro, A., Aoki, M., Hishiki, M., Yamada, T., Tanaka, R., Sakamoto, H., Yoshida, T., Bekki, K., Inaba, Y., Nakagome, H., and Kunugita, N.: Gaseous chemical compounds in indoor and outdoor air of 602 houses throughout Japan in winter and 805 summer, *Environmental Research*, 137, 364–372, <https://doi.org/10.1016/j.envres.2014.12.005>, 2015.
- Uhde, E. and Salthammer, T.: Impact of reaction products from building materials and furnishings on indoor air quality-A review of recent advances in indoor chemistry, *Atmospheric Environment*, 41, 3111–3128, <https://doi.org/10.1016/j.atmosenv.2006.05.082>, 2007.
- Vichi, F., Mašková, L., Frattoni, M., Imperiali, A., and Smolík, J.: Simultaneous measurement of nitrous acid, nitric acid, and nitrogen dioxide by means of a novel multipollutant diffusive sampler in libraries and archives, *Heritage Science*, 4, 1–8, [https://doi.org/10.1186/s40494-016-0074-5](https://doi.org/10.1186/s40494-810 016-0074-5), 2016.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, I., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., van Mulbregt, P., Vijaykumar, A., Bardelli, A. P., Rothberg, A., Hilboll, A., Kloeckner, 815 A., Scopatz, A., Lee, A., Rokem, A., Woods, C. N., Fulton, C., Masson, C., Häggström, C., Fitzgerald, C., Nicholson, D. A., Hagen, D. R., Pasechnik, D. V., Olivetti, E., Martin, E., Wieser, E., Silva, F., Lenders, F., Wilhelm, F., Young, G., Price, G. A., Ingold, G.-L., Allen, G. E., Lee, G. R., Audren, H., Probst, I., Dietrich, J. P., Silterra, J., Webber, J. T., Slavič, J., Nothman, J., Buchner, J., Kulick, J., Schönberger, J. L., de Miranda Cardoso, J. V., Reimer, J., Harrington, J., Rodríguez, J. L. C., Nunez-Iglesias, J., Kuczynski, J., Tritz, K., Thoma, M., Neville, M., Kümmerer, M., Bolingbroke, M., Tartre, M., Pak, M., Smith, N. J., Nowaczyk, N., Shebanov, N., Pavlyk,

- 820 O., Brodtkorb, P. A., Lee, P., McGibbon, R. T., Feldbauer, R., Lewis, S., Tygier, S., Sievert, S., Vigna, S., Peterson, S., More, S., Pudlik, T., Oshima, T., Pingel, T. J., Robitaille, T. P., Spura, T., Jones, T. R., Cera, T., Leslie, T., Zito, T., Krauss, T., Upadhyay, U., Halchenko, Y. O., and Vázquez-Baeza, Y.: SciPy 1.0: fundamental algorithms for scientific computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Wang, H. and Morrison, G.: Ozone-surface reactions in five homes: Surface reaction probabilities, aldehyde yields, and trends, *Indoor Air*, 825 20, 224–234, <https://doi.org/10.1111/j.1600-0668.2010.00648.x>, 2010.
- Wang, H. and Morrison, G. C.: Ozone-Initiated Secondary Emission Rates of Aldehydes from Indoor Surfaces in Four Homes, *Environmental Science & Technology*, 40, 5263–5268, <https://doi.org/10.1021/es060080s>, 2006.
- Wang, Z., Shaw, D., Kahan, T., Schoemaeker, C., and Carslaw, N.: A modeling study of the impact of photolysis on indoor air quality, *Indoor Air*, 32, <https://doi.org/10.1111/ina.13054>, 2022.
- 830 Weschler, C. J.: Ozone in indoor environments: Concentration and chemistry, *Indoor Air*, 10, 269–288, <https://doi.org/10.1034/j.1600-0668.2000.010004269.x>, 2000.
- Weschler, C. J.: Changes in indoor pollutants since the 1950s, *Atmospheric Environment*, 43, 153–169, <https://doi.org/10.1016/j.atmosenv.2008.09.044>, 2009.
- Weschler, C. J., Wisthaler, A., Cowlin, S., Tamás, G., Strøm-Tejse, P., Hodgson, A. T., Destailats, H., Herrington, J., Zhang, J., and 835 Nazaroff, W. W.: Ozone-initiated chemistry in an occupied simulated aircraft cabin, *Environmental Science and Technology*, 41, 6177–6184, <https://doi.org/10.1021/es0708520>, 2007.
- WHO Air quality and health Guidelines Review Committee: World Health Organization Guidelines for Indoor Air Quality, Selected Pollutants, Tech. rep., World Health Organisation, <https://www.who.int/publications/i/item/9789289002134>, 2010.
- Wisthaler, A. and Weschler, C. J.: Reactions of ozone with human skin lipids: Sources of carbonyls, dicarbonyls, and hydroxy- 840 carbonyls in indoor air, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 6568–6575, <https://doi.org/10.1073/pnas.0904498106>, 2010.
- Wong, J. P., Carslaw, N., Zhao, R., Zhou, S., and Abbatt, J. P.: Observations and impacts of bleach washing on indoor chlorine chemistry, *Indoor Air*, 27, 1082–1090, <https://doi.org/10.1111/ina.12402>, 2017.
- World Health Organisation: WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide 845 and carbon monoxide, Tech. rep., 2001.
- Yao, M., Ke, L., Liu, Y., Luo, Z., and Zhao, B.: Measurement of ozone deposition velocity onto human surfaces of Chinese residents and estimation of corresponding production of oxidation products, *Environmental Pollution*, 266, 115215, <https://doi.org/10.1016/j.envpol.2020.115215>, 2020.
- Ye, W., Wang, H., Chen, Z., and Zhang, X.: Ozone deposition on free-running indoor materials and the corresponding volatile organic compound emissions: Implications for ventilation requirements, *Applied Sciences (Switzerland)*, 10, <https://doi.org/10.3390/APP10124109>, 850 2020.
- Zhou, S., Liu, Z., Wang, Z., Young, C. J., Vandenboer, T. C., Guo, B. B., Zhang, J., Carslaw, N., and Kahan, T. F.: Hydrogen Peroxide Emission and Fate Indoors during Non-bleach Cleaning: A Chamber and Modeling Study, <https://doi.org/10.1021/acs.est.0c04702>, 2020.