

Insights into secondary organic aerosol formation from the day- and nighttime oxidation of PAHs and furans in an oxidation flow reactor

Abd El Rahman El Mais^{1,2}, Barbara D'Anna², Luka Drinovec³, Andrew T. Lambe⁴, Zhe Peng⁵, Jean-Eudes Petit⁶, Olivier Favez¹, Selim Aït-Aïssa¹, Alexandre Albinet¹

5 ¹Ineris, Parc Technologique Alata, Verneuil-en-Halatte, 60550, France

²Aix Marseille Univ, CNRS, LCE, Marseille, France

³Center for Atmospheric Research, University of Nova Gorica, Nova Gorica, Slovenia


⁴Aerodyne Research, Inc. (ARI), Billerica, Massachusetts, USA

⁵CIRES and Department of Chemistry, University of Colorado, Boulder, Colorado 80309, USA

10 ⁶Laboratoire des Sciences du Climat et de l'Environnement (LSCE), 91190 Gif sur Yvette, France

Correspondence to: Alexandre Albinet (alexandre.albinet@ineris.fr)

Graphical abstract.



Abstract. Secondary organic aerosols (SOA) formed by oxidation of typical precursors largely emitted by biomass burning, such as PAHs and furans, are still poorly characterized in terms of formation yields, physical and light absorption properties, particularly those generated at night following reaction with nitrate radicals (NO₃). In the present study, we evaluated and compared the formation yields, effective density (ρ_{eff}), absorption Ångström exponent (α), and mass absorption coefficient (MAC) of laboratory-generated SOA from three furan compounds (furan, 2-methylfuran, and 2,5-dimethylfuran) and four PAHs (naphthalene, acenaphthylene, fluorene, and phenanthrene). SOA were generated in an oxidation flow reactor from the reaction between hydroxyl radicals (OH; 0.1 - 20 equivalent aging days) or NO₃ radicals (0.05 - 6 equivalent aging nights of 14 h) with single furan or PAH. The ρ_{eff} , formation yields, α , and MAC of the generated SOA varied depending on the precursor and oxidant considered. The ρ_{eff} of SOA formed with OH and NO₃ tended to increase with particle size before reaching a “plateau”. This was particularly evident for the nighttime chemistry experiments with NO₃ radicals (1.2 to 1.6 on average for particles ≥ 100 nm). Such results highlighted potential differences in the chemical composition of the SOA, as well as possibly in their morphology, according to the particle size. Three times lower SOA formation yields were obtained with NO₃ compared

15

20

25

to OH. The yields of PAH SOA (18 to 76%) were 5 to 6 times higher than those obtained for furans (3 - 12%). While furan SOA showed low or negligible light absorption properties, PAH SOA was found to have a significant impact in the UV-Visible region, implying a significant contribution to atmospheric brown carbon (BrC). No increase in the MAC values was observed from OH to NO₃ oxidation processes, probably due to a low formation of nitrogen-containing chromophores with NO₃ only (without NO_x). Overall, the results obtained in this work demonstrated that PAHs are significant precursors of SOA emitted by biomass burning, through both, day- and nighttime processes, and have a substantial impact on the aerosol light absorption properties and so probably on climate.

Keywords: Secondary organic aerosols (SOA), Formation yields, Density, Brown carbon, Biomass burning.

1 Introduction

Airborne particles (particulate matter, PM) have strong impacts on air quality, and climate (Monks et al., 2009; Fuzzi et al., 2015; IPCC, 2022). They have also been associated with adverse health effects, including respiratory and cardiovascular diseases, mortality, and morbidity (Pope et al., 2002; Brook et al., 2010; Thurston et al., 2016; Rajagopalan et al., 2018; Pope et al., 2020). Biomass burning is one of the major sources of fine PM (PM_{2.5}) in ambient air, particularly in the winter season due to residential wood combustion (RWC) used for heating purposes (Vicente and Alves, 2018; Crippa et al., 2013; Weber et al., 2019; Zhang et al., 2019, 2020; Chen et al., 2017; Srivastava et al., 2018b; Denier van der Gon et al., 2015; Chen et al., 2022a; Viana et al., 2016). This source also emits large amounts of volatile and semi-volatile organic compounds (VOCs and SVOCs) (Akherati et al., 2020; Bruns et al., 2016; Ahern et al., 2019; Hartikainen et al., 2018; Růžičková et al., 2022; Hatch et al., 2015, 2017, 2018; Baudic et al., 2016) that can undergo (photo-) chemical oxidation processes involving atmospheric oxidants, such as ozone (O₃), hydroxyl (OH) or nitrate (NO₃) radicals, resulting in the formation of Secondary Organic Aerosols (SOA) (Kroll and Seinfeld, 2008; Hallquist et al., 2009; Jimenez et al., 2009; Ziemann and Atkinson, 2012; Carlton et al., 2009; Heald and Kroll, 2020). OA constitute a significant fraction of fine PM (Bressi et al., 2021; Zhang et al., 2007, 2011), and SOA account for a large proportion of OA (up to 90%, depending on the location) (Zhang et al., 2007, 2011; Srivastava et al., 2018a; Kroll and Seinfeld, 2008). Identifying the major SOA precursors and investigating the physicochemical properties, formation yields, and chemical composition of SOA are crucial to implement efficient air quality policies.

Among the different VOCs and SVOCs emitted from RWC, some have been identified as major SOA precursors, such as monoaromatic hydrocarbons (benzene, toluene, xylenes...), phenols, furans, and polycyclic aromatic hydrocarbons (PAHs) (Yee et al., 2013; Bruns et al., 2016; Tiitta et al., 2016; Ahern et al., 2019; Akherati et al., 2020; Růžičková et al., 2022). Aromatic hydrocarbons (including PAHs), oxygenated aromatic (including phenolic species) and furans constitute a significant fraction of the total VOCs emitted by biomass burning [10 - 50% (with 10 - 20% of PAHs), 5 - 20%, and 5 - 40%, respectively]

(Hatch et al., 2015; Schauer et al., 2001; Bruns et al., 2016; Bhattu et al., 2019). The reactivity of PAHs and furans through homogeneous (in gaseous phase) or heterogenous (gas/particle) processes with different oxidants is well documented in the literature (Keyte et al., 2013; Jiang et al., 2020; Li et al., 2018; Ringuet et al., 2012; Aschmann et al., 2011, 2014; Al Ali et al., 2022; Bierbach et al., 1995; Newland et al., 2022; Kind et al., 1996; Zheng et al., 2006). However, the level of scientific understanding in terms of SOA formation from such species is still limited. Different studies available in the literature have been focused on the formation yields, chemical composition, and physicochemical properties of SOA formed from PAHs and furans (Lee and Lane, 2009; Chan et al., 2009; Lee and Lane, 2010; Lee et al., 2012; Shakya and Griffin, 2010; Kleindienst et al., 2012; Zhou and Wenger, 2013a, b; Chen et al., 2016; Riva et al., 2017; Kautzman et al., 2010; Gómez Alvarez et al., 2009; Strollo and Ziemann, 2013; Jiang et al., 2019b; Joo et al., 2019a; Tajuelo et al., 2021; Srivastava et al., 2022; Joo et al., 2019b; Jiang et al., 2019b; Chen et al., 2022b; Jiang et al., 2019a). However, most of these studies have been conducted with OH radicals and/or ozone, and only a small number of them have investigated the SOA formation with nitrate radicals (NO₃). While OH and O₃ are the key atmospheric oxidants during the day, NO₃ is known to be the major one during the night (Brown and Stutz, 2012). The study of NO₃ radical chemistry is crucial for describing winter pollution, when RWC is more relevant, and when the night lasts longer than the day. Several authors have reported its significance in the formation of SOA from RWC emissions (Kodros et al., 2020; Jorga et al., 2021; Kodros et al., 2022; Tiitta et al., 2016). The current underestimation of OA concentrations by air quality models by a factor of 3 to 5 in winter might be partly due to the neglected nighttime chemistry (Fountoukis et al., 2016; Mircea et al., 2019; Tsimpidi et al., 2014). Recent simulation results including this nighttime chemistry have shown that more than 70% of the OA from biomass combustion are significantly influenced by the oxidation processes that take place in the absence of light, i.e., involving nitrate radical (Kodros et al., 2020).

Finally, atmospheric brown carbon (BrC) aerosols, which primarily absorb light in the shorter visible to the ultraviolet (UV) wavelengths, have been recognized to play a critical role in the Earth's radiative balance (Hems et al., 2021; Laskin et al., 2015; Moise et al., 2015; Andreae and Gelencsér, 2006). BrC can also impact black carbon (BC) light absorption due to the so-called lensing effect under internal mixing conditions (Zhang et al., 2018; Saleh et al., 2015). On the other hand, the aging of BrC-containing OA may result in decreasing their absorptivity (Sumlin et al., 2017). Quantifying the BrC contribution to light absorption is therefore essential for an accurate interpretation of the aerosol optical depth (AOD), the atmospheric column's light extinction due to both scattering and absorption, and regional climate. Numerous studies have quantitatively characterized the parameters governing the optical absorption properties (e.g. absorption angström exponent (α), Mass Absorption Coefficient (MAC), and Refractive index) of SOA formed from various biogenic and anthropogenic precursors (Lambe et al., 2013; Liu et al., 2016; Xie et al., 2017a; Dingle et al., 2019; Siemens et al., 2022; Laskin et al., 2015; Jiang et al., 2019a; Metcalf et al., 2013; Klodt et al., 2023; He et al., 2022; Chen et al., 2022b; Hems et al., 2021; Moise et al., 2015). However, the information regarding SOA formed from PAHs (other than naphthalene) and furans is still far from complete (Cheng et al., 2020). Besides, as particle density is linked to optical properties due to its dominant role in the effects on refractive index, its accurate determination is crucial for the evaluation of SOA radiative forcing (Liu and Daum, 2008).

Overall, density is a key physical property of particles because it influences transport properties and so, the fate of particles in both the atmosphere and the human respiratory system (Seinfeld and Pandis, 1998; Finlayson-Pitts and Pitts Jr, 2000).

The main objectives of this study were to evaluate and compare the formation yields, physical (granulometry and effective densities (ρ_{eff}), and light absorption properties (α and MAC) of the SOA formed from typical precursors emitted by biomass burning, namely PAHs and furans following OH and NO₃-initiated oxidation and aging processes.

2 Experimental methods

2.1 Generation of radicals and SOA using OFR

Most of the previous studies focusing on the SOA formation from PAHs and furans have been conducted in environmental (smog) chambers. Oxidation Flow Reactors (OFR) are an alternative tool to simulate atmospheric aging (Peng and Jimenez, 2020). Several works have shown the reliability, comparability, and complementary of the results provided using such systems in the study of SOA formation processes (Bruns et al., 2015; Peng and Jimenez, 2020). Their application in the study of SOA formation from key precursors emitted by biomass burning is of prime interest (Hodshire et al., 2019; Srivastava et al., 2022). In our study, the stable (up to 17 hours) generation of SOA (Figure S1 of the Supplementary Material, SM) has been carried out at ambient temperature ($T = 24 - 32$ °C) and environmentally relevant relative humidity ($RH = 25 - 62\%$) in a Potential Aerosol Mass Oxidation Flow Reactor (PAM-OFR, Aerodyne Research) (Kang et al., 2007; Lambe et al., 2011), which is a 13.3 L aluminum horizontal cylindrical chamber (46 cm long \times 22 cm ID) operated in a continuous flow mode (Figure 1). Three furans (furan (99% purity), 2-methylfuran (2-MF, 99%) and 2,5-dimethylfuran (2,5-DMF, 99%)), and four PAHs (naphthalene (Naph, 99%), acenaphthylene (Acy, 75%, the remaining 25% are acenaphthene (Ace)), phenanthrene (Phe, 99%), and fluorene (Flu, 98%)) have been selected as SOA precursors and studied here. All were purchased from Sigma Aldrich (France) and injected into the PAM-OFR in their pure state, in the absence of NO_x, and without seed particles. The injection method was determined by the state of the different substances under ambient temperature conditions. Liquid furans were injected, at a constant flow, through a 0.0152 cm i.d. Teflon tubing using a microliter syringe pump (TriContinent C24000, 100 μ L, syringe plunger velocity of 3 to 17 μ steps s⁻¹ (= 5.63 to 31.88 μ L min⁻¹) depending on the compound studied and concentration generated) and subsequently nebulized/vaporized using a clean air flow of 2 L min⁻¹ adjusted by a Mass Flow Controller (MFC, MKS G series) before being introduced into the PAM-OFR. PAHs were vaporized by circulating heated clean air through a metallic tube packed with the corresponding solid compounds (50 - 3500 mL min⁻¹, depending on the PAH studied and concentration generated, regulated at 30 °C using a CROCO-CIL liquid chromatography column oven). A filtered air supply (TSI 3074B) combined with an air zero generator (Claind AZ air purifier 2010) were used to deliver clean-dry air and maintain the total carrier gas flow inside the PAM-OFR at 9.5 L min⁻¹.




Figure 1. Simplified schematic diagram of the experimental set-up used to generate and monitor the resulting SOA.

120

To simulate the daytime chemistry, OH radicals were produced by photolysis of O₂ by four UV lamps emitting at $\lambda = 185$ and 254 nm (OFR185 method) (Li et al., 2015). Briefly, the photolysis of O₂ at 185 nm produces O₃ which is then photolyzed at 254 nm to produce O(¹D). O(¹D) then reacted with water vapor (introduced using a Nafion membrane humidifier, Perma Pure LLC, FC100-80-6-MKK) to generate OH radicals. The lamps' voltage was set, depending on the precursor studied, between
 125 1.8 and 3 V, and the corresponding irradiance (I_r), ranging between 6 and 126 $\mu\text{W cm}^{-2}$ (Table 1) was continuously measured by the PAM-OFR photodiode (Tocon_C6, sglux). A UV photometric ozone analyzer (Model 202, 2B Technologies) has been used to monitor the O₃ concentrations in the PAM-OFR.

For nighttime chemistry, NO₃ radicals were generated from the thermal decomposition of N₂O₅ at room temperature in a dark OFR (OFR-iN₂O₅ method) (Lambe et al., 2020). Briefly, NO₃ and N₂O₅ were generated in the gas phase following the reaction of separate reagent flows containing approximately 100 ppm NO₂ (1% in N₂, Air Liquide) and 200 ppm O₃ in a 152.4 cm long × 2.22 cm ID perfluoroalkoxy (PFA) laminar flow reactor (LFR) coupled to the PAM-OFR. To achieve these reagent concentrations, the NO₂ flow (1% in N₂) was set at 20 cm³ min⁻¹, and 2 L min⁻¹ of pure O₂ (99.9995%, Air Liquide) was passed through an ozone chamber housing a mercury fluorescent lamp. The O₃ mixing ratio inside the LFR was measured using an additional ozone analyzer (Model 202, 2B Technologies). In such conditions, NO₂ concentrations in the PAM-OFR (NO_x
 130 Analyzer 42i-HL, Thermo Scientific) were below the **quantification limit of 1 ppm (in the range used of 0 - 200 ppm)**.

135

Table 1. Experimental conditions used to study the SOA formation from selected PAHs and furans under daytime conditions (OH radical, OFR185).

SOA precursor	SOA precursor concentration [VOC] ₀		Lamps voltage (V)	Ir (μW cm ⁻²)	RH (%)	T (°C)	[O ₃] _{PAM} (ppm)	Equivalent aging days	
	(mg m ⁻³)	(ppmv)						Experimental evaluation	KinSim calculations
2-methylfuran	8.99 ± 0.20	2.73 ± 0.05	3.0	71	38	30	15	6	9
2,5-dimethylfuran	17.68 ± 0.02	4.58 ± 0.01	2.8	63	38	29	12	3	6
Furan	18.41 ± 0.05	6.73 ± 0.03	2.5	34	34	29	7	nd ¹	0.1
Naphthalene	2.60 ± 0.17	0.50 ± 0.03	1.8	6	41	26	3	3	2
Acenaphthylene	0.33 ± 0.02	0.05 ± 0.00	1.8	14	39	30	4	8	12
Fluorene	0.43 ± 0.01	0.06 ± 0.00	3.0	70	39	29	11	17	20
Phenanthrene	0.19 ± 0.02	0.03 ± 0.00	3.0	58	27	29	10	11	15

¹not determined.

Table 2. Experimental conditions used to study the SOA formation from selected PAHs and furans under nighttime conditions (NO₃ radical, OFR-iN₂O₅).

140

SOA precursor	SOA precursor concentration [VOC] ₀		[O ₃] _{0,LFR} (ppm)	[NO ₂] _{0,LFR} : [O ₃] _{0,LFR}	RH (%)	T (°C)	[O ₃] _{PAM} (ppm)	Equivalent aging nights of 14 h
	(mg m ⁻³)	(ppmv)						
2-methylfuran	41.80 ± 0.10	12.52 ± 0.02	194	0.52	49	26	11	0.1
2,5-dimethylfuran	50.23 ± 0.11	12.82 ± 0.03	179	0.56	53	25	6	0.1
Furan	18.44 ± 0.00	6.67 ± 0.00	200	0.5	62	26	10	*
Naphthalene	15.10 ± 0.11	2.89 ± 0.02	194	0.51	43	25	12	5.7
Acenaphthylene	3.05 ± 0.10	0.49 ± 0.02	183	0.55	49	25	13	1.7
Fluorene	1.07 ± 0.10	0.16 ± 0.01	198	0.5	45	26	8	**
Phenanthrene	0.26 ± 0.01	0.04 ± 0.00	200	0.5	43	25	10	5.6

*Unstable furan injection inducing an unstable SOA generation.

**No SOA generation observed.

2.2 Estimation of the injected SOA precursor concentrations

145 Furans and PAH concentrations introduced into the PAM-OFR were established in the range of 8 to 50 mg m⁻³ (3 to 13 ppmv) and 0.19 to 15.1 mg m⁻³ (0.03 to 3 ppmv), respectively (Tables 1 and 2). These concentrations were relatively high and may represent concentrations encountered in large forest fire plumes. Besides the study of the SOA formation, our objective was to produce and collect enough quantity of PM (collected on filters, sampling flow of 6.5 L min⁻¹, Figure 1) to further perform *in vitro* biological assessment (El Mais, 2023).

150 Furan concentrations were calculated from the syringe pump injection flow rate, temperature, analyte molecular weight, density, and dilution ratio into the OFR carrier gas (Eq. S1). PAH concentrations were evaluated during spare experiments under the same conditions as for SOA generation, but without oxidants (PAM-OFR lights off, and no O₃ and NO₂ injected into

the LFR, only O₂). Particulate and gaseous phases were collected at the PAM-OFR exit on quartz fiber filters (Pallflex Tissuquartz, Ø = 47 mm) and polyurethane foams (PUFs; Tisch Environmental, 1.5 × 3 inches), respectively. Prior to sampling, filters were pre-heated at 500 °C for 12 h, while PUFs were pre-cleaned using pressurized solvent extraction (ASE 350, Thermo; one hexane cycle followed by two acetone cycles: 80 °C, 100 bars, 5 min heat time, 15 min static time) (Zielinska, 2008). Sampling durations were 15 and 30 min, and experiments were performed in duplicate for each duration. One or two experimental blanks (one for each sampling duration), for each PAH and oxidant, were also collected with no PAH injection into the PAM-OFR. After collection, samples were wrapped in aluminum foils, placed in zip bags, and then stored at -20 °C until analysis. The filter and its associated PUF (28 samples and 12 blanks in total) were extracted together using pressurized liquid extraction (Dionex, ASE 200, 80 °C, 100 bars, 5 min heat time, 15 min static time, 2 cycles) with acetonitrile as the solvent (VWR, HPLC grade). Prior to extraction, a known amount of 6-methylchrysene was added to the samples and used as a surrogate standard to check the extraction efficiency (ranging from 88 to 108%). The extracts were then directly quantified (no reduction step to avoid any loss of PAH by evaporation, only a 10 times dilution was applied) by UPLC-Fluorescence-UV (Ultimate 3000, Thermo Scientific) using a C18 UPLC column (Zorbax Eclipse PAH, 2.1 mm × 150 mm × 1.8 µm, Agilent, 1 µL injected).

2.3 Online SOA characterization

The SOA generated have been monitored and characterized using a set of different online instrumentations (Figure 1). Organic aerosol mass and chemical composition of the non-refractory aerosol fraction were measured using a Quadrupole Aerosol Chemical Speciation Monitor (Q-ACSM; Aerodyne Research Inc.), equipped with a PM_{2.5} aerodynamic lens, using a 1 min time resolution. The aerosols were dried before analysis using a Nafion dryer system. Calibration of the detector response factor was performed by using ammonium nitrate and sulfate solutions (Ng et al., 2011; Crenn et al., 2015; Freney et al., 2019). A Relative ion efficiency (RIE) of 1.4 was applied to organic matter. The instrument was equipped with a capture vaporizer so that a fixed collection efficiency (CE) of 1 could be used for the whole ACSM dataset (Xu et al., 2017). Connected in parallel with the Q-ACSM, a Condensation Particle Counter (CPC; Grimm version 5.403), and a Scanning Mobility Particle Sizer (SMPS, composed of a Differential Mobility Analyzer (DMA) 3081, an electrostatic classifier 3080, and a CPC 3775; TSI, USA) were used to monitor the total particle number concentration (in the range of 4.5 nm to 3 µm, 1 s time resolution) and the particle size distribution (from 14.6 to 661.2 nm, 5 min time resolution). SOA effective density (ρ_{eff}) was evaluated by combining, in series, a DMA, a Centrifugal Particle Mass Analyzer (CPMA, Cambustion) (Olfert and Collings, 2005; Olfert et al., 2006; Johnson et al., 2013) and a CPC, all connected with the Q-ACSM (instead the SMPS). The mass of the generated SOA was determined for a given particle size (mobility diameter) over a range of 30 to 200 nm allowing the evaluation of the SOA ρ_{eff} according to the particle size. Such an approach has already been used to determine the effective density of particles of different types of primary emissions (vehicular, flame soot) and SOA (Peng et al., 2021; Malloy et al., 2009), but only few data exist for SOA formed from anthropogenic and biogenic precursors. Finally, a 7-wavelengths (370, 470, 520, 590, 660,

880, and 950 nm) aethalometer (AE33, Magee scientific, 1 min time resolution) was also connected at the exit of the PAM-
185 OFR. Such online filter tape-based optical measurement method is commonly used for the measurement of black carbon
concentrations (BC, at 880 nm) in ambient air as well as for the determination of BrC aerosol fraction (Drinovec et al., 2015;
Zhang et al., 2020; Drinovec et al., 2017). Here it was used to tentatively assess the SOA light absorption properties and to
compare the results obtained with the most conventional techniques in the literature such as photoacoustic spectrometry or on
aerosol sample extract by UV-Vis spectroscopy (Moise et al., 2015).

190 2.4 Calculations

2.4.1 OH and NO₃ exposures

The OH exposure (OH_{exp}) was determined experimentally by continuously measuring the decay of SO₂ injected into the PAM-
OFR (200 ppb as initial concentration) during spare experiments with the same conditions as for SOA generation, using a SO₂
analyzer (AF 21 M Environment S.A) (Lambe et al., 2015) and applying Eq. (1)

$$OH_{exp} = -\frac{1}{k_{SO_2}^{OH}} \times \ln\left(\frac{[SO_2]}{[SO_2]_0}\right) \quad (1)$$

195 where $k_{SO_2}^{OH}$ is the rate coefficient of the reaction between OH and SO₂, equal to $9.4 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Davis et al.,
1979). $[SO_2]_0$ was the initial SO₂ concentration injected into the PAM-OFR, and $[SO_2]$ was the final concentration after
oxidation. The OH exposure ranged from 3.74×10^{11} to 2.19×10^{12} molecules $\text{cm}^{-3} \text{ s}$ (Table S1), corresponding to 3 - 17
equivalent aging days, respectively (Table 1), assuming a daily average OH radical concentrations of 1.5×10^6 molecules cm^{-3}
(Finlayson-Pitts and Pitts Jr, 2000; Mao et al., 2009).

200 In addition to the experimental evaluation, OH_{exp} was evaluated using the KinSim chemical kinetic solver (Peng and Jimenez,
2019). Inputs to the OFR KinSim model included Pressure (P), T, RH, $[VOC]_0$ (Table 1), total residence time ($\tau = 84 \text{ s}$), photon
fluxes at 185 and 254 nm (I_{185} and I_{254} , respectively) (Table S2), bimolecular rate coefficients of each precursor with OH or
O₃, and photoabsorption cross sections (σ_i) of each precursor at 185 and 254 nm (Table S3).

The obtained OH_{exp} ranged from 1.05×10^{10} to 2.57×10^{12} molecule $\text{cm}^{-3} \text{ s}$ (Table S1), corresponding to about 0.1 - 20
205 equivalent aging days, and were comparable, even though larger, with the experimental values (Table 1). These differences
seemed to be due to a significant rise in OH concentrations after the VOC consumption, which proportionally influenced the
modeled OH_{exp} values (Figure S2).

NO_{3exp} was estimated theoretically using a KinSim mechanism adapted from previous work (Lambe et al., 2020). Model
parameters included P, T, RH, $[VOC]_0$, $[O_3]_{0, LFR}$ (Table 2), τ of 109 s, $[NO_2]$ of 100 ppm, and the bimolecular rate coefficients
210 of each precursor with NO₃ or O₃ (Table S3). NO_{3exp} values ranged from 1.68×10^{12} to 2.10×10^{14} molecule $\text{cm}^{-3} \text{ s}$ (Table S1),
corresponding to about 0.05 to 5.7 equivalent nights of 14 h (Table 2) at an average nighttime NO₃ mixing ratio of 7.36×10^8
molecules cm^{-3} (Asaf et al., 2010). Again, most of the NO₃ exposure was due to the modeled increase of NO₃ concentrations
after VOC consumption (especially in the cases of Acy, 2-MF, and 2,5-DMF, Figure S2).

2.4.2 Effective density

215 The effective density (ρ_{eff}) of SOA particles was calculated using Eq. (2) as the ratio of the measured particle mass (corresponding to the mode in the scan data (m_p)) to the calculated particle volume, assuming a spherical particle (shape factor = 1) with a diameter equals to the electrical equivalent mobility diameter (d_m) selected by the DMA (McMurry et al., 2002; DeCarlo et al., 2004):

$$\rho_{eff} = \frac{6}{\pi} \frac{m_p}{(d_m)^3} \quad (2)$$

2.4.3 SOA yields

220 SOA yields were calculated using Eq. (3):

$$SOA \text{ yield} = \frac{\text{mass of SOA produced}}{\text{mass of VOC reacted}} \quad (3)$$

where the mass of VOC reacted with the corresponding precursor was obtained from the KinSim model (see section 3.1), and the mass of SOA was determined from Q-ACSM measurements as the SOA generation was stable (Figure S1). SOA was considered equivalent to OM for OH radical experiments, and equal to OM + NO₃ for NO₃ radical experiments. Measurement of particle transmission through the PAM-OFR, in the range of 30 - 200 nm, using ammonium sulfate aerosols showed particle
225 loss below 5% for all PM sizes, therefore no particle wall-loss correction was applied.

2.4.4 Optical absorption coefficient (b_{abs}), absorption Ångström exponent (α), and Mass absorption coefficient (MAC)

The AE33 has been designed for automatic correction of the so-called filter loading effect (Drinovec et al., 2015, 2017). Briefly, the sampled ambient air is divided, and the sample is deposited onto two filter spots at different flow rates, leading to uneven loadings on the respective filter spots. A compensation parameter $k(\lambda)$ is retrieved from the different loading effect
230 magnitudes influencing these two spots, and $k(\lambda)$ is further used to determine the light attenuation due to carbonaceous aerosols. However, initial results obtained, mainly for the PAH SOA, showed a poor compensation efficiency and large jumps during the spot changes. Manual compensation of AE33 data was therefore applied using a fixed filter loading compensation parameter $k(\lambda)$ (Table S4) to get usable data (Figure S3).

Light absorption coefficients (b_{abs}) were calculated from the manually compensated BC values according to Eq. (4).

$$b_{abs}(\lambda) = BC(\lambda) \times \sigma_{air}(\lambda) \times \frac{C_{Manufacturer}}{C_{new}} \quad (4)$$

235 where σ_{air} is the air mass absorption efficiency (Table S5). Comparison between AE33 and photothermal interferometer PTAAM-2 λ showed that a higher value of multiple scattering parameter C should be used (Drinovec et al., 2022) than the one provided by the manufacturer ($C_{Manufacturer} = 1.41$). A new C value (C_{new}) of 3.75 was selected for the measured particle sizes. The corrected absorption is smaller by the factor $C_{new}/C_{Manufacturer}$.


Absorption Ångström exponents (α) were determined as the absolute value of the slope of $\ln(b_{\text{abs}})$ as a function of $\ln(\lambda)$ from 240 370 to 590 nm. The fitting parameters are summarized in Table S6.

Mass absorption coefficients (MAC) were finally calculated as the ratio of b_{abs} to the SOA mass concentrations evaluated from Q-ACSM measurements using Eq. (5).

$$MAC(\lambda) = \frac{b_{\text{abs}}(\lambda)}{\text{SOA mass concentration}} \quad (5)$$

3 Results and discussion

3.1 Understanding of oxidants (and photolysis) competition in the PAM-OFR for both oxidation methods



245

Figure 2. Relative importance of PAH and furan loss pathways inside the PAM-OFR during both OFR185 (daytime) and OFR-iN₂O₅ (nighttime) modes. No results are shown for Furan and Flu with NO₃ radicals due to an unstable SOA generation or no SOA formed.

Figure 2 shows the competition between the oxidants, as well as direct photolysis, on the reactivity of the PAHs and furans 250 studied inside the OFR.

In the case of OFR185 (OH reactivity), the predominant reaction of PAHs was observed with OH radicals (> 92%). The remaining photolyzed at 254 nm. Both ozonolysis and photolysis at 185 nm had negligible impact on the reactivity of PAHs. For furans, photolysis at both wavelengths was negligible (1%), and the competition between OH and O₃ differed depending on the compound considered. In the case of furan, ozonolysis was negligible, and the majority (96%) reacted with OH. This
255 contrasted with 2-MF and 2,5-DMF, where O₃ competed with OH to consume 11% of 2-MF, and nearly half of 2,5-DMF. In the OFR-iN₂O₅ mode (NO₃ reactivity), only O₃ and NO₃ were in competition. For Naph and Phe, O₃ had no impact. However, ozonolysis is significant for Acy (+ 25% of Ace) (20%) or even predominant for 2-MF (46%) and 2,5-DMF (70%). These results showed that in our experimental conditions, the reaction with OH or NO₃ were largely the predominant ones, except for 2-MF and 2,5-DMF. These results were used to calculate the exact concentrations of the SOA precursor that reacted
260 to get the most accurate evaluation of the SOA yields.

3.2 SOA granulometry and effective density

Figure 3 shows the average number-weighted mobility size distributions obtained for the generated SOA under both day- and nighttime oxidation processes.

Overall, the observed particle size distributions were all monomodal and showed a shift towards larger particle sizes according
265 to the precursor concentrations injected into the PAM-OFR. For instance, for daytime chemistry, the injected concentrations of Phe, Acy, and Flu were about 0.19 to 0.43 mg m⁻³, and the resulting SOA particle distributions were centered around 32 nm; while the particle size distributions for Naph, 2-MF, 2,5-DMF, and Furan were centered around 55 and up to 100 nm due to the larger precursor concentrations injected (2.6 to 18 mg m⁻³) (Table 1). Similarly, for NO₃ radical experiments, as the precursor concentrations of Naph, Acy, and 2-MF were 4.6 to 10 times larger than during OH radical experiments (Table 2),
270 a strong shift in the size distribution towards larger particles, centered around 200 nm, was observed. Interestingly, for similar precursor concentrations, the SOA formed from Phe by reaction with NO₃ also shifted to a large particle size of about 85 nm. Moreover, the number of aerosol particles produced from Naph, Acy, Phe, and 2-MF by OH oxidation was 10 - 100 times larger compared to NO₃ oxidation. The aerosol particles from NO₃ oxidation grew substantially larger since fewer of them were available for semi-volatiles species produced to condense onto. For 2,5-DMF, the particle size distribution shifted towards
275 smaller particles (D_p ~ 35 nm) even though the concentration injected was about 3 times higher than during OH exposure studies. This might be also linked to the lower particle concentrations in the system observed with OH oxidation.




Figure 3. Comparison of the particle size distributions, in number, of the PAHs and furans SOA formed under day and nighttime conditions (OH and NO₃ radicals respectively). No results are shown for Furan and Flu with NO₃ radicals due to unstable SOA generation or no SOA formed.

280

The effective densities of the generated SOA with OH and NO₃ radicals as a function of the aerosol mobility diameter are presented in Figure 4.

Generally, the SOA ρ_{eff} tended to increase with the particle size in the range of 30 to 100 or 150 nm for both reactivities. A “plateau” was then observed especially in the case of the SOA formed by reaction with NO₃ radical. These results highlighted potential differences in the chemical composition of the SOA, as well as possibly in terms of their morphology, with the particle size which should be further investigated.

The average ρ_{eff} for each precursor are presented in Table 3. Values for particle diameter ≥ 100 nm were considered according to the “plateau” observed and assuming that this range of particles contributes significantly to the particle mass. Relatively large variations can be observed depending on the precursor and oxidant being studied, ranging from 1.23 to 1.61 g cm⁻³. Between OH and NO₃ oxidation, the effective densities of Naph and Acy SOA increased from 1.33 to 1.39 g cm⁻³ and from 1.23 to 1.47 g cm⁻³, respectively. In contrast, the effective densities of Phe and 2-MF SOA decreased from 1.52 to 1.29 and to 1.38 g cm⁻³, respectively, while they remained comparable for 2,5-DMF. The measured Naph SOA ρ_{eff} were consistent with previously reported values (1.35 - 1.55 g cm⁻³) from smog chamber experiments at low and high NO_x conditions (Chan et al., 2009; Chen et al., 2016, 2018; He et al., 2022). In addition, densities of 1.24 and 1.40 g cm⁻³ have been respectively reported


290

295 for 3-methylfuran SOA (Joo et al., 2019a) and furan SOA (Chen et al., 2022b), both formed by reaction with NO₃. Overall the values determined here were in agreement with previously reported results for SOA formed from different anthropogenic precursors (1.06 - 1.65 g cm⁻³) (Nakao et al., 2013; Kostenidou et al., 2007; Hallquist et al., 2009). In the absence of density measurements, an SOA density of 1.4 g cm⁻³ is usually assumed for smog chamber experiments (Shakya and Griffin, 2010; Riva et al., 2017; Jiang et al., 2019b; Hallquist et al., 2009). Our results confirm this default value as a fairly good
 300 approximation, but also that it may need to be refined according to the given investigated compounds, for instance, for determining the particle transport properties, and converting PM size distributions into mass concentrations (McMurry et al., 2002). Therefore, and as reported before, our study shows the importance of figuring out the density for individual experiments to **reduce the potential biases in quantitative analyses and SOA yields evaluation** (Chen et al., 2016), as well as to assess the optical properties of the studied SOA.

305

Table 3. Average effective densities of the PAHs and furans SOA formed under day and nighttime conditions (OH and NO₃ radicals respectively). Results obtained from the mean of the densities corresponding to SOA with d_m > 100 nm.

Precursor	Aerosol effective densities (g cm ⁻³)	
	OH radical	NO ₃ radical
Naphthalene	1.33 ± 0.11	1.39 ± 0.06
Acenaphthylene	1.23 ± 0.05	1.47 ± 0.13
Fluorene	1.60 ± 0.10	No SOA formation
Phenanthrene	1.52 ± 0.02	1.29 ± 0.11
2-methylfuran	1.52 ± 0.06	1.38 ± 0.06
2,5-dimethylfuran	1.35 ± 0.08	1.35 ± 0.08
Furan	1.61 ± 0.05	Not determined



310 **Figure 4.** Comparison of the effective density of the PAHs and furans SOA formed under day and nighttime conditions (OH and NO₃ radicals respectively) as a function of the aerosol mobility diameter. The error bars correspond to the standard deviations (2σ) on the number of measurements replicates ($n=2$ to 4). No results are shown for Furan and Flu with NO₃ radicals due to unstable SOA generation or no SOA formed.

3.3 SOA yields

315 Figure 5 presents the SOA yields for the studied PAHs and furans under day- and nighttime oxidations as a function of the radical exposure used for each precursor.

The SOA yields from the oxidation of PAHs with OH radicals were all comparable, especially for Naph, Acy, and Flu, ranging from 62 to 76%. The differences observed, including the highest SOA yield for Phe, could be related to the differences in OH exposures ranging from 2.78×10^{11} to 2.57×10^{12} molecules cm^{-3} s (Table S1). The same has been observed for furans with comparable SOA yields (9 - 12%) even if the range of tested OH exposures was largely variable (from 1.05×10^{10} to 1.13×10^{12} molecule cm^{-3} s). For NO_3 -initiated SOA formation, approximately 3 times lower SOA yields were observed relative to OH-initiated SOA formation for most precursors. Only Acy showed an average SOA yield of 44% with NO_3 comparable to that observed with OH radical chemistry (68%) even though the NO_3 exposure tested was the lowest of the PAH group. Table 4 compares the calculated SOA yield with the literature. Most of the previous works reported smog chamber experiments with OH radicals, under low or high NO_x regimes and dry conditions (RH < 10%) (Chan et al., 2009; Shakya and Griffin, 2010; Kleindienst et al., 2012; Chen et al., 2016; Riva et al., 2017; Chen et al., 2018; Gómez Alvarez et al., 2009; Jiang et al., 2019b; Tajuelo et al., 2021). For the NO_3 radical with furan, only one previous study has reported an SOA yield of 7% (Jiang et al., 2019a). Joo et al. (2019a) reported SOA yields of 1.6 to 2.4% for 3-methylfuran in smog chamber dry experiments. As the experimental conditions are quite different, the reported SOA yields are also highly variable, yet our results lie on the upper range of previously reported works. Our results are 2 to 3 times higher than those of the single study performed using an OFR (Wang et al., 2018) for which no information on OH exposure is available. Several other factors have an influence the SOA yields such as the NO_x concentrations (or ratio VOC/ NO_x), the use and type of seed particles, PM (organic) mass concentrations, temperature and RH conditions (Srivastava et al., 2022; Lambe et al., 2015; Zhang et al., 2023). It has also been shown recently that smog chamber and OFR studies probably overestimate SOA formation yields compared to atmospheric conditions due to high peroxy radical (RO_2) concentrations leading to overemphasis of cross reactions of RO_2 compared to the reaction with HO_2 (or NO_3) (Schervish and Donahue, 2021). Thus, the high VOC and PM (OA) concentrations together with RH% and $\text{RO}_2 + \text{RO}_2$, $\text{RO}_2 + \text{HO}_2$ or $\text{RO}_2 + \text{NO}_3$ reaction processes, may have enhanced the SOA formation here, explaining the differences observed with literature data (Table 4). Overall, SOA yields from furans oxidation were about 5 - 6 times lower than those from PAHs, implying that the contribution of furans to SOA formation from biomass burning emissions seems limited, whereas the PAHs contribution is probably significant during both, day- and nighttime periods.




Figure 5. Comparison of the PAHs and furans SOA yields obtained from the day- and nighttime oxidation processes (with OH and NO₃ radicals respectively). Results evaluated using Q-ACSM measurements and expressed as a function of OH or NO₃ exposures obtained from KinSim simulations. No results are shown for Furan and Flu with NO₃ radicals due to unstable SOA generation or no SOA formed.

Table 4. Comparison of the SOA yields obtained for the different PAHs and furans studied and oxidation conditions (using the PAM-OFR) with values reported in the literature (experimental conditions are specified).

SOA precursor	Experimental conditions	PM mass ($\mu\text{g m}^{-3}$)	SOA yield (%)	References
Naphthalene	OH, PAM-OFR, RH = 41%	1473 - 1973	61.5 - 69.9	This study
	NO ₃ , PAM-OFR, RH = 43 %	2347 - 2911	15.7 - 19.5	This study
	OH, smog chamber, low NO _x , 5 < RH < 8%, seed	10 - 29	73	Chan et al., (2009)
	OH, smog chamber, high NO _x , 5 < RH < 8%, seed	8 - 28	19 - 30	Chan et al., (2009)
	OH, smog chamber, low NO _x , RH < 5%	4 - 18	8 - 16	Shakya and Griffin, (2010)
	OH, smog chamber, low NO _x , RH < 3%, seed	22 - 158	18 - 36	Kleindienst et al., (2012)
	OH, smog chamber, high NO _x , RH = 30%, seed	39 - 132	11 - 29	Kleindienst et al., (2012)
	OH, smog chamber, low NO _x , RH < 0.1%	6 - 50	4 - 31	Chen et al., (2016)
	OH, smog chamber, high NO _x , RH < 0.1%	3 - 95	3 - 60	Chen et al., (2016)
	OH, smog chamber, low NO _x , RH < 0.1%	40 - 80	21 - 50	Chen et al., (2018)
	OH, OFR, RH = 57%	not indicated	28 ± 6.7	Wang et al., (2018)
Acenaphthylene	OH, PAM-OFR, RH = 39%	557 - 1340	62.4 - 78.2	This study
	NO ₃ , PAM-OFR, RH = 49%	1177 - 1502	39.0 - 49.8	This study
	OH, smog chamber, low NO _x , RH < 5%	1 - 15	4 - 13	Shakya and Griffin, (2010)
	OH, smog chamber, low NO _x , RH < 1%	1120	61	Riva et al., (2017)
	OH, smog chamber, high NO _x , RH < 1%	730	46	Riva et al., (2017)
Fluorene	OH, PAM-OFR, RH = 39%	174 - 331	54.9 - 67.9	This study
Phenanthrene	OH, PAM-OFR, RH = 27%	108 - 161	59.1 - 88.1	This study
	NO ₃ , PAM-OFR, RH = 43%	44 - 53	16.7 - 20.1	This study
	OH, OFR, RH = 57%	not indicated	12 ± 2.6	Wang et al., (2018)
2-methylfuran	OH, PAM-OFR, RH = 38%	500 - 985	12.3 - 12.5	This study
	NO ₃ , PAM-OFR, RH = 49%	739 - 899	3.3 - 3.5	This study
	OH, smog chamber, high NO _x , dry conditions	13	5.5 ± 1.6	Gómez Alvarez et al., (2009)
2,5-dimethylfuran	OH, PAM-OFR, RH = 38%	810 - 884	9.0 - 9.8	This study
	NO ₃ , PAM-OFR, RH = 53%	598 - 721	4.0 - 4.4	This study
	OH, smog chamber, 25 < RH < 60%	4 - 15	0.4 - 6.2	Tajuelo et al., (2021)
	OH, smog chamber, high NO _x , RH < 10%	2	0.9 - 1.2	Tajuelo et al., (2021)
Furan	OH, PAM-OFR, RH = 34%	1667 - 2030	9.4 - 11.5	This study
	OH, smog chamber, high NO _x , 5 < RH < 88%, seed	1 - 119	0.04 - 5	Jiang et al., (2019b)
	OH, smog chamber, high NO _x , dry conditions	69 - 74	1.9 - 7.2	Gómez Alvarez et al., (2009)
	NO ₃ , smog chamber, RH = 2 - 16%	17 - 47	7	Jiang et al., (2019a)


3.4 SOA light absorption properties (b_{abs} , α , and MAC)

Figure 6a shows the wavelength-dependent MAC values (from 370 to 590 nm) for SOA derived from the 4 PAHs and 3 furans studied here with OH and NO₃ reactivity. A comparison with MAC values, from 300 to 550 nm, reported in the literature for laboratory-generated SOA from the oxidation of furan, monoaromatic and phenolic compounds (Lambe et al., 2013; Liu et al., 2016; Xie et al., 2017a; Jiang et al., 2019a) is also presented. MAC values alone together with b_{abs} are shown in Figure S4. Overall, the shape of the spectra is characteristic of atmospheric BrC materials, with higher absorption in the UV range (Laskin et al., 2015; Hems et al., 2021; Liu et al., 2016; Siemens et al., 2022; Moise et al., 2015). Our results showed significantly lower light absorption for furans SOA than PAHs SOA for both, day- and nighttime oxidation conditions. No light absorption was observed for furans SOA for wavelengths > 370 nm (Ångström exponents were then not calculated for furans SOA). At this wavelength of 370 nm, the highest MAC value was observed for Acy with OH radicals ($0.97 \pm 0.06 \text{ m}^2 \text{ g}^{-1}$). Other PAHs SOA generated with OH radicals showed comparable MAC values in the range of 0.36 - 0.44 $\text{m}^2 \text{ g}^{-1}$. Several studies have reported MAC values for Naph SOA, at 400 - 405 nm, in the range of 0.02 - 0.35 $\text{m}^2 \text{ g}^{-1}$ with low NO_x (Updyke et al., 2012; Lambe et al., 2013; Lee et al., 2014; Xie et al., 2017a; Siemens et al., 2022; Metcalf et al., 2013; Klodt et al., 2023; He et al., 2022) (Table S7). Results at 400 nm, with an average MAC value of 0.22 $\text{m}^2 \text{ g}^{-1}$, were in good agreement with the literature data obtained from filter extracts. The same applies to the absorption Ångström exponent (α) for which the value of 5.43 ± 0.15 obtained was comparable to the literature data (ranging from 5.2 to 8.9). These results show that the filter tape-based optical method (aethalometer) applied here can provide a good estimate of such aerosol optical properties. However, using AE33 data, the characterization of the UV region is limited to 370 nm while major light absorption for SOA has been mainly reported from 350 nm and below.

Over the PAH SOA formed with OH radicals, Naph showed the lowest α value (Figure S5). Acy SOA had the highest α of 8.35 ± 0.25 , while Phe and Flu showed comparable values (6.92 ± 0.48 and 6.08 ± 0.51 , respectively). With NO₃, the α values for Naph (5.57 ± 0.12) and Phe SOA (7.05 ± 0.88) were similar as with OH exposure while a significantly lower α value was obtained for Acy (6.78 ± 0.13), and comparable to Phe. Similarly, the MAC value (at 370 nm) of Acy SOA formed with NO₃ radical ($0.41 \pm 0.12 \text{ m}^2 \text{ g}^{-1}$) was largely lower than with OH radical and comparable to Phe ($0.43 \pm 0.10 \text{ m}^2 \text{ g}^{-1}$) and, in a lesser extent, to Naph ($0.32 \pm 0.03 \text{ m}^2 \text{ g}^{-1}$). The MAC values for furans SOA were significantly lower and about 0.010 $\text{m}^2 \text{ g}^{-1}$ for 2-MF, 2,5-DMF, and furan with OH radicals, and ranging from 0.005 to 0.021 $\text{m}^2 \text{ g}^{-1}$ for 2-MF and 2,5-DMF with NO₃ radicals. A similar range has been reported for furan SOA formed by oxidation with NO₃ radicals (0.08 - 0.11 $\text{m}^2 \text{ g}^{-1}$) (Jiang et al., 2019a; Chen et al., 2022b). While furans SOA exhibits a low light absorption in the UV region, PAH SOA absorption properties are significant. Only m-cresol and benzene SOA formed under high NO_x concentrations show higher MAC values in the UV-visible region (Figure 6a). MAC values determined here are comparable to those from ambient air biomass burning organic aerosols (BBOA) (Figure 6b) (Moise et al., 2015; Hoffer et al., 2006; Lack et al., 2012; Washenfelder et al., 2015; Zhang et al., 2016; Xie et al., 2017b; Cappa et al., 2019; Soleimanian et al., 2020; Mbengue et al., 2021). PAHs emissions from biomass burning (except naphthalene which is in the same range) are about 4 to 10 times lower than benzene, toluene,

phenol or furans and comparable to other monoaromatics and phenolic species (Bruns et al., 2016; Schauer et al., 2001; Bhattu et al., 2019; Hatch et al., 2015). Considering the SOA formation yields of these species and their contribution to the total SOA produced from biomass burning emissions (Bruns et al., 2016) (e.g., comparable for naphthalene and benzene, twice lower for phenol, and 4 times higher for m-cresol) and considering the light absorption properties of the different SOA (Figure 6), we can figure out that PAH SOA are a significant BrC component of the emissions following biomass burning.

Finally, MAC values obtained for PAH and furans SOA with NO₃ radicals were similar, or even lower for Acy SOA, to values observed with OH radicals. Only the NO₃ radical with 2,5-DMF gave higher MAC values. Usually, higher light absorption values are observed for anthropogenic SOA formed under a high NO_x regime due to the formation of stronger nitrogen-containing chromophores (nitroaromatic mainly) (Moise et al., 2015; Hems et al., 2021; He et al., 2022; Liu et al., 2016; Klodt et al., 2023; Laskin et al., 2015; Siemens et al., 2022; Xie et al., 2017a). The SOA formation with NO₃ radical alone does not induce necessarily a large formation of nitro-organic species. For PAHs, a significant formation of nitro-chromophores species is usually observed upon heterogeneous oxidation processes with NO₃ radical (gas/particle) but not for homogenous reactions in the gaseous phase (Cheng et al., 2020; Kwamena and Abbatt, 2008; Lu et al., 2011; Keyte et al., 2013). Once the aerosols formed in the PAM-OFR, heterogenous reactions may occur but reactions involving NO₃ interaction with substituent groups through H-atom abstraction, for example for Ace, Acy, and Flu, are not expected to induce an addition of NO₂ on the by-products formed (Keyte et al., 2013; Zhou and Wenger, 2013a, b). The reaction of Naph with NO₃ alone induces a formation of nitronaphthalenes and nitronaphthols (Keyte et al., 2013) that are mainly in the gaseous phase (Tomaz et al., 2016; Nalin et al., 2016; Reisen et al., 2003) (two-ring compounds) and are not associated with the SOA formed.



400

Figure 6. Comparison of the mass absorption cross section (MAC) from the day- and nighttime oxidation processes (with OH and NO_3 radicals respectively) with literature data for laboratory-generated SOA from anthropogenic precursors (A) and with data for ambient air biomass burning organic aerosols (BBOA) (B).

4 Conclusion

405 This study provides new key information on the properties of SOA formed from the oxidation of typical precursors emitted by biomass burning, namely PAHs and furans, under both day- and nighttime chemistry. Overall, PAH SOA formation yields were 5 - 6 times greater than those evaluated for furans (3 - 12%), illustrating the significant potential of PAHs in terms of SOA formation. Under our conditions, without NO_x , the SOA yields observed with NO_3 radicals were 3 times lower than with OH. The SOA formation yields obtained here from OFR experiments were either consistent with or higher than those reported
410 in previous studies, essentially from environmental (smog) chamber experiments. This might be the result of the different experimental conditions applied in terms of oxidant, SOA precursor, **PM (OA) mass or NO_x concentrations**, **cross reactions of RO_2** , and humidity. However, our results showed again that OFR is a relevant alternative and complementary tool to smog chambers to simulate atmospheric aging (Bruns et al., 2015; Peng and Jimenez, 2020). The SOA effective density showed an increasing trend with particle size for both reactivities before reaching a “plateau”, highlighting potential differences in terms
415 of chemical composition and morphology with the particle size, and thus a need to investigate them in future studies. PAHs SOA showed a higher light absorption in the range of 370 to 590 nm by comparison to furans that do not show significant absorption at higher wavelengths than 370 nm. A significant contribution of PAH SOA to ambient air BrC linked to biomass burning emissions is then expected. Finally, no increase in the MAC values was observed from OH to NO_3 oxidation processes, probably due to a low formation **of nitrogen-containing chromophores with NO_3 only** (without NO_x). As it is well known that
420 NO_x has a significant impact on the SOA formation (in terms of yields for instance) and their optical properties, with the subsequent formation of strong absorbing chromophores, experiments performed under high NO_x conditions, with both OH and NO_3 radicals, will be the focus of future works.

Competing interests

The authors declare that they have no conflict of interest.

425 Author contribution

AA designed and led the research. AEM and AA performed the experiments. AEM and AA analyzed the data. AEM, AA, BD, ZP, OF, JEP, LD and ATL contributed to the interpretation of the data and results. AA was responsible for funding acquisition. AA, SAA and BD supervised AEM PhD work. AEM and AA wrote the manuscript with inputs from all co-authors.

Acknowledgments

- 430 This work has been supported by the French Ministry of the Environment. The authors gratefully acknowledge Serguei Stavrovski, Ahmad El Masri, Robin Aujay, Nicolas Karoski, Tanguy Amodeo and Laurent Meunier (Ineris) for their help on the different instrumentations and for sample preparation and PAH analyses.

References

- 435 Ahern, A. T., Robinson, E. S., Tkacik, D. S., Saleh, R., Hatch, L. E., Barsanti, K. C., Stockwell, C. E., Yokelson, R. J., Presto, A. A., Robinson, A. L., Sullivan, R. C., and Donahue, N. M.: Production of Secondary Organic Aerosol During Aging of Biomass Burning Smoke From Fresh Fuels and Its Relationship to VOC Precursors, *Journal of Geophysical Research: Atmospheres*, 124, 3583–3606, <https://doi.org/10.1029/2018JD029068>, 2019.
- 440 Akherati, A., He, Y., Coggon, M. M., Koss, A. R., Hodshire, A. L., Sekimoto, K., Warneke, C., de Gouw, J., Yee, L., Seinfeld, J. H., Onasch, T. B., Herndon, S. C., Knighton, W. B., Cappa, C. D., Kleeman, M. J., Lim, C. Y., Kroll, J. H., Pierce, J. R., and Jathar, S. H.: Oxygenated Aromatic Compounds are Important Precursors of Secondary Organic Aerosol in Biomass-Burning Emissions, *Environ. Sci. Technol.*, 54, 8568–8579, <https://doi.org/10.1021/acs.est.0c01345>, 2020.
- Al Ali, F., Coeur, C., Houzel, N., Bouya, H., Tomas, A., and Romanias, M. N.: Rate Coefficients for the Gas-Phase Reactions of Nitrate Radicals with a Series of Furan Compounds, *J. Phys. Chem. A*, 126, 8674–8681, <https://doi.org/10.1021/acs.jpca.2c03828>, 2022.
- 445 Andreae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmospheric Chemistry and Physics*, 6, 3131–3148, <https://doi.org/10.5194/acp-6-3131-2006>, 2006.
- Asaf, D., Tas, E., Pedersen, D., Peleg, M., and Luria, M.: Long-Term Measurements of NO₃ Radical at a Semiarid Urban Site: 2. Seasonal Trends and Loss Mechanisms, *Environ. Sci. Technol.*, 44, 5901–5907, <https://doi.org/10.1021/es100967z>, 2010.
- 450 Aschmann, S. M., Nishino, N., Arey, J., and Atkinson, R.: Kinetics of the Reactions of OH Radicals with 2- and 3-Methylfuran, 2,3- and 2,5-Dimethylfuran, and *E*- and *Z*-3-Hexene-2,5-dione, and Products of OH + 2,5-Dimethylfuran, *Environ. Sci. Technol.*, 45, 1859–1865, <https://doi.org/10.1021/es103207k>, 2011.
- Aschmann, S. M., Nishino, N., Arey, J., and Atkinson, R.: Products of the OH Radical-Initiated Reactions of Furan, 2- and 3-Methylfuran, and 2,3- and 2,5-Dimethylfuran in the Presence of NO, *J. Phys. Chem. A*, 118, 457–466, <https://doi.org/10.1021/jp410345k>, 2014.
- 455 Baudic, A., Gros, V., Sauvage, S., Locoge, N., Sanchez, O., Sarda-Estève, R., Kalogridis, C., Petit, J.-E., Bonnaire, N., Baisnée, D., Favez, O., Albinet, A., Sciare, J., and Bonsang, B.: Seasonal variability and source apportionment of volatile organic compounds (VOCs) in the Paris megacity (France), *Atmospheric Chemistry and Physics*, 16, 11961–11989, <https://doi.org/10.5194/acp-16-11961-2016>, 2016.
- 460 Bhattu, D., Zotter, P., Zhou, J., Stefenelli, G., Klein, F., Bertrand, A., Temime-Roussel, B., Marchand, N., Slowik, J. G., Baltensperger, U., Prévôt, A. S. H., Nussbaumer, T., El Haddad, I., and Dommen, J.: Effect of Stove Technology and Combustion Conditions on Gas and Particulate Emissions from Residential Biomass Combustion, *Environ. Sci. Technol.*, 53, 2209–2219, <https://doi.org/10.1021/acs.est.8b05020>, 2019.

- 465 Bierbach, A., Barnes, I., and Becker, K. H.: Product and kinetic study of the OH-initiated gas-phase oxidation of Furan, 2-methylfuran and furanaldehydes at ≈ 300 K, *Atmospheric Environment*, 29, 2651–2660, [https://doi.org/10.1016/1352-2310\(95\)00096-H](https://doi.org/10.1016/1352-2310(95)00096-H), 1995.
- 470 Bressi, M., Cavalli, F., Putaud, J. P., Fröhlich, R., Petit, J.-E., Aas, W., Äijälä, M., Alastuey, A., Allan, J. D., Aurela, M., Berico, M., Bougiatioti, A., Bukowiecki, N., Canonaco, F., Crenn, V., Dusanter, S., Ehn, M., Elsasser, M., Flentje, H., Graf, P., Green, D. C., Heikkinen, L., Hermann, H., Holzinger, R., Hueglin, C., Keernik, H., Kiendler-Scharr, A., Kubelová, L., Lunder, C., Maasikmets, M., Makeš, O., Malaguti, A., Mihalopoulos, N., Nicolas, J. B., O'Dowd, C., Ovadnevaite, J., Petralia, E., Poulain, L., Priestman, M., Riffault, V., Ripoll, A., Schlag, P., Schwarz, J., Sciare, J., Slowik, J., Sosedova, Y., Stavroulas, I., Teinmaa, E., Via, M., Vodička, P., Williams, P. I., Wiedensohler, A., Young, D. E., Zhang, S., Favez, O., Minguillón, M. C., and Prevot, A. S. H.: A European aerosol phenomenology - 7: High-time resolution chemical characteristics of submicron particulate matter across Europe, *Atmospheric Environment: X*, 10, 100108, <https://doi.org/10.1016/j.aeaoa.2021.100108>, 2021.
- 475 Brook, R. D., Rajagopalan, S., Pope, C. A., Brook, J. R., Bhatnagar, A., Diez-Roux, A. V., Holguin, F., Hong, Y., Luepker, R. V., Mittleman, M. A., Peters, A., Siscovick, D., Smith, S. C., Whitsel, L., and Kaufman, J. D.: Particulate Matter Air Pollution and Cardiovascular Disease, *Circulation*, 121, 2331–2378, <https://doi.org/10.1161/CIR.0b013e3181d8ce1>, 2010.
- Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, *Chem. Soc. Rev.*, 41, 6405–6447, <https://doi.org/10.1039/C2CS35181A>, 2012.
- 480 Bruns, E. A., El Haddad, I., Keller, A., Klein, F., Kumar, N. K., Pieber, S. M., Corbin, J. C., Slowik, J. G., Brune, W. H., Baltensperger, U., and Prévôt, A. S. H.: Inter-comparison of laboratory smog chamber and flow reactor systems on organic aerosol yield and composition, *Atmos. Meas. Tech.*, 8, 2315–2332, <https://doi.org/10.5194/amt-8-2315-2015>, 2015.
- 485 Bruns, E. A., El Haddad, I., Slowik, J. G., Kilic, D., Klein, F., Baltensperger, U., and Prévôt, A. S. H.: Identification of significant precursor gases of secondary organic aerosols from residential wood combustion, *Scientific Reports*, 6, 27881, <https://doi.org/10.1038/srep27881>, 2016.
- 490 Cappa, C. D., Zhang, X., Russell, L. M., Collier, S., Lee, A. K. Y., Chen, C.-L., Betha, R., Chen, S., Liu, J., Price, D. J., Sanchez, K. J., McMeeking, G. R., Williams, L. R., Onasch, T. B., Worsnop, D. R., Abbatt, J., and Zhang, Q.: Light Absorption by Ambient Black and Brown Carbon and its Dependence on Black Carbon Coating State for Two California, USA, Cities in Winter and Summer, *Journal of Geophysical Research: Atmospheres*, 124, 1550–1577, <https://doi.org/10.1029/2018JD029501>, 2019.
- Carlton, A. G., Wiedinmyer, C., and Kroll, J. H.: A review of Secondary Organic Aerosol (SOA) formation from isoprene, *Atmospheric Chemistry and Physics*, 9, 4987–5005, <https://doi.org/10.5194/acp-9-4987-2009>, 2009.
- 495 Chan, A. W. H., Kautzman, K. E., Chhabra, P. S., Surratt, J. D., Chan, M. N., Crouse, J. D., Kürten, A., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from photooxidation of naphthalene and alkylnaphthalenes: implications for oxidation of intermediate volatility organic compounds (IVOCs), *Atmospheric Chemistry and Physics*, 9, 3049–3060, <https://doi.org/10.5194/acp-9-3049-2009>, 2009.
- Chen, C.-L., Kacarab, M., Tang, P., and Cocker, D. R.: SOA formation from naphthalene, 1-methylnaphthalene, and 2-methylnaphthalene photooxidation, *Atmospheric Environment*, 131, 424–433, <https://doi.org/10.1016/j.atmosenv.2016.02.007>, 2016.
- 500 Chen, C.-L., Li, L., Tang, P., and Cocker, D. R.: SOA formation from photooxidation of naphthalene and methylnaphthalenes with m-xylene and surrogate mixtures, *Atmospheric Environment*, 180, 256–264, <https://doi.org/10.1016/j.atmosenv.2018.02.051>, 2018.

- Chen, G., Canonaco, F., Tobler, A., Aas, W., Alastuey, A., Allan, J., Atabakhsh, S., Aurela, M., Baltensperger, U., Bougiatioti, A., De Brito, J. F., Ceburnis, D., Chazeau, B., Chebaicheb, H., Daellenbach, K. R., Ehn, M., El Haddad, I., Eleftheriadis, K., Favez, O., Flentje, H., Font, A., Fossum, K., Freney, E., Gini, M., Green, D. C., Heikkinen, L., Herrmann, H., Kalogridis, A.-C., Keernik, H., Lhotka, R., Lin, C., Lunder, C., Maasikmets, M., Manousakas, M. I., Marchand, N., Marin, C., Marmureanu, L., Mihalopoulos, N., Močnik, G., Nećki, J., O'Dowd, C., Ovadnevaite, J., Peter, T., Petit, J.-E., Pikridas, M., Matthew Platt, S., Pokorná, P., Poulain, L., Priestman, M., Riffault, V., Rinaldi, M., Róžański, K., Schwarz, J., Sciare, J., Simon, L., Skiba, A., Slowik, J. G., Sosedova, Y., Stavroulas, I., Styszko, K., Teinmaa, E., Timonen, H., Tremper, A., Vasilescu, J., Via, M., Vodička, P., Wiedensohler, A., Zografou, O., Cruz Minguillón, M., and Prévôt, A. S. H.: European aerosol phenomenology – 8: Harmonised source apportionment of organic aerosol using 22 Year-long ACSM/AMS datasets, *Environment International*, 166, 107325, <https://doi.org/10.1016/j.envint.2022.107325>, 2022a.
- Chen, J., Li, C., Ristovski, Z., Milic, A., Gu, Y., Islam, M. S., Wang, S., Hao, J., Zhang, H., He, C., Guo, H., Fu, H., Miljevic, B., Morawska, L., Thai, P., Lam, Y. F., Pereira, G., Ding, A., Huang, X., and Dumka, U. C.: A review of biomass burning: Emissions and impacts on air quality, health and climate in China, *Science of The Total Environment*, 579, 1000–1034, <https://doi.org/10.1016/j.scitotenv.2016.11.025>, 2017.
- Chen, K., Mayorga, R., Raeofy, N., Lum, M., Woods, M., Bahreini, R., Zhang, H., and Lin, Y.-H.: Effects of Nitrate Radical Levels and Pre-Existing Particles on Secondary Brown Carbon Formation from Nighttime Oxidation of Furan, *ACS Earth and Space Chemistry*, <https://doi.org/10.1021/acsearthspacechem.2c00244>, 2022b.
- Cheng, Z., Atwi, K. M., Yu, Z., Avery, A., Fortner, E. C., Williams, L., Majluf, F., Krechmer, J. E., Lambe, A. T., and Saleh, R.: Evolution of the light-absorption properties of combustion brown carbon aerosols following reaction with nitrate radicals, *Aerosol Science and Technology*, 54, 849–863, <https://doi.org/10.1080/02786826.2020.1726867>, 2020.
- Crenn, V., Sciare, J., Croteau, P. L., Verlhac, S., Fröhlich, R., Belis, C. A., Aas, W., Äijälä, M., Alastuey, A., Artiñano, B., Baisnée, D., Bonnaire, N., Bressi, M., Canagaratna, M., Canonaco, F., Carbone, C., Cavalli, F., Coz, E., Cubison, M. J., Esser-Gietl, J. K., Green, D. C., Gros, V., Heikkinen, L., Herrmann, H., Lunder, C., Minguillón, M. C., Močnik, G., O'Dowd, C. D., Ovadnevaite, J., Petit, J.-E., Petralia, E., Poulain, L., Priestman, M., Riffault, V., Ripoll, A., Sarda-Estève, R., Slowik, J. G., Setyan, A., Wiedensohler, A., Baltensperger, U., Prévôt, A. S. H., Jayne, J. T., and Favez, O.: ACTRIS ACSM intercomparison – Part 1: Reproducibility of concentration and fragment results from 13 individual Quadrupole Aerosol Chemical Speciation Monitors (Q-ACSM) and consistency with co-located instruments, *Atmospheric Measurement Techniques*, 8, 5063–5087, <https://doi.org/10.5194/amt-8-5063-2015>, 2015.
- Crippa, M., DeCarlo, P. F., Slowik, J. G., Mohr, C., Heringa, M. F., Chirico, R., Poulain, L., Freutel, F., Sciare, J., Cozic, J., Di Marco, C. F., Elsasser, M., Nicolas, J. B., Marchand, N., Abidi, E., Wiedensohler, A., Drewnick, F., Schneider, J., Borrmann, S., Nemitz, E., Zimmermann, R., Jaffrezo, J.-L., Prévôt, A. S. H., and Baltensperger, U.: Wintertime aerosol chemical composition and source apportionment of the organic fraction in the metropolitan area of Paris, *Atmospheric Chemistry and Physics*, 13, 961–981, <https://doi.org/10.5194/acp-13-961-2013>, 2013.
- Davis, D. D., Ravishankara, A. R., and Fischer, S.: SO₂ oxidation via the hydroxyl radical: Atmospheric fate of HSO_x radicals, *Geophysical Research Letters*, 6, 113–116, <https://doi.org/10.1029/GL006i002p00113>, 1979.
- DeCarlo, P. F., Slowik, J. G., Worsnop, D. R., Davidovits, P., and Jimenez, J. L.: Particle Morphology and Density Characterization by Combined Mobility and Aerodynamic Diameter Measurements. Part 1: Theory, *Aerosol Science and Technology*, 38, 1185–1205, <https://doi.org/10.1080/027868290903907>, 2004.
- Denier van der Gon, H. a. C., Bergström, R., Fountoukis, C., Johansson, C., Pandis, S. N., Simpson, D., and Visschedijk, A. J. H.: Particulate emissions from residential wood combustion in Europe – revised estimates and an evaluation, *Atmospheric Chemistry and Physics*, 15, 6503–6519, <https://doi.org/10.5194/acp-15-6503-2015>, 2015.

- 545 Dingle, J. H., Zimmerman, S., Frie, A. L., Min, J., Jung, H., and Bahreini, R.: Complex refractive index, single scattering albedo, and mass absorption coefficient of secondary organic aerosols generated from oxidation of biogenic and anthropogenic precursors, *Aerosol Science and Technology*, 53, 449–463, <https://doi.org/10.1080/02786826.2019.1571680>, 2019.
- 550 Drinovec, L., Močnik, G., Zotter, P., Prévôt, A. S. H., Ruckstuhl, C., Coz, E., Rupakheti, M., Sciare, J., Müller, T., Wiedensohler, A., and Hansen, A. D. A.: The “dual-spot” Aethalometer: an improved measurement of aerosol black carbon with real-time loading compensation, *Atmospheric Measurement Techniques*, 8, 1965–1979, <https://doi.org/10.5194/amt-8-1965-2015>, 2015.
- Drinovec, L., Gregorič, A., Zotter, P., Wolf, R., Bruns, E. A., Prévôt, A. S. H., Petit, J.-E., Favez, O., Sciare, J., Arnold, I. J., Chakrabarty, R. K., Moosmüller, H., Filep, A., and Močnik, G.: The filter-loading effect by ambient aerosols in filter absorption photometers depends on the coating of the sampled particles, *Atmospheric Measurement Techniques*, 10, 1043–1059, <https://doi.org/10.5194/amt-10-1043-2017>, 2017.
- 555 Drinovec, L., Jagodič, U., Pirker, L., Škarabot, M., Kurtjak, M., Vidović, K., Ferrero, L., Visser, B., Röhrbein, J., Weingartner, E., Kalbermatter, D. M., Vasilatou, K., Bühlmann, T., Pascale, C., Müller, T., Wiedensohler, A., and Močnik, G.: A dual-wavelength photothermal aerosol absorption monitor: design, calibration and performance, *Atmospheric Measurement Techniques*, 15, 3805–3825, <https://doi.org/10.5194/amt-15-3805-2022>, 2022.
- 560 El Mais, A. E. R.: Primary and secondary residential wood combustion emissions: physicochemical and bio-analytical characterization and effect-directed analysis (EDA), PhD Thesis, Université Aix Marseille, 2023.
- Finlayson-Pitts, B. J. and Pitts Jr, J. N.: *Chemistry of the upper and lower atmosphere*, Academic Press, 969 pp., 2000.
- Fountoukis, C., Megaritis, A. G., Skyllakou, K., Charalampidis, P. E., Denier van der Gon, H. A. C., Crippa, M., Prévôt, A. S. H., Fachinger, F., Wiedensohler, A., Pilinis, C., and Pandis, S. N.: Simulating the formation of carbonaceous aerosol in a European Megacity (Paris) during the MEGAPOLI summer and winter campaigns, *Atmospheric Chemistry and Physics*, 16, 3727–3741, <https://doi.org/10.5194/acp-16-3727-2016>, 2016.
- 570 Freney, E., Zhang, Y., Croteau, P., Amodeo, T., Williams, L., Truong, F., Petit, J.-E., Sciare, J., Sarda-Estève, R., Bonnaire, N., Arumae, T., Aurela, M., Bougiatioti, A., Mihalopoulos, N., Coz, E., Artinano, B., Crenn, V., Elste, T., Heikkinen, L., Poulain, L., Wiedensohler, A., Herrmann, H., Priestman, M., Alastuey, A., Stavroulas, I., Tobler, A., Vasilescu, J., Zanca, N., Canagaratna, M., Carbone, C., Flentje, H., Green, D., Maasikmets, M., Marmureanu, L., Minguillon, M. C., Prevot, A. S. H., Gros, V., Jayne, J., and Favez, O.: The second ACTRIS inter-comparison (2016) for Aerosol Chemical Speciation Monitors (ACSM): Calibration protocols and instrument performance evaluations, *Aerosol Science and Technology*, 53, 830–842, <https://doi.org/10.1080/02786826.2019.1608901>, 2019.
- 575 Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., and Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs, *Atmospheric Chemistry and Physics*, 15, 8217–8299, <https://doi.org/10.5194/acp-15-8217-2015>, 2015.
- Gómez Alvarez, E., Borrás, E., Viidanoja, J., and Hjorth, J.: Unsaturated dicarbonyl products from the OH-initiated photo-oxidation of furan, 2-methylfuran and 3-methylfuran, *Atmospheric Environment*, 43, 1603–1612, <https://doi.org/10.1016/j.atmosenv.2008.12.019>, 2009.
- 580 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski,

- R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmospheric Chemistry and Physics*, 9, 5155–5236, <https://doi.org/10.5194/acp-9-5155-2009>, 2009.
- 585 Hartikainen, A., Yli-Pirilä, P., Tiitta, P., Leskinen, A., Kortelainen, M., Orasche, J., Schnelle-Kreis, J., Lehtinen, K. E. J., Zimmermann, R., Jokiniemi, J., and Sippula, O.: Volatile Organic Compounds from Logwood Combustion: Emissions and Transformation under Dark and Photochemical Aging Conditions in a Smog Chamber, *Environ. Sci. Technol.*, 52, 4979–4988, <https://doi.org/10.1021/acs.est.7b06269>, 2018.
- 590 Hatch, L. E., Luo, W., Pankow, J. F., Yokelson, R. J., Stockwell, C. E., and Barsanti, K. C.: Identification and quantification of gaseous organic compounds emitted from biomass burning using two-dimensional gas chromatography–time-of-flight mass spectrometry, *Atmospheric Chemistry and Physics*, 15, 1865–1899, <https://doi.org/10.5194/acp-15-1865-2015>, 2015.
- 595 Hatch, L. E., Yokelson, R. J., Stockwell, C. E., Veres, P. R., Simpson, I. J., Blake, D. R., Orlando, J. J., and Barsanti, K. C.: Multi-instrument comparison and compilation of non-methane organic gas emissions from biomass burning and implications for smoke-derived secondary organic aerosol precursors, *Atmospheric Chemistry and Physics*, 17, 1471–1489, <https://doi.org/10.5194/acp-17-1471-2017>, 2017.
- Hatch, L. E., Rivas-Ubach, A., Jen, C. N., Lipton, M., Goldstein, A. H., and Barsanti, K. C.: Measurements of I/SVOCs in biomass-burning smoke using solid-phase extraction disks and two-dimensional gas chromatography, *Atmospheric Chemistry and Physics*, 18, 17801–17817, <https://doi.org/10.5194/acp-18-17801-2018>, 2018.
- 600 He, Q., Li, C., Siemens, K., Morales, A. C., Hettiyadura, A. P. S., Laskin, A., and Rudich, Y.: Optical Properties of Secondary Organic Aerosol Produced by Photooxidation of Naphthalene under NO_x Condition, *Environ. Sci. Technol.*, 56, 4816–4827, <https://doi.org/10.1021/acs.est.1c07328>, 2022.
- Heald, C. L. and Kroll, J. H.: The fuel of atmospheric chemistry: Toward a complete description of reactive organic carbon, *Sci. Adv.*, 6, eaay8967, <https://doi.org/10.1126/sciadv.aay8967>, 2020.
- 605 Hems, R. F., Schnitzler, E. G., Liu-Kang, C., Cappa, C. D., and Abbatt, J. P. D.: Aging of Atmospheric Brown Carbon Aerosol, *ACS Earth Space Chem.*, 5, 722–748, <https://doi.org/10.1021/acsearthspacechem.0c00346>, 2021.
- Hodshire, A. L., Akherati, A., Alvarado, M. J., Brown-Steiner, B., Jathar, S. H., Jimenez, J. L., Kreidenweis, S. M., Lonsdale, C. R., Onasch, T. B., Ortega, A. M., and Pierce, J. R.: Aging Effects on Biomass Burning Aerosol Mass and Composition: A Critical Review of Field and Laboratory Studies, *Environ. Sci. Technol.*, 53, 10007–10022, <https://doi.org/10.1021/acs.est.9b02588>, 2019.
- 610 Hoffer, A., Gelencsér, A., Guyon, P., Kiss, G., Schmid, O., Frank, G. P., Artaxo, P., and Andreae, M. O.: Optical properties of humic-like substances (HULIS) in biomass-burning aerosols, *Atmospheric Chemistry and Physics*, 6, 3563–3570, <https://doi.org/10.5194/acp-6-3563-2006>, 2006.
- 615 IPCC: Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change., edited by: Pörtner, H.-O., Roberts, D. C., Tignor, M. M. B., Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A., and Rama, B., Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009325844>., 2022.
- 620 Jiang, H., Frie, A. L., Lavi, A., Chen, J. Y., Zhang, H., Bahreini, R., and Lin, Y.-H.: Brown Carbon Formation from Nighttime Chemistry of Unsaturated Heterocyclic Volatile Organic Compounds, *Environ. Sci. Technol. Lett.*, 6, 184–190, <https://doi.org/10.1021/acs.estlett.9b00017>, 2019a.

- Jiang, J., Carter, W. P. L., Cocker, D. R. I., and Barsanti, K. C.: Development and Evaluation of a Detailed Mechanism for Gas-Phase Atmospheric Reactions of Furans, *ACS Earth Space Chem.*, 4, 1254–1268, <https://doi.org/10.1021/acsearthspacechem.0c00058>, 2020.
- 625 Jiang, X., Tsona, N. T., Jia, L., Liu, S., Zhang, H., Xu, Y., and Du, L.: Secondary organic aerosol formation from photooxidation of furan: effects of NO_x and humidity, *Atmospheric Chemistry and Physics*, 19, 13591–13609, <https://doi.org/10.5194/acp-19-13591-2019>, 2019b.
- 630 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., E., Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529, <https://doi.org/10.1126/science.1180353>, 2009.
- 635 Johnson, T. J., Symonds, J. P. R., and Olfert, J. S.: Mass–Mobility Measurements Using a Centrifugal Particle Mass Analyzer and Differential Mobility Spectrometer, *Aerosol Science and Technology*, 47, 1215–1225, <https://doi.org/10.1080/02786826.2013.830692>, 2013.
- 640 Joo, T., Rivera-Rios, J. C., Takeuchi, M., Alvarado, M. J., and Ng, N. L.: Secondary Organic Aerosol Formation from Reaction of 3-Methylfuran with Nitrate Radicals, *ACS Earth Space Chem.*, 3, 922–934, <https://doi.org/10.1021/acsearthspacechem.9b00068>, 2019a.
- Joo, T., Rivera-Rios, J. C., Takeuchi, M., Alvarado, M. J., and Ng, N. L.: Secondary Organic Aerosol Formation from Reaction of 3-Methylfuran with Nitrate Radicals, *ACS Earth Space Chem.*, 3, 922–934, <https://doi.org/10.1021/acsearthspacechem.9b00068>, 2019b.
- 645 Jorga, S. D., Florou, K., Kaltsonoudis, C., Kodros, J. K., Vasilakopoulou, C., Cirtog, M., Fouqueau, A., Picquet-Varrault, B., Nenes, A., and Pandis, S. N.: Nighttime chemistry of biomass burning emissions in urban areas: A dual mobile chamber study, *Atmospheric Chemistry and Physics*, 21, 15337–15349, <https://doi.org/10.5194/acp-21-15337-2021>, 2021.
- Kang, E., Root, M. J., Toohey, D. W., and Brune, W. H.: Introducing the concept of Potential Aerosol Mass (PAM), *Atmos. Chem. Phys.*, 18, 2007.
- 650 Kautzman, K. E., Surratt, J. D., Chan, M. N., Chan, A. W. H., Hersey, S. P., Chhabra, P. S., Dalleska, N. F., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Chemical Composition of Gas- and Aerosol-Phase Products from the Photooxidation of Naphthalene, *J. Phys. Chem. A*, 114, 913–934, <https://doi.org/10.1021/jp908530s>, 2010.
- Keyte, I. J., Harrison, R. M., and Lammel, G.: Chemical reactivity and long-range transport potential of polycyclic aromatic hydrocarbons – a review, *Chem. Soc. Rev.*, 42, 9333, <https://doi.org/10.1039/c3cs60147a>, 2013.
- 655 Kind, I., Berndt, T., Böge, O., and Rolle, W.: Gas-phase rate constants for the reaction of NO₃ radicals with furan and methyl-substituted furans, *Chemical physics letters*, 256, 679–683, 1996.
- Kleindienst, T. E., Jaoui, M., Lewandowski, M., Offenberg, J. H., and Docherty, K. S.: The formation of SOA and chemical tracer compounds from the photooxidation of naphthalene and its methyl analogs in the presence and absence of nitrogen oxides, *Atmospheric Chemistry and Physics*, 12, 8711–8726, <https://doi.org/10.5194/acp-12-8711-2012>, 2012.

- 660 Klodt, A., Aiona, P., MacMillan, A., Lee, H. J. (Julie), Zhang, X., Helgestad, T., Novak, G., Lin, P., Laskin, J., Laskin, A., Bertram, T., Cappa, C., and Nizkorodov, S. A.: Effect of Relative Humidity, NO_x, and Ammonia on Physical Properties of Naphthalene Secondary Organic Aerosol, *Environ. Sci.: Atmos.*, <https://doi.org/10.1039/D3EA00033H>, 2023.
- Kodros, J. K., Papanastasiou, D. K., Paglione, M., Masiol, M., Squizzato, S., Florou, K., Skyllakou, K., Kaltsonoudis, C., Nenes, A., and Pandis, S. N.: Rapid dark aging of biomass burning as an overlooked source of oxidized organic aerosol, *Proceedings of the National Academy of Sciences of the United States of America*, 117, 33028–33039, 2020.
- 665 Kodros, J. K., Kaltsonoudis, C., Paglione, M., Florou, K., Jorga, S., Vasilakopoulou, C., Cirtog, M., Cazaunau, M., Picquet-Varrault, B., Nenes, A., and Pandis, S. N.: Secondary aerosol formation during the dark oxidation of residential biomass burning emissions, *Environ. Sci.: Atmos.*, 2, 1221–1236, <https://doi.org/10.1039/D2EA00031H>, 2022.
- Kostenidou, E., Pathak, R. K., and Pandis, S. N.: An Algorithm for the Calculation of Secondary Organic Aerosol Density Combining AMS and SMPS Data, *Aerosol Science and Technology*, 41, 1002–1010, <https://doi.org/10.1080/02786820701666270>, 2007.
- 670 Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmospheric Environment*, 42, 3593–3624, <https://doi.org/10.1016/j.atmosenv.2008.01.003>, 2008.
- Kwamena, N.-O. A. and Abbatt, J. P. D.: Heterogeneous nitration reactions of polycyclic aromatic hydrocarbons and n-hexane soot by exposure to NO₃/NO₂/N₂O₅, *Atmospheric Environment*, 42, 8309–8314, <https://doi.org/10.1016/j.atmosenv.2008.07.037>, 2008.
- 675 Lack, D. A., Langridge, J. M., Bahreini, R., Cappa, C. D., Middlebrook, A. M., and Schwarz, J. P.: Brown carbon and internal mixing in biomass burning particles, *Proceedings of the National Academy of Sciences*, 109, 14802–14807, <https://doi.org/10.1073/pnas.1206575109>, 2012.
- 680 Lambe, A. T., Ahern, A. T., Williams, L. R., Slowik, J. G., Wong, J. P. S., Abbatt, J. P. D., Brune, W. H., Ng, N. L., Wright, J. P., Croasdale, D. R., Worsnop, D. R., Davidovits, P., and Onasch, T. B.: Characterization of aerosol photooxidation flow reactors: heterogeneous oxidation, secondary organic aerosol formation and cloud condensation nuclei activity measurements, *Atmospheric Measurement Techniques*, 4, 445–461, <https://doi.org/10.5194/amt-4-445-2011>, 2011.
- 685 Lambe, A. T., Cappa, C. D., Massoli, P., Onasch, T. B., Forestieri, S. D., Martin, A. T., Cummings, M. J., Croasdale, D. R., Brune, W. H., Worsnop, D. R., and Davidovits, P.: Relationship between Oxidation Level and Optical Properties of Secondary Organic Aerosol, *Environ. Sci. Technol.*, 47, 6349–6357, <https://doi.org/10.1021/es401043j>, 2013.
- Lambe, A. T., Chhabra, P. S., Onasch, T. B., Brune, W. H., Hunter, J. F., Kroll, J. H., Cummings, M. J., Brogan, J. F., Parmar, Y., Worsnop, D. R., Kolb, C. E., and Davidovits, P.: Effect of oxidant concentration, exposure time, and seed particles on secondary organic aerosol chemical composition and yield, *Atmos. Chem. Phys.*, 15, 3063–3075, <https://doi.org/10.5194/acp-15-3063-2015>, 2015.
- 690 Lambe, A. T., Wood, E. C., Krechmer, J. E., Majluf, F., Williams, L. R., Croteau, P. L., Cirtog, M., Féron, A., Petit, J.-E., Albinet, A., Jimenez, J. L., and Peng, Z.: Nitrate radical generation via continuous generation of dinitrogen pentoxide in a laminar flow reactor coupled to an oxidation flow reactor, *Atmospheric Measurement Techniques*, 13, 2397–2411, <https://doi.org/10.5194/amt-13-2397-2020>, 2020.
- 695 Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chem. Rev.*, 115, 4335–4382, <https://doi.org/10.1021/cr5006167>, 2015.

- Lee, H. J. (Julie), Aiona, P. K., Laskin, A., Laskin, J., and Nizkorodov, S. A.: Effect of Solar Radiation on the Optical Properties and Molecular Composition of Laboratory Proxies of Atmospheric Brown Carbon, *Environ. Sci. Technol.*, 48, 10217–10226, <https://doi.org/10.1021/es502515r>, 2014.
- 700 Lee, J. and Lane, D. A.: Formation of oxidized products from the reaction of gaseous phenanthrene with the OH radical in a reaction chamber, *Atmospheric Environment*, 44, 2469–2477, <https://doi.org/10.1016/j.atmosenv.2010.03.008>, 2010.
- Lee, J. Y. and Lane, D. A.: Unique products from the reaction of naphthalene with the hydroxyl radical, *Atmospheric Environment*, 43, 4886–4893, <https://doi.org/10.1016/j.atmosenv.2009.07.018>, 2009.
- 705 Lee, J. Y., Lane, D. A., Heo, J. B., Yi, S.-M., and Kim, Y. P.: Quantification and seasonal pattern of atmospheric reaction products of gas phase PAHs in PM_{2.5}, *Atmospheric Environment*, 55, 17–25, <https://doi.org/10.1016/j.atmosenv.2012.03.007>, 2012.
- Li, M., Liu, Y., and Wang, L.: Gas-phase ozonolysis of furans, methylfurans, and dimethylfurans in the atmosphere, *Phys. Chem. Chem. Phys.*, 20, 24735–24743, <https://doi.org/10.1039/C8CP04947E>, 2018.
- 710 Li, R., Palm, B. B., Ortega, A. M., Hlywiak, J., Hu, W., Peng, Z., Day, D. A., Knote, C., Brune, W. H., de Gouw, J. A., and Jimenez, J. L.: Modeling the Radical Chemistry in an Oxidation Flow Reactor: Radical Formation and Recycling, Sensitivities, and the OH Exposure Estimation Equation, *J. Phys. Chem. A*, 119, 4418–4432, <https://doi.org/10.1021/jp509534k>, 2015.
- Liu, J., Lin, P., Laskin, A., Laskin, J., Kathmann, S. M., Wise, M., Caylor, R., Imholt, F., Selimovic, V., and Shilling, J. E.: Optical properties and aging of light-absorbing secondary organic aerosol, *Atmos. Chem. Phys.*, 16, 12815–12827, <https://doi.org/10.5194/acp-16-12815-2016>, 2016.
- 715 Liu, Y. and Daum, P. H.: Relationship of refractive index to mass density and self-consistency of mixing rules for multicomponent mixtures like ambient aerosols, *Journal of Aerosol Science*, 39, 974–986, <https://doi.org/10.1016/j.jaerosci.2008.06.006>, 2008.
- Lu, J. W., Flores, J. M., Lavi, A., Abo-Riziq, A., and Rudich, Y.: Changes in the optical properties of benzo[a]pyrene-coated aerosols upon heterogeneous reactions with NO₂ and NO₃, *Phys. Chem. Chem. Phys.*, 13, 6484–6492, <https://doi.org/10.1039/C0CP02114H>, 2011.
- 720 Malloy, Q. G. J., Nakao, S., Qi, L., Austin, R., Stothers, C., Hagino, H., and Cocker, D. R.: Real-Time Aerosol Density Determination Utilizing a Modified Scanning Mobility Particle Sizer—Aerosol Particle Mass Analyzer System, *Aerosol Science and Technology*, 43, 673–678, <https://doi.org/10.1080/02786820902832960>, 2009.
- Mao, J., Ren, X., Brune, W. H., Olson, J. R., Crawford, J. H., Fried, A., Huey, L. G., Cohen, R. C., Heikes, B., Singh, H. B., Blake, D. R., Sachse, G. W., Diskin, G. S., Hall, S. R., and Shetter, R. E.: Airborne measurement of OH reactivity during INTEX-B, *Atmospheric Chemistry and Physics*, 9, 163–173, <https://doi.org/10.5194/acp-9-163-2009>, 2009.
- Mbengue, S., Zikova, N., Schwarz, J., Vodička, P., Šmejkalová, A. H., and Holoubek, I.: Mass absorption cross-section and absorption enhancement from long term black and elemental carbon measurements: A rural background station in Central Europe, *Science of The Total Environment*, 794, 148365, <https://doi.org/10.1016/j.scitotenv.2021.148365>, 2021.
- 730 McMurry, P. H., Wang, X., Park, K., and Ehara, K.: The Relationship between Mass and Mobility for Atmospheric Particles: A New Technique for Measuring Particle Density, *Aerosol Science and Technology*, 36, 227–238, <https://doi.org/10.1080/027868202753504083>, 2002.

- 735 Metcalf, A. R., Loza, C. L., Coggon, M. M., Craven, J. S., Jonsson, H. H., Flagan, R. C., and Seinfeld, J. H.: Secondary Organic Aerosol Coating Formation and Evaporation: Chamber Studies Using Black Carbon Seed Aerosol and the Single-Particle Soot Photometer, *Aerosol Science and Technology*, 47, 326–347, <https://doi.org/10.1080/02786826.2012.750712>, 2013.
- 740 Mircea, M., Bessagnet, B., D'Isidoro, M., Pirovano, G., Aksoyoglu, S., Ciarelli, G., Tsyro, S., Manders, A., Bieser, J., Stern, R., Vivanco, M. G., Cuvelier, C., Aas, W., Prévôt, A. S. H., Aulinger, A., Briganti, G., Calori, G., Cappelletti, A., Colette, A., Couvidat, F., Fagerli, H., Finardi, S., Kranenburg, R., Rouïl, L., Silibello, C., Spindler, G., Poulain, L., Herrmann, H., Jimenez, J. L., Day, D. A., Tiitta, P., and Carbone, S.: EURODELTA III exercise: An evaluation of air quality models' capacity to reproduce the carbonaceous aerosol, *Atmospheric Environment: X*, 2, 100018, <https://doi.org/10.1016/j.aeaoa.2019.100018>, 2019.
- Moise, T., Flores, J. M., and Rudich, Y.: Optical Properties of Secondary Organic Aerosols and Their Changes by Chemical Processes, *Chem. Rev.*, 115, 4400–4439, <https://doi.org/10.1021/cr5005259>, 2015.
- 745 Monks, P. S., Granier, C., Fuzzi, S., Stohl, A., Williams, M. L., Akimoto, H., Amann, M., Baklanov, A., Baltensperger, U., Bey, I., Blake, N., Blake, R. S., Carslaw, K., Cooper, O. R., Dentener, F., Fowler, D., Fragkou, E., Frost, G. J., Generoso, S., Ginoux, P., Grewe, V., Guenther, A., Hansson, H. C., Henne, S., Hjorth, J., Hofzumahaus, A., Huntrieser, H., Isaksen, I. S. A., Jenkin, M. E., Kaiser, J., Kanakidou, M., Klimont, Z., Kulmala, M., Laj, P., Lawrence, M. G., Lee, J. D., Liousse, C., Maione, M., McFiggans, G., Metzger, A., Mieville, A., Moussiopoulos, N., Orlando, J. J., O'Dowd, C. D., Palmer, P. I., Parrish, D. D., Petzold, A., Platt, U., Pöschl, U., Prévôt, A. S. H., Reeves, C. E., Reimann, S., Rudich, Y., Sellegri, K., 750 Steinbrecher, R., Simpson, D., ten Brink, H., Theloke, J., van der Werf, G. R., Vautard, R., Vestreng, V., Vlachokostas, Ch., and von Glasow, R.: Atmospheric composition change – global and regional air quality, *Atmospheric Environment*, 43, 5268–5350, <https://doi.org/10.1016/j.atmosenv.2009.08.021>, 2009.
- Nakao, S., Tang, P., Tang, X., Clark, C. H., Qi, L., Seo, E., Asa-Awuku, A., and Cocker, D.: Density and elemental ratios of secondary organic aerosol: Application of a density prediction method, *Atmospheric Environment*, 68, 273–277, 755 <https://doi.org/10.1016/j.atmosenv.2012.11.006>, 2013.
- Nalin, F., Golly, B., Besombes, J.-L., Pelletier, C., Aujay-Plouzeau, R., Verlhac, S., Dermigny, A., Fievet, A., Karoski, N., Dubois, P., Collet, S., Favez, O., and Albinet, A.: Fast oxidation processes from emission to ambient air introduction of aerosol emitted by residential log wood stoves, *Atmospheric Environment*, 143, 15–26, <https://doi.org/10.1016/j.atmosenv.2016.08.002>, 2016.
- 760 Newland, M. J., Ren, Y., McGillen, M. R., Michelat, L., Daële, V., and Mellouki, A.: NO₃ chemistry of wildfire emissions: a kinetic study of the gas-phase reactions of furans with the NO₃ radical, *Atmospheric Chemistry and Physics*, 22, 1761–1772, <https://doi.org/10.5194/acp-22-1761-2022>, 2022.
- 765 Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P. L., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., Sun, Y. L., and Jayne, J. T.: An Aerosol Chemical Speciation Monitor (ACSM) for Routine Monitoring of the Composition and Mass Concentrations of Ambient Aerosol, *Aerosol Science and Technology*, 45, 780–794, <https://doi.org/10.1080/02786826.2011.560211>, 2011.
- Olfert, J. S. and Collings, N.: New method for particle mass classification—the Couette centrifugal particle mass analyzer, *Journal of Aerosol Science*, 36, 1338–1352, <https://doi.org/10.1016/j.jaerosci.2005.03.006>, 2005.
- 770 Olfert, J. S., Reavell, K. StJ., Rushton, M. G., and Collings, N.: The experimental transfer function of the Couette centrifugal particle mass analyzer, *Journal of Aerosol Science*, 37, 1840–1852, <https://doi.org/10.1016/j.jaerosci.2006.07.007>, 2006.

- Peng, L., Li, Z., Zhang, G., Bi, X., Hu, W., Tang, M., Wang, X., Peng, P., and Sheng, G.: A review of measurement techniques for aerosol effective density, *Science of The Total Environment*, 146248, <https://doi.org/10.1016/j.scitotenv.2021.146248>, 2021.
- 775 Peng, Z. and Jimenez, J. L.: KinSim: A Research-Grade, User-Friendly, Visual Kinetics Simulator for Chemical-Kinetics and Environmental-Chemistry Teaching, *J. Chem. Educ.*, 96, 806–811, <https://doi.org/10.1021/acs.jchemed.9b00033>, 2019.
- Peng, Z. and Jimenez, J. L.: Radical chemistry in oxidation flow reactors for atmospheric chemistry research, *Chem. Soc. Rev.*, 49, 2570–2616, <https://doi.org/10.1039/C9CS00766K>, 2020.
- 780 Pope, C. A., Burnett, R. T., Thun, M. J., Calle, E. E., Krewski, D., Ito, K., and Thurston, G. D.: Lung Cancer, Cardiopulmonary Mortality, and Long-term Exposure to Fine Particulate Air Pollution, *JAMA*, 287, 1132–1141, <https://doi.org/10.1001/jama.287.9.1132>, 2002.
- Pope, C. A., Coleman, N., Pond, Z. A., and Burnett, R. T.: Fine particulate air pollution and human mortality: 25+ years of cohort studies, *Environmental Research*, 183, 108924, <https://doi.org/10.1016/j.envres.2019.108924>, 2020.
- Rajagopalan, S., Al -Kindi Sadeer G., and Brook, R. D.: Air Pollution and Cardiovascular Disease, *Journal of the American College of Cardiology*, 72, 2054–2070, <https://doi.org/10.1016/j.jacc.2018.07.099>, 2018.
- 785 Reisen, F., Wheeler, S., and Arey, J.: Methyl- and dimethyl-/ethyl-nitronaphthalenes measured in ambient air in Southern California, *Atmospheric Environment*, 37, 3653–3657, [https://doi.org/10.1016/S1352-2310\(03\)00469-2](https://doi.org/10.1016/S1352-2310(03)00469-2), 2003.
- Ringuet, J., Albinet, A., Leoz-Garziandia, E., Budzinski, H., and Villenave, E.: Reactivity of polycyclic aromatic compounds (PAHs, NPAHs and OPAHs) adsorbed on natural aerosol particles exposed to atmospheric oxidants, *Atmospheric Environment*, 61, 15–22, <https://doi.org/10.1016/j.atmosenv.2012.07.025>, 2012.
- 790 Riva, M., Healy, R. M., Flaud, P.-M., Perraudin, E., Wenger, J. C., and Villenave, E.: Gas- and particle-phase products from the photooxidation of acenaphthene and acenaphthylene by OH radicals, *Atmospheric Environment*, 151, 34–44, <https://doi.org/10.1016/j.atmosenv.2016.11.063>, 2017.
- Růžičková, J., Raclavská, H., Juchelková, D., Kucbel, M., and Slamová, K.: The origin of potential precursors of secondary organic aerosols during combustion of biochar and softwood in residential heating, *Process Safety and Environmental Protection*, 161, 147–161, <https://doi.org/10.1016/j.psep.2022.03.036>, 2022.
- 795 Saleh, R., Marks, M., Heo, J., Adams, P. J., Donahue, N. M., and Robinson, A. L.: Contribution of brown carbon and lensing to the direct radiative effect of carbonaceous aerosols from biomass and biofuel burning emissions, *Journal of Geophysical Research: Atmospheres*, 120, 10,285–10,296, <https://doi.org/10.1002/2015JD023697>, 2015.
- 800 Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of Emissions from Air Pollution Sources. 3. C1–C29 Organic Compounds from Fireplace Combustion of Wood, *Environ. Sci. Technol.*, 35, 1716–1728, <https://doi.org/10.1021/es001331e>, 2001.
- Schervish, M. and Donahue, N. M.: Peroxy radical kinetics and new particle formation, *Environ. Sci.: Atmos.*, 1, 79–92, <https://doi.org/10.1039/D0EA00017E>, 2021.
- 805 Seinfeld, J. H. and Pandis, S. N.: Atmospheric chemistry and physics: From air pollution to climate change, Wiley-Interscience, 1152 pp., 1998.

- Shakya, K. M. and Griffin, R. J.: Secondary Organic Aerosol from Photooxidation of Polycyclic Aromatic Hydrocarbons, *Environ. Sci. Technol.*, 44, 8134–8139, <https://doi.org/10.1021/es1019417>, 2010.
- Siemens, K., Morales, A., He, Q., Li, C., Hettiyadura, A. P. S., Rudich, Y., and Laskin, A.: Molecular Analysis of Secondary Brown Carbon Produced from the Photooxidation of Naphthalene, *Environ. Sci. Technol.*, 56, 3340–3353, <https://doi.org/10.1021/acs.est.1c03135>, 2022.
- 810 Soleimanian, E., Mousavi, A., Taghvaei, S., Shafer, M. M., and Sioutas, C.: Impact of secondary and primary particulate matter (PM) sources on the enhanced light absorption by brown carbon (BrC) particles in central Los Angeles, *Science of The Total Environment*, 705, 135902, <https://doi.org/10.1016/j.scitotenv.2019.135902>, 2020.
- Srivastava, D., Favez, O., Perraudin, E., Villenave, E., and Albinet, A.: Comparison of Measurement-Based Methodologies to Apportion Secondary Organic Carbon (SOC) in PM_{2.5}: A Review of Recent Studies, *Atmosphere*, 9, 452, <https://doi.org/10.3390/atmos9110452>, 2018a.
- 815 Srivastava, D., Tomaz, S., Favez, O., Lanzafame, G. M., Golly, B., Besombes, J.-L., Alleman, L. Y., Jaffrezo, J.-L., Jacob, V., Perraudin, E., Villenave, E., and Albinet, A.: Speciation of organic fraction does matter for source apportionment. Part 1: A one-year campaign in Grenoble (France), *Science of The Total Environment*, 624, 1598–1611, <https://doi.org/10.1016/j.scitotenv.2017.12.135>, 2018b.
- 820 Srivastava, D., Vu, T. V., Tong, S., Shi, Z., and Harrison, R. M.: Formation of secondary organic aerosols from anthropogenic precursors in laboratory studies, *npj Clim Atmos Sci*, 5, 1–30, <https://doi.org/10.1038/s41612-022-00238-6>, 2022.
- Strollo, C. M. and Ziemann, P. J.: Products and mechanism of secondary organic aerosol formation from the reaction of 3-methylfuran with OH radicals in the presence of NO_x, *Atmospheric Environment*, 77, 534–543, <https://doi.org/10.1016/j.atmosenv.2013.05.033>, 2013.
- 825 Sumlin, B. J., Pandey, A., Walker, M. J., Pattison, R. S., Williams, B. J., and Chakrabarty, R. K.: Atmospheric Photooxidation Diminishes Light Absorption by Primary Brown Carbon Aerosol from Biomass Burning, *Environ. Sci. Technol. Lett.*, 4, 540–545, <https://doi.org/10.1021/acs.estlett.7b00393>, 2017.
- Tajuelo, M., Rodríguez, D., Rodríguez, A., Escalona, A., Viteri, G., Aranda, A., and Diaz-de-Mera, Y.: Secondary organic aerosol formation from the ozonolysis and oh-photooxidation of 2,5-dimethylfuran, *Atmospheric Environment*, 245, 118041, <https://doi.org/10.1016/j.atmosenv.2020.118041>, 2021.
- 830 Thurston, G. D., Ahn, J., Cromar, K. R., Shao, Y., Reynolds, H. R., Jerrett, M., Lim, C. C., Shanley, R., Park, Y., and Hayes, R. B.: Ambient Particulate Matter Air Pollution Exposure and Mortality in the NIH-AARP Diet and Health Cohort, *Environ Health Perspect*, 124, 484–490, <https://doi.org/10.1289/ehp.1509676>, 2016.
- 835 Tiitta, P., Leskinen, A., Hao, L., Yli-Pirilä, P., Kortelainen, M., Grigonyte, J., Tissari, J., Lamberg, H., Hartikainen, A., Kuusalo, K., Kortelainen, A.-M., Virtanen, A., Lehtinen, K. E. J., Komppula, M., Pieber, S., Prévôt, A. S. H., Onasch, T. B., Worsnop, D. R., Czech, H., Zimmermann, R., Jokiniemi, J., and Sippula, O.: Transformation of logwood combustion emissions in a smog chamber: formation of secondary organic aerosol and changes in the primary organic aerosol upon daytime and nighttime aging, *Atmos. Chem. Phys.*, 16, 13251–13269, <https://doi.org/10.5194/acp-16-13251-2016>, 2016.
- 840 Tomaz, S., Shahpoury, P., Jaffrezo, J.-L., Lammel, G., Perraudin, E., Villenave, E., and Albinet, A.: One-year study of polycyclic aromatic compounds at an urban site in Grenoble (France): Seasonal variations, gas/particle partitioning and cancer risk estimation, *Science of The Total Environment*, 565, 1071–1083, <https://doi.org/10.1016/j.scitotenv.2016.05.137>, 2016.

- 845 Tsimpidi, A. P., Karydis, V. A., Pozzer, A., Pandis, S. N., and Lelieveld, J.: ORACLE (v1.0): module to simulate the organic aerosol composition and evolution in the atmosphere, *Geoscientific Model Development*, 7, 3153–3172, <https://doi.org/10.5194/gmd-7-3153-2014>, 2014.
- Updyke, K. M., Nguyen, T. B., and Nizkorodov, S. A.: Formation of brown carbon via reactions of ammonia with secondary organic aerosols from biogenic and anthropogenic precursors, *Atmospheric Environment*, 63, 22–31, <https://doi.org/10.1016/j.atmosenv.2012.09.012>, 2012.
- 850 Viana, M., Alastuey, A., Querol, X., Guerreiro, C., Vogt, M., Colette, A., Collet, S., Albinet, A., Fraboulet, I., Lacombe, J.-M., Tognet, F., and De Leeuw, F.: Contribution of residential combustion to ambient air pollution and greenhouse gas emissions, European Environment Agency (EEA). [online] Available from: https://www.eionet.europa.eu/etc/etcatni/products/etc-atni-reports/etcacm_tp_2015_1_residential_combustion, 2016.
- Vicente, E. D. and Alves, C. A.: An overview of particulate emissions from residential biomass combustion, *Atmospheric Research*, 199, 159–185, <https://doi.org/10.1016/j.atmosres.2017.08.027>, 2018.
- 855 Wang, S., Ye, J., Soong, R., Wu, B., Yu, L., Simpson, A. J., and Chan, A. W. H.: Relationship between chemical composition and oxidative potential of secondary organic aerosol from polycyclic aromatic hydrocarbons, *Atmos. Chem. Phys.*, 18, 3987–4003, <https://doi.org/10.5194/acp-18-3987-2018>, 2018.
- 860 Washenfelder, R. A., Attwood, A. R., Brock, C. A., Guo, H., Xu, L., Weber, R. J., Ng, N. L., Allen, H. M., Ayres, B. R., Baumann, K., Cohen, R. C., Draper, D. C., Duffey, K. C., Edgerton, E., Fry, J. L., Hu, W. W., Jimenez, J. L., Palm, B. B., Romer, P., Stone, E. A., Wooldridge, P. J., and Brown, S. S.: Biomass burning dominates brown carbon absorption in the rural southeastern United States, *Geophysical Research Letters*, 42, 653–664, <https://doi.org/10.1002/2014GL062444>, 2015.
- Weber, S., Salameh, D., Albinet, A., Alleman, L. Y., Waked, A., Besombes, J.-L., Jacob, V., Guillaud, G., Meshbah, B., Rocq, B., Hulin, A., Dominik-Sègue, M., Chrétien, E., Jaffrezo, J.-L., and Favez, O.: Comparison of PM10 Sources Profiles at 15 French Sites Using a Harmonized Constrained Positive Matrix Factorization Approach, *Atmosphere*, 10, 310, <https://doi.org/10.3390/atmos10060310>, 2019.
- 865 Xie, M., Chen, X., Hays, M. D., Lewandowski, M., Offenberg, J., Kleindienst, T. E., and Holder, A. L.: Light Absorption of Secondary Organic Aerosol: Composition and Contribution of Nitroaromatic Compounds, *Environ. Sci. Technol.*, 51, 11607–11616, <https://doi.org/10.1021/acs.est.7b03263>, 2017a.
- 870 Xie, M., Hays, M. D., and Holder, A. L.: Light-absorbing organic carbon from prescribed and laboratory biomass burning and gasoline vehicle emissions, *Sci Rep*, 7, 7318, <https://doi.org/10.1038/s41598-017-06981-8>, 2017b.
- Xu, W., Croteau, P., Williams, L., Canagaratna, M., Onasch, T., Cross, E., Zhang, X., Robinson, W., Worsnop, D., and Jayne, J.: Laboratory characterization of an aerosol chemical speciation monitor with PM2.5 measurement capability, *Aerosol Science and Technology*, 51, 69–83, <https://doi.org/10.1080/02786826.2016.1241859>, 2017.
- 875 Yee, L. D., Kautzman, K. E., Loza, C. L., Schilling, K. A., Coggon, M. M., Chhabra, P. S., Chan, M. N., Chan, A. W. H., Hersey, S. P., Crouse, J. D., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from biomass burning intermediates: phenol and methoxyphenols, *Atmospheric Chemistry and Physics*, 13, 8019–8043, <https://doi.org/10.5194/acp-13-8019-2013>, 2013.
- 880 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D.

- R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophysical Research Letters*, 34, L13801, <https://doi.org/10.1029/2007GL029979>, 2007.
- 885 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N. L., Worsnop, D. R., and Sun, Y.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: a review, *Anal Bioanal Chem*, 401, 3045–3067, <https://doi.org/10.1007/s00216-011-5355-y>, 2011.
- Zhang, X., Kim, H., Parworth, C. L., Young, D. E., Zhang, Q., Metcalf, A. R., and Cappa, C. D.: Optical Properties of Wintertime Aerosols from Residential Wood Burning in Fresno, CA: Results from DISCOVER-AQ 2013, *Environ. Sci. Technol.*, 50, 1681–1690, <https://doi.org/10.1021/acs.est.5b04134>, 2016.
- 890 Zhang, Y., Favez, O., Canonaco, F., Liu, D., Močnik, G., Amodeo, T., Sciare, J., Prévôt, A. S. H., Gros, V., and Albinet, A.: Evidence of major secondary organic aerosol contribution to lensing effect black carbon absorption enhancement, *npj Climate and Atmospheric Science*, 1, 47, <https://doi.org/10.1038/s41612-018-0056-2>, 2018.
- Zhang, Y., Favez, O., Petit, J.-E., Canonaco, F., Truong, F., Bonnaire, N., Crenn, V., Amodeo, T., Prévôt, A. S. H., Sciare, J., Gros, V., and Albinet, A.: Six-year source apportionment of submicron organic aerosols from near-continuous highly time-resolved measurements at SIRTa (Paris area, France), *Atmospheric Chemistry and Physics*, 19, 14755–14776, <https://doi.org/10.5194/acp-19-14755-2019>, 2019.
- 895 Zhang, Y., Albinet, A., Petit, J.-E., Jacob, V., Chevrier, F., Gille, G., Pontet, S., Chrétien, E., Dominik-Sègue, M., Levigoureux, G., Močnik, G., Gros, V., Jaffrezo, J.-L., and Favez, O.: Substantial brown carbon emissions from wintertime residential wood burning over France, *Science of The Total Environment*, 743, 140752, <https://doi.org/10.1016/j.scitotenv.2020.140752>, 2020.
- 900 Zhang, Y., Cheng, M., Gao, J., and Li, J.: Review of the influencing factors of secondary organic aerosol formation and aging mechanism based on photochemical smog chamber simulation methods, *Journal of Environmental Sciences*, 123, 545–559, <https://doi.org/10.1016/j.jes.2022.10.033>, 2023.
- Zheng, H.-Y., Zhu, Y.-L., Teng, B.-T., Bai, Z.-Q., Zhang, C.-H., Xiang, H.-W., and Li, Y.-W.: Towards understanding the reaction pathway in vapour phase hydrogenation of furfural to 2-methylfuran, *Journal of Molecular Catalysis A: Chemical*, 246, 18–23, <https://doi.org/10.1016/j.molcata.2005.10.003>, 2006.
- 905 Zhou, S. and Wenger, J. C.: Kinetics and products of the gas-phase reactions of acenaphthene with hydroxyl radicals, nitrate radicals and ozone, *Atmospheric Environment*, 72, 97–104, <https://doi.org/10.1016/j.atmosenv.2013.02.044>, 2013a.
- Zhou, S. and Wenger, J. C.: Kinetics and products of the gas-phase reactions of acenaphthylene with hydroxyl radicals, nitrate radicals and ozone, *Atmospheric Environment*, 75, 103–112, <https://doi.org/10.1016/j.atmosenv.2013.04.049>, 2013b.
- 910 Zielinska, B.: Analysis of Semi-Volatile Organic Compound by GC/MS, DRI Standard Operating Procedure, Desert Research Institute, Reno (NV), 2008.
- Ziemann, P. J. and Atkinson, R.: Kinetics, products, and mechanisms of secondary organic aerosol formation, *Chemical Society Reviews*, 41, 6582, <https://doi.org/10.1039/c2cs35122f>, 2012.