

RC1, Sjoukje de Lange (main comments)

Thanks for writing this great manuscript. You did a great job in exploring various experimental conditions for sand-clay ripples by current/wave dominated conditions. I think you did a very thorough job with a lot of literature research. It is generally well written (although the amount of numbers in the text can be distracting), and the figures are nice and clear. It won't be much work to get this manuscript ready for publication! It will be a very valuable contribution to the scientific community.

We welcome the Reviewer's generally very favourable comments on our manuscript and thank her for the time taken in reading through it so carefully.

However, I should provide some critical notes.

(1) First of all, on first sight I find it hard to differentiate this paper from Wu's earlier publication in 2022 on deep cleaning. It might be worth it to state this difference a bit more, already in the abstract. In the introduction the difference becomes clear, but readers might be confused and mix up the two papers after only browsing through the abstract.

We agree and have now made a clearer distinction from Wu et al. (2022) concerning winnowing (deep cleaning) in the abstract.

[L12-14: '...in coastal and estuarine environments. While laboratory research on ripples forming in these types of mixed-bed environments is relatively limited, it has identified deep cleaning, the removal of clay below the ripple troughs, as an important factor controlling ripple development. New large scale flume experiments ...']

(2) Following up on that, it is good that the conclusions in the 2022 paper are largely confirmed. However, this causes repetition of many of the findings. For example, the decrease in height/length etc with clay content already known from 2022 paper, and so is the notion of deep cleaning, and the longer time till onset/equilibrium of the ripples. The new finding of this publication is the phase diagram. It would be good to focus more on this, rather than a data analysis of the raw results. Maybe it is possible to compile it in more comprehensive graphs instead of rough visualization of data, such as done in the 2022 publication.

We accept that there is overlap between this paper and the 2022 deep-cleaning paper. The idea in the present paper is that it is a synthesis, which pulls together the 2018 and 2022 papers as well as the fine sand Baas et al. (2013) paper and consolidates the findings. The 2022 and 2013 papers certainly showed a reduction in the dimensions with clay concentration. However, here we take this a step further by determining the constant of proportionality for the wave and wave-current ripples based on the orbital diameter and identifying the drop in hydraulic conductivity as a potential mechanism. While the main result is the phase diagram, there are a number of prerequisites to the discussion that are required before the phase diagram can be set forth. We hope that the revised abstract, introduction and conclusions, and re-ordering of the discussions, make this clearer.

(3) Regarding readability, it's generally very easy to read. However, in the results a lot of numbers are given, including many abbreviations. Additionally, many comparisons are made. It causes me to get lost in your data, and I miss the main message. Maybe it would help to start giving the main take away, and support this with numbers, rather than giving a lot of numbers from which the reader has to derive the conclusion themselves, making it hard to read and hard to concentrate.

We have now incorporated nomenclature and abbreviation lists after the conclusions (L537-539), to facilitate readers' comprehension of symbols used throughout the text, ensuring ease of understanding. In addition, we have streamlined reference to T_η and T_λ , which are not relevant to the discussion section (see RC2.1). We have also substantially reduced reference to specific numbers in the conclusions, to help emphasise the main message, in addition to the restructuring listed above. We believe these modifications will significantly enhance the readability and clarity of the paper.

All smaller comments can be found in the attached pdf document. Please contact me if you have any questions regarding my feedback!

We have included the highlighted sections of text from the pdf in our responses below.

RC1, Sjoukje de Lange (smaller comments)

(4) L14: ‘experiments seek to address **this** by considering ...’

what exactly? For me it sounds like you want to do lab research on mixed-ripples, simply because it has not been done before. I'm not sure what "this" is that requires addressing

We agree that further clarification is necessary here and have made the following changes to the text:

[L14-18: ‘... New large-scale flume experiments seek to address this **sparsity in data** by considering two wave–current conditions with initial clay content, C_0 , ranging from 0 to 18.3%. The experiments record ripple development and pre- and post-experiment bed clay contents, to quantify clay winnowing. The **present** experiments are combined with previous **wave-only, wave–current and current-only experiments** to produce a consistent’]

(5) L18-19: ‘... $C_0 > 10.6\%$, likely associated with **hydraulic conductivity**.’

I would specify this to “an increase in hydraulic conductivity”

We accept this, but it is a decrease not an increase in hydraulic conductivity. The text has been changed:

[L20-21: ‘... for $C_0 > 10.6\%$, likely associated with **a three-orders of magnitude decrease in** hydraulic conductivity.’]

(6) L19-20: ‘**Accompanying the sudden change in steepness is a gradual linear decrease in wavelength with C_0 for $C_0 > 7.4\%$** , which may ...’

just wondering (without having read the rest of the paper), is this also related to this hydraulic conductivity?

We do not think so, as it is a gradual rather than sudden change.

(7) L26-27: ‘**Winnowing occurs for both flat and rippled beds, but the rate is two orders of magnitude smaller for flat beds.**’

this line feels a bit misplaced, as it states a result, at a location where you would expect a discussion/implication

We agree with the Reviewer and have relocated this sentence about winnowing to an earlier position in the abstract, as shown below, so that the abstract ends with the previous sentence.

[L22-24: ‘is a gradual linear decrease in wavelength with C_0 for $C_0 > 7.4\%$. **Ultimately, for the highest values of C_0 , the bed remains flat, but clay winnowing still takes place, albeit at a rate two orders of magnitude lower than for rippled beds. For a given flow, the initiation time, ...** which has important implications for morphodynamic modelling.’]

(8) **L28: ‘1 Introduction’**

I think you did a great job explaining the importance of phase diagrams under these specific conditions. However, why do we care about clay winnowing? It would be good to add a few sentences about this.

The winnowing of clay causes mixed sand–clay ripples to return to clean sand ripples so as a mechanism it is important. Deep cleaning, which removes clay from below the active layer provides a greater supply of clean sand to form ripples. Also on a more practical level, we make use of winnowing to go from high to low clay content experiments (L112-113). The importance of winnowing has been addressed in the updated introduction, which includes a contrast between the 2018 and 2022 papers (L50-59).

(9) L36-37: ‘... vice versa ~~for coastal and estuarine management and ecological balance maintenance,~~ especially as such areas may’

We are not sure why this has been deleted as there is no comment, but this phrase has been removed from the modified introduction in any case (L39-40).

(10) L61-63: ‘under combined flows and revealed the formation of small flat ripples with an initial clay content, C_0 , greater than 10.6% under hydrodynamic conditions that would generate large equilibrium ripples in clean sand.’

I find this sentence a little bit confusing. I understand you want to say that the ripples with clay are smaller than the clean-sand equivalent (right?), but I would suggest rephrasing.

We accept this and have rephrased the sentence:

[L47-50: ‘... Wu et al. (2022) were the first to consider ripple dynamics in sand–clay mixtures under combined flows. Their study showed that small flat ripples formed for initial clay contents, C_0 , greater than 10.6%, under hydrodynamic conditions that formed large equilibrium ripples in clean sand.’]

(11) L66-67: ‘Wu et al. (2022) also found that the clay loss rate from the bed was much lower when $C_0 > 10.6\%$, as the stronger cohesion resisted further ripple development.’

It might be worth it to briefly relate this to Schindler's Sticky Stuff paper (although it is unidirectional flow focussed on dunes).

Schindler et al. (2015) showed that increasing C_0 reduces dune size in unidirectional flow, but they did not quantify the clay loss, so this reference cannot be used here. Schindler et al. (2015) is discussed in the context of the 3D phase diagram (L459), but was mistakenly omitted from the reference list. This has now been corrected (L671-674).

(12) L75-77: ‘... under different hydrodynamic conditions; (2) to compare clay winnowing efficiency, based on quantifying bed clay content during ripple development, under different flow conditions;’

Based on the introduction, it is unclear to me why we care about clay winnowing. Would it be possible to expand a bit on this?

We have now linked these three points more closely together (see also RC1.8):

[L82-86: ‘... The three specific objectives of this study were: (1) to compare ripple development and occurrence on beds with similar initial clay contents under different hydrodynamic conditions; (2) to compare clay winnowing efficiency during ripple development, under different flow conditions; and (3) to propose a new phase diagram for bedforms generated in sand–clay substrates to systematise the experiments, using the knowledge gained from objectives 1 and 2.’]

(13) L94: ‘... flow conditions of increased relative current strength, with $U_c \approx 0.3 \text{ m s}^{-1}$ and’
do I understand correctly that U_c does not vary between the two new experiments?

Yes, that is correct, only the wave strength was changed.

(14) L103: ‘..., with C_0 of 9.9%, 13.1%, and 18.3% in channels 1 to 3, were under WC3 flow’
maybe write "strong" flow conditions or something like that, so that the reader does not need to flip back and forth to the table?

The idea is that the numbering system represents the relative strength of the current from weakest WC1 to strongest WC3, see L105-108 as well as L158-159 of the table, precisely so that it is not necessary to keep looking at the table, but perhaps this was not clear enough so we have made the text more explicit:

[L105-108: ‘... The wave–current flow conditions (WC1, in Table A1 from Wu et al., 2022; WC2 and WC3, in Table 1) are numbered according to increasing relative current strength, from the weakest, WC1, to the strongest, WC3. The experiments involved ...’

L158-159: ‘The flow code represents the relative current strength, where WC1 (Wu et al., 2022, Table A1) is the weakest and WC3 is the strongest.’]

(15) L104: ‘... (Table 1). The beds in Run 2 remained essentially flat after 650 min. In Run 3, ...’
this seems like it belongs in the results?

We can see how the Reviewer might think this, but the sentence is really only describing what is stated in Table 1, which describes the flow conditions and the general nature of the bed: rippled or flat.

(16) L111: ‘in each channel before and after the run, **after the water had been drained.**’
did the drainage cause any sediment to be remobilized? And if not and drainage was very slow, how did you prevent the clay depositing in the ripples troughs?
The draining of the flume was sufficiently slow, and the deposition of material had a minimal effect as it was only a surface dusting of clay.

(17) L111-112: ‘Furthermore, the flow was **temporarily stopped during the runs** at pre-set times ...’
I got a bit a similar question here - was the stopping time short enough to prevent deposition of the suspended material?
The cores were taken before the clay had a chance to settle out, so this was not really an issue either. A sentence has been added to explain this:
[L123-125: ‘... length of 90 mm. **These short pauses in the experiments resulted in negligible bed deposition of suspended clay particles due to their slow settling velocity.** Details of the ...’]

(18) L115-116: ‘... was excluded from the analysis, **because the sediment in this channel was not sufficiently well mixed.**’
how did you discover this issue?
This was determined through core samples, which has now been clarified in the text:
[L126-127: ‘..., because **core samples collected from the flat bed showed that** the sediment in this channel ...’]

(19) L135: ‘were **co-linear.** Whilst θ_0 is clearly affected by grain size, it is assumed that the other ...’
I'm not sure what you mean with this? What are co-linear experimental flows?
This simply means the wave and current are aligned.

(20) L145: ‘**Table** 1 Experimental parameters’
I'm kinda missing the logic behind how this table is ordered. It seems to be on run/channel, which has very little physical meaning to me. Why not organize it either on C_0 content, flow code, or current intensity?
These are organised in the order in which the experiments were conducted to take advantage of natural winnowing even when the bed remained flat (see L112-113).

(21) L168-169: ‘... growth. **Finally, the sediment concentration profiles were characterised by a Gaussian-type function.**’
why was this done?
This was done to be consistent with the results in Wu et al. (2022) and also because the core results were sometimes quite noisy and these help as trend lines (see Figure 5).

(22) L175: ‘marked **C - current-dominant, WC - wave-current, W - wave-dominant**, NM - no motion’
out of curiosity: is there a way to discriminate between current ripples, wave ripples and mixed ripples in terms of geometry? And if so, does the height/length/etc of the ripples observed in these WO/WC/C regimes match the expected geometry?
The Reviewer is jumping the gun here, as we discuss this in the results section. The categorisation here is based solely on the wave and current strength with reference to Baas et al. (2021).

(23) L176: ‘**WO – washout.** The threshold of motion ...’
what will be in the upper right corner of this diagram, for even stronger currents than washout conditions?

In principle flat featureless upper stage plane beds, as found by Baas et al. (2021), but we could not generate these in the present experiments. In angular wave–current cases there is also the occurrence of lunate ripples to consider. We have added a sentence to the text to point this out:

[L151-153: ‘... threshold of motion, $\theta_c + \theta_w = \theta_0$. It is anticipated that ultimately above washout there are flat featureless upper stage plane beds (for waves and currents at angles this may be complicated by the occurrence of lunate ripples, Baas et al., 2021). The reason that not all conditions’]

(24) L184: ‘147.9 mm, required a longer $T_\lambda = 302 \text{ min}$ (Figure 3a and b; Table 2). As shown in ...’
I would love to see a figure that plots equilibrium time (and/or initiation time?) with clay content!
The reason we have not shown these is because similar plots were shown in Wu et al. (2018, 2022) and they are not relevant to the discussion. Also, Figure 7g (now Figure 8g) already shows the initiation time versus θ_{0E} .

(25) L189: ‘Figure 3. Development ...’
why are figure a and b not having the same width as the other figures? I would either make them the same width, or scale the x-axis similarly (thereby decreasing the size of them even more).
We agree that this makes the figure look untidy so have made the Figure 3a and b subplots the same width.

(26) L193: ‘intervals. The yellow lines are the best-fit curves for clean sand (a, b).’
in c and d, right?
The yellow lines are in c and d, but they have been taken from (a, b). This has been clarified in the text:

[L205: ‘The yellow lines in c and d are the best-fit curves for clean sand from (a, b).’]

(27) L195: ‘Figure 4. Plan view ...’
love this figure! Would it be possible to use the same scale bar for each figure though? That would clean up the figure a bit.

Unfortunately, if all of the scales are the same, definition will be lost in e and f, but we have reduced it to the two-colour bars with common scales for a-d and e and f in the revised Figure 4, and changed the caption to explain this:

[L209-210: ‘Note the elevation scale for d also applies to a – c, and the elevation scale for f also applies to e.’]

(28) L223: ‘Table 2. Bedform characteristics’
this table is differently organised than table 1
That is correct. Unlike Table 1, which is based on the experimental order (see above), this is grouped according to flow type to aid in the description of bedform types in the results.

(29) L237-238: ‘active layer (Figure 5e). The equivalent clean-sand depth (black horizontal dashed line), $d_c = I/C_0$, was 57 mm, more than two times the ripple height (25 mm).’
interesting! I hope to read something about this in the discussion!
This is the deep cleaning that is shown in Figure 7 in the revised manuscript.

(30) L252: ‘... for WC3 conditions (pre- a–d and post- e–h) and for WC2 conditions (pre- f–h ...’
since the headers of figure a-d indicate "initial" in it, why not indicating "end" (or something equivalent) in the headers of e-k?

We accept that this figure is a little confusing and have changed ‘initial’ to (pre-) in the titles of Figures 5a-d and added (post-) to the titles of Figures 5e-k, so that the terminology is consistent with the caption. However, it is important to stress that Figures 5f-h are the pre-experiment cores for WC2 (L259). To emphasise this point, we have added the depth-mean post-experiment value in blue for Figures 5f-g, as this represents C_0 values for Figure 5i-k.

(31) L264: ‘... In the present experiments under WC2 conditions, this discontinuity in equilibrium’
So why not under WC3 conditions? (maybe you talk about it later, but this is the first question popping up when reading this).

This is because, for WC3 conditions, the higher clay contents are below threshold, so no ripples formed, which we discuss in the threshold of motion section.

(32) L284-285: ‘Other than this reduction in steepness, current strength was found to have a modest influence on ripple geometry;’

Interesting!

(33) L290: ‘..., $D_{50} = 0.45$ mm, may have resulted in a greater tendency towards 2D symmetric ripples,’

Didn't Baas show in his 1994 paper that ripples always develop towards 3D linguoid structures when in equilibrium? How would you use that finding compared to yours?

Sedimentology (1994) 41, 185-209.

The Baas (1994) ripples are for unidirectional flow, which tend to be far more asymmetric and linguoid. The present ripples, while formed by waves and currents, are wave-dominated (below the diagonal $\theta_c = \theta_w$ line in Figure 2). So, the present experiments are much more 2D and straight-crested in character than Baas' linguoid ripples, with only subtle changes in the ripple character as a result of the current (L290-291).

(34) L304-306: ‘described by $\lambda_e = \alpha d_{wc}$, where α is the constant of proportionality (= 0.62, according to Wiberg and Harris, 1994) and d_{wc} is the orbital diameter enhanced by the current. Appendix B explains how d_{wc} is determined based on a sinusoidal wave. The ...’

Although you explained this in a bit more detail in the appendix, I think the storyline would benefit from a longer explanation here. The result is beautiful, and leads to an important figure in your paper. It's a shame it remains unclear what αd_{wc} exactly is until doing a deeper dive into the appendix. additional question - is there a physical explanation behind this equation?

We accept that this could be made clearer in the main text. The orbital diameter is an important parameter for wave ripples as it represents the distance a neutrally buoyant particle will move in half a wave cycle and controls wave ripple spacing. Thus, when a wave and current are added together the particle will move further and d_{wc} will be larger. We have added a sentence to the text to reflect this: [L311-314: ‘1994) and d_{wc} is the orbital diameter enhanced by the current. The orbital diameter scales the ripple spacing, as it represents the distance a neutrally buoyant particle can move in half a wave cycle (Clifton and Dingler, 1984). With the addition of a current, this particle can move further and Appendix B explains how d_{wc} is determined based on a sinusoidal wave.’]

(35) L314: ‘where $C_{0m} = 7.4\%$, such that $\alpha = 0.31$ when ...’

what would happen if you would choose 8% here?

While making $C_{0m} = 8\%$ might seem tempting based on L295, it worsens the fit, as $C_{0m} = 7.4\%$, which corresponds to the upper limit of Wu et al.'s (2018) experiments, best describes the data in Figure 6b. We therefore would prefer to keep it as 7.4%. A phrase has been added to L295 to explain where the 7.4% value comes from.

[L300-301: ‘... for $C_0 > 7.4\%$ (the upper limit of Wu et al.'s (2018) experiments). Interestingly, ...’]

(36) L318: ‘Figure 6. Equilibrium ...’

great figure, love how it all collapses!

Thank you.

(37) L323: ‘4.2 The enhanced threshold of motion’

how would this story hold if you would use the calculations as proposed in van Rijns 2019 paper [https://doi.org/10.1061/\(ASCE\)HY.1943-7900.0001677](https://doi.org/10.1061/(ASCE)HY.1943-7900.0001677) It might be quite applicable to your story.

For coarse sand, van Rijn (2019) used the exact same equation for the threshold of motion (see L140-141 and his eq. 1a). However, here we are inferring what the enhanced threshold would have to be, based on whether or not ripples form, so his enhancement might be different, but on L445-452 we are using his explanation for the different coarse and fine threshold enhancements (in L446 and L690 the year has been changed to 2019 as above since it is the same reference).

(38) L369-370: ‘Figure 7 clearly shows that for the wave-only and wave-current cases, the initiation time is related to the fraction of the wave cycle when shear stresses are above threshold.’

is this the take-away from the figure? I had a hard time reading through the text trying to find why you were suddenly looking at wave-orbit-time scales. Maybe you can clarify the aim before diving in the analysis?

This section is all about choosing a value for P_θ in eq. (7) for the enhanced threshold that is consistent with the results: below threshold for flat featureless beds and above threshold for rippled beds (L387-402) based on the size of the maximum stress in the wave cycle ($= \theta_c + \theta_w$). This may not have been made clear enough, so a phrase has been added to L394-395. This is a necessary first step that is required before Figure 7 (Figure 8 in the revised manuscript, L418) can be shown, as it determines the position of the yellow lines in each case. Figures 8a-f confirm the choice of P_θ , but provide more detail by showing the time varying signal, eq. (8). Thus, Figure 8g is the main additional information provided by the figure that summarises the time-varying behaviour.

[L394-395: ‘For combined co-linear wave–current flow, the threshold line is described by $\theta_{0E} = \theta_c + \theta_w$, where $\theta_c + \theta_w$ is the maximum stress in the wave cycle. This results ...’]

(39) L379: ‘.... $1+6.4C_0$, $D_{50} = 0.143$ mm,’

is the D_{50} really a good measure to describe clay-sand mixtures? It really just consists of two modi, and the D_{50} is a value that does not mean much in this case.

While we accept that D_{50} of the mixture will be distinct from D_{50} of the sand, D_{50} of the mixture, or any other representative measure of the mixture, will vary in time and between experiments and so will not help in their characterisation. D_{50} of the sand remains constant in the experiments and provides a useful way of grouping them together, and there is also a clear sand-grain size dependence in the enhancement in eq. (6). Also, in L369-376, we do use D_{10} of the mixture to explain the sudden potential drop off of hydraulic conductivity and ripple steepness.

(40) L386-387: ‘Thus, the thicker clay layer around the coarser grains likely causes increased enhancement of the threshold of motion, which has also been found in field observations (Harris et al., 2016).’

So like the hiding-exposure effect? Or does this have nothing to do with it?

No, this is opposite to the hiding-exposure effect, which tends to make coarser grains easier to move and finer grains harder to move.

(41) L395: ‘4.3 The potential factors controlling the deep cleaning of clay’

I'm still not entirely sure why this is important?

We hope that our responses and suggested changes to the introduction and conclusions (RC1.1, RC1.8) have helped to make this clearer. However, we recognise that this section requires an opening sentence to set out the significance of winnowing:

[L332-333: ‘Winnowing is an important mechanism in ripple development, because it provides the necessary supply of clean sand in the active layer from which ripples can grow. Significant clay ...’]

(42) L440: ‘This reduction in hydraulic conductivity alone, does not fully explain the behaviour ...’
so you did not measure this right? it's purely from the theoretical analysis?

That is correct, we did not measure the hydraulic conductivity.

(43) L446: 'Figure 8. Average clay mass flux ...'

I'm not so sure about this figure. I find it pretty hard to interpret, mostly because it's hard to interpret all the colours and type of markers. Would there be a way to make this clearer? For example by shortly describing the experiments, rather than using their abbreviations? All the colours and abbreviations distract me, causing that I don't really understand what the purpose of this figure is.

The idea of this figure is to try to quantify the clay loss, as it relates to deep cleaning, which is important for providing the supply of clean sand available to build ripples. It is really only an order of magnitude plot, showing the difference between ripples and no ripples, and should be viewed as such. Throughout the manuscript we have tried to use consistent colours for the different cases WC1-WC3, Wu et al. (2018) and Baas et al. (2013) and symbols for large, small and no ripples. We feel this is the most concise way to describe the different conditions and ripple types, since using weak, medium and strong wave will soon get very confusing and affect readability (see RC1.14). However, we accept that this is distracting in Figure 8 (Figure 7 in the revised manuscript, L383), so have reduced the legend to the three colours representing WC1–WC3 in the new Figure 7, explained the different symbols without abbreviations in the figure caption instead and labelled the two main groupings: ripples and no ripples. Also, for clarity the caption no longer lists the values of C_0 for WC1. While this does not completely address the Reviewer's concerns, we feel the various colours and symbols should be retained so that the cases can be distinguished from one another.

[L383-385: 'Figure 7. Average clay mass flux out of the bed, F_b , against initial clay content, C_0 , for WC1 (red; Wu et al., 2022), WC2 (light blue) and WC3 (dark blue). Here solid circles, asterisks and open circles signify large, small and no ripples.']

(44) L449: '4.4 Implications for paleowave climate predictions and bedform phase diagrams'

great section!

Thank you.

(45) L469: 'dominant (W and SW; $\theta_c < 0.2\theta_w$), wave-current ...'

what is the S in those subdivisions?

'S' stands for small as it says on the previous line in the text (L471).

(46) L480: 'and therefore form roughness, which is proportional to η^2/λ .'

Form roughness? Where did you do an analysis on this? (sorry if I simply missed it, you all did a lot!)

No analysis has been done, but it is generally accepted that form roughness relates to the ripple dimensions (η^2/λ) and a reference has been added to clarify this:

[L498: '..., which is proportional to η^2/λ (Soulsby, 1997). In the specific case ...']

(47) L490: 'Figure 9. Orthographic projection of the 3D phase ...'

would it be possible to present these figures without all the abbreviations (NM, WO, SC...) in it? I would improve readability.

We accept that these abbreviations make the figures more difficult to read, but the problem is that these are needed to demarcate the regions and writing them out in full will make the figures far busier.

(48) L503-508: '... quasi-symmetric in geometry (ripple steepness, $RS = 0.11$; ripple symmetry index, $RSI = 1.5$), with equilibrium heights and wavelengths, $\eta_e \approx 15$ mm, $\lambda_e \approx 148$ mm. Under WC2 conditions, for $C_0 = 8.4\%$, less steep, 2D and quasi-asymmetric ($RS = 0.09$ and $RSI = 1.4$) equilibrium ripples developed, with $\eta_e = 13.9$ mm, $\lambda_e = 137$ mm. However, for $C_0 = 10.8\%$, equilibrium ripple dimensions drastically decreased to $\eta_e = 5.6$ mm, $\lambda_e = 125$ mm, and the ripples transformed to flatter, 3D and asymmetric geometries ($RS = 0.04$ and $RSI = 1.6$).'

is it really needed to give all those numbers here? I think a description would suffice, improving readability.

We have restructured the conclusions (L508-535), shortening some of the preceding description (L508-514). Now all the results of the present experiments are summed up in the first paragraph, reducing a lot of numbers and definitions. The following two paragraphs pull together all the data and then the phase diagram results.

(49) L529-532: ‘and fine ($D_{50} = 0.143$ mm) mixed sand–clay motion. On the basis of these enhancements, new 3D phase diagrams, involving the non-dimensional wave and current shear stresses and C_0 are proposed to characterise the two ripple size groupings under different flow conditions. This new 3D phase diagram framework should prove important to the morphodynamic modelling community.’

For me, this is the main result of your work! I think this deserves way more attention than to be tucked away at the end of your conclusions.

We agree that this is the most important result of the paper, which is precisely why it is at the end of the conclusions. However, we accept that it needs more prominence in the rest of the paper so have moved the 3D phase diagram and the bulk of the associated text from the implications section to a new section (4.4 Bedform phase diagrams) at the end of the discussion section (L458-489). There is now a shorter implications section with the same title as before (L490-506) outlining the wider implications of the changes in bedform dimensions and 3D phase diagrams. Also, we have swapped sections 4.3 and 4.2, so that the threshold of motion leads on to the phase diagram. We now finish on the most important point according to both the Reviewers and the order is consistent with the abstract and conclusions. In addition, that way our reduction in hydraulic conductivity argument then relates back more immediately to the sudden drop in steepness discussed in section 4.1.

RC2, Anonymous (main comments)

The submitted manuscript provides rigorous analysis on wave-current driven ripple development in cohesive clay bed. The analysis is based on a series of laboratory experiments. The discussed topic is interesting and important to advance our understanding of coastal morphodynamics.

We are grateful for the Reviewer's positive feedback regarding our contributions to the field of coastal morphodynamics.

(1) However, I find it difficult to read this paper. The main reasons of the difficulties in reading this paper are mainly associated with too many presented numeric and abbreviation when things are explained in the result section.

We have added nomenclature and abbreviation lists for readers to more easily understand the symbols. In addition, since equilibrium time, T_η and T_λ , are not really relevant to the discussion section, we have streamlined their mention in section 3.1 (L194-198; L218-224) and instead stated that they show similar trends to Wu et al. (2022):

[L226-227: ‘... and $RS = 0.04$ (Figure 4e, Table 2). **The general trends in equilibrium time for ripples height, T_η , and wavelength, T_λ , with C_0 were similar to those found by Wu et al. (2022).** For the $C_0 = 16.3\%$ (WC2) case, the bed ...’]

(2) In addition, the authors refer earlier studies regarding experiment set-up and results from Wu et al. (2022). I often get lost what the main messages are in the texts or figures. Thus, some clarifications and rework in writing are needed before publication. I suggest the author focus on telling the main messages from their experiments and each figure in the revised manuscript. I also have some concerns about configurations of their experiment.

In our response to Reviewer 1, we hope that we have addressed Reviewer 2's concerns about the restructuring of the text (see RC1.2, 3, 48 and 49) and clarifying the figures (see RC1.30, 43). As regards the reviewer's concerns about the experimental configurations, these are addressed below.

RC2, Anonymous (main comments)

I am sure that the authors meticulously design and run the experiments, but I have some concerns about their experiment configurations, which can affect experimental condition characterization and results. Please see below for the details.

(1) The experiments were conducted in a recirculating flume tank, where two different types of sediment are placed in the streamwise direction (mixed sand clay in upstream and clean sand downstream). In such configuration, only clean sand is likely to be recirculated during the experiments, which can change clay – sand mixture ratio in the test section. Although the authors argue the decreases in clay content after the experiments in Figure 5 are mainly associated with clay winnowing, the decreases in clay content can be also affected by the recirculated sand. Was the recirculated sand transport rate measured and negligibly small? More explanations are needed in this matter.

We agree with the Reviewer that a fraction of clean sand was recirculated during the course of the experiments. Based on the largest grains in suspension, determined by a settling velocity associated with the skin friction (Fredsoe and Deigaard, 1992, doi: 10.1142/1546), we estimate it to be about 1% of the sand grains on the bed in the clean-sand section. We believe that the recirculation of this clean-sand fraction has a negligible effect on the dilution of clay content in the mixed sand-clay section. This is demonstrated by the vertical consistency in clay content observed in post-experiment sediment cores collected from the flat beds (f – h, and k in Figure 5), where the mixed sections were below threshold, but the clean-sand section was above threshold. Had there been a substantial effect in these cases the cores would have shown some vertical variation.

(2) I presume the flume is separated with the brick walls to maximize the number of experiments in one run. The height of the brick walls (0.2 m) is a half of the water depth (0.4 m). Why is it not entirely separated (the height of the brick walls > 0.4 m)? I can imagine that some three-dimensional flow

structures are generated at the interface between top (> 0.2 m) and bottom (< 0.2 m) layers, which would not exist in normal flow conditions (i.e. without flume separations). Did authors check vertical velocity profiles before flume separation are comparable with the ones obtained after flume separation? If there are substantial differences between the velocity profiles, quite different bed shear stress can be obtained even with the same experiment conditions. This can affect your 3D phase diagram.

The brick walls were indeed used to maximise the number of experiments for each run. In these experiments and those of Wu et al. (2022), short walls were chosen for two reasons: (i) to help ensure flow was more consistent across the tank as a whole between channels and (ii) so that it possible to take sediment cores during the experiment when the flow was temporarily paused.

We acknowledge that this short-wall configuration could potentially induce the development of 3D flow structures, but we did not take any velocity measurements prior to putting the walls in place for practical reasons (the walls were built before the sediment was added). Nonetheless, we presumed that this 3D flow pattern would be minimal. This presumption is supported by the consistent and conventional, cross-channel ripple structure for both 2D and 3D ripples (see Figure 4). Moreover, our results show a greater tendency towards 2D straight-crested ripples than other experiments (L286-292). As far as the 3D phase diagram is concerned, each channel had its own hydrodynamic measurement determined along the centre line where any 3D flow effect would be the least possible.

(3) Also, regarding the flume separation, there seems to be a preferential bed slope that can affect ripple development. For example, more sediment aggregation (larger ripples) is observed in the upper part of the panel c,f (channel 3) in Figure 4. Similarly, more sediment aggregation (larger ripples) is observed in the lower part of the panel b,d (channel 1) in Figure 4. This can influence ripple morphology characterization (particularly height and steepness). Some explanations are needed in this matter.

We believe that sediment aggregation was mainly due to a degree of wave reflection despite wave absorption at the downstream end of the flume by the artificial beach. It may also be related to competing patches, where wave ripples first form, spreading out and interacting with one another (Sekiguchi and Sunamura, 2004, doi: 10.1016/j.coastaleng.2003.11.002). Similar behaviours have been reported previously within this wave tank (Wu et al., 2018, doi:10.1029/2018JF004681). However, our approach involved the recording of ripple data through URS sensors across the entire channel within a 2-meter URS-scanning area to capture the average ripple development characteristics. This methodology effectively mitigated the potential influence of uneven sediment aggregation. A phrase has been added to L166 to make this clearer. Most importantly, as depicted in Figure 6a, there is a consistent trend in ripple steepness for larger ripples as the clay content increased from 0 to 10.8%. It also suggests that the impact of both wall-induced 3D flow structures, as mentioned above, and wave reflections on ripple geometries is negligible.

[L165-166: ‘... by calculation of the mean, across-channel values of λ_t and η_t at a bed scanning ...’]

(4) Lastly, the test section is located fairly close to the current inlet (only ~ 2 times of the channel width excluding the gravel section). I wonder whether the space is enough for the flow to fully develop. If the flow is still in transition (not fully developed), this can explain why slightly inhomogeneous ripple fields are generated in different channels that are described above. Please address this matter if the authors have checked the flow is fully developed.

Based on the work of Elliot (1958), doi:10.1029/TR039i006p01048, in response to a change in roughness from skin friction, $z_0 = 2.5D_{50}/30 = D_{50}/12$, to form drag, $z_0 \sim \eta_e^2/\lambda_e$, the distance along the tank for full boundary layer development in water depth of 0.4 m is 0.28 m, so the 1-m gravel section is sufficient.

RC2, Anonymous (specific comments)

(5) In the abstract, please avoid speculative words, such as “likely associated with hydraulic conductivity” or “which may be significant for paleoenvironmental reconstruction”. I further wonder

whether readers can understand the reason of your findings at this stage. It could be worth skipping such explanations and leaving them for the discussion section.

We argue that the first example using ‘likely’ to describe an effect is not speculative, particularly as we have now made it more specific by adding that there is a ‘3-orders of magnitude reduction in hydraulic conductivity’ in response to Reviewer 1 (RC1.5). However, we accept that the second example is speculative and so have removed it:

[L22: ‘is a gradual linear decrease in wavelength with C_0 for $C_0 > 7.4\%$. Ultimately, for the highest values of C_0 ,’]

(6) L29-30: *In the introduction, please define combined wave-current flows before using it.*

We have introduced the definition of combined flow at the beginning of the introduction:

[L31-34: ‘In coastal and estuarine environments, the superimposition of waves and currents, termed combined wave–current flows, are a common occurrence. In these environments, which include continental shelves, the shoreface, and tidal flats, combined wave–current flows frequently form ripples on the seabed (e.g., Osborne and Greenwood, 1993; ...’]

(7) L74-77: *Importance of point (1) and (2) is not clearly stated in the introduction. I also think point (3) is the main scientific contribution of this paper. Please consider putting point (3) first.*

We have restructured the introduction to emphasize the significance of points 1 and 2. The revised introduction also underscores the importance of understanding points 1 and 2 as fundamental prerequisites for addressing point 3 (see RC1.8, 12).

(8) L83: *temporal resolution of the velocity measurement looks low. From my understanding, Vectrino profiler can collect velocity samples up to 200 Hz. Is there any reason of low temporal resolution? If there is, please explain it in the revised manuscript.*

The purpose of measuring the velocity in the experiments is to obtain the average velocity necessary for determining skin friction using Malarkey and Davies’ (2013) method as we mentioned in L132. Consequently, variations in the ADV sampling frequency do not impact the results of the mean data. Hence, we opted to use a sampling frequency of 25 Hz in these experiments.

(9) L83: *please also provide the velocity measurement duration.*

We have added this information:

[L92-93: ‘and flow velocities in each channel were measured by a 25 Hz Vectrino profiler throughout the entire duration of the experiments. Freshwater ...’]

(10) *Please consider change notation of wave velocity magnitude U_0 into U_w for the sake of readability.*

We accept this, particularly as it is more consistent with θ_w , particularly as θ_0 is used for the clean-sand threshold of motion, and so we have changed U_0 to U_w where it appears in the text and figures.

(11) L103: *please elaborate how C_0 is measured? Is it based on mass or volume? If it is based on the volume, how the bed porosity is considered?*

C_0 is based on mass. This has been clarified in the text:

[L113-114: ‘..., and Run 2, with C_0 of 9.9%, 13.1%, and 18.3% by dry weight in channels 1 to 3...’]

(12) L115-116: *please explain how you define “sufficiently not well mixed channel” Is it based on visual inspection? Or is it based on inhomogeneous ripple development?*

This was determined through core samples which has now been added to the text, as explained above (RC1.18).

(13) Please consider rearrange the order of experiments in Table 1. Consistent grouping based on experimental parameters seems to me more important than chronological order. To be specific, it would be better to present results of WC3 in an order of increasing C_0 .

We believe presenting the experimental process in chronological order is essential for the reader's understanding, as it helps illustrate how the experiments were conducted; relying on the natural winnowing of the bed, as explained above in response to Reviewer 1. Table 2, and indeed Tables A1-A3, all of which list ripple characteristic rather than flow parameters, are arranged by flow type and in increasing order of C_0 , so that trends can be picked out more readily.

(14) Please consider putting experimental parameters of the WC1 experiments in Table 1. Please clearly state that the WC1 results are from Wu et al. (2022).

We believe Table A1 is a more suitable location for the WC1 experiments as it helps clarify that the WC1 experiments are taken from Wu et al. (2022) and separate from the present experiments but with the same experimental setup. However, we agree that incorporating a summarized overview of the parameters used in the WC1 experiments within Table 1 would significantly enhance readability. Consequently, we have expanded note 1 following Table 1 to accommodate this additional information (see also RC1.14):

[L159-160: '... and WC3 is the strongest. For WC1 conditions (Table A1), corresponding to $0 \leq C_0 \leq 12.6$, $0.31 \leq U_w \leq 0.33$ m/s, $0.15 \leq U_c \leq 0.2$ m/s, $0.15 \leq \theta_w \leq 0.16$ and $0.007 \leq \theta_c \leq 0.13$, ripples always developed.']

(15) Figure 2: If I am not mistaken, WC1 results are from Wu et al. (2022). Please clearly state this. Also, last tick of 0.25 in the x-axis is cut out in Figure 2.

Both of these issues have been rectified in the revised Figure 2 and caption:

[L186: '(Wu et al., 2018), wave-current, WC1 (Wu et al., 2022), WC2 and WC3, and current- ...']

(16) L260: Please discuss how the threshold of $C_0 \sim 10\%$ will change under saline water (more realistic flow condition in coastal regions), not freshwater.

The data of Wu et al. (2018) experiments, see Table A3, were performed in salt water. While these experiments do not cover a very wide a range of clay concentrations, $0 \leq C_0 \leq 7.4\%$, the fact that these experiments are consistent with all the other freshwater experiments leads us to believe that if there is an effect it is likely to be very weak. In addition, our argument that hydraulic conductivity drops significantly at $\sim 10\%$ does not depend on the presence of salt or fresh water.