

The role of upper ocean heat content in the regional variability of Arctic sea ice at sub-seasonal time scales

Elena Bianco^{1,2}, Doroteaciro Iovino¹, Simona Masina¹, Stefano Materia^{3,4}, and Paolo Ruggieri⁵

¹Ocean modeling and Data Assimilation Division, Centro Euro-Mediterraneo Sui Cambiamenti Climatici (CMCC), Bologna, Italy

²Department of Environmental Sciences, Informatics and Statistics, Ca' Foscari University, Venice, Italy

³Climate Simulations and Predictions Division, Centro Euro-Mediterraneo Sui Cambiamenti Climatici (CMCC), Bologna, Italy

⁴Barcelona Supercomputing Center, Barcelona, Spain

⁵Department of Physics and Astronomy, University of Bologna, Bologna, Italy

Correspondence: Elena Bianco (elena.bianco@cmcc.it)

Abstract. In recent decades, the Arctic Ocean has undergone changes associated with enhanced poleward inflow of Atlantic and Pacific waters and increased heat flux exchange with the atmosphere in seasonally ice-free regions. The associated changes in upper ocean heat content can alter the exchange of energy at the ocean-ice interface. Yet, the role of ocean heat content in modulating Arctic sea ice variability at sub-seasonal timescales is still poorly documented. We analyze ocean heat transports and surface heat fluxes between 1980-2021 using two eddy-permitting global ocean reanalyses, C-GLORSv5 and ORAS5, to assess the surface energy budget of the Arctic Ocean and its regional seas. We then assess the role of upper ocean heat content, computed in the surface mixed layer (Q_{ml}) and in the 0-300 m layer (Q_{300}), as a sub-seasonal precursor of sea ice variability by means of lag correlations. Our results reveal that in the Pacific Arctic regions, sea ice variability in autumn is linked with Q_{ml} anomalies leading by 1 to 3 months, and this relationship has strengthened in the Laptev and East Siberian seas during 2001-2021 relative to 1980-2000, primarily due to reduced surface heat loss since the mid-2000s. Q_{300} anomalies act as a precursor for wintertime sea ice variability in the Barents and Kara seas, with considerable strengthening and expansion of this link from 1980-2000 and 2001-2021 in both reanalyses. Our results highlight the role played by upper ocean heat content in modulating the interannual variability of Arctic sea ice at sub-seasonal timescales. Heat stored in the ocean has important implications for the predictability of sea ice, calling for improvements in forecast initialization and focus upon regional predictions in the Arctic region.

1 Introduction

Satellite observations have documented a rapid decline of Arctic sea ice in all seasons (Simmonds, 2015; Stroeve and Notz, 2018; Onarheim et al., 2018). The mean sea ice state is transitioning to a new regime of thinner, more fractured and mobile ice (e.g., Graham et al., 2019; Sumata et al., 2023). This results in increased vulnerability to dynamic and thermodynamic forcing mechanisms, including pulses of ocean heat from the lower latitudes (Holland et al., 2006), atmospheric temperature fluctuations (Olonscheck et al., 2019) and enhanced ocean mixing and ocean-ice heat fluxes (e.g., Duarte et al., 2020; Ricker

et al., 2021). The interplay between the atmosphere, ocean and sea ice is central to explaining the pronounced temporal and spatial variability of the Arctic ice cover observed at a range of time scales (e.g., England et al., 2019; Ding et al., 2019). These complex interactions pose a challenge for accurate sea ice predictions, particularly at the regional level.

25 Sea ice variability in the Arctic is strongly region-dependent (e.g., Onarheim et al., 2018). The Atlantic sector (including the Greenland, Barents, Kara and Laptev seas) and in particular the Barents Sea have experienced larger sea ice loss than other areas of the Arctic Ocean (Onarheim et al., 2014, 2018; Lind et al., 2018), due to enhanced heat advection by the warm and saline Atlantic Water (AW) entering the basin at intermediate depths (200-800m, Aagaard 1989). Heat from the Atlantic Ocean reaches the Arctic through the Barents Sea Opening (Smedsrud et al., 2010) and Fram Strait (Schauer and Beszczynska-Möller, 30 2009). The combined effects of increased volume inflow (Smedsrud et al., 2022) and warmer temperatures (Wang et al., 2019) of the AW current have been linked to sea ice decline in the Barents Sea (e.g., Årthun et al., 2012; Smedsrud et al., 2013) and more recently in the eastern Eurasian Basin (Polyakov et al., 2020b). In the latter, the interaction between AW and sea ice has also been examined in relation to the weakening of the stratified, cold halocline layer and consequent increase in vertical heat fluxes towards the surface (Polyakov et al., 2010, 2017). These changes are collectively referred to as the ‘Atlantification’ of 35 the Arctic Ocean (Årthun et al., 2012; Asbjørnsen et al., 2020) and are thought to be the primary cause of thermodynamic ice melt during winter, when incoming solar radiation is absent (Onarheim et al., 2014; Tsubouchi et al., 2021; Ivanov et al., 2012). The expansion of the Barents Sea open water region leads to enhanced heat release from the ocean to the colder atmosphere, a phenomenon known as the “Barents Sea cooling machine” (Skagseth et al., 2020). In this part of the Arctic, complex feedbacks exist between sea ice retreat, anomalous atmospheric circulation, and induced changes in ocean heat transport by wind-driven 40 currents (e.g., Mohamed et al., 2022). Observations over the past 20 years have shown reduced heat loss in ice-free areas of the Barents Sea, which exceeds the increase driven by the sea-ice retreat (Skagseth et al., 2020).

The Pacific sector of the Arctic (including the Beaufort, Chukchi and East Siberian seas) has also been the epicenter of some of the most remarkable episodes of sea ice loss in recent years (Comiso et al., 2017). While small in comparison to the inflow through Fram Strait (about 10 times smaller in volume and with a heat flux that is 1/3 of the Fram Strait heat flux; Woodgate 45 et al. 2012), the inflow of Pacific Water (PW) through Bering Strait has also shown a warming trend that can accelerate sea ice loss (MacKinnon et al., 2021; A Woodgate and Peralta-Ferriz, 2021) and act as a trigger for early melt onset (Woodgate et al., 2010).

Enhanced poleward heat transport from the lower latitudes or reduced ocean heat loss contribute to altering the Arctic’s energy budget, causing the amplification of upper ocean warming (Shu et al., 2022; Asbjørnsen et al., 2020). Mayer et al. (2019) 50 estimated an energy imbalance of the Arctic Ocean of the order of 1 Wm^{-2} between 2001–17, with 1/3 of the accumulated heat going to the sea ice. This finding is corroborated by remarkable upward trends in upper ocean heat content over the past four decades, particularly during summer and autumn (Li et al., 2022). The temperature increase in the Arctic Ocean is at the core of the ice-albedo feedback (Perovich et al., 2007; Perovich and Polashenski, 2012). As more areas of open ocean become exposed to solar radiation due to sea ice break-up and early retreat, the seasonal upper ocean heat uptake increases and a surplus 55 of energy becomes stored in the surface layers (Stammerjohn et al., 2012; Serreze and Meier, 2019), where it is immediately available to the ice through lateral and bottom melting (e.g., Carmack et al., 2015).

To date, studies addressing oceanic drivers of sea ice variability have mainly focused on quantifying the effect of northward heat transport from the Pacific and Atlantic Oceans (e.g., Onarheim et al., 2015; Nummelin et al., 2017; Dörr et al., 2021; Aylmer et al., 2022). This generally entails a focus on interannual timescales of variability, owing to the delayed response of sea ice to heat anomalies that originate from lower latitudes (e.g., Woodgate et al., 2010). In comparison, the role of ocean heat content as a precursor of sea ice anomalies at sub-seasonal timescales has not been sufficiently explored. An improved characterization of the co-variability of sea ice and upper ocean heat content is of practical value to stakeholder groups, who are primarily interested in short-term predictability. Moreover, the accelerating trends towards a thinner sea ice cover implicate a higher vulnerability to regional scale forcing (Perovich and Polashenski, 2012).

Additionally, the role of the Arctic surface mixed layer, which represents the link between the ocean, sea ice and atmosphere (Toole et al., 2010), has received little attention in the literature, despite recent assessments of changes in the thermal state of the upper ocean have documented warming trends, especially in regions of maximum sea ice retreat (e.g., Polyakov et al., 2020a). While the relative amounts of upper ocean heat that are lost to the atmosphere and to the overlying sea ice remain hard to quantify, there is evidence that a surplus of energy absorbed by the mixed layer in summer can act to delay autumn ice growth (Perovich et al., 2007; Steele et al., 2008; Ivanov et al., 2016). However, our understanding of the effects of mixed layer heat on sea ice variability is still limited, partly due to the scarcity of sustained subsurface observations (e.g., Yang, 2006).

In this work, we address these gaps by assessing the role of upper ocean heat content on regional sea ice variability at sub-seasonal time scales, placing our results in the context of recent changes to the Arctic Ocean heat budget. We achieve this by using two eddy-permitting global ocean reanalyses, CMCC C-GLORSv5 and ECMWF ORAS5. Ocean reanalyses provide uniformly gridded reconstructions of the state of the ocean climate by combining ocean models driven by atmospheric forcing and available observational data via advanced data assimilation methods (Balmaseda et al., 2015). While they have become an established tool for a variety of climate services and science-driven studies (Storto et al., 2019), ocean reanalyses are especially valuable for the assessment and monitoring of the state of the Arctic Ocean, where hydrographic observations are severely limited by perennial ice cover (e.g., Uotila et al., 2019). Furthermore, the performance of different reanalysis products in polar regions has been widely evaluated with encouraging results (e.g., Iovino et al., 2022; Lien et al., 2016; Ilıcak et al., 2016). With respect to fully modeled data, errors and uncertainties in the representation of ocean variables in reanalyses are effectively reduced through the assimilation of observational fields. The CMCC C-GLORSv5 and ECMWF ORAS5 products specifically have also been employed in several applications (e.g. Takahashi et al., 2021; Carton et al., 2019) including in polar regions (e.g., Mayer et al., 2019; Shu et al., 2021; Nie et al., 2022). Their extensive temporal coverage (1980-present) and their high horizontal and vertical resolution, which allows for a high level of detail in the surface layers, make these products particularly suited for the study of the polar upper ocean. However, very few studies using ocean reanalyses have investigated changes in pan-Arctic heat budget and ocean heat content and, to our knowledge, no previous study has explored these aspects in relation to sea ice variability with a regional focus.

The remainder of this article is structured as follows: Section 2 introduces the data and methodological approach of the study. Section 3 presents the results, which are further divided into three parts: Section 3.1, which provides an overview of the dominant changes in the Arctic Ocean between 1980-2021; Section 3.2, where we perform a heat budget analysis to ensure the

accuracy of our results; and Section 3.3, where we examine the coupling between ocean heat content and sea ice variability in the past 40 years. In Section 4, we discuss our findings and their implications for the future climate.

2 Data and Methods

95 2.1 Sea ice and ocean reanalyses

Monthly means of sea ice concentrations between 1980 and 2021 are derived from the fifth generation atmospheric global reanalysis of the European Center for Medium-Range Weather Forecasts (ECMWF ERA5, Hersbach et al. 2020). ERA5 data are available from the Copernicus Climate Change Service (C3S) Data Store (Thépaut et al., 2018) on a regular latitude-longitude grid with $0.25^\circ \times 0.25^\circ$ horizontal resolution, from 1950 to present day. Sea ice concentration in the ERA5 data set after 1979 relies on the EUMETSAT Ocean and Sea Ice Application Facilities (OSI-SAF) Climate data Record version 1.2 (OSI-409a) and operational OSI-SAF (OSI-430) products (1979-2007 and 2007-present, respectively), which are based on combined passive microwave observations from the Defense Meteorological Satellite Program (DMSP) Special Sensor Microwave/Imager (SSM/I) and the Special Sensor Microwave Imager/Sounder (SSMIS).

Table 1. Overview of Ocean Reanalyses used in this study

Product Name	Model / DA Scheme	Resolution (horizontal, vertical)	Atmospheric forcing	Observations (SST and SIC)	Reference
C-GLORSv5	NEMO 3.2-LIM2 / OceanVAR	$0.25^\circ \times 0.25^\circ$, 75L	ERA-Interim	NOAA and NSIDC DMSP ²	Storto and Masina (2016)
ORAS5	NEMO 3.4.1-LIM2 / 3D-Var FGAT	$0.25^\circ \times 0.25^\circ$, 75L	ERA-Interim/ ECMWF OPS ¹	HadISSTv2 and OSTIA ³	Zuo et al. (2019)

¹ up to 2015/ After 2015

² SST is assimilated from the NOAA SST 1/4° analyses (Reynolds et al., 2007). With respect to sea ice data assimilation, C-GLORSv5 introduces a nudging scheme to weakly constrain sea ice thickness in the Arctic from PIOMAS.

³ SST is assimilated from HadISSTv2 pentad before 2008 and from OSTIA analysis from 2008. SIC data come from the OSTIA reprocessed analysis before 2008 and OSTIA analysis after.

We derive temperature, mixed layer depth (MLD), surface heat fluxes and velocity fields at monthly resolution between 1980-2021 from the Global Ocean Reanalysis System version 5 from the Euro-Mediterranean Center on Climate Change (CMCC)




Figure 1. Arctic Ocean bathymetry in C-GLORSv5. Dashed lines indicate the boundaries of the regional domains considered in this study (Barents-Kara, BK; Laptev-East Siberian, LES; Chukchi Sea, CHU and Beaufort Sea, BEA). Solid lines mark the location of the main Arctic gateways: Fram Strait (blue), Barents Sea Opening (BSO; red), Bering Strait (magenta) and Davis Strait (green). Note the irregular depth intervals in the colorbar.

C-GLORSv5; Storto and Masina 2016) and the Ocean ReAnalysis System 5 from the European Center for Medium-Range Weather Forecasts (ECMWF ORAS5; Zuo et al. 2019), which is available for download on the C3S Data Store. An overview of the key product characteristics is provided in Table 1. Both C-GLORSv5 and ORAS5 are produced with the NEMO ocean model (Madec et al., 2017) coupled to the Louvain-la-Neuve Sea ice Model (LIM2, Fichefet and Maqueda 1997). Both models
 110 adopt a tripolar grid at eddy-permitting configuration with $0.25^\circ \times 0.25^\circ$ horizontal resolution. The original vertical resolution is 50 and 75 vertical levels for C-GLORSv5 and ORAS5, respectively, however we adopt a version of C-GLORSv5 with 75 vertical levels (C-GLORSv5e025L75, herein referred to as simply C-GLORSv5). In both reanalyses, the ocean is forced by the ECMWF ERA-Interim atmospheric reanalysis (Dee et al., 2011); in ORAS5, this is replaced by ECMWF operational NWP after 2015. The two products differ in their assimilation schemes in terms of input observational datasets, frequency of analysis,
 115 assimilation time-windows and bias correction schemes. A full description of the assimilated satellite and in-situ observations as well as the specifics of data assimilation schemes can be found in the reference papers listed in Table 1.

2.2 Regional and pan-Arctic heat budget

We first estimate the ocean heat budget in the C-GLORSv5 and ORAS5 reanalyses for the whole Arctic and for four sub-domains: the Barents-Kara region (BK); the Laptev-East Siberian region (LES); the Chukchi Sea (CHU) and the Beaufort Sea

120 (BEA, Fig. 1). These regional domains are largely consistent with those of previous studies (e.g., Stroeve et al., 2016; Bliss et al., 2019; Lenetsky et al., 2021).

Considering a control ocean volume with surface A and vertical section S , where mass and salinity are conserved, the heat budget is given by the balance between advective, vertical and diffusive heat flux terms (1):

$$\underbrace{\frac{\partial Q}{\partial t}}_{Qt} = \underbrace{\rho_0 C_p \int_S VT dS}_{OHT} + \underbrace{\int_A Q_s dA}_{SSHf} + Q_{diff} \quad (1)$$

125 where Qt is the ocean heat content tendency; OHT represents the advective ocean heat transport through S , the reference density ρ_0 (1026 kg m^{-3}) and specific heat capacity C_p ($3996 \text{ J kg}^{-1} \text{ K}^{-1}$) of sea water are both assumed constant, V and T are the cross-sectional velocity and potential temperature, respectively; SSHF (positive downward) indicates the net sea surface heat flux, Q_s , over the surface A ; Q_{diff} represents the lateral heat diffusion.

We define the pan-Arctic domain as the area enclosed by the four ocean gateways: the Fram, Bering, and Davis straits and
 130 the Barents Sea Opening (BSO) (Fig. 1), as in previous observational (e.g., Tsubouchi et al., 2012) and modeling studies (e.g., Lique and Steele, 2013). The net ocean heat transport into the Arctic is computed along each section (on the native grid) at monthly frequency for C-GLORSv5 and ORAS5 (2).

$$OHT = \rho_0 C_p \int_{-Z(\lambda)}^{\eta} \int_{\lambda_1}^{\lambda_2} V(T - T_{ref}) d\lambda dz \quad (2)$$

135 Where λ_1 and λ_2 are the coordinates of the section line, $Z(\lambda)$ is the depth at each section, and T is taken relative to a reference temperature T_{ref} of 0°C , following previous studies (Årthun et al., 2012; Lique and Steele, 2013). It is widely recognised that heat transports are sensitive to the choice of reference temperatures. Therefore, the reader should note that when considering partial sections, the estimated values of OHT are always dependent on the arbitrary choice of T_{ref} , since there is a non-zero net volume flux (i.e., the volume transport across individual sections is not balanced; see Schauer and Beszczynska-Möller 2009).

140 We consider the net sea surface heat flux term (SSHf, Wm^{-2}) over the box area enclosed by the four gates. SSHf is defined positive downward in both reanalyses and represents the exchange of solar and non-solar fluxes between the ocean and atmosphere. Note that in considering the heat budget components in (1) we omit the lateral heat diffusion term as this is negligible compared to the advective and surface flux terms (Lique and Steele, 2013). For a detailed analysis of the coupled ocean-ice-atmosphere surface energy budget of the Arctic Ocean, which is beyond the scope of this study, the reader can refer
 145 to Mayer et al. (2019). We further compute the same heat budget terms for each individual Arctic sub-region by considering the total advected heat as the sum of net OHT along each section line bounding the given region (solid coloured lines and white dashed lines in Fig. 1). Given the full seasonal sea ice coverage (SIC > 90%) in the Laptev-East Siberian and Beaufort regions, we consider the net surface heat flux in each region as the sum of the ocean-ice heat flux (OIHF, Wm^{-2}) for the ice-covered

fraction and SSHF (Wm^{-2}) for the remaining ice-free fraction of each grid cell. The regional heat budget analysis is computed from C-GLORSv5 data only, as the ocean-ice heat flux is not available for download from the ORAS5 reanalysis.

2.3 Ocean heat content

We next estimate ocean heat content per unit area (Q , Jm^{-2}) by vertically integrating monthly potential temperature over two target depths: from the surface to the seasonally-varying mixed layer depth (Q_{ml}), to account for the energy stored at the atmosphere-ice-ocean interface, and the 0-300 m depth layer (Q_{300}), to capture the intermediate depth layer characterized by the greatest AW warming (e.g., Shu et al., 2022) (3).

$$Q = \rho_0 C_p \int_{-Z}^{\eta} (T - T_{freeze}) dz \quad (3)$$

where T_{freeze} is the freezing temperature of seawater = -1.8°C and Z represents the maximum depth boundary. Here, we adopt the constant of freezing as reference temperature following previous studies (e.g., Mayer et al., 2019; Korhonen et al., 2013) to obtain the heat available for potential sea ice melt. In calculating Q_{ml} , Z corresponds to the time-varying MLD, which is defined with the threshold method where potential density exceeds the surface reference value by 0.01 kg m^{-3} (Peralta-Ferriz and Woodgate, 2015). The density difference criterion and the 0.01 kg m^{-3} threshold have been widely tested in the Arctic (e.g., Toole et al., 2010; Gimbert et al., 2012; Timmermans et al., 2012; Thomson and Fine, 2003). We do not find any substantial differences in the spatial distribution and seasonality of MLD when adopting a different density criterion (e.g., 0.03 kg m^{-3} ; not shown). In calculating Q_{300} , we consider Z to be fixed at 300 m, or the ocean bottom if the water column is shallower than 300 m.

Finally, we obtain monthly anomaly time series of SIC, Q_{ml} and Q_{300} by removing a monthly climatological mean from each time series. To investigate how upper ocean heat content at both target depths influences sea ice variability on sub-seasonal time scale, lagged correlations between region-averaged anomalies of Q and SIC are computed for each month, with the ocean leading sea ice by a lead time of one to three months. The time series used for the lagged correlation analysis are split between an earlier period (1980-2000) and a later period (2001-2021) to assess the role of upper ocean warming and sea ice loss in recent decades. For comparison, we also show the lagged auto-correlation of SIC anomalies at each month as an indicator of the inherent sea ice predictability and how this varies across the two periods.

3 Results

3.1 Changes in the Arctic upper ocean between 1980-2021

Figure 2 shows annual mean trends in Arctic Ocean SIC, MLD, Q_{ml} , Q_{300} , SSHF and OIHF between 1980 and 2021 for the C-GLORSv5 reanalysis. Given the strong seasonality of SSHF, trends for this variable are shown separately for the seasons of melt-onset (MJJ) and freeze-onset (OND). Warming of the upper ocean is most evident in the Barents and Kara regions, both




Figure 2. Annual trends (1980-2021) in (a) sea ice concentration (SIC, %), (b) mixed layer depth (MLD, m) (c) ocean heat content in the mixed layer and in (d) the 0-300 m layer (Q_{ml} and Q_{300} , 10^8 Jm^{-2}), (e) sea surface heat flux (SSHF, in May-June-July; MJJ, Wm^{-2}) and in (f) October-November-December; OND, and (g) ocean-ice heat flux (OIHF, Wm^{-2}) for the C-GLORSv5 ocean reanalysis. SSHF is defined as positive downward. Stippling denotes 99% significance.

in terms of SIC reductions and positive trends in Q_{ml} and Q_{300} , especially along the pathways of AW inflow (Fig. 2 a,c,d). Increased springtime SSHF (Fig. 2e) is consistent with enhanced solar absorption in regions of reduced ice cover. During the
180 autumn season (OND), the positive SSHF trend in the southern Barents Sea suggests reduced ocean cooling in this part of the region (Fig. 2f). Conversely, ocean heat loss to the atmosphere during autumn has been intensifying in the Chukchi Sea and in large parts of the the northern Barents and Kara seas, particularly along the pathway of Fram Strait inflow. These changes suggest that the area of effective cooling in the Barents Sea has expanded towards the northern Barents and Kara seas (Shu et al., 2021). Significant reductions in OIHF are consistent with regions of sea ice loss along coastlines and in the southwest
185 Barents Sea (Fig. 2g). The overall weakening in stratification in the Atlantic Arctic sector (Fig. 2b) is opposed to a modest decrease in MLD in the Canada basin, as corroborated by recent literature (e.g., Muilwijk et al., 2022).




Figure 3. Time series of SIC (blue), Q_{300} (black) and Q_{ml} (grey) in C-GLORSv5 and ORAS5 for the Arctic regional seas and for the total Arctic. The pan-Arctic region (e) is defined as the area bounded by the Fram, BSO, Bering and Davis straits. Note the different y-axis ranges.

Figure 3 illustrates annual time series of Q_{ml} , Q_{300} and SIC for the four Arctic regional seas and for the pan-Arctic region. The interannual variability of SIC is intrinsically linked with that of Q . Reductions in SIC and an increase in Q_{300} between 1980-2021 are evident for all regions as well as for the pan-Arctic mean (Fig. 3e), though trends are most pronounced after the early/mid-2000s. Among all sub-regions, the Beaufort Sea presents the weakest decline in SIC, presumably due to stronger upper ocean stratification. The Barents-Kara region experienced the largest increase in Q_{300} ($0.11 / 0.15 \text{ } 10^8 \text{ J m}^{-2} \text{ y}^{-1}$ for C-GLORSv5 / ORAS5, respectively) and Q_{ml} ($0.08 / 0.06 \text{ } 10^8 \text{ J m}^{-2} \text{ y}^{-1}$). Both the trend and variability of Q_{300} and Q_{ml} are comparatively small in the Laptev-East Siberian and Beaufort seas, as these regions are fully ice-covered for most of the year with limited surface mixing. It is worth mentioning the close agreement between the two reanalyses, with exception for a minor discrepancy in the Laptev-East Siberian Q_{300} , which is greater in C-GLORSv5 than ORAS5.

3.2 Assessment of budget closure

The net annual heat transport through the main Arctic gates is shown in Fig. 4 for C-GLORSv5 and ORAS5, and complemented in Table 2. The largest contribution to the total ocean heat transport into the Arctic comes from the BSO ($62\text{-}99 \text{ TW}$ C-GLORSv5; $57\text{-}96 \text{ TW}$ ORAS5, values referring to annual means), followed by Fram Strait ($14\text{-}47 \text{ TW}$ C-GLORSv5; $19\text{-}38 \text{ TW}$ ORAS5), Davis Strait ($8\text{-}16 \text{ TW}$ C-GLORSv5; $10\text{-}19 \text{ TW}$ ORAS5) and Bering Strait ($4\text{-}12 \text{ TW}$ C-GLORSv5; $5\text{-}14 \text{ TW}$ ORAS5). Transports along all sections show an upward trend for both reanalyses except for Davis Strait, where the net transport




Figure 4. Annual time series of (a) ocean heat transport (OHT, TW) via the four main Arctic gates and their sum (red lines) and (b) pan-Arctic heat budget for C-GLORSv5 (solid) and ORAS5 (dotted). The locations of the Arctic gates are marked in Figure 1. Surface heat flux (SSHF) is weighted on the area bounded by the four gates and the negative sign indicates ocean heat loss. Black lines in panel (b) denote the sum of total ocean heat transport and surface heat flux.

has been decreasing (-0.29 GW y^{-1} in C-GLORSv5 and -0.24 GW y^{-1} in ORAS5). While there is a satisfactory agreement between the two reanalyses, C-GLORSv5 generally simulates smaller net transports via the Fram and Davis straits and a larger net transport via the BSO relative to ORAS5. As a result, the total Arctic OHT is slightly lower in C-GLORSv5 than in ORAS5 (99-150 TW and 103-155 TW, respectively).

We find OHT in C-GLORSv5 and ORAS5 to be generally consistent with the existing literature, though a direct comparison with previous studies, especially observational ones, is challenging given the sensitivity of heat transport estimates to AW temperatures, temporal coverage, and chosen reference temperature values (Table 2). Our 1980-2000 estimates for Fram Strait and Davis Strait are in close agreement with the 1948-2007 multimodel mean of 22.9 TW and 13.4 TW, respectively, in CORE-II models (Ilıcak et al., 2016) and with NorESM model estimates by Muilwijk et al. (2018) (both studies adopt $T_{ref} = 0^\circ\text{C}$). The BSO OHT during 1980-2000 (70.1 and 77.4 TW in ORAS5 and C-GLORSv5, respectively) is larger than the 1948-2007 multimodel mean of 55.96 TW (Ilıcak et al., 2016) and the observed estimate of 64.5 TW for 2005-2009 (Tsubouchi et al., 2019) referenced to 0°C . Albeit referenced to a different temperature ($T_{ref} = -0.1^\circ\text{C}$), we find close agreement with Maslowski et al. (2004), who reported 74 TW through the BSO in the same period (1979-2001).

Between 1980-2021, the total net OHT into the Arctic Ocean shows a positive trend in both reanalyses (1.21 GW y^{-1} in C-GLORSv5 and 1.92 GW y^{-1} in ORAS5). The increase in heat transport can also be appreciated by comparing monthly OHT estimates via each gateway averaged over the two halves of the timeseries (1980-2000 and 2001-2021; Table 2). The upward

Table 2. Average Net Ocean Heat Transport (referenced to $T_{ref} = 0^{\circ}\text{C}$) via Arctic Gates [TW] for the C-GLORSv5 and ORAS5 reanalyses and for existing observational and model estimates¹

Source	Section/Gate	Period	Full depth	Mixed layer	0-300 m
C-GLORSv5	Fram Strait	1980-2000	20.7	9.1	12.2
	BSO	1980-2000	77.4	37	72.9
	Bering Strait	1980-2000	7.1	3.1	/
	Davis Strait	1980-2000	13.4	3	/
C-GLORSv5	Fram Strait	2001-2021	25.3	10.7	15.4
	BSO	2001-2021	83.2	39.1	78.2
	Bering Strait	2001-2021	8	3.5	/
	Davis Strait	2001-2021	10.9	2.6	/
ORAS5	Fram Strait	1980-2000	27.6	14.5	16
	BSO	1980-2000	70.1	34.5	65.4
	Bering Strait	1980-2000	8.5	3.9	/
	Davis Strait	1980-2000	17.1	3.7	/
ORAS5	Fram Strait	2001-2021	30	16.8	18.6
	BSO	2001-2021	83.1	35.7	78.2
	Bering Strait	2001-2021	9.4	4.2	/
	Davis Strait	2001-2021	15.1	3.2	/
Observations					
Tsubouchi et al. 2019	Fram Strait	2005-2009 ²	61.2	/	/
	BSO	2005-2009	64.5	/	/
	Bering Strait	2005-2009	2.14	/	/
	Davis Strait	2005-2009	23.67	/	/
Model Simulations					
Mulwijk et al. 2018	Fram Strait	1890-2009	10-30	/	/
	BSO	1890-2009	30-60	/	/
	Bering Strait	1890-2009	0-5	/	/
	Davis Strait	1890-2009	5-15	/	/

¹ Averages for C-GLORSv5 and ORAS5 are computed from monthly net heat transports. OHT for the 0-300 m layer is marked as '/' if the section depth is shallower than 300 m and values therefore correspond to the full depth OHT.

² Values refer to Jan 2005 - Dec 2009 averages based on mooring data (details in Tsubouchi et al. 2018).

trend in OHT is offset by an overall decline in surface heat fluxes averaged over the pan-Arctic region (SSHF; -2.3 GW y^{-1} in C-GLORSv5; -1.95 GW y^{-1} in ORAS5), which is associated with enhanced ocean heat release to the atmosphere owing to sea ice loss (Smedsrud et al., 2022). With the opposing trends of OHT and SSHF, the surface heat budget is closed around zero over the historical time series (Fig. 4b), with good agreement between the two reanalyses.

Annual time series of regional heat budgets (Fig. 5) further demonstrate the realistic representation of ocean heat advection and surface heat fluxes in the C-GLORSv5 reanalysis. Net heat transport into the Barents-Kara region has been increasing at a rate of 0.93 GW y^{-1} , chiefly due to the upward trend in OHT through the BSO. The increase in OHT to this region is offset by a weak decline in surface heat fluxes (SSHF+OIHF, -0.2 GW y^{-1}), which is however non-significant and levels off after 2010, indicating a reduction in the overall cooling efficiency of the Barents-Kara region relative to the previous decades. Similarly, in all other regions, the sum of advective and surface heat flux terms after the late 2000s marks a shift towards warmer conditions due to reduced ocean heat loss. We find this to be largely driven by a sharp drop in OIHF in all regions, concurrent with a decline in sea ice concentration (Fig. 5e-h).

230 3.3 Ocean heat content as a precursor of SIC variability

Building on the results of the heat budget analysis, we explore the role of upper ocean heat content as a precursor of SIC anomalies in the Arctic's regional seas and how this has changed between the first (1980-2000) and second half (2001-2021) of our time series. Figure 6 shows 1-month lag correlations between SIC anomalies at each month and ocean heat content anomalies at both target depths (Q_{300} and Q_{ml}), along with 1-month lag SIC anomaly auto-correlations, for the C-GLORSv5 reanalysis. The equivalent figure for ORAS5 is presented in Fig. A1 in the Appendix. We focus on the Barents-Kara and Laptev-East Siberian seas as these regions have undergone the most remarkable changes in the Q-SIC interaction between the two periods. Regional differences in the seasonal patterns of correlations are largely consistent with the seasonal cycle of sea ice in each region (dashed green lines in Fig. 6). In the Barents-Kara region, SIC variability is closely linked to Q_{300} and the seasonal peak of correlations occurs in the freezing season (Jan-Feb), consistent with the timing of strongest AW inflow towards the Arctic (e.g., Beszczynska-Möller et al., 2012).

During the second half of the time series (2001-2021, Fig. 6e-h), changes in ocean warming and melt season duration are reflected on SIC predictability. The 1-month lagged Q-SIC correlation is increased, particularly in the autumn (note the shift in the maximum Q_{ml} -SIC correlation from August to October-November in the Laptev-East Siberian region). Relative to 1980-2000, the influence of Q_{300} on SIC variability in the Barents-Kara region during winter (Nov-Jan) 2001-2021 is considerably more negative: for instance, the correlation of January SIC to December Q_{300} increased from $r=-0.72$ in 1980-2000 to $r=-0.83$ in 2001-2021 in C-GLORSv5 (from $r=-0.73$ to $r=-0.86$ in ORAS5), while the December to January SIC auto-correlation is not significant in either period. Changes in the intensity and seasonality of the Q-SIC coupling between the two periods are also noticeable in the Chukchi Sea, where the correlation between SIC and Q anomalies at both target depths appears intensified in the autumn months (Oct-Dec) during 2001-2021 (not shown). No substantial changes are observed in the Beaufort Sea (not shown). Though with considerable regional differences, correlations between sea ice and ocean heat content anomalies are also significant at 2 months lag (Fig. A2 and A3, which show 2-month lagged Q-SIC correlations for C-GLORSv5 and




Figure 5. Time series of regional ocean heat budgets in the C-GLORSv5 reanalysis. **(a-d)** Annual time series of ocean heat transport, surface ocean heat fluxes and their sum. Ocean heat transport into a given region is defined as the sum of the net ocean heat transport along each section bounding the region, as marked in Figure 1. Surface heat fluxes (navy blue lines in **a-d**) are defined as the sum of net sea surface heat flux (SSHF) for the ice-free fraction of the grid cell and the ocean-ice heat flux (OIHF) for the ice-covered fraction of the grid cell. **(e-h)** shows the individual contribution of SSHF and OIHF in each region. Time series of sea ice concentration are shown in light blue. Note that here the sign of OIHF, which is positive from the ocean to sea ice, is reversed to match SSHF, indicating ocean heat loss.

ORAS5, respectively) and at 3 months lag (Fig. 9 and A4, which show 3-month lagged Q-SIC correlations for C-GLORSv5 and ORAS5, respectively).

Overall, the 1-month lagged SIC auto-correlation also increased in the second half of the time series, particularly in regions that have experienced the largest sea ice loss, i.e., the Barents-Kara region and Chukchi Sea (not shown). In the remainder of this section, we focus on the months where the 1-month lagged Q-SIC correlation is greater than the SIC auto-correlation, and thus where the predictive potential of ocean heat content is highest.

3.3.1 Q_{ml} and autumn sea ice in the Pacific Arctic

The regional seas of the Pacific Arctic (East Siberian, Chukchi and Beaufort seas) are characterized by complete or near-complete winter freeze-up and are considered to be areas of greater vulnerability to warming and lengthening of the melt




Figure 6. Lag 1 month correlations between SIC anomalies and ocean heat content anomalies in the 0-300 m layer and in the mixed layer, in the Barents-Kara and Laptev-East Siberian regions for the 1980-2000 (a-b) and 2001-2021 (e-f) periods, with the ocean leading sea ice. The 1-month lag SIC anomaly auto-correlation is shown in the bottom panels for comparison (c-d, g-h). Green dashed lines indicate the SIC seasonal cycle in each region and period. Note that the sign of SIC auto-correlation is inverted. All values shown in color are significant at the 95% level; empty boxes denote non-significant correlations. For brevity, only C-GLORSv5 is shown. The equivalent figure for ORAS5 is provided in the Appendix.

season (Peng and Meier, 2018; Serreze et al., 2016). In these regions, the Q_{ml} -SIC anticorrelation maximizes in autumn (sea ice advance, Fig. 7) at 1-month lag time, and correlations are generally significant up to 3 months lag (not shown).

Figure 7 shows spatial correlation maps of regionally-averaged Q_{ml} in October and the grid-point SIC anomaly in November in both reanalyses for 1980-2000 (Fig. 7 a,b) and 2001-2021 (Fig. 7 c,d). In all three regions, the anticorrelation between

265 October Q_{ml} and November SIC is higher than the October to November SIC auto-correlation (not shown). While this is true
for both periods, there is a noticeable increase in the strength of the correlation in the Laptev-East Siberian region during
2001-2021, consistent with the shift in the timing of maximum ice-ocean coupling from summer to autumn (Figure 6 b-
f). Interestingly, the spatial pattern of correlation in the northeastern Beaufort Sea indicates a reduction of the area of SIC
prediction skill associated with Q_{ml} during 2001-2021 (Fig. 7c,d). This area corresponds to the region of MLD shoaling
270 during 1980-2021 (Fig. 2b).

The negative co-variability of Q and SIC anomalies in the shelf seas of the Pacific Arctic at the time of ice formation is
consistent with the influence of the relatively warm summer Pacific water inflow through the shallow (~ 50 m) and narrow (\sim
85 km) Bering Strait (Rudels, 2015; Koenigk and Brodeau, 2014). However, we find greater difference in the predictability
of November SIC between the first and second period in the Laptev-East Siberian region (from $r=-0.52$ to $r=-0.86$ in C-
275 GLORSv5; from $r=-0.51$ to $r=-0.78$ in ORAS5) than in regions of greater PW influence, i.e., the Chukchi Sea (no change in
C-GLORSv5; from $r=-0.87$ to $r=-0.88$ in ORAS5). This is consistent with the statistically non-significant OHT trend in the
Chukchi Sea (Fig. 5c) and the weak increase in Bering Strait inflow from 1980-2000 to 2001-2021 (Table 2). Furthermore, the
time evolution of heat budget components in the Laptev-East Siberian region would suggest that the shift to warmer conditions
after the mid-2000s is due to a decrease in ocean-ice fluxes associated with sea ice loss rather than an increase in OHT (Fig.
280 5f).

3.3.2 Q_{300} and winter sea ice in the Barents-Kara region

In the Barents-Kara region, SIC and Q exhibit the highest negative correlation in the winter months, in agreement with existing
evidence of the link between AW inflow and winter sea ice variability (Årthun et al., 2019). December Q_{ml} (not shown) and
 Q_{300} anomalies are both strongly anticorrelated with January SIC anomalies, although the correlation is higher for Q_{300} (Fig.
285 8). An intensification of the Q_{300} -SIC link emerges in the 2001-2021 period (Fig. 8c,d), with correlations extending further
east into the Kara Sea and over large part of the marginal ice zone in proximity of the St. Anna Trough. During 2001-2021, the
southern Barents Sea is fully ice-free as the sea ice edge retreats north of 80°N and Atlantic water entering through the BSO is
advected a longer distance before encountering sea ice. This change is consistent with the notion that a warming trend in AW
inflow leads to less sea ice formation in the cold season (e.g., Long and Perrie, 2017), as previously demonstrated by the link
290 between BSO heat transport and sea ice area ($r=-0.76$, Shu et al. 2021; $r=-0.8$, Li et al. 2017).

The predictability of regionally-averaged winter (Nov-Jan) SIC associated with Q_{300} anomalies in the Barents-Kara region
is maintained above $r=-0.8$ up to a lead time of 3 months in both reanalyses (Fig. 9a, A4a), highlighting the importance of
ocean heat content as a precursor of SIC variability in this region, especially in months with lower or non-significant SIC
auto-correlation.




Figure 7. Correlation between the region-averaged Q_{ml} in October and the grid-point SIC in November for the Laptev-East Siberian, Chukchi and Beaufort regions in 1980-2000 (a,b) and 2001-2021 (c,d) for C-GLORSv5 (a,c) and ORAS5 (b,d). Black and magenta contour lines indicate the November climatological pack ice extension (80% SIC) and sea ice edge (15% SIC), respectively, over each period.

295 4 Discussion and conclusions

The analysis presented in this study aims to shed light on the implications of upper ocean warming for regional sea ice variability. We used two eddy-permitting global ocean reanalyses, C-GLORSv5 and ORAS5, to investigate changes in ocean heat transport and surface heat fluxes through a heat budget analysis of the Arctic Ocean and its regional seas. Secondly, we assessed the role of Q anomalies in preconditioning sea ice variability by means of lag correlations. Results showed that while the total OHT into the Arctic has increased substantially between 1980-2021 (1.21 GW y^{-1} in C-GLORSv5 and 1.92 GW y^{-1} in ORAS5), only the Barents-Kara region is affected by a significant positive trend in ocean heat advection (0.93 GW y^{-1}), originating from the BSO. In this region, reduced ocean heat loss to the atmosphere after the mid-2000s also contributed to

300




Figure 8. As in Figure 7, but showing the correlation between the region-averaged Q_{300} in December and the grid-point SIC in January for the Barents-Kara region in 1980-2000 (a,b) and 2001-2021 (c,d) for C-GLORSv5 (a,c) and ORAS5 (b,d). Black and magenta contour lines indicate the January climatological pack ice extension (80% SIC) and sea ice edge (15% SIC), respectively, over each period.

the warming trend (Fig. 5a,e), in agreement with what found by Asbjørnsen et al. (2020). We have shown that Q anomalies in the Barents-Kara seas, particularly in the 0-300 m layer, act as an important precursor of wintertime sea ice variability on sub-




Figure 9. As in Figure 6, but for a lag time of 3 months, with the ocean leading sea ice.

305 seasonal timescales. Relative to 1980-2000, this link intensified and expanded northwards and eastward during 2001-2021 (Fig. 7c,d), with close agreement between the two reanalyses ($r[Q_{300DEC}, SIC_{JAN}] = -0.83$ in C-GLORSv5 and -0.86 in ORAS5).

In light of these findings, it becomes apparent that the evolution of sea ice variability in the Atlantic sector will depend on the opposing contribution of poleward heat advection and surface heat loss; in other words, whether the 'Barents Sea cooling machine' will gradually lose its efficiency (Skagseth et al., 2020) or rather expand it to accommodate changes in AW inflow.

310 For instance, CMIP6 simulations by Shu et al. (2021) showed that under the RCP8.5 scenario AW warming will increase and trends in winter surface heat loss and sea ice concentration will expand poleward, together with mixed layer deepening in the

northern Barents Sea and Kara Sea. According to this scenario, the Q_{300} -SIC coupling in the Kara Sea that emerged from our analysis between 2001-2021 ($r < -0.5$ in the western Kara Sea; Fig. 8) will likely intensify and expand northward and eastward. This will likely allow for more local heat loss (negative SSHF trend) and enhanced convection as larger areas of open ocean become exposed to heat exchange with the atmosphere. According to Smedsrud et al. (2022), more heat loss will in turn accommodate for larger poleward AW inflow. However, substantial uncertainty remains around future projections of Arctic OHT, partly due to biases in AW representation in coupled climate models (e.g., Heuzé et al., 2023). It remains unclear how changes in large-scale weather patterns, which this study did not address, may have contributed to the observed strengthening of the upper ocean-sea ice coupling in the Barents-Kara region. It was previously demonstrated (Sandø et al., 2010) that OHT through the BSO, non-solar heat fluxes (i.e., heat loss to the atmosphere) and ocean heat content in the Barents Sea are all correlated to the North Atlantic Oscillation (NAO). On decadal timescales, variations in the NAO can induce changes in the westerlies and, as a result, oceanic heat transports (Mulwijk et al., 2018). Mikhailova et al. (2021) showed that the correlation between the NAO and winter upper ocean temperatures in the Barents Sea weakened during 1995-2009, however there is currently limited data to infer how this link might have evolved during the last decade (2010-present).

In the Laptev-East Siberian, Chukchi and Beaufort seas, we found no evidence of OHT having a substantial effect on upper ocean warming (i.e., OHT trends between 1980-2021 are not significant). Regional heat budgets point to a larger contribution of SSHF and OIHF to the overall warming, especially since the mid-2000s (Fig. 5). The variability of SIC during the warmer months is strongly linked to that of Q at 1-3 lead months, particularly Q_{ml} . During 2001-2021, the timing of maximum correlation appears shifted from summer to autumn and the co-variability is heightened in the Laptev-East Siberian (notably, $r[Q_{mlOCT}, SIC_{NOV}]$ in this region increased from -0.52 to -0.86 in C-GLORSv5 and from -0.51 to -0.78 in ORAS5). This finding is consistent with recent results by Sumata et al. (2023), who highlight a regime shift after 2007 consisting in a drastic reduction in September SIC in the area of sea ice formation of the Laptev-East Siberian region, with the consequent triggering of a widespread ice-albedo feedback. The timing of the regime shift agrees with the sharp drop in SIC anomaly in 2007 in the Laptev-East Siberian region ($< -10\%$), concurrent with a shift to positive Q anomalies in both reanalyses (Fig. 3b). Because extra solar heating during summer is absorbed at a higher rate than it can be released, the thermodynamic response of the ice-ocean system to warming is the shift of the ice-free season into autumn (Lebrun et al., 2019). Hence, we conclude that enhanced summer absorption of atmospheric heat associated with sea ice loss and the resulting Q_{ml} anomaly contributed to strengthening and delaying the peak of Q_{ml} -SIC anticorrelation during 2001-2021. While it is possible that an increase in Bering Strait inflow and the expansion of Fram Strait influence along the Siberian shelf may have additionally contributed to the stronger coupling, we found no evidence that this is the case.

It is worth noting that despite the long temporal coverage of C-GLORSv5 and ORAS5, the time series analyzed in this study are insufficient to infer patterns of decadal and multi-decadal variability in OHT and SSHF. It has been suggested that the recent increase in AW inflow could be in part associated with multidecadal fluctuations (e.g., Smedsrud et al., 2013), as earlier studies showed a similar warming during the 1930s-1940s (ETCW), followed by a period of relative cooling (Polyakov et al., 2004, 2005; Mulwijk et al., 2018). Internal climate variability indeed remains a source of substantial uncertainty for the long-term evolution of sea ice.

350 Additionally, the limited availability of observational datasets to be used as a benchmark implies that caution should be taken when interpreting the variability and trends of reanalysis data, especially when it comes to ocean-ice heat fluxes, which here could not be compared against other sources. Given the current trends in upper ocean warming, further work is necessary to better quantify the processes underlying the sea ice response to ocean heat surplus, particularly in regions where sea ice is transitioning to a state of higher vulnerability, i.e. the Kara and Chukchi Sea (Bliss et al., 2019). For instance, changes in the strength of stratification and halocline stability, which this study did not address, remain a large source of uncertainty due to substantial regional variability and model spread (e.g., Muilwijk et al., 2022; Pan et al., 2023; Shu et al., 2022). Several physical mechanisms can influence the exchange of heat fluxes between the ocean, atmosphere and sea ice, including changes in wind forcing (Onarheim et al., 2015; Lien et al., 2017) and freshwater fluxes (Karami et al., 2021). For instance, there is evidence that wind-driven mixing will increase under ice-depleted conditions (Perovich, 2011), which in turn will accelerate sea ice loss (Liang and Losch, 2018). At the same time, the surplus of freshwater associated with sea ice decline is expected to promote stratification, thus leading to limited vertical mixing (Davis et al., 2016). Because of these opposing contributions, predicting changes in the Arctic upper ocean stratification remains a difficult task.

360 Despite the aforementioned uncertainties, our results provide clear evidence that recent changes to the Arctic Ocean's surface heat budget have induced a strengthening of upper ocean-sea ice interactions in the Arctic regional seas. The strong coupling between ocean heat content and sea ice anomalies that emerged from our analysis has important implications for the sub-seasonal predictability of sea ice, which is of practical value to local communities and stakeholder groups, including for the navigability of Arctic shipping routes such as the Northern Sea Route (e.g., D'Angelo et al., 2021). Given the ongoing transition of the Arctic system to thinner, younger and more mobile sea ice, it is reasonable to expect increased vulnerability to continued warming of the upper ocean. Among other applications, ocean reanalyses have been successfully employed for the initialization of sea ice and ocean components of seasonal retrospective forecasts (e.g., Johnson et al., 2019; McAdam et al., 2022). Hence, continued efforts in the representation of upper ocean variables through improved accuracy of ocean models, atmospheric forcing and data assimilation schemes, together with the current expansion of the observational network (e.g., Tsubouchi et al., 370 2012), are crucial to help us address open questions while ensuring consistent monitoring of the Arctic Ocean climate.

Data availability. All data analyzed in this study are freely available online. ECMWF ERA5 can be downloaded from the Copernicus Climate Change Service (C3S) Data Store. The CMCC C-GLORSv5 Reanalysis can be downloaded from the PANGAEA repository. The ECMWF ORAS5 Reanalysis is available from C3S. Both C-GLORSv5 and ORAS5 reanalyses can also be found on the Copernicus Marine Service website (CMEMS) as part of the Global Ocean Ensemble Physics Reanalysis (Product Identifier: GLOBAL REANALYSIS PHY 375 001 031).




Figure A1. Lag 1 month correlations between SIC anomalies and ocean heat content anomalies in the 0-300 m layer and in the mixed layer, in the Barents-Kara and Laptev-East Siberian regions for the 1980-2000 (a-b) and 2001-2021 (e-f) periods, with the ocean leading sea ice for the ORAS5 reanalysis. The 1-month lag SIC anomaly auto-correlation is shown in the bottom panels for comparison (c-d, g-h). Green dashed lines indicate the SIC seasonal cycle in each region and period. Note that the sign of SIC auto-correlation is inverted. All correlations are significant at the 95% level.




Figure A2. Lag 2 month correlations between SIC anomalies and ocean heat content anomalies in the 0-300 m layer and in the mixed layer, in the Barents-Kara and Laptev-East Siberian regions for the 1980-2000 (a-b) and 2001-2021 (e-f) periods, with the ocean leading sea ice for the C-GLORSv5 reanalysis. The 2-month lag SIC anomaly auto-correlation is shown in the bottom panels for comparison (c-d, g-h). Green dashed lines indicate the SIC seasonal cycle in each region and period. Note that the sign of SIC auto-correlation is inverted. All correlations are significant at the 95% level.




Figure A3. As in A1, but for lag 2 months.




Figure A4. As in A1 and A3, but for lag 3 months.

Author contributions. EB processed and analyzed the data, made the figures and wrote the manuscript. DI conceived the study and contributed to the interpretation of findings. SMas, SMat and PR equally contributed to interpreting, discussing and improving the results of the study. All authors reviewed the manuscript.

380 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. We gratefully acknowledge the three anonymous reviewers for their suggestions on how to improve the manuscript. We thank Ronan McAdam and Deep S. Banerjee for providing valuable insight into the calculation of ocean heat content and transport in the C-GLORSv5 and ORAS5 reanalyses.

References

- 385 A Woodgate, R. and Peralta-Ferriz, C.: Warming and Freshening of the Pacific Inflow to the Arctic from 1990-2019 implying dramatic shoaling in Pacific Winter Water ventilation of the Arctic water column, *Geophysical Research Letters*, 48, e2021GL092 528, 2021.
- Aagaard, K.: A synthesis of the Arctic Ocean circulation, Le Conseil, 1989.
- Årthun, M., Eldevik, T., Smedsrud, L. H., Skagseth, and Ingvaldsen, R. B.: Quantifying the influence of atlantic heat on barents sea ice variability and retreat, *Journal of Climate*, 25, 4736–4743, <https://doi.org/10.1175/JCLI-D-11-00466.1>, 2012.
- 390 Årthun, M., Eldevik, T., and Smedsrud, L. H.: The role of Atlantic heat transport in future Arctic winter sea ice loss, *Journal of Climate*, 32, 3327–3341, <https://doi.org/10.1175/JCLI-D-18-0750.1>, 2019.
- Asbjørnsen, H., Årthun, M., Skagseth, Ø., and Eldevik, T.: Mechanisms underlying recent arctic atlantification, *Geophysical Research Letters*, 47, e2020GL088 036, 2020.
- Aylmer, J., Ferreira, D., and Feltham, D.: Different mechanisms of Arctic and Antarctic sea ice response to ocean heat transport, *Climate*
- 395 *Dynamics*, <https://doi.org/10.1007/s00382-021-06131-x>, 2022.
- Balmaseda, M. A., Hernandez, F., Storto, A., Palmer, M. D., Alves, O., Shi, L., Smith, G. C., Toyoda, T., Valdivieso, M., Barnier, B., Behringer, D., Boyer, T., Chang, Y. S., Chepurin, G. A., Ferry, N., Forget, G., Fujii, Y., Good, S., Guinehut, S., Haines, K., Ishikawa, Y., Keeley, S., Köhl, A., Lee, T., Martin, M. J., Masina, S., Masuda, S., Meyssignac, B., Mogensen, K., Parent, L., Peterson, K. A., Tang, Y. M., Yin, Y., Vernieres, G., Wang, X., Waters, J., Wedd, R., Wang, O., Xue, Y., Chevallier, M., Lemieux, J. F., Dupont, F., Kuragano,
- 400 T., Kamachi, M., Awaji, T., Caltabiano, A., Wilmer-Becker, K., and Gaillard, F.: The ocean reanalyses intercomparison project (ORA-IP), *Journal of Operational Oceanography*, 8, s80–s97, <https://doi.org/10.1080/1755876X.2015.1022329>, 2015.
- Beszczynska-Möller, A., Fahrbach, E., Schauer, U., and Hansen, E.: Variability in Atlantic water temperature and transport at the entrance to the Arctic Ocean, 1997–2010, *ICES Journal of Marine Science*, 69, 852–863, 2012.
- Bliss, A. C., Steele, M., Peng, G., Meier, W. N., and Dickinson, S.: Regional variability of Arctic sea ice seasonal change climate indicators
- 405 from a passive microwave climate data record, *Environmental Research Letters*, 14, <https://doi.org/10.1088/1748-9326/aafb84>, 2019.
- Carmack, E., Polyakov, I., Padman, L., Fer, I., Hunke, E., Hutchings, J., Jackson, J., Kelley, D., Kwok, R., Layton, C., Melling, H., Perovich, D., Persson, O., Ruddick, B., Timmermans, M. L., Toole, J., Ross, T., Vavrus, S., and Winsor, P.: Toward quantifying the increasing role of oceanic heat in sea ice loss in the new arctic, *Bulletin of the American Meteorological Society*, 96, 2079–2105, <https://doi.org/10.1175/BAMS-D-13-00177.1>, 2015.
- 410 Carton, J. A., Penny, S. G., and Kalnay, E.: Temperature and salinity variability in the SODA3, ECCO4r3, and ORAS5 ocean reanalyses, 1993–2015, *Journal of Climate*, 32, 2277–2293, 2019.
- Comiso, J. C., Meier, W. N., and Gersten, R.: Variability and trends in the Arctic Sea ice cover: Results from different techniques, *Journal of Geophysical Research: Oceans*, 122, 6883–6900, <https://doi.org/10.1002/2017JC012768>, 2017.
- D’Angelo, G., Hunke, E. C., Pitts, T. R., Price, S. F., and Roberts, A. F.: Sea ice evolution along the Northern Sea Route and implications for
- 415 trans-Arctic shipping from 2021 through 2060, Tech. rep., Los Alamos National Lab.(LANL), Los Alamos, NM (United States), 2021.
- Davis, P. E., Lique, C., Johnson, H. L., and Guthrie, J. D.: Competing effects of elevated vertical mixing and increased freshwater input on the stratification and sea ice cover in a changing Arctic Ocean, *Journal of Physical Oceanography*, 46, 1531–1553, 2016.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haim-
- 420 berger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz,

- B. M., Morcrette, J. J., Park, B. K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J. N., and Vitart, F.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 425 Ding, Q., Schweiger, A., L'Heureux, M., Steig, E. J., Battisti, D. S., Johnson, N. C., Blanchard-Wrigglesworth, E., Po-Chedley, S., Zhang, Q., Harnos, K., Bushuk, M., Markle, B., and Baxter, I.: Fingerprints of internal drivers of Arctic sea ice loss in observations and model simulations, *Nature Geoscience*, 12, 28–33, <https://doi.org/10.1038/s41561-018-0256-8>, 2019.
- Dörr, J., Årthun, M., Eldevik, T., and Madonna, E.: Mechanisms of regional winter sea-ice variability in a warming arctic, *Journal of Climate*, 34, 8635–8653, <https://doi.org/10.1175/JCLI-D-21-0149.1>, 2021.
- Duarte, P., Sundfjord, A., Meyer, A., Hudson, S. R., Spreen, G., and Smedsrud, L. H.: Warm Atlantic Water Explains Observed Sea Ice Melt Rates North of Svalbard, *Journal of Geophysical Research: Oceans*, 125, <https://doi.org/10.1029/2019JC015662>, 2020.
- 430 England, M., Jahn, A., and Polvani, L.: Nonuniform contribution of internal variability to recent Arctic sea ice loss, *Journal of Climate*, 32, 4039–4053, 2019.
- Fichefet, T. and Maqueda, M. A.: Sensitivity of a global sea ice model to the treatment of ice thermodynamics and dynamics, *Journal of Geophysical Research: Oceans*, 102, 12 609–12 646, <https://doi.org/10.1029/97JC00480>, 1997.
- 435 Gimbert, F., Jourdain, N. C., Marsan, D., Weiss, J., and Barnier, B.: Recent mechanical weakening of the Arctic sea ice cover as revealed from larger inertial oscillations, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/10.1029/2011JC007633>, 2012.
- Graham, R. M., Itkin, P., Meyer, A., Sundfjord, A., Spreen, G., Smedsrud, L. H., Liston, G. E., Cheng, B., Cohen, L., Divine, D., Fer, I., Fransson, A., Gerland, S., Haapala, J., Hudson, S. R., Johansson, A. M., King, J., Merkouriadi, I., Peterson, A. K., Provost, C., Randelhoff, A., Rinke, A., Rösel, A., Sennéchaël, N., Walden, V. P., Duarte, P., Assmy, P., Steen, H., and Granskog, M. A.: Winter storms accelerate the demise of sea ice in the Atlantic sector of the Arctic Ocean, *Scientific Reports*, 9, <https://doi.org/10.1038/s41598-019-45574-5>, 2019.
- 440 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 445 Heuzé, C., Zanowski, H., Karam, S., and Muilwijk, M.: The deep Arctic Ocean and Fram Strait in CMIP6 models, *Journal of Climate*, 36, 2551–2584, 2023.
- Holland, M. M., Bitz, C. M., and Tremblay, B.: Future abrupt reductions in the summer Arctic sea ice, *Geophysical research letters*, 33, 2006.
- 450 Ilıcak, M., Drange, H., Wang, Q., Gerdes, R., Aksenov, Y., Bailey, D., Bentsen, M., Biastoch, A., Bozec, A., Böning, C., et al.: An assessment of the Arctic Ocean in a suite of interannual CORE-II simulations. Part III: Hydrography and fluxes, *Ocean Modelling*, 100, 141–161, 2016.
- Iovino, D., Selivanova, J., Masina, S., and Cipollone, A.: The Antarctic Marginal Ice Zone and Pack Ice Area in CMEMS GREP Ensemble Reanalysis Product, *Frontiers in Earth Science*, 10, 95, 2022.
- 455 Ivanov, V., Alexeev, V., Koldunov, N. V., Repina, I., Sandø, A. B., Smedsrud, L. H., and Smirnov, A.: Arctic Ocean heat impact on regional ice decay: A suggested positive feedback, *Journal of Physical Oceanography*, 46, 1437–1456, 2016.
- Ivanov, V. V., Alexeev, V. A., Repina, I., Koldunov, N. V., and Smirnov, A.: Tracing atlantic water signature in the arctic sea ice cover east of svalbard, *Advances in Meteorology*, 2012, <https://doi.org/10.1155/2012/201818>, 2012.

- Johnson, S. J., Stockdale, T. N., Ferranti, L., Balmaseda, M. A., Molteni, F., Magnusson, L., Tietsche, S., Decremer, D., Weisheimer, A.,
460 Balsamo, G., et al.: SEAS5: the new ECMWF seasonal forecast system, *Geoscientific Model Development*, 12, 1087–1117, 2019.
- Karami, M. P., Myers, P. G., de Vernal, A., Tremblay, L. B., and Hu, X.: The role of Arctic gateways on sea ice and circulation in the Arctic
and North Atlantic Oceans: a sensitivity study with an ocean-sea-ice model, *Climate Dynamics*, 57, 2129–2151, 2021.
- Koenigk, T. and Brodeau, L.: Ocean heat transport into the Arctic in the twentieth and twenty-first century in EC-Earth, *Climate Dynamics*,
42, 3101–3120, 2014.
- 465 Korhonen, M., Rudels, B., Marnela, M., Wisotzki, A., and Zhao, J.: Time and space variability of freshwater content, heat content and
seasonal ice melt in the Arctic Ocean from 1991 to 2011, *Ocean Science*, 9, 1015–1055, 2013.
- Lebrun, M., Vancoppenolle, M., Madec, G., and Massonnet, F.: Arctic sea-ice-free season projected to extend into autumn, *The Cryosphere*,
13, 79–96, 2019.
- Lenetsky, J. E., Tremblay, B., Brunette, C., and Meneghello, G.: Subseasonal predictability of Arctic Ocean sea ice conditions: Bering Strait
470 and Ekman-driven ocean heat transport, *Journal of Climate*, 34, 4449–4462, 2021.
- Li, D., Zhang, R., and Knutson, T. R.: On the discrepancy between observed and CMIP5 multi-model simulated Barents Sea winter sea ice
decline, *Nature Communications*, 8, 14991, 2017.
- Li, Z., Ding, Q., Steele, M., and Schweiger, A.: Recent upper Arctic Ocean warming expedited by summertime atmospheric processes,
Nature Communications, 13, <https://doi.org/10.1038/s41467-022-28047-8>, 2022.
- 475 Liang, X. and Losch, M.: On the effects of increased vertical mixing on the Arctic Ocean and sea ice, *Journal of Geophysical Research: Oceans*,
123, 9266–9282, 2018.
- Lien, V. S., Hjøllø, S. S., Skogen, M. D., Svendsen, E., Wehde, H., Bertino, L., Counillon, F., Chevallier, M., and Garric, G.: An assessment
of the added value from data assimilation on modelled Nordic Seas hydrography and ocean transports, *Ocean Modelling*, 99, 43–59, 2016.
- Lien, V. S., Schlichtholz, P., Skagseth, Ø., and Vikebø, F. B.: Wind-Driven Atlantic Water Flow as a Direct Mode for Reduced Barents Sea
480 Ice Cover, *Journal of Climate*, 30, 803–812, <https://doi.org/10.1175/jcli-d-16-0025.1>, 2017.
- Lind, S., Ingvaldsen, R. B., and Furevik, T.: Arctic warming hotspot in the northern Barents Sea linked to declining sea-ice import, *Nature
Climate Change*, 8, 634–639, <https://doi.org/10.1038/s41558-018-0205-y>, 2018.
- Lique, C. and Steele, M.: Seasonal to decadal variability of Arctic Ocean heat content: A model-based analysis and implications for au-
tonomous observing systems, *Journal of Geophysical Research: Oceans*, 118, 1673–1695, 2013.
- 485 Long, Z. and Perrie, W.: Changes in ocean temperature in the Barents Sea in the twenty-first century, *Journal of Climate*, 30, 5901–5921,
2017.
- MacKinnon, J. A., Simmons, H. L., Hargrove, J., Thomson, J., Peacock, T., Alford, M. H., Barton, B. I., Boury, S., Brenner, S. D., Couto,
N., et al.: A warm jet in a cold ocean, *Nature communications*, 12, 1–12, 2021.
- Madec, G., Bourdallé-Badie, R., Bouttier, P.-A., Bricaud, C., Bruciaferri, D., Calvert, D., Chanut, J., Clementi, E., Coward, A., Delrosso, D.,
490 et al.: NEMO ocean engine, 2017.
- Maslowski, W., Marble, D., Walczowski, W., Schauer, U., Clement, J. L., and Semtner, A. J.: On climatological mass, heat, and salt transports
through the Barents Sea and Fram Strait from a pan-Arctic coupled ice-ocean model simulation, *Journal of Geophysical Research: Oceans*,
109, 2004.
- Mayer, M., Tietsche, S., Haimberger, L., Tsubouchi, T., Mayer, J., and Zuo, H.: An improved estimate of the coupled Arctic energy budget,
495 *Journal of Climate*, 32, 7915–7934, 2019.

- McAdam, R., Masina, S., Balmaseda, M., Gualdi, S., Senan, R., and Mayer, M.: Seasonal forecast skill of upper-ocean heat content in coupled high-resolution systems, *Climate Dynamics*, 58, 3335–3350, 2022.
- Mikhailova, N., Bayankina, T., and Sizov, A.: Two modes of atmosphere–ocean interaction in the Atlantic sector of the Arctic Basin, *Oceanology*, 61, 443–449, 2021.
- 500 Mohamed, B., Nilsen, F., and Skogseth, R.: Marine heatwaves characteristics in the barents sea based on high resolution satellite data (1982–2020), *Frontiers in Marine Science*, 9, 821 646, 2022.
- Muilwijk, M., Smedsrud, L. H., Ilicak, M., and Drange, H.: Atlantic Water heat transport variability in the 20th century Arctic Ocean from a global ocean model and observations, *Journal of Geophysical Research: Oceans*, 123, 8159–8179, 2018.
- Muilwijk, M., Smedsrud, L. H., Polyakov, I. V., Nummelin, A., Heuzé, C., and Zanowski, H.: Divergence in CMIP6 projections of future
505 Arctic Ocean stratification, Tech. rep., Copernicus Meetings, 2022.
- Nie, Y., Uotila, P., Cheng, B., Massonnet, F., Kimura, N., Cipollone, A., and Lv, X.: Southern Ocean sea ice concentration budgets of five ocean-sea ice reanalyses, *Climate Dynamics*, 59, 3265–3285, 2022.
- Nummelin, A., Li, C., and Hezel, P. J.: Connecting ocean heat transport changes from the midlatitudes to the Arctic Ocean, *Geophysical Research Letters*, 44, 1899–1908, <https://doi.org/10.1002/2016GL071333>, 2017.
- 510 Olonscheck, D., Mauritsen, T., and Notz, D.: Arctic sea-ice variability is primarily driven by atmospheric temperature fluctuations, *Nature Geoscience*, 12, 430–434, <https://doi.org/10.1038/s41561-019-0363-1>, 2019.
- Onarheim, I. H., Smedsrud, L. H., Ingvaldsen, R. B., and Nilsen, F.: Loss of sea ice during winter north of Svalbard, *Tellus, Series A: Dynamic Meteorology and Oceanography*, 66, <https://doi.org/10.3402/tellusa.v66.23933>, 2014.
- Onarheim, I. H., Eldevik, T., Årthun, M., Ingvaldsen, R. B., and Smedsrud, L. H.: Skillful prediction of Barents Sea ice cover, *Geophysical
515 Research Letters*, 42, 5364–5371, <https://doi.org/10.1002/2015GL064359>, 2015.
- Onarheim, I. H., Eldevik, T., Smedsrud, L. H., and Stroeve, J. C.: Seasonal and regional manifestation of Arctic sea ice loss, *Journal of Climate*, 31, 4917–4932, <https://doi.org/10.1175/JCLI-D-17-0427.1>, 2018.
- Pan, R., Shu, Q., Wang, Q., Wang, S., Song, Z., He, Y., and Qiao, F.: Future Arctic Climate Change in CMIP6 Strikingly Intensified by NEMO-Family Climate Models, *Geophysical Research Letters*, 50, e2022GL102 077, 2023.
- 520 Peng, G. and Meier, W. N.: Temporal and regional variability of Arctic sea-ice coverage from satellite data, *Annals of Glaciology*, 59, 191–200, 2018.
- Peralta-Ferriz, C. and Woodgate, R. A.: Seasonal and interannual variability of pan-Arctic surface mixed layer properties from 1979 to 2012 from hydrographic data, and the dominance of stratification for multiyear mixed layer depth shoaling, *Progress in Oceanography*, 134, 19–53, 2015.
- 525 Perovich, D. K.: The changing Arctic sea ice cover, *Oceanography*, 24, 162–173, 2011.
- Perovich, D. K. and Polashenski, C.: Albedo evolution of seasonal Arctic sea ice, *Geophysical Research Letters*, 39, 2012.
- Perovich, D. K., Light, B., Eicken, H., Jones, K. F., Runciman, K., and Nghiem, S. V.: Increasing solar heating of the Arctic Ocean and adjacent seas, 1979-2005: Attribution and role in the ice-albedo feedback, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL031480>, 2007.
- 530 Polyakov, I., Alekseev, G., Timokhov, L., Bhatt, U., Colony, R., Simmons, H., Walsh, D., Walsh, J., and Zakharov, V.: Variability of the intermediate Atlantic water of the Arctic Ocean over the last 100 years, *Journal of Climate*, 17, 4485–4497, 2004.
- Polyakov, I., Bhatt, U., Simmons, H., Walsh, D., Walsh, J., and Zhang, X.: Multidecadal variability of North Atlantic temperature and salinity during the twentieth century, *Journal of Climate*, 18, 4562–4581, 2005.

- Polyakov, I. V., Timokhov, L. A., Alexeev, V. A., Bacon, S., Dmitrenko, I. A., Fortier, L., Frolov, I. E., Gascard, J. C., Hansen, E., Ivanov, V. V., Laxon, S., Mauritzen, C., Perovich, D., Shimada, K., Simmons, H. L., Sokolov, V. T., Steele, M., and Toole, J.: Arctic ocean warming contributes to reduced polar ice cap, *Journal of Physical Oceanography*, 40, 2743–2756, <https://doi.org/10.1175/2010JPO4339.1>, 2010.
- Polyakov, I. V., Pnyushkov, A. V., Alkire, M. B., Ashik, I. M., Baumann, T. M., Carmack, E. C., Goszczko, I., Guthrie, J., Ivanov, V. V., Kanzow, T., et al.: Greater role for Atlantic inflows on sea-ice loss in the Eurasian Basin of the Arctic Ocean, *Science*, 356, 285–291, 2017.
- Polyakov, I. V., Alkire, M. B., Bluhm, B. A., Brown, K. A., Carmack, E. C., Chierici, M., Danielson, S. L., Ellingsen, I., Ershova, E. A., Gårdfeldt, K., et al.: Borealization of the Arctic Ocean in response to anomalous advection from sub-Arctic seas, *Frontiers in Marine Science*, 7, 491, 2020a.
- Polyakov, I. V., Rippeth, T. P., Fer, I., Alkire, M. B., Baumann, T. M., Carmack, E. C., Ingvaldsen, R., Ivanov, V. V., Janout, M., Lind, S., Padman, L., Pnyushkov, A. V., and Rember, R.: Weakening of cold halocline layer exposes sea ice to oceanic heat in the eastern arctic ocean, *Journal of Climate*, 33, 8107–8123, <https://doi.org/10.1175/JCLI-D-19-0976.1>, 2020b.
- Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., and Schlax, M. G.: Daily high-resolution-blended analyses for sea surface temperature, *Journal of climate*, 20, 5473–5496, 2007.
- Ricker, R., Kauker, F., Schweiger, A., Hendricks, S., Zhang, J., and Paul, S.: Evidence for an increasing role of ocean heat in arctic winter sea ice growth, *Journal of Climate*, 34, 5215–5227, <https://doi.org/10.1175/JCLI-D-20-0848.1>, 2021.
- Rudels, B.: Arctic Ocean circulation, processes and water masses: A description of observations and ideas with focus on the period prior to the International Polar Year 2007–2009, *Progress in Oceanography*, 132, 22–67, 2015.
- Sandø, A. B., Nilsen, J. Ø., Gao, Y., and Lohmann, K.: Importance of heat transport and local air-sea heat fluxes for Barents Sea climate variability, *Journal of Geophysical Research: Oceans*, 115, 2010.
- Schauer, U. and Beszczynska-Möller, A.: Problems with estimating oceanic heat transport—conceptual remarks for the case of Fram Strait in the Arctic Ocean, *Ocean Science Discussions*, 6, 1007–1029, 2009.
- Serreze, M. C. and Meier, W. N.: The Arctic’s sea ice cover: trends, variability, predictability, and comparisons to the Antarctic, *Annals of the New York Academy of Sciences*, 1436, 36–53, 2019.
- Serreze, M. C., Crawford, A. D., Stroeve, J. C., Barrett, A. P., and Woodgate, R. A.: Variability, trends, and predictability of seasonal sea ice retreat and advance in the Chukchi Sea, *Journal of Geophysical Research: Oceans*, 121, 7308–7325, <https://doi.org/10.1002/2016JC011977>, 2016.
- Shu, Q., Wang, Q., Song, Z., and Qiao, F.: The poleward enhanced Arctic Ocean cooling machine in a warming climate, *Nature Communications*, 12, <https://doi.org/10.1038/s41467-021-23321-7>, 2021.
- Shu, Q., Wang, Q., Årthun, M., Wang, S., Song, Z., Zhang, M., and Qiao, F.: Arctic Ocean Amplification in a warming climate in CMIP6 models, *Science Advances*, 8, eabn9755, 2022.
- Simmonds, I.: Comparing and contrasting the behaviour of Arctic and Antarctic sea ice over the 35 year period 1979–2013, *Annals of Glaciology*, 56, 18–28, <https://doi.org/10.3189/2015AoG69A909>, 2015.
- Skagseth, Ø., Eldevik, T., Årthun, M., Asbjørnsen, H., Lien, V. S., and Smedsrud, L. H.: Reduced efficiency of the Barents Sea cooling machine, *Nature Climate Change*, 10, 661–666, 2020.
- Smedsrud, L. H., Ingvaldsen, R., Nilsen, J. Ø., and Skagseth, Ø.: Heat in the Barents Sea: Transport, storage, and surface fluxes, *Ocean Science*, 6, 219–234, 2010.

- Smedsrud, L. H., Esau, I., Ingvaldsen, R. B., Eldevik, T., Haugan, P. M., Li, C., Lien, V. S., Olsen, A., Omar, A. M., Risebrobakken, B., Sandø, A. B., Semenov, V. A., and Sorokina, S. A.: The role of the Barents Sea in the Arctic climate system, *Reviews of Geophysics*, 51, 415–449, <https://doi.org/10.1002/rog.20017>, 2013.
- 575 Smedsrud, L. H., Muilwijk, M., Brakstad, A., Madonna, E., Lauvset, S. K., Spensberger, C., Born, A., Eldevik, T., Drange, H., Jeansson, E., et al.: Nordic Seas heat loss, Atlantic inflow, and Arctic sea ice cover over the last century, *Reviews of Geophysics*, 60, e2020RG000725, 2022.
- Stammerjohn, S., Massom, R., Rind, D., and Martinson, D.: Regions of rapid sea ice change: An inter-hemispheric seasonal comparison, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL050874>, 2012.
- Steele, M., Ermold, W., and Zhang, J.: Arctic Ocean surface warming trends over the past 100 years, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2007GL031651>, 2008.
- 580 Storto, A. and Masina, S.: C-GLORSv5: An improved multipurpose global ocean eddy-permitting physical reanalysis, *Earth System Science Data*, 8, 679–696, <https://doi.org/10.5194/essd-8-679-2016>, 2016.
- Storto, A., Alvera-Azcárate, A., Balmaseda, M. A., Barth, A., Chevallier, M., Counillon, F., Domingues, C. M., Drevillon, M., Drillet, Y., Forget, G., et al.: Ocean reanalyses: recent advances and unsolved challenges, *Frontiers in Marine Science*, 6, 418, 2019.
- 585 Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, *Environmental Research Letters*, 13, 103001, 2018.
- Stroeve, J. C., Crawford, A. D., and Stammerjohn, S.: Using timing of ice retreat to predict timing of fall freeze-up in the Arctic, *Geophysical Research Letters*, 43, 6332–6340, <https://doi.org/10.1002/2016GL069314>, 2016.
- Sumata, H., de Steur, L., Divine, D. V., Granskog, M. A., and Gerland, S.: Regime shift in Arctic Ocean sea ice thickness, *Nature*, 615, 443–449, 2023.
- 590 Takahashi, N., Richards, K. J., Schneider, N., Annamalai, H., Hsu, W. C., and Nonaka, M.: Formation Mechanism of Warm SST Anomalies in 2010s Around Hawaii, *Journal of Geophysical Research: Oceans*, 126, <https://doi.org/10.1029/2021JC017763>, 2021.
- Thépaut, J.-N., Dee, D., Engelen, R., and Pinty, B.: The Copernicus programme and its climate change service, in: *IGARSS 2018-2018 IEEE International Geoscience and Remote Sensing Symposium*, pp. 1591–1593, IEEE, 2018.
- Thomson, R. E. and Fine, I. V.: *Estimating Mixed Layer Depth from Oceanic Profile Data*, Tech. rep., 2003.
- 595 Timmermans, M. L., Cole, S., and Toole, J.: Horizontal density structure and restratification of the Arctic Ocean surface layer, *Journal of Physical Oceanography*, 42, 659–668, <https://doi.org/10.1175/JPO-D-11-0125.1>, 2012.
- Toole, J. M., Timmermans, M. L., Perovich, D. K., Krishfield, R. A., Proshutinsky, A., and Richter-Menge, J. A.: Influences of the ocean surface mixed layer and thermohaline stratification on Arctic Sea ice in the central Canada Basin, *Journal of Geophysical Research: Oceans*, 115, <https://doi.org/10.1029/2009JC005660>, 2010.
- 600 Tsubouchi, T., Bacon, S., Naveira Garabato, A., Aksenov, Y., Laxon, S. W., Fahrbach, E., Beszczynska-Möller, A., Hansen, E., Lee, C. M., and Ingvaldsen, R.: The Arctic Ocean in summer: A quasi-synoptic inverse estimate of boundary fluxes and water mass transformation, *Journal of Geophysical Research: Oceans*, 117, 2012.
- Tsubouchi, T., von Appen, W., Schauer, U., Kanzow, T., Lee, C., Curry, B., de Steur, L., Ingvaldsen, R., and Woodgate, R.: The Arctic Ocean volume, heat and fresh water transports time series from October 2004 to May 2010, *PANGAEA* <https://doi.org/10.1594/PANGAEA.909966>, 2019.
- 605 Tsubouchi, T., Våge, K., Hansen, B., Larsen, K. M. H., Østerhus, S., Johnson, C., Jónsson, S., and Valdimarsson, H.: Increased ocean heat transport into the Nordic Seas and Arctic Ocean over the period 1993–2016, *Nature Climate Change*, 11, 21–26, <https://doi.org/10.1038/s41558-020-00941-3>, 2021.

- Uotila, P., Goosse, H., Haines, K., Chevallier, M., Barthélemy, A., Bricaud, C., Carton, J., Fučkar, N., Garric, G., Iovino, D., et al.: An
610 assessment of ten ocean reanalyses in the polar regions, *Climate Dynamics*, 52, 1613–1650, 2019.
- Wang, Q., Wang, X., Wekerle, C., Danilov, S., Jung, T., Koldunov, N., Lind, S., Sein, D., Shu, Q., and Sidorenko, D.: Ocean heat transport
into the Barents Sea: Distinct controls on the upward trend and interannual variability, *Geophysical Research Letters*, 46, 13 180–13 190,
2019.
- Woodgate, R. A., Weingartner, T., and Lindsay, R.: The 2007 Bering Strait oceanic heat flux and anomalous Arctic sea-ice retreat, *Geophys-
615 ical Research Letters*, 37, <https://doi.org/10.1029/2009GL041621>, 2010.
- Woodgate, R. A., Weingartner, T. J., and Lindsay, R.: Observed increases in Bering Strait oceanic fluxes from the Pacific to the Arctic from
2001 to 2011 and their impacts on the Arctic Ocean water column, *Geophysical Research Letters*, 39, 2012.
- Yang, J.: The seasonal variability of the Arctic Ocean Ekman transport and its role in the mixed layer heat and salt fluxes, *Journal of Climate*,
19, 5366–5387, 2006.
- 620 Zuo, H., Balmaseda, M. A., Tietsche, S., Mogensen, K., and Mayer, M.: The ECMWF operational ensemble reanalysis-analysis system for
ocean and sea ice: A description of the system and assessment, *Ocean Science*, 15, 779–808, <https://doi.org/10.5194/os-15-779-2019>,
2019.