

Causal ~~associations~~ relationships and predictability of the summer East Atlantic teleconnection

Julianna Carvalho-Oliveira^{1,2,3,a}, Giorgia Di Capua^{4,5}, Leonard Borchert¹, Reik V. Donner^{5,4}, and Johanna Baehr¹

¹Institute of Oceanography, Center for Earth System Research and Sustainability (CEN), Universität Hamburg, Hamburg, Germany

²International Max Planck Research School on Earth System Modelling, Max Planck Institute for Meteorology, Hamburg, Germany

³Helmholtz-Zentrum Hereon, Institute of Coastal Systems - Analysis and Modeling, Geesthacht, Germany

⁴Potsdam Institute for Climate Impact Research, Member of the Leibniz Association, Potsdam, Germany

⁵Department of Water, Environment, Construction and Safety, Magdeburg-Stendal University of Applied Sciences, Magdeburg, Germany

^apresent address: Leipzig Institute for Meteorology, Universität Leipzig, Leipzig, Germany

Correspondence: Julianna Carvalho-Oliveira (julianna.carvalho.oliveira@uni-hamburg.de)

Abstract.

We apply Causal Effect Networks to evaluate the influence of spring North Atlantic extratropical sea surface temperatures (~~SSTNA-SST~~) on the summer East Atlantic Pattern (EA) seasonal predictability during the 20th-century period of 1908-2008. We find in the ERA-20C reanalysis that a meridional ~~SST-NA-SST~~ gradient in spring (SST index) causally influences the summer EA, with an estimated causal effect expressed by a β -coefficient of about 0.2 (a 1 standard deviation change in spring SST index causes a 0.2 standard deviation change in the EA 3-4 months later). ~~We only find this link to be causal, however,~~ Notably, this causal link is only evident during the period from 1958 -2008- to 2008, and is not observed throughout the entirety of the investigated period. When performing the analysis on 45-year-long ~~timeseries-time series~~ randomly sampled in this late period, we find the strength of the causal link to be affected by interannual variability, suggesting a potential modulation by an external physical mechanism. In addition to the summer EA, we find that ~~spring-SST~~ the spring SST index has an estimated causal effect of about -0.2 on summer 2-metre air temperatures over northwestern Europe, possibly mediated by summer EA. We then use a pre-industrial and a historical simulation, as well as a 30-member initialised seasonal prediction ensemble with MPI-ESM-MR to test the model performance in reproducing the detected causal links in ERA-20C and to evaluate whether this performance might leave an imprint in the model predictive skill of European summer climate. We find that while both the pre-industrial and historical simulations using MPI-ESM-MR ~~is-are~~ mostly unable to reproduce the causal link between spring SST and the summer EA among the different datasets, the 30-member initialised ensemble can ~~moderately~~ reproduce a causal link between spring SST and summer 2-metre air temperatures over a region west of the British Isles. We perform a predictive skill assessment conditioned on the spring SST index causal links for July-August sea level pressure, 500 hPa geopotential height and 2-metre air temperatures for predictions initialised in May. Our results suggest that MPI-ESM-MR's performance in reproducing the spring SST index causal links constrains the seasonal prediction skill of European summer climate.

1 Introduction

The summer East Atlantic Pattern (EA) is an important atmospheric teleconnections influencing weather and climate in the Euro-Atlantic region (e.g. Comas-Bru and McDermott (2014); Bastos et al. (2016)). Along with the summer North Atlantic Oscillation (NAO), these teleconnections are often used to describe the combined changes in latitude and speed of the North Atlantic jet stream (Woollings et al., 2010) – one of the major modulators of mid-latitude weather extremes (e.g. Rousi et al. (2022)). Understanding the predictability associated with these teleconnections is therefore of paramount importance. Although several recent studies have focused on predictability of the NAO (Domeisen et al., 2018; O'Reilly et al., 2019; Athanasiadis et al., 2020; Klavans et al., 2021), the EA has received less attention. Here, we apply ~~a causal inference-based tool~~ [the Peter and Clark momentary conditional independence \(PCMCI\) causal discovery algorithm](#) to evaluate the influence of North Atlantic extratropical surface temperatures (~~SST~~[NA-SST](#)) on the predictability of EA at seasonal timescales.

The most common description of the EA pattern features a well-defined sea level pressure (SLP) centre of action south of Iceland and west of the British Isles, usually defined as the second leading empirical orthogonal function (EOF) of SLP in the Euro-Atlantic region (e.g. Moore et al. (2013)). Wallace and Gutzler (1981) define a positive phase of the EA as characterised by the centre of action exhibiting anticyclonic conditions, featuring the northward extension of the Azores High. A positive EA has been associated with below-average surface temperatures (Cassou et al., 2005; Comas-Bru and Hernández, 2018) and dry spells in parts of Europe (Rousi et al., 2021). Conversely, anomalous cyclonic conditions offshore of Ireland have been suggested to influence heatwaves in Europe for a negative EA phase (e.g. Duchez et al. (2016)). Using a clustering approach (e.g. Cassou et al. (2004); Carvalho-Oliveira et al. (2022)), a positive EA phase is reminiscent of an Atlantic Ridge, whereas a negative EA phase resembles the Atlantic Low. A common feature amongst the different EA definitions is that its centre of action is positioned along the NAO nodal line, thus ultimately modulating the location and strength of the NAO dipole and the North Atlantic storm track (Woollings et al., 2010). That is, summer climate predictability in the Euro-Atlantic region is closely linked to EA variability.

While there is no consensus on the physical processes driving the EA, spring ~~North Atlantic sea surface temperatures (SSTs)~~ [NA-SST](#) have been proposed to influence EA variability and predictability. Gastineau and Frankignoul (2015) suggested that summer 500-hPa geopotential height anomalies in the Euro-Atlantic significantly co-vary with a spring ~~North Atlantic SST~~ [NA-SST](#) tripole pattern in observations over the 20th century. Moreover, Carvalho-Oliveira et al. (2022) suggested that spring North Atlantic SSTs can influence predictive skill of summers dominated by EA in initialised simulations. Based on linear regression analyses of the period 1979–2017, Ossó et al. (2018) and Ossó et al. (2020) proposed a physical mechanism whereby anomalous extratropical North Atlantic SSTs in spring may persist into summer and influence shifts in the eddy-driven jet stream, imprinting at the surface an SLP pattern that resembles the EA. These studies suggest that this mechanism is forced by changes in baroclinicity of the lower troposphere associated with a strong meridional ~~SST~~ [NA-SST](#) gradient in spring located between subpolar and subtropical ~~North Atlantic gyres~~. The authors hypothesised that the delayed atmospheric response in summer, and not in spring, could be explained by the seasonal evolution of both ~~SST~~ [NA-SST](#) gradient and jet stream position, modulated by a positive coupled ocean–atmosphere feedback that operates primarily in summer.

55 Nevertheless, while the linear regression-based analysis provided in Ossó et al. (2018) suggests a contribution of spring ~~SST-NA-SST~~ on the summer SLP variability, this approach does not imply causation. Disentangling the complex causal-effect pathways underlying the mechanism proposed in Ossó et al. (2020) over a long observational record is a crucial step to evaluate EA predictability in dynamical climate models. Hence, in this paper we use Causal Effect Network (~~CEN~~; ~~Runge et al. (2015); Kretschmer et al. (2016)~~ based on PCMCI (hereafter: CEN, Kretschmer et al. (2016)) to test the hypothesis of spring ~~SST-NA-SST~~ causally driving a response in the summer SLP and temperature fields in the Euro-Atlantic sector during the 20th century. CEN overcomes spurious correlations due to autocorrelation, indirect effects, or common drivers (~~Runge et al., 2019~~), and (~~Runge et al., 2014, 2019~~). It has been successfully used to complement hypothesis testing for other ~~teleconnections~~ tropical and mid-latitude teleconnections in the Atlantic-Pacific region (e.g. Karmouche et al. (2023) and Indian Ocean (e.g. Di Capua et al. (2020a)), as well as in the Arctic region (e.g. Siew et al. (2020); Kretschmer et al. (2020)).

65 Although dynamical seasonal forecasts of European summer climate usually show very little skill (e.g. Mishra et al. (2019)), recent studies suggest that improving the representation of teleconnections can increase forecast skill (Oliveira et al., 2020; Carvalho-Oliveira et al., 2022; Schuhen et al., 2022). The physical mechanism connecting ~~SST-NA-SST~~ variability and jet stream dynamics proposed in Ossó et al. (2020) thus offers a framework to more generally assess the influence of ~~SST-NA-SST~~ on seasonal predictability of the EA – the aim of the present study.

Here, we use CEN to firstly investigate under which circumstances spring extratropical North Atlantic SSTs causally influence the summer EA and its associated impact on surface climate. Secondly, we analyse pre-industrial, historical and initialised simulations with the Max Planck Institute Earth System Model in its mixed-resolution setup MPI-ESM-MR (MPI-ESM-MR, Dobrynin et al. (2018)). We specifically test the model performance in reproducing the observed ~~SST-NA-SST~~ - EA link, in order to identify how this performance might constrain the seasonal prediction skill of European summer climate.

2 Methodology

2.1 Reanalysis and model data

We investigate the ~~SST-NA-SST~~ - EA link first using ERA-20C reanalysis (Poli et al., 2016), and then using model simulations with MPI-ESM-MR (Dobrynin et al., 2018)). ~~We analyse sea level pressure, sea surface temperature (SST), The physical~~ variables analysed are NA-SST, SLP and air temperature at 2 metre height (T2m). We use monthly means for each variable as we are testing mechanisms which are expected to act on monthly timescales. We focus our analysis on the 101-year long period spanning 1908-2008 in the model and observations.

In MPI-ESM-MR, the atmospheric component ECHAM6 (Stevens et al., 2013) has a resolution of T63L95, with a nominal horizontal resolution of 200 km (1.875°) and 95 vertical layers up to 0.01 hPa. The oceanic component MPI-OM (Jungclaus et al., 2013) is coupled to ECHAM6 and has a resolution of TP04L40, with an approximate horizontal resolution of 40 km (0.4°) and 40 vertical layers. External forcing is taken from CMIP5 (Giorgetta et al., 2013).

We investigate how MPI-ESM-MR performs in reproducing the SST-NA-SST - EA link among three independent sets of MPI-ESM-MR simulations. The datasets comprise a pre-industrial control run (piControl), a historical run, and a 30-member seasonal initialised hindcast ensemble (MR-30). Comparing the performance of each set against reanalysis enables us to distinguish the role of forcing (from piControl to historical), and of assimilation (historical to initialised ensemble) on the model skill.

The pre-industrial coupled atmosphere/ocean control run piControl has a total length of 1000 years (period 1850-2849) (Giorgetta et al., 2011), with forcing constant in time: orbital parameters and greenhouse gases concentration are fixed at 1850 values; spectral solar irradiance remains constant as the solar cycle average over 1844-1856, and monthly ozone concentrations are fixed at the 11-year average over 1850-1860 (Mauritsen et al., 2012). The historical simulations run from 1850 to 2005 under natural and anthropogenic forcing following CMIP5 protocol (Dobrynin et al., 2018).

Lastly, the hindcast ensemble MR-30 is initialised on 1st of May every year from 1902-2008, with initial conditions taken from an assimilation experiment (Oliveira et al., 2020). In the assimilation experiment, Newtonian relaxation (*nudging*) is used in full-field mode towards all atmospheric and ocean levels except in the boundary layer. The atmosphere conditions of vorticity, divergence, three-dimensional temperature and two-dimensional pressure are assimilated with ERA-20C data. In the ocean, three-dimensional daily mean salinity and temperature anomalies are nudged at a relaxation time of approximately 10 days. The To help reduce initialisation shock, the ocean state is derived in from an ocean-only simulation performed with MPI-OM forced with the atmospheric variables from ERA-20C, thus maintaining consistency in model physics. The three-dimensional atmospheric and ocean fields of the assimilation experiment form the initial conditions, from which 30 ensemble members are generated by perturbing the atmospheric state with slightly disturbed diffusion coefficients in the uppermost layer.

2.2 Data-processing and climate indices

We compute anomalies at every gridpoint by removing mean seasonal cycle and linear trend, satisfying data input requirements for the CEN algorithm (Kretschmer et al., 2016). We analyse bimonthly means in March-April (MA) and April-May (AM) for spring SST-NA-SST and July-August (JA) SLP and T2m. In MR-30, we use the assimilation experiment to obtain spring SST-NA-SST fields, and the hindcast ensemble at lead times of 3-4 months to obtain summer SLP, T2m and 500 hPa geopotential height (Z500). We apply area-weighting by multiplying each value with the cosine of its latitudinal location to take into account the dependence of the gridpoint density on latitude.

We calculate the EA index to analyse the summer EA teleconnection. As a first step, we define a reference EA index as the second principal component (PC) of the leading-empirical-orthogonal-function-(EOF)-EOF of JA anomalies of sea-level pressure-SLP over the Euro-Atlantic sector 70°W-40°E, 25°-80°N calculated from the ERA-20C reanalysis data (e.g. Comas-Bru and McDermott (2014)). Next, EA index values in the model simulations from MPI-ESM-MR are calculated by projecting each ensemble member onto the EA reference EOF pattern. We consider a positive phase of the EA index when characterised by a centre of positive SLP anomalies that lies south of Iceland and west of the British Isles (e.g. Wallace and Gutzler (1981); Comas-Bru and McDermott (2014), Fig.1a).

120 We further test the influence of spring extratropical North Atlantic SSTs on the summer EA using the SST index proposed in Ossó et al. (2018). We calculate the SST index by subtracting the average SST_{NA-SST} over the eastern box (35°W-20°W, 35°-42°N) from the average SST_{NA-SST} over the western box (52°W-40°W, 42°-52°N), represented by green boxes in Fig. 1b. We analyse the SST index for both March-April and April-May means.

To ~~fully analyse the impact of spring SST on the~~ comprehensively investigate the influence of spring NA-SST on summer SLP variability, we ~~include-incorporate~~ the SLP index ~~proposed in Ossó et al. (2018)~~, ~~in addition to the EA index~~ introduced by Ossó et al. (2018) alongside the EA index. This approach aims to address the broader significance of pressure dynamics in the region, particularly in relation to the physical mechanism proposed by Ossó et al. (2020). The SLP index is calculated as JA SLP anomalies averaged over the region 45°N-55°N; 25°W-5°W indicated by a blue box in Fig 1b. In a second step, we analyse the impact of NA-SST on summer T2m using two additional indices, i.e. T2m_{CE} and T2m_{Ridge} (Sects.3.2,3.4). The T2m_{CE} index is calculated as JA T2m anomalies averaged over the region 45°N-55°N; 10°E-35°E (indicated by a red box in Fig.2f), and the T2m_{Ridge} index is calculated over the region 40°N-55°N; 15°W-34°W (indicated by a black box in Fig.7b).

All climate indices are standardised to have mean of zero and ~~SD~~ standard deviation (SD) of 1 to allow for comparison. Using the aforementioned climate indices, we perform linear regressions and correlations to analyse the linear relationship between the predictor spring SST_{NA-SST} and the target variables summer EA, SLP, and T2m_{CE} indices. ~~Wherever useful,~~ we use a two-tailed Student's t-test to calculate the statistical significance of ~~correlations~~ point-wise correlations maps. We provide a description of the indices used in the analysis in Table 1.

Table 1. Summary of the indices used in our analysis.

<u>Index</u>	<u>Variable name</u>	<u>Region used for calculation</u>
<u>SST_ind</u>	<u>Sea surface temperature index</u>	<u>Eastern box (35°W-20°W, 35°-42°N); Western box (52°W-40°W, 42°-52°N)</u>
<u>SLP_ind</u>	<u>Sea level pressure index</u>	<u>45°N-55°N; 25°W-5°W</u>
<u>EA</u>	<u>East Atlantic Pattern</u>	<u>70°W-40°E, 25°-80°N</u>
<u>T2m_CE</u>	<u>Air temperature at 2m height for central Europe</u>	<u>46°N-55°N; 11°E-34°E</u>
<u>T2m_Ridge</u>	<u>Air temperature at 2m height for Atlantic Ridge region</u>	<u>40°N-55°N; 15°W-34°W</u>

2.3 Causal effect networks

We use Causal Effect Network analysis (CEN, Runge et al. (2015); Kretschmer et al. (2016)) to test whether spring SST_{NA-SST} anomalies causally influences the variability of summer SLP and temperature fields in the Euro-Atlantic sector during the 20th century. ~~CEN analysis is a causal discovery tool which implements the so-called~~, investigating the mechanism

proposed in Ossó et al. (2018, 2020). CEN allow to represent the output of the Peter and Clark momentary conditional independence algorithm (PC-MCI, Runge et al. (2019) (PCMCI) causal discovery algorithm (Runge et al., 2019; Spirtes et al., 2000). We specifically use the PCMCI version 4.2 from the Python package Tigramite (<https://github.com/jakobrunge/tigramite>). This algorithm iteratively calculates partial correlations amongst a set of ~~time-series~~ time series (actors) to test if a link between a potential precursor and a target variable at a certain time lag is: *i*) considered spurious, i.e. can be explained by the linear combination of other ~~time-series~~ time series at different lags (i.e. conditional independence); or *ii*) considered causal, i.e. cannot be explained by the combined influence of other investigated variables (i.e. conditional dependence). ~~We stress, however, In the algorithm, this calculation is performed for a minimum and maximum time lags, denoted τ_{min} and τ_{max} .~~

We emphasise that the term ~~causal~~ "causal" should be interpreted ~~as causal~~ cautiously within the context of this study. When we refer to causality, we mean causality relative to the set of investigated variables ~~, under the and under the specific assumptions considered in the PC-MCI algorithm (e.g. stationarity of time series).~~ We refer the reader to Runge (2018) for ~~a thorough description~~ PCMCI algorithm (such as the stationarity of time series data). As a consequence, the possibility of remaining spurious correlations cannot be entirely ruled out. The choices of variables included in the analysis is another crucial aspect for determining the causality of the identified links. Yet, this poses a challenge as including more variables enhances the credibility of causal discoveries but introduces complexities. For instance, accommodating numerous variables and significant time lags to address physical delays, like identifying atmospheric teleconnections, leads to high dimensionality. This, in turn, can significantly affect the reliability of statistical outcomes. Hence, a successful application of CEN requires (such as for any data-driven method), expert knowledge of the underlying physical processes, including relevant variables, time-scales and temporal resolution. For a more detailed understanding of the CEN analysis and the ~~PC-MCI algorithm~~ PCMCI algorithm, we refer the reader to Runge (2018), which provides a comprehensive description of these techniques.

We visualise the output of ~~CEN analysis as a process graph, where circles~~ PCMCI in a CEN, i.e. a causal graph where nodes represent the investigated variables, ~~and~~ arrows indicate the ~~strength and the~~ direction of the causal links, ~~and colours denote the strength of these links~~. The strength is expressed by the standardised linear regression coefficient, denoted β -coefficient, and defined as the expected change of Y_t in units of its ~~standard deviation (SD)~~ SD induced by raising $X_{t-\tau}$ by 1 SD, while keeping all other potential precursors constant. Moreover, CEN analysis outputs the autocorrelation path coefficient, which represents the causal influence of a variable on itself, as opposed to the Pearson autocorrelation.

We apply causal maps (Di Capua et al., 2020b) to investigate the causal effects of a specific variable on a given atmospheric field along latitude, longitude and time dimensions. This tool builds upon the ~~PC-MCI~~ PCMCI algorithm and CEN approach, and provides a powerful visualisation of spatial patterns. Causal maps display β -coefficients calculated with the time series of a potential precursor and each grid point of a target atmospheric field. We refer the reader to Di Capua et al. (2020b) for a detailed explanation of this method.

Lastly, the PCMCI parameters are chosen as follows: ~~pc~~ alpha = 0.2, alpha level to print results = 0.1, τ_{min} = 3 months and τ_{max} = 4 months, independence test = parcorr, significance='analytic', masking type 'y'.

2.4 Bootstrapping and ensemble subsampling

175 We perform bootstrapping and ensemble subsampling (e.g. Dobrynin et al. (2018)) to investigate the sensitivity of the causal links to data sampling. When analysing 1958-2008 using observations in Sect.3.3, we randomly select 500 samples of 45-years long, i.e. excluding 6 random years at each iteration. Each of these time series are then analysed with CEN using the same hyperparameters (see Sect.2.3). We perform a similar bootstrapping using MR-30, but additionally include a second step of ensemble subsampling. That is, we first randomly exclude 6 years from the analysed period. Next, we randomly select 1
180 ensemble member from the 30-member set for each selected year. Each time series is then analysed with CEN. This process is repeated at least 500 times. It is important to note that reducing the length of the time series in this way increases the variability and hence lowers the significance of the obtained β -coefficients. However, this should not by itself lower the strength of the coefficients themselves.

2.5 Predictive skill assessment

185 In Sect.3.4, we perform a predictive skill assessment for SLP, T2m and Z500 at lead times of 3-4 months in MR-30 against ERA-20C. For this assessment we use point-wise detrended anomaly correlation coefficient (ACC, Collins (2002)). We are interested in assessing the predictive skill conditioned to the strength of significant β -coefficients (p-value < 0.1). Our hypothesis is that the predictive skill in summer is likely to increase in cases where MR-30 is able to capture the causal link between spring SST index and summer EA, as opposed to cases where the model fails to capture the observed causal link. We refer to these
190 time series as *MR-30 bootstrap ensemble*. For example, we shall assume that we are interested in calculating the conditioned predictive skill of JA Z500. To accomplish this task, we first identify the specific years and ensemble members that correspond to significant β -coefficients for the spring SST and summer EA. With this information, we can then sample JA Z500 to create a time series of similar length. In case more than one ensemble member is randomly selected in a given year, we calculate an ensemble mean. We then determine the ACC between the MR-30 bootstrap and ERA-20C.

195 3 Results

3.1 Characteristics of the observed link: temporal and spatial variability

The spatial pattern of the summer EA in its positive phase is characterised by large-scale cyclonic conditions, except at the anticyclonic centre of action located south of Iceland and west of the British Isles (Fig.1a). A typical surface climate imprint of the summer EA in positive phase ~~shows~~ correlates with below-average temperatures in continental Europe (Fig.1c) and
200 below-average precipitation in the British Isles and northwestern Europe (Fig.1d). As a first approach to evaluate the influence of spring extratropical SSTs in the summer EA, we use the SST index defined in Ossó et al. (2018). A ~~pearson correlation analysis suggests that the strength of the~~ Pearson correlation analysis reveals a time-dependent relationship between the SST index in spring AM SST index and the EA index in summer is time dependent in summer (Fig.1f). ~~Considering a period of observations spanning e).~~ Over a span of 101 years (1908-2008), this relationship is ~~appears~~ weak ($r = 0.2$, ~~Fig.1e~~ 0.22, $p <$
205 0.05). However, ~~this relationship changes over time: considering only the latest~~ examining the most recent 51 years (1958-2008)

~~correlation reaches significant~~ shows a doubling of correlation values ($r = 0.5043$, $p < 0.05$). Furthermore, focusing on the latest 30 years (the period analysed in Ossó et al. (2018)) results in correlation values increasing even further to 0.60 ($p < 0.05$). The temporal variability of this relationship is well illustrated for correlations calculated using a 20-year running window, which shows a reversal in the sign of correlations starting from 1945, and highlights an increase in the strength beyond 1958 (Fig.1f). This analysis suggests that the spring ~~SST-NA-SST~~ - summer EA relationship is nonstationary. Hence, we distinguish the following three periods to scope the remaining analysis: i) *early period*: 1908 - 1957; ii) *late period*: 1958-2008, and iii) *full period*: 1908-2008.

We assess the spatial features of the SST index influence on the summer atmospheric circulation in the different periods to further explore the variability of the spring ~~SST-NA-SST~~ - summer EA relationship. We analyse bimonthly averages, i.e. April-May (AM) ~~SST-NA-SST~~ and July-August (JA) SLP means, to observe the seasonal evolution of anomalies. Correlation maps show distinct patterns in early and late periods. We find significant correlations between the precursor SST index and ~~the summer sea level pressure (SLP) - summer SLP~~ over a region in the North Atlantic which reasonably coincides with the location of the EA teleconnection centre of action during the late period (Fig.2b, e-f-Fig.1a). The location of this region seems to oscillate about 45°N, remaining south of this latitude in the early period (Fig.2a), while located northwards in the late one (Fig.2b). Surrounding this high correlation region, the sign of correlations is opposite between early ~~and late (Fig.2a) and late (Fig.2b)~~ periods. We find similar results using March-April (MA) ~~SST-NA-SST~~ means, only in weaker strength ~~-(not shown)~~.

Regression maps further suggest that spring ~~SST-NA-SST~~ anomalies persist into summer and then influence atmospheric circulation (Fig.2d-fg-i). Positive values of the AM SST index in spring are associated with warm summer anomalies east of Newfoundland and cool anomalies west of Iberia, leading to concomitant anticyclonic conditions in the ocean located south of Greenland. In the late period (Fig.2h), these anticyclonic conditions coincide specifically with the position of the EA centre of action, whereas this association is absent in the early period.

Moreover, we test whether the SST index influences JA T2m via the EA. We find significant correlations between the AM SST index and JA T2m, showing a similar pattern of significant positive correlations west of the British Isles, as in Fig.1b-c corresponding to JA EA - T2m. We find that correlations between AM SST index and JA T2m show distinct patterns between early and late periods (Fig.2g-hd,e). A positive phase of the SST index in spring precedes a positive phase of the summer EA (e.g. Fig.2eh), which in turn can be associated with below-average temperatures, primarily over central Europe. ~~For To~~ further investigate this relationship, we calculate a $T2m_{CE}$ index, defined as the average summer ~~air temperatures T2m~~ over the central European region 46°N-55°N; 11°E-34°E, represented by the red box in Fig.2if. In summary, this analysis reveals that spring extratropical oceanic forcing of the summer atmospheric circulation has a marked temporal and spatial variability over the 20th century, only projecting onto the EA pattern over the late period. This variability might pose a constraint on the predictive skill of European summer climate based on spring extratropical ~~SST-NA-SST~~ during certain periods of time.

3.2 Investigating causality

To further test the robustness of the SST-EA relationship in ERA-20C, we evaluate whether spring extratropical SSTs and summer EA are conditionally dependent. Specifically, we test the hypothesis that spring ~~SST-NA-SST~~ is a causal driver for




Figure 1. Variability and linear relationships of EA in ERA-20C. a) Positive phase of the EA teleconnection, defined as the second EOF of July-August (JA) SLP. b) Regions used to calculate the SST-NA-SST and SLP indices proposed in Ossó et al. (2018). c) Pointwise correlation of EA index with concurrent July-August JA anomalies of 2-metre air temperatures in the full period (1908-2008). d) Same as c), for July-August JA anomalies of total precipitation. e) Time series of April-May (AM) SSTA (blue) and JA EA (grey) indices in ERA-20C for 1908-2008, smoothed by a 3-year running mean. f) Running-correlation between AM SSTA and JA EA indices for a 20-year window. Coloured markers indicate significant correlations at the 95% confidence interval, illustrated by dashed lines.




Figure 2. Distinct spatial characteristics of the spring SST influence on the summer circulation over the 20th century (for ERA-20C). **Top row (a-c)** shows point-wise correlation coefficients for the April-May SST index and July-August SLP means considering **early** (1908-1957, **left**), **late** (1958-2008, **middle**) and **full** (1908-2008), **right** column). **Middle-Top row (d-f)** shows linear regression maps of July-August SST anomalies (shading) and SLP (contours) against point-wise correlation coefficients for the precursor April-May SST index (normalised by the standard deviation σ). Contour interval is $0.2 \text{ hPa}\sigma^{-1}$ and July-August SLP. **Bottom-Middle row (g-i)** shows point-wise correlation coefficients for the April-May SST index and July-August air temperature at 2 metre height. Stippling indicates correlations significant at the 95% confidence level, calculated with a Student's t-test. **Bottom row (g-i)** shows linear regression maps of July-August NA-SST anomalies (shading) and SLP (contours) against the precursor SST index (normalised by the SD). Contour interval is 0.2 hPaSD^{-1} . Box in Fig.2f illustrates the region used to calculate the $T2m_{CE}$ index, as described in the text.

240 the summer EA, thus excluding autocorrelation effects or common drivers which could lead to spurious links. **We perform a causality analysis using the PC-MCI algorithm (Runge et al., 2015).**

First, we build one CEN for each of the three investigated periods in ERA-20C. Besides the EA and SST indices, we include two additional indices in the CEN. The first is the SLP index, defined in Ossó et al. (2018) and illustrated by the blue box in Fig.1b. **We thus** ~~Thus, we~~ test whether differences between early and late periods (e.f. Sec.3.1) are reflected in distinct timing or strength among the EA and SLP indices with SST. The second index concerns summer air temperatures averaged over the region represented by the red box in Fig.2f ($T2m_{CE}$), which shows significant **correlations** ~~anticorrelations~~ with SST. We test whether the spring SST index causally drives changes in **air temperature** ~~summer T2m~~ over central Europe and under which circumstances this holds true. **Our** ~~Therefore, our~~ CEN analysis focuses on $\tau_{min} = 3$ ~~and months~~ and $\tau_{max} = 4$ months ~~lag only~~, **which for simplicity we refer to as 3 and 4-month lags.**

245




Figure 3. Causal effect network analysis for the late period (1958-2008) in ERA-20C. Causal graphs between a) causal-graph-between-SST index, EA teleconnection, SLP index and T2m_{CE} (left) and b) SST index, EA teleconnection and schematic with illustration of the causal pathways in the Euro-Atlantic region SLP index only. The strength and direction of the causal links is given by the β -coefficient and is represented by the arrows, whereas the auto-correlation path coefficient is represented for each variable by the respective circle colour. The numbers over each arrow represent the time tag (in months) when the strongest causal link between each variable pair is detected. b-ec-f Sensitivity of the causal links shown as the PDF of β -coefficients calculated for a random sample selection of 45 years, iterated 500 times, between the variables: SST and SLP indices at lag 3 (bc) and lag 4 (ed), and SST index and EA at lag 3 (de) and lag 4 (ef). Only causal links with p-value < 0.1 are shown. Red lines show the correspondent β -coefficients represented in (a).

250 Over the late period, we confirm that the spring SST index is a causal driver for both the summer EA \rightarrow and the summer SLP index, at distinct time lags (Fig.3a). The strength of the causal link is expressed by the standardised regression coefficient, denoted β -coefficient in CEN. At a 4-month lag, we find $\beta_{SST \rightarrow EA} \approx 0.22$, $\beta_{SST \rightarrow EA} \approx 0.22$, which means that a change of 1 standard deviation (SD) in the March-April SST index leads to a change of 0.22 SD in July-August EA. We find a causal link of similar strength at a 3-month lag $\beta_{SST \rightarrow SLP} \approx 0.21$, $\beta_{SST \rightarrow SLP} \approx 0.21$, between April-May SST index and July-August EA, as well as -0.2 $\beta_{SST \rightarrow T2m_{CE}} \approx -0.2$ between April-May SST index and July-August T2m_{CE}. Although we speculate that the link $SST \rightarrow T2m_{CE} \rightarrow SST \rightarrow T2m_{CE}$ is mediated via the summer EA, we are unable to confirm this mediation with a CEN analysis focusing on 3-4 months lag. We While PCMCI cannot handle the nonstationarity identified in the full period (e.g. Runge (2018)), we find no significant causal links when analysing the full or in the early period.

260 Next, we test the sensitivity of the detected causal links between spring SST index and summer SLP to slight differences in the analysed years. By removing 6 randomly selected years (12% of tested years in the late period) in each new CEN

over 500 iterations, we test whether the causal links are particularly subjected to interannual variability (Fig.3b-ec-f). We find high variability in the strength of the links $\beta_{SST \rightarrow EA}$ and $\beta_{SST \rightarrow SLP}$ (Fig.3b, ec, f), ranging from zero (i.e. no causal link) to 0.5, with median values corresponding to β -coefficients calculated in Fig.3a. This sensitivity in the causal link strength due to sampling suggests that the spring SST index - summer SLP relationship might be modulated by an external physical mechanism, i.e. an additional ~~actor-excluded-variable not included~~ from this CEN.

3.3 Does MPI-ESM reproduce the observed link?

We now test whether the causal links detected in ERA-20C during the late period can be reproduced by MR-30. As a first step, we compare the model ability to reproduce the temporal variability of the observed summer EA. We find that MPI-ESM ~~is overall able to reproduce~~ generally captures the range of variability ~~but shows different levels of skill in reproducing,~~ although its performance in replicating the summer EA ~~amongst the~~ varies across different simulation sets. (Fig.4a). Historical simulations show low agreement with ERA-20C ($r = 0.14$), whereas MR-30 initialised simulations tend to mostly encompass the observed variability (Fig.4c).

Next, we evaluate the model skill in reproducing the spring SST index - summer EA relationship. We find that the model shows limited skill, ~~with MR-30 capturing the temporal variability of the relationship in the early, but not particularly~~ in the late period (Figs.4b, 5). A comparison between correlation maps computed for the evaluated periods shows that while historical simulations do not show agreement in the spatial pattern of the spring SST - summer EA relationship against observations (Fig.2a-c), the MR-30 ensemble mean shows an improvement in reproducing the mechanism (Fig.5d-f). These results motivate us to assess whether the model is able to reproduce any of the observed causal links, or whether it shows different causal paths than those observed.

~~We build~~ The observed disparities between the model and observations, as highlighted in the spatial correlations and time series analyses depicted in Figs.4-5, prompt further investigation into the causal relationships within MR-30. To address this, we proceed to assess whether the model reproduces any of the observed causal links or presents alternative causal pathways. We construct three different CEN sets to evaluate, respectively, pi-Control, historical and initialised simulations with MR-30. The variables analysed in the CEN sets are ~~first~~-SST, EA and SLP indices and the time lag of interest is spring - summer (3 and 4 months lag). While no causal links are found in the historical simulations, we find opposite causal links than those in ERA-20C for the pi-Control simulation, suggesting an atmospheric forcing ~~into the ocean from EA into the extratropical North Atlantic~~ (e.g. $\beta_{EA \rightarrow SST} \approx 0.22$), but no detected causal influence from the ocean on the atmosphere (Fig.6c).

Analysing ~~Moving on to~~ the initialised simulations, we ~~first-exploit the full-leverage the entire~~ 30-member ensemble of MR-30 to ~~build a CEN for construct a comprehensive CEN spanning~~ the full period (1908-2008), ~~where each constructed timeseries thus comprises~~ resulting in each constructed time series comprising 3030 years. We find that MR-30 is able to reproduce a weakly positive SST index - EA link (i.e. $\beta_{SST \rightarrow EA|SLP} = 0.030.04$) at 3-month lag (Fig.6a), but not at 4-month lag as detected in ERA-20C during the late period, and in much weaker strength (i.e. $\beta_{SST \rightarrow EA|SLP} = 0.22$, ERA-20C). Moreover, we find a weak negative causal link from SST index to SLP index in the model (i.e. $\beta_{SST \rightarrow SLP|EA} = -0.02$), as opposed to observations (i.e. $\beta_{SST \rightarrow SLP|EA} = 0.21$, Fig.3b). This finding aligns with Fig.5d-f, which shows that the area




Figure 4. Model skill in reproducing summer EA and its link with spring SST. a) probability density functions (PDF) of the summer EA, b) running-correlation between SST and EA indices for a 20-year window, c) time series of the summer EA. Light grey colours in a) and e) represent individual ensemble members, and dashed grey line shows black represents ERA-20C, green represents the pi-Control historical simulation. b) Running-correlation between SST and EA indices for In a 20-year window, for ERA-20C (black line), the ensemble mean (dashed grey line) and shows the historical simulation (green line pi-Control and in b). Coloured the coloured markers indicate significant correlations at the 95% confidence interval, illustrated by the horizontal dashed lines.

295 of positive correlations in MR-30 is displaced southwestwards with respect to ERA-20C. No causal links from SST index to EA or SLP indices are found when analysing only the late period (1958-2008). Next, we therefore investigate the causal link sensitivity to the sample size and focus on 45-year long timeseries-time series covering the late period, allowing a direct comparison with the sensitivity analysis performed in ERA-20C (Fig.6b-eb-d).

3.4 Sensitivity analysis and impact on predictive skill

300 We perform a two-step sampling method in our sensitivity analysis with MR-30. First, 45-years are randomly selected in the late period (1958-2008). Second, one ensemble member amongst the full 30-member ensemble is randomly selected in every year. We iterate this process 2000 times, thus generating 2000 45-year-long timeseries-time series for each SST, EA and SLP variables. In each iteration, we build one CEN to analyse whether any causal associations-relationships are detected for the sampled SST, SLP and EA timeseries-time series (Sect.2.4). Our sensitivity results suggest that the model does-mostly-not
 305 predominantly fails to reproduce the observed links between SST index and EA or SLP indices (Fig.6b), showing only in very rare cases β -coefficients in the positive range as in ERA-20C (Fig.3).




Figure 5. Comparison of the spatial characteristics of the SST-SLP relationship over the 20th century (model against ERA-20C) in MPI-ESM-MR. Correlation maps show point-wise correlation coefficients for the April-May SST index and July-August SLP means considering *early* (1908-1957; a,d,g), *late* (1958-2008; b,e,h) and *full* periods (1908-2008; c,f,i), respectively. Top row shows results for the MPI-ESM-MR historical simulation, middle and bottom row for MPI-ESM-MR 30-member ensemble, and bottom row. The reader may refer to Figs. 2a-c for a comparison with ERA-20C.

We hypothesise that this MR-30 limitation in reproducing the causal links detected in ERA-20C might constrain the skillful prediction of European summers a season ahead. As a first test, we focus on two particular values of the β -coefficients, namely $\beta_1 = -0.28$ and $\beta_2 = 0.36$, corresponding to the link SST index \rightarrow SLP at 3-month lag illustrated by orange arrows in Fig. 6b.

310 In other words, we analyse two cases with strong causal link strength but in opposite signs, with β_2 lying in the observed ERA-20C range.

We perform a predictive skill assessment for each the MR-30 causal timeseries bootstrap ensemble respective to β_1 and β_2 against ERA-20C, checking whether the strength of the causal link has a fingerprint in the predictive skill of JA SLP (Sect. 2.5). We find a better agreement between model and reanalysis for β_2 than for β_1 , with significant anomaly correlation coefficients (ACC) particularly over the region where spring SST is significantly correlated to summer SLP in ERA-20C (e.g. Fig 2b). However, since positive causal links are only rarely present in MR-30, we are unable to identify a robust fingerprint in the predictive skill related to any of the links between SST and EA or SLP indices.

Nevertheless, identifying

3.5 Forecasts of opportunity: could causality help?

320 We aim to identify a robust fingerprint of spring SST-NA-SST on summer predictive skill could be an important step towards targeting forecasts of opportunity (Mariotti et al., 2020). The correlation analysis, which could potentially enhance targeted forecasting opportunities (Mariotti et al., 2020). Our correlation analysis, as depicted in Fig. 2 suggests that spring SST could




Figure 6. Causal effect networks for MPI-ESM. a) Causal graph-CEN between SST index, EA teleconnection and SLP index for the MPI-ESM-MR 30-member ensemble (MR-30) considering the full period. b) Sensitivity of the causal links between SST, SLP and EA indices at 3 and 4-month lags in the late period. Boxplots show β -coefficients calculated for a random selection of 45 years, sampling one random ensemble member amongst the 30-member set per year. This process is repeated 2000 times and only significant β -coefficients different from zero are shown (here denoted MR-30-causal-ensemble-p-value < 0.1). Orange "x" markers represent the β -coefficient calculated from ERA-20C (red lines in Fig.3). c) Same as (a) for a 1000-year long pi-Control simulation with MPI-ESM-MR. Only causal links with p-value < 0.1 are shown. d) Comparison of the impact on SLP predictive skill for 3-4 month in lead time-times of 3-4 months in MR-30 against ERA-20C for timeseries-time series showing opposite β -coefficient strengths: a MR-30 causal-timeseries-bootstrap ensemble with (left) $\beta_1 = -0.28$, and (right) $\beta_2 = 0.36$. Predictive skill is quantified with anomaly correlation coefficients for the late period. β_1 and β_2 are highlighted in (b) by orange arrows.




Figure 7. Spatial features of the causal influence of spring SST index on summer climate. a) observed causal links between SST index, EA teleconnection and $T2m_{CE}$ in the late period (1958-2008); b) respective causal map for 3-month lag, showing causal links between April-May SST index and July-August temperature in shading $\beta_{SST \rightarrow T}$ and April-May SST index and July-August SLP $\beta_{SST \rightarrow SLP}$ in contours. Black box highlights the region of strongest causal influence and represents the area used to calculate the T2m index denoted $T2m_{Ridge}$ in the text.

influence, indicates the potential influence of spring NA-SST on summer T2m variability over-across the Euro-Atlantic region in ERA-20C during the late period (Fig.7a). Therefore, we perform an additional causality. Thus, we conduct an additional causal analysis in ERA-20C to highlight where in pinpoint the regions within the T2m field a causal influence of spring SST is expected, and where a causal relationship with spring NA-SST is anticipated. We also explore whether this causal relationship could be used to investigate an effect on might impact the predictive skill of MR-30's predictive skill.

Spring extratropical SST causal associations and impact on MR-30 predictive skill. a) observed causal links between SST index, EA teleconnection and $T2m_{CE}$ in the late period (1958-2008); b) respective causal map for 3-month lag, showing causal links between April-May SST index and July-August temperature in shading $\beta_{SST \rightarrow T}$ and April-May SST index and July-August SLP $\beta_{SST \rightarrow SLP}$ in contours. Black box highlights the region of strongest causal influence and represents the area used to calculate the T2m index denoted $T2m_{Ridge}$ in the text. c) Sensitivity of CEN built with the SST, SLP and T2m indices for MR-30. Boxplots show β -coefficients calculated for a random selection of 45 years, sampling one random ensemble member amongst the 30-member set per year. This process is repeated 500 times and only β -coefficients different from zero are shown. Orange "x" markers represent the β -coefficient calculated for a CEN built with the SST, SLP and T2m indices from ERA-20C for the late period. Only causal links with p-value < 0.1 are shown. d) Comparison of the impact on summer surface climate predictive skill in MR-30 against ERA-20C for causal and non-causal MR-30 timeseries. Mean ACCs are shown for

July–August sea level pressure (SLP), 2-metre air temperature (T2m) and 500 hPa geopotential height (Z500), averaged over the region highlighted by the grey box. See the text for further description.

340 We compute a causal map (Di Capua et al., 2020b) that represents the β -coefficients calculated for the link between AM SST index and each grid point of JA T2m and SLP fields, i.e. $\beta_{SST \rightarrow T2m}$ and $\beta_{SST \rightarrow SLP}$ (Fig.7b, shading and contours, respectively). We find two causal regions of opposite signs. The first region shows negative causal links and is located in northwestern Europe, partly encompassing the area used to calculate the $T2m_{CE}$ index expressed in the causal graph in Fig.3a. This can be interpreted as an increase of 1 SD in the spring SST index (e.g. warming over subpolar, and cooling over subtropical
345 North Atlantic) causally driving a decrease of about 0.3 SD in the summer T2m field in northwestern Europe. The second region shows a positive causal influence on both T2m and SLP fields, reaching strong values above 0.5 for the T2m field. A black box illustrates this causal region (40°N – 55°N ; 15°W – 34°W), denoted *Ridge*, and used to calculate the index $T2m_{Ridge}$.

Targeting this causal region, we test the hypothesis that predictive skill of the summer surface climate in MR-30 might be higher for ~~timeseries~~ time series able to reproduce the causal link strength in ERA-20C ($\beta_{SST \rightarrow T2m_{Ridge}} > 0.5$), than for
350 those unable to reproduce the link ($\beta_{SST \rightarrow T2m_{Ridge}} = 0$). To test this hypothesis, we ~~perform a two-step sampling method~~ first perform bootstrapping and ensemble subsampling to generate 500 ~~timeseries~~ time series consisting of 45-years randomly selected in the ensemble space during the late period ~~—similarly to the analysis performed for SST, EA and SLP(Sect.2.4)~~. At 3-month lag, we find that MR-30 is able to reproduce a range of β -coefficients for SST index $\rightarrow T2m_{Ridge}$, encompassing the observed link 16% of the times (Fig.7e??a). That is, 16% of random combinations in the MR-30 ensemble space result
355 in ~~a MR-30 causal timeseries which represents~~ time series which represent a causal influence of the SST index in spring (April–May) onto the $T2m_{Ridge}$ in summer (JA).

Next, we evaluate whether the strength of this causal link is imprinted on MR-30’s skill in predicting summer SLP, T2m and Z500 for the *Ridge* region a season ahead. We quantify the predictive skill with ACC using ERA-20C as a reference, for two opposite cases in MR-30: i) ~~timeseries~~ time series showing strong β -coefficients lying in the range $0.6 < \beta < 0.8$ (denoted
360 "causal" in Fig.??) and ii) ~~timeseries~~ time series showing β -coefficients = 0 ~~, i.e. (denoted "non-causal")~~. We find 25 samples in i), and we therefore randomly select 25 samples in ii) to enable a direct comparison. We calculate the ACC for each of the total 50 samples, averaging over the Ridge region (Fig.7d??b). We find that a random selection in the ensemble space tends to show higher median and maximum values for the predictive skill of SLP, T2m and Z500 when MR-30 reproduces the causal link

365 The framework of *forecasts of opportunity* (Mariotti et al., 2020) in seasonal prediction has been increasingly explored to identify physical processes which lead to enhanced predictability and forecast skill. Such a strategy has been particularly useful for summer (Carvalho-Oliveira et al., 2022) and winter (Dobrynin et al., 2018) seasonal predictions in the European region, where predictive skill is limited. Here, we target the summer EA to understand how its seasonal predictability is influenced by spring North Atlantic SSTs using the causal inference-based tool CEN based on PCMCI algorithm.

370 Using ERA-20C, our CEN analysis confirms that the spring SST index proposed in Ossó et al. (2018) causally influences the variability of summer SLP in the Euro-Atlantic region with a 3-4 months delay during the late period (1958–2008). Specifically,

we find that a 1 SD change in the spring SST index first drives a 0.2 SD change in the summer SLP index at 3-month lag (e.g. March-April SST → June-July SLP index), and then drives a 0.2 SD change a month later in the summer EA (e.g. March-April SST → July-August EA, Fig.3a). While EA and SLP indices are highly correlated ($r = 0.82$), the position of the area used to calculate the SLP index (Fig.2c) only partly overlaps the EA centre of action (Fig.1a), which extends further northwest. We speculate that the northward migration of the North Atlantic jet stream during summer (e.g. Hallam et al. (2022)) could explain the delay of a month between the causal link of SST index and EA/SLP indices.

Besides extratropical SSTs, ENSO-related tropical forcing has been suggested to influence the summer EA over more recent decades (1979 - 2016, e.g. Wulff et al. (2017); O'Reilly et al. (2018)). As opposed to the mechanism proposed in Ossó et al. (2018), Wulff et al. (2017) suggested that the summer EA is forced by diabatic heating anomalies in the tropical Pacific and Caribbean, and it is characterised by an extratropical Rossby wave train with a centre of action west of the British Isles. The CEN analysis proposed in this paper could therefore be extended to include tropical SST predictors, thus testing how the causal links discussed here could be affected by the influence of additional drivers.

Our findings suggest that the causal links detected in ERA-20C are nonstationary during the 20th century, being present only in the late period (1958-2008). Nonstationarity in teleconnections has been reported by several studies (e.g. Woollings et al. (2015); Weisheimer et al. (2019)). In particular, Rieke et al. (2021) used a 700-year pre-industrial control run with MPI-ESM-LR to investigate the tropical link of the summer EA (Wulff et al., 2017) with a statistical model, and showed that the link had a nonstationary behaviour, being present in some multidecadal epochs but not in others. Detecting nonstationarity in the causal links discussed here has an important consequence for the application on predictive skill in seasonal forecasting, implying a limited use of such causal links to target forecasts of opportunity.

Yet, our ~~causality~~-causal analysis with CEN offers an alternative assessment of MPI-ESM-MR's performance, enabling a direct comparison of the causal links reproduced by the model with those detected in reanalysis. We find that the causal links between spring SST index and summer EA and T2m are absent in pi-Control and historical simulations, but appear in some 45-year-long ~~timeseries~~-time series sampled in the initialised ensemble MR-30, thus suggesting a role of initialisation (Fig.6). ~~We use a random ensemble subsampling to perform a predictive skill assessment conditioned to MR-30's performance in reproducing causal links between spring SST and both summer EA and $T2m_{\text{Ridge}}$. As a result, one ensemble member is randomly chosen among the 30-member per year. Alternatively, performing an ensemble subsampling to calculate an ensemble mean over a subset of ensemble members could provide a better analysis of MR-30's potential.~~ Nevertheless, our results suggest that MR-30's limited performance in reproducing these causal links, in particular between spring SST index and the summer EA, might explain its low skill in predicting summer seasonal European climate (e.g. Neddermann et al. (2018); Carvalho-Oliveira et al. (2022)).

5 Conclusions

We apply the causal inference-based tool CEN based on PCMCI algorithm to evaluate the influence of spring North Atlantic extratropical SSTs on the predictability of summer EA and its associated impact on surface climate at seasonal timescales. Our
405 main findings are:

- Analysing ERA-20C, we find that the observed relationship between spring SST index and summer EA is nonstationary during the 20th century, showing distinct spatial patterns between early (1902-1957) and late (1958-2008) periods. The estimated causal influence of spring SST index on summer EA is of $\beta \approx 0.2$, ~~meaning a 0.2 SD increase in EA when SST increases by 1 SD.~~
- 410 – We find that this relationship in ERA-20C is only causal over the late period. A sensitivity analysis of its strength during the late period shows high variability, suggesting that the presence or absence of specific years plays an important role in the quantification of the causal link. This implies that an external physical mechanism not included in our analysis might modulate the spring SST - summer EA causal link.
- In addition to summer EA, we find in ERA-20C that the spring SST index causally influences summer T2m ($\beta \approx -0.2$)
415 over a region in northwestern Europe, and the Ridge region located west of the British Isles ($\beta \approx 0.5$). This causal influence is possibly mediated by the EA.
- We find that pre-industrial and historical simulations of the MPI-ESM-MR do not reproduce the causal links detected in ERA-20C. In contrast, our CEN analysis with the full initialised ensemble MR-30 reveals a weakly weak positive causal link between spring SST index and summer EA ($\beta \approx 0.030.04$).
- 420 – However, for 45-year-long ~~timeseries~~ time series randomly sampled in MR-30, we find that the initialised ensemble is mostly unable to reproduce the spring SST index - summer EA link.
- In contrast, MR-30 shows a moderate performance in reproducing the spring SST index - summer $T2m_{Ridge}$ causal link. We find that MR-30 tends to show improved predictive skill for summer surface climate predictions over the Ridge region when the spring SST index - summer $T2m_{Ridge}$ causal link is correctly reproduced by the model.
- 425 In this analysis, we demonstrate that MPI-ESM-MR has limited performance in reproducing a causal link between spring SST NA-SST (SST index) and summer EA amongst uninitialised and initialised model datasets. Our causality analysis therefore sheds light on the limitations of this model in providing skillful seasonal predictions of summer climate, particularly over areas which undergo a significant EA influence. Exploring the causes behind the model’s deficiencies in this aspect—such as whether they stem from inadequacies in representing crucial coupled ocean-atmosphere feedbacks or other processes—will be
430 a significant focus for future research. Finally, our results for the initialised ensemble MR-30 show that ensemble members able to reproduce a causal link to spring SST have a potential for regional skill improvement. Our findings thereby illustrate how a causality framework could be used to target forecasts of opportunity, and highlight the importance of improving the representation of teleconnections in climate models.

Acknowledgements. The authors would like to thank Efi Rousi and Eduardo Zorita for insightful discussions at the start phase of this research.

435 The authors also thank the Climate Modelling group at Universität Hamburg and the Climate Extremes group at VU Amsterdam for helpful discussions. Model simulations were performed using the high-performance computer at the German Climate Computing Center (DKRZ). This research was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy—EXC 2037 'CLICCS—Climate, Climatic Change, and Society'—Project Number: 390683824, contribution to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg (J.B and L.F.B), and by the German Federal Ministry for Education and

440 Research of Germany (BMBF) via the JPI Climate/JPI Oceans project ROADMAP (grant no. 01LP2002B) (G.D.C. and R.V.D.). J.B. was supported by Copernicus Climate Change Service, funded by the EU, under contracts C3S-330, C3S2-370.

References

- Athanasiadis, P. J., Yeager, S., Kwon, Y.-O., Bellucci, A., Smith, D. W., and Tibaldi, S.: Decadal predictability of North Atlantic blocking and the NAO, *NPJ Climate and Atmospheric Science*, 3, 1–10, 2020.
- 445 Bastos, A., Janssens, I. A., Gouveia, C. M., Trigo, R. M., Ciais, P., Chevallier, F., Penuelas, J., Rödenbeck, C., Piao, S., Friedlingstein, P., et al.: European land CO₂ sink influenced by NAO and East-Atlantic Pattern coupling, *Nature communications*, 7, 1–9, 2016.
- Carvalho-Oliveira, J., Borchert, L. F., Zorita, E., and Baehr, J.: Self-organizing maps identify windows of opportunity for seasonal European summer predictions, *Frontiers in Climate*, 4, 2022.
- Cassou, C., Terray, L., Hurrell, J. W., and Deser, C.: North Atlantic winter climate regimes: Spatial asymmetry, stationarity with time, and 450 oceanic forcing, *Journal of Climate*, 17, 1055–1068, 2004.
- Cassou, C., Terray, L., and Phillips, A. S.: Tropical Atlantic influence on European heat waves, *Journal of climate*, 18, 2805–2811, 2005.
- Collins, M.: Climate predictability on interannual to decadal time scales: The initial value problem, *Climate dynamics*, 19, 671–692, 2002.
- Comas-Bru, L. and Hernández, A.: Reconciling North Atlantic climate modes: revised monthly indices for the East Atlantic and the Scandinavian patterns beyond the 20th century, *Earth System Science Data*, 10, 2329–2344, 2018.
- 455 Comas-Bru, L. and McDermott, F.: Impacts of the EA and SCA patterns on the European twentieth century NAO–winter climate relationship, *Quarterly Journal of the Royal Meteorological Society*, 140, 354–363, 2014.
- Di Capua, G., Kretschmer, M., Donner, R. V., van den Hurk, B., Vellore, R., Krishnan, R., and Coumou, D.: Tropical and mid-latitude teleconnections interacting with the Indian summer monsoon rainfall: a theory-guided causal effect network approach, *Earth System Dynamics*, 11, 17–34, 2020a.
- 460 Di Capua, G., Runge, J., Donner, R. V., van den Hurk, B., Turner, A. G., Vellore, R., Krishnan, R., and Coumou, D.: Dominant patterns of interaction between the tropics and mid-latitudes in boreal summer: causal relationships and the role of timescales, *Weather and Climate Dynamics*, 1, 519–539, 2020b.
- Dobrynin, M., Domeisen, D. I., Müller, W. A., Bell, L., Brune, S., Bunzel, F., Düsterhus, A., Fröhlich, K., Pohlmann, H., and Baehr, J.: Improved teleconnection-based dynamical seasonal predictions of boreal winter, *Geophysical Research Letters*, 45, 3605–3614, 2018.
- 465 Domeisen, D. I., Badin, G., and Koszalka, I. M.: How predictable are the Arctic and North Atlantic Oscillations? Exploring the variability and predictability of the Northern Hemisphere, *Journal of Climate*, 31, 997–1014, 2018.
- Duchez, A., Frajka-Williams, E., Josey, S. A., Evans, D. G., Grist, J. P., Marsh, R., McCarthy, G. D., Sinha, B., Berry, D. I., and Hirschi, J. J.: Drivers of exceptionally cold North Atlantic Ocean temperatures and their link to the 2015 European heat wave, *Environmental Research Letters*, 11, 074004, 2016.
- 470 Gastineau, G. and Frankignoul, C.: Influence of the North Atlantic SST variability on the atmospheric circulation during the twentieth century, *Journal of Climate*, 28, 1396–1416, 2015.
- Gorgetta, M., Jungclaus, J., Reick, C., Legutke, S., Brovkin, V., Crueger, T., Esch, M., Fieg, K., Glushak, K., Gayler, V., et al.: CMIP5 simulations of the Max Planck Institute for Meteorology (MPI-M) based on the MPI-ESM-LR model: The piControl experiment, served by ESGF, 2011.
- 475 Gorgetta, M. A., Jungclaus, J., Reick, C. H., Legutke, S., Bader, J., Böttinger, M., Brovkin, V., Crueger, T., Esch, M., Fieg, K., et al.: Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5, *Journal of Advances in Modeling Earth Systems*, 5, 572–597, 2013.

- Hallam, S., Josey, S. A., McCarthy, G. D., and Hirschi, J. J.-M.: A regional (land–ocean) comparison of the seasonal to decadal variability of the Northern Hemisphere jet stream 1871–2011, *Climate Dynamics*, pp. 1–22, 2022.
- 480 Jungclaus, J., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and Von Storch, J.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *Journal of Advances in Modeling Earth Systems*, 5, 422–446, 2013.
- Karmouche, S., Galytska, E., Runge, J., Meehl, G. A., Phillips, A. S., Weigel, K., and Eyring, V.: Regime-oriented causal model evaluation of Atlantic–Pacific teleconnections in CMIP6, *Earth System Dynamics*, 14, 309–344, 2023.
- 485 Klavans, J. M., Cane, M. A., Clement, A. C., and Murphy, L. N.: NAO predictability from external forcing in the late 20th century, *NPJ climate and atmospheric science*, 4, 1–8, 2021.
- Kretschmer, M., Coumou, D., Donges, J. F., and Runge, J.: Using causal effect networks to analyze different Arctic drivers of midlatitude winter circulation, *Journal of climate*, 29, 4069–4081, 2016.
- Kretschmer, M., Zappa, G., and Shepherd, T. G.: The role of Barents–Kara sea ice loss in projected polar vortex changes, *Weather and*
490 *Climate Dynamics*, 1, 715–730, 2020.
- Mariotti, A., Baggett, C., Barnes, E. A., Becker, E., Butler, A., Collins, D. C., Dirmeyer, P. A., Ferranti, L., Johnson, N. C., Jones, J., et al.: Windows of opportunity for skillful forecasts subseasonal to seasonal and beyond, *Bulletin of the American Meteorological Society*, 101, E608–E625, 2020.
- Mauritsen, T., Stevens, B., Roeckner, E., Crueger, T., Esch, M., Giorgetta, M., Haak, H., Jungclaus, J., Klocke, D., Matei, D., et al.: Tuning
495 the climate of a global model, *Journal of advances in modeling Earth systems*, 4, 2012.
- Mishra, N., Prodhomme, C., and Guemas, V.: Multi-model skill assessment of seasonal temperature and precipitation forecasts over Europe, *Climate Dynamics*, 52, 4207–4225, 2019.
- Moore, G., Renfrew, I., and Pickart, R. S.: Multidecadal mobility of the North Atlantic oscillation, *Journal of Climate*, 26, 2453–2466, 2013.
- Neddermann, N.-C., Müller, W. A., Dobrynin, M., Düsterhus, A., and Baehr, J.: Seasonal predictability of European summer climate re-
500 assessed, *Climate Dynamics*, pp. 1–18, 2018.
- Oliveira, J. C., Zorita, E., Koul, V., Ludwig, T., and Baehr, J.: Forecast opportunities for European summer climate ensemble predictions using Self-Organising Maps, in: *Proceedings of the 10th International Conference on Climate Informatics*, pp. 67–71, 2020.
- O’Reilly, C. H., Weisheimer, A., Woollings, T., Gray, L. J., and MacLeod, D.: The importance of stratospheric initial conditions for winter North Atlantic Oscillation predictability and implications for the signal-to-noise paradox, *Quarterly Journal of the Royal Meteorological*
505 *Society*, 145, 131–146, 2019.
- Ossó, A., Sutton, R., Shaffrey, L., and Dong, B.: Observational evidence of European summer weather patterns predictable from spring, *Proceedings of the National Academy of Sciences*, 115, 59–63, 2018.
- Ossó, A., Sutton, R., Shaffrey, L., and Dong, B.: Development, Amplification, and Decay of Atlantic/European Summer Weather Patterns Linked to Spring North Atlantic Sea Surface Temperatures, *Journal of Climate*, 33, 5939–5951, 2020.
- 510 O’Reilly, C. H., Woollings, T., Zanna, L., and Weisheimer, A.: The impact of tropical precipitation on summertime Euro-Atlantic circulation via a circumglobal wave train, *Journal of Climate*, 31, 6481–6504, 2018.
- Poli, P., Hersbach, H., Dee, D. P., Berrisford, P., Simmons, A. J., Vitart, F., Laloyaux, P., Tan, D. G., Peubey, C., Thépaut, J.-N., et al.: ERA-20C: An atmospheric reanalysis of the twentieth century, *Journal of Climate*, 29, 4083–4097, 2016.
- Rieke, O., Greatbatch, R. J., and Gollan, G.: Nonstationarity of the link between the Tropics and the summer East Atlantic pattern, *Atmospheric Science Letters*, p. e1026, 2021.
- 515

- Rousi, E., Selten, F., Rahmstorf, S., and Coumou, D.: Changes in North Atlantic atmospheric circulation in a warmer climate favor winter flooding and summer drought over Europe, *Journal of Climate*, 34, 2277–2295, 2021.
- Rousi, E., Kornhuber, K., Beobide-Arsuaga, G., Luo, F., and Coumou, D.: Accelerated western European heatwave trends linked to more-persistent double jets over Eurasia, *Nature communications*, 13, 1–11, 2022.
- 520 Runge, J.: Causal network reconstruction from time series: From theoretical assumptions to practical estimation, *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 28, 075 310, 2018.
- Runge, J., Petoukhov, V., and Kurths, J.: Quantifying the strength and delay of climatic interactions: The ambiguities of cross correlation and a novel measure based on graphical models, *Journal of climate*, 27, 720–739, 2014.
- Runge, J., Petoukhov, V., Donges, J. F., Hlinka, J., Jajcay, N., Vejmelka, M., Hartman, D., Marwan, N., Paluš, M., and Kurths, J.: Identifying
525 causal gateways and mediators in complex spatio-temporal systems, *Nature communications*, 6, 1–10, 2015.
- Runge, J., Nowack, P., Kretschmer, M., Flaxman, S., and Sejdinovic, D.: Detecting and quantifying causal associations in large nonlinear time series datasets, *Science Advances*, 5, eaau4996, 2019.
- Schuhen, N., Schaller, N., Bloomfield, H. C., Brayshaw, D. J., Lledó, L., Cionni, I., and Sillmann, J.: Predictive Skill of Teleconnection Patterns in Twentieth Century Seasonal Hindcasts and Their Relationship to Extreme Winter Temperatures in Europe, *Geophysical Research
530 Letters*, p. e2020GL092360, 2022.
- Siew, P. Y. F., Li, C., Sobolowski, S. P., and King, M. P.: Intermittency of Arctic–mid-latitude teleconnections: stratospheric pathway between autumn sea ice and the winter North Atlantic Oscillation, *Weather and Climate Dynamics*, 1, 261–275, 2020.
- Spirtes, P., Glymour, C. N., and Scheines, R.: *Causation, prediction, and search*, MIT press, 2000.
- Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., et al.: Atmospheric
535 component of the MPI-M Earth system model: ECHAM6, *Journal of Advances in Modeling Earth Systems*, 5, 146–172, 2013.
- Wallace, J. M. and Gutzler, D. S.: Teleconnections in the geopotential height field during the Northern Hemisphere winter, *Monthly weather review*, 109, 784–812, 1981.
- Weisheimer, A., Decremier, D., MacLeod, D., O’Reilly, C., Stockdale, T. N., Johnson, S., and Palmer, T. N.: How confident are predictability estimates of the winter North Atlantic Oscillation?, *Quarterly Journal of the Royal Meteorological Society*, 145, 140–159, 2019.
- 540 Woollings, T., Hannachi, A., and Hoskins, B.: Variability of the North Atlantic eddy-driven jet stream, *Quarterly Journal of the Royal Meteorological Society*, 136, 856–868, 2010.
- Woollings, T., Franzke, C., Hodson, D., Dong, B., Barnes, E. A., Raible, C., and Pinto, J.: Contrasting interannual and multidecadal NAO variability, *Climate Dynamics*, 45, 539–556, 2015.
- Wulff, C. O., Greatbatch, R. J., Domeisen, D. I., Gollan, G., and Hansen, F.: Tropical forcing of the summer East Atlantic pattern, *Geophysical
545 Research Letters*, 44, 11–166, 2017.