

The agricultural expansion in South America’s Dry Chaco: Regional hydroclimate effects

M. Agostina Bracalenti^{1,2}, Omar V. Müller^{1,2}, Miguel A. Lovino^{1,2}, E. Hugo Berbery³

¹Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Santa Fe, Argentina.

²Centro de Estudios de Variabilidad y Cambio Climático (CEVARCAM), Facultad de Ingeniería y Ciencias Hídricas (FICH), Universidad Nacional del Litoral (UNL), Santa Fe, Argentina.

³Earth System Science Interdisciplinary Center (ESSIC)/Cooperative Institute for Satellite Earth System Studies (CISESS), University of Maryland, College Park, MD, USA

Correspondence to: Omar V. Müller (ovmuller@unl.edu.ar)

Abstract. The Gran Chaco ecoregion is South America’s largest remaining continuous stretch of dry forest. It has experienced intensive deforestation, mainly in the western part known as Dry Chaco, resulting in the highest rate of dry forest loss globally between 2000 and 2012. The replacement of natural vegetation with other land uses modifies the surface’s biophysical properties, affecting heat and water fluxes and modifying the regional climate. This study examines land use and land cover changes (LULCCs) in Dry Chaco from 2001 to 2015, their effects on local and non-local climate, and explores the potential impacts of future agricultural expansion in the region. To this end, Weather Research and Forecasting (WRF) model simulations are performed for two scenarios: the first one evaluates the observed land cover changes between 2001 and 2015 that covered 8% of the total area of Dry Chaco; the second scenario assumes an intensive agricultural expansion within the Dry Chaco. ~~In both scenarios, deforestation processes lead to decreases in LAI, reductions in stomatal resistance, and increases in albedo, thus reducing the net surface radiation and, correspondingly, decreasing the turbulent fluxes, suggesting a decline in available energy in the boundary layer.~~ The result is an overall weakening of the water cycle in the Dry Chaco and, most prominently, ~~implying~~ a reduction in precipitation. A feedback loop develops since dry soil absorbs significantly less solar radiation than moist soil. Finally, the simulations suggest that the Dry Chaco would intensify its aridity, extending the drier and hotter conditions into the Humid Chaco.

1 Introduction

The expansion of the agricultural frontier in southern South American countries (Brazil, Paraguay, Argentina, and Uruguay) has been favoured by natural, technological, economic, and socio-cultural factors. Large areas of South America have experienced a significant increase in land use and land-cover changes (LULCCs), mostly of anthropogenic origin (Richards et al., 2012; De Sy et al. 2015, Stanimirova et al. 2022). The regional increase in annual average rainfall in arid regions (Barros et al. 2015), combined with biotechnological advances (Ribichich et al. 2020; Bulacio et al. 2023), strengthened the crops’ adaptability to adverse climate conditions and enhanced the availability of productive lands. Moreover, the availability of

Formatted: Font: (Default) Times New Roman, 10 pt, Font colour: Auto

Deleted: In both scenarios, deforestation processes lead to decreases in LAI, increases in albedo, and reductions in stomatal resistance, reducing the net surface radiation and, correspondingly, a decrease in turbulent fluxes suggesting a decline in available energy in the boundary layer.

Deleted: implies

Deleted: Gasparri and Grau, 2009

cheap land and labour, the lack of environmental regulations or their inadequate enforcement, and the increased food demand from new international markets in the last decades contributed to the agricultural expansion advancing on native vegetation

40 (Paruelo et al., 2005; Volante et al. 2016).

The Gran Chaco, comprised of the arid Dry Chaco in the west and Humid Chaco in the east, stands as the largest remaining continuous stretch of dry forest in South America (Portillo-Quintero and Sanchez-Azofeifa, 2010). This dry forest ecoregion combines all the previous factors that increase the crop productivity, leading to an intensive deforestation process,

45 predominantly in the Dry Chaco subregion (Fehlenberg et al., 2017; Mosciaro et al., 2022). Currently, only 9 % of Gran Chaco is protected (Nori, 2016), exposing it to the highest rate of dry forest loss worldwide and positioning the region as one of the largest global deforestation hotspots (Hansen et al., 2013). The Gran Chaco underwent deforestation of 78,000 km² between 2001 and 2012, with varying deforestation rates, from ~2,900 km² year⁻¹ during 2001-2002 to ~9,200 km² year⁻¹ during 2007-2008 (Fehlenberg et al., 2017). In particular, dry forest deforestation rates in the Argentinian part of Gran Chaco during the

50 2000s became up to three times higher than in the 1980s (Piquer-Rodríguez et al., 2015).

LULCCs affect the soil state and the overlying atmosphere through the influence of the soil conditions on the land-atmosphere interactions (Dirmeyer et al., 2000). This is particularly evident in southern South America, where the land-atmosphere coupling shows strong relationships between the water cycle components (Spennemann and Saulo, 2015; Ruscica et al., 2016; Martínez et al., 2016). The replacement of natural vegetation by pastures, crops, or other types of land use modifies biophysical properties related to vegetation (albedo, stomatal resistance, surface roughness, among others), altering the ecosystem functioning and the surface fluxes (Lee and Berbery, 2012; Müller et al., 2014; Baldi et al., 2015). For instance, when crops replace forests, groundwater uptake and transpiration are reduced due to the lack of deep roots in crops. This mechanism may induce the rise of the water table level, facilitating soil saturation, increasing surface runoff and, therefore, the probability of

55 occurrence of floods or waterlogging (Miguez-Macho et al., 2007a; Martínez et al., 2016).

In brief, the changes in soil conditions affect the energy and water fluxes on the surface, which in turn modify the atmospheric fluxes (e.g., moisture transport) (Lee and Berbery, 2012; Mahmood et al., 2014). Thus, LULCCs inevitably alter the local climate state, but they may also impact adjacent or remote areas due to changes in river flow or atmospheric dynamics

65 (Mahmood et al., 2010). A suitable way to comprehensively investigate the impact of LULCCs on the regional climate is by using regional climate model (RCM) simulations. RCMs have two components -the land surface and the atmosphere- that interact at each time step through surface fluxes, enabling feedback between the atmosphere and the soil, thus, uncovering the atmospheric response to LULCCs. Land cover types and their associated biophysical properties are usually prescribed in Land Surface Models (LSMs), the land component of RCMs. This feature hinders LSMs from simulating ongoing LULCCs

70 (Maertens et al., 2021). However, sensitivity experiments can be conducted by updating the land cover map to simulate the impact of observed LULCCs on the regional hydroclimate (e.g. Lal et al. 2021), or artificially modifying the land cover to

Deleted: Dros, 2004;

Deleted: by

Deleted: o

Deleted: The

Deleted: Gran Chaco ecoregion

Deleted: mainly

Deleted: western part known as

Deleted: Boletta et al., 2006; Zak et al., 2008

Deleted: The eastern part is known as Humid Chaco. The Gran Chaco is the largest remaining continuous stretch of dry forest in South America (Gasparri and Grau, 2009; Portillo-Quintero and Sanchez-Azofeifa, 2010; Fehlenberg et al., 2017). O

Deleted: it

Deleted: currently

Deleted: and consequently, it is one of the most threatened ecoregions worldwide (Kummerle, 2017). Due to agricultural expansion, the Gran Chaco is

Deleted: one of the largest global deforestation hotspots, reaching

Deleted: ; Volante et al. 2016

Deleted: ; Müller et al., 2021a

Deleted: s

Deleted: Jobbagy et al., 2008;

Deleted: A meta-analysis performed by Sirimarco et al. (2018) based on more than 100 scientific papers about land cover changes in different parts of the world found that replacing grasslands with forests (afforestation) may increase infiltration and evapotranspiration by up to 65 % and 30 %, respectively. Still, when crops replace pastures, water infiltration is significantly reduced without major changes in evapotranspiration.

Deleted: i

simulate possible scenarios of LULCCs and their potential impacts (e.g., Lee and Berbery 2012, Georgescu et al. 2013, Flanagan et al. 2021).

In this context, and as a follow-on of the previous studies, the questions that motivate this paper are: How do LULCCs in the Dry Chaco alter the land surface processes locally? And what are the local and non-local effects of LULCCs on the Gran Chaco? To answer the raised questions we adopt a comprehensive three-step methodology. First, we evaluate the dominant LULCC in Dry Chaco during the current century using state-of-the-art satellite information (the Moderate Resolution Imaging Spectroradiometer, MODIS). Second, we analyse the local effects of current LULCCs in Dry Chaco by assessing simulations performed for the same period but using MODIS land cover maps of different years with the Weather Research and Forecasting (WRF) model. Third, we investigate the remote effects of LULCCs in Humid Chaco by analysing WRF simulations that assume a progressive expansion of the agricultural frontier in Dry Chaco. In this way, this study considers both observed and realistic expansions of LULCCs, offering a nuanced understanding of the actual and potential impacts on regional climate. By considering more realistic scenarios, it is possible to hypothesize how agricultural expansion in one region influences the hydroclimate of another, which is crucial for effective regional land use planning on a topic prone to developing socio-ecological conflicts associated with LULCCs. This focus on uncovering the processes behind such remote effects is a relatively unexplored aspect of existing literature.

The paper is organized as follows: Section 2 describes the region of interest, the observational data, the experiments, and the evaluation datasets. Section 3 evaluates the performance of the WRF control simulations. Section 4 presents the analysis of the model simulation responses to different LULC scenarios. Finally, a discussion is provided in Section 5 and the concluding remarks are summarized in Section 6.

2 Data, model, and experiments

2.1 Study Region

2.1.1 Geographical features

The Gran Chaco is a natural region of about 1,100,000 km² stretching from northern Argentina to southeastern Bolivia and northwestern Paraguay (Fig. 1). In South America, it is the second-largest forest after the Amazon and the largest remaining continuous stretch of dry forest (Bucher and Huszar, 1999; Portillo-Quintero and Sánchez-Azofeifa, 2010). The Gran Chaco is characterised by a flat topography with elevations varying between 100 and 500 m. The region is part of the subtropical belt, with a latitudinal south-north thermal gradient with a mean annual temperature ranging from 19 °C to 24 °C (Garreaud et al., 2009; Almazroui et al., 2021). The region shows a pronounced longitudinal gradient in precipitation, dividing the Gran Chaco into two distinct subregions: the Dry Chaco and the Humid Chaco (as indicated by the borders in Fig. 1). The Dry Chaco is dominated by a semi-arid climate with annual precipitation ranging from 450 to 900 mm. The Humid Chaco presents a humid

Deleted: s

Deleted: ¶

This study addresses the following questions:

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Right: -0.99 cm, Space Before: 18 pt, After: 18 pt, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Deleted: And,

Deleted: A suitable way to comprehensively tackle these questions is using regional climate model (RCM) simulations. RCMs have two components - the land surface and the atmosphere - that interact at each time step through surface fluxes, enabling feedback between the atmosphere and the soil. This feature is key to uncovering the atmospheric response to LULCCs. Land cover types and their associated biophysical properties are usually prescribed in Land Surface Models (LSMs), the land component of RCMs. Then, sensitivity experiments can be conducted by updating the land cover map or artificially modifying the land cover to simulate potential scenarios of LULCCs.

To address the questions raised above, we

Deleted: in

Deleted: and

Deleted: and

Deleted: 3

Deleted: the

Deleted: detailed

Deleted: presented

Deleted: 4

Deleted: ,

Deleted: Bucher

Deleted: 1982.

Deleted: Almazroui et al., 19822021).

subtropical climate with annual precipitation of about 900–1200 mm (Garreaud et al., 2009; Almazroui et al., 2021). Note that the Gran Chaco region exhibits minimal sensitivity to large-scale phenomena such as the El Niño Southern Oscillation (ENSO) and the Madden Julian Oscillation (MJO). Extreme ENSO phases predominantly induce precipitation excesses or deficits in eastern SESA, with minimal influence observed in the study region (Cai et al., 2020). The MJO has non-significant impact on precipitation anomalies over the Gran Chaco region (Grimm et al., 2019).

The Gran Chaco is one of the most dramatic global deforestation hotspots, with the highest rate of dry forest loss in the world in the 2000-2012 period (Hansen et al., 2013), due to the expansion of soybean production and cattle ranching (Fehlenberg et al., 2017). The Humid Chaco has been intensively transformed into agriculture, while the Dry Chaco remained largely forested because agriculture methods depended on rain, making it unprofitable despite high soil fertility and flat terrain (Bucher and Huszar, 1999). This historical behaviour has changed in recent decades due to the westward shift of the isohyets in the Gran Chaco and the availability of climate-resistant seeds (Dros et al., 2004).

2.1.2 Land cover and its evolution in Dry Chaco

The dominant land cover in the Dry Chaco and its evolution from 2001 to 2015 are estimated using land use maps derived from the MODIS Land Cover Climate Modeling Grid Product MCD12C1. The product is provided as a global mosaic at 0.05° grid spacing and follows the International Geosphere–Biosphere Program (IGBP) classification.

The top land cover categories in the region, as observed in 2015, are savanna, woody savanna, forest, grassland, shrubland, and cropland (Fig. 2a). Figure 2b shows how the dominant land cover categories in Dry Chaco evolved from 2001 to 2015. For simplicity, some of the IGBP types were grouped into major land cover classes on this panel. For instance, forest includes evergreen broadleaf, deciduous broadleaf, and mixed forests; shrubland includes closed and open shrublands, while cropland includes a cropland and cropland/natural vegetation mosaic. The land cover evolution indicates that agricultural lands (cropland and grassland) increased mainly up to 2007. Between 2001 and 2015, agriculture coverage increased by 15.4 %, representing 2.6 % of the total area of the Dry Chaco. On the other hand, forests and woody savanna decreased by 1.5 % and 16.4 %, respectively, representing 4.3 % of the Dry Chaco area.

Figure 2c indicates that 18 % of the Dry Chaco area suffered LULCCs from 2001 to 2015. These changes are quantified per category in Fig. 2d. Most of these changes (52 %) involve transitioning from categories characterised by higher tree coverage to others with lower coverage, which could be associated with deforestation processes. According to Fig. 2d, they are:

- from woody savanna to savanna, 23,328 km²,
- from savanna to agriculture, 12,672 km² (40 % to cropland and 60 % to grassland),
- from woody savanna to agriculture, 10,368 km² (50 % to cropland and 50 % to grassland)
- from forest to savanna, 8,064 km²,

Deleted: ,

Deleted: Almazroui et al.

Deleted: ,

Deleted: (Bucher, 1982)

Deleted: It is important to n

Formatted: Font: (Default) Times New Roman, 10 pt, Font colour: Auto

Deleted: Cai et al. (2020) shows the insensitivity of Gran Chaco to ENSO phases, while Grimm (2019) demonstrates that the various MJO modes presents non-significant small anomalies in our region of interest. Regarding ENSO, Cai et al. (2020) shows the insensitivity of Gran Chaco to ENSO phases.

Deleted: ; Boletta et al., 2006; Zak et al., 2008

Deleted: evaluated

- from forest to woody savanna, 6,336 km²,
- from forest to agriculture, 6,192 km² (67 % to cropland and 33 % to grassland), and
- from open shrubland to grassland, 5,904 km².

Note that about 31 % of the area corresponds to changes called “others” that consist in changes that are not linked to

210 deforestation processes (e.g., from open shrubland to savanna or to barren, from grassland to cropland, among others).

Curiously, about 17 % of the changes are from savanna and woody savanna to deciduous broadleaf forests. This transformation between natural land covers in a short period (14 years) is unthinkable in the real world. The rapid and natural conversion to deciduous broadleaf forests contradicts established ecological dynamics and is likely an artifact of the computation algorithms of land cover maps (Liang and Gong, 2010; Cai et al., 2014). Following the terminology defined by Cai et al. (2014), these implausible rapid transformations are named “illogical” in Fig. 2c.

215

2.2 The WRF model and its configuration

The numerical simulations were performed with the WRF model in its Advanced Research WRF (ARW) version 3.9 (Skamarock et al., 2008). The WRF was run with a horizontal grid spacing of 12 km and 38 vertical levels over a domain that covers the southern part of South America, which includes the Gran Chaco (see Fig. 1). A lateral boundary relaxation zone

220 spanning ten grid-points was implemented, and spectral nudging within the domain was intentionally omitted to allow the

atmosphere more freedom in responding to surface forcing (Pohl and Crétat, 2014). The model physics configuration is summarised in Table 1. The physical parameterisations follow the selection of schemes and options suggested by Lee (2010) and Lee and Berbery (2012), who developed a comprehensive set of ten WRF simulations combining different surface layer schemes, atmospheric boundary layer schemes, cumulus parameterization schemes, and microphysical schemes to identify the configuration that exhibited the highest skill for southern South America. This optimal configuration was subsequently employed in Müller et al. (2014) to assess droughts and in Müller et al. (2016) to evaluate the model's performance in short-term forecasts. In both cases, the selected parameterisations were well-suited to represent the hydroclimate in South America.

225

The model was forced by the initial and 6-hr boundary conditions obtained from the National Centers for Environmental Prediction (NCEP) Climate Forecast System Version 2 (CFSv2, Saha et al., 2014).

230

The WRF atmospheric component is coupled with the Noah-MP LSM (Niu et al., 2011), which solves the surface energy and water balances linking the surface conditions with the atmosphere. Noah-MP is an improved version of the Noah LSM (Chen et al. 1996; Chen and Dudhia, 2001), including multiple parameterisation options for selected physical processes (see selected schemes in Table 2). Noah-MP comprises four soil layers with a thickness from top to bottom of 10, 30, 60 and 100 cm (2 m total depth), and includes representations of the root zone, vegetation categories, monthly vegetation fraction, soil hydraulic properties, among others. It simulates soil moisture, soil temperature, skin temperature, canopy water content, and the energy flux and water flux terms of the surface energy balance and surface water balance. Different options of schemes for various physical processes that are key in the soil-atmosphere interaction, are available in Noah-MP. These processes include dynamic

235

Deleted: This

Deleted: and may be an artefact of the computation algorithms (Liang and Gong, 2010; Cai et al., 2014). T

Deleted: changes

Deleted: as

Deleted: 1

Deleted: tested

Deleted: reported they minimise biases in simulations for South America. Moreover, this configuration demonstrated high performance in long-term simulations (Müller et al. 2014) and short-term weather forecasts (Müller et al. 2016) in southern South America....

Deleted: It has four soil layers with a corresponding thickness of 10, 30, 60, and 100 cm from top to bottom (2 m total depth)

Deleted: texture

255 vegetation; canopy interception; soil moisture factor controlling stomatal resistance, b Factor; runoff and groundwater; surface
exchange coefficient for heat; and radiation transfer. The vegetation and soil components are closely coupled and interact with
each other via complex energy, water, and biochemical processes.

Noah-MP's thermodynamics resolves energy budgets and processes by separating the canopy layer from the ground surface
using the semitile subgrid scheme. Within this scheme, shortwave radiation transfer considers gap probabilities across the
entire grid cell, avoiding the overlap of shadows. Longwave radiation, latent heat, sensible heat, and ground heat fluxes are
independently calculated over two tiles: a fractional vegetated area ($Fveg$) and a fractional bare ground area ($1 - Fveg$),
where $Fveg$ depends on leaf area index (LAI). Regarding the model hydrology, the multi-layer soil structure fed by
precipitation allows for the simulation of soil moisture dynamics (water movement and storage). In the vertical column, Noah-
MP considers processes such as infiltration, evaporation, transpiration, and groundwater recharge. It simulates the
redistribution of water within the soil profile, accounting for the interplay between precipitation input, soil moisture storage,
and water fluxes. The runoff in Noah-MP is constituted by surface and groundwater runoff. Surface runoff is mainly saturation-
excess, while groundwater runoff mainly depends on the depth of the water table. Further details on model thermodynamics
and hydrology are found in Chen and Dudhia 2001, Niu et al. 2011, and He et al. 2023.

Focusing on the representation of vegetation in Noah-MP, the LSM assigns a dominant land cover type to each grid point, and
this assignment remains constant over time (Li et al., 2013). In turn, each land cover is associated with a set of 15 biophysical
properties. The properties can either be values fixed on time or can vary seasonally or dynamically when vegetation dynamics
is activated. Our simulations enable vegetation dynamics, i.e., the model simulates changes in vegetation properties, such as
LAI, surface roughness, and other land surface characteristics, as they naturally evolve over the simulated period due to
seasonal changes and vegetation growth cycles. These dynamic properties allow the model to capture the seasonality of
vegetation and its impact on land-atmosphere interactions. By default, the model employs the land cover map derived from
MODIS, which classifies the land cover following the 21 categories proposed by the IGBP classification. Then, the land cover
changes in our experiments are implicitly imposed by the change of land cover map among the various ensembles. Noah-MP
simulates isolated columns, thus grid cells with changed land cover will exert a direct and immediate local impact on surface
fluxes, while surrounding grid cells will remain unchanged in the initial time-step. However, as Noah-MP is coupled to WRF,
the surface change generates horizontal gradients that give rise to advective processes, effectively facilitating nonlocal effects
of the LULCCs.

Deleted:). Further characteristics are listed in Table 2.

Formatted: Font: Times New Roman

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: Not Bold

2.3 Experimental design

285 Sensitivity experiments were performed to determine how LULCC in Dry Chaco may influence the regional hydroclimate.
Three sets of simulations (ensembles) were conducted for the same period (January 2014 to June 2016), using different land

cover maps (see Fig. 3). Each ensemble has four members with identical parameterisations, with the only difference being the initial conditions that are 24-hr apart, following the lagged average forecasting method (Hoffman and Kalnay, 1983).

The CONTROL ensemble employs the MODIS land cover map corresponding to 2015 (Fig. 2a). The PAST ensemble uses the MODIS land cover map for the year 2001 (Fig. 4a). The FUTURE ensemble assumes an intensive agricultural expansion within Dry Chaco (Fig. 4b), mimicking what could be expected in the future if current expansion trends are to continue in a global-low scenario. This scenario considers a strong global market opening with low state regulation of LUCC, where the conditions favour great agricultural expansion (Mosciaro et al. 2022).

The corresponding FUTURE land cover map is a modified version of the 2015 land cover map in which all crop/grassland areas are expanded through a dilation process (Gonzalez and Woods, 1993). Dilation is a morphological method commonly used in digital image processing to expand target areas by adding surrounding pixels to boundaries and filling gaps. In this study, dilation is first applied to the crop category by adding ten lines, and subsequently, dilation is applied to grassland pixels by the addition of five lines. In the resulting land cover map, croplands cover 29.4% while grasslands cover 19.1% of the Dry Chaco area, meaning that the areas occupied by crops and pastures have been expanded by 5 and 2.5 times, respectively. Based on the rate of change of the global-low scenario proposed by Mosciaro et al. (2022), which is $\sim 7800 \text{ km}^2 \text{ yr}^{-1}$, the time horizon of our FUTURE scenario is 2065.

The differences between the CONTROL and the PAST ensembles will be called the ‘observed LULCCs’ scenario (OBS_LULCC). This scenario is employed to assess the impact of the land cover changes that occurred since 2001 on the 2014-2016 climate of the Dry Chaco. The differences between the FUTURE and the CONTROL ensembles are referred to as a scenario of ‘agricultural intensification’ (AG_INT). This scenario is used to evaluate the response of the 2014-2016 climate had there been a preceding large agricultural expansion in Dry Chaco. It is well known that land cover changes have been extensive over the south of South America outside the Gran Chaco, so other effects may come into play affecting the climate of Humid Chaco (Salazar et al., 2015), however these other factors are not part of the current study.

Lastly, there are two important aspects related to the 30 months long (2.5 years) simulation period. First, the validation of the CONTROL ensemble involves a comparison of simulated and observed variables for the entire period. However, the evaluation of changes has a specific focus on the austral summer season (DJF), encompassing the 2014/2015 and 2015/2016 summers, when vegetation is highly active, and land-atmosphere interactions are intensified in the Gran Chaco ecoregion (Müller et al., 2021a). The first 11-months are dismissed from the analysis to avoid any spin-up issues and ensuring the stabilization of all variables. Second, the simulation period includes an El Niño event, developed between Sep 2014 and March 2016. However, Cai et al. (2020) demonstrate that historical El Niño events did not lead to significant precipitation and temperature anomalies over the Gran Chaco. Moreover, Vera and Osman (2018) reported that the impact of the El Niño 2015 event has been weakened

Deleted: .

Deleted: a dilation method

Deleted: is applied over agricultural grid cells within the Dry Chaco....

Deleted: This method expands all crop/grassland areas by adding surrounding grid cells to the boundaries of the agriculture areas.

Deleted: It implies an

Deleted: s

Deleted: pin-up period that is dismissed

Deleted: ,

by the Southern Annular Mode (SAM). Thus, the resulting differences among ensembles are directly attributed to the proposed changes in land cover.

2.4 Evaluation datasets

335 The performance of the CONTROL ensemble is evaluated by comparing simulated precipitation, soil moisture, and 2m
temperature against observational data sets, either from satellite sources or gridded observations. The choice of these variables
serves a deliberate purpose. Temperature and precipitation, being truly independent variables with relatively low uncertainty
due to multiple monitoring sources, are commonly used in climate model validation (e.g., Sánchez et al. 2015, Ortega et al.
2021, Lovino et al. 2021, among others). Soil moisture is included in the evaluation due to its crucial role in land cover
340 dynamics and land-atmosphere interactions, although we acknowledge the uncertainty in its estimates based on remotely
sensed signals.

Three different precipitation datasets are employed. First, the NCEP’s Climate Prediction Center (CPC) Unified Gauge-Based
Analysis of Global Daily Precipitation (Chen et al., 2008; Xie et al., 2010). This dataset consists of daily rain gauge
345 observations interpolated to a $0.5^\circ \times 0.5^\circ$ grid. The second dataset, also at 0.5° grid spacing, is the Climate Research Unit
(CRU) monthly precipitation dataset. The CRU TS v. 4.03 dataset is based on analysing more than 4000 meteorological stations
(Harris et al., 2020). Lastly, we use the monthly precipitation at 0.25° grid spacing from the European Centre for Medium-
Range Weather Forecasts (ECMWF) reanalysis version 5 (ERA5) (Hersbach et al., 2020). Note that the three datasets overlap
in their sources.

350 Soil moisture is assessed at two model layers: 0-10 cm and 0-100 cm. The top layer (10 cm) is compared against the Soil
Moisture Operational Products System (SMOPS; Liu et al., 2016). This product combines soil moisture retrievals from
multiple satellite sensors producing global soil moisture maps as a volumetric content near the surface (top 1-5 cm) at daily
intervals and at a 0.25° grid spacing (Liu et al., 2016). Observed and simulated near-surface soil moisture are compared in
355 terms of volumetric water content (m^3m^{-3}) to minimise the impact of the thickness differences. The root zone soil moisture
(100 cm), crucial for vegetation growth, is evaluated with the operational product H14-SM-DAS-2 developed at the
EUMETSAT Satellite Application Facility on Support to Operational Hydrology and Water Management (HSAF). This
product gives estimates of root zone soil moisture at daily intervals on a $25 \text{ km} \times 25 \text{ km}$ grid (Albergel et al., 2012).

360 Temperature is evaluated with two products: the CRU TS v. 4.03 global monthly surface air temperature data set and the ERA5
monthly temperature. They are provided at a grid spacing of 0.5° and 0.25° respectively.

Deleted: 2.4 Model evaluation

Deleted: .1

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Text 1

Formatted: Font colour: Text 1

Formatted: Font: (Default) Times New Roman, Font colour: Text 1

Formatted: Font: (Default) +Body (Times New Roman), Font colour: Text 1

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Deleted:

Formatted: Font: (Default) Times New Roman, 10 pt, Font colour: Auto

Deleted: three hourly precipitation in the WATCH Forcing Data methodology applied to ERA-Interim reanalysis data (WFDEI)

Deleted: Weedon

Deleted: 14

Formatted: Font: (Default) Times New Roman, 10 pt, Font colour: Auto

Deleted: WFDEI

Deleted: three hourly

Deleted: Both

3 Model evaluation

3.1 Precipitation

375 The spatial pattern of the observation-based precipitation for the simulation period presents maximum values in southern
Brazil, eastern Paraguay, and northeastern Argentina (Figure 5a), where the precipitation rate is ~ 150 mm month⁻¹. This
380 magnitude gradually decreases westward to less than ~ 25 mm month⁻¹ on top of the Andes Mountains and the desert of
Atacama. The mean WRF precipitation field (Fig. 5b) reveals a similar spatial structure to the observations but with weaker
values in the wet part, where the model simulates less rainfall (see Fig. 5c). On the other hand, WRF simulates wetter conditions
over the Altiplano (high plateau in Bolivia). Wet anomalies over mountainous regions are often attributed to model limitations
and uncertainties in gridded observations and satellite estimates (Adam et al., 2006; Beck et al., 2017; Müller et al., 2021).

Focusing on the Gran Chaco, the model simulations show almost no biases in the Dry Chaco and present dry biases of ~ 30 – 50
mm month⁻¹ in the Humid Chaco. The drier conditions arise from underestimated late summer and autumn rain, as shown in
385 Figure 5d. Conversely, WRF notably improves the estimation in winter (the dry season) and spring, showing almost the same
peaks as observations for all years. The correlation and the error estimates confirm the good performance of WRF in the Gran
Chaco. The monthly temporal correlation between the area-averaged observed and simulated precipitation is $r=0.93$ for the
Dry Chaco and $r=0.85$ for the Humid Chaco. The RMSEs are 21.6 mm month⁻¹ and 56.5 mm month⁻¹, respectively.
390 Interestingly, the ensemble spread reveals relevant uncertainty only during the first month of simulation, after which the
dispersion consistently remains in a narrow range, indicating that the model uncertainty is comparable or even smaller than
the observational uncertainty.

3.2 Soil Moisture

The SMOPS satellite estimate of near-surface soil moisture exhibits high values to the east and a decreasing gradient toward
the west (Fig. 6a), following the precipitation pattern. The simulated soil moisture content shows a remarkable spatial structure
395 and magnitude agreement in most of the domain, except over the Altiplano where the wet precipitation biases produce wet
soils (Fig. 6b). The resemblance is even more evident over the Gran Chaco (Fig. 6c). In terms of temporal evolution, the area-
averaged simulated soil moisture presents similarities with SMOPS ($r=0.61$ for Humid Chaco and $r=0.55$ for Dry Chaco) but
with a slight systematic overestimation (Fig. 6g).

400 The simulated root-zone soil moisture also presents a notable overall resemblance to the HSAF satellite estimates, with slight
dry biases towards the east (Fig. 6d–e). Focusing on the Gran Chaco, the area-averaged WRF soil moisture presents a similar
evolution to HSAF but with a systematic negative bias (Fig. 6g). This bias is explained by the differences in Humid Chaco,
where RMSE reaches 0.091 m³m⁻³, likely in response to the underestimated precipitation (Fig. 5d). Instead, the Dry Chaco
presents a strong spatial agreement with HSAF (RMSE= 0.035 m³m⁻³).

- Deleted: ¶
2.4
- Deleted: 2
- Deleted: day
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: 1
- Deleted: day
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: drier
- Deleted: b
- Deleted: 1
- Deleted: 2
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: day
- Formatted: Not Highlight
- Deleted: 3
- Deleted: 0
- Deleted: day
- Deleted: 1.9
- Deleted: day
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Formatted: Not Highlight
- Deleted: 2.4
- Deleted: 3
- Deleted: (Fig. 6d–e)
- Deleted: , likely derived from the dry precipitation biases sh... [1]
- Deleted: ,
- Deleted:
- Deleted: g)
- Deleted: , that can also
- Deleted: be attributed to precipitation biases
- Deleted: .
- Deleted: It is important to highlight that the performance is ... [2]
- Deleted: , but weak temporal correlation ($r=0.33$), while the ... [3]

440

Notably, the simulated time series of near-surface and root-zone soil moisture reveals minimal internal variability throughout the entire simulation period. This consistency aligns with the findings of Sörensson and Berbery (2015), who observed that the initialization of WRF/Noah-MP in wet months (January in our simulations) favours a rapid stabilization of soil conditions. The limited internal variability of precipitation also contributes to the small ensemble members dispersion.

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Deleted: 2.4

Deleted: 4

445 3.3 Temperature

The spatial distribution of the time-mean observed temperature shows a latitudinal gradient with maximum values over the central-northern domain decreasing toward the south (Fig. 7a). The warmer region has an average temperature above 25 °C that gradually decreases southward to less than 15 °C. The coldest region over the high-altitude Andes Mountains has average temperatures below 5 °C.

450

Figure 7b shows that the simulated temperature's spatial pattern captures the observation-based products' main features in terms of distribution and magnitude (Fig. 7c), especially in flat areas (see Fig. 1). In high-altitude regions, the model produces cold biases. For the region of interest, WRF shows slight warm and cool biases (about ±1 °C) in parts of Dry Chaco and Humid Chaco, respectively. The simulated temperature evolution in the Gran Chaco closely aligns with observations ($r=0.96$ for Humid Chaco and $r=0.97$ for Dry Chaco), with slightly higher seasonal variability (Fig. 7d). As for precipitation and soil moisture, the simulated temperature presents insignificant sensitivity to initial conditions, reinforcing the robustness of the model simulations.

Deleted: Gran Chaco

Deleted: is remarkably similar to

Deleted: 7

Deleted: 8

455

4 Experiments' results

This section examines the two scenarios (OBS_LULCC and AG_INT) presented in Section 2.3. In both cases, simulations were carried out for 2014-2016 just changing the corresponding land cover maps, and the analysis is conducted during the austral summer months (DJF). Throughout the analysis, we will refer to local effects as those that pertain to the grid points within Dry Chaco that experienced LULCCs (purple grid cells in Figs. 4c-d). Non-local effects are those found at grid points that did not undergo any land cover change in the remaining Dry Chaco area (blank areas in Fig. 4c). Lastly, remote effects encompass those observed in the Humid Chaco area (blank areas within the blue limits in Fig. 4d). The subsections 4.1 and 4.2 focus on the description of the experiments' results, while plausible interpretation of the process behind the changes is offered in subsection 4.3.

Deleted: 3

Deleted: . In both scenarios

Deleted: but

Deleted: ,

Deleted: , when vegetation is highly active, and land-atmosphere interactions are intensified in the Gran Chaco ecoregion (Müller et al., 2021a)...

460

465

4.1 Scenario OBS_LULCC: Actual LULC changes from 2001 to 2015

Almost 20 % of the Dry Chaco's land cover changed from 2001 to 2015. Figures 8a-c show the changes in biophysical properties that are particularly sensitive to the observed LULCCs: leaf area index (LAI), albedo, and stomatal resistance. The

Deleted: 3

Deleted: (see Section 2.1.2).

485 changes in properties are spread over the small fragmented areas that underwent LULCCs. The decrease in LAI, increase in
albedo, and reduction in stomatal resistance in the Dry Chaco's centre-east, northeast, and west boundaries are associated with
deforestation processes (orange grid cells in Fig. 2c). On the other hand, other areas showing increases in LAI and stomatal
resistance, and decreases in albedo are not associated with deforestation as discussed in Section 2.1.2.

4.1.1 Effects on the energy budget

490 The changes in biophysical properties lead to the uneven distribution of positive and negative changes in the radiation fluxes
(Figs. 9a-c). The net radiation pattern is primarily influenced by variations in shortwave radiation, while changes in longwave
radiation are minimal due to negligible emissivity variations. On average, the areas that underwent local LULCCs experienced
a ~2 % reduction in net radiation, which aligns with the overall increase in albedo (Fig. 8b) and the consequent enhancement
of outgoing shortwave radiation. In the remaining areas of Dry Chaco, where land cover was not changed, positive and negative
495 changes up to $\pm 10 \text{ W m}^{-2}$ balance each other, resulting in a near-zero areal average.

The changes in net radiation alter the main components of the energy balance, i.e., the sensible heat flux and the latent heat
flux. The spatial distributions of sensible and latent heat fluxes play important roles in determining near-surface temperature.
Figures 9d-e reveal that changes in sensible heat and latent heat have a similar spatial pattern os but with opposite signs. ▼ As
500 with the radiation terms, the changes are of the order of $\pm 10 \text{ W m}^{-2}$. Furthermore, Fig. 9f shows that the deforested areas in the
northern and central eastern zones of Dry Chaco experience warming, while the southern area exhibits cooling. Both of these
changes can be attributed to variations in sensible heat flux. On average, the changes in net radiation lead to an average decrease
of -3.5 % in sensible heat at the local level and 0.6 % in the remaining Dry Chaco (Fig. 9e). Latent heat exhibits minor or
negligible changes over the whole Dry Chaco (Fig. 9d). Despite the presence of locally strong signals (up to $\pm 0.4 \text{ }^{\circ}\text{C}$), the
505 spatially averaged temperature changes present a slight rise due to the compensatory warming and cooling effects.

4.1.2 Effects on the hydrological response

The changes in biophysical properties lead to overall drier conditions in Dry Chaco, with precipitation decreasing by about -
1.7 % on average. However, there is a significant spatial heterogeneity with precipitation changes ranging from -29 mm month⁻¹
510 to +35 mm month⁻¹, even in areas that conserved the land cover (Fig. 10a). Soil moisture (Fig. 10b) follows the spatial pattern
of precipitation, with net negative changes in both local (-2.6 %) and non-local (-0.7 %) areas. Evapotranspiration also follows
the patterns of precipitation and soil moisture, although the net changes are negligible (see inset in Fig 10c). Note that stomatal
resistance is strongly reduced in deforested areas, which would facilitate plant transpiration and consequently higher
evapotranspiration. Our interpretation is that plants cannot take advantage of the open stomata due to the dry soils, given that
Dry Chaco is a region where soil moisture availability limits the evapotranspiration regime. The reduced precipitation and
515 drier soils end up affecting the total runoff, which presents a net reduction in the Dry Chaco of -2.1 % in local areas and -5.5

Deleted: 3

Deleted: for changes in sensible heat and latent heat

Deleted: 3

Deleted: water cycle

520 % in non-local areas (Fig. 10d). These results suggest that the observed LULCCs in just 14 years can weaken the hydrological cycle in Dry Chaco, leading this arid region to even drier summers.

Deleted: water

Deleted: , eventually increasing its exposure and vulnerability to extremes...

Deleted: 3

4.2 Scenario AG_INT: Intensive expansion of the agricultural land

The changes in surface properties in the OBS_LULCC scenario have a granular structure that illustrates the realistic ways deforestation progresses. Unlike the OBS_LULCC scenario, the AG_INT scenario shows changes in LAI, albedo, and stomatal resistance that exhibit spatial continuity, consistent with the assumed idealised LULCCs. In this AG_INT scenario, the expansion of crops and grasslands covers 64 % of the Dry Chaco. Figs. 11a-c illustrate that replacing native vegetation with grasslands and croplands leads to a general reduction in LAI and stomatal resistance, as well as an increase in albedo. Interestingly, the anthropisation of certain native covers results in “greener” conditions, in line with the prescribed values of the biophysical properties applied by the LSM. For instance, Noah-MP prescribes a higher LAI for grassland (1.7) than open shrubland (1.6).

Deleted: 3

4.2.1 Effects on the energy budget

The intensive expansion of agriculture simulated in Dry Chaco resulted in an overall decrease in net total radiation, which is reduced by -9.5 % (Fig. 12c). This reduction is primarily due to changes in shortwave radiation (Fig. 12a). Areas where LULCCs are imposed present higher albedo (Fig. 11b), reducing the net shortwave radiation by -15.6 % (Fig. 12a) due to the increased outgoing shortwave radiation. Changes in LAI, longwave radiation, and near-surface temperature follow similar patterns (Figs. 11a, 12b, and 12f). In the southwest, where LAI is increased, there is a decrease in net longwave radiation and cooler near-surface temperature. Conversely, regions with reduced LAI in the rest of Dry Chaco result in increased net longwave radiation and warmer temperature. This can be explained by the reduced vegetation cover, which diminishes shading and leads to surface warming and increased outgoing longwave radiation. The opposite effect occurs in areas with higher LAI. Although area-averaged values at local level tend to be small due to compensating positive and negative changes, temperature changes can reach values of up to ± 0.6 °C. It is worth noting that the overall reduction in net total radiation is also evident in the generally weaker turbulent surface fluxes (Fig. 12d-e). Latent heat decays by -4.2 %, while sensible heat decreases by -11 % on average.

545 The agricultural expansion imposed in Dry Chaco has weak remote effects on the energy budget of Humid Chaco (Fig. 12). The net radiation presents a slight increase consistent with the overall rise in the net shortwave radiation (Fig. 12a-c). The spatial pattern of changes in latent heat and sensible heat are similar but of opposite signs, with latent heat decreasing by -1.1 % and sensible heat increasing by 2.7 % (Figs. 12d-e). The near-surface temperature shows a slight warming that can be with increased sensible heat (Fig. 12f).

555 **4.2.2 Effects on the hydrological response**

The extensive expansion of crops and grasslands in Dry Chaco results in generally drier summers in the Gran Chaco (Fig. 13). The areas that experienced direct LULCCs (local effects) present a decay in summer precipitation of -7 % on average, and -9.3 % in soil moisture. The overall drier soils cause a reduction in evapotranspiration of -4.2 % with a spatial distribution of changes similar to those observed for precipitation and soil moisture, except in the southeast of the Dry Chaco. Similarly, runoff is reduced by -5 % on average. While the average reductions in water balance components represent less than 10 % in magnitude, nearly half of the area with LULCCs exhibit significant negative changes of approximately -30 %. These substantial changes have the potential for a significant impact on the region, considering its arid nature.

The overall remote effects in the Humid Chaco region result in drier summers, similar to those observed in Dry Chaco but with lesser intensity (Fig. 13). The changes in precipitation indicate a general tendency towards increased dryness in most parts of the Humid Chaco, except for isolated areas experiencing significant increases (Fig. 13a). On average, summer precipitation decreases by -4 %, although there are areas with changes of up to ± 30 mm month⁻¹. These changes in precipitation are accompanied by corresponding patterns in soil moisture, which also show a slight average decrease of about -1.2 % (Fig. 13b). Furthermore, evapotranspiration experiences an average reduction of -1.1 %, with localised decreases of up to -8 % in specific parts of the Humid Chaco. The overall drier conditions contribute to a reduction in runoff by about -6.2 % on average. Interestingly, this reduction is greater in the Humid Chaco than the Dry Chaco (-4.5 %). In summary, our results suggest that the expansion of agriculture in Dry Chaco intensifies the aridity of this already dry region and extends drier conditions into the Humid Chaco.

575 **4.3 Process-based analysis**

The results of the experiments consistently demonstrate that agricultural expansion in the Dry Chaco region leads to drier and warmer summers, regardless of the magnitude of the agricultural expansion. It is always a challenge to elucidate the processes that explain a specific change in a given variable due to the several factors acting together, as shown in the Fig. 1 of Santanello et al. (2018). Thus, our goal is to identify the most relevant mechanisms that explain the effects of LULCCs in Gran Chaco hydroclimate, while acknowledging that other processes may also have certain impact on our results. Figure 14 presents a schematic diagram summarising our interpretation of the key processes involved.

The intensification of agriculture in the Dry Chaco region brings about significant variations in biophysical properties, particularly LAI and albedo. While stomatal resistance is also greatly reduced, which would facilitate plant transpiration, its impact remains limited due to the water-limited regime. Consequently, the main focus lies on the effect of reduced LAI, leading to increased surface temperature due to decreased vegetation radiation sheltering. This rise in temperature increases the outgoing longwave radiation, resulting in a decrease in net longwave radiation. Conversely, the increased albedo decreases the

Deleted: 3

Deleted: water cycle

Deleted: -

Deleted: 3

Deleted: te

Deleted:

net shortwave radiation by reflecting a greater portion of incoming shortwave radiation. These changes in the energy budget consequently lead to a reduction in net surface radiation, aligning with the feedback mechanisms proposed by Eltahir (1998), Seneviratne et al. (2010), and Santanello et al. (2018).

Deleted: lead

The changes in net surface radiation are counterbalanced by the sum of latent and sensible fluxes, with soil heat flux considered negligible over longer time periods. Thus, the decrease in net radiation is equivalent to a decrease in the combined latent and sensible heat fluxes. This total energy flux from the land surface into the atmosphere represents the moist static energy (MSE), characterising the overall energy within the atmospheric boundary layer. Consequently, a reduction in net radiation, and therefore, a decline in available energy at the land surface, can be associated with less energetic conditions in the boundary layer. This, in turn, influences the dynamics of the planetary boundary layer and diminishes the generation of convective precipitation (Eltahir and Pal, 1996; Eltahir, 1998). Our interpretation is that the reduced energy in the atmosphere combined with the reduced precipitable water (PW, not shown) contributes to the overall stabilization of the boundary layer, resulting in less convective rainfall. This decrease in precipitation subsequently contributes to a decline in soil moisture. This creates a feedback loop since dry soil absorbs significantly less solar radiation than moist soil.

Deleted: Moreover, as suggested by Williams and Renno (1993), an increase in the planet boundary layer MSE should result in higher Convective Available Potential Energy (CAPE) directly linked to storm development. In our simulations, we observe the opposite scenario: reduced MSE implies lower CAPE, which can lead to reduced convective rainfall.

5 Discussion

Deleted: 4

The significant deforestation in the Dry Chaco, driven by agricultural expansion, is widely recognized and several studies focused on its causes and environmental consequences. This study goes beyond by exploring how the dominant LULCCs in the Dry Chaco from 2001 to 2015 impact on the regional hydroclimate, and the potential non-local effects that may develop in Humid Chaco if the intensive agricultural expansion within Dry Chaco continues. To this end, three ensembles of four simulations using different land cover maps were conducted using the WRF model from January 2014 to June 2016. While longer simulations could offer understanding under diverse large-scale atmospheric conditions, the length of these high-resolution simulations (determined by the computational capacity) provides valuable insights for assessing the land-atmosphere processes, which occur on daily to monthly time scales. The first ensemble employed the MODIS land cover map corresponding to 2015 (CONTROL). The second one used the MODIS land cover map corresponding to 2001 (PAST). The third one employed a modified version of the 2015 land cover map where the existing crop and pasture areas in 2015 within Dry Chaco were expanded (FUTURE).

Deleted: and

Deleted: conclusions

Deleted: The Gran Chaco region has two dominant ecoregions: Dry Chaco and Humid Chaco. The first one, particularly in the last decades, has been experiencing an intensive deforestation process due to agricultural expansion with important environmental consequences. ...

Deleted: explores

Deleted: and their effect

Deleted:

Deleted: regional climate

Deleted: . Furthermore, we examine other

Deleted: (of crop and grassland)

Deleted: , using different land cover maps,

Deleted: sets

Deleted: (ensembles)

Formatted: Font: (Default) +Body (Times New Roman)

The performance of the CONTROL simulation was assessed by comparing simulated precipitation, near-surface and root zone soil moisture, and 2-m temperature with gridded observations or satellite estimates. Other variables (radiation, heat fluxes, runoff, etc) could be included in the evaluation. However, their derivation involves algorithms or models with inherent approximations and uncertainties, which cannot be verified by in-situ measurements given the lack of field observations of non-conventional variables in the Gran Chaco region. Thus, the validation of such variables would be reduced to a comparison

Deleted: may have estimates and

Deleted: is

655 between tow disparate estimates without the inclusion of ground truth. The evaluation of the selected variables demonstrates good skills of the WRF Model for the region of interest. Simulated precipitation shows a marked resemblance with observations in Gran Chaco, with minor negative biases in the Humid Chaco. Near-surface and root-zone soil moisture closely matches the products derived from remotely sensed data in the Gran Chaco, although, consistent with negative precipitation biases, drier biases are observed in the Humid Chaco. The correlation coefficients for near-surface temperature indicate a strong agreement with observations in the Gran Chaco, although with somewhat larger seasonal variability. These results reinforce the effectiveness of the WRF model and the selected parameterisations in simulating southern South America hydroclimate, aligning with the results of our prior research efforts (Lee and Berbery, 2012, Müller et al. 2014, Sörensson and Berbery 2015, Müller et al. 2016).

Deleted: good

660 The MODIS maps reveal that the dominant land cover categories in the Dry Chaco include savanna, woody savanna, forest, grassland, shrubland, and cropland. Over the period from 2001 to 2015, land cover changes were observed in 18% of the entire Chaco region. These changes can be categorised as deforestation (52 %), illogical (17 %), and other (31 %). Agricultural lands, specifically cropland and grassland, experienced a sustained increase in the region particularly until 2007. Land cover changes due to deforestation processes, primarily characterized by the loss of woodlands, have been observed along the central-east, northeast, and western boundaries of the Dry Chaco, consistent with the finding of recent studies (Maertens et al. 2021, Mosciaro et al. 2022). The changes were observed in small fragmented areas, with deforestation primarily associated with decreases in LAI, increases in albedo, and reduced stomatal resistance. These findings align with other studies (e.g., Jiang et al. 2021), which reported that deforestation leads to lower LAI and higher surface albedo. In other areas where different land cover changes occurred (from open shrubland to barren, from grassland to cropland, among others), some biophysical properties show opposite trends, resulting in a heterogeneous pattern of property changes.

Deleted: are

Deleted: found in t

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Deleted: leaf area index

Deleted: (LAI)

675 The simulations show that the heterogeneous LULCCs from 2001 to 2015 and the associated changes in biophysical properties led to uneven distribution of positive and negative changes in radiation fluxes, particularly affecting net radiation, sensible heat flux, and latent heat flux. While there were average reductions in net radiation and sensible heat at the local level, latent heat changes were minor. Spatially, changes in sensible heat influenced near-surface temperature, leading to warming in deforested areas and cooling in areas with other LULCCs. The changes in biophysical properties and in the energy budget also affected the water cycle, resulting in overall drier conditions in the Dry Chaco. Precipitation exhibited significant changes, with both decreases and increases observed in different areas. Soil moisture followed the pattern of precipitation, displaying net local negative changes. The changes in precipitation and soil moisture are reflected in the evapotranspiration and the total runoff, all experiencing net negative changes. The findings suggest that the observed land cover changes weakened the water cycle in the Dry Chaco, contributing to drier summers and increased vulnerability to extreme events in this arid region.

Deleted: (green grid cells in Fig. 2c)

At present, there is no indication that deforestation will slow down. Assuming a scenario of strong global market opening with minimal regulation (Mosciaro et al. 2022), we designed a set of simulations where agricultural expansion of the existing agricultural areas within Dry Chaco and evaluated the possible consequences for the regional climate. Consistent with the effects observed in the OBS LULCC scenario, a general decrease in LAI and stomatal resistance, and an increase in albedo was observed at the local level. The expansion of agriculture in the Dry Chaco region resulted in significant changes in the energy budget, with further reductions in net radiation and weakening of the turbulent surface fluxes. These changes had local effects on the water cycle, causing drier conditions with decreased precipitation, soil moisture, evapotranspiration, and runoff. The extensive expansion of crops and grasslands in the Dry Chaco led to generally drier summers in the simulations, particularly in areas that experienced land use and land cover changes, with significant reductions in precipitation, soil moisture, evapotranspiration, and runoff. In this context, the aridity of the already dry Dry Chaco region is further intensified, with potential implications for its ecosystems and water resources. Furthermore, the expansion of agriculture has remote effects on the Humid Chaco, extending the drier conditions into this region, although to a lesser extent than the Dry Chaco. These findings highlight the complex interactions between land use, land cover changes, and the energy and water cycles, emphasising the need for sustainable land management practices in the Chaco region to mitigate the impacts of agricultural expansion and preserve these ecosystems' ecological and hydrological integrity.

Lastly, we acknowledge the inherent uncertainties associated with models that may limit the full representation of complex physical processes related to deforestation. However, we highlight that climate models offer distinctive advantages, including the ability to maintain physical consistency within the Earth System, simulate a wide range of processes, and facilitate controlled experiments by modifying initial and boundary conditions. These features make climate models the only tool capable of exploring the intricate chain of processes affected by changes in initial and boundary conditions, an analysis that cannot be replicated using observed data or reanalysis. Moreover, the minimal internal variability of the WRF simulations, coupled with the high level of skill of the WRF model in simulating fundamental variables—a recognition echoed in prior studies—strengthens our confidence in its ability to capture underlying processes. Thus, our experiments provide a valid and unique approach to analyzing the sensitivity of land-atmosphere physical interactions to different land cover scenarios.

6. Concluding remarks

Our investigation demonstrates that (a) the WRF model exhibited strong performance, effectively simulating fundamental variables in the Gran Chaco, reinforcing its reliability for assessing land-atmosphere processes. Additionally, the remote sensed data shows that the predominant form of LULCCs in the Dry Chaco since 2000 involves the conversion of natural vegetation, primarily woodlands and forests, into agricultural patches in lands adjacent to previously deforested areas, typically along the central-east, northeast, and western boundaries of the Dry Chaco.

Formatted: Font: Times New Roman

Deleted: The possible consequences for the regional climate are explored with a n

Formatted: Font: Times New Roman

Formatted: Font: Times New Roman

Deleted: ew set of simulations using a land cover map that assumes an agricultural expansion of the existing crop and pasture areas over savannas, woody savannas, forests, and shrublands, within Dry Chaco.

Deleted: for the LULCCs associated with deforestation

Deleted: ¶

Deleted: . These changes had substantial negative impacts on the water balance components

Deleted: The

Deleted: was

Deleted: d

Deleted: . The Humid Chaco also experienced reduced precipitation, soil moisture, evapotranspiration, and runoff

Deleted:

Formatted: No underline

Formatted: Font: (Default) Times New Roman, Font colour: Auto

Deleted: our results indicate that the expansion of agriculture and corresponding deforestation processes alter the regional hydroclimate through local, non-local, and remote changes of different magnitudes, leading to austral summers with lower precipitation and drier and warmer soils. Although our simulations are subject to uncertainties, the results suggest that the current land-use practices over Dry Chaco may not be sustainable in the long-term.

Deleted: c

Deleted: sions

750 The observed LULCCs from 2001 to 2015 modifies the biophysical properties, affecting the energy and water budget in Dry Chaco summers. The simulations indicate an average reduction of net surface radiation and surface energy fluxes, which in turn, lead to a hydrological response of the water cycle with overall drier conditions. Although changes in hydroclimate variables are found in the entire Dry Chaco, they are larger on areas that undergo land cover changes.

755 Simulations assuming continued agricultural expansion in the Dry Chaco, indicate further reductions in net radiation and turbulent surface fluxes and drier conditions, with the noteworthy outcome of extending these impacts remotely into the Humid Chaco. Thus, the expansion of agriculture alters the regional hydroclimate through local, non-local, and remote changes of different magnitudes. Although our simulations are subject to uncertainties, the results suggest that the current land-use practices over Dry Chaco may have not be sustainable in the long-term, as it exposes an already arid and hot region to even drier and warmer conditions, increasing its vulnerability to extreme events.

760 **Data availability**

The observational datasets used in this study are freely available online. The MODIS MCD12C1 product is available at <https://lpdaac.usgs.gov/products/mcd12c1v006/>. The CPC dataset is available at <https://psl.noaa.gov/data/gridded/data.cpc.globalprecip.html>. The CRU TS v. 4.03 dataset is available at <https://dx.doi.org/10.5285/10d3e3640f004c578403419aac167d82>. WFDEI dataset is available at <https://rda.ucar.edu/datasets/ds314.2/dataaccess>, SMOPS dataset is available at <https://www.ospo.noaa.gov/Products/land/smops>, H14-SM-DAS-2 dataset can be required at <https://hsaf.meteoam.it/User/UserSupport>. Simulations are available upon request.

Author contribution

770 MAB, OVM, and EHB designed the study. MAB performed the simulations, computed the results, and prepared the original manuscript. All authors analysed and discussed the results and contributed to the subsequent versions of the paper. OVM, MAL, and EHB acquired funding and resources.

Competing interests

The authors declare that they have no conflict of interest.

Deleted: Our investigation yields the following concluding remarks:

Deleted: The WRF model exhibited strong performance, effectively simulating fundamental variables in the Gran Chaco, reinforcing its reliability for assessing land-atmosphere processes. The predominant form of LULCCs in the Dry Chaco since 2000 involves the conversion of natural vegetation, primarily woodlands and forests, into agricultural patches in lands adjacent to previously deforested areas, typically along the central-east, northeast, and western boundaries of the Dry Chaco. The observed LULCCs from 2001 to 2015 modifies the biophysical properties, affecting the energy and water budget in Dry Chaco summers. The simulations indicate an average reduction of net surface radiation and surface energy fluxes, which in turn, lead to a hydrological response of the water cycle with overall drier conditions. Although changes in hydroclimate variables are found in the entire Dry Chaco, they are larger on areas that undergo land cover changes. Simulations assuming continued agricultural expansion in the Dry Chaco, indicate further reductions in net radiation and turbulent surface fluxes and drier conditions, with the noteworthy outcome of extending these impacts remotely into the Humid Chaco. The expansion of agriculture and corresponding deforestation processes alter the regional hydroclimate through local, non-local, and remote changes of different magnitudes. Although our simulations are subject to uncertainties, the results suggest that the current land-use practices over Dry Chaco may have not be sustainable in the long-term, as it exposes an already arid and hot region to even drier and warmer conditions, increasing its vulnerability to extreme events.our results indicate that the expansion of agriculture and corresponding deforestation processes alter the regional hydroclimate through local, non-local, and remote changes of different magnitudes, leading to austral summers with lower precipitation and drier and warmer soils. Although our simulations are subject to uncertainties, the results suggest that the current land-use practices over Dry Chaco may not be sustainable in the long-term.

Acknowledgements

We thank Prof. José Paruelo for his comments on the MODIS classification of land covers. This research was supported by the projects CAI+D-2016-50020150100005LI from the UNL. OVM and MAL acknowledge further support from the grants PICT-2019-2019-03982 and PICT-2019-2019-00481 from the ANPCYT, PIP 11220200102257CO from the CONICET, and PEICID-2021-028 from the ASaCTeI. EHB was supported by NOAA grant NA19NES4320002 (Cooperative Institute for Satellite Earth System Studies -CISESS) at the University of Maryland/ESSIC. He also thanks the Fulbright Commission for facilitating the last stages of this research.

References

Albergel, C., De Rosnay, P., Gruhier, C., Muñoz-Sabater, J., Hasenauer, S., Isaksen, L., Kerr Y. and Wagner, W.: Evaluation of remotely sensed and modelled soil moisture products using global ground-based in situ observations, Remote Sens. Environ., 118, 215-226, doi:10.1016/j.rse.2011.11.017, 2012.

Almazroui, M., Ashfaq, M., Islam, M., Rashid, I., Kamil, S., Abid, M., O'Brien, E., Ismail, M., Reboita, M., Sörensson, A., Arias, P., Muniz Alves, L., Tippet, M., Saeed, S., Haarsma, R., Doblas-Reyes, F., Saeed, F., Kucharski, F., Nadeem, I., Silva-Vidal, Y., Rivera, J., Ehsan, M., Martínez-Castro, D., Muñoz, A., Ali, M., Coppola, E., Bamba Sylla, M.: Assessment of CMIP6 Performance and Projected Temperature and Precipitation Changes Over South America, Earth Syst. Environ., 5(2), 155-183, doi: 10.1007/s41748-021-00233-6, 2021.

Baldi, G., Houspanossian, J., Murray, F., Rosales, A., Rueda, C., and Jobbágy, E.: Cultivating the dry forests of South America: Diversity of land users and imprints on ecosystem functioning, J. Arid Environ., 123, 47-59, doi:10.1016/j.jaridenv.2014.05.027, 2015.

Barros, V. R., Boninsegna, J. A., Camilloni, I. A., Chidiak, M., Magrín, G. O., and Rusticucci, M.: Climate change in Argentina: trends, projections, impacts and adaptation, WIREs Clim. Change, 6(2), 151-169, doi:10.1002/wcc.316, 2015.

Bucher, E. H., and Huszar, P. C.: Sustainable management of the Gran Chaco of South America: ecological promise and economic constraints, J. Environ. Manage., 57(2), 99-108, doi:10.1006/jema.1999.0290, 1999.

Bulacio, E.M., Romagnoli, M., Otegui, M.E., Chan, R.L., and Portapila, M.: OSTRICH-CROPGRO multi-objective optimization methodology for calibration of the growing dynamics of a second-generation transgenic soybean tolerant to high temperatures and dry growing conditions, Agric. Syst., 205, 103583, doi:10.1016/j.agsy.2022.103583, 2023.

Cai, S., Liu, D., Sulla-Menashe, D., and Friedl, M. A.: Enhancing MODIS land cover product with a spatial-temporal modeling algorithm, Remote Sens. Environ., 147, 243-255, doi:10.1016/j.rse.2014.03.012, 2014.

Cai, W., McPhaden, M. J., Grimm, A. M., Rodrigues, R. R., Taschetto, A. S., Garreaud, R. D., Dewitte, B., Poveda, G., Ham, Y.-G., Santoso, A., Ng, B., Anderson, W., Wang, G., Geng, T., Jo, H.-S., Marengo, J. A., Alves, L. M., Osman, M.,

Deleted:	https://
Deleted:	.org/
Deleted:	Almazroui, M., Ashfaq, M., Islam, M.N., et al. ... [4]
Deleted:	.
Deleted:	https://
Deleted:	.org/
Formatted	... [5]
Formatted	... [7]
Formatted	... [8]
Formatted	... [9]
Formatted	... [6]
Formatted	... [10]
Formatted	... [11]
Formatted	... [12]
Formatted	... [13]
Formatted	... [14]
Formatted	... [15]
Deleted:	Boletta, P. E., Ravelo, A. C., Planchuelo, A. M. ... [16]
Deleted:	Bucher, E.H.: Chaco and Caatinga — South Am ... [17]
Deleted:	https://
Deleted:	.org/
Formatted	... [18]
Formatted	... [20]
Formatted	... [19]
Formatted	... [21]
Formatted	... [22]
Formatted	... [23]
Formatted	... [24]
Formatted	... [25]
Deleted:	https://doi.org/
Formatted	... [26]
Formatted	... [28]
Formatted	... [29]
Formatted	... [30]
Formatted	... [31]
Formatted	... [32]
Formatted	... [27]
Formatted	... [33]
Formatted	... [34]
Formatted	... [35]

Li, S., Karamperidou, C., Takahashi, K., and Vera, C.: Climate impacts of the El Niño–southern oscillation on South America, *Nat. Rev. Earth Environ.*, 1(4), 215–231, doi:10.1038/s43017-020-0040-3, 2020.

Chen, F., and Dudhia J.: Coupling an advanced land surface hydrology model with the Penn State–NCAR MM5 modeling system. Part I: Model implementation and sensitivity, *Mon. Weather. Rev.*, 129, 569–585, doi:10.1175/1520-0493(2001)129<0569:CAALSH>2.0.CO;2, 2001.

Chen, F., Mitchell, K., Schaake, J., Xue, Y., Pan, H. L., Koren, V., Duan, Q. Y., Ek, M. and Betts, A.: Modeling of land surface evaporation by four schemes and comparison with FIFE observations. *J. Geophys. Res-Atmos.*, 101(D3), 7251–7268, doi:10.1029/95JD02165, 1996.

De Sy, V., Herold, M., Achard, F., Beuchle, R., Clevers, J.G., Lindquist, E., and Verchot, L.: Land use patterns and related carbon losses following deforestation in South America. *Environ. Res. Lett.*, 10(12), 124004, doi:10.1088/1748-9326/10/12/124004, 2015.

Dirmeyer, P.: Using a global soil wetness dataset to improve seasonal climate simulation, *J. Climate*, 13, 2900–2922, doi:10.1175/1520-0442(2000)013<2900:UAGSWD>2.0.CO;2, 2000.

Dros, J. M.: Managing the Soy Boom: Two scenarios of soy production, Amsterdam, AID Environment, 63 pp., 2004.

Eltahir, E. A. B., and Pal, J. S.: Relationship between surface conditions and subsequent rainfall in convective storms, *J. Geophys. Res.*, 101(D21), 26237– 26245, https://despecially thankD01380, 1996.

Eltahir, E. A. B.: A Soil Moisture–Rainfall Feedback Mechanism: 1. Theory and observations, *Water Resour. Res.*, 34(4), 765– 776, doi:10.1029/97WR03499, 1998.

Fehlenberg, V., Baumann, M., Gasparri, N. I., Piquer-Rodriguez, M., Gavier-Pizarro, G., and Kuemmerle, T.: The role of soybean production as an underlying driver of deforestation in the South American Chaco, *Global Environ. Chang.*, 45, 24–34, doi:10.1016/j.gloenvcha.2017.05.001, 2017.

Garreaud, R.D., Vuille, M., Compagnucci, R. and Marengo, J.: Present-day south American climate. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 281(3–4), 180–195, doi:10.1016/j.palaeo.2007.10.032, 2009.

Gasparri, N. I., and Grau, H. R.: Deforestation and fragmentation of Chaco dry forest in NW Argentina (1972–2007), *Forest Ecol. Manag.*, 258(6), 913–921, doi:10.1016/j.foreco.2009.02.024, 2009.

Georgescu, M., Lobell, D. B., Field, C. B., and Mahalov, A.: Simulated hydroclimatic impacts of projected Brazilian sugarcane expansion, *Geophys. Res. Lett.*, 40(5), 972–977, doi:10.1002/grl.50206, 2013.

González, R. and Woods, R.: *Digital Image Processing*. Addison-Wesley Longman Publishing Co, Boston, United States, 503 pp., ISBN:978-0-201-11026-5, 1993.

Grimm, A. M.: Madden–Julian Oscillation impacts on South American summer monsoon season: precipitation anomalies, extreme events, teleconnections, and role in the MJO cycle. *Clim. Dyn.*, 53(1-2), 907–932, doi:10.1007/s00382-019-04622-6, 2019.

Formatted: Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Formatted: Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Formatted ... [36]

Formatted ... [37]

Deleted: https://doi.org/

Deleted:

Deleted: https://doi.org/

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Formatted: Font: 10 pt, Not Italic, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, Not Italic, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted ... [38]

Deleted: https://doi.org/

Deleted: https://doi.org/

Deleted: https://doi.org/

Deleted: raphy...Palaeoclimatol.ogy,...Palaeoecol.ogy... 281(3–

Deleted: ogy,

Deleted: ogy

Deleted: .

Deleted: https://doi.org/

Deleted: https://doi.org/

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, Not Italic, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US), Pattern: Clear

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: Font: 10 pt, English (US)

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Formatted: Font: 10 pt, Not Italic, English (US)

Formatted: Font: 10 pt, English (US)

Formatted ... [39]

Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thaus, D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O. and Townshend, J.: High-resolution global maps of 21st-century forest cover change, *Science*, 342(6160), 850-853, doi: 10.1126/science.1244693, 2013.

915 Harris, I., Osborn, T.J., Jones, P. and Lister, D.: Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset, *Scientific Data*, 7(109), 1–18, doi:10.1038/s41597-020-0453-3, 2020.

He, C., Valayamkunnath, P., Barlage, M., Chen, F., Gochis, D., Cabell, R., Schneider, T., Rasmussen, R., Niu, G.-Y., Yang, Z.-L., Niyogi, D., and Ek, M.: Modernizing the open-source community Noah with multi-parameterization options (Noah-MP) land surface model (version 5.0) with enhanced modularity, interoperability, and applicability. *Geosci. Model Dev.*, 16(17), 5131-5151, doi:10.5194/gmd-16-5131-2023, 2023.

920 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999-2049, doi:10.1002/qj.3803, 2020.

925 Hoffman, R. N., and Kalnay, E.: Lagged average forecasting, an alternative to Monte Carlo forecasting, *Tellus A*, 35(2), 100-118, doi:10.3402/tellusa.v35i2.11425, 1983.

Jiang, Y., Wang, G., Liu, W., Erfanian, A., Peng, Q., and Fu, R.: Modeled response of South American climate to three decades of deforestation, *J. of Climate*, 34(6), 2189-2203, doi:10.1175/JCLI-D-20-0380.1, 2021.

930 Lal, P., Shekhar, A., and Kumar, A.: Quantifying temperature and precipitation change caused by land cover change: a case study of India using the WRF model, *Front. Environ. Sci.*, 9, 766328, doi:10.3389/fenvs.2021.766328, 2021.

Lee, S.-J.: Impact of land surface vegetation change over the La Plata Basin on the regional climatic environment: A study using conventional land-cover/land-use and newly developed ecosystem functional types, Ph.D. dissertation, University of Maryland, <http://hdl.handle.net/1903/10831>, 153 pp, 2010.



935 Lee, S. J., and Berbery, E. H.: Land cover change effects on the climate of the La Plata Basin, *J. Hydrometeorol.*, 13 (1), 84-102, doi:10.1175/JHM-D-11-021.1, 2012.

Li, D., Bou-Zeid, E., Barlage, M., Chen, F., and Smith, J. A.: Development and evaluation of a mosaic approach in the WRF-Noah framework, *J. Geophys. Res. Atmos.*, 118(21):11–918, doi:10.1002/2013JD020657, 2013.

940 Liang L, and Gong, P.: An assessment of MODIS Collection 5 global land cover product for biological conservation studies. In: 2010 18th international conference on geoinformatics, Beijing, China, 18–20 June 2010, pp 1–6, doi:10.1109/GEOINFORMATICS.2010.5567991, 2010.

Liu, J., Zhan, X., Hain, C., Yin, J., Fang, L., Li, Z., and Zhao, L.: NOAA soil moisture operational product system (SMOPS) and its validations. In 2016 IEEE International Geoscience and Remote Sensing Symposium (IGARSS) (pp. 3477-3480). IEEE, doi:10.1109/IGARSS.2016.7729899, 2016.

945

- Deleted:**  González, R. and Woods, R.: Digital Image Processing. Addison-Wesley Longman Publishing Co, Boston, United States, 503 pp., ISBN:978-0-201-11026-5, 1993, 
- Deleted:** <https://doi.org/>
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: 10 pt, Not Italic, English (US)
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: (Default) Times New Roman, Font colour: Auto, English (US), Pattern: Clear
- Formatted:** Indent: Left: 0 cm, Hanging: 1 cm
- Formatted:** Font: (Default) Times New Roman, Font colour: Auto, English (US), Pattern: Clear
- Formatted:** Font: (Default) Times New Roman, Font colour: Auto, English (US), Pattern: Clear
- Deleted:** <https://doi.org/>
- Deleted:** <https://doi.org/>
- Formatted:** Font: 10 pt, Font colour: Auto, English (US), Not Highlight
- Formatted:** Font: 10 pt, Font colour: Auto, English (US), Not Highlight
- Formatted:** Font: 10 pt, Font colour: Auto, English (US), Not Highlight
- Formatted:** English (US)
- Formatted:** English (US)
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Indent: Left: 0 cm, Hanging: 1 cm
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: 10 pt, English (US)
- Deleted:** Jobbágy, E., Nosetto, M., Santoni, C. and Baldi, G.: El desafío ecohidrológico de las transiciones entre sistemas l... [40]
- Deleted:** <https://doi.org/>
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Indent: Left: 0 cm, Hanging: 1 cm
- Formatted:** Font: 10 pt, English (US)
- Formatted:** Font: 10 pt, Not Italic, English (US)
- Formatted:** Font: 10 pt, English (US)
- Deleted:** <https://doi.org/>
- Deleted:** <https://doi.org/>

975 [Lovino, M. A., Pierrestegui, M. J., Müller, O. V., Berbery, E. H., Müller, G.V., and Pasten, M.: Evaluation of historical CMIP6 model simulations and future projections of temperature and precipitation in Paraguay, *Clim. Change*, 164, 1-24, doi:10.1007/s10584-021-03012-4, 2021.](#)

980 [Flanagan, P.X., Mahmood, R., Sohl, T., Svoboda, M., Wardlow, B., Hayes, M., and Rappin, E.: Simulated Atmospheric Response to Four Projected Land-Use Land-Cover Change Scenarios for 2050 in the North-Central United States, *Earth Interact.*, 25\(1\), 177-194, doi:10.1175/EI-D-20-0019.1, 2021.](#)

985 [Maertens, M., De Lannoy, G. J. M., Apers, S., Kumar, S. V., and Mahanama, S. P. P.: Land surface modeling over the Dry Chaco: the impact of model structures, and soil, vegetation and land cover parameters, *Hydrol. Earth Syst. Sci.*, 25, 4099–4125, doi:10.5194/hess-25-4099-2021, 2021.](#)

Mahmood, R., Pielke Sr, R. A., Hubbard, K. G., Niyogi, D., Bonan, G., Lawrence, P., McNider, R., McAlpine, C., Etter, A., Gameda, S., Qian, B., Carleton, A., Beltran-Przekurat, A., Chase, T., Quintanar, A., Adegoke, J., Vezhapparambu, S., Conner, G., Asefi, S, Sertel, E., Legates, D., Wu, Y., Hale, R., Frauenfeld, O., Watts, A., Shepherd, M., Mitra, C., Anantharaj, V., Fall, S., Lund, R., Treviño, A., Blanken, P., Du, J., Chang, H-I., Leeper, R., Nair, U., Dobler, S., Deo, R., and Syktus, J.: Impacts of land use/land cover change on climate and future research priorities, *Bull. Am. Meteorol. Soc.*, 91(1), 37-46, doi:10.1175/2009BAMS2769.1, 2010.

990 Mahmood, R., Pielke, R.A., Sr., Hubbard, K.G., Niyogi, D., Dirmeyer, P.A., McAlpine, C., Carleton, A.M., Hale, R., Gameda, S., Beltrán-Przekurat, A., Baker, B., McNider, R., Legates, D.R., Shepherd, M., Du, J., Blanken, P.D., Frauenfeld, O.W., Nair, U.S. and Fall, S.: Land cover changes and their biogeophysical effects on climate, *Int. J. Climatol.*, 34(4), 929-953, doi:10.1002/joc.3736, 2014.

Martinez, J., Dominguez, F. and Miguez-Macho, G.: Impacts of a Groundwater Scheme on Hydroclimatological Conditions over Southern South America, *J. Hydrometeorol.*, 17(11), 2959-2978, doi:10.1175/JHM-D-16-0052.1, 2016.

995 Miguez-Macho, G., Fan, Y., Weaver, C., Walko, R. and Robock, A.: Incorporating water table dynamics in climate modeling: 1. Water table observations and equilibrium water table simulations, *J. Geophys. Res.*, 112, D10125, doi:10.1029/2006JD008111, 2007.

1000 [Mosciaro, M.J., Calamari, N.C., Peri, P.L., Flores Montes, N., Seghezzo, L., Ortiz, E., Rejalaga, L., Barral, P., Villarino, S., Mastrangelo, M., Volante, J.: Future scenarios of land use change in the Gran Chaco: how far is zero-deforestation?, *Reg. Environ. Change*, 22, 115, doi:10.1007/s10113-022-01965-5, 2022.](#)

Müller, O. V., Berbery, E. H., Alcaraz Segura, D., and Ek, M. B.: Regional model simulations of the 2008 drought in southern South America using a consistent set of land surface properties, *J. Clim.*, 27(17), 6754-6778, doi:10.1175/JCLI-D-13-00463.1, 2014.

1005 Müller, O. V., Lovino, M. A., and Berbery, E. H.: Evaluation of WRF model forecasts and their use for hydroclimate monitoring over southern South America, *Weather Forecast.*, 31(3), 1001-1017., doi:10.1175/WAF-D-15-0130.1, 2016.

Formatted: Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Formatted: Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Formatted: Font: 10 pt, Not Italic, Font colour: Auto, English (US)

Formatted: English (US)

Formatted: English (US)

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Indent: Left: 0 cm, Hanging: 1 cm, Right: -0.05 cm, Space Before: 2 pt, After: 2 pt

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, Not Italic, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: (Default) Times New Roman, 10 pt, Font colour: Auto, English (US), Pattern: Clear

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Deleted: https://doi.org/

Deleted: https://doi.org/

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Formatted: English (US)

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 10 pt, Not Italic, English (US), Not Highlight

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted: Font: 12 pt, English (UK)

Deleted:

Deleted: https://doi.org/

Deleted: https://doi.org/

- Müller, O., Vidale, P. L., Vannière, B., Schiemann, R., and McGuire, P.: Does the HadGEM3-GC3.1 GCM overestimate land precipitation at high resolution? A constraint based on observed river discharge. *J. Hydrometeorol.*, 22, 8, 2131-2151. doi:10.1175/JHM-D-20-0290.1, 2021.
- Niu, G. Y., Yang, Z. L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning, K., Niyogi, D., Rosero, E., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-scale measurements. *J. Geophys. Res. Atmos.*, 116(D12), doi:10.1029/2010JD015139, 2011.
- Nori, J., Torres, R., Lescano, J. N., Cordier, J. M., Periago, M. E., and Baldo, D.: Protected areas and spatial conservation priorities for endemic vertebrates of the Gran Chaco, one of the most threatened ecoregions of the world. *Divers. Distrib.*, 22(12), 1212-1219, doi:10.1111/ddi.12497, 2016.
- Ortega, G., Arias, P. A., Villegas, J. C., Marquet, P. A., and Nobre, P.: Present-day and future climate over central and South America according to CMIP5/CMIP6 models. *Int. J. Climat.*, 41(15), 6713-6735, doi:10.1002/joc.7221, 2021.
- Paruelo, J. M., Guerschman, J. P., and Verón, S. R.: Agricultural expansion and changes in land use, *Ciencia Hoy*, 15, 14-23, 2005 (in Spanish).
- Piquer-Rodríguez, M., Torella, S., Gavier-Pizarro, G., Volante, J., Somma, D., Ginzburg, R., and Kuemmerle, T.: Effects of past and future land conversions on forest connectivity in the Argentine Chaco. *Landscape Ecol.*, 30, 817-833, doi:10.1007/s10980-014-0147-3, 2015.
- Pohl, B., and Crétat, J.: On the use of nudging techniques for regional climate modeling: Application for tropical convection. *Climate Dyn.*, 43, 1693-1714, doi:10.1007/s00382-013-1994-3, 2014.
- Portillo-Quintero, C. A., and Sánchez-Azofeifa, G. A.: Extent and conservation of tropical dry forests in the Americas. *Biol. Conserv.*, 143(1), 144-155, doi:10.1016/j.biocon.2009.09.020, 2010.
- Ribichich, K. F., Chiozza, M., Ávalos-Britez, S., Cabello, J. V., Arce, A. L., Watson, G., Arias, C., Portapila, M., Trucco, F., Otegui, M. E., and Chan, R. L.: Successful field performance in warm and dry environments of soybean expressing the sunflower transcription factor HB4. *J. Exp. Bot.*, 71(10), 3142-3156, doi:10.1093/jxb/eraa064, 2020.
- Richards, P. D., Myers, R. J., Swinton, S. M., and Walker, R. T.: Exchange rates, soybean supply response, and deforestation in South America. *Glob. Environ. Change*, 22(2), 454-462, doi:10.1016/j.gloenvcha.2012.01.004, 2012.
- Ruscica, R., Menéndez, C. and Sörensson, A.: Land surface-atmosphere interaction in future South American climate using a multi-model ensemble. *Atmos. Sci. Lett.*, 17: 141-147, https://doi.org/10.1002/asl.635, 2016.
- Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y., Chuang, H., Iredell, M., Ek, M., Meng, J., Yang, R., Mendez, M. P., van den Dool, H., Zhang, Q., Wang, W., Chen, M., and Becker, E.: The NCEP Climate Forecast System Version 2. *J. Clim.*, 27(6), 2185-2208, doi:10.1175/JCLI-D-12-00823.1, 2014.
- Salazar, A., Baldi, G., Hirota, M., Syktus, J., and McAlpine, C.: Land use and land cover change impacts on the regional climate of non-Amazonian South America: A review. *Glob. Planet. Change.*, 128, 103-119, doi:10.1016/j.gloplacha.2015.02.009, 2015.

Deleted: Müller, O. V., Vidale, P. L., Vannière, B., Schieman, R., Senan, R., Haarsma, R. J., and Jungclaus, J. H.: Land-Atmosphere Coupling Sensitivity to GCMs Resolution: A Multimodel Assessment of Local and Remote Processes in the Sahel Hot Spot, *J. Clim.*, 34(3), 967-985, doi:10.1175/JCLI-D-20-0303.1, 2021a.

Formatted: Font: 10 pt, English (US)

Formatted

Formatted: Indent: Left: 0 cm. Hanging: 1 cm

Deleted: <https://doi.org/>

Formatted: Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Formatted

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Deleted: . . . Landscape Ecol., 30, 817–833, <https://doi.org/10.1007/s10980-017-0637-1> [44]

Formatted: Font: 10 pt, English (US)

Formatted

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Deleted: <https://doi.org/>

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted

Formatted: Indent: Left: 0 cm, Hanging: 1 cm, Right: 0 cm, Space Before: 0 pt, After: 0 pt, Border: Top: (No border), Bottom: (No border), Left: (No border), Right: (No border), Between : (No border)

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Deleted: <https://doi.org/>

Deleted: <https://doi.org/>

070 [Sánchez, E., Solman, S., Remedio, A. R. C., Berbery, H., Samuelsson, P., Da Rocha, R. P., Mourão, C., Li, L., Samuelsson, P., Da Rocha, R. P., de Castro, M., Jacob, D.: Regional climate modelling in CLARIS-LPB: a concerted approach towards twentyfirst century projections of regional temperature and precipitation over South America. *Clim. Dyn.*, 45, 2193-2212, doi:10.1007/s00382-014-2466-0, 2015.](#)

1075 Santanello, J. A., Dirmeyer, P. A., Ferguson, C. R., Findell, K. L., Tawfik, A. B., Berg, A., Ek, M., Gentine, P., Guillod, B. P., van Heerwaarden, C., Roundy, J., and Wulfmeyer, V.: Land-atmosphere interactions: The loco perspective. *Bull. Am. Meteorol. Soc.*, 99(6), 1253–1272, doi:10.1175/bams-d-17-0001.1, 2018.

Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating soil moisture-climate interactions in a changing climate: A Review. *Earth Sci. Rev.*, 99(3–4), 125–161, doi:10.1016/j.earscirev.2010.02.004, 2010.

080 [Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Wang, W., and Powers, J.G.: A description of the Advanced Research WRF version 3, Technical report, NCAR, doi:10.5065/D68S4MVH, 2008.](#)

[Sörensson, A.A., and Berbery, E. H.: A note on soil moisture memory and interactions with surface climate for different vegetation types in the La Plata basin. *J. Hydrometeorol.*, 16\(2\), 716-729, doi:10.1175/JHM-D-14-0102.1, 2015.](#)

Spennemann, P. and Saulo, A.: An estimation of the land-atmosphere coupling strength in South America using the Global Land Data Assimilation System, *Int. J. Climatol.*, 35, 4151-4166, doi:10.1002/joc.4274, 2015.

085 [Stanimirova, R., Graesser, J., Olofsson, P., and Friedl, M. A.: Widespread changes in 21st century vegetation cover in Argentina, Paraguay, and Uruguay. *Remote Sens. Environ.*, 282, doi:10.1016/j.rse.2022.113277, 2022.](#)

University of East Anglia Climatic Research Unit; Harris, I.C.; Jones, P.D. (2020): CRU TS4.03: Climatic Research Unit (CRU) Time-Series (TS) version 4.03 of high-resolution gridded data of month-by-month variation in climate (Jan. 1901- Dec. 2018). Centre for Environmental Data Analysis, 22 January 2020.

1090 [doi:10.5285/10d3e3640f004c578403419aac167d82, 2020.](#)

[Vera, C.S., and Osman, M.: Activity of the Southern Annular Mode during 2015–2016 El Niño event and its impact on Southern Hemisphere climate anomalies, *Int. J. of Climatol.*, 38, e1288-e1295, doi:10.1002/joc.5419, 2018.](#)

Volante, J. N., Mosciaro, M. J., Gaviera-Pizarro, G. I., and Paruelo, J. M.: Agricultural expansion in the Semiarid Chaco: Poorly selective contagious advance, *Land use policy*, 55, 154-165, doi:10.1016/j.landusepol.2016.03.025, 2016.

095 Weedon, G. P., Balsamo, G., Bellouin, N., Gomes, S., Best, M. J., and Viterbo, P.: The WFDEI Meteorological Forcing Data. Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, doi: 10.5065/486N-8109, 2018.

Williams, E., and Renno, N.: An Analysis of the Conditional Instability of the Tropical Atmosphere. *Mon. Wea. Rev.*, 121, 21–36, doi:10.1175/1520-0493(1993)121<0021:AAOTCI>2.0.CO;2, 1993.

1100

Formatted: Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Formatted ... [48]

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Deleted: https://doi.org/

Deleted:Earth Sci. Rev., 99(3–4), 125–161, https://d... [49]

Formatted: Font: 10 pt, English (US)

Formatted ... [50]

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Deleted: Sirimarco, X., Barral, M. P., Villarino, S. H., and Laterra, P.: Water regulation by grasslands: A global meta-analysis, *Ecohydrology*, e19-34, https://doi.org/10.1002/eco.1934, 2018.

Formatted ... [51]

Deleted: https://doi.org/

Formatted: Font: 10 pt, English (US), Not Highlight

Formatted ... [52]

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Formatted ... [53]

Deleted: https://dx.doi.org/

Formatted: Font: 10 pt, Font colour: Auto, Not Highlight

Formatted ... [54]

Formatted: Indent: Left: 0 cm, Hanging: 1 cm

Formatted ... [55]

Deleted: https://doi.org/

Deleted: https://doi.org/

▼

Parameter	Description
Region	Southeastern South America
Grid resolution	12 km
Grid size	320 x 283 grid points
N° of vertical levels	38
Period	January 2014 – June 2016
Integration time step	60 s
Dynamic solver	ARW
Boundary conditions	CFSv2 (Saha et al. 2011)
Microphysics	Eta (Ferrier) (Rogers y otros, 2001)
Cumulus convection	Betts-Miller-Janjic scheme (Janjic, 1994, 2000)
Surface layer	MM5 Monin-Obukhov scheme (Jiménez et al, 2012)
Land surface model	Noah-MP (Niu et al, 2011)
Planet boundary layer	No boundary-layer
Shortwave radiation	Dudhia scheme (Dudhia, 1989)
Longwave radiation	Rapid Radiative Transfer Model (Mlawer y otros, 1997)

Table 1. Summary of WRF configuration.

Parametrization	Selected Option
Land cover classification	MODIS - IGBP 21 Categories
Stomatal resistance	Ball–Berry (Ball et al. 1987)
Surface layer drag coefficient calculation	Monin–Obukhov (Brutsaert, 1982)
Soil moisture factor for stomatal resistance	Noah type (Chen and Dudhia, 2001)
Runoff and groundwater	TOPMODEL with groundwater (Niu et al. 2011)
Supercooled liquid water	Standard freezing point depression (Niu and Yang 2006)
Soil permeability	Linear effect, more permeable (Niu and Yang 2006)
Radiative transfer	Modified two-stream
Ground surface albedo	CLASS (Canadian Land Surface Scheme) (Verseghy, 1991)
Precipitation partitioning between snow and rain	Jordan (Jordan, 1991)
Soil temp lower boundary condition	TBOT at 8 m from input file
Snow/soil temperature time scheme	Semi implicit

Table 2. NOAH-MP selected parametrization.

Deleted: [Zak, M. R., Cabido, M., Cáceres, D., and Díaz, S.: What drives accelerated land cover change in central Argentina? Synergistic consequences of climatic, socioeconomic, and technological factors. *Enviro. Manage.*, 42\(2\), 181-189, <https://doi.org/10.1007/s00267-008-9101-y>, 2008.](#)

Formatted Table

Deleted: é

Deleted:

Deleted: y otros

Formatted Table

Deleted: ía

Deleted: 07

1125

1130

1135




Figure 1: Model domain and topography. The coloured lines highlight the Gran Chaco subregions: Dry Chaco (red) and Humid Chaco (green).

1150





Figure 2: (a) 2015 MODIS land cover map using IGBP classification. (b) Temporal evolution of land cover in Dry Chaco. For simplicity land cover types are aggregated in fewer classes (see text). (c) LULCCs in Dry Chaco. (d) Quantification (in 1000 km²) of LULCCs in Dry Chaco from 2001 to 2015. The cell shade is proportional to its value.



Deleted:

Deleted: colour





Figure 3: Experimental design.



1160 **Figure 4: Land cover map of (a) the PAST and (b) the FUTURE ensemble, and (c-d) their respective differences with the land cover map of the CONTROL ensemble.**





Figure 5: Time average of (a) the observed precipitation (mean between WFDEI, CRU, and CPC), (b) the CONTROL ensemble precipitation, and (c) their differences. (d) Precipitation time series averaged in the Gran Chaco region. The bands represent the dispersion among observational (grey) and ensemble members (pink), while the solid lines depict their respective means.



Deleted:



Deleted: Time

Deleted: the

Formatted: Font: (Default) Times New Roman, Bold, Font colour: Black

Formatted: Font: (Default) Times New Roman, Bold, Font colour: Black

Formatted: Font: (Default) Times New Roman, Bold, Font colour: Black



Deleted:

Figure 6: Time average of near-surface soil moisture for (a) the SMOPS product, (b) the CONTROL ensemble, (c) their differences, and time average of **DJF** root-zone soil moisture for (d) the HSAF product, (e) the CONTROL ensemble, and (f) their differences. (g) Observed and simulated soil moisture time series averaged over the Gran

Chaco. The pink band represent the dispersion among ensemble members, while the solid red line depicts the respective mean.






Figure 7: Time average of (a) the observed 2m temperature, (b) the CONTROL ensemble 2m temperature, and (c) their differences. (d) Temperature time series averaged over the Gran Chaco. The bands represent the dispersion among observational (grey) and ensemble members (pink), while the solid lines depict their respective means.



Deleted:

Deleted: Time

Figure 8: Differences in the biophysical properties for the OBS_LULCC scenario (CONTROL-PAST): (a) LAI, (b) albedo, and (c) stomatal resistance. Opacity was used in all panels to highlight the region of interest.







Figure 9: Differences in the summer energy budget components in the OBS_LULCC scenario (CONTROL-PAST): (a) net shortwave radiation, (b) net longwave radiation, (c) net radiation, (d) latent heat, (e) sensible heat, and (f) 2m temperature. Opacity was used to highlight the region of interest. The boxplots in the insets show the percentage differences of grid cells with land cover changes (Local) and without land cover changes in Dry Chaco (Non-Local).



Deleted: averaged
Deleted: over



Deleted:

Deleted: The insets show the percentage differences averaged over grid cells with land cover changes (Local) and without land cover changes in Dry Chaco (Non-Local).




Figure 11: Differences in the biophysical properties for the AG_INT scenario (FUTURE-CONTROL): (a) LAI, (b) albedo, and (c) stomatal resistance. Opacity was used in all panels to highlight the region of interest.






Figure 12: Differences in the summer energy budget components in the AG_INT scenario (FUTURE-CONTROL): (a) net shortwave radiation, (b) net longwave radiation, (c) net radiation, (d) latent heat, (e) sensible heat, and (f) 2m temperature. Opacity was used to highlight the region of interest. The boxplots in the insets show the percentage differences of grid cells with land cover changes in Dry Chaco (Local) and over the Humid Chaco (Rem).




Deleted:

Deleted: The insets show the percentage differences averaged over grid cells with land cover changes in Dry Chaco (Local) and





1215 **Figure 13: Differences in the summer water budget components in the AG_INT scenario (FUTURE-CONTROL): (a) precipitation, (b) soil moisture, (c) evapotranspiration, and (d) runoff for Past experiment. Opacity was used to highlight the region of interest. The boxplots in the insets show the percentage differences of grid cells with land cover changes in Dry Chaco (Local) and over the Humid Chaco (Rem).**



Deleted:

Deleted: The insets show the percentage differences averaged over grid cells with land cover changes in Dry Chaco (Local) and over the Humid Chaco (Rem).



Deleted:

Figure 14: Schematic diagram of the possible land-atmosphere feedback pathway caused by agricultural expansion. Upward arrows represent increases, while downward arrows depict decreases.

▼
▲
Page 9: [1] Deleted Omar Müller 17/01/2024 11:44:00

▼
▲
Page 9: [2] Deleted Omar Müller 17/01/2024 11:39:00

▼
▲
Page 9: [3] Deleted Omar Müller 17/01/2024 11:40:00

▼
▲
Page 18: [4] Deleted OVM 19/02/2024 09:03:00

▼
▲
Page 18: [5] Formatted Omar Müller 26/12/2023 11:40:00

Font: 10 pt, English (UK), Not Highlight

▼
▲
Page 18: [6] Formatted Omar Müller 26/12/2023 11:35:00

Indent: Left: 0 cm, Hanging: 1 cm

▼
▲
Page 18: [7] Formatted Omar Müller 26/12/2023 11:40:00

English (UK)

▼
▲
Page 18: [8] Formatted Omar Müller 26/12/2023 11:40:00

Font: 10 pt, English (UK), Not Highlight

▼
▲
Page 18: [9] Formatted Omar Müller 26/12/2023 11:35:00

Font: 10 pt, English (US), Not Highlight

▼
▲
Page 18: [10] Formatted Omar Müller 26/12/2023 11:35:00

Font: 10 pt, English (US), Not Highlight

▼
▲
Page 18: [11] Formatted Omar Müller 26/12/2023 11:38:00

Font: 10 pt, Not Italic, English (US), Not Highlight

▼
▲
Page 18: [12] Formatted Omar Müller 26/12/2023 11:38:00

Font: 10 pt, English (US), Not Highlight

▼
▲
Page 18: [13] Formatted Omar Müller 26/12/2023 11:35:00

Font: 10 pt, Not Italic, English (US), Not Highlight

▼
▲
Page 18: [14] Formatted Omar Müller 26/12/2023 11:35:00

Font: 10 pt, English (US), Not Highlight

▼
▲
Page 18: [15] Formatted Omar Müller 26/12/2023 11:40:00

English (US)

Page 18: [16] Deleted Omar Müller 26/12/2023 10:03:00

Page 18: [17] Deleted OVM 14/02/2024 09:49:00

Page 18: [18] Formatted Omar Müller 26/12/2023 11:44:00

Font: 10 pt, English (US), Not Highlight

Page 18: [19] Formatted Omar Müller 26/12/2023 11:44:00

Indent: Left: 0 cm, Hanging: 1 cm

Page 18: [20] Formatted Omar Müller 26/12/2023 11:44:00

Font: 10 pt, English (US), Not Highlight

Page 18: [21] Formatted Omar Müller 26/12/2023 11:44:00

Font: 10 pt, English (US), Not Highlight

Page 18: [22] Formatted Omar Müller 26/12/2023 11:44:00

Font: 10 pt, Not Italic, English (US), Not Highlight

Page 18: [23] Formatted Omar Müller 26/12/2023 11:44:00

Font: 10 pt, English (US), Not Highlight

Page 18: [24] Formatted Omar Müller 26/12/2023 11:44:00

Font: 10 pt, Not Italic, English (US), Not Highlight

Page 18: [25] Formatted Omar Müller 26/12/2023 11:44:00

Font: 10 pt, English (US), Not Highlight

Page 18: [26] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Page 18: [27] Formatted Omar Müller 27/12/2023 13:51:00

Indent: Left: 0 cm, Hanging: 1 cm

Page 18: [28] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Page 18: [29] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Page 18: [30] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

Page 18: [31] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 18: [32] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 18: [33] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 18: [34] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 18: [35] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 19: [36] Formatted Omar Müller 27/12/2023 13:53:00

Font: (Default) Times New Roman, Font colour: Auto, English (US), Pattern: Clear

▲
Page 19: [37] Formatted Omar Müller 27/12/2023 13:51:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 19: [38] Formatted Omar Müller 26/12/2023 11:48:00

Font: (Default) Times New Roman, 10 pt, Font colour: Auto, English (US)

▲
Page 19: [39] Formatted Omar Müller 27/12/2023 13:55:00

Font: Times New Roman, Font colour: Auto, English (US), Pattern: Clear

▲
Page 20: [40] Deleted Omar Müller 27/12/2023 08:43:00

✖
Page 22: [41] Deleted Omar Müller 26/12/2023 11:24:00

✖
Page 22: [41] Deleted Omar Müller 26/12/2023 11:24:00

✖
Page 22: [41] Deleted Omar Müller 26/12/2023 11:24:00

✖
Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲
Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲
Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲
Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲ Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲ Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲ Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲ Page 22: [42] Formatted Omar Müller 19/01/2024 10:49:00

Font: 10 pt, English (US)

▲ Page 22: [43] Formatted Omar Müller 15/01/2024 09:23:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 22: [43] Formatted Omar Müller 15/01/2024 09:23:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 22: [43] Formatted Omar Müller 15/01/2024 09:23:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 22: [43] Formatted Omar Müller 15/01/2024 09:23:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 22: [43] Formatted Omar Müller 15/01/2024 09:23:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 22: [44] Deleted Omar Müller 15/01/2024 09:24:00

▼

▲ Page 22: [44] Deleted Omar Müller 15/01/2024 09:24:00

▼

▲ Page 22: [45] Formatted Omar Müller 18/01/2024 15:42:00

Font: 10 pt, English (US)

▲ Page 22: [45] Formatted Omar Müller 18/01/2024 15:42:00

Font: 10 pt, English (US)

▲ Page 22: [45] Formatted Omar Müller 18/01/2024 15:42:00

Font: 10 pt, English (US)

▲ Page 22: [45] Formatted Omar Müller 18/01/2024 15:42:00

Font: 10 pt, English (US)

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [46] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [47] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [47] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲
Page 22: [47] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲ Page 22: [47] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲ Page 22: [47] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲ Page 22: [47] Formatted Omar Müller 26/12/2023 12:01:00

Font: 10 pt, English (US), Not Highlight

▲ Page 23: [48] Formatted Omar Müller 15/01/2024 09:30:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 23: [48] Formatted Omar Müller 15/01/2024 09:30:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 23: [48] Formatted Omar Müller 15/01/2024 09:30:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 23: [48] Formatted Omar Müller 15/01/2024 09:30:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 23: [48] Formatted Omar Müller 15/01/2024 09:30:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 23: [48] Formatted Omar Müller 15/01/2024 09:30:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲ Page 23: [49] Deleted Omar Müller 26/12/2023 12:05:00

▼

▲ Page 23: [49] Deleted Omar Müller 26/12/2023 12:05:00

▼

▲ Page 23: [50] Formatted Omar Müller 29/12/2023 10:10:00

Font: 10 pt, English (US)

▲ Page 23: [50] Formatted Omar Müller 29/12/2023 10:10:00

Font: 10 pt, English (US)

▲ Page 23: [50] Formatted Omar Müller 29/12/2023 10:10:00

Font: 10 pt, English (US)

▲ Page 23: [50] Formatted Omar Müller 29/12/2023 10:10:00

Font: 10 pt, English (US)

▲
Page 23: [50] Formatted Omar Müller 29/12/2023 10:10:00

Font: 10 pt, English (US)

▲
Page 23: [50] Formatted Omar Müller 29/12/2023 10:10:00

Font: 10 pt, English (US)

▲
Page 23: [50] Formatted Omar Müller 29/12/2023 10:10:00

Font: 10 pt, English (US)

▲
Page 23: [51] Formatted Omar Müller 15/01/2024 10:16:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 23: [51] Formatted Omar Müller 15/01/2024 10:16:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 23: [51] Formatted Omar Müller 15/01/2024 10:16:00

Font: 10 pt, Font colour: Auto, English (US), Not Highlight

▲
Page 23: [52] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [52] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [52] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [52] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [52] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [52] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [53] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [53] Formatted Omar Müller 26/12/2023 12:04:00

Font: 10 pt, English (US), Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [54] Formatted Omar Müller 27/12/2023 13:56:00

Font: 10 pt, Font colour: Auto, Not Highlight

▲
Page 23: [55] Formatted Omar Müller 27/12/2023 13:57:00

English (UK)

▲
Page 23: [55] Formatted Omar Müller 27/12/2023 13:57:00

English (UK)

▲
Page 23: [55] Formatted Omar Müller 27/12/2023 13:57:00

English (UK)

▲