



Identification of compound drought and heatwave events on a daily scale and across four seasons

Baoying Shan^{1,2}, Niko E.C. Verhoest², and Bernard De Baets¹

¹KERMIT, Department of Data Analysis and Mathematical Modelling, Ghent University, 9000 Ghent, Belgium

²Hydro-Climatic Extremes Lab, Ghent University, 9000 Ghent, Belgium

Correspondence: Baoying Shan (baoying.shan@ugent.be)

Abstract. Compound drought and heatwave (CDHW) events can incur intensified damage to ecosystems, economies, and societies, especially on a warming planet. Although it has been reported that CDHW events in the winter season can also affect insects, birds, and wildfires, the literature generally focuses on the summer season exclusively. Moreover, the coarse resolution of droughts as determined on a monthly scale may hamper a precise identification of the start and/or end dates of CDHW events. Therefore, we propose a method to identify CDHW events on a daily scale that is applicable across the four seasons. More specifically, we use standardized indices calculated on a daily scale to identify four types of compound events in a systematic way. Based on the hypothesis that droughts or heatwaves should be statistically extreme and independent, we remove minor dry or warm spells and merge mutually dependent ones. To demonstrate our method, we make use of 120 years of daily precipitation and temperature observed at Uccle, Brussels-Capital Region, Belgium. Our method yields more precise start and end dates for droughts and heatwaves than what can be obtained with a classical approach acting on a monthly scale, thus also allowing for a better identification of CDHW events. Consistent with existing literature, we find an increase in the occurrence of the number of days in CDHW events at Uccle, mainly due to the increasing frequency of heatwaves. Our results also reveal a seasonality of CDHW events as droughts and heatwaves are negatively dependent in the winter season at Uccle, while they are positively dependent in the other seasons. Overall, the method proposed in this study is shown to be robust and shows potential for exploring how year-round CDHW events influence ecosystems.

1 Introduction

Compound events are defined as a combination of multiple drivers and/or hazards that could lead to extreme ecological and socioeconomic impacts of larger magnitude than the sum of those of the individual ones (Zscheischler et al., 2018; Masson-Delmotte et al., 2021). Under the challenge of climate change, the extremes, especially those related to temperature, are observed at an increasing pace (Perkins et al., 2012; Sharma and Mujumdar, 2017; Mukherjee and Mishra, 2021; Kruger and Sekele, 2013) and are expected to become even more severe in the future (Byrne, 2021; Wu et al., 2021). It is critical to study heatwaves concurrently with droughts, because of the intensification of negative impacts, such as exacerbating water shortage (Coffel et al., 2019), crop failure, and gross primary production reduction (Ciais et al., 2005; Mishra et al., 2020; Zampieri et al., 2017), wildfire and tree mortality (Libonati et al., 2022; Brando et al., 2014; Reichstein et al., 2013), and so on.



25 Recently, the increasing frequency of compound drought and heatwave (CDHW) events (or compound hot and dry events),
has received attention, both globally (Mukherjee and Mishra, 2021; Zhang et al., 2022) and more regionally, e.g. in the United
States (Mazdiyasnani and AghaKouchak, 2015; Alizadeh et al., 2020), India (Sharma and Mujumdar, 2017), Europe (Manning
et al., 2019), China (Kong et al., 2020) and Southeast Brazil (Geirinhas et al., 2021).

The identification of CDHW events lies at the core of temporal trend and frequency analyses. However, the coarse resolution
30 and scale inconsistency of the indices used hamper the reliability of existing studies. Generally, droughts are identified on the
basis of monthly magnitude variations in historical climate variables (McKee et al., 1993), and therefore completely ignore
the intramonthly distribution, while heatwaves are usually obtained on a daily scale. For example, the *3-month* Standardized
Precipitation Index (SPI) for meteorological droughts and the *daily* temperature index for heatwaves are a popular pair of
indices that are used for identifying compound events (see, e.g., Geirinhas et al. (2021)). This coarse scale of the drought index
35 and the mismatch of scales between the drought and heatwave indices could entail some bias on the actual start and/or end
dates as well as the severity of droughts, further affecting the identification of CDHW events. A recent study by Mukherjee and
Mishra (2021) upscaled the resolution to weekly droughts and daily heatwaves. They identified a drought week when the self-
calibrated Palmer Drought Severity Index (Wells et al., 2004) falls below the 10th percentile, thus regarding short dry spells
(7 d) as drought events. Although the mismatch of scales is partially solved this way, too short dry spells might be identified as
40 drought events.

Most studies on CDHW events exclusively focus on the summer or extended summer season, mainly because heatwaves
that lead to deadly heat stress for human beings and ecosystems are only expected in this period of the year (McKechnie
and Wolf, 2010; Ummenhofer and Meehl, 2017; Stillman, 2019; Soroye et al., 2020). While being overlooked in the study of
CDHW events, ecologists have studied the (ecological) impact of relatively high temperatures (and droughts) in non-summer
45 seasons. For example, warmer winters and earlier springtime drying of soils and forest fuels (fallen leaves) are probably linked
to increasing numbers of large wildfires and total area burned in western US forests (Westerling et al., 2006; Littell et al., 2009;
Williams et al., 2010), which could weaken the resistance of surviving trees against bark beetle attack (Raffa et al., 2008).
Laboratory experiments conducted by Radchuk et al. (2013) and Oliver et al. (2015), and a long-term countrywide data set
analysis by McDermott Long et al. (2017) all found that elevated temperatures during the overwintering period could increase
50 the rates of mortality for UK butterflies, possibly due to the increased incidence of diseases and fungi (Radchuk et al., 2013)
or decoupling from photoperiod cues (Wiklund et al., 1996). More studies on the wild passerine in France (Marrot et al.,
2017), the Great tit (a songbird) in UK (Cole et al., 2021) and bee colonies across Austria (Switanek et al., 2017) indicate the
significant ecological influence of relatively high temperatures. Hence, from an ecological point of view, a tool that is able to
identify CDHW events across all seasons is badly needed.

55 Zscheischler et al. (2020) proposed a typology of compound hazard events focusing on four themes: preconditioned, multi-
variate, temporally compounding, and spatially compounding. For the multivariate events where multiple hazards occur at the
same time (for example, droughts and heatwaves in this study), existing literature generally identifies such compound events as
the intersection of the periods of the co-occurring hazards (Zscheischler and Seneviratne, 2017; Ridder et al., 2018; Mukherjee
and Mishra, 2021). However, from a more systematic point of view, we can distinguish various ways to determine the start and



60 end dates of such compound events: the mentioned intersection, but also the union, or one of the periods conditioned on the other.

The overall objective of this study is to develop a method to identify CDHW events on a daily scale across four seasons by addressing the following research questions:

- (a) How can we define heatwaves across four seasons?
- 65 (b) Can we consistently define droughts and heatwaves?
- (c) How can we avoid the fact that the use of indices calculated on a daily scale could produce a large number of small-scale events and/or mutually dependent events when using fixed-threshold-based identification approaches?
- (d) How should we interpret the term ‘compound’ and how can we retrieve CDHW events from the droughts and heatwaves themselves?
- 70 (e) Do CDHW events in non-summer seasons have similar characteristics as those in the summer season?
- (f) Are the temporal and frequency trends observed consistent with existing studies?

This study is structured as follows. In Section 2, we describe the method developed and the data used. We make use of 120 years of daily data of precipitation and temperature observed at the Belgian meteorological institute in Uccle, Brussels-Capital Region, Belgium. In Section 3, we present and discuss the results obtained, followed by some conclusions in Section 4.

75 **2 Materials and methods**

In Subsection 2.1, we describe the data we make use of in this study. In Subsection 2.2, we define what a heatwave is and provide equations for calculating daily heatwave and drought indices across the four seasons, while in Subsection 2.3 we develop a method to identify extreme and independent droughts and heatwaves. In Subsection 2.4, we give the definition of compound events, not only defined through the intersection of periods but also in three alternative ways. Finally, in Subsection 2.5, we
80 describe how to estimate the probabilities of droughts, heatwaves, and CDHW events.

2.1 Data

Daily minimum/maximum temperatures from 1901 to 2020 were acquired from the climatological station of the Royal Meteorological Institute (RMI) of Belgium in Uccle (50°47'55" N, 4°21'29" E, 100 m a.s.l.), while daily rainfall was obtained from 10-minute precipitation time series recorded at the same site. These high-quality observations have been subjected to many
85 studies, see e.g. Verhoest et al. (1997); Ntegeka and Willems (2008); Vandenberghe et al. (2010); Pham et al. (2018).

2.2 Daily drought and heatwave indices

In this study, a heatwave is defined in a relative sense, *i.e.*, as a period of excessively high temperatures compared to the expected normal temperatures in this period. Heatwaves defined in this manner could capture anomalous warm periods during



summer as well as during winter. We use the daily SPI (McKee et al., 1993) and SHI (Standardized Heatwave Index) (Raei et al., 2018) as indices to identify droughts and heatwaves. Inspired by the SPI, the SHI describes hot spells according to the probability of occurrence in a reference period. Given the non-stationarity of temperature records due to global warming, the historical reference period and climate normals for calculating such probabilities must be carefully selected. Accounting for the fact that people and ecosystems adapt to a changing climate, we use the past 30 years as the historical reference period and average the values on every day during this period. This average is then regarded as the expected normal temperature on this day, instead of the average during the longest climatology (the period of record). This 30-year moving window approach has been suggested to deal with the climate non-stationarity bias (Hoylman et al., 2022). To examine the impact of this approach, we apply the Mann–Kendall (MK) test for the daily mean temperature in the period 1901–2020 (the period of record) and in the 30-year moving windows (91 windows per day) based on the data described in Subsection 2.1. The test results (see Fig. S1 for details) point out that over the period of record, 40.27% of calendar days (147 out of 365 days) show a significant increase, while not a single day shows a significant decrease. However, for the 30-year moving windows, only 5.59% of windows (33215 windows in total, 91 per day) show a significant trend, 4.38% with an increase and 1.21% with a decrease, indicating that this approach better accounts for climate non-stationarity.

After having overcome the non-stationarity problem in the above way, we calculate SHI values for each day as:

$$\hat{T}_{m,i} = \begin{cases} \frac{1}{N_h} \sum_{j=i-N_h+1}^i T_{m,j} & , \text{ if } j \geq N_h \\ \frac{1}{N_h} \left(\sum_{j=365-N_h+i+1}^{365} T_{m-1,j} + \sum_{j=1}^i T_{m,j} \right) & , \text{ if } j < N_h \end{cases} \quad (1)$$

$$\text{SHI}_{m,i} = \Phi^{-1}(F_{m,i}^h(\hat{T}_{m,i})) \quad (2)$$

where i and j represent calendar days ($i, j \in \{1, 2, \dots, 365\}$), and $m \in \{1, 2, \dots, M\}$ the year considered, with M the number of years of historical observations; $T_{m,i}$ (°C) is the daily mean temperature at day i of year m , which is obtained by averaging the daily maximum and minimum temperatures. For leap years, we average the daily mean temperature on the 28th and the 29th February and assign it to the 28th February, and then remove the leap days. N_h (d) is the length of the accumulation period for a heatwave. $\hat{T}_{m,i}$ (°C) is the mean temperature over the accumulation period up to and including day i ; $F_{m,i}^h$ is the cumulative distribution function (CDF) fitted to $\{\hat{T}_{m,i}, \hat{T}_{m-1,i}, \dots, \hat{T}_{m-29,i}\}$ if $m \geq 30$, otherwise to $\{\hat{T}_{30,i}, \hat{T}_{29,i}, \dots, \hat{T}_{1,i}\}$; Φ^{-1} is the inverse standard normal CDF; $\text{SHI}_{m,i}$ is the estimated SHI value at day i of year m .

While not relevant for SHI values, a critical problem in the calculation of SPI values is how to deal with the common zeros in precipitation time series. Indeed, strictly positive distributions (like the gamma distribution) are undefined at zero, while for a great deal of positive distributions (like the Weibull distribution), the density function at 0 is always 0, no matter how many zero values are present in the data to which they are fitted. If these distributions were used to fit precipitation time series with many zeros, zero and small rainfall amounts (usually indicating dry conditions) would be not well accounted for, which might affect the drought analysis. To resolve this issue, we split the precipitation values into zero values and positive values and estimate their probabilities separately (Naresh Kumar et al., 2009), as expressed in Eq. (4). In this way, we calculate SPI



120 values for each day as:

$$SP_{m,i} = \begin{cases} \sum_{j=i-N_d+1}^i P_{m,j} & , \text{ if } j \geq N_d \\ \sum_{j=365-N_d+i+1}^{365} P_{m-1,j} + \sum_{j=1}^i P_{m,j} & , \text{ if } j < N_d \end{cases} \quad (3)$$

$$F_i^d(SP_{m,i}) = \begin{cases} F_i^d(0) & , \text{ if } SP_{m,i} = 0 \\ F_i^d(0) + (1 - F_i^d(0)) \times F_i^Y(SP_{m,i}) & , \text{ if } SP_{m,i} > 0 \end{cases} \quad (4)$$

$$F_i^d(0) = \frac{Z_i}{M} \quad (5)$$

$$SPI_{m,i} = \Phi^{-1}(F_i^d(SP_{m,i})) \quad (6)$$

125 where $P_{m,i}$ (mm) is the daily precipitation at day i of year m . For leap years, we add the precipitation on leap days to that on
the 28th February, and then remove the leap days. N_d (d) is the length of the accumulation period for a drought. $SP_{m,i}$ (mm)
is the precipitation over the accumulation period up to and including day i ; F_i^Y is the CDF fitted to the positive precipitation
values $\{SP_{m,i} \mid SP_{m,i} > 0, m = 1, 2, \dots, M\}$; F_i^d is the CDF of all precipitation values and $F_i^d(0)$ is the probability of zero
precipitation. With Z_i representing the number of zero $SP_{m,i}$ values over the M years, $F_i^d(0)$ is the fraction of such zero
130 values for day i .

The choice of the distribution to be fitted is an important source of uncertainty in the SPI values, given that, depending on
the case considered, other commonly used distributions are valid (Laimighofer and Laaha, 2022). In this study, we consider
10 commonly used distributions from which we select the one with the lowest value for the Akaike Information Criterion
(AIC) (Akaike, 1974) in the calculation of both the SPI and SHI values. The distributions considered are the normal, exponen-
135 tial, gamma, generalized extreme value (GEV), inverse Gaussian, logistic, log-logistic, log-normal, Burr, and extreme value
distributions.

2.3 Extreme and independent events

Droughts (resp. heatwaves) are commonly identified by a fixed-threshold approach as periods with index values consecutively
below (resp. above) a certain threshold (e.g., -1 for SPI), even though some studies use different thresholds for the onset and
140 ending of an extreme event. In this study, we select one threshold and refer to it as the pre-identification threshold and to the
resulting events as dry (resp. warm) spells.

Yet, this fixed-threshold approach, especially when applied to daily indices, may result in a large number of small-scale
and mutually dependent spells, causing the identified droughts or heatwaves to be not extreme or not independent. In turn,
this could affect the analysis of temporal trends and frequencies. Procedures for excluding minor spells and merging mutually
145 dependent ones have already been explored in literature (Fleig et al., 2006). However, it is still tricky to discern objectively
whether spells are minor or whether two neighboring spells are dependent.

In this study, we develop a method to obtain objective thresholds for excluding minor spells and merging mutually dependent
ones. First, two indicators are selected: the *duration* (d) and the *total deficiency*. The duration expresses how long a spell



lasts, and equals the number of days in a dry (D_d) or a warm spell (D_h), while the total deficiency describes how close two
150 neighboring spells are (Fig. 1). In particular, the *total drought deficiency* (TD_d) corresponds to the area enclosed between the
SPI curve and the pre-identification threshold line in the period between two dry spells:

$$TD_d = \sum_{i=a}^b (SPI_{m,i} - SPI_d),$$

where $[a, b]$ is the interval between two spells and SPI_d is the pre-identification threshold for droughts. Similarly, the *total heat*
deficiency (TD_h) is defined as:

155 $TD_h = \sum_{i=a}^b (SHI_h - SHI_{m,i}),$

where SHI_h is the pre-identification threshold for heatwaves. The total deficiency TD_d (resp. TD_h) is put forward as it provides
a comprehensive view of the period between two spells, accounting for both duration and how far it is from being dry (resp.
warm). For example, a small value of TD_d means that the interval is short and/or the SPI values are near the pre-identification
threshold. In that case, the neighboring spells are more likely to be mutually dependent, suggesting to merge the neighboring
160 spells and the interval between them into one longer spell.

More importantly, in view of our aim of droughts and heatwaves to be extreme and independent, we assume that their severity
should follow a Generalized Extreme Value (GEV) distribution (to guarantee that only extreme-impact events are included)
and their arrivals to occur according to a Poisson process (to guarantee that they occur independently in time). Severity is the
characteristic used to quantify droughts (resp. heatwaves), usually calculated by means of drought (resp. heatwave) indices.
165 The above hypotheses could be used as objective criteria for determining appropriate thresholds for removing and merging
spells, *i.e.*, thresholds that make the severity of the final identified events follow a GEV distribution, as well as the inter-arrival
times to follow an exponential distribution at the same time.

Concretely, we remove a spell when its duration is shorter than a chosen removal threshold ($D_d < R_d$, resp. $D_h < R_h$).
After removing such spells, we check whether any total deficiency is smaller than a chosen merging threshold ($TD_d < M_d$,
170 resp. $TD_h < M_h$). If this is the case, the two neighboring spells are considered to belong to one longer spell. Therefore, they
are merged, and thus the period between both initial spells is now considered to be part of a longer spell. These processes are
illustrated in Fig. 1.

The specific steps to obtain the proper thresholds for removing and merging spells, taking the identification of heatwaves as
an example, are explained next:

- 175 (1) Given an accumulation period length N_h and pre-identification threshold SHI_h , we calculate the daily SHI values using
Eqs. (1)–(2).
- (2) We label all warm spells potentially belonging to a heatwave (days consecutively above SHI_h).
- (3) We consider all possible combinations of removal and merging thresholds R_h and M_h , *i.e.*, $R_h \in \{1, \dots, 5N_h\}$ and $M_h \in$
 $\{-\infty, 0, 1, \dots, 5N_h\}$, where $-\infty$ means no merging is carried out. For each combination the following steps are carried
180 out.



(3.1) We remove the warm spells with a duration shorter than R_h and relabel the remaining warm spells. Subsequently, we merge neighboring spells if the corresponding total deficiency TD_h is less than M_h and again relabel the warm spells. After removing and merging, the severities (sum of SHI values during a warm spell) and inter-arrival times (d) for this combination are calculated as shown in Fig. 1.

185 (3.2) We test whether the inter-arrival times follow an exponential distribution by means of a Kolmogorov–Smirnov test (KS test) (Massey, 1951) to meet the hypothesis of heatwaves being independent. The KS test is chosen because it does not assume any particular underlying distribution. If the KS test rejects the null hypothesis, we proceed to the next combination; otherwise, we go to step (3.3).

190 (3.3) We test whether the GEV distribution fits the severities to meet the hypothesis that heatwaves should be extreme. If the KS test rejects the null hypothesis, we proceed to the next combination; otherwise, we go to step (3.4).

(3.4) We check whether other commonly used distributions also fit the severities by means of the KS test. The distributions considered are the same as those used in the calculation of the SPI and SHI values. If the KS test rejects all other distributions, we take a note for this combination that the GEV distribution is the only fit, and proceed to the next combination; otherwise, we go to step (3.5);

195 (3.5) We compute the AIC and Root Mean Square Deviation (RMSD) values for all distributions that pass the KS test. We record whether the GEV distribution has the lowest AIC or the lowest RMSD.

(4) Among all combinations for which the GEV distribution is the only fit or results in both the lowest AIC and the lowest RMSD values, we select the one leading to the largest number of events in order to increase the probability of having more compound events and benefit the further analyses. If there is no such combination, we consider the combinations resulting in the lowest AIC only. If no combination at all has been identified, then the accumulation period length and pre-identification threshold considered are discarded.

200

(5) If a combination of thresholds has been identified, then the warm spells obtained after applying the removal and merging procedures are finally labeled as heatwaves, meeting the hypothesis that they are extreme and independent.

The identification of drought events follows a similar procedure.

205 2.4 Identification of compound events

After having identified droughts and heatwaves, we proceed with the identification of CDHW events. In literature, a CDHW event is generally defined as the period of overlap, *i.e.*, the intersection, of a drought and a heatwave (Fig. 2(d)). However, this is not the only way in which a compound event could be defined. In general, we can distinguish at least four different ways to define a compound event. These compound events consist of:

210 (i) the days belonging to both the drought and the heatwave, *i.e.*, the intersection (d-and-h). Note that a drought might overlap with multiple heatwaves, thus resulting in multiple compound events; the same applies to a heatwave;




Figure 1. Conceptual illustration of the identification of droughts and heatwaves. (a) and (c) show the dry spells and warm spells obtained after applying the pre-identification threshold; (b) and (d) show the final identified droughts and heatwaves after applying the removal and merging procedures. Duration refers to the length of a warm spell or dry spell, while inter-arrival time is the length of the period between the arrivals of two spells. The blue area represents the total deficiency, while the yellow area represents the severity, calculated as the sum of the SHI (resp. negative SPI) values in a warm (resp. dry) spell.

(ii) the days belonging to the drought or the heatwave, *i.e.*, the union (d-or-h), provided the events overlap. Note that a drought might overlap with multiple heatwaves, thus resulting in multiple overlapping compound events, that need to be merged subsequently;

215 (iii) the days belonging to the drought, denoted as d-cond-h, provided the events overlap. Note that a drought overlap with multiple heatwaves. Thus, in the d-cond-h case, only those droughts are preserved that overlap with at least one heatwave;

(iv) the days belonging to the heatwave, denoted as h-cond-d, provided the events overlap. Note that a heatwave might overlap with multiple droughts. A heatwave that does not overlap with at least one heatwave is not considered in the h-cond-d case.

220 Figure 2 displays a conceptual diagram of the above four different types of CDHW events. A CDHW event is characterized by three variables: the duration, the marginal drought severity, and the marginal heatwave severity. The marginal drought severity is computed as the sum of the negative SPI values within the CDHW event, while the marginal heatwave severity is computed as the sum of the SHI values within the CDHW event.



It could be of relevance to ecologists to investigate whether compound events have a higher impact than droughts or heat-
 225 waves only. For certain species, it is known that heatwaves have an impact (Pipoly et al., 2021), yet it might be interesting
 to investigate the possible intensified or weakened impacts of heatwaves during drought conditions (*i.e.*, h-cond-d). Similarly,
 droughts occurring during a heatwave may have an increased impact (*i.e.*, d-cond-h). For species that suffer from both droughts
 and heatwaves, a compound event during which a drought and/or a heatwave occur (*i.e.*, d-or-h) may have a stronger impact
 than when droughts or heatwaves occur independently of each other.




Figure 2. Conceptual diagram of four types of CDHW events: intersection (d-and-h), union (d-or-h), drought conditioned on heatwave (d-
 cond-h), and heatwave conditioned on drought (h-cond-d). Grey shadows represent the days in droughts or heatwaves; Arrows indicate the
 duration of CDHW events.

230 2.5 Probabilities of occurrence

The probabilities of occurrence of droughts (P_d), heatwaves (P_h) and CDHW events (*i.e.*, $P_{d\text{-and-h}}$, $P_{d\text{-or-h}}$, $P_{d\text{-cond-h}}$, and $P_{h\text{-cond-d}}$)
 are estimated empirically as the fractions of days belonging to the corresponding events within the study period. By comparing
 $P_{d\text{-and-h}}$ with the probability of having a compound event assuming that droughts and heatwaves occur independently of each
 other, *i.e.*, by comparing $P_{d\text{-and-h}}$ with $P_d P_h$, we can examine whether there is a positive dependence ($P_{d\text{-and-h}} > P_d P_h$), negative
 235 dependence ($P_{d\text{-and-h}} < P_d P_h$), or no dependence at all ($P_{d\text{-and-h}} \approx P_d P_h$) between droughts and heatwaves (Ridder et al., 2020).

3 Results and discussion

3.1 Removal and merging thresholds

We identify the droughts and heatwaves for various accumulation period lengths ($N_d = 15, 30, 45, 60, 90$ d; $N_h = 3, 5, 7, 10, 15$ d)
 and pre-identification thresholds ($SPI_d = -0.5, -1, -1.3$; $SHI_h = 0.5, 1, 1.3$). Table 1 shows the resulting removal and merging



Table 1. Removal thresholds, R_d , merging thresholds, M_d , average number of days per year in an event, and average number of events per year, for droughts (a) and heatwaves (b).

(a)	R_d			M_d			Number of days per yr			Number of events per yr		
SPI_d	-0.5	-1	-1.3	-0.5	-1	-1.3	-0.5	-1	-1.3	-0.5	-1	-1.3
$N_d = 15$	23	14	10	1	15	11	37.8	28.3	21.2	1.12	1.13	1.18
$N_d = 30$	42	22	20	8	46	1	45.7	33.6	17.9	0.65	0.68	0.55
$N_d = 45$	44	31	25	1	$-\infty$	1	59.8	30.8	18.7	0.78	0.60	0.43
$N_d = 60$	58	45	24	0	$-\infty$	51	62.1	28.9	24.9	0.58	0.43	0.39
$N_d = 90$	81	59	32	1	9	17	67.2	30.2	23.6	0.43	0.25	0.28

(b)	R_h			M_h			Number of days per yr			Number of events per yr		
SHI_h	0.5	1	1.3	0.5	1	1.3	0.5	1	1.3	0.5	1	1.3
$N_h = 3$	13	9	6	0	0	3	34.9	19.2	16.4	1.83	1.56	1.88
$N_h = 5$	16	10	8	0	$-\infty$	$-\infty$	40.6	25.5	16.4	1.73	1.83	1.47
$N_h = 7$	27	11	10	4	$-\infty$	6	23.5	29.9	16.5	0.49	1.90	1.09
$N_h = 10$	23	13	12	$-\infty$	$-\infty$	$-\infty$	41.1	33.6	18.6	1.23	1.76	1.08
$N_h = 15$	29	17	18	2	$-\infty$	11	49.1	33.6	15.4	1.03	1.30	0.58

240 thresholds, the corresponding average number of days per year in an event, and the average number of events per year. We consider, for example, the case $N_d = 15$ d and $SPI_d = -1$. In this case, the combination $R_d = 14$ d and $M_d = 15$, results in 28.3 drought days per year and 1.13 drought events per year on average. The values $R_d = 14$ d and $M_d = 15$ indicate that all dry spells shorter than 14 d are removed, and that all neighboring spells with a total deficiency of less than 15 are merged, respectively, to meet the hypothesis that drought events are extreme and independent.

245 The proposed removal and merging method shows the desired behavior. As could be expected, more extended accumulation periods (*i.e.*, larger N_d or N_h values) cause the SPI and SHI values to evolve more gradually over time, resulting in dry or warm spells being longer. Besides, larger pre-identification thresholds for droughts (*i.e.*, larger SPI_d) or smaller pre-identification thresholds for heatwaves (*i.e.*, smaller SHI_h) tend to lead to the inclusion of more days in dry or warm spells, hence also resulting in their duration being longer. We can expect the removal threshold needed to exclude minor spells to increase with
250 larger N_d or N_h values as well as with larger SPI_d or smaller SHI_h values.

As shown in Table 1, for example, if $SPI_d = -1$, then R_d takes as values 14, 22, 31, 45, 59 d when N_d ranges over 15, 30, 45, 60, 90 d. If $N_d = 15$, then dry spells shorter than 10, 14, and 23 d are removed when SPI_d ranges over -1.3, -1, and -0.5. In the same way, if $SHI_h = 1$, then R_h takes as values 9, 10, 11, 13, 17 d when N_h ranges over 3, 5, 7, 10, 15 d. In case $N_h = 3$, then warm spells shorter than 6, 9, and 13 d are removed when SHI_h ranges over 0.5, 1.0, and 1.3. The same behavior is also
255 observed for other accumulation period lengths and other pre-identification thresholds. In summary, the way how R_d and R_h change with varying accumulation period lengths and pre-identification thresholds in this study is consistent with expectations.



In contrast to the removal thresholds, no clear patterns can be distinguished for the merging thresholds when varying the accumulation period lengths or pre-identification thresholds in Table 1. Interestingly, the merging thresholds influence the number of events. What stands out in Table 1 is the case with $SHI_h = 1.3$ and $N_h = 15$ d. It then holds that $M_h = 11$, leading to 0.58 heatwaves per year on average, which is considerably less than in other cases with the same accumulation period length or pre-identification threshold. This can be explained by the fact that a larger merging threshold results in neighboring spells being combined, ultimately leading to fewer events.

We also apply the fixed-threshold method for various accumulation period lengths and pre-identification thresholds. The number of days and events is strongly reduced when using our method compared to the fixed-threshold one, as only extreme and independent events are selected. We find an average 28.3–33.6 d/yr and 17.9–24.9 d/yr in droughts for $SPI_d = -1$ and $SPI_d = -1.3$, respectively. Without the removal and merging procedure, there are 56.8–60.6 d/yr and 35.0–38.4 d/yr for $SPI_d = -1$ and $SPI_d = -1.3$, respectively (see Table S1 for details). Similarly, there are 19.2–33.6 d/yr in heatwaves when $SHI_h = 1$, while there are 68.2–68.8 d/yr in heatwaves before removing and merging. This way, only about half of the days are left in extreme events. Moreover, 35.5%–50.7% of dry spells last less than five days, and 5.6%–22.3% of warm spells only last one day in all cases (Fig. S2 in appendix) if the fixed-threshold method is applied. We doubt that such very short spells influence society or ecosystems, and thus suggest to exclude them from the analysis of climate extremes.

3.2 Identification results for CDHW events

Different accumulation period lengths and/or pre-identification thresholds for droughts and heatwaves could result in different CDHW events. We consider, for instance, the CDHW events corresponding to $N_d = 15$ d, $SPI_d = -1$ and $N_h = 3$ d and $SHI_h = 1$. The corresponding removal and merging thresholds are given by $R_d = 14$ d, $M_d = 15$ and $R_h = 9$ d, $M_h = 0$.

Next, we discuss the results for the four different types of compound events. In Uccle, in the period 1901–2020 (the period of record), in total 33 CDHW events of types d-and-h and h-cond-d are identified, and 29 CDHW events of types d-cond-h and d-or-h. Hence, their frequency of occurrence ranges from once every 3.6 years to once every 4.1 years (3.6 to 4.1 year return period). Furthermore, the corresponding numbers of days in these CDHW events are given by 2.9 (type d-and-h), 9.1 (type d-or-h), 8.3 (type d-cond-h), and 3.7 (type h-cond-d) d/yr on average. The low numbers of events and numbers of days illustrate that the identification of CDHW events only considers rare events.

Given that we work on a daily scale, a more precise identification of the start and end dates of an event is possible. To demonstrate this, we take the year 1976 as an example, in which extraordinary droughts and heatwaves occurred in Belgium, leading to several forest fires and even requiring restrictions on the use of potable water (KMI, 2017). Our method identifies a heatwave spanning from June 23 to July 10, lasting more than half a month, which is exceptionally long for a heatwave in Belgium (shown on the left of Fig. 3). At the same time, there was a severe drought event lasting more than four months, starting on February 28 and ending on July 11 (due to an intense precipitation event (31.7 mm daily rainfall) on July 12). Moreover, a CDHW event of types d-or-h and d-cond-h is identified starting on February 28 and ending on July 11, while a CDHW event of the other two types is identified from June 23 to July 10. Applying a 1-month SPI (shown on the right of Fig. 3), the drought is identified as ending on June 30. Consequently, the corresponding compound event of type d-and-h then




Figure 3. Identification results in 1976 for droughts, heatwaves, and CDHW events (of types d-and-h, d-or-h, d-cond-h, and h-cond-d) based on the daily SPI (left) and 1-month SPI (right). Grey areas indicate CDHW events. Identification results in other years are shown in Fig. S3.

starts on June 23 and ends on June 30, lasting 8 d only, in contrast to 18 d in case the daily SPI is used. Comparing these two results reveals that the use of the monthly index could significantly bias the estimation of the duration of CDHW events. The proposed method offers a more precise identification as it is able to capture the start and end dates of droughts and CDHW events intramonthly thanks to the use of daily indices.

295 Moreover, the method proposed also allows to identify heatwaves and CDHW events in the winter season because the SPI and SHI treat each day in the same way by evaluating extreme anomalies. Figure 4 shows an example of the winter of 2015 when Uccle had an exceptionally mild winter. An average temperature of 6.6 °C was recorded from November 1 until the end of December, while the average temperature in this period over the past 30 years was 3.8 °C. A CDHW event of type d-and-h is identified from November 3 to November 17. Given that CDHW events in non-summer months are also of interest to ecologists
300 (Williams et al., 2010), our method could be of high interest to them.

CDHW events can be described by the following numerical characteristics: the duration, the marginal drought severity, and the marginal heatwave severity. The pairwise relationships between these three characteristics are explored (see Fig. 5): all show positive correlations for the four types of CDHW events. Especially for type d-and-h, the three characteristics are highly correlated, with Pearson correlation coefficients larger than 0.9. Interestingly, CDHW events of type d-or-h and d-cond-h tend




Figure 4. Identification results for a winter event in 2015. The orange color is used for droughts, the green color for heatwaves.




Figure 5. Scatter plots for the duration and marginal severity of CDHW events of four types. r stands for the Pearson correlation coefficient, with the subscripts 1, 2, and 3 representing the duration, marginal drought severity, and marginal heatwave severity, respectively. The orange color is used for droughts, the green color for heatwaves.

305 to have a longer duration (with a maximum duration of 148 d), while the duration of CDHW events of type h-cond-d and d-and-h is much shorter (with a maximum duration of 32 d and 28 d, respectively), because the length of heatwaves limits the duration of these events.



3.3 Temporal trends

In this subsection, we specifically analyze the CDHW events corresponding to $N_d = 15$ d, $SPI_d = -1$ and $N_h = 3$ d, $SHI_h = 1$.
310 However, the trends observed are in line with those for other accumulation period lengths and pre-identification thresholds (see Table S2).

We can observe that the number of CDHW events and the number of days in CDHW events in Uccle are increasing over the period 1901–2020 (see Fig. 6). In particular, the number of CDHW events shows a large increase from 0.167 events/yr in the period 1901–1960 to 0.317 events/yr in the period 1961–2020 for types d-or-h and d-cond-h (or from 0.217 to 0.333 events/yr
315 for types d-and-h and h-cond-d), nearly the double. Also the numbers of days in the four types of CDHW events are higher in the period 1960–2020 compared to the period 1901–1960, such as a rise from 7.78 d/yr to 10.4 d/yr for CDHW events of type d-or-h and from 2.23 d/yr to 3.50 d/yr for CDHW events of type d-and-h. We further applied the MK test to check whether there is a statistically significant trend for the number of days per year in CDHW events. Despite the positive trends for the CDHW events of all four types considered, they do not appear to be significant at the 95% confidence level.

320 The increasing frequency of heatwaves contributes to the increasing frequency of CDHW events. It should be underlined that heatwaves are obtained through comparison with the expected normal temperatures during the past 30 years instead of the period of record. Nevertheless, there has been a substantial rise in the number of days in heatwaves per year and the number of heatwaves, increasing from 15.6 d/yr and 76 events in the period 1901–1960 to 22.8 d/yr and 111 events in the period 1961–2020. In addition, the MK test for the number of days in heatwaves per year confirms a significant increasing trend at the
325 95% confidence level. For droughts, there is a slight yet non-significant increase in the number of days in droughts per year, despite the rise in the average number of days in droughts per year and the number of droughts from 27.3 d/yr and 63 events in the period 1901–1960 to 29.3 d/yr and 72 droughts in the period 1961–2020. Thus, the increases for the CDHW events are mainly due to the corresponding increases for heatwaves. This is consistent with the study of Manning et al. (2019) analyzing dry and hot events over Europe in the summer season.

340 3.4 Seasonality

Next, we examine the seasonality of CDHW events (Fig. 7). Their probabilities of occurrence show monthly characteristics. For instance, CDHW events in April, August, and October all have relatively high probabilities of occurrence compared to the other months, regardless of the type of CDHW event considered (d-and-h, d-or-h, d-cond-h, or h-cond-d). Contrarily, the probability of experiencing a compound event is low in December, January and February. For the month of January in
335 particular, no CDHW events have been identified in the period 1901–2020. The limited number of events may preclude the reliable estimation of the monthly probability of occurrence. Alternatively, we can also calculate the probability of occurrence at a seasonal scale. The results show that the probability of occurrence of CDHW events in winter is smaller than in the other three seasons, but this difference is not apparent for droughts or heatwaves.

We further compare the probability of occurrence of CDHW events of type d-and-h ($P_{d\text{-and-h}}$) and the probability of having a
340 compound event assuming that droughts and heatwaves are independent ($P_d P_h$). As can be seen from Fig. 7, $P_{d\text{-and-h}}$ is smaller




Figure 6. Temporal trends of droughts, heatwaves, and CDHW events of four types. Two periods are considered: 1901–1960 (gray) and 1961–2020 (yellow). The superimposed text indicates the number of days in these events per year and the number of events per year averaged over the corresponding period.

than $P_d P_h$ in winter, while it is greater in spring, summer, and autumn. This reveals that droughts and heatwaves in Uccle tend to coincide in spring, summer, and autumn, while CDHW events are less likely in winter.

Atmospheric and land-atmosphere interactions could explain the above seasonal differences. It has been widely reported and interpreted in literature (see, e.g., Geirinhas et al. (2021); Schumacher et al. (2019); Miralles et al. (2014)) that droughts and heatwaves in summer are positively dependent. First, this phenomenon has been interpreted in the context of soil moisture–atmosphere coupling, with soil moisture deficits related to reduced precipitation leading to enhanced surface sensible heating and higher surface temperatures (Berg et al., 2015). Second, heatwaves could reduce the total energy transfer to the atmosphere, possibly decreasing convective precipitation (Zaitchik et al., 2006). This, in turn, leads to a soil-precipitation feedback loop that tends to extend or intensify drought conditions (Mazdiyasi and AghaKouchak, 2015). Third, the synoptic-scale weather systems favorable for extreme heat are also unfavorable for rain, which increases the probability of co-occurrence (Berg et al., 2015). Precipitation in Uccle is characterized by cyclonal and convective rainfall in the summer months (Verhoest et al., 1997). The above three mechanisms are possible explanations for the simultaneous occurrence of CDHW events in the non-winter seasons at Uccle. In winter, negative correlations between droughts and heatwaves dominate in Uccle as warm moist advection in extratropical cyclones favors precipitation (Trenberth and Shea, 2005). Uccle is located in the westerlies area with the prevailing wind from west to east. The westerlies are strongest in the winter hemisphere; the pressure is lower over the poles during this time. The warm winter west wind in Uccle favors precipitation causing the CDHW events to be less frequent. Furthermore, due to the radiative influence of clouds, cloudy (often wet) days in winter are generally warmer while sunny (dry)




Figure 7. Seasonal characteristics. The first three rows show the probabilities of occurrence of droughts, heatwaves, and CDHW events of types d-and-h, d-or-h, d-cond-h, and h-cond-d, in each month and each season. The bottom two figures compare $P_{d\text{-and-}h}$ to $P_d P_h$ in each month and each season, respectively.

days are colder. Finally, CDHW events develop less likely in winter as soil-precipitation feedbacks play no role as the low atmospheric demand can easily be met by the typically wet soils (Seneviratne et al., 2010; Schumacher et al., 2022).

360 4 Conclusion

In this study, we have extended the identification of compound drought and heatwave events on a daily scale and across four seasons based on daily SPI and SHI values. Given the hypothesis that droughts or heatwaves are extreme and independent, we developed a convenient and objective method to remove minor spells and merge mutually dependent ones. We argued that there are four ways to define a CDHW event, *i.e.*, the intersection, the union, droughts conditioned on heatwaves, and heatwaves
 365 conditioned on droughts. We demonstrated the method on daily observations from 1901 to 2020 at Uccle, Brussels-Capital Region.

The results confirmed the effectiveness of the proposed method. Operating on a daily scale facilitates the identification of intra-month or intra-week events, thus providing more precise start and end dates. We found that the frequency of occurrence of heatwaves and CDHW events has increased and that the increasing temperatures contribute to the increase in CDHW events,



370 which aligns with the current literature. Another significant finding is that the occurrence of CDWH events shows seasonal patterns, with the occurrences of droughts and heatwaves being negatively dependent in winter but positively dependent in the other three seasons at Uccle.

Ecologists have demonstrated the impact of extremes during winter. The method proposed in this study, allowing for a better identification of droughts, heatwaves, and their joint occurrence, could provide them with a more appropriate looking glass.

375

380

Acknowledgements. The support provided by the China Scholarship Council (CSC) of Baoying Shan at Ghent University is acknowledged.



385 References

- Akaike, H.: A new look at the statistical model identification, *IEEE Transactions on Automatic Control*, 19, 716–723, 1974.
- Alizadeh, M. R., Adamowski, J., Nikoo, M. R., AghaKouchak, A., Dennison, P., and Sadegh, M.: A century of observations reveals increasing likelihood of continental-scale compound dry-hot extremes, *Science Advances*, 6, eaaz4571, 2020.
- Berg, A., Lintner, B. R., Findell, K., Seneviratne, S. I., van den Hurk, B., Ducharne, A., Cheruy, F., Hagemann, S., Lawrence, D. M.,
390 Malyshev, S., et al.: Interannual coupling between summertime surface temperature and precipitation over land: Processes and implications for climate change, *Journal of Climate*, 28, 1308–1328, 2015.
- Brando, P. M., Balch, J. K., Nepstad, D. C., Morton, D. C., Putz, F. E., Coe, M. T., Silvério, D., Macedo, M. N., Davidson, E. A., Nóbrega, C. C., et al.: Abrupt increases in Amazonian tree mortality due to drought–fire interactions, *Proceedings of the National Academy of Sciences*, 111, 6347–6352, 2014.
- 395 Byrne, M. P.: Amplified warming of extreme temperatures over tropical land, *Nature Geoscience*, 14, 837–841, 2021.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A., et al.: Europe-wide reduction in primary productivity caused by the heat and drought in 2003, *Nature*, 437, 529–533, 2005.
- Coffel, E. D., Keith, B., Lesk, C., Horton, R. M., Bower, E., Lee, J., and Mankin, J. S.: Future hot and dry years worsen Nile Basin water scarcity despite projected precipitation increases, *Earth's Future*, 7, 967–977, 2019.
- 400 Cole, E. F., Regan, C. E., and Sheldon, B. C.: Spatial variation in avian phenological response to climate change linked to tree health, *Nature Climate Change*, 11, 872–878, 2021.
- Fleig, A. K., Tallaksen, L. M., Hisdal, H., and Demuth, S.: A global evaluation of streamflow drought characteristics, *Hydrology and Earth System Sciences*, 10, 535–552, 2006.
- Geirinhas, J. L., Russo, A., Libonati, R., Sousa, P. M., Miralles, D. G., and Trigo, R. M.: Recent increasing frequency of compound summer
405 drought and heatwaves in Southeast Brazil, *Environmental Research Letters*, 16, 034036, 2021.
- Hoylman, Z. H., Bocinsky, R. K., and Jencso, K. G.: Drought assessment has been outpaced by climate change: empirical arguments for a paradigm shift, *Nature Communications*, 13, 1–8, 2022.
- KMI: 1975-1976 drought and heatwaves, <https://www.meteo.be/nl/info/nieuwsbrief/artikels-2017/de-droogte-van-2016-2017-het-archief>, accessed: 2022.3.15, 2017.
- 410 Kong, Q., Guerreiro, S. B., Blenkinsop, S., Li, X.-F., and Fowler, H. J.: Increases in summertime concurrent drought and heatwave in Eastern China, *Weather and Climate Extremes*, 28, 100242, 2020.
- Kruger, A. and Sekele, S.: Trends in extreme temperature indices in South Africa: 1962–2009, *International Journal of Climatology*, 33, 661–676, 2013.
- Laimighofer, J. and Laaha, G.: How standard are standardized drought indices? Uncertainty components for the SPI & SPEI case, *Journal of*
415 *Hydrology*, p. 128385, <https://doi.org/https://doi.org/10.1016/j.jhydrol.2022.128385>, 2022.
- Libonati, R., Geirinhas, J. L., Silva, P. S., Russo, A., Rodrigues, J. A., Belém, L. B., Nogueira, J., Roque, F. O., DaCamara, C. C., Nunes, A. M., et al.: Assessing the role of compound drought and heatwave events on unprecedented 2020 wildfires in the Pantanal, *Environmental Research Letters*, 17, 015005, 2022.
- Littell, J. S., McKenzie, D., Peterson, D. L., and Westerling, A. L.: Climate and wildfire area burned in western US ecoprovinces, 1916–2003,
420 *Ecological Applications*, 19, 1003–1021, 2009.



- Manning, C., Widmann, M., Bevacqua, E., Van Loon, A. F., Maraun, D., and Vrac, M.: Increased probability of compound long-duration dry and hot events in Europe during summer (1950–2013), *Environmental Research Letters*, 14, 094 006, 2019.
- Marrot, P., Garant, D., and Charmantier, A.: Multiple extreme climatic events strengthen selection for earlier breeding in a wild passerine, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372, 20160 372, 2017.
- 425 Massey, F. J.: The Kolmogorov-Smirnov Test for Goodness of Fit, *Journal of the American Statistical Association*, 46, 68–78, <http://www.jstor.org/stable/2280095>, 1951.
- Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., et al.: IPCC, 2021: climate change 2021: the physical science basis. Contribution of working group I to the sixth assessment report of the Intergovernmental Panel on Climate Change, Tech. rep., Cambridge University Press, 2021.
- 430 Mazdiyasn, O. and AghaKouchak, A.: Substantial increase in concurrent droughts and heatwaves in the United States, *Proceedings of the National Academy of Sciences*, 112, 11 484–11 489, 2015.
- McDermott Long, O., Warren, R., Price, J., Brereton, T. M., Botham, M. S., and Franco, A. M.: Sensitivity of UK butterflies to local climatic extremes: which life stages are most at risk?, *Journal of Animal Ecology*, 86, 108–116, 2017.
- McKechnie, A. E. and Wolf, B. O.: Climate change increases the likelihood of catastrophic avian mortality events during extreme heat waves, 435 *Biology Letters*, 6, 253–256, 2010.
- McKee, T. B., Doesken, N. J., Kleist, J., et al.: The relationship of drought frequency and duration to time scales, in: *Proceedings of the 8th Conference on Applied Climatology*, vol. 17, pp. 179–183, California, 1993.
- Miralles, D. G., Teuling, A. J., Van Heerwaarden, C. C., and Vilà-Guerau de Arellano, J.: Mega-heatwave temperatures due to combined soil desiccation and atmospheric heat accumulation, *Nature Geoscience*, 7, 345–349, 2014.
- 440 Mishra, V., Thirumalai, K., Singh, D., and Aadhar, S.: Future exacerbation of hot and dry summer monsoon extremes in India, *npj Climate and Atmospheric Science*, 3, 1–9, 2020.
- Mukherjee, S. and Mishra, A. K.: Increase in compound drought and heatwaves in a warming world, *Geophysical Research Letters*, 48, e2020GL090 617, 2021.
- Naresh Kumar, M., Murthy, C., Sessa Sai, M., and Roy, P.: On the use of Standardized Precipitation Index (SPI) for drought intensity 445 assessment, *Meteorological Applications: A Journal of Forecasting, Practical Applications, Training Techniques and Modelling*, 16, 381–389, 2009.
- Ntegeka, V. and Willems, P.: Trends and multidecadal oscillations in rainfall extremes, based on a more than 100-year time series of 10 min rainfall intensities at Uccle, Belgium, *Water Resources Research*, 44, 2008.
- Oliver, T. H., Marshall, H. H., Morecroft, M. D., Brereton, T., Prudhomme, C., and Huntingford, C.: Interacting effects of climate change 450 and habitat fragmentation on drought-sensitive butterflies, *Nature Climate Change*, 5, 941–945, 2015.
- Perkins, S., Alexander, L., and Nairn, J.: Increasing frequency, intensity and duration of observed global heatwaves and warm spells, *Geophysical Research Letters*, 39, L20 714, 2012.
- Pham, M. T., Vernieuwe, H., De Baets, B., and Verhoest, N. E.: A coupled stochastic rainfall–evapotranspiration model for hydrological impact analysis, *Hydrology and Earth System Sciences*, 22, 1263–1283, 2018.
- 455 Pipoly, I., Preiszner, B., Sándor, K., Sinkovics, C., Seress, G., Vincze, E., Bókony, V., and Liker, A.: Extreme hot weather has stronger impacts on avian reproduction in forests than in cities, *BioRxiv*, pp. 2020–01, 2021.
- Radchuk, V., Turlure, C., and Schtickzelle, N.: Each life stage matters: the importance of assessing the response to climate change over the complete life cycle in butterflies, *Journal of Animal Ecology*, 82, 275–285, 2013.



- Raei, E., Nikoo, M. R., AghaKouchak, A., Mazdiyasi, O., and Sadegh, M.: GHWR, a multi-method global heatwave and warm-spell record and toolbox, *Scientific data*, 5, 1–15, 2018.
- Raffa, K. F., Aukema, B. H., Bentz, B. J., Carroll, A. L., Hicke, J. A., Turner, M. G., and Romme, W. H.: Cross-scale drivers of natural disturbances prone to anthropogenic amplification: the dynamics of bark beetle eruptions, *Bioscience*, 58, 501–517, 2008.
- Reichstein, M., Bahn, M., Ciais, P., Frank, D., Mahecha, M. D., Seneviratne, S. I., Zscheischler, J., Beer, C., Buchmann, N., Frank, D. C., et al.: Climate extremes and the carbon cycle, *Nature*, 500, 287–295, 2013.
- 465 Ridder, N., De Vries, H., and Drijfhout, S.: The role of atmospheric rivers in compound events consisting of heavy precipitation and high storm surges along the Dutch coast, *Natural Hazards and Earth System Sciences*, 18, 3311–3326, 2018.
- Ridder, N. N., Pitman, A. J., Westra, S., Ukkola, A., Do, H. X., Bador, M., Hirsch, A. L., Evans, J. P., Di Luca, A., and Zscheischler, J.: Global hotspots for the occurrence of compound events, *Nature communications*, 11, 1–10, 2020.
- Schumacher, D. L., Keune, J., Van Heerwaarden, C. C., Vilà-Guerau de Arellano, J., Teuling, A. J., and Miralles, D. G.: Amplification of 470 mega-heatwaves through heat torrents fuelled by upwind drought, *Nature Geoscience*, 12, 712–717, 2019.
- Schumacher, D. L., Keune, J., Dirmeyer, P., and Miralles, D. G.: Drought self-propagation in drylands due to land–atmosphere feedbacks, *Nature Geoscience*, 15, 262–268, 2022.
- Seneviratne, S. I., Corti, T., Davin, E. L., Hirschi, M., Jaeger, E. B., Lehner, I., Orlowsky, B., and Teuling, A. J.: Investigating soil moisture–climate interactions in a changing climate: A review, *Earth-Science Reviews*, 99, 125–161, 2010.
- 475 Sharma, S. and Mujumdar, P.: Increasing frequency and spatial extent of concurrent meteorological droughts and heatwaves in India, *Scientific Reports*, 7, 1–9, 2017.
- Soroye, P., Newbold, T., and Kerr, J.: Climate change contributes to widespread declines among bumble bees across continents, *Science*, 367, 685–688, 2020.
- Stillman, J. H.: Heat waves, the new normal: summertime temperature extremes will impact animals, ecosystems, and human communities, 480 *Physiology*, 34, 86–100, 2019.
- Switanek, M., Crailsheim, K., Truhetz, H., and Brodschneider, R.: Modelling seasonal effects of temperature and precipitation on honey bee winter mortality in a temperate climate, *Science of the Total Environment*, 579, 1581–1587, 2017.
- Trenberth, K. E. and Shea, D. J.: Relationships between precipitation and surface temperature, *Geophysical Research Letters*, 32, L14 703, 2005.
- 485 Ummenhofer, C. C. and Meehl, G. A.: Extreme weather and climate events with ecological relevance: a review, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 372, 20160 135, 2017.
- Vandenbergh, S., Verhoest, N., Buyse, E., and De Baets, B.: A stochastic design rainfall generator based on copulas and mass curves, *Hydrology and Earth System Sciences*, 14, 2429–2442, 2010.
- Verhoest, N., Troch, P. A., and De Troch, F. P.: On the applicability of Bartlett–Lewis rectangular pulses models in the modeling of design 490 storms at a point, *Journal of Hydrology*, 202, 108–120, 1997.
- Wells, N., Goddard, S., and Hayes, M. J.: A self-calibrating Palmer drought severity index, *Journal of climate*, 17, 2335–2351, 2004.
- Westerling, A. L., Hidalgo, H. G., Cayan, D. R., and Swetnam, T. W.: Warming and earlier spring increase western US forest wildfire activity, *Science*, 313, 940–943, 2006.
- Wiklund, C., Lindfors, V., and Forsberg, J.: Early male emergence and reproductive phenology of the adult overwintering butterfly 495 *Gonepteryx rhamni* in Sweden, *Oikos*, 75, 227–240, 1996.



- Williams, A. P., Allen, C. D., Millar, C. I., Swetnam, T. W., Michaelsen, J., Still, C. J., and Leavitt, S. W.: Forest responses to increasing aridity and warmth in the southwestern United States, *Proceedings of the National Academy of Sciences*, 107, 21 289–21 294, 2010.
- Wu, Y., Miao, C., Sun, Y., AghaKouchak, A., Shen, C., and Fan, X.: Global observations and CMIP6 simulations of compound extremes of monthly temperature and precipitation, *GeoHealth*, 5, e2021GH000 390, 2021.
- 500 Zaitchik, B. F., Macalady, A. K., Bonneau, L. R., and Smith, R. B.: Europe’s 2003 heat wave: a satellite view of impacts and land–atmosphere feedbacks, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 26, 743–769, 2006.
- Zampieri, M., Ceglar, A., Dentener, F., and Toreti, A.: Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales, *Environmental Research Letters*, 12, 064 008, 2017.
- Zhang, Y., Hao, Z., Zhang, X., and Hao, F.: Anthropogenically forced increases in compound dry and hot events at the global and continental
505 scales, *Environmental Research Letters*, 17, 024 018, 2022.
- Zscheischler, J. and Seneviratne, S. I.: Dependence of drivers affects risks associated with compound events, *Science advances*, 3, e1700 263, 2017.
- Zscheischler, J., Westra, S., Van Den Hurk, B. J., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch, D. N., Leonard, M., Wahl, T., et al.: Future climate risk from compound events, *Nature Climate Change*, 8, 469–477, 2018.
- 510 Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., van den Hurk, B., AghaKouchak, A., Jézéquel, A., Mahecha, M. D., et al.: A typology of compound weather and climate events, *Nature reviews earth & environment*, 1, 333–347, 2020.