

# Air Mass Transport to the Tropical West Pacific Troposphere inferred from Ozone and Relative Humidity Balloon Observations above Palau

Katrin Müller<sup>1</sup>, Peter von der Gathen<sup>1</sup>, and Markus Rex<sup>1,2</sup>

<sup>1</sup>Alfred-Wegener-Institute, Helmholtz Center for Polar and Marine Research, Telegrafenberg A45, 14473 Potsdam, Germany

<sup>2</sup>Universität Potsdam, Institut für Physik und Astronomie, Potsdam, Germany

**Correspondence:** Katrin Müller (katrin.mueller@awi.de)

## Abstract.

The transport history of tropospheric air masses above the remote tropical West Pacific (TWP) is reflected by the local ozone ( $O_3$ ) and relative humidity (RH) characteristics. In boreal winter, the TWP is the main global entry point for air masses into the stratosphere and therefore a key region of atmospheric chemistry and dynamics. However, a long-term in situ monitoring of tropospheric  $O_3$  to assess the variability of TWP air masses and the controlling processes has yet been missing. Our study aims to identify air masses with different origins and pathways to the TWP using and their seasonality using the new time series of ozone and radio soundings ~~from measured in~~ Palau from 2016-2019. ~~Supported by backward trajectory calculations, we found~~ The Palau time series confirms a year-round presence of clean, low  $O_3$  background air masses typical for the region and a seasonal mid-tropospheric cycle in  $O_3$  with a prominent anti-correlation between  $O_3$  volume mixing ratios and RH. We propose five different types of air masses ~~–We further with different tracer characteristics and pathways that can be observed above Palau. For a quantitative assessment of air masses, we~~ defined locally controlled ~~ozone-~~ $O_3$  and RH background profiles for the free troposphere based on monthly statistics and analyzed corresponding anomalies in the ~~5–10~~ 5–10 km altitude range. This analysis was complemented by backward trajectory calculations with the transport module of the Lagrangian chemistry and transport model ATLAS. Our results show a bimodality in RH anomalies. ~~Humid~~ High RH is typically observed for local air masses and associated with convective uplift. Most air masses of low RH experienced descent and indicate a non-local origin. The seasonal occurrence of high  $O_3$  observations is tied to the position of the Intertropical Convergence Zone which controls the transport pathway from potential source regions.

We conclude, that humid and ozone-poor ~~background~~ air masses are of local or Pacific convective origin and occur year-round, but dominate from August until October. Anomalously dry and ozone-rich air masses indicate a non-local origin in Tropical Asia and are subsequently transported to the TWP via an anticyclonic route, mostly from February to April. The geographic location of origin suggests anthropogenic pollution or biomass burning as a cause for ~~ozone production~~  $O_3$  production at the source region. We propose large-scale descent within the tropical troposphere and radiative cooling in connection with the Hadley circulation as ~~being~~ responsible for the dehydration during transport. ~~The~~ Our trajectory analysis revealed no indication of a stratospheric influence. ~~–Our study thus presents a valuable contribution to the discussion about anomalous layers of dry for the dry,~~ ozone-rich air ~~observed in ozone-poor background profiles in the TWP.~~

## 1 Introduction

The Tropical West Pacific (TWP), an area extending from the Maritime Continent to the International Date Line, is considered the major air mass transport pathway from the troposphere to the stratosphere during boreal winter (e.g., Newell and Gould-  
30 Stewart, 1981; Fueglistaler et al., 2004; Krüger et al., 2008). Air masses entering the stratosphere largely originate in the local TWP boundary layer and free troposphere (Rex et al., 2014). Thus, the local tropospheric air composition has a key impact on concentrations of various chemical species within the global stratosphere, even up to a potential impact on polar ozone depletion.

Understanding (1) the variability of composition and transport of air masses to this region as well as (2) the unique air  
35 chemistry are therefore of great relevance. The monitoring of tropospheric ozone ( $O_3$ ) concentrations sheds light on both aspects: (1)  $O_3$  is a chemical tracer for both local convective ("low"  $O_3$  in clean, maritime air, e.g. Kley et al. (1996); Pan et al. (2015)) and long-range transport processes to this region ("high"  $O_3$  from polluted or stratospheric origin, e.g. Anderson et al. (2016); Randel et al. (2016); Tao et al. (2018)). (2) The abundance of ozone is important since the hydroxyl (OH) radical essentially defines the oxidizing capacity of the local troposphere. The close coupling of  $O_3$  and OH in the very low  
40  $NO_x$  environment of the TWP (Levy, 1971) allows estimation of the OH abundance from  $O_3$  and relative humidity (RH) measurements and thus an assessment of chemical lifetimes (e.g., Rex et al., 2014; Nicely et al., 2016; Bozem et al., 2017).

Major research activities focusing on tropospheric air chemistry and  $O_3$  observations have been conducted in the wider region since the late 1980s (e.g., Pan et al., 2017 and overview in Müller, 2020). The typical tropospheric composition of the TWP has been related to the local humid, marine, and pollutant-free environment, favouring  $O_3$  destruction (e.g., Kley et al.,  
45 1996; Levy, 1971):



Observations in the TWP found low  $NO_x$  concentrations inhibiting  $O_3$  production and thus facilitating an  $O_3$  loss rate via the above reactions of 3.4% per day for the tropospheric column (Crawford et al., 1997). For boundary layer  $O_3$  in the  
50 equatorial Pacific the efficiency of this loss mechanism results in a lifetime of around five days (e.g., Liu et al., 1983; Kley et al., 1997). Deep convective outflow and overturning processes lift the clean boundary layer air to the Tropical Tropopause Layer (TTL). In conjunction with a lack of in situ net  $O_3$  production this creates a well-mixed, humid tropospheric profile with a uniform vertical  $O_3$  distribution (e.g., Pan et al., 2015). These typical dynamical conditions are conducive to a respective zonal wave one pattern with a persistent tropospheric  $O_3$  minimum in the TWP in particular (e.g., Thompson et al., 2003b;  
55 Rex et al., 2014).


Many studies highlight dry intrusions of enhanced  $O_3$  volume mixing ratios (VMR) against the humid, ozone-poor background as characteristic features of the mid-troposphere for the wider tropical Pacific (e.g., Newell et al., 1999; Stoller et al.,

1999; Thouret et al., 2000; Browell et al., 2001; Oltmans et al., 2001; Hayashi et al., 2008; Pan et al., 2015; Anderson et al., 2016). The importance of these distinct, anomalous layers for local air composition and climate forcing are often acknowledged (e.g., Mapes and Zuidema, 1996; Kley et al., 1997; Yoneyama and Parsons, 1999). Their genesis, source region and impact on the local radiative budget and oxidizing capacity are, however, the subject of an ongoing debate (Anderson et al., 2016; Randel et al., 2016; Nicely et al., 2016; Tao et al., 2018). Most studies agree, that they have been advected from remote regions and thus indicate a departure from dominating local conditions, i.e. an absence of the local imprint on air composition (see Anderson et al., 2016, for an overview).

65 A long-term monitoring of tropospheric  $O_3$  in the TWP as a key region of stratospheric entry has yet been missing (Smit and Thompson, 2021), but is essential to clarify the source regions of air masses and their annual and interannual variability. All relevant major research campaigns in the TWP were temporarily limited to specific seasons and years (e.g. PEM Tropics/West (Newell et al., 1996), CEPEX (Kley et al., 1996), TransBrom (Krüger and Quack, 2013; Rex et al., 2014), CONTRAST/CAST (Pan et al., 2017; Harris et al., 2017)). To fill the observational gap, the Palau Atmospheric Observatory (PAO) has been  
70 established in 2015 and has since been providing an unprecedented regular balloon sounding program for  $O_3$  and RH. It is located in the center of the warm pool on the island state Palau, 1000 km east of the Philippines (7.34° N, 134.47° E). The instrumental setup and meteorological conditions for the time series used in this study are introduced in more detail in a companion study by Müller et al. (2023) (cf. Müller, 2020).

The aim of this study was to examine major air mass transport processes and pathways to the TWP troposphere by using  
75 balloon-borne  $O_3$  and RH measurements from the PAO. Based on our process understanding explained in Section 3.1.3, we propose five major air mass pathways to Palau resulting in distinct relations between the  $O_3$  abundance and RH in the column of air above the TWP. The continuous 4 years ozone- and radiosondes time series (2016-2019) enables a seasonal analysis of the source regions, reflected mainly in the  $O_3$  variability. We distinguished between air mass categories first using a statistical, data-based approach and performed a seasonal analysis. Lagrangian backward trajectories were then used to examine according  
80 differences in air mass origin. A potential vorticity (PV) analysis was included to investigate a possible stratospheric origin.

In the following, we first introduce the observational data (Sect. 2) and methods (Sect. 3) used to define different categories of air masses, which we relate to different pathways to Palau (Sect. 3.1.1). Main results of this study are the characterization of seasonal tropospheric air mass variability (Sect. 4.1) and the verification of our process-based understanding of air mass transport using trajectory modelling (Sect. 4.2). After a discussion (Sect. 5), we conclude (Sect. 6) that the maximum of the  
85 mid-tropospheric seasonal  $O_3$  cycle can be associated with long-range transport mainly from potentially polluted areas in South East Asia. We found no evidence of transport from the extra-tropical stratosphere. Additional figures with more details on the full seasonal extent are in the Appendix.



**Figure 1.** Tropospheric (0–20 km) time-height cross-section of  $O_3$  VMR (in color-filled contours) and coinciding RH observations below 30 % (hatched areas enclosed in black contours) derived from PAO sounding data; arrows on top indicate individual soundings; data is linearly interpolated between soundings; measurement gaps longer than 20 days are in white, beginning 7 days after/before the last/next sounding; note the non-linear scaling;  $O_3$  VMR are calculated using a pressure dependent background current correction, see Müller et al. (2023) for more details.

## 2 Data

### 2.1 Sondes

90 Our analysis is based on Electrochemical Concentration Cell (ECC) ozonesonde and radiosonde observations from 01/2016–10/2019 conducted at the PAO. Balloon-borne measurements with ECC ozonesondes (Komhyr, 1969; Smit, 2014) are the most practical way to observe and continuously monitor  $O_3$  in situ, especially at more remote sites (e.g., Thompson et al., 2019). The specific instrumentation (models SPC 6A and Vaisala RS92/41) and dataset are introduced and described in detail by Müller et al. (2023), who provide additional meteorological and climatological context for the Palau time series. While carefully following the Standard Operating Procedures (SOP) as recommended by Smit (2014), the  $O_3$  VMR are calculated using a pressure dependent background current correction (see Müller et al., 2023, for details). Figure 1 shows a time-height cross-section of the dataset with individual soundings marked by arrows, ozone in color-filled contours and RH observations below 30 % enclosed in hatched contours. Fortnightly measurements, mostly during mid-day, are complemented by one or two intensive campaigns per year. The onset of measurements in the beginning of 2016 coincides with a very strong El Niño event (e.g., 100 Huang et al., 2016; Diallo et al., 2018). As a consequence and in compliance with acknowledged El Niño Southern Oscillation (ENSO) indices (cf. Müller et al., 2023), data before August 2016 was disregarded in some of the following statistical seasonal analysis and will be referred to as “excluding El Niño 2016”. The 2019 El Niño episode, weaker in nature, is still included.

The sparsity of the  $O_3$  data with a sounding frequency of zero to eleven launches per month and the resulting non-uniform distribution of the data affects the assessment of the temporal variability in our 4-year time series, which is still short for climatological studies. Details about our statistical averaging calculations can be found in the Appendix B. For the trajectory 105 analysis, the dataset was limited to 138 soundings due to missing meta data (timestamps) in some of the soundings and the

vertical profile resolution was reduced by selecting every 10th sonde reading. This dataset will be referred to as the trajectory dataset, including 13627 individual observations in the 5–10 km altitude range.

## 2.2 Meteorological Data

110 Back trajectory calculations were driven by 3D meteorological fields and diabatic heating rates from the European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis data ERA5 (Hersbach et al., 2020) retrieved in a  $1.125^\circ \times 1.125^\circ$  horizontal and 3 h temporal resolution. ERA5 uses a parameterization for convection on the subgrid-scale. Isolated deep convection is not captured, but the net wind flow from each grid cell is zero (Hersbach et al., 2020). While our trajectory model does not calculate convective transport, the ERA5 input therefore implicitly allows us to capture synoptic-scale convective processes.

## 115 3 Methods and Analysis


The identification of differences in air mass origin by local  $O_3$  and RH profile measurements naturally depends on the air mass definition itself. We chose air mass categories derived from statistical analysis of the two tracers,  $O_3$  and RH, from the PAO balloon-borne time series, which reflect our process understanding in this particular location. The chemical composition of the tropospheric column above Palau is governed by the interplay of local and non-local atmospheric processes in time and  
120 space. Our air mass categories represent these differences in chemical or dynamical properties and allow attribution to local or non-local source regions. The tropospheric column of a single day can consist of a variety of air masses, sometimes visible as distinct layers within the tracer profiles.

In the following, we use  $O_3 \pm RH \pm$  as a qualitative notation for air masses of low ("-") or high ("+")  $O_3$  or RH respectively (compare e.g. Stoller et al., 1999). This leads to four qualitative categories, which we use to explain our process understanding  
125 and for comparison with previous studies (Sect. 3.1.1). The " $\Delta$ " symbol denotes quantitative air mass categories by anomalies (positive, "+" or negative, "-") from atmospheric background profiles (i.e. close to zero anomalies, denoted as "o") as defined by our statistical approach. Within this anomaly space, we established a quantitative grid with nine categories with boundaries derived from the distributions of  $O_3$  and RH anomalies,  $\Delta O_3$  and  $\Delta RH$  (Sect. 4.1.3). With this approach, we particularly targeted the questions of local or non-local genesis of background air masses ( $\Delta O_3 \circ \Delta RH \circ$ ), which are humid and ozone-poor,  
130 and anomalously dry  $O_3$ -rich air ( $\Delta O_3 + \Delta RH -$ ) together with their respective seasonality.

### 3.1 Air Mass Definition

#### 3.1.1 Tracers and Process Understanding

$O_3$  is a common tracer in transport studies within the TWP, with typical dynamical conditions being conducive to a tropospheric column of low  $O_3$  VMR attributed to local air masses (e.g., Kley et al., 1996; Pan et al., 2015). Dominant marine convection  
135 leads to low  $NO_x$  concentrations and subsequently low  $O_3$  production rates (e.g., Crawford et al., 1997; Rex et al., 2014). In this environment the  $HO_x$  driven loss of  $O_3$  is favored (see reaction R2) resulting in a lifetime of about five days for boundary




**Figure 2.** Example tropospheric O<sub>3</sub> VMR (black lines) and RH (blue lines) profiles from the PAO for the background atmosphere (a) and anomalously dry ozone-rich layers (b). The ideal shape for the background atmosphere in (a) is illustrated in green and blue shading; two layers disrupting the respective background are indicated by yellow arrows in (b).

layer O<sub>3</sub> (e.g., Liu et al., 1983; Kley et al., 1997). The low NO<sub>x</sub> environment is fostered by dominating easterly winds in the annual mean and most of the troposphere (Müller et al., 2023). Most of the air reaching Palau has crossed the Pacific and has thus been cleaned of O<sub>3</sub> precursors from anthropogenic or other continental pollution. Palau has a hot, humid and wet climate all year, resulting in a continuously high convective activity (Gettelman and Forster, 2002; Müller et al., 2023). This persistent high convective activity creates a well-mixed profile of uniformly low O<sub>3</sub> VMR throughout the free troposphere, as illustrated in Fig. 2a. We refer to this as the clean state of the atmosphere or simply "background".

Layered structures of higher O<sub>3</sub> VMR in the mid-troposphere disturbing the uniform low O<sub>3</sub> profile are often observed and a signature of non-local air masses (e.g., Pan et al., 2015; Anderson et al., 2016; Randel et al., 2016). Figure 2b shows an example profile with two distinct layers of enhanced O<sub>3</sub> levels (> 50 ppb) around 4 and 8 km, interrupting otherwise quite constant levels of low O<sub>3</sub> VMR (~ 25 ppb). Enhanced tropospheric O<sub>3</sub> levels are generated either by photo-chemical processes in high tropospheric NO<sub>x</sub> regimes, i.e. in polluted air masses, or by photo-dissociation of O<sub>2</sub> in the stratosphere and subsequent transport to the troposphere. In situ net O<sub>3</sub> production is unlikely at this altitude in the remote TWP, which is far from pollution sources and shows a lack of lightning activity (Christian, 2003; Cecil et al., 2014). Free tropospheric O<sub>3</sub> life times are in the order of weeks and comparable to the dynamical timescales of long-range transport within the tropics and from the subtropics (e.g., Folkins, 2002; Ploeger et al., 2011). Thus, air masses transported either from polluted areas elsewhere or the extra-tropical stratosphere can retain their high O<sub>3</sub> VMR until arrival in the deep tropics.

To complement O<sub>3</sub> as an indicator of air mass origin, we made use of RH as a measure of vertical displacement of air masses. The benefit of using RH is the readily available data from the combined sonde measurements. The dominant anti-correlation between O<sub>3</sub> and RH in the PAO dataset further suggests closer examination of RH (Figs. 1 and 2). RH allows us to distinguish between different underlying processes affecting either local or non-local air masses (e.g., Hayashi et al., 2008; Schoeberl et al.,



**Figure 3.** Schematic for transport pathways to Palau and the TWP on the zonal plane; major dynamical drivers are marked with arrows (blue colors), transport pathways are color-coded according to the air masses'  $O_3$ / $RH$  characteristics shown in the qualitative grid (turquoise colors for  $O_3^-$ , brown colors for  $O_3^+$ , hue for  $RH_{\pm}$ );  $O_3+RH^-$  air (light brown) masses above Palau may have followed two of the five shown pathways, indicated by a solid and dotted line in the schematic (for details see text).

2015). Local air masses of low  $O_3$  VMR often show high  $RH$  values ( $O_3-RH^+$ ) in accordance with the assumed dominance of convective activity and uplift. The example in Fig. 2a shows, that  $RH$  values characteristically decrease with altitude, but remain greater than 45% throughout the mid-troposphere (cf. Mapes, 2001). Non-local, ozone-rich air masses often correspond to low  $RH$  compared to the air masses above and below the humid tropical column ( $O_3+RH^-$ ). The depressed  $RH$  levels can be explained by either stratospheric origin or various dynamical processes during transport, mostly involving a descent of air masses towards Palau which results in adiabatic compression and heating of the air, reducing  $RH$  at constant absolute humidity (Cau et al., 2007; Dessler and Minschwaner, 2007; Anderson et al., 2016). The study of  $RH$  data thus supports our understanding of processes governing  $O_3$  variability above Palau.

165 The competing hypotheses for the genesis and origin of  $O_3+RH^-$  air demand the use of an additional tracer (e.g. Stoller et al., 1999). In this study, we use potential vorticity (PV) as a dynamical tracer to identify a potential stratospheric origin of air masses. We assume a significant in-mixing of extra-tropical stratospheric air for air masses with an absolute PV greater than 1.5 PVU for at least one day during ten days before arrival in Palau (Waugh and Polvani, 2000; Kunz et al., 2011).

In conclusion, we propose four qualitative categories of air masses and five different pathways identifiable by our tracers, as illustrated in Fig. 3. Ozone-depleted air masses ( $O_3^-$ , turquoise colors) are of local or Pacific, convective origin and ozone-rich air masses ( $O_3^+$ , brown colors) originate from non-local pollution or the stratosphere. High  $RH$  results from a dominant convective uplift of air masses ( $RH^+$ , darker hues), while low  $RH$  indicates a stratospheric origin or dehydration of previously lifted air masses during transport due to clear sky cooling and descent ( $RH^-$ , lighter hues).  $O_3+RH^-$  air is characteristic for two different pathways (solid and dotted light brown lines in Fig. 3), thus representing two different types of air masses.

175 As discussed above, the two important air mass categories are tied to the dominant anti-correlation of  $O_3$  and RH above Palau:  $O_3$ -RH+ background air and  $O_3$ +RH- air masses occurring mostly in layers interrupting the background. Air masses with positively correlated  $O_3$  and RH are observed less often and more difficult to assess.  $O_3$ +RH+ air masses (dark brown) are potentially caused by non-local pollution convectively lifted in the source region and transported rapidly towards Palau on the same altitude, conserving the high humidity. We further suggest, that  $O_3$ -RH- air (light turquoise) consists of clean  
180 boundary layer air convectively lifted in the Pacific vicinity of Palau, dehydrated due to clear sky cooling during transport. Other processes, like remoistening or in-mixing of mid-latitude air during transit to the TWP could also play a role (Cau et al., 2007; Sherwood et al., 2010; Schoeberl et al., 2015; Tao et al., 2018) and are difficult to assess just using the tracer data.

### 3.1.2 Previous Approaches

To detect and assess anomalous layers from balloon and aircraft profiles different methods have been applied (Stoller et al.,  
185 1999; Thouret et al., 2000; Hayashi et al., 2008; Pan et al., 2015). Stoller et al. (1999) used data from three NASA campaigns (PEM-Tropics, PEM West A and B) to calculate a free tropospheric background mode for various atmospheric constituents ( $O_3$ ,  $H_2O$ ,  $CO$ ,  $CH_4$ ) and for individual profiles. Using a spike detection approach, they produced extensive statistics for the frequency of anomalous layers in the whole tropical Pacific region for two different seasons. The same methodology was applied to MOZAIC aircraft data by Thouret et al. (2000). Its main disadvantage lies within an arbitrary threshold used to de-  
190 termine the background mode. Despite the statistical uncertainty, Stoller et al. (1999) recognized the importance of anomalous layers in tropical Pacific profiles and emphasized their role in atmospheric chemistry modeling due to their frequent occurrence.

A similar analysis of longer time series from three SHADOZ sites (Southern Hemispheric Additional Ozonesondes Network Thompson et al., 2003a, b, 2012, 2017) confirms the frequent occurrence of dry enhanced  $O_3$  layers below 12 km (Hayashi et al., 2008). Using a statistical method for the layer detection, the study found respective layers in approximately 50 % of  
195 profiles per year and station with differing seasonal variations. The study relies on the separation of a wet and distinct dry season at the respective stations, which is not applicable for the Palau site (cf. Müller et al., 2023).


Pan et al. (2015) isolated the low  $O_3$  background or “primary mode” by removing all “dry” data with RH less than 45 % for all observations during the CONTRAST campaign (Pan et al., 2017). They thus refrained from resolving the individual vertical structure of layers, but proposed the RH threshold as an overall criterion in the free troposphere to separate local and  
200 non-local air masses. Pan et al. (2015) emphasized the possible variability of air mass origin within the individual vertical column, which is obscured in seasonal or even annual profile averages, i.e. layered structures in the  $O_3$  profile hidden within the typical so-called “S”-shape (cf. Sect. 4.1). Their method, however, seems limited to the specific sites and season of the CONTRAST campaign, as it does not yield a robust output for the Palau data during all seasons (Müller et al., 2023).

### 3.1.3 Statistical Definition of Background and Anomalies

205 For a quantitative definition of air masses above Palau, we first determined background profiles in the free troposphere from the PAO  $O_3$  and RH time series. These profiles represent humid, ozone-poor, local air masses that are controlled by convective influence and not by long-range transport. This was performed on a statistical and monthly basis and aimed to identify profile



### March (16 Profiles)



**Figure 4.** O<sub>3</sub> VMR and RH statistics for the month of March as example for Palau free tropospheric profiles as a function of altitude: the 20th quantile (Q20) for O<sub>3</sub> is shown in green and the 83.3th quantile (Q83) for RH in blue (both vertically smoothed using exponentially weighted moving averages); the median is marked in grey and the central 66.6 % range in grey horizontal bars; number of included individual profiles in brackets, for orientation 20 ppb O<sub>3</sub> VMR and 45 % RH are marked in black (see also Figs. A3 and A4).

shapes similar to the signature profiles in Fig. 2a. For the monthly O<sub>3</sub> VMR background profiles we chose the 20th quantile (Q20) profiles. Q20 guarantees sufficiently low O<sub>3</sub> values to avoid a “belly” shape of the profiles, typical for tropical average  
 210 O<sub>3</sub> profiles. For the monthly RH background profiles we chose the 83.3th quantile (Q83) profiles. Q83 is the upper boundary of the central 66.6 % range, guaranteeing high humidity. Fig. 4 shows an example of these background profiles for March. The monthly quantile profiles have been vertically smoothed using first a 1-km-binning and then exponentially weighted moving averages with altitude (see also Figs. A3 and A4). Fig. 4 includes the median and central 66.6 % ranges for both tracers separately and is calculated from 16 individual profiles (see Müller, 2020, for more details).

215 Our method roughly followed the approach of Hayashi et al. (2008). However, Hayashi et al. defined air masses above the 83.3th quantile as O<sub>3</sub> enhanced layers. With the Q20 limit for O<sub>3</sub> we used a less conservative approach, because the Palau background atmosphere is characterized as a uniform, well-mixed low O<sub>3</sub> profile, caused by uplift of ozone-poor boundary layer air in active convection. We chose this particular O<sub>3</sub> quantile to yield the most uniform, “straight-line” profile in the free troposphere for all monthly averages of the given time series. This essentially smoothes out any interrupting layers occurring  
 220 at varying altitudes in individual profiles. While Hayashi et al. analyzed RH in ozone-enhanced anomalous layers in contrast to the air masses above and below, we applied the statistical approach to both tracers and chose Q83 for RH to detect typical convection associated with the background. The resemblance of both the particular low O<sub>3</sub> and high RH quantiles with the example of an individual background profile in Fig. 2a is evident. The monthly averaging accounts for the seasonal variability of the background as the uniform O<sub>3</sub> profile shifts towards higher base VMR from summer/fall to winter/spring (compare Figs.  
 225 2 and A3).

In a second step, we determined the anomalies against the monthly background profiles, denoted in the following by  $\Delta O_3$  and  $\Delta RH$ . These are calculated for individual measured O<sub>3</sub>/RH pairs within each sounding and in their respective month-

altitude bin. Within the anomaly space, we define a new background air mass category, " $\Delta O_3 \circ \Delta RH \circ$ ", with close to zero anomalies in both tracers. We further denote eight other quantitative air mass categories with combinations of positive ("+"), negative ("-") or close to zero ("o") anomalies for both  $\Delta O_3$  and  $\Delta RH$  individually (Sect. 4.1.3). In this study however, we limited further analyses to the two most interesting quantitative air mass categories,  $\Delta O_3 \circ \Delta RH \circ$  and  $\Delta O_3 + \Delta RH -$ , i.e. per definition humid, ozone-poor background air masses and anomalously dry, ozone-rich air masses. In contrast to Hayashi et al. we did not assess the vertical structure of the anomalous layers in individual profiles. Hence, neither the layer thickness nor the layer position within the profile were considered. Any measured data point of a profile was quantitatively attributed to one of nine air mass categories, which were then analyzed "in bulk". In our following analysis, we further focused on the altitude region between 5 and 10 km, where the weakest cloud-mass divergence occurs, i.e. the weakest convective detrainment (Folkins and Martin, 2005) and the greatest frequency of anomalous layers.


### 3.2 Trajectory Analysis

We deployed the trajectory module of the fully Lagrangian chemistry and transport model ATLAS (Wohltmann and Rex, 2009; Wohltmann et al., 2010) driven by ERA5 data (see Sect. 2). The model uses a hybrid vertical coordinate, which gradually transforms from pressure at the surface to potential temperature in the stratosphere. The corresponding vertical velocities change from vertical winds in pressure coordinates to diabatic heating rates, respectively.

10-days backward trajectories with a time step of ten minutes were initialized at the location and time of every tenth sonde reading of a profile (approximately every 50 m, i.e. 20 trajectories per km) for all profiles within the trajectory dataset. They are considered as a footprint for transport pathways towards Palau. We assume  $O_3$  to be a passive tracer during transport, i.e. did not implement chemical reactions. This assumption is justified for dry, mid- to upper tropospheric air because of photochemical lifetimes of  $O_3$  in the order of several weeks to months under these conditions. In the wet lower troposphere and marine boundary layer this approach is only valid for a few days, because of  $O_3$  lifetimes in the order of days (e.g., Thompson et al., 1997). Convection is not treated explicitly in the model setup, but partly reflected in the reanalysis data. Anderson et al. (2016) found air parcel ages of around 10 days in the TWP in winter 2014, when stopping their trajectories at the point of last precipitating convection based on satellite observations of cloud top height and precipitation. We therefore assume no "reset" of the composition of our air mass within five days and stop our trajectories five days before arrival to identify the origin.

## 4 Results

We first present the variability of the PAO tropospheric  $O_3$  time series to assess air mass seasonality and the application of our method to determine different groups of air masses. Then, our results from the trajectory analysis are shown, first investigating general seasonal tracer variability, and second combining these results with our method to identify different air masses (Fig. 3). A physical interpretation of the results follows in the discussion (Sect. 5).



**Figure 5.** Monthly means (a), anomalies from the annual mean (b) and the annual mean profiles (c) for individual years (thin colored lines) and for the whole time series excluding El Niño 2016 (thick black line) with standard deviation for the whole time series (grey dashed lines) and relative standard deviation (RSD) (grey dotted line); individual soundings are marked as arrows above with different colors for different years; non-linear scaling is used in (a) to match the different orders of magnitude in the lower and upper troposphere (compare Müller et al., 2023, Fig. 6)

## 4.1 Air Mass Variability


### 4.1.1 Annual O<sub>3</sub> Variability

260 Figure 5 shows time-height cross-sections of the annual variability of tropospheric O<sub>3</sub>. Figure 5 (a) shows monthly means, (b) anomalies from the annual mean profile and (c) the annual mean profile of the overall time series (black solid line), excluding El Niño 2016. Panel (c) also shows the standard deviation (SD) and the relative standard deviation (RSD) of the overall annual mean profile (dashed and dotted lines, respectively), as well as the annual mean profiles of individual years (colored lines). As the fraction of the SD relative to the annual mean, the RSD can be considered as a measure of variability with altitude (cf.

265 Ogino et al., 2013). The annual mean tropospheric O<sub>3</sub> VMR profile (Fig. 5c) has an "S"-shape typical for tropical sounding stations (e.g., Folkins, 2002; Thompson et al., 2003a, b; Pan et al., 2015). O<sub>3</sub> VMR are lowest in the boundary layer (< 20 ppb) and also low between 10 and 12 km (< 30 ppb). They increase in the mid-troposphere to about 35 ppb and again above 12 km towards their stratospheric maximum, which is an order of magnitude higher. The interannual variability of the annual mean O<sub>3</sub> profile is low for all years with the exception of 2016.

270 The monthly O<sub>3</sub> VMR (Figs. 5a and b) reveal two dominant signals: an annual cycle in the 5–10 km altitude range with the maximum from February until April (30–60 ppb) 2–3 times greater than the minimum from July until October (10–30 ppb), and a reverse, strong cycle in the TTL with maximum anomalies (> 40 ppb) from June until September. The peak in RSD of 50 % at about 17 km reflects the strong TTL cycle. The enhanced variability in mid-tropospheric O<sub>3</sub> (RSD ~ 30 %) corresponds with the cycle revealed by the monthly means. Annual variations in O<sub>3</sub> observations are smallest between 10-12

275 km and especially in the boundary layer (RSD ~ 20 %), thus coincidental with the O<sub>3</sub> minima of the annual profile. The regionally typical ozone-poor background is clearly subject to seasonal variations.



**Figure 6.** Seasonal mean O<sub>3</sub> VMR profiles (colored solid lines) in comparison to the annual mean (dashed black line), all excluding El Niño 2016, for Palau seasons November–December–January (NDJ), February–March–April (FMA), May–June–July (MJJ) and August–September–October (ASO).

#### 4.1.2 Seasonal O<sub>3</sub> Profiles

A more detailed analysis of the seasonal O<sub>3</sub> variability helps to identify recurring seasonal characteristics and differences in controlling processes like synoptic or meteorological conditions. Therefore, we divided our time series into four seasons, shifted  
 280 by one month compared to the temperate climate seasons: November–December–January (NDJ), February–March–April (FMA), May–June–July (MJJ) and August–September–October (ASO). These Palau seasons reflect different influences dependent on the time of the year and were chosen empirically by sorting monthly O<sub>3</sub> profiles (Müller et al., 2023, Fig. A1) by similar shape, considering the full free-troposphere. The resulting four most different types of profile shapes are shown in Fig. 6. The Palau seasons turned out to be centered around the equinoxes, which is reasonable for a tropical station.

285 In the 5–10 km altitude range FMA and ASO show the largest differences and represent the extremes of the mid-tropospheric cycle. The "S"-shape of the annual mean profile prevails in the winter seasons, i.e. in NDJ and most pronounced in FMA. The interannual variability is particularly high during FMA from 5 to 10 km with generally enhanced O<sub>3</sub> VMR between 40 and 50 ppb (Fig. A1). The ASO profile is closest to the uniform background profile (20–25 ppb O<sub>3</sub> VMR) with little interannual variability, while having the highest sampling rate (59 profiles). We thus refer to ASO as the background season. In the TTL,  
 290 both extreme seasons, ASO and FMA, show a pronounced O<sub>3</sub> minimum, except during the 2016 El Niño FMA season (Fig. A1). The steep onset of higher levels of (stratospheric) O<sub>3</sub> VMR in ASO occurs at lower altitudes compared to the NDJ and FMA seasons.

MJJ and NDJ can be considered as intermediate seasons with respect to the mid-tropospheric cycle, i.e.  $O_3$  values here are in between the minimum and maximum. The NDJ profile resembles the annual mean profile up to about 14 km, while the MJJ profile diverges from the annual mean above 10 km towards higher values. The resulting tilted line shape during MJJ, compared to the straight line profile during ASO, implies a more gradual increase of  $O_3$  in the UTLS (cf. Müller et al., 2023) and a lack of a pronounced  $O_3$  minimum in the UT. In MJJ, the tropopause is at its lowest altitude of the year, thus is the occurrence of high levels of stratospheric  $O_3$  VMR.

#### 4.1.3 $O_3$ and RH: Background and Anomalies


The application of our method to separate air masses on the full PAO time series for the free troposphere (3–14 km) yields the distribution of anomalies from the background profiles,  $\Delta RH$  and  $\Delta O_3$ , given in Fig. 7 (see also Sect. 3.1.3). The 2D-histogram in Fig. 7a shows  $\Delta O_3$  versus  $\Delta RH$  from all months and altitudes as percentages of the total count of all measured free-tropospheric  $O_3/RH$  data pairs in grey-scale. Data along the zero lines within this anomaly space indicate zero anomalies and thus an attribution to the respective background profiles, Q20 of monthly  $O_3$  VMR and Q83 of monthly RH values. The marginal 1D-histograms show the distributions of both tracers individually normalized to unity.

Both  $\Delta RH$  and  $\Delta O_3$  show heavily skewed distributions, with a long tail in the  $\Delta O_3$  distribution towards less frequent high  $O_3$  occurrences and even a bimodal distribution for  $\Delta RH$  with a secondary maximum in the tail towards dry air masses. These distributions motivate the separation of air masses into three domains for both parameters, one corresponding to the low tail, one to the bulk of the observations and one corresponding to the high tail. Overall this leads to nine quantitative air mass categories in a three times three grid referred to as the quantitative grid (Fig. 7b). The boundaries were chosen empirically with respect to the shape of the distributions with a range of  $-5/+15$  ppb  $\Delta O_3$  VMR and  $-20/+5$  % for  $\Delta RH$  for the central group.


In particular, the correlated occurrence of air masses with high  $O_3$  and low RH ( $\Delta O_3+\Delta RH-$ ) stands out as a low, but separate population in Fig. 7a (lower right quadrant), while the central group ( $\Delta O_3\circ\Delta RH\circ$ ) represents the most frequent conditions and is referred to as background category (not to be confused with the background profile), since anomalies from the background profiles are smallest. In this study, we focus on the distinct population of  $\Delta O_3+\Delta RH-$  air masses in contrast to this background category.  $\Delta O_3\circ\Delta RH\circ$  air masses are characterized by significantly higher RH and lower  $O_3$  and are therefore also referred to simply as the humid, ozone-poor background.

The anomalies are mostly clustered within 20 ppb along the  $\Delta O_3$  zero line indicating an overall dominance of ozone-poor air masses. Apart from the tail towards less frequent higher  $\Delta O_3$ , the narrow distribution is nearly Gaussian and centered around 0 ppb  $\Delta O_3$  VMR with a tendency towards positive values. The majority of  $\Delta RH$  values appears between  $-60$  and  $20$  %. The bimodality of the  $\Delta RH$  distribution is persistent in all seasons with the exception of MJJ (Fig. A5). The primary, dominant mode is centered slightly off the  $\Delta RH$  zero line towards negative values, i.e. the majority of measured RH values matches the Q83 or slightly lower values of the respective month-altitude bin. The secondary mode occurs for  $\Delta RH$  values 40 % lower than their background profiles estimates.

Figure 8 shows the  $\Delta O_3/\Delta RH$  distribution for the seasons FMA and ASO. As expected from the previous analysis,  $\Delta O_3+\Delta RH-$  are almost absent in ASO (Fig. 8a), emphasizing that the season represents an overall ozone-poor, mostly humid




**Figure 7.** (a) Free-tropospheric (3–14 km) relation between tracer anomalies  $\Delta O_3$  VMR and  $\Delta RH$  from monthly background profiles defined in Section 3.1; color shading indicates percentages of total count of measured  $O_3/RH$  data pairs per grid point; marginal plots for individual tracer anomaly distributions are normalized to unity; dashed lines refer to quantitative air mass categories in the three times three quantitative grid as illustrated in (b), with boundary values for the central grid box ( $\Delta O_3\circ\Delta RH\circ$ ) of  $-5/+15$  ppb for  $\Delta O_3$  VMR and  $-20/+5$  % for  $\Delta RH$ ; two target categories of the analysis are highlighted in color in (b), compare to Fig. 3.



**Figure 8.** Seasonal free-tropospheric (3-14 km) relation between tracer anomalies  $\Delta O_3$  VMR and  $\Delta RH$ , total number of datapoints in brackets, colors according to Fig. 6, more details in Fig. 7 and 6.

background in the free troposphere. In contrast to the seasonal mean FMA  $O_3$  profile with elevated  $O_3$  levels in the mid-troposphere (Fig. 6), the FMA anomalies distribution (Fig. 8b) reveals a dominance of background category air masses over  $\Delta O_3+\Delta RH-$  air masses within the free troposphere. A statistical view on the seasonal occurrence of different air masses in the mid-troposphere emphasizes the year-round dominance of  $\Delta O_3\circ\Delta RH\circ$  air, which were observed in more than 70 % of profiles within each season (Fig. A6).




**Figure 9.** Geographical footprint of air masses: 10-days backward trajectories from ATLAS arriving at Palau in the 5-10 km altitude range by season (2016-2019); colored segments represent time periods of maximal three (green), five (purple) and 10 (orange) days backwards, respectively; the x marks the location of Palau; only a subset (every 20th data point of a profile, roughly one per km) of the trajectory dataset is shown, see Fig. 6 for seasons.

## 4.2 Air Mass Transport and Processes

### 4.2.1 Trajectory Origin by Season

In the following, we focus on the 5–10 km altitude range with the most frequent occurrence of  $\Delta O_3 + \Delta RH$ - air masses and the largest seasonal differences in  $O_3$  VMR (cf. Fig. 6 and Sect. 3.1.3). A footprint of air mass transport to Palau is analyzed using 10-days backward trajectories for the study period (2016–2019) sorted by season arriving in Palau in the 5–10 km altitude range (Fig. 9). For better visualization, a representative subset was chosen, displaying only every 20th trajectory of a profile (roughly 1 trajectory per km). Colored line-segments show three different time periods backwards, three, five and 10 days, thus indicating the velocity of air mass transport.

Most air masses reach Palau from the east, but have traveled two different paths on different timescales. The seasonal distinction of trajectories roughly separates an Eastern Pacific pathway, dominating in MJJ and ASO, from an anticyclonic route in NDJ and FMA, connecting Palau with South East Asia and some remote areas as far as the African continent. The



**Figure 10.** Origin of air masses by season: Location of air masses five days before their measurement in Palau in the 5-10 km altitude range inferred from trajectories for Palau seasons FMA (upper row) and ASO (lower row); color-coded by  $O_3$  VMR (a), RH (b) and by difference in pressure altitude, between measurement and five days before measurement (c); black contours in (a) show distribution density and schematic arrows in (b) indicate the pathway as shown by the footprint in Fig. 9; see Fig. 6 for seasons.

anticyclonic route can be associated with long-range transport on shorter timescales compared to the Pacific route. As these transport patterns are most pronounced in FMA and ASO, respectively, which also exhibit the greatest tracer differences in the mid-troposphere (compare Fig. 6), we focus on these two Palau seasons in the following.

The geo-spatial extent of the footprints differs significantly for season FMA and ASO. While all trajectories stay mostly within the tropical zone between  $0^\circ$  to  $30^\circ$  N, trajectories during FMA have a wider longitudinal extent than during ASO. Most ASO air masses never left the Pacific Ocean area within 10 days before arrival in Palau and have traveled shorter distances. Only a few trajectories on a Southern hemispheric cyclonic route show the influence of the Western Pacific monsoon, which is active from July to October, but governs air mass transport mostly below 5 km altitude (Müller et al., 2023).

Figure 10 shows the location of air masses five days before their measurement in Palau inferred from trajectories for the seasons FMA (top) and ASO (bottom) as an indication of the origin of the air masses. The geo-spatial probability density function of data points is highlighted as black contour lines in Fig. 10a. Here, the colors indicate  $O_3$  VMR as observed upon arrival in Palau. Figures 10b and c show the same locations as Fig. 10a, but colored-coded by RH (b) or difference in pressure altitude (c). Red colors in panel (c) indicate descent, blue colors ascent towards Palau.

The dominant source region of free-tropospheric Palauan air masses, indicated by the center of the density distribution in Fig. 10 a, is located east of Palau during both seasons and corresponds to low  $O_3$  VMR ( $\leq 30$  ppb). During FMA, this eastern center of the distribution is actually split in two parts along a latitudinal axis. The dominant part is located further south on the equator, exhibiting low  $O_3$  VMR. The northern part of the eastern center is of higher  $O_3$  VMR. A separate cluster of trajectory points showing enhanced  $O_3$  VMR ( $\geq 60$  ppb) exists northwest of Palau during FMA, roughly extending from India to Taiwan. As revealed by the full 10-days trajectories in Fig. 9b, all air masses of higher  $O_3$  VMR during FMA can be related to the anticyclonic route, though some took longer than five days to travel from South East Asia to Palau. During ASO the overall



unimodal distribution is centered on Palau's latitude, 20° to the East, and observations of enhanced O<sub>3</sub> VMR originate outside the main cluster. For both seasons, air masses with O<sub>3</sub> VMR greater than 60 ppb rarely originate south of Palau.

365 The distinction between the two pathways is also reflected in RH (Fig. 10b) and even more clearly in the vertical displacement of air masses from origin to destination (Fig. 10c). In FMA, Fig. 10c shows descended air masses originating north and ascended air south of Palau. Here, the descent of air masses corresponds with greater O<sub>3</sub> VMR and lower RH, although the pairwise correlation, R, of these two parameters with the pressure height difference is not strong (R=-0.53 for O<sub>3</sub>VMR and R=0.44 for RH, for the latter cf. Fig. A7). We can, however, associate these dry, descending air masses with the anticyclonic  
370 route shown in the 10-days backward trajectory footprint.


For ASO air masses, the picture is not as clear, but ascent and humid air masses with low O<sub>3</sub> VMR are slightly dominating (Figs. 10c and A7b) and have likely followed the Pacific pathway. The pairwise correlation for pressure height difference and O<sub>3</sub> VMR or RH, respectively, is even lower than during FMA (R=-0.36 for O<sub>3</sub>, R=0.35 for RH). However, there are clearly two  
375 main groups of air masses, one with RH below 40 %, descended mostly only 1 km, and one group with RH centered around 70 %, ascended mostly 1 km or more.

The investigation of the history of PV for Palauan air masses revealed no stratospheric pathway. Below 1 % of all 13627 trajectories stayed above 1.5 PVU for more than a day during 10 days transit to Palau. Within five days before arrival only few trajectories crossed the 1.5 PVU threshold.

#### 4.2.2 Trajectory Origin by Air Mass

380 While we have looked at the origin of the trajectories as a function of season in the previous section, we now turn to the origin as a function of air mass. The classification of trajectories by quantitative air mass category as defined in Section 3.1 yields the distribution of source regions as shown in Fig. 11. Background  $\Delta O_3 \circ \Delta RH \circ$  air masses originate within a compact region in the Pacific, mostly east of Palau stretching as far as 180° E. The origin pattern for  $\Delta O_3 + \Delta RH$ - air masses clearly show transport from the South East Asian region and along the anticyclonic path to Palau. In comparison to the seasonal analysis, the  
385 classification of trajectories by air masses shows a clear distinction for the history of vertical displacement of the air masses. On their way towards Palau most  $\Delta O_3 \circ \Delta RH \circ$  air masses have been ascending, while  $\Delta O_3 + \Delta RH$ - air has descended.

A seasonal analysis of the trajectories sorted by air mass anomalies reveals some differences in the source regions of  $\Delta O_3 + \Delta RH$ - air masses (Fig. A8). The origin hot spots around India and the Southeast Asian peninsula are almost exclusively observed during FMA. The Southern Philippines and the area slightly North East of Palau are equally dominant source  
390 regions in this season. In ASO,  $\Delta O_3 + \Delta RH$ - air masses are less frequently observed and their origins are clustered around three main areas: 1. north east of Palau in the West Pacific, 2. between Borneo and the Southern Philippines and 3. in the New Guinea region, extending towards Northern Australia.



**Figure 11.** Origin of air masses by air mass anomalies: Location of air masses five days before their measurement in Palau in the 5–10 km altitude range inferred from trajectories for  $\Delta O_3 \circ \Delta RH \circ$  (a) and  $\Delta O_3 + \Delta RH$ - (b) air masses, as indicated by the pictograms; color-coded by difference in pressure altitude between measurement and five days before measurement, see also Figs. 7 and 10.

## 5 Discussion

The central goal of our study was to identify the air mass origin of TWP air and its seasonality by means of the observed  $O_3/RH$  relation above the PAO. We proposed a qualitative transport scheme for different types of air masses ( $O_3 \pm RH \pm$ , Fig. 3) that distinguishes between local and non-local processes. Lagrangian backward trajectories support this hypothesis. Trajectories with different air mass characteristics, defined by their  $O_3$  and RH anomalies from statistically defined background profiles, show different geographical origins and transit properties. The PAO 4-year time series reveals the seasonality of air mass types and thus origin. Our analysis thereby confirms the usefulness of  $O_3$  and RH as combined tracers for air mass origin in the TWP.

### 400 5.1 Seasonality and Air Mass Definition

The PAO time series provides unprecedented insight into the seasonality of tropospheric air composition in the TWP (Müller, 2020; Müller et al., 2023). An important observation from the PAO dataset is the dominant anti-correlation between  $O_3$  and RH (Fig. 1). The annual cycles of  $O_3$  in the mid- and upper troposphere were expected from satellite observations and previous studies, especially the zonal wave one pattern in the TTL (e.g., Thompson et al., 2003b; Randel et al., 2007), but had not been monitored continuously with in situ measurements before. The high temporal and vertical resolution of the ozonesonde dataset revealed day-to-day variations, but also the prevalence of significant seasonal signals modulating these. Especially the increased sampling frequency during the extremes of the mid-tropospheric cycle shows the robustness of this signal for Palau and the dominance of the local low  $O_3$  mode during late summer (Fig. 6). Currently, the shortness of the time series and statistical bias due to the differences in monthly sampling frequencies have to be taken into account. A future analysis of the growing time series will be better able to assess the influence of interannual variability. The ENSO cycle will likely play the most important role in this.

The mid-tropospheric  $O_3$  cycle is in accordance with the annual movement of the Intertropical Convergence Zone (ITCZ). The occurrence frequency of either a clean convective (local) state or a non-local state with long-range transport to the TWP

is regulated by the two major barriers ITCZ and Southern Pacific Convergence Zone (SPCZ). Müller et al. (2023) showed  
415 the correspondence of the ITCZ's position north of Palau with the minimum in  $O_3$  measurements (cf. Fig. 5). This relates to  
previous studies on air mass transport to the region: during the PEM-Tropics West B campaign, Browell et al. (2001) found  
that layers of high  $O_3$  VMR advected to the TWP from remote polluted areas are mostly confined north of the ITCZ. Very  
low  $O_3$  values were only present in the "equatorial wedge" between ITCZ and SPCZ and associated with enhanced vertical  
mixing in convection in the absence of gross ground pollution. Palau is enclosed between these bands during ASO (Sun et al.,  
420 2023), therefore the best season to observe the clean air background. During FMA, when the ITCZ is located furthest south,  
transport from higher northern latitudes is possible. MJJ and NDJ can be considered as intermediate seasons with respect to  
both, the mid-tropospheric cycle and the similar location of the ITCZ during its annual crossing. This feature explains, why  
studies related to the CONTRAST campaign which took place in January/February 2014 did not see low  $O_3$  extremes (Pan  
et al., 2015; Newton et al., 2016; Nicely et al., 2016; Tao et al., 2018), and why background conditions of the region need to  
425 be assessed by long-term measurements. Its unique location and year-round operation makes the PAO an excellent background  
site to study the influence of dynamics (cf. Thompson et al., 2021; Sun et al., 2023).

The strong TTL cycle as a zonal phenomenon can be mostly explained by the interplay between Brewer-Dobson and Hadley  
circulation (cf. Müller et al., 2023). Mass fluxes for both circulation regimes become comparable at around 16 km altitude  
(e.g. Folkins, 2002; Pan et al., 2014). The minimum in the vertical  $O_3$  profile in the UT (the so called chemopause) can be  
430 attributed to deep convective outflow. Recognized as a characteristic feature of tropical profiles, low UT  $O_3$  is often used as an  
indicator for deep convective detrainment (e.g., Kley et al., 1996; Folkins, 2002; Solomon et al., 2005; Gettelman et al., 2009;  
Paulik and Birner, 2012). Satellite observations confirm year-round convective activity in Palau, with some variability due to  
the traverse of the ITCZ twice a year (Gettelman and Forster, 2002; Müller et al., 2023). Müller et al. (2023) associate the  
occurrence of the strongest winds below 15 km when the ITCZ is furthest south or north of Palau with the poleward branch of  
435 the Hadley circulation. The low variability in UT  $O_3$  (Fig. 5c) can hence be explained by the persistence of deep convection  
and the level of convective outflow in association with the Hadley and Walker circulation (e.g., Folkins, 2002; Solomon et al.,  
2005; Takashima et al., 2008). The months of May and June stand out in this context with higher  $O_3$  levels between 12 and  
15 km thus lacking a chemopause (Fig. 5). The dry environment at this altitude increases  $O_3$  lifetimes and makes in situ  $O_3$   
destruction unlikely, although it is discussed as a consequence of convectively injected water vapor (Schoeberl et al., 2015;  
440 Anderson et al., 2016). The PAO dataset is predestined for future studies on the seasonality of deep convective outflow and  
stratosphere-troposphere exchange (STE) processes in the TTL in this region of major entry into the stratosphere in boreal  
winter.

The specific classification of seasons used in this study corresponds to distinct dynamical conditions. In particular, the  
Palau seasons can be related to the movement of the ITCZ, the main driver of dynamical variability. Our interpretation of  
445 the profile shapes is supported by the separation of trajectory footprints in Fig. 9, i.e. different dominating pathways for the  
different seasons, and a comparison with an analysis of SHADOZ station profiles from Java and American Samoa by Stauffer  
et al. (2018). They used a sophisticated clustering technique to characterize ozonesonde profile variability related to different  
controlling processes. Out of four cluster profiles, the low  $O_3$  cluster associated with convective lifting of low  $O_3$  VMR

compares well with the Palau ASO profile. The cluster with highest average  $O_3$  VMR exhibits a similar "tilted line" shape like  
450 Palau MJJ with a lower tropopause and weak gradient, explained by Stauffer et al. (2018) with STE. This corresponds well  
with the minimum annual tropopause height and a shift in wind regimes due to the crossing of the ITCZ around June (Müller  
et al., 2023). The horizontal wind is weakest throughout the entire tropospheric column in May, as the subtropical ridge shifts  
south with increasing altitude, which could favor quasi-horizontal transport of extra-tropical stratospheric air into the TTL.  
July can be seen as a transition month, as the tropopause is comparable with May and June, but mid-tropospheric  $O_3$  VMR are  
455 as low as observed in August and September (cf. Fig. A1 in Müller et al., 2023). In July, the Western Pacific monsoon already  
reaches Palau in lower altitudes in the form of equatorial westerlies, but the tropopause is still low. While the construction  
of 3-month-seasons remains arbitrary in any case, it could be reconsidered especially for the summer months (MJJ, ASO) to  
further differentiate between these processes.

Our new definition of humid, ozone-poor background and anomalously dry ozone-rich air masses in relation to statistically  
460 defined background profiles allows a separation of air mass origin based on backward trajectories. Looking at anomalies,  $\Delta O_3$   
and  $\Delta RH$ , our method provides an improved criterion compared to using fixed thresholds based on absolute tracer values, as  
have been used in previous studies (e.g. Anderson et al., 2016; Nicely et al., 2016; Randel et al., 2016; Tao et al., 2018), who  
all defined  $O_3+RH$ - air masses with  $RH < 20\%$  and  $O_3 \text{ VMR} > 40$  ppb. Our approach using quantiles based on a 4-year time  
series focuses on a physically motivated background definition. A comparison between air mass categories based on absolute  
465 thresholds and our quantitative air mass categories for the PAO data set shows an improved separation of processes in the  
trajectory analysis, namely between ascent and descent of air masses (not shown here). The main advantage of our method  
compared to algorithms of layer detection in individual profiles (e.g., Stoller et al., 1999) lies within its simplicity and the  
consideration of the identified bimodality in the RH anomalies. In principle, it is also applicable for sounding data time series  
from other tropical stations. However, it may not yield the best results as atmospheric processes and mechanisms determining  
470 air composition might be specific to this geographic region.

The bimodality in RH anomalies is a striking result of our study. It further justifies our separation between  $\Delta RH_0$  and  
 $\Delta RH$ - air masses in the quantitative grid classification (dashed lines in Figs. 7 and 8), although the exact definition of the  
grid boundaries could be revisited. Global probability distribution functions of RH from observations have already been shown  
as bimodal for different tropospheric altitudes in tropical regions, in particular within the ascending branch of the Hadley  
475 circulation (e.g., Zhang et al., 2003; Ruzmaikin et al., 2014). Our results fit into the findings of these studies. Other studies  
e.g. by Sherwood et al. (2006, 2010) earlier disputed the original claim of a common bimodality in RH distributions by Zhang  
et al. (2003). The PAO dataset now adds to this discussion. However, a comparison of the different methods arriving at RH  
distributions would be necessary to draw further conclusions.

A classification of free-tropospheric air mass anomalies by seasons in Fig. 8 shows the absence of  $\Delta O_3+\Delta RH$ - air in ASO  
480 and MJJ, as already inferred from the seasonal mean profiles (Fig. 6), compared to their prominent appearance in NDJ and  
FMA (see marginal 1D histograms for  $\Delta RH$  in Figs. A5 and 8). During the NDJ and MJJ, both maxima in  $\Delta RH$  are of similar  
amplitude, which emphasizes the importance of dry air masses intruding into the otherwise humid troposphere (cf. Stoller et al.,

1999). While the relevance of  $\Delta O_3 + \Delta RH$ - filaments is undisputed, other categories of air mass anomalies are encountered less often and are thus more difficult to relate to underlying atmospheric processes (cf. Sects. 1 and 3.1.1).

485 The selection of Q83 for RH could be revisited for the analysis of a larger time series, also in context of the  $\Delta RH$  bimodality, as the maximum peak of the  $\Delta RH$  distribution is mostly just below zero, i.e. background values (Figs. 7, 8 and A5). This is also reflected in the dominant occurrence of both  $\Delta O_3 \circ \Delta RH \circ$  and  $\Delta O_3 \circ \Delta RH$ - found for the 5–10 km subset of the Palau data (Fig. A6). An adjustment of the RH quantile for the background profile and a corresponding shift onto the center of the  $\Delta O_3 \circ \Delta RH \circ$  group could still improve the air mass selection. The monthly  $O_3$  profiles are not uniform throughout the column, which is  
490 proposed as the ideal, purely convective profile (compare Figs. 2 and A3, A4). During the months of increased occurrence of dry ozone-rich layers, i.e. FMA and NDJ, the Q20 profile still incorporates the “S”-shape, presumably caused by these layers. This effect could possibly be reduced by changing the temporal resolution from monthly to seasonal statistics, which has not been assessed yet. A growing time series will help to validate our approach as it reduces possible biases caused by different sampling frequencies per season.

## 495 **5.2 Air Mass Transport and Processes**

The trajectories reflect the dominating general circulation patterns, namely the Hadley and Walker circulation and the trade winds, with most air masses eventually reaching Palau from the East (Fig. 9). The influence of the Western Pacific Monsoon, active from July until October, becomes most apparent below 5 km altitude (cf. Müller et al., 2023). We identified two dominating routes for air mass transport to Palau, the local, Pacific route and the anticyclonic route from Asia. On the anticyclonic  
500 route, air masses can travel from the ascending branch of the Hadley circulation towards the descending branch along the subtropical ridge (Anderson et al., 2016). Once air parcels reach the West to Central Pacific during winter they are either diverted eastwards by the subtropical high anticyclone or are picked up by the general trade wind circulation, subsequently reaching Palau from the East. In contrast, with a subtropical ridge further north during summer, transport is governed only by the trade winds and monsoon circulation, bringing air masses from the convectively active Pacific region to Palau.

505 The anticyclonic flow pattern resembles the planetary wave response to tropical diabatic heating as shown for the National Centers for Environmental Prediction (NCEP) reanalysis data by Dima et al. (2005). Following the theoretical concepts of Matsuno (1966), Gill (1980) and Van Tuyl (1986), Dima et al. (2005) identified a pair of upper tropospheric anticyclonic Rossby gyres located west of and at a maximum of latent heating over the western Pacific and Indian oceans. An analysis of the equatorially symmetric component of the circulation pattern reveals the year-round presence of the Rossby wave couplet  
510 at 150 hPa, located at the same longitude on the Northern and Southern hemisphere, with a seasonal shift in longitude for the centers of the anticyclones from approx. 160° E in January/ February to approx. 80° E in July/ August. At Palau, the anticyclonic flow pattern is indeed dominating for back trajectories above 14 km altitude (not shown here). The equatorially symmetric nature of the Rossby wave couplet is potentially captured by some Palau back trajectories between 5 and 10 km during ASO reaching Palau from the Southern hemisphere (Fig. 9d). The lack of the anticyclonic route for midtropospheric  
515 Palau back trajectories for MJJ and ASO and the dominance of the Eastern pathway, seems to be supported by the seasonal

shift in longitude of the centers of the Rossby gyres in the summer months and subsequent stronger easterly flow near the equator over the western Pacific region. However, Dima et al. (2005) focused their analysis on the upper troposphere.

The results of the backward trajectory analysis support our assumptions on combined  $O_3$  and RH as tracers to identify local and non-local air masses in the 5–10 km altitude range. They further allow conclusions on air mass origin, frequency  
520 of occurrence and controlling processes. Seasonal variability in  $O_3$  VMR, RH and the vertical movement already separate between pathways to the TWP and air mass origin. The additional quantitative tracer anomaly categorization emphasizes the link between air mass origin and tracer variability directly tied to different processes.

The two quantitative air mass categories  $\Delta O_3 \circ \Delta RH \circ$  and  $\Delta O_3 + \Delta RH -$  can be related to the two governing transport patterns for Palau mid-tropospheric air masses, the clean Pacific and the polluted anticyclonic pathway (Fig. 11). The anticyclonic  
525 route for  $\Delta O_3 + \Delta RH -$  air masses occurs particularly during FMA. These air masses originate in tropical South East Asia, where biomass burning is a potent source of pollution (e.g., Anderson et al., 2016; Yadav et al., 2017; Ogino et al., 2022). They experience large-scale clear-sky subsidence associated with the Hadley circulation and dehydration during their transport within the tropical troposphere (Dessler and Minschwaner, 2007; Anderson et al., 2016). The PV values along the 10 days backward trajectories remain below 1.5 PVU (with some very rare exceptions). That means, that they do not originate in the  
530 extra-tropical stratosphere, which is characterized by PV values above 1.5 PVU. During ASO, these air masses are missing and we therefore see the undisturbed, extremely ozone-poor background (Fig. 8a). The  $\Delta O_3 \circ \Delta RH \circ$  background air masses ascend towards Palau, consistent with convective uplift. They do not leave the convectively active local Pacific region in the 10 days before arrival in Palau.

Within the debate on origin and genesis of  $O_3 + RH -$  air masses in the TWP, our analysis therefore supports the results of  
535 Anderson et al. (2016), i.e. a tropospheric origin of  $O_3 + RH -$  air. Other studies, some using the same data from the CONTRAST campaign as Anderson et al., come to the conclusion of a dominant stratospheric origin (e.g., Stoller et al., 1999; Hayashi et al., 2008; Randel et al., 2016; Tao et al., 2018). Tao et al. (2018) conducted a quantitative study of the origin of  $O_3 + RH -$  layers for the CONTRAST campaign using an artificial stratospheric tracer in the Lagrangian transport model CLaMS. They found a stratospheric influence in 60% of  $O_3 + RH -$  air masses due to in-mixing during isentropic transport and point out the limitations  
540 of calculating pure Lagrangian trajectories without chemistry. According to our analysis, the anticyclonic route identified for  $\Delta O_3 + \Delta RH -$  air masses reaching Palau is indeed a pathway along the subtropical ridge, i.e. in close proximity to mid-latitude UTLS air masses. But a lack of air masses with high PV and the clustering of the trajectory ending points near centers of pollution sources on the ground are strong evidence for a tropical tropospheric origin. This is supported by the seasonality of  $\Delta O_3 + \Delta RH -$  layers coinciding with the annual low in convective activity in FMA. We, however, cannot fully rule out a possible  
545 contribution of in-mixing of extratropical stratospheric air along the way to Palau. The additional use of aerosol observations from the co-located lidar instrument ComCAL and potentially more tracer observations during the ACCLIP campaign in late summer 2022 will contribute to the debate. Further insights are expected during the ongoing El Niño cycle with a potentially higher frequency of  $O_3 + RH -$  air mass observations.

We propose biomass burning or anthropogenic pollution as a source of  $O_3$  production in dry, ozone-rich layers at their  
550 remote origin. Their seasonal occurrence tied to the position of the Intertropical Convergence Zone indeed opens a pathway

from potential source regions which is confirmed by the trajectory analysis. If the attribution to tropical biomass burning holds, this might become a strong argument for policy makers. Anderson et al. (2016) pointed out, that due to the contribution of tropospheric O<sub>3</sub> to radiative forcing, present legislation aiming at the limitation of O<sub>3</sub> precursor emissions in the extra-tropics might not be enough to mitigate climate change.


## 555 6 Conclusions

Our study sheds light on air mass transport to the TWP as a key region of stratospheric entry and sets a valuable contribution to the discussion about anomalous layers of dry ozone-rich air observed in ozone-poor background profiles in the TWP (e.g., Anderson et al., 2016; Randel et al., 2016; Tao et al., 2018). We complemented a seasonal and statistical analysis of the balloon-borne PAO O<sub>3</sub> and RH time series (2016-19) with Lagrangian backward trajectory calculations. This approach  
560 allowed us to differentiate between air masses in the 5–10 km altitude range above Palau regarding their origin, frequency and underlying processes. We conclude, that humid, ozone-poor ( $\Delta O_3 \circ \Delta RH \circ$ ) air masses are of local or Pacific convective origin and occur year-round, but dominate from August until October. Anomalously dry ozone-rich ( $\Delta O_3 + \Delta RH -$ ) air originates in Tropical Asia and is subsequently transported to the TWP via an anticyclonic route, mostly from February to April. The origin in Tropical Asia suggests different sources of ground pollution as a cause for high O<sub>3</sub> values. No evidence for a  
565 potential stratospheric origin was found by investigating potential vorticity on the backward trajectories or by analyzing the geographical distribution of their origin. We propose that large-scale descent within the tropical troposphere and subsequent radiative cooling in connection with the Hadley circulation is responsible for the vertical displacement and dehydration. In the future, an extended time series incorporating more ENSO cycles, coinciding measurements from the ACCLIP campaign and aerosol observations by the co-located lidar system ComCAL will be applied to further validate the non-stratospheric origin of  
570 anomalous layers.


*Code and data availability.* All code used to produce the data and results is available upon request. The ozonesonde dataset is available under <https://doi.org/10.5281/zenodo.6920648> and will be included in the SHADOZ database in the future. The trajectory dataset calculated by ATLAS is available under <https://doi.org/10.5281/zenodo.8038600>. ECMWF ERA5 data used for trajectory modelling is accessible via the Copernicus Climate Change Service Climate Data Store (CDS): <https://cds.climate.copernicus.eu/>.

Figures A1, A2, A3 and A4 all give detailed insight into the seasonal or monthly variability of  $O_3$  VMR and RH measured at the PAO. The following Figs. A5, A6 and A8 refer to the anomaly categories  $\Delta O_3 \Delta RH$  as defined in Sect. 3.1.3. Figure A7 shows data from the trajectory dataset for the seasons FMA and ASO.







**Figure A1.** Variability of seasonal mean  $O_3$  VMR profiles for Palau (black solid lines) in comparison to the long-term annual mean (dashed black line) in individual panels per season (a-d), all excluding the El Niño 2016 event, i.e. starting 08/2016; the grey shaded area encloses all observations, minimum to maximum, thin colored lines indicate seasonal means per year, number (#) of profiles included in the calculations given in brackets (compare Fig. 6).




**Figure A2.** Statistical O<sub>3</sub> VMR measures for the 5–10 km altitude range highlighting the minimum O<sub>3</sub> season: monthly means from individual soundings' 25 % quantiles (red dashed line), mean (black dashed line) and minimum (black dotted line).




**Figure A3.** Monthly O<sub>3</sub> VMR statistics for Palau free tropospheric profiles as a function of altitude: 20th quantile (Q20) in blue, vertically smoothed using exponentially weighted averages in green, median in grey and central 66.6 % range in grey horizontal bars; number of included individual profiles in brackets, for orientation 20 ppb O<sub>3</sub> VMR is marked as a black vertical line (compare Figs. 4 and A4).



**Figure A4.** Monthly RH statistics for Palau free tropospheric profiles as a function of altitude: 83.3th quantile (Q83) in blue, for more details see Fig. A3; 45 % RH is highlighted by a black vertical line (compare Figs. 4 and A3).




**Figure A5.** Seasonal free-tropospheric (3-14 km) relation between tracer anomalies  $\Delta O_3$  VMR and  $\Delta RH$  for November-December-January (NDJ) (a) and May-June-July (MJJ) (b), colors according to Fig. 6, more details in Figs. 8 and 7.




**Figure A6.** Heat map for the seasonal occurrence of air mass anomaly categories  $\Delta O_3/\Delta RH$  based on the definition given in Sect. 3.1.3 and presented in Fig. 7 for Palau observations in the 5–10 km altitude range (trajectory dataset, see Sect. 2); panel (a) shows the occurrence relative to the total number (#) of seasonal profiles and panel (b) relative to the total number of all data points within the season; total numbers are given in brackets per season.

For the trajectory dataset in the 5–10 km altitude range, we observed  $\Delta O_3 \circ \Delta RH \circ$  air masses in 74 % of FMA profiles and 88 % of ASO profiles (Fig. A6).  $\Delta O_3 + \Delta RH_-$  air masses occur in 71 % of all individual FMA profiles and only 25 % of ASO profiles. In relation to the total number of seasonal data points, this contrast is even stronger: 40 % of all observed FMA data points and only 9 % of all ASO data points are identified as  $\Delta O_3 + \Delta RH_-$  air.



**Figure A7.** Relative humidity (RH) from observations above Palau versus difference in pressure altitude (between measurement date in Palau and five days before measurement) inferred from the backward trajectory analysis for the seasons FMA (a) and ASO (b); grey "+" indicate individual measurements from the trajectory dataset; colored contours show the distribution density; the black line indicates a fit by linear regression with a grey shaded 95 % confidence interval; the red dashed line highlights the zero line for the altitude difference with positive values indicating ascent and negative values descent towards Palau; the correlation coefficient R is shown in the plot; the marginal plots show univariate histograms and kernel-density estimated curves.

Figure A7 visualizes the correlation between RH and ascent/descent in a 2D histogram with a linear regression line. For FMA (Fig. A7a) the distribution reveals a cluster of very dry (<20 % RH) air masses that descended towards Palau ( $\approx 3$  km).  
 585 This cluster at the lower end of the physically possible scale is responsible for the low correlation between the two parameters, which is surprisingly low despite the clear geographical separation visible in Fig. 10c.



**Figure A8.** Seasonal distributions of the location of  $\Delta O_3 + \Delta RH$ - air masses five days before their measurement in Palau in the 5–10 km altitude range inferred from trajectories, according to the definition shown in Fig. 7; numbers in the plot from left to right: absolute number of individual air masses (trajectories) per season and as percentage per season in brackets, absolute number of profiles per season given and as percentage per season in brackets; differences to numbers shown in Fig. A6 due to rounding differences; see Fig. 6 for seasons.



## Appendix B: Statistical Averages

There are different ways to derive a climatological seasonal average and choosing the best-suited definition is not trivial with our given data set. Data from an individual sounding are first averaged in 300 m height bins using an arithmetic mean (hereafter referred to as “mean”), which essentially complies with the vertical measurement resolution, introducing only a slightly greater degree of smoothing (cf. Müller et al., 2023). In the following temporal binning, the order of required steps must be considered as they lead to different values in our time series. There are four different possible sequences for (i) monthly, (ii) seasonal and (iii) annual averaging (Müller, 2020). In our case, we chose to first directly average individual profiles  $x_i(m, y)$  of month  $m$  of all years  $y$  to a climatological month mean,  $\text{climmonmean}(m)$ , then to a climatological season mean over all years,  $\text{climseasmean}(s)$ :

$$\text{climmonmean}(m) = \frac{1}{\sum_{y=1}^Y N(m, y)} * \sum_{y=1}^Y \sum_{i=1}^{N(m, y)} x_i(m, y) \quad (\text{B1})$$

$$\text{climseasmean}(s) = \frac{1}{M(s)} * \sum_{i=1}^{M(s)} \text{climmonmean}(i), \quad (\text{B2})$$

with  $M(s)$  as the months in season  $s$ ,  $N(m, y)$  as the number of profiles in month  $m$  of year  $y$  and  $Y$  as the number of years.  $\text{climseasmean}(s)$  will not overestimate single months with very few and/or outlying profiles, assuming the interannual variability of the months is small compared to variations between different months.

*Author contributions.* KM wrote the original draft of this work and performed the analysis. PvdG and MR supported the analysis and provided effective and constructive comments to improve the manuscript. KM and others performed the measurements. All authors contributed to writing the paper.

*Competing interests.* None of the authors has any financial or personal relationships that could inappropriately influence or bias the content of the paper.

*Acknowledgements.* The setup of the PAO and this study was mostly funded by the European Union under the 7th Framework Program as part of the StratoClim project ([www.stratoclim.org](http://www.stratoclim.org)), grant number 603557. The authors thank Ingo Wohltmann (AWI) for support with the trajectory model and valuable feedback for the manuscript, Dr. Patrick Tellei, President of the Palau Community College, for provision of space, German Honorary Consul Thomas Schubert for overall support, and various people and institutions for operations at the PAO: Sharon Patris (CRRF), Pat and Lori Colin (CRRF), Gerda Ucharm (CRRF), Ingo Beninga (impres GmbH), Wilfried Ruhe (impres GmbH), Winfried Markert (Uni Bremen), Tine Weinzierl (formerly Uni Bremen), Jordis Tradowsky (NMet) and Jürgen "Egon" Graeser (AWI). The authors

want to further thank Herman Smit (FZJ), Ross Salawitch (UMD), Laura Pan (NCAR), Anne Thompson (NASA/GSFC) and many others from the international ozone research community for discussions and encouragement.

## 615 References

- Anderson, D. C., Nicely, J. M., Salawitch, R. J., Canty, T. P., Dickerson, R. R., Hanisco, T. F., Wolfe, G. M., Apel, E. C., Atlas, E., Bannan, T., Bauguutte, S., Blake, N. J., Bresch, J. F., Campos, T. L., Carpenter, L. J., Cohen, M. D., Evans, M., Fernandez, R. P., Kahn, B. H., Kinnison, D. E., Hall, S. R., Harris, N. R., Hornbrook, R. S., Lamarque, J.-F., Le Breton, M., Lee, J. D., Percival, C., Pfister, L., Pierce, R. B., Riemer, D. D., Saiz-Lopez, A., Stunder, B. J., Thompson, A. M., Ullmann, K., Vaughan, A., and Weinheimer, A. J.: A Pervasive Role for Biomass  
620 Burning in Tropical High Ozone/Low Water Structures, *Nature Communications*, 7, 10 267, <https://doi.org/10.1038/ncomms10267>, 2016.
- Bozem, H., Butler, T. M., Lawrence, M. G., Harder, H., Martinez, M., Kubistin, D., Lelieveld, J., and Fischer, H.: Chemical Processes Related to Net Ozone Tendencies in the Free Troposphere, *Atmospheric Chemistry and Physics*, 17, 10 565–10 582, <https://doi.org/10.5194/acp-17-10565-2017>, 2017.
- Browell, E. V., Fenn, M. A., Butler, C. F., Grant, W. B., Ismail, S., Ferrare, R. A., Kooi, S. A., Brackett, V. G., Clayton, M. B., Avery, M. A.,  
625 Barrick, J. D. W., Fuelberg, H. E., Maloney, J. C., Newell, R. E., Zhu, Y., Mahoney, M. J., Anderson, B. E., Blake, D. R., Brune, W. H., Heikes, B. G., Sachse, G. W., Singh, H. B., and Talbot, R. W.: Large-scale Air Mass Characteristics Observed over the Remote Tropical Pacific Ocean during March-April 1999: Results from PEM-Tropics B Field Experiment, *Journal of Geophysical Research: Atmospheres*, 106, 32 481–32 501, <https://doi.org/10.1029/2001JD900001> @ 10.1002/(ISSN)2169-8996.PEMTROP2, 2001.
- Cau, P., Methven, J., and Hoskins, B.: Origins of Dry Air in the Tropics and Subtropics, *Journal of Climate*, 20, 2745–2759,  
630 <https://doi.org/10.1175/JCLI4176.1>, 2007.
- Cecil, D. J., Buechler, D. E., and Blakeslee, R. J.: Gridded Lightning Climatology from TRMM-LIS and OTD: Dataset Description, *Atmospheric Research*, 135–136, 404–414, <https://doi.org/10.1016/j.atmosres.2012.06.028>, 2014.
- Christian, H. J.: Global Frequency and Distribution of Lightning as Observed from Space by the Optical Transient Detector, *Journal of Geophysical Research*, 108, 4005, <https://doi.org/10.1029/2002JD002347>, 2003.
- 635 Crawford, J. H., Davis, D. D., Chen, G., Bradshaw, J., Sandholm, S., Kondo, Y., Merrill, J., Liu, S., Browell, E., Gregory, G., Anderson, B., Sachse, G., Barrick, J., Blake, D., Talbot, R., and Pueschel, R.: Implications of Large Scale Shifts in Tropospheric NO<sub>x</sub> Levels in the Remote Tropical Pacific, *Journal of Geophysical Research: Atmospheres*, 102, 28 447–28 468, <https://doi.org/10.1029/97JD00011>, 1997.
- Dessler, A. E. and Minschwaner, K.: An Analysis of the Regulation of Tropical Tropospheric Water Vapor, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD007683>, 2007.
- 640 Diallo, M., Riese, M., Birner, T., Konopka, P., Müller, R., Hegglin, M. I., Santee, M. L., Baldwin, M., Legras, B., and Ploeger, F.: Response of Stratospheric Water Vapor and Ozone to the Unusual Timing of El Niño and the QBO Disruption in 2015–2016, *Atmospheric Chemistry and Physics*, 18, 13 055–13 073, <https://doi.org/10.5194/acp-18-13055-2018>, 2018.
- Dima, I. M., Wallace, J. M., and Kraucunas, I.: Tropical Zonal Momentum Balance in the NCEP Reanalyses, *Journal of the Atmospheric Sciences*, 62, 2499 – 2513, <https://doi.org/10.1175/JAS3486.1>, 2005.
- 645 Folkens, I.: Tropical Ozone as an Indicator of Deep Convection, *Journal of Geophysical Research*, 107, 4184, <https://doi.org/10.1029/2001JD001178>, 2002.
- Folkens, I. and Martin, R. V.: The Vertical Structure of Tropical Convection and Its Impact on the Budgets of Water Vapor and Ozone, *Journal of the Atmospheric Sciences*, 62, 1560–1573, <https://doi.org/10.1175/JAS3407.1>, 2005.
- Fueglistaler, S., Wernli, H., and Peter, T.: Tropical Troposphere-to-Stratosphere Transport Inferred from Trajectory Calculations, *Journal of*  
650 *Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004069>, 2004.

- Gottelman, A. and Forster, P. d. F.: A Climatology of the Tropical Tropopause Layer., *Journal of the Meteorological Society of Japan*, 80, 911–924, <https://doi.org/10.2151/jmsj.80.911>, 2002.
- Gottelman, A., Lauritzen, P. H., Park, M., and Kay, J. E.: Processes Regulating Short-Lived Species in the Tropical Tropopause Layer, *Journal of Geophysical Research*, 114, D13 303, <https://doi.org/10.1029/2009JD011785>, 2009.
- 655 Gill, A. E.: Some simple solutions for heat-induced tropical circulation, *Quarterly Journal of the Royal Meteorological Society*, 106, 447–462, 1980.
- Harris, N. R. P., Carpenter, L. J., Lee, J. D., Vaughan, G., Filus, M. T., Jones, R. L., OuYang, B., Pyle, J. A., Robinson, A. D., Andrews, S. J., Lewis, A. C., Minaeian, J., Vaughan, A., Dorsey, J. R., Gallagher, M. W., Le Breton, M., Newton, R., Percival, C. J., Ricketts, H. M. A., Bauguitte, S. J.-B., Nott, G. J., Wellpott, A., Ashfold, M. J., Flemming, J., Butler, R., Palmer, P. I., Kaye, P. H., Stopford, C.,
- 660 Chemel, C., Boesch, H., Humpage, N., Vick, A., MacKenzie, A. R., Hyde, R., Angelov, P., Meneguz, E., and Manning, A. J.: Coordinated Airborne Studies in the Tropics (CAST), *Bulletin of the American Meteorological Society*, 98, 145–162, <https://doi.org/10.1175/BAMS-D-14-00290.1>, 2017.
- Hayashi, H., Kita, K., and Taguchi, S.: Ozone-Enhanced Layers in the Troposphere over the Equatorial Pacific Ocean and the Influence of Transport of Midlatitude UT/LS Air, *Atmospheric Chemistry and Physics*, p. 14, 2008.
- 665 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 Global Reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 670 Huang, B., L'Heureux, M., Hu, Z.-Z., and Zhang, H.-M.: Ranking the Strongest ENSO Events While Incorporating SST Uncertainty, *Geophysical Research Letters*, 43, 9165–9172, 2016.
- Kley, D., Crutzen, P. J., Smit, H. G. J., Vomel, H., Oltmans, S. J., Grassl, H., and Ramanathan, V.: Observations of Near-Zero Ozone Concentrations Over the Convective Pacific: Effects on Air Chemistry, *Science*, 274, 230–233, <https://doi.org/10.1126/science.274.5285.230>, 1996.
- 675 Kley, D., Smit, H. G. J., Vömel, H., Grassl, H., Ramanathan, V., Crutzen, P. J., Williams, S., Meywerk, J., and Oltmans, S. J.: Tropospheric Water-Vapour and Ozone Cross-Sections in a Zonal Plane over the Central Equatorial Pacific Ocean, *Quarterly Journal of the Royal Meteorological Society*, 123, 2009–2040, <https://doi.org/10.1002/qj.49712354312>, 1997.
- Komhyr, W. D.: Electrical Concentration Cells for Gas Analysis, *Annales Geophysicae*, 25, 203–210, 1969.
- Krüger, K. and Quack, B.: Introduction to Special Issue: The TransBrom Sonne Expedition in the Tropical West Pacific, *Atmospheric Chemistry and Physics*, 13, 9439–9446, <https://doi.org/10.5194/acp-13-9439-2013>, 2013.
- 680 Krüger, K., Tegtmeier, S., and Rex, M.: Long-Term Climatology of Air Mass Transport through the Tropical Tropopause Layer (TTL) during NH Winter, *Atmospheric Chemistry and Physics*, 8, 813–823, <https://doi.org/10.5194/acp-8-813-2008>, 2008.
- Kunz, A., Konopka, P., Müller, R., and Pan, L. L.: Dynamical Tropopause Based on Isentropic Potential Vorticity Gradients, *Journal of Geophysical Research*, 116, D01 110, <https://doi.org/10.1029/2010JD014343>, 2011.
- 685 Levy, H.: Normal Atmosphere: Large Radical and Formaldehyde Concentrations Predicted, *Science*, 173, 141–143, <https://doi.org/10.1126/science.173.3992.141>, 1971.
- Liu, S. C., McFarland, M., Kley, D., Zafiriou, O., and Huebert, B.: Tropospheric NO<sub>x</sub> and O<sub>3</sub> Budgets in the Equatorial Pacific, *Journal of Geophysical Research*, 88, 1360, <https://doi.org/10.1029/JC088iC02p01360>, 1983.

- 690 Mapes, B. E.: Water's Two Height Scales: The Moist Adiabatic and the Radiative Troposphere, *Quarterly Journal of the Royal Meteorological Society*, 127, 2353–2366, <https://doi.org/10.1002/qj.49712757708>, 2001.
- Mapes, B. E. and Zuidema, P.: Radiative-Dynamical Consequences of Dry Tongues in the Tropical Troposphere, *Journal of the Atmospheric Sciences*, 53, 620–638, [https://doi.org/10.1175/1520-0469\(1996\)053<0620:RDCODT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1996)053<0620:RDCODT>2.0.CO;2), 1996.
- Matsuno, T.: Quasi-geostrophic motions in the equatorial area, *Journal of the Meteorological Society of Japan. Ser. II*, 44, 25–43, 1966.
- 695 Müller, K.: Characterization of Ozone and the Oxidizing Capacity of the Tropical West Pacific Troposphere, Ph.D. thesis, Universität Bremen, <https://doi.org/10.26092/ELIB/463>, 2020.
- Müller, K., Tradowsky, J. S., von der Gathen, P., Ritter, C., Patris, S., Notholt, J., and Rex, M.: Measurement Report: The Palau Atmospheric Observatory and Its Ozonesonde Record - Continuous Monitoring of Tropospheric Composition and Dynamics in the Tropical West Pacific, *EGUsphere*, 2023, 1–35, <https://doi.org/10.5194/egusphere-2023-1023>, 2023.
- Newell, R. E. and Gould-Stewart, S.: A Stratospheric Fountain?, *Journal of the Atmospheric Sciences*, 38, 2789–2796, 1981.
- 700 Newell, R. E., Zhu, Y., Browell, E. V., Ismail, S., Read, W. G., Waters, J. W., Kelly, K. K., and Liu, S. C.: Upper Tropospheric Water Vapor and Cirrus: Comparison of DC-8 Observations, Preliminary UARS Microwave Limb Sounder Measurements and Meteorological Analyses, *Journal of Geophysical Research: Atmospheres*, 101, 1931–1941, 1996.
- Newell, R. E., Thouret, V., Cho, J. Y. N., Stoller, P., Marengo, A., and Smit, H. G.: Ubiquity of Quasi-Horizontal Layers in the Troposphere, *Nature*, 398, 316–319, <https://doi.org/10.1038/18642>, 1999.
- 705 Newton, R., Vaughan, G., Ricketts, H. M. A., Pan, L. L., Weinheimer, A. J., and Chemel, C.: Ozonesonde Profiles from the West Pacific Warm Pool: Measurements and Validation, *Atmospheric Chemistry and Physics*, 16, 619–634, <https://doi.org/10.5194/acp-16-619-2016>, 2016.
- Nicely, J. M., Anderson, D. C., Canty, T. P., Salawitch, R. J., Wolfe, G. M., Apel, E. C., Arnold, S. R., Atlas, E. L., Blake, N. J., Bresch, J. F., Campos, T. L., Dickerson, R. R., Duncan, B., Emmons, L. K., Evans, M. J., Fernandez, R. P., Flemming, J., Hall, S. R., Hanisco, T. F., Honomichl, S. B., Hornbrook, R. S., Huijnen, V., Kaser, L., Kinnison, D. E., Lamarque, J.-F., Mao, J., Monks, S. A., Montzka, D. D., Pan, L. L., Riemer, D. D., Saiz-Lopez, A., Steenrod, S. D., Stell, M. H., Tilmes, S., Turquety, S., Ullmann, K., and Weinheimer, A. J.: An Observationally Constrained Evaluation of the Oxidative Capacity in the Tropical Western Pacific Troposphere: Observationally Constrained OH in TWP, *Journal of Geophysical Research: Atmospheres*, 121, 7461–7488, <https://doi.org/10.1002/2016JD025067>, 2016.
- 710 Ogino, S.-Y., Fujiwara, M., Shiotani, M., Hasebe, F., Matsumoto, J., T. Hoang, T. H., and T. Nguyen, T. T.: Ozone Variations over the Northern Subtropical Region Revealed by Ozonesonde Observations in Hanoi, *Journal of Geophysical Research: Atmospheres*, 118, 3245–3257, <https://doi.org/10.1002/jgrd.50348>, 2013.
- Ogino, S.-Y., Miyazaki, K., Fujiwara, M., Nodzu, M. I., Shiotani, M., Hasebe, F., Matsumoto, J., Witte, J., Thompson, A. M., Nguyen-Thi, H. A., and Nguyen, T. V.: Cause of a Lower-Tropospheric High-Ozone Layer in Spring Over Hanoi, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035727, <https://doi.org/https://doi.org/10.1029/2021JD035727>, e2021JD035727 2021JD035727, 2022.
- 720 Oltmans, S. J., Johnson, B. J., Harris, J. M., Vömel, H., Thompson, A. M., Koshy, K., Simon, P., Bendura, R. J., Logan, J. A., Hasebe, F., Shiotani, M., Kirchhoff, V. W. J. H., Maata, M., Sami, G., Samad, A., Tabuadravu, J., Enriquez, H., Agama, M., Cornejo, J., and Paredes, F.: Ozone in the Pacific Tropical Troposphere from Ozonesonde Observations, *Journal of Geophysical Research: Atmospheres*, 106, 32503–32525, [https://doi.org/10.1029/2000JD900834@10.1002/\(ISSN\)2169-8996.PEMTROP2](https://doi.org/10.1029/2000JD900834@10.1002/(ISSN)2169-8996.PEMTROP2), 2001.
- 725 Pan, L. L., Paulik, L. C., Honomichl, S. B., Munchak, L. A., Bian, J., Selkirk, H. B., and Vömel, H.: Identification of the Tropical Tropopause Transition Layer Using the Ozone-Water Vapor Relationship, *Journal of Geophysical Research: Atmospheres*, 119, 3586–3599, <https://doi.org/10.1002/2013JD020558>, 2014.

- Pan, L. L., Honomichl, S. B., Randel, W. J., Apel, E. C., Atlas, E. L., Beaton, S. P., Bresch, J. F., Hornbrook, R., Kinnison, D. E., Lamarque, J.-F., Saiz-Lopez, A., Salawitch, R. J., and Weinheimer, A. J.: Bimodal Distribution of Free Tropospheric Ozone over the Tropical Western Pacific Revealed by Airborne Observations, *Geophysical Research Letters*, 42, 7844–7851, <https://doi.org/10.1002/2015GL065562>, 2015.
- 730 Pan, L. L., Atlas, E. L., Salawitch, R. J., Honomichl, S. B., Bresch, J. F., Randel, W. J., Apel, E. C., Hornbrook, R. S., Weinheimer, A. J., Anderson, D. C., Andrews, S. J., Baidar, S., Beaton, S. P., Campos, T. L., Carpenter, L. J., Chen, D., Dix, B., Donets, V., Hall, S. R., Hanisco, T. F., Homeyer, C. R., Huey, L. G., Jensen, J. B., Kaser, L., Kinnison, D. E., Koenig, T. K., Lamarque, J.-F., Liu, C., Luo, J., Luo, Z. J., Montzka, D. D., Nicely, J. M., Pierce, R. B., Riemer, D. D., Robinson, T., Romashkin, P., Saiz-Lopez, A., Schauffler, S., Shieh, O., Stell, M. H., Ullmann, K., Vaughan, G., Volkamer, R., and Wolfe, G.: The Convective Transport of Active Species in the Tropics (CONTRAST) Experiment, *Bulletin of the American Meteorological Society*, 98, 106–128, <https://doi.org/10.1175/BAMS-D-14-00272.1>, 2017.
- 735 Paulik, L. C. and Birner, T.: Quantifying the Deep Convective Temperature Signal within the Tropical Tropopause Layer (TTL), *Atmospheric Chemistry and Physics*, 12, 12 183–12 195, <https://doi.org/10.5194/acp-12-12183-2012>, 2012.
- Ploeger, F., Fueglistaler, S., Grooß, J.-U., Günther, G., Konopka, P., Liu, Y., Müller, R., Ravegnani, F., Schiller, C., Ulanovski, A., and Riese, M.: Insight from Ozone and Water Vapour on Transport in the Tropical Tropopause Layer (TTL), *Atmospheric Chemistry and Physics*, 11, 407–419, <https://doi.org/10.5194/acp-11-407-2011>, 2011.
- 740 Randel, W. J., Park, M., Wu, F., and Livesey, N.: A Large Annual Cycle in Ozone above the Tropical Tropopause Linked to the Brewer–Dobson Circulation, *Journal of the Atmospheric Sciences*, 64, 4479–4488, <https://doi.org/10.1175/2007JAS2409.1>, 2007.
- Randel, W. J., Rivoire, L., Pan, L. L., and Honomichl, S. B.: Dry Layers in the Tropical Troposphere Observed during CONTRAST and Global Behavior from GFS Analyses: Dry Layers in the Tropical Troposphere, *Journal of Geophysical Research: Atmospheres*, 121, 14,142–14,158, <https://doi.org/10.1002/2016JD025841>, 2016.
- 745 Rex, M., Wohltmann, I., Ridder, T., Lehmann, R., Rosenlof, K., Wennberg, P., Weisenstein, D., Notholt, J., Krüger, K., Mohr, V., and Tegtmeier, S.: A Tropical West Pacific OH Minimum and Implications for Stratospheric Composition, *Atmospheric Chemistry and Physics*, 14, 4827–4841, <https://doi.org/10.5194/acp-14-4827-2014>, 2014.
- 750 Ruzmaikin, A., Aumann, H. H., and Manning, E. M.: Relative Humidity in the Troposphere with AIRS, *Journal of the Atmospheric Sciences*, 71, 2516 – 2533, <https://doi.org/10.1175/JAS-D-13-0363.1>, 2014.
- Schoeberl, M. R., Selkirk, H. B., Vömel, H., and Douglass, A. R.: Sources of Seasonal Variability in Tropical Upper Troposphere and Lower Stratosphere Water Vapor and Ozone: Inferences from the Ticosonde Data Set at Costa Rica, *Journal of Geophysical Research: Atmospheres*, 120, 9684–9701, <https://doi.org/10.1002/2015JD023299>, 2015.
- 755 Sherwood, S. C., Kursinski, E. R., and Read, W. G.: A Distribution Law for Free-Tropospheric Relative Humidity, *Journal of Climate*, 19, 6267 – 6277, <https://doi.org/10.1175/JCLI3978.1>, 2006.
- Sherwood, S. C., Roca, R., Weckwerth, T. M., and Andronova, N. G.: Tropospheric Water Vapor, Convection, and Climate, *Reviews of Geophysics*, 48, 1, <https://doi.org/10.1029/2009RG000301>, 2010.
- Smit, H. and Thompson, A.: Ozone-sonde Measurement Principles and Best Operational Practices: ASOPOS 2.0 (Assessment of Standard Operating Procedures for Ozone-sondes), WMO, World Meteorological Organization, GAW Report, p. 173, 2021.
- 760 Smit, H. G. J.: Quality Assurance and Quality Control for Ozone-sonde Measurements in GAW, GAW Report No. 201, WMO, 2014.
- Solomon, S., Thompson, D. W. J., Portmann, R. W., Oltmans, S. J., and Thompson, A. M.: On the Distribution and Variability of Ozone in the Tropical Upper Troposphere: Implications for Tropical Deep Convection and Chemical-Dynamical Coupling, *Geophysical Research Letters*, 32, L23 813, <https://doi.org/10.1029/2005GL024323>, 2005.

- 765 Stauffer, R. M., Thompson, A. M., and Witte, J. C.: Characterizing Global Ozonesonde Profile Variability From Surface to the UT/LS With a Clustering Technique and MERRA-2 Reanalysis, *Journal of Geophysical Research: Atmospheres*, 123, 6213–6229, <https://doi.org/10.1029/2018JD028465>, 2018.
- Stoller, P., Cho, J. Y. N., Newell, R. E., Thouret, V., Zhu, Y., Carroll, M. A., Albercook, G. M., Anderson, B. E., Barrick, J. D. W., Browell, E. V., Gregory, G. L., Sachse, G. W., Vay, S., Bradshaw, J. D., and Sandholm, S.: Measurements of Atmospheric Layers  
770 from the NASA DC-8 and P-3B Aircraft during PEM-Tropics A, *Journal of Geophysical Research: Atmospheres*, 104, 5745–5764, <https://doi.org/10.1029/98JD02717>, 1999.
- Sun, X., Palm, M., Müller, K., Hachmeister, J., and Notholt, J.: Determination of the Chemical Equator from GEOS-Chem Model Simulation: A Focus on the Tropical Western Pacific Region, *Atmospheric Chemistry and Physics*, 23, 7075–7090, <https://doi.org/10.5194/acp-23-7075-2023>, 2023.
- 775 Takashima, H., Shiotani, M., Fujiwara, M., Nishi, N., and Hasebe, F.: Ozonesonde Observations at Christmas Island (2°N, 157°W) in the Equatorial Central Pacific, *Journal of Geophysical Research*, 113, D10 112, <https://doi.org/10.1029/2007JD009374>, 2008.
- Tao, M., Pan, L. L., Konopka, P., Honomichl, S. B., Kinnison, D. E., and Apel, E. C.: A Lagrangian Model Diagnosis of Stratospheric Contributions to Tropical Midtropospheric Air, *Journal of Geophysical Research: Atmospheres*, 123, 9764–9785, <https://doi.org/10.1029/2018JD028696>, 2018.
- 780 Thompson, A. M., Tao, W.-K., Pickering, K. E., Scala, J. R., and Simpson, J.: Tropical Deep Convection and Ozone Formation, *Bulletin of the American Meteorological Society*, 78, 1043–1054, [https://doi.org/10.1175/1520-0477\(1997\)078<1043:TDCAOF>2.0.CO;2](https://doi.org/10.1175/1520-0477(1997)078<1043:TDCAOF>2.0.CO;2), 1997.
- Thompson, A. M., Witte, J. C., McPeters, R. D., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V. W. J. H., Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami, S., Toshihiro, O., Johnson, B. J., Vömel, H., and Labow, G. J.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 Tropical Ozone Climatology 1. Comparison with Total Ozone Mapping Spectrometer (TOMS) and  
785 Ground-Based Measurements, *Journal of Geophysical Research*, 108, 8238, <https://doi.org/10.1029/2001JD000967>, 2003a.
- Thompson, A. M., Witte, J. C., Oltmans, S. J., Schmidlin, F. J., Logan, J. A., Fujiwara, M., Kirchhoff, V. W. J. H., Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami, S., Ogawa, T., Fortuin, J. P. F., and Kelder, H. M.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–2000 Tropical Ozone Climatology 2. Tropospheric Variability and the Zonal Wave-One, *Journal of Geophysical Research*, 108, 8241, <https://doi.org/10.1029/2002JD002241>, 2003b.
- 790 Thompson, A. M., Miller, S. K., Tilmes, S., Kollonige, D. W., Witte, J. C., Oltmans, S. J., Johnson, B. J., Fujiwara, M., Schmidlin, F. J., Coetzee, G. J. R., Komala, N., Maata, M., bt Mohamad, M., Nguyo, J., Mutai, C., Ogino, S.-Y., Da Silva, F. R., Leme, N. M. P., Posny, F., Scheele, R., Selkirk, H. B., Shiotani, M., Stübi, R., Levrat, G., Calpini, B., Thouret, V., Tsuruta, H., Canossa, J. V., Vömel, H., Yonemura, S., Diaz, J. A., Tan Thanh, N. T., and Thuy Ha, H. T.: Southern Hemisphere Additional Ozonesondes (SHADOZ) Ozone Climatology (2005-2009): Tropospheric and Tropical Tropopause Layer (TTL) Profiles with Comparisons to OMI-based Ozone Products: SHADOZ  
795 OZONE DURING AURA, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2011JD016911>, 2012.
- Thompson, A. M., Witte, J. C., Sterling, C., Jordan, A., Johnson, B. J., Oltmans, S. J., Fujiwara, M., Vömel, H., Allaart, M., Piders, A., Coetzee, G. J. R., Posny, F., Corrales, E., Diaz, J. A., Félix, C., Komala, N., Lai, N., Ahn Nguyen, H. T., Maata, M., Mani, F., Zainal, Z., Ogino, S.-y., Paredes, F., Penha, T. L. B., Silva, F. R., Sallons-Mitro, S., Selkirk, H. B., Schmidlin, F. J., Stübi, R., and Thiongo, K.: First Reprocessing of Southern Hemisphere Additional Ozonesondes (SHADOZ) Ozone Profiles (1998–2016): 2. Comparisons With Satellites  
800 and Ground-Based Instruments, *Journal of Geophysical Research: Atmospheres*, 122, <https://doi.org/10.1002/2017JD027406>, 2017.
- Thompson, A. M., Smit, H. G. J., Witte, J. C., Stauffer, R. M., Johnson, B. J., Morris, G., von der Gathen, P., Van Malderen, R., Davies, J., Piders, A., Allaart, M., Posny, F., Kivi, R., Cullis, P., Hoang Anh, N. T., Corrales, E., Machinini, T., da Silva, F. R., Paiman, G., Thiong’o,

- K., Zainal, Z., Brothers, G. B., Wolff, K. R., Nakano, T., Stübi, R., Romanens, G., Coetzee, G. J. R., Diaz, J. A., Mitro, S., Mohamad, M., and Ogino, S.-Y.: Ozonesonde Quality Assurance: The JOSIE–SHADOZ (2017) Experience, *Bulletin of the American Meteorological Society*, 100, 155–171, <https://doi.org/10.1175/BAMS-D-17-0311.1>, 2019.
- 805 Thompson, A. M., Stauffer, R. M., Wargan, K., Witte, J. C., Kollonige, D. E., and Ziemke, J. R.: Regional and Seasonal Trends in Tropical Ozone From SHADOZ Profiles: Reference for Models and Satellite Products, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2021JD034691>, 2021.
- Thouret, V., Cho, J. Y. N., Newell, R. E., Marengo, A., and Smit, H. G. J.: General Characteristics of Tropospheric Trace  
810 Constituent Layers Observed in the MOZAIC Program, *Journal of Geophysical Research: Atmospheres*, 105, 17 379–17 392, <https://doi.org/10.1029/2000JD900238>, 2000.
- Van Tuyl, A. H.: Advective influences on forced tropical motions, *Journal of the atmospheric sciences*, 43, 141–161, 1986.
- Waugh, D. W. and Polvani, L. M.: Climatology of Intrusions into the Tropical Upper Troposphere, *Geophysical Research Letters*, 27, 3857–3860, <https://doi.org/10.1029/2000GL012250>, 2000.
- 815 Wohltmann, I. and Rex, M.: The Lagrangian Chemistry and Transport Model ATLAS: Validation of Advective Transport and Mixing, *Geoscientific Model Development*, 2, 153–173, <https://doi.org/10.5194/gmd-2-153-2009>, 2009.
- Wohltmann, I., Lehmann, R., and Rex, M.: The Lagrangian Chemistry and Transport Model ATLAS: Simulation and Validation of Stratospheric Chemistry and Ozone Loss in the Winter 1999/2000, *Geoscientific Model Development*, 3, 585–601, <https://doi.org/10.5194/gmd-3-585-2010>, 2010.
- 820 Yadav, I. C., Linthoingambi Devi, N., Li, J., Syed, J. H., Zhang, G., and Watanabe, H.: Biomass burning in Indo-China peninsula and its impacts on regional air quality and global climate change - a review, *Environmental Pollution*, 227, 414–427, <https://doi.org/https://doi.org/10.1016/j.envpol.2017.04.085>, 2017.
- Yoneyama, K. and Parsons, D. B.: A Proposed Mechanism for the Intrusion of Dry Air into the Tropical Western Pacific Region, *Journal of the Atmospheric Sciences*, 56, 1524–1546, [https://doi.org/10.1175/1520-0469\(1999\)056<1524:APMFTI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<1524:APMFTI>2.0.CO;2), 1999.
- 825 Zhang, C., Mapes, B. E., and Soden, B. J.: Bimodality in tropical water vapour, *Quarterly Journal of the Royal Meteorological Society*, 129, 2847–2866, <https://doi.org/https://doi.org/10.1256/qj.02.166>, 2003.