



Observational relationships between ammonia, carbon dioxide and water vapor under a wide range of meteorological and turbulent conditions: RITA-2021 campaign

Ruben B. Schulte¹, Jordi Vilà-Guerau de Arellano¹, Shelley van der Graaf², Susanna Rutledge-Jonker², Jun Zhang³, and Margreet C. van Zanten^{1,2}

¹Meteorology and Air Quality group, Wageningen University & Research, P.O. Box 47, 6700, AA Wageningen, the Netherlands

²National Institute for Public Health and the Environment (RIVM), Antonie van Leeuwenhoeklaan 9, 3721, MA Bilthoven, the Netherlands

³TNO, Postbus 15, 1755 ZG, Petten, the Netherlands

Correspondence: Margreet van Zanten (margreet.van.zanten@rivm.nl)

Abstract. We present a comprehensive observational approach, aiming to establish relations between the surface-atmosphere exchange of ammonia (NH_3) and the CO_2 uptake and transpiration by vegetation. In doing so, we study relationships useful for the the improvement and development of NH_3 flux representations and their dependences. The NH_3 concentration and flux are measured using a novel open-path miniDOAS measurement setup, taken during the five week RITA-2021 campaign (25 August until 12 October 2021) at the Ruisdael Observatory at Cabauw, the Netherlands. After filtering for unobstructed flow, sufficient turbulent mixing and CO_2 uptake, we find the diurnal variability of the NH_3 flux to be characterized by daytime emissions ($0.05 \mu\text{g m}^{-2}\text{s}^{-1}$ on average) and deposition at sunrise and sunset ($-0.05 \mu\text{g m}^{-2}\text{s}^{-1}$ on average). We first compare the NH_3 flux to the observed gross primary production (GPP), representing CO_2 uptake, and latent heat flux (L_vE), representing to evapotranspiration. Next we study the observations following the main drivers of the dynamic vegetation response, which are photosynthetically active radiation (PAR), temperature (T) and the water vapor pressure deficit (VPD). Our findings show indication of the dominance of stomatal emission of NH_3 , with high correlation between the observed emissions and both net L_vE (0.70) and PAR (0.72), as well as close similarities in the diurnal variability of the NH_3 flux and GPP. However, the efforts to establish relationships are hampered due to the amount of diversity of NH_3 sources of the active agricultural region and low data availability after filtering. Our findings show the need to collocate meteorological, carbon and nitrogen studies to advance on our understanding of NH_3 deposition and its representation.

1 Introduction

While nitrogen is an essential nutrient for the growth of plants, acting as a fertilizer, excess nitrogen deposition causes environmental damage and leads to an increased public health risk through the formation of particulate matter (Bobbink et al., 2003; Behera et al., 2013; Erisman and Schaap, 2004; Erisman et al., 2013; Smit and Heederik, 2017). When nitrogen critical loads are exceeded, excess nitrogen deposition threatens biodiversity through acidification and eutrophication of soils. When mitigation of the harmful effects of nitrogen fails, there can be serious political, economic and societal consequences, as demonstrated by the current Dutch nitrogen crisis (Stokstad, 2019). Atmospheric ammonia (NH_3) plays a key role in the deposition of nitrogen, mainly originating from agricultural activity. This is especially true in the Netherlands where NH_3 deposition accounts for about three-quarters of all nitrogen deposition (Wichink Kruit and van Pul, 2018; RIVM et al., 2019).



Efforts to mitigate the harmful effects of nitrogen deposition heavily rely on models representing the concentration and deposition of nitrogen compounds, supported by a network of concentration and surface-atmosphere exchange measurements. The surface-atmosphere exchange in such models is represented by parameterizations, which are developed, validated and improved based on advanced high-resolution observations. In the case of atmospheric ammonia, taking accurate high-resolution measurements is notoriously difficult, due to the reactive nature of gaseous NH₃ causing the gas to "stick" to inlet walls of conventional instruments (Parrish and Fehsenfeld, 2000; von Bobruzki et al., 2010). These challenges are amplified when measuring the NH₃ surface-atmosphere exchange flux (deposition or emission), where high precision is particularly important (Nemitz et al., 2004; Whitehead et al., 2008).

Recent developments in advanced instrumental techniques resolve these inlet issues by using optical open-path analyzers. Swart et al. (2023) presents an intercomparison of two novel open-path measurement setups, aimed at measuring the NH₃ flux at half-hourly resolution: the RIVM-miniDOAS 2.2D and the commercial Healthy Photon HT8700E. The two setups showed very similar results, despite being widely different in their measurement principle and approach to derive the flux from concentrations, as the Healthy Photon uses the eddy covariance technique while the miniDOAS applies the flux-gradient method to line average concentration measurements over a 22 m open-path at two heights. In this study, we continue the analysis of the observations of the miniDOAS system presented by (Swart et al., 2023), as the system provides reliable measurements of both the concentration and flux with a high operational uptime.

In a previous study, based on measurements from the predecessor of the miniDOAS system at the Veenkampen meteorological site in the Netherlands, we identified that the mechanisms behind stomatal exchange of NH₃ are not yet fully understood (Schulte et al., 2021). Here, we continue to study this stomatal exchange pathway by linking the observed NH₃ flux (F_{NH_3}) to photosynthesis, i.e. the stomatal exchange of CO₂ and water vapor (plant transpiration). The similarities between the stomatal exchange of NH₃ and CO₂ have long been recognized (San José et al., 1991; Schrader et al., 2020). However, there are very few parallel measurements of NH₃ and CO₂ fluxes, and research into the two gases is generally conducted by separate scientific communities (Milford et al., 2001). Milford et al. (2001) performed one of the few attempts to develop a simple parameterization for both the CO₂ and NH₃ flux, but was unsuccessful to find such relations for NH₃ as the observed NH₃ flux over Scottish Heathland was dominated by non-stomatal exchange. Further, Zöll et al. (2019) performed an analysis to study whether the biosphere-atmosphere exchange of total reactive nitrogen was driven by the same variables as carbon dioxide.

Our aim is to relate NH₃ and CO₂ fluxes to advance in our understanding of NH₃ stomatal exchange. These surface exchange require to be related to the sensible and latent heat fluxes, and the diurnal boundary layer dynamics (Vilà-Guerau de Arellano et al., 2023). Utilizing the recent developments in NH₃ measurement techniques, we combine the high-quality miniDOAS F_{NH_3} observations with measurements of both CO₂ and water vapor fluxes, as well as other meteorological variables. We first describe the observations, after which we link the observed F_{NH_3} to stomatal exchange, with the intention to establish relationships between the stomatal exchange of ammonia and the processes of CO₂ uptake and transpiration by vegetation. As these processes of photosynthesis are well understood, we explore how this understanding can lead to further improving the parameterization of the NH₃ stomatal exchange.

2 Characterizing the RITA-2021 campaign observations

2.1 Site description and measurement strategy

In September 2021, the Ruisdael Land-Atmosphere Interactions Intensive Trace-gas and Aerosol measurement campaign, known as RITA-2021, took place at the Cabauw Observatory (<https://ruisdael-observatory.nl/cabauw/>). The Cabauw Observatory, one of the 6 sites within the Ruisdael Observatory, is located on flat grassland in the Netherlands (51.971°N, 4.927°E), with an average grass height of 0.1 m. The site provides a unique set of surface and upper air observations, matched by only a very few station world-wide. This includes measurements of thermodynamic variables along the 213 m mast, radiation, surface fluxes, clouds and trace gasses. Surface elevation changes are at most a few meters over 20 km and the nearby region is agricultural. An overview of the Cabauw site, the instruments stationed at the site and its 50-years of observations is given in Bosveld et al. (2020).

During the campaign, 48 days of ammonia measurements are taken using the miniDOAS (Differential Optical Absorption Spectroscopy) flux measurement setup (Berkhout et al., 2017), starting on 25 August until 12 October. The measurement setup and more details on the measurement campaign are described in Swart et al. (2023). In short, the miniDOAS is an optical instrument, measuring the line average concentration (mass density) over a 22 m open path from instrument to its retroreflector. The 30 minute average NH₃ concentrations have an accuracy of 0.01 $\mu\text{g m}^{-3}$. The flux measurement setup uses two miniDOAS instruments, which measure the concentration over parallel paths at different heights, i.e. 0.76 m and 2.29 m respectively. F_{NH_3} is then inferred using the flux-gradient method, based on the Monin–Obukhov similarity theory (Moene and Van Dam, 2014).



Ruben B.Schulte: Observational relationships between NH₃, CO₂ and evapotranspiration

3




Figure 1. The area surrounding the Cabauw Observatory (left) and the setup of the instruments at the measurement site (right). The transparent white circle represents a distance of 500 m from the NH₃ measurements and the color coded dots represent the locations of nearby farms, i.e. emission sources of NH₃ (Source: Emissieregistratie, www.emissieregistratie.nl, last access 21 January 2022). The colored circle in both panels indicates the wind directions where the airflow towards the miniDOAS light-path is obstructed by either other instruments (yellow) or larger structures (red). (Source left panel: "Cabauw Observatory", 51.971°N, 4.927°E. © Google Earth, 27 January 2022, Image by Landsat / Copernicus. Source right panel: Swart et al. (2023), modified)

The flux-gradient method combines the observed vertical NH₃ gradient with turbulent measurements of a sonic anemometer (model Gill WindMasterPro™ Gill Instruments, Lymington, UK) (Wyers et al., 1993; Nemitz et al., 2004; Wichink Kruit et al., 2007; Schulte et al., 2021). The sonic anemometer was mounted at 2.8 m above the ground alongside the mini-DOAS measurement path. The 10 Hz open-path H₂O and CO₂ analyser (LI-7500DS, LI-COR Biosciences, Lincoln, USA) was placed at a similar height, 15 cm away from the sonic (for more details see information over sonic #1 in (Swart et al., 2023)) The CO₂ and water vapor fluxes and other micrometeorological parameters were calculated using EddyPro software (LI-COR Biosciences) at 30-min intervals using the 10 Hz raw data. The general flux calculation procedure followed the standard FluxNet methodology (McDermitt et al., 2011).

The measurement field and its surroundings are shown in Fig. 1. The miniDOAS light paths are aimed in north-northwestern direction (right panel) to ensure unobstructed flow for wind coming from the west, which is the dominant wind direction in the Netherlands. North of the light path, shown in yellow in Fig 1, the flow of air is obstructed by several instruments, including the aforementioned sonic anemometer. To the east and south, the airflow is obstructed by a trailer, the 213 m high meteorological tower and the container which houses the miniDOAS instruments. The unobstructed region west of the measurement field, is mainly characterized by actively managed agricultural grassland and the small town of Cabauw (about 750 inhabitants), as shown left in Fig. 1. Several farms can be seen northwest and west of the measurement field, with varying emission strengths up to over 1200 kg NH₃ year⁻¹. Sheep and cattle graze on these agricultural fields, which are actively maintained and fertilized. These activities were not documented and sporadic fertilization events do affect the NH₃ measurements, as will be discussed later.

2.2 Data filtering

We apply several filter criteria to the RITA-2021 observations, which are shown in Table 1 with acceptance rates for each individual filter criterion. The miniDOAS flux setup requires several days of intercalibration measurements, as described in Swart et al. (2023). No ammonia flux can be inferred from these intercalibration measurements, leaving 65% of the campaign observations suitable for flux measurements. We furthermore discard observations from 11 - 12 September, as these NH₃ emission fluxes are outliers with respect to the average observed NH₃ flux, indicating towards a fertilization event in close proximity to the measurement site.



Table 1. Filter criteria with the filter acceptance rates (in percentages and hours).

#	Filter	Criterion	Acceptance	
			[%]	[hours]
-	Unfiltered observations	-	100 %	1152
Discard 1:	miniDOAS intercalibration	-	65 %	746
Discard 2:	Fertilization events 11 - 12 September	-	61 %	698
Filter 1:	Wind direction	$331^\circ \geq U_d \geq 201^\circ$	16 %	188
Filter 2:	Rain duration	$t_{rain} \leq 5 \text{ min}$	16 %	179.5
Filter 3:	Turbulent mixing	$u_* > 0.1 \text{ m s}^{-1}$	13 %	151
Filter 4:	Gross primary production	$GPP > 0 \text{ mgC m}^{-2} \text{ s}^{-1}$	11 %	123.5
Filter 5:	Incoming short-wave radiation	$SW_{in} > 10 \text{ W m}^{-2}$	9 %	102

The remaining measurements are processed by applying 5 filters in total. The use of the flux-gradient method requires unobstructed upwind air flow with sufficient turbulent mixing. Figure 1 shows that the instruments were positioned anticipating winds from the south-west (green), with the obstacles located east (red) and north (yellow) of the miniDOAS optical path. We therefore apply a criterion filtering for wind directions between 201° and 331°. This filter leads to a large reduction data available for analysis, decreasing the available data from 61 % to 16 % as the prevalent wind direction during the campaign was from the north-east. As a secondary effect of this filter, the available observations are taken under synoptic weather conditions characterized frontal passages with some rain events. The second filter excludes rain events lasting more than 5 min, as rain droplets can obstruct the light path of the miniDOAS. Finally, sufficient turbulent mixing is one of the main requirements for flux measurement using the flux-gradient method. The third filter therefore requires the friction velocity to have a value of at least 0.1 m s⁻¹ ($u_* \geq 0.1 \text{ m s}^{-1}$). With these three filters, we ensure the quality of the ammonia measurements, observing the NH₃ flux with an accuracy of 0.01 μg m⁻²s⁻¹.

The fourth and fifth filter criteria focus on the ammonia surface-atmosphere exchange pathways. The NH₃ flux follows three pathways: the stomatal pathway, external leaf surface pathway and the soil pathway (Nemitz et al., 2001; Massad et al., 2010; van Zanten et al., 2010). The latter is generally assumed to be negligible for the F_{NH₃} over grass, as the dense vegetation completely covers the soil. The external leaf pathways, represents exchange of ammonia with a thin film of water and leaf surface waxes on the leaf surface, depending on the relative humidity (RH) (Van Hove et al., 1989). Finally, the stomatal pathway represents exchange of NH₃ through the plant stomata with ammonium dissolved in the apoplast fluids of the plant (Farquhar et al., 1980; Wichink Kruit et al., 2010). While these processes happen at very small scales, at micrometer or milimeter level, they can be represented at the canopy level, i.e. at field scale (Vilà-Guerau de Arellano et al., 2020).

The NH₃ exchange through the stomatal pathway is governed by the dynamic response of vegetation to meteorological conditions and is closely related to photosynthesis. The stomata open during the day in response to solar radiation, as the vegetation uses energy for photosynthesis, particularly the photosynthetically active radiation (PAR) (Hsiao, 1973; Cowan and Farquhar, 1977; Papaioannou et al., 1996; Ronda et al., 2001). Plants ingest CO₂ through the stomata, but water from inside the plant can evaporate as the stomata are opened. The plant can reduce this loss of water by (partly) closing the stomata in case of high water vapor pressure deficit (VPD), or increase the evaporation rate by actively opening the stomata. Increasing the evaporation rate provides cooling, lowering the leaf temperature to reach optimal conditions to perform photosynthesis (Jacobs and de Bruin, 1997; Takagi et al., 1998; de Groot et al., 2019; Vilà-Guerau de Arellano et al., 2020). As the temperature and VPD are often highest in the late afternoon, the stomata often partly close to manage the loss of water. During the night, there is no PAR for photosynthesis, so the stomata are closed. As a result, the characteristics of ammonia surface-atmosphere exchange of NH₃ differ between day and night, with the stomatal pathway being dominant during the day and the external leaf pathway being the dominant pathway during the night and in the early morning.

The uptake of CO₂ is represented by the Gross Primary Production (GPP), in mgC m⁻²s⁻¹. The GPP and the (soil) CO₂ respiration (RESP) combined define the Net Ecosystem Exchange (NEE) of CO₂, following $NEE = GPP + RESP$, which is measured as the CO₂ flux. The CO₂ respiration is estimated by taking the average campaign nighttime ($Q_{net} < 0$) CO₂ flux, which is approximately 2.2 mgCO₂ m⁻² s⁻¹. The GPP is then estimated by subtracting this respiration estimate from the observed CO₂ flux, or $GPP = NEE - RESP$. Note that a positive GPP (carbon uptake) corresponds to a observed negative CO₂ flux (CO₂ deposition).

To capture observations with active stomatal exchange, Filter 4 is set to only accept $GPP > 0 \text{ mgCO}_2 \text{ m}^{-2} \text{ s}^{-1}$. Due to the uncertainty of our GPP estimate, there are still some night-time observations which pass the filter. We therefore add an additional 5th filter using incoming shortwave radiation (SW_{in}). Only measurements with $SW_{in} > 10 \text{ W m}^{-2}$ will pass, in order to filter out these last remaining night-time observations.



Ruben B.Schulte: Observational relationships between NH₃, CO₂ and evapotranspiration

5

Table 2. A characterization of the meteorology of the 17 unique days at which any number of observations passes the filters, with the 17 day average and the range of the diurnal minimum/maximum of several (meteorological) variables. A requirement of non-zero net radiation is set for the GPP and footprint anemometer, due to the measurement uncertainty of these variables during stable nights.

Variable	Symbol	Diurnal minimum/maximum	
		17-day average	17-day range
Daily minimum temperature	T_{\min}	11.5 °C	5.6 - 16.7 °C
Daily maximum temperature	T_{\max}	19.7 °C	13.6 - 25.5 °C
Daily maximum wind speed	u	4.5 m s ⁻¹	2.2 - 7.2 m s ⁻¹
Daily maximum net radiation	Q_{net}	295 W m ⁻²	137 - 400 W m ⁻²
Daily maximum sensible heat flux	H	99 W m ⁻²	27 - 173 W m ⁻²
Daily maximum latent heat flux	L, E	145 W m ⁻²	83 - 230 W m ⁻²
Daily maximum gross primary production	GPP	0.78 mgC m ⁻² s ⁻¹	0.57 - 1.4 mgC m ⁻² s ⁻¹
Daily maximum water vapor pressure deficit	VPD	966 Pa	365 - 1420 Pa
Daytime maximum sonic anemometer footprint (70 %)	$fp_{70\%}$	148 m	88 - 255 m

After filtering, 102 hours (9 %) of all RITA-2021 observations, or 18 % of all daytime RITA-2021 observations, are available for analysis. These observations are taken over 17 unique days, spanning 29 August to 30 September, with an average of 6 hours and a maximum of 12 hours of accepted measurement per day.

2.3 Characterization of the campaign meteorology

The summer months (June, July and August) leading up to the RITA-2021 campaign are characterized as an average Dutch summer, with average temperatures (17.7 °C), above average precipitation (244 mm accumulated) and below average hours of sunshine (618 hours). Additionally the ground and surface water levels are actively managed for the agricultural activity in the area (Brauer et al., 2014). It is therefore expected that the role of long-term vegetation stress on stomatal exchange is negligible during the RITA-2021 campaign.

As discussed in Section 2.2, high temperatures or VPD can induce vegetation stress during the campaign. We therefore characterize the meteorological conditions of the 17 unique days in which the 102 hours of filtered measurements were taken. The meteorological conditions of these days are summarized in Table 2, which shows the 17 day average and the observed range of the diurnal minimum/maximum of several variables. The 17-day average values provide a characterization of mild meteorological conditions with no indication that the vegetation is under stress. Additionally, Table 2 includes an estimate of the maximum daytime footprint of the sonic anemometer, as a first-order approximation of the footprint of the NH₃ flux measurements. The footprint shown in Table 2 is the 70 % footprint of the sonic anemometer at a height of 2.8 m, following the method from Kljun et al. (2015).

As the filtered campaign measurements are characterized by frontal passages, the weather conditions range from clear sky summer conditions with moderately high temperatures, to colder cloudy days with short precipitation events (not shown). Furthermore, the atmospheric stability for the 102 hours of filtered measurements is classified using the measured Obukhov length (L) and the height of the sonic anemometer ($z = 2.8$ m). In total, 4.5 hours (4 %) can be classified as stable ($z/L > 0.05$), 61 hours (60 %) as neutral ($-0.05 \leq z/L \leq 0.05$) and 36.5 hours (36 %) as unstable conditions ($z/L < -0.05$). This variation leads to a large spread in all variables shown in Table 2, as indicated by the column showing the 17-day range.

2.4 Characterization of the NH₃ observations

The variety in meteorological conditions could be an explanation of the large day-to-day difference in the observed NH₃ concentrations, shown in Fig. 2b. The histogram shows that most observed NH₃ concentrations are below 7 μg m⁻³, with a long tail leading to a maximum concentration of 24.7 μg m⁻³. Still, the mean (solid line) and median (dotted line) concentrations do indicate that the concentration decreases during the day, until the late afternoon. This would be in line with observations at several other sites, both in the Netherlands (Wichink Kruit et al., 2007; Schulte et al., 2021) and in other countries, e.g. Scotland (von Bobruzki et al., 2010) or Italy (Ferrara et al., 2021). The large day-to-day differences in the NH₃ measurements could be a result of the changing meteorological conditions, the nearby agricultural activity, or a combination of both.

Despite the high variability in the NH₃ concentration measurements, a consistent diurnal variability is observed in the NH₃ gradient (ΔNH_3) and corresponding flux in Fig. 2c and 2d. Both Fig. 2b and 2c indicate that the observed ΔNH_3 is independent of the absolute NH₃ concentration, i.e. high absolute concentrations do not lead to a large concentration difference between the two miniDOAS instruments. The average diurnal variability is characterized by negative ΔNH_3 (deposition) in the early



morning and late afternoon and positive (emission) ΔNH_3 during the afternoon, with a typical range of about $0.5 \mu\text{g m}^{-3}$ in both directions. In total 79 % of the filtered observations have a positive ΔNH_3 , corresponding to NH_3 emissions.

As F_{NH_3} is directly inferred from ΔNH_3 , the diurnal variability in Fig. 2d is very similar. The NH_3 flux typically reaches its maximum around noon at little over $0.05 \mu\text{g m}^{-2}\text{s}^{-1}$ on average, with individual noon observations ranging from $-0.01 \mu\text{g m}^{-2}\text{s}^{-1}$ to $0.14 \mu\text{g m}^{-2}\text{s}^{-1}$. Note that the measurements taken on 11 - 12 September, the aforementioned fertilization event, are approximately a factor 4 larger than the mean campaign values. Despite the large observed F_{NH_3} on these days, the observed concentrations are only slightly larger than the campaign averages. These two days will not be included in the analysis presented in this study, but they are shown as an illustration of how fertilization events can impact our analysis.

2.5 Characterization of the ammonia flux

10 The main drivers of ammonia surface-atmosphere exchange are turbulent mixing, as well as the difference between the atmospheric NH_3 concentration and the so called canopy compensation point, i.e. an internal NH_3 concentration (Farquhar et al., 1980; Nemitz et al., 2001; Wichink Kruit et al., 2007). A compensation point lower than the atmospheric concentration leads to deposition of ammonia and vice-versa. The canopy compensation point consists of individual compensation points for each of the individual flux pathways. Following parameterizations of the stomatal compensation point, we find it related to the (leaf) 15 temperature and some form of nitrogen input parameter (actual or long-term NH_3 concentration). The stomatal compensation point increases non-linearly with increasing temperature or nitrogen input (Nemitz et al., 2001; Massad et al., 2010; van Zanten et al., 2010). As the stomatal compensation point increases and becomes larger than the atmospheric NH_3 concentration, or as the atmospheric concentration drops below this compensation point, emission will be observed from the stomatal exchange pathway.

20 In Fig. 3a, we show the observed ammonia flux against the observed temperature of the sonic anemometer, with the colors indicating the atmospheric NH_3 concentration at 2.29 m. Despite our efforts to filter for observations where the stomatal pathway is dominant, it is possible that the external leaf pathway still plays an important role in the morning, through deposition onto morning dew at canopy level (van Zanten et al., 2010; Wentworth et al., 2016). We therefore use black circles to mark observations taken before 12:00 UTC with $\text{RH} > 80\%$ in Fig. 3. These highlighted observations indeed generally correspond 25 with measurements of deposition or weak emission, indicating that NH_3 exchange through the external leaf pathway is still significant for these observations. While the inclusion complicate our analysis of stomatal NH_3 exchange, they are still included in the analysis as it also offers an opportunity to test if the relations found in the filtered dataset differ for the marked- and unmarked observations. If that is the case, it shows we are indeed able to attribute the unmarked observations to stomatal exchange.

30 Figure 3a shows that the observed F_{NH_3} increases with temperature for low atmospheric concentration ($2 \mu\text{g m}^{-3} \leq \text{NH}_{3, 2.29\text{m}} \leq 7 \mu\text{g m}^{-3}$). We attribute this increase in NH_3 emissions to the change in ΔNH_3 for increasing temperature, i.e. the difference between the approximately constant atmospheric NH_3 concentration and stomatal compensation point. This is indeed shown in Fig. 3b, where the theoretical stomatal compensation point (dotted line) increases with temperature. Here, the compensation point is calculated following the DEPosition of Acidifying Compounds (DEPAC) parameterization (van Zanten et al., 2010), 35 using the sonic anemometer temperature and campaign median $\text{NH}_{3, 2.29\text{m}}$ ($7.7 \mu\text{g m}^{-3}$).

This relation is scattered for measurements taken at high temperatures ($> 21^\circ\text{C}$). While there were only small variations in the NH_3 concentration for temperatures below 21°C , NH_3 concentration for these warmer temperatures higher than the campaign average ($> 7 \mu\text{g m}^{-3}$) and highly variable. As the NH_3 flux is directly related to the difference between the atmospheric NH_3 and the stomatal compensation point, the variability in the atmospheric concentration lead to the scatter shown in Fig. 3a, 40 where higher NH_3 observations correspond to weaker emission fluxes.

3 Ammonia flux relationships to dynamic vegetation responses

The diurnal pattern of F_{NH_3} in Fig. 2d shows similarities with the diurnal variability of the GPP in Fig. 2e. To further study the role of stomatal exchange during the campaign, we link the observed F_{NH_3} to the dynamic vegetation responses. First, we relate the ammonia flux to the GPP, the latent heat flux (L_vE) and the sensible heat flux (H). The GPP and L_vE are directly 45 governed by the opening and closing of the stomata and represent stomatal exchange. Given the low data availability (9%), we are aware that the analysis could be dominated by variations resulting from the diurnal variability of the fluxes. We therefore also include H in our analysis. The sensible heat flux is only indirectly related to the dynamic vegetation response through the surface energy balance, as the available energy from (solar) radiation and the soil heat flux is split between L_vE and H . If the observed fluxes are indeed regulated through the opening and closing of stomata, the analysis for F_{NH_3} to L_vE and GPP should 50 differ from the comparison with H .



Ruben B.Schulte: Observational relationships between NH_3 , CO_2 and evapotranspiration




Figure 2. The diurnal variability from sunrise (6:00 UTC) to sunset (18:00 UTC) of the filtered NH_3 concentration (b), NH_3 gradient (c), F_{NH_3} (d) and the GPP with the corresponding histogram to the right. At each moment in time, the multi-day mean (solid line) and median (dotted line) are calculated. Highlighted are observations from the fertilization event at 11 - 12 September (open circles). The N number of observations over which these averages are calculated are displayed at the top (a). ΔNH_3 is defined so the sign matches that of F_{NH_3} .




Figure 3. The 2.8 m temperature plotted against F_{NH_3} (a) and observed NH_3 , 2.29 m (b). The color coding in (a) represents the NH_3 concentration, observed at 2.29 m. In (b), the dotted line represents the theoretical stomatal compensation point (χ_s) for a long-term NH_3 concentration of $7.7 \mu\text{g m}^{-3}$. Highlighted with black circles are observations with $\text{RH} > 80\%$, taken before noon, where NH_3 exchange through the external leaf pathway can play a significant role.

Next, we organize the observations following current dynamic vegetation models, based on temperature, radiation and moisture (Jarvis et al., 1976; Stewart, 1988; Ronda et al., 2001). Here, we compare the responses of the four individual fluxes to temperature (T), PAR and VPD. As these three variables control the stomatal response at canopy level, we will use the responses of the fluxes to these variables as a guidance to better understand the diurnal variability of the ammonia flux. Note that 5 measurements taken on 11 - 12 September are not used to calculate correlation coefficients, but are shown in the figures and included in the visual analysis.

3.1 Relating the ammonia flux to photosynthesis

Plotting F_{NH_3} against the GPP in Fig. 4a shows a low positive correlation between the two fluxes, with a correlation coefficient of 0.48. There is a large spread in the data, particularly for GPP values larger than $0.125 \text{ mgC m}^{-2}\text{s}^{-1}$. Part of this spread is 10 attributed to the high relative humidity (black circles), where F_{NH_3} is not yet dominated by stomatal exchange and the external leaf pathway is still expected to be significant. Note that the atmospheric stability (color coded) plays an important role for the GPP as unstable conditions are typically characterized by clear skies and high PAR values, which favors photosynthesis (discussed in Section 3.2). This relation is not found in the observed F_{NH_3} , as there is a large spread in F_{NH_3} for both neutral and unstable conditions.

15 A moderate positive correlation is found in Fig. 4b between F_{NH_3} and L_vE . Our interpretation of this moderate correlation is that both evaporation and stomatal NH_3 emissions follow a similar process. The opening of the stomata for photosynthesis allows for the exchange of several gasses, including water vapor and ammonia, depending on VPD or the difference between atmosphere NH_3 and the stomatal compensation point (Cowan and Farquhar, 1977; Hsiao, 1973; Farquhar et al., 1980; Wichink Kruit et al., 2010). Note that L_vE represents the net L_vE since evaporation from the soil plays a role as well. Assuming a 20 vegetation cover of 90% for grass, soil evaporation could be between 10 and 30%. Note further that the observations with high relative humidity generally correspond to low L_vE and that again unstable conditions correspond with high L_vE values, related to the VPD between the leaf and stomata, and the atmosphere.

When plotting F_{NH_3} against H, two branches are found in the spread of the data, with a third branch being formed by the filtered out fertilization event on 11 - 12 September (black crosses). The smaller branch, with $F_{\text{NH}_3} > 0.1 \mu\text{g m}^{-2}\text{s}^{-1}$, could point 25 towards another (weaker) fertilization event. Still, the second highest positive correlation is found at 0.65, indicating that the natural diurnal variability indeed plays an important role. Note that most of the measurements with high relative humidity are clustered around $H = 0 \text{ W m}^{-2}$, i.e. there is little transfer of heat between the surface and atmosphere.

Based on the three scatter plots we find the highest correlation between F_{NH_3} and L_vE . Together with the diurnal variability of F_{NH_3} , transitioning from nighttime deposition to daytime emission from 8:30 to 16:30 UTC, this is the second indication 30 towards stomatal emission of NH_3 , opposed to emission from fertilization or animal droppings. However, the moderate corre-



Ruben B.Schulte: Observational relationships between NH₃, CO₂ and evapotranspiration

9




Figure 4. Scatter plots of F_{NH_3} against the GPP (a), L_vE (b) and H (c), with the colors indicating the ABL stability. Highlighted by black circles are observations with a $RH > 80\%$, where deposition through the external leaf path can still play an important role. The black crosses are observations from the fertilization event observed on 11 - 12 September.

lation between F_{NH_3} and H indicates that the diurnal variability of the fluxes influences the correlation coefficient. Finally, we want to mention the observations on 11 - 12 September, which support the interpretation of the scatter plots in showing how fertilization events affect our analysis.

3.2 The dynamic vegetation response to varying meteorological conditions

3.2.1 The dynamic response to temperature

We further investigate the stomatal exchange of NH_3 by analyzing the response of F_{NH_3} to varying meteorological conditions. The optimal conditions (PAR, T, VPD) for photosynthesis are different for different vegetation types (Gates, 1980; Jacobs, 1994; Vilà-Guerau de Arellano et al., 2015). Starting with the 2.8 meter temperature (T) in Fig. 5, we find a large spread for all four surface fluxes, resulting in low positive correlations (0.33 - 0.51). The lowest correlation coefficients are found for GPP and L_vE , indicating that temperature has little impact on the opening and closing of the stomata. Slightly higher correlation is found for F_{NH_3} , which we attribute to the relation between the stomatal compensation point and the NH_3 flux, discussed in Section 2.5. Note that the NH_3 emissions on 11 - 12 September stand out as outliers in Fig. 5a, while being average for the other three subplots.

5

10




Figure 5. Scatter plots of the temperature against F_{NH_3} (a), GPP (b), L_vE (c) and H (d), with the colors indicating the ABL stability (see Figure 4 for legend). Highlighted by black circles are observations with a $\text{RH} > 80\%$. The black crosses are observations from the fertilization event on 11 - 12 September.



Ruben B.Schulte: Observational relationships between NH_3 , CO_2 and evapotranspiration

11




Figure 6. Scatter plots of the VPD against F_{NH_3} (a), GPP (b), L_vE (c) and H (d), with the colors indicating the ABL stability (see Figure 4 for legend). Highlighted by black circles are observations with a $\text{RH} > 80\%$. The black crosses are observations from the fertilization event on 11 - 12 September.

3.2.2 The dynamic response to VPD

Moving on to analyzing the response of the four fluxes to the VPD, we find moderate correlation coefficients (0.42 - 0.53) in Fig. 6. L_vE shows in Fig. 6c a non-linear relationship with the VPD, called the evaporation hysteresis (Zhang et al., 2014; de Groot et al., 2019). This hysteresis is driven by both the vegetation regulating the loss of water through evaporation, described in Section 2.2, and the time difference when the maximum values for L_vE (12:00 UTC) and VPD (15:00 UTC) are reached. The same holds true for the other three fluxes (F_{NH_3} , GPP and H), as all three reach their maximum around noon.

Note that the observations of 11 - 12 September again are clear outliers in Fig. 6a, forming two branches in the scatter plot. Also standing out are several observations with $F_{\text{NH}_3} > 0.1 \mu\text{g m}^{-2}\text{s}^{-1}$. These are the observations that appear as the small upper branch in the $H - F_{\text{NH}_3}$ scatter plot in Fig. 4c and, again, form their separate branch here in Fig. 6a. This further indicates that there is a second (weak) fertilization event in the filtered dataset of the RITA-2021 campaign.




Figure 7. Scatter plots of PAR against F_{NH_3} (a), GPP (b), L_vE (c) and H (d), with the colors indicating the ABL stability (see Figure 4 for legend). Highlighted by black circles are observations with a RH > 80 %. The black crosses are observations from the fertilization event on 11 - 12 September.



3.2.3 The dynamic response to PAR

When relating F_{NH_3} to PAR, we find high positive correlation coefficients for all four surface fluxes (0.72 - 0.93) in Fig. 7. The highest value is found for H (Fig. 7d), as the solar radiation is the main source of energy for the surface-atmosphere exchange of heat. Where H exhibits a linear relation to PAR, non-linear effects are observed in the response of the two photosynthesis fluxes GPP and L_vE . This difference, together with the high correlation coefficients, indicates that PAR is the main driver of the dynamic vegetation response.

The GPP has a strongly non-linear response to PAR, as the GPP appears to reach a plateau for $\text{PAR} > 150 \text{ W m}^{-2}$. There are several reasons for this GPP maximum. At constant temperature and PAR, the stomatal uptake of CO_2 will increase the concentration within the plant to the point that the CO_2 supply is no longer the limiting factor. The GPP then reaches a plateau of maximum photosynthesis rate (see Figure 6.13a Moene and Van Dam (2014)), similar to the observations in Fig. 7b. Additionally, the photosynthesis system can become light saturated for high PAR values, at constant temperature. Following this latter process, the GPP is expected to level off more gradually, compared to the plateau that is reached by CO_2 saturation (see Figure 6.13b Moene and Van Dam (2014)). Finally, the (partly) closing of the stomata in response to high VPD could also reduce the GPP. However, as the VPD typically reaches its maximum around 15:00 UTC (not shown), it is unlikely that this is a limiting factor for GPP at high PAR values, which peaks around noon. All these processes depend on the temperature, VPD and PAR, and can explain the vertical spread in Fig. 7b.

When taking a close look at the response of the L_vE to PAR, it is possible to distinguish two phases in Fig. 7c. First, for PAR values up to about 100 W m^{-2} , the L_vE increases linearly to roughly 75 W m^{-2} , related to the opening/closing of the stomata around sunrise/sunset. The second phase shows a more gradual linear increase of L_vE to PAR. From the linear response and the small spread in Fig. 7c, we conclude that opening and closing of the stomata during the RITA-2021 campaign is governed by PAR and that the role of the VPD or temperature is small.

Similar to the L_vE , the NH_3 flux generally shows a linear response; transitioning from weak deposition to emission as the stomata open in response to increasing PAR. The spread in the F_{NH_3} response is larger compared to the L_vE response, which results in the lowest correlation coefficient at 0.72. We attribute this spread to three factors: the relation between temperature and the stomatal compensation point, the variations in the NH_3 concentration and the measurements where $\text{RH} > 80\%$ (black circles). Furthermore, observations where $F_{\text{NH}_3} > 0.1 \mu\text{g m}^{-2}\text{s}^{-1}$, i.e. the possible (weak) fertilization event, again appear to form a second branch in the scatter plot. Based on the strong similarities between F_{NH_3} and L_vE in their response to PAR, we conclude that the observed NH_3 emission is indeed stomatal (re)emission from vegetation.

4 Discussion

Observations of the NH_3 flux after filtering, taken over 17 individual days during the RITA-2021 campaign, are characterized by day-time emissions. The measurement site at Cabauw is located on flat grassland in an agricultural area, with the nearby fields being actively managed and/or grazed upon. It is therefore possible that the observed NH_3 emissions originate from sources like fertilization events (e.g. manure application) or animal droppings. We therefore first aim to establish relationships between the NH_3 emissions to stomatal exchange and follow up by attempting to find relations between F_{NH_3} and observations of the CO_2 and water vapor flux.

The first indication towards stomatal emission is found in the diurnal variability of F_{NH_3} . The flux transitions from deposition to emission in the early morning around 8:00 UTC, reaches maximum emission around 12:00 UTC and transitions to deposition again just before sunset, around 16:30 UTC, as shown in Fig. 2c. Our interpretation of this diurnal cycle is the flux transitions from (nighttime) NH_3 deposition, through the external leaf path, towards emission through the stomatal path during the day. This diurnal variability of F_{NH_3} shares similarities to the diurnal variability of the CO_2 flux. As the stomata open for photosynthesis in response to PAR, the CO_2 flux transitions from CO_2 respiration to stomatal uptake of CO_2 . High correlation between F_{NH_3} and L_vE (0.7) and between F_{NH_3} and PAR (0.72), further point towards stomatal NH_3 emission and a possible relation between F_{NH_3} and the photosynthesis fluxes.

4.1 Critical analysis of RITA-2021 dataset

The conditions during the RITA-2021 campaign present a challenge for the analysis conducted in this study. The site is located in an active agricultural region, with several potential emission sources within only a few hundred meters to a couple of kilometers distance upwind of the measurement site. The fields next to the site are actively managed and nitrogen contents of the soil and vegetation can differ on a field-to-field basis. This high level of surface heterogeneity within the estimated footprint of the flux measurements (up to about 250 m, Table 2) adds an additional level of complexity to the analysis (Swart et al., 2023). Furthermore, there are several farms located within two kilometers of the site, some of which with yearly NH_3 emissions up



to 12000 kg year⁻¹. Studies on the blending distance (i.e. the distance at which a plume can be considered well-mixed with respect to the background) indicate that emission plumes from such strong local NH₃ sources can affect flux measurements over distances of a couple of kilometers (Schulte et al., 2022). In this study, at least one instance of strong local emissions has been identified as the fertilization event on 11 - 12 September. Other potential weaker events have been shown and discussed as well in Fig. 4a, 4c and 6a.

The analysis is further complicated by the complex meteorological conditions, characterized by frontal passages. As the miniDOAS setup was positioned anticipating winds from the south-west, the meteorology of the filtered data is characterized by frontal passages. As a result, most observations are taken under neutral stability conditions (60 %), with clouds and some rain showers. While rain events are filtered out, wet deposition by rain does lead to a sudden change in the NH₃ concentration and can lead to re-emission of NH₃ as the rainwater evaporates.

Finally, the south-western orientation of the instruments leads to a significant loss in the availability of data suitable for analysis. Historically, southwestern winds tend to be most common in September, but the wind direction during the campaign was highly variable. Filtering for unobstructed wind directions reduces the availability of viable data by 510 hours, i.e. 44 % of all measurement data. As a result, the observed range in the measurements presented in the Figures is strongly influenced by the natural diurnal variability of the variables. While we do address the role of the natural diurnal variability by including the sensible heat flux in our analysis, it does make the observed relations between F_{NH₃} and the other variables somewhat speculative.

The high level of heterogeneity due to complex emission sources, the low data availability after filtering and the complex weather conditions make the RITA-2021 dataset unfavorable for establishing relationships between F_{NH₃} and the CO₂ or water vapor flux. It highlights the importance of homogeneity of the NH₃ surface characteristics and that proximity of NH₃ emission sources should be considered as well when selecting a measurement site, in addition to the availability of high quality meteorological observations. Despite the challenges, the NH₃ measurements are of unprecedented high quality (Swart et al., 2023) and analyzing the unique dataset following our approach is still worthwhile. While we do not manage to develop new relationships, we are able to link the observed F_{NH₃} emission to stomatal exchange, following an approach where we organize the observations following current dynamic vegetation models.

4.2 Recommendations

Following the results presented in this study, we recommend a comprehensive approach to future NH₃ flux measurements, including observations of the CO₂ and water vapor flux as auxiliary measurements. The opening of the stomata for CO₂ uptake through photosynthesis allows for the exchange of several other gasses, including water vapor and ammonia. The process representations of photosynthesis have been widely researched and it has been better tested against sub-diurnal observations under different scales (Vilà-Guerau de Arellano et al., 2020) and such auxiliary observations can be used to further our understanding of NH₃ exchange through the individual exchange pathways, as was done for ozone deposition by Visser et al. (2021).

Furthermore, we recommend to analyze and compare observations of the NH₃ flux at different measurement (grassland) sites, similar to the intercomparison of CO₂ exchange measurements by Jacobs et al. (2007). For example, the F_{NH₃} diurnal variability presented in this study significantly differs from measurements in 2013 at the Veenkampen meteorological site near the city of Wageningen (<https://www.wur.nl/en/show/Weather-Station-De-Veenkampen.htm>). Located only 50 km east, the diurnal variability of F_{NH₃} at Veenkampen is characterized by weak morning deposition and strong afternoon deposition, up to about -0.3 μg m⁻²s⁻¹, for clear sky conditions over unfertilized grassland (Schulte et al., 2021). At the Haarweg meteorological site, the predecessor to the Veenkampen, chemical wet denuder measurements of F_{NH₃} in 2004 were characterized by strong deposition in the early morning, attributed to morning dew, and weak stomatal emissions in the afternoon (Wichink Kruit et al., 2007). The differences between observed diurnal variability in these three studies stress the high variability at the local and regional scales and highlight the need for long term high-resolution F_{NH₃} observations at multiple locations.

Efforts to further our understanding of the NH₃ exchange and its diurnal variability are already being made. The miniDOAS setup used in the RITA-2021 will be taking long-term (> 1 year) observations of the NH₃ flux at the Veenkampen meteorological site, starting in the spring of 2023. This year-long record of high-resolution F_{NH₃} observations will be analyzed, alongside a wide range of meteorological and turbulent measurements, including the CO₂ and water vapor flux, aiming to improve the parameterization of the NH₃ surface-atmosphere exchange. The collocation of surface and upper-atmospheric observations (Vilà-Guerau de Arellano et al., 2023) is key to obtain a comprehensive and complete understanding of NH₃ flux. The analysis can be taken one step further in the context of the Ruisdael Observatory project, following a process analysis combining the observations with both conceptual (Schulte et al., 2021) and high-resolution turbulent resolved models (Schulte et al., 2022).



5 Conclusions

We analyze over a month of ammonia flux measurements (F_{NH_3}), taken during the RITA-2021 campaign at the Ruisdael Observatory at Cabauw. The analysis is centered around observations from the miniDOAS flux measurement setup, which applies the flux-gradient method to line average concentration measurements over a 22 m open-path at two heights. Our objective was to find relationships between the observed NH₃ flux and the other turbulent surface fluxes such as sensible heat flux and photosynthesis, i.e. the stomatal exchange of CO₂ and water vapor (plant transpiration). The process of photosynthesis has been more widely studied and such auxiliary observations can be used to advance our understanding of NH₃ surface-atmosphere exchange through the individual exchange pathways, e.g. stomatal exchange.

After filtering, the observed F_{NH_3} is characterized by daytime emissions, averaging at about $0.05 \mu\text{g m}^{-2}\text{s}^{-1}$, and nighttime deposition of about $-0.05 \mu\text{g m}^{-2}\text{s}^{-1}$. We compare the NH₃ flux to the CO₂ uptake by vegetation and the exchange of water vapor, represented by the gross primary production (GPP) and net latent heat flux (L_vE), as well as the sensible heat flux (H) which is only indirectly related to the dynamic vegetation response. Here, we find high correlation between the observed daytime NH₃ emissions and L_vE (0.70) and the photosynthetically active radiation (PAR, 0.72). These results provide a first-order quantification of stomatal emission of NH₃. It shows that auxiliary and collocated flux measurements of CO₂ and water vapor are appropriate variables to distinguish stomatal NH₃ exchange from non-stomatal exchange.

The analysis presented in this study is hampered by the challenging conditions during the RITA-2021 campaign. The measurement site is located in a highly agricultural region, with actively managed (i.e. fertilized) grassland and emission sources in close proximity (< 2 km) of the NH₃ measurements. These heterogeneous NH₃ surface characteristics lead to outlier observations caused by at least one fertilization event and there are indications of other (weaker) fertilization events. Additional layers of complexity are added by the varying meteorological conditions, characterized by frontal passages, and the small subset of the campaign measurements being suitable for analysis after filtering (9 % or 102 hours). The latter is largely the outcome of filtering out unfavorable wind directions, where the airflow is obstructed. As a result, the observed range in the measurements is largely governed by the natural diurnal variability, hampering the efforts to establish clear relationships.

Despite the challenging conditions, the comprehensive approach presented in this study paves the way for the potential of combining high-quality NH₃ observations with auxiliary flux measurements of CO₂, water vapor and other meteorological variables. By organizing the observations following current dynamic vegetation models, we managed to attribute the observed NH₃ emission to stomatal exchange and identify outliers. In order to establish relations between the NH₃ and the photosynthesis fluxes, the approach should be applied to measurements that are still representative of the sources and sinks, but at the same time are well mixed and outside the blending distance of nearby emission sources. Further, longer time series are needed in order to make a better distinction between days with and without the influences of nearby sources. The results presented in this study already indicate that there is room to find such patterns.

Author Contributions

RS: Conceptualization, Formal Analysis, Investigation, Methodology, Visualization, Writing - original draft preparation. JVGdA: Conceptualization, Writing - Review and Editing, Supervision MCvZ: Conceptualization, Funding Acquisition, Writing - Review and Editing. JZ: Investigation (CO₂ flux and meteorological data). SvdG and SJR Investigation (NH₃ flux data). MCvZ: Conceptualization, Funding Acquisition, Writing - Review and Editing.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements. Ruben B. Schulte PhD project was supported by RIVM within the framework of project 36.7: Monitoring of dry ammonia deposition; financed by the Dutch Ministry of Agriculture, Nature and Food Quality. We thank the Royal Netherlands Meteorological Institute (KNMI) for site access and assistance during the RITA -2021 campaign. All people of RIVM and TNO involved in this campaign are gratefully acknowledged for their support and fruitful discussions during the analysis of the data. This article has been accomplished by using data generated in the Ruisdael Observatory, a scientific research infrastructure which is (partly) financed by the Dutch Research Council (NWO, grant number 184.034.015).



References

- Behera, S. N., Sharma, M., Aneja, V. P., and Balasubramanian, R.: Ammonia in the atmosphere: a review on emission sources, atmospheric chemistry and deposition on terrestrial bodies, *Environmental Science and Pollution Research*, 20, 8092 – 8131, <https://doi.org/10.1007/s11356-013-2051-9>, 2013.
- 5 Berkhout, A. J. C., Swart, D. P. J., Volten, H., Gast, L. F. L., Haaïma, M., Verboom, H., Stefess, G., Hafkenscheid, T., and Hoogerbrugge, R.: Replacing the AMOR with the miniDOAS in the ammonia monitoring network in the Netherlands, *Atmospheric Measurement Techniques*, 10, 4099–4120, <https://doi.org/10.5194/amt-10-4099-2017>, 2017.
- Bobbink, R., Hornung, M., and Roelofs, J. G. M.: The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation, *Journal of Ecology*, 86, 717 – 738, <https://doi.org/10.1046/j.1365-2745.1998.8650717.x>, 2003.
- 10 Bosveld, F., Baas, P., Beljaars, A., Holtslag, A., Vilà-Guerau de Arellano, J., and van de Wiel, B.: Fifty Years of Atmospheric Boundary-Layer Research at Cabauw Serving Weather, Air Quality and Climate, *Boundary-Layer Meteorology*, 177, 583–612, <https://doi.org/10.1007/s10546-020-00541-w>, 2020.
- Brauer, C. C., Torfs, P. J. J. F., Teuling, A. J., and Uijlenhoet, R.: The Wageningen Lowland Runoff Simulator (WALRUS): application to the Hupsel Brook catchment and the Cabauw polder, *Hydrology and Earth System Sciences*, 18, 4007–4028, <https://doi.org/10.5194/hess-18-4007-2014>, 2014.
- 15 Cowan, I. and Farquhar, G.: Stomatal function in relation to leaf metabolism and environment, Cambridge: at the university press, pp. 471 – 505, 1977.
- de Groot, G. E., Vilà-Guerau de Arellano, J., de Roode, S. R., Coenders-Gerrits, A. M. J., and van de Wiel, B. J. H.: Evaporation Hysteresis over Vegetation: The Impact of Surface Processes and Boundary Layer Dynamics, Master's thesis, Delft University of Technology, the Netherlands, <http://resolver.tudelft.nl/uuid:8284f137-2c64-48ab-ba8d-bf099d70c70c>, 2019.
- 20 Erisman, J. and Schaap, M.: The need for ammonia abatement with respect to secondary PM reductions in Europe, *Environmental Pollution*, 129, 159–163, <https://doi.org/10.1016/j.envpol.2003.08.042>, 2004.
- Erisman, J. W., Galloway, J. N., Seitzinger, S., Bleeker, A., Dise, N. B., Petrescu, A. M. R., Leach, A. M., and de Vries, W.: Consequences of human modification of the global nitrogen cycle, *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368, 20130 116, <https://doi.org/10.1098/rstb.2013.0116>, 2013.
- 25 Farquhar, G. D., Firth, P. M., Wetselaar, R., and Weir, B.: On the Gaseous Exchange of Ammonia between Leaves and the Environment: Determination of the Ammonia Compensation Point, *Plant Physiology*, 66, 710–714, <https://doi.org/10.1104/pp.66.4.710>, 1980.
- Ferrara, R., Di Tommasi, P., Famulari, D., and Rana, G.: Limitations of an Eddy-Covariance System in Measuring Low Ammonia Fluxes, *Boundary-Layer Meteorology*, 180, 173–186, <https://doi.org/10.1007/s10546-021-00612-6>, 2021.
- 30 Gates, D. M.: *Biophysical ecology*, Springer-Verlag, 1980.
- Hsiao, T.: Plant responses to water stress, *Annu. Rev. Plant Physiol.*, 24, 519–570, 1973.
- Jacobs, C.: Direct impact of atmospheric CO₂ enrichment on regional transpiration, Ph.D. thesis, Wageningen University, <https://wur.on.worldcat.org/oclc/906644518>, 1994.
- Jacobs, C. M. J. and de Bruin, H. A. R.: Predicting Regional Transpiration at Elevated Atmospheric CO₂: Influence of the PBL–Vegetation Interaction, *Journal of Applied Meteorology*, 36, 1663 – 1675, [https://doi.org/10.1175/1520-0450\(1997\)036<1663:PRTAEA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1997)036<1663:PRTAEA>2.0.CO;2), 1997.
- 35 Jacobs, C. M. J., Jacobs, A. F. G., Bosveld, F. C., Hendriks, D. M. D., Hensen, A., Kroon, P. S., Moors, E. J., Nol, L., Schrier-Uijl, A., and Veenendaal, E. M.: Variability of annual CO₂ exchange from Dutch grasslands, *Biogeosciences*, 4, 803–816, <https://doi.org/10.5194/bg-4-803-2007>, 2007.
- 40 Jarvis, P. G., Monteith, J. L., and Weatherley, P. E.: The interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field, *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 273, 593–610, <https://doi.org/10.1098/rstb.1976.0035>, 1976.
- Kljun, N., Calanca, P., Rotach, M. W., and Schmid, H. P.: A simple two-dimensional parameterisation for Flux Footprint Prediction (FFP), *Geoscientific Model Development*, 8, 3695–3713, <https://doi.org/10.5194/gmd-8-3695-2015>, 2015.
- 45 Massad, R.-S., Nemitz, E., and Sutton, M. A.: Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere, *Atmospheric Chemistry and Physics*, 10, 10 359–10 386, <https://doi.org/10.5194/acp-10-10359-2010>, 2010.
- McDermitt, D. K., Burba, G., Xu, L., Anderson, T. G., Komissarov, A. V., Riensche, B., Schedlbauer, J. L., Starr, G., Zona, D., Oechel, W. C., Oberbauer, S. F., and Hastings, S. J.: A new low-power, open-path instrument for measuring methane flux by eddy covariance, *Appl. Phys. B*, 102, 391–405, <https://doi.org/https://doi.org/10.1007/s00340-010-4307-0>, 2011.
- 50 Milford, C., Hargreaves, K. J., Sutton, M. A., Loubet, B., and Cellier, P.: Fluxes of NH₃ and CO₂ over upland moorland in the vicinity of agricultural land, *Journal of Geophysical Research: Atmospheres*, 106, 24 169–24 181, <https://doi.org/10.1029/2001JD900082>, 2001.
- Moene, A. F. and Van Dam, J. C.: *Transport in the Atmosphere-Vegetation-Soil Continuum*, Cambridge University Press, 2014.
- Nemitz, E., Milford, C., and Sutton, M.: A two-layer canopy compensation point model for describing bi-directional biosphere-atmosphere exchange of ammonia, *Quarterly Journal of the Royal Meteorological Society*, 127, 815–833, <https://doi.org/10.1002/qj.49712757306>, 2001.
- 55 Nemitz, E., Sutton, M., Wyers, G., and Jongejan, P.: Gas-particle interactions above a Dutch heathland: I. Surface exchange fluxes of NH₃, SO₂, HNO₃ and HCl, *Atmospheric Chemistry and Physics*, 4, 989–1005, <https://doi.org/10.5194/acp-4-989-2004>, 2004.
- Papaioannou, G., Nikolidakis, G., Asimakopoulos, D., and Retalis, D.: Photosynthetically active radiation in Athens, *Agricultural and Forest Meteorology*, 81, 287–298, [https://doi.org/10.1016/0168-1923\(95\)02290-2](https://doi.org/10.1016/0168-1923(95)02290-2), 1996.



Ruben B.Schulte: Observational relationships between NH₃, CO₂ and evapotranspiration

17

- Parrish, D. D. and Fehsenfeld, F. C.: Methods for gas-phase measurements of ozone, ozone precursors and aerosol precursors, *Atmospheric Environment*, 34, 1921–1957, [https://doi.org/10.1016/S1352-2310\(99\)00454-9](https://doi.org/10.1016/S1352-2310(99)00454-9), 2000.
- RIVM, CBS, PBL, and WUR: Nitrogen deposition, 1990-2018, Available at <https://www.clo.nl/indicatoren/en018918>, (Indicator 0189, version 18, 25 November 2019) RIVM National Institute for Public Health and the Environment, Bilthoven; Statistics Netherlands (CBS), The Hague; PBL Netherlands Environmental Assessment Agency, The Hague; and Wageningen University and Research, Wageningen. (Last access: 17 August 2022), 2019. 5
- Ronda, R. J., de Bruin, H. A. R., and Holtslag, A. A. M.: Representation of the Canopy Conductance in Modeling the Surface Energy Budget for Low Vegetation, *Journal of Applied Meteorology*, 40, 1431–1444, [https://doi.org/10.1175/1520-0450\(2001\)040<1431:ROTCI>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<1431:ROTCI>2.0.CO;2), 2001.
- San José, J. J., Montes, R., and Nikonova-Crespo, N.: Carbon Dioxide and Ammonia Exchange in the Trachypogon Savannas of the Orinoco Llanos, *Annals of Botany*, 68, 321–328, <https://doi.org/10.1093/oxfordjournals.aob.a088259>, 1991. 10
- Schrader, F., Erisman, J. W., and Brümmner, C.: Towards a coupled paradigm of NH₃-CO₂ biosphere–atmosphere exchange modelling, *Global Change Biology*, 26, 4654–4663, <https://doi.org/10.1111/gcb.15184>, 2020.
- Schulte, R., van Zanten, M., Rutledge-Jonker, S., Swart, D., Wichink Kruit, R., Krol, M., van Pul, W., and Vilà-Guerau de Arellano, J.: Unraveling the diurnal atmospheric ammonia budget of a prototypical convective boundary layer, *Atmospheric Environment*, 249, 118 153, <https://doi.org/10.1016/j.atmosenv.2020.118153>, 2021. 15
- Schulte, R. B., van Zanten, M. C., van Stratum, B. J. H., and Vilà-Guerau de Arellano, J.: Assessing the representativity of NH₃ measurements influenced by boundary-layer dynamics and the turbulent dispersion of a nearby emission source, *Atmospheric Chemistry and Physics*, 22, 8241–8257, <https://doi.org/10.5194/acp-22-8241-2022>, 2022.
- Smit, L. A. M. and Heederik, D.: Impacts of Intensive Livestock Production on Human Health in Densely Populated Regions, *GeoHealth*, 1, 272–277, <https://doi.org/10.1002/2017GH000103>, 2017. 20
- Stewart, J.: Modelling surface conductance of pine forest, *Agricultural and Forest Meteorology*, 43, 19–35, [https://doi.org/10.1016/0168-1923\(88\)90003-2](https://doi.org/10.1016/0168-1923(88)90003-2), 1988.
- Stokstad, E.: Nitrogen crisis threatens Dutch environment- and economy, *Science*, 366, 1180–1181, <https://doi.org/10.1126/science.366.6470.1180>, 2019. 25
- Swart, D., Zhang, J., van der Graaf, S., Rutledge-Jonker, S., Hensen, A., Berkhout, S., Wintjen, P., van der Hoff, R., Haaima, M., Frumau, A., van den Bulk, P., Schulte, R., van Zanten, M. C., and van Goethem, T.: Field comparison of two novel open-path instruments that measure dry deposition and emission of ammonia using flux-gradient and eddy covariance methods, *Atmospheric Measurement Technique*, 2023, 529–546, <https://doi.org/10.5194/amt-16-529-2023>, 2023.
- Takagi, K., Tsuboya, T., and Takahashi, H.: Diurnal hystereses of stomatal and bulk surface conductances in relation to vapor pressure deficit in a cool-temperate wetland, *Agricultural and Forest Meteorology*, 91, 177–191, [https://doi.org/10.1016/S0168-1923\(98\)00078-1](https://doi.org/10.1016/S0168-1923(98)00078-1), 1998. 30
- Van Hove, L., Adema, E., Vredenberg, W., and Pieters, G.: A study of the adsorption of NH₃ and SO₂ on leaf surfaces, *Atmospheric Environment* (1967), 23, 1479–1486, [https://doi.org/10.1016/0004-6981\(89\)90407-1](https://doi.org/10.1016/0004-6981(89)90407-1), 1989.
- van Zanten, M. C., Sauter, F. J., Wichink Kruit, R. J., van Jaarsveld, J. A., and van Pul, W. A. J.: Description of the DEPAC module: Dry deposition modelling with DEPAC-GCN2010, Tech. Rep. 680180001, National Institute for Public Health and the Environment (RIVM), <https://www.rivm.nl/bibliotheek/rapporten/680180001.pdf>, 2010. 35
- Vilà-Guerau de Arellano, J., Ney, P., Hartogensis, O., de Boer, H., van Diepen, K., Emin, D., de Groot, G., Klosterhalfen, A., Langensiepen, M., Matveeva, M., Miranda-García, G., Moene, A. F., Rascher, U., Röckmann, T., Adnew, G., Brüggemann, N., Rothfuss, Y., and Graf, A.: CloudRoots: integration of advanced instrumental techniques and process modelling of sub-hourly and sub-kilometre land–atmosphere interactions, *Biogeosciences*, 17, 4375–4404, <https://doi.org/10.5194/bg-17-4375-2020>, 2020. 40
- Vilà-Guerau de Arellano, J., van Heerwaarden, C. C., van Stratum, B. J. H., and van den Dries, K.: *Atmospheric Boundary Layer: Integrating Air Chemistry and Land Interactions*, Cambridge University Press, <https://doi.org/10.1017/CBO9781316117422>, 2015.
- Vilà-Guerau de Arellano, J., Hartogensis, O., Benedict, I., de Boer, H., Bosman, P. J. M., Botía, S., Cecchini, M. A., Faassen, K. A. P., González-Armas, R., van Diepen, K., Heusinkveld, B. G., Janssens, M., Lobos-Roco, F., Luijkx, I. T., Machado, L. A. T., Mangan, M. R., Moene, A. F., Mol, W. B., van der Molen, M., Moonen, R., Ouwensloot, H. G., Park, S.-W., Pedruzo-Bagazgoitia, X., Röckmann, T., Adnew, G. A., Ronda, R., Sikma, M., Schulte, R., van Stratum, B. J. H., Veerman, M. A., van Zanten, M. C., and van Heerwaarden, C. C.: Advancing understanding of land–atmosphere interactions by breaking discipline and scale barriers, *Annals of the New York Academy of Sciences*, 00, 1–24, <https://doi.org/https://doi.org/10.1111/nyas.14956>, 2023. 45
- Visser, A. J., Ganzeveld, L. N., Goded, I., Krol, M. C., Mammarella, I., Manca, G., and Boersma, K. F.: Ozone deposition impact assessments for forest canopies require accurate ozone flux partitioning on diurnal timescales, *Atmospheric Chemistry and Physics*, 21, 18 393–18 411, <https://doi.org/10.5194/acp-21-18393-2021>, 2021. 50
- von Bobrutzki, K., Braban, C. F., Famulari, D., Jones, S. K., Blackall, T., Smith, T. E. L., Blom, M., Coe, H., Gallagher, M., Ghalaieny, M., McGillen, M. R., Percival, C. J., Whitehead, J. D., Ellis, R., Murphy, J., Mohacsi, A., Pogany, A., Junninen, H., Rantanen, S., Sutton, M. A., and Nemitz, E.: Field inter-comparison of eleven atmospheric ammonia measurement techniques, *Atmospheric Measurement Techniques*, 3, 91–112, <https://doi.org/10.5194/amt-3-91-2010>, 2010. 55
- Wentworth, G., Murphy, J., Benedict, K., Bangs, E., and Collett Jr., J.: The role of dew as a night-time reservoir and morning source for atmospheric ammonia, *Atmospheric Chemistry and Physics*, 16, 7435–7449, <https://doi.org/10.5194/acp-16-7435-2016>, 2016.



- Whitehead, J. D., Twigg, M., Famulari, D., Nemitz, E., Sutton, M. A., Gallagher, M. W., and Fowler, D.: Evaluation of Laser Absorption Spectroscopic Techniques for Eddy Covariance Flux Measurements of Ammonia, *Environmental Science and Technology*, 42, 2041–2046, <https://doi.org/10.1021/es071596u>, 2008.
- Wichink Kruit, R. J. and van Pul, W. A. J.: Ontwikkelingen in de stikstofdepositie, Briefrapport 2018-0117, National Institute for Public Health and the Environment (RIVM), <https://doi.org/10.21945/RIVM-2018-0117>, 2018.
- 5 Wichink Kruit, R. J., van Pul, W. A. J., Otjes, R. P., Hofschreuder, P., Jacobs, A. F. G., and Holtslag, A. A. M.: Ammonia fluxes and derived canopy compensation points over non-fertilized agricultural grassland in The Netherlands using the new GRadient Ammonia - High Accuracy - Monitoring (GRAHAM), *Atmospheric Environment*, 41, 1275 – 1287, <https://doi.org/https://doi.org/10.1016/j.atmosenv.2006.09.039>, 2007.
- 10 Wichink Kruit, R. J., van Pul, W. A. J., Sauter, F. J., van den Broek, M., E., N., Sutton, M. A., Krol, M., and Holtslag, A. A. M.: Modeling the surface–atmosphere exchange of ammonia, *Atmospheric Environment*, 44, 945 – 957, <https://doi.org/10.1016/j.atmosenv.2009.11.049>, 2010.
- Wyers, G., Otjes, R., and Slanina, J.: A continuous-flow denuder for the measurement of ambient concentrations and surface-exchange fluxes of ammonia, *Atmospheric Environment. Part A. General Topics*, 27, 2085–2090, [https://doi.org/10.1016/0960-1686\(93\)90280-C](https://doi.org/10.1016/0960-1686(93)90280-C), 1993.
- 15 Zhang, Q., Manzoni, S., Katul, G., Porporato, A., and Yang, D.: The hysteretic evapotranspiration — Vapor pressure deficit relation, *Journal of Geophysical Research: Biogeosciences*, 119, 125–140, <https://doi.org/https://doi.org/10.1002/2013JG002484>, 2014.
- Zöll, U., Lucas-Moffat, A., Wintjen, P., Schrader, F., Beudert, B., and Bruemmer, C.: Is the biosphere-atmosphere exchange of total reactive nitrogen above forest driven by the same factors as carbon dioxide? An analysis using artificial neural networks, *Atmospheric Environment*, 206, 108–118, <https://doi.org/10.1016/j.atmosenv.2019.02.042>, 2019.