



Diagnosing drivers of PM_{2.5} simulation biases from meteorology, chemical composition, and emission sources using an efficient machine learning method

Shuai Wang¹, Mengyuan Zhang¹, Yueqi Gao¹, Peng Wang^{2,3}, Qingyan Fu⁴, Hongliang Zhang^{1,3,5}

5 ¹Department of Environmental Science and Engineering, Fudan University, Shanghai 200438, China

²Department of Atmospheric and Oceanic Sciences, and Institute of Atmospheric Sciences, Fudan University, Shanghai, 200438, China

³IRDR ICoE on Risk Interconnectivity and Governance on Weather/Climate Extremes Impact and Public Health, Fudan University, Shanghai, China;

10 ⁴Shanghai Environmental Monitoring Center, Shanghai 200235, China

⁵Institute of Eco-Chongming (IEC), Shanghai 200062, China

Correspondence to: Hongliang Zhang (zhanghl@fudan.edu.cn)

Abstract. Chemical transport models (CTMs) are widely used for air pollution modeling, which suffer from significant biases due to uncertainties in simplified parameterization, meteorological fields, and emission inventories. Accurate diagnosis of simulation biases is critical for improvement of models, interpretation of results, and efficient air quality management, especially for the simulation of fine particulate matter (PM_{2.5}). In this study, an efficient method based on machine learning (ML) was designed to diagnose the drivers of the Community Multiscale Air Quality (CMAQ) model biases in simulating PM_{2.5} concentrations from three perspectives of meteorology, chemical composition, and emission sources. The source-oriented CMAQ were used to diagnose influences of different emission sources on PM_{2.5} biases. The ML models showed good fitting ability with small performance gap between training and validation. The CMAQ model underestimates PM_{2.5} by -19.25 to -2.66 µg/m³ in 2019, especially in winter and spring and high PM_{2.5} events. Secondary organic components showed the largest contribution to PM_{2.5} simulation bias for different regions and seasons (13.8 - 22.6%) among components. Relative humidity, cloud cover, and soil surface moisture were the main meteorological factors contributing to PM_{2.5} bias in the North China Plain, Pearl River Delta, and northwestern, respectively. Both primary and secondary inorganic components from residential sources showed the largest contribution (12.05 % and 12.78 %), implying large uncertainties in this sector. The ML-based methods provide valuable complements to traditional mechanism-based methods for model improvement, with high efficiency and low reliance on prior information.

1 Introduction

Fine particulate matter (PM_{2.5}) is a complex mixture of primary (PPM) and secondary inorganic/organic components (SIA/SOA), with adverse effects on public health and ecosystems. Ambient levels of PM_{2.5} are influenced by meteorological conditions, emission from different sources, and atmospheric chemical processes (Organization, 2021; Xiao et al., 2021a; Yang et al., 2016; Liu et al., 2021b; Zhai et al., 2019). China has experienced severe PM_{2.5} pollution over the past two decades (Bai et al., 2022; Liang et al., 2020a). For effective air quality management, accurate PM_{2.5} modeling is essential. Chemical transport models (CTMs) like the Community Multiscale Air Quality (CMAQ) model, have been widely developed and applied to PM_{2.5} simulations through atmospheric processes of dispersion, deposition, and chemical reactions (Qiao et al., 2018; Wang et al., 2021; Hu et al., 2017a). Application of CTMs simulations is often limited by the biases due to uncertainties of simplified parameterization, meteorological prediction, emission inventories, initial and boundary conditions (Binkowski and Roselle, 2003; Hu et al., 2014; Hu et al., 2016; Wang et al., 2023c; Wang et al., 2021). Thus, it is essential to diagnose specific sources of simulation biases according to specific model applications, including grid resolution, parameterization, mechanisms,



40 meteorological inputs, and emission inventories.

Traditional bias diagnosis approaches for CTM models usually rely on empirical and prior assumptions with extensive sensitivity testing and high demands on computational resources, such as Monte Carlo methods or Latin hypercube sampling (Beekmann and Derognat, 2003; Hanna et al., 2005; Aleksankina et al., 2019). Recently machine learning (ML) methods have been widely used in environmental science researches due to their simple structure, fast speed and ability to deal with non-linear relationships (Liu et al., 2022). Many studies used ML to predict air pollutant concentrations like PM_{2.5} and ozone (Wei et al., 2021a; Sun et al., 2021; Zhu et al., 2022; Bai et al., 2022), improve the accuracy of CTMs simulations (Wang et al., 2023; Wei et al., 2020), and explain the prediction results using interpretable ML techniques (Hou et al., 2022; Li et al., 2023; Stirnberg et al., 2021). To date, few studies have used ML to diagnose the simulation bias of CTMs. A study has shown the potential of machine learning in explaining the simulation bias of ozone (Ye et al., 2022). However, as a complex multi-phase mixture, it is still challenging to diagnose biases in PM_{2.5} simulations using ML methods (Liu and Xing, 2022). Moreover, due to the strong influence of emissions, it is instructive to diagnose CTMs biases of PM_{2.5} based on a source-appointment perspective.

In this study, we use the Light Gradient Boosting Machine (lightGBM) model, an efficient ensemble ML approach, to diagnose the drivers of CMAQ biases in simulating PM_{2.5} concentrations. A source-oriented version of CMAQ is used to track sectoral source contributions to PM_{2.5}. Model biases are diagnosed from multiple perspectives, including meteorology, chemical components, and emission sources.

2 Materials and methods

2.1 Surface PM_{2.5} observations

This study focused on 2019 because of the large number of observations, the reliability of the emission inventories, and without interference of COVID19. Hourly PM_{2.5} observations for 2019 are collected from the China National Environmental Monitoring Centre (CNEMC, <http://www.cnemc.cn/>). The hourly data from each station was sanity checked to exclude problematic data points, and the 24-hour average concentrations of PM_{2.5} with less than 20 hourly valid records are excluded. For observation sites located on the same simulation grid, average PM_{2.5} concentrations were calculated to compare with CMAQ simulation. The distribution of observation sites is shown in Figure S1. Analysis focuses on nationwide as well as several interested regions (Figure S1), including Beijing-Tianjin-Hebei (BTH); the Yangtze River Delta (YRD); the Pearl River Delta (PRD); the Sichuan Basin (SCB); and Northwestern China (NWCHN).

2.2 CMAQ simulation

The Weather Research & Forecasting Model (WRF v4.2) was used to generate meteorological fields driven by National Centers for Environmental Prediction (NCEP) FNL Operational Model Global Tropospheric Analyses dataset (<http://rda.ucar.edu/datasets/ds083.2/>) (Commerce, 2000). Several meteorological factors (Table S1) that are highly relevant to aerosol concentrations are selected for ML model building (Xiao et al., 2021b; Chen et al., 2020b; Meng et al., 2019). The CMAQ v5.0.2 with a modified SAPRC-11 photochemical mechanism and AERO6 aerosol module was applied for aerosol simulations (Carter and Heo, 2013; Ying et al., 2015; Binkowski and Roselle, 2003). Details of CMAQ model improvement can be found in previous studies (Hu et al., 2017b; Ying et al., 2015). The Multi-resolution Emission Inventory for China (MEIC) was used as anthropogenic emission (<http://meicmodel.org/>), and the Model for Emissions of Gases and Aerosols from Nature (MEGAN) version 2.1 was used to generate biogenic emissions (Guenther et al., 2012; Guenther et al., 2006). The Fire INventory from NCAR (FINN) based on satellite were used to generate open burning emission (Wiedinmyer et al., 2011). The list of PM_{2.5} components simulated by CMAQ is shown in Table S1. The CMAQ simulation (36 km×36 km) was conducted over mainland China and surrounding regions in 2019.



80 The source apportionment method was used to quantify the contributions of industry, energy, residential, transportation, agriculture, open burning, and biogenic sources to PPM and SIA through a modified version of CMAQ (Zhang et al., 2012; Ma et al., 2021; Qiao et al., 2018). PPM from different source sectors are tracked by non-reactive tracers (10^{-5} of the PPM emission rates). The concentrations of PPM from given sources are then calculated by multiplying the tracer with 10^5 . The contributions of source sectors to SIA are quantified using specific reactive tagged tracers. NO_x , SO_2 , and NH_3 from different
85 sources were tracked separately through a series of chemical and physical processes involving in SIA formation. Details of source apportionment can be found in previous studies (Zhang et al., 2012; Ma et al., 2021; Qiao et al., 2018; Ying et al., 2014). The contributions of source sectors to SOA were not tracked due to insufficient knowledge of its precursors and incomplete formation mechanisms (Yang et al., 2019; Carlton et al., 2007; Zhang et al., 2011).

2.3 Machine learning method

90 The lightGBM model was used to diagnose $\text{PM}_{2.5}$ simulation biases in this study (Ke et al., 2017). Our previous studies have shown its reliability for $\text{PM}_{2.5}$ modeling (Wang et al., 2023c; Wang et al., 2023b). The target variable was set to be the difference between observed and simulated daily $\text{PM}_{2.5}$ concentrations, and the key contributors to the simulation bias were then determined by the relative importance of the input features (Ye et al., 2022). LightGBM is an optimised gradient boosting decision tree (GBDT) that can handle non-linear relationships between multiple variables and has shown accurate performance
95 in many fields (Wei et al., 2021b; Yan et al., 2021; Sun et al., 2020; Liang et al., 2020b).

Three combinations of input variables were designed to fit the simulation biases: meteorological factors, chemical components, and emission sources. Meteorological factors, including wind speed, wind direction, temperature, humidity, surface pressure, cloud fraction and boundary layer height, are used to investigate the impact of meteorology on the CMAQ simulation biases. The components of $\text{PM}_{2.5}$ are divided into SIA (sulfate, nitrate, ammonium), primary/secondary organic
100 aerosols (POA/SOA), elemental carbon (EC), and other components. Seven sectoral sources (industry, energy, residential, transportation, agriculture, open burning, and biogenic emission) were used to quantify the contribution to the simulation bias. Five-fold cross-validation method was used to evaluate the model fit and prediction ability (Browne, 2000).

3 Results and discussion

3.1 Observation and simulation of $\text{PM}_{2.5}$

105 Figure 1a shows the time series of observed and simulated daily surface $\text{PM}_{2.5}$ concentrations in China and five regions (BTH, YRD, PRD, SCB, and NWCHN) over 2019. Basically, the CMAQ model successfully simulates the daily variation of $\text{PM}_{2.5}$ with some biases. $\text{PM}_{2.5}$ concentrations were higher in BTH ($51.172 \mu\text{g}/\text{m}^3$) and lower in PRD ($28.273 \mu\text{g}/\text{m}^3$). The CMAQ model underestimates $\text{PM}_{2.5}$ concentrations of $-8.59 \mu\text{g}/\text{m}^3$, $-2.66 \mu\text{g}/\text{m}^3$, $-6.21 \mu\text{g}/\text{m}^3$, and $-19.25 \mu\text{g}/\text{m}^3$ in the BTH, YRD, PRD, and NWCHN, respectively (Figure 1b). Moreover, the underestimation occurred mainly in winter and spring (Figure
110 1c), as well as high $\text{PM}_{2.5}$ events (Figure 1d) (Hu et al., 2016; Huang et al., 2017).

Table S2 shows the validation of CMAQ simulations against observations in different regions. Four indicators (MNB: mean normalized bias; MNE: mean normalized error; MFB: mean fractional bias; MFE: mean fractional error) were used to systematically evaluate the performance of the CMAQ simulations. The $\text{PM}_{2.5}$ simulations in the BTH, YRD, and PRD regions were in better agreement with observations, with average MNB of -0.08, -0.07, and -0.08 respectively (within the standard of
115 0.66). The $\text{PM}_{2.5}$ simulations in SCB and NWCHN regions show large biases with MNB of 0.46 and -0.42 respectively. The differences of CMAQ performance between regions can be attributed to multiple factors including emission inventories, dominant mechanisms of $\text{PM}_{2.5}$ generation, topographic, and meteorology conditions (Ma et al., 2021; Xue et al., 2019; Hu et al., 2014).

Annual and monthly mean $\text{PM}_{2.5}$ components (SIA, POA, SOA, EC, and other components) were calculated for China



120 and five key regions (Figure 2). $PM_{2.5}$ and its components show similar spatial distribution, with high concentrations occurring in the eastern regions (SCB, BTH, and central YRD). SOA showed high concentrations in summer over China ($6.80 \mu\text{g}/\text{m}^3$), which could be related to enhanced solar radiation and atmospheric oxidation capacity in summer (precursors of SOA such as isoprene are highly dependent on temperature and light) (Yang et al., 2019; Chen et al., 2020a; Liu et al., 2021b). Nitrate and POA were the dominate components in winter ($10.14 \mu\text{g}/\text{m}^3$ and $9.11 \mu\text{g}/\text{m}^3$ respectively). In BTH and SCB, POA accounts
135 for a higher proportion than nitrate in winter, whereas nitrate has a higher proportion in YRD. Nitrate showed higher concentration than sulfate in most regions and seasons due to the implementation of coal combustion control policies (Shang et al., 2021; Liu et al., 2021b; Xu et al., 2019).

The results of the $PM_{2.5}$ sectoral source appointment (Figure 3 and Figure S2) show that industries and residential sources were the main contributors to daily $PM_{2.5}$ concentrations for all regions and seasons, with seasonal fractional contributions of
130 25.31 - 31.92 % and 11.13 - 30.64 %, respectively). The seasonal average fractional contributions from energy, transportation, and agricultural NH_3 in whole China were 3.26 - 5.67%, 6.82 - 11.26 %, and 7.50 - 8.67 %, respectively. The contributions from biogenic source were negligible in all region and seasons (< 1 %). In contrast to the contributions from energy, transportation, industrial, and agricultural sources, significant seasonal variations occurred from residential source in all five regions, with high contribution in winter (17.60 - 30.90 %) and low contribution in summer (5.53 - 16.46 %).

135 As the secondary component makes up a large proportion of the total $PM_{2.5}$, the source sectors of SIA were analyzed for five regions (Figure S2). High concentrations of SIA were found in winter ($12.36 - 34.08 \mu\text{g}/\text{m}^3$), with large contribution from industrial, agricultural, and transportation sources (31.49 - 36.41 %, 20.40 - 22.40 %, and 19.77 - 22.46 %). The low contribution of residential sector to secondary $PM_{2.5}$ but high contribution to total $PM_{2.5}$ indicates that most residential emission sources emit PPM directly, with a small fraction of secondary generation. The contributions from biogenic and open burning
140 sector to SIA were relatively low in all region and seasons (< 10 %).

3.2 Drivers of $PM_{2.5}$ simulation bias

The contribution of each $PM_{2.5}$ components to the simulation bias was further quantified by relative importance. The ML models were trained separately for different regions and seasons, and a 5-fold cross-validation was used to measure the model performance (Figure 4). The models explain most part of simulation bias, with some differences across seasons and regions
145 (mean $CV R^2$ of 0.49, 0.65, 0.52, 0.49 for spring, summer, autumn, and winter).

Among the $PM_{2.5}$ components (Figure S4), SOA showed the largest contribution to $PM_{2.5}$ simulation bias for different regions and seasons (13.8 - 22.6%), which is consistent with previous studies (Liu et al., 2021b; Yang et al., 2019; Fry et al., 2014). The inorganic aerosols (e.g. sulphates) are produced mainly by chemical pathways, while SOA is produced by the condensation of numerous partially oxidized gases and is therefore influenced by complex precursor concentrations and multi-
150 stage oxidation processes. The incomplete description of SOA formation pathways in CTMs models (simplified SOA parameterization) leads to significant differences between simulations and observations (Carlton et al., 2007; Zhang et al., 2018; Yang et al., 2019). In addition, biogenic emissions play an important role in SOA formation, with biogenic SOA accounting for more than 70% of total SOA in China during summer (Hu et al., 2017b; Wu et al., 2020), so uncertainty in biogenic emissions can further contribute to uncertainty in SOA. Nitrate showed large contribution to $PM_{2.5}$ simulation bias in
155 winter at BTH, which is consistent with previous study.(Liu and Xing, 2022) Nitrate contribution to simulation bias further implies the inaccuracy of nitrate simulations, which can relate to the imperfect pathways of nitrate production (e.g., non-homogeneous oxidation) and the uncertainties of nitrate precursor emission inventories in winter (Xu et al., 2019; Zhang et al., 2018).

The main meteorological factors were used for modeling and were able to explain most simulation biases ($CV R^2$ of 0.62,
160 0.80, 0.69, 0.70 for spring, summer, autumn, and winter, Figure 4). The contribution of meteorological factors to the simulation bias varies across regions and seasons (Figure 5). In the BTH region, surface pressure and relative humidity contribute the



most to the simulation bias. In the PRD region, relative humidity, cloud cover, and wind direction were the main contributors in all four seasons.

Humidity positively or negatively influences $PM_{2.5}$ concentrations through several mechanisms. By enhancing $PM_{2.5}$ hygroscopic increase, promoting the secondary formation, and facilitating the gas-to-particle partitioning, high humidity positively influences $PM_{2.5}$ concentrations and contributes significantly to the haze pollution (Chen et al., 2020b; Cheng et al., 2015; Zhang et al., 2011). The contribution of humidity to CMAQ simulation biases can partly attributed to the uncertain of WRF simulation. The mean RMSE of relative humidity from WRF simulations versus observations was 20.38% in this study (Table S3). In addition, imperfections in the mechanism of humidity-promoted secondary particle formation (e.g., non-homogeneous catalysis of SOA) can also lead to simulation biases (Zhang et al., 2011; Liu et al., 2021b). Atmospheric pressure indirectly influences $PM_{2.5}$ concentrations through other meteorological factors (e.g., humidity, wind, etc.). High pressure systems are connected to the stationary weather, which is unfavorable for $PM_{2.5}$ dispersion. On the other hand, low pressure is usually accompanied by high humidity, influencing $PM_{2.5}$ nucleation, condensation and coagulation, leading to increased $PM_{2.5}$ concentrations (Chen et al., 2020b). Therefore, influence of atmospheric pressure on the CMAQ simulation biases in the BTH region maybe be attributed to the uncertain of meteorological field (Bei et al., 2017; Zhang et al., 2015). Contribution of wind direction in YRD may also related to the uncertain of WRF simulation (mean RMSE: 90.39 °). Aerosols have feedbacks on meteorology (Qu et al., 2021). In addition to directly changing the Earth's radiation receipts through scattering and absorbing (direct radiation effect), $PM_{2.5}$ can also influence radiation through aerosol-cloud interactions (indirect radiation effect) (Zhao et al., 2017; Yang et al., 2016). Moreover, $PM_{2.5}$ can act as cloud condensation and nucleation sites, contributing to clouds' microphysical development and precipitation formation process (Wang et al., 2020). However, the aerosol-to-meteorological feedback mechanism is missing in CMAQ used in this study. Previous study shown the dominant role of cloud chemistry in aerosol-cloud interactions in southern China (Zhao et al., 2017). Therefore, the influence of cloud cover on simulation biases in YRD can attributed to the missing of aerosol feedback mechanism.

In the NWCHN region, soil surface moisture and stomatal conductance contributed significantly to the simulation bias. These factors can associate with ground-level sand rise and dust emission (Liu et al., 2021c). Underestimation of dust aerosol in NWCHN can attributed to emission (both natural and anthropogenic sources), an accurate emission inventory (empirical- or physical-based numerical models) should be established in Northwest China by detailed activity data and emission factors (Hu et al., 2016; Liu et al., 2021a). In addition, the parameterization and mechanism for dust aerosol simulation in CMAQ should be further examined and improved.

Dry and wet days were divided to analyze influence of humidity on the simulation biases (Table S4). In most areas of China, CMAQ underestimates $PM_{2.5}$ more severely in dry days than in wet days, with larger absolute biases ($-14.56 \mu\text{g}/\text{m}^3$, $-7.09 \mu\text{g}/\text{m}^3$, $-7.11 \mu\text{g}/\text{m}^3$, and $-27.87 \mu\text{g}/\text{m}^3$ in spring, summer, autumn and winter respectively). In dry days, BTH showed severe underestimation in winter ($-22.86 \mu\text{g}/\text{m}^3$), while PRD showed large simulation bias in spring ($-21.55 \mu\text{g}/\text{m}^3$). Severe underestimation of $PM_{2.5}$ was observed in both wet and dry days at NWCHN. These underestimates of $PM_{2.5}$ in dry days can related to the dry deposition process. Dry deposition is a critical but highly uncertain sink for aerosols, which depends on the chemical and physical properties of aerosols, and influenced by subsurface and meteorological conditions (Shu et al., 2022). The CMAQ model in this study used the dry deposition scheme PR11 from Pleim and Ran (Pleim and Ran, 2011). This study shows that the PR11 scheme underestimates $PM_{2.5}$ concentrations in China. Recent studies for the United States also showed the underestimates for PM_{10} concentrations (Shu et al., 2022). Therefore, it is necessary to further develop and optimize the dry deposition scheme, especially for $PM_{2.5}$. $PM_{2.5}$ underestimation in wet days may attributed to the biases of wet deposition and secondary organic aerosol formation under high humidity conditions (Wu et al., 2018; Ryu and Min, 2022; Liu et al., 2021b; Zhang et al., 2011).

Source sectors contributions of PPM and SIA (obtained from the source-oriented CMAQ) were used to build the model and diagnose influences of different emissions sources on $PM_{2.5}$ simulation biases (Figure 6). Cross validation results of model



205 training by source sectors are shown in Figure 4. In China and in five key regions, sectoral sources were able to fit the simulation bias well, with mean R^2 and RMSE of 0.53 % and 20.18 $\mu\text{g}/\text{m}^3$. The PPM and SIA from residential showed the largest contribution (12.05 % and 12.78 %) to $\text{PM}_{2.5}$ simulation bias. The same conclusion was obtained when building model with total $\text{PM}_{2.5}$ concentrations from different source sectors (Table S5). $\text{PM}_{2.5}$ from residential emission is the main contributor to the CMAQ simulation bias, accounting for 20.2% of the total bias.

210 In China, residential sector consumed fossil fuels (coal, oil, and natural gas) and biofuels (wood and crop straw) with low combustion efficiency for cooking and heating and emitted large amounts of air pollutants (Li et al., 2017). However, due to the lack of reliable data (locally accurate emission factor and fuel consumption data), residential sector has been recognized as a major uncertainty source in anthropogenic emission inventories (Liu et al., 2021d; Shen et al., 2021), which is consistent with the results identified by machine learning in this study. Therefore, developing an accurate residential sector emissions
215 inventory is essential for accurate $\text{PM}_{2.5}$ modeling, which requires reliable data of fuel consumption and emission factors based on fuel type, fuel characteristics and combustion conditions (Liu et al., 2021d).

3.3 Uncertainties

Although we filtered the features according to their relative importance and priori knowledge, collinearity still exists among the input features. Features collinearity does not affect the performance of tree-based models like Random Forest and
220 LightGBM (Belgiu and Drăguț, 2016; Ke et al., 2017; Chen et al., 2016), but the contribution of a single feature might be influenced by other features, which is unfavorable for the interpretation of bias diagnosis results. In addition, the main objective of this study was diagnosing the contributors to CMAQ simulation biases using machine learning, therefore we did not pursue a very good model performance. The models showed good fitting ability with small performance gap between training and validation (Figure 4). Therefore, we consider such machine learning models to be reliable. Besides, the ML bias diagnosis
225 model constructed in this study is based entirely on local data and some regional processes influencing $\text{PM}_{2.5}$ concentrations are not considered in this study, such as vertical transport, long distance transport, which should be better diagnosed in future work, and the main bias contributors of identified by variable importance are in good agreement with the current findings.

4 Conclusion

Based on artificial intelligence technology, this study systematically diagnoses the possible drivers of biases in $\text{PM}_{2.5}$
230 simulation from three perspectives of meteorology, chemical components and emission sources. The relative importance of multiple factors helps to understand the sources of simulation bias and the deficiencies of the CMAQ mechanisms. SOA is the main contributor to simulation biases among chemical components. $\text{PM}_{2.5}$ is more underestimated in dry weather. Among source sectors, residential contributed the most simulation bias for both PPM and SIA. These results provide valuable information for CMAQ model improvement from SOA and dust aerosol underestimation, meteorological field uncertainties,
235 imperfection of the dry deposition scheme, and inaccurate residential emission inventories. As an efficient bias diagnosis method, machine learning based methods provide valuable complements to traditional mechanism-based methods. This approach also greatly reduces the prior information for diagnosing simulation bias and efficiently identifies the important contributors, so it can be easily extended to other CTMs models as well as other pollutants.



240 **Supporting**

Additional descriptions of the study domain, WRF-CMAQ simulation performance, concentrations and biases contribution of PM_{2.5} components and sectoral sources.

Code and data availability

The data and code (<https://zenodo.org/record/7907626>) are available on Zenodo (Wang et al., 2023a), including machine
245 learning code for diagnosing CMAQ simulation bias and the corresponding training dataset. CMAQ is an open-source chemical transport model developed by the US Environmental Protection Agency, which can be downloaded at <https://github.com/USEPA/CMAQ>.

Author contribution

Shuai Wang: Methodology, Software, Writing - original draft. **Mengyuan Zhang:** Software, Validation. **Yueqi Gao:** Data
250 curation, Visualization. **Peng Wang:** Methodology, Writing - reviewing and editing. **Qingyan Fu:** Writing - reviewing and editing. **Hongliang Zhang:** Conceptualization, Supervision, Writing - reviewing and editing.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

255



Reference


- Aleksankina, K., Reis, S., Vieno, M., and Heal, M. R.: Advanced methods for uncertainty assessment and global sensitivity analysis of an Eulerian atmospheric chemistry transport model, *Atmos. Chem. Phys.*, 19, 2881-2898, 2019.
- 260 Bai, K., Li, K., Ma, M., Li, K., Li, Z., Guo, J., Chang, N. B., Tan, Z., and Han, D.: LGHAP: the Long-term Gap-free High-resolution Air Pollutant concentration dataset, derived via tensor-flow-based multimodal data fusion, *Earth Syst. Sci. Data*, 14, 907-927, 10.5194/essd-14-907-2022, 2022.
- Beekmann, M. and Derognat, C.: Monte Carlo uncertainty analysis of a regional-scale transport chemistry model constrained by measurements from the atmospheric pollution over the Paris area (ESQUIF) campaign, *J. Geophys. Res.-Atmos.*, 108, 2003.
- 265 Bei, N., Wu, J., Elser, M., Feng, T., Cao, J., El-Haddad, I., Li, X., Huang, R., Li, Z., Long, X., Xing, L., Zhao, S., Tie, X., Prévôt, A. S. H., and Li, G.: Impacts of meteorological uncertainties on the haze formation in Beijing–Tianjin–Hebei (BTH) during wintertime: a case study, *Atmos. Chem. Phys.*, 17, 14579-14591, 10.5194/acp-17-14579-2017, 2017.
- Belgiu, M. and Drăguț, L.: Random forest in remote sensing: A review of applications and future directions, *ISPRS-J. Photogramm. Remote Sens.*, 114, 24-31, 2016.
- Binkowski, F. S. and Roselle, S. J.: Models-3 Community Multiscale Air Quality (CMAQ) model aerosol component 1. Model description, *J. Geophys. Res.-Atmos.*, 108, 2003.
- 270 Brown, M. W.: Cross-validation methods, *Journal of mathematical psychology*, 44, 108-132, 2000.
- Carlton, A. G., Turpin, B. J., Altieri, K. E., Seitzinger, S., Reff, A., Lim, H.-J., and Ervens, B.: Atmospheric oxalic acid and SOA production from glyoxal: Results of aqueous photooxidation experiments, *Atmos. Environ.*, 41, 7588-7602, <https://doi.org/10.1016/j.atmosenv.2007.05.035>, 2007.
- 275 Carter, W. P. and Heo, G.: Development of revised SAPRC aromatics mechanisms, *Atmos. Environ.*, 77, 404-414, 2013.
- Chen, S., Wang, H., Lu, K., Zeng, L., Hu, M., and Zhang, Y.: The trend of surface ozone in Beijing from 2013 to 2019: Indications of the persisting strong atmospheric oxidation capacity, *Atmos. Environ.*, 242, 117801, 2020a.
- Chen, T. Q., Guestrin, C., and Assoc Comp, M.: XGBoost: A Scalable Tree Boosting System, 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining (KDD), San Francisco, CA, Aug 13-17, WOS:000485529800092, 785-794, 10.1145/2939672.2939785, 2016.
- 280 Chen, Z. Y., Chen, D. L., Zhao, C. F., Kwan, M. P., Cai, J., Zhuang, Y., Zhao, B., Wang, X. Y., Chen, B., Yang, J., Li, R. Y., He, B., Gao, B. B., Wang, K. C., and Xu, B.: Influence of meteorological conditions on PM_{2.5} concentrations across China: A review of methodology and mechanism, *Environ. Int.*, 139, 10.1016/j.envint.2020.105558, 2020b.
- Cheng, Y., He, K. B., Du, Z. Y., Zheng, M., Duan, F. K., and Ma, Y. L.: Humidity plays an important role in the PM_{2.5} pollution in Beijing, *Environ. Pollut.*, 197, 68-75, 10.1016/j.envpol.2014.11.028, 2015.
- 285 Commerce, N. C. f. E. P. N. W. S. N. U. D. o.: NCEP FNL operational model global tropospheric analyses, continuing from July 1999, Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, 2000.
- Fry, J. L., Draper, D. C., Barsanti, K. C., Smith, J. N., Ortega, J., Winkler, P. M., Lawler, M. J., Brown, S. S., Edwards, P. M., 290 Cohen, R. C., and Lee, L.: Secondary Organic Aerosol Formation and Organic Nitrate Yield from NO₃ Oxidation of Biogenic Hydrocarbons, *Environ. Sci. Technol.*, 48, 11944-11953, 10.1021/es502204x, 2014.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, 2006.
- 295 Guenther, A., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Emmons, L., and Wang, X.: The Model of Emissions of Gases and Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated framework for modeling biogenic emissions, *Geosci. Model Dev.*, 5, 1471-1492, 2012.
- Hanna, S., Russell, A., Wilkinson, J., Vukovich, J., and Hansen, D.: Monte Carlo estimation of uncertainties in BEIS3 emission outputs and their effects on uncertainties in chemical transport model predictions, *J. Geophys. Res.-Atmos.*, 110, 2005.
- Hou, L. L., Dai, Q. L., Song, C. B., Liu, B. W., Guo, F. Z., Dai, T. J., Li, L. X., Liu, B. S., Bi, X. H., Zhang, Y. F., and Feng, 300 Y. C.: Revealing Drivers of Haze Pollution by Explainable Machine Learning, *Environmental Science & Technology Letters*, 9, 112-119, 10.1021/acs.estlett.1c00865, 2022.
- Hu, J., Chen, J., Ying, Q., and Zhang, H.: One-year simulation of ozone and particulate matter in China using WRF/CMAQ modeling system, *Atmos. Chem. Phys.*, 16, 10333-10350, 2016.
- Hu, J., Wang, Y., Ying, Q., and Zhang, H.: Spatial and temporal variability of PM_{2.5} and PM₁₀ over the North China Plain and the Yangtze River Delta, China, *Atmos. Environ.*, 95, 598-609, 2014.
- 305 Hu, J., Huang, L., Chen, M., Liao, H., Zhang, H., Wang, S., Zhang, Q., and Ying, Q.: Premature mortality attributable to particulate matter in China: source contributions and responses to reductions, *Environ. Sci. Technol.*, 51, 9950-9959, 2017a.
- Hu, J., Wang, P., Ying, Q., Zhang, H., Chen, J., Ge, X., Li, X., Jiang, J., Wang, S., and Zhang, J.: Modeling biogenic and anthropogenic secondary organic aerosol in China, *Atmos. Chem. Phys.*, 17, 77-92, 2017b.
- 310 Huang, Z., Hu, Y., Zheng, J., Yuan, Z., Russell, A. G., Ou, J., and Zhong, Z.: A New Combined Stepwise-Based High-Order Decoupled Direct and Reduced-Form Method To Improve Uncertainty Analysis in PM(2.5) Simulations, *Environ Sci Technol*, 51, 3852-3859, 10.1021/acs.est.6b05479, 2017.
- Ke, G. L., Meng, Q., Finley, T., Wang, T. F., Chen, W., Ma, W. D., Ye, Q. W., and Liu, T. Y.: LightGBM: A Highly Efficient Gradient Boosting Decision Tree, 31st Annual Conference on Neural Information Processing Systems (NIPS), Long Beach, CA, Dec 04-09, WOS:000452649403021, 2017.
- 315 Li, M., Liu, H., Geng, G., Hong, C., Liu, F., Song, Y., Tong, D., Zheng, B., Cui, H., Man, H., Zhang, Q., and He, K.: Anthropogenic emission inventories in China: a review, *National Science Review*, 4, 834-866, 10.1093/nsr/nwx150, 2017.
- Li, T., Zhang, Q., Peng, Y., Guan, X., Li, L., Mu, J., Wang, X., Yin, X., and Wang, Q.: Contributions of Various Driving



- Factors to Air Pollution Events: Interpretability Analysis from Machine Learning Perspective, *Environ. Int.*, 107861, 2023.
- 320 Liang, F., Xiao, Q., Huang, K., Yang, X., Liu, F., Li, J., Lu, X., Liu, Y., and Gu, D.: The 17-y spatiotemporal trend of PM_{2.5} and its mortality burden in China, *Proc. Natl. Acad. Sci. U. S. A.*, 117, 25601-25608, 2020a.
- Liang, W., Luo, S., Zhao, G., and Wu, H.: Predicting hard rock pillar stability using GBDT, XGBoost, and LightGBM algorithms, *Mathematics*, 8, 765, 2020b.
- Liu, J. and Xing, J.: Identifying Contributors to PM_{2.5} Simulation Biases of Chemical Transport Model Using Fully Connected Neural Networks, *Journal of Advances in Modeling Earth Systems*, 15, 2022.
- 325 Liu, J., Ding, J., Rexiding, M., Li, X., Zhang, J., Ran, S., Bao, Q., and Ge, X.: Characteristics of dust aerosols and identification of dust sources in Xinjiang, China, *Atmos. Environ.*, 262, 118651, <https://doi.org/10.1016/j.atmosenv.2021.118651>, 2021a.
- Liu, J., Chu, B., Chen, T., Zhong, C., Liu, C., Ma, Q., Ma, J., Zhang, P., and He, H.: Secondary organic aerosol formation potential from ambient air in Beijing: effects of atmospheric oxidation capacity at different pollution levels, *Environ. Sci. Technol.*, 55, 4565-4572, 2021b.
- 330 Liu, S., Xing, J., Sahu, S. K., Liu, X., Liu, S., Jiang, Y., Zhang, H., Li, S., Ding, D., Chang, X., and Wang, S.: Wind-blown dust and its impacts on particulate matter pollution in Northern China: current and future scenarios, *Environ. Res. Lett.*, 16, 114041, [10.1088/1748-9326/ac31ec](https://doi.org/10.1088/1748-9326/ac31ec), 2021c.
- Liu, X., Lu, D., Zhang, A., Liu, Q., and Jiang, G.: Data-driven machine learning in environmental pollution: Gains and problems, *Environ. Sci. Technol.*, 56, 2124-2133, 2022.
- 335 Liu, X., Shen, G., Chen, L., Qian, Z., Zhang, N., Chen, Y., Chen, Y., Cao, J., Cheng, H., Du, W., Li, B., Li, G., Li, Y., Liang, X., Liu, M., Lu, H., Luo, Z., Ren, Y., Zhang, Y., Zhu, D., and Tao, S.: Spatially Resolved Emission Factors to Reduce Uncertainties in Air Pollutant Emission Estimates from the Residential Sector, *Environ. Sci. Technol.*, 55, 4483-4493, [10.1021/acs.est.0c08568](https://doi.org/10.1021/acs.est.0c08568), 2021d.
- 340 Ma, J., Shen, J., Wang, P., Zhu, S., Wang, Y., Wang, P., Wang, G., Chen, J., and Zhang, H.: Modeled changes in source contributions of particulate matter during the COVID-19 pandemic in the Yangtze River Delta, China, *Atmos. Chem. Phys.*, 21, 7343-7355, 2021.
- Meng, C., Cheng, T. H., Gu, X. F., Shi, S. Y., Wang, W. N., Wu, Y., and Bao, F. W.: Contribution of meteorological factors to particulate pollution during winters in Beijing, *Science of the Total Environment*, 656, 977-985, [10.1016/j.scitotenv.2018.11.365](https://doi.org/10.1016/j.scitotenv.2018.11.365), 2019.
- Organization, W. H.: WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide: executive summary, 2021.
- Pleim, J. and Ran, L.: Surface flux modeling for air quality applications, *Atmosphere*, 2, 271-302, 2011.
- 350 Qiao, X., Ying, Q., Li, X., Zhang, H., Hu, J., Tang, Y., and Chen, X.: Source apportionment of PM_{2.5} for 25 Chinese provincial capitals and municipalities using a source-oriented Community Multiscale Air Quality model, *Science of the Total Environment*, 612, 462-471, [10.1016/j.scitotenv.2017.08.272](https://doi.org/10.1016/j.scitotenv.2017.08.272), 2018.
- Qu, Y., Voulgarakis, A., Wang, T., Kasoar, M., Wells, C., Yuan, C., Varma, S., and Mansfield, L.: A study of the effect of aerosols on surface ozone through meteorology feedbacks over China, *Atmos. Chem. Phys.*, 21, 5705-5718, [10.5194/acp-21-5705-2021](https://doi.org/10.5194/acp-21-5705-2021), 2021.
- 355 Ryu, Y.-H. and Min, S.-K.: Improving Wet and Dry Deposition of Aerosols in WRF-Chem: Updates to Below-Cloud Scavenging and Coarse-Particle Dry Deposition, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002792, <https://doi.org/10.1029/2021MS002792>, 2022.
- Shang, D., Peng, J., Guo, S., Wu, Z., and Hu, M.: Secondary aerosol formation in winter haze over the Beijing-Tianjin-Hebei Region, China, *Frontiers of Environmental Science & Engineering*, 15, 1-13, 2021.
- 360 Shen, H., Luo, Z., Xiong, R., Liu, X., Zhang, L., Li, Y., Du, W., Chen, Y., Cheng, H., Shen, G., and Tao, S.: A critical review of pollutant emission factors from fuel combustion in home stoves, *Environ. Int.*, 157, 106841, <https://doi.org/10.1016/j.envint.2021.106841>, 2021.
- Shu, Q., Murphy, B., Schwede, D., Henderson, B. H., Pye, H. O. T., Appel, K. W., Khan, T. R., and Perlinger, J. A.: Improving the particle dry deposition scheme in the CMAQ photochemical modeling system, *Atmos. Environ.*, 289, 119343, <https://doi.org/10.1016/j.atmosenv.2022.119343>, 2022.
- 365 Stirnberg, R., Cermak, J., Kotthaus, S., Haeffelin, M., Andersen, H., Fuchs, J., Kim, M., Petit, J. E., and Favez, O.: Meteorology-driven variability of air pollution (PM₁) revealed with explainable machine learning, *Atmos. Chem. Phys.*, 21, 3919-3948, [10.5194/acp-21-3919-2021](https://doi.org/10.5194/acp-21-3919-2021), 2021.
- Sun, H., Shin, Y. M., Xia, M., Ke, S., Wan, M., Yuan, L., Guo, Y., and Archibald, A. T.: Spatial Resolved Surface Ozone with Urban and Rural Differentiation during 1990–2019: A Space–Time Bayesian Neural Network Downscaler, *Environ. Sci. Technol.*, 56, 7337-7349, 2021.
- Sun, X., Liu, M., and Sima, Z.: A novel cryptocurrency price trend forecasting model based on LightGBM, *Finance Research Letters*, 32, 101084, 2020.
- 375 Wang, P., Qiao, X., and Zhang, H.: Modeling PM_{2.5} and O₃ with aerosol feedbacks using WRF/Chem over the Sichuan Basin, southwestern China, *Chemosphere*, 254, 126735, <https://doi.org/10.1016/j.chemosphere.2020.126735>, 2020.
- Wang, S., Zhang, M., Gao, Y., Wang, P., and Zhang, H.: Diagnosing drivers of PM_{2.5} simulation biases from meteorology, chemical composition, and emission sources using an efficient machine learning method, *Zenodo [code]*, <https://doi.org/10.5281/zenodo.7907626>, 2023a.
- 380 Wang, S., Wang, P., Zhang, R., Meng, X., Kan, H., and Zhang, H.: Estimating particulate matter concentrations and meteorological contributions in China during 2000–2020, *Chemosphere*, 330, 138742, <https://doi.org/10.1016/j.chemosphere.2023.138742>, 2023b.
- Wang, S., Wang, P., Qi, Q., Wang, S., Meng, X., Kan, H., Zhu, S., and Zhang, H.: Improved estimation of particulate matter



- in China based on multisource data fusion, *Science of The Total Environment*, 161552, 2023c.
- 385 Wang, S. Y., Zhang, Y. L., Ma, J. L., Zhu, S. Q., Shen, J. Y., Wang, P., and Zhang, H. L.: Responses of decline in air pollution and recovery associated with COVID-19 lockdown in the Pearl River Delta, *Science of the Total Environment*, 756, 10.1016/j.scitotenv.2020.143868, 2021.
- Wei, J., Li, Z., Lyapustin, A., Sun, L., Peng, Y., Xue, W., Su, T., and Cribb, M.: Reconstructing 1-km-resolution high-quality PM_{2.5} data records from 2000 to 2018 in China: spatiotemporal variations and policy implications, *Remote Sensing of Environment*, 252, 112136, 2021a.
- 390 Wei, J., Li, Z., Pinker, R. T., Wang, J., Sun, L., Xue, W., Li, R., and Cribb, M.: Himawari-8-derived diurnal variations in ground-level PM_{2.5} pollution across China using the fast space-time Light Gradient Boosting Machine (LightGBM), *Atmos. Chem. Phys.*, 21, 7863-7880, 10.5194/acp-21-7863-2021, 2021b.
- Wei, J., Li, Z. Q., Cribb, M., Huang, W., Xue, W. H., Sun, L., Guo, J. P., Peng, Y. R., Li, J., Lyapustin, A., Liu, L., Wu, H., and Song, Y. M.: Improved 1 km resolution PM_{2.5} estimates across China using enhanced space-time extremely randomized trees, *Atmos. Chem. Phys.*, 20, 3273-3289, 10.5194/acp-20-3273-2020, 2020.
- 395 Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625-641, 10.5194/gmd-4-625-2011, 2011.
- Wu, K., Yang, X. Y., Chen, D., Gu, S., Lu, Y. Q., Jiang, Q., Wang, K., Ou, Y. H., Qian, Y., Shao, P., and Lu, S. H.: Estimation of biogenic VOC emissions and their corresponding impact on ozone and secondary organic aerosol formation in China, *Atmospheric Research*, 231, 10.1016/j.atmosres.2019.104656, 2020.
- 400 Wu, Y., Liu, J., Zhai, J., Cong, L., Wang, Y., Ma, W., Zhang, Z., and Li, C.: Comparison of dry and wet deposition of particulate matter in near-surface waters during summer, *PloS one*, 13, e0199241, 2018.
- Xiao, Q., Geng, G., Xue, T., Liu, S., Cai, C., He, K., and Zhang, Q.: Tracking PM_{2.5} and O₃ Pollution and the Related Health Burden in China 2013–2020, 2021a.
- 405 Xiao, Q., Zheng, Y., Geng, G., Chen, C., Huang, X., Che, H., Zhang, X., He, K., and Zhang, Q.: Separating emission and meteorological contributions to long-term PM_{2.5} trends over eastern China during 2000–2018, *Atmos. Chem. Phys.*, 21, 9475-9496, 10.5194/acp-21-9475-2021, 2021b.
- Xu, Q., Wang, S., Jiang, J., Bhattarai, N., Li, X., Chang, X., Qiu, X., Zheng, M., Hua, Y., and Hao, J.: Nitrate dominates the chemical composition of PM_{2.5} during haze event in Beijing, China, *Science of the Total Environment*, 689, 1293-1303, 2019.
- 410 Xue, T., Zheng, Y., Tong, D., Zheng, B., Li, X., Zhu, T., and Zhang, Q.: Spatiotemporal continuous estimates of PM_{2.5} concentrations in China, 2000–2016: A machine learning method with inputs from satellites, chemical transport model, and ground observations, *Environ. Int.*, 123, 345-357, 2019.
- Yan, J., Xu, Y., Cheng, Q., Jiang, S., Wang, Q., Xiao, Y., Ma, C., Yan, J., and Wang, X.: LightGBM: accelerated genomically designed crop breeding through ensemble learning, *Genome Biology*, 22, 1-24, 2021.
- 415 Yang, W., Li, J., Wang, W., Li, J., Ge, M., Sun, Y., Chen, X., Ge, B., Tong, S., Wang, Q., and Wang, Z.: Investigating secondary organic aerosol formation pathways in China during 2014, *Atmos. Environ.*, 213, 133-147, <https://doi.org/10.1016/j.atmosenv.2019.05.057>, 2019.
- Yang, X., Zhao, C. F., Guo, J. P., and Wang, Y.: Intensification of aerosol pollution associated with its feedback with surface solar radiation and winds in Beijing, *J. Geophys. Res.-Atmos.*, 121, 4093-4099, 10.1002/2015jd024645, 2016.
- 420 Ye, X., Wang, X., and Zhang, L.: Diagnosing the Model Bias in Simulating Daily Surface Ozone Variability Using a Machine Learning Method: The Effects of Dry Deposition and Cloud Optical Depth, *Environ. Sci. Technol.*, 56, 16665-16675, 10.1021/acs.est.2c05712, 2022.
- Ying, Q., Li, J., and Kota, S. H.: Significant contributions of isoprene to summertime secondary organic aerosol in eastern United States, *Environ. Sci. Technol.*, 49, 7834-7842, 2015.
- 425 Ying, Q., Wu, L., and Zhang, H.: Local and inter-regional contributions to PM_{2.5} nitrate and sulfate in China, *Atmos. Environ.*, 94, 582-592, 2014.
- Zhai, S., Jacob, D. J., Wang, X., Shen, L., Li, K., Zhang, Y., Gui, K., Zhao, T., and Liao, H.: Fine particulate matter (PM_{2.5}) trends in China, 2013–2018: Separating contributions from anthropogenic emissions and meteorology, *Atmos. Chem. Phys.*, 19, 11031-11041, 2019.
- 430 Zhang, H., Surratt, J. D., Lin, Y. H., Bapat, J., and Kamens, R. M.: Effect of relative humidity on SOA formation from isoprene/NO photooxidation: enhancement of 2-methylglyceric acid and its corresponding oligoesters under dry conditions, *Atmos. Chem. Phys.*, 11, 6411-6424, 10.5194/acp-11-6411-2011, 2011.
- Zhang, H., Li, J., Ying, Q., Yu, J. Z., Wu, D., Cheng, Y., He, K., and Jiang, J.: Source apportionment of PM_{2.5} nitrate and sulfate in China using a source-oriented chemical transport model, *Atmos. Environ.*, 62, 228-242, 2012.
- 435 Zhang, Q., Quan, J., Tie, X., Li, X., Liu, Q., Gao, Y., and Zhao, D.: Effects of meteorology and secondary particle formation on visibility during heavy haze events in Beijing, China, *Science of the Total Environment*, 502, 578-584, 2015.
- Zhang, R., Sun, X. S., Shi, A. J., Huang, Y. H., Yan, J., Nie, T., Yan, X., and Li, X.: Secondary inorganic aerosols formation during haze episodes at an urban site in Beijing, China, *Atmos. Environ.*, 177, 275-282, 10.1016/j.atmosenv.2017.12.031, 2018.
- 440 Zhao, B., Liou, K.-N., Gu, Y., Li, Q., Jiang, J. H., Su, H., He, C., Tseng, H.-L. R., Wang, S., Liu, R., Qi, L., Lee, W.-L., and Hao, J.: Enhanced PM_{2.5} pollution in China due to aerosol-cloud interactions, *Sci Rep*, 7, 4453, 10.1038/s41598-017-04096-8, 2017.
- Zhu, Q., Bi, J., Liu, X., Li, S., Wang, W., Zhao, Y., and Liu, Y.: Satellite-based long-term spatiotemporal patterns of surface ozone concentrations in China: 2005–2019, *Environ. Health Perspect.*, 130, 027004, 2022.



450 Figure 1. (a) The time series of observed (black) and CMAQ simulated (red) daily surface PM_{2.5} concentrations in China and five
 455 regions. Mean concentrations of the observed and simulated PM_{2.5} and MNB also shown in the inset. (b) Box plots of CMAQ
 simulated biases (simulated minus observed) for different regions. Crosses indicate average values and outliers are determined to
 be > 1.5 times of the upper quartile and < 1.5 times of the lower quartile. (c) Same as (b) but for four seasons. Spring, summer,
 autumn and winter are defined as March to May, June to August, September to November, December January and February,
 respectively. (d) Same as (b) but for different PM_{2.5} concentration levels (L1: [0, 35], L2: [35, 75], L3: [75, 115], L4: [115, 150], L5:
 [150, 1000]).




Figure 2. Annual mean concentrations map (a - g) and monthly mean concentrations (h-m) of $PM_{2.5}$ and its components (SIA, POA, SOA, EC, and other components) for China and five key regions in 2019. Dotted lines in h-m indicate $PM_{2.5}$ observations

460





Figure 3. Seasonal average fractional contributions from different sources to $PM_{2.5}$ concentrations (white circle on the right-hand axis) in China and five regions.



465

Figure 4. Cross validation (5-fold) results of CMAQ bias model training by meteorology (a), $PM_{2.5}$ components (b), and source sectors (c). RSME unit: $\mu g/m^3$.



470 Figure 5. Contribution (%) of each meteorological factor to CMAQ simulation biases by region and season.




Figure 6. Contribution (%) of each source sectors to CMAQ biases by region and season. res: residential, ene: energy, tra: transportation, agr: agriculture, ind: industry, AEC: elemental carbon, Other: other components.