

## **Author's reply to referee comments**

Miettunen, E., Tuomi, L., Westerlund, A., Kanarik, H., and Myrberg, K.: Transport dynamics in a complex coastal archipelago, EGUsphere [preprint], <https://doi.org/10.5194/egusphere-2023-1547>, 2023.

Below, referee comments are displayed with italic font, highlighted with a grey background. Author replies are without highlighting. When listing the changes in the revised manuscript, line numbers refer to the "Author's tracked changes" file.

### **RC1, Anonymous Referee #1, 30 Aug 2023**

*> Review of MS by Elina Miettunen et al.*

*> Transport dynamics in a complex coastal archipelago*

*> The MS is aimed to study the currents and volume transports of water in the Archipelago Sea using results from the high-resolution (0.25 nautical miles) 3D NEMO ocean model. From the same model experiments, results about the neighboring Åland Sea have been published by Westerlund et al. (2022). Model results in the Archipelago Sea are validated with acceptable results by the observations of temperature, salinity and currents using the data from the archives. Results part of the MS present (chapter 4) statistics of modelled currents in relation to winds, using mainly directional "roses", and (chapter 5) time series of monthly mean volume transports across the selected 3 west-east transects. In particular, the study reveals that currents are steered by the geometry of the islands and straits and the bottom topography. Net transport in the upper 20 m layer was southward. Monthly volume transport had maximum southward direction in spring and northward direction in autumn and winter. Thus, the study and its results are generally interesting and could be published.*

*> In the following, I elaborate the background in order to give recommendations how the MS might be made more significant and interesting.*

We would like to thank the referee for taking the time to review the manuscript and providing recommendations on how to make it more significant and interesting. We greatly appreciate the feedback and think that the manuscript is now improved after addressing these comments. Please find our replies to the comments below.

*> A. The modelling results of the Archipelago Sea with a grid step of 0.25 nautical miles have been published earlier by Tuomi et al. (2018) and Miettunen et al. (2020). They used the 3D COHERENS model. Present MS should also reference to the earlier model. It should be interesting to know is there an improvement from COHERENS to NEMO. Oceanographic results of the two earlier studies are referenced in the Introduction. In my understanding, the oceanographic results of present MS does not go much beyond these earlier studies. Perhaps this feeling is subjective and fed by massive use of the term "complex" (15 cases vs 2 and 6 cases in earlier studies). Authors are encouraged to look how to include more oceanography and reduce information-poor terms/formulations like "complex" and "resolution" (26 cases).*

Thank you for this comment. We agree and have added reference to the earlier COHERENS-based model setup to the introduction (lines 74–76 in the "Author's tracked changes" file) and discussion (lines 327–329) sections.

Thank you for pointing out our excessive use of the term “complex”. In the revised manuscript, we use more diverse language and more descriptive formulations (e.g., lines 36–37, 316–317). Regarding the resolution, we have included definitions of what we mean when talking about “high” and “coarse” resolution models (lines 45–48).

*> B. The study reveals steering of currents along closely spaced isobaths in straits, channels and trenches. This issue of topographically constrained currents is generally known and could be more presented and discussed, including more references to the theoretical studies and observations in nearby Baltic regions. Is the role of islands to guide the flow, without significant frictional slow-down?*

Topographically restricted currents are quite common in the Baltic Sea due to its varying topography, elongated gulfs and archipelagos. We think that theoretical discussion on this is out of scope of this manuscript. However, we have added a bit more information about this to the introduction, concentrating on the specific features of the Archipelago Sea where the flow is guided both by the islands and bathymetry (lines 26–33).

*> C. Complementary to the current roses in Figs. 4-5, it should be interesting to see (seasonally?) mean current maps (perhaps together with persistency contours).*

When studying circulation in the model region, we have analysed the monthly and seasonal currents by drawing both mean current maps and current roses. The bi-modal nature of the currents, especially in the northern and central parts of the area, makes the current means misleading. While for some periods, the mean current field can actually represent the most dominant current direction, in some cases the mean does not show either of the dominant directions. This is especially the case for autumn: the mean current field calculated from the vector components shows eastward currents also in the northern part of the region where the roses show clearly that northward and southward directions dominate (see the figure below). That is why we have decided to present the circulation with current roses only.




Figure: Current roses for selected grid points (left) and mean current field (right) for the 5 m surface layer in autumn (Oct–Dec), 2013–2017.

> D. Section of volume transports is interesting, but more information on water budget, transect areas and forcing factors could be presented. (a) Time series of monthly mean transports as shown in Fig. 7 are similar on surface and intermediated layers. Nevertheless, they are also similar on different transects, with correlation above 0.9 (I made this check). This indicates large-scale forcing of volume transport. (b) Forcing of volume transports has been discussed but not evaluated. Ambjörn and Gidhagen (1979) have concluded: "Main driving force on the net current, when the vertical stratification is weak, is the surface slope along the channel. Local acceleration and bottom friction are also important." This can be directly evaluated from the monthly mean model results. It should also be interesting to know what wind stress projections (to what angle) cause sea level slopes across the Archipelago Sea favoring northward or southward flows. For example, SE and NNW winds in 2014 created larger volume transports than in other years; was it related to the larger sea level slopes?

(a) It is true that the surface layer transports are similar at all the three transects. However, the lower layer transports are different at different transects. This was not perhaps clearly visible in the original figures. In response to the 12th minor comment about duplication of data in Figs. 7 and 8, we have replaced those figures in the revised manuscript. The new ones show the differences between the upper and lower layer at the different transects more clearly.

(b) This is the first time that this kind of model analysis of volume transports is conducted for the Archipelago Sea. We chose the same method used in Westerlund et al. (2022) in the Åland Sea so that we can compare the results for these two regions. We agree that the water budget and other forcing factors are important to analyse to better understand the dynamics of these regions and the water exchange between the different basins. However, as both the Åland Sea and the Archipelago sea act as a pathway for the water exchange between the Baltic proper and the Bothnian Sea, the analysis of the forcing should include both areas. This kind of analysis is out of scope of this manuscript and will be continued in the future. In our future work, we aim to use longer model simulations, preferably using a two-way nested model setup, and also include measurements conducted in the region during the past years.

> E. The MS emphasizes further need to increase the resolution. How many details are reasonable? Thousands of islands are making already some statistical entity. For example, flows in the porous media (e.g. Pratt, L.J. and Spall, M.A., 2003. A porous-medium theory for barotropic flow through ridges and archipelagos. *Journal of physical oceanography*, 33(12), pp.2702-2718.) can be modelled without counting each individual grain and/or pore. (Consider also Darcy law).

We think that the current model resolution of 0.25 nautical miles is enough for our studies. In general, the resolution is enough to describe the main waterways in sufficient detail. However, some of the channels are quite narrow and oriented diagonally with respect to the model grid. Higher grid resolution could improve the description of those channels and their orientation and thus improve especially the estimates for the lower layer transports.

Impression that further increase in resolution would be needed for this study was not intended. We have rephrased the parts that might have given this impression to avoid misinterpretation (e.g., lines 324, 338).

*> F. The title of MS is too general for the present content. Archipelago dynamics in general oceanographic sense is not presented and discussed. References to the other archipelago sea studies focus mainly on technical details, such as model setup, need for higher model resolution and more dense monitoring network. I recommend to rephrase the title.*

We chose to use a more generic title for this manuscript, since although these types of archipelago areas are rare, they still exist also outside our study area. We also think that our study and the methods we use can be of interest also to researchers focusing on other coastal archipelago regions in and outside the Baltic Sea.

*> I include also some minor remarks.*

*> 1) The term “high-resolution” (counted 11 times) could be specified.*

It is true that this term is a bit vague and it depends on the modelled region whether a model resolution can be considered high or not. As mentioned in our reply to the comment A, we have now specified in the revised manuscript what we mean when talking about high and coarse resolution models (in the introduction, lines 45–48).

*> 2) The term “area” is used as a synonym for “region”. It could be useful to present and discuss actual geometrical areas of the transects, hypsographic curves of the regions etc.*

We included plots of the topography along the transects in Fig. 1. These show how the thickness of the upper and lower layers vary along the transects and how the lower layer is mostly present only in the channels crossing the area.

*> 3) Line 4: It has to be specified what NEMO is, even in the abstract (an oceanographic model?)*

This is now specified in the abstract (lines 4–5) and in the introduction (line 71).

*> 4) Lines 40-44 say that “situations where substances are transported through the Archipelago Sea occur rarely” and “there is constant exchange of water”. How water exchange occurs without transport of substances? The role of salt exchange is not figured out, although there should be long-term salt flux based on the Knudsen formulae.*

These sentences were perhaps not clear in the original manuscript. What we mean is that there is constant water exchange between the outer parts of the northern and southern Archipelago Sea with their neighbouring basins (Bothnian Sea in the north and Baltic Proper in the south). However, there is not constant water exchange between the Baltic Proper and the Bothnian Sea through the Archipelago Sea because the exchange is limited by the archipelago. We have now rephrased this part in the revised manuscript so that it is more clear (lines 50–57).

*> 5) Lines 78-79: open boundary data were taken from the Baltic Sea Physical Reanalysis Product. This data set has daily mean values for currents, temperature and salinity. How the boundary values with periods shorter than a day were taken into account? The reader could be interested to understand the main features without reading Westerlund et al. (2022).*

Currents, temperature and salinity in the open boundary data are indeed daily mean values, but sea surface heights are hourly values. This was not mentioned in the model description in the original manuscript but it is now included in the revised version (lines 100–101). Thank you for pointing this out.

> 6) Line 87: *“temperature, salinity and currents are saved as 6 h averages” is nearly able to cover the daily cycle. How shorter period processes like 14-hour inertial oscillations, sea level variations can be taken into account? // Considered in the discussion, lines 273-276 but could be brought in earlier, in the methods.*

The model results are indeed saved as 6 h averages, but the model of course accounts for short-term variability. As we analyse the transports as monthly sums and net values, we see that the output frequency is enough for this study.

Modelled sea level was saved at 1 h intervals and validated in our earlier paper (Westerlund et al., 2022) using tide gauge data from Föglö station which is located in the western Archipelago Sea. This validation was not clearly referenced in the original manuscript. In the revised manuscript, we now summarise the sea level validation done in Westerlund et al. (2022) at the beginning of Section 3 (lines 140–143).

> 7) Lines 128-129: *“The model grid is too shallow to reproduce halocline in this area. However, this does not affect our study of currents and transports, as we focus on the shallower archipelago areas with no halocline.” It should be better justified. For example, presenting the fraction of halocline-covered area to the area of whole transect.*

Halocline is seen only at the southern edge of the Archipelago Sea where the IU7 measurement station is located. The other IU stations north of this show no halocline, not even the closest one, IU6, that is only 15 km away from IU7, and is 121 m deep. So we argue that even though the model cannot reproduce the occasional halocline seen in IU7, this has no effect on the main analysis of our study, as there is no halocline northward from this and thus possible sub-halocline transports would not be able to propagate northward. We have modified this part in the revised manuscript so that it is more clear what we mean (lines 155–159).

> 8) Lines 306-307: *the statement “Archipelago Sea as a buffer zone between the Baltic Proper and the Bothnian Sea” needs explanation. It cannot be directly deduced from the synchronous monthly mean transports presented in Fig. 7. Regarding spreading of tracers, indeed Miettunen et al. (2020) have shown by integration of Lagrangian transport that “only a small percentage of the particles released in the southern and northern parts of the model area entered the middle and inner archipelagos.” Perhaps the flow speed corresponding to monthly mean transport is so small that water cannot be transported through the all sections during a month. Distance between northern and southern transects is about 85 km, there should be the speed 3.3 cm/s to cover such distance.*

We agree that our statement about the Archipelago Sea being a buffer zone cannot be deduced from the results presented in this manuscript. Thank you for pointing this out. In addition to the work done in this paper, we base the statement on the earlier work done in this area by ourselves and others. We have modified the conclusions accordingly (lines 370–371).

> 9) Lines 307-308: *there is a statement “The transport dynamics in the Archipelago Sea are complicated so that no single transect can be chosen to represent the transport through the whole area.” On the other hand, Fig. 7 shows that monthly transports across the three sections are similar. There seems to be some controversy; please explain in the revised text.*

It is true that the monthly transports across the three transects are similar when looking at the surface layer or the whole water column, and thus the dominant southward net transport is similar everywhere. However, transports in the lower layer are generally towards opposite directions in

north and south. We have edited the discussion part in the revised manuscript to explain this (lines 297–306) and also rephrased the abstract (lines 16–17) the conclusions (lines 371–372) accordingly.

> 10) *Figures 4 and 5 could be combined together as (a) and (b) since their only difference is in the selection of layer: uppermost 5 m vs bottommost 5 m.*

This is a good idea. We will discuss this with the technical editor if the manuscript is accepted for publication.

> 11) *Figures 3 and 6 are very similar and contain repeated information. Perhaps to keep only one figure.*

We decided to keep both figures but modified Fig. 6 to avoid repetition. Fig. 3 is part of validation, showing the differences between the wind forcing and observations. The new version of Fig. 6 now shows the wind roses for the 30-year period as well as for the years 2013–2017 separately, to show how 2014 differs from the other years and from the long-term wind distribution.

> 12) *Net transport in Fig. 8 is the same as already presented in Fig. 7. Please try to avoid duplication.*

We agree that our choice of figures was not the best. Fig. 7 showed the mean transport in the upper and lower layers separately and we included it to enable comparison with the figures in Westerlund et al. (2022). Fig. 8, on the other hand, showed the monthly net transport in the whole water column but included also the southward and northward components of the net transport. To avoid duplication but still show the difference between the upper and lower layers, we decided to change these figures in the revised manuscript. Now the time series plot that was Fig. 7 is removed and the new Figs. 7 and 8 show the monthly net transports as well as the southward and northward components for the upper and lower layers, respectively. Parts of Section 5 are modified/rearranged accordingly (lines 216–239, line 262).

> 13) *Figures 8 and 9 contain the same information, only for the two different transects. By such presentation, comparison of transports is not straightforward. Please consider some other reader-friendly presentation.*

We agree that it was not a reader-friendly presentation. Now that we changed Figs. 7 and 8 in the revised version, we included the panel (a) from the old Fig. 8 to Fig. 9, making the comparison between the two transects easier.

> *I recommend an editorial revision of the MS.*